

Potential impacts of marine fuel regulations on an Arctic stratocumulus case and its radiative response

Luís Filipe Escusa dos Santos^{1,a}, Hannah C. Frostenberg², Alejandro Baró Pérez², Annica M. L. Ekman^{3,4}, Luisa Ickes², and Erik S. Thomson¹

¹Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg, Sweden.

²Department of Space, Earth and Environment, Chalmers University of Technology, Gothenburg, Sweden.

³Department of Meteorology, Stockholm University, Stockholm, Sweden.

⁴Bolin Centre for Climate Research, Stockholm, Sweden.

^anow at: Department of Space, Earth and Environment, Chalmers University of Technology, Gothenburg, Sweden.

Correspondence: Luis F. E. d. Santos (luis.santos@chalmers.se) and Luisa Ickes (luisa.ickes@chalmers.se)

Abstract. Increased surface warming over the Arctic, triggered by increased greenhouse gas concentrations and feedback processes in the climate system, has been causing a steady decline in sea-ice extent and thickness. With the retreating sea-ice, shipping activity will likely increase in the future driven by economic activity and the potential for realizing time and fuel savings from transiting shorter trade routes. Moreover, over the last decade, the global shipping sector has been subject to regulatory changes, that affect the physicochemical properties of exhaust particles. International regulations aiming to reduce SO_x and particulate matter (PM) emissions, mandate ships to burn fuels with reduced sulfur content or alternatively, use wet scrubbing as exhaust after-treatment when using fuels with sulfur contents exceeding regulatory limits. Compliance measures affect the physicochemical properties of exhaust particles and their cloud condensation nuclei (CCN) activity in different ways, with the potential to have both direct and indirect impacts on atmospheric processes such as the formation and lifetime of clouds. Given the relatively pristine Arctic environment, ship exhaust particle emissions could be a large perturbation to natural baseline Arctic aerosol concentrations. Low-level stratiform mixed-phase clouds cover large areas of the Arctic region and play an important role in the regional energy budget. Results from laboratory marine engine measurements, which investigated the impact of fuel sulfur content (FSC) reduction and wet scrubbing on exhaust particle properties, motivate the use of large eddy simulations to further investigate how such particles may influence the micro- and macrophysical properties of a stratiform mixed-phase cloud case observed during the Arctic Summer Cloud Ocean Study campaign. Simulations with diagnostic ice crystal number concentrations revealed that enhancements of ship exhaust particles predominantly affected the liquid-phase properties of the cloud and led to a decrease in liquid surface precipitation, increased cloud albedo and increased longwave surface warming. The magnitude of the impact strongly depended on ship exhaust particle concentration, hygroscopicity, and size where the effect of particle size dominated the impact of hygroscopicity. While low FSC exhaust particles were mostly observed to affect cloud properties at exhaust particle concentrations of 1000 cm⁻³, exhaust wet scrubbing already led to significant changes at concentrations of 100 cm⁻³. Additional simulations with cloud ice water path increased from ≈5.5 g m⁻² to ≈9.3 g m⁻², show more muted responses to ship exhaust perturbations but revealed that exhaust perturbations may even lead to a slight radiative cooling effect depending on the microphysical state of the cloud. The regional impact

of shipping activity on Arctic cloud properties may, therefore, strongly depend on ship fuel type, whether ships utilize wet scrubbers, and ambient thermodynamic conditions that determine prevailing cloud properties.

1 Introduction

Maritime shipping is a significant source of atmospheric pollutants with wide-ranging impacts on human health (Corbett et al., 2007; Jonson et al., 2020; Liu et al., 2016) and the climate system (Lauer et al., 2007; Eyring et al., 2010; Lund et al., 2012, 2020). Quantifying the net impact of ship exhaust emissions on Earth's radiative budget is a challenging task, due to large spatial variability in atmospheric conditions and heterogeneity in air exhaust composition. While ships emit a substantial amount of greenhouse gases, such as CO₂, the remaining constituents can vary substantially with the propulsion system and fuel type used by the individual vessel (Lack et al., 2009; Lack and Corbett, 2012; Lehtoranta et al., 2019). Whereas CO₂ emissions contribute to climate warming, the overall impact of particulate matter and SO₂ exhaust emissions is subject to a much larger uncertainty envelope.

Ship exhaust emissions of primary and secondary particles have been identified to lead to tens of thousands of premature deaths worldwide (Corbett et al., 2007). Regions that are particularly affected by these emissions include coastal areas and port cities, with high population densities, including parts of Europe and East Asia. Motivated by the harmful effects that ship exhaust particles have for human health, the International Maritime Organization (IMO) decided to introduce international marine fuel regulations, which primarily target a reduction of sulfur oxides (SO_x). These regulations mandate that ship operators use marine fuels with fuel sulfur content (FSC) lower than 0.5 wt % effective globally and lower than 0.1 wt % in designated sulfur emission control areas (SECA), or utilize exhaust treatment systems to reduce emissions (IMO, 2008). Since low FSC fuels are generally associated with a higher cost than conventional, high FSC residual fuel oils (UNCTAD, 2022) wet scrubbing systems pose an economically attractive treatment alternative, which allows stakeholders to continue to use marine fuels with FSCs exceeding regulatory limits (IMO, 2008). Wet scrubbers are exhaust after-treatment systems, that utilize mists of seawater or chemically treated freshwater to remove SO_x from ships' exhaust, and thus, prevent the formation of sulfur-containing, secondary aerosol particles (Oikawa et al., 2003; Andreasen and Mayer, 2007). While recent studies demonstrate that utilization of FSC-lean marine fuels generally reduces the amount of particles emitted by ships (Zetterdahl et al., 2016; Kuittinen et al., 2021; Seppälä et al., 2021), the impact of wet scrubbing on particle exhaust emissions is less well understood and subjected to a large variability (Fridell and Salo, 2016; Lehtoranta et al., 2019; Winnes et al., 2020; Yang et al., 2021; Jeong et al., 2023). Moreover, compliance alternatives, such as exhaust after-treatment systems, have been found to affect the physicochemical properties of exhaust particles in different ways, which also have implications for atmospheric processes and the net climate effect of shipping activity. Combustion of low FSC fuels often results in the emission of predominantly hydrophobic soot particles, leading to reduced emissions of cloud condensation nuclei (CCN) compared to higher CCN emissions from conventional, high FSC fuel combustion (Lack et al., 2009; Yu et al., 2020, 2023). In contrast, wet scrubbing has been found to alter the physicochemical properties of the particle emissions (Lieke et al., 2013; Santos et al., 2023, 2024b). This can yield larger fractions of water-soluble content in the exhaust particle phase and a shift in particle size distributions to larger

particles compared to exhaust particle emissions from conventional, high FSC fuel combustion. Combustion particles from wet scrubbing require relatively low supersaturations to be activated into liquid droplets (Santos et al., 2023) which can lead to enhanced CCN number emissions at given supersaturations (Santos et al., 2023, 2024b).

60 Shipping emissions are currently estimated to have a net cooling effect on the climate; higher exhaust particle number concentrations can lead to increased cloud reflectivity which dominates the warming effect of shipping-related CO₂ emissions (Lauer et al., 2007; Eyring et al., 2010; Lund et al., 2012, 2020). Ship tracks are the visible manifestation of ship exhaust perturbations on cloud properties, resulting in persistent, regionally constrained marine stratiform cloud features with increased cloud albedo (Coakley et al., 1987; Hobbs et al., 2000; Possner et al., 2018). The extent of ship tracks depends on the back-
65 ground state of the boundary layer including meteorological parameters, the cloud fraction, and aerosol particle and CCN number concentrations (Coakley et al., 1987; Durkee et al., 2000; Hobbs et al., 2000). Observations along the coast of California have shown that the 0.1 wt % FSC limit, introduced in 2015 in SECAs, led to strong reductions in visible ship track formation (Gryspeerd et al., 2019; Watson-Parris et al., 2022). While ship sulfate emissions are one key driver to ship track formation, FSC reduction policies may still lead to cloud perturbations. These cloud perturbations may be undetectable for some
70 analysis techniques, resulting in an underestimate of shipping-induced radiative forcing (Gryspeerd et al., 2019; Manshausen et al., 2022). With the introduction of the global 0.5 wt % FSC cap in 2020 and associated implications for exhaust particles, radiative cooling induced by ship exhaust emissions may be diminished. Studies investigating the impact of the 2020 0.5 wt % FSC cap have reported lower ship track formation frequencies and highlight the reduction in SO₂ emissions as key drivers for this observation (Gryspeerd et al., 2019; Yuan et al., 2022; Watson-Parris et al., 2022). Therefore, IMO FSC regulations may
75 imply a diminished radiative cooling from shipping emissions. However, the magnitude of diminished cooling may be subject to a systematic underestimate, as ship track visibility is strongly dependent on the clouds' background states (Gryspeerd et al., 2019; Yuan et al., 2022; Watson-Parris et al., 2022).

One region where future shipping activity might lead to a strong climate feedback is the Arctic. The Arctic is experiencing unprecedented amplified surface warming compared to the global average, caused by a complex system of interacting processes
80 within its climate system (Serreze and Francis, 2006; Serreze and Barry, 2011; Rantanen et al., 2022). Low-level mixed-phase clouds play a key role in the Arctic climate system (Morrison et al., 2012). Whereas low-level clouds generally lead to surface cooling, they tend to enhance surface warming in the Arctic throughout most of the year by trapping and re-emitting longwave radiation (Intrieri et al., 2002; Shupe and Intrieri, 2004). Enhanced surface warming in the Arctic promotes ice and snow melting and as a consequence, Arctic sea-ice extent and thickness have been in decline for the past decades (Screen and
85 Simmonds, 2010; Serreze and Barry, 2011). This will likely grant ships easier access to exploration and extraction of natural resources and may enable the use of shorter trading routes through Arctic waterways, deviating from the more conventional and longer routes through the Suez and Panama Canals. The economic feasibility of Arctic shipping routes compared to traditional routes is debated (Lasserre and Pelletier, 2011). Nonetheless, shipping activity and related exhaust emissions are expected to increase significantly within the near future (Corbett et al., 2010; Paxian et al., 2010; Peters et al., 2011). In the Arctic, ambient
90 particle number concentrations are relatively low compared to other regions of the Earth and thus, relatively small absolute increases in aerosol concentrations can substantially impact cloud formation and properties (Mauritsen et al., 2011; Bulatovic

et al., 2021). Ship emissions may therefore become a strong, localized aerosol source that could alter the properties of Arctic clouds and thereby the radiative budget.

Several studies have investigated the potential impacts of increased Arctic shipping activity on Arctic cloud properties (Christensen et al., 2014; Possner et al., 2017; Gilgen et al., 2018; Eirund et al., 2019). Ship aerosol emissions were observed to generate a shift towards the ice phase, reducing precipitation and increasing cloud albedo (Christensen et al., 2014; Possner et al., 2017). Possner et al. (2017) observed a noteworthy rise in liquid water content (LWC) when ship-emitted CCN surpassed 1000 cm^{-3} . However, results were inconclusive in determining whether ship emission-related changes were sufficient to impact Arctic warming rates (Christensen et al., 2014; Possner et al., 2017). Gilgen et al. (2018) modeled significant impacts on Arctic cloud properties from shipping when exaggerated future Arctic ship emission inventories were used, i.e., when Arctic shipping emissions for 2050 were increased by a factor of 10. In contrast, Stephenson et al. (2018) investigated the total climate impact from trans-Arctic shipping and found an increase in total cloud fraction and cloud liquid water path (LWP) due to CCN-enhancements from ship emissions, diminishing Arctic warming rates and exerting cooling rates on the order of 1° C by the end of the 21st century. Eirund et al. (2019) highlight how underlying surfaces influence the properties of mixed-phase clouds and thus, the impact of additional CCN from ship exhaust emissions may be weakened or strengthened, depending on the ice cover.

Ship exhaust emissions also have the potential to exert radiative forcing via direct interactions between emitted particles and solar radiation, or via reduction of the surface albedo due to deposition of light-absorbing black carbon (BC) particles onto snow. A modelling study by Dalsøren et al. (2013) examined a number of direct and indirect processes related to shipping emissions and radiative processes. The study found significant seasonal variability for all processes and that direct sulfate aerosol interactions exert the largest radiative forcing (positive) out of all processes, i.e., a larger forcing than aerosol-cloud interactions. Given IMO's marine fuel policies, the impact of ship-related sulfate contributions may be subject to large uncertainties. In contrast, Gilgen et al. (2018) and Stephenson et al. (2018) found that radiative forcing induced by aerosol-cloud interactions outweighs forcing exerted by direct aerosol-radiation interaction and BC deposition onto snow. Similarly, Browse et al. (2013) and Li et al. (2021) report only minor contributions of BC deposition from shipping activity which would yield insignificant changes in radiative forcing and not contribute to accelerated sea ice loss. While these findings apply for the Arctic in general, surface albedo adjustments due to BC deposition may have stronger local constrained impacts, for example, in the sub-Arctic region (Browse et al., 2013).

The aim of our study is to investigate how ship exhaust particle perturbations influence the microphysical structure of an Arctic mixed-phase cloud and thereby its climate effect. We elaborate on the differences in different ship exhausts based on laboratory results (Santos et al., 2022, 2023, 2024b). In this study, we use large-eddy simulation (LES) to simulate a well-characterized mixed-phase stratocumulus cloud observed during the Arctic Summer Cloud Ocean Study (ASCOS) campaign (Tjernström et al., 2012, 2014). We systematically perturb the aerosol concentrations in the model domain to explore the effect of different types of ship exhausts. Whereas previous studies investigating cloud perturbations caused by ship exhaust emissions, used simplistic representations of physicochemical properties of ship exhaust particles, herein, we utilize detailed exhaust particle information obtained from laboratory marine engine experiments where the impact of FSC reduction and

exhaust wet scrubbing on ship exhaust particle properties was examined (Santos et al., 2022, 2023, 2024b). The model is initially run with an ambient background aerosol concentration only. Subsequent model simulations utilize several potential ship aerosol concentrations with different particle size distributions, densities, and hygroscopicities, mirroring the effects of FSC reduction and wet scrubbing. We evaluate the role of ship aerosol properties in affecting the cloud's LWP and ice water path (IWP), and the concentration of cloud droplets and raindrops. The results are used to calculate changes in surface precipitation, cloud drop effective radius, cloud albedo, and other cloud properties, which have implications for the radiative surface budget. Potential Arctic climate feedbacks from increased shipping activity, in the context of the adaption of different fuel types and propulsion technologies by ships, are discussed.

135 2 Methods

2.1 Laboratory pre-study - Physicochemical properties of ship exhaust aerosol

The experimental results used in this study are based on a series of laboratory experiments that were performed between 2019 and 2022 in Gothenburg, Sweden. More details on the laboratory experiments can be found in Santos et al. (2022, 2023, 2024b). Engine experiments were performed using stationary, marine test-bed diesel engines, fuel types of varying sulfur content, a laboratory wet scrubber, and a range of gas and aerosol instrumentation quantifying physicochemical properties of exhaust particles. For the simulations with the MISU MIT Cloud and Aerosol (MIMICA) LES model, the following parameters are needed as input to describe the aerosol perturbation: particle size distributions, particle effective densities, and hygroscopicities. Particle size distributions were measured using Scanning Mobility Particle Sizers (SMPS). We describe average particle size distributions using the count median diameter (CMD) and the geometric standard deviation (σ_g). Effective particle densities (ρ_{eff}) were determined by coupled SMPS and Aerodynamic Aerosol Classifier (AAC) measurements and calculated following Tavakoli and Olfert (2014) and Santos et al. (2022, 2024b). Exhaust particle hygroscopicities (κ) were determined from size-selected CCN measurements, using a CCN counter (CCNc; CCN-100, Droplet Measurement Technologies (Roberts and Nenes, 2005)) and parameterizations by Petters and Kreidenweis (2007).

From Santos et al. (2022, 2023) we use results from measurements with high FSC fuel (HiS; HGO in the respective studies), one low FSC fuel (LoS; MGO in the respective studies), and seawater scrubbing experiments, performed in combination with HiS fuel combustion (WS; SWS in the respective studies). For the LES experiments, the results were simplified by assigning identical size distributions to HiS and LoS which did not display substantial differences during the respective measurement campaigns. The particle size distributions used in the model are shown in Fig. 1 a. Other results, which were used as input parameters, such as average case-dependent ρ_{eff} and κ values are listed in Table 1 and discussed further in Sect. 2.3.




Figure 1. (a) Particle size distributions of high (HiS) and low sulfur content fuel (LoS), and wet scrubbed (WS) exhaust particles from Santos et al. (2022, 2023) and (b) the sulfate particle modes of HiS fuel (HiS_sul) and scrubbed exhaust particles (WS_sul) from Santos et al. (2024b). The dashed lines in panel (a) represent the two individual modes of the bimodal WS case. The data shown in the figure represent size distributions measured during the respective measurement campaigns that have been averaged and simplified to be parameterized within MIMICA.

155 Engine experiments summarized in Santos et al. (2024b) utilized a different engine with higher power output and fuels with different properties compared to Santos et al. (2022, 2023) and therefore, resulted in different emission characteristics. From Santos et al. (2024b), only results obtained for high FSC fuel combustion and exhaust wet scrubbing at engine load points of 50% were implemented in this study. Both cases resulted in emissions of bimodal size distributions consisting of a dominant, hygroscopic sulfate mode and a smaller, relatively hydrophobic soot mode. Here, corresponding bimodal particle
 160 size distributions were simplified to the respective unimodal, hygroscopic sulfate modes. The dominant sulfate modes (Fig.1 b) with their respective averaged ρ_{eff} and κ values are summarized in Table 1. The high FSC case is referred to as HiS_sul and the seawater wet scrubbing case as WS_sul.

The main findings from Santos et al. (2022, 2023) showed that FSC reduction and exhaust wet scrubbing led to substantial impacts on particulate emissions from ship engines. A switch to marine fuels with reduced FSC did not significantly affect
 165 particle size distributions and total number emissions but decreased the exhaust particles' ρ_{eff} and κ values. On the other hand, wet scrubbing was found to lead to the formation of a dominant particle mode around 20 nm, and to increased ρ_{eff} and κ values, due to changes in the chemical mixing state. Similarly, Santos et al. (2024b) investigated the impact of different fuel types and seawater exhaust wet scrubbing on exhaust particle properties, but used a different test-bed engine, with higher total power output. One key difference compared to Santos et al. (2022, 2023) was, that the combustion of non-compliant, high FSC fuel
 170 resulted in bimodal particle size distributions with a dominant sulfate mode around 20 nm. When the high FSC fuel exhaust was scrubbed, the sulfate mode was shifted towards larger sizes, likely due to the coagulation of particles inside the scrubber.

2.2 MIMICA model description and case setup

LES experiments were conducted with the MIMICA model. This model was originally designed to study high-latitude mixed-phase clouds and has been thoroughly documented and evaluated against observations (see e.g., Savre et al. (2015); Stevens

Table 1. Properties of marine background (BG) and ship aerosol used as model input parameters, including the count median diameter (CMD) and geometric standard deviation (σ_g) of the size distributions, particle density (ρ_{eff}) and aerosol hygroscopicity (κ). HiS, LoS, and WS data were obtained from experiments outlined in Santos et al. (2022, 2023) and refer to combustion of high (HiS) and low FSC fuels (LoS), and wet scrubbed HiS exhaust (WS). HiS_sul and WS_sul represent the sulfate particle modes measured for high FSC fuel and wet scrubbed exhaust in Santos et al. (2024b), respectively. The WS case is composed of a bimodal distribution, hence, the two separate aerosol modes are listed in the table. For each ship exhaust sensitivity test, two sets of simulations with either low or high ship aerosol number concentrations were performed. Corresponding simulations with low and high concentrations (N_p) are additionally labeled with *_lo* and *_hi* respectively.

Case	CMD [nm]	σ_g	ρ_{eff} [g cm ⁻³]	κ	N_p [cm ⁻³]
BG (Ait)	32	1.1	2.18	1	30
BG (Acc)	93	1.5	2.18	1	30
LoS	45	1.6	0.91	0.04	100/1000
HiS	45	1.6	1.02	0.11	100/1000
WS (Mode 1)	22	1.2	1.18	0.22	131.3/1313
WS (Mode 2)	64	1.3	1.09	0.16	36.7/367
HiS_sul	18	1.15	1.6	0.64	100/1000
WS_sul	39	1.22	1.6	0.64	100/1000

175 et al. (2018); Bulatovic et al. (2023)). Herein, only a brief description of the model is provided. For more detailed information, see Savre et al. (2014).

MIMICA solves a set of anelastic, non-hydrostatic governing equations and uses a two-moment bulk microphysics scheme to predict mass mixing ratios (Q) and number densities (N) of five hydrometeor classes, including cloud droplets, raindrops, ice crystals, graupel, and snow. Growth of liquid-phase hydrometeors via auto-conversion, self-collection, and collision-
180 coalescence are treated following Seifert and Beheng (2001) and Seifert and Beheng (2006). Interactions between liquid- and ice-phase hydrometeors are treated according to the two-moment bulk microphysics by Wang and Chang (1993). Hygroscopic growth of aerosol particles and activation into cloud droplets is calculated according to κ -Köhler theory (Petters and Kreidenweis, 2007). While MIMICA does include options for heterogeneous ice nucleation, here diagnostic ice crystal number concentrations (N_i) are utilized as in Ovchinnikov et al. (2011, 2014). This means that in grid cells where the temperature
185 (T) is less than 0° C and sufficient supercooled cloud water is present ($Q_c \geq 2 \times 10^{-7}$ g m⁻³), N_i is relaxed towards a pre-determined, constant value. As a default, this value for N_i was set to 200 m⁻³ based on the control simulations in Stevens et al. (2018). The decision to use a constant diagnostic N_i instead of an interactive heterogeneous ice nucleation scheme was motivated by findings showing that typical engine exhaust particles, such as BC, are inefficient ice nucleators in the immersion freezing regime (Mahrt et al., 2018; Kanji et al., 2020). In addition, Santos et al. (2024b) found no significant differences in
190 the ice nucleation behavior of exhaust particles emerging from low and high FSC fuel combustion, and exhaust wet scrubbing.

The simulated stratocumulus case is based on observations made during ASCOS on 31.08.2008 at approximately 87°N, 11°W (see Appendix A with the detailed vertical profiles used to initialize the model; Tjernström et al., 2012, 2014). Note that meteorological conditions from ASCOS were only used to initialize the model. Potential temperature and total water mixing ratio are prognostic variables influenced by sources and sinks in the model (e.g., radiation, microphysical phase changes, and precipitation), but do not necessarily represent the temporal evolution of the real atmospheric state. The case study represents a stable mixed-phase cloud that has previously been investigated using MIMICA (Igel et al., 2017; Stevens et al., 2018; Christiansen et al., 2020; Sotiropoulou et al., 2021; Frostenberg et al., 2023). For more details on the setup of MIMICA please see the aforementioned references, and for more extensive information on the ASCOS campaign and the experimental results see Tjernström et al. (2012, 2014).

The MIMICA 3-D domain consists of $96 \times 96 \times 128$ grid cells with periodic boundaries. The horizontal resolution is uniform $dx = dy = 62.5$ m while the grid spacing in the vertical z -direction is variable $7.5 \text{ m} \leq dz \leq 25 \text{ m}$. Higher vertical resolution is applied to grid cells near the surface and within the cloud layer, whereas a sinusoidal function is used to define the vertical spacing of grid cells at other altitudes. The total domain is $6 \text{ km} \times 6 \text{ km}$ in the horizontal direction and 1.7 km in the vertical direction. All simulations were run for 16 h. The first 4 h are considered as spin-up and are thus excluded from the presented results.

The radiation solver used in this study is based on Fu and Liou (1992). It is important to note that while radiation is affected by cloud hydrometeors it is not affected by aerosols. Surface temperature and pressure have prescribed values of 269.8 K and 1026.3 hPa, respectively. The surface albedo is set to 0.844 and the surface roughness to 0.0004 m. Sensible and latent heat fluxes at the surface are both set to 0 W m^{-2} based on the small values reported in Tjernström et al. (2014). A large-scale divergence of $1.5 \times 10^{-6} \text{ s}^{-1}$ is imposed over the whole domain. Large-scale advection is turned off in the model.

Additional simulations of no_ship, HiS_sul, and WS_sul were performed with N_i increased from 200 to 600 m^{-3} . The aim of these additional simulations was to investigate the susceptibility of a thinner mixed-phase cloud, i.e., with reduced LWP and cloud depth, towards ship exhaust particle perturbations. A maximum value of $N_i = 600 \text{ m}^{-3}$ as a large enough reduction in LWP was induced to perform aforementioned sensitivity tests and simultaneously, simulated LWP and IWP values agree well with observational data (see Sect. 3.4). Additional testing revealed that further increases in N_i would lead to dissipation of the cloud. Associated model runs are named as previous model runs but with an appended _ni600, e.g., no_ship_ni600.

2.3 Aerosol implementation in MIMICA

Aerosol particles in MIMICA were represented as aerosol modes that follow lognormal distributions, described by a CMD and σ_g . To each aerosol mode, values of the aerosol effective density ρ_{eff} , the aerosol hygroscopicity expressed via κ , and the aerosol number concentrations (N_p) were assigned. For all simulations, aerosol number concentrations and properties were set to be uniform and constant in time over the entire 3-D domain. Aerosol particles can be activated into cloud droplets according to κ -Köhler theory but are modeled without additional sources and sinks during the simulations. Several aerosol modes can co-exist. In our simulations, we describe the total aerosol by natural background aerosol modes and ship exhaust aerosol modes (for the cases with additional ship exhaust particles).

225 Natural background aerosol (BG) concentrations were present in all model runs (Table 1). These BG aerosols were assumed to have hygroscopicity values in agreement with marine seaspray ($\kappa = 1$) and were included in both the Aitken (Ait) and accumulation mode (Acc). The N_p of the BG aerosol was chosen to be 30 cm^{-3} in both modes based on aerosol measurements during the ASCOS expedition (Kupiszewski et al., 2013). In the baseline simulation (referred to as no_ship), only BG Ait and Acc mode aerosol were present.

230 For the sensitivity experiments, different aerosol concentrations and types were added to represent different ship exhaust perturbations (HiS, LoS and WS from Santos et al. (2022, 2023), and HiS_sul and WS_sul from Santos et al. (2024b)). Ship aerosol properties are summarized in Table 1 and Sect. 2.1. For each case, ship exhaust perturbation experiments were performed at two concentration levels, $N_{p,\text{ship}} = 100 \text{ cm}^{-3}$ (labeled with _lo) and 1000 cm^{-3} (labeled with _hi), respectively. An exception is the WS case where the concentration levels are increased by a factor of ≈ 1.7 following the increase in particle
 235 number concentration that has been observed in the experiments when using the wet scrubber. This increase in $N_{p,\text{ship}}$ was accounted for in the two particle modes comprising the WS case. The same _lo and _hi labeling, signifying the low and high concentration simulations, was used for WS cases.

2.4 Calculations of the cloud drop effective radius and cloud albedo

To examine the difference in cloud radiative properties between the simulations, we calculate the effective droplet radius (r_e),
 240 cloud optical depth (τ), and cloud albedo (α) from the model output. To calculate the effective cloud droplet radius r_v we use the relationship as suggested by Freud and Rosenfeld (2012), who found that r_e is on average a factor 1.08 larger than the volume mean cloud droplet radius r_v ,

$$r_e \approx 1.08 r_v . \quad (1)$$

The volume mean cloud droplet radius r_v is defined as,

$$245 \quad r_v = \left(\frac{3}{4} \frac{Q_c}{\pi \rho_w N_c} \right)^{1/3} , \quad (2)$$

where Q_c is the cloud liquid water content, ρ_w is the density of water (1000 kg m^{-3}) and N_c is the cloud droplet number concentration. The cloud's optical depth can be approximated by

$$\tau = \frac{3 \text{ LWP}}{2 r_e \rho_w} , \quad (3)$$

where LWP is the liquid water path, i.e., the vertically integrated amount of liquid cloud water, in kg m^{-2} (Stephens, 1978).

250 The cloud albedo can be approximated with

$$\alpha = \frac{(1-g)\tau}{1+(1-g)\tau} , \quad (4)$$

where g is the scattering asymmetry factor, i.e., the average value of the cosine of the scattering angle, and equals 0.85 for the scattering of solar radiation by clouds (Meador and Weaver, 1980).

3 Results

255 3.1 Influence of ship aerosol on LWP and IWP

In Fig. 2, the time evolution of the domain-averaged LWP and IWP are shown for all the simulations in comparison to the observation from the ASCOS campaign. Note that ASCOS observations used to initialize the model do not change with time during the remaining simulation period. In all simulations, MIMICA simulates an LWP that exceeds the 75th percentile of the observations (Fig. 2 a - b) which was also observed by Bulatovic et al. (2021), who used MIMICA to simulate the same
260 ASCOS case. MIMICA was previously reported to generate greater LWP when prescribed instead of interactive aerosol particle concentrations are used (Stevens et al., 2018). Additional sensitivity tests with reduced LWP are discussed in Sect. 3.4. The addition of ship aerosol tends to increase the LWP of the cloud compared to the reference no_ship case (Fig. 2 a - b; Table 2). This effect is found to be dependent on the ship exhaust aerosol concentrations and the hygroscopicity of the ship exhaust aerosol. The LWP increase is most pronounced for sensitivity tests with high ship aerosol concentrations (HiS_hi, WS_hi, and
265 WS_sul_hi), where LWP increases by up to $\approx 13\%$. The increase is less pronounced for both LoS cases due to the hydrophobic nature of the added particles ($\kappa = 0.04$). Despite having comparatively large κ -values, both HiS_sul cases do not yield any substantial increase in LWP, suggesting that the ship exhaust aerosol are too small (CMD = 18 nm) to induce a pronounced effect. The identified LWP response for the mixed-phase cloud perturbed by ship exhaust agrees with Possner et al. (2017), who reported substantial increases in LWP when ship-related CCN concentrations exceeded 1000 cm^{-3} in their simulations.

270 The simulated IWP is close to the 25th percentile of the observations for all simulations (Fig. 2 c - d). In contrast to results for the LWP, additional ship exhaust particles have no substantial effect on the modeled IWP which is due to the implementation of diagnostic N_i , meaning ship aerosol can not directly impact N_i . However, in our simulations, ship exhaust aerosol can affect the properties of precipitating ice-phase hydrometeors (graupel and snow) by influencing the accretion of hydrometeors and the availability of water vapor. Sporadic spikes in the temporal evolution of IWP are in all cases caused by increased graupel
275 formation rates at the expense of raindrops. Similar features in IWP evolution are reported by Bulatovic et al. (2021) who used MIMICA with a similar setup. Small increases in IWP for some simulations are mainly caused by an increasing graupel number (N_g) and mass concentration (Q_g ; Fig. D1).

3.2 Impact of ship aerosol on hydrometeors

The effect of ship exhaust aerosol perturbations on the temporal evolution of cloud hydrometeors and cloud depth is investigated
280 by examining horizontally averaged number and mass concentrations of cloud droplets (N_c and Q_c) and raindrops (N_r and Q_r). Corresponding contour plots for the reference simulation (no_ship) and all sensitivity simulations with high ship aerosol concentrations (_hi; 1000 cm^{-3} or 1680 cm^{-3} (WS)) are shown in Fig. 3. In addition, each subplot features the horizontally averaged cloud bottom and cloud top height, depicted by the black dashed lines.

In general, impacts of ship aerosol particles on liquid-phase hydrometeors are mostly observed in ship aerosol simulations
285 with the highest $N_{p,\text{ship}}$ in accordance with the results presented in Sect. 3.1. Ice-phase hydrometeors have been excluded from Fig. 3, as changes in ice crystal concentrations remain more or less unaffected due to the prescribed ice parameterization scheme




Figure 2. Time evolution of the simulated domain-averaged (a and b) liquid water path (LWP) and (c and d) ice water path (IWP). `no_ship` refers to the reference case with background aerosol only. HiS, LoS, and WS represent ship aerosol from measurements of high and low sulfur content fuels and wet scrubbing respectively (Santos et al., 2022, 2023). The HiS_sul and WS_sul cases represent sulfate particle modes of high FSC fuel combustion and exhaust gas wet scrubbing from Santos et al. (2024b). The label additions `_lo` and `_hi` signify the ship aerosol concentrations used in the individual model runs. Significant differences between ship exhaust cases and `no_ship` were assessed using two-sided t tests at a confidence level of 95%. Model runs with significant differences are marked with star icons in inset legends. For the statistical tests the last 4 simulation hours were used. The blue shaded area refers to the retrieved LWP and IWP from microwave radiometer measurements (median over the observation period; the corresponding dashed lines are the 25th/75th percentiles) during the ASCOS campaign (Tjernström et al., 2012, 2014) which were used to initialize the model. The first four hours are considered a spin-up period of the model and are removed from the figures.

used in this study (see Sect. 2.2), but can be found in Appendix B. Whereas cloud droplet numbers are evenly distributed within the cloud layer, Q_c is highest near the cloud top, which is typical for stratocumulus clouds with near-adiabatic conditions. The distribution of N_r displays a more dynamic behavior and reaches its maximum after around 8 hours of simulation. In contrast to cloud droplets, Q_r is concentrated towards the lower regions of the cloud. All simulations show similar cloud depths and evolutions (Fig. 3). After 4 h (spin-up) the clouds have a depth of around 500 m. The cloud depth increases in all simulations and ranges between 535 m (`no_ship`) and 570 m (`WS_sul_hi`) at the end of the simulations due to steady increases in cloud top height.

With the addition of ship aerosol, more aerosol particles are activated into cloud droplets as can be seen from increased N_c values for `LoS_hi`, `HiS_hi`, `WS_hi`, `HiS_sul_hi` and `WS_hi` (Fig. 3 and Fig. 4). The largest increase is observed for `WS_hi` and `WS_sul_hi`, where the vertically integrated N_c averaged over the last 4 simulation hours increases by $\approx 57\%$. Note that Q_c is

Table 2. Overview of mean LWP, IWP, surface precipitation rates, cloud effective radius (r_e), cloud albedo (α), and net long- (Net LW) and shortwave radiation at the surface (Net SW) averaged over the last four hours of simulation time. Statistical significance, determined by performing two-side t-tests, is highlighted by asterisks. The label additions *_lo* and *_hi* signify the ship aerosol concentrations used in the individual model runs.

Case	LWP [g m ⁻²]	IWP [g m ⁻²]	Surface precip. (Total) [mm d ⁻¹]	Surface precip. (Rain) [mm d ⁻¹]	r_e [μ m]	α	Net LW [W m ⁻²]	Net SW [W m ⁻²]
no_ship	139.3	5.5	0.32	0.06	16.56	0.65	-14.6	6.5
LoS_lo	140.7*	5.3	0.31	0.05	16.50	0.65*	-14.5	6.5
LoS_hi	143.6*	5.5	0.30	0.04*	16.16*	0.66*	-14.4*	6.5
HiS_lo	142.5*	5.1	0.31	0.06	16.41*	0.66*	-14.5	6.5
HiS_hi	152.8*	5.6	0.28*	0.03*	15.41*	0.68*	-14.2*	6.4
WS_lo	145.4*	5.1	0.30	0.06	16.22*	0.66*	-14.4*	6.5
WS_hi	158.6*	5.9	0.27*	0.02*	14.68*	0.70*	-13.9*	6.3
HiS_sul_lo	141.0*	5.0	0.31	0.06*	16.57	0.65*	-14.6	6.5
HiS_sul_hi	139.8	5.4	0.31	0.06	16.52	0.65	-14.6	6.5
WS_sul_lo	144.6*	5.4	0.30	0.05*	16.10*	0.67*	-14.4*	6.5
WS_sul_hi	158.0*	6.2*	0.29	0.02*	14.69*	0.70*	-14.0*	6.3
no_ship_ni600	88.6	9.3	0.35	0.00	15.51	0.57	-17.6	6.9
HiS_sul_lo_ni600	86.6*	9.4	0.36	0.00	15.47*	0.57*	-17.9*	6.9
HiS_sul_hi_ni600	85.8*	9.4	0.35	0.00	15.46*	0.57*	-18.0*	6.9
WS_sul_lo_ni600	88.0*	9.2	0.34	0.00	14.94*	0.58*	-17.9*	6.9
WS_sul_hi_ni600	86.8*	9.6	0.35	0.00	13.59*	0.60*	-17.9*	6.9

almost unaffected by the added ship exhaust aerosol due to the low precipitation rates in the no_ship case. Despite the relatively large κ value of HiS_sul_hi ship exhaust particles ($\kappa = 0.64$), N_c and N_r are not strongly affected, even when ship exhaust aerosol concentrations are set to $N_p = 1000 \text{ cm}^{-3}$. This implies that HiS_sul exhaust particles were too small (CMD = 18 nm) to act as CCN. The observed increase in N_c is also observed for LoS_hi, suggesting that additional aerosol particles of low hygroscopicity ($\kappa = 0.04$) can impact cloud properties, given their CMD is sufficiently large. Differences in modeled N_c between HiS_sul_hi and LoS_hi ship exhaust aerosol imply that the size of aerosol particles plays a more dominant role in inducing changes in cloud properties than particle hygroscopicity. This observation agrees with Christiansen et al. (2020), who found microphysical cloud properties were not affected by aerosol particles' hygroscopicities if accumulation mode particles were present in the model domain (Christiansen et al., 2020).

Vertical profiles of N_c , Q_c , N_r , and Q_r averaged over the last four simulation hours reveal a more detailed picture of how ship perturbations affect concentrations of cloud droplets and raindrops (Fig.4). The sensitivity tests that show substantial




Figure 3. Temporal evolution of horizontal domain averaged cloud droplet number concentrations (N_c ; $[N_c] = \text{m}^{-3}$), cloud droplet mixing ratios (Q_c ; $[Q_c] = \text{g m}^{-3}$), raindrop number concentrations (N_r) and raindrop mixing ratios (Q_r) simulated for the reference case (no_ship) and the high ship aerosol concentration cases LoS_hi, HiS_hi, WS_hi, HiS_sul_hi and WS_sul_hi. The black dashed lines represent case-specific, horizontally averaged cloud bottom and cloud top heights. The spin-up period (0 to 4 h) is removed from all figures.

increases in N_c also show reduced raindrop formation in the cloud (Fig. 3 and Fig. 4). Whereas no_ship produces substantial amounts of raindrops near the cloud top after about 6 to 7 h of simulation, ship cases with high exhaust particle concentrations lead to general reductions in N_r and Q_r by up to 58% and 63% respectively (Fig. 3 and Fig. 4). The magnitude of this response

310

is dependent on the CMD and κ , where the CMD effect dominates the cloud response. The strongest reduction in N_r and Q_r is observed for WS_hi and WS_sul_hi, where both quantities are reduced by about 52 to 58% (N_r) and 56 to 63% (Q_r) compared to no_ship. Results for raindrop formation coincide with a general reduction in r_e for relevant ship exhaust cases. For both 315 WS_hi and WS_sul_hi, r_e is reduced from 16.56 μm (no_ship) to $\approx 14.68 \mu\text{m}$ (Fig. 5 and Table 2). A reduction in r_e indicates reduced self-collection (autoconversion) and coalescence/accretion of cloud droplets by raindrops or other hydrometeors.




Figure 4. Vertical profiles of horizontally averaged (a and e) N_c , (b and f) Q_c , (c and g) N_r and (d and h) Q_r averaged over the last four simulation hours. The light blue, dashed line represents the average cloud bottom and top height calculated for the reference case (no_ship). HiS, LoS, and WS represent ship aerosol from measurements of high and low sulfur content fuels and wet scrubbing respectively (Santos et al., 2022, 2023). The HiS_sul and WS_sul cases represent sulfate particle modes of high FSC fuel combustion and exhaust gas wet scrubbing from Santos et al. (2024b). The label additions *_lo* and *_hi* signify the ship aerosol concentrations used in the individual model runs. Significant differences between ship exhaust cases and no_ship were assessed using two-sided t tests at a confidence level of 95%. Model runs with significant differences are marked with star icons in inset legends.

3.3 Impact of ship aerosol on surface precipitation and cloud radiative properties

The sensitivity tests that led to reductions in N_r and Q_r are not found to impact total surface precipitation rates compared to no_ship (Fig. 5 a - b). The majority of surface precipitation is dominated by graupel which is not found to be affected by the addition of ship exhaust aerosol. Liquid rain typically constitutes less than 5% of the total surface precipitation (Table 2). Despite the relatively low absolute rates, rain surface precipitation rates are found to be reduced with additional ship exhaust aerosol, agreeing with tendencies in ship exhaust cases to produce smaller r_e (Fig. 5 c - d and Table 2). Changes in surface precipitation rates may significantly change with more realistic ice formation parametrizations. As a result, we cannot exclude whether emissions associated with shipping activity may extend cloud lifetimes due to potential reductions in total precipitation rates (Albrecht, 1989).

In order to estimate the potential climatic impact of increased Arctic shipping activity, α , net short- (SW) and longwave (LW) radiative fluxes at the surface are characterized (Fig. 5 e - f and Table 2). Net radiative fluxes are calculated by subtracting net upwelling fluxes from net downwelling fluxes, hence, a negative value implies net outgoing radiation. At high latitudes, LW radiation generally has a larger influence on the surface energy budget compared to SW radiation, as solar radiation is limited outside the summer months. However, since the ASCOS case used in this study is based on observations from August, the net SW at the surface is also investigated.

With the exception of HiS_sul_hi, all ship sensitivity simulations tend to significantly increase α compared to no_ship (Fig. 5 e - f). This observation agrees with results shown in Fig. 4 and Fig. 5 c - d where tendencies towards generating larger N_c and reduced r_e values are shown. The largest increase in α is observed for WS_hi and WS_sul_hi where cloud albedo increases from 0.65 to 0.70. The changes in LWP and r_e induced by ship aerosol perturbations are also seen in the LW net radiative fluxes at the surface (Fig. 5 g - h).

Net LW fluxes at the surface are negative, meaning the net radiative LW flux is upwelling and therefore, cooling the surface. After about 5 h of simulation, net LW surface fluxes reach values of $\approx -13 \text{ W m}^{-2}$ and eventually decrease to $\approx -15 \text{ W m}^{-2}$ (no_ship) and $\approx -14.3 \text{ W m}^{-2}$ (WS_hi) towards the end of the simulation. Ship cases, which are found to lead to the largest increase in LWP (His_hi, WS_hi, and WS_sul_hi; Fig. 2 a - b), also reduce net LW cooling at the surface compared to no_ship, i.e., net LW becomes less negative. Our results suggest that ship exhaust perturbations may lead to diminished surface LW radiative cooling and could therefore lead to enhanced surface warming, that is if the concentrations and size of the associated exhaust particle size distributions are sufficiently large to act as CCN. Similar relationships between increased LWP and a reduced LW radiative cooling were also noted by Christiansen et al. (2020).

The net SW radiation is positive in all simulations, meaning the net flux is downwelling. In all simulations, the net SW fluxes initially increase until 6 h into the respective simulations where a maximum of around 14 W m^{-2} is reached. By the end of the simulations, net SW decreases to $\approx 5 \text{ W m}^{-2}$. The temporal trends in LW and SW radiation both coincide with the solar angle. The results indicate that WS cases tend to slightly decrease the net SW (Table 2 and Figure C1), yet, none of the ship sensitivity tests are found to significantly impact net SW fluxes at the surface, despite associated increases in α (Fig. 5 i - j). Changes in cloud properties induced by ship exhaust perturbations are expected to only lead to small changes in SW surface

fluxes, due to the reduced solar fluxes based on the geographical location, and the comparatively large LWP, which leads to a substantial extinction of incoming SW radiation. Relatively small changes in α are therefore only expected to lead to minor changes in SW surface fluxes.




Figure 5. Time evolution of the simulated domain-averaged (a and b) surface precipitation, (c and d) r_e , (e and f) α , (g and h) net longwave radiation at the surface (Net LW) and (i and j) net shortwave radiation at the surface (Net SW) for the set of simulations. Net radiative fluxes are calculated by subtracting the upwelling radiative flux from the downwelling flux (e.g., $LW_{\text{down}} - LW_{\text{up}}$), hence, a negative value implies net outgoing radiation. no_ship refers to the reference case with background aerosol only. HiS, LoS, and WS represent ship aerosol from measurements of high and low sulfur content fuels and wet scrubbing respectively (Santos et al., 2022, 2023). The HiS_sul and WS_sul cases represent sulfate particle modes of high FSC fuel combustion and exhaust gas wet scrubbing from Santos et al. (2024b). The label additions *_lo* and *_hi* signify the ship aerosol concentrations used in the individual model runs. Significant differences between ship exhaust cases and no_ship were assessed using two-sided t tests at a confidence level of 95%. Model runs with significant differences are marked with star icons in right-hand side legends. The last 4 simulation hours were used to perform statistical tests.

3.4 Sensitivity to different levels of N_i

355 Additional simulations of no_ship, HiS_sul, and WS_sul were performed with N_i increased from 200 to 600 m^{-3} . The aim of these additional simulations was to investigate the susceptibility of a thinner mixed-phase cloud, i.e., with reduced LWP and cloud depth, towards ship exhaust particle perturbations. When N_i is increased to 600 m^{-3} , the LWP is reduced by $\approx 45\%$ compared to no_ship and is close to the 75th percentile of the observational data (Fig. 6 a). IWP is increased by 3.8 g m^{-2} ($\approx 69\%$; comparison between no_ship and no_ship_ni600) and is also within the observed range (Fig. 6 b). With an increased

360 N_i , ship exhaust particles do not lead to as strong perturbations in LWP ($|\Delta\text{LWP}| < 3.2\%$) compared to sensitivity tests performed with $N_i = 200 \text{ m}^{-3}$. In fact, the ship exhaust sensitivity simulations display a tendency towards reducing the LWP compared to no_ship_ni600 which is in contrast to the model runs with $N_i = 200 \text{ m}^{-3}$ shown in Fig. 2. Such a muted response in LWP adjustments from additional ship exhaust particles was also reported by Possner et al. (2017) who found a suppressed LWP response from ship-related CCN emissions when the number of ice nucleating particles was increased. The IWP, on

365 the other hand, is increased for no_ship_ni600, HiS_sul_hi_ni600 and WS_sul_hi_ni600 compared to no_ship (Fig. 6 b) as all ice-phase hydrometeors increase substantially in number and mass concentrations (Fig. 7 c - e and h - j). As with the first set of simulations, ship exhaust perturbations do not significantly impact the IWP compared to the respective reference case, no_ship_ni600 which, as previously discussed, is mainly due to the implementation of diagnostic N_i .




Figure 6. Time evolution of the simulated domain-averaged (a) liquid water path (LWP) and (b) ice water path (IWP). Figures show results for model runs where N_i was increased from 200 to 600 m^{-3} . Model runs with increased N_i are labeled with _ni600. The simulations are compared to the no_ship case where $N_i = 200 \text{ m}^{-3}$. The blue shaded area refers to the retrieved LWP and IWP from microwave radiometer measurements (median over the observation period; the corresponding dashed lines are the 25th/75th percentiles) during the ASCOS campaign (Tjernström et al., 2012, 2014). Only ship exhaust sensitivity cases with high concentrations (_hi) are shown in the figure. Significant differences between ship exhaust cases and no_ship_ni600 were assessed using two-sided t tests at a confidence level of 95%. Model runs with significant differences are marked with star icons in inset legends. The last 4 simulation hours were used to perform statistical tests. The first 4 hours are considered a spin-up period of the model and are removed from the figures.

Fig. 7 shows the impact on cloud depth and all hydrometeor classes for the _ni600 simulations compared to no_ship

370 (with $N_i = 200 \text{ m}^{-3}$). The cloud depth is decreased, mostly due to adjustments in cloud bottom height, which increases

by ≈ 80 m (Fig. 7). Taking into account the increase in cloud bottom height, N_c values are either similar (no_ship_ni600, HiS_sul_hi_ni600) or increased (WS_sul_hi_ni600) compared to the respective no_ship values (Fig. 7 a). Simultaneously, Q_c values of all _ni600 cases are reduced compared to no_ship, suggesting smaller cloud droplets in the cloud (Fig. 7 f). Raindrop number N_r and mass concentrations Q_r are significantly reduced compared to no_ship (Fig. 7 b - g), resulting from reduced auto-conversion of cloud droplets to raindrops and droplet coalescence efficiency, as well as enhanced scavenging of raindrops by graupel and snow, which display substantial concentration increases in all _ni600 runs (Fig. 7 d - e and Fig. 7 i - j).




Figure 7. Vertical profiles of horizontally averaged (a) N_c , (b) N_r , (c) N_i , (d) N_g , (e) N_s , (f) Q_c , (g) Q_r , (h) Q_i , (i) Q_g and (j) Q_s averaged over the last four simulation hours for no_ship_ni600, HiS_sul_hi_ni600, WS_sul_hi_ni600 and no_ship. The light blue, dashed line represents the average cloud bottom and top height calculated for no_ship_ni600 (green, dotted line for no_ship respectively). The HiS_sul and WS_sul cases represent sulfate particle modes of high FSC fuel combustion and exhaust gas wet scrubbing. Significant differences between ship exhaust cases and no_ship_ni600 were assessed using two-sided t tests at a confidence level of 95%. Model runs with significant differences are marked with star icons in inset legends.

Surface precipitation rates are slightly increased in the _ni600 simulations compared to no_ship ($< 10\%$; Table 2 and Fig. 8 a) which is due to enhancements in graupel formation (Fig. 7 d and i). Liquid-phase surface precipitation rates are small and negligible for all _ni600 simulations. Average cloud drop effective radii r_e values are reduced by $\approx 1 \mu\text{m}$ compared to the respective simulations with $N_i = 200 \text{ m}^{-3}$ agreeing with observed reductions in LWP (Fig. 6 a) and Q_c (Fig. 7 f). The reductions in LWP and α are both reflected in the net radiative fluxes at the surface. Net LW radiative fluxes at the surface are reduced by at least

2 W m^{-2} compared to no_ship, meaning that reductions in LWP and cloud depth lead to reduced re-emission of LW radiation to the surface (Fig. 8 d). Respective _ni600 ship exhaust cases tend to reduce net LW at the surface further, suggesting that in these instances ship exhaust perturbations would yield a slight net cooling effect. This is in contrast to the results shown in
385 Sect. 3.3 where ship exhaust perturbations reduced net outgoing LW radiation at the surface. Relative reductions in α for _ni600 cases are reflected in net SW fluxes at the surface. In comparison to no_ship, net SW is on average increased by $\approx 0.5 \text{ W m}^{-2}$ compared to no_ship (Fig. 8 e and Table 2). Nevertheless, additional ship exhaust particles do not significantly alter net SW radiative surface fluxes (Figure C2).

In summary, the results shown in this section demonstrate that the impact of ship exhaust particles on clouds and the surface
390 radiative budget does not only depend on the ship exhaust particles themselves. The sensitivity is also strongly dependent on the background state of the atmosphere and the background cloud properties, such as cloud thickness. Interestingly, ship exhaust perturbations can have opposite effects on certain cloud parameters and net radiative fluxes. The most striking difference is that ship emissions tend to decrease the LWP and enhance net outgoing LW radiative surface fluxes when $N_i = 600 \text{ m}^{-3}$, whereas the opposite is obtained when $N_i = 200 \text{ m}^{-3}$.




Figure 8. Time evolution of the simulated domain-averaged (a) surface precipitation rates, (b) r_e , (c) α , (d) net longwave radiation at the surface (Net LW) and (e) net shortwave radiation at the surface (Net SW) for a set of simulations with $N_i = 600 \text{ m}^{-3}$ (_ni600) and the no_ship case ($N_i = 200 \text{ m}^{-3}$). Net radiative fluxes are calculated by subtracting the upwelling radiative flux from the downwelling flux (e.g., $\text{LW}_{\text{down}} - \text{LW}_{\text{up}}$), hence, a negative value implies net outgoing radiation. no_ship refers to the reference case with background aerosol only. The HiS_sul and WS_sul cases represent sulfate particle modes of high FSC fuel combustion and exhaust gas wet scrubbing from Santos et al. (2024b). Only ship exhaust sensitivity cases with high concentrations (_hi) are shown in the figure. Significant differences between ship exhaust cases and no_ship_ni600 were assessed using two-sided t tests at a confidence level of 95%. Model runs with significant differences are marked with star icons in inset legends. The last 4 simulation hours were used to perform statistical tests.

In general, ship emissions can lead to more but smaller liquid droplets in the mixed-phase clouds studied here. This means that even if the clouds contain more liquid water, rain surface precipitation is reduced. Nevertheless, this response was strongly coupled to the cloud IWP which is an indicator for the cloud thickness. In our case study, total surface precipitation rates are dominated by graupel which is not significantly affected by additional ship aerosol particles. Moreover, ship exhaust emissions have the potential to affect cloud radiative processes that play a vital role in the Arctic climate system. While the shortwave radiative budget is mostly unaffected, ship exhaust perturbations can lead to both reductions and increases in net longwave radiative cooling at the surface and potentially impact the net surface radiative budget. The magnitude of the ship exhaust-induced cloud perturbations is strongly dependent on the number concentrations of the particle emissions. It is also affected more by the size of the exhaust particles than by their hygroscopicity. This means that the impact of ship emissions on Arctic cloud properties would depend strongly on the fuel types used and whether exhaust after-treatment systems like scrubbers are used. Uncertainties in Arctic fuel type projections increase the challenge of constraining regional climate impacts from Arctic shipping. Transitions towards fuels with reduced sulfur content have been shown to lead to substantial reductions in CCN number emissions, which potentially could reduce radiative effects from ship aerosol-cloud interactions. This is due to a reduction in hygroscopicity of the ship exhaust particles (Santos et al., 2023, 2024b) and a shift in the particle size distribution towards smaller sizes (decreasing CMD; Lack et al., 2011; Yu et al., 2020, 2023). From June 2025 ships will no longer be allowed to carry and use fuel oils with densities and viscosities exceeding predefined limits (IMO, 2021), which could have ramifications for wet scrubber usage, as these systems are mainly designed for use with high FSC residual fuels. Moreover, due to environmental concerns associated with increased Arctic shipping activity, Canada and several organizations have proposed BC emission control areas for Arctic waters, that include mandates for low FSC distillate fuel usage for ships operating in the Arctic (IMO, 2023b, a). Assuming that these proposals are ratified, they would limit the allowed Arctic ship fuels and thus, facilitate estimating the climate impact from shipping. In this case, other environmental impacts from shipping such as BC deposition on snow surfaces, which reduces the surface albedo and enhances surface warming, could play a larger role and dominate the climate impact from Arctic shipping.

Ship exhaust particle concentrations above 1000 cm^{-3} are realistic if one considers narrower and more localized regions where ship exhaust particles perturb clouds (Hobbs et al., 2000; Possner et al., 2018). Once emitted by a transiting ship, exhaust particles become dispersed in the atmosphere, resulting generally in smaller particle number concentrations. As a result, perturbations would likely exert changes in cloud properties more akin to our low concentration model runs. One area of focus for future research could be to implement ship plume dispersion and to use more realistic vertical exhaust particle concentration profiles. Moreover, exhaust particles will undergo chemical and physical transformations in the atmosphere associated with ship exhaust plume aging. In aged plumes, changes in particle size distributions are often observed due to coagulation of exhaust particles, and condensation and evaporation of water vapor and other atmospheric substances (Petzold et al., 2008; Celik et al., 2020). If, for example, fewer but larger particles are present, it could have a stronger impact on cloud hydrometeors compared to the ship exhaust particle size distributions and number concentrations implemented in this study.

While this study focuses on Arctic shipping and clouds, IMO FSC regulations apply worldwide. This means that global
430 ship exhaust emissions are subject to changes and thus, the global radiative forcing exerted by ship exhaust emissions will
likely change. It is therefore important to improve our general understanding of the potential effects of FSC reduction and wet
scrubbing on particulate matter emissions and what this implies for cloud and climate processes.

5 Conclusions

In this study, we used LES together with aerosol data from laboratory experiments to examine the potential impact of ship
435 exhaust particles on Arctic mixed-phase cloud properties. The laboratory experiments investigated the impacts of fuel sul-
fur content reduction and exhaust wet scrubbing on the physicochemical properties of ship exhaust particles (Santos et al.,
2022, 2023, 2024b). Wet scrubbing and FSC reduction represent regulatory compliance measures in the maritime shipping
sector, which affect ship exhaust particle emissions and could potentially be utilized by ships in the Arctic. Given the projected
increase in Arctic shipping activity due to strongly declining Arctic sea-ice extent and the availability of shorter trans-Arctic
440 transportation routes, we have sought to illuminate how ship emissions may impact Arctic clouds and thereby affect the regional
radiative balance.

The simulations were done for a persistent stratiform mixed-phase cloud, based on observations from the ASCOS campaign
(Tjernström et al., 2012, 2014) and previous simulations (Igel et al., 2017; Stevens et al., 2018; Christiansen et al., 2020;
Sotiropoulou et al., 2021; Frostenberg et al., 2023). The simulated cloud was subsequently perturbed by adding ship exhaust
445 particle profiles into the model domain. A selected number of model runs were repeated with increased pre-fixed N_i to study
the impact of ship exhaust perturbations on a thinner baseline cloud with increased IWP and reduced LWP.

Ship exhaust simulations revealed potential impacts on cloud droplet and raindrop concentrations, affecting the LWP and
decreasing the cloud drop effective radius. Total surface precipitation was found to be mostly unaffected; liquid-phase precip-
itation was reduced, but it was only a minor constituent of total surface precipitation. Moreover, the cloud albedo increased
450 marginally in all ship exhaust experiments. Our first set of simulations, with N_i concentrations in line with observations and
IWP values at the lower end of the retrieved values, demonstrated that ship exhaust perturbations can lead to a reduction in
longwave radiative cooling at the surface of up to 4.8 W m^{-2} . This result implies that ship emissions may lead to a net warm-
ing effect compared to our baseline simulation without ship exhaust aerosol. The magnitude of the surface radiation change
depended on the hygroscopicity and the CMD of the added ship aerosol particles, where the effect of the CMD was most
455 important. Additional sensitivity tests with N_i increased to 600 m^{-3} , with reduced LWP and increased IWP (both in line with
retrieved values), revealed that ship exhaust perturbations may lead to enhanced surface radiative cooling. This demonstrates
that the net effect of ship exhaust emissions on the radiative forcing exerted by Arctic low-level clouds would not only strongly
depend on the prevalent fuel types, and whether ships in the Arctic utilize wet scrubbers for exhaust after-treatment, but also
460 strongly coupled to the surface properties, and that sea-ice free conditions can lead to generally larger cloud fractions and

increased LWP (Barton and Veron, 2012; Taylor and Monroe, 2023). It is therefore likely that future Arctic low-level cloud properties may be more similar to our first case ($N_i = 200 \text{ m}^{-2}$).

465 Bulatovic et al. (2021) showed large variations in cloud microphysical properties for different background aerosol concentrations and sizes. Results of ship exhaust perturbations may therefore vary substantially with different background aerosol concentrations and thus, the microphysical structure of the perturbed cloud, as demonstrated by our additional set of simulations with increased N_i . Consequently, it is important to highlight that the case study used in this study is based on observations made in the high Arctic. Most of Arctic shipping activity will likely occur closer to coastal regions where air masses are likely to be more strongly influenced by anthropogenic and biogenic activity (see, for example, Smith and Stephenson (2013)). This means that the atmospheric background conditions and cloud properties may vary from the mixed-phase cloud case studied here and will likely affect the impact of ship exhaust perturbations on cloud properties. Moreover, the enhanced warming that 470 the Arctic is experiencing will likely change the state of ambient aerosol concentrations due to biogenic and anthropogenic processes (Schmale et al., 2021). For example, more open sea surface area will lead to enhanced new particle formation due to marine biogenic emissions (Dall'Osto et al., 2017). General low ambient aerosol number concentrations mean that already small increases in concentrations can have large impacts on cloud properties (Mauritsen et al., 2011). It is therefore expected 475 that these changes will also affect the properties of Arctic clouds.

Our collective results do show evidence that Arctic shipping emissions can lead to alterations in the micro- and macrophysical state of persistent Arctic low-level mixed-phase clouds. While a stronger tendency towards enhanced surface warming from ship exhaust emissions was obtained, this effect was mostly observed when ship aerosol concentrations were increased by $N = 1000 \text{ cm}^{-3}$. When low ship exhaust particle concentrations ($N = 100 \text{ cm}^{-3}$) were utilized, only wet scrubbing model 480 runs were found to alter cloud radiative properties significantly compared to the baseline. However, given the ban on carriage and usage of high-density/viscosity residual fuel oils in 2025, wet scrubbing might not be utilized by a large fraction of ships in the Arctic (IMO, 2021).

This study may help in constraining possible climate feedbacks from a projected increase in Arctic shipping activity. However, our results show that more information on future Arctic shipping activity, including fuel types, traffic volume and associated emissions characteristics, prevalent meteorological conditions, and cloud types is required for more accurate estimates. 485

Data availability. The model output data presented in this study are available at <https://doi.org/10.5281/zenodo.12166747> (Santos et al., 2024a). The experimental data utilized in this study has been previously published and made available through the cited publications, Santos et al. (2022, 2023, 2024b).

Appendix A: Meteorological parameter of ASCOS case




Figure A1. Radiosonde observations of (a) temperature, (b) potential temperature, (c) specific humidity and (d) derived cloud liquid water based on radiometer measurements performed on August 31st 2008 during the ASCOS campaign (Tjernström et al., 2012, 2014). The data was used to initialize MIMICA in this study.




Figure B1. Temporal evolution of horizontally averaged number concentrations of ice crystals (N_i), graupel (N_g) and snow (N_s) simulated for the reference case (no_ship) and the high ship aerosol concentration cases LoS_hi, HiS_hi, WS_hi, HiS_sul_hi and WS_sul_hi. The black dashed lines represent case-specific, horizontally averaged cloud bottom and cloud top heights. The spin-up period (0 to 4 h) is removed from all figures.




Figure B2. Temporal evolution of horizontally averaged mixing ratios of ice crystals (Q_i), graupel (Q_g) and snow (Q_s) simulated for the reference case (no_ship) and the high ship aerosol concentration cases LoS_hi, HiS_hi, WS_hi, HiS_sul_hi and WS_sul_hi. The black dashed lines represent case-specific, horizontally averaged cloud bottom and cloud top heights. The spin-up period (0 to 4 h) is removed from all figures.

Appendix C: Net shortwave surface fluxes




Figure C1. (a) Time evolution of the simulated domain-averaged net shortwave radiation at the surface (Net SW) for the set of ship exhaust sensitivity cases with $N_i = 200 \text{ m}^{-3}$ and the corresponding no_ship case. Only ship exhaust sensitivity cases with high concentrations (_hi) are shown in the figure. (b) Corresponding absolute changes in net SW surface fluxes calculated as $(\text{Net SW})_{\text{ship}} - (\text{Net SW})_{\text{no_ship}}$.




Figure C2. (a) Time evolution of the simulated domain-averaged net shortwave radiation at the surface (Net SW) for the set of ship exhaust sensitivity cases with $N_i = 600 \text{ m}^{-3}$ and the corresponding no_ship_ni600 case. Only ship exhaust sensitivity cases with high concentrations (_hi) are shown in the figure. (b) Corresponding absolute changes in net SW surface fluxes calculated as $(\text{Net SW})_{\text{ship}} - (\text{Net SW})_{\text{no_ship}}$.

Appendix D: Averaged profiles of ice-phase hydrometeors




Figure D1. Vertical profiles of horizontally averaged (a and g) N_i , (b and h) Q_i , (c and i) N_g , (d and j) Q_g , (e and k) N_s and (f and l) Q_s averaged over the last four simulation hours. The light blue, dashed line represents the average cloud bottom and top height calculated for the reference case (no_ship). HiS, LoS, and WS represent ship aerosol from measurements of high and low sulfur content fuels and wet scrubbing respectively (Santos et al., 2022, 2023). The HiS_sul and WS_sul cases represent sulfate particle modes of high FSC fuel combustion and exhaust gas wet scrubbing from Santos et al. (2024b). The label additions *_lo* and *_hi* signify the ship aerosol concentrations used in the individual model runs. Significant differences between ship exhaust cases and no_ship were assessed using two-sided *t* tests at a confidence level of 95 %. Model runs with significant differences are marked with star icons in inset legends.

Author contributions. Idea: EST and AE. Conceptualization: LI, HF, AE. Model implementation and simulations: LS, HF, LI, and ABP. Analysis of the data: LS, HF, LI, ABP. Visualization: LS. Writing, review and editing: All co-authors.

495 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. This research was funded by the Swedish Research Councils FORMAS (2017-00564) and VR (2020-03497 & 2021-04042). EST, LS, HF, ABP, and LI thank the Swedish Strategic Research Area MERGE for their support. HF and LI were supported by Chalmers Gender Initiative for Excellence (Genie). The computations were enabled by resources provided by the National Academic Infrastructure for Supercomputing in Sweden (NAISS) at the National Supercomputer Centre (NSC) partially funded by the Swedish Research Council through grant agreement no. 2022-06725.

References

- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.
- Andreasen, A. and Mayer, S.: Use of seawater scrubbing for SO₂ removal from marine engine exhaust gas, *Energy and Fuels*, 21, 3274–3279, <https://doi.org/10.1021/ef700359w>, 2007.
- 505 Barton, N. and Veron, D.: Response of clouds and surface energy fluxes to changes in sea-ice cover over the Laptev Sea (Arctic Ocean), *Climate Research*, 54, 69–84, <https://doi.org/10.3354/cr01101>, 2012.
- Browse, J., Carslaw, K. S., Schmidt, A., and Corbett, J. J.: Impact of future Arctic shipping on high-latitude black carbon deposition, *Geophysical Research Letters*, 40, 4459–4463, <https://doi.org/10.1002/grl.50876>, 2013.
- 510 Bulatovic, I., Igel, A. L., Leck, C., Heintzenberg, J., Riipinen, I., and Ekman, A. M. L.: The importance of Aitken mode aerosol particles for cloud sustenance in the summertime high Arctic – a simulation study supported by observational data, *Atmos. Chem. Phys.*, 21, 3871–3897, <https://doi.org/10.5194/acp-21-3871-2021>, 2021.
- Bulatovic, I., Savre, J., Tjernström, M., Leck, C., and Ekman, A. M. L.: Large-eddy simulation of a two-layer boundary-layer cloud system from the Arctic Ocean 2018 expedition, *Atmospheric Chemistry and Physics*, 23, 7033–7055, <https://doi.org/10.5194/acp-23-7033-2023>,
515 2023.
- Celik, S., Drewnick, F., Fachinger, F., Brooks, J., Darbyshire, E., Coe, H., Paris, J.-D., Eger, P. G., Schuladen, J., Tadic, I., Friedrich, N., Dienhart, D., Hottmann, B., Fischer, H., Crowley, J. N., Harder, H., and Borrmann, S.: Influence of vessel characteristics and atmospheric processes on the gas and particle phase of ship emission plumes: in situ measurements in the Mediterranean Sea and around the Arabian Peninsula, *Atmospheric Chemistry and Physics*, 20, 4713–4734, <https://doi.org/10.5194/acp-20-4713-2020>, 2020.
- 520 Christensen, M. W., Suzuki, K., Zambri, B., and Stephens, G. L.: Ship track observations of a reduced shortwave aerosol indirect effect in mixed-phase clouds, *Geophysical Research Letters*, 41, 6970–6977, <https://doi.org/10.1002/2014GL061320>, 2014.
- Christiansen, S., Ickes, L., Bulatovic, I., Leck, C., Murray, B. J., Bertram, A. K., Wagner, R., Gorokhova, E., Salter, M. E., Ekman, A. M., and Bilde, M.: Influence of Arctic Microlayers and Algal Cultures on Sea Spray Hygroscopicity and the Possible Implications for Mixed-Phase Clouds, *Journal of Geophysical Research: Atmospheres*, 125, <https://doi.org/10.1029/2020JD032808>, 2020.
- 525 Coakley, J. A., Bernstein, R. L., and Durkee, P. A.: Effect of Ship-Stack Effluents on Cloud Reflectivity, *Science*, 237, 1020–1022, <https://doi.org/10.1126/science.237.4818.1020>, 1987.
- Corbett, J. J., Winebrake, J. J., Green, E. H., Kasibhatla, P., Eyring, V., and Lauer, A.: Mortality from Ship Emissions: A Global Assessment, *Environmental Science & Technology*, 41, 8512–8518, <https://doi.org/10.1021/es071686z>, 2007.
- Corbett, J. J., Lack, D. A., Winebrake, J. J., Harder, S., Silberman, J. A., and Gold, M.: Arctic shipping emissions inventories and future
530 scenarios, *Atmospheric Chemistry and Physics*, 10, 9689–9704, <https://doi.org/10.5194/acp-10-9689-2010>, 2010.
- Dall’Osto, M., Beddows, D. C. S., Tunved, P., Krejci, R., Ström, J., Hansson, H.-C., Yoon, Y. J., Park, K.-T., Becagli, S., Udisti, R., Onasch, T., O’Dowd, C. D., Simó, R., and Harrison, R. M.: Arctic sea ice melt leads to atmospheric new particle formation, *Scientific Reports*, 7, 3318, <https://doi.org/10.1038/s41598-017-03328-1>, 2017.
- Dalsøren, S. B., Samset, B. H., Myhre, G., Corbett, J. J., Minjares, R., Lack, D., and Fuglestedt, J. S.: Environmental Impacts of Shipping in
535 2030 with a Particular Focus on the Arctic Region, *Atmospheric Chem. Phys.*, 13, 1941–1955, <https://doi.org/10.5194/acp-13-1941-2013>, 2013.

- Durkee, P. A., Noone, K. J., Ferek, R. J., Johnson, D. W., Taylor, J. P., Garrett, T. J., Hobbs, P. V., Hudson, J. G., Bretherton, C. S., Innis, G., Frick, G. M., Hoppel, W. A., O'Dowd, C. D., Russell, L. M., Gasparovic, R., Nielsen, K. E., Tessmer, S. A., Öström, E., Osborne, S. R., Flagan, R. C., Seinfeld, J. H., and Rand, H.: The Impact of Ship-Produced Aerosols on the Microstructure and Albedo of Warm Marine Stratocumulus Clouds: A Test of MAST Hypotheses Ii and Iii, *Journal of the Atmospheric Sciences*, 57, 2554–2569, [https://doi.org/10.1175/1520-0469\(2000\)057<2554:TIOSPA>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<2554:TIOSPA>2.0.CO;2), 2000.
- 540 Eirund, G. K., Possner, A., and Lohmann, U.: Response of Arctic mixed-phase clouds to aerosol perturbations under different surface forcings, *Atmospheric Chemistry and Physics*, 19, 9847–9864, <https://doi.org/10.5194/acp-19-9847-2019>, 2019.
- Eyring, V., Isaksen, I. S., Bernsten, T., Collins, W. J., Corbett, J. J., Endresen, O., Grainger, R. G., Moldanova, J., Schlager, H., and Stevenson, D. S.: Transport impacts on atmosphere and climate: Shipping, *Atmospheric Environment*, 44, 4735–4771, <https://doi.org/10.1016/j.atmosenv.2009.04.059>, 2010.
- 545 Freud, E. and Rosenfeld, D.: Linear relation between convective cloud drop number concentration and depth for rain initiation, *Journal of Geophysical Research: Atmospheres*, 117, n/a–n/a, <https://doi.org/10.1029/2011JD016457>, 2012.
- Fridell, E. and Salo, K.: Measurements of abatement of particles and exhaust gases in a marine gas scrubber, *Proceedings of the Institution of Mechanical Engineers, Part M: Journal of Engineering for the Maritime Environment*, 230, 154–162, <https://doi.org/10.1177/1475090214543716>, 2016.
- 550 Frostenberg, H. C., Welti, A., Luhr, M., Savre, J., Thomson, E. S., and Ickes, L.: The chance of freezing – a conceptual study to parameterize temperature-dependent freezing by including randomness of ice-nucleating particle concentrations, *Atmospheric Chemistry and Physics*, 23, 10 883–10 900, <https://doi.org/10.5194/acp-23-10883-2023>, 2023.
- 555 Fu, Q. and Liou, K. N.: On the Correlated k-Distribution Method for Radiative Transfer in Nonhomogeneous Atmospheres, *Journal of Atmospheric Sciences*, 49, 2139 – 2156, [https://doi.org/10.1175/1520-0469\(1992\)049<2139:OTCDMF>2.0.CO;2](https://doi.org/10.1175/1520-0469(1992)049<2139:OTCDMF>2.0.CO;2), 1992.
- Gilgen, A., Huang, W. T. K., Ickes, L., Neubauer, D., and Lohmann, U.: How important are future marine and shipping aerosol emissions in a warming Arctic summer and autumn?, *Atmospheric Chemistry and Physics*, 18, 10 521–10 555, <https://doi.org/10.5194/acp-18-10521-2018>, 2018.
- 560 Gryspeerdt, E., Smith, T. W. P., O’Keeffe, E., Christensen, M. W., and Goldsworth, F. W.: The Impact of Ship Emission Controls Recorded by Cloud Properties, *Geophysical Research Letters*, 46, 12 547–12 555, <https://doi.org/10.1029/2019GL084700>, 2019.
- Hobbs, P. V., Garrett, T. J., Ferek, R. J., Strader, S. R., Hegg, D. A., Frick, G. M., Hoppel, W. A., Gasparovic, R. F., Russell, L. M., Johnson, D. W., O’Dowd, C., Durkee, P. A., Nielsen, K. E., and Innis, G.: Emissions from ships with respect to their effects on clouds, *Journal of the Atmospheric Sciences*, 57, 2570–2590, [https://doi.org/10.1175/1520-0469\(2000\)057<2570:EFSWRT>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<2570:EFSWRT>2.0.CO;2), 2000.
- 565 Igel, A. L., Ekman, A. M. L., Leck, C., Tjernström, M., Savre, J., and Sedlar, J.: The free troposphere as a potential source of arctic boundary layer aerosol particles, *Geophysical Research Letters*, 44, 7053–7060, <https://doi.org/10.1002/2017GL073808>, 2017.
- IMO: RESOLUTION MEPC.176(58) - Revised MARPOL Annex VI, 2008.
- IMO: RESOLUTION MEPC.329(76) - Prohibition on the use and carriage for use as fuel of heavy fuel oil by ships in Arctic waters, 2021.
- IMO: MEPC 80/9/1 - Reducing Black Carbon emissions in the Arctic, 2023a.
- 570 IMO: MEPC 80/16/2 - Development of a proposal to designate a Canadian Arctic Emission Control Area, 2023b.
- Intrieri, J. M., Fairall, C. W., Shupe, M. D., Persson, P. O. G., Andreas, E. L., Guest, P. S., and Moritz, R. E.: An annual cycle of Arctic surface cloud forcing at SHEBA, *Journal of Geophysical Research: Oceans*, 107, <https://doi.org/10.1029/2000JC000439>, 2002.
- Jeong, S., Bendl, J., Saraji-Bozorgzad, M., Käfer, U., Etzien, U., Schade, J., Bauer, M., Jakobi, G., Orasche, J., Fisch, K., Cwierz, P. P., Rüger, C. P., Czech, H., Karg, E., Heyen, G., Krausnick, M., Geissler, A., Geipel, C., Streibel, T., Schnelle-Kreis, J., Sklorz, M., Schulz-Bull,

- 575 D. E., Buchholz, B., Adam, T., and Zimmermann, R.: Aerosol emissions from a marine diesel engine running on different fuels and effects of exhaust gas cleaning measures, *Environmental Pollution*, 316, 120 526, <https://doi.org/10.1016/j.envpol.2022.120526>, 2023.
- Jonson, J. E., Gauss, M., Schulz, M., Jalkanen, J.-P., and Fagerli, H.: Effects of global ship emissions on European air pollution levels, *Atmospheric Chemistry and Physics*, 20, 11 399–11 422, <https://doi.org/10.5194/acp-20-11399-2020>, 2020.
- Kanji, Z. A., Welti, A., Corbin, J. C., and Mensah, A. A.: Black Carbon Particles Do Not Matter for Immersion Mode Ice Nucleation, *Geophysical Research Letters*, 47, 1–9, <https://doi.org/10.1029/2019GL086764>, 2020.
- 580 Kuittinen, N., Jalkanen, J.-P., Alanen, J., Ntziachristos, L., Hannuniemi, H., Johansson, L., Karjalainen, P., Saukko, E., Isotalo, M., Aakko-Saksa, P., Lehtoranta, K., Keskinen, J., Simonen, P., Saarikoski, S., Asmi, E., Laurila, T., Hillamo, R., Mylläri, F., Lihavainen, H., Timonen, H., and Rönkkö, T.: Shipping Remains a Globally Significant Source of Anthropogenic PN Emissions Even after 2020 Sulfur Regulation, *Environmental Science & Technology*, 55, 129–138, <https://doi.org/10.1021/acs.est.0c03627>, 2021.
- 585 Kupiszewski, P., Leck, C., Tjernström, M., Sjogren, S., Sedlar, J., Graus, M., Müller, M., Brooks, B., Swietlicki, E., Norris, S., and Hansel, A.: Vertical profiling of aerosol particles and trace gases over the central Arctic Ocean during summer, *Atmospheric Chemistry and Physics*, 13, 12 405–12 431, <https://doi.org/10.5194/acp-13-12405-2013>, 2013.
- Lack, D. A. and Corbett, J. J.: Black carbon from ships: A review of the effects of ship speed, fuel quality and exhaust gas scrubbing, *Atmospheric Chemistry and Physics*, 12, 3985–4000, <https://doi.org/10.5194/acp-12-3985-2012>, 2012.
- 590 Lack, D. A., Corbett, J. J., Onasch, T., Lerner, B., Massoli, P., Quinn, P. K., Bates, T. S., Covert, D. S., Coffman, D., Sierau, B., Herndon, S., Allan, J., Baynard, T., Lovejoy, E., Ravishankara, A. R., and Williams, E.: Particulate emissions from commercial shipping: Chemical, physical, and optical properties, *J. Geophys. Res.: Atmos.*, 114, 1–16, <https://doi.org/10.1029/2008JD011300>, 2009.
- Lack, D. A., Cappa, C. D., Langridge, J., Bahreini, R., Buffaloe, G., Brock, C., Cerully, K., Coffman, D., Hayden, K., Holloway, J., Lerner, B., Massoli, P., Li, S.-M., McLaren, R., Middlebrook, A. M., Moore, R., Nenes, A., Nuaaman, I., Onasch, T. B., Peischl, J., Perring, A., Quinn, P. K., Ryerson, T., Schwartz, J. P., Spackman, R., Wofsy, S. C., Worsnop, D., Xiang, B., and Williams, E.: Impact of Fuel Quality Regulation and Speed Reductions on Shipping Emissions: Implications for Climate and Air Quality, *Environmental Science & Technology*, 45, 9052–9060, <https://doi.org/10.1021/es2013424>, 2011.
- 595 Lasserre, F. and Pelletier, S.: Polar super seaways? Maritime transport in the Arctic: an analysis of shipowners' intentions, *Journal of Transport Geography*, 19, 1465–1473, <https://doi.org/10.1016/j.jtrangeo.2011.08.006>, 2011.
- 600 Lauer, A., Eyring, V., Hendricks, J., Jöckel, P., and Lohmann, U.: Global model simulations of the impact of ocean-going ships on aerosols, clouds, and the radiation budget, *Atmospheric Chemistry and Physics*, 7, 5061–5079, <https://doi.org/10.5194/acp-7-5061-2007>, 2007.
- Lehtoranta, K., Aakko-Saksa, P., Murtonen, T., Vesala, H., Ntziachristos, L., Rönkkö, T., Karjalainen, P., Kuittinen, N., and Timonen, H.: Particulate Mass and Nonvolatile Particle Number Emissions from Marine Engines Using Low-Sulfur Fuels, Natural Gas, or Scrubbers, *Environmental Science & Technology*, 53, 3315–3322, <https://doi.org/10.1021/acs.est.8b05555>, 2019.
- 605 Li, X., Lynch, A. H., Bailey, D. A., Stephenson, S. R., and Veland, S.: The impact of black carbon emissions from projected Arctic shipping on regional ice transport, *Climate Dynamics*, 57, 2453–2466, <https://doi.org/10.1007/s00382-021-05814-9>, 2021.
- Lieke, K. I., Rosenørn, T., Pedersen, J., Larsson, D., Kling, J., Fuglsang, K., and Bilde, M.: Micro- and Nanostructural Characteristics of Particles Before and After an Exhaust Gas Recirculation System Scrubber, *Aerosol Science and Technology*, 47, 1038–1046, <https://doi.org/10.1080/02786826.2013.813012>, 2013.
- 610 Liu, H., Fu, M., Jin, X., Shang, Y., Shindell, D., Faluvegi, G., Shindell, C., and He, K.: Health and climate impacts of ocean-going vessels in East Asia, *Nature Climate Change*, 6, 1037–1041, <https://doi.org/10.1038/nclimate3083>, 2016.

- Lund, M. T., Eyring, V., Fuglestedt, J., Hendricks, J., Lauer, A., Lee, D., and Righi, M.: Global-Mean Temperature Change from Shipping toward 2050: Improved Representation of the Indirect Aerosol Effect in Simple Climate Models, *Environmental Science & Technology*, 46, 8868–8877, <https://doi.org/10.1021/es301166e>, 2012.
- 615 Lund, M. T., Aamaas, B., Stjern, C. W., Klimont, Z., Berntsen, T. K., and Samset, B. H.: A continued role of short-lived climate forcers under the Shared Socioeconomic Pathways, *Earth System Dynamics*, 11, 977–993, <https://doi.org/10.5194/esd-11-977-2020>, 2020.
- Mahrt, F., Marcolli, C., David, R. O., Grönquist, P., Barthazy Meier, E. J., Lohmann, U., and Kanji, Z. A.: Ice nucleation abilities of soot particles determined with the Horizontal Ice Nucleation Chamber, *Atmospheric Chemistry and Physics*, 18, 13363–13392, <https://doi.org/10.5194/acp-18-13363-2018>, 2018.
- 620 Manshausen, P., Watson-Parris, D., Christensen, M. W., Jalkanen, J.-P., and Stier, P.: Invisible Ship Tracks Show Large Cloud Sensitivity to Aerosol, *Nature*, 610, 101–106, <https://doi.org/10.1038/s41586-022-05122-0>, 2022.
- Mauritsen, T., Sedlar, J., Tjernström, M., Leck, C., Martin, M., Shupe, M., Sjogren, S., Sierau, B., Persson, P. O., Brooks, I. M., and Swietlicki, E.: An Arctic CCN-limited cloud-aerosol regime, *Atmos. Chem. Phys.*, 11, 165–173, <https://doi.org/10.5194/acp-11-165-2011>, 2011.
- Meador, W. E. and Weaver, W. R.: Two-Stream Approximations to Radiative Transfer in Planetary Atmospheres: A Unified Description of Existing Methods and a New Improvement, *Journal of Atmospheric Sciences*, 37, 630 – 643, [https://doi.org/10.1175/1520-0469\(1980\)037<0630:TSATRT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<0630:TSATRT>2.0.CO;2), 1980.
- Morrison, H., De Boer, G., Feingold, G., Harrington, J., Shupe, M. D., and Sulia, K.: Resilience of persistent Arctic mixed-phase clouds, *Nature Geoscience*, 5, 11–17, <https://doi.org/10.1038/ngeo1332>, 2012.
- Oikawa, K., Yongsiri, C., Takeda, K., and Harimoto, T.: Seawater flue gas desulfurization: Its technical implications and performance results, 630 *Environmental Progress*, 22, 67–73, <https://doi.org/10.1002/ep.670220118>, 2003.
- Ovchinnikov, M., Korolev, A., and Fan, J.: Effects of ice number concentration on dynamics of a shallow mixed-phase stratiform cloud, *Journal of Geophysical Research*, 116, D00T06, <https://doi.org/10.1029/2011JD015888>, 2011.
- Ovchinnikov, M., Ackerman, A. S., Avramov, A., Cheng, A., Fan, J., Fridlind, A. M., Ghan, S., Harrington, J., Hoose, C., Korolev, A., McFarquhar, G. M., Morrison, H., Paukert, M., Savre, J., Shipway, B. J., Shupe, M. D., Solomon, A., and Sulia, K.: Intercomparison of 635 large-eddy simulations of Arctic mixed-phase clouds: Importance of ice size distribution assumptions, *Journal of Advances in Modeling Earth Systems*, 6, 223–248, <https://doi.org/10.1002/2013MS000282>, 2014.
- Paxian, A., Eyring, V., Beer, W., Sausen, R., and Wright, C.: Present-day and future global bottom-up ship emission inventories including polar routes, *Environmental Science and Technology*, 44, 1333–1339, <https://doi.org/10.1021/es9022859>, 2010.
- Peters, G. P., Nilssen, T. B., Lindholt, L., Eide, M. S., Glomsrød, S., Eide, L. I., and Fuglestedt, J. S.: Future emissions from shipping and 640 petroleum activities in the Arctic, *Atmospheric Chemistry and Physics*, 11, 5305–5320, <https://doi.org/10.5194/acp-11-5305-2011>, 2011.
- Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmospheric Chemistry and Physics*, 7, 1961–1971, <https://doi.org/10.5194/acp-7-1961-2007>, 2007.
- Petzold, A., Hasselbach, J., Lauer, P., Baumann, R., Franke, K., Gurk, C., Schlager, H., and Weingartner, E.: Experimental studies on particle emissions from cruising ship, their characteristic properties, transformation and atmospheric lifetime in the marine boundary 645 layer, *Atmospheric Chemistry and Physics*, 8, 2387–2403, <https://doi.org/10.5194/acp-8-2387-2008>, 2008.
- Possner, A., Ekman, A. M. L., and Lohmann, U.: Cloud response and feedback processes in stratiform mixed-phase clouds perturbed by ship exhaust, *Geophysical Research Letters*, 44, 1964–1972, <https://doi.org/10.1002/2016GL071358>, 2017.

- Possner, A., Wang, H., Wood, R., Caldeira, K., and Ackerman, T. P.: The efficacy of aerosol–cloud radiative perturbations from near-surface emissions in deep open-cell stratocumuli, *Atmospheric Chemistry and Physics*, 18, 17 475–17 488, <https://doi.org/10.5194/acp-18-17475-2018>, 2018.
- 650 Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., and Laaksonen, A.: The Arctic has warmed nearly four times faster than the globe since 1979, *Communications Earth & Environment*, 3, 168, <https://doi.org/10.1038/s43247-022-00498-3>, 2022.
- Roberts, G. C. and Nenes, A.: A Continuous-Flow Streamwise Thermal-Gradient CCN Chamber for Atmospheric Measurements, *Aerosol Science and Technology*, 39, 206–221, <https://doi.org/10.1080/027868290913988>, 2005.
- 655 Santos, L. F. E. d., Salo, K., and Thomson, E. S.: Quantification and physical analysis of nanoparticle emissions from a marine engine using different fuels and a laboratory wet scrubber, *Environ. Sci.: Processes Impacts*, 24, 1769–1781, <https://doi.org/10.1039/D2EM00054G>, 2022.
- Santos, L. F. E. d., Salo, K., Kong, X., Noda, J., Kristensen, T. B., Ohigashi, T., and Thomson, E. S.: Changes in CCN activity of ship exhaust particles induced by fuel sulfur content reduction and wet scrubbing, *Environmental Science: Atmospheres*, 3, 182–195, <https://doi.org/10.1039/D2EA00081D>, 2023.
- 660 Santos, L. F. E. D., Frostenberg, H., Baró Pérez, A., Ekman, A. M. L., Ickes, L., and Thomson, E. S.: Model output from "Potential impacts of marine fuel regulations on an Arctic stratocumulus case and its radiative response", <https://doi.org/10.5281/zenodo.12166747>, 2024a.
- Santos, L. F. E. D., Salo, K., Kong, X., Hartmann, M., Sjöblom, J., and Thomson, E. S.: Marine Fuel Regulations and Engine Emissions: Impacts on Physicochemical Properties, Cloud Activity and Emission Factors, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD040 389, <https://doi.org/https://doi.org/10.1029/2023JD040389>, 2024b.
- 665 Savre, J., Ekman, A. M. L., and Svensson, G.: Technical note: Introduction to MIMICA, a large-eddy simulation solver for cloudy planetary boundary layers, *Journal of Advances in Modeling Earth Systems*, 6, 630–649, <https://doi.org/10.1002/2013MS000292>, 2014.
- Savre, J., Ekman, A. M. L., Svensson, G., and Tjernström, M.: Large-eddy simulations of an Arctic mixed-phase stratiform cloud observed during ISDAC: sensitivity to moisture aloft, surface fluxes and large-scale forcing, *Quarterly Journal of the Royal Meteorological Society*, 141, 1177–1190, <https://doi.org/10.1002/qj.2425>, 2015.
- 670 Schmale, J., Zieger, P., and Ekman, A. M. L.: Aerosols in current and future Arctic climate, *Nature Climate Change*, 11, 95–105, <https://doi.org/10.1038/s41558-020-00969-5>, 2021.
- Screen, J. A. and Simmonds, I.: The central role of diminishing sea ice in recent Arctic temperature amplification, *Nature*, 464, 1334–1337, <https://doi.org/10.1038/nature09051>, 2010.
- 675 Seifert, A. and Beheng, K. D.: A double-moment parameterization for simulating autoconversion, accretion and selfcollection, *Atmospheric Research*, 59-60, 265–281, [https://doi.org/10.1016/S0169-8095\(01\)00126-0](https://doi.org/10.1016/S0169-8095(01)00126-0), 2001.
- Seifert, A. and Beheng, K. D.: A Two-Moment Cloud Microphysics Parameterization for Mixed-Phase Clouds. Part 1: Model Description, *Meteorol. Atmospheric Phys.*, 92, 45–66, <https://doi.org/10.1007/s00703-005-0112-4>, 2006.
- 680 Seppälä, S. D., Kuula, J., Hyvärinen, A.-P., Saarikoski, S., Rönkkö, T., Keskinen, J., Jalkanen, J.-P., and Timonen, H.: Effects of marine fuel sulfur restrictions on particle number concentrations and size distributions in ship plumes in the Baltic Sea, *Atmospheric Chemistry and Physics*, 21, 3215–3234, <https://doi.org/10.5194/acp-21-3215-2021>, 2021.
- Serreze, M. C. and Barry, R. G.: Processes and impacts of Arctic amplification: A research synthesis, *Global and Planetary Change*, 77, 85–96, <https://doi.org/10.1016/j.gloplacha.2011.03.004>, 2011.

- 685 Serreze, M. C. and Francis, J. A.: The Arctic Amplification Debate, *Climatic Change*, 76, 241–264, <https://doi.org/10.1007/s10584-005-9017-y>, 2006.
- Shupe, M. D. and Intrieri, J. M.: Cloud Radiative Forcing of the Arctic Surface: The Influence of Cloud Properties, Surface Albedo, and Solar Zenith Angle, *Journal of Climate*, 17, 616–628, [https://doi.org/10.1175/1520-0442\(2004\)017<0616:CRFOTA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<0616:CRFOTA>2.0.CO;2), 2004.
- Smith, L. C. and Stephenson, S. R.: New Trans-Arctic shipping routes navigable by midcentury, *Proceedings of the National Academy of Sciences*, 110, <https://doi.org/10.1073/pnas.1214212110>, 2013.
- 690 Sotiropoulou, G., Ickes, L., Nenes, A., and Ekman, A. M. L.: Ice Multiplication from Ice–Ice Collisions in the High Arctic: Sensitivity to Ice Habit, Rimed Fraction, Ice Type and Uncertainties in the Numerical Description of the Process, *Atmospheric Chem. Phys.*, 21, 9741–9760, <https://doi.org/10.5194/acp-21-9741-2021>, 2021.
- Stephens, G. L.: Radiation Profiles in Extended Water Clouds. II: Parameterization Schemes, *Journal of Atmospheric Sciences*, 35, 2123 – 2132, 1978.
- 695 Stephenson, S. R., Wang, W., Zender, C. S., Wang, H., Davis, S. J., and Rasch, P. J.: Climatic Responses to Future Trans-Arctic Shipping, *Geophysical Research Letters*, 45, 9898–9908, <https://doi.org/10.1029/2018GL078969>, 2018.
- Stevens, R. G., Loewe, K., Dearden, C., Dimitrellos, A., Possner, A., Eirund, G. K., Raatikainen, T., Hill, A. A., Shipway, B. J., Wilkinson, J., Romakkaniemi, S., Tonttila, J., Laaksonen, A., Korhonen, H., Connolly, P., Lohmann, U., Hoose, C., Ekman, A. M. L., Carslaw, K. S., and Field, P. R.: A model intercomparison of CCN-limited tenuous clouds in the high Arctic, *Atmospheric Chemistry and Physics*, 18, 11 041–11 071, <https://doi.org/10.5194/acp-18-11041-2018>, 2018.
- 700 Tavakoli, F. and Olfert, J. S.: Determination of particle mass, effective density, mass-mobility exponent, and dynamic shape factor using an aerodynamic aerosol classifier and a differential mobility analyzer in tandem, *Journal of Aerosol Science*, 75, 35–42, <https://doi.org/10.1016/j.jaerosci.2014.04.010>, 2014.
- 705 Taylor, P. C. and Monroe, E.: Isolating the Surface Type Influence on Arctic Low-Clouds, *Journal of Geophysical Research: Atmospheres*, 128, e2022JD038 098, <https://doi.org/10.1029/2022JD038098>, 2023.
- Tjernström, M., Birch, C. E., Brooks, I. M., Shupe, M. D., Persson, P. O. G., Sedlar, J., Mauritsen, T., Leck, C., Paatero, J., Szczodrak, M., and Wheeler, C. R.: Meteorological Conditions in the Central Arctic Summer during the Arctic Summer Cloud Ocean Study (ASCOS), *Atmospheric Chem. Phys.*, 12, 6863–6889, <https://doi.org/10.5194/acp-12-6863-2012>, 2012.
- 710 Tjernström, M., Leck, C., Birch, C. E., Bottenheim, J. W., Brooks, B. J., Brooks, I. M., Bäcklin, L., Chang, R. Y.-W., de Leeuw, G., Di Liberto, L., de la Rosa, S., Granath, E., Graus, M., Hansel, A., Heintzenberg, J., Held, A., Hind, A., Johnston, P., Knulst, J., Martin, M., Matrai, P. A., Mauritsen, T., Müller, M., Norris, S. J., Orellana, M. V., Orsini, D. A., Paatero, J., Persson, P. O. G., Gao, Q., Rauschenberg, C., Ristovski, Z., Sedlar, J., Shupe, M. D., Sierau, B., Sirevaag, A., Sjogren, S., Stetzer, O., Swietlicki, E., Szczodrak, M., Vaattovaara, P., Wahlberg, N., Westberg, M., and Wheeler, C. R.: The Arctic Summer Cloud Ocean Study (ASCOS): Overview and Experimental Design, *Atmospheric Chem. Phys.*, 14, 2823–2869, <https://doi.org/10.5194/acp-14-2823-2014>, 2014.
- 715 UNCTAD: Review of Maritime Transport 2023, no. 2022 in Review of maritime transport / United Nations Conference on Trade and Development, Geneva, United Nations, Geneva, 2022.
- Wang, C. and Chang, J. S.: A three-dimensional numerical model of cloud dynamics, microphysics, and chemistry: 1. Concepts and formulation, *Journal of Geophysical Research: Atmospheres*, 98, 14 827–14 844, <https://doi.org/10.1029/92JD01393>, 1993.
- 720 Watson-Parris, D., Christensen, M. W., Laurenson, A., Clewley, D., Gryspeerd, E., and Stier, P.: Shipping Regulations Lead to Large Reduction in Cloud Perturbations, *Proc. Natl. Acad. Sci.*, 119, e2206885 119, <https://doi.org/10.1073/pnas.2206885119>, 2022.

- Winnes, H., Fridell, E., and Moldanová, J.: Effects of Marine Exhaust Gas Scrubbers on Gas and Particle Emissions, *Journal of Marine Science and Engineering*, 8, 299, <https://doi.org/10.3390/jmse8040299>, 2020.
- 725 Yang, J., Tang, T., Jiang, Y., Karavalakis, G., Durbin, T. D., Wayne Miller, J., Cocker, D. R., and Johnson, K. C.: Controlling emissions from an ocean-going container vessel with a wet scrubber system, *Fuel*, 304, 121 323, <https://doi.org/10.1016/j.fuel.2021.121323>, 2021.
- Yu, C., Pasternak, D., Lee, J., Yang, M., Bell, T., Bower, K., Wu, H., Liu, D., Reed, C., Bauguitte, S., Cliff, S., Trembath, J., Coe, H., and Allan, J. D.: Characterizing the Particle Composition and Cloud Condensation Nuclei from Shipping Emission in Western Europe, *Environmental Science & Technology*, 54, 15 604–15 612, <https://doi.org/10.1021/acs.est.0c04039>, 2020.
- 730 Yu, C., Pasternak, D., Lee, J., Yang, M., Bell, T., Bower, K., Wu, H., Liu, D., Reed, C., Bauguitte, S., Cliff, S., Trembath, J., Coe, H., and Allan, J. D.: Correction to “Characterizing the Particle Composition and Cloud Condensation Nuclei from Shipping Emission in Western Europe”, *Environmental Science & Technology*, 57, 7888–7889, <https://doi.org/10.1021/acs.est.3c03005>, 2023.
- Yuan, T., Song, H., Wood, R., Wang, C., Oreopoulos, L., Platnick, S. E., von Hippel, S., Meyer, K., Light, S., and Wilcox, E.: Global Reduction in Ship-Tracks from Sulfur Regulations for Shipping Fuel, *Sci. Adv.*, 8, eabn7988, <https://doi.org/10.1126/sciadv.abn7988>, 2022.
- 735 Zetterdahl, M., Moldanová, J., Pei, X., Pathak, R. K., and Demirdjian, B.: Impact of the 0.1% fuel sulfur content limit in SECA on particle and gaseous emissions from marine vessels, *Atmospheric Environment*, 145, 338–345, <https://doi.org/10.1016/j.atmosenv.2016.09.022>, 2016.