

High-Resolution Snow Water Equivalent Estimation: A Data-Driven Method for Localized Downscaling of Climate Data

Fatemeh Zakeri ^{a,b}, Gregoire Mariethoz ^a, Manuela Girotto ^b

^a Institute of Earth Surface Dynamics, Faculty of Geosciences and Environment, University of Lausanne, Lausanne, Switzerland

^b Department of Environmental Science, Policy, and Management, University of California, Berkeley, Berkeley, CA, USA

Correspondence to: Fatemeh Zakeri (fatemeh.zakeri@unil.ch; fatemeh.zakeri@berkeley.edu)

Abstract. Estimating high-resolution daily Snow Water Equivalent (SWE) in mountainous regions is challenging due to geographical complexity and the irregular availability of high-resolution meteorological data. This study introduces a method for downscaling SWE based on low-resolution climate models. It is based on the dependence between meteorological estimators and SWE, and the fact that while SWE can change rapidly within days, its patterns may exhibit year-day-to-year-day analogies under similar meteorological conditions. We implement this principle to downscale SWE to at a 500 m resolution using a K-nearest neighbor algorithm with a customized distance metric.

To evaluate the performance of our approach, we conduct tests in two regions of interest California's Sierra Nevada and Colorado's Upper Colorado River Basin in the western United States using different low-resolution climate models ('ec-earth3-veg', 'mpi-esm1-2', and 'cnrm-esm2-1') at both 100 km and 9 km scales. A cross-validation analysis is performed, and comparisons are made with commonly used gridded SWE datasets as well as against in-situ data through point-scale time-series comparisons. The results demonstrate that our approach enables the generation of downscaled SWE that, which closely matches that observed in reanalysis data in terms of statistical properties. The outputs demonstrate that, for each region, performance depends on the choice and accuracy of the climate model inputs, such as precipitation and temperature data. Overall, the 'cnrm-esm2-1' model demonstrates superior accuracy in Colorado, outperforming other models at both 100 km and 9 km resolutions. Conversely, the 'ec-earth3-veg' model achieves the best performance in California with 9 km climate data. Across models, a 9 km resolution typically provides slightly better accuracy compared to a 100 km resolution. This opens up possibilities for applications in regions with limited in-situ data or meteorological data measurements. The approach also has the potential to recreate unmeasured historical SWE values and could be extended to future periods using climate projections.

Keywords: Downscaling; Climate Reanalysis Models; Time Series; Snow Water equivalent; k-Nearest Neighbors; Data-Driven Method; Machine Learning; Spatial and Temporal Analysis

30 1. Introduction

The snowpack in high-elevation regions plays a crucial role as a primary source of streamflow, particularly during the spring and summer seasons (Bales et al., 2006). Rapidly changing weather conditions (Ranzi et al., 2024) and extreme events, such as atmospheric rivers, can lead to significant snowmelt and generate extreme runoff, posing threats to both water supply and infrastructure (Henn et al., 2020). Understanding snow and its role in the hydrological cycle is not merely a scientific question, but also ~~a matter of economic significance~~ economically significant as it directly relates to water resources, agriculture, and energy production (Sturm et al., 2017). Therefore, accurate and detailed information on Snow Water Equivalent (SWE) with high temporal and spatial resolution is crucial for effective water resource management and decision-making (Bales et al., 2006; Fiddes et al., 2019; Siirila-Woodburn et al., 2021b).

Although ground stations are valuable for collecting SWE data, their limited presence in certain regions affects their representativeness. ~~Furthermore~~ Moreover, variations in topography, land cover, and environmental conditions in mountainous areas make point-scale data insufficient for capturing the overall spatial ~~scale~~ characteristics of a watershed (Bales et al., 2006; Alonso-González et al., 2023). To ~~mitigate~~ address this lack of data, physically based snow models ~~use~~ utilize an energy balance approach to estimate snowmelt. ~~Various~~ These models ~~of different complexities have been introduced, but~~ range in complexity, with more advanced models integrating detailed processes such as wind-induced snow transport, interactions with topography, and vegetation impacts. While complex models, such as those incorporating advection-diffusion equations or three-dimensional wind fields, provide more accurate representations of snow properties, they often require extensive input data ~~that, which~~ may not always be available (Liston and Sturm, 1998; Lehning et al., 2006; Vionnet et al., 2014) ~~readily accessible, while~~. Simpler models, on the other hand, may ~~not fully~~ fail to capture ~~the complexity~~ critical aspects of ~~the snowmelt process~~ snow dynamics (Bair et al., 2016; Clow et al., 2012). Moreover, to achieve high-resolution SWE (HR-SWE) estimates using these models, it is necessary to use meteorological and land cover-related data that ~~matches~~ match the desired output resolution of the SWE. However, obtaining high-resolution data in mountainous regions remains challenging (Wundram and Löffler, 2008). Although generating HR-SWE with physical models can be time-consuming, recent studies have focused on reducing this computational burden. For instance, advances in SWE modeling have been achieved by implementing parallelized versions of snow models. This approach maintains the integrity of physical processes while utilizing parallelization to manage the computational demands of fine-resolution datasets over large domains (Mower et al., 2024). However, it is important to clarify that the computational expense becomes significant when operating at high spatial and temporal resolutions, particularly in an ensemble context, which is often required for robust climate predictions. Additionally, regardless of the computational aspects, generating high-resolution snow data using physical models typically necessitates high-resolution climate inputs, such as temperature and precipitation fields, to ensure the quality of the downscaled outputs.

~~Reflecting on~~ Given these challenges in obtaining high-resolution climate data, remote sensing ~~is becoming~~ has become increasingly important for monitoring and predicting snowpack conditions and their impacts on water resources (USBR, 2021).

Satellites equipped with optical sensors, such as MODIS and Landsat, provide data on snow cover (Dietz et al., 2012; Painter et al., 2012; Langeron et al., 2020; Wu et al., 2021) and temperature (Lundquist et al., 2018), while lidar and microwave sensors (Tsang et al., 2021; Saberi et al., 2020) offer insights into snow depth or SWE (Lievens et al., 2019; Shi and Dozier, 2000; Pflug et al., 2024; Ma et al., 2023), and wetness (Shi and Dozier, 1995; Snapir et al., 2019). However, limitations such as revisit time and cloud cover restrict the availability of daily data.

To overcome the limitations of models and observations, data assimilation has emerged as a promising approach because it capitalizes on the strengths of both observations and models, minimizing their respective uncertainties. It has proven useful in improving the accuracy of snow state estimates, snow physics, model parameters, and identifying sources of uncertainty. Data assimilation is particularly effective in harmonizing the different temporal and spatial resolutions of in-situ and remotely sensed observations and bridging the scale gap between observations and models. In general, the assimilation of satellite and airborne observations leads to enhanced estimates of seasonal snow and related variables (Fang et al., 2022; Margulis et al., 2016). However, a major shortcoming of snow data assimilation has its shortcomings, the largest one being that it cannot inability to provide daily high-resolution SWE (HR-SWE) for periods when no without satellite data are available. Key areas of ongoing research include understanding the impact of underlying spatial error correlations in data assimilation to improve the spatial estimates of SWE, and the potential integration of multiple observations to boost snow model accuracy. Despite these challenges, the snow science community continues to make strides in enhancing improve the accuracy of seasonal snow estimation (Giroto et al., 2020).

Acknowledging the complexities involved in modeling snow processes and assimilating diverse datasets, we propose a localized climate data downscaling method to estimate HR-SWE. Statistical downscaling methods have demonstrated their capability to act as a bridge between large-scale climate forecasts and local-scale climate impacts (Abatzoglou and Brown, 2012; Tabari et al., 2021; Rettie et al., 2023). One of the statistical downscaling methods is bias-correction spatial disaggregation (BCSD) (Wood et al., 2004), which effectively reduces uncertainties in climate model outputs by adjusting biases based on high-resolution observational data. These methods are particularly useful in non-mountainous regions, where data availability is typically higher. Their primary strength lies in their ability to correct model outputs while capturing local variability. However, their reliance on high-quality in-situ data significantly restricts their applicability in remote or data-scarce areas. In contrast, our method overcomes these limitations by utilizing low-resolution climate data without requiring ground-based observations, making it well-suited for a wider range of conditions, including regions with limited data availability.

Another widely used statistical downscaling method in climatology is based on a pattern known as the analog method (Zorita and Von Storch, 1999). These methods identify patterns in historical data that closely match the patterns simulated by atmosphere-ocean general circulation models. The observed surface climate conditions corresponding to these historical matches are then used as downscaled predictions. Analog methods have seen extensive application, as highlighted in studies such as those by Abatzoglou and Brown (2012), who demonstrated their effectiveness in wildfire assessments through the

95 multivariate adapted constructed analog, and outperformed traditional spatial downscaling methods. Similarly, Pons et al. (2010) utilized analog-based downscaling to analyze snow trends in Northern Spain, successfully replicating observed variability and trends for seasonal and climate change projections. The study by Caillouet et al. (2016) demonstrated the utility of probabilistic downscaling in reconstructing high-resolution precipitation and temperature fields over France, effectively addressing seasonal biases. The AtmoSwing software by Horton (2019) demonstrates the flexibility of analog methods for
100 operational forecasting and climate impact studies.

Additionally, an analog-type method, named nearest neighbor resampling, also used by Lall and Sharma (1996), relies on identifying patterns in historical point-based time series data and resampling them using a nearest-neighbor approach to preserve the serial dependence structure of the data. The k-nearest neighbors (k-NN) algorithm is a simple, non-parametric
105 machine learning technique commonly used for classification and regression (Cover and Hart, 1967). It works by identifying the 'k' most similar data points (neighbors) to a target data point based on a chosen distance metric, such as the Manhattan distance. The k-NN downscaling by Gangopadhyay et al. (2005) extends the analog method by weighting several similar historical analogs to create predictive ensembles, adding further flexibility to this approach. Building on this concept, Rajagopalan and Lall (1999) extended the methodology to multivariate weather simulations, incorporating variables such as
110 precipitation, temperature, and wind speed to simulate daily weather sequences. Later, Yates et al. (2003) used an adapted version of these methods to generate daily weather sequences and alternative climate scenarios. In the weather generator models based on k-NN, in general, the day directly succeeding the identified analog day is selected as the next day in the generated sequence, and this process continues iteratively (Gangopadhyay et al., 2005). Similarly, recent advancements, such as the study by Yiou and Déandréis (2019)–, have extended analog methods to ensemble-based probabilistic forecasts,
115 demonstrating skill in predicting variables like the NAO index and temperature measurements at European stations. These innovations highlight the adaptability and growing utility of analog-based and k-NN approaches in climate and hydrological modeling.

Recognizing the potential for snow spatiotemporal patterns to repeat ~~in years~~ on days with similar ~~climate~~ climatological characteristics (Zakeri and Mariethoz, 2024; Pflug and Lundquist, 2020), this study aims to provide daily, high-spatial-
120 resolution SWE data. Our proposed approach is based on establishing a statistical relationship between daily global low-resolution climate data, such as temperature, precipitation, low-resolution SWE (LR-SWE), and local reanalysis HR-SWE images as a training dataset. Then, based on this learned relationship, embedded in a k-~~nearest neighbor~~ NN algorithm, a unique daily HR-SWE dataset is obtained for the historical period (1950 to present) based on low-resolution climate data.

This method offers several key advancements over existing k-NN downscaling techniques. First, the adaptation made to the
125 k-NN downscaling method, specifically by introducing far and near temporal intervals of climate data, is highly flexible to dynamic variables undergoing significant changes due to climate variability. Second, unlike most analog methods that restrict analog candidates to a specific temporal window near the query date, this approach does not impose such limitations. This

flexibility is crucial for three reasons: (1) the inclusion of far and near temporal intervals makes such restrictions unnecessary, as the most suitable candidates are selected based on their match within the temporal window; (2) it is essential for preserving extreme events, as restricting candidates to a narrow date range risks losing matches that represent rare but important extreme events; and (3) it enables downscaling for future periods where exact analogs may not exist in the historical record within a specific date range. However, suitable analogs may still be found in historical observations but on different dates. For example, with climate change, a specific snow day in winter may no longer match the query day, but an analog might be found in another season, such as fall or spring, due to warmer climatic conditions.

Third, the method can reconstruct HR-SWE data for historical periods where only low-resolution climate data is available, providing valuable insights into past snow conditions. Additionally, the method excels at capturing fine-scale SWE patterns in complex terrains, such as mountainous regions, significantly improving upon traditional statistical downscaling models that often struggle in such environments. Finally, our method does not require high-resolution climate input data. This substantially reduces computational demands while maintaining the high quality of the downscaled SWE data.

The paper is structured as follows: Section 2 details the methodology; Section 3 describes the study area, database, and parameters; Sections 4 and 5 present the evaluation approaches and the results; and ~~Section 5 concludes~~ Sections 6 and 7 ~~conclude~~ with discussions and conclusions.

2. Methodology

2.1 Overview of the Algorithm

The algorithm involves two primary datasets: ~~the training dataset and the target dataset~~. The training dataset consists of HR-SWE images ~~and~~, low-resolution (LR) climate data, ~~and LR-SWE~~. In contrast, the target dataset includes LR-SWE images alongside LR climate data, for which we aim to estimate HR-SWE images.

The fundamental strategy of the proposed method involves ranking the ~~data in the training dataset~~ data to estimate an HR-SWE image for a given target date. This ranking is based on the Manhattan distance between each date in the training dataset and the selected target date, as detailed in Section 2.3. The Manhattan distance is chosen as it is more robust against outliers and often computationally ~~cheaper~~ less expensive than the Euclidean distance. Then, the algorithm estimates the downscaled SWE for the target date based on the K-nearest SWE candidates. As detailed in the recent work by Zakeri and Mariethoz (2024), this approach was initially designed to create synthetic satellite snow cover images for dates with no satellite data, based on the relationship between meteorological estimators from the ERA5-Land reanalysis dataset and available clear sky Landsat/Sentinel-2 snow cover images. Here, we have adapted this approach to downscale SWE. The innovation lies in the focus on downscaling the LR daily SWE data at the spatial scale of climate model simulations to produce much higher-resolution daily SWE estimates. This output is particularly important in areas with complex terrain, such as the Western United

States, where global models face challenges in accurately representing the regional climate. Indeed, the resolution of global climate simulations is often insufficient for capturing local influences (e.g., topography or vegetation) on SWE patterns. On the other hand, physical models are computationally expensive. Therefore, developing a tool to estimate HR-SWE that closely aligns with established regional HR-SWE reanalysis is essential for scientific and management purposes, offering a faster and less computationally demanding solution. Further details of the proposed methodology ~~will be~~are provided in the subsequent sections.

2.2 Input Data and Preprocessing

2.2.1 Definition of Temporal Intervals for Climate Variables

SWE, representing the amount of water stored in the snowpack, is influenced by various factors, which can be classified into two main categories: climatic variables (such as temperature, precipitation, and surface downwelling shortwave radiation) and environmental variables (including land cover, topography, and the presence of topography shadows). While environmental variables can impact SWE distribution and accumulation, their effects are assumed to be consistent and not subject to significant temporal variations within the specified regions. ~~Therefore, for downscaled SWE estimation, we primarily focus on climatic variables and do not explicitly incorporate topography and land cover information.~~Although the effects of environmental variables like terrain shading may vary, driven by the sun's position throughout the year, these can be captured in the baseline SWE models used to generate training data. In contrast, climatic variables exhibit significant temporal and spatial variability, making them the primary drivers of SWE dynamics. Consequently, the downscaled SWE estimation focuses primarily on climatic variables. Their dominant influence ensures that the proposed model can capture the spatiotemporal variability essential for SWE estimation. The effects of environmental variables, while excluded from the direct downscaling process, can be accounted for in the baseline SWE models that provide the training data.

SWE is also affected by preceding meteorological conditions, such as the temperature and precipitation patterns of previous days. For instance, the amount of SWE today may vary depending on the conditions experienced in the preceding ~~periods~~days. To represent this, we introduce two distinct climate temporal intervals: a near interval and a far interval (as shown in Fig. 1~~Fig. 1refer to Fig. 1).~~). These intervals consider climate variables ~~such as~~including minimum temperature, maximum temperature, precipitation, and surface downwelling shortwave radiation. Considering both far and near-term variations allows accounting for complex relationships between the climate dynamics and snow accumulation and melting. By incorporating these intervals, we can capture the influence of climate variables over different timescales and ~~improve the accuracy of~~enhance SWE estimation accuracy. The specific lengths of these ~~far~~near and ~~near~~far climate intervals are determined through an optimization procedure outlined in Section 2.4.1.




Fig. 1. Illustration of the definition of Near (shown in gray) and Far (shown in white) daily intervals for climate variables, highlighting the Far Temporal Interval (FI) and Near Temporal Interval (NI) of Climate Data at a Specific Time (C_t).

2.2.2 Input Climate Variables

The downscaling process relies on low-resolution climate information and LR-SWE data obtained from global or regional climate models. As a result, the ~~SWE~~ downscaling procedure can be described by Equation (1)(1)(1):

$$\widehat{SWE}_{HR}(t) \approx f(T_{min,LR}^{FI}, T_{min,LR}^{NI}, T_{max,LR}^{FI}, T_{max,LR}^{NI}, P_{LR}^{FI}, P_{LR}^{NI}, RSDS_{LR}^{FI}, RSDS_{LR}^{NI}, SWE_{LR}(t)) \quad (1)$$

190

Here, $\widehat{SWE}_{HR}(t)$ and $SWE_{LR}(t)$ are the downscaled SWE and the LR-SWE, both at the query time (t), and T_{min} and T_{max} represent the minimum and maximum temperature, while P and $RSDS$ denote precipitation and surface downwelling shortwave radiation. The superscript " FI " or " NI " indicates the far or near temporal intervals ~~that are~~ introduced in Fig. 1 Fig. 1 Fig. 1. The subscript notation (" LR ", " HR ") explicitly indicates the resolution (Low Resolution, High Resolution). All climate variables (T_{min} , T_{max} , P , $RSDS$, SWE) are daily measurements. To ensure compatibility among datasets with varying units, all input data are rescaled to a range of 0-1 based on the absolute minimum and maximum values observed in the training dates. Hereafter, ~~SWE refer to downscaled SWE.~~ refers to downscaled SWE. Details on the experiments conducted to identify the most effective meteorological predictors within these datasets are available in the Supplementary (Table S1 and Fig. S1).

195

2.3 Downscaled SWE Estimation

200

Estimating \widehat{SWE} for a date when ~~HR-SWE data~~ is not available relies on the existing HR-SWE data from the training dates. The training dates ~~consist of~~include the input variables described in Equation (1)(1)(1) and their corresponding HR-SWE images. By utilizing a vector of inputs, we estimate \widehat{SWE} by selecting the K-nearest candidates in the input space. ~~The downscaling process follows these steps:~~

225 In this study, the k-NN algorithm is applied to downscale low-resolution climate data to HR-SWE estimates by selecting learning days with similar climate conditions. The flowchart (Fig. 2) illustrates the proposed downscaling method for estimating HR-SWE, and the “steps” are as follows:

1. Gather the input variables, including the far and near intervals of temperature, precipitation, shortwave radiation, and the LR-SWE for both the target date and the training dates.
- 230 2. Calculate the ~~similarity or~~ distance between the input vector of the target date and the input vectors of the training dates.
3. Select the K-nearest training dates based on their proximity to the target date in the input space.
4. Retrieve the corresponding HR-SWE images associated with the selected K-nearest training dates.
5. Aggregate the retrieved HR-SWE images to estimate \hat{SWE} for the target date.

235 The metric used to identify the nearest neighbors between a given target date (t_1) and a training date (t_2) is the multivariate Manhattan distance, described in Equation (2)(2)(2):

$$d\{t_1, t_2\} = \sum_{i=1}^E \alpha_i |x_i(t_2) - x_i(t_1)| \quad (2)$$

The weights α_i assigned to the input variables ~~add up sum~~ to 1 and are used to balance their influence in the selection of the K-nearest observations. The determination of optimal weights is further discussed in Section 2.4.2. Here, x_i represents each of the E estimators introduced in Equation (1)(1)Here, x_i represent each of the E estimators introduced in Equation (1). Unlike the Euclidean distance, which calculates the shortest straight-line distance, the Manhattan distance computes the sum of the absolute differences between variables. This makes it more robust against outliers and better suited to handling high-dimensional datasets, such as those containing multiple climate variables. In our method, this distance is used to rank the training dates based on their similarity to the target date.

245




Fig. 2. Flowchart depicting the proposed downscaling algorithm for estimating high-resolution Snow Water Equivalent (HR-SWE) on a target date using low-resolution (LR) climate data: 1. Gather Input Variables: Collect the input variables, including Near (NI) and Far (FI) temporal intervals of climate data (e.g., temperature, precipitation) for both the target date and the training dataset. 2. Calculate Distances: Compute the distance (d) between the input vector of the target date and the input vectors of each candidate in the training dataset using a defined distance metric (Manhattan distance). 3. Identify k-Nearest Neighbors (k-NNs): Rank the training data in order of increasing distances to identify the k-nearest neighbors. 4. Aggregate HR-SWE Images: Retrieve the HR-SWE images corresponding to the k-NNs and aggregate them to estimate the HR-SWE for the target date.

2.4 Estimating Parameters and Weights

To optimize parameter estimation and ensure better convergence, the parameters are estimated either using sensitivity analysis or an optimization algorithm. In this paper, the determination of the far and near intervals (FI and NI) for climate variables (Fig. 1 Fig. 1 (Fig. 1)), as well as the number of K-nearest observations, are achieved through a sensitivity analysis.

250 The weights assigned to the input features are established using an optimization algorithm.

2.4.1 Parameter Sensitivity Analysis

The measure of Dissimilarity, denoted as ε , is critical for determining the optimal near (NI) and far (FI) temporal intervals, as well as the optimal number of nearest observations (K), for estimating \hat{SWE} . This dissimilarity is quantified using the Root-Mean-Squared Error (RMSE), a standard measure of accuracy that reflects the average magnitude of the square of errors between estimated and observed values. The RMSE is defined as follows:

$$RMSE = \sqrt{\frac{\sum_{i=1}^N (\hat{SWE}_i - SWE_i)^2}{N}} \quad (3)$$

where \hat{SWE}_i is the downscaled SWE at the i^{th} pixel, SWE_i is the ~~observed (reference)~~ HR-SWE at the i^{th} pixel, and N is the total number of pixels. This formula underpins the evaluation of ε across different configurations of temporal intervals and K-nearest observation counts.

To identify the optimal temporal intervals (NI^* and FI^*), we conduct a sensitivity analysis within predefined ranges, set by $NI \in [1, 7]$ ~~days~~ and $FI \in [1, 90]$ ~~days~~, aiming to minimize ε . Similarly, the optimal number of nearest observations (K^*) is determined by evaluating ε as a function of K, within a range 1 to the maximum number of available training dates ($K \in [1, K_{max}]$) to identify the configuration that yields the smallest dissimilarity between \hat{SWE} and reference HR-SWE. The maximum number of available training dates (K_{max}) can vary for each case study from ~~several~~ a few days to several months.

Mathematically, the optimization processes are represented as:

$$\begin{aligned} NI^* &= \underset{NI}{\operatorname{argmin}} \varepsilon_{NI} \\ FI^* &= \underset{FI}{\operatorname{argmin}} \varepsilon_{FI} \\ K^* &= \underset{K}{\operatorname{argmin}} \varepsilon_K \end{aligned} \quad (4)$$

where ε_{FI} , ε_{NI} , and ε_K are the sum of RMSEs for respective configurations, reflecting the dissimilarity between \hat{SWE} (using different sizes of NI, FI, and K) and the reference HR-SWE. The only distinction between $RMSE_{FI}$, $RMSE_{NI}$, and $RMSE_K$ is based on the varying elements (FI, NI, or K), illustrating how each influences the \hat{SWE} estimate. The determination of these optimal parameters enables the refinement of our SWE estimation model, enhancing its accuracy by minimizing ε .

The parameters are selected through a sensitivity analysis to minimize ε , but they do not necessarily correspond a global optimum. This means that while they perform well in the studied regions, they may not generalize well to other locations or climatic conditions. Therefore, we recommend performing a sensitivity analysis for each region. Additionally, the sensitivity analysis assumes that the influence of these intervals is consistent across different temporal scales, which may not always be valid, particularly in regions with highly variable climate patterns. Despite these assumptions, the chosen parameters strike a

290 balance between computational efficiency and accuracy for the downscaling task. Moreover, the subsequent weight optimization further mitigates the impact of non-global optimal parameter selection.


2.4.2 Weights Optimization

295 The weights α_i defined in Equation ~~(2)(2)(2)~~ are determined by minimizing the RMSE between the SWE and the reference HR-SWE using Bayesian optimization. In this model-based optimization, the global maximum or minimum of an unknown objective function is found sequentially (Shahriari et al., 2015; Snoek et al., 2012). The key aspect of this approach is the use of a probabilistic model of the response function, which is evaluated at a minimal cost through the acquisition function. The Bayesian optimization framework employs a Gaussian process prior over the objective function and iteratively refines the model through Bayesian posterior updating to determine the optimal solution.

3. Study Areas, Datasets, and Parameters

300 3.1 Study Areas

To evaluate the effectiveness of the proposed methodology, we selected two regions in the Western United States as indicated in ~~Figure 2~~Fig. 3: the California Sierra Nevada region (referred to as California) and the Upper Colorado River Basin (referred to as Colorado). These regions are heavily rely dependent on snowmelt as a vital source of water resources (Dawadi and Ahmad, 2012; Siirila-Woodburn et al., 2021a).



(a) Western U.S.




Fig. 3. a) Map of elevation (in meters) across the Western United States, highlighting two regions of interest (red squares). b) Map of elevation and locations of in-situ sites in the Sierra Nevada, California. c) Map of elevation and locations of in-situ sites in the Upper Colorado River Basin, Colorado.

In the California region, elevation ranges from a minimum of 1200 m to a maximum of 3700 m, with an average elevation of 2200 m. In the Colorado region, elevation ranges from a minimum of 1900 m to a maximum of 4300 m, with an average elevation of 3000 m. By focusing on these specific regions, we can assess the performance and applicability of the proposed methodology in other areas where snow water resources are an important component of the hydrology system.

The California Sierra Nevada snowpack plays a critical role in water resource management, contributing approximately 30% of the state's water supply through snowmelt. However, its sensitivity to warming temperatures is evident, with the 2015 SWE on April 1 dropping to just 5% of historical averages. This dramatic decline underscores the combined effects of reduced precipitation and higher temperatures, which exacerbate drought severity and shift the timing of water availability (Belmecheri et al., 2016).

In the Upper Colorado River Basin, snowmelt accounts for 70–90% of annual streamflow, making snowpack dynamics essential to hydrological processes and water management. Heldmyer et al. (2023) identify three distinct snow drought types: 'warm', 'dry', and 'warm-and-dry', which differentially impact SWE and streamflow timing. Warm droughts tend to reduce SWE at lower elevations, while dry conditions cause uniform SWE reductions across elevations. These droughts advance peak streamflow timing by 7–13 days, emphasizing the region's sensitivity to climatic changes in temperature and precipitation (Heldmyer et al., 2023).

3.2 Datasets

The reference ~~high-resolution HR~~-SWE data, which this study aims to produce, ~~is sourced~~ comes from the reanalyzed SWE dataset for the Western United States (Fang et al., 2022). This dataset, captured at a 16 arcsecond (~500 m) resolution, covers the period from the water year (i.e., 1st October to 30th September) 1984 to 2021 and is updated daily. It combines high-resolution remotely-sensed data with a Bayesian data assimilation (Margulis et al., 2019; Margulis et al., 2015; Margulis et al., 2016) framework. The dataset is derived from Landsat-based fractional snow-covered area observations, updated daily, and incorporates a land surface model to estimate SWE and snow depth. This approach enables spatially and temporally continuous SWE estimates, which are verified against in situ and lidar-derived SWE measurements for accuracy. It offers ensemble median values of SWE, calculated from a discrete probability distribution of posterior weights, thus providing an extensive and detailed view of the snow water content over the years. The dataset can be accessed at <https://doi.org/10.5067/PP7T2GBI52I2>.

Hereafter, we refer to this dataset as the UCLA SWE.

To obtain the necessary climatic estimators, ~~such as~~ including daily minimum and maximum air temperature, daily total precipitation, and shortwave radiation, we utilize two distinct sources: the Coupled Model Intercomparison Project (CMIP) version 6 simulations (Meehl et al., 2014) and the downscaled CMIP6 over the Western United States using the Weather Research and Forecasting (WRF) model datasets (referred to as WRF-CMIP6) (Rahimi et al., 2024; Rahimi et al., 2022). The CMIP6 simulations provide daily climate data at a spatial resolution of 100 km, while the WRF-CMIP6 datasets offer high-resolution daily climate simulations at a resolution of 9 km for the Western United States.

Rahimi et al. (2024) created climate data utilizing reanalysis data based on dynamical downscaling to refine global climate models through regional climate models, specifically using the WRF model. They used ERA5 reanalysis data for calibrating and validating the WRF model setup. This method effectively enhances the representation of local weather phenomena by increasing resolution, thereby improving the accuracy of climate projections, particularly in areas with complex terrain like the Western United States.

While CMIP6 models capture long-term climate variability and trends, including seasonality that might influence snow patterns, their resolution is relatively coarse. In other words, they simulate the general variability and stochastic nature of the climate. However, WRF-CMIP6 incorporates ERA5 reanalysis data ~~in the process of to~~ dynamically ~~downscaling~~ downscale

CMIP6 [dataoutputs](#), using a configuration that [also considers temporal and spatial](#) [enhances the representation of local climate](#) variability [in the selection of global climate models for specific to](#) the Western United States. This [process](#) results in more accurate regional climate models. Utilizing both CMIP6 and WRF-CMIP6 at different resolutions enables us to investigate the impact of using a regional configuration of a climate model that dynamically downscales CMIP6 data, incorporating reanalysis data, alongside a global climate model with different spatial resolutions [on, in](#) the accuracy of our approach.

To illustrate the effect of resolution on [the SWE image](#) accuracy [of the SWE image](#), Fig. [34](#) depicts the UCLA SWE, WRF-CMIP6, and CMIP6 images for a day in the California region.




Fig. [4](#). Comparative illustration of SWE images highlighting the effect of spatial resolution on accuracy. Displayed from left to right are the UCLA SWE, WRF-CMIP6, and CMIP6 images captured over the California region (Fig. [23](#) (b), lon: [-120](#) to [-119](#), lat: 38 to 39) on January 8, 1995. [The](#) Distinct variations between these images underscore the influence of scaling on image accuracy.

Among the available models with resolutions of both 9 km and 100 km, we have selected three models for testing our methodology ([Table 1](#) [Table 1](#) ([Table 1](#))). These models have been utilized in other studies ([Thrasher et al., 2022](#); [Kouki et al., 2022](#)); Although our methodology is not limited to these models, they were selected merely as examples. Hereafter, we will

refer to them as 'cnrm-esm2-1', 'ec-earth3-veg', and 'mpi-esm1-2'.

Table 1. The climate models that are used in this study.

Climate model	Developer	100 km resolution	9 km resolution
cnrm-esm2-1	The Earth system model of CNRM of 2nd generation was developed by the CNRM/CERFACS.		
ec-earth3-veg	A configuration extending the Earth System Model EC-Earth3 developed by the EC-Earth Consortium.	CMIP6 data: https://esgf-node.ipsl.upmc.fr/search/cmip6-ipsi/ ;	WRF-CMIP6 data: https://registry.opendata.aws/wrf-cmip6/ ;

mpi-esm1-2-1r	Part of the family of Earth System Models (ESMs) developed by the Max Planck Institute (MPI).	Historical period (1985-2014) downloaded in January 2023.	Historical period (1985-2014) downloaded in January 2023.
----------------------	---	---	---

To ensure that the test ~~dates are~~period is not influenced by the training ~~dates~~period, the periods 1985-2004 and 2011-2014 are designated ~~as~~for training dates, while the years 2005 to 2010 are allocated for testing. Additionally, to confirm that our testing dates are not grouped into nearly identical wet or dry categories, we employed the CONUS Drought Indices dataset (Abatzoglou, 2013). This ~~dataset~~ includes drought indices derived from the 4 -km daily Gridded Surface Meteorological (GRIDMET) dataset. The Standardized Precipitation Evapotranspiration Index (SPEI) is the specific drought index used in our analysis. We calculated the average aggregated SPEI for each year across each region to categorize the testing and training years in Supplementary Table ~~S1. As is clear,~~S2. The testing years are different in terms of the drought index.

3.3 Parameters for Both Case Studies

Through a sensitivity analysis, we determine the NI^* and FI^* using Equation ~~(4)(4)(4).~~ This analysis is carried out for both datasets: CMIP6 (100 km) and WRF-CMIP6 (9 km). The results of this analysis are presented in supplementary Figures ~~S1S2~~ to ~~S4S5~~, indicating that a FI^* of 60 days and a NI^* of 4 days are sufficient for accurate SWE estimation in all examined scenarios. Even though these values might not represent a global optimum, potential suboptimal effects are deemed negligible. Furthermore, these effects are addressed in subsequent optimization stages when determining weights.

Next, we conduct a sensitivity analysis for the parameter K^* , the number of nearest observations, using the methodology described in Section 2.4.1 and Equation ~~(4)(4)(4).~~ Fig. ~~S5S6~~ and Fig. ~~S6S7~~ illustrate that, on average, a value of $K^*=130$ is appropriate for both the California and Colorado regions. Considering the relatively small variations in accuracy, we decided not to optimize K for each dataset. Note that the value of 130 nearest observations is high. This suggests the presence of frequently repeating situations or patterns within the ~~dataset~~data. This is advantageous for statistical analysis, as it provides a ~~substantial volume~~large number of data points, thereby enhancing the robustness and reliability of our statistical inferences. The optimized weights α obtained from the Bayesian optimization algorithm after 500 iterations are summarized in ~~Table 2~~Table 2. ~~Both regions exhibit~~ similar weight patterns ~~are observed in both regions. In general,~~ with relatively high weights ~~are~~ assigned to variables $T_{min,LR}^{NI}$, P_{LR}^{NI} , SWE_{LR} , and P_{LR}^{FI} .

Table 2. Weight optimization (α , in Equation ~~(2)(2)(2))~~ results using Bayesian optimization. The numbers greater than 0.1 are ~~bolded~~highlighted in bold. This value, 0.1, is chosen because it almost represents the scenario where weights are distributed equally. cnrm refers to 'cnrm-esm2-1', earth3 to 'ec-earth3-veg', and mpi to 'mpi-esm1-2'.

Region of Interests		California						Colorado					
Climate Model		cnrm		earth3		mpi		cnrm		earth3		mpi	
Climate resolution (km)		100	9	100	9	100	9	100	9	100	9	100	9

$T_{max,LR}^{NI}$	0.03	0.08	0.24	0.12	0.12	0.17	0.03	0.02	0.01	0.21	0.01	0.24
$T_{min,LR}^{NI}$	0.16	0.08	0.13	0.03	0.08	0.10	0.10	0.13	0.18	0.35	0.18	0.15
P_{LR}^{NI}	0.27	0.04	0.15	0.21	0.15	0.15	0.24	0.14	0.32	0.04	0.33	0.09
$RSDS_{LR}^{NI}$	0.03	0.05	0.04	0.13	0.04	0.02	0.25	0	0	0.01	0	0
$T_{max,LR}^{FI}$	0.03	0.03	0.15	0.02	0.27	0.17	0	0.02	0.01	0.01	0.01	0.08
$T_{min,LR}^{FI}$	0.03	0.05	0.04	0.03	0.03	0.18	0	0.31	0	0.01	0	0.33
P_{LR}^{FI}	0.27	0.31	0.03	0.19	0.03	0.13	0.01	0.36	0.33	0.26	0.32	0
$RSDS_{LR}^{FI}$	0.03	0.04	0.11	0.20	0.03	0.02	0.30	0	0	0	0	0
SWE_{LR}	0.14	0.32	0.11	0.07	0.25	0.07	0.07	0	0.15	0.10	0.15	0.11

From a physical perspective, this weighting aligns with the processes governing SWE distribution and dynamics. Minimum temperature ($T_{min,LR}^{NI}$) significantly influences freezing and melting thresholds, which are critical for snowpack accumulation or reduction. Precipitation variables, both near and far intervals (P_{LR}^{NI} , P_{LR}^{FI}), directly contribute to SWE through their impact on the volume of snowfall. Meanwhile, the inclusion of SWE_{LR} as a highly weighted variable underscores its importance role as a baseline indicator of existing snowpack conditions.

4. Evaluation Approach

Our method is evaluated through various accuracy metrics and independent datasets. Firstly, we generate two quality metrics in conjunction with the \hat{SWE} images. These metrics provide a quick assessment of the quality of the \hat{SWE} products. For each target date, two quality metrics are provided: the average and the standard deviation of the similarity metric across the k best-selected SWE candidates. The lower the average and standard deviation of the similarity metric, the better the \hat{SWE} estimation.

4.1 Time-Series Visualization

To further demonstrate the efficacy of our method, we showcase some SWE time series spanning six years at in-situ locations that are obtained from the Snow Telemetry (SNOTEL) network (<https://www.wcc.nrcs.usda.gov/snow/>). First, the pixels nearest to the in-situ locations are identified. Then, the SWE values at these locations in both the UCLA SWE data and other well-established SWE datasets (explained in Section 4.3) in the Western United States are compared with the \hat{SWE} values at these locations. We also present scatter plots colored by density (density scatter plots) illustrating the correlation between reference (UCLA SWE) and \hat{SWE} values at in-situ locations. As in Fang et al. (2022)’s study, SWE values below 10 mm are excluded from the comparison. This screening is done because very small SWE values (less than 10 mm) can be the result of measurement errors, minimal snow presence, or other factors that might not provide meaningful data insights.

While the SNOTEL network provided the spatial locations for these comparisons, we did not use direct in-situ SNOTEL SWE measurements in this analysis. The UCLA SWE dataset has been previously validated against in-situ observations, including SNOTEL, and has been widely recognized as a benchmark for SWE validation in subsequent studies. For example, Ma et al. (2023) used the UCLA SWE dataset as a reference to assess the performance of machine learning approaches for estimating spatiotemporally continuous SWE. Similarly, Fang et al. (2023) used the UCLA SWE dataset as a baseline for evaluating snow water storage uncertainty. These applications highlight the dataset's reliability and its critical role in advancing snow hydrology research.

4.2 Cross-validation

We also validate our results using cross-validation. This involves removing six years of the UCLA SWE from the training dataset. Then, We estimate \hat{SWE} based on the remaining training data and compare the results with the reference UCLA SWE image using correlation, mean difference, and RMSE as evaluation criteria. These assessment metrics are summarized in Table 3.

Table 3. A summary of accuracy assessment criteria.

Mean difference	A lower value indicates a better estimation	$MD = \text{Mean}(SWE - \hat{SWE})$
Correlation	A higher value indicates a better estimation	$R = \frac{\sum_{i=1}^N (\hat{SWE}_i - \overline{\hat{SWE}})(SWE_i - \overline{SWE})}{\sqrt{(\sum_{i=1}^N (\hat{SWE}_i - \overline{\hat{SWE}})^2)(\sum_{i=1}^N (SWE_i - \overline{SWE})^2)}}$
RMSE	A lower value indicates a better estimation	Equation (3)

where \hat{SWE}_i is the downscaled SWE at the i th pixel, SWE_i is the observed (reference) SWE at the i th pixel, and N is the total number of pixels.

In our quantitative comparison, we calculate the accuracy metrics at different elevation bands (elevation < 2000 m, 2000 m < elevation < 3000 m, and elevation > 3000 m) and across various land cover types, including forest and non-forest areas. We utilize the MODIS Land Cover Type (MCD12Q1) Version 6.1 data for land cover information (https://lpdaac.usgs.gov). This data, as of 2018, is maintained by the NASA EOSDIS Land Processes Distributed Active Archive Center (LP DAAC) at the USGS Earth Resources Observation and Science (EROS) Center in Sioux Falls, South Dakota. Elevations are derived from the NASA SRTM Digital Elevation 30 m data (Farr et al., 2007).

4.3 Comparison with Independent SWE Datasets

450 To enhance the reliability of the SWE, additional verification is performed through comparisons with the three following independent SWE datasets. These comparisons are essential for validating the accuracy and consistency of our SWE images, using the rigorous evaluation criteria detailed in Table [3S3](#).

SNODAS 1 km SWE Product

455 The first independent dataset includes the 1 km SWE data from the Snow Data Assimilation System (SNODAS), developed by the NOAA National Weather Service's National Operational Hydrologic Remote Sensing Center (NOHRSC) ([Center, 2004](#)). SNODAS is a comprehensive modeling and data assimilation system designed to provide estimates of snow cover; ~~which~~ [that](#) are crucial for hydrologic modeling and analysis. The dataset covers the period from September 28, 2003, to the present.

460 We first resample the SWE images by aggregating the values of all pixels that intersect each 1 km SNODAS pixel. This ensures that both datasets have the same spatial resolution. Subsequently, we compare the SNODAS SWE products with the SWE images using the evaluation criteria introduced in Table [4S3](#). SWE pixel values less than ~~4 cm~~ [10 mm](#) are not included in the comparison.

Daymet 1 km SWE Product

The second independent dataset includes the 1 km SWE data from the Daymet V4 dataset: Daily Surface Weather and Climatological Summaries ([Thornton et al., 2022](#)). Daymet provides gridded estimates of daily weather parameters for North America, including minimum and maximum temperatures, precipitation, and SWE, with data spanning from January 1, 1980. We follow a similar procedure as in the comparison with the SNODAS SWE products.

470

University of Arizona (UA) 4 km SWE Product

The first independent dataset is the 4 km SWE data provided by the University of Arizona (UA) ([Broxton et al., 2019](#); [Zeng et al., 2018](#)). It assimilates in-situ snow measurements from the SNOTEL and Cooperative Observer Network (COOP) networks ([Fleming et al., 2023](#); [Council and Committee, 1998](#)), incorporating detailed ~~snow data~~ [snowpack measurements](#). It
475 also integrates modeled, gridded temperature and precipitation data, ~~ensuring~~ [providing](#) a comprehensive representation of SWE over the conterminous United States since 1981. For this comparison, we employ a similar procedure ~~as~~ [to that used](#) in the comparisons with the previous two independent SWE products.

5. Results

5.1 Time-Series Visualizations

480 To showcase the effectiveness of the proposed technique, SWE time series that encompass a six-year duration at the three
highest-elevation in-situ locations (see [Table 3](#)~~Table 3~~[Table 4](#)) in California are presented in Fig. [4.5](#) (and Supplementary Fig.
[S7](#)[S8](#) for Colorado). These time series visually represent the SWE values over time, enabling a comparison between the SWE
values, and the UCLA SWE, Daymet, SNODAS, and UA SWE data. The red line, representing our ~~estimation~~[estimated values](#),
generally tracks the blue line (UCLA SWE) well across the six-year period for the three high-elevation locations. However, in
485 some years and locations, such as at the 'Leavitt Lake' and 'Lobdell Lake' stations in 2005, [where](#) the red line diverges slightly
from the blue line, indicating variability in ~~estimation~~[the accuracy of the estimates](#) across different time periods. Nevertheless,
even in 2005, our estimation is still within the range of other well-established SWE estimations. Moreover, in 2005, the other
datasets ~~also~~ do not [agree](#)[align](#) on the amount of SWE, indicating that this year is intrinsically challenging. This demonstrates
that the proposed method can closely estimate the UCLA SWE ~~at~~[on](#) a point scale in most years.

490

495

Table 3. Names and locations of in-situ sites.

California				Colorado			
Station Name	Elevation (m)	Lat	Lon	Station Name	Elevation (m)	Lat	Lon
1. 'Blue Lakes'	2438	38.61	-119.92	1. 'Beaver Ck Village'	2612	39.60	-106.51
2. 'Burnside Lake'	2478	38.72	-119.89	2. 'Buckskin Joe'	3404	39.30	-106.11
3. 'Carson Pass'	2547	38.69	-120.00	3. 'Copper Mountain'	3207	39.49	-106.17
4. 'Ebbetts Pass'	2639	38.55	-119.80	4. 'Hoosier Pass'	3539	39.36	-106.06
5. 'Forestdale Creek'	2441	38.68	-119.96	5. 'Independence Pass'	3230	39.08	-106.61
6. 'Hagans Meadow'	2360	38.85	-119.94	6. 'Middle Fork Camp'	2734	39.80	-106.03
7. 'Heavenly Valley'	2601	38.92	-119.92	7. 'Nast Lake'	2660	39.30	-106.61
8. 'Horse Meadow'	2607	38.84	-119.89	8. 'Rough And Tumble'	3179	39.03	-106.08
9. 'Leavitt Lake'	2927	38.28	-119.61	9. 'Summit Ranch'	2856	39.72	-106.16
10. 'Leavitt Meadows'	2195	38.30	-119.55	10. 'Vail Mountain'	3142	39.62	-106.38
11. 'Lobdell Lake'	2820	38.44	-119.37				
12. 'Monitor Pass'	2533	38.67	-119.61				





Fig. 5. The SWE time series spans a six-year period at the three highest ~~elevations-elevation~~ in-situ locations (Table 3 Table 3 (Table 4)) in California. "UA" represents the University of Arizona SWE datasets. The downscaled SWE (\hat{SWE}) results were obtained from the 'ec-earth3-veg' climate data at a 9 km resolution.

Additionally, density scatter plots are obtained using the 'dscatter' function (Henson, 2024) and are illustrated in Fig. 56 for California (and Supplementary Fig. S8S9 for Colorado), depicting the relationship between UCLA SWE and \hat{SWE} values at in-situ locations (see Table 3 Table 3 (Table 4)) obtained from three climate models as estimators: 'ec-earth3-veg', 'mpi-esm1-2', and 'cnrm-esm2-1' at both 9 km and 100 km resolution.

Referencing Fig. 56, using 'ec-earth3-veg' at a 9 km resolution as an estimator, the derived \hat{SWE} in California shows a stronger correlation with the UCLA SWE at in-situ locations, achieving an R-value of approximately 0.87. This model also demonstrates improved accuracy, with a lower RMSE and MD (approximately 0.14 (m) for RMSE and about 0.02 (m) for MD). In contrast, as shown in Supplementary Fig. S8S9, using 'cnrm-esm2-1' at a 100 km resolution as an estimator, the derived \hat{SWE} in Colorado exhibits a stronger correlation with the UCLA SWE at in-situ locations, marked by an R-value close to 0.85, and the RMSE estimated at around 0.09 m.






Fig. 6. A density scatter plot in California from 2005 to 2010 compares in-situ data locations (Table 3) from UCLA SWE images to the downscaled SWE (\hat{SWE}) values. The solid black line represents the 1:1 line. The displayed metrics include the correlation coefficient (R), mean difference (MD), and Root-Mean-Squared Error ($RMSE$). Only data with SWE values greater than 4 mm are included in the comparison.

Fig. 67 presents a comparison of the mean \hat{SWE} and mean UCLA SWE over a six-year period in California and Colorado. Among the models, the 'cnrm-esm2-1' at 100 km resolution, in most months, outperforms 'ec-earth3-veg' and 'mpi-esm1-2' in Colorado. In contrast, in California, 'ec-earth3-veg' mostly exhibits superior performance at both 9 km and 100 km resolutions. However, all the models underestimate SWE in California, suggesting that testing other climate models for this region may be beneficial. Collectively, these findings emphasize the significance of selecting an appropriate model and its optimal resolution for SWE estimation.




Fig. 7. The average of the UCLA SWE (black dotted line, reference) and the downscaled SWEs (SWE; blue and green lines) for each area over the six-year period from 2005 to 2010. For each area, the three climate models used in the downscaling are represented by a unique color: light blue for 'ec-earth3-veg', green for 'cnrm-esm2-1', and dark blue for 'mpi-esm1-2'. Within each color group, line styles indicate the spatial resolution of the climate data: solid lines correspond to results based on a 9 km climate estimators resolution, and dashed lines represent results based on a 100 km climate estimators resolution. The dotted black line represents the average UCLA SWE, which is the reference.

Additionally, two videos (one for each study region) have been created to showcase a side-by-side comparison of UCLA SWE and SWE images for corresponding dates: California (SWE obtained from 'ec-earth3-veg' at 100 km), Colorado (SWE obtained from 'cnrm-esm2-1' at 9 km).

Several factors explain why certain models outperform others in specific locations. While a detailed comparison between model performances across various regions is beyond the scope of this study, other studies have explored this area. For example, Kouki et al. (2022) evaluated the ability of CMIP6 models to estimate SWE across the Northern Hemisphere and found that different models perform variably in specific regions based on their ability to simulate particular climatic and geographical conditions. In terms of the contribution of temperature and precipitation to SWE biases, precipitation plays a more dominant role, especially in winter. However, temperature becomes more significant during spring, when snowmelt occurs. In regions where temperatures are closer to 0°C, biases in temperature can substantially affect snowmelt. Overall, the results underscore that precipitation is the primary driver of SWE biases in winter. However, temperature plays a crucial role during the snowmelt season in spring, particularly in regions with more temperate climates, such as the southern parts of North America and Europe.

5.2 Cross-validation

Cross-validation accuracy assessments of \hat{SWE} are outlined in Supplementary Tables S2S4 and S3. Fig. 7S5. Figs. 8 and 89, and Figs. S9S10 and S10S11 in the supplementary material further present these assessments considering variations in elevation and land cover for California and Colorado. Overall, for most months, using climate data with a higher resolution (9 km) as an estimator yields only slightly higher or identical accuracies compared to climate data with a lower resolution (100 km). In a broad comparison across resolutions in Colorado and California, the 'cnrm-esm2-1' and 'ec-earth3-veg' models generally exhibit the best performance. Nonetheless, it is crucial to recognize the variability in model performance across different regions. For instance, in both California and Colorado, all models generally exhibit perform better or close performance comparably at higher elevations (elevation > 3000 m) compared to medium elevations (2000 m < elevation < 3000 m). Although errors at lower elevations (elevation < 2000 m) are less fewer than at higher elevations, the number of pixels with elevations less than 2000 m is also fewer smaller. This is significant because accurately obtaining snow data at higher elevations is more critical than at other elevations in certain studies and regions, such as. For example, in water management for areas where, in summer, most of the summer water supply originates from higher elevations.

In Colorado, all models exhibit comparable performance in both forested and non-forested areas. In contrast, in California, models tend to perform slightly better in non-forested areas. Moreover, Fig. 89 and S10 illustrate the average spatial error in March and May, respectively, for California and Colorado. In California, the \hat{SWE} underestimates reference SWE, with the extent of underestimation varying across different climate models. For instance, in California, 'cnrm-esm2-1' at 9 km achieves the lowest underestimation. In general, the 'cnrm-esm2-1' model at 9 km exhibits superior performance for Colorado as an estimator, with a mean RMSE of 0.07 m and a standard deviation of 0.05 m. Conversely, for California, the 'ec-earth3-veg' model at 9 km demonstrates provides the best highest accuracy as an estimator, yielding a mean RMSE of 0.13 m and a standard deviation of 0.1 m.




Fig. 8. Accuracy assessment of downscaling cross-validation, conducted in California, over the six-year period from 2005 to 2010, conducted in California. Column titles indicate the results for downscaled SWE (SWE) using respective climate models as estimators. '9 km' and '100 km' refer to the resolutions of the climate data. The evaluation metrics include the Correlation Coefficient (R), Mean Difference (MD), and Root-Mean-Squared Error (RMSE). These metrics are considered in different elevation values and different land covers. In the legend, 2000 m < Elevation < 3000 m is shown as Elevation < 3000 m.




Fig. 9. Spatial assessment of the downscaling. The first three rows represent the mean difference between UCLA SWE and downscaled SWE (SWE - \hat{SWE}) over California throughout the 6-year testing period (2005-2010) for two different months (March and May). The last row represents the averaged reference SWE (UCLA SWE) or averaged downscaled SWE (\hat{SWE}) throughout the 6-year testing period (2005-2010) for two different months (March and May). The evaluation metric includes Mean Difference (MD).

5.3 Comparison with 1 km SWE SNODAS Dataset

The comparison of UCLA SWE and \hat{SWE} with the 1 km resolution SNODAS dataset is detailed presented in Supplementary Table S4S6, Figs. 910 and 4011, and Figs. S41S12 and S42S13. The results align with previous sections: all models exhibit comparable accuracy and correlation levels with the SNODAS SWE data, with distinctions arising based on resolutions, elevation ranges, and land cover types.

In California, during the snowiest months, the 'ec-earth3-veg' estimator excels in terms of RMSE, correlation, and mean difference at both resolutions. In Colorado, the 'mpi-esml1-2' estimator mostly demonstrates superior performance. Interestingly, in Colorado, the 100 km resolution slightly outperforms the 9 km resolution across some models. However, but in some other models others, the 100 km is marginally surpassed by the 9 km resolution marginally surpasses the 100 km resolution. In other words, the performance of different resolutions is nearly similar across all models, suggesting that using

the coarse-scale CMIP6 resolution can yield results comparable to those of finer resolutions. According to Fig. 4011 and S12 generally, in California, the 'ec-earth3-veg' model outperforms others at both 9 km and 100 km resolutions in both March and May, although it tends to mostly underestimate the SWE. In contrast, in Colorado, the 'mpi-esm1-2' model at 9 km performs better in March and May, mostly with an overestimation tendency overestimates. In general, in Colorado, the 'mpi-esm1-2' model at 100 km showed the most favorable results, with a mean RMSE of 0.08 m and a standard deviation of 0.07 m. In California, the 'ec-earth3-veg' model at 9 km outperformed others achieved the best performance, with a mean RMSE of 0.23 m and a standard deviation of 0.20 m.




Fig. 10. The average mean difference of SWE over California for the 6-year testing period (2005-2010) compares UCLA SWE (black dotted line) and downscaled SWE (SWE) data (blue and green lines) against SNODAS. The SWE data, derived from various different climate estimators/models, are identified in the legend represented by their names and associated distinct colors: light blue for 'ec-earth3-veg', green for 'cnrm-esm2-1', and dark blue for 'mpi-esm1-2'. Within each color group, line styles distinguish the spatial resolutions of the climate data: Solid lines indicate climate estimators estimates at a 9 km resolution, while dashed lines represent climate estimators at correspond to a 100 km resolution, distinguishing the spatial resolutions of the climate data.

of UCLA SWE and \hat{SWE} against the 1 km SWE Daymet dataset. The results show that all three models display comparable
 580 levels of accuracy and correlation with the Daymet SWE data.

In California, the 'ec-earth3-veg' model leads in performance at both 9 km and 100 km resolutions, boasting the lowest RMSE;
 and highest correlation. For Colorado, the 'cnrm-esm2-1' estimator predominantly demonstrates superior performance. When
 comparing both the UCLA SWE and \hat{SWE} to the 1 km SWE Daymet dataset, the results also indicate a close alignment
 between the two sets of data. Fig. 4213 and S13 demonstrate that in California and Colorado during the months of March and
 585 May, the 'cnrm-esm2-1' estimator yields the best results. In both regions, this model predominantly tends to overestimates SWE
 in March and underestimates it in May. In general, the 'cnrm-esm2-1' estimator at 100 km is the most accurate for Colorado,
 presenting a mean RMSE of 0.11 m and a standard deviation of 0.04 m. For California, the 'mpi-esm1-2' model at 100 km
achieves the performs best performance, with a mean RMSE of 0.19 m and a standard deviation of 0.06 m. However, 'ec-
 earth3-veg' achieves the same mean RMSE, with a standard deviation of 0.07 m.




Fig.-12. Same as Fig. 910, but for comparison against the Daymet data.




Fig. -13. Same as Fig. 4011, but for comparison against the Daymet data.

5.5 Comparison with 4 km SWE of the University of Arizona Dataset

This section compares the SWE from three models against the 4 km SWE data from the University of Arizona (UA). Supplementary Table S6S8, Figs. 1314 and 1415, and Figs. S14S16 and S15 illustrate the comparison between of UCLA SWE and SWE against with the 4 km SWE data from the UA dataset. The findings indicate that all three models provide comparable accuracy and correlation with the UCLA SWE, underscoring the effectiveness and reliability of our estimation approach in representing SWE in both California and Colorado.

A deeper analysis of model performance ~~reveals subtle~~highlights nuanced differences. In Colorado, the 'cnrm-esm2-1' estimator stands out at both 9 km and 100 km resolutions. Meanwhile, in California, mostly the 'ec-earth3-veg' model surpasses its counterparts, achieving the lowest mean RMSE at both resolutions.

Overall, the 9 km resolution ~~models~~ slightly ~~outperform~~outperforms the 100 km resolution ~~models~~ in both states. This suggests that a finer resolution might yield slightly more precise \hat{SWE} estimates. Additionally, when comparing ~~both the~~ UCLA SWE and \hat{SWE} to the 4 km UA SWE, the results demonstrate a tight congruence between the two datasets. Furthermore, Fig. ~~14~~15 and Fig. S15 show that in California, for March and May, the 'ec-earth3-veg' model at a 100 km resolution performs ~~optimally better~~. Meanwhile, in Colorado, the 'cnrm-esm2-1' estimator ~~achieves the~~performs best ~~performance~~ in these two months.

The 'ec-earth3-veg' model tends to underestimate SWE in California, while in Colorado, 'cnrm-esm2-1' generally overestimates SWE in May. In general, the 'cnrm-esm2-1' estimator at the 9 km scale provides the best results for Colorado, having a mean RMSE of 0.07 m and a standard deviation of 0.05 m. In California, the 'ec-earth3-veg' estimator at the 9 km scale ~~proves to be~~is the most precise, with a mean RMSE of 0.14 m and a standard deviation of 0.13 m.




Fig. -14. Same as Fig. 910, but for comparison against the University of Arizona (UA) data.




Fig. 15. Same as Fig. 4011, but for comparison against the University of Arizona (UA) data.

6. Discussion

In general, using the 'cnrm-esm2-1' model as an estimator results in better accuracy in Colorado at both 100 km and 9 km resolutions compared to other models. For instance, in Colorado, the use of the 'cnrm-esm2-1' model at a 9 km resolution demonstrated close agreement with observed SWE, with an average RMSE of 0.07 meters. This performance highlights the model's strong compatibility with the climatic and geographical complexities of Colorado. Conversely, in California, the 'ec-earth3-veg' model excels at a 9 km resolution, providing the most accurate results with an average RMSE of 0.13 meters

compared to the reference datasets. This suggests that its higher resolution better captures the region's complex environmental and topographical variations.

It also appears that a finer resolution of 9 km provides slightly better accuracy than a 100 km resolution across all models, although the difference is not substantial. This underscores the importance of selecting the appropriate climate model for SWE estimation, which can have a more significant impact than merely choosing a higher-resolution model. Moreover, CMIP6 models are designed for long-term climate projections and capture broad climate trends rather than predicting specific weather events. Despite this, the downscaled SWE using the proposed approach based on CMIP6 is comparable to that of WRF-CMIP6, which dynamically downscales CMIP6. This is largely because the proposed methodology relies on long-term climate data through the use of far and near temporal intervals, and CMIP6 effectively captures broad climatic trends and seasonality, including changes in temperature and precipitation patterns. Accordingly, achieving accurate HR-SWE estimation relies significantly on the choice and accuracy of the climate model inputs, such as precipitation and temperature data, which may introduce biases into the SWE estimates. For example, precipitation biases are a dominant factor influencing SWE estimation errors, while temperature biases become more significant during transitional periods, such as the spring melt season.

The following comparison provides a broader perspective on how our proposed method compares with other statistical downscaling techniques. BCSD methods are effective in reducing uncertainties in climate model outputs by adjusting model biases using high-resolution observations. These methods are particularly valuable for ensuring that model outputs align with observed climatology and capture local variability. However, they depend heavily on the availability of high-quality in-situ data, which limits their application in remote or data-sparse regions. In contrast, our method excels in areas with sparse data, as it uses low-resolution climate data without requiring ground observations, making it adaptable to a broader range of conditions.

Analog-type statistical downscaling approaches offer a simple, computationally efficient way to project high-resolution data based on historical relationships between large-scale climate patterns and local climate variables. These methods are useful in regions where historical climate patterns are stable and well-documented. On the other hand, our method introduces several key improvements over traditional k-NN downscaling techniques. First, the adaptation of the k-NN approach through the incorporation of far and near temporal intervals of climate data enhances its ability to handle dynamic variables, such as snow, which are subject to significant changes due to climate variability. Unlike conventional analog methods, which constrain analog candidates to a specific temporal window near the query date, this method eliminates such restrictions.

This flexibility is crucial for three reasons: (1) the inclusion of far and near temporal intervals allows the selection of the most suitable candidates across a broader temporal range, eliminating the need for narrow constraints; (2) it ensures the preservation of extreme events, as restricting candidates to a narrow temporal window may exclude matches that represent rare but important extreme events; and (3) it facilitates downscaling for future periods where no exact analogs exist in the historical record within

a specific date range. Instead, suitable analogs may still be found in historical data but during different periods. For instance, as climate change progresses, a future day in winter may no longer have a match on the same calendar day in the past. Instead, an analog might be found on another calendar day, such as in a warmer season like fall or spring.

Using very low-resolution climate data as input reduces both memory requirements and computation time compared to physical snow models, which also require high-resolution climate data as input to estimate HR-SWE information. This efficiency enables the generation of HR-SWE estimates over large spatial domains with reduced computational overhead. This is particularly beneficial when applying the method to large areas, long temporal scales, or ensembles of climate data.

6.7. Conclusion

This research ~~aims to propose~~proposes a methodology to downscale low-resolution daily Snow Water Equivalent (LR-SWE) by utilizing low-resolution daily climate data to produce daily, high-resolution SWE (HR-SWE) data, covering the period from 1950 to the present. We test our approach in two distinct areas (California and Colorado) in the western United States. Utilizing existing low-resolution meteorological factors obtained from global CMIP6 climate data (100 km) and downscaled CMIP6 for the Western United States (WRF-CMIP6, 9 km) and available 500-meter HR-SWE images from 1984 to 2021 (UCLA SWE dataset) for the Western United States, we implement a downscaling algorithm to estimate HR-SWE images. WRF-CMIP6 is a regional configuration of a climate model that dynamically downscales CMIP6 data, ~~incorporating ERA5 reanalysis data~~ to enhance spatial ~~and temporal~~ resolution. Utilizing both CMIP6 and WRF-CMIP6 at different resolutions ~~allows~~enables us to ~~investigate~~assess the ~~effects~~impact of using a regional climate model that dynamically downscales CMIP6 data and incorporates reanalysis data, alongside a global climate model with different spatial resolutions, on the accuracy of our approach. We also select three climate models ('ec-earth3-veg', 'mpi-esm1-2', and 'cnrm-esm2-1') for testing the proposed method using different models.

To perform a comprehensive accuracy assessment, we compared the downscaled SWE (\hat{SWE}) with established datasets like the 1 km SWE Daymet, 1 km SWE SNODAS, and the 4 km SWE from the University of Arizona. Moreover, we performed cross-validation accuracy assessments. Our \hat{SWE} data closely mirrored reference HR-SWE conditions. Even when ~~utilizing~~using lower-resolution climate datasets like CMIP6 (100 km), the method remained robust, leveraging recurring SWE patterns and climate data to generate more detailed SWE images.

A comprehensive analysis over all months reveals notable patterns in the performance of various climate models at different resolutions in California and Colorado. ~~In general, using~~Overall, the 'cnrm-esm2-1' model ~~as an estimator results in better~~tends to provide higher accuracy in Colorado ~~when used as an estimator, outperforming other models~~ at both 100 km and 9 km resolutions ~~compared to using other models. This performance highlights the model's strong compatibility with the climatic and geographical complexities of Colorado.~~ Conversely, in California, In contrast, the 'ec-earth3-veg' model ~~exceeds~~performs

best in California at a 9 km resolution, ~~offering the most accurate results. This suggests that its higher resolution better captures the state's complex environmental and topographical variations. It also appeared that~~. A finer resolution of ~~(9 km provided)~~ generally offers slightly better accuracy than a 100 km resolution across ~~all~~ models, ~~although though~~ the difference ~~was not substantial~~. This underscores the importance of selecting the appropriate climate model for SWE estimation, which can have a more significant impact than merely choosing a higher resolution model. Moreover, CMIP6 models are designed for long-term climate projections and capture broad climate trends rather than predicting specific weather events. Despite this, the downscaled SWE using the proposed approach based on CMIP6 is comparable to that of WRF-CMIP6, which dynamically downscales CMIP6 data with the incorporation of ERA5 reanalysis data. This is largely because the proposed methodology relies on long-term climate data, and CMIP6 effectively captures broad climatic trends and seasonality, including changes in temperature and precipitation patterns is modest, emphasizing the importance of model selection over merely increasing resolution.

Ongoing research will focus on applying our downscaling framework to predict ~~high-resolution HR~~-SWE based on future climate predictions. The data generated could be invaluable not only in climate change studies but also for water resources management. This underscores the need for precise model selection and the potential for applying our findings to future climate scenarios, ~~enhancing our understanding and~~ improving water resource management ~~of water resources in the face of~~ amid climate change. Moreover, we anticipate that our approach could be applied in areas ~~lacking where HR-SWE data is available for some years but~~ high-resolution climate models are lacking for past or future periods.

Code and Data Availability

The code for generating downscaled SWE maps, along with some sample data, is available in the assets section.

Video Supplement

The videos of High-Resolution SWE estimations for California and Colorado are available in the assets section.

Supplement Link

It is uploaded while submitting, and the link to the supplement will be included by Copernicus.

705 **Author Contribution**

FZ: Conceptualization, Methodology, Software, Validation, Investigation, Data curation, Formal analysis, Writing – original draft.

GM and MG: Conceptualization, Methodology, Validation, Formal analysis, Writing – review & editing, Supervision, Project administration, and Resources.

710 **FZ and GM:** Funding acquisition.

Competing Interests

The authors declare that they have no conflict of interest.

Acknowledgments

Our study utilized datasets including [the Western United States snow reanalysis dataset](#), Daymet V4 [dataset](#), SNODAS [dataset](#),
715 University of Arizona's SWE [dataset](#), [SNOTEL dataset](#), MODIS Land Cover Type MCD12Q1 [dataset](#), NASA SRTM Digital Elevation 30 m, CMIP6 climate predictions, and downscaled CMIP6 data using the Weather Research and Forecasting (WRF) model. We thank the respective institutions and researchers for providing these data sets.

This work was primarily carried out as part of the scientific project “Deep-time synthetic data cubes to enable long-term hydrological modeling”, funded by The Swiss National Science Foundation (SNSF), with project number 200021_204130.

720 The authors would like to gratefully acknowledge the SNSF. Moreover, this project was carried out as a joint effort between the University of Lausanne (UNIL) and the University of California, Berkeley (UCB), and was made possible thanks to the UNIL Mobility Fellowship with fellowship number MD0012. We express our gratitude to UNIL for providing the funding and to UCB for hosting the first author.

References

725 U.S. Bureau of Reclamation (USBR): Emerging Technologies in Snow Monitoring,, Report to Congress, 2021.

Abatzoglou, J. T.: Development of gridded surface meteorological data for ecological applications and modelling, International Journal of Climatology, 33, 121-131, 2013.

Abatzoglou, J. T. and Brown, T. J.: A comparison of statistical downscaling methods suited for wildfire applications, International journal of climatology, 32, 772-780, 2012.

- 730 Alonso-González, E., Aalstad, K., Pirk, N., Mazzolini, M., Treichler, D., Leclercq, P., Westermann, S., López-Moreno, J. I., and Gascoin, S.: Spatio-temporal information propagation using sparse observations in hyper-resolution ensemble-based snow data assimilation, *Hydrology Earth System Sciences*, 27, 4637-4659, 2023.
- Bair, E. H., Rittger, K., Davis, R. E., Painter, T. H., and Dozier, J.: Validating reconstruction of snow water equivalent in California's Sierra Nevada using measurements from the NASA Airborne Snow Observatory, *Water Resources Research*, 52, 8437-8460, 2016.
- 735 Bales, R. C., Molotch, N. P., Painter, T. H., Dettinger, M. D., Rice, R., and Dozier, J.: Mountain hydrology of the western United States, *Water Resources Research*, 42, 2006.
- Belmecheri, S., Babst, F., Wahl, E. R., Stahle, D. W., and Trouet, V.: Multi-century evaluation of Sierra Nevada snowpack, *Nature Climate Change*, 6, 2-3, 2016.
- 740 Broxton, P., Zeng, X., and Dawson, N.: Daily 4 km Gridded SWE and Snow Depth from Assimilated In-Situ and Modeled Data over the Conterminous US, Version 1 [Data Set]. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active [dataset], 2019.
- Caillouet, L., Vidal, J.-P., Sauquet, E., and Graff, B.: Probabilistic precipitation and temperature downscaling of the Twentieth Century Reanalysis over France, *Climate of the Past*, 12, 635-662, 2016.
- 745 Center, N. O. H. R. S.: Snow Data Assimilation System (SNODAS) Data Products at NSIDC, Version 1, National Snow and Ice Data Center, Boulder, Colorado USA2004.
- Clow, D. W., Nanus, L., Verdin, K. L., and Schmidt, J.: Evaluation of SNODAS snow depth and snow water equivalent estimates for the Colorado Rocky Mountains, USA, *Hydrological Processes*, 26, 2583-2591, 2012.
- Council, N. R. and Committee, N. W. S. M.: Future of the National Weather Service Cooperative Observer Network, National Academies Press1998.
- 750 Cover, T. and Hart, P.: Nearest neighbor pattern classification, *IEEE transactions on information theory*, 13, 21-27, 1967.
- Dawadi, S. and Ahmad, S.: Changing climatic conditions in the Colorado River Basin: Implications for water resources management, *Journal of Hydrology*, 430, 127-141, 2012.
- Dietz, A. J., Wohner, C., and Kuenzer, C.: European Snow Cover Characteristics between 2000 and 2011 Derived from Improved MODIS Daily Snow Cover Products, *Remote Sensing*, 4, 2432-2454, 10.3390/rs4082432, 2012.
- 755 Fang, Y., Liu, Y., and Margulis, S. A.: A western United States snow reanalysis dataset over the Landsat era from water years 1985 to 2021, *Scientific Data*, 9, 677, 2022.
- Fang, Y., Liu, Y., Li, D., Sun, H., and Margulis, S. A.: Spatiotemporal snow water storage uncertainty in the midlatitude American Cordillera, *The Cryosphere*, 17, 5175-5195, 2023.
- 760 Farr, T. G., Rosen, P. A., Caro, E., Crippen, R., Duren, R., Hensley, S., Kobrick, M., Paller, M., Rodriguez, E., and Roth, L.: The shuttle radar topography mission, *Reviews of geophysics*, 45, 2007.
- Fiddes, J., Aalstad, K., and Westermann, S.: Hyper-resolution ensemble-based snow reanalysis in mountain regions using clustering, *Hydrology Earth System Sciences*, 23, 4717-4736, 2019.

- 765 Fleming, S. W., Zukiewicz, L., Strobel, M. L., Hofman, H., and Goodbody, A. G.: SNOTEL, the Soil Climate Analysis Network, and water supply forecasting at the Natural Resources Conservation Service: Past, present, and future, *JAWRA Journal of the American Water Resources Association*, 2023.
- Gangopadhyay, S., Clark, M., and Rajagopalan, B.: Statistical downscaling using K-nearest neighbors, *Water Resources Research*, 41, 2005.
- 770 Girotto, M., Musselman, K. N., and Essery, R. L.: Data assimilation improves estimates of climate-sensitive seasonal snow, *Current Climate Change Reports*, 6, 81-94, 2020.
- Heldmyer, A. J., Bjarke, N. R., and Livneh, B.: A 21st-Century perspective on snow drought in the Upper Colorado River Basin, *Journal of the American Water Resources Association*, 59, 396-415, 2023.
- 775 Henn, B., Musselman, K. N., Lestak, L., Ralph, F. M., and Molotch, N. P.: Extreme runoff generation from atmospheric river driven snowmelt during the 2017 Oroville Dam spillways incident, *Geophysical Research Letters*, 47, e2020GL088189, 2020.
- Henson, R.: Flow Cytometry Data Reader and Visualization, MATLAB Central File Exchange [code], 2024.
- Horton, P.: AtmoSwing: Analog technique model for statistical weather forecastING and downscaling (v2. 1.0), *Geoscientific Model Development*, 12, 2915-2940, 2019.
- 780 Kouki, K., Räisänen, P., Luojus, K., Luomaranta, A., and Riihelä, A.: Evaluation of Northern Hemisphere snow water equivalent in CMIP6 models during 1982–2014, *The Cryosphere*, 16, 1007-1030, 2022.
- Lall, U. and Sharma, A.: A nearest neighbor bootstrap for resampling hydrologic time series, *Water resources research*, 32, 679-693, 1996.
- 785 Llargeron, C., Dumont, M., Morin, S., Boone, A., Lafaysse, M., Metref, S., Cosme, E., Jonas, T., Winstral, A., and Margulis, S. A.: Toward Snow Cover Estimation in Mountainous Areas Using Modern Data Assimilation Methods: A Review, *Frontiers in Earth Science*, 8, 10.3389/feart.2020.00325, 2020.
- Lehning, M., Völsch, I., Gustafsson, D., Nguyen, T. A., Stähli, M., and Zappa, M.: ALPINE3D: a detailed model of mountain surface processes and its application to snow hydrology, *Hydrological Processes: An International Journal*, 20, 2111-2128, 2006.
- 790 Lievens, H., Demuzere, M., Marshall, H. P., Reichle, R. H., Brucker, L., Brangers, I., de Rosnay, P., Dumont, M., Girotto, M., Immerzeel, W. W., Jonas, T., Kim, E. J., Koch, I., Marty, C., Saloranta, T., Schober, J., and De Lannoy, G. J. M.: Snow depth variability in the Northern Hemisphere mountains observed from space, *Nature Communications*, 10, 10.1038/s41467-019-12566-y, 2019.
- Liston, G. E. and Sturm, M.: A snow-transport model for complex terrain, *Journal of Glaciology*, 44, 498-516, 1998.
- 795 Lundquist, J. D., Chickadel, C., Cristea, N., Currier, W. R., Henn, B., Keenan, E., and Dozier, J.: Separating snow and forest temperatures with thermal infrared remote sensing, *Remote Sensing of Environment*, 209, 764-779, 10.1016/j.rse.2018.03.001, 2018.
- Ma, X., Li, D., Fang, Y., Margulis, S. A., and Lettenmaier, D. P.: Estimating spatiotemporally continuous snow water equivalent from intermittent satellite observations: an evaluation using synthetic data, *Hydrology Earth System Sciences*, 27, 21-38, 2023.

- 800 Margulis, S. A., Liu, Y., and Baldo, E.: A joint landsat-and modis-based reanalysis approach for midlatitude montane seasonal snow characterization, *Frontiers in Earth Science*, 7, 272, 2019.
- Margulis, S. A., Cortés, G., Giroto, M., and Durand, M.: A Landsat-era Sierra Nevada snow reanalysis (1985–2015), *Journal of Hydrometeorology*, 17, 1203-1221, 2016.
- 805 Margulis, S. A., Giroto, M., Cortés, G., and Durand, M.: A particle batch smoother approach to snow water equivalent estimation, *Journal of Hydrometeorology*, 16, 1752-1772, 2015.
- Meehl, G. A., Moss, R., Taylor, K. E., Eyring, V., Stouffer, R. J., Bony, S., and Stevens, B.: Climate model intercomparisons: Preparing for the next phase, *Eos, Transactions American Geophysical Union*, 95, 77-78: The data was downloaded in January 2023., 2014.
- 810 Mower, R., Gutmann, E. D., Liston, G. E., Lundquist, J., and Rasmussen, S.: Parallel SnowModel (v1. 0): a parallel implementation of a distributed snow-evolution modeling system (SnowModel), *Geoscientific Model Development*, 17, 4135-4154, 2024.
- Painter, T. H., Bryant, A. C., and Skiles, S. M.: Radiative forcing by light absorbing impurities in snow from MODIS surface reflectance data, *Geophysical Research Letters*, 39, 10.1029/2012gl052457, 2012.
- 815 Pflug, J. and Lundquist, J.: Inferring distributed snow depth by leveraging snow pattern repeatability: Investigation using 47 lidar observations in the Tuolumne watershed, Sierra Nevada, California, *Water Resources Research*, 56, e2020WR027243, 2020.
- Pflug, J. M., Wrzesien, M. L., Kumar, S. V., Cho, E., Arsenault, K. R., Houser, P. R., and Vuyovich, C. M.: Extending the utility of space-borne snow water equivalent observations over vegetated areas with data assimilation, *Hydrology Earth System Sciences*, 28, 631-648, 2024.
- 820 Pons, M., San-Martín, D., Herrera, S., and Gutiérrez, J. M.: Snow trends in Northern Spain: analysis and simulation with statistical downscaling methods, *international journal of climatology*, 2010.
- Rahimi, S., Krantz, W., Lin, Y. H., Bass, B., Goldenson, N., Hall, A., Lebo, Z. J., and Norris, J.: Evaluation of a Reanalysis-Driven Configuration of WRF4 Over the Western United States From 1980 to 2020, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035699, 2022.
- 825 Rahimi, S., Huang, L., Norris, J., Hall, A., Goldenson, N., Krantz, W., Bass, B., Thackeray, C., Lin, H., and Chen, D.: An overview of the Western United States Dynamically Downscaled Dataset (WUS-D3), *Geoscientific Model Development*, 17, 2265-2286, 2024.
- Rajagopalan, B. and Lall, U.: A k-nearest-neighbor simulator for daily precipitation and other weather variables, *Water resources research*, 35, 3089-3101, 1999.
- 830 Ranzi, R., Colosio, P., and Galeati, G.: Climatology of snow depth and water equivalent measurements in the Italian Alps (1967–2020), *Hydrology Earth System Sciences*, 28, 2555-2578, 2024.
- Rettie, F. M., Gayler, S., Weber, T. K., Tesfaye, K., and Streck, T.: High-resolution CMIP6 climate projections for Ethiopia using the gridded statistical downscaling method, *Scientific data*, 10, 442, 2023.
- 835 Saberi, N., Kelly, R., Flemming, M., and Li, Q.: Review of snow water equivalent retrieval methods using spaceborne passive microwave radiometry, *International Journal of Remote Sensing*, 41, 996-1018, 2020.

- Shahriari, B., Swersky, K., Wang, Z., Adams, R. P., and De Freitas, N.: Taking the human out of the loop: A review of Bayesian optimization, *Proceedings of the IEEE*, 104, 148-175, 2015.
- Shi, J. and Dozier, J.: Inferring snow wetness using C-band data from SIR-C's polarimetric synthetic aperture radar, *IEEE transactions on geoscience remote sensing*, 33, 905-914, 1995.
- 840 Shi, J. and Dozier, J.: Estimation of snow water equivalence using SIR-C/X-SAR. II. Inferring snow depth and particle size, *IEEE Transactions on Geoscience Remote sensing*, 38, 2475-2488, 2000.
- Siirila-Woodburn, E. R., Rhoades, A. M., Hatchett, B. J., Huning, L. S., Szinai, J., Tague, C., Nico, P. S., Feldman, D. R., Jones, A. D., and Collins, W. D.: A low-to-no snow future and its impacts on water resources in the western United States, *Nature Reviews Earth and Environmental*, 2, 800-819, 2021a.
- 845 Siirila-Woodburn, E. R., Rhoades, A. M., Hatchett, B. J., Huning, L. S., Szinai, J., Tague, C., Nico, P. S., Feldman, D. R., Jones, A. D., Collins, W. D. J. N. R. E., and Environment: A low-to-no snow future and its impacts on water resources in the western United States, *Nature Reviews Earth Environment*, 2, 800-819, 2021b.
- 850 Snapir, B., Momblanch, A., Jain, S., Waive, T. W., and Holman, I. P.: A method for monthly mapping of wet and dry snow using Sentinel-1 and MODIS: Application to a Himalayan river basin, *International Journal of Applied Earth Observation Geoinformation*, 74, 222-230, 2019.
- Snoek, J., Larochelle, H., and Adams, R. P.: Practical bayesian optimization of machine learning algorithms, *Advances in neural information processing systems*, 25, 2012.
- Sturm, M., Goldstein, M. A., and Parr, C.: Water and life from snow: A trillion dollar science question, *Water Resources Research*, 53, 3534-3544, 2017.
- 855 Tabari, H., Paz, S. M., Buekenhout, D., and Willems, P.: Comparison of statistical downscaling methods for climate change impact analysis on precipitation-driven drought, *Hydrology Earth System Sciences*, 25, 3493-3517, 2021.
- Thornton, M., Shrestha, R., Wei, Y., Thornton, P., Kao, S., and Wilson, B.: Daymet: daily surface weather data on a 1-km grid for North America, Version 4 R1. ORNL DAAC, Oak Ridge, Tennessee, USA, <https://doi.org/10.3334/ORNLDAAAC/2129>, 2022.
- 860 Thrasher, B., Wang, W., Michaelis, A., Melton, F., Lee, T., and Nemani, R.: NASA global daily downscaled projections, CMIP6, *Scientific data*, 9, 262, 2022.
- Tsang, L., Durand, M., Derksen, C., Barros, A. P., Kang, D.-H., Lievens, H., Marshall, H.-P., Zhu, J., Johnson, J., and King, J.: Global Monitoring of Snow Water Equivalent using High Frequency Radar Remote Sensing, *The Cryosphere Discussions*, 2021, 1-57, 2021.
- 865 Vionnet, V., Martin, E., Masson, V., Guyomarc'h, G., Naaïm-Bouvet, F., Prokop, A., Durand, Y., and Lac, C.: Simulation of wind-induced snow transport and sublimation in alpine terrain using a fully coupled snowpack/atmosphere model, *The Cryosphere*, 8, 395-415, 2014.
- Wood, A. W., Leung, L. R., Sridhar, V., and Lettenmaier, D.: Hydrologic implications of dynamical and statistical approaches to downscaling climate model outputs, *Climatic change*, 62, 189-216, 2004.

- 870 Wu, X., Naegeli, K., Premier, V., Marin, C., Ma, D., Wang, J., and Wunderle, S.: Evaluation of snow extent time series derived from Advanced Very High Resolution Radiometer global area coverage data (1982–2018) in the Hindu Kush Himalayas, *The Cryosphere*, 15, 4261-4279, 2021.
- Wundram, D. and Löffler, J.: High-resolution spatial analysis of mountain landscapes using a low-altitude remote sensing approach, *International Journal of Remote Sensing*, 29, 961-974, 2008.
- 875 Yates, D., Gangopadhyay, S., Rajagopalan, B., and Strzepek, K.: A technique for generating regional climate scenarios using a nearest-neighbor algorithm, *Water resources research*, 39, 2003.
- Yiou, P. and Déandréis, C.: Stochastic ensemble climate forecast with an analogue model, *Geoscientific Model Development*, 12, 723-734, 2019.
- 880 Zakeri, F. and Mariethoz, G.: Synthesizing long-term satellite imagery consistent with climate data: Application to daily snow cover, *Remote Sensing of Environment*, 300, 113877, 2024.
- Zeng, X., Broxton, P., and Dawson, N.: Snowpack change from 1982 to 2016 over conterminous United States, *Geophysical Research Letters*, 45, 12,940-912,947, 2018.
- Zorita, E. and Von Storch, H.: The analog method as a simple statistical downscaling technique: Comparison with more complicated methods, *Journal of climate*, 12, 2474-2489, 1999.
- 885