

# Partitioning anthropogenic and natural methane emissions in Finland during 2000–2021 by combining bottom-up and top-down estimates

Maria K. Tenkanen<sup>1</sup>, Aki Tsuruta<sup>1</sup>, Hugo Denier van der Gon<sup>2</sup>, Lena Höglund-Isaksson<sup>3</sup>, Antti Leppänen<sup>1,4</sup>, Tiina Markkanen<sup>1</sup>, Ana Maria Roxana Petrescu<sup>5</sup>, Maarit Raivonen<sup>4</sup>, Hermanni Aaltonen<sup>1</sup>, and Tuula Aalto<sup>1</sup>

<sup>1</sup>Climate System Research, Finnish Meteorological Institute, 00560 Helsinki, Finland

<sup>2</sup>Department of Air Quality and Emissions Research, TNO, 3584 CB Utrecht, the Netherlands

<sup>3</sup>International Institute for Applied Systems Analysis (IIASA), 2361 Laxenburg, Austria

<sup>4</sup>Institute for Atmospheric and Earth System Research/Physics, Faculty of Science, University of Helsinki, 00014 Helsinki, Finland

<sup>5</sup>Department of Earth Sciences, Vrije Universiteit Amsterdam, 1081HV, Amsterdam, the Netherlands

**Correspondence:** Maria K. Tenkanen (maria.tenkanen@fmi.fi)

**Abstract.** Accurate national methane (CH<sub>4</sub>) emission estimates are essential for tracking progress towards climate goals. This study investigated Finnish CH<sub>4</sub> emissions from ~~2000–2021~~ 2000–2021 using bottom-up and top-down approaches. We evaluated the ability of a global atmospheric ~~inversion model's ability~~ inverse model CarbonTracker Europe - CH<sub>4</sub> to estimate CH<sub>4</sub> emissions within a single country, ~~focusing on how the choice of priors and uncertainties affected optimised emissions. The~~ 5 We focused on how different priors and their uncertainties affect the optimised emissions and showed that the optimised anthropogenic and natural CH<sub>4</sub> emissions ~~strongly depended~~ were strongly dependent on the prior emissions. ~~While~~ However, while the range of CH<sub>4</sub> estimates was large, the optimised emissions were more constrained than the bottom-up estimates. Further analysis ~~of CarbonTracker Europe – CH<sub>4</sub> results showed that~~ showed that the optimisation aligned the trends of anthropogenic and natural CH<sub>4</sub> ~~emission and improved~~ emissions and improved the modelled seasonal cycles of natural emissions. Comparison of atmospheric CH<sub>4</sub> observations with model results showed no clear preference between anthropogenic inventories (EDGAR v6 and CAMS-REG), but results using the ~~largest~~ highest natural prior (JSBACH-HIMMELI) ~~best agreed~~ agreed best with observations, suggesting that process-based models may underestimate CH<sub>4</sub> emissions from Finnish peatlands or unaccounted sources such as freshwater emissions. Additionally, using ~~a process-model spread-based uncertainty estimate an~~ uncertainty estimate based on a process-based model ensemble for natural CH<sub>4</sub> emissions seemed to be advantageous compared to the standard ~~constant estimate~~ uncertainty definition. The average total posterior emission of the ensemble from one ~~inversion~~ inverse model with different priors was similar to the average of the ensemble ~~including different inversion~~ including different inverse models but similar priors. Thus, a single inverse model using a range of priors can be used to reliably estimate CH<sub>4</sub> emissions when an ensemble of different models is unavailable.

## 1 Introduction

20 Methane (CH<sub>4</sub>) is the second most important anthropogenic greenhouse gas (GHG) after carbon dioxide (CO<sub>2</sub>). Its atmospheric concentration has nearly tripled since pre-industrial times, ~~largely primarily~~ due to human activities (Canadell et al., 2023). ~~In recent years, especially in 2020 and 2021~~ Since atmospheric CH<sub>4</sub> measurements began in the late 1970s (Rice et al., 2016), the growth rate of atmospheric CH<sub>4</sub> has ~~been remarkably high~~ (varied considerably, with periods of rapid growth as well as a plateau (Nisbet et al., 2023); The growth rate of CH<sub>4</sub> was close to zero from 2000 to 2006, after which the atmospheric CH<sub>4</sub> levels began to rise again (Nisbet et al., 2014; Mikaloff-Fletcher and Schaefer, 2019), reaching remarkably high increases of 15.15 ppb in 2020 and 17.97 ppb in 2021 ) (Lan et al., 2024), continuing the renewed growth that began in 2007 (Nisbet et al., 2014; Mikaloff-Fletcher and Schaefer, 2019) (Lan et al., 2024). The reasons for this renewed growth and the record-high CH<sub>4</sub> growth rates in 2020 and 2021 are still under discussion (Nisbet et al., 2023), which reflects the large uncertainties in CH<sub>4</sub> emissions. Reducing CH<sub>4</sub> emissions is an effective way to mitigate climate change ~~given CH<sub>4</sub>'s~~ (Nisbet et al., 2020; Collins et al., 2018), ~~given the~~ short atmospheric lifetime ~~and strong of CH<sub>4</sub>~~ (9.1 years; Canadell et al., 2023) and its high global warming potential (Forster et al., 2023; Nisbet et al., 2020; Collins et al., 2018) ~~but~~ (82.5 times higher than CO<sub>2</sub> on a 20-year time scale; Forster et al., 2023). However, in order to assess the success of the CH<sub>4</sub> emission reductions, we need to ~~be able to improve our ability~~ to quantify CH<sub>4</sub> emissions and their changes ~~better than we currently do~~.

In the Paris Agreement ~~, the participating countries pledged to report~~ (UNFCCC, 2015), participating countries committed to ~~reporting~~ their GHG emissions and removals coherently and transparently by compiling national ~~GHG inventories (NGHGI)~~ (UNFCCC, 2015) ~~greenhouse gas inventories (NGHGs)~~. The NGHGs are evaluated jointly every five years in the global stocktake (~~global~~) (UNFCCC, 2023) which was completed for the first time in 2023. They are based on a bottom-up approach ~~, starting from that starts at~~ the sources and ~~estimating how much each source emits GHGs~~ estimates how much GHG is emitted by each source. The main ~~aim objective~~ is to capture trends caused by (direct) anthropogenic activities in order to track the effect of ~~climate change~~ mitigation efforts put into practice, and thus, the NGHGs report ~~the~~ emissions and sinks as annual ~~country-totals~~ country totals. In addition to the NGHGs, which ~~each country compiles independently~~ are compiled independently by each country, there are other bottom-up anthropogenic GHG inventories ~~composed that are compiled~~ for larger regions or even ~~globally on a global scale~~. Such inventories relevant ~~for to~~ Finland are, for example, the Emissions Database for Global Atmospheric Research (EDGAR, ~~European Commission and Joint Research Centre et al. (2023)~~); ~~European Commission and Joint Research Centre et al., 2023~~ ~~), the~~ Copernicus Atmosphere Monitoring Service Regional inventory (CAMS-REG, ~~(Kuenen et al., 2022)~~) and ~~); Kuenen et al., 2022~~ ~~) and the~~ Greenhouse gas and Air pollution Interactions and Synergies (GAINS, ~~Höglund-Isaksson et al. (2020)~~); Höglund-Isaksson et al., 2020). EDGAR is a widely used global inventory with regular updates, while CAMS-REG and GAINS (the version used in this study) ~~cover only only cover~~ Europe. However, both CAMS-REG and GAINS use more specific country-level data, while EDGAR uses globally consistent methods. The main uncertainties in the bottom-up inventories result from the estimated ~~magnitudes magnitude~~ of each source category and ~~their used emission factors~~ the emission factors used. Nevertheless, they provide estimates for each source category separately.

Another way to estimate GHG emissions is to use a top-down approach, or atmospheric ~~inversion~~. With inverse modelling. Using a combination of an atmospheric chemical transport ~~models and~~ model and measurements of atmospheric GHG mole ~~fraction measurements~~ fractions, they revise the assumed prior emissions. The atmospheric ~~inversion~~ inverse models of GHGs  
55 are becoming ~~ever more important in relation to our climate politics~~ increasingly important in the context of our climate policies (Leip et al., 2018). Until now, the assessment of national GHG budgets has relied on bottom-up-based inventories, especially on the NGHGs. ~~In the~~ The 2019 ~~Refinement to~~ refinement of the 2006 IPCC Guidelines for National Greenhouse Gas Inventories ~~, the inversion models were highlighted~~ (Maksyutov et al., 2019) highlighted inverse models as a potential way to support and verify the NGHGs ~~(Maksyutov et al., 2019)~~. ~~A couple of countries (e.g. the UK~~ . Some countries, such as the  
60 United Kingdom (Manning et al., 2021; Lunt et al., 2021), Switzerland (Henne et al., 2016), Germany (Integrated Greenhouse Gas Monitoring System for Germany ITMS, 2024), Australia (Luhar et al., 2020) and New Zealand (Geddes et al., 2021)) ~~already utilise inversion~~ , already use inverse modelling in their NGHGs, either as an appendix or ~~correcting to correct~~ the methods used in the NGHGI. All of these countries have certain advantages, for example, being an island and having several atmospheric observation sites, which ~~makes make~~ it easier for ~~inversion~~ inverse models to estimate GHGs within their national  
65 borders. However, without such advantages, ~~partitioning inversion model results on country-level~~ the partitioning of inverse model results at the country level is still uncertain and shows more differences between different models and model ~~setups~~ (Deng et al., 2022; McGrath et al., 2023; Petrescu et al., 2023; ?) set-ups (Deng et al., 2022; McGrath et al., 2023; Petrescu et al., 2023, 20

Atmospheric ~~inversion models are at their~~ inverse models are strongest when estimating ~~the~~ total emissions, including both  
70 ~~the~~ anthropogenic emissions reported in the NGHGs and natural sources. The further ~~partition to~~ partitioning into source categories is ~~more complex~~ , however, more complex, but there are ~~several methods for how a number of ways in which~~ this can be ~~achieved~~ done. One way is to take advantage of prior distributions and optimise different source categories individually but simultaneously (e.g. Tsuruta et al., 2017; Segers and Nanni, 2023; Janardanan et al., 2024). ~~With this method, since the~~ Since this method relies on the prior distribution to partition between different source categories ~~relies on the prior distribution~~, the optimisation is prone to ~~miscategorise the misclassifying~~ CH<sub>4</sub> emissions if there are several multiple sources in the same area and if the relative magnitudes of the priors are not correct. This uncertainty can be quantified to some extent by using different prior emissions and assessing how different prior emission estimates affect the optimised emissions. ~~Additionally~~ In addition, analytical inversions can be used (Cusworth et al., 2021; Worden et al., 2023). To reduce the dependence on prior distributions, ~~usage of carbon isotopes~~ the use of carbon isotope measurements has been intensively studied (e.g. Thompson et al., 2018; McNorton et al., 2018; Basu et al., 2022; Haghnegahdar et al., 2023; Chandra et al., 2024; ?). The models rely on different CH<sub>4</sub> sources having different ~~isotope signatures~~ , for example, isotopic signatures ( $\delta^{13}\text{C}$ ), e.g. emissions from wetlands have lower  $\delta^{13}\text{C}$  than fossil fuel emissions. However, the sparse number of isotope measurements limits the ~~usage~~ use of isotope measurements in the inversions as an additional constraint. Furthermore, ~~the isotope signatures~~  $\delta^{13}\text{C}$  values have uncertainties (Thanwerdas et al., 2024), although ~~, the~~ largest uncertainty has been attributed to uncertainties in the atmospheric chemistry (Basu et al.,  
85 2022). Similar to using CH<sub>4</sub> isotope measurements as an additional constraint, we can also use co-emitted species and the ratio

of them to CH<sub>4</sub> emitted from specific sources, such as ethane (Rice et al., 2016; Ramsden et al., 2022; Thompson et al., 2018)

~  
Although only anthropogenic emissions are reported in the NGHGs, ~~as our climate change mitigation efforts since our efforts to mitigate climate change~~ can be targeted ~~to~~ at them, natural GHG emissions also have an impact on climate change.

90 Thus, it is equally important to quantify natural emissions. In Finland, large ~~peatland areas are a significant areas of peatlands~~ are an important source of CH<sub>4</sub>, and the magnitude of natural CH<sub>4</sub> emissions is high compared to anthropogenic CH<sub>4</sub> emissions ~~as~~ estimated by the Finnish NGHGI (Tenkanen et al., 2024). Peatlands are concentrated in northern Finland, while the majority of the Finnish population lives in the south. Consequently, anthropogenic CH<sub>4</sub> emissions originate from the south. Different bottom-up estimates of CH<sub>4</sub> estimates, including both anthropogenic inventories ~~and process~~ (0.19–0.76 Tg yr<sup>-1</sup>; Section 3.1)  
95 and process-based models estimating the soil CH<sub>4</sub> balance (0.08–0.39 Tg yr<sup>-1</sup>; Section 3.2), vary considerably in Finland. ~~It is important~~ When interpreting the inverse model results, it's essential to identify the reasons for these inconsistencies ~~Furthermore, when interpreting the inversion model results, it is important~~ and to understand the extent to which the ~~used prior emissions~~ prior emissions used cause uncertainties in the optimised emission estimates, especially in regions where both anthropogenic and natural CH<sub>4</sub> emissions are abundant.

100 We studied CH<sub>4</sub> emissions in Finland during the ~~past~~ last two decades (2000–2021) using both bottom-up and top-down approaches, and discuss how ~~the~~ estimates from the two approaches differ. We aim to separate anthropogenic emissions from natural peatland emissions and estimate their relative magnitudes in Finland. Our focus is on ~~the~~ CH<sub>4</sub> emission estimates from the ~~inversion~~ inverse model CarbonTracker Europe - CH<sub>4</sub> (CTE-CH<sub>4</sub>) (Tsuruta et al., 2017), ~~used by previous studies~~ which previous studies have used to estimate Finnish CH<sub>4</sub> emissions (Tsuruta et al., 2019; Tenkanen et al., 2024), but here we  
105 extend the study period and investigate the results in more detail. Our study can be divided into four parts: First, we compare different anthropogenic ~~emissions~~ emission inventories (the Finnish NGHGI, GAINS, CAMS-REG and EDGAR v6, v7 and v8), both their total emission estimates and the magnitudes of different source categories. Second, we study the estimates from our ~~inversion~~ inverse model using different prior and uncertainty estimates and compare our ensemble with 13 inversion estimates collected in the VERIFY project (<https://verify.lsce.ipsl.fr/>). Third, we study the seasonal cycles of CH<sub>4</sub> emissions  
110 to see how ~~the use of~~ using atmospheric CH<sub>4</sub> observations affects ~~the~~ this seasonal cycle of ~~the~~ CH<sub>4</sub> estimates. Finally, we compare the modelled atmospheric mole fractions with observations and rank our ~~inversion~~ inverse model estimates based on this comparison, attempting to disentangle which ~~inversion~~ inverse model setup agrees ~~the~~ best with observations.

## 2 Materials and methods

### 2.1 Anthropogenic methane emission inventories

#### 115 2.1.1 ~~Finnish NGHGI~~

##### Finnish NGHGI

Finnish anthropogenic CH<sub>4</sub> emissions are reported on an annual basis by Statistics Finland (Statistics Finland, 2023) following ~~. The reporting follows~~ the IPCC 2006 reporting guidelines with refinements in 2019 (IPCC, 2019). The ~~NGHGI Fi~~ Finnish NGHGI (NGHGI Fi) includes CH<sub>4</sub> emissions from agriculture, waste, energy, industry ~~;~~ and land use, land use change and  
120 forestry (LULUCF) ~~;~~ and uses a mix of Tiers 1 (emission factors from IPCC reports), 2 ~~and 3.~~ The emissions (country-specific emission factors) and 3 (more advanced methods such as process-based modelling). Emissions from the fifth reporting category, ~~(LULUCF)~~ LULUCF, are not ~~studied here because the~~ analysed here because NGHGI Fi was the only studied inventory ~~which reported the~~ that reported LULUCF emissions. However, CH<sub>4</sub> emissions from the LULUCF sector in Finland have been  
125 discussed in detail by Tenkanen et al. (2024) ~~;~~ and were on average 0.03 Tg yr<sup>-1</sup> during 2000–2021 according to ~~the~~ NGHGI Fi.

#### 2.1.1 ~~EDGAR~~

##### EDGAR

EDGAR (<https://edgar.jrc.ec.europa.eu/>) is a global emission inventory developed by the Joint Research Centre of the European Commission, which provides estimates in a globally consistent way and ~~does not often~~ often does not use country-specific  
130 details. CH<sub>4</sub> estimates are provided by sector from agriculture, waste, energy and industry ~~with a detailed subdivision further~~ with further partitioning into subcategories. ~~Emissions are estimated fully based on statistical data from 1970.~~ The latest version, EDGAR v8 (European Commission and Joint Research Centre et al., 2023), ~~has estimates until~~ provides estimates up to 2022. The spatial resolution is 0.1° × 0.1° (latitude × longitude) and the temporal resolution is monthly.

#### 2.1.1 ~~CAMS-REG~~

135 ~~CAMS-REG-v5 is a regional~~

##### CAMS-REG

CAMS-REG v5 is a European anthropogenic emission inventory ~~for the years 2005–2018 which~~ covering the period from 2005 to 2018. It builds on the ~~officially reported emission data by the countries in the year~~ emission data reported officially in 2020 ~~under the LRTAP Convention and the NEC Directive~~ by countries under the convention on long-range transboundary air  
140 pollution (UNECE, 2012) and the EU national emission ceilings directive (European Commission, 2016) for the air pollutants

and, similarly, the reported ~~greenhouse gas emissions~~ GHG emissions by the countries to UNFCCC. The structure of the dataset, the harmonisation and gap-filling approach, ~~and proxies used to distribute the emissions spatially and the proxies used for the spatial distribution of emissions~~ are described in detail by Kuenen et al. (2022). The spatial resolution of CAMS-REG is  $0.05^\circ \times 0.1^\circ$ . The dataset provides ~~annual-total~~ total annual emissions by sector and is accompanied by temporal profiles by  
145 country ~~by~~ and sector to construct hourly and monthly emissions that can be used as model input.

### 2.1.1 GAINS

~~The methodology~~

#### GAINS

The methodology used in GAINS (Höglund-Isaksson et al., 2020) to estimate anthropogenic CH<sub>4</sub> emissions is based on the  
150 ~~recommendation recommendations~~ in the IPCC (2006) guidelines. For most sectors, GAINS uses country-specific information in such a way that the estimated emissions are consistent and comparable across geographic geographical and temporal scales. More advanced methods are ~~use~~ used to estimate emissions from the solid waste sector (Gómez-Sanabria et al., 2018) and fossil and gas systems (Höglund-Isaksson, 2017). In addition to past estimates, GAINS can be used to estimate future emissions based on ~~abatement measure mitigation~~ scenarios. The version of GAINS used in this study includes estimates for the countries that  
155 are part of the European Union, Norway, the ~~UK~~ United Kingdom and Switzerland. ~~The emissions~~ Emissions are estimated monthly ~~for 1990–2021 from 1990 to 2021~~ and the spatial resolution is  $0.1^\circ \times 0.1^\circ$ . The emissions from energy (upstream and downstream sources in fossil fuel extraction and use), agriculture (livestock, rice cultivation and agricultural waste burning) and waste handling (solid waste and wastewater) are estimated.

## 2.2 Atmospheric ~~inversion inverse~~ model CTE-CH<sub>4</sub>

160 The atmospheric ~~inversion inverse~~ model CTE-CH<sub>4</sub> (Tsuruta et al., 2017) is based on the Bayesian inversion approach, where optimised CH<sub>4</sub> fluxes are obtained by minimising the ~~mismatch between modelled and observed atmospheric observations, depending also on prior knowledge and uncertainties~~ cost function

$$J = (\mathbf{x} - \mathbf{x}^b)^T \mathbf{P}^{-1} (\mathbf{x} - \mathbf{x}^b) + (\mathbf{y} - H(\mathbf{x}))^T \mathbf{R}^{-1} (\mathbf{y} - H(\mathbf{x})), \quad (1)$$

where the first part of the right-hand side contains the state vector  $\mathbf{x}$  [dimension  $N \times 1$ ] of the scaling factors used to multiply  
165 by the CH<sub>4</sub> surface fluxes, the prior state vector  $\mathbf{x}^b$  and the prior error covariance matrix  $\mathbf{P}$  [ $N \times N$ ]. The second part contains the observation vector  $\mathbf{y}$  [dimension  $M \times 1$ ] and its error covariance matrix  $\mathbf{R}$  [ $M \times M$ ].  $H$  is the observation operator that samples atmospheric CH<sub>4</sub> at the measured location and times based on the states  $\mathbf{x}$ , which are then compared with the observations.

### 2.2.1 Data Assimilation

170 To optimise CH<sub>4</sub> fluxes, we use the CarbonTracker Data Assimilation Shell (CTDAS) (van der Laan-Luijckx et al., 2017) which  
is based on the fixed-lag ensemble Kalman filter (EnKF) (Evensen, 2003; Peters et al., 2005). Using EnKF, we represent the  
true state as an ensemble of sample states randomly drawn from a mean state ( $\bar{x}$ ) and based on covariance  $P$ . Each ensemble  
member is then optimised independently similarly to the Kalman filter (Kalman, 1960) (Peters et al., 2005). In this study, the  
size of the ensemble is 500 and we use a time lag of 5 weeks (Peters et al., 2005; Tsuruta et al., 2017). CTE-CH<sub>4</sub> provides  
175 optimises anthropogenic and natural CH<sub>4</sub> emission estimates fluxes simultaneously but separately at a spatial resolution of  
 $1^\circ \times 1^\circ$  (approximately  $110 \text{ km} \times 40\text{--}60 \text{ km}$  in Finland) over the northern high latitudes and at a temporal resolution of  
Europe, Russia, Canada and the USA, and regionally elsewhere (Supplementary Fig. S1). The spatial correlation is defined  
using an exponential decay model (Peters et al., 2005), with correlation lengths of 100 km for  $1^\circ \times 1^\circ$  grid-based optimisation  
domains, 500 km for other land domains and 900 km for oceanic domains. The anthropogenic and natural CH<sub>4</sub> fluxes are  
180 assumed uncorrelated, and domains between land and ocean are also assumed uncorrelated (see also Section 2.2.5). The  
temporal optimisation resolution is one week. CTE-CH<sub>4</sub> consists of prior flux maps representing different emission sources,  
the TM5

### 2.2.2 Atmospheric chemistry transport model TM5

We use the atmospheric chemistry transport model (Krol et al., 2005), and an ensemble Kalman filter data assimilation scheme  
185 (Peters et al., 2005; van der Laan-Luijckx et al., 2017). The size of the ensemble is 500 with a time lag of 5 weeks. The global  
horizontal resolution of TM5 is TM5 (Krol et al., 2005) as the observation operator. TM5 is an Eulerian model with a two-way  
nested zoom capability, i.e. we can model atmospheric transport at a higher resolution in a region of interest and have a  
coarser resolution globally. In this study, we simulate atmospheric transport in  $4^\circ \times 6^\circ$  (latitude  $\times$  longitude) but it includes  
resolution globally with a  $1^\circ \times 1^\circ$  zoom grid over Europe with  $(24^\circ \text{ N} \text{--} 74^\circ \text{ N}, 21^\circ \text{ W} \text{--} 45^\circ \text{ E})$  including a  $2^\circ \times 3^\circ$  zone  
190 around it intermediate zone around the zoom  $(14^\circ \text{ N} \text{--} 82^\circ \text{ N}, 36^\circ \text{ W} \text{--} 54^\circ \text{ E})$ . The vertical domain is divided into 25 hybrid sigma  
pressure levels from the surface to the upper atmosphere. We use ECMWF ERA5 meteorological data are used at with a 3-  
hour resolution (Hersbach et al., 2020). Calculations The calculations include atmospheric sinks from photochemical reactions  
involving with OH, Cl and O(<sup>1</sup>D). The reactions with OH is are calculated based on Houweling et al. (2014). For reactions  
with Cl and O(<sup>1</sup>D), we use two schemes: using prescribed reaction rates calculated from the atmospheric chemistry general  
195 circulation model ECHAM5/MESSy1 (Jöckel et al., 2006; Kangasaho et al., 2022) and reaction rates based on Brühl and  
Crutzen (1993). The atmospheric sink varies from month to month but does not include interannual variability.

### 2.2.3 Observations

We use observations from a global in situ measurement network that includes the NOAA National Oceanic and Atmospheric  
Administration GLOBALVIEWplus ObsPack v4.0 dataset (Schuldt et al., 2021) and data from the National Institute for Envi-  
200 ronmental Studies (JR-STATION: Japan-Russia Siberian Tall Tower Inland Observation Network, Ver1.2 (Sasakawa et al., 2010)  
; Sasakawa et al., 2010) and the Finnish Meteorological Institute (Tsuruta et al., 2019). Within Finland, measurements were  
collected at six sites located from southern to northern Finland, including urban, natural and marine areas (: KMP, PAL

**Table 1.** List of the Finnish surface observation sites used in inversions. Observation uncertainty (Obs. unc.) is used to define diagonal values in the observation covariance matrix. The data type is categorised into two measurements (discrete (D) and continuous (C)). All stations had data until the end of 2021. The contribution abbreviations are: the Finnish Meteorological Institute (FMI), National Oceanic and Atmospheric Administration Earth System Research Laboratories (NOAA ESRL), Integrated Carbon Observation System - Atmosphere Thematic Centre (ICOS-ATC), University of Eastern Finland (UEF) and University of Helsinki (UHELS).

Sitecode	Site Name	Contributor	Longitude [°E]	Latitude [°N]	Sampling height [m a.s.l.]	Obs. unc. [ppb]	Data type D/C	Date min. [year/month]
KMP	Kumpula	FMI	24.96	60.20	53.00	30.0	C	2010/01
PAL	Pallas-Sammaltunturi, GAW Station	NOAA ESRL	24.12	67.97	570.00	15.0	D	2001/12
PAL	Pallas-Sammaltunturi, GAW Station	ICOS-ATC, FMI	24.12	67.97	577.00	15.0	C	2004/02
PUI	Puijo	ICOS-ATC, UEF	27.66	62.91	84.00	30.0	C	2011/11
SMR	Hyytiälä	ICOS-ATC, UHELS	24.29	61.85	306.00	25.0	C	2012/12
SOD	Sodankylä	FMI	26.64	67.36	227.00	25.0	C	2012/01
UTO	Utö	ICOS-ATC, FMI	21.37	59.78	65.00	25.0	C	2012/03

(Hatakka, 2024), PUI (Lehtinen and Leskinen, 2024), SMR (Levula and Mammarella, 2024), SOD and UTO (Hatakka and Laurila, 2024) (Table 1 and Fig. 1). Globally, our dataset included 175 stations from 2000 to 2021 (Supplementary Fig. S1 and Table S1). Both weekly discrete air samples and hourly continuous measurements are filtered based on quality flags established by the respective institutions. To standardise the dataset, hourly continuous observations representing well-mixed atmospheric conditions were converted to daily averages, calculated from 12 to 4 pm local time, with exceptions except for high mountain sites, where averages were calculated from 0 to 4 am local time, following Tsuruta et al. (2017). Observational uncertainties were

Observational uncertainties, also referred to as "model-data mismatches", are quantified for each site, taking into account by considering site-specific characteristics and measurement accuracy, and also the model's ability to predict the ability of TM5 to simulate atmospheric CH<sub>4</sub> mole fractions (Bruhwiler et al., 2014; Tsuruta et al., 2017, 2019). Discrepancies between modelled and observed mole fractions are expected due to the resolution of TM5 and transport errors. For example, TM5 performs better in simulating mole fractions from remote marine background sites compared to sites influenced by strong local emissions. We classify the sites into different categories such as marine boundary layer (4.5 ppb), terrestrial (25 ppb), mixed marine and terrestrial (15 ppb) and strong local influence (30 ppb). The uncertainties range from 4.5 to 75 ppb between global sites for global sites (Supplementary Table S1) and from 15 to 30 ppb at for the Finnish sites (Table 1).

CTE-CH<sub>4</sub> optimises anthropogenic and natural fluxes simultaneously but separately at 1° × 1° resolution in Canada, the USA, Europe and Russia, and regionally elsewhere. The spatial correlation is defined using an exponential decay model (Peters et al., 2005), with correlation lengths of 100 km for 1° × 1° grid-based optimisation domains, 500 km for other land domains, and 900 km for oceanic domains.



## 2.2.4 Prior emissions

As an anthropogenic prior, EDGAR v6 (European Commission and Joint Research Centre et al., 2021) ~~was used. Additionally is used. In addition,~~ a modified version of EDGAR v6 is used, where emissions in Europe ~~were replaced with~~ are replaced by emissions from CAMS-REG, ~~was used~~. In Finland, ~~the~~ CAMS-REG emissions ~~were~~ are redistributed based on Statistics Finland's national GHG inventories of livestock and landfill ~~distributions~~ distribution (see details in ~~Tenkanen et al. (2024)~~ Tenkanen et al., 2024). For natural prior emissions, we ~~used~~ use estimates from two ecosystem models: Jena Scheme for Biosphere-Atmosphere Coupling in Hamburg with the Helsinki Model of MEthane buiLd-up and emISSION for peatlands module (JSBACH-HIMMELI (Raivonen et al., 2017; Kleinen et al., 2020) and-) (Raivonen et al., 2017; Kleinen et al., 2020) and the Land surface Processes and eXchanges with the Dynamical Peatland Model Based on TOPMODEL (LPX-Bern DYP-TOP version 1.4) v1.4 (Lienert and Joos, 2018; Stocker et al., 2014; Spahni et al., 2011, 2013), which include CH<sub>4</sub> emissions from peatlands and mineral ~~lands~~ soils as well as the soil sink. LPX-Bern DYPTOP also simulates emissions from inundated ~~lands~~ soils. In addition, ~~natural prior from Saunois et al. (2024) was used to the emissions from JSBACH-HIMMELI and LPX-Bern DYPTOP, the wetland prior (monthly averages from the 11 models used by Poulter et al., 2017) combined with the soil sink from Saunois et al. (2024) is used and referred here as the Global Carbon Project (GCP) prior.~~ For other a priori sources, ~~we used~~ we use estimates from GFED v4.1s (van der Werf et al., 2017) for fire, VISIT (Ito and Inatomi, 2012) or Saunois et al. (2020) for termites, those calculated ~~based on from~~ ECMWF data for ocean sources (Tsuruta et al., 2017) or from Weber et al. (2019), ~~and~~ Etiope et al. (2019) for geological emissions. We recognise that the emission categories referred to here as "other" are also from natural sources. However, as the optimised emissions using wetland emissions as the prior emissions are likely to include emissions from other sources (e.g. freshwater) as well as the soil sink, we refer to these emissions as "natural" in this study.

## 2.2.5 Prior uncertainty estimates

As default prior uncertainties, ~~we used for both anthropogenic and natural emissions, we use~~ 80 % for terrestrial fluxes and 20 % for oceanic fluxes, assuming uncorrelated uncertainties, following the practice established in previous studies (e.g. Tsuruta et al., 2017; Bruhwiler et al., 2014). Since the uncertainty depends on the prior flux, this means that when the prior flux is small, the uncertainty is also small, i.e. we trust the ~~emission~~ prior fluxes more. If we not only optimise the total emissions but also try to separate different categories such as anthropogenic and natural emissions, this could lead to a misallocation, especially in regions where both anthropogenic and natural sources are prominent. The optimisation may not be able to change the correct category because the uncertainties are relatively too small, or because the uncertainties in the other category are too large (and therefore the optimisation is more likely to change those emissions). ~~The process-model-based studies~~ Studies based on process-based models have shown that estimates of CH<sub>4</sub> emissions from wetlands vary substantially and inhomogeneously (e.g., a global annual average was 119–203 Tg yr<sup>-1</sup> from 2010 to 2019; Saunois et al., 2024) and thus have large uncertainties (Melton et al., 2013; Saunois et al., 2020; Ito et al., 2023; Chang et al., 2023). ~~Therefore, we took advantage of this large range of different estimates of wetland methane emissions and defined the uncertainty of the natural prior emissions based on a~~

255 ~~process-model-ensemble~~(Melton et al., 2013; Ito et al., 2023; Chang et al., 2023). We therefore let this guide our uncertainty estimates, assigning larger uncertainties to areas and time periods where the process-based models disagreed most, and smaller uncertainties where they agreed most.

To calculate the uncertainties, we ~~used-use~~ an ensemble of six process-based models (~~Global Carbon Project, Saunois et al. (2020)~~ from the GCP (Saunois et al., 2020). This ensemble ~~used prognostic runs in which consists of prognostic runs, i.e.~~ the models used their own internal approach to estimate the area ~~and dynamics~~ of wetlands. This ~~enabled-enables~~ us to account for the differences in the location of wetlands, which is one of the largest uncertainties in modelling regional wetland emissions. The following models ~~were included :- are included~~ LPX-Bern, JULES, ORCHIDEE, ELM, VISIT and LPJ-WSL. There ~~was-is~~ also a prognostic version of CLASS-CTEM, but we ~~excluded-exclude~~ it because of its coarse resolution and anomalously high values in the tropics. The average monthly values of the used process-based models in northern and southern Finland are shown in Supplementary Fig. ~~S1S2~~.

~~We made the test for the post-2010 period, so we calculated monthly averages for the 2010–2017 period . The uncertainty was then defined monthly and also independently for each optimisation~~ The uncertainty estimate based on a process-based model ensemble is calculated as follows: For each month  $t$  from January to December, we first calculate the monthly average fluxes  $\bar{F}(m, t, r)$  from the process-based model estimates  $F(m, t_j, r)$ , where  $m$  is a model,  $t_j$  is a month in a year  $j$  and  $r$  is a region. The inverse model run with these new uncertainty estimates extends from 2010 to 2021, but the process-based models only have estimates up to 2017. Thus, we calculate monthly averages from the process-based model estimates for the period 2010–2017:

$$\bar{F}(m, t, r) = \frac{1}{J} \sum_{j=1}^J F(m, t_j, r), \quad J = 8. \quad (2)$$

The quantile range  $U_q$  from the six model ensemble  $\mathbf{m}$  is then calculated for each month and optimisation region (see Section 2.2.1). The quantile range for each month and region ~~, i.e. at the resolution of  $1^\circ \times 1^\circ$  in the high northern latitudes and regionally in the other regions.~~ For the calculation of the uncertainties, we took the second and third quartiles from  $U_q(t, r)$  is defined as the range from the second ( $Q_{0.25}$ ) to the third ( $Q_{0.75}$ ) quartile of the monthly estimates of the process-based model, e.g., the range of the lowest and highest 25 %. ~~This value was then :-~~

$$U_q(t, r) = Q_{0.75}(\bar{F}(\mathbf{m}, t, r)) - Q_{0.25}(\bar{F}(\mathbf{m}, t, r)). \quad (3)$$

280 Lastly,  $U_q(t, r)$  is divided by the monthly averages of the natural prior used (LPX-Bern DYPTOP) in the inversion to obtain the ~~proportional values needed final uncertainty values  $U$ , which are used~~ to calculate the prior covariance matrix. ~~The maximum uncertainty was set at~~ However, we set the maximum uncertainty to 500 % and the minimum ~~at-to~~ 10 % ~~of the prior fluxes:~~

$$U(t, r) = \frac{U_q(t, r)}{\bar{F}(\text{LPX-Bern DYPTOP}, t, r)}, \quad U(t, r) \in [0.1, 5]. \quad (4)$$

**Table 2.** List of inversion setups. [The anthropogenic and natural priors used, the natural prior uncertainty calculation scheme used, and the simulation periods are specified.](#)

Inversion	Anthropogenic prior	Natural prior	Natural unc	Years
InvJSBACH_CAMSREG	CAMS-REG	JSBACH-HIMMELI	80 %	2004–2020
InvLPX_CAMSREG	CAMS-REG	LPX-Bern DYPTOP	80 %	2004–2020
InvLPX_EDGAR	EDGAR v.6	LPX-Bern DYPTOP	80 %	2000–2021
InvLPX_EDGAR_UNC	EDGAR v.6	LPX-Bern DYPTOP	varying	2010–2021
InvGCP_EDGAR	EDGAR v.6	Saunois et al. (2024)	80 %	2000–2021

## 285 2.2.6 Ensemble of CTE-CH<sub>4</sub> inversions

In this study, we formed an ensemble of five CTE-CH<sub>4</sub> inversion runs ~~which had with~~ different anthropogenic and natural priors, as well as different uncertainty estimates for the natural prior emissions. The ~~inversion time periods covered by each inverse model run also differed depending on the priors used and the time periods they covered.~~ The inverse model setups are listed in Table 2. The name of an inversion setup includes the ~~used prior prior used~~ (Inv<sub>natural\_anthropogenic</sub>). The experiment with varying natural uncertainty estimates was done ~~using with~~ the same priors as Inv<sub>LPX\_EDGAR</sub> and is noted ~~with by~~ adding a subscript "UNC" ~~at the end setups to the end of the setup~~ name (Inv<sub>LPX\_EDGAR\_UNC</sub>). Other differences between the setups (~~used priors priors used~~ and atmospheric sinks) are listed in Supplementary Table ~~S1 S2~~.


## 2.3 Ensemble of VERIFY inversions ~~Auxiliary CH<sub>4</sub> data~~

~~To help us interpret the CH<sub>4</sub> emissions estimated by the inverse model, we use auxiliary CH<sub>4</sub> datasets introduced below.~~

## 295 Ensemble of VERIFY inversions

The VERIFY project (<https://verify.lsce.ipsl.fr/>) was a Research and Innovation project funded by the European Commission under the H2020 programme in ~~2018–2022.~~ ~~Within the project, a system to estimate GHG 2018–2022.~~ The project developed ~~a GHG estimation system~~ to support NGHGI reporting ~~was developed.~~ ~~The project focused,~~ focusing on the three major anthropogenic GHGs: carbon dioxide, methane and nitrous oxide. Here ~~, we used~~ ~~we use the results of the CH<sub>4</sub> inversion model results inverse model~~ from the VERIFY project, ~~of which some were conducted some of which were performed~~ within the project and the rest of the estimates were ~~gathered collected~~ from other projects, such as ~~Global Carbon Project the GCP~~ (Saunois et al., 2020).

The VERIFY ensemble ~~consisted consists~~ of 14 CH<sub>4</sub> ~~inversion model estimates inverse model estimates,~~ but since one of them ~~was is~~ Inv<sub>GCP\_EDGAR</sub>, we ~~excluded that exclude it~~ from the VERIFY ensemble as it ~~was is~~ already included in our CTE-CH<sub>4</sub> ensemble. We only ~~investigated the examine~~ total CH<sub>4</sub> ~~emission emissions~~ because only 3 of the 13 VERIFY ensemble members ~~gave the partition to provide the partitioning between~~ natural and anthropogenic emissions. ~~There were three inversion~~



**Figure 1.** The land mask for Finland. Locations of six Finnish atmospheric observation sites (~~black-white dots and old bold text~~) and flux measurement sites (blue dots). The ~~squares show orange line shows~~ the division boundary between ~~northern and southern and northern~~ Finland.

~~model estimates which did~~ Three inverse model estimates do not include prior emission emissions estimates: two inversion runs with FLEXINVERT provided by the Norwegian research institute NILU, and one inversion run with FLExKF provided by the Swiss institute EMPA.

## 310 2.4 **Auxiliary CH<sub>4</sub> data**

### 2.3.1 **Eddy covariance CH<sub>4</sub> flux measurements**

#### Eddy covariance CH<sub>4</sub> flux measurements

To verify the ~~inversion model~~ CH<sub>4</sub> emissions, ~~we used of the inverse model, we use~~ eddy covariance measurements from two Finnish pristine open peatland sites, Lompolojankkä and Siikaneva. Lompolojankkä (68.0° N, 24.2° E) is located in northern Finland and Siikaneva (61.8° N, 24.2° E) in southern Finland (Fig. 1). A more detailed description of Lompolojankkä ~~has been is~~ given by Aurela et al. (2015) and ~~of~~ Siikaneva by Rinne et al. (2018). Eddy covariance is an atmospheric measurement technique ~~in which that frequently measures~~ vertical turbulent fluxes ~~are measured frequently~~ within the atmospheric surface layer. The footprint of the measurement, i.e., ~~where the where the measured~~ CH<sub>4</sub> fluxes ~~measured~~ originate, varies depending on the prevailing meteorological conditions but ~~is aimed aims~~ to cover the ~~whole entire~~ peatland ecosystem. The ~~measurement~~

315

320 ~~were measurements are~~ taken from the FLUXNET-CH<sub>4</sub> dataset (Delwiche et al., 2021), and the ~~gapfilled daily values were~~  
gap-filled daily values are used here to calculate monthly averages. Lompolojänkka ~~had has~~ data from 2006–2010 and Siikaneva  
from 2013–2018.

### 2.3.1 ~~Freshwater CH<sub>4</sub> emissions~~

~~Freshwaters were defined here similar to Saunois et al. (2020), including-~~

### 325 Freshwater CH<sub>4</sub> emissions

Freshwater is defined here similarly to Saunois et al. (2020) and includes lakes, ponds, reservoirs, streams and rivers. The  
freshwater CH<sub>4</sub> emission estimates ~~studied here were~~ examined here are from Stavert et al. (2022), which estimated the global  
annual freshwater CH<sub>4</sub> to be ~~of~~ 53 Tg yr<sup>-1</sup>.

### 2.3.1 ~~Biomass-burning CH<sub>4</sub> emissions~~

330 ~~Two-~~

### Biomass-burning CH<sub>4</sub> emissions


Two estimates of biomass-burning CH<sub>4</sub> ~~emission estimates were~~ emissions are used: GFED v4.1s (van der Werf et al., 2017),  
which ~~was is~~ also used as a prior in the inversions, and the Copernicus Atmosphere Monitoring Service (CAMS) Global Fire  
Assimilation System (GFAS) (Kaiser et al., 2012). GFED is provided in resolutions of 0.25° × 0.25° and GFAS in resolutions  
335 of 0.1° × 0.1° resolutions. GFED has ~~monthly and~~ GFAS a monthly temporal resolution and GFAS has a daily temporal  
resolution. We ~~aggregated both dataset~~ aggregate both datasets to 1° × 1° and monthly resolutions.

## 3 Results

### 3.1 Anthropogenic methane emission inventory estimates in Finland

In this section, we study the annual CH<sub>4</sub> emission estimates in Finland from six anthropogenic inventories. The spatial  
340 distributions of CAMS-REG, GAINS and EDGAR v7 are shown in Supplementary Fig. S4.

The annual emission estimates for the four main source categories defined in the 2006 IPCC Guidelines for National Green-  
house Gas Inventories (Eggleston et al., 2006) (energy, industrial processes and product use, agriculture and waste) are shown  
for each inventory in Fig. 2. ~~The spatial distribution of CAMS-REG, GAINS and EDGAR v6 can be found from Supplementary~~  
~~Fig S3.~~ Of the six inventories examined, NGHGI Fi, CAMS-REG and GAINS were in good agreement (Fig. 2a). CAMS-REG  
345 used reported Finnish national data ~~, and , overall , and overall~~ the emissions in the two inventories had similar magnitudes  
and trends. The small differences between the values examined here could be ~~because~~ explained by the fact that CAMS-REG  
has gridded estimates and the values in Finland were obtained using an area mask, ~~whereas while~~ NGHGI Fi is not spatially



**Figure 2.** Annual anthropogenic CH<sub>4</sub> emissions per source category in Finland reported by a) NGHGI Fi, CAMS-REG and GAINS, and b) EDGAR v6, v7 and v8. Note the different y-axis ranges. c) Annual total CH<sub>4</sub> emissions in all the six inventories.

distributed. GAINS emissions were at the same level as the other two, but there were some differences: waste emissions had a larger decreasing trend and agricultural emissions were ~~larger~~ higher from 2000 to 2015. The ~~driving cause for the~~ trend in  
350 GAINS was ~~declining driven by the decreasing~~ number of cows and cattle, ~~as was the case also in~~. This was also the case for  
NGHGI Fi. According to NGHGI Fi, the number of cattle decreased by more than ~~one-a~~ third between 1995 and 2021, but  
the decline slowed down after 2010. The ~~decline decrease in the~~ number of cattle has ~~also been counterbalanced by increased~~  
been offset by an increase in animal weight, growth and milk production ~~which has lead to larger emission factors, and thus,~~  
~~the magnitude agriculture,~~ which has led to higher emission factors so that the magnitude of agricultural CH<sub>4</sub> emissions has  
355 remained the same ~~during the latest years.~~ in recent years.

The three EDGAR versions differed from the other inventories but were similar ~~with~~ to each other. The magnitudes of  
CH<sub>4</sub> emissions of different emission categories were the same in EDGAR v6 and v7 ~~were the same~~ (Fig. 2b). The difference  
between the two versions was that v7 had a longer time series until 2021, while v6 ended in 2018. Agricultural and industrial  
emissions in all EDGAR versions were similar to those in the other inventories. Energy emissions in EDGAR v8 were ~~smaller~~  
360 lower than in v6 or v7 and showed no trend, ~~similar to NGHGI Fi, CAMS-REG and GAINS~~. However, the absolute magnitude  
was still three times higher in EDGAR v8 (0.03 Tg yr<sup>-1</sup>) than in ~~the other inventories~~ NGHGI Fi, CAMS-REG and GAINS  
(0.01 Tg yr<sup>-1</sup>). In EDGAR v6 and v7, energy emissions increased significantly from 2004 onwards. This increase was due to  
~~increased~~ higher estimates of fugitive methane emissions from oil refining and methane emissions from natural gas processing,  
transmission and distribution (Supplementary Fig. 2, S3). Waste emissions in EDGAR inventories, although decreasing over the  
365 period ~~2000–2021~~ 2000–2021, were about 4–5 times higher ~~in the EDGAR inventories~~ than in the other inventories, although  
they were lower in v8 than in v6 and v7.


Waste emissions were the largest source in all inventories at the beginning of the study period. Due to the decreasing trend  
of waste emissions, agriculture ~~was~~ became the largest source after 2008 ~~according in~~ GAINS and after 2009 ~~according to in~~  
NGHGI Fi. In CAMS-REG, ~~the~~ emissions from waste were higher than ~~the emissions~~ those from agriculture in 2006, otherwise  
370 ~~the emissions from agriculture were the highest~~ agriculture had the highest emissions.

The EDGAR estimates ~~stood out when examining the~~ stand out when looking at total annual emissions (Fig. 2c). While the  
average from ~~2000–2020~~ 2000 to 2020 was about 0.19 Tg yr<sup>-1</sup> in NGHGI Fi and GAINS, it was 0.76 Tg yr<sup>-1</sup> in EDGAR  
v7 and 0.58 Tg yr<sup>-1</sup> in EDGAR v8. The EDGAR inventories also showed a ~~larger interannual~~ higher inter-annual variability,  
especially in the waste sector, than the other inventory estimates.

## 375 3.2 Atmospheric ~~inversion~~ inverse model emission estimates in Finland

### 3.2.1 Annual estimates from CTE-CH<sub>4</sub>

In this section, we study the annual total, anthropogenic and natural CH<sub>4</sub> emissions in Finland and how the posterior emissions  
differed from the prior emissions depending on the ~~inversion~~ inverse model setup. The spatial distribution of the prior and  
posterior emissions ~~can be found from Supplementary Fig. S3–S6~~ is shown in Supplementary Figs. S4–S7.




**Figure 3.** Annual total CH<sub>4</sub> emission estimates from the five CTE-CH<sub>4</sub> inversion model runs. Prior estimates are shown with dashed lines and posterior estimates with emissions are shown as solid lines. The shaded areas around the posterior emissions show one standard deviation of the ensemble distributions. The right panel shows the mean prior and posterior estimates from emissions over the whole study period.

380 **Finland's annual total emission estimates.** The annual total emissions of Finland from the five CTE-CH<sub>4</sub> inverse model runs are shown in Fig. 3. The prior emissions had evidently a strong influence on the posterior emissions. The range of the total prior emissions was large, but the range of the optimised emissions was smaller after 2009 and until 2020 (the average range was, with an average range of 0.57 Tg yr<sup>-1</sup> in 2009–2020) than, while the range of prior emissions (the prior emissions was 0.69 Tg yr<sup>-1</sup>) in the same period. The inversions using LPX-Bern DYPTOP as the natural prior estimates had the highest (InVLPX\_EDGAR, 1.1 Tg yr<sup>-1</sup> on average) and lowest (InVLPX\_CAMSREG, 0.51 Tg yr<sup>-1</sup> on average) total emission estimates, while the three estimates between them agreed well, especially after 2016. To better explain the differences between the total emission estimates, the anthropogenic and natural CH<sub>4</sub> emissions are studied separately below.

Magnitudes of the two anthropogenic posterior emission estimates using CAMS-REG were similar and slightly higher than CAMS-REG (Fig. 4). The optimised results using EDGAR v6 varied more but all posterior emissions were higher than EDGAR v6 until 2009 and lower thereafter until 2020 or 2021, bringing the posterior estimates of the five inversions closer together compared to their prior estimates. The two anthropogenic posterior emissions using EDGAR v6 combined with the default uncertainty estimate for the natural prior (InVLPX\_EDGAR and InVGCP\_EDGAR) had similar anthropogenic emission es-





**Figure 4.** Annual anthropogenic CH<sub>4</sub> emission estimates from the five CTE-CH<sub>4</sub> inversion-inverse model runs. Prior estimates emissions are shown with as dashed lines and posterior estimates with emissions as solid lines. Shaded The shaded areas around the posterior emissions show one standard deviation of the ensemble distributions. The right panel shows the mean prior and posterior estimates from emissions over the whole study period.


timates, but the inversion with varying uncertainty estimates for the natural prior (InvLPX\_EDGAR\_UNC) had lower optimised anthropogenic emission estimates than the other two, especially after 2016.

395 ~~Natural~~ The natural posterior emissions were always higher than the prior used, regardless of the natural prior, except in 2005 and 2006 when JSBACH-HIMMELI was higher (Fig. 5). The order of the emission estimates was also preserved after the optimisation: maintained after optimisation: the posterior emissions of InvJSBACH\_CAMSREG were the highest, and InvGCP\_EDGAR the smallest and were the lowest. The three posterior emissions using LPX-Bern DYPTOP as the prior, were in between the two estimates with the varying uncertainty estimates a prior lay between these two estimates, with the inversion using the prior uncertainty estimates based on the ensemble of process-based models (InvLPX\_EDGAR\_UNC) being the smallest estimates giving the lowest estimates of the three. Since the estimates using the GCP prior did not change a lot much, but the optimised emissions with JSBACH-HIMMELI were increased, the range of natural posterior emissions (0.08–0.44 Tg yr<sup>-1</sup>) was larger than the range of prior emissions (0.08–0.39 Tg yr<sup>-1</sup>).

400

Even though the absolute magnitudes of total CH<sub>4</sub> emissions and the partition between anthropogenic and natural emissions differed between inversion runs, the trends of the emission estimates were more aligned similar after the optimisation. All the (Table 3). All anthropogenic posterior emissions had decreasing trends, even though, although there was no significant trend

405



**Figure 5.** Annual natural CH<sub>4</sub> emission estimates from the five CTE-CH<sub>4</sub> inversion model runs. Prior estimates are shown with dashed lines and optimised estimates with posterior emissions as solid lines. The shaded areas around the posterior emissions show one standard deviation of the ensemble distributions. The right panel shows the mean prior and optimised estimates from posterior emissions over the whole study period.

**Table 3.** Linear trends [Gg yr<sup>-1</sup>] and their p-values (in brackets) for anthropogenic, natural and total CH<sub>4</sub> emission estimates in Finland. Values for prior and optimised estimates from the five CTE-CH<sub>4</sub> inversion runs are shown. Statistically significant trends (p-value smaller than 0.05) are **bolded**.

	Anthropogenic		Natural		Total	
	Prior	Optimised	Prior	Optimised	Prior	Optimised
InvJSBACH_CAMSREG	<b>-2.9 (0.00)</b>	<b>-2.2 (0.00)</b>	0.8 (0.45)	<b>11.3 (0.00)</b>	-2.0 (0.07)	<b>9.1 (0.00)</b>
InvLPX_CAMSREG	<b>-2.9 (0.00)</b>	<b>-1.7 (0.01)</b>	-1.9 (0.16)	1.1 (0.64)	<b>-4.7 (0.00)</b>	-0.6 (0.78)
InvLPX_EDGAR	1.8 (0.36)	<b>-5.6 (0.03)</b>	0.1 (0.87)	<b>4.9 (0.00)</b>	2.0 (0.36)	-0.7 (0.84)
InvLPX_EDGAR_UNC	<b>-9.6 (0.03)</b>	-14.5 (0.07)	0.2 (0.92)	1.2 (0.64)	-9.3 (0.06)	-13.2 (0.13)
InvGCP_EDGAR	1.2 (0.56)	<b>-7.4 (0.03)</b>	-0.2 (0.12)	0.3 (0.15)	1.0 (0.63)	<b>-7.2 (0.03)</b>


in EDGAR v6 ~~in-for~~ 2000–2021. ~~Likewise~~Similarly, there were no significant trends in the natural prior emissions, but ~~in the optimised natural emissions~~, there were small positive trends in the optimised natural emissions with InvJSBACH\_CAMSREG (0.01 Tg yr<sup>-1</sup>, p-value 0.0003) and InvLPX\_EDGAR (0.005 Tg yr<sup>-1</sup>, p-value 0.004). Decreasing anthropogenic emissions and  
410 increasing natural emissions cancelled each other out, so that the only statistically significant trends in ~~the~~ total emissions were ~~noted-found~~ in InvGCP\_EDGAR (-0.007 Tg yr<sup>-1</sup>, p-value 0.03) and InvJSBACH\_CAMSREG (0.009 Tg yr<sup>-1</sup>, p-value 0.001). The signs of the trends were ~~the~~ opposite, reflecting the ~~partition~~partitioning between natural and anthropogenic emissions: InvGCP\_EDGAR had the highest anthropogenic emissions and the ~~smallest natural emissions~~lowest natural emissions, while InvJSBACH\_CAMSREG had the highest natural emissions and the ~~smallest~~lowest anthropogenic emissions.

### 415 3.2.2 Years 2020 and 2021

During the last two years of the study period, 2020–2021, the growth rate of the global atmospheric CH<sub>4</sub> was remarkably high (Lan et al., 2024; Nisbet et al., 2023). Although our inversion results did not show exceptionally high CH<sub>4</sub> emissions during this period, there were still some consistent signals from the inversion estimates. In 2020, all total posterior emissions were higher than in 2019. This increase was due to ~~an-increase~~increases in both anthropogenic and natural emissions, except for  
420 InvGCP\_EDGAR, where the increase was due to anthropogenic emissions alone. However, its natural prior and posterior emissions were ~~small~~low compared to the other estimates. In contrast to InvGCP\_EDGAR, the natural posterior emissions using JSBACH-HIMMELI as the natural prior (InvJSBACH\_CAMSREG) were ~~the-largest~~highest in 2020.

In 2021~~there were~~, there were only three posterior emission estimates: InvLPX\_EDGAR, InvLPX\_EDGAR\_UNC and InvGCP\_EDGAR. ~~All The three~~ optimised total emissions were still higher than in 2019, but ~~the differences between 2020 and emissions from~~  
425 ~~InvGCP\_EDGAR were lower in 2021~~ diverged and the differences in than in 2020, while inversions with LPX-Bern DYPTOP were higher in 2021. The partitioning to natural or anthropogenic ~~partitioning were also inconsistent. However~~ was also inconsistent across the three inversion estimates: InvGCP\_EDGAR had both lower anthropogenic and natural emissions in 2021 than in 2020, while in InvLPX\_EDGAR\_UNC it was the other way round. At the same time, the anthropogenic emissions of InvLPX\_EDGAR were higher and the natural emissions were lower in 2021 than in 2020. However, the differences between 2021 and 2020 were  
430 similar in magnitude to previous years.


In 2021, all estimates were higher than the prior total emissions, and in particular, the anthropogenic posterior estimates were close to the anthropogenic prior estimates. Part of the reason for this may be due to the high biomass-burning ~~emission estimates~~  
emissions in GFEDv4.1s, which seemed to affect at least the emissions in InvLPX\_EDGAR\_UNC. The natural posterior emission estimates of InvLPX\_EDGAR\_UNC in high northern latitudes (north of 50° N) were substantially higher than the emissions from  
435 LPX-Bern DYPTOP in 2016–2020, but in 2021 the emissions decreased by 15 Tg ~~from~~compared to 2020 (Supplementary Fig. ~~S7~~S7). ~~In the GFED, S8~~, CH<sub>4</sub> emissions of GFED in the high northern latitudes were 8.6 Tg in 2021 compared to 3.3 Tg yr<sup>-1</sup> in 2016–2020 (Supplementary Fig. ~~S8~~S8). The high ~~biomass-burning emissions in~~emissions from biomass-burning in the high northern latitudes most likely also affected the methane budget estimates in Finland, although there were no major forest fires in Finland. ~~The emission~~Emission estimates from another biomass-burning dataset, GFAS, were also high in 2021, but only  
440 about half of the GFED estimates (4.9 Tg, Supplementary Fig. ~~S8~~S8).



**Figure 6.** Estimated total annual CH<sub>4</sub> emissions in Finland in ~~2000–2021~~2000–2021, estimated using ~~inversion~~inverse models. Two ensembles are included: CTE-CH<sub>4</sub> and VERIFY. The circle (prior) and diamond (optimised) symbols indicate the ensemble means. The lowest and highest emission estimates are indicated by the lower and upper ends of the bars. Values for the priors are shown in yellow (VERIFY) and red (CTE-CH<sub>4</sub>) and values for the optimised estimates are shown in light blue (VERIFY) and dark blue (CTE-CH<sub>4</sub>). The mean values of the priors are connected to the mean values of the posteriors by a line. The colour of the line indicates whether ~~the~~ emissions have increased (red) or decreased (blue) compared to the prior. The number below the year ~~shows~~gives the number of ensemble members ~~of in the~~ CTE-CH<sub>4</sub> (dark blue) and VERIFY (light blue) ensembles for that year. The right panel shows the mean ~~mean~~mean-values ~~from over~~from over the whole study period.

### 3.2.3 Comparison of CTE-CH<sub>4</sub> and VERIFY ensembles

We also compared the CTE-CH<sub>4</sub> emission estimates ~~to~~with the inversion results from the VERIFY project (Fig. 6). In the CTE-CH<sub>4</sub> ensemble, the average of the total CH<sub>4</sub> emissions from ~~2000–2021~~2000–2021 was 0.83 Tg yr<sup>-1</sup> (average minimum and maximum range was 0.51–1.10 Tg yr<sup>-1</sup>) in the prior and 0.85 Tg yr<sup>-1</sup> (0.59–1.08 Tg yr<sup>-1</sup>) in the posterior  
445 estimates. In the VERIFY ensemble, the prior emissions were 1.50 Tg yr<sup>-1</sup> (0.78–2.28 Tg yr<sup>-1</sup>), almost twice that of the CTE-CH<sub>4</sub> ensemble, but the posterior emissions were reduced to 0.98 Tg yr<sup>-1</sup> (0.61–1.43 Tg yr<sup>-1</sup>), bringing the total CH<sub>4</sub> emission estimates close to the CTE-CH<sub>4</sub> ensemble and within the range of the posterior CTE-CH<sub>4</sub> ensemble estimates. The ranges of the posterior emission were large, but the range was ~~notably~~considerably smaller than the range of ~~prior estimates,~~ especially with the VERIFY ensemble. The average of the posteriors in the VERIFY ensemble was close to the average of  
450 the posteriors in the CTE-CH<sub>4</sub> ensemble and within the range of the posterior CTE-CH<sub>4</sub> ensemble the prior estimates. The lowest ~~emission estimates~~posterior emissions from both ensembles were approximately 0.6 Tg yr<sup>-1</sup> but the upper limit, but the highest emissions differed by 0.3 Tg yr<sup>-1</sup>.



**Figure 7.** Average monthly natural  $\text{CH}_4$  emission estimates from the five CTE- $\text{CH}_4$  inversion-inverse model runs in a) northern and b) southern Finland in 2010-2020. Prior estimates are shown with as dashed lines and optimised estimates with are shown as solid lines. The shaded areas show the smallest minimum and the largest maximum monthly posterior emission estimates. Freshwater emissions from Stavert et al. (2022) are shown with the dash-dot red dash-dotted lines.

### 3.3 Seasonal cycle of methane emissions

Methane emissions, and especially emissions especially those from natural sources, have a strong seasonal cycle, and in addition to estimating the magnitude of the emissions, it is essential important to have a correct estimate of the timing of the  $\text{CH}_4$  emissions. We calculated the monthly  $\text{CH}_4$  emissions to see how the optimisation affected the seasonal cycle and compared the seasonal cycles of the natural emission estimates with the flux measurements from two Finnish pristine peatlands. Since the climate in southern and northern Finland is different, and thus the timing of the natural  $\text{CH}_4$  emissions in southern and northern Finland is also different also differs, we divided the emissions from  $64^\circ\text{N}$  according to the division used, for example, in the Finnish NGHGI (Fig. 1). We focused on studying the emissions during the common time period between all inversion runs, 2010–2020.


The average, maximum and minimum monthly optimised natural emissions and the average of the prior emissions in 2010-2020 are shown in Fig. 7. There was a clear difference between the natural emissions in northern and southern Finland, as the maximum monthly emission estimate was almost  $0.12 \text{ Tg month}^{-1}$  in the north and only half of that in the south, although both occurred in July. In addition, the timing of the maxima of the posterior emissions differed between south and the south

and the north, whereas the maxima in the priors ~~differed only in LPX. In LPX~~ were different only in the LPX-Bern DYPTOP: In LPX-Bern DYPTOP, the maximum was either in August or September in the north and in September or October in the south, while in JSBACH-HIMMELI and the natural prior GCP, the maximum was always in July. In northern Finland, the maximum of the ~~Inv~~Inv<sub>GCP\_EDGAR</sub> emissions did not change from July, but the maximum of the ~~Inv~~Inv<sub>JSBACH\_CAMSREG</sub> emissions was in August rather than in July. In southern Finland, the timing of the emissions did not change much from the priors, except ~~in~~ for Inv<sub>JSBACH\_CAMSREG</sub>, where the posterior emissions were ~~shifted slightly~~ shifted towards late summer.

In northern Finland, the posterior emissions using LPX-Bern DYPTOP (Inv<sub>LPX\_CAMSREG</sub>, Inv<sub>LPX\_EDGAR</sub>, Inv<sub>LPX\_EDGAR\_UNC</sub>) had the largest increase from the ~~prior in July–September~~ priors in July to September so that the maximum was also shifted earlier towards late summer, although September was still the maximum in half of the years (Fig. 7a). This shift was less pronounced in Inv<sub>LPX\_EDGAR\_UNC</sub>. The relative uncertainty estimates of the natural prior emissions in Inv<sub>LPX\_EDGAR\_UNC</sub> varied monthly and were defined independently for each 1°×1° grid cell in Finland. This meant that whether the assigned uncertainty was larger or smaller than the constant 80 % used in the other inversions also depended on the month and location. During the winter months (From November to January), the uncertainty was smaller almost everywhere in Inv<sub>LPX\_EDGAR\_UNC</sub> than in Inv<sub>LPX\_EDGAR</sub>. In February and March, both the natural Inv<sub>LPX\_EDGAR\_UNC</sub> and Inv<sub>LPX\_EDGAR</sub> had low natural prior CH<sub>4</sub> emissions and the uncertainties were small regardless of the uncertainty estimates used small uncertainties. From April to June, the uncertainty in Inv<sub>LPX\_EDGAR\_UNC</sub> was larger in northern Finland and smaller in southern Finland, but this did not have a strong effect on the posterior emissions which stayed, which remained close to the prior (Fig. 7a). From July to October, the uncertainty in the north was much smaller in Inv<sub>LPX\_EDGAR\_UNC</sub> than in the other inversions, especially in grid cells where the natural prior emissions were high. Thus, the optimisation was more constrained by the prior than when the constant default uncertainty was used. However, southern Finland had a larger uncertainty in summer and autumn. As the optimisation had more freedom to adjust the CH<sub>4</sub> emissions in southern Finland in Inv<sub>LPX\_EDGAR\_UNC</sub>, it could also give more weight to the observation in the south. The natural posterior CH<sub>4</sub> emissions in the south did not differ between Inv<sub>LPX\_EDGAR\_UNC</sub> and Inv<sub>LPX\_EDGAR</sub>, but the anthropogenic posterior emissions were smaller lower in Inv<sub>LPX\_EDGAR\_UNC</sub> than in Inv<sub>LPX\_EDGAR</sub>, especially in July–October (Supplementary Fig. S9S10). The smaller lower natural emissions in the north and anthropogenic emissions in the south led to smaller resulted in lower total emissions in Inv<sub>LPX\_EDGAR\_UNC</sub> compared to Inv<sub>LPX\_EDGAR</sub>, and brought the seasonal cycle of total CH<sub>4</sub> emissions of Inv<sub>LPX\_EDGAR\_UNC</sub> close to the seasonal cycle of Inv<sub>JSBACH\_CAMSREG</sub> (Fig. 8a).

A comparison of the seasonal cycles of the total CH<sub>4</sub> emissions between the CTE-CH<sub>4</sub> and VERIFY ensembles (Fig. 8b) shows similar results to the comparison of the annual totals: the VERIFY prior emissions were larger higher on average and had a wider range than the prior prior emissions in the CTE-CH<sub>4</sub> ensemble, but the amplitudes of the seasonal cycles of the average posterior emissions agreed well. However, the phases of the average posterior emissions differed: the VERIFY ensemble was consistently one month ahead of the CTE-CH<sub>4</sub> ensemble.


In Fig. 7, there are also shown also shows the freshwater CH<sub>4</sub> emissions from Stavert et al. (2022). These emissions were not included in the prior emissions of Inv<sub>GCP\_EDGAR</sub> or in the other inversions. In northern Finland, the estimated freshwater emissions were considerably smaller lower than the posterior natural emission estimates (except for Inv<sub>GCP\_EDGAR</sub>), but in southern Finland, where there are many shallow lakes, they were larger than almost as high as the posterior natural emis-



**Figure 8.** a) Average monthly total CH<sub>4</sub> emission estimates from the five CTE-CH<sub>4</sub> inversion-inverse model runs in Finland from 2010-2020. Prior estimates are shown with as dashed lines and optimised estimates with are shown as solid lines. The shaded areas show the smallest lowest and the largest highest monthly posterior emission estimates. b) Average over of the monthly total CH<sub>4</sub> emission estimates from the CTE-CH<sub>4</sub> and VERIFY ensembles in Finland from 2010-2020. The shaded areas show the smallest minimum and the largest maximum monthly estimates.

sions: the freshwater emissions were 0.180.15 Tg yr<sup>-1</sup>, while the JSBACH-HIMMELI emissions were 0.13 Tg yr<sup>-1</sup> and the optimised natural emissions in InvJSBACH\_CAMSREG were 0.16 Tg yr<sup>-1</sup> on average in 2010-2020.

We also compared the seasonal cycles of the natural emission estimates with flux measurements from two Finnish pristine peatlands. Figure 9 shows the measured CH<sub>4</sub> fluxes from two Finnish pristine peatlands: Lompolojännkä (northern Finland) and Siikaneva (southern Finland). In At Lompolojännkä, the different years had very similar seasonal cycles and the maximum was in August, except in 2008 when the maximum was in July. In Siikaneva, the At Siikaneva, CH<sub>4</sub> fluxes had more annual variation varied more from year to year. Nevertheless, the maximum of the fluxes was relatively consistent in being in July, except in 2015 and 2016 when it was or in August. In both peatlands, July and August were the months with the highest CH<sub>4</sub> fluxes. Due to the alterations made by the inversion-inverse model, the seasonal cycles of the optimised emission estimates were more consistent with the flux measurements than the seasonal cycles of the prior estimates. However, it should be noted, however, that the inversion that the inverse model estimates aggregate much larger areas and a variety of sources rather than a single peatland, so a direct comparison between the flux measurements and inversion-the inverse model results is not possible.



**Figure 9.** Average monthly CH<sub>4</sub> flux measurements at a) Lompolojankkä (northern Finland) and b) Siikaneva (southern Finland). The shaded areas show a 95 % confidence interval.

### 3.4 Comparison of modelled methane mole fractions to observations in Finnish sites

The CH<sub>4</sub> emission estimates in Finland varied depending on the priors and prior uncertainty estimates used. To **have an insight** **into understand** which inversion best estimated CH<sub>4</sub> emissions in Finland, we compared modelled **mole fractions with and** observed mole fractions at the six Finnish **in-situ sites that were in situ sites** also used in the optimisation. We examined only the years 2010-2020, as these were common to all inversion runs. Observations from Utö were included from March 2018 and Hyytiälä from December 2016, **since as** all inversions included observations from these two sites. The effect of including all available years was also examined, but **there was no significant difference** **it did not change the result significantly**. In addition **to the optimised mole fractions**, we also studied the mole fractions modelled with the prior emissions using a so-called forward run mode, i.e. using only the TM5 transport model without the data assimilation. The term "prior" refers to these modelled mole fractions in this section. Similarly, the term "posterior" is used to refer to the mole fractions obtained using the optimised emissions.

In Fig. 10, correlation coefficients, detrended root mean square errors (RMSE) and standard deviations are shown **for all** **Finnish sites** as Taylor's diagrams (Taylor, 2001) **for all Finnish sites**, comparing both the prior and posterior mole fractions. The correlation coefficient describes the linear relationship between the modelled and observed mole fractions, with values close to one indicating good agreement between the two. The detrended RMSE quantifies and summarises the differences between the modelled and observed mole fractions. From the RMSE alone, it is not possible to determine whether the differences




are due to different phases in the ~~datasets or due data sets or~~ to differences in the amplitudes of the variations. The standard deviation provides additional information ~~, and it tells and shows~~ how much variation ~~there~~ is in each dataset.

The statistics of the priors varied more than the statistics of the posterior mole fractions (Fig. 10, see also Supplementary Fig. ~~S10~~S11). The prior values of  $\text{Inv}_{\text{GCP\_EDGAR}}$  were in better agreement with the observations than the other priors ~~in at~~ almost all observation sites, especially ~~in Puijo, Hyytiälä and Pallas~~ at PUI, SMR and PAL. However, in contrast to the other ~~inversion inverse~~ model setups, the posterior statistics of  $\text{Inv}_{\text{GCP\_EDGAR}}$  improved only slightly from the priors at ~~Pallas and Sodankylä~~ PAL and SOD, two northern stations surrounded by natural  $\text{CH}_4$  sources. Overall, the posterior mole fraction from different ~~inversion inverse~~ model runs showed similar statistics, especially at the ~~Utö and Hyytiälä~~ UTO and SMR stations.

To summarise the statistics of the optimised mole fractions, we ranked selected statistics as follows: For each site and inversion run, the bias, the detrended RMSE and the detrended linear correlation coefficient R were calculated. The bias was used instead of the standard deviations used in Taylor's diagrams to emphasise any systematic errors in the modelled mole fractions. Detrending the data removes long-term variations and allows us to examine short-term variations. ~~The detrending was done~~ We detrended the data using the method introduced by Thoning and Tans (1989), which takes into account a seasonal cycle. The absolute value of each variable was then ranked between the inversion runs from one to five, with the smallest being the best value for the bias and the detrended RMSE and the largest being the best value for the detrended R. The average of the three rankings for each inversion run, as well as the average of all six stations, is shown in Fig. 11. The same figure, ~~but with~~ but with prior statistics, is shown in Supplementary Fig. ~~S10~~S11.

Based on the average rankings, there was no single inversion setup that stood out as the best across all sites. Most inversion runs had both better and worse rankings ~~, depending on the site~~. However,  $\text{Inv}_{\text{LPX\_EDGAR\_UNC}}$  had the best rankings in general (average 2.11) and especially in the southern sites (~~Utö, Kumpula and Hyytiälä~~ UTO, KMP and SMR). In the northern sites (~~Pallas and Sodankylä~~ PAL and SOD), where natural  $\text{CH}_4$  sources dominate,  $\text{Inv}_{\text{JSBACH\_CAMSREG}}$  had the best rankings.  $\text{Inv}_{\text{JSBACH\_CAMSREG}}$  also had the ~~second best~~ second best overall ranking (2.61). The seasonal cycles of the optimised total  $\text{CH}_4$  emissions of  $\text{Inv}_{\text{LPX\_EDGAR\_UNC}}$  and  $\text{Inv}_{\text{JSBACH\_CAMSREG}}$  were also quite similar (Fig. 8a).  $\text{Inv}_{\text{GCP\_EDGAR}}$  had the best rankings in ~~Puijo and in Hyytiälä~~ PUI and SMR, where the prior statistics already showed a good agreement with the observations, but the worst rankings in the northern peatland sites.  $\text{Inv}_{\text{LPX\_CAMSREG}}$ , which had the ~~smallest lowest~~ total posterior emissions, had the worst rankings in general, and  $\text{Inv}_{\text{LPX\_EDGAR}}$ , which had the ~~largest highest~~ total posterior emissions, had average rankings across all sites.



**Figure 10.** Taylor's diagram of the results of the five CTE-CH<sub>4</sub> inversion-inverse model runs against the measured mole fractions from the Finnish stations: a) UtöUTO, b) KumpulaKMP, c) PuijoPUI, d) HyytiäläSMR, e) PaHas-SammaltunturiPAL and f) SodankyläSOD. Smaller circles respond-correspond to values from forward modelling results using the TM5 transport model TM5 and the prior emissions. Prior-The prior values of Inv<sub>LPX\_EDGAR\_UNC</sub> is the same for Inv<sub>LPX\_EDGAR</sub> and Inv<sub>LPX\_EDGAR\_UNC</sub> as they had the same prior emissions.

## 4 Discussion

### 4.1 Total methane emission estimates

We estimated methane emissions in Finland using the atmospheric inversion-inverse model CTE-CH<sub>4</sub>. As a global model, it was able to constrain the global total emissions well (on average 525 Tg yr<sup>-1</sup> with a minimum and maximum range of 560 3.2 %). However, the ratio of the range to the average total emissions in Finland was much larger at 58 % (71 % in the priors), which shows the difficulty of constraining emissions at the country level and also how the underlying prior emissions and their distribution affect emission estimates at a smaller scale. Nonetheless, by using a global model, optimising-the optimisation

Inversion run	Site						
	UTO	KMP	PUI	SMR	PAL	SOD	Mean
Inv <sub>JSBACH_CAMSREG</sub>	5.00	3.33	2.00	3.33	1.00	1.00	2.61
Inv <sub>LPX_CAMSREG</sub>	3.00	4.33	5.00	5.00	2.67	3.67	3.94
Inv <sub>LPX_EDGAR</sub>	3.33	2.33	3.67	3.67	3.00	2.33	3.06
Inv <sub>LPX_EDGAR_UNC</sub>	1.00	1.33	2.33	1.67	3.33	3.00	2.11
Inv <sub>GCP_EDGAR</sub>	2.67	3.67	2.00	1.33	5.00	5.00	3.28

**Figure 11.** The average rank calculated for each site-inverse model run for each inversion-run-site is shown. The bias, ~~the~~-detrended RMSE and ~~the~~-detrended R of the modelled and measured mole fractions were calculated with each inversion model run in each site, and the values were then ranked between the model estimates (the smallest-lowest being the best with-in bias and RMSE and the highest being the best with-in R). AdditionallyIn addition, the right-most-rightmost column is the average over-of all sites-site averages.

of emissions in a region of interest is not separated from ~~emissions occurring outside~~ the emissions that occur outside of the region.

565 The range of posterior CH<sub>4</sub> emissions in Finland was ~~also~~-large in the VERIFY ensemble (Figure 6), which included estimates from different inversion-inverse models and model runs, some of which used the same priors and observations, and some of which had their own setups. The range of prior estimates was even wider, and ~~since-as~~ the optimisation always reduced the estimates, the largest-highest prior estimates were most likely too high. Furthermore, the largest-highest CTE-CH<sub>4</sub> estimate (Inv<sub>LPX\_EDGAR</sub>), which was lower than the largest-highest estimate in the VERIFY ensemble, ~~and~~-showed only moderate agreement with atmospheric CH<sub>4</sub> observations, indicating ~~probably-too-high-that~~ CH<sub>4</sub> emissions were probably too high. Thus, the CTE-CH<sub>4</sub> ensemble range seemed ~~to-be~~-more reliable, especially ~~if-its-largest-estimate-were-excluded-when~~ excluding its highest estimate. Although the ranges of posterior emissions were large, the averages of the VERIFY and CTE-CH<sub>4</sub> ensembles ~~agreed-well-were~~ in good agreement. This is consistent with previous model intercomparisons which have shown that inversions can constrain emissions on a larger scale and that ensembles of inversion-inverse model estimates are more reliable and robust than estimates from a single inversion run (~~?Saunois et al., 2020; Stavert et al., 2022~~) (Petrescu et al., 2024; Saunois et al., 2020; Stavert et al., 2022). Further partitioning into independent countries still relies on the distributions of the priors.

575

## 4.2 Partition to anthropogenic and natural emissions

When comparing CH<sub>4</sub> estimates from the ~~inversion-inverse~~ model with national GHG inventories, it is important to understand not only the total CH<sub>4</sub> budget ~~but~~ also the partitioning of emissions reported in the inventories. As these inventories only ~~account for cover~~ anthropogenic activities, the share of anthropogenic emissions in the total CH<sub>4</sub> estimates is particularly important. In CTE-CH<sub>4</sub>, emissions from anthropogenic and natural ~~emissions-sources~~ were optimised separately but simultaneously. The emissions from both categories were analysed as they were refined by CTE-CH<sub>4</sub>.

The anthropogenic emission inventories gave two drastically different estimates of Finnish CH<sub>4</sub> emissions, with EDGAR ~~v6, v7 and v8~~ giving much higher estimates than the other ~~inventories~~ ~~three, NGHGI Fi, CAMS-REG and GAINS~~ (Figure 2c). Olhoff et al. (2022) compared the ~~NGHGIs-NGHGI~~ estimates with EDGAR v6 in the Nordic countries and showed that the CH<sub>4</sub> estimates from EDGAR v6 were much higher than the NGHGI estimates from Finland, Norway and Sweden. They showed that the discrepancies were due to fugitive emissions (in Norway and Finland) and waste emissions (in Sweden and Finland). ~~Additionally~~ ~~In addition~~, using Bayesian ~~inversion-inverse~~ modelling, Worden et al. (2022) estimated Finland's waste emissions to be  $0.11 \pm 0.29$  Tg in 2019 instead of the prior value of  $0.60 \pm 0.36$  Tg (EDGAR v4.3.2 in 2012), which is much more consistent with the NGHGI Fi (0.06 Tg). Saboya et al. (2022) compared modelled mole fractions using an older version of EDGAR (v4.3.2) with observations in London and found that emissions from the waste sector were ~~large-high~~ and inconsistent with their estimates. As a global product, EDGAR uses globally consistent methodologies. ~~As a result, there may have been a lack of consideration of,~~ ~~meaning~~ country-specific mitigation strategies ~~may not have been considered~~. For example, fugitive emissions from the oil and gas sector in EDGAR v6 ~~followed~~ followed the trend of activity data in the Nordic countries, indicating that efforts to reduce emissions ~~have not been were not~~ taken into account (Janssens-Maenhout et al., 2019; Olhoff et al., 2022). However, in the latest update of EDGAR v8 (European Commission and Joint Research Centre et al., 2023), there seems to be an improvement in the estimates for the energy sector, as they show the same trend as the other inventories in Finland (European Commission and Joint Research Centre et al., 2023) (Figure 2c).

Based on the comparison between atmospheric mole fractions modelled with CTE-CH<sub>4</sub> and observations from Finnish sites, neither ~~the~~ EDGAR v6 nor ~~the~~ CAMS-REG seemed to be better than the other ~~Since~~ (Figure 11). ~~As~~ the comparison with the atmospheric mole fraction does not ~~tell directly directly provide information on~~ whether the split between anthropogenic and natural emissions is correct, this ~~could may~~ indicate that the ~~inversion-inverse~~ model had difficulties in separating anthropogenic and natural emissions. This is particularly likely in southern Finland, ~~which has where there are~~ both anthropogenic and natural CH<sub>4</sub> sources. ~~In addition, it can~~ ~~It may~~ also reflect the complexity of modelling urban fluxes. To improve ~~the~~ estimates of anthropogenic emissions, it would be interesting to combine city-scale estimates with our larger-scale inversions.

The three natural CH<sub>4</sub> priors used in this study differed in absolute magnitude and in spatial and temporal distribution. The comparison with the atmospheric observations from northern Finland gave a clear ranking of the three priors: INV<sub>JSBACH\_CAMSREG</sub>, with the highest emission estimates, seemed to have the most accurate natural estimates in Finland, inversion runs with LPX-Bern DYPTOP had the second best ~~rankings-ranking~~ and INV<sub>GCP\_EDGAR</sub>, with the lowest emission estimates, had the worst ~~rankings-ranking~~ (Figure 11). The natural posterior emissions were always ~~larger-higher~~ than their priors, even from

JSBACH-HIMMELI, and the largest increases were in 2016 ~~;~~ when the summer was warm and rainy (Finnish Meteorological Institute, 2016), and in 2019–2020 (Figure 5). Our results indicate that Finnish natural CH<sub>4</sub> emissions from peatlands ~~are~~ might be underestimated by the process-based models, although the ~~large~~ high natural posterior emissions could also be due to sources other than peatlands. In particular, ~~freshwater emissions~~ emissions from freshwater sources are relevant. We compared the freshwater emission estimates from Stavert et al. (2022) with the natural CH<sub>4</sub> prior and posterior emissions in Fig. 7 and showed that in southern Finland these estimates were ~~larger~~ higher than even the highest optimised natural emissions (Inv<sub>JSBACH\_CAMSREG</sub>). The spatial distribution of the freshwater emission estimates ~~and~~ coincided with the JSBACH-HIMMELI estimates ~~coincided~~ (Supplementary Figure S12), so the inversion would most likely have included freshwater emissions in the posterior natural emission estimates. However, as there were still methane-emitting peatlands in southern Finland, it is not expected that optimised Inv<sub>JSBACH\_CAMSREG</sub> emission estimates would have included only freshwater emissions. Therefore, the freshwater emission estimates in Finland seemed to ~~have been~~ be too high.

### 4.3 Years 2020 and 2021

The reasons for the high atmospheric CH<sub>4</sub> growth rates in recent years, especially in 2020–2021, have been ~~discussed~~ In 2020, ~~part~~ under discussion. Part of the high growth rate in 2020 has been attributed to a weaker OH sink caused by a decrease in NO<sub>x</sub> emissions due to Covid-19 lockdowns (Stevenson et al., 2022; Qu et al., 2022; Peng et al., 2022; Feng et al., 2023). ~~The~~ However, the weaker OH sink could not explain all of the increase in atmospheric CH<sub>4</sub>, and wetlands, especially at high latitudes and in the tropics, were also suggested to be responsible for the increase (~~Qu et al., 2022; Peng et al., 2022; Zhang et al., 2023; Feng et al., 2023~~) (Qu et al., 2022; Peng et al., 2022; Zhang et al., 2023; Feng et al., 2023; Qu et al., 2024). In Finland, total CH<sub>4</sub> emissions were higher in 2020 than in 2019 in all CTE-CH<sub>4</sub> inversions, and the increase was attributed to both anthropogenic and natural emissions, but the posterior natural emissions in Inv<sub>JSBACH\_CAMSREG</sub>, which probably gave the most realistic ~~natural emission estimates~~ estimates of natural emissions, were highest in 2020.

The increase in the CH<sub>4</sub> growth rate in 2021 has also been attributed to wetlands (~~Feng et al., 2023; Zhang et al., 2023~~) (Feng et al., 2023; Zhang et al., 2023; Qu et al., 2024). The natural CH<sub>4</sub> emission estimates ~~from the of~~ CTE-CH<sub>4</sub> in Finland ~~were higher~~ in 2021 ~~were higher~~ 2021 than in 2019, but at the same level or lower than in 2020 ~~;~~ but to understand the model results (Figure 5). To understand the Finnish emissions estimates in 2021, it is beneficial to study the emissions in the whole northern high latitudes. The biomass-burning product used in the CTE-CH<sub>4</sub> runs, GFEDv4.1s, estimated the CH<sub>4</sub> emissions in the boreal forests (north of 50° N) to be 8.6 Tg while they were 4.2 Tg in ~~2019–2019~~ (Supplementary Fig. S9). According to Feng et al. (2023), the global CH<sub>4</sub> emissions should have been 20.8 Tg higher in 2021 than in 2019 to reproduce the observed atmospheric methane, meaning ~~the that~~ emissions from biomass-burning in boreal forests would account for 21 % of the increase in global CH<sub>4</sub> emissions. Zheng et al. (2023) showed that CO<sub>2</sub> emissions from boreal forests have been increasing in recent decades ~~;~~ and that and that CO<sub>2</sub> emissions were at a record high in 2021. They also used GFEDv4.1s in their analysis, but unlike our inversions, they specifically optimised ~~emissions from~~ biomass-burning emissions. The record-high biomass-burning CH<sub>4</sub> emissions in the boreal forests were probably the cause ~~for of~~ the large decrease in the optimised wetland emissions of Inv<sub>LPX\_EDGAR\_UNC</sub> in the high northern latitudes from 2020 to ~~2021–2021~~ (Supplementary Fig. S8). They most likely also

constrained the optimisation in Finland, keeping the posterior emissions close to the prior emissions. However, ~~this to verify this, it~~ would require further investigation and, for example, an ~~inversion-inverse~~ model setup where ~~the~~ biomass-burning CH<sub>4</sub> emissions are also optimised. There are also uncertainties in the biomass-burning emission estimates ~~, for example and as discussed,~~ GFAS had much lower emission estimates ~~in boreal forests north of 50° N~~ in 2021 (4.9 Tg, Supplementary Fig. S8S9). The increase in 2021 compared to 2019 was still relatively ~~large-high~~ in GFAS (2.2 Tg), i.e. it would have explained 11 % of the global increase in CH<sub>4</sub> emissions in ~~2021, 2021 estimated by Feng et al. (2023).~~

#### 4.4 Uncertainty estimations

In addition to different prior emissions, we also investigated how different prior uncertainties affected the emission estimates in Finland. Inv<sub>LPX\_EDGAR\_UNC</sub>, where the natural prior uncertainty was defined based on a ~~process-process-based~~ model ensemble, showed better agreement with ~~the~~ observations at the southern sites than Inv<sub>LPX\_EDGAR</sub>, which ~~had-used~~ the default prior uncertainties but ~~had-otherwise-otherwise had~~ the same setup (Figure 11). In northern Finland, Inv<sub>LPX\_EDGAR</sub> had larger uncertainties and performed better than Inv<sub>LPX\_EDGAR\_UNC</sub>. ~~Additionally~~In addition, Inv<sub>JSBACH\_CAMSREG</sub>, which had the ~~largest-highest~~ natural prior emissions and thus the largest uncertainties in the north, performed ~~the-best-in-best at~~ the northern sites. One might therefore ~~assume-expect~~ that large uncertainties would give the optimisation the freedom to fit the posterior emissions to the observations and, with enough observations ~~would,~~ lead to better emission estimates. However, this only seemed to be the case for natural emissions. The anthropogenic prior, EDGAR v6, had ~~large-high~~ emissions and therefore large uncertainties, so in theory, the optimisation could have reduced ~~the~~ anthropogenic emissions more than it did. The largest reduction from EDGAR v6 was seen with Inv<sub>LPX\_EDGAR\_UNC</sub> (Figure 4), even though its anthropogenic prior uncertainty was the same as in the other runs. Thus, simply having large uncertainties and a relatively large number of sites does not guarantee a better estimate, but it is important to know where the uncertainties lie. It can be complicated to determine realistic uncertainty ranges, and even using an ensemble of ~~multiple individual estimates might~~ ~~several individual estimates may~~ not capture the true magnitude of CH<sub>4</sub> emissions. ~~It is good to keep in mind, though, that we used-~~

~~Our uncertainty estimates were based on~~ the process-based models from the ~~latest-latest~~ published GCP-CH<sub>4</sub> (Saunois et al., 2020). The ongoing effort to update the global CH<sub>4</sub> budget (Saunois et al., 2024) used ~~an~~ updated wetland extent product (WAD2M v2.0(Zhang et al., 2021); Zhang et al., 2021), and also 12 models had prognostic versions, almost doubling the number of model estimates ~~from Saunois et al. (2020)~~. It would be interesting to see how the uncertainty estimates would change if the process-based models from Saunois et al. (2024) were used.

The optimisation in our ~~inversion-inverse~~ model is based on the ~~Ensemble Kalman Filter~~ensemble Kalman filter, which creates an ensemble of 500 members based on the priors and their uncertainties. By default, this method gives us a range of estimates that represent the uncertainties in the emission estimates. With these uncertainties we can, for example, calculate the uncertainty reduction from prior to posterior, which indicates how well the optimisation is able to constrain emissions. Another fairly robust way ~~to estimate-of estimating~~ the uncertainties is to use different model ensembles and obtain a range of estimates (~~?Saunois et al., 2020; Stavert et al., 2022)~~. ~~As demonstrated~~(Petrescu et al., 2024; Saunois et al., 2020; Stavert et al., 2022). ~~As shown~~ here, using a single ~~inversion-inverse~~ model with different setups can constrain and produce a comparable range

680 of CH<sub>4</sub> emission estimates at the country-level as an ensemble ~~consisting of different inversion of different inverse~~ models. As it is easier for an individual researcher or a research group to maintain one ~~inversion-inverse~~ model at a time, it would be recommended to use different priors to produce more constrained and reliable CH<sub>4</sub> emission estimates.

## 5 Conclusions

~~This study investigated a wide range of~~ In this study, CH<sub>4</sub> ~~emission estimates in Finland using emissions in Finland were~~ investigated using a range of bottom-up (inventories and process-based models) and top-down (~~inversion models~~) approaches ~~inverse models~~ estimates. We studied how ~~the optimised emissions were affected by the setup of the inversion model runs and showed that the choice of priors strongly influences~~ different bottom-up emission estimates used as priors in the inverse model affected the posterior CH<sub>4</sub> ~~emission estimates. Furthermore, our results indicate that the choice of priors affected the emission estimates as much~~ emissions. This choice not only strongly influenced the posterior emissions, but was as important as the choice of ~~inversion model, and that the ensembles~~ inverse model: the ensemble of inversion runs using the same ~~inversion-inverse~~ model but different priors resulted in similar average total posterior emissions as when using different ~~inversion-inverse~~ models but similar priors.

Even though the CH<sub>4</sub> emission estimates in Finland had a large range, the range of the total posterior emissions was smaller than the range of the prior emissions. The optimisation was also able to ~~align-reconcile~~ the trends of ~~the~~ anthropogenic and ~~the~~ natural CH<sub>4</sub> emissions, and the seasonal cycles of ~~the~~ natural CH<sub>4</sub> emissions were altered by the optimisation ~~so that they~~ matched better with the to better match flux measurements from peatland sites. However, the spatial distributions were not radically changed from the prior emissions.

The comparison of atmospheric ~~CH<sub>4</sub>-CH<sub>4</sub>~~ observations with model results showed no clear preference between the anthropogenic inventories (EDGAR v6 and CAMS-REG), but the comparison seemed to favour the ~~largest-highest~~ natural prior (JSBACH-HIMMELI). The optimised natural emissions were ~~larger-higher~~ than their prior emissions, which could be due to missing emissions in the prior estimates, such as freshwater. Estimates of freshwater emissions are still highly uncertain, and the estimates examined in this study (Stavert et al., 2022) ~~seemed-appeared~~ to be too ~~large-high~~ for Finland. We also found evidence that CH<sub>4</sub> emissions from biomass-burning ~~CH<sub>4</sub>~~, which were not optimised in CTE-CH<sub>4</sub>, were likely to have a large impact on the optimised anthropogenic and natural emissions in Finland and ~~the~~ high northern latitudes, especially in 2021. ~~The biomass-burning~~ estimates-emissions used in our inversions (GFEDv4.1s) had much higher emissions than ~~estimates from~~ GFAS, which highlights those in GFAS, highlighting the uncertainties in the biomass-burning CH<sub>4</sub> emission estimates.

The optimised CH<sub>4</sub> emissions were shown to depend on the choice of prior emissions. This choice was particularly important for the optimisation of the different emission components, ~~since-as~~ the optimisation of the different emission components was based on the spatial and temporal distribution of the priors. Currently, there are six stations in Finland where atmospheric ~~CH<sub>4</sub>~~ is measured. Adding more stations would most likely help to better constrain the different emission components. In addition to more stations, we also need more reliable prior estimates and realistic uncertainty estimates. In this study, we ~~tried-a~~ process-model spread-based uncertainty estimate ~~tested an uncertainty estimate based on a process-based model ensemble~~ for

natural CH<sub>4</sub> emissions, which appeared to be an advantageous method compared to the standard uncertainty estimates (80 % of the prior emissions). This type of uncertainty estimation could also be used for anthropogenic emissions, although many of the anthropogenic inventories use the same statistics and activity data. However, as shown here, the choices made in compiling the inventories affect the ~~emissions-estimated~~ estimated emissions, and the differences between them can help us to identify where the largest uncertainties lie.

The absolute magnitude of CH<sub>4</sub> emissions from Finland, ~~particularly~~ especially anthropogenic emissions, is relatively small compared to global totals. Consequently, these magnitudes are primarily relevant in the context of methodological comparisons or for verifying the NGHGI. The broader relevance of this study emerges from our assessment of the ability of a global model ~~'s-ability~~ to estimate CH<sub>4</sub> emissions within a single country. Such objectives are becoming increasingly relevant, as highlighted by initiatives such as the World Meteorological ~~Organization~~ Organisation's Global Greenhouse Gas Watch (G3W). This initiative aims to have global ~~inversion-inverse~~ models operationally running that could be used to assess country-specific GHG emissions. Under ~~the~~-G3W, ~~the inversion-inverse~~ model results will be available in common standard formats, making them more accessible and easier to use. This will likely also encourage their use in future studies ~~by individuals unfamiliar with inversion~~ also by those unfamiliar with inverse models. As discussed in this study, ~~interpreting inversion~~ the interpretation of inverse model results requires careful consideration of the model setup and, in particular, posterior estimates should be considered alongside prior emissions rather than as ~~standalone~~ stand-alone, definitive results. Ideally, those conducting the model runs would also provide uncertainty estimates (e.g. ~~spatial and temporal uncertainty reductions from prior to posterior~~, or an ensemble of inversions using different priors) and ~~offer~~ guidance on how to interpret the results and what factors to consider. Although ~~preparing~~ the preparation of a comprehensive interpretation guide is challenging due to the possible diverse applications of ~~the~~-model results, ~~establishing~~ the establishment of some common guidelines would be ~~advantageous~~ beneficial (Peters et al., 2023).

*Code and data availability.* All the model results, inputs and code will be provided on request from the corresponding author (Maria Tenkanen, maria.tenkanen@fmi.fi)

*Author contributions.* MKT, AT and TA participated in the study design. MKT performed the data processing, prepared and performed the model runs, and prepared the visualisations for the manuscript. AT helped to set up the CTE-CH<sub>4</sub> and TM5 model runs, and performed the INV<sub>GCP\_EDGAR</sub> model run. MKT was responsible for the analysis and preparation of the original manuscript together with TA, AT and AMRP. HDvdG provided the CMAS-REG-v5 estimates, LHI the GAINS estimates and AMRP the VERIFY inverse model estimates. HDvdG, LHI and AMRP assisted in the analysis of the results. AL, TM and MR performed the JSBACH-HIMMELI model runs and provided the CH<sub>4</sub> emission estimates used in the inversion. HA provided the atmospheric CH<sub>4</sub> measurements from KMP and SOD. All authors have read and approved the published version of the manuscript.



*Competing interests.* The authors declare no conflicts of interest. Also, the funders had no role in the design of the study; in the collection, analysis, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

745 *Acknowledgements.* We thank the [teams team](#) behind the LPX-Bern DYPTOP v1.4 [for providing the CH<sub>4</sub> emission estimates](#). We would also like to thank the people who worked on the VERIFY project and the modellers who performed the inversion runs and provided the model results used in this study. ~~We thank the European Union's project FPCUP Action 2019-2-49 "Developing supports for monitoring and reporting of GHG emissions and removals from land use, land use change and forestry" (219/SI2.818795/07 (CLIMA)) for their support~~ [The authors would like to thank the ICOS and ICOS-Finland PIs for providing the data on CH<sub>4</sub> mole fractions. We thank the Finnish Meteorological](#)  
750 [Institute \(PAL, UTO, KMP, SOD\), University of Eastern Finland \(PUI\) and University of Helsinki \(SMR\) for providing the methane data in Finland.](#) We are grateful for CSIRO Oceans and Atmosphere, Climate Science Centre (CSIRO), Environment and Climate Change Canada (~~EECCC~~), the Hungarian Meteorological Service (HMS), the Institute for Atmospheric Sciences and Climate (ISAC), the Institute on Atmospheric Pollution of the National Research Council (IIA), the Institute of Environmental Physics, University of Heidelberg (IUP), Laboratoire des Sciences du Climat et de l'Environnement (LSCE), Lawrence Berkeley National Laboratory (LBNL-ARM), the Environment  
755 Division Global Environment and Marine Department Japan Meteorological Agency (JMA), the Main Geophysical Observatory (MGO), the Max Planck Institute for Biogeochemistry (MPIBGC), National Institute for Environmental Studies (NIES), Norwegian Institute for Air Research (NILU), [National Oceanic and Atmospheric Administration Earth System Research Laboratories \(NOAA ESRL\)](#), the Pennsylvania State University (PSU), Swedish University of Agricultural Sciences (SLU) ~~Marklund et al. (2022)~~, the Swiss Federal Laboratories for Materials Science and Technology (EMPA), Umweltbundesamt Germany/Federal Environmental Agency (UBA), Umweltbundesamt Austria/Environment Agency Austria (EAA) as the data provider for Sonnblick, University of Bristol (UNIVBRIS), University of Exeter (~~Elena Kozlova~~ [Univ. Exeter](#)), and University of Urbino (UNIURB) for performing high-quality CH<sub>4</sub> measurements at global sites and making them available through the [Global Atmosphere Watch - World Data Centre for Greenhouse Gases \(GAW-WDCGG\)](#) and personal communications. We also thank ~~ICOS PIs for providing the data/facilities on the atmospheric CH<sub>4</sub> concentration products. In situ observations collected over the US Southern Great Plains were supported by the Office of Biological and Environmental Research of the US Department of Energy under contract no. DE-AC02-05CH11231 as part of the Atmospheric Radiation Measurement (ARM) Program, ARM Aerial Facility (AAF), and Terrestrial Ecosystem Science (TES) Program. Measurements at Jungfraujoch were supported by ICOS-Switzerland~~ [Juha Leskinen for preprocessing the VERIFY inversion dataset](#). The following AI tools were used to revise and proofread parts of the text: Grammarly (<https://grammarly.com/>), DeepL Write (<https://www.deepl.com/en/write>) and Microsoft's Copilot. Copilot was also used to assist in the plotting of the figures. Taylor's diagram in Fig. 10 were plotted using code provided by Rochford (2016).

770 *Financial support.* We thank the EU-H2020 VERIFY (776810), EU-Horizon EYE-CLIMA (101081395), Academy of Finland Center of Excellence (272041), FIRI - ICOS Finland (345531), and GHGSUPER (351311), Flagships ACCC and FAME (337552 and 359196), CSC-IT Center for Science (FICOCOSS) and JTF-VISIO for financial support.

## References

- UNFCCC Global Stocktake (<https://unfccc.int/topics/global-stocktake>, accessed on 7th May, 2024), <https://unfccc.int/topics/global-stocktake>.  
775
- Aurela, M., Lohila, A., Tuovinen, J.-P., and Hatakka, J.: Carbon dioxide and energy flux measurements in four northern-boreal ecosystems at Pallas, *Boreal environment research*, 20, 455–473, 2015.
- Basu, S., Lan, X., Dlugokencky, E., Michel, S., Schwietzke, S., Miller, J. B., Bruhwiler, L., Oh, Y., Tans, P. P., Apadula, F., Gatti, L. V., Jordan, A., Necki, J., Sasakawa, M., Morimoto, S., Di Iorio, T., Lee, H., Arduini, J., and Manca, G.: Estimating emissions of methane  
780 consistent with atmospheric measurements of methane and  $\delta^{13}\text{C}$  of methane, *Atmospheric Chemistry and Physics*, 22, 15 351–15 377, <https://doi.org/10.5194/acp-22-15351-2022>, 2022.
- Brühl, C. and Crutzen, P. J.: MPIC two-dimensional model, Tech. rep., NASA, Washington, DC, 1993.
- Bruhwiler, L., Dlugokencky, E., Masarie, K., Ishizawa, M., Andrews, A., Miller, J., Sweeney, C., Tans, P., and Worthy, D.: CarbonTracker-CH<sub>4</sub>: An assimilation system for estimating emissions of atmospheric methane, *Atmospheric Chemistry and Physics*, 14, 8269–8293,  
785 <https://doi.org/10.5194/acp-14-8269-2014>, 2014.
- Canadell, J., Monteiro, P., Costa, M., Cunha, L. C. d., Cox, P., Eliseev, A., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S., and K. Zickfeld: Global Carbon and Other Biogeochemical Cycles and Feedbacks, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N.,  
790 Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., pp. 673–816, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, ISBN 9781009157896, <https://doi.org/10.1017/9781009157896.007>, 2023.
- Chandra, N., Patra, P. K., Fujita, R., Höglund-Isaksson, L., Umezawa, T., Goto, D., Morimoto, S., Vaughn, B. H., and Röckmann, T.: Methane emissions decreased in fossil fuel exploitation and sustainably increased in microbial source sectors during 1990–2020, *Communications Earth & Environment*, 5, 147, <https://doi.org/10.1038/s43247-024-01286-x>, 2024.  
795
- Chang, K.-Y., Riley, W. J., Collier, N., McNicol, G., Fluet-Chouinard, E., Knox, S. H., Delwiche, K. B., Jackson, R. B., Poulter, B., Saunio, M., Chandra, N., Gedney, N., Ishizawa, M., Ito, A., Joos, F., Kleinen, T., Maggi, F., McNorton, J., Melton, J. R., Miller, P., Niwa, Y., Pasut, C., Patra, P. K., Peng, C., Peng, S., Segers, A., Tian, H., Tsuruta, A., Yao, Y., Yin, Y., Zhang, W., Zhang, Z., Zhu, Q., Zhu, Q., and Zhuang, Q.: Observational constraints reduce model spread but not uncertainty in global wetland methane emission estimates, *Global Change Biology*, pp. 1–15, <https://doi.org/10.1111/gcb.16755>, 2023.  
800
- Collins, W. J., Webber, C. P., Cox, P. M., Huntingford, C., Lowe, J., Sitch, S., Chadburn, S. E., Comyn-Platt, E., Harper, A. B., Hayman, G., and Powell, T.: Increased importance of methane reduction for a 1.5 degree target, *Environmental Research Letters*, 13, 054 003, <https://doi.org/10.1088/1748-9326/aab89c>, 2018.
- Cusworth, D. H., Bloom, A. A., Ma, S., Miller, C. E., Bowman, K., Yin, Y., Maasackers, J. D., Zhang, Y., Scarpelli, T. R., Qu, Z., Jacob, D. J., and Worden, J. R.: A Bayesian framework for deriving sector-based methane emissions from top-down fluxes, *Communications Earth & Environment*, 2, 242, <https://doi.org/10.1038/s43247-021-00312-6>, 2021.  
805
- Delwiche, K. B., Knox, S. H., Malhotra, A., Fluet-Chouinard, E., McNicol, G., Feron, S., Ouyang, Z., Papale, D., Trotta, C., Canfora, E., Cheah, Y.-W., Christianson, D., Alberto, M. C. R., Alekseychik, P., Aurela, M., Baldocchi, D., Bansal, S., Billesbach, D. P., Bohrer, G., Bracho, R., Buchmann, N., Campbell, D. I., Celis, G., Chen, J., Chen, W., Chu, H., Dalmagro, H. J., Dengel, S., Desai, A. R., Detto,

- 810 M., Dolman, H., Eichelmann, E., Euskirchen, E., Famulari, D., Fuchs, K., Goeckede, M., Gogo, S., Gondwe, M. J., Goodrich, J. P., Gottschalk, P., Graham, S. L., Heimann, M., Helbig, M., Helfter, C., Hemes, K. S., Hirano, T., Hollinger, D., Hörtnagl, L., Iwata, H., Jacotot, A., Jurasinski, G., Kang, M., Kasak, K., King, J., Klatt, J., Koebisch, F., Krauss, K. W., Lai, D. Y. F., Lohila, A., Mammarella, I., Belelli Marchesini, L., Manca, G., Matthes, J. H., Maximov, T., Merbold, L., Mitra, B., Morin, T. H., Nemitz, E., Nilsson, M. B., Niu, S., Oechel, W. C., Oikawa, P. Y., Ono, K., Peichl, M., Peltola, O., Reba, M. L., Richardson, A. D., Riley, W., Runkle, B. R. K., Ryu, Y., Sachs, T., Sakabe, A., Sanchez, C. R., Schuur, E. A., Schäfer, K. V. R., Sonntag, O., Sparks, J. P., Stuart-Haëntjens, E., Sturtevant, C., Sullivan, R. C., Szutu, D. J., Thom, J. E., Torn, M. S., Tuittila, E.-S., Turner, J., Ueyama, M., Valach, A. C., Vargas, R., Varlagin, A., Vazquez-Lule, A., Verfaillie, J. G., Vesala, T., Vourlitis, G. L., Ward, E. J., Wille, C., Wohlfahrt, G., Wong, G. X., Zhang, Z., Zona, D., Windham-Myers, L., Poulter, B., and Jackson, R. B.: FLUXNET-CH4: a global, multi-ecosystem dataset and analysis of methane seasonality from freshwater wetlands, *Earth System Science Data*, 13, 3607–3689, <https://doi.org/10.5194/essd-13-3607-2021>, 2021.
- 815
- 820 Deng, Z., Ciais, P., Tzompa-Sosa, Z. A., Saunio, M., Qiu, C., Tan, C., Sun, T., Ke, P., Cui, Y., Tanaka, K., Lin, X., Thompson, R. L., Tian, H., Yao, Y., Huang, Y., Lauerwald, R., Jain, A. K., Xu, X., Bastos, A., Sitch, S., Palmer, P. I., Lauvaux, T., D’Aspremont, A., Giron, C., Benoit, A., Poulter, B., Chang, J., Petrescu, A. M. R., Davis, S. J., Liu, Z., Grassi, G., Albergel, C., Tubiello, F. N., Perugini, L., Peters, W., and Chevallier, F.: Comparing national greenhouse gas budgets reported in UNFCCC inventories against atmospheric inversions, *Earth System Science Data*, 14, 1639–1675, <https://doi.org/10.5194/essd-14-1639-2022>, 2022.
- 825 Eggleston, S., Buendia, L., Miwa, K., Ngara, T., and Tanabe, K.: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, <https://www.ipcc-nggip.iges.or.jp/public/2006gl/>, 2006.
- Etiopie, G., Ciotoli, G., Schwietzke, S., and Schoell, M.: Gridded maps of geological methane emissions and their isotopic signature, *Earth System Science Data*, 11, 1–22, <https://doi.org/10.5194/essd-11-1-2019>, 2019.
- European Commission: DIRECTIVE (EU) 2016/2284 OF THE EUROPEAN PARLIAMENT AND OF THE COUNCIL of 14 December
- 830 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC, *Official Journal of the European Union*, L 344, 1–31, [https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=uriserv:OJ.L\\_.2016.344.01.0001.01.ENG&toc=OJ:L:2016:344:TOC](https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=uriserv:OJ.L_.2016.344.01.0001.01.ENG&toc=OJ:L:2016:344:TOC), 2016.
- European Commission and Joint Research Centre, Olivier, J., Guizzardi, D., Schaaf, E., Solazzo, E., Crippa, M., Vignati, E., Banja, M., Muntean, M., Grassi, G., Monforti-Ferrario, F., and Rossi, S.: GHG emissions of all world – 2021 report, Publications Office of the
- 835 European Union, <https://doi.org/10.2760/173513>, 2021.
- European Commission and Joint Research Centre, Crippa, M., Guizzardi, D., Schaaf, E., Monforti-Ferrario, F., Quadrelli, R., Riskez Martin, A., Rossi, S., Vignati, E., Muntean, M., Brandao De Melo, J., Oom, D., Pagani, F., Banja, M., Taghavi-Moharamli, P., Köykkä, J., Grassi, G., Branco, A., and San-Miguel, J.: GHG emissions of all world countries – 2023, Publications Office of the European Union, <https://doi.org/doi/10.2760/953322>, 2023.
- 840 Evensen, G.: The Ensemble Kalman Filter: Theoretical formulation and practical implementation, *Ocean Dynamics*, 53, 343–367, <https://doi.org/10.1007/s10236-003-0036-9>, 2003.
- Feng, L., Palmer, P. I., Parker, R. J., Lunt, M. F., and Bösch, H.: Methane emissions are predominantly responsible for record-breaking atmospheric methane growth rates in 2020 and 2021, *Atmospheric Chemistry and Physics*, 23, 4863–4880, <https://doi.org/10.5194/acp-23-4863-2023>, 2023.
- 845 Finnish Meteorological Institute: Vuoden 2016 sää, <https://www.ilmatieteenlaitos.fi/vuosi-2016>, Accessed 21 May 2024, <https://www.ilmatieteenlaitos.fi/vuosi-2016>, 2016.

- Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M., and Zhang, H.: The Earth's Energy Budget, Climate Feedbacks and Climate Sensitivity, in: *Climate Change 2021 – The Physical Science Basis*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., pp. 923–1054, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, ISBN 9781009157896, <https://doi.org/10.1017/9781009157896.009>, 2023.
- 850 Geddes, A., Mikaloff-Fletcher, S., Schaefer, H., Smale, D., Brailsford, G., Law, R., and Ausseil, A.-G.: Atmospheric Methane Modelling: Evaluating Regional Methane Emission Using Inverse Modelling, MPI Technical Paper No: 2021/02, vol. 1, Ministry for Primary Industries, ISBN 978-1-99-100333-1, <https://www.mpi.govt.nz/dmsdocument/46222/direct>, 2021.
- 855 Gómez-Sanabria, A., Höglund-Isaksson, L., Rafaj, P., and Schöpp, W.: Carbon in global waste and wastewater flows-its potential as energy source under alternative future waste management regimes, *Advances in Geosciences*, 45, 105–113, <https://doi.org/10.5194/adgeo-45-105-2018>, 2018.
- Haghnegahdar, M. A., Sun, J., Hultquist, N., Hamovit, N. D., Kitchen, N., Eiler, J., Ono, S., Yarwood, S. A., Kaufman, A. J., Dickerson, R. R., Bouyon, A., Magen, C., and Farquhar, J.: Tracing sources of atmospheric methane using clumped isotopes, *Proceedings of the National Academy of Sciences*, 120, 2017, <https://doi.org/10.1073/pnas.2305574120>, 2023.
- 860 Hatakka, J.: ICOS ATC CH4 Release, Pallas (12.0 m), 2017-09-16–2024-03-31, <https://hdl.handle.net/11676/IMY9pSZLevM3UXmNVKiKi4WH>, 2024.
- Hatakka, J. and Laurila, T.: ICOS ATC CH4 Release, Utö - Baltic sea (57.0 m), 2018-03-09–2024-03-31, <https://hdl.handle.net/11676/ExAsED0VdOT1oAM4ZG3mNXT7>, 2024.
- 865 Henne, S., Brunner, D., Oney, B., Leuenberger, M., Eugster, W., Bamberger, I., Meinhardt, F., Steinbacher, M., and Emmenegger, L.: Validation of the Swiss methane emission inventory by atmospheric observations and inverse modelling, *Atmospheric Chemistry and Physics*, 16, 3683–3710, <https://doi.org/10.5194/acp-16-3683-2016>, 2016.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 870 Höglund-Isaksson, L.: Bottom-up simulations of methane and ethane emissions from global oil and gas systems 1980 to 2012, *Environmental Research Letters*, 12, <https://doi.org/10.1088/1748-9326/aa583e>, 2017.
- Höglund-Isaksson, L., Gómez-Sanabria, A., Klimont, Z., Rafaj, P., and Schöpp, W.: Technical potentials and costs for reducing global anthropogenic methane emissions in the 2050 timeframe –results from the GAINS model, *Environmental Research Communications*, 2, 025 004, <https://doi.org/10.1088/2515-7620/ab7457>, 2020.
- 880 Houweling, S., Krol, M., Bergamaschi, P., Frankenberg, C., Dlugokencky, E. J., Morino, I., Notholt, J., Sherlock, V., Wunch, D., Beck, V., Gerbig, C., Chen, H., Kort, E. A., Röckmann, T., and Aben, I.: A multi-year methane inversion using SCIAMACHY, accounting for systematic errors using TCCON measurements, *Atmospheric Chemistry and Physics*, 14, 3991–4012, <https://doi.org/10.5194/acp-14-3991-2014>, 2014.

- Integrated Greenhouse Gas Monitoring System for Germany ITMS: Available online (accessed on 18 April 2024), <https://www.itms-germany.de/en>, 2024.
- 885 IPCC: Wetlands, 2019.
- Ito, A. and Inatomi, M.: Use of a process-based model for assessing the methane budgets of global terrestrial ecosystems and evaluation of uncertainty, *Biogeosciences*, 9, 759–773, <https://doi.org/10.5194/bg-9-759-2012>, 2012.
- Ito, A., Li, T., Qin, Z., Melton, J. R., Tian, H., Kleinen, T., Zhang, W., Zhang, Z., Joos, F., Ciais, P., Hopcroft, P. O., Beerling, D. J., Liu, X., Zhuang, Q., Zhu, Q., Peng, C., Chang, K. Y., Fluet-Chouinard, E., McNicol, G., Patra, P., Poulter, B., Sitch, S., Riley, W., and Zhu, Q.: Cold-Season Methane Fluxes Simulated by GCP-CH<sub>4</sub> Models, *Geophysical Research Letters*, 50, 1–12, <https://doi.org/10.1029/2023GL103037>, 2023.
- 890 Janardanan, R., Maksyutov, S., Wang, F., Nayagam, L., Sahu, S. K., Mangaraj, P., Saunio, M., Lan, X., and Matsunaga, T.: Country-level methane emissions and their sectoral trends during 2009–2020 estimated by high-resolution inversion of GOSAT and surface observations, *Environmental Research Letters*, 19, <https://doi.org/10.1088/1748-9326/ad2436>, 2024.
- 895 Janssens-Maenhout, G., Crippa, M., Guizzardi, D., Muntean, M., Schaaf, E., Dentener, F., Bergamaschi, P., Pagliari, V., Olivier, J. G. J., Peters, J. A. H. W., van Aardenne, J. A., Monni, S., Doering, U., Petrescu, A. M. R., Solazzo, E., and Oreggioni, G. D.: EDGAR v4.3.2 Global Atlas of the three major greenhouse gas emissions for the period 1970–2012, *Earth System Science Data*, 11, 959–1002, <https://doi.org/10.5194/essd-11-959-2019>, 2019.
- 900 Jöckei, P., Tost, H., Pozzer, A., Brühl, C., Buchholz, J., Ganzeveld, L., Hoor, P., Kerkweg, A., Lawrence, M. G., Sander, R., Steil, B., Stiller, G., Tanarhte, M., Taraborrelli, D., Van Aardenne, J., and Lelieveld, J.: The atmospheric chemistry general circulation model ECHAM5/MESSy1: Consistent simulation of ozone from the surface to the mesosphere, *Atmospheric Chemistry and Physics*, 6, 5067–5104, <https://doi.org/10.5194/acp-6-5067-2006>, 2006.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J. J., Razinger, M., Schultz, M. G., Suttie, M., and Van Der Werf, G. R.: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.
- 905 Kalman, R. E.: A New Approach to Linear Filtering and Prediction Problems, *Journal of Basic Engineering*, 82, 35–45, <https://doi.org/10.1115/1.3662552>, 1960.
- Kangasaho, V., Tsuruta, A., Backman, L., Mäkinen, P., Houweling, S., Segers, A., Krol, M., Dlugokencky, E. J., Michel, S., White, J. W. C., and Aalto, T.: The Role of Emission Sources and Atmospheric Sink in the Seasonal Cycle of CH<sub>4</sub> and  $\delta^{13}\text{C-CH}_4$ : Analysis Based on the Atmospheric Chemistry Transport Model TM5, *Atmosphere*, 13, 888, <https://doi.org/10.3390/atmos13060888>, 2022.
- 910 Kleinen, T., Mikolajewicz, U., and Brovkin, V.: Terrestrial methane emissions from the Last Glacial Maximum to the preindustrial period, *Climate of the Past*, 16, 575–595, <https://doi.org/10.5194/cp-16-575-2020>, 2020.
- Krol, M., Houweling, S., Bregman, B., van den Broek, M., Segers, A., van Velthoven, P., Peters, W., Dentener, F., Bergamaschi, P., Broek, M. V. D., Segers, A., Velthoven, P. V., Peters, W., Dentener, F., van den Broek, M., Segers, A., van Velthoven, P., Peters, W., Dentener, F., Bergamaschi, P., Broek, M. V. D., Segers, A., Velthoven, P. V., Peters, W., and Dentener, F.: The two-way nested global chemistry-transport zoom model TM5: algorithm and applications, *Atmospheric Chemistry and Physics*, 5, 417–432, <https://doi.org/10.5194/acp-5-417-2005>, 2005.
- 915 Kuenen, J., Dellaert, S., Visschedijk, A., Jalkanen, J.-P., Super, I., and Denier van der Gon, H.: CAMS-REG-v4: a state-of-the-art high-resolution European emission inventory for air quality modelling, *Earth System Science Data*, 14, 491–515, <https://doi.org/10.5194/essd-14-491-2022>, 2022.
- 920

- Lan, X., Thoning, K., and Dlugokencky, E.: Trends in globally-averaged CH<sub>4</sub>, N<sub>2</sub>O, and SF<sub>6</sub> determined from NOAA Global Monitoring Laboratory measurements. Version 2024-04, <https://doi.org/10.15138/P8XG-AA10>, 2024.
- Lehtinen, K. and Leskinen, A.: ICOS ATC CH<sub>4</sub> Release, Puijo (84.0 m), 2020-11-10–2024-03-31, <https://hdl.handle.net/11676/925> JsWZ0-rT9MBzDC-Euc-IIFZ7, 2024.
- Leip, A., Skiba, U., Vermeulen, A., and Thompson, R. L.: A complete rethink is needed on how greenhouse gas emissions are quantified for national reporting, *Atmospheric Environment*, 174, 237–240, <https://doi.org/10.1016/j.atmosenv.2017.12.006>, 2018.
- Levula, J. and Mammarella, I.: ICOS ATC CH<sub>4</sub> Release, Hyttiälä (125.0 m), 2016-12-13–2024-03-31, <https://hdl.handle.net/11676/A2dMBXD1QB1opynipabmJtWm>, 2024.
- 930 Lienert, S. and Joos, F.: A Bayesian ensemble data assimilation to constrain model parameters and land-use carbon emissions, *Biogeosciences*, 15, 2909–2930, <https://doi.org/10.5194/bg-15-2909-2018>, 2018.
- Luhar, A. K., Etheridge, D. M., Loh, Z. M., Noonan, J., Spencer, D., Smith, L., and Ong, C.: Quantifying methane emissions from Queensland’s coal seam gas producing Surat Basin using inventory data and a regional Bayesian inversion, *Atmospheric Chemistry and Physics*, 20, 15 487–15 511, <https://doi.org/10.5194/acp-20-15487-2020>, 2020.
- 935 Lunt, M. F., Manning, A. J., Allen, G., Arnold, T., Bauguitte, S. J., Boesch, H., Ganesan, A. L., Grant, A., Helfter, C., Nemitz, E., O’Doherty, S. J., Palmer, P. I., Pitt, J. R., Rennick, C., Say, D., Stanley, K. M., Stavert, A. R., Young, D., and Rigby, M.: Atmospheric observations consistent with reported decline in the UK’s methane emissions (2013–2020), *Atmospheric Chemistry and Physics*, 21, 16 257–16 276, <https://doi.org/10.5194/acp-21-16257-2021>, 2021.
- Maksyutov, S., Eggleston, S., Woo, J. H., Fang, S., Witi, J., Gillenwater, M., Goodwin, J., and Tubiello, F.: 2019 Refinement to the 2006 IPCC
- 940 Guidelines for National Greenhouse Gas Inventories, Volume 1, IPCC, Switzerland, <https://www.ipcc-nggip.iges.or.jp/public/2019rf/index.html>, 2019.
- Manning, A. J., Redington, A. L., Say, D., O’Doherty, S., Young, D., Simmonds, P. G., Vollmer, M. K., Mühle, J., Arduini, J., Spain, G., Wisher, A., Maione, M., Schuck, T. J., Stanley, K., Reimann, S., Engel, A., Krummel, P. B., Fraser, P. J., Harth, C. M., Salameh, P. K., Weiss, R. F., Gluckman, R., Brown, P. N., Watterson, J. D., and Arnold, T.: Evidence of a recent decline in UK emissions of
- 945 hydrofluorocarbons determined by the InTEM inverse model and atmospheric measurements, *Atmospheric Chemistry and Physics*, 21, 12 739–12 755, <https://doi.org/10.5194/acp-21-12739-2021>, 2021.
- Marklund, P., Ottosson-Löfvenius, M., and Smith, P.: ICOS ATC CH<sub>4</sub> Release, Svartberget (150.0 m), 2017-06-01–2022-02-28, <https://hdl.handle.net/11676/atargbJjvxhGDR2Yov09hD-E>, 2022.
- McGrath, M. J., Petrescu, A. M. R., Peylin, P., Andrew, R. M., Matthews, B., Dentener, F., Balkovič, J., Bastrikov, V., Becker, M., Broquet, G., Ciais, P., Fortems-Cheiney, A., Ganzenmüller, R., Grassi, G., Harris, I., Jones, M., Knauer, J., Kuhnert, M., Monteil, G., Munassar, S., Palmer, P. I., Peters, G. P., Qiu, C., Schelhaas, M.-J., Tarasova, O., Vizzarri, M., Winkler, K., Balsamo, G., Berchet, A., Briggs, P., Brockmann, P., Chevallier, F., Conchedda, G., Crippa, M., Dellaert, S. N. C., Denier van der Gon, H. A. C., Filipek, S., Friedlingstein, P., Fuchs, R., Gauss, M., Gerbig, C., Guizzardi, D., Günther, D., Houghton, R. A., Janssens-Maenhout, G., Lauerwald, R., Lerink, B., Luijkx, I. T., Moulas, G., Muntean, M., Nabuurs, G.-J., Paquirissamy, A., Perugini, L., Peters, W., Pilli, R., Pongratz, J., Regnier, P., Scholze, M.,
- 955 Serengil, Y., Smith, P., Solazzo, E., Thompson, R. L., Tubiello, F. N., Vesala, T., and Walther, S.: The consolidated European synthesis of CO<sub>2</sub> emissions and removals for the European Union and United Kingdom: 1990–2020, *Earth System Science Data*, 15, 4295–4370, <https://doi.org/10.5194/essd-15-4295-2023>, 2023.

- McNorton, J., Wilson, C., Gloor, M., Parker, R. J., Boesch, H., Feng, W., Hossaini, R., and Chipperfield, M. P.: Attribution of recent increases in atmospheric methane through 3-D inverse modelling, *Atmospheric Chemistry and Physics*, 18, 18 149–18 168, <https://doi.org/10.5194/acp-18-18149-2018>, 2018.
- 960 Melton, J. R., Wania, R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C. A., Beerling, D. J., Chen, G., Eliseev, A. V., Denisov, S. N., Hopcroft, P. O., Lettenmaier, D. P., Riley, W. J., Singarayer, J. S., Subin, Z. M., Tian, H., Zürcher, S., Brovkin, V., Van Bodegom, P. M., Kleinen, T., Yu, Z. C., and Kaplan, J. O.: Present state of global wetland extent and wetland methane modelling: Conclusions from a model inter-comparison project (WETCHIMP), *Biogeosciences*, 10, 753–788, [https://doi.org/10.5194/bg-10-753-](https://doi.org/10.5194/bg-10-753-2013)
- 965 2013, 2013.
- Mikaloff-Fletcher, S. and Schaefer, H.: Rising methane: A new climate challenge, *Science*, 364, 932–933, <https://doi.org/10.1126/science.aax1828>, 2019.
- Nisbet, E. G., Dlugokencky, E. J., and Bousquet, P.: Methane on the Rise—Again, *Science*, 343, 493–495, <https://doi.org/10.1126/science.1247828>, 2014.
- 970 Nisbet, E. G., Fisher, R. E., Lowry, D., France, J. L., Allen, G., Bakkaloglu, S., Broderick, T. J., Cain, M., Coleman, M., Fernandez, J., Forster, G., Griffiths, P. T., Iverach, C. P., Kelly, B. F., Manning, M. R., Nisbet-Jones, P. B., Pyle, J. A., Townsend-Small, A., al Shalaan, A., Warwick, N., and Zazzeri, G.: Methane Mitigation: Methods to Reduce Emissions, on the Path to the Paris Agreement, *Reviews of Geophysics*, 58, 1–51, <https://doi.org/10.1029/2019RG000675>, 2020.
- Nisbet, E. G., Manning, M. R., Dlugokencky, E. J., Michel, S. E., Lan, X., Röckmann, T., Denier van der Gon, H. A. C., Schmitt, J., Palmer, P. I., Dyonisius, M. N., Oh, Y., Fisher, R. E., Lowry, D., France, J. L., White, J. W. C., Brailsford, G., and Bromley, T.: Atmospheric
- 975 Methane: Comparison Between Methane’s Record in 2006–2022 and During Glacial Terminations, *Global Biogeochemical Cycles*, 37, <https://doi.org/10.1029/2023GB007875>, 2023.
- Olhoff, A., Rocha Romero, J., Hans, F., Kuramochi, T., Höhne, N., Peters, G. P., Andrew, R. M., Dafnomilis, I., den Elzen, M., Chen, H.-H., de Boer, H.-S., Daioglou, V., and Edelenbosch, O.: The impact of COVID-19 and recovery packages on emission pathways to 2030: Inputs to the UNEP Emissions Gap Report 2021 Final project report, TemaNord, Nordic Council of Ministers, ISBN 9789289373265, <https://doi.org/10.6027/temanord2022-530>, 2022.
- 980 Peng, S., Lin, X., Thompson, R. L., Xi, Y., Liu, G., Hauglustaine, D., Lan, X., Poulter, B., Ramonet, M., Saunois, M., Yin, Y., Zhang, Z., Zheng, B., and Ciais, P.: Wetland emission and atmospheric sink changes explain methane growth in 2020, *Nature*, 612, 477–482, <https://doi.org/10.1038/s41586-022-05447-w>, 2022.
- 985 Peters, G., Andrew, R., Børke, R., Lucia, P., Petrescu, A. M. R., and Engelen, R.: CoCO2: Prototype system for a Copernicus CO2 service. Decision Support Blueprint, <https://coco2-project.eu/node/408>, 2023.
- Peters, W., Miller, J. B., Whitaker, J., Denning, A. S., Hirsch, A., Krol, M. C., Zupanski, D., Bruhwiler, L., and Tans, P. P.: An ensemble data assimilation system to estimate CO2 surface fluxes from atmospheric trace gas observations, *Journal of Geophysical Research Atmospheres*, 110, 1–18, <https://doi.org/10.1029/2005JD006157>, 2005.
- 990 Petrescu, A. M. R., Qiu, C., McGrath, M. J., Peylin, P., Peters, G. P., Ciais, P., Thompson, R. L., Tsuruta, A., Brunner, D., Kuhnert, M., Matthews, B., Palmer, P. I., Tarasova, O., Regnier, P., Lauerwald, R., Bastviken, D., Höglund-Isaksson, L., Winiwarter, W., Etiope, G., Aalto, T., Balsamo, G., Bastrikov, V., Berchet, A., Brockmann, P., Ciotoli, G., Conchedda, G., Crippa, M., Dentener, F., Groot Zwaaftink, C. D., Guizzardi, D., Günther, D., Haussaire, J.-M., Houweling, S., Janssens-Maenhout, G., Kouyate, M., Leip, A., Leppänen, A., Lugato, E., Maisonnier, M., Manning, A. J., Markkanen, T., McNorton, J., Muntean, M., Oreggioni, G. D., Patra, P. K., Perugini, L., Pison, I.,
- 995 Raivonen, M. T., Saunois, M., Segers, A. J., Smith, P., Solazzo, E., Tian, H., Tubiello, F. N., Vesala, T., van der Werf, G. R., Wilson, C., and

- Zaehle, S.: The consolidated European synthesis of CH<sub>4</sub> and N<sub>2</sub>O emissions for the European Union and United Kingdom: 1990–2019, *Earth System Science Data*, 15, 1197–1268, <https://doi.org/10.5194/essd-15-1197-2023>, 2023.
- 1000 Petrescu, A. M. R., Peters, G. P., Engelen, R., Houweling, S., Brunner, D., Tsuruta, A., Matthews, B., Patra, P. K., Belikov, D., Thompson, R. L., Höglund-Isaksson, L., Zhang, W., Segers, A. J., Etiope, G., Ciotoli, G., Peylin, P., Chevallier, F., Aalto, T., Andrew, R. M., Bastviken, D., Berchet, A., Broquet, G., Conchedda, G., Dellaert, S. N. C., Denier van der Gon, H., Gütschow, J., Haussaire, J.-M., Lauerwald, R., Markkanen, T., van Peet, J. C. A., Pison, I., Regnier, P., Solum, E., Scholze, M., Tenkanen, M., Tubiello, F. N., van der Werf, G. R., and Worden, J. R.: Comparison of observation- and inventory-based methane emissions for eight large global emitters, *Earth System Science Data*, 16, 4325–4350, <https://doi.org/10.5194/essd-16-4325-2024>, 2024.
- 1005 Poulter, B., Bousquet, P., Canadell, J. G., Ciais, P., Peregón, A., Saunio, M., Arora, V. K., Beerling, D. J., Brovkin, V., Jones, C. D., Joos, F., Gedney, N., Ito, A., Kleinen, T., Koven, C. D., McDonald, K., Melton, J. R., Peng, C., Peng, S., Prigent, C., Schroeder, R., Riley, W. J., Saito, M., Spahni, R., Tian, H., Taylor, L., Viovy, N., Wilton, D., Wiltshire, A., Xu, X., Zhang, B., Zhang, Z., and Zhu, Q.: Global wetland contribution to 2000–2012 atmospheric methane growth rate dynamics, <https://doi.org/10.1088/1748-9326/aa8391>, 2017.
- 1010 Qu, Z., Jacob, D. J., Zhang, Y., Shen, L., Varon, D. J., Lu, X., Scarpelli, T., Bloom, A., Worden, J., and Parker, R. J.: Attribution of the 2020 surge in atmospheric methane by inverse analysis of GOSAT observations, *Environmental Research Letters*, 17, <https://doi.org/10.1088/1748-9326/ac8754>, 2022.
- Qu, Z., Jacob, D. J., Bloom, A. A., Worden, J. R., Parker, R. J., and Boesch, H.: Inverse modeling of 2010–2022 satellite observations shows that inundation of the wet tropics drove the 2020–2022 methane surge, *Proceedings of the National Academy of Sciences*, 121, 2017, <https://doi.org/10.1073/pnas.2402730121>, 2024.
- 1015 Raivonen, M., Smolander, S., Backman, L., Susiluoto, J., Aalto, T., Markkanen, T., Mäkelä, J., Rinne, J., Peltola, O., Aurela, M., Lohila, A., Tomasic, M., Li, X., Larmola, T., Juutinen, S., Tuittila, E.-S., Heimann, M., Sevanto, S., Kleinen, T., Brovkin, V., and Vesala, T.: HIMMELI v1.0: Helsinki Model of Methane build-up and emission for peatlands, *Geoscientific Model Development*, 10, 4665–4691, <https://doi.org/10.5194/gmd-10-4665-2017>, 2017.
- 1020 Ramsden, A. E., Ganesan, A. L., Western, L. M., Rigby, M., Manning, A. J., Foulds, A., France, J. L., Barker, P., Levy, P., Say, D., Wisher, A., Arnold, T., Rennick, C., Stanley, K. M., Young, D., and O’Doherty, S.: Quantifying fossil fuel methane emissions using observations of atmospheric ethane and an uncertain emission ratio, *Atmospheric Chemistry and Physics*, 22, 3911–3929, <https://doi.org/10.5194/acp-22-3911-2022>, 2022.
- Rice, A. L., Butenhoff, C. L., Teama, D. G., Röger, F. H., Khalil, M. A. K., and Rasmussen, R. A.: Atmospheric methane isotopic record favors fossil sources flat in 1980s and 1990s with recent increase, *Proceedings of the National Academy of Sciences*, 113, 10 791–10 796, <https://doi.org/10.1073/pnas.1522923113>, 2016.
- 1025 Rinne, J., Tuittila, E. S., Peltola, O., Li, X., Raivonen, M., Alekseychik, P., Haapanala, S., Pihlatie, M., Aurela, M., Mammarella, I., and Vesala, T.: Temporal Variation of Ecosystem Scale Methane Emission From a Boreal Fen in Relation to Temperature, Water Table Position, and Carbon Dioxide Fluxes, *Global Biogeochemical Cycles*, 32, 1087–1106, <https://doi.org/10.1029/2017GB005747>, 2018.
- Rochford, P. A.: SkillMetrics: A Python package for calculating the skill of model predictions against observations, <http://github.com/PeterRochford/SkillMetrics>, 2016.
- 1030 Saboya, E., Zazzeri, G., Graven, H., Manning, A. J., and Englund Michel, S.: Continuous CH<sub>4</sub> and  $\delta^{13}\text{C}_{\text{CH}_4}$  measurements in London demonstrate under-reported natural gas leakage, *Atmospheric Chemistry and Physics*, 22, 3595–3613, <https://doi.org/10.5194/acp-22-3595-2022>, 2022.



- Sasakawa, M., Shimoyama, K., Machida, T., Tsuda, N., Suto, H., Arshinov, M., Davydov, D., Fofonov, A., Krasnov, O., Saeki, T., Koyama, Y., and Maksyutov, S.: Continuous measurements of methane from a tower network over Siberia, *Tellus, Series B: Chemical and Physical Meteorology*, 62, 403–416, <https://doi.org/10.1111/j.1600-0889.2010.00494.x>, 2010.
- 1035 Saunois, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., Raymond, P. A., Dlugokencky, E. J., Houweling, S., Patra, P. K., Ciais, P., Arora, V. K., Bastviken, D., Bergamaschi, P., Blake, D. R., Brailsford, G., Bruhwiler, L., Carlson, K. M., Carrol, M., Castaldi, S., Chandra, N., Crevoisier, C., Crill, P. M., Covey, K., Curry, C. L., Etiope, G., Frankenberg, C., Gedney, N., Hegglin, M. I., Höglund-Isaksson, L., Hugelius, G., Ishizawa, M., Ito, A., Janssens-Maenhout, G., Jensen, K. M., Joos, F., Kleinen, T., Krummel, P. B., Langenfelds, R. L., Laruelle, G. G., Liu, L., Machida, T., Maksyutov, S., McDonald, K. C., McNorton, J., Miller, P. A., Melton, J. R., Morino, I., Müller, J., Murguia-Flores, F., Naik, V., Niwa, Y., Noce, S., O'Doherty, S., Parker, R. J., Peng, C., Peng, S., Peters, G. P., Prigent, C., Prinn, R., Ramonet, M., Regnier, P., Riley, W. J., Rosentreter, J. A., Segers, A., Simpson, I. J., Shi, H., Smith, S. J., Steele, L. P., Thornton, B. F., Tian, H., Tohjima, Y., Tubiello, F. N., Tsuruta, A., Viovy, N., Voulgarakis, A., Weber, T. S., van Weele, M., van der Werf, G. R., Weiss, R. F., Worthy, D., Wunch, D., Yin, Y., Yoshida, Y., Zhang, W., Zhang, Z., Zhao, Y., Zheng, B., Zhu, Q., Zhu, Q., and Zhuang, Q.: The Global Methane Budget 2000–2017, *Earth System Science Data*, 12, 1561–1623, <https://doi.org/10.5194/essd-12-1561-2020>, 2020.
- 1040 Saunois, M., Martinez, A., Poulter, B., Zhang, Z., Raymond, P. A., Pierre Regnier, Canadell, J. G., Jackson, R. B., Patra, P. K., Bousquet, P., Philippe Ciais, Dlugokencky, E. J., Lan, X., Allen, G. H., Bastviken, D., Beerling, J., D., Belikov, D. A., Blake, D. R., Castaldi, S., Crippa, M., Deemer, B. R., Dennison, F., Etiope, G., Gedney, N., Höglund-Isaksson, L., Holgerson, M. A., Hopcroft, P. O., Hugelius, G., Ito, A., Jain, A. K., Janardanan, R., Johnson, M. S., Kleinen, T., Krummel, P. B., Lauerwald, R., Li, T., Liu, X., McDonald, K. C., Melton, J. R., Mühle, J., Müller, J., Murguia-Flores, F., Niwa, Y., Noce, S., Pan, S., Parker, R. J., Peng, C., Ramonet, M., Riley, W. J., Rocher-Ros, G., Rosentreter, J. A., Sasakawa, M., Segers, A., Smith, S. J., Stanley, E. H., Thanwerdas, J., Tian, H., Tsuruta, A., Tubiello, F. N., Weber, T. S., Werf, G. R. v. d., Worthy, D. E. J., Xi, Y., Yoshida, Y., Zhang, W., Bo Zheng, Zhu, Q., Zhu, Q., and Zhuang, Q.: Global Methane Budget 2000-2020, *Earth System Science Data Discussions*, pp. 1–147, <https://doi.org/10.5194/essd-2024-115>, 2024.
- 1050 Schult, K. N., Aalto, T., Andrews, A., Aoki, S., Arduini, J., Baier, B., Bergamaschi, P., Biermann, T., Biraud, S. C., Boenisch, H., Brailsford, G., Chen, H., Colomb, A., Conil, S., Cristofanelli, P., Cuevas, E., Daube, B., Davis, K., Mazière, M. D., Delmotte, M., Desai, A., DiGangi, J. P., Dlugokencky, E., Elkins, J. W., Emmenegger, L., Fischer, M. L., Gatti, L. V., Gehrlein, T., Gerbig, C., Gloor, E., Goto, D., Haszpra, L., Hatakka, J., Heimann, M., Heliasz, M., Hermanssen, O., Hints, E., Holst, J., Ivakhov, V., Jaffe, D., Joubert, W., Kang, H.-Y., Karion, A., Kazan, V., Keronen, P., Ko, M.-Y., Kominkova, K., Kort, E., Kozlova, E., Krummel, P., Kubistin, D., Labuschagne, C., Langenfelds, R., Laurent, O., Laurila, T., Lauvaux, T., Lee, J., Lee, H., Lee, C.-H., Lehner, I., Leppert, R., Leuenberger, M., Lindauer, M., Loh, Z., Lopez, M., Machida, T., Mammarella, I., Manca, G., Marek, M. V., Martin, M. Y., Matsueda, H., McKain, K., Miles, N., Miller, C. E., Miller, J. B., Moore, F., Morimoto, S., Munro, D., Myhre, C. L., Mölder, M., Müller-Williams, J., Nichol, S., Niwa, Y., O'Doherty, S., Obersteiner, F., Piacentino, S., Pichon, J. M., Pittman, J., Plass-Duelmer, C., Ramonet, M., Richardson, S., Rivas, P. P., Saito, K., Santoni, G., Sasakawa, M., Scheeren, B., Schuck, T., Schumacher, M., Seifert, T., Sha, M. K., Shepson, P., Sloop, C. D., Smith, P., Steinbacher, M., Stephens, B., Sweeney, C., Timas, H., Torn, M., Trisolino, P., Turnbull, J., Tørseth, K., Viner, B., Vitkova, G., Watson, A., Wofsy, S., Worsley, J., Worthy, D., Zahn, A., and Sarra, A. G. d.: Multi-laboratory compilation of atmospheric carbon dioxide data for the period 1983-2020; *obspack\_ch4\_1\_GLOBALVIEWplus\_v4.0\_2021-10-14*, <https://doi.org/10.25925/20211001>, 2021.
- 1060 Segers, A. and Nanni, R.: CAMS2\_55\_CEA – Provision of global inversion- optimised greenhouse gas fluxes and concentrations, Contribution to documentation of products and services as provided within the scope of this contract – 2023 – Part CH4, <https://atmosphere.copernicus.eu/greenhouse-gases-supplementary-products>, 2023.
- 1070

- Spahni, R., Wania, R., Neef, L., Van Weele, M., Pison, I., Bousquet, P., Frankenberg, C., Foster, P. N., Joos, F., Prentice, I. C., and Van Velthoven, P.: Constraining global methane emissions and uptake by ecosystems, *Biogeosciences*, 8, 1643–1665, <https://doi.org/10.5194/bg-8-1643-2011>, 2011.
- Spahni, R., Joos, F., Stocker, B. D., Steinacher, M., and Yu, Z. C.: Transient simulations of the carbon and nitrogen dynamics in northern peatlands: From the Last Glacial Maximum to the 21st century, *Climate of the Past*, 9, 1287–1308, <https://doi.org/10.5194/cp-9-1287-2013>, 2013.
- Statistics Finland: Greenhouse Gas Emissions in Finland 1990 to 2021. National Inventory Report under the UNFCCC and the Kyoto Protocol., 2023.
- Stavert, A. R., Saunio, M., Canadell, J. G., Poulter, B., Jackson, R. B., Regnier, P., Lauerwald, R., Raymond, P. A., Allen, G. H., Patra, P. K., Bergamaschi, P., Bousquet, P., Chandra, N., Ciais, P., Gustafson, A., Ishizawa, M., Ito, A., Kleinen, T., Maksyutov, S., McNorton, J., Melton, J. R., Müller, J., Niwa, Y., Peng, S., Riley, W. J., Segers, A., Tian, H., Tsuruta, A., Yin, Y., Zhang, Z., Zheng, B., and Zhuang, Q.: Regional trends and drivers of the global methane budget, *Global Change Biology*, 28, 182–200, <https://doi.org/10.1111/gcb.15901>, 2022.
- Stevenson, D. S., Derwent, R. G., Wild, O., and Collins, W. J.: COVID-19 lockdown emission reductions have the potential to explain over half of the coincident increase in global atmospheric methane, *Atmospheric Chemistry and Physics*, 22, 14 243–14 252, <https://doi.org/10.5194/acp-22-14243-2022>, 2022.
- Stocker, B. D., Spahni, R., and Joos, F.: DYPTOP: a cost-efficient TOPMODEL implementation to simulate sub-grid spatio-temporal dynamics of global wetlands and peatlands, *Geoscientific Model Development*, 7, 3089–3110, <https://doi.org/10.5194/gmd-7-3089-2014>, 2014.
- Taylor, K. E.: Summarizing multiple aspects of model performance in a single diagram, *Journal of Geophysical Research: Atmospheres*, 106, 7183–7192, <https://doi.org/10.1029/2000JD900719>, 2001.
- Tenkanen, M. K., Tsuruta, A., Tyystjärvi, V., Törmä, M., Autio, I., Haakana, M., Tuomainen, T., Leppänen, A., Markkanen, T., Raivonen, M., Niinistö, S., Arslan, A. N., and Aalto, T.: Using Atmospheric Inverse Modelling of Methane Budgets with Copernicus Land Water and Wetness Data to Detect Land Use-Related Emissions, *Remote Sensing*, 16, 124, <https://doi.org/10.3390/rs16010124>, 2024.
- Thanwerdas, J., Saunio, M., Berchet, A., Pison, I., and Bousquet, P.: Investigation of the renewed methane growth post-2007 with high-resolution 3-D variational inverse modeling and isotopic constraints, *Atmospheric Chemistry and Physics*, 24, 2129–2167, <https://doi.org/10.5194/acp-24-2129-2024>, 2024.
- Thompson, R. L., Nisbet, E. G., Pisso, I., Stohl, A., Blake, D., Dlugokencky, E. J., Helmig, D., and White, J. W. C.: Variability in Atmospheric Methane From Fossil Fuel and Microbial Sources Over the Last Three Decades, *Geophysical Research Letters*, 45, 499–11, <https://doi.org/10.1029/2018GL078127>, 2018.
- Thoning, K. W. and Tans, P. P.: Atmospheric carbon dioxide at Mauna Loa Observatory. 2. Analysis of the NOAA GMCC data, 1974–1985, *Journal of Geophysical Research*, 94, 8549–8565, <https://doi.org/10.1029/JD094iD06p08549>, 1989.
- Tsuruta, A., Aalto, T., Backman, L., Hakkarainen, J., van der Laan-Luijkx, I. T., Krol, M. C., Spahni, R., Houweling, S., Laine, M., Dlugokencky, E., Gomez-Pelaez, A. J., van der Schoot, M., Langenfelds, R., Ellul, R., Arduini, J., Apadula, F., Gerbig, C., Feist, D. G., Kivi, R., Yoshida, Y., and Peters, W.: Global methane emission estimates for 2000–2012 from CarbonTracker Europe-CH<sub>4</sub> v1.0, *Geoscientific Model Development*, 10, 1261–1289, <https://doi.org/10.5194/gmd-10-1261-2017>, 2017.
- Tsuruta, A., Aalto, T., Backman, L., Krol, M. C., Peters, W., Lienert, S., Joos, F., Miller, P. A., Zhang, W., Laurila, T., Hatakka, J., Leskinen, A., Lehtinen, K. E. J., Peltola, O., Vesala, T., Levula, J., Dlugokencky, E., Heimann, M., Kozlova, E., Aurela, M., Lohila, A., Kauhaniemi,

- M., and Gomez-Pelaez, A. J.: Methane budget estimates in Finland from the CarbonTracker Europe-CH<sub>4</sub> data assimilation system, *Tellus B: Chemical and Physical Meteorology*, 71, 1–20, <https://doi.org/10.1080/16000889.2018.1565030>, 2019.
- 1110 UNECE: 1999 Protocol to Abate Acidification, Eutrophication and Ground-level ozone to the Convention on Long-range Transboundary Air Pollution, as amended on 4 May 2012, p. ECE/EB. AIR/114, [https://unece.org/fileadmin/DAM/env/documents/2013/air/eb/ECE.EB.AIR.114\\_ENG.pdf](https://unece.org/fileadmin/DAM/env/documents/2013/air/eb/ECE.EB.AIR.114_ENG.pdf), 2012.
- UNFCCC: The Paris Agreement, United Nations Treaty Collection, Chapter XXVII 7. d, <https://unfccc.int/process-and-meetings/the-paris-agreement/>, 2015.
- 1115 UNFCCC: Technical Dialogue of the First Global Stocktake: Synthesis Report by the Co-facilitators on the Technical Dialogue, <http://unfccc.int/resource/docs/2009/cop15/eng/11a01.pdf>, 2023.
- van der Laan-Luijkx, I. T., van der Velde, I. R., van der Veen, E., Tsuruta, A., Stanislawski, K., Babenhauerheide, A., Zhang, H. F., Liu, Y., He, W., Chen, H., Masarie, K. A., Krol, M. C., and Peters, W.: The CarbonTracker Data Assimilation Shell (CTDAS) v1.0: implementation and global carbon balance 2001–2015, *Geoscientific Model Development*, 10, 2785–2800, <https://doi.org/10.5194/gmd-10-2785-2017>, 2017.
- 1120 van der Werf, G. R., Randerson, J. T., Giglio, L., van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997–2016, *Earth System Science Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
- 1125 Weber, T., Wiseman, N. A., and Kock, A.: Global ocean methane emissions dominated by shallow coastal waters, *Nature Communications*, 10, 4584, <https://doi.org/10.1038/s41467-019-12541-7>, 2019.
- Worden, J. R., Cusworth, D. H., Qu, Z., Yin, Y., Zhang, Y., Bloom, A. A., Ma, S., Byrne, B. K., Scarpelli, T., Maasackers, J. D., Crisp, D., Duren, R., and Jacob, D. J.: The 2019 methane budget and uncertainties at 1° resolution and each country through Bayesian integration of GOSAT total column methane data and a priori inventory estimates, *Atmospheric Chemistry and Physics*, 22, 6811–6841, <https://doi.org/10.5194/acp-22-6811-2022>, 2022.
- 1130 Worden, J. R., Pandey, S., Zhang, Y., Cusworth, D. H., Qu, Z., Bloom, A. A., Ma, S., Maasackers, J. D., Byrne, B., Duren, R., Crisp, D., Gordon, D., and Jacob, D. J.: Verifying Methane Inventories and Trends With Atmospheric Methane Data, *AGU Advances*, 4, <https://doi.org/10.1029/2023AV000871>, 2023.
- Zhang, Z., Fluet-Chouinard, E., Jensen, K., McDonald, K., Hugelius, G., Gumbrecht, T., Carroll, M., Prigent, C., Bartsch, A., and Poulter, B.: Development of the global dataset of Wetland Area and Dynamics for Methane Modeling (WAD2M), *Earth System Science Data*, 13, 2001–2023, <https://doi.org/10.5194/essd-13-2001-2021>, 2021.
- 1135 Zhang, Z., Poulter, B., Feldman, A. F., Ying, Q., Ciais, P., Peng, S., and Li, X.: Recent intensification of wetland methane feedback, *Nature Climate Change*, 13, 430–433, <https://doi.org/10.1038/s41558-023-01629-0>, 2023.
- Zheng, B., Ciais, P., Chevallier, F., Yang, H., Canadell, J. G., Chen, Y., van der Velde, I. R., Aben, I., Chuvieco, E., Davis, S. J., Deeter, M., Hong, C., Kong, Y., Li, H., Li, H., Lin, X., He, K., and Zhang, Q.: Record-high CO<sub>2</sub> emissions from boreal fires in 2021, *Science*, 379, 912–917, <https://doi.org/10.1126/science.ade0805>, 2023.
- 1140