

Model analysis of biases in satellite diagnosed aerosol effect on cloud liquid water path

Harri Kokkola^{1,2}, Juha Tonttila³, Silvia Calderón¹, Sami Romakkaniemi¹, Antti Lipponen¹, Aapo Peräkorpä⁴, Tero Mielonen¹, Edward Gryspeerdt⁵, Timo Henrik Virtanen¹, Pekka Kolmonen¹, and Antti Arola¹

¹Atmospheric Research Centre of Eastern Finland, Finnish Meteorological Institute, Kuopio, Finland

²University of Eastern Finland, Kuopio, Finland

³CSC - IT Center for Science, Espoo, Finland

⁴Datrix S.p.A., Milan, Italy

⁵Grantham Institute - Climate Change and the Environment, Imperial College London, London, UK

Correspondence: Harri Kokkola (harri.kokkola@fmi.fi)

Abstract. The response in cloud water content to changes in cloud condensation nuclei remains one of the major uncertainties in determining how aerosols can perturb cloud properties. In this study, we used an ensemble of large eddy simulations of marine stratocumulus clouds to investigate the correlation between cloud liquid water path and the amount of cloud condensation nuclei. We compare this correlation directly from the model to the correlation derived using equations which are used to retrieve liquid water path from satellite observations. Our comparison shows that spatial variability in cloud properties and instrumental noise in satellite retrievals of cloud optical depth and cloud effective radii result in bias in satellite-derived liquid water path. In depth investigation of high-resolution model data shows that in large part of a cloud, the assumption of adiabaticity does not hold which results in a similar bias in LWP-CDNC relationship as seen in satellite data. In addition, our analysis shows a significant positive bias of between 18 % and 40 % in satellite-derived cloud droplet number concentration. However, for the individual ensemble members, the correlation between the cloud condensation nuclei and the mean of the liquid water path was very similar between the methods. This suggests that if cloud cases are carefully chosen for similar meteorological conditions and it is ensured that cloud condensation nuclei concentrations are well-defined, changes in liquid water can be confidently determined using satellite data.

1 Introduction

Clouds are in a crucial role in the Earth's climate affecting the radiative balance of the Earth as they cover majority of the Earth surface having high reflectivity on the incoming solar radiation and absorbing outgoing thermal radiation (Bellouin et al., 2020; Forster et al., 2021). As aerosol can perturb the cloud properties, accurate knowledge of how aerosol-cloud interactions affect clouds will allow for better estimation on how changes in anthropogenic emissions affect the Earth's radiative balance and thus the climate. Satellite based estimates of aerosol effects on clouds have proven to be challenging to interpret as they have not always supported the theoretical assumptions of decreasing cloud droplet sizes with increasing number of cloud droplets (Twomey, 1974; Jia et al., 2019) or the increase of cloud liquid water content with increasing number of cloud




Figure 1. Cloud properties of a stratocumulus cloud deck west of Peru and Chile over South Pacific on Aug 30th, 2003. The upper left panel shows the CDNC calculated according to Quaas et al. (2006) and lower left panel shows the retrieved LWP from Moderate Resolution Imaging Spectroradiometer (MODIS) Level-2 (L2) Collection 6.1 (Platnick et al., 2015). Right panels show a magnification of the structure of a cloud cell within the cloud field denoting the cloud effective radius and the cloud reflectance for the corresponding cloud cell. Small squares are approximately $3\text{ km} \times 3\text{ km}$ and large rectangles are approx. $500\text{ km} \times 6\text{ km}$

droplets (Albrecht, 1989; Gryspeerdt et al., 2019). These mixed results have been attributed to several counteracting physical processes, for example the effects of solar heating, cloud-top mixing, and variability in moisture on LWP (Feingold et al., 2022; Gryspeerdt et al., 2022; Glassmeier et al., 2021; Zhang et al., 2024), but also challenges in satellite retrievals (Feingold et al., 2022; Arola et al., 2022).

Arola et al. (2022) showed that variability in positively correlated cloud droplet number concentration (CDNC) and liquid water path (LWP) data will dilute the correlation and can even result in an apparent strongly negative correlation between CDNC and LWP. However, in that study, the causes of variability were not studied further. Such variability can come from 1) internal variability in clouds originating from the circulation within clouds, e.g., in updrafts at the center of the cloud cells and downdrafts at the edges of the cell, 2) mesoscale variability in meteorological conditions and phase of the cloud evolution, 3) instrumental noise in satellite retrievals.

Figure 1 shows the cloud properties in a stratocumulus deck west of Peru and Chile, South America. In the figure, the wide rectangles show regions in the cloud field which show mesoscale variability in CDNC and LWP while the small squares indicate internal variability in CDNC and LWP within a cloud cell. The smaller squares are magnified in the right-hand side

35 panels, showing both the effective radius and the reflectivity of the cloud at visible wavelengths. The wide rectangle in the figure shows a region with significant mesoscale variabilities in cloud properties.

As for the small squares in Figure 1, we can see that the retrieved cloud effective radius decreases towards the cloud cell edges. The decrease in retrieved cloud effective radius results from the entrainment mixing at the cloud top and downdrafts in the cloud cell boundaries which both reduce the liquid water path. At the cloud cell edges this is in conflict with the assumptions
40 made in the calculation of CDNC. Calculation of CDNC based on the effective radius, and assuming constant sub-adiabaticity, would lead to overestimation in retrieved CDNC values compared to the real CDNC values at the cell boundaries (see Equation (2) in Section 2). In addition to actual variability in physical properties of clouds, satellite retrievals include uncertainties and instrument noise causing another potential source of bias in the satellite-derived correlation between CDNC and LWP. All these different sources of variability are potential causes for biasing the estimate of aerosol effect on LWP as shown by Arola et al.
45 (2022)

In this study we use a cloud resolving large eddy simulations (LES) model to investigate the relative contribution of these sources of variability (cloud condensation nuclei, cloud structure, and noise in satellite retrievals) to the correlation between CDNC and LWP. We will analyse how the diagnosed response in liquid water path (LWP) to perturbed aerosol concentrations differ when calculated with equations used in the satellite retrievals of LWP, compared to LWP diagnosed directly from the
50 LES model.

2 METHODS

2.1 Model description

We simulated the effect of aerosol concentration on cloud properties, especially the liquid water content using UCLALES-SALSA large eddy simulations model (Stevens et al., 2005; Kokkola et al., 2008; Tonttila et al., 2017; Ahola et al., 2020).
55 In this model setup, UCLALES which simulates the dynamics of the boundary layer is coupled to the aerosol-cloud model SALSA which simulates aerosol and cloud droplet microphysics. SALSA has a sectional description for aerosol particles, cloud and precipitation droplets, and ice crystals.

The model is built around UCLALES, a platform for idealized cloud simulations. The model resolves the turbulent flow in a three-dimensional cartesian grid with cyclic boundary conditions. The main prognostic scalar variables include the liquid
60 water potential temperature and tracer variables describing water vapor and liquid water mixing ratios. When coupled with the sectional aerosol-cloud microphysics model SALSA, the set of prognostic scalars is vastly extended, now including the number and mass mixing ratios for each size section of four particle categories, comprising aerosol particles, cloud droplets, drizzle/precipitation and ice (the latter not used in this study). The size resolved framework is used to describe particle growth via condensation and coalescence processes in all categories. Aerosol cloud activation is determined directly from the resolved
65 particle growth whereas the transition between cloud droplets and drizzle is diagnosed from the resolved collision-coalescence process. UCLALES-SALSA has been validated against observations in liquid phase clouds and fogs and found to reproduce

the observed droplet distributions (e.g. Boutle et al., 2018; Calderón et al., 2022). More details on the model and simulations are given in Supplementary Information Sections S1 and S2.

2.1.1 Experiment setup

70 To better understand the features of the LWP response to changes in CDNC seen in the satellite data, UCLALES-SALSA was configured for a typical marine stratocumulus cloud setup (nocturnal drizzling stratocumulus cloud DYCOMS-II RF02 of the Second Dynamics and Chemistry of Marine Stratocumulus field campaign (Ackerman et al., 2009)), representing a very commonly occurring cloud type which allows for disentangling the potential underlying numerical biases related to satellite retrievals. The horizontal model domain size was 51×51 km with a resolution of 75 m with vertical domain extending up to
75 1.4 km with vertical resolution of 20 m. Vertical profiles of atmospheric variables used for model initialization are shown in Figure S1 in Supporting Information.

For aerosol, we used a bi-modal size distribution. The Aitken mode is centered at $0.022 \mu\text{m}$ with a standard deviation of 1.2 and total number concentration of 150 cm^{-3} . The accumulation mode is centered at $0.120 \mu\text{m}$ with standard deviation of 1.7. To determine the effect of aerosol on cloud properties, we ran a set of 3 simulations with cloud condensation nuclei
80 (CCN) equivalent to initial accumulation mode aerosol concentrations of 65, 150 and 300 cm^{-3} (corresponding to the total number concentrations of 78, 180 and 360 cm^{-3} for the whole size distribution). Size distributions are illustrated in Figure S2 in Supporting Information.

The simulations span over 14 h and the model output was sampled 2 h, 6 h, and 10 h from the start of the run. These time intervals allow us to draw samples from different cloud structures, as the microphysical properties and circulation structures
85 are allowed to evolve freely in the model. In particular, the simulations with the lowest initial aerosol concentrations exhibit a clear drizzle induced transition from closed stratocumulus cells to an open cell structure within 10 hours of model time.

It is well-known that satellite-based CDNC values are biased because radiances used in the CER retrievals correspond to an optically thicker region below cloud top. Platnick (2000) established that infrared radiance fluxes from liquid clouds include all reflected photons that penetrate to a maximum optical depth equivalent to up to 3.5 units below cloud top (Grosvenor et al.,
90 2018b) depending on the viewing geometry and cloud heterogeneity (Grosvenor et al., 2018a). With CER values that are smaller than those expected at cloud top, the cloud-top based pseudo-adiabatic model inevitably fails, producing satellite-retrievals of CDNC and LWP that are different from real ones (Grosvenor et al., 2018a).

In this study, we followed a sampling methodology that mimics this so-called penetration depth bias (Grosvenor et al., 2018a). We determined the CER and CDNC values for the top part of a cloud based on as many model layers as needed to
95 reduce the cloud optical thickness (COT) in the infrared region by three units. The infrared COT was calculated using the wavelength band of $2.38 \mu\text{m}$ - $4.00 \mu\text{m}$ to match the MODIS retrievals done at the $2.13 \mu\text{m}$ and $3.7 \mu\text{m}$ channels. Both CER and CDNC were calculated as extinction coefficient b_{ext} weighted average values that consider all model layers in the top part of the cloud (Equations (S.1) and (S.2)). For the sake of simplicity we refer to this region as the *extended cloud top*. COT was calculated for the visible wavelength band of $0.25 \mu\text{m}$ - $0.69 \mu\text{m}$ as a surrogate of MODIS retrievals done at $0.66 \mu\text{m}$. More
100 details on the sampling methodology can be found in section S2 of the supporting information.

Values for CDNC (N_d) were calculated with CER (r_e) and COT (τ_c) obtained for the extended cloud top in each cloud column from the following equation

$$N_d = \frac{\sqrt{5}}{2\pi k} \left(\frac{f_{ad} c_w \tau_c}{Q_{ext} \rho_w r_e^5} \right)^{\frac{1}{2}}, \quad (1)$$

where k is the the relation of volume mean radius and effective radius of the droplet size distribution, f_{ad} is the adiabaticity factor, c_w is the rate of increase of liquid water content with height in a moist adiabatically ascending air parcel, Q_{ext} is the Mie extinction efficiency, and ρ_w is the density of water (Grosvenor et al., 2018b). The parameters k , f_{ad} , c_w , Q_{ext} , τ_c , and r_e were diagnosed from the UCLALES-SALSA model.

The cloud parameters k , f_{ad} , c_w , and Q_{ext} vary with time along the cloud structure. However the actual values cannot be directly derived from MODIS observations and thus they are assumed to be constant and denoted by α for which an often used value for marine stratiform clouds is $1.37 \times 10^{-5} \text{ m}^{-\frac{1}{2}}$ (Quaas et al., 2006; Grosvenor et al., 2018b; Gryspeerd et al., 2022; Arola et al., 2022). Estimates of f_{ad} could possibly be improved combining MODIS/CALIOP observations. Consequently, CDNC values can be obtained from

$$N_d = \alpha \tau_c^{\frac{1}{2}} r_e^{-\frac{5}{2}}. \quad (2)$$

LWP values were calculated with the following equation (Wood, 2006)

$$LWP = 5/9 \rho_w r_e \tau_c. \quad (3)$$

In the analysis, we filtered the data so that we only considered cloudy columns where $\tau > 4$ and $4 \mu\text{m} < r_e < 15 \mu\text{m}$ similar to Gryspeerd et al. (2019) and Arola et al. (2022). An example of cloud field properties can be seen in Figure S3.

2.2 RESULTS

2.2.1 The effect of cloud internal variability on retrieved CDNC and LWP

First, to get an indication on how the cloud cell level variability affects the satellite retrieved LWP adjustment, we compared model predicted CDNC and LWP values with CDNC and LWP values calculated using Equations (1)-(3). We carried out an ensemble of UCLALES-SALSA simulations, varying the conditions for cloud formation and the number concentrations of aerosol particles, and then analysed the simulated CDNC and LWP values with the approaches detailed below.

As an example, Figure 2 shows the cloud droplet number concentration over the model domain of a simulation where the model was initialized with total aerosol number concentration of 300 cm^{-3} . The leftmost panel in Figure 2 represents CDNC values diagnosed directly from the UCLALES-SALSA model. In the middle panel, CDNC was calculated from Equation (1), using LES simulated values for k , f_{ad} , c_w , Q_{ext} , τ_c , and r_e in the equation. In the rightmost panel CDNC was calculated from Equation (2) assuming constant α of $1.37 \times 10^{-5} \text{ m}^{-\frac{1}{2}}$, and using simulated τ_c and r_e .

The leftmost panel shows a closed cell type structure in the cloud with lower values for CDNC at the boundaries of the cells. A snapshot of LWP is shown in Supplementary Material, Figure S3. Note that the simulated cloud in Figure 2 is fully overcast,




Figure 2. CDNC at the cloud top a) from the direct output of UCLALES-SALSA, b) calculated using Equation (1) with UCLALES-SALSA simulated values for all parameters, c) using Equation (2), d) relative biases in CDNC between UCLALES-SALSA and Equation (1), e) relative biases in CDNC between UCLALES-SALSA and Equation (2).

also at the cloud cell edges. Comparing Figures 2a and b, we can see that when all the parameters in Equation (1) are from LES simulations, CDNC corresponds quite well with the model values, showing similar structure although overestimating the CDNC throughout the model domain. However, the averaging of satellite data will mitigate this since spatial aggregation of the data will reduce the maximum CDNC values (see the differences between Figures S9 and S10 in Supporting Information) making the CDNC distribution more narrow (Figure S11). This is also in line with observations where aircraft and satellite observed CDNC are compared (Gryspeerd et al., 2022).

Comparing Figures 2a and c, we can see that when we use a fixed value for α , the satellite equation exhibits inverse behaviour at cloud cell boundaries compared to the direct output of the model, i.e., CDNC increases towards the boundaries of the cloud cells. This indicates that the assumptions of e.g., adiabaticity does not hold at the cloud cell boundaries. This is also in line with

140 a previous study by Feingold et al. (2022). Biases in LWP also occur differently across cloudy areas (Figure S5). Cloud cell boundaries tend to have low biased LWP values while cloud cell centers are biased high. In cloud cell boundaries processes such as entrainment and lateral mixing leads to sub-adiabaticity. Since these sources of variability are not considered in the formulation of satellite retrieval equations, there are important deviations from the assumptions of vertically constant values for droplet number concentration, droplet size distribution breadth and adiabaticity. A more detailed analysis of CDNC biases related to changes in LWP, cloud effective radius and adiabatic factor has been included in Figure S15.

145




Figure 3. Joint and marginal histograms for LWP and CDNC values using a) UCLALES-SALSA and b) Equation (2) at a time instance of 6 hours. Simulations are colour coded according to CCN concentrations used in the model initialization. The intensity of colour in joint histograms increases when the probability increases. The probability is represented as a density function calculated as counts/sum(counts)/bin area. Continuous lines indicate the arithmetic mean.

To see how the discrepancy between the cloud properties diagnosed directly from UCLALES-SALSA and using Equation (3) translate to differences in the correlation between CDNC and LWP, we calculated this relation for all ensemble members. Figure 3 illustrates LWP as a function of CDNC for direct model input and calculated using Eqs (2) and (3) for the three different initial CCN concentrations at 6 hours into the simulation. For all initial CCN concentrations, the direct model output indicates an almost linear correlation in the log-log scale within single cloud scenes. For an individual simulation, the positive slope between CDNC and LWP reflects the horizontal structure of the cell, where air flows from the core, characterized by high LWP and high CDNC, outward toward the cell edges with lower LWP and CDNC. The lowest CCN case exhibits a drop in LWP at highest CDNC values. However, the amount of data points is very low at highest CDNC values. In contrast, when

150

using Equations (2) and (3) the variability and the unphysical behaviour in N_d at the boundaries of cloud cells yield in curves which reach a local maximum and for higher CDNC values exhibit a downward slope with increasing CDNC. In addition, CDNC values have a clear high bias. In this case, satellite derived CDNC values are at least two times higher than the direct LES values. Values are positively biased due to the assumption of vertically uniform cloud columns which is not valid in thin cloud layers such as those observed at cloud cell edges. Although the spatial resolution in LES is much higher than in satellite data and it has been shown that spatial and temporal averaging affects the CCN-LWP correlation (Rosenfeld et al., 2023), this behaviour is similar to what is seen in satellite data and what was also demonstrated with synthetic data by Arola et al. (2022). This same behaviour was seen in all of the ensemble members over all analysed time instances and all CCN concentrations (Figures S6-S8).

The probability distributions of CDNC and LWP data are shown at the top and to the right of the coordinate frame, respectively in Figure 3. From these distributions we can see that LES derived data is skewed towards higher CDNC values while the LWP probability distributions look very similar for both LES and satellite equations. The sharp cut off of data at low LWP and CDNC values are caused by filtering of the data to include only values where $\tau > 4$ and $4\mu\text{m} < r_e < 15\mu\text{m}$. Earlier studies on satellite data limit this filtering to CDNC but not to LWP (Gryspeerd et al., 2019). However, due to doing pixel-by-pixel analysis for CDNC-LWP correlation, both CDNC and LWP data are filtered here.

Within individual ensemble members, the cloud internal variability contributes to the CDNC-LWP correlation and cannot be considered to be an aerosol effect on clouds, also shown by Zhou and Feingold (2023). In addition, in the analysis of satellite data, it is a common practice to avoid non-adiabatic clouds (e.g. Grosvenor et al., 2018b), so that issues in satellite retrieved cloud properties at, for example, cloud cell edges can be avoided. Previous studies have shown that selecting adiabatic pixels in a model and satellite analysis bring their results closer to each other (Dipu et al., 2022). Varble et al. (2023) also showed that removing the differences between the adiabaticity in an Earth System Model and satellite retrievals brings the observed and satellite retrieved LWP adjustment closer to each other.

2.2.2 The effect of combining different aerosol and time instances on CCN-LWP correlation

In addition to internal variability within clouds due to the dynamics affecting the cloud structure, cloud scenes can include clouds at different phases, e.g., transitioning between closed cell structure to open cell structure. Cloud scenes can also include large scale variability in the cloud geometric thickness and water content which are due to differences in meteorological conditions rather than caused by differences in aerosol concentration. To get an indication on how such variability affects the correlation between LWP and CDNC, we analysed simulated cloud scenes at different points in time (2 h, 6h, and 10 h). During the simulation, the closed cell structure transformed to an open cell structure for the case with the lowest initial aerosol load, and for the cases of higher aerosol load the size of convective cells increased.

Figure 4 shows the correlation between CDNC and LWP from direct model output and calculated using Equations (2) and (3) when all the cloud scenes are aggregated. Figure 4a shows that the model produces an overall positive correlation while satellite equations produce a similar shape correlation as shown in Figure 3b for one ensemble member, only spreading over a wider CDNC range due to the variability in CDNC concentrations. The direct model output resembles the shape of the




Figure 4. Joint histogram of LWP as a function of CDNC a) from the direct output of UCLALES-SALSA, b) calculated using Equations (2) and (3) assuming a constant α . Black continuous lines indicate the 25th, 50th and 75th percentiles of LWP per bin. The color scale indicate the probability density calculated as counts/sum(counts)/bin area.

correlation between CDNC and LWP simulated by the ICON model in Fons et al. (2024), while the satellite equation exhibits a decrease in LWP at CDNC values higher than 300 cm^{-3} .

190 To further investigate the differences between the LES model and satellite equations, we compared the mean LWPs of the LES domain using both direct output and satellite equation derived LWP for different initial aerosol concentrations. Figure 5 represents a proxy for satellite aggregation. It shows the LES domain mean LWP at three different time instances into the simulation for three different runs as a function of the initial CCN concentration. Solid lines denote the mean LWP in the domain and the shading indicates the standard deviation in the data.

195 Early into the simulation, we expect the simulated clouds to be close to a very similar phase of the cloud cycle and thus the difference in LWP between the different aerosol cases comes mainly from the differences in CCN concentrations. During the first two hours into the simulation there has not been enough time for precipitation to develop, and thus the LWP decreases slightly as a function of CCN concentration. Previous LES studies of the same DYCOMS case have produced qualitatively similar results (Ackerman et al., 2009; Bulatovic et al., 2019). Later in the simulation, precipitation is initiated with the lowest
 200 CCN concentration, and a typical shape of CCN-LWP correlation is reached where LWP first increases with CCN, and then decreases due to increased entrainment rate (Ackerman et al., 2009; Bulatovic et al., 2019). However, the difference in LWP between CCN concentrations of 180 cm^{-3} and 360 cm^{-3} does not change significantly from 6 to 10 hours into the simulation, which is opposite to the findings in Glassmeier et al. (2021), and can be characteristics to DYCOMS2 input profile with

quite small moisture inversion. The figure also illustrates, that although we also included non-adiabatic model grid points
205 in calculations where Equations (2) and (3) are used, changes in LWP with increasing CCN are strikingly similar to those
diagnosed from the LES model. There is a slight bias in satellite equation derived LWP (Figure S5) in all the cases, but the
relative changes correspond well with LES diagnosed relative changes in LWP. This indicates that if cloud condensation nuclei
concentrations are well-defined, changes in liquid water path due to changes in aerosol can be confidently determined using
satellite data.

210 This analysis indicates that, when using domain averaged CCN and LWP values, the non-adiabaticity of the cloud cell edges
does not contribute significantly to the "inverted v" shaped correlation between CDNC and LWP seen in satellite data. Although
there are issues in using Equations (2)-(3), coarse resolution of satellite data will reduce these issues significantly. Due to its
coarse resolution, satellite data can include both cloud cell centers and edges and could therefore, introduce bias in the retrieved
LWP values. However, based on Figure 5, in our simulated cases this aggregation of different cloud structures does not affect
215 the derived response in LWP to changes in CCN. We also tested this by using spatial averaging of 1.425 km by 1.425 km to
correspond to the spatial resolution of satellite data. Since radiances are directly proportional to cloud optical thickness, we use
COT values in cloudy columns as a weighting factor to perform horizontal averaging operations along subdomains. Figures
S9 and S10 in Supporting Information illustrate that spatially averaged data shows very similar cloud field properties with less
frequent large CDNC relative deviations because averaging reduces the variability in r_e (Figure S11). The CCN-LWP shaped
220 correlation for different time instances and CCN scenarios lack of the inverted v-shape (Figures S12, S13, and S14).

2.2.3 The effect of satellite instrument uncertainty or variability on retrieved CDNC and LWP

In addition to variability in cloud properties that originate from variability in aerosol concentrations and meteorological con-
ditions, satellite instruments also include uncertainty originating from instrument noise as well as three-dimensional radiative
effects. Such variability or noise will further affect the correlation between CDNC and LWP. Here we repeated the analysis
225 combining all analysed cases and adding 20% variability in both τ_c and r_e then calculating CDNC and LWP from Equations
(2) and (3). The level of variability was chosen to be in line with those used in Arola et al. (2022).

Figure 6 shows that adding noise produces a CCN-LWP correlation which increases, reaches a local maximum and is
followed by a decreasing CCN-LWP correlation. However, the additional noise does not affect the correlation between CDNC
and LWP compared to Figure 4b. Both Figures 3b) and 4b) exhibit a similar "inverted v" behaviour of the same magnitude. At
230 high CDNC values the negative correlation becomes less pronounced due to the variability in x-scale, i.e., increased variability
in CDNC.

3 Conclusions

Our LES simulations show that variability in cloud properties when including different cloud types, CCN concentrations,
and clouds in different phases of their cycle will bias satellite derived correlation between CDNC and LWP similar to Arola
235 et al. (2022). The root cause for this is that variability in cloud effective radius causes stronger positive bias in cloud droplet




Figure 5. LWP as a function of CCN from the direct output of UCLALES-SALSA and calculated using Equation (3). Time instances are color coded. Shaded areas indicate the spread of values in terms of the standard deviation.

number concentration when using the retrieval equation (2). Although our LES simulations include a detailed description of aerosol-cloud interactions, they do not consider the following potential error sources: 3D radiative effects in broken cloud fields, viewing geometry effects on the penetration depth bias, cloud heterogeneity at regional scale, and changes in surface reflectivity induced by changes in cloud coverage (Grosvenor et al., 2018a). Nonetheless, we could not find evidence in our model results to validate the persistent negative LWP adjustment predicted by satellite-equations for stratocumulus clouds affected by aerosol perturbations.

Cloud cell level variability in cloud effective radius and cloud optical thickness caused significantly different response in LWP with respect to changes in CDNC within individual simulations with different aerosol loads. However, when comparing the direct output of LWP from LES and those derived using Equation (3) for different CCN values, both show remarkably similar response between CCN and LWP. This indicates that although adiabaticity assumption near cloud edges causes error in CDNC and LWP, on the average Equation (3) derived LWP corresponds well with averaged LWP diagnosed from the LES output (see Figure 5). Furthermore, the relatively coarse resolution of satellite retrievals mitigates the impact of cloud cell level variability through averaging. However, CDNC is overestimated when using Equation (2) especially at the lower LWP values and this highlights the need to obtain a good constraint for CCN instead of using satellite derived CDNC as a proxy for CCN. Based on this, determining LWP requires careful selection of clouds, estimating the mean LWP for different aerosol loads, e.g., over regions where there has been a clear change in aerosol emissions. Since this study focuses only on one cloud

Satellite retrieval equations + 20% variability in CER and COT




Figure 6. Joint histogram of LWP as a function of CDNC calculated using Equations (2) and (3) assuming a constant α and including 20% variability in τ_c and r_e . Black continuous lines indicate the 25th, 50th and 75th percentiles of LWP per bin. The color scale indicate the probability density calculated as counts/sum(counts)/bin area.

case, analysis could be extended to cover wider variability in cloud conditions using for example large scale aerosol-aware high resolution climate models, which would capture also the mesoscale variability. The "inverted v" shape functionality of LWP adjustments is seen in simulations of GCMs of current generation and could be introducing confounding effects into the effective radiative forcing of aerosol-cloud-interactions (ERFaci) (Mülmenstädt et al., 2024). This clearly highlights that the behaviour of cloud water in response to changes in aerosol remains an open question and current knowledge does not support modifying the climate model cloud schemes to produce the "inverted v" behaviour.

Code availability. Large-eddy-simulations were performed with UCLALES-SALSA (DEV branch version November, 2023) available from <https://github.com/UCLALES-SALSA/UCLALES-SALSA/tree/DEV>. Input files used to initialise the model can be built as it is shown in section S1 of the supporting information.

Data availability. Datasets of cloud properties derived from simulations and equations for satellite retrievals used are available at <https://fmi.b2share.csc.fi/records/8fc77f2c6a8a4deab3de2efd46683010> (Kokkola et al., 2024)

Supplement. There is supplementary information available for this study including:

1. Model initial settings
 - 265 2. Conditional sampling of modeled cloud properties
 3. Surrogates of satellite-retrievals for CDNC and r_e
 4. Liquid water path susceptibility to changes in CDNC
 5. Spatial aggregation of cloud properties
 6. Spatial aggregation effects on liquid water path susceptibility to changes in CDNC
- 270 *Author contributions.* Conceptualization: HK, AA, AP, SC, JT, SR, TM, formal analysis: HK, AP, JT, SC, funding acquisition: HK, SR, TM, investigation: HK, AP, JT, SC, methodology: HK, AA, AL, AP, JT, PK, SC, SR, THV, software: HK, JT, SR, SC, supervision: HK, AA, SR, validation: HK, SC, visualization: HK, SC, writing - original draft preparation: HK, SC, SR, writing - review and editing: HK, AA, AL, AP, JT, SC, SR, THV, TM, EG

Competing interests. Authors declare that no competing interests are present.

- 275 *Acknowledgements.* This project has received funding from Horizon Europe programme under Grant Agreement No 101137680 via project CERTAINTY (Cloud-aERosol inTeractions & their impActs IN The earth sYstem), Marie Skłodowska-Curie Actions (MSCA) of the European Union (EU) under Grant Agreement No 101072354, Horizon 2020 programme under Grant Agreement No 821205 via project FORCeS (Constrained aerosol forcing for improved climate projections), Research Council of Finland grants No 337549 (Atmosphere and Climate Competence Centre, ACCC) and No 339885 BSOA-BORE (Are Biogenic Secondary Organic Aerosols Climatically Significant in the Boreal
- 280 Region?) and a Royal Society University Research Fellowship (URF/R1/191602)

References

- Ackerman, A. S., vanZanten, M. C., Stevens, B., Savic-Jovicic, V., Bretherton, C. S., Chlond, A., Golaz, J.-C., Jiang, H., Khairoutdinov, M., Krueger, S. K., Lewellen, D. C., Lock, A., Moeng, C.-H., Nakamura, K., Petters, M. D., Snider, J. R., Weinbrecht, S., and Zulauf, M.: Large-Eddy Simulations of a Drizzling, Stratocumulus-Topped Marine Boundary Layer, *Monthly Weather Review*, 137, 1083–1110, 285 <https://doi.org/10.1175/2008MWR2582.1>, 2009.
- Ahola, J., Korhonen, H., Tonttila, J., Romakkaniemi, S., Kokkola, H., and Raatikainen, T.: Modelling mixed-phase clouds with the large-eddy model UCLALES–SALSA, *Atmospheric Chemistry and Physics*, 20, 11 639–11 654, <https://doi.org/10.5194/acp-20-11639-2020>, 2020.
- Albrecht, B. A.: Aerosols, cloud microphysics, and fractional cloudiness, *Science*, 245, 1227–1230, 1989.
- Arola, A., Lipponen, A., Kolmonen, P., Virtanen, T. H., Bellouin, N., Grosvenor, D. P., Gryspeerdt, E., Quaas, J., and Kokkola, H.: Aerosol 290 effects on clouds are concealed by natural cloud heterogeneity and satellite retrieval errors, *Nature Communications*, 13, 7357, 2022.
- Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A.-L., Dufresne, J.-L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle, F., Mauritsen, T., McCoy, D. T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding Global Aerosol Radiative Forcing of Climate Change, *Reviews 295 of Geophysics*, 58, e2019RG000 660, <https://doi.org/https://doi.org/10.1029/2019RG000660>, e2019RG000660 10.1029/2019RG000660, 2020.
- Boutle, I., Price, J., Kudzotsa, I., Kokkola, H., and Romakkaniemi, S.: Aerosol–fog interaction and the transition to well-mixed radiation fog, *Atmospheric Chemistry and Physics*, 18, 7827–7840, <https://doi.org/10.5194/acp-18-7827-2018>, 2018.
- Bulatovic, I., Ekman, A. M. L., Savre, J., Riipinen, I., and Leck, C.: Aerosol Indirect Effects in Marine Stratocumulus: The Importance of Explicitly Predicting Cloud Droplet Activation, *Geophysical Research Letters*, 46, 3473–3481, 300 <https://doi.org/https://doi.org/10.1029/2018GL081746>, 2019.
- Calderón, S. M., Tonttila, J., Buchholz, A., Joutsensaari, J., Komppula, M., Leskinen, A., Hao, L., Moisseev, D., Pullinen, I., Tiitta, P., Xu, J., Virtanen, A., Kokkola, H., and Romakkaniemi, S.: Aerosol–stratocumulus interactions: towards a better process understanding using closures between observations and large eddy simulations, *Atmospheric Chemistry and Physics*, 22, 12 417–12 441, 305 <https://doi.org/10.5194/acp-22-12417-2022>, 2022.
- Dipu, S., Schwarz, M., Ekman, A. M. L., Gryspeerdt, E., Goren, T., Sourdeval, O., Mülmenstädt, J., and Quaas, J.: Exploring Satellite-Derived Relationships between Cloud Droplet Number Concentration and Liquid Water Path Using a Large-Domain Large-Eddy Simulation, *Tellus B: Chemical and Physical Meteorology*, <https://doi.org/10.16993/tellusb.27>, 2022.
- Feingold, G., Goren, T., and Yamaguchi, T.: Quantifying albedo susceptibility biases in shallow clouds, *Atmospheric Chemistry and Physics*, 310 22, 3303–3319, <https://doi.org/10.5194/acp-22-3303-2022>, 2022.
- Fons, E., Naumann, A. K., Neubauer, D., Lang, T., and Lohmann, U.: Investigating the sign of stratocumulus adjustments to aerosols in the global storm-resolving model ICON, *EGUsphere*, 2024, 1–30, <https://doi.org/10.5194/egusphere-2024-195>, 2024.
- Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M., and Zhang, H.: *The Earth’s Energy Budget, Climate Feedbacks, and Climate Sensitivity*, p. 923–1054, Cambridge University Press, 315 Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.009>, 2021.
- Glassmeier, F., Hoffmann, F., Johnson, J. S., Yamaguchi, T., Carslaw, K. S., and Feingold, G.: Aerosol-cloud-climate cooling overestimated by ship-track data, *Science*, 371, 485–489, <https://doi.org/10.1126/science.abd3980>, 2021.

- Grosvenor, D. P., Sourdeval, O., and Wood, R.: Parameterizing cloud top effective radii from satellite retrieved values, accounting for vertical photon transport: quantification and correction of the resulting bias in droplet concentration and liquid water path retrievals, *Atmospheric Measurement Techniques*, 11, 4273–4289, <https://doi.org/10.5194/amt-11-4273-2018>, 2018a.
- Grosvenor, D. P., Sourdeval, O., Zuidema, P., Ackerman, A., Alexandrov, M. D., Bennartz, R., Boers, R., Cairns, B., Chiu, J. C., Christensen, M., Deneke, H., Diamond, M., Feingold, G., Fridlind, A., Hünerbein, A., Knist, C., Kollias, P., Marshak, A., McCoy, D., Merk, D., Painemal, D., Rausch, J., Rosenfeld, D., Russchenberg, H., Seifert, P., Sinclair, K., Stier, P., van Diedenhoven, B., Wendisch, M., Werner, F., Wood, R., Zhang, Z., and Quaas, J.: Remote Sensing of Droplet Number Concentration in Warm Clouds: A Review of the Current State of Knowledge and Perspectives, *Reviews of Geophysics*, 56, 409–453, <https://doi.org/https://doi.org/10.1029/2017RG000593>, 2018b.
- Gryspeerd, E., Goren, T., Sourdeval, O., Quaas, J., Mülmenstädt, J., Dipu, S., Unglaub, C., Gettelman, A., and Christensen, M.: Constraining the aerosol influence on cloud liquid water path, *Atmospheric Chemistry and Physics*, 19, 5331–5347, <https://doi.org/10.5194/acp-19-5331-2019>, 2019.
- Gryspeerd, E., McCoy, D. T., Crosbie, E., Moore, R. H., Nott, G. J., Painemal, D., Small-Griswold, J., Sorooshian, A., and Ziemba, L.: The impact of sampling strategy on the cloud droplet number concentration estimated from satellite data, *Atmospheric Measurement Techniques*, 15, 3875–3892, <https://doi.org/10.5194/amt-15-3875-2022>, 2022.
- Jia, H., Ma, X., Quaas, J., Yin, Y., and Qiu, T.: Is positive correlation between cloud droplet effective radius and aerosol optical depth over land due to retrieval artifacts or real physical processes?, *Atmospheric Chemistry and Physics*, 19, 8879–8896, <https://doi.org/10.5194/acp-19-8879-2019>, 2019.
- Kokkola, H., Korhonen, H., Lehtinen, K. E. J., Makkonen, R., Asmi, A., Järvenoja, S., Anttila, T., Partanen, A.-I., Kulmala, M., Järvinen, H., Laaksonen, A., and Kerminen, V.-M.: SALSA: a Sectional Aerosol module for Large Scale Applications, *Atmospheric Chemistry and Physics*, 8, 2469–2483, <https://doi.org/10.5194/acp-8-2469-2008>, 2008.
- Kokkola, H., Tonttila, J., Calderón, S. M., Romakkaniemi, S., Lipponen, A., Peräkörpi, A., Mielonen, T., Virtanen, T. H., Kolmonen, P., and Arola, A.: Datasets used in Kokkola et al. (2024) "Model analysis of biases in satellite diagnosed aerosol effect on cloud liquid water path", <https://doi.org/10.57707/FMI-B2SHARE.8FC77F2C6A8A4DEAB3DE2EFD46683010>, 2024.
- Mülmenstädt, J., Gryspeerd, E., Dipu, S., Quaas, J., Ackerman, A. S., Fridlind, A. M., Tornow, F., Bauer, S. E., Gettelman, A., Ming, Y., Zheng, Y., Ma, P.-L., Wang, H., Zhang, K., Christensen, M. W., Varble, A. C., Leung, L. R., Liu, X., Neubauer, D., Partridge, D. G., Stier, P., and Takemura, T.: General circulation models simulate negative liquid water path–droplet number correlations, but anthropogenic aerosols still increase simulated liquid water path, *EGUsphere*, 2024, 1–29, <https://doi.org/10.5194/egusphere-2024-4>, 2024.
- Platnick, S.: Vertical photon transport in cloud remote sensing problems, *Journal of Geophysical Research: Atmospheres*, 105, 22 919–22 935, <https://doi.org/https://doi.org/10.1029/2000JD900333>, 2000.
- Platnick, S., King, M. D., Meyer, K. G., Wind, G., Amarasinghe, N., Marchant, B., Arnold, G. T., Zhang, Z., Hubanks, P. A., Ridgway, B., et al.: MODIS cloud optical properties: User guide for the Collection 6 Level-2 MOD06/MYD06 product and associated Level-3 Datasets, Version, 1, 145, 2015.
- Quaas, J., Boucher, O., and Lohmann, U.: Constraining the total aerosol indirect effect in the LMDZ and ECHAM4 GCMs using MODIS satellite data, *Atmospheric Chemistry and Physics*, 6, 947–955, 2006.
- Rosenfeld, D., Kokhanovsky, A., Goren, T., Gryspeerd, E., Hasekamp, O., Jia, H., Lopatin, A., Quaas, J., Pan, Z., and Sourdeval, O.: Frontiers in Satellite-Based Estimates of Cloud-Mediated Aerosol Forcing, *Reviews of Geophysics*, 61, e2022RG000 799, <https://doi.org/https://doi.org/10.1029/2022RG000799>, e2022RG000799 2022RG000799, 2023.

- 355 Stevens, B., Moeng, C., Ackerman, A. S., Bretherton, C. S., Chlond, A., de Roode, S., Edwards, J., Golaz, J., Jiang, H., Khairoutdinov, M., Kirkpatrick, M. P., Lewellen, D. C., Lock, A., Müller, F., Stevens, D. E., Whelan, E., and Zhu, P.: Evaluation of Large-Eddy Simulations via Observations of Nocturnal Marine Stratocumulus, *Monthly Weather Review*, 133, 1443 – 1462, <https://doi.org/10.1175/MWR2930.1>, 2005.
- Tonttila, J., Maalick, Z., Raatikainen, T., Kokkola, H., Kühn, T., and Romakkaniemi, S.: UCLALES–SALSA v1.0: a large-eddy
360 model with interactive sectional microphysics for aerosol, clouds and precipitation, *Geoscientific Model Development*, 10, 169–188, <https://doi.org/10.5194/gmd-10-169-2017>, 2017.
- Twomey, S.: Pollution and the planetary albedo, *Atmospheric Environment* (1967), 8, 1251–1256, 1974.
- Varble, A. C., Ma, P.-L., Christensen, M. W., Mülmenstädt, J., Tang, S., and Fast, J.: Evaluation of liquid cloud albedo susceptibility in E3SM using coupled eastern North Atlantic surface and satellite retrievals, *Atmospheric Chemistry and Physics*, 23, 13 523–13 553,
365 <https://doi.org/10.5194/acp-23-13523-2023>, 2023.
- Wood, R.: Relationships between optical depth, liquid water path, droplet concentration and effective radius in an adiabatic layer cloud, https://atmos.uw.edu/~robwood/papers/chilean_plume/optical_depth_relations.pdf, 2006.
- Zhang, J., Chen, Y.-S., Yamaguchi, T., and Feingold, G.: Cloud water adjustments to aerosol perturbations are buffered by solar heating in non-precipitating marine stratocumuli, *EGUsphere*, 2024, 1–27, <https://doi.org/10.5194/egusphere-2024-1021>, 2024.
- 370 Zhou, X. and Feingold, G.: Impacts of Mesoscale Cloud Organization on Aerosol-Induced Cloud Water Adjustment and Cloud Brightness, *Geophysical Research Letters*, 50, e2023GL103 417, <https://doi.org/https://doi.org/10.1029/2023GL103417>, e2023GL103417 2023GL103417, 2023.