

Can large-scale tree cover change negate climate change impacts on future water availability?

Freek Engel¹, Anne J. Hoek van Dijke², Caspar T.J. Roebroek^{3, 4, 5}, and Imme Benedict¹

¹Meteorology and Air Quality Group, Wageningen University and Research, Wageningen, The Netherlands

²Max Planck Institute for biogeochemistry, Jena, Germany

³European Commission, Joint Research Centre (JRC), Ispra, Italy

⁴Hydrology and Water Management group, Wageningen University and Research, Wageningen, The Netherlands

⁵Institute for Atmospheric and Climate Science, Eidgenössische Technische Hochschule (ETH) Zurich, Zurich, Switzerland

Correspondence: Freek Engel (freek.engel@wur.nl) and Imme Benedict (imme.benedict@wur.nl)

Abstract. The availability of fresh water over land may become increasingly scarce under climate change, and natural and human-induced tree cover changes can further enhance or negate the water scarcity. Previous studies showed that global tree cover change can have large impacts on water availability under current climate conditions, but did not touch upon the implications of global tree cover change under climate change. Here, we study the hydrological impacts of large-scale tree cover

5 change (climate-induced changes in combination with large-scale afforestation) in a future climate (SSP3-7.0) following an interdisciplinary approach. By combining data from five CMIP6 climate models with a future potential tree cover dataset, six Budyko models, and the UTrack moisture recycling dataset, we can disentangle the impacts of climate change and future tree cover change on evaporation, precipitation, and runoff. We quantify per grid cell and for five selected river basins (Yukon, Mississippi, Amazon, Danube, and Murray-Darling) if tree cover changes enhance or counteract the climate-driven changes
10 in runoff due to their impact on evapotranspiration and moisture recycling. Globally averaged, the impacts of climate change and large-scale tree cover change on runoff are of similar magnitude with opposite signs. While climate change increases the global runoff, the changing tree cover reverses this effect which overall results in a limited net impact on runoff relative to the present climate and current tree cover. Nevertheless, locally the change in runoff due to tree cover change and climate change can be substantial with increases and decreases of more than 100 mm yr^{-1} . We show that for approximately 16 % of the land
15 surface, tree cover change can increase the water availability significantly. However, ~~we also find that~~, for 14 % of the land surface, both tree cover change and climate change ~~might~~ could decrease water availability ~~with~~ by more than 5 mm yr^{-1} . For each of the selected catchments, the direction and magnitude of the impacts of climate change and tree cover change vary, with dominating climate change impacts in all basins except the Mississippi River basin. Our results show that ecosystem restoration projects targeting an altered tree cover should consider the corresponding hydrological impacts to limit unwanted (non-)local
20 reductions in water availability.

1 Introduction

Forests play an important role in, among others, the conservation of biodiversity, reduction of soil erosion, and mitigation of climate change (Herrick et al., 2019). Therefore, there have been many local and global initiatives to increase tree cover, such as the Bonn challenge, Grain for Green, the 20x20 Initiative, Billion Trees Campaign, and AFR100. Climate change, especially 25 climatic drying, has consequences for the ability to restore forests. At the same time, large-scale forest restoration has major impacts for local and global water availability, and can reduce water availability in water scarce regions (Hoek van Dijke et al., 2022).

Over the past decades, both climate warming and land cover changes have impacted global freshwater availability by changing evaporation and precipitation (Fahrländer et al., 2024). While recent climate warming likely enhanced global evaporation 30 and precipitation over land (Douville et al., 2021), the simultaneous changes in land cover showed contrasting impacts on evaporation. For example, ~~land-cover~~ land cover changes that occurred between 1950 – 2000 decreased evaporation by 5 % (Sterling et al., 2013), whereas from the 1980s onwards, global vegetation greening increased evaporation with by 3.7 % (Yang et al., 2023). Teuling et al. (2019) showed that both climate warming and land cover changes impacted evaporation over Europe with large regional differences. For instance, while evaporation increased with by more than 15 % between 1960 – 2010, 35 it decreased for parts of Southern Europe (Teuling et al., 2019). Furthermore, the greening of vegetation in China enhanced the yearly precipitation due to an increase in convective precipitation (Yu et al., 2020). These changes in evaporation and precipitation have impacted streamflow globally (Sterling et al., 2013; Teuling et al., 2019; Piao et al., 2007; Wang-Erlansson et al., 2018), and thereby affected the occurrence of floods and droughts, production of hydropower, and availability of drinking and 40 irrigation water. However, the projected increases in climate warming and land cover changes in the (near) future could impact the water fluxes and availability even further.

Since climate warming can increase the energy available for evaporation and enhances the moisture holding capacity of air, these effects are expected to further increase evaporation, mean precipitation, and extreme precipitation events (Trenberth, 2011). At the same time, the future holds longer periods of dry spells and droughts in many regions (Milly and Dunne, 2016), and therefore water scarcity will likely increase for a growing fraction of land, impacting many lives and ecosystems (Caretta 45 et al., 2022).

The combination of climate warming and human interventions, such as afforestation and deforestation, can affect the global tree cover in a future climate. Roebroek (2023) recently calculated a future potential tree cover, representing the maximum number of trees that could grow on Earth given the climate and soil characteristics, for different climate pathways. The study found that a warmer climate can support tree growth in the colder high latitudes, whereas increasing aridity is expected to 50 reduce the forest cover in the American and African tropics (see Fig. 1, showing the differences between the present and future potential tree covers). At the same time, there is a strong global incentive to increase tree cover due to the benefits for climate mitigation, ecosystem restoration, and conservation of species (Bastin et al., 2019; Griscom et al., 2017). Roebroek (2023) showed that global climate change decreases the overall global potential for tree restoration, however, the tree carrying

capacity is 55 % above the current tree cover. This estimation of potential tree cover change under climate change allows us to
55 study the combined effects of future global warming and tree cover change on water availability.




Figure 1. Tree cover change (TCC) in percentages, the TCC is defined as the difference between the potential tree cover in a future climate (2041-2060) and the tree cover in a present climate (2000). For the future potential tree cover we use the average potential tree cover data based on bioclimatic variables, derived from the outputs of five CMIP6 models (CMCC-ESM2; INM-CM5-0; IPSL-CM6A-LR; MIROC6; and UKESM1-0-LL) under climate change pathway SSP3-7.0.

Changes in tree cover impact local and regional water availability through their effect on evapotranspiration and precipitation (Ellison et al., 2017; Hoek van Dijke et al., 2022; King et al., 2024; Zhang et al., 2022a). Forest evapotranspiration is higher compared to evapotranspiration from short vegetation and bare land (Zhang et al., 2001) since trees increase the availability of energy (through albedo), and water (through their deep rooting systems) for evapotranspiration, and increase the aerodynamic 60 conductance. In addition, trees can impact precipitation directly by locally increasing convection and turbulence in the atmosphere, while trees can also affect precipitation indirectly by enhancing the recycling of evaporated moisture both locally or regionally (Meier et al., 2021; Ellison et al., 2017; De Hertog et al., 2023; Wang-Erlundsson et al., 2018). Therefore, future forest restoration projects can potentially be used to increase water availability in water scarce regions (Staal et al., 2024b).

Previous literature has already explored the effects of climate change and large-scale tree cover change, nevertheless, uncertainty persists regarding the separate and combined impacts on global water availability in a future climate. Tuinenburg et al. (2022) analysed how the current potential tree cover on a global scale could mitigate future drying trends, however, this study only focused on global precipitation and did not consider an altered potential tree cover in a future climate. King et al. (2024) applied a global forestation scenario in one earth system model and found regional reductions in water availability up to 15 %. More local studies showed that in mountainous catchment areas, climate-driven changes in vegetation could mitigate or 70 reverse the climate-driven increases in runoff (Rasouli et al., 2019). Another example is that in Northern China the soil water

availability decreased before 2000 and increased afterwards due to climate change and vegetation greening, although the effect of vegetation greening was minor compared to the climate change effect (Douville et al., 2021). Given the remaining uncertainties, a global comprehensive analysis regarding the impacts of tree cover and climate change, and their separate contributions, is needed which can help to secure freshwater availability in the future, and to support strategic tree restoration planning.

75

Here, we study the combined and separate impacts of climate change and large-scale tree cover change on terrestrial evaporation, precipitation, and runoff worldwide. We take a data-driven interdisciplinary approach, by combining tree cover maps with Budyko models and projected climate data from the Coupled Model Intercomparison Project phase 6 (CMIP6), to calculate present and future water fluxes. The future climate data is obtained for the Shared Socioeconomic Pathway (SSP) with 80 an intermediate to strong climate change signal (SSP3-7.0). This climate pathway involves high greenhouse gas emissions which result in an average increase in global surface air temperature of 1.4° for 2041–2060 relative to 1995–2014 (Lee et al., 2021). Our implemented tree cover change scenario represents the tree cover carrying capacity under climate pathway SSP3-7.0 and includes natural changes in tree cover, as well as, human-induced large-scale tree planting. In this study we compare 85 the impacts of climate change and tree cover change on water fluxes over land and analyse where these effects can enhance or counteract each other on grid cell level and at catchment scale. For five selected catchments we provide insight whether climate or large-scale tree cover change can be a dominant driver of water availability change in a future climate.

2 Methodology

There are three research scenarios that guide this work to investigate the impact of climate change and tree cover change on the hydrological fluxes precipitation (P), evapotranspiration (ET), and runoff (Q, also referred to as water availability). ~~These~~ 90 ~~These~~ three scenarios are: 1) scenario present climate, 2) scenario climate change (CC), and 3) scenario climate change with tree cover change and moisture recycling change (CC + TCC) (Fig. 2, Table 1). ~~All research scenarios~~ ~~We calculate the effects of climate change as the difference between scenario present climate and scenario CC, and calculate the effects of tree cover change as the difference between scenario CC and scenario CC + TCC (Fig. 2). For each research scenario, we~~ use P and potential evapotranspiration (PET) datasets from five CMIP6 climate models (Sect. 2.1), ~~along with~~ a tree cover dataset (Sect. 95 2.2), ~~and Budyko models as input for the Budyko model calculations~~ (Sect. 2.3) ~~for the calculation of~~ to generate ET and Q fluxes. Furthermore, scenario CC + TCC includes the (non-)local indirect effects of changes in ET and P by accounting for an altered moisture recycling, obtained with the moisture tracking dataset UTrack (Sect. 2.4). ~~We therefore build on the general research methodology from Hoek van Dijke et al. (2022), extending their approach to assess the hydrological effects of climate and tree cover change under future climate conditions.~~ All datasets in ~~this~~ ~~our~~ study are reprojected to a spatial resolution of 1° 100 by 1° to ensure compatibility with the spatial resolution of the UTrack dataset. In addition, datasets with a time dimension are aggregated to yearly averaged data for the use of the Budyko models. ~~An A simplified~~ overview of the ~~research scenarios with the corresponding input and output datasets and calculations~~ ~~main research steps is illustrated in Fig. 2 and a detailed research~~

overview is shown in Table 1. The analyses in this study are performed at a global scale and for five river basins: Yukon River basin, Mississippi River basin, Amazon River basin, Danube River basin, and the Murray-Darling River basin.




Figure 2. Simplified overview of the research methodology. This figure shows the research scenarios: 1) scenario present climate; 2) scenario climate change (CC); 3) scenario climate change with tree cover change with changed moisture recycling (CC + TCC); as well as the intermediate research steps. A detailed overview of the input and output data for each research step can be found in Table 1.

105 2.1 Climate input data from CMIP6 climate models

Five global climate model simulations from CMIP6 are selected to provide P and PET datasets for the different research scenarios in this study. For the present climate scenario, climate data is obtained from historical CMIP6 simulations for the time interval 1985–2014 and for the future climate scenarios, data is used from ScenarioMIP simulations under climate change pathway SSP3-7.0 for 2035–2064 (Table 1) (O’Neill et al., 2016). The P datasets are retrieved directly from the CMIP6 models, 110 however, due to the limited availability of global PET output from CMIP6 models for climate pathway SSP3-7.0, we use the PET datasets provided by Bjarke et al. (2023). Bjarke et al. (2023) calculated PET with the Priestley-Taylor method (Priestley and Taylor, 1972) based on the sensible and latent heat fluxes, mean surface air temperature, and surface air pressure for which data was directly retrieved from selected CMIP6 climate models for both the historical and ScenarioMIP simulations.

The five CMIP6 models in this study are selected based on the following criteria; 1) the model should have PET output 115 provided by Bjarke et al. (2023) and a future tree cover dataset for ScenarioMIP SSP3-7.0 from Roebroek (2023) (Sect. 2.2); 2) the spatial resolution of the CMIP6 model data should be close to the 1° by 1° spatial resolution used in this study; 3) the absolute percentual bias of the calculated historical PET compared to the ERA5Land dataset, as provided in Table 2 from Bjarke et al. (2023), should be smaller than 10 %; and 4) from each institute only one CMIP6 model is selected. The aforementioned

Table 1. [Overview](#) [Detailed overview](#) of the research methodology and showing the input and output data for each of the research steps. The abbreviations represent; P: precipitation from CMIP6, PET: potential evapotranspiration from CMIP6, ET: evapotranspiration, Q: runoff. Furthermore, 'mean' refers to the mean flux over the CMIP6-Budyko models and 'std' refers to the standard deviation of the flux over the CMIP6-Budyko models. Note that five CMIP6 models and six Budyko models result in 30 CMIP6-Budyko combinations for ET and Q. The general research methodology is adopted from Hoek van Dijke et al. (2022). For more information about the datasets and their sources, Table A1 provides an overview of the datasets used within our study.

Research scenario	Tree cover	Climate pathway	Calculations with six Budyko models		UTrack	
			Climate input data	Budyko output data	Input	Output
1. Present climate	Present tree cover (2000)	Historical (1985–2014)	5x P, 5x PET	30x ET, → ET _{mean} , ET _{std} 30x Q → Q _{mean} , Q _{std}		
2. Climate change (CC)		SSP3-7.0 (2035–2064)	5x P, 5x PET	30x ET, ↘ 30x Q 30x ΔET → ΔET _{mean} 30x ET, ↗ 30x Q	ΔET _{mean}	ΔP _{mean}
Climate change with tree cover change (intermediate step)	Future potential tree cover (2041–2060)		5x P+ ΔP _{mean} , 5x PET	30x ET, → ET _{mean} , ET _{std} 30x Q → Q _{mean} , Q _{std}		
3. Climate change with tree cover change with changed moisture recycling (CC+TCC)						

criteria result in the selection of the following five CMIP6 models: CMCC-ESM2; INM-CM5-0; IPSL-CM6A-LR; MIROC6; and UKESM1-0-LL, for which an overview is provided in Table 2.

The monthly P ($\text{kg m}^{-2} \text{s}^{-1}$) and PET (mm d^{-1}) datasets are aggregated to yearly data and averaged over a 30 year period (mm yr^{-1}) as indicated in Table 1. These yearly mean P and PET datasets are used as input for the Budyko calculations (Table 1 and Sect. 2.3). Note that large water bodies, the Antarctic region, and most of Greenland are masked out.

To gain insight into the differences between the five climate models, Fig. 3a shows PET and P over land in relation to the aridity index (PET / P) for a future climate. The variability between the models is highest for terrestrial P over humid regions, ranging from 837 mm yr^{-1} in the INM model to 2782 mm yr^{-1} in the CMCC model (Fig. 3a). The high variability in P between CMIP6 models over wet regions is also underlined by Yazdandoost et al. (2021) for historical data. The model spread for humid regions is also present for PET, however, the largest variability for PET is found over arid regions.

Table 2. Overview of the models from phase 6 of the Coupled Model Intercomparison Project (CMIP6) from which climate data for precipitation (P) and potential evapotranspiration (PET) was used in this study. The climate data is obtained for variant label 'r1i1p1f1' for all models, except for UKESM1-0-LL for which 'r1i1p1f2' is used due to data availability restrictions, meaning that a different forcing dataset was used for this model simulation in UKESM1. The lon x lat resolution in the table refers to the number of gridcells present for the longitude and latitude dimensions within the global dataset.

CMIP6 model	Institution (institution ID)	Original resolution (lon x lat) for P and PET	Model type*	Reference
CMCC-ESM2	Fondazione Centro Euro-Mediterraneo (CMCC)	288 x 192	Earth system	Lovato et al. (2022)
INM-CM5-0	Institute for Numerical Mathematics (INM)	180 x 120	Atmosphere-ocean general circulation	Volodin et al. (2017)
IPSL-CM6A-LR	Institut Pierre Simon Laplace (IPSL)	144 x 143	Earth system	Boucher et al. (2020)
MIROC6	Atmosphere and Ocean Research Institute, The University of Tokyo and Japan Agency for Marine-Earth Science and Technology (MIROC)	256 x 128	Atmosphere-ocean general circulation	Tatebe et al. (2019)
UKESM1-0-LL	Met Office Hadley Centre (MOHC)	192 x 144	Earth system	Sellar et al. (2019)

* As classified by Kuma et al. (2023)

130 2.2 Tree cover datasets

In this study, two tree cover datasets are applied: 1) a present tree cover dataset used for scenario present climate and scenario CC, and 2) a future potential tree cover dataset, representing the maximum tree carrying capacity under the studied climate pathway, which is used for scenario CC+TCC (Table 1). For the present tree cover, we use the global tree cover dataset for the 135 year 2000 provided by Hansen et al. (2013), which is generated with Landsat satellite data. For the future potential tree cover dataset, we use a dataset by Roebroek (2023) that predicts the tree cover carrying capacity for different climate scenarios, and is 140 adjusted for the natural occurrence of disturbances (e.g., forest fires, wind throws, insect outbreaks etc.). This tree cover data, here referred to as 'future potential tree cover' for simplicity, is created by integrating the previously mentioned tree cover (Hansen et al., 2013) and climate characterisations from the WorldClim V2 dataset (Fick and Hijmans, 2017) in a machine learning framework. Note that the dataset represents the natural capacity of the earth to support trees, and thus would allow for tree restoration on agricultural or urban land. In this study, we retrieved the future potential tree cover datasets for the time 145 range 2041–2060 under climate pathway SSP3-7.0 for each of the five selected CMIP6 models (Sect. 2.1) to create an averaged future potential tree cover dataset for scenario CC+TCC. The future potential tree cover maps represent a shorter time period (20 years) than the climatological data (30 years), as the future potential tree cover maps are only available for twenty-year




Figure 3. Illustration of variability over the CMIP6 and Budyko models in relation to the aridity index (PET / P); a) variability in terrestrial precipitation (P) and potential evapotranspiration (PET) fluxes for five CMIP6 models (CMCC-ESM2; INM-CM5-0; IPSL-CM6A-LR; MIROC6; and UKESM1-0-LL) under climate change pathway SSP3-7.0. b) variability in the evapotranspiration over precipitation ratio (ET / P) for tree cover and non-tree cover for six Budyko (Bud) models, the numbering of these models is consistent with the numbering in Table A2.

periods (Roebroek, 2023). We assume that the tree cover datasets for both the present and the future time period consist of 145 mature trees.

The tree cover change illustrates the differences between the two tree cover datasets, which is calculated by subtracting the present tree cover from the future potential tree cover (Fig. 1). ~~Note that the tree-cover change is not used for calculations and hence is solely used to visualize the differences in tree-cover between the scenarios present climate and CC, and scenario CC+TCC.~~ Globally, there is an average increase in tree cover of 15.5 %. Decreases in tree cover are found over the Amazon, 150 in northwestern North America, and in middle north Eurasia.

2.3 Budyko model calculation

For each research scenario, yearly mean ET and Q are calculated using the Budyko framework (Budyko, 1974), following the approach used by Hoek van Dijke et al. (2022); Teuling et al. (2019). This approach was tested by Hoek van Dijke et al. (2022) whereby the mean streamflow (Q) obtained with Budyko model calculations for the present climate was validated with observational data for 19 large river basins. A Budyko model describes the long-term mean partitioning of P into ET and Q, as a function of aridity (Fig. 3b) and has a general form like:

$$\frac{ET}{P} = f\left(\frac{PET}{P}, \omega\right) \quad (1)$$

where $\frac{ET}{P}$ is the fraction of precipitation partitioned into evaporation, $\frac{PET}{P}$ is the aridity index and ω is a model parameter. Previous studies have shown that ω is closely related to vegetation type (e.g. Zhang et al., 2001). In our study, we use an ensemble of six different Budyko models, that have a different model formulation, but represent a similar curve (Fig. 3, model equations in Table A2; Zhang et al. (2001, 2004); Zhou et al. (2015); Teuling et al. (2019); Oudin et al. (2008)). Each of these Budyko models was calibrated with lysimeter or streamflow data originating from different river basins, thereby representing different vegetation types (e.g. plantation or natural vegetation, or deciduous or evergreen forest) and climate conditions. Therefore, we include multiple Budyko models for our global scale calculations to represent the spread between different models and minimize potential biases related to e.g. climate conditions. After calculating the $\frac{ET}{P}$ fraction with a Budyko model, the $\frac{Q}{P}$ fraction is obtained by rewriting a simplified water balance for a multi-year timescale ($Q = P-ET$) to $\frac{Q}{P} = 1 - \frac{ET}{P}$. In this study, ET and Q are calculated by using P and PET datasets from the five CMIP6 models in combination with the fractional tree cover from the present and the future potential tree cover datasets following Table 1. For each grid cell, values for ET and Q are calculated as a fraction of tree-covered and non-tree-covered surfaces relative to their occurrence in the grid cell.

Fig. 3b shows the variability between the six Budyko models for a theoretical ET/P fraction in relation to a theoretical aridity index (PET/P) for (non-)tree-covered surfaces. The figure shows that there can be pronounced variability between the different Budyko models, however, it is also shown that for all models a larger fraction of P is evaporated for tree-covered surfaces compared to non-tree-covered surfaces. The combined variability of the climate input data from the CMIP6 models (Fig. 3a) and the variability between the Budyko models (Fig. 3b) provides an uncertainty estimate for the calculated ET and Q in this study.

2.4 UTrack moisture tracking dataset

Following large-scale tree cover change, the increase in terrestrial ET will increase the (terrestrial) P both locally and remotely through moisture recycling. In scenario CC+TCC we account for this local and remote change in P by using the UTrack dataset, created by Tuinenburg et al. (2020), in which the moisture recycling is quantified per grid cell (please note that there is a potential issue with the water balance, as noted by (De Petrillo et al., 2024)). This dataset is generated by combining ERA5 reanalysis data with UTrack, a Lagrangian model that tracks the transport of moisture through the atmosphere (Tuinenburg and Staal, 2020). The UTrack dataset contains atmospheric moisture trajectories, averaged over 2008-2017 (Tuinenburg et al., 2020), that show the transport of evaporated moisture from a source location to precipitated moisture at a target location. We retrieved the dataset at a spatial resolution of 0.5° by 1° as the datasets of the selected CMIP6 models in our study are only available at relatively coarse spatial resolutions (Table 2). We retrieved the data for each month of the year and we aggregated these monthly moisture trajectories to the yearly timescale. The yearly average moisture trajectories were used to calculate how the mean tree cover-driven change in ET affects the global P flux (Table 1). The use of yearly trajectories was preferred over monthly trajectories since the Budyko approach is only valid on multi-year timescale. Hoek van Dijke et al. (2022) showed that temporal aggregation of UTrack, rather than temporal disaggregation of Budyko, resulted in similar patterns for moisture recycling. It

should be noted that altering the tree cover creates a feedback loop where a change in terrestrial ET can in turn affect the (terrestrial) P, which again impacts ET, although the impact on both fluxes becomes increasingly smaller for each cycle. In this 195 study, the changed moisture recycling is calculated twice for scenario CC+TCC, and the impacts of the changed moisture recycling on the P, ET, and Q fluxes are relatively small after the UTrack dataset is applied the second time (Table A3 and Table A4).

We underline that the UTrack dataset represents current atmospheric conditions (Tuinenburg et al., 2020) and therefore, this 200 study does not account for possible changes in moisture recycling due to changes in circulation which result from climate change. The UTrack dataset also does not include the feedbacks of an altered tree cover, even though tree cover change can impact the atmospheric conditions and circulation (De Hertog et al., 2023; Portmann et al., 2022; Davin and de Noblet-Ducoudré, 2010; Duveiller et al., 2018). Nevertheless, most moisture tracking models that are currently available rely on meteorological reanalysis data and thus these models are only valid under current climate and ~~land-cover~~land cover conditions. Hence, given the unavailability of a moisture tracking dataset for a future climate, and the approach of combining climate 205 datasets instead of actively running a global climate model, this study relies on the UTrack dataset for the future moisture recycling. The implications of this approach are further discussed in Sect. 4.

3 Results and Discussion

3.1 Global mean impact of climate change and future tree cover change on hydrological fluxes

First, we compare our global mean hydrological fluxes over land for the present climate with multi-model outcomes from 210 CMIP6, after which we discuss the impacts of scenarios climate change (CC) and climate change with tree cover change (CC+TCC). For the present climate scenario, the mean fluxes in this study are 565 mm yr^{-1} for ET over land and 386 mm yr^{-1} for Q (Fig. 4a). Our mean value of ET exceeds the 90 % confidence range for the multi-model averages over land by IPCC of $482 - 544 \text{ mm yr}^{-1}$, obtained with 32 CMIP6 models for 1995 – 2014, whereas our mean Q value is consistent with its corresponding range of $179 - 460 \text{ mm yr}^{-1}$ (Douville et al., 2021). Note that our mean terrestrial P flux of 952 mm yr^{-1} also 215 slightly exceeds the multi-model range of $723 - 942 \text{ mm yr}^{-1}$ by Douville et al. (2021). This indicates that the hydrological fluxes for our present climate scenario (1985 – 2014) for the five selected CMIP6 models in this study are at the higher end of the multi-model range.

Under climate change (scenario CC), the terrestrial P increases with $33 \pm 55 \text{ mm yr}^{-1}$ and terrestrial ET increases with $22 \pm 24 \text{ mm yr}^{-1}$, resulting in an average increase for Q of $11 \pm 39 \text{ mm yr}^{-1}$ (Fig. 4b, Table A3). These flux differences relative 220 to the present climate were calculated for each of the 30 CMIP6(-Budyko) model combinations, generating 30 Δflux values for each land grid cell after which the mean and standard deviation were computed for each grid cell. Although the mean changes for the hydrological fluxes are consistent with the climate change impacts described by Douville et al. (2021), in which 32 CMIP6 models are compared, our study shows larger corresponding standard deviations. These high standard deviations arise from deviations in both the climate input data as well as the use of six different Budyko models (Fig. 3). Especially the Q flux

shows a relatively large variability compared to the other hydrological fluxes, which is also mentioned by Li and Li (2022) for
225 SSP1-2.6, SSP2-4.5, and SSP5-8.5.

Implementing the future potential tree cover scenario by Roebroek (2023) in a future climate (scenario CC+TCC) generates
226 can generate mean hydrological impacts that are of similar magnitude as the effects of climate change (scenario CC). The
changing tree cover enhances We estimate the mean global ET to enhance with $28 \pm 19 \text{ mm yr}^{-1}$ (4.7 %) relative to scenario
227 CC due to an average global tree cover following an average increase of 15.5 % in global tree cover potential. This increase in
ET includes both the direct and indirect tree cover change effects, of which the latter relates to the altered P flux. The enhanced
228 ET can affect affects P over both land and ocean surfaces (Fig. A1), enhancing which could enhance P with 16 mm yr^{-1} over
land. Please note that the standard deviation for the altered P remains unchanged across scenarios CC and CC+TCC since
229 this standard deviation only shows the (constant) variability over the CMIP6 models. We find that approximately 60 % of the
additional evaporated moisture precipitates over land. The in scenario CC+TCC. In addition, the projected change in tree
230 cover can generate a shift in Q from a climate-driven net increase in Q is shifted to a slight net decrease after tree cover change
(scenario CC+TCC). Hence, globally. Therefore, on a global scale, the contrasting effects of climate change and tree cover
231 change generate large-scale forestation may result in a limited net effect for Q relative to the present climate on Q (Fig. 4b).
A similarity in magnitude of impact for Rasouli et al. (2019) also showed that the impacts of future vegetation change and
232 climate change on Q is also demonstrated regionally by Rasouli et al. (2019) regional Q can be of similar magnitude for moun-
tainous catchments in North America, although the impact extents can net effects vary among catchments (Rasouli et al., 2019).
233 However, in In our study, the climate- and tree cover-driven impacts on the Q flux can differ substantially on a regional level
as well and can therefore deviate substantially from the global effect, as further discussed in Sect. 3.2 and Sect. 3.3.

3.2 Spatial impact of climate change and future tree cover change on hydrological fluxes

Under climate change, there is generally an increase our study shows general increases in P and ET fluxes over land, however,
245 there are also regions in e.g., South America, southern Europe, Africa and Australia where decreases in both P and ET are
found (Fig. 5a, 5b). We find that P and ET can experience pronounced climate-driven absolute changes in the tropical belt,
246 although it should be noted but note that the magnitude of relative changes can be comparatively smaller in these regions (not
shown). These spatial trends for P and ET over land agree with the multi-model ensemble projections for climate pathways
SSP2-4.5 and SSP5-8.5 (Zhao and Dai, 2021; Li and Li, 2022). When including the potential changes in tree cover, the patterns
247 for tree cover change (Fig. 1) are mirrored in the tree cover-driven changes in ET (Fig. 5f); hence an increase in tree cover
enhances ET and a decrease in tree cover reduces ET. This causal relationship for large-scale tree cover increase and ET is also
248 shown under present climate conditions (Tuinenburg et al., 2022; De Hertog et al., 2023). The change in ET per change in tree
cover is largest in wet regions, which is why we find a large increase in ET over Southeast Asia, tropical Africa, and eastern
249 South America. Since only a small part of the recycled moisture generally precipitates within 100 km of the evaporation source
(Cui et al., 2022; Theeuwen et al., 2023), a change in local ET will mainly affect the downwind P at regional (or larger) spatial
250 scales. HenceTherefore, the tree cover change patterns are less distinguishable for the change in P which is, for example,
251 illustrated with the absence of negative P change over land despite the negative tree cover changes (Fig. 5e). This absence of




Figure 4. Overview of the terrestrial hydrological fluxes precipitation (P), evapotranspiration (ET), and runoff (Q) for the scenarios; present climate, climate change (CC), and climate change with tree cover change (CC + TCC). a) Average fluxes over land for each scenario, averaged over the CMIP6(-Budyko) models, in mm yr^{-1} . b) Average change (Δ) in fluxes for each scenario relative to the present climate in mm yr^{-1} . The corresponding standard deviations display the variability over the CMIP6(-Budyko) models. The flux changes were calculated for each of the 30 CMIP6(-Budyko) model combinations, generating 30 Δ flux values for each land grid cell from which the mean and standard deviation were computed. Note that this figure shows the weighted averages over the total land surface area and these averages are displayed as rounded values (for more details see Table A3).

negative P could be attributed to a compensation effect where small negative local P changes are compensated by increased P originating from other (upwind) areas, as also indicated by Tuinenburg et al. (2022).

260

For the Q flux, we find that the spatial changes in Q due to climate change (Fig. 6a) are roughly consistent with the multi-model trends for SSP5-8.5 by Wang et al. (2022); Li and Li (2022), and the changes in Q typically follow the trends for P. However, in certain regions, such as in parts of Europe and Northern South America, the climate-driven decrease in Q seems to be more pronounced than the limited change in P, as was also shown by Cook et al. (2020) for SSP3-7.0. The difference 265 between the P and Q trends that we found in these areas could be attributed to the elevated climate-driven increase in ET. Analyzing the impacts of an altered tree-cover large-scale global forestation on Q (Fig. 6e) shows an apparent inverse relationship between Q and tree cover change, whereby an increased (decreased) tree cover generally results in a decreased (increased) Q, because of an enhanced (reduced) ET. However, the magnitude of the overall tree-cover-driven impacts on extent to which tree cover change can impact Q may vary locally due to the corresponding indirect effects (altered moisture recycling) following 270 tree-cover change. Note that for both scenarios CC and CC+TCC, the variability between the CMIP6-Budyko models regarding the change in Q is most pronounced in areas that experience larger flux changes. As the high model variability is mainly




Figure 5. Average precipitation change (ΔP ; left) and evapotranspiration change (ΔET ; right) in a future climate due to the impact of climate change (CC; a and b) and the combined impacts of climate change and tree cover change (CC + TCC; c and d). The hydrological changes in these figures are relative to the present climate. Figures e and f only display the impact of tree cover change (TCC; difference between CC and CC+TCC).

concentrated in the wet lower latitudes, the variability for these areas could be (partly) explained by the large spread in P for very wet climate regions as seen in Fig. 3a. Li and Li (2022) also show a higher uncertainty in Q for the regions around the equator for SSP1-2.6, SSP2-4.5, and SSP5-8.5.

275

To analyse the spatial significance of the Q change due to a changing tree cover, we use the Wilcoxon Signed-Rank Test (Wilcoxon, 1945) (i.e. a non-parametric version of the paired samples Student's t-test) for every land grid cell with a p-value threshold of 0.05. For most of the land surface ~~there is a significant change in Q due to tree cover change~~, the significant




Figure 6. Average runoff change (ΔQ : a, c, e, and f) and the corresponding standard deviation (b and d) in a future climate due to the impact of climate change (CC; a and b) and the combined impacts of climate change and tree cover change (CC + TCC; c and d). The hydrological changes in these figures are relative to the present climate. Figures e and f both display the impact of tree cover change (TCC; difference between CC and CC+TCC), whereby figure f illustrates the spatial significance for ΔQ due to TCC. The spatial significance is obtained by applying a Wilcoxon Signed-Rank Test to every land grid cell and the stippling in the figure indicates which grid cells have a significant p-value (<0.05).

changes in Q are associated with changes in potential tree cover (Fig. 6f), and for however in some regions, such as the Sahara 280 desert, the very small absolute change in Q is changes in Q are also marked as significant. We find that, for 16 % of the land surface, By setting a minimal flux change threshold of 5 mm yr^{-1} to exclude such regions, we find that tree cover change (under climate change) significantly increases Q with can significantly increase Q by more than 5 mm yr^{-1} for 16 % of the

land surface.

285 As the impacts of tree cover change can amplify, mitigate, or even reverse the effects of climate change, Fig. 7 shows the spatial distribution of the separate impacts of tree cover change and climate change on Q . The figure includes nine different color combinations, illustrating whether the sign of change due to a changing tree cover aligns with or contradicts the climate change effect on Q . In addition, we use a threshold of 5 mm yr^{-1} in the figure to indicate regions with a low absolute change in Q . For approximately 6 % of the land surface, Q decreases with-by more than 5 mm yr^{-1} due to climate change and increases 290 with-by more than 5 mm yr^{-1} following our large-scale tree cover change scenario (pink color in Fig. 7). Hence, for only a relatively small fraction of the land surface, the tree cover change change in tree cover can mitigate the drying trend under climate change. In contrast, it is more common that the altered tree cover can appears to be more common for tree restoration 295 to mitigate the climate-driven increase in water availability, which as this occurs for approximately 21 % of the land surface (light blue color in Fig. 7). Although the altered tree cover can counteract climate change impacts in certain regions, we also find estimate that part of the land surface (14 %) experiences a decrease in water availability as a result of both climate change and tree cover change effects, such as in Southern Europe and Eastern South America. For the higher latitudes on the Northern Hemisphere (around 60° latitude) there is our study shows a clear climate-driven increase in Q whereas the effects of tree cover change vary. We find that most areas that experience an increasing (decreasing) Q with both climate change and tree 300 cover change are situated in areas where there is a potential decrease (increase) in tree cover.

300

So far, we show that analysing hydrological changes on grid cell level can provide insights about the impact patterns of climate change and tree cover change on the global water fluxes and water availability. However, as impact patterns can differ extensively on a local level, the overall impact on the hydrological fluxes in a catchment remains unclear. By aggregating the results to catchment level (Sect. 3.3), we can also obtain insights about the changes in water availability for e.g., communities, 305 shipping, and agriculture.

3.3 Catchment responses to climate change and future tree cover change

We zoom in on five catchments to take a closer look at the impacts of climate change and tree cover change on integrated hydrological fluxes. The following five catchments are selected; Yukon River basin, Mississippi River basin, Amazon River 310 basin, Danube River basin, and the Murray-Darling River basin. These river basins are chosen since they are situated in different climate zones (Table 3), and because the catchments have a large surface area (Table 3) which is suitable for the coarse spatial resolution of our results analyses. Furthermore, all basins encounter different impacts on Q due to changes in tree cover and climate (Fig. 8).

315 By only looking at Focusing only on the climate change effects, the Yukon River basin experiences can experience an increase in P of approximately 93 mm yr^{-1} (14.5 %) relative to the present climate, which is would be the largest climate-driven change in P amongst among the five river basins (Fig. 8, Table A4). The pronounced increase in yearly P under climate change




Figure 7. Global overview for the changes in runoff (Q) due to climate change (CC) and tree cover change (TCC). The sign of change for Q due to CC and TCC effects can be contradictory or complementary depending on the region. The symbols '+' and '-' in the legend correspond to a positive and negative change in Q, respectively, where these changes exceed the specified absolute threshold of 5 mm yr^{-1} . The '0' in the legend represents small changes in runoff in the absolute range of 0 to 5 mm yr^{-1} . The legend should be read as the following; when there is '+' for TCC and '-' for CC this means that the TCC increases the Q flux whereas CC reduces the Q flux, thus the TCC can mitigate the negative CC effect on Q. The bar graph inside the figure shows the percentages of land surface taken up by each of the CC and TCC combinations.

Table 3. Characteristics of the five chosen catchments in this study. The climate classification according to Köppen (1936).

River basin	Location	Surface area* (km^2)	Climate zone
Yukon	North America; Canada, United States (Alaska)	832819.3	polar climate
Mississippi	North America; Canada, United States	3240616.8	continental climate
Amazon	South America	5912922.8	tropical climate
Danube	Europe	795318.4	temperate climate
Murray-Darling	Australia	1055416.2	arid / temperate climate

* According to the catchment shapefiles retrieved from <https://www.hydrosheds.org/products/hydrobasins>

for this catchment agrees with the projections of Hay and McCabe (2010); Bush and Lemmen (2019), and can lead to a relatively larger increase for Q compared to ET (Hay and McCabe, 2010), which is in line with our results. This larger impact on annual Q can be attributed to a smaller increase in ET under climate change as temperatures remain relatively cold for this region (Hay and McCabe, 2010). We find that altering the tree cover enhances an overall increase in tree cover of 19.7 % could enhance ET (directly) and P (indirectly) within the river basin, thereby amplifying the climate change effect for these two fluxes. The higher tree cover in the basin causes makes that a slightly larger fraction of P to be evaporated can evaporate which subsequently reduces Q and thus partly counteracts the climate-driven increase in Q. For the Wolf Creek basin, located in

325 within the Yukon catchment region, Rasouli et al. (2019) also showed that future vegetation increase can mitigate the climate-driven increase in Q. The In this smaller basin the vegetation effects could even largely offset the climate change impacts in the Wolf Creek basin as the climate and vegetation effects were found to be of similar magnitude (Rasouli et al., 2019). In contrast, we find our study suggests that, for the larger catchment, the climate change impacts on water availability within the Yukon River basin are clearly are dominant over the impacts of a changing tree cover within the Yukon River basin.

330 Similarly to the Yukon River basin, the hydrological fluxes in the Mississippi River basin can increase under climate change, although the extent of increase is much smaller for the latter. However, contrasting with the Yukon basin, the Mississippi catchment shows a more substantial climate-driven increase in ET compared to Q, which could be attributed to the different dominant driving factors for ET in both regions. As the ET flux in the Mississippi River basin is primarily water-limited (Li et al., 2022a), the majority of the additional P resulting from climate change is evaporated and therefore Q encounters can evaporate and therefore we estimate only a small increase of 3 mm yr^{-1} for Q (Fig. 8, Table A4). For this catchment the dominating terrestrial moisture sources for P are situated within or to the southwest of the catchment (Benedict et al., 2020), both of which experience an increase in tree cover under scenario CC+TCC and subsequently an increase in ET. Hence, by accounting for the tree cover change estimated increase in tree cover, the P flux received by the basin is further enhanced. Since more moisture is would be ‘lost’ from the catchment through increased ET than is ‘gained’ through enhanced P following tree 340 cover change restoration, the overall Q shifts may shift to a decrease of 16 mm yr^{-1} (-5%) relative to the present climate. Therefore, when focusing solely on the water availability, the our results indicate that tree cover-driven impacts are generally dominant over the impacts of climate change in the Mississippi River basin when focusing solely on the water availability.

345 For the Amazon River basin, the climate change effects deviate from those in the previous previously discussed catchments as there is can be a decrease in P under pathway SSP3-7.0, in line with Almazroui et al. (2021); Cook et al. (2020). In most of the Amazon region, ET is limited by the available energy (Li et al., 2022a) and therefore the strong increase in temperature for this region under SSP3-7.0 (Almazroui et al., 2021) can enhance ET. Under scenario CC+TCC, the Amazon River basin experiences a widespread (slight) tree cover decrease decrease in potential tree cover due to reduced water availability and extended periods of drought, leading to higher tree mortality (Tavares et al., 2023; Wey et al., 2022). However, the Amazon also contains areas with an increasing tree cover there are also areas with increasing potential tree cover in the Amazon, for example along the southern catchment boundary, that offset the decreases on catchment scale, resulting in an overall tree cover estimated net increase of 4% in tree cover (Fig. 8). This compensation effect on catchment scale also occurs for ET since there is seems to occur for ET as well since we find an overall tree cover-driven increase in ET regardless of the widespread reduction following tree cover change (Fig. 5f). According to Tuinenburg et al. (2020), the Amazon River basin has the highest local moisture recycling of our five selected catchments (with 63 % of the locally evaporated moisture rains raining out within the basin). Hence, the increasing tree cover within (and upwind of) the Amazon basin can enhance ET (directly) and P (indirectly) on catchment level, thereby amplifying and mitigating the climate change impacts on these fluxes, respectively. The changing We estimate that the slight increase in tree cover can also enhance Q with 2 mm yr^{-1} (0.3 %) in the basin. According to Guimberteau et al.




Figure 8. Mean change for the hydrological fluxes precipitation (P), evapotranspiration (ET), and runoff (Q) due to climate change (CC) and climate change with tree cover change (CC + TCC), relative to the present climate. The changes are in mm yr^{-1} . On the left the Yukon (North America; Canada, United States (Alaska)), Mississippi (North America; Canada, United States), and Amazon (South America) River basins are displayed with the corresponding flux changes and on the right the Danube (Europe) and Murray-Darling (Australia) River basins are shown with the corresponding flux changes. The river basins are indicated with a dark blue line. The TCC_{mean} value in the upper left of each bar plot shows the average tree cover change in the catchments and the $Q_{\text{CC+TCC}}$ shows the total Q in the catchment for research scenario CC + TCC. Note that the values presented in the bar graphs are rounded, for more details see Table A4.

(2017), deforestation effects in a future climate could substantially mitigate the climate-driven decreases for Q at a sub-basin scale in the Amazon. **However** **In contrast**, we find that, **on catchment level, the tree cover change only has a limited effect on Q. Therefore, the climate change impacts dominate over the effects only a limited net impact of tree cover change regarding the water availability on Q in the Amazon River basin, suggesting that climate change has a more pronounced influence on water availability compared to tree cover change in this region.**

365 In the Danube River basin the hydrological trends under climate change mirror those shown for the Amazon River basin as both catchments ~~can~~ experience a decreasing P and Q while ET increases (Fig. 8). Interestingly, ~~this-the~~ decreasing P and Q that we find ~~for-in~~ in the Danube basin ~~is-in~~ contrast with the enhanced P and subsequently increased Q under climate pathways RCP2.6 and RCP8.5 ~~as~~ shown by Probst and Mauser (2023). ~~However, it should be noted that in the This difference could be partly attributed to the large uncertainties in the projections of future P in the study of Probst and Mauser (2023) as well as in our study (Table A4), there can be large uncertainties for the projections of future P. Accounting for the changing tree cover, the~~ The Danube River basin ~~experiences~~ has an estimated potential for a tree cover increase of 37.6 % and thus has the largest tree restoration potential ~~over the five~~ across the five selected river basins. In addition, the dominant terrestrial moisture sources of the Danube basin (Central and Eastern Europe as well as the catchment itself (Ciric et al., 2016)) also ~~experience an enhanced tree cover. As a result, have a high forestation potential. Under scenario CC+TCC~~ the P flux received by the catchment ~~increases~~ could therefore increase (indirectly) and consequently ~~shifts~~ shift the climate-driven P decrease to an overall increase. Following the enhanced tree cover, a larger fraction (71.5 %) of the incoming P ~~is evaporated~~ can evaporate and thus a smaller P fraction ends up in the Q flux, which contributes to an amplification of the negative climate change effect on Q. Overall, we find that the impacts of climate change and tree ~~ever change are~~ restoration on water availability can be of similar magnitude for the ~~water availability in the~~ Danube River basin.

380 ~~Last~~ Finally, focusing on the Murray-Darling River basin, we ~~see~~ find that climate change ~~causes a decrease for all~~ can decrease all three hydrological fluxes, ~~contrasting to the other catchments discussed~~. This deviation which is in contrast with the previously discussed catchments and could be attributed to the more arid climate conditions within the basin (Table 3). The Since ET in this catchment is constrained by water availability (Li et al., 2022a), the climate-driven decrease in P resulting from 385 ~~climate change can subsequently lead to reduced~~ can subsequently reduce ET (Fig. 8) as the ET in this catchment is constrained by water availability (Li et al., 2022a). Including Accounting for the (limited) tree cover increase increase in potential tree cover (scenario CC+TCC) can shift ET in the Murray-Darling River basin ~~can shift the ET~~ from a climate-driven decrease to a small increase. However, the incoming P ~~is enhanced~~ can enhance only slightly since the region shows a relatively low moisture recycling whereby only 11 % of the locally evaporated moisture ~~again~~ precipitates in the basin (Tuinenburg et al., 390 2020). In addition, the catchment receives little precipitation recycled from upwind terrestrial regions (Holgate et al., 2020), indicating that upwind changes in tree cover have minimal impact on the P received by this catchment. The We find that the small increase in P due to a locally enhanced tree cover cannot compensate for the additional moisture 'loss' from the basin through enhanced ET, thereby amplifying the impact of climate change on Q. Overall, we find Hence, our results suggest that the climate change impacts ~~are~~ can be dominant over the effects of tree ~~ever change~~ restoration for the water availability in 395 the Murray-Darling River basin.

In conclusion, we show for five large catchments distributed over different climate zones that the impacts of climate change and potential future tree restoration can deviate substantially on a regional scale. These results provide insights into how climate-driven changes in tree cover or tree restoration measures can enhance or offset unwanted climate change effects in each

400 catchment. However, it should be kept in mind that these results correspond with noteworthy uncertainties due to divergent CMIP6 model projections, [uncertainties in our methodology](#), and the use of various Budyko models.

4 Discussion of methodology

405 This study provides a first estimate of the impacts of tree cover change under climate change on global hydrological fluxes over land. To study the combined and separate impacts of tree cover change under climate change, we took an interdisciplinary approach combining state-of-the-art datasets and methods following the methodology from Hoek van Dijke et al. (2022), adapted for future climate scenarios. We would like to stress that other studies also applied the Budyko method and UTrack dataset under different climate and land cover scenarios Kazemi et al. (2019); Tuinenburg et al. (2022); Teuling et al. (2019); Li et al. (2022a). In the discussion below we reflect on the implications of the methodological constraints and their potential impacts on the results.

410 4.1 Implications of the SSP3-7.0 pathway, tree cover change map, and Budyko method

415 The SSP3-7.0 climate pathway was selected, because it enabled us to study strong climate change effects, without stretching our methods too much towards unknown conditions. Both the tree cover map and Budyko models rely on the change for time principle: in a warmer climate we ~~will~~ [would](#) find a similar distribution of trees as in the current climate, however in a different spatial region. For a stronger climate change scenario, more frequent extrapolation outside of current conditions would introduce uncertainties in the future potential tree cover map and make the use of the Budyko framework more uncertain. However, some uncertainties and limitations remain.

420 The SSP3-7.0 pathway describes a resource intensive world with, unlike the other SSP scenarios, a strong reduction in tree cover in the coming decades (Hurtt et al., 2020; Shiogama et al., 2023). The land use changes corresponding to SSP3-7.0 can affect temperature and precipitation extremes, whereby the ~~land-cover~~ [land cover](#) impacts may be more pronounced on a regional level compared to the global level (Hong et al., 2022). In our study, we do not ~~consider the climatic impacts of these land-cover changes~~ [distinguish between the SSP3-7.0 impacts of land use](#) (which can vary ~~between~~ [across](#) CMIP6 models). ~~Hence, this assumption may~~ [and the impacts of climate change, thus attributing the effects only to climate change. This assumption may therefore](#) result in the under- or overestimation of climate change effects on a regional level. ~~For example, the large-scale deforestation in tropical regions presented in SSP3-7.0 could decrease the regional mean and extreme precipitation~~ (Hong et al., 2022). ~~Hence, the precipitation increase in tropical Africa, which we attribute solely to climate change, may actually consist of a stronger increase driven by climate change combined with a decrease driven by deforestation, meaning that for this region the climate change effects could be underestimated.~~

425 To create the future potential tree cover map for SSP3-7.0 (Roebroek, 2023), no ~~feedback~~ [feedbacks](#) between changing tree cover and the climate in SSP3-7.0 ~~was~~ [were](#) included. By using the potential tree cover map for SSP3-7.0 and assuming large-scale tree cover change, we deviate from this climate pathway, which subsequently should alter climate characteristics and therefore the future potential tree cover. Additionally, the potential tree cover map describes the tree cover that could

be established given certain climate conditions. However, three decades would not suffice to reach this level of tree cover everywhere, especially in areas that currently do not contain trees. ~~Furthermore, we~~ We also assume a static tree cover, and do not consider temporal variability in water fluxes that result from e.g., forest disturbance and forest succession stages (Goeking and Tarboton, 2020; Teuling and Hoek van Dijke, 2020). To explicitly model all feedbacks between climate, tree cover, and tree growth, one would need a fully integrated earth system model.

It should be kept in mind that although our study is hypothetical, the realization of the potential tree cover in a future climate can have widespread consequences for biodiversity and agricultural production. For example, tree planting based on the future potential tree cover map could negatively impact biodiversity as the map permits afforestation in grassy biomes, for example in the Mississippi River basin. These regions can naturally support trees but host very different species and therefore afforestation would lead to large losses in biodiversity (Veldman et al., 2015). In addition, the future potential tree cover map shows a high potential for tree cover changes on agricultural (and urban) land (Section 2.2), while actual reforestation in these areas is likely limited to maintain agricultural production (Roebroek, 2023).

The Budyko models used were calibrated under current climate conditions, but the evapotranspiration to precipitation ratio (ET/P) could change in the future. The parameterisation of the models reflects the catchment-integrated effects of differences in interception, plant available water, evaporation, water use efficiency and soil water storage capacity. Some of those characteristics could potentially change the evapotranspiration to precipitation ratio (ET/P) under future climate conditions. For example, the Budyko models do not take into account the CO₂ fertilization effect on reduced surface conductance and increased vegetation greenness (Zhu et al., 2016), which changes the albedo and water use efficiency (Bala et al., 2006). Although these opposing effects are of similar importance on a global scale, the regional effect can be highly positive or negative (Zhang et al., 2022b), and therefore, the Budyko parameters and our calculation of ET, will be non-uniformly affected. Also, the Budyko vegetation parameters are sensitive for, among others, tree species and short vegetation coverage (Chen et al., 2021; Liu et al., 2018; Ding et al., 2022), and the hydrological impacts of forest restoration are therefore highly dependent on the characteristics of forest restoration (e.g. (monoculture) plantations versus natural regrowth of vegetation, and coniferous tree species versus deciduous species).

4.2 Implications of missing the feedback of climate change and tree cover change on recycling

The UTrack dataset used in this study is based on ERA5 reanalysis data and thus represents moisture recycling for current climate and land cover characteristics. However, in this study we apply the UTrack dataset for a future climate and therefore we do not account for the impact of climate change and tree cover change on moisture recycling. ~~So far, There are no moisture tracking datasets available for an Earth with high tree cover and / or climate change, which is why the current UTrack moisture recycling dataset is used. Below, we describe the feedbacks that were omitted, and the effect that it has on our results. This research does not account for the feedbacks of tree cover change on moisture recycling via changes in e.g. the surface albedo, cloud cover, atmospheric carbon dioxide concentrations, surface temperatures, length of moisture transport pathways, and global circulation. These feedbacks can impact the energy balance, e.g. increased tree cover can lower the surface albedo and subsequently enhance the surface temperature, and impact global atmospheric circulation and moisture~~

transport (Portmann et al., 2022). However, these feedbacks are complex and can be contrasting, depending on the location of the land use change. De Hertog et al. (2023) studied the impact of land cover change on atmospheric moisture recycling. They found trends of general wetting and increased local moisture recycling following afforestation in two earth-system models (De Hertog et al., 2023). This indicates that regions with increased evapotranspiration due to an enhanced tree cover will receive more precipitation than currently estimated, thus limiting the decrease in local water availability. The afforestation impacts found by De Hertog et al. (2023) are opposite to the effect of climate change decreasing the local recycling, however those opposite effects are not quantified.

Findell et al. (2019) and Staal et al. (2024a) determined moisture recycling ratios globally in a future climate, both for one climate model, and found that continental moisture recycling ratios decrease with 2–3 % for each degree of warming. Or differently said, the land-to-land water vapor transport decreases. This signal is explained by the fact that evapotranspiration over land is moisture-limited, allowing oceans to have a relatively larger role in the hydrological cycle. Although included in the analyses, Findell et al. (2019) and Staal et al. (2024a) did not discuss in detail the impacts of changes in future circulation on moisture recycling. There are clear indications that climate warming impacts atmospheric circulation, possibly resulting in a poleward shift of the Hadley cells and storm tracks (Shaw, 2019; Francis and Skific, 2015; Vecchi et al., 2006). Future studies applying moisture tracking on future climate simulations ~~will could~~ provide regional insights on the impact of shifting weather patterns on moisture recycling.

Projecting the reported decrease in future recycling on this study would mean that less precipitation due to additional evapotranspiration (due to tree cover change) ~~will would~~ return over land, ~~which will thus decrease thus decreasing~~ the water availability. This indicates that under climate change and tree cover change (scenario CC+TCC), the runoff ~~and water availability~~ may become lower compared to the results in our study. Findell et al. (2019) also shows that under climate change, the contribution of oceanic evaporation to precipitation over land is larger, as was confirmed for the Mississippi River basin by Benedict et al. (2020). It should be noted that the higher evaporation rates over the ocean and the resulting feedback of increased precipitation over land are part of the CMIP6 datasets for a future climate, and these climate change effects are thus included in this study.

Moreover, this research does not account for tree cover change feedbacks on moisture recycling via changes in e.g. the surface albedo, cloud cover, atmospheric carbon dioxide concentrations, surface temperatures, length of moisture transport pathways, and global circulation. These feedbacks are complex and can be contrasting, depending on the location of the land-use change, and the same holds for representing those feedbacks in earth-system models (Portmann et al., 2022; De Hertog et al., 2023). Trends of general wetting and increased local moisture recycling are shown by De Hertog et al. (2023) following afforestation in two earth-system models, indicating that regions with increasing evapotranspiration due to an enhanced tree cover will receive more precipitation than currently estimated, thus limiting the decrease in local water availability. The afforestation impacts found by De Hertog et al. (2023) are opposite to the effect of climate change decreasing the local recycling, however those opposite effects are not quantified.

To conclude, the approach of this study allows to disentangle the impacts of climate change and future tree cover change on hydrological fluxes, but does not include all feedbacks in the earth system in a future climate. To include all those feed-

backs, one would need to run coupled earth system models including tree cover change such as done for current climate by Portmann et al. (2022); De Hertog et al. (2023). Similar model simulations would be needed for future climate conditions and future potential tree cover, ideally for multiple earth system models that actively couple the biosphere and atmosphere. ~~Such, but such~~ simulations are not widely available, to our knowledge a global simulation by King et al. (2024) is available for yet widely available. King et al. (2024) did a global analyses using one earth system model, and Buechel et al. (2024) run a regional simulation by Buechel et al. (2024) for the UK using one convection permitting model. Besides, previous research shows that coupled models do not agree on the implementation of land-atmosphere processes in earth system models, resulting in uncertainty in the sensitivity of vegetation to changing water availability (Denissen et al., 2022; Li et al., 2022b; Baker et al., 2021). Overall, earth-system model studies and data-driven studies, like this one, face different sources of uncertainty and therefore have specific strengths and shortcomings. We believe that our study complements earth-system model studies by contributing to the diversity of methodological approaches presented in scientific literature. Different approaches can improve our understanding of uncertainties and thus enable the most robust scientific progress, which is important when advising society and policymakers. Given the constraints mentioned above, our study provides a first estimate of water availability under climate change and future tree cover change.

515 5 Conclusions

In this study we ~~analysed analyse~~ the impacts of climate change and global tree cover change on ~~the~~ terrestrial precipitation, evapotranspiration, and runoff. To do so, we took an interdisciplinary approach and combined multiple datasets and models. The hydrological fluxes under climate change and future tree cover change were compared ~~to with~~ present climate fluxes to analyse the magnitude and direction of changes in water availability.

520 ~~We find Our findings show~~ that, globally averaged, climate change and large-scale tree ~~ever change restoration~~ can exert similar absolute impacts on runoff. ~~However, these impacts have opposite signs and thus generate a net limited effect relative to the present climate. Following, but with opposite signs. Namely, under~~ climate change, ~~there is an overall larger the overall~~ increase in precipitation ~~than could exceed that of~~ evapotranspiration over land, resulting in enhanced runoff. ~~On the contrary, large scale~~ In contrast, accounting for large-scale tree restoration (15.5%, see Fig. 1) ~~will increase evapotranspiration more than the increase in evapotranspiration over land could exceed that of~~ the recycled precipitation (which partly rains out over water bodies) ~~and thus decreases~~, thus decreasing the runoff. ~~Hence, we find a limited net effect of climate and tree cover changes on the global runoff relative to the present climate.~~

525 ~~While~~ Although the average impact on global runoff ~~is can be~~ limited, the effects of tree cover change and climate change on runoff can be substantial on a regional scale, resulting in enhanced and decreased trends in local runoff up to 100 mm yr^{-1} . For example, the Amazon region ~~will could~~ experience a strong climate-driven decrease in runoff, whereas the related reduction in tree cover could potentially mitigate these impacts. ~~The~~ We also find a drying trend under climate change ~~also applies to~~ for Southern Europe, however, in this region the natural or human-induced forestation would exacerbate rather than mitigate the drying. We ~~find estimate~~ that 14 % of the global land area could experience a pronounced decrease in water availability

due to both climate change and tree cover change. In contrast, the higher latitudes on the Northern Hemisphere (around 60° latitude) show climate-driven ~~enhances increases~~ in runoff and varying effects of tree cover change. Overall, for approximately 535 16 % of the land surface, ~~tree cover changes~~ changes in tree cover could increase water availability (in)directly with more than 5 mm yr⁻¹. Analysing the river basin results in this study shows that the ~~effects of~~ climate change and tree cover change ~~effects~~ can diverge substantially per catchment, whereby four out of five ~~catchments~~ ~~studied catchments could~~ encounter dominant impacts of climate change on the regional water availability.

540 This is the first study to disentangle the effects of climate change and large-scale tree cover change ~~regarding on~~ the future water availability on a global scale and for selected river catchments. We show that climate-driven or human-induced changes in tree cover can mitigate as well as exacerbate climate-induced drying or wetting trends. Ecosystem restoration projects should consider these long-term hydrological effects to limit unintended reductions for local, downstream, and downwind water availability. ~~However, in our study there are many potential sources of uncertainty, due to the omission of important feedbacks.~~ As a next step, we ~~therefore~~ recommend the use of local coupled modelling studies whereby e.g., different afforestation scenarios can be implemented in a regional weather model under future climate conditions, such as ~~(Buechel et al., 2024)~~ 545 ~~Buechel et al. (2024)~~ for the UK. Such studies would enable the analysis of direct local feedbacks and sensitivities of tree cover changes to evaporation and precipitation, based on local atmospheric conditions. Hence, these local studies could address and verify the hydrological responses that we find in our study ~~and thereby guide local and global forest restoration projects. By~~ 550 ~~following a diversity of approaches to study the effects of climate change and tree cover change, we can make the most robust scientific progress, which is needed to inform society and policymakers.~~

555 *Data availability.* The CMIP6 climate simulations are publicly available from <https://esgf-node.llnl.gov/search/cmip6/>. Potential evapotranspiration (PET) data derived from the CMIP6 simulations is provided by Bjarke et al. (2023). The UTrack dataset is publicly available from <https://doi.pangaea.de/10.1594/PANGAEA.912710>. The tree cover dataset for the present climate is provided by Hansen et al. (2013) and the future potential tree cover datasets are provided by Roebroek (2023).

610 *Author contributions.* FE carried out the study and created all figures. The idea of this study was conceived by IB and AHvD. FE, IB, and AHvD wrote the first draft of the paper. All authors interpreted the results, contributed to the discussion and were involved in writing the final paper.

660 *Competing interests.* The authors declare no competing interests.

660 *Acknowledgements.* AHvD acknowledges funding by the German Research Foundation (Emmy Noether grant 391059971). We would like to thank ~~two anonymous~~ ~~the~~ reviewers for their feedback on an earlier version of this manuscript.




Figure A1. Average global change in precipitation (ΔP) in a future climate (2041-2060) due to the indirect impacts of a changing tree cover which affects the moisture recycling. This figure shows the ΔP averaged over the CMIP6-Budyko model combinations and the moisture recycling change was applied twice.

Table A1. Datasets used in this study with corresponding characteristics and the sources from which these datasets can be obtained. The datasets with an asterisk symbol (*) were retrieved for five models selected from phase 6 of the Coupled Model Intercomparison Project (CMIP6). The datasets were used in the following scenarios; present climate, climate change, climate change with tree cover change (TCC), climate change with TCC and moisture recycling change (MRC) applied once, and climate change with TCC and MRC applied twice. Note that scenario 'Climate change' is presented as the 'CC' scenario outside of the appendix, while 'Climate change + TCC + 2x MRC' is presented as the 'CC + TCC' scenario.

Data type	Dataset (climate pathway)	Time period (temporal resolution)	Source	Used for research scenario
CMIP6 climate data	Precipitation* (Historical pathway)	1985–2014 (monthly)	https://esgf-node.llnl.gov/search/cmip6/	Present climate
	Potential evapotranspiration* (Historical pathway)	1985–2014 (monthly)	(Bjarke et al., 2023)	
	Precipitation* (SSP3-7.0)	2035–2064 (monthly)	https://esgf-node.llnl.gov/search/cmip6/	Climate change, Climate change + TCC, Climate change + TCC + 1x MRC, Climate change + TCC + 2x MRC
	Potential evapotranspiration* (SSP3-7.0)	2035–2064 (monthly)	(Bjarke et al., 2023)	
Tree cover data	Tree cover present climate	2000	(Hansen et al., 2013)	Present climate, Climate change
	Potential tree cover future climate* (SSP3-7.0)	2041–2060	(Roebroek, 2023)	Climate change + TCC, Climate change + TCC + 1x MRC, Climate change + TCC + 2x MRC
Moisture tracking data	UTrack moisture trajectories at $1^\circ \times 1^\circ$ resolution	2008–2017 (monthly)	https://doi.pangaea.de/10.1594/PANGAEA.912710	Climate change + TCC + 1x MRC, Climate change + TCC + 2x MRC

* The datasets are retrieved for five selected CMIP6 models; CMCC-ESM2, INM-CM5-0, IPSL-CM6A-LR, MIROC6, and UKESM1-0-LL.

Table A2. The six Budyko models used in this study for which the evapotranspiration (ET) and runoff (Q) fluxes are calculated from the CMIP6 precipitation (P) and potential evapotranspiration (PET) datasets. The ω for these models is calibrated for forest and non-forest vegetation, based on yearly mean streamflow or lysimeter data from various climatic regions. Streamflow Q is calculated as $\frac{Q}{P} = 1 - \frac{ET}{P}$.

Model	Functional form	Calibrated ω	Details
1	$\frac{ET}{P} = \frac{1 + \omega \frac{PET}{P}}{1 + \omega \frac{PET}{P} + \frac{P}{PET}}$	$\omega_g = 0.5$ $\omega_f = 2.0$	Model developed and calibrated by Zhang et al. (2001). Calibrated using 240 global river basins. ω_f is calculated using natural and plantations data, and ω_g is calibrated using grassland and cropland data.
2	$\frac{ET}{P} = \frac{1 + \omega \frac{PET_z}{P}}{1 + \omega \frac{PET_z}{P} + \frac{P}{PET_z}}$	$\omega_g = 0.5$ $\omega_f = 2.0$	As Model 1. PET is also calibrated (PET_z). $PET_{z,grass} = 1100$, $PET_{z,trees} = 1410$
3	$\frac{ET}{P} = 1 + \frac{PET}{P} - (1 + (\frac{PET}{P})^\omega)^{\frac{1}{\omega}}$	$\omega_g = 2.55$ $\omega_f = 2.84$	Model developed by Fu in 1981, calibrated by Zhang et al. (2004). Calibrated based on 200 Australian and 270 worldwide river basins. ω_f and ω_g are calibrated using river basins $\leq 75\%$ forest and grassland cover.
4	$\frac{ET}{P} = 1 + \frac{PET}{P} - (1 + (\frac{PET}{P})^\omega)^{\frac{1}{\omega}}$	$\omega_g = 2.28$ $\omega_f = 2.83$	Model developed by Fu in 1981, calibrated by Zhang et al. (2004). Calibrated with 1420 river basins with forest (ω_f), and grassland and cropland (ω_g) cover.
5	$\frac{ET}{P} = 1 + \frac{aPET}{P} - (1 + (\frac{aPET}{P})^\omega)^{\frac{1}{\omega}}$	$\omega_g = 1.7$ $\omega_f = 3.1$	Model developed by Fu in 1981, calibrated by Teuling et al. (2019). Calibrated based on European Lysimeter data. Teuling et al. (2019) introduced the adjusted potential evaporation ($aPET = 1.6PET$) to account for lysimeter observations above the energy line.
6	a) $\frac{ET}{P} = 1 - \exp(-\frac{\omega PET}{P})$ b) $\frac{ET}{P} = \omega \frac{PET}{P} \tanh((\omega \frac{PET}{P})^{-1})$ c) $\frac{ET}{P} = \frac{1}{(1 + (\omega \frac{PET}{P})^{-2})^{0.5}}$ d) $\frac{ET}{P} = (\frac{PET}{P} (1 - \exp(-\omega \frac{PET}{P})) \tanh(\frac{P}{PET}))^{0.5}$ e) $\frac{ET}{P} = \frac{1 + \omega \frac{PET}{P}}{1 + \omega \frac{PET}{P} + \omega + \frac{P}{PET}}$	$\omega_g = 0.977$ $\omega_f = 1.248$ $\omega_g = 0.767$ $\omega_f = 0.910$ $\omega_g = 0.831$ $\omega_f = 1.025$ $\omega_g = 0.762$ $\omega_f = 1.125$ $\omega_g = 0.682$ $\omega_f = 1.404$	Mean ET calculated from five Budyko equations of: a) Schreiber, b) Ol'DeKop, c) Turc, d) Budyko, and e) Zhang. Oudin et al. (2008) introduced ω in these equations to capture the vegetation effects. The formulas are calibrated using data from 1508 river basins in United States, United Kingdom, Sweden and France.

Table A3. Overview of the global terrestrial hydrological flux values for precipitation (P), evapotranspiration (ET), and runoff (Q) for the scenarios; present climate, climate change, climate change with tree cover change (TCC), climate change with TCC and moisture recycling change (MRC) applied once, and climate change with TCC and MRC applied twice. The mean of the variables represents the mean over the CMIP6(-Budyko) models and the corresponding standard deviations display the variability over the CMIP6(-Budyko) models. Note that scenario 'Climate change' is presented as the 'CC' scenario outside of the appendix, while 'Climate change + TCC + 2x MRC' is presented as the 'CC + TCC' scenario.

Variable	Units	Present climate land total	Change in variable relative to present climate			
			Climate change	Climate change + TCC	Climate change + TCC + 1x MRC	Climate change + TCC + 2x MRC
Tree cover mean	%	23.1	+0.0	+15.5	+15.5	+15.5
P mean \pm std	mm yr ⁻¹	951.5 \pm 198.7	+32.8 \pm 55.3	+32.8 \pm 55.3	+45.8 \pm 55.3	+48.6 \pm 55.3
ET mean \pm std		565.2 \pm 103.1	+22.3 \pm 23.8	+44.6 \pm 33.5	+48.8 \pm 33.8	+49.7 \pm 33.9
Q mean \pm std		386.3 \pm 158.8	+10.5 \pm 39.3	-11.8 \pm 45.0	-3.0 \pm 45.7	-1.1 \pm 45.8

Table A4. Overview of the catchment hydrological flux values for precipitation (P), evapotranspiration (ET), and runoff (Q) for the scenarios; present climate, climate change, climate change with tree cover change (TCC), climate change with TCC and moisture recycling change (MRC) applied once, and climate change with TCC and MRC applied twice. The values of the variables represent the mean over the CMIP6(-Budyko) models and the corresponding standard deviations, the latter of which display the variability over the CMIP6(-Budyko) models. Note that scenario 'Climate change' is presented as the 'CC' scenario outside of the appendix, while 'Climate change + TCC + 2x MRC' is presented as the 'CC + TCC' scenario.

Catchment	Variable	Units	Present climate land total	Change in variable relative to present climate			
				Climate change	Climate change + TCC	Climate change + TCC + 1x MRC	Climate change + TCC + 2x MRC
Amazon	Tree cover	%	82.7	+0.0	+4.0	+4.0	+4.0
	P \pm std	mm yr ⁻¹	2191.4 \pm 457.0	-38.3 \pm 105.6	-38.3 \pm 105.6	-25.0 \pm 105.6	-21.5 \pm 105.6
	ET \pm std		1325.4 \pm 218.4	+15.1 \pm 42.2	+23.8 \pm 49.6	+28.1 \pm 49.4	+29.3 \pm 49.3
	Q \pm std		866.0 \pm 376.0	-53.4 \pm 80.8	-62.1 \pm 86.3	-53.1 \pm 86.4	-50.8 \pm 86.4
Danube	Tree cover	%	30.8	+0.0	+37.6	+37.6	+37.6
	P \pm std	mm yr ⁻¹	778.0 \pm 140.9	-7.4 \pm 39.1	-7.4 \pm 39.1	+15.2 \pm 39.1	+20.4 \pm 39.1
	ET \pm std		503.8 \pm 68.3	+18.8 \pm 21.0	+54.6 \pm 30.0	+64.6 \pm 30.3	+66.9 \pm 30.4
	Q \pm std		274.1 \pm 113.5	-26.2 \pm 29.9	-62.0 \pm 36.1	-49.3 \pm 36.5	-46.5 \pm 36.7
Mississippi	Tree cover	%	21.6	+0.0	+21.5	+21.5	+21.5
	P \pm std	mm yr ⁻¹	929.9 \pm 79.8	+34.7 \pm 40.4	+34.7 \pm 40.4	+47.9 \pm 40.4	+50.4 \pm 40.4
	ET \pm std		603.4 \pm 54.1	+31.8 \pm 16.0	+60.9 \pm 25.5	+65.8 \pm 25.9	+66.7 \pm 26.0
	Q \pm std		326.6 \pm 68.9	+2.9 \pm 30.5	-26.2 \pm 35.7	-17.9 \pm 36.3	-16.4 \pm 36.4
Murray-Darling	Tree cover	%	8.8	+0.0	+7.1	+7.1	+7.1
	P \pm std	mm yr ⁻¹	737.7 \pm 115.1	-23.2 \pm 34.7	-23.2 \pm 34.7	-20.6 \pm 34.7	-20.3 \pm 34.7
	ET \pm std		551.4 \pm 67.1	-5.9 \pm 21.0	+1.6 \pm 23.4	+2.9 \pm 23.3	+3.1 \pm 23.3
	Q \pm std		186.3 \pm 62.3	-17.3 \pm 16.3	-24.8 \pm 18.0	-23.6 \pm 18.0	-23.4 \pm 18.0
Yukon	Tree cover	%	36.0	+0.0	+19.7	+19.7	+19.7
	P \pm std	mm yr ⁻¹	640.8 \pm 104.6	+92.8 \pm 40.5	+92.8 \pm 40.5	+101.4 \pm 40.5	+102.1 \pm 40.5
	ET \pm std		299.5 \pm 105.8	+40.2 \pm 21.6	+54.2 \pm 28.1	+55.8 \pm 29.1	+56.0 \pm 29.1
	Q \pm std		341.3 \pm 141.6	+52.6 \pm 26.7	+38.6 \pm 32.1	+45.5 \pm 33.0	+46.1 \pm 33.1

References

Almazroui, M., Ashfaq, M., Islam, M. N., Rashid, I. U., Kamil, S., Abid, M. A., O'Brien, E., Ismail, M., Reboita, M. S., Sörensson, A. A.,
565 Arias, P. A., Alves, L. M., Tippett, M. K., Saeed, S., Haarsma, R., Doblas-Reyes, F. J., Saeed, F., Kucharski, F., Nadeem, I., Silva-Vidal,
Rivera, J. A., Ehsan, M. A., Martínez-Castro, D., Muñoz, G., Ali, M. A., Coppola, E., and Sylla, M. B.: Assessment of CMIP6
Performance and Projected Temperature and Precipitation Changes Over South America, *Earth Systems and Environment*, 5, 155–183,
https://doi.org/10.1007/s41748-021-00233-6, 2021.

Baker, J., Garcia-Carreras, L., Buermann, W., De Souza, D. C., Marsham, J., Kubota, P., Gloor, M., Coelho, C., and Spracklen, D.: Robust
570 Amazon precipitation projections in climate models that capture realistic land–atmosphere interactions, *Environmental Research Letters*,
16, https://doi.org/10.1088/1748-9326/abfb2e, 2021.

Bala, G., Caldeira, K., Mirin, A., Wickett, M., Delire, C., and Phillips, T.: Biogeophysical effects of CO₂ fertilization on global climate,
Tellus, Series B: Chemical and Physical Meteorology, 58, 620 – 627, https://doi.org/10.1111/j.1600-0889.2006.00210.x, 2006.

Bastin, J.-F., Finegold, Y., Garcia, C., Mollicone, D., Rezende, M., Routh, D., Zohner, C. M., and Crowther, T. W.: The global tree restoration
575 potential, *Science*, 365, 76–79, https://doi.org/10.1126/science.aax0848, 2019.

Benedict, I., van Heerwaarden, C. C., van der Ent, R. J., Weerts, A. H., and Hazeleger, W.: Decline in Terrestrial Moisture Sources of the
Mississippi River Basin in a Future Climate, *Journal of Hydrometeorology*, 21, 299–316, https://doi.org/10.1175/JHM-D-19-0094.1, 2020.

Bjarke, N., Barsugli, J., and Livneh, B.: Ensemble of CMIP6 derived reference and potential evapotranspiration with radiative and advective
580 components, *Scientific Data*, 10, 417, https://doi.org/10.1038/s41597-023-02290-0, 2023.

Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot,
585 P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D'Andrea, F., Davini, P., de Lavergne, C., Denvil,
S., Deshayes, J., Devilliers, M., Ducharne, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L., Falletti, L., Flavoni, S., Foujols, M.-A.,
Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez, Lionel, E., Guilyardi, E., Guimbertea, M., Hauglustaine,
D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M., Khodri, M., Krinner, G., Lebas, N., Levavasseur, G., Lévy, C., Li, L., Lott,
F., Lurton, T., Luysaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meurdesoif, Y., Mignot,
590 J., Musat, I., Ottlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D.,
Thiéblemont, R., Traore, A. K., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N.: Presentation and Evaluation of the
IPSL-CM6A-LR Climate Model, *Journal of Advances in Modeling Earth Systems*, 12, https://doi.org/10.1029/2019MS002010, 2020.

Budyko, M.: Climate and Life, Academic Press, New York, NY, USA, ISBN 9780121394509, 1974.

Buechel, M., Berthou, S., Slater, L., Keat, W., Lewis, H., and Dadson, S.: Hydrometeorological response to afforestation in the UK: findings
595 from a kilometer-scale climate model, *Environmental Research Letters*, 19, https://doi.org/10.1088/1748-9326/ad4bf6, 2024.

Bush, E. and Lemmen, D.: Canada's Changing Climate Report, Government of Canada, Ottawa, ON, https://publications.gc.ca/collections/
collection_2019/eccc/En4-368-2019-eng.pdf, 2019.

Caretta, M., Mukherji, A., Arfanuzzaman, M., Betts, R., Gelfan, A., Hirabayashi, Y., Lissner, T., Liu, J., Gunn, E. L., Morgan, R., Mwanga,
595 S., and Supratid, S.: Water. In: Climate Change 2022: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the
Sixth Assessment Report of the Intergovernmental Panel on Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska,
K. Mintenbeck, A. Alegria, M. Craig, S. Langsdorf, S. Löschke, V. Möller, A. Okem, B. Rama (eds.)]. Cambridge University Press,
Cambridge, UK and New York, NY, USA, p. 551–712, https://doi.org/10.1017/9781009325844.006, 2022.

Chen, Z., Wang, W., Woods, R. A., and Shao, Q.: Hydrological effects of change in vegetation components across global catchments, *Journal of Hydrology*, 595, <https://doi.org/10.1016/j.jhydrol.2020.125775>, 2021.

600 Ceric, D., Stojanovic, M., Drumond, A., Nieto, R., and Gimeno, L.: Tracking the Origin of Moisture over the Danube River Basin Using a Lagrangian Approach, *Atmosphere*, 7, 162, <https://doi.org/10.3390/atmos7120162>, 2016.

Cook, B. I., Mankin, J. S., Marvel, K., Williams, A. P., Smerdon, J. E., and Anchukaitis, K. J.: Twenty-First Century Drought Projections in the CMIP6 Forcing Scenarios, *Earth's Future*, 8, <https://doi.org/10.1029/2019EF001461>, 2020.

605 Cui, J., Lian, X., Huntingford, C., Gimeno, L., Wang, T., Ding, J., He, M., Xu, H., Chen, A., Gentine, P., and Piao, S.: Global water availability boosted by vegetation-driven changes in atmospheric moisture transport, *Nature Geoscience*, 15, 982–988, <https://doi.org/10.1038/s41561-022-01061-7>, 2022.

Davin, E. L. and de Noblet-Ducoudré, N.: Climatic Impact of Global-Scale Deforestation: Radiative versus Nonradiative Processes, *Journal of Climate*, 23, 97–112, <https://doi.org/10.1175/2009JCLI3102.1>, 2010.

610 De Hertog, S. J., Lopez-Fabara, C. E., van der Ent, R., Keune, J., Miralles, D. G., Portmann, R., Schemm, S., Havermann, F., Guo, S., Luo, F., et al.: Effects of idealised land cover and land management changes on the atmospheric water cycle, *EGUphere* [preprint], 2023, 1–36, <https://doi.org/10.5194/egusphere-2023-953>, 2023.

De Petrillo, E., Fahrländer, S., Tuninetti, M., Andersen, L., L., M., Ridolfi, M., and Laio, F.: Reconciling tracked atmospheric water flows to close the global freshwater cycle, *PREPRINT*, <https://doi.org/10.21203/rs.3.rs-4177311/v2>, 2024.

615 Denissen, J. M. C., Teuling, A. J., Pitman, A. J., Koirala, S., Migliavacca, M., Li, W., Reichstein, M., Winkler, A. J., Zhan, C., and Orth, R.: Widespread shift from ecosystem energy to water limitation with climate change, *Nature Climate Change*, 12, 677 – 684, <https://doi.org/10.1038/s41558-022-01403-8>, 2022.

Ding, B., Zhang, Y., Yu, X., Jia, G., Wang, Y., Wang, Y., Zheng, P., and Li, Z.: Effects of forest cover type and ratio changes on runoff and its components, *International Soil and Water Conservation Research*, 10, 445–456, <https://doi.org/10.1016/j.iswcr.2022.01.006>, 2022.

620 Douville, H., Raghavan, K., Renwick, J., Allan, R., Arias, P., Barlow, M., Cerezo-Mota, R., Cherchi, A., Gan, T., Gergis, J., Jiang, D., Khan, A., Mba, W. P., Rosenfeld, D., Tierney, J., and Zolina, O.: Water Cycle Changes. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 1055–1210, <https://doi.org/10.1017/9781009157896.010>, 2021.

625 Duveiller, G., Hooker, J., and Cescatti, A.: The mark of vegetation change on Earth's surface energy balance, *Nature Communications*, 9, 679, <https://doi.org/10.1038/s41467-017-02810-8>, 2018.

Ellison, D., Morris, C. E., Locatelli, B., Sheil, D., Cohen, J., Murdiyarso, D., Gutierrez, V., van Noordwijk, M., Creed, I. F., Pokorny, J., Gaveau, D., Spracklen, D. V., Tobella, A. B., Ilstedt, U., Teuling, A. J., Gebrehiwot, S. G., Sands, D. C., Muys, B., Verbist, B., Springgay, E., Sugandi, Y., and Sullivan, C. A.: Trees, forests and water: Cool insights for a hot world, *Global Environmental Change*, 43, 51–61, <https://doi.org/10.1016/j.gloenvcha.2017.01.002>, 2017.

Fahrländer, S. F., Wang-Erlandsson, L., Pramindita, A., and Jaramillo, F.: Hydroclimatic Vulnerability of Wetlands to Upwind Land Use Changes, *Earth's Future*, 12, e2023EF003837, <https://doi.org/10.1029/2023EF003837>, e2023EF003837 2023EF003837, 2024.

630 Fick, S. E. and Hijmans, R. J.: WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas, *International journal of climatology*, 37, 4302–4315, <https://doi.org/10.1002/joc.5086>, 2017.

635 Findell, K. L., Keys, P. W., Van Der Ent, R. J., Lintner, B. R., Berg, A., and Krasting, J. P.: Rising temperatures increase importance of oceanic evaporation as a source for continental precipitation, *Journal of Climate*, 32, 7713–7726, <https://doi.org/10.1175/JCLI-D-19-0145.1>, 2019.

Francis, J. and Skific, N.: Evidence linking rapid Arctic warming to mid-latitude weather patterns, *Philosophical Transactions of the Royal Society A*, 373, <https://doi.org/10.1098/rsta.2014.0170>, 2015.

640 Goeking, S. A. and Tarboton, D. G.: Forests and Water Yield: A Synthesis of Disturbance Effects on Streamflow and Snowpack in Western Coniferous Forests, *Journal of Forestry*, 118, 172–192, <https://doi.org/10.1093/jofore/fvz069>, 2020.

Griscom, B. W., Adams, J., Ellis, P. W., Houghton, R. A., Lomax, G., Miteva, D. A., Schlesinger, W. H., Shoch, D., Siikamäki, J. V., Smith, P., Woodbury, P., Zganjar, C., Blackman, A., Campari, J., Conant, R. T., Delgado, C., Elias, P., Gopalakrishna, T., Hamsik, M. R., Herrero, M., Kiesecker, J., Landis, E., Laestadius, L., Leavitt, S. M., Minnemeyer, S., Polasky, S., Potapov, P., Putz, F. E., Sanderman, J., Silvius, M., Wollenberg, E., and Fargione, J.: Natural climate solutions, *Proceedings of the National Academy of Sciences of the United States of America*, 114, 11 645–11 650, <https://doi.org/10.1073/pnas.1710465114>, 2017.

645 Guimbarteau, M., Ciais, P., Ducharne, A., Boisier, J. P., Dutra Aguiar, A. P., Biemans, H., De Deurwaerder, H., Galbraith, D., Kruijt, B., Langerwisch, F., Poveda, G., Rammig, A., Rodriguez, D. A., Tejada, G., Thonicke, K., Von Randow, C., Von Randow, R. C. S., Zhang, K., and Verbeeck, H.: Impacts of future deforestation and climate change on the hydrology of the Amazon Basin: a multi-model analysis with a new set of land-cover change scenarios, *Hydrology and Earth System Sciences*, 21, 1455–1475, <https://doi.org/10.5194/hess-21-1455-2017>, 2017.

650 Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., Thau, D., Stehman, S. V., Goetz, S. J., Loveland, T. R., Kommareddy, A., Egorov, A., Chini, L., Justice, C. O., and Townshend, J. R. G.: High-Resolution Global Maps of 21st-Century Forest Cover Change, *Science*, 342, 850–853, <https://doi.org/10.1126/science.1244693>, 2013.

655 Hay, L. E. and McCabe, G. J.: Hydrologic effects of climate change in the Yukon River Basin, *Climatic Change*, 100, 509–523, <https://doi.org/10.1007/s10584-010-9805-x>, 2010.

Herrick, J. E., Abrahamse, T., Abhilash, P. C., Ali, S. H., Alvarez-Torres, P., Barau, A. S., Branquinho, C., Chhatre, A., Chotte, J. L., Cowie, A. L., Davis, K. F., Edrissi, S. A., Fennelly, M. S., Fletcher, S., Flores-Díaz, A. C., Franco, I. B., Ganguli, A. C., Ifejika Speranza, C., Kamar, M. J., Kaudia, A. A., Kimiti, D. W., Luz, A. C., Matos, P., Metternicht, G., Neff, J., Nunes, A., Olaniyi, A. O., Pinho, P., Primmer, E., Quandt, A., Sarkar, P., Scherr, S. J., Singh, A., Sudoi, V., von Maltitz, G. P., Wertz, L., and Zeleke, G.: Land restoration for achieving the sustainable development goals: An international resource panel think piece, United Nations Environment Programme, <https://www.resourcepanel.org/reports/land-restoration-achieving-sustainable-development-goals>, 2019.

660 Hoek van Dijke, A. J., Herold, M., Mallick, K., Benedict, I., Machwitz, M., Schlerf, M., Pranindita, A., Theeuwen, J. J. E., Bastin, J.-F., and Teuling, A. J.: Shifts in regional water availability due to global tree restoration, *Nature Geoscience*, 15, 363–368, <https://doi.org/10.1038/s41561-022-00935-0>, 2022.

665 Holgate, C. M., Evans, J. P., van Dijk, A. I. J. M., Pitman, A. J., and Virgilio, G. D.: Australian Precipitation Recycling and Evaporative Source Regions, *Journal of Climate*, 33, 8721 – 8735, <https://doi.org/10.1175/JCLI-D-19-0926.1>, 2020.

Hong, T., Wu, J., Kang, X., Yuan, M., and Duan, L.: Impacts of Different Land Use Scenarios on Future Global and Regional Climate Extremes, *Atmosphere*, 13, <https://doi.org/10.3390/atmos13060995>, 2022.

670 Hurt, G. C., Chini, L., Sahajpal, R., Frolking, S., Bodirsky, B. L., Calvin, K., Doelman, J. C., Fisk, J., Fujimori, S., Klein Goldewijk, K., Hasegawa, T., Havlik, P., Heinemann, A., Humpenöder, F., Jungclaus, J., Kaplan, J. O., Kennedy, J., Krisztin, T., Lawrence, D., Lawrence, P., Ma, L., Mertz, O., Pongratz, J., Popp, A., Poulter, B., Riahi, K., Shevliakova, E., Stehfest, E., Thornton, P., Tubiello, F. N., van

Vuuren, D. P., and Zhang, X.: Harmonization of global land use change and management for the period 850–2100 (LUH2) for CMIP6, *Geoscientific Model Development*, 13, 5425–5464, <https://doi.org/10.5194/gmd-13-5425-2020>, 2020.

Kazemi, H., Sarukkalige, R., and Badrzadeh, H.: Evaluation of streamflow changes due to climate variation and human activities using the Budyko approach, *Environmental earth sciences*, 78, 713, 2019.

King, J. A., Weber, J., Lawrence, P., Roe, S., Swann, A. L. S., and Val Martin, M.: Global and regional hydrological impacts of global forest expansion, *Biogeosciences*, 21, 3883–3902, <https://doi.org/10.5194/bg-21-3883-2024>, 2024.

Kuma, P., Bender, F. A.-M., and Jönsson, A. R.: Climate Model Code Genealogy and Its Relation to Climate Feedbacks and Sensitivity, *Journal of Advances in Modeling Earth Systems*, 15, <https://doi.org/10.1029/2022MS003588>, 2023.

Köppen, W.: Das geographische System der Klimate, *Handbuch der Klimatologie*, Gebrüder Borntraeger, Berlin, 1936.

Lee, J.-Y., Marotzke, J., Bala, G., Cao, L., Corti, S., Dunne, J., Engelbrecht, F., Fischer, E., Fyfe, J., Jones, C., Maycock, A., Mutemi, J., Ndiaye, O., Panickal, S., and Zhou, T.: Future Global Climate: Scenario-Based Projections and Near-Term Information. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* [Masson-Delmotte, V., P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 553–672, <https://doi.org/10.1017/9781009157896.006>, 2021.

Li, S., Wang, G., Zhu, C., Lu, J., Ullah, W., Hagan, D. F. T., Kattel, G., and Peng, J.: Attribution of global evapotranspiration trends based on the Budyko framework, *Hydrology and Earth System Sciences*, 26, 3691–3707, <https://doi.org/10.5194/hess-26-3691-2022>, 2022a.

Li, W., Migliavacca, M., Forkel, M., Denissen, J. M. C., Reichstein, M., Yang, H., Duveiller, G., Weber, U., and Orth, R.: Widespread increasing vegetation sensitivity to soil moisture, *Nature Communications*, 13, <https://doi.org/10.1038/s41467-022-31667-9>, 2022b.

Li, X. and Li, Z.: Global water availability and its distribution under the Coupled Model Intercomparison Project Phase Six scenarios, *International Journal of Climatology*, 42, 5748–5767, <https://doi.org/10.1002/joc.7559>, 2022.

Liu, C. L. C., Kuchma, O., and Krutovsky, K. V.: Mixed-species versus monocultures in plantation forestry: Development, benefits, ecosystem services and perspectives for the future, *Global Ecology and Conservation*, 15, <https://doi.org/10.1016/j.gecco.2018.e00419>, 2018.

Lovato, T., Peano, D., Butenschön, M., Materia, S., Iovino, D., Scoccimarro, E., Fogli, P. G., Cherchi, A., Bellucci, A., Gualdi, S., Masina, S., and Navarra, A.: CMIP6 Simulations With the CMCC Earth System Model (CMCC-ESM2), *Journal of Advances in Modeling Earth Systems*, 14, <https://doi.org/10.1029/2021MS002814>, 2022.

Meier, R., Schwaab, J., Seneviratne, S., Sprenger, M., Lewis, E., and Davin, E.: Empirical estimate of forestation-induced precipitation changes in Europe, *Nature Geoscience*, 14, 1–6, <https://doi.org/10.1038/s41561-021-00773-6>, 2021.

Milly, P. C. and Dunne, K. A.: Potential evapotranspiration and continental drying, *Nature Climate Change*, 6, 946–949, <https://doi.org/10.1038/nclimate3046>, 2016.

O'Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Kriegler, E., Lamarque, J.-F., Lowe, J., Meehl, G. A., Moss, R., Riahi, K., and Sanderson, B. M.: The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6, *Geoscientific Model Development*, 9, 3461–3482, <https://doi.org/10.5194/gmd-9-3461-2016>, 2016.

Oudin, L., Andréassian, V., Lerat, J., and Michel, C.: Has land cover a significant impact on mean annual streamflow? An international assessment using 1508 catchments, *Journal of Hydrology*, 357, 303–316, <https://doi.org/10.1016/j.jhydrol.2008.05.021>, 2008.

Piao, S., Friedlingstein, P., Ciais, P., de Noblet-Ducoudré, N., Labat, D., and Zaehle, S.: Changes in Climate and Land Use Have a Larger Direct Impact Than Rising CO₂ on Global River Runoff Trends, *Proceedings of the National Academy of Sciences of the United States of America*, 104, 15 242–15 247, <https://doi.org/10.1073/pnas.0707213104>, 2007.

710 Portmann, R., Beyerle, U., Davin, E., Fischer, E. M., De Hertog, S., and Schemm, S.: Global forestation and deforestation affect remote climate via adjusted atmosphere and ocean circulation, *Nature Communications*, 13, <https://doi.org/10.1038/s41467-022-33279-9>, 2022.

Priestley, C. H. B. and Taylor, R. J.: On the assessment of surface heat flux and evaporation using large-scale parameters, *Monthly weather review*, 100, 81–92, [https://doi.org/10.1175/1520-0493\(1972\)100<0081:OTAOSH>2.3.CO;2](https://doi.org/10.1175/1520-0493(1972)100<0081:OTAOSH>2.3.CO;2), 1972.

715 Probst, E. and Mauser, W.: Climate Change Impacts on Water Resources in the Danube River Basin: A Hydrological Modelling Study Using EURO-CORDEX Climate Scenarios, *Water*, 15, <https://doi.org/10.3390/w15010008>, 2023.

Rasouli, K., Pomeroy, J. W., and Whitfield, P. H.: Are the effects of vegetation and soil changes as important as climate change impacts on hydrological processes?, *Hydrology and Earth System Sciences*, 23, 4933–4954, <https://doi.org/10.5194/hess-23-4933-2019>, 2019.

Roebroek, C. T.: Exploring the limits of forest carbon storage for climate change mitigation, Doctoral thesis, ETH Zurich, <https://doi.org/10.3929/ethz-b-000655156>, 2023.

720 Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O'Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., de Mora, L., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahaan, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G., and Zerroukat, M.: UKESM1: Description and Evaluation of the U.K. Earth System Model, *Journal of Advances in Modeling Earth Systems*, 11, 4513–4558, <https://doi.org/10.1029/2019MS001739>, 2019.

725 Shaw, T. A.: Mechanisms of Future Predicted Changes in the Zonal Mean Mid-Latitude Circulation, *Current Climate Change Reports*, 5, 345–357, <https://doi.org/10.1007/s40641-019-00145-8>, 2019.

Shiogama, H., Fujimori, S., Hasegawa, T., Hayashi, M., Hirabayashi, Y., Ogura, T., Iizumi, T., Takahashi, K., and Takemura, T.: Important 730 distinctiveness of SSP3-7.0 for use in impact assessments, *Nature Climate Change*, 13, 1276–1278, <https://doi.org/10.1038/s41558-023-01883-2>, 2023.

Staal, A., Meijer, P., Nyasulu, M. K., Tuinenburg, O. A., and Dekker, S. C.: Global terrestrial moisture recycling in Shared Socioeconomic Pathways, *EGUphere*, 2024, 1–31, 2024a.

735 Staal, A., Theeuwen, J. J. E., Wang-Erlandsson, L., Wunderling, N., and Dekker, S. C.: Targeted rainfall enhancement as an objective of forestation, *Global Change Biology*, 30, <https://doi.org/10.1111/gcb.17096>, 2024b.

Sterling, S., Ducharne, A., and Polcher, J.: The impact of global land-cover change on the terrestrial water cycle, *Nature Climate Change*, 3, 385–390, <https://doi.org/10.1038/nclimate1690>, 2013.

740 Tatebe, H., Ogura, T., Nitta, T., Komuro, Y., Ogochi, K., Takemura, T., Sudo, K., Sekiguchi, M., Abe, M., Saito, F., Chikira, M., Watanabe, S., Mori, M., Hirota, N., Kawatani, Y., Mochizuki, T., Yoshimura, K., Takata, K., O’ishi, R., Yamazaki, D., Suzuki, T., Kurogi, M., Kataoka, T., Watanabe, M., and Kimoto, M.: Description and basic evaluation of simulated mean state, internal variability, and climate sensitivity in MIROC6, *Geoscientific Model Development*, 12, 2727–2765, <https://doi.org/10.5194/gmd-12-2727-2019>, 2019.

Tavares, J., Oliveira, R., Mencuccini, M., and et al.: Basin-wide variation in tree hydraulic safety margins predicts the carbon balance of Amazon forests, *Nature*, 617, 111–117, <https://doi.org/10.1038/s41586-023-05971-3>, 2023.

745 Teuling, A. J. and Hoek van Dijke, A. J.: Forest age and water yield, *Nature*, 578, E16–E18, <https://doi.org/10.1038/s41586-020-1941-5>, 2020.

Teuling, A. J., de Badts, E. A. G., Jansen, F. A., Fuchs, R., Buitink, J., Hoek van Dijke, A. J., and Sterling, S. M.: Climate change, reforestation/afforestation, and urbanization impacts on evapotranspiration and streamflow in Europe, *Hydrology and Earth System Sciences*, 23, 3631–3652, <https://doi.org/10.5194/hess-23-3631-2019>, 2019.

Theeuwen, J. J., Staal, A., Tuinenburg, O. A., Hamelers, B. V., and Dekker, S. C.: Local moisture recycling across the globe, *Hydrology and Earth System Sciences*, 27, 1457–1476, <https://doi.org/10.5194/hess-27-1457-2023>, 2023.

Trenberth, K. E.: Changes in precipitation with climate change, *Climate Research*, 47, 123–138, <https://doi.org/10.3354/cr00953>, 2011.

Tuinenburg, O. A. and Staal, A.: Tracking the global flows of atmospheric moisture and associated uncertainties, *Hydrology and Earth System Sciences*, 24, 2419–2435, <https://doi.org/10.5194/hess-24-2419-2020>, 2020.

Tuinenburg, O. A., Theeuwen, J. J. E., Staal, A., and Environmental, S.: High-resolution global atmospheric moisture connections from evaporation to precipitation, *Earth System Science Data*, 12, 3177–3188, <https://doi.org/10.5194/essd-12-3177-2020>, 2020.

Tuinenburg, O. A., Bosmans, J. H. C., and Staal, A.: The global potential of forest restoration for drought mitigation, *Environmental Research Letters*, 17, 1–8, <https://doi.org/10.1088/1748-9326/ac55b8>, 2022.

Vecchi, G. A., Soden, B. J., Wittenberg, A. T., Held, I. M., Leetmaa, A., and Harrison, M. J.: Weakening of tropical Pacific atmospheric circulation due to anthropogenic forcing, *Nature*, 441, 73–76, <https://doi.org/10.1038/nature04744>, 2006.

Veldman, J. W., Overbeck, G. E., Negreiros, D., Mahy, G., Le Stradic, S., Fernandes, G. W., Durigan, G., Buisson, E., Putz, F. E., and Bond, W. J.: Where Tree Planting and Forest Expansion are Bad for Biodiversity and Ecosystem Services, *BioScience*, 65, 1011–1018, <https://doi.org/10.1093/biosci/biv118>, 2015.

Volodin, E. M., Mortikov, E. V., Kostrykin, S. V., Galin, V. Y., Lykossov, V. N., Gritsun, A. S., Diansky, N. A., Gusev, A. V., and Iakovlev, N. G.: Simulation of the present-day climate with the climate model INMCM5, *Climate Dynamics: Observational, Theoretical and Computational Research on the Climate System*, 49, 3715–3734, <https://doi.org/10.1007/s00382-017-3539-7>, 2017.

Wang, A., Miao, Y., Kong, X., and Wu, H.: Future Changes in Global Runoff and Runoff Coefficient From CMIP6 Multi-Model Simulation Under SSP1-2.6 and SSP5-8.5 Scenarios, *Earth's Future*, 10, <https://doi.org/10.1029/2022EF002910>, 2022.

Wang-Erlansson, L., Fetzer, I., Keys, P. W., Van Der Ent, R. J., Savenije, H. H., and Gordon, L. J.: Remote land use impacts on river flows through atmospheric teleconnections, *Hydrology and Earth System Sciences*, 22, 4311–4328, <https://doi.org/10.5194/hess-22-4311-2018>, 2018.

Wey, H.-W., Pongratz, J., Nabel, J. E. M. S., and Naudts, K.: Effects of Increased Drought in Amazon Forests Under Climate Change: Separating the Roles of Canopy Responses and Soil Moisture, *Journal of Geophysical Research: Biogeosciences*, 127, <https://doi.org/10.1029/2021JG006525>, 2022.

Wilcoxon, F.: Individual Comparisons by Ranking Methods, *Biometrics Bulletin*, 1, 80–83, <https://doi.org/10.2307/3001968>, 1945.

Yang, Y., Roderick, M., Guo, H., Miralles, D., Zhang, L., Fatichi, S., Luo, X., Zhang, Y., McVicar, T., Tu, Z., Keenan, T., Fisher, J., Gan, R., Zhang, X., Piao, S., Zhang, B., and Yang, D.: Evapotranspiration on a greening Earth, *Nature Reviews Earth Environment*, 4, 626–641, <https://doi.org/10.1038/s43017-023-00464-3>, 2023.

Yazdandoost, F., Moradian, S., Izadi, A., and Aghakouchak, A.: Evaluation of CMIP6 precipitation simulations across different climatic zones: Uncertainty and model intercomparison, *Atmospheric Research*, 250, <https://doi.org/10.1016/j.atmosres.2020.105369>, 2021.

Yu, L., Liu, Y., Liu, T., and Yan, F.: Impact of recent vegetation greening on temperature and precipitation over China, *Agricultural and Forest Meteorology*, 295, <https://doi.org/10.1016/j.agrformet.2020.108197>, 2020.

Zhang, B., Tian, L., Yang, Y., and He, X.: Revegetation Does Not Decrease Water Yield in the Loess Plateau of China, *Geophysical Research Letters*, 49, <https://doi.org/10.1029/2022GL098025>, 2022a.

Zhang, L., Dawes, W. R., and Walker, G. R.: Response of mean annual evapotranspiration to vegetation changes at catchment scale, Water
785 Resources Research, 37, 701–708, <https://doi.org/10.1029/2000wr900325>, 2001.

Zhang, L., Hickel, K., Dawes, W. R., Chiew, F. H. S., Western, A. W., and Briggs, P. R.: A rational function approach for estimating mean
annual evapotranspiration, Water Resources Research, 40, <https://doi.org/10.1029/2003wr002710>, 2004.

Zhang, X., Zhang, Y., T. J., Ma, N., and Wang, Y.-P.: CO₂ fertilization is spatially distinct from stomatal conductance reduction in controlling
ecosystem water-use efficiency increase, Environmental Research Letters, <https://doi.org/10.1088/1748-9326/ac6c9c>, 2022b.

790 Zhao, T. and Dai, A.: CMIP6 Model-projected Hydroclimatic and Drought Changes and Their Causes in the Twenty-First Century, Journal
of Climate, 35, 897–921, <https://doi.org/10.1175/JCLI-D-21-0442.1>, 2021.

Zhou, S., Yu, B., Huang, Y., and Wang, G.: The complementary relationship and generation of the Budyko functions, Geophysical Research
Letters, 42, 1781–1790, <https://doi.org/10.1002/2015GL063511>, 2015.

Zhu, Z., Piao, S., Myneni, R. B., Huang, M., Zeng, Z., Canadell, J. G., Ciais, P., Sitch, S., Friedlingstein, P., Arneth, A., Cao, C., Cheng, L.,
795 Kato, E., Koven, C., Li, Y., Lian, X., Liu, Y., Liu, R., Mao, J., Pan, Y., Peng, S., Peuelas, J., Poulter, B., Pugh, T. A., Stocker, B. D., Viovy,
N., Wang, X., Wang, Y., Xiao, Z., Yang, H., Zaehle, S., and Zeng, N.: Greening of the Earth and its drivers, Nature Climate Change, 6,
791 – 795, <https://doi.org/10.1038/nclimate3004>, 2016.