

Silurian syn- and post-collision granitic magmatism in the western section of the North Qinling Orogen: Implications for collisional orogenic processes

Hao Lin¹, Zuochen Li^{1,2}, Xianzhi Pei^{1,2}, Shaowei Zhao^{1,2}, Meng Wang³, Hai Zhou^{1,2}, Feng Gao⁴, Mao Wang¹, Li Qin¹

¹School of Earth Science and Resources, Chang'an University, Xi'an 710054, China

²Xi'an Key Laboratory for Mineralization and Efficient Utilization of Critical Metals, Xi'an 710054, China

³Geophysical Exploration Academy of China Metallurgical Geology Bureau, Baoding 071051, China

⁴School of Geology and Mining Engineering, Xinjiang University, Urumqi 830046, China

Correspondence to: Zuochen Li (lizuoichen@chd.edu.cn)

Abstract. The Liqiao and Xianping plutons can provide crucial evidence for the collision-orogeny process of the Proto-Tethys Ocean in the western section of the North Qinling Orogen. In this study, we present petrological, zircon U-Pb geochronological, geochemical, and zircon Lu-Hf isotopic data for these plutons. Both the Liqiao and Xianping plutons are characterized as high-K, calc-alkaline, metaluminous to weakly peraluminous granites, with ages of 429 Ma and 421 Ma, respectively. The Liqiao pluton was classified as I-type granite, displaying positive $\varepsilon_{\text{Hf}}(t)$ values ranging from -0.1 to +3.4, and high $\text{Mg}^\#$ values from 37.86 to 48.25. We interpret this to indicate that it was generated by the partial melting of juvenile felsic lower crust, with a contribution from mantle-derived material. In contrast, the Xianping pluton exhibited lower $\text{Mg}^\#$ values (20.40 to 35.11) and negative $\varepsilon_{\text{Hf}}(t)$ values (-18.0 to -13.9), consistent with the geochemical characteristics of highly fractionated I-type granite. This suggests that the Xianping pluton formed through the partial melting and **extensive fractional crystallization** of ancient felsic crust. We propose that the Liqiao pluton originated in a syn-collisional setting, while the Xianping pluton formed in a post-collisional environment. Both plutons are products of the collisional orogeny between the Yangtze Block and the North Qinling Orogen, which were associated with the closure of the Wushan-Shangdan Ocean, the northern of the Proto-Tethys Ocean.

1 Introduction

It is generally believed that the Proto-Tethys Ocean was a giant ocean, which was located between the northern Laurasia continent and the southern Gondwana continent (Stampfli and Borel, 2002; Li et al., 2016a; Wu et al., 2020). The Central China Orogenic Belt (CCOB) is one of the most developed areas of the Proto-Tethys tectonic domain, it includes the West Kunlun Orogen, East Kunlun Orogen, Altyn Orogen, North Qaidam Orogen, Qilian Orogen and Qinling-Dabie Orogen, which preserved abundant information on the process of orogeny of various blocks in the North China, South China, Qaidam,

Tarim, and Qiangtang regions from the Paleozoic-Early Mesozoic (Dong et al., 2021, 2022a, 2022b). The Qinling Orogen (~~QO~~) is an important component of the CCOB, which was located between the Yangtze Block (YB) and the North China Block (NCB), it consists of two sutures (the Wushan-Shangdan Suture (~~WSS~~) and the Mianlue Suture (~~MS~~)) and three blocks (the North Qinling Orogen (~~NQO~~), the South Qinling micro-block, and the northern margin of the YB) (Zhang et al., 2001; Wang et al., 2013, 2015; Dong et al., 2021, 2022a, 2022b; Fig. 1a). Previous scholars have conducted extensive studies on the Qinling Orogen (~~QO~~) (Zhang et al., 2001, 2019; Pei et al., 2004, 2009; Wu et al., 2006; Wang et al., 2006, 2007; Xu et al., 2008; Dong et al., 2011a, 2011b, 2011c, 2021), the Early Paleozoic was the main ocean-continent transformation stage of the Qinling Orogen (~~QO~~) (Xia et al., 1998; Dong et al., 2008; Wang et al., 2015; Ren et al., 2019; Dong et al., 2021). As an important part of the Qinling Orogen (~~QO~~), the NQO North Qinling Orogen has complex tectonic evolution and large-scale magmatism. It is a key for exploring the collision of the YB and the NQO North Qinling Orogen. Also, it is an important window for studying the structural evolution of the Proto-Tethys Ocean.

Existing studies indicate that the WSS Wushan-Shangdan Suture was formed during the Caledonian period., which is the result of the collision orogeny between the YB and the NCB (Ren et al., 2019). However, there is still considerable controversy over the duration of the subduction-collision orogeny process of the northern branch of the Proto-Tethys Ocean represented by the Wushan-Shangdan Ocean (~~WSO~~). There are two main views: some scholars believe that the main orogenic stage of the NQO North Qinling Orogen was the late Neoproterozoic to the Mid-Late Triassic, the WSO Wushan-Shangdan Ocean was subducted into the NCB during the Early Paleozoic, by the Middle-Late Devonian, this process transitioned into a continent-continent collision between the YB with the NCB, culminating in the closure along the WSS Wushan-Shangdan Suture (Zhang et al., 2001, 2019; Dong et al., 2011a, 2011c). Another scholar believes that the WSO Wushan-Shangdan Ocean was formed in the NQO North Qinling Orogen during the Late Cambrian, the ocean subduction and the development of ancient island arc occurred in the Ordovician, until to the Silurian-Late Devonian periods, there were continent-continent or continent-arc collision orogeny (Pei et al., 2009; Wang et al., 2009; Ren et al., 2019).

In the western section of the NQO North Qinling Orogen, there are lots of Caledonian basic-intermediate pluton and granite distributed in the Liushuigou-Shuangchangxia and Baihua-Liqiao areas (Pei et al., 2007a; Fig. 1b). These are primarily magmatic rocks related to subduction-collision (Dong et al., 2011a; Li et al., 2018a; Ren et al., 2018, 2021; Yang et al., 2018a). Research on the Paleozoic magmatic rocks of different ages and origins in the western section of the NQO North Qinling Orogen can provide important evidence for the subduction-collision-post-collision orogeny process of the WSO Wushan-Shangdan Ocean in the western section of the NQO North Qinling Orogen in the Early Paleozoic, there by constraining the process and timing of Early Paleozoic orogeny in eastern of the Proto-Tethys Ocean. For a clearer understanding of the tectonic evolution of the western section of NQO North Qinling Orogen, we focus on the Liqiao pluton (LP) and Xianping pluton (XP) in the western section of the NQO North Qinling Orogen (Fig. 1, 2). We conduct studies on petrology, geochemistry, zircon U-Pb geochronology, and zircon Lu-Hf isotope to determine the petrogenesis and age of the rocks, trace the magma source and tectonic background, establish a sequence of tectonic and magmatic evolution events, and

65 explore the relationship between the Caledonian orogeny in the ~~NQO~~North Qinling Orogen and the tectonic evolution of the Proto-Tethys Ocean.




Figure 1: (a) Simplified geological map of the division of tectonic units in Qinling Orogen (Mao et al. 2017; Dong et al. 2022b) (b) Simplified geological map of the conjunction zone between the North Qinling Orogen and the North Qilian Orogen showing the distribution of Early Paleozoic plutons (Pei et al. 2004; Xu et al. 2012).




Figure 1: (a) Simplified geological map of the division of tectonic units in Qinling Orogen (Dong et al. 2022) (b) Simplified geological map of the conjunction zone between the NQO and NQLO showing the distribution of Early Paleozoic plutons (modified from 1:500,000 geological map of Qinling metallogenic belt, and Bottom image from Map World)

75 2 Geological Settingbackground

2.1 Regional geological background

The western section of the NQO is located in the central and western part of China, and is positioned at the conjunction between the NQO and the North Qilian Orogen (NQLO) in the middle section of the CCOB (Zhang et al., 2001; Dong et al., 2022a; Fig. 1a). It is limited in the Tethys tectonic domain, the Paleo-Asian Ocean tectonic domain, and the Pacific tectonic domain, which has developed an active continental margin trench arc basin system in the Early Paleozoic (Pei et al., 2009; Dong et al., 2011a; Zhang et al., 2011; Dong and Santosh, 2016). The ophiolite with Late Cambrian N-MORB type basic volcanic rocks developed in the Guanzizhen area, which extends eastward to the Liqiao area and westward to the Wushan-Yuanyangzhen area, representing the material record of ancient oceanic crust (Pei et al., 2004, 2007a; Dong et al., 2008). In the Late Cambrian-early Ordovician, the WSO (represented by the Guanzizhen-Wushan ophiolite) was subducted from south to north, and formed the island arc fore arc basin represented by the Liziyuan Group metamorphic sedimentary volcanic rocks (Pei et al., 2006; Yang et al., 2018b). With the continued subduction of the WSO, the metavolcanic rocks of the Caotangou Group and the corresponding volcanoclastic and shallowly metamorphosed clastic rocks, which are typical of island arc, were formed in the Middle-Late Ordovician (Pei et al., 2009). Concurrently, intermediate basic igneous complex of Liushuigou and Baihua were formed (Pei et al., 2007b; Gao et al., 2012), as well as subduction type pluton such as Tangzang quartz diorite, Honghuapu tonalite, Yangjiazhuang quartz diorite, and Sanchahe quartz diorite (Chen et al., 2002, 2008; Wang et al., 2006; Ren et al., 2018; Qin et al., 2022). The tectonic background began to transition to the continental-continental or arc-continental collision orogeny after the subduction of the ancient oceanic crust and developed Caledonian collision type pluton such as Dangehuan granite, and entered into an extension environment until the end of the orogeny in the Late Silurian-early Devonian (Wang et al., 2008; Wang, 2013; Ren et al., 2018, 2021; Qin et al., 2022; Xin and Huang, 2023).

2.2 Regional tectonic and stratigraphic features

The western section of the NQO is bounded by the NQLO and NCB to the north and the South Qinling tectonic belt to the south (Fig. 1b). The exposed strata in this area range from the Paleoproterozoic to the Early Paleozoic (Pei et al., 2009). The Precambrian crystalline basement in the western section of the NQO are composed by Paleoproterozoic Qinling Group, which was composed by felsic gneisses, aluminous gneiss and marble calc-silicate (Pei et al., 2009; Diwu et al., 2014). Previous zircon U-Pb dating indicate that it was formed in 2298 to 1867 Ma (Zhang et al., 2001). The high pressure and ultra-high pressure (HP-UHP) metamorphic rock are exposed in the Qinling Group, which are the products of exhumation of continental crust experienced subduction-deep subduction during the Early Paleozoic (Gong et al., 2016; Liu et al., 2020). The Kuanping, Liziyuan and Caotangou Group (Fig. 1b) are consisted by metamorphic volcanic sediment sequence. The previous studies of the Kuanping Group have mainly focused on the central and eastern parts of the NQOB, and have reported Meso-Neoproterozoic ages from 1974 to 813 Ma for the meta-volcanic rocks which with normal type mid-ocean

ridge basalt (N-MORB) affinities, and it was considered to represent oceanic crust (Xue et al., 1996; Zhang et al., 2001; He et al., 2007a; Pei et al., 2009; Diwu et al., 2010; Gao et al., 2015; Dong et al., 2014, 2015, 2021; Zeng et al., 2023). The Paleozoic Liziyuan Group are composed by metamorphic clastic rocks and carbonates sedimentary facies and metamorphic basalt, metamorphic basalt andesites, and metamorphic andesites volcanic facies, while the volcanic displayed typical island-arc or fore arc affinity (Pei et al., 2006). The Ordovician Caotangou Group was composed by metamorphic volcanic sediment, and it was divided into the lower Honghuapu Formation, the middle Zhangjiazhuang Formation, and the upper Longwanggou Formation (Song et al., 1991; Sun and Dong, 1995; Pei et al., 2009; Chen et al., 2019), while the volcanic rocks showed island arc affinity (Yan et al., 2007; Zhu et al., 2008; Xu et al., 2014; Xie et al., 2020). These Paleozoic volcanic sediment sequence in the western section of the NQO indicated that the Early Paleozoic tectonic evolution of the NQO (Xu et al., 2008; Pei et al., 2009; Dong et al., 2015, 2021).

The western section of the North Qinling Orogen is limited in the Tethys tectonic domain, the Paleo-Asian Ocean tectonic domain, and the Pacific tectonic domain (Pei et al., 2009). It is located in the middle section of the CCOB (Zhang et al., 2001; Dong et al., 2022a; Fig. 1a). It is bounded by the North Qilian Orogen and NCB to the north, and the South Qinling tectonic belt to the south (Fig. 1b), which has developed an active continental margin trench-arc-basin system in the Early Paleozoic (Pei et al., 2009; Dong et al., 2011a; Zhang et al., 2011; Dong and Santosh, 2016). The exposed strata in this area range from the Paleoproterozoic to the Early Paleozoic (Pei et al., 2009). The Precambrian crystalline basement is composed by Paleoproterozoic Qinling Group, which was composed by felsic gneisses, aluminous gneiss and marble-calc-silicate (Pei et al., 2009; Diwu et al., 2014). Previous zircon U-Pb dating indicate that it was formed in 2298 to 1867 Ma (Zhang et al., 2001). The high-pressure and ultra-high-pressure (HP-UHP) metamorphic rock are exposed in the Qinling Group, which are the products of exhumation of continental crust experienced subduction-deep subduction during the Early Paleozoic (Gong et al., 2016; Liu et al., 2020). The previous studies of the Kuanping Group have mainly focused on the central and eastern parts of the North Qinling Orogen, and have reported Meso-Neoproterozoic ages from 1974 to 813 Ma for the meta-volcanic rocks which with normal-type mid-ocean ridge basalt (N-MORB) affinities, and it was considered to represent oceanic crust (Xue et al., 1996; Zhang et al., 2001; He et al., 2007a; Pei et al., 2009; Diwu et al., 2010; Gao et al., 2015; Dong et al., 2014, 2015, 2021; Zeng et al., 2023). The ophiolite with Late Cambrian N-MORB type basic volcanic rocks developed in the Guanzizhen area, which extends eastward to the Liqiao area and westward to the Wushan-Yuanyangzhen area, representing the material record of ancient oceanic crust (Pei et al., 2004, 2007a; Dong et al., 2008). In the Late Cambrian-Early Ordovician, the Wushan-Shangdan Ocean (represented by the Guanzizhen-Wushan ophiolite) was subducted from south to north, and formed the island arc-forearc basin represented by the Liziyuan Group metamorphic sedimentary-volcanic rocks (Pei et al., 2006; Yang et al., 2018b). The Paleozoic Liziyuan Group are composed by metamorphic clastic rocks and carbonates sedimentary facies and metamorphic basalt, metamorphic basalt andesites, and metamorphic andesites volcanic facies (Pei et al., 2006). With the continued subduction of the Wushan-Shangdan Ocean, the metavolcanic rocks of the Caotangou Group and the corresponding volcanoclastic and shallowly metamorphosed clastic rocks, which are typical of island arc, were formed in the Middle-Late Ordovician (Yan et al., 2007; Zhu et al., 2008; Pei et al., 2009; Xu et al., 2014;

145 Xie et al., 2020). The Ordovician Caotangou Group was composed by metamorphic volcanic-sediment, and it was divided into the lower Honghuapu Formation, the middle Zhangjiazhuang Formation, and the upper Longwanggou Formation (Song et al., 1991; Sun and Dong, 1995; Pei et al., 2009; Chen et al., 2019). Concurrently, intermediate-basic igneous complex of Liushuigou and Baihua were formed (Pei et al., 2007b; Gao et al., 2012), as well as subduction-type pluton such as Tangzang quartz diorite, Honghuapu tonalite, Yangjiazhuang quartz diorite, and Sanchahe quartz diorite (Chen et al., 2002, 2008; Wang et al., 2006; Ren et al., 2018; Qin et al., 2022). The tectonic background began to transition to the continental-continental or arc-continental collision orogeny after the subduction of the ancient oceanic crust and developed Caledonian collision-type pluton such as Dangchuan granite, and entered into an extension environment until the end of the orogeny in the Late Silurian-Early Devonian (Wang et al., 2008; Wang, 2013; Ren et al., 2018, 2021; Qin et al., 2022; Xin and Huang, 2023).

150




Figure 2: Geological map showing the LP and XP in the western section of North Qinling Orogen (Pei et al. 2004; Xu et al. 2012).




Figure 2: Geological map showing the LP and XP in the western section of NQO

3 Field geology and petrography

3.1 Liqiao pluton

The LP is located in the Xiongshangou-Shaping-Yujiaping-Liqiao area of the western section of the **NQO North Qinling Orogen** (Fig. 1b, Fig. 2), and the pluton is distributed in the NWW direction, which is consistent with the regional tectonic line. In spatial distribution, it intruded into the Paleoproterozoic complex (e.g. Qinling Group), the Early Silurian Baihua intermediate-basic igneous complex, the Ordovician metasedimentary clastic strata (e.g. the Honghuapu Formation of the Caotangou Group). **The LP contacted with Baihua Complex by faults. In addition, there is a large number of diorite xenoliths of Baihua Complex in the LP. It was consisted by** monzogranite and a few syenogranite (Fig. 3a-b). The red monzogranite shows medium- to coarse-grained, some with pseudoporphyritic granite, and massive structure (Fig. 3a). Samples from the monzogranite are composed by K-feldspar (35~45 vol%), plagioclase (35~45 vol%), quartz (20~25 vol%), biotite (~5 vol%). The accessory minerals are mainly apatite, sphene, ilmenite, monazite. It shows distinct myrmekitic structure (Fig. 3c). K-feldspar is euhedral, with development of gridiron twinning, ranging from 1.0 to 5.0 mm in size. Plagioclase is generally euhedral or subhedral, with development of polysynthetic twin and carlsbad—albite compound twin, ranging from 2.0 to 3.0 mm in size. The plagioclase commonly **has crystallization of subhedral columnar**. Quartz is allotriomorphic granular structure with undulatory extinction. Biotite is yellow-brown, block-like structure with significant pleochroism, local

chloritization. To contrast with the monzogranite, the syenogranite (Fig. 3b) has higher K-feldspar (45~55 vol%), lower plagioclase (20~25 vol%), similar quartz (20~30 vol%), biotite (~5 vol%), and a small amount of hornblende (Fig. 3d).




Figure 3: Representative field photographs and photomicrographs of the LP and XP in the western section of North Qinling Orogen. Kfs = K-feldspar, Pl = Plagioclase, Bt = biotite, Qz = Quartz, Hbl = Hornblende, Spn = Sphene.

3.2 Xianping pluton

The XP locates in the east-nouth of the LP (Fig. 2). It also contacted with Baihua Complex by faults. The pluton intruded into the Meso-Neoproterozoic metamorphic volcanic-sediment (e.g. Kuanping Group), the Early Silurian Baihua intermediate-basic igneous complex. It is bounded by the Qinlingdabao granite to the north, which is in contact with a fault.

185 The original structures observed in XP are believed to have been formed during magmatic crystallisation process. During the process of magmatic crystallisation, the magma at the edge of the XP remains thermoplastic due to the influence of its melt. The upwelling magma, along with the pressure of surrounding rocks, results in the formation of local shear, leading to mylonitization of surrounding rocks, the minerals are flattened and elongated. This causes the minerals to become flattened and elongated, and it also leads to the formation of augen structures and banded structures. The pluton is mainly syenogranite, which shows medium-coarse-grained and massive structure (Fig. 3e-f). There was a distinct coarse-medium-grained transition zone can be seen in the pluton (Fig. 3e). The syenogranites are comprising K-feldspar (45~60 vol%), plagioclase (10~20 vol%), quartz (25~30 vol%), biotite (~5 vol%), and minor accessory minerals include zircon, apatite, sphene, monazite, hornblende (Fig. 3g-h). K-feldspar is ranging from 2.0 to 4.0 mm. Plagioclase is ranging from 2.0 to 5.0 mm. Quartz is subhedral and allotriomorphic granular structure with undulatory extinction, it fills in the spaces between other minerals in a granular form. Biotite is gray-brown, block-like structure, form with significant pleochroism, local chloritization.

3.3 Samples

195 Two zircon U-Pb dating samples were collected from the LP, including a monzogranite (TS19001-3) and a syenogranite (TS19007-2). The locations of the two samples are 34°14'04.05"N, 106°25'09.71"E, and 34°13'22.92"N, 106°24'27.68"E in the Liqiao-Yanghe-Wuhe area. Two zircon U-Pb dating samples were collected from the XP, including two syenogranite (TS19014-1 and TS19014-3). The sampling location is in the Xianpinggou-Shangxianping-Nianziping area, which located in 34°16'28.15"N, 106°24'26.88"E, and 34°16'28.21"N, 106°24'26.95"E, respectively. We also collected 15 whole-rock geochemical samples from the LP and 12 samples from the XP. All samples were fresh and free of corrosion, and the sampling locations are shown in Fig. 2.


200



4 Sample collected and analytical methods

4.1 Sample collected

Two zircon U-Pb dating samples were collected from the LP, including a monzogranite (TS19001-3) and a syenogranite (TS19007-2). The locations of the two samples are 34°14'04.05"N, 106°25'09.71"E, and 34°13'22.92"N, 106°24'27.68"E in the Liqiao-Yanghe-Wuhe area. Two zircon U-Pb dating samples were collected from the XP, including two syenogranite (TS19014-1 and TS19014-3). The sampling location is in the Xianpinggou-Shangxianping-Nianziping area, which located in 34°16'28.15"N, 106°24'26.88"E, and 34°16'28.21"N, 106°24'26.95"E, respectively. We also collected 15 whole rock geochemical samples from the LP and 12 samples from the XP. All samples were fresh and free of corrosion, and the sampling locations are shown in Fig. 2.



~~Figure 3: Representative field photographs and photomicrographs of the LP and XP in the western section of NQO. Kfs = K-feldspar, Pl = Plagioclase, Bt = biotite, Qz = Quartz, Hbl = Hornblende, Spn = Sphene.~~

4.24.1 Zircon U-Pb age

215 The samples used for geochronological research were crushed and zircon separated by Xi'an Ruishi Geological Technology Co., Ltd. Zirconium target and cathodoluminescence (CL) imaging were completed by Beijing Zircon Nian Linghang Technology Co., Ltd. Zircon U-Pb isotope testing was conducted on the German Jena PQMS ICP-MS instrument using the laser ablation system NWR193. The 91500 standard sample and GJ-1 control sample were used in this analysis. Data
220 processing was done using the ICPMSDataCal program (Liu et al., 2010). Weighted mean age calculations and concordia diagrams were produced using the Isoplot program (Ludwig, 2012). For a detailed description of the analysis methods and instrument parameters, refer to Li et al., (2009).

4.34.2 Whole-rock major and trace elements

The testing of major, rare earth, and trace elements in whole-rock samples was conducted in the Key Laboratory of Western China's Mineral Resources and Geological Engineering, Ministry of Education, Chang'an University. The analysis of major
225 elements was performed using X-ray fluorescence spectroscopy (XRF) method. The XRF fusion method was carried out in accordance with the national standard GB/T 14506.28-1993, with an analysis precision better than 2% to 3%. The samples were weighed after being heated at 1000 °C for 90 minutes in an oven to determine the loss on ignition (LOI). Rare earth and trace elements were analyzed using a Thermo-X7 Inductively Coupled Plasma Mass Spectrometer (ICP-MS).




Figure 4: CL image of zircons and the U-Pb zircon concordia diagrams from the LP and XP in the western section of NQO.

4.4.4.3 Zircon Lu-Hf isotope

The zircon Lu-Hf isotope dating was carried out at Langfang Fengzeyuan Rock Mine Detection Technology Co., Ltd. The selected Zircon Lu-Hf isotope analysis point was located in the in-situ region of the zircon U-Pb dating site, and the zircon was ablated by the Resolution SE 193 nm excimer laser ablation system of ASI (Applied Spectra Inc.), and the spot beam diameter of laser ablation was generally 38 μm , the energy density was 7~8 J/cm^2 , and the frequency was 10 Hz. The analytical system used is the multi-collector inductively coupled plasma mass spectrometer (Neptune Plus) from the American company Thermo Fisher, with laser ablation material introduced into the Neptune Plus (MC-ICPMS) using high-purity He as carrier gas. The temperature requirement for the detection environment is 18~22°C, with a relative humidity of less than 65%. For detailed experimental principles, analytical techniques, and experimental procedures, refer to Wu et al., (2007) and Geng et al., (2011).




Figure 5: Zircons U-Pb age-Th/U (a) and Chondrite-normalized REE patterns (b) diagram of the LP and XP in the western section of NQO. The chondrite and Primitive Mantle values are from Sun and McDonough (1989).

5 Results of analyses

The data for zircon U-Pb ages, zircon trace elements, whole-rock major and trace elements, and zircon Lu-Hf isotopes are shown in [Supplementary Tables 1, 2, 3 and 4](#), respectively.

5.1 Zircon U-Pb age

The zircon CL image and concordant U-Pb diagrams are shown in [Fig. 4](#). Zircons (TS19001-3) from the Liqiao monzogranite have length of 80 to 180 μm , with respect ratios from 1:1 to 3:1, most grains display developed oscillatory zoning ([Fig. 4a](#)). Except for 4 discordant spots, the remaining 21 concordant spots have $\text{U} = 1020$ to $2806 \mu\text{g/g}$, $\text{Th} = 875$ to $4928 \mu\text{g/g}$, with variable Th/U ratios of 0.48 to 1.98, consistent with the characteristics of typical magmatic zircons ([Fig. 5](#)). They have $^{206}\text{Pb}/^{238}\text{U}$ ages of 417 to 435 Ma ([Fig. 4a](#)) and yielding a weighted mean age of 429.9 ± 2.6 Ma (MSWD = 0.77, $n = 21$).




Figure 4: CL image of zircons and the U-Pb zircon concordia diagrams from the LP and XP in the western section of North Qinling Orogen.

Zircons (TS19007-2) from the Liqiao syenogranite have length of 60 to 170 μm , with respect ratios of 1:1 to 2:1, most grains display developed oscillatory zoning (Fig. 4b). 23 out of 25 spots have U = 341 to 2939 $\mu\text{g/g}$, Th = 290 to 3153 $\mu\text{g/g}$, with Th/U ratios of 0.21 to 1.63, consistent with the characteristics of typical magmatic zircons (Fig. 5). One spot (#21) has $^{206}\text{Pb}/^{238}\text{U}$ age of $489 \pm 6.39\text{Ma}$, suggesting inherited origin. The remaining 22 spots have $^{206}\text{Pb}/^{238}\text{U}$ ages of 421 to 432 Ma (Fig. 4b) and yield a weighted mean age of $429.4 \pm 2.4\text{ Ma}$ (MSWD = 0.22, n = 22).




Figure 6: Geochemical plots for rocks of the LP and XP of the western section of NQO: (a) Q-A-P diagram (Streckeisen 1976); (b) TAS classification diagram (Peccerillo and Taylor 1976); (c) K₂O versus SiO₂ diagram (Middlemost 1994); (d) A/CNK (molar Al₂O₃/(CaO+Na₂O+K₂O)) versus A/NK (molar Al₂O₃/(Na₂O+K₂O)) diagram (Maniar and Piccoli 1989).

270

Zircons (TS19014-1) from the Xianping syenogranite have length of 50 to 130 μm , with respect ratios of 1:1 to 3:1, most grains display developed oscillatory zoning (Fig. 4b). Except for one discordant spots, the remaining 24 spot have high U and Th contents (U = 692 to 9684 $\mu\text{g/g}$, Th = 342 to 3117 $\mu\text{g/g}$, with variable Th/U ratios of 0.22 to 0.70). They are consistent with the characteristics of typical magmatic zircons (Fig. 5). These 24 spots have $^{206}\text{Pb}/^{238}\text{U}$ ages of 409 to 432 Ma (Fig. 4b), yield a weighted mean age of 421.8 ± 2.2 Ma (MSWD=1.00, n=24).




Figure 5: Zircons U-Pb age-Th/U (a) and Chondrite-normalized REE patterns (b) diagram of the LP and XP in the western section of North Qinling Orogen. The chondrite and Primitive Mantle values are from Sun and McDonough (1989).

Zircons (TS19014-3) from the Xianping syenogranite have length of 40 to 170 μm , with respect ratios of 1:1 to 4:1, most grains display developed oscillatory zoning (Fig. 4c). In the 25 analysis spots, 18 spots display concordant U-Pb ages. They have high U and Th contents (U = 1244 to 12052 $\mu\text{g/g}$, Th = 714 to 2351 $\mu\text{g/g}$, with variable Th/U ratios of 0.15 to 0.59), consistent with the characteristics of typical magmatic zircons (Fig. 5). These 18 spots have $^{206}\text{Pb}/^{238}\text{U}$ ages of 408 to 428 Ma (Fig. 4d), yield a weighted mean age of 421.5 ± 2.5 Ma (MSWD = 1.04, n = 18).

5.2 Whole-rock major and trace element geochemistry

5.2.1 Liqiao pluton

Liqiao monzogranite and syenogranite have similar geochemistry characteristics. The LP display high SiO_2 (71.09 to 76.45 wt%) contents, Al_2O_3 = 12.15 to 15.26wt%, as shown in the Q-A-P diagram (Fig. 6a), these sample plotted in the field of monzogranite and syenogranite. They have low $\text{Fe}_2\text{O}_3^{\text{T}}$ (0.90 to 1.77 wt%) contents, MgO (0.30 to 0.55 wt%) contents and $\text{Mg}^{\#}$ (37.86 to 48.25) values, high differentiation index (DI= 89.21 to 93.24) values, high K_2O (4.02 to 5.76 wt%) contents, high total alkali($\text{K}_2\text{O}+\text{Na}_2\text{O}$ = 8.02 to 9.83 wt%) contents and low $\text{Na}_2\text{O}/\text{K}_2\text{O}$ (0.68 to 1.00) ratios. In the TAS diagram (Fig. 6b), they plotted in the subalkaline granite field. They have low rittmann index(σ) values of 1.94 to 3.43. As shown in the SiO_2 versus K_2O diagram (Fig. 6c), most samples display high-K calc-alkaline series. All the samples have moderate A/CNK values of 0.95 to 1.05 (Fig. 6d).

All samples have high total REE contents of 180.35 to 412.32 ppm. They show variable enrichment in LREE with $(\text{La}/\text{Yb})_{\text{N}}$ ratios = 27.46 to 51.51, $(\text{La}/\text{Sm})_{\text{N}}$ ratios = 6.94 to 12.72, $(\text{Gd}/\text{Yb})_{\text{N}}$ ratios = 2.49 to 3.48, and display insignificant Eu anomalies with $\text{Eu}/\text{Eu}^* = 0.54$ to 1.03, $\text{Ce}/\text{Ce}^* = 0.90$ to 0.98 (Fig. 7a). On the primitive mantle normalized trace element spider diagram (Fig. 7b), all these samples show enrichment in Rb, Th, K and Pb, and significantly negative Nb, Ta, Zr, Sr, P, Ce and Ti anomalies, with high Sr (229 to 336 ppm), Ba (688 to 1445-1877 ppm) and Rb (202 to 267 ppm) contents, characterizing crust-derived melts.




Figure 6: Geochemical plots for rocks of the LP and XP of the western section of North Qinling Orogen: (a) Q-A-P diagram (Streckeisen 1976); (b) TAS classification diagram (Middlemost 1994); (c) K_2O versus SiO_2 diagram (Peccerillo and Taylor, 1976); (d) A/CNK (molar $Al_2O_3/[CaO+Na_2O+K_2O]$) versus A/NK (molar $Al_2O_3/[Na_2O+K_2O]$) diagram (Maniar and Piccoli, 1989).

5.2.2 Xianping pluton

Xianping syenogranite have $SiO_2 = 73.92$ to 78.28 wt%, $Al_2O_3 = 11.27$ to 14.14 wt%, $SiO_2 = 71.09$ to 76.45 wt%, $Al_2O_3 = 12.15$ to 15.26 wt%, $Fe_2O_3^T = 0.56$ to 1.91 wt%, $MgO = 0.13$ to 0.26 wt% and $Mg^\# = 20.40$ to 35.11 , differentiation index (DI) = 91.69 to 94.47 . Samples from the XP plot into the syenogranite field on the Q-A-P diagram (Fig. 6a). In the TAS diagram (Fig. 6b), all the samples plotted in the subalkaline granite field. They are rich in K_2O ($K_2O = 3.57$ to 5.34 wt%, $Na_2O/K_2O = 0.71$ to 0.91) and have total alkali ($K_2O + Na_2O$) contents of 6.83 to 9.37 wt%. They have low rittmann index(σ) values of 1.32 to 2.81 . The Xianping syenogranite show calc-alkaline affinity. As shown in the SiO_2 versus K_2O diagram (Fig. 6c), most samples display high-K calc-alkaline series. Xianping syenogranite are metaluminous-weakly peraluminous with A/CNK ratios of 0.98 to 1.04 (Fig. 6d).




Figure 7: Chondrite-normalized REE patterns (a, c) and primitive-mantle-normalized trace element spider diagrams (b, d) of the LP and XP in the western section of NQO. The chondrite and Primitive Mantle values are from Sun and McDonough (1989). Syn-Collision plutons from the western section of NQO after Wang et al. 2008; Ren et al. 2021; Qin et al. 2022; Xin and Huang 2023. Beixinggou pluton from the western section of NQO after Ren et al. 2021.

These samples have REE contents of 81.26 to 205.05 ppm and show fractionated REE patterns expressed by LREE enrichment and HREE depletion with $(La/Yb)_N = 1.46$ to 8.99 , $(La/Sm)_N = 2.43$ to 4.66 , $(Gd/Yb)_N = 0.55$ to 1.39 . They show significant negative Eu anomalies with $Eu/Eu^* = 0.18$ to 0.33 (Fig. 7c). On the spider diagram (Fig. 7d), they are enriched in Rb, K, Pb, Th, and U and depleted in Nb, Ta, Ti, P, Zr, Ce and Ba, with low Sr (39 to 74 ppm), Ba (159 to 271 ppm) and Rb (189 to 477 ppm) contents.




Figure 7: Chondrite-normalized REE patterns (a, c) and primitive mantle normalized trace element spider diagrams (b, d) of the LP and XP in the western section of North Qinling Orogen. The chondrite and Primitive Mantle values are from Sun and McDonough (1989). Syn-Collision plutons from the western section of North Qinling Orogen after Wang et al., 2008; Ren et al., 2021; Qin et al., 2022; Xin and Huang, 2023. Beixinggou pluton from the western section of North Qinling Orogen after Ren et al., 2021.

5.3 Zircon Lu-Hf isotope

Fifteen Lu-Hf spots from the Liqiao monzogranite (TS19001-3, 429 Ma) have positive $\varepsilon_{\text{Hf}}(t)$ values of +0.5 to +3.4, corresponding two-stage model ages of 1197 to 1383 Ma. Sixteen zircon Lu-Hf analysis spots from the Liqiao syenogranite (TS19007-2, 429 Ma) have variable Lu-Hf isotopic compositions, 14 spots display positive $\varepsilon_{\text{Hf}}(t)$ values of +0.6 to +3.3, corresponding two-stage model ages of 1202 to 1374 Ma, only one spot (#1) has negative $\varepsilon_{\text{Hf}}(t)$ value of -0.1 with corresponding two-stage model ages of 1411 Ma. And one inherited origin spot (#21) has positive $\varepsilon_{\text{Hf}}(t)$ values of +0.3, corresponding two-stage model ages of 1269 Ma.

Fifteen Lu-Hf spots from the Xianping syenogranite (TS19014-1, 421 Ma) have negative $\varepsilon_{\text{Hf}}(t)$ values of -18.0 to -13.9, corresponding two-stage model ages of 2276 to 2530 Ma. Fifteen Lu-Hf spots from the Xianping syenogranite (TS19014-3, 421 Ma) have negative $\varepsilon_{\text{Hf}}(t)$ values of -18.5 to -13.6 with corresponding two-stage model ages of 2259 to 2560 Ma.




Figure 8: Pluton-type discrimination diagram of the LP and XP in the western section of NQO: (a) $\text{FeO}^{\text{T}}/\text{MgO}$ versus $(\text{Zr}+\text{Nb}+\text{Ce}+\text{Y})$ (Whalen et al. 1987); (b) $(\text{K}_2\text{O}+\text{Na}_2\text{O})/\text{CaO}$ versus $(\text{Zr}+\text{Nb}+\text{Ce}+\text{Y})$ (Whalen et al. 1987); (c) $100 \times (\text{MgO}+\text{FeO}^{\text{T}}+\text{TiO}_2)/\text{SiO}_2$ versus $(\text{Al}_2\text{O}_3+\text{CaO})/(\text{FeO}^{\text{T}}+\text{Na}_2\text{O}+\text{K}_2\text{O})$ (Sylvester 1989); (d) $10000\text{Ga}/\text{Al}$ versus Zn (Whalen et al. 1987); (e) $10000\text{Ga}/\text{Al}$ versus Zr (Whalen et al. 1987); (f) Zr versus TiO_2 .

345 6 Discussion

6.1 Types of rock genesis

Currently, the most widely used classification scheme for granite types is the MISA classification, with M-type granite derived from mantle magmas being rare, and I-type, S-type, and A-type granite being predominant (Chappell and White,

1974, 2001; Coleman and Peterman, 1975; Whalen et al., 1987; Amri et al., 1996; Wu et al., 2007). Hornblende, cordierite
350 and alkaline dark minerals are considered to be the most important and effective markers for the identification of I-type, S-
type, and A-type granite, respectively (Miller, 1985). Fractionated granite is commonly characterized by the presence of the
following minerals: beryl, tantalum-niobium ores, elbaite, lepidolite, or lithium muscovite (Zhu et al., 2002; Chudik et al.,
2008; Merino et al., 2013; Wu et al., 2017).

Not only do the LP exhibit low concentrations of Zr (143 to 253 ppm), Ce (83.8 to 201.3 ppm), Zr + Nb + Ce + Y
355 (258.2 to 507.68 ppm), and FeO^T/MgO ratios (2.25 to 3.44), but the XP also display low levels of Zr (85.3 to 187.2 ppm), Ce
(26.7 to 90.5 ppm), Zr + Nb + Ce + Y (201.99 to 345.07 ppm), and FeO^T/MgO ratios (3.88 to 8.18). Both are distinct from
typical A-type granites (Zr > 250 ppm, Zr + Nb + Ce + Y > 350 ppm, Ce > 100 ppm, FeO^T/MgO > 16; Whalen et al., 1987).
Furthermore, the zircon saturation temperatures for the LP (T_{Zm} : 770 to 825 °C) and XP (T_{Zm} : 729 to 807 °C) are notably
lower than those of typical A-type granites (> 850 °C, Watson and Harrison, 1983; Eby, 1990, 1992; Siégel et al., 2018). In
360 addition, all compositions of the XP fall in the region of FG, while the LP plotted into the OGT in the discrimination
diagrams (Fig. 8a-b). Consequently, the LP and XP may not be classified as A-type granites. The LP exhibits lower Rb
contents (159 to 267 ppm, mean 201 ppm) compared to highly fractionated granites (Rb > 270 ppm, King et al., 1997). In
contrast, the XP displays high Rb contents (189 to 477 ppm, mean 307 ppm), indicating that the XP is a highly fractionated
granite (Fig. 8c), whereas the LP is not. The 10000Ga/Al ratio is elevated in late-crystallizing granite when I- and S-type
365 granites undergo highly fractional crystallization of plagioclase (Dahlquist et al., 2014). Meanwhile, samples will be
positioned within the A-type granite field in the 10000Ga/Al discriminant diagrams (Whalen et al., 1987). The LP exhibits
low 10000Ga/Al ratios (2.14 to 2.78), while the XP displays high 10000Ga/Al ratios (2.46 to 3.29). It is consistent with the
observation that the samples are categorized as I & S-type granite and A-type granite, respectively (Fig. 8d-e). The LP is
characterized by a low A/CNK ratio (0.95 to 1.05, less than 1.1), low P_2O_5 content (0.07 to 0.14 wt%, less than 0.20 wt%),
370 and high Na_2O content (3.27 to 4.22 wt%, greater than 3.26 wt%). Similarly, the XP exhibits low P_2O_5 contents (less than
0.20 wt%) and high Na_2O contents (greater than 3.26 wt%), and these characteristics differ from those of S-type granites
(Chappell and White, 1974; Chappell, 1999). Moreover, all samples are positioned in the I-type granite field in the Zr versus
TiO₂ discrimination diagrams (Fig. 8f). The presence of potential I-type granite indicators, such as hornblende (Fig. 3d and g,
Miller, 1985), suggests that the LP should be classified as I-type granite and the XP as highly fractionated I-type granite.




Figure 8: Pluton type discrimination diagram of the LP and XP in the western section of North Qinling Orogen: (a) FeO^T/MgO versus $(\text{Zr}+\text{Nb}+\text{Ce}+\text{Y})$ (Whalen et al., 1987); (b) $(\text{K}_2\text{O}+\text{Na}_2\text{O})/\text{CaO}$ versus $(\text{Zr}+\text{Nb}+\text{Ce}+\text{Y})$ (Whalen et al., 1987); (c) $100 \times (\text{MgO}+\text{FeO}^T+\text{TiO}_2)/\text{SiO}_2$ versus $(\text{Al}_2\text{O}_3+\text{CaO})/(\text{FeO}^T+\text{Na}_2\text{O}+\text{K}_2\text{O})$ (Sylvester 1989); (d) $10000\text{Ga}/\text{Al}$ versus Zn (Whalen et al., 1987); (e) $10000\text{Ga}/\text{Al}$ versus Zr (Whalen et al., 1987); (f) Zr versus TiO_2 .

6.2 Petrogenesis of the late Silurian intrusions

It is generally believed that there are three main origins of granite: (1) fractional crystallization of primary basaltic magma (Cawthorn and Brown, 1976; Wyborn et al., 1987; Turner et al., 1992; Mushkin et al., 2003), (2) partial melting and fractional crystallization of crustal material (Barbarin, 1988; Skjerlie and Johnston, 1992; Turpin et al., 1990; Patiño Douce,

1997; Chappell et al., 2012), (3) The mixing of mantle- and crustal-derived materials (Petford and Atherton, 1996; Chappell, 1999; Harris et al., 1999; Yang et al., 2006).




Figure 9: Age versus $\epsilon_{\text{Hf}}(t)$ diagram (a) and Age versus $^{176}\text{Hf}/^{177}\text{Hf}$ diagram (b) of the LP and XP in the western section of NQO.

The LP and XP consist of K-feldspar, plagioclase, quartz, biotite and a few accessory minerals. They differ in the following details: (1) the LP has higher Al_2O_3 (12.15 to 15.26 wt%), K_2O (4.02 to 5.76 wt%) and $\text{Na}_2\text{O}/\text{K}_2\text{O}$ values (0.68 to 1.00) than the XP (11.27 to 14.14 wt%, 3.57 to 5.34 wt% and 0.71 to 0.91, respectively); (2) the LP has higher Sr (276 to 575 ppm) and Ba (688 to 1877 ppm) contents and Sr/Y ratios (15.67 to 53.85), and lower Y contents (8.12 to 19.49 ppm) and Rb/Sr ratios (0.35 to 0.85) than the XP (Sr = 39 to 74 ppm; Ba = 159 to 271 ppm; Sr/Y = 0.97 to 2.62; Y = 20.43 to 42.41 ppm; Rb/Sr = 2.57 to 11.64); (3) the LP has higher La contents (44.80 to 108.25 ppm), lower Yb contents (1.89 to 19.35 ppm), and more fractional REEs ($(\text{La}/\text{Yb})_{\text{N}} = 27.5$ to 51.5; $\text{LREE}/\text{HREE} = 18.9$ to 30.0) than the XP (La = 12.17 to 44.17 ppm; Yb = 2.53 to 5.99 ppm; $(\text{La}/\text{Yb})_{\text{N}} = 1.5$ to 9.0; $\text{LREE}/\text{HREE} = 2.3$ to 8.9), and the Eu anomalies are slightly negative ($\text{Eu}/\text{Eu}^* = 0.54$ to 1.03) and significant negative ($\text{Eu}/\text{Eu}^* = 0.18$ to 0.33), respectively; and (4) LP has positive $\epsilon_{\text{Hf}}(t)$ values of -0.1 to +3.4 and XP has negative $\epsilon_{\text{Hf}}(t)$ values of -18.5 to -13.6. These differences indicate that the LP and XP formed by different process.




Figure 9: Age versus $\epsilon_{\text{Hf}}(t)$ diagram (a) and Age versus $^{176}\text{Hf}/^{177}\text{Hf}$ diagram (b) of the LP and XP in the western section of North Qinling Orogen.




Figure 10: The source region diagrams of the LP and XP in the western section of NQO: (a) molar $\text{K}_2\text{O}/\text{Na}_2\text{O}$ versus molar $\text{CaO}/(\text{MgO}+\text{FeO}^\text{T})$ (Altherr and Siebel 2002); (b) Nb/Y versus Th/Y (Boztuğ et al. 2007); (c) CaO versus Sr (He et al. 2011); (d) $(\text{La}/\text{Yb})_{\text{N}}$ versus Sr/Y (He et al. 2011).

410 6.2.1 Liqiao pluton

On the one hand, LP have low $\text{Na}_2\text{O}/\text{K}_2\text{O}$ (0.68 to 1.00) and middle A/CNK (0.95 to 1.05) ratios, indicating that it was formed by biotite dehydration melting (Patiño Douce and Beard, 1995). In general, a positive value of $\varepsilon_{\text{Hf}}(t)$ can be interpreted as the source rock being from the juvenile crust or depleted mantle, whereas a negative value of $\varepsilon_{\text{Hf}}(t)$ indicates that the source rock is ancient crustal components (Taylor and McLennan, 1985; Wu et al., 2007). The $\varepsilon_{\text{Hf}}(t)$ values of the LP range from -0.1 to +3.4 (Fig. 9), which indicating that the source rock is juvenile crustal components. On the other hand, LP has high Sr (276 to 575) and La (44.8 to 108.3) contents, low Y (8.12 to 19.49) and Yb (0.88 to 1.89) contents indicated that the LP might be partial melting of thickened crust. The negative Eu anomalies ($\text{Eu}^*/\text{Eu} = 0.54$ to 1.03) indicate that plagioclase is the predominant residue phase in the melting process (Patiño Douce and Beard, 1995). The Nb/Ta ratios (mean 14.28) and Zr/Hf ratios (mean 37.42) of the LP were close to the continental crust (Nb/Ta = 13.4, Zr/Hf = 35.7), significantly lower than melted magmas from the mantle source (Nb/Ta = 17.5) (McDonough and Sun, 1995; Rudnick, 1995; Weyer et al., 2003). The Rb/Sr ratios of the crust source magmas is generally greater than 0.5, while the Ti/Zr ratios were usually less than 20 and the Cr contents of the primitive basaltic magma were 500 to 600 $\mu\text{g/g}$ (Wilson, 1989). LP has low Cr content (2.89 to 7.27 $< 500 \mu\text{g/g}$), Ti/Zr ratios (6.12 to 7.85 < 20) and high Rb/Sr ratios (0.35 to 0.85 > 0.5), which indicated the LP were fundamentally different from mantle source magma and similar to crustal source magma. Also, In the LP, no mafic intrusions have been found, and it has high SiO_2 contents (71.09 to 76.45 wt%), which indicated that the fractional crystallization of mantle source magma model is impossible (Defant and Drummond, 1990).




Figure 10: The source region diagrams of the LP and XP in the western section of North Qinling Orogen: (a) molar K_2O/Na_2O versus molar $CaO/(MgO+FeO^I)$ (Altherr and Siebel 2002); (b) Nb/Y versus Th/Y (Boztuğ et al. 2007); (c) CaO versus Sr (He et al. 2011); (d) $(La/Yb)_N$ versus Sr/Y (He et al. 2011).




Figure 11: The evolution and fractional diagrams of the LP and XP in the western section of NQO: (a) La versus La/Sm (b) Zr versus Zr/Sm (c) Sr versus Ba (Rudnick 1995) (d) La versus (La/Yb)_N (Rudnick 1995). Pl: Plagioclase; Kfs: K-feldspar; Amp: Amphibole; Bt: biotite; Zr: Zircon; Spn: Spene; Ap: Apatite; Mon: Monazite; Allan: Allanite.

As shown in the K_2O/Na_2O versus $CaO/(MgO+FeO^T)$ diagram (Fig. 10a), the samples plotted in the field of metaandesites and metabasalts. In the Nb/Y versus Th/Y diagram (Fig. 10b), they plotted in middle-lower field. In the CaO versus Sr and $(La/Yb)_N$ versus Sr/Y diagrams (Fig. 10c-d), they plotted in crust field ~~and there is a clear tendency to evolve towards to the thickened crust~~. The positive correlations of La and La/Sm, Zr and Zr/Sm suggest that the LP exhibits partial melting trend (Fig. 11a-b). The $Mg^\#$ value can serve as a basis for determining whether mantle-derived material has been added to crustal source magma (Frost, 2001). Melt produced by crustal melting shows $Mg^\# < 40$, and $Mg^\#$ does not change with the degree of partial melting, and it only displays $Mg^\# > 40$ when mantle-derived material is added (Frey et al., 1978; Rapp and Watson, 1995; Hawkes and Kemp, 2006). The $Mg^\#$ values of LP range from 37.86 to 48.25 (mean 41.81 > 40), indicating the involvement of mantle-derived material in the magmatic source area. In summary, we propose that the LP was formed by partial melting of juvenile felsic crust with the involvement of mantle-derived material.




Figure 11: The evolution and fractional diagrams of the LP and XP in the western section of North Qinling Orogen: (a) La versus La/Sm (b) Zr versus Zr/Sm (c) Sr versus Ba (Rudnick 1995) (d) La versus $(La/Yb)_N$ (Rudnick 1995). Pl: Plagioclase; Kfs: K-feldspar; Amp: Amphibole; Bt: biotite; Zr: Zircon; Spn-Sphene; Ap: Apatite; Mon: Monazite; Allan-Allanite.

6.2.2 Xianping pluton

There are two main sources of highly fractionated I-type granite: (1) Formation of highly fractionated I-type granite as a result of partial melting of crustal material due to magmatic underplating of the fractionated mantle source (Wu et al., 2003; Wang et al., 2014). (2) The fractionated basic magma from mantle source underplated the lower crust and mixed with crust-derived felsic magma. They formed shallow-source hybrid magma chambers and undergo fractional crystallization to form highly fractionated I-type granite in later stage (Qiu et al., 2008). The high SiO_2 contents (73.92 to 78.28 wt%) of the XP suggests that it was not derived from mantle-derived magmas (Defant and Drummond, 1990). The XP showed zircon Hf isotopic compositions ($\epsilon_{Hf}(t) = -18.5$ to -13.6), with corresponding two-stage model ages of 2259 to 2560 Ma (Fig. 9), these features clearly differ from the LP, suggesting a matured continental source region (Taylor and McLennan, 1985; Wu et al., 2007). It has higher K_2O contents (3.57 to 5.34 wt%), low Nb/Ta ratios (mean 11.61) and Zr/Hf ratios (mean 27.93), similar to crustal source magmas (Nb/Ta=13.4, Zr/Hf=35.7), different from mantle source magmas (Nb/Ta= 17.5) (McDonough and Sun, 1995; Rudnick, 1995; Weyer et al., 2003). It also has low Cr content (1.42 to 5.36 $< 500 \mu g/g$), Ti/Zr ratio (2.63 to 5.73 < 20) and high Rb/Sr ratio (2.57 to 11.64 > 0.5), suggesting that the XP was crustal source origin (Wilson, 1989). As shown in the K_2O/Na_2O versus $CaO/(MgO+FeO^T)$ diagram (Fig. 10a), the samples plotted in the field of

metaandesites and metabasalts. In the Nb/Y versus Th/Y diagram (Fig. 10b), they plotted in middle-upper crust field. In the CaO versus Sr and (La/Yb)_N versus Sr/Y diagrams (Fig. 10c-d), they plotted in crust field. The Mg[#] values of XP range from 20.40 to 35.11 (<40), indicating the absence of mantle-derived material in the magmatic source area (Frey et al., 1978; Rapp and Watson, 1995; Hawkes and Kemp, 2006). Characterization of partial melting in La versus La/Sm and Zr versus Zr/Sm diagrams (Fig. 11a-b). The XP was characterized by a high differentiation index (91.69-94.47, average 93.22) as well as significant depleting in HFSEs (i.e., Nb, Ta, P, Ti), suggesting that its source magma underwent significant fractional crystallization. Negative anomalies of Eu (Eu*/Eu = 0.18 to 0.33) and depleted in Sr and Ba in the samples were also indicative the fractional crystallization of plagioclase and K-feldspar (Rudnick, 1995; Patiño Douce and Beard, 1995). The Sr versus Ba diagram (Fig. 11c) reflects fractional crystallization of K-feldspar and plagioclase with a predominance of K-feldspar, which is consistent with the occurrence of K-feldspar and plagioclase phenocrysts in the samples. The depleted in P and Ti, Nb, and Ta were due to fractional crystallization of apatite and Ti-rich minerals, respectively, and as shown in the La versus (La/Yb)_N diagram (Fig. 11d), also reflected fractional crystallization of apatite and sphene. In summary, XP may be the formed by partial melting and highly fractional crystallization of the ancient felsic crust.




Figure 12: The tectonic setting diagrams of the LP and XP in the western section of NQO: (a) Zr versus $(\text{Nb}/\text{Zr})_N$ (Batchelor and Bowden 1985); (b) SiO_2 versus $\text{FeO}^{\text{F}}/(\text{FeO}^{\text{F}}+\text{MgO})$ (Maniar and Piccoli 1989); (c) Y versus Nb (Pearce et al. 1984); (d) Yb versus Ta (Pearce et al. 1984); (e) $(\text{Y}+\text{Nb})$ versus Rb (Pearce et al. 1984); (f) R1 versus R2 (Batchelor and Bowden 1985). Syn Collision plutons from the western section of NQO after Wang et al. 2008; Ren et al. 2021; Qin et al. 2022; Xin and Huang 2023. Post-Collision pluton from the western section of NQO after Ren et al. 2021.

6.3 Tectonic background

The orogeny is typically used to describe a range of geological processes, including the magmatism of island and continental arcs, oceanic subduction and closure, continents collision and subduction, slab rollback during the syn-collision, delamination and orogenic collapse during the post-collision (Song et al., 2015; Xu et al., 2021; Zhang and Hou 2015; Zheng et al., 2015, 2022; Zhu et al., 2022). The initiation of continental collision occurs when two continents converge and the

oceanic crust between them is fully subducted, and the demise of this oceanic crust can be inferred from the ages of the high-pressure (HP) metamorphic rocks, arc volcanic rocks, and youngest ophiolites (Song et al., 2015). The syn-collision is characterized by a continental collision and subduction process. Despite the ocean having closed, subduction is not terminated, this is because it requires a significant amount of time for one continent to be subducted beneath another and achieve ultra-high-pressure (UHP) metamorphic conditions. During the process of continental collision and subduction, both the previously subducted oceanic crust and the continental crust undergo decompression melting as they are exhumed and the crust thickens, and accompanied by partial melting of crustal material to produce tonalites and peraluminous granites, without input of mantle materials (Chen et al., 2013; Liu et al., 2014; Song et al., 2014). Additionally, crustal melts derived from the deep subducted continental crust during exhumation could metasomatize the overlying lithospheric mantle, leading to the formation of mafic magmas (Zhao et al., 2012). The post-collisional is used to describe magmatism that occurs after the major collisional event, which typically marks the end of orogeny. It occurred in the relaxation phase following the main orogeny, characterized by the large-scale horizontal movement of plate boundaries in the early phase and, in the late phase, by the delamination and collapse of orogeny, extension of the lithosphere, and crust-mantle interactions, such as the post-collisional magmas in the North Qaidam UHP metamorphic belt (Wu et al., 2014; Wang et al., 2014).




Figure 12: The tectonic setting diagrams of the LP and XP in the western section of North Qinling Orogen: (a) Zr versus $(\text{Nb}/\text{Zr})_N$ (Batchelor and Bowden, 1985); (b) SiO_2 versus $\text{FeO}^T/(\text{FeO}^T+\text{MgO})$ (Maniar and Piccoli, 1989); (c) Y versus Nb (Pearce et al. 1984); (d) Yb versus Ta (Pearce et al., 1984); (e) $(\text{Y}+\text{Nb})$ versus Rb (Pearce et al., 1984); (f) R1 versus R2 (Batchelor and Bowden, 1985). Syn-Collision plutons from the western section of North Qinling Orogen after Wang et al., 2008; Ren et al., 2021; Qin et al., 2022; Xin and Huang, 2023. Post-Collision pluton from the western section of North Qinling Orogen after Ren et al., 2021.

According to the variations in chronology and geochemical features in the western section of the North Qinling Orogen, Ren et al. (2021) proposed that three magmatic pulses indicate a tectonic switch from slab rollback (ca. 441–434 Ma), slab break-off (ca. 430–423 Ma), to post-collision (ca. 415–409 Ma) from 441 to 409 Ma. In contrast, Qin et al. (2022) identified three main Paleozoic magmatic pulses from ~500 to 410 Ma, which correspond to oceanic subduction (ca. 500 to 450 Ma), slab rollback & break-off (ca. 440 to 430 Ma), and extension in a post-collision setting (ca. 420 to 410 Ma), respectively.


The zircon U-Pb ages of the LP was 429 Ma, while the XP was 421 Ma, both representing the period of collision orogeny (Pei et al., 2009; Wang, 2013; Liu, 2013; Xie et al., 2020). They are consistent with the second and third magma pulses, respectively, that have been studied by previous authors (Ren et al., 2021; Qin et al., 2022). The Qinling Complex in the western section of the North Qinling Orogen underwent late Silurian (433–424 Ma) granulite-facies metamorphism with a clockwise *P-T* path, and followed by 411–402 Ma amphibolite-facies or retrograde overprinting metamorphism, suggesting that the granulite-facies rocks may have been formed by the continent-continent collisional orogeny in the Paleozoic (Guo et al., 2022; Mao et al., 2017, 2018). In contrast, the ages of granulite-facies metamorphism in the Qinling Complex, located in the eastern part of the North Qinling Orogen is mainly concentrated between 430 and 450 Ma (Liu et al., 2013; Xiang et al., 2014; Zhang et al., 2009, 2011). The zircon U-Pb age of the LP is 429 Ma, coinciding with the timeframe of magmatism during the second pulse, specifically the slab break-off (Ren et al., 2021; Qin et al., 2022). This aligns with the granulite-facies metamorphism period at 433–424 Ma (Mao et al., 2017, 2018), suggesting that the LP was likely formed during the intermediate stage of slab break-off in the syn-collisional process. Pay attention, the collision of YB and North Qinling Orogen along the Wushan-Shangdan Ocean margin might not terminate subduction (Li et al., 2018b, 2018c; Allen et al., 2023). The zircon U-Pb age of the XP is 421 Ma, coinciding with the post-collisional orogenic process within the error range (420–409 Ma; Ren et al., 2021; Qin et al., 2022). It is also close to the timeframe limit of the amphibolite-facies or retrograde overprinting metamorphism from 411 to 402 Ma, suggesting that the XP was formed during the early stage of the post-collisional phase. Therefore, both of the LP and XP represent periods of collisional orogeny (Pei et al., 2009; Wang, 2013; Liu, 2013; Xie et al., 2020).

In the Zr versus (Nb/Zr)_N diagram (Fig. 12a), LP and XP plotted into the region of the collision related environment, indicating that they were formed in the collision-related setting, and the XP should be later than LP. In the SiO₂ versus FeO^T/(FeO^T+MgO) diagram (Fig. 12b), most samples from the LP plotted into the IAG-CAG-CCG region, while the XP fell into the POG region. In the Y versus Nb and Yb versus Ta diagrams (Fig. 12c-d), the LP mostly fell into the syn-collision field. In the (Y+Nb) versus Rb diagram (Fig. 12e), the XP mainly fell into the post-collision field. The LP shows syn-collision-related features, while the XP shows late-orogenic-related features (Fig. 12f). Combining with previous research on syn-collision and post-collision granite in the western section of the ~~NQO~~North Qinling Orogen (Wang et al., 2008; Ren et al., 2021; Qin et al., 2022; Xin and Huang, 2023), chondrite-normalized REE patterns and primitive mantle normalized trace element spider diagrams of the LP (Fig. 7a-b) was similar to the syn-collision granite, while the XP (Fig. 7c-d) was similar to the post-collisional granite, and there was a very clear linear evolutionary relationship from the LP to the XP, namely the trend from syn-collision to post-collision (Fig. 12). It can be inferred that the LP formed in syn-collision stage (Fig. 13a), as well as the XP formed in a post-collision stage (Fig. 13b). Both were products of the collision-orogeny between the northern margin of the YB and the ~~NQO~~North Qinling Orogen. The genesis of the LP and XP suggests that the heat required for partial melting of the lower-middle crust is likely to have originated from upwelling of the asthenosphere and the underplating of basaltic magma within an extension setting (Turnbull et al., 2016), and ridge subduction, slab rollback, slab break-off, and lower crustal delamination are the main deep geodynamic mechanisms that trigger extensional activity. Slab

windows can provide enhanced heat flux during ridge subduction and form a large number of high-temperature granites (e.g., charnockite and alkali-feldspar granite; Geng et al., 2009). However, the North Qinling Orogen does not expose any high-temperature granites from the Ordovician and Silurian periods. The K-rich High-Ba-Sr intrusions (439-438 Ma) formed in a slab subduction setting, and the Na-rich intrusion (429-425 Ma) was associated with slab break-off (Ren et al., 2018). Moreover, the LP has a similar age to the Na-rich high-Ba-Sr intrusion.

Therefore, we suggest that the slab break-off following continental collision is the most likely cause of the magmatism that formed the LP (Fig. 13a). Although exhumation of a thickened lower crust generally occurs in post-collisional setting (Song et al., 2015), it may therefore be the main mechanism for the formation of the magmatism of the XP (Fig. 13b).

(b) ~421 Ma



(a) ~429 Ma




Figure 13: Schematics geodynamic and petrogenetic model illustrating the evolution of the western section of the North Qinling Orogen at ca. 429 Ma and 421Ma, and the generation of the LP (a) and XP (b) during Late Silurian.

6.4 Tectonic significance

The rock types and geochemical characteristics of the intermediate-acid intrusive are closely related to the composition of the crustal and mantle (Chen et al., 2014). Furthermore, different tectonic setting can form the same type of granitic rocks, whereas the same tectonic environment can also give rise to different types of granitic rocks (Li et al., 2013; Wang et al.,

2014). Therefore, any discussion on the magmatism and formation mechanism of orogeny must be based on the tectonic framework. The Proto-Tethys Ocean was a giant ocean developed during Neoproterozoic to Early Paleozoic, which was located between the northern Laurasia and the southern Gondwana, and it generated by the breakup of the Rodinia Supercontinent (Mattern and Schneider, 2000; Stampfli and Borel, 2002; Raumer and Stampfli, 2008; Li et al., 2016b, 2018b; Yang et al., 2018; Wu et al., 2020). The northern boundary of the Qinling-Qilian Orogen in the CCOB is regarded as the northern boundary of the Early Paleozoic Proto-Tethys Ocean closure (Li et al., 2016b, 2017). The NQQNorth Qinling Orogen preserves abundant Early Paleozoic records, revealing the tectonic evolution of the northern Proto-Tethys Ocean on the periphery of the northern margin of eastern Gondwana (Dong et al., 2021; Mark et al., 2023). The WSSWushan-Shangdan Suture is considered to represent an oceanic suture zone formed by closure of the WSOWushan-Shangdan Ocean, which formed the northern part of the Proto-Tethys Ocean along the northern margin of the eastern Gondwana during the Early Paleozoic (Zhang et al., 2001; Pei et al., 2009; Chen et al., 2024). The WSSWushan-Shangdan Suture is the most important tectonic zone separating the NCB from the YB (Zhang et al., 2001; Yang et al., 2002; Xu et al., 2006; Pei et al., 2009). The mafic volcanic rocks within Guanzizhen and Wushan ophiolite exhibited features of N-MORB and E-MORB, respectively, and yield zircon U-Pb ages of 534 to 489 Ma, which were both represented the remnants of the WSOWushan-Shangdan Ocean lithosphere (Hou et al., 2006; Li et al., 2007; Li, 2008; Pei et al., 2004, 2007a, 2009; Dong et al., 2008, 2011a). The northward subduction of the WSOWushan-Shangdan Ocean in the western section of the NQQNorth Qinling Orogen in the Ediacaran-Ordovician resulted in the collision of the YB and the NQQNorth Qinling Orogen in the southern part of the NCB, which reflected the relationship between the Caledonian orogeny in the NQQNorth Qinling Orogen and the tectonic evolution of the Proto-Tethys Ocean (Pei et al., 2005, 2009). In combination with the results of previous research, the tectonic evolution of the western section of the NQQNorth Qinling Orogen can be divided into three stages as follows.





Figure 13: Schematics geodynamic and petrogenetic model illustrating the evolution of the western section of the NQO at ca. 429 Ma and 421 Ma, and the generation of the LP (a) and XP (b) during Late Silurian. MORB—mid-ocean ridge basalt.

Stage 1 from 472 Ma to 438 Ma. Northward subduction of the WSO Wushan-Shangdan Ocean.

With the northward subduction of the NQO North Qinling Orogen, the Liziyuan Group represented by the metamorphic-volcanic-sedimentary rock system of the island arc-forearc basin during 472 to 451 Ma, and the Caotangou Group represented by the island-arc-type volcano-sedimentary rock system was formed in the fore-arc basin, which located between the island arc and the subduction zone at the southern margin of the NQO North Qinling Orogen (Pei et al., 2006; He et al., 2007; Yan et al., 2007; Yang et al., 2018a). The Liushuigou gabbro, located north of the Guanzizhen ophiolite, has a zircon U-Pb age of 471 Ma, which was formed in island-arc tectonic setting (Pei et al., 2005, 2009; Yang et al., 2006). Additionally,

the Baihua intermediate-basic igneous complex (449.7 Ma) in the western section of the ~~NQ~~North Qinling Orogen, the Hualingou meta-gabbro (440 Ma) and the Yuanyangzhen meta-gabbroic amphibolite (456 Ma) in the Wushan area were formed in an island-arc tectonic setting (Pei et al., 2007b; Li, 2008). The intermediate-basic volcanic (456.4 Ma, e.g. the Zhangjiazhuang Formation) and the acidic tuffs (457.4 Ma, e.g. the Longwanggou Formation) showed arc magmatism characterize, which recorded the subduction of ~~WSO~~Wushan-Shangdan Ocean in the Early Paleozoic (Wang et al., 2007; Chen et al., 2019; Zhu et al., 2008; Xu et al., 2014). There were Honghuapu tonalite (450.5 Ma), Tangzang quartz diorite (454.7 Ma), and the Sanchahe quartz diorite (459 Ma) in the Fengxian area (Wang et al., 2006, Chen et al., 2008, Qing et al., 2022), as well as the Yangjiazhuang and Honghuapu K-rich, high-Ba-Sr granitoids (439-438 Ma, Ren et al., 2018), were formed in subduction-related setting. In this stage, the slab roll-back of the ~~WSO~~Wushan-Shangdan Ocean initiated asthenosphere decompression melting in the mantle wedge and generated corresponding magmatic activities. Subsequently, asthenosphere convection induced partial melting of enriched mantle that had interacted with slab-derived fluids and produced basaltic magma that evolved into the Early Ordovician-Early Silurian gabbro, diorite and quartz diorite by fractional crystallization. Slab roll-back and its consequences strongly increased the geothermal gradient of the lower crust, which formed the early Silurian granites. These arc-related and subduction-related magmatic rocks define the age for the northward subduction of the ~~WSO~~Wushan-Shangdan Ocean from 472 to 438 Ma.

Stage 2 from 438-~~Ma~~ to 423Ma. Initial continental collision of the YB with the ~~NQ~~North Qinling Orogen.

Previous studies have shown that slab break-off usually occurs very soon after continental collision (Duretz et al., 2011; van de Zedde and Wortel, 2001). Continuous subduction ultimately gave rise to the consumption of the ~~WSO~~Wushan-Shangdan Ocean oceanic crust and final collision between the YB and the ~~NQ~~North Qinling Orogen in the Early Silurian which resulted in the significantly thickened the crust, and there was no typical sedimentary state exposed in the area during this stage, but the intrusion activities of collisional-related magmas was violent extensive development of Caledonian pluton (Pei et al., 2009). Due to buoyancy differences, slab break-off might have taken place, followed by subsequent asthenosphere upwelling, as well as the asthenosphere mantle-derived magma supplied both heat and mantle material, which underplated the lower part of the accretionary prism and triggered partial melting of juvenile crustal material accompanied by crust-mantle interaction (Davies and von Blanckenburg, 1995). The Dangchuan granite was formed by the partial melting of the thickened lower crust, with a U-Pb age from 438 to 432 Ma (Wang et al., 2008; Qing et al., 2022; Xin and Huang, 2022), as well as the Xiongshangou syn-collision granite (named LP of this paper) formed at 438 Ma (Wang, 2013), and the Jiguanya syn-collision granite at 435 Ma (Xu et al., 2018). The Tangzang Na-rich high-Ba-Sr granite, Zhangjiazhuang two-mica granite and muscovite granite formed during the slab break-off from 430 to 423 Ma (Ren et al., 2018, 2021). The formation of the LP (429 Ma) studied in this paper was attributed to partial melting of lower crust dominated by accreted ~~WSO~~Wushan-Shangdan Ocean oceanic crust and sediments, accompanied by local magma mixing with the mantle-derived magma. The presence of these intrusive rocks suggested that the ~~WSO~~Wushan-Shangdan Ocean closed before 438 Ma, and that the northern margin of the YB and the ~~NQ~~North Qinling Orogen were underwent continent-continent collision from 438 to 423 Ma. During the Late Silurian (433 to 424 Ma), the western section of the

~~NQO~~North Qinling Orogen experienced granulite-facies metamorphism and anatexis (Mao et al., 2017), which may have occurred in a thickened lower crustal setting during the early stage of collision orogeny (Yu et al., 2013; Peng et al., 2021). This reflects the existence of an Early Paleozoic crustal thickening in the western section of the ~~NQO~~North Qinling Orogen, which comprehensively indicates that the northern margin of the YB and the ~~NQO~~North Qinling Orogen were in a docking and collision stage during the period of 438 to 423 Ma.

Stage 3 from 421-~~Ma~~ to 409 Ma. Post-collision extension phase.

Given that slab break-off occurred before 423 Ma in the western section of the ~~NQO~~North Qinling Orogen (Ren et al., 2021), delamination of thickened crust might have caused extension during the Late Silurian-Early Devonian. Continued continental compression from 438 to 423 Ma would cause shortening and thickening of the continental crust and underlying lithospheric mantle to produce a high-density lithospheric root (eclogitic root) that protruded into the asthenosphere (Houseman and Molnar, 1997). The delamination was characterized by the sinking of eclogitic root and subsequent upwelling of extensive hot asthenosphere, accompanied by crustal extension (Deng et al., 2015; Ma et al., 2015). Huoyanshan granite was formed by high-temperature melting of the heterogeneous continental crust under extensional environment, with a zircon U-Pb age of 415 Ma (Qing et al., 2022). Leijayuan quartz diorite (415 Ma), Beixinggou black mica granite (412 Ma) and Yanwan dolomite granite (409 to 414 Ma) were formed in a post-collision tectonic setting (Wang et al., 2009; Ren et al., 2021). They record the tectonic evolution of the post-collision phase of the ~~NQO~~North Qinling Orogen. The XP (421 Ma) in this paper was product of the post-collision orogeny stage of extension between the northern margin of the YB and the ~~NQO~~North Qinling Orogen. The upwelling of hot asthenosphere underplated the ancient crust and supplied heat, not only caused the crust to thin and decompress, but also resulted in widespread partial melting of the ancient intermediate-basic lower crust. Later, under the control of mineral separation crystallization processes such as K-feldspar, plagioclase, and apatite, it underwent highly fractional crystallization to form the Xianping highly fractionated I-type granite. Reflecting the information on the differentiated evolution of crustal materials during the Late Caledonian post-collision in the western section of the ~~NQO~~North Qinling Orogen, it provides further evidence to reveal the subduction-collision timeframe of the ~~WSO~~Wushan-Shangdan Ocean in the ~~NQO~~North Qinling Orogen during the Early Paleozoic.

When the continental crust was disconnected from the subducting oceanic crust, the continental crust stopped subducting, and UHP metamorphic rocks began to emerge due to the buoyancy (Davies and van Blanckenburg, 1995). The continental subduction channel model posits that deep-subducted crustal rocks may be detached at mantle depth and subsequently rollback to crustal levels or even to the surface, driven by buoyancy and corner flow (Zheng et al., 2013). This process can be observed in the *P-T* paths of UHP metamorphic rocks (Li et al., 2014). The *P-T* paths of high-grade metamorphic rocks are typically closely related to the tectonic background in which they occur. Clockwise *P-T* paths are predominantly associated with tectonic contexts involving subduction, arc-continent collision, and continent-continent collision at convergent margins (Brown, 1995; England and Thompson, 1984; Thompson and England, 1984). In contrast, anticlockwise *P-T* paths are usually associated with island-arc, back-arc, continental rift or mantle plume tectonic environment (Sandiford and Powell, 1986; Bohlen, 1987). Furthermore, the geochronological sequence of Caledonian

igneous and metamorphic rocks in the North Qinling Orogen offers significant insights into the temporal framework of the YB with NCB. The Qinling Complex in the eastern part of the North Qinling Orogen has been identified as containing HP-UHP metamorphic rocks between 507 Ma and 485 Ma (Chen et al., 2004, 2011; Liu et al., 2013; Su et al., 2004; Wang et al., 2013; Yang et al., 2002; Zhang et al., 2011), medium- to low-pressure granulite-facies metamorphic rocks between 430 Ma and 450 Ma (Liu et al., 2013; Xiang et al., 2014; Zhai et al., 1998; Zhang et al., 2009, 2011), and the amphibolite-facies metamorphic rocks overprinting between 420 Ma and 426 Ma (Liu et al., 2013; Zhang et al., 2009, 2011). Significant anatexis associated with granulite-amphibolite facies metamorphism is also identified (Liu et al., 2014; Xiang et al., 2014). However, there are still controversies about the genesis of the medium- to low-pressure granulite-facies metamorphic rocks as follows: (1) related to the northward subduction of the Shangdan Ocean (Wang et al., 2011, 2013; Wu et al., 2013; Xiang et al., 2014); (2) related to the collisional orogeny of the Early Paleozoic (Liu et al., 2016; Zhai et al., 1998; Zhang et al., 2011, 2013). The garnet amphibolite probably experienced eclogite-facies metamorphism at 497 Ma and two stages retrograde overprint at 452 Ma and 423 Ma in the western section of the North Qinling Orogen, which are basically consistent with the continental-type HP-UHP rocks of the eastern section of North Qinling Orogen, suggesting that both western and eastern section of North Qinling Orogen experienced similar continental deep subduction and exhumation processes (Tang et al., 2022). Combined with regional geological data, this suggests that the granulite-facies metamorphic rocks in the western section of the North Qinling Orogen may be the result of the Early Paleozoic continent-continent collision orogenic process.

In the global context, the Proto-Tethys Ocean was located between the North American Laurentia-Tarim-North China and Gondwana continent, formed by the breakup of the Rodinia supercontinent (Stampfli and Borel, 2002; Wu et al., 2020; Dong et al., 2022). The Rodinia supercontinent underwent a breakup during the Neoproterozoic, and the Laurentia and Baltic continent separated to form the Iapetus Oceans in the Late Proterozoic, which face west towards the continent of West Gondwana in the Southern Hemisphere (van Staal et al., 2012). In the Late Ordovician, the Avalonia Terrane drifted northwards to form the Rheic Ocean in the active continental margin of the western Gondwana, which continued to expand and shrink the Iapetus Ocean, while the collision of the Avalonia Terrane with the Baltic continent caused the disappearance of the Tornquist Ocean and the formation of the Thor Suture (Torsvik and Rehnstrom, 2003). Subsequently, the collision of the Avalonia Terrane-Baltic continent with the Laurentia continent formed the Appalachian-Caledonian orogen in southern North America-Britain-Norway, which resulted in the disappearance of the Iapetus Ocean between the Laurentia and Baltic continents, it means the closure of the Proto-Tethys Ocean (McKerrow et al., 2000; Dewey et al., 2015; Torsvik, 2019). In the northern part of the British Caledonian Belt, between the Baltic and Greenland continent, the famous Norwegian HP-UHP metamorphic rocks were formed due to the subduction of the Baltic under the Laurentia continent, which indicated that the Proto-Tethys Ocean had disappeared at ca. 435 Ma (Wu et al., 2020).

The CCOB is one of the most developed locations in the Proto-Tethys tectonic domain. Through long-term research on this area, it has been found that the ages of ophiolites in the North Altyn-North Qilian-Kuanping Suture, South Altyn-North

Qaidam-Shandan Suture, and Kudi-Kunzhong Suture were mainly concentrated between 520–490 Ma (Xiao et al., 2003; Zhang et al., 2004, 2015; Song et al., 2019). The Kangxiwar-A'nyemaqen-Mianlue Suture zone developed a large number of Early Paleozoic ophiolites of 530–480 Ma and Late Paleozoic ophiolites. A small amount of Proto-Tethys ophiolites also developed in the Longmuco-Shuanghu-Changning-Menglian belt (Pan et al., 2020; Wang et al., 2021). These ophiolites reflected the formation of the Proto-Tethys Ocean in China, while the formation of the European Iapetus Ocean is basically consistent with or slightly earlier than this (Xu et al., 2010a, 2010b; Song et al., 2013, 2019; Wu et al., 2020; Li et al., 2022). The NQO North Qinling Orogen marks the northernmost boundary of the Proto-Tethys Ocean in the eastern part of the CCOB (Li et al., 2016a), as well as the WSO Wushan-Shangdan Ocean is located in the northern part of the Proto-Tethys Ocean (Zhang et al., 2001). The closure of the WSO Wushan-Shangdan Ocean resulted in the collision of the northern margin of the YB with the NQO North Qinling Orogen in the Silurian, which ultimately merged with the eastern part of the Gondwana continent, and were closely related to the evolution of the Proto-Tethys Ocean (Zhao et al., 2018; Li et al., 2018b).

7 Conclusions

(1) LP and XP both were high-K calc-alkaline metaluminous-weakly peraluminous granite series, which showed high K_2O , low Na_2O , $Fe_2O_3^T$, and exhibited distinct enrichment in LFSE (e.g. Rb, K, and Pb), and depleted in HFSE (e.g. Nb, Ta, Ti, Zr, and Ce). The LP has higher REE contents (180.35 to 412.32 $\mu g/g$), more fractionated REEs ($(La/Yb)_N = 27.5$ to 51.5; $LREE/HREE = 18.9$ to 30.0) than the XP ($REE = 81.26$ to 205.05 $\mu g/g$, $(La/Yb)_N = 1.5$ to 9.0, $LREE/HREE = 2.3$ to 8.9), and the Eu anomalies are slightly negative ($Eu/Eu^* = 0.54$ to 1.03) and significant negative ($Eu/Eu^* = 0.18$ to 0.33), respectively;

(2) The zircon U-Pb age of the LP and XP were 429 Ma and 421 Ma, respectively. The LP was I-type granite with positive $\varepsilon_{Hf}(t)$ values of -0.1 to +3.4, which was formed by partial melting of the intermediate-basic juvenile lower crust, accompanied by magma mixing with the mantle material. The formation of the Xianping highly fractionated I-type granite ($\varepsilon_{Hf}(t) = -18.5$ to -13.6) was attributed to partial melting of mature lower crust with highly fractional crystallization in extensional setting.

(3) From the Liqiao I-type granite to Xianping highly fractionated I-type granite in the western section of the NQO North Qinling Orogen, reflecting the transformation of the thickening of the crust and the magma mixing of crust-mantle materials in syn-collision stage between the YB and NQO North Qinling Orogen to post-collision stage with fractional crystallization of crustal material. They were closely related to the evolution of the WSO Wushan-Shangdan Ocean, northern of the Proto-Tethys Ocean.

Data availability

725 The data presented in the supplement.

Author contribution

Hao Lin: Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Writing – original draft. Zuochen Li: Conceptualization, Formal analysis, Investigation, Methodology, Visualization, Funding acquisition. Xianzhi Pei: Investigation, Methodology, Funding acquisition. Meng Wang: Investigation. Shaowei Zhao: Formal analysis, Methodology. 730 Hai Zhou: Formal analysis, Methodology. Feng Gao: Investigation. Mao Wang: Writing – review & editing, Methodology. Li Qin: Investigation, Methodology.

Acknowledgements

We thanks are extended to the chief editor and the two anonymous reviewers for their constructive reviews which have greatly improved our manuscript. We wish to acknowledge Yifeng Wang and others for their help during the fieldwork and 735 sample preparations. We also extremely thankful to Dr. Yinchuan Wang and Dr. Xiao Wang provided valuable comments on the language of this paper.

Financial support

This work was supported by the National Natural Science Foundation of China [Grant number 41872235, 42172236, 41872234]; Fundamental Research Funds for the Central Universities [Grant number 300102270202, 300103183081, 740 300104282717]; and Youth Innovation Team of Shaanxi Universities.

Competing interest

The authors declare that they have no conflict of interest.

Disclaimer

Publisher’s note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published 745 maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

References

- Miller, B. B. and Carter, C.: The test article, J. Sci. Res., 12, 135–147, doi:10.1234/56789, 2015.
- Smith, A. A., Carter, C., and Miller, B. B.: More test articles, J. Adv. Res., 35, 13–28, doi:10.2345/67890, 2014.
- 750 Allen, M.B., Song, S.G., Wang, C., Zeng, R.Y., and Wen, T.: An oblique subduction model for closure of the Proto-Tethys and Palaeo-Tethys oceans and creation of the Central China Orogenic Belt, Earth-Sci. Rev., 240, 104385, <https://doi.org/10.1016/j.earscirev.2023.104385>, 2023.
- Altherr, R. and Siebel, W.: I-type plutonism in a continental back-arc setting: Miocene granitoids and monzonites from the central Aegean Sea, Greece, Contrib. Mineral. Petrol., 143, 397-415, <https://doi.org/10.1007/s00410-002-0352-y>, 2002.
- 755 Amri, I., Benoit, M., and Ceuleneer, G.: Tectonic setting for the genesis of oceanic plagiogranites: Evidence from a paleo-spreading structure in the Oman ophiolite, Earth Planet. Sc. Lett, 139, 177-194, [https://doi.org/10.1016/0012-821X\(95\)00233-3](https://doi.org/10.1016/0012-821X(95)00233-3), 1996.
- Barbarin, B.: Field evidence for successive mixing and mingling between the Piolard Diorite and the Saint-Julien-la-Vêtre Monzogranite (Nord-Foréz, Massif Central, France), Can. J. Earth Sci., 25, 49-59, <https://doi.org/10.1139/e88-005>,
760 1988.
- Batchelor, R.A. and Bowden, P.: Petrogenetic interpretation of granitoid rock series using multicationic parameters, Chem. Geol., 48, 43-55, [https://doi.org/10.1016/0009-2541\(85\)90034-8](https://doi.org/10.1016/0009-2541(85)90034-8), 1985.
- Bohlen, S.R.: Pressure-temperature-time paths and a tectonic model for the evolution of granulites, J. Geol., 95, 617-632, <https://doi.org/10.1086/629159>, 1987.
- 765 Boztuğ, D., Harlavan, Y., Archart, G.B., Satir, M., and Avci, N.: K-Ar age, whole-rock and isotope geochemistry of A-type granitoids in the Divriği-Sivas region, eastern-central Anatolia, Turkey, Lithos, 97, 193-218, <https://doi.org/10.1016/j.lithos.2006.12.014>, 2007.
- Brown, M.: P-T-t evolution of orogenic belts and the causes of regional metamorphism, Geological Society, 16, 67-81, <https://doi.org/10.1144/GSL.MEM.1995.016.01.09>, 1995.
- 770 Cawthorn, R.G. and Brown, P.A.: A model for the formation and crystallization of corundum normative calc-alkaline magmas through amphibole fractionation, The Journal of Geology, 84, 467-476, <https://doi.org/10.1086/628212>, 1976.
- Chappell, B.W. and White, A.J.R.: Two contrasting granite type, Pacific Geology, 8, 173-174, 1974.
- Chappell, B.W. and White, A.J.R.: Two contrasting granite types: 25 years later, Aust. J. Earth Sci., 48, 489-499, <https://doi.org/10.1046/j.1440-0952.2001.00882.x>, 2001.
- 775 Chappell, B.W.: Aluminium saturation in I- and S-type granites and the characterization of fractionated haplogranites, Lithos, 46, 535-551, [https://doi.org/10.1016/S0024-4937\(98\)00086-3](https://doi.org/10.1016/S0024-4937(98)00086-3), 1999.
- Chappell, B.W., Bryant, C.J., and Wyborn, D.: Peraluminous I-type granites, Lithos, 153, 142-153, <https://doi.org/10.1016/j.lithos.2012.07.008>, 2012.

- Chen, B., Jahn, B.M., and Suzuki, K.: Petrological and Nd-Sr-Os isotopic constraints on the origin of high-Mg adakitic rocks from the North China Craton: Tectonic implications, *Geology*, 41: 91–94, <https://doi.org/10.1130/G33472.1>, 2013.
- Chen, D.L., Liu, L., Sun, Y., Zhang, A.D., Liu, X.M., and Luo, J.H.: LA-ICP-MS zircon U-Pb dating for high-pressure basic granulite from North Qinling and its geological significance, *Chin. Sci. Bull.*, 49, 2296–2304, <https://doi.org/10.1360/03wd0544>, 2004.
- Chen, D.L. and Liu, L.: New data on the chronology of eclogite and associated rock from Guanpo area, North Qinling orogeny and its constraint on nature of North Qinling HP-UHP eclogite terrane, *Earth. Sci. Front.*, 18, 158-169 (in Chinese with English abstract), 2011.
- Chen, G.C., Zhang, Y.F., Pei, X.Z., Li, Z.C., Ji, X.J., Wei, J.Q., Mo, R.H., and Wang, C.: Age and geochemical characteristics of acidic tuff in the Longwanggou Formation of Caotangou Group in the Fengxian area, western part of the North Qinling orogenic belt, and their tectonic implications, *Acta Geol. Sin.*, 93, 1968-1984 (in Chinese with English abstract), <https://doi.org/10.19762/j.cnki.dizhixuebao.2019133>, 2019.
- Chen, G.C., Zhang, X.F., Pei, X.Z., Li, R.B., Li, Z.C., Chen, X.Z., and Zhang, R.Z.: Petrogenesis and geodynamic settings of the Devonian adakites and A-type granites in the North Qinling orogen, *Lithos*, 478-479, 107606, <https://doi.org/10.1016/j.lithos.2024.107606>, 2024.
- Chen, J.L., Chen, Y.C., Li, H.P., and Zhao, X.S.: Correlation between the Longshan rock group and the Qinling rock group at the junction of Qilian mountain and north Qinling mountain, *Geology of Shaanxi*, 20, 39-49 (in Chinese with English abstract), 2002.
- Chen, J.L., Xu, X.Y., Wang, H.L., Wang, Z.Q., Zeng, Z.X., Wang C., and Li P.: LA-ICPMS Zircon U-Pb Dating of Tangzang Quartz-diorite Pluton in the West Segment of North Qinling Mountains and Its Tectonic Significance, *Geoscience*, 22, 45-52 (in Chinese with English abstract), 2008.
- Chen, Y.X., Song, S.G., Niu, Y.L., and Wei, C.J.: Melting of continental crust during subduction initiation: A case study from the Chaidanuo peraluminous granite in the North Qilian suture zone, *Geochim. Cosmochim. Acta*, 132, 311-336, <https://doi.org/10.1016/j.gca.2014.02.011>, 2014.
- Chudík, P., Uher, P., Kohút, M., and Bačík, P.: Accessory columbite to tantalite, tapiolite and zircon: products of extreme fractionation in highly peraluminous pegmatitic granite from the Považský Inovec Mountains, Western Carpathians, Slovakia, *J. Geosci-Czech.*, 53, 323-334, <https://doi.org/10.3190/jgeosci.031>, 2008.
- Coleman, R.G. and Peterman, Z.E.: Oceanic plagiogranite, *J. Geophys. Res.*, 80, 1099-1108, <https://doi.org/10.1029/JB080i008p01099>, 1975.
- Dahlquist, J.A., Alasino, P.H., and Bello, C.: Devonian F-rich peraluminous A-type magmatism in the proto-Andean foreland (Sierras Pampeanas, Argentina): geochemical constraints and petrogenesis from the western-central region of the Achala batholith, *Miner. Petrol.*, 108, 391-417, <https://doi.org/10.1007/s00710-013-0308-0>, 2014.

- Davies, J.H. and von Blanckenburg, F.: Slab breakoff: A model of lithosphere detachment and its test in the magmatism and deformation of collisional orogens, *Earth Planet. Sc. Lett.*, 129, 85-102, [https://doi.org/10.1016/0012-821X\(94\)00237-S](https://doi.org/10.1016/0012-821X(94)00237-S), 1995.
- Defant, M.J. and Drummond, M.S.: Derivation of some modern arc magmas by melting of young subducted lithosphere, *Nature*, 347, 662-665, <https://doi.org/10.1038/347662a0>, 1990.
- Deng, Y.F., Song, X.Y., Hollings, P., Zhou, T.F., Yuan, F., Chen, L.M., and Zhang, D.Y.: Role of asthenosphere and lithosphere in the genesis of the Early Permian Huangshan mafic-ultramafic intrusion in the Northern Tianshan, NW China, *Lithos*, 227, 241-254, <https://doi.org/10.1016/j.lithos.2015.04.014>, ~~1990~~, 2015.
- Dewey, J.F., Dalziel, I.W.D., Reavy, R.J., and Strachan, R.A.: The Neoproterozoic to Mid-Devonian evolution of Scotland: A review and unresolved issues, *Scot. J. Geol.*, 51, 5-30, <https://doi.org/10.1144/sjg2014-007>, 2015.
- Diwu, C.R., Sun, Y., Liu, L., Zhang, C.L., and Wang, H.L.: The disintegration of Kuanping Group in North inling orogenic belts and Neo-proterozoic N-MORB, *Acta Petrol. Sin.*, 26, 2025-2038 (in Chinese with English abstract), 2010.
- Diwu, C.R., Sun, Y., Zhao, Y., Liu, B.X., and Lai, S.C.: Geochronological, geochemical, and Nd-Hf isotopic studies of the Qinling Complex, central China: Implications for the evolutionary history of the North Qinling Orogenic Belt, *Geosci. Front.*, 5, 499-513, <https://doi.org/10.1016/j.gsf.2014.04.001>, 2014.
- Dong, Y.P., Yang, Z., Zhang, G.W., Zhao, X., Xu, J.G., and Yao, A.P.: Geochemistry of the ophiolite in the Guanzizhen Area, West Qinling and its tectonic implications, *Acta Geol. Sin.*, 82, 1186-1194 (in Chinese with English abstract), 2008.
- Dong, Y.P., Zhang, G.W., Hauzenberger, C., Neubauer, F., Yang, Z., and Liu, X.M.: Palaeozoic tectonics and evolutionary history of the Qinling Orogen: Evidence from geochemistry and geochronology of ophiolite and related volcanic rocks, *Lithos*, 122, 39-56, <https://doi.org/10.1016/j.lithos.2010.11.011>, 2011a.
- Dong, Y.P., Genser, J., Neubauer, F., Zhang G.W., Liu, X.M., Yang, Z., and Heberer, B.: U-Pb and $^{40}\text{Ar}/\text{Ar}^{39}\text{Ar}$ geochronological constraints on the exhumation history of the North Qinling terrane, China, *Gondwana Res.*, 19, 881-893, <https://doi.org/10.1016/j.gr.2010.09.007>, 2011b.
- Dong, Y.P., Zhang, G.W., Neubauer, F., Liu, X.M., Genser, J., and Hauzenberger, C.: Tectonic evolution of the Qinling Orogen, China: Review and synthesis, *J. Asian Earth Sci.*, 41, 213-237, <https://doi.org/10.1016/j.jseaes.2011.03.002>, 2011c.
- Dong, Y.P., Yang, Z., Liu, X.M., Zhang, X.N., He, D.F., Li, W., Zhang, F.F., Sun, S.S., Zhang, H.F., and Zhang, G.W.: Neoproterozoic amalgamation of the Northern Qinling terrain to the North China Craton: Constraints from geochronology and geochemistry of the Kuanping ophiolite, *Precambrian Res.*, 255, 77-95, <https://doi.org/10.1016/j.precamres.2014.09.008>, 2014.
- Dong, Y.P., Zhang, X.N., Liu, X.M., Li, W., Chen, Q., Zhang, G.W., Zhang, H.F., Yang, Z., Sun, S.S., and Zhang, F.F.: Propagation tectonics and multiple accretionary processes of the Qinling Orogen, *J. Asian Earth Sci.*, 104, 84-98, <https://doi.org/10.1016/j.jseaes.2014.10.007>, 2015.

- 845 Dong, Y.P. and Santosh, M.: Tectonic architecture and multiple orogeny of the Qinling Orogenic Belt, Central China, *Gondwana Res.*, 29, 1-40, <https://doi.org/10.1016/j.gr.2015.06.009>, 2016.
- Dong, Y.P., Sun, S.S., Santosh, M., Zhao, J., Sun, J.P., He, D.F., Shi, X.H., Hui, B., Cheng, C., and Zhang, G.W.: Central China Orogenic Belt and amalgamation of East Asian continents, *Gondwana Res.*, 100, 131-194, <https://doi.org/10.1016/j.gr.2021.03.006>, 2021.
- 850 Dong, Y.P., Sun S.S., Santosh, M., Hui, B., Sun, J.P., Zhang, F.F., Cheng, B., Yang, Z., Shi, X.H., He, D.F., Yang, L., Cheng, C., Liu, X.M., Zhou, X.H., Wang, W., and Qi, N.: Cross Orogenic Belts in Central China: Implications for the tectonic and paleogeographic evolution of the East Asian continental collage, *Gondwana Res.*, 109, 18-88, <https://doi.org/10.1016/j.gr.2022.04.012>, 2022a.
- Dong Y.P., Hui B., Sun S.S., Yang Z., Zhang F.F., He D.F., Sun J.P., and Shi X.H.: Multiple orogeny and geodynamics from Proto-Tethys to Paleo-Tethys of the Central China Orogenic Belt, *Acta Geol. Sin.*, 96, 3426-3448 (in Chinese with English abstract), <https://doi.org/10.19762/j.cnki.dizhixuebao.2022245>, 2022b.
- 855 Duretz, T., Gerya, T.V., and May, D.A.: Numerical modelling of spontaneous slab breakoff and subsequent topographic response, *Tectonophysics*, 502, 244-256, <https://doi.org/10.1016/j.tecto.2010.05.024>, 2011.
- Eby, G.N., 1990: The A-type granitoids: A review of their occurrence and chemical characteristics and speculations on their petrogenesis, *Lithos*, 26, 115-134, [https://doi.org/10.1016/0024-4937\(90\)90043-Z](https://doi.org/10.1016/0024-4937(90)90043-Z).
- 860 Eby, G.N., 1992: Chemical subdivision of the A-type granitoids: Petrogenetic and tectonic implications, *Geology*, 20, 641-644, [https://doi.org/10.1130/0091-7613\(1992\)020<0641:CSOTAT>2.3.CO;2](https://doi.org/10.1130/0091-7613(1992)020<0641:CSOTAT>2.3.CO;2).
- England, P.C. and Thompson, A.B.: Pressure-Temperature-Time Paths of Regional Metamorphism I. Heat Transfer during the Evolution of Regions of Thickened Continental Crust, *J. Petrol.*, 25, 894-928, <https://doi.org/10.1093/petrology/25.4.894>, 1984.
- 865 Frey, F.A., Green, D.H., and Roy, S.D.: Integrated models of basalt petrogenesis: A study of quartz tholeiites to olivine melilites from South Eastern Australia utilizing geochemical and experimental petrological data, *J. Petrol.*, 19, 463-513, <https://doi.org/10.1093/petrology/19.3.463>, 1978.
- Frost, B.R., Barnes, C.G., Collins, W.J., Arculus, R.J., Ellis, D.J., and Frost, C.D.: A geochemical classification for granitic rocks, *J. Petrol.*, 42, 2033-2048, <https://doi.org/10.1093/petrology/42.11.2033>, 2001.
- 870 Gao, J.M., Pei, X.Z., Li, Z.C., Li, R.B., Pei, L., Wei, F.H., Wu, S.K., Liu, C.J., Wang, Y.C., and Chen, Y.X.: LA-ICP-MS zircon U-Pb dating and geochemical characteristics of the Liushuigou igneous complex, Tianshui area West Qinling Mountains, *Geological Bulletin of China*, 31, 1482-1495 (in Chinese with English abstract), 2012.
- Gao, S., Chen, D.L., Gong, X.K., Ren, Y.F., and Li, H.P.: Zircon U-Pb dating of clastic rocks and granites of Kuanping Group in Dongcha areas of Tianshui, and its geological implications, *Earth Science Frontiers*, 22, 255-264 (in Chinese with English abstract), <https://doi.org/10.13745/j.esf.2015.04.026>, 2015.
- 875

- Geng, H.Y., Sun, M., Yuan, C., Xiao, W.J., Xian, W.S., Zhao, G.C., Zhang, L.F., Wong, K., and Wu, F.Y.: Geochemical, Sr-Nd and zircon U-Pb-Hf isotopic studies of Late Carboniferous magmatism in the West Junggar, Xinjiang: Implications for ridge subduction?, *Chem. Geol.*, 266, 364-389, <https://doi.org/10.1016/j.chemgeo.2009.07.001>, 2009.
- 880 Geng, J.Z., Li, H.K., Zhang, Jian, Zhou, H.Y., and Li, H.M.: Zircon Hf isotope analysis by means of LA-MC-ICP-MS, *Geological Bulletin of China*, 30, 1508-1513 (in Chinese with English abstract), 2011.
- Gong, X.K., Chen, D.L., Ren, Y.F., Liu, L., Gao, S., and Yang, S.J.: Identification of coesite-bearing amphibolite in the North Qinling and its geological significance, *Chinese Science Bulletin*, 61, 1365-1381 (in Chinese with English abstract), 2016.
- 885 Guo, Q., Mao, X.H., Zhang, J.X., Lu, Z.L., Zhou, G.S., Teng, X., and Wu, Y.W.: Granulite-facies metamorphism in the northern part of West Qinling: Constraints from phase equilibrium modeling and in-situ U-Pb dating of monazite, *Acta Petrol. Sin.*, 38, 3259-3280, <https://doi.org/10.18654/1000-0569/2022.11.03>, 2022.
- Harris, C., Marsh, J.S., and Milner, S.C.: Petrology of the alkaline core of the essum igneous complex, Namibia: Evidence for the progressively decreasing effect of crustal contamination, *J. Petrol.*, 40, 1377-1397, <https://doi.org/10.1093/petroj/40.9.1377>, 1999.
- 890 Hawkes, W.C.J. and Kemp, A.I.S.: The differentiation and rates of generation of the continental crust, *Chem. Geol.*, 226, 134-143, <https://doi.org/10.1016/j.chemgeo.2005.09.017>, 2006.
- He, S.P., Wang, H.L., Chen, J.L., Xu, X.Y., Zhang, H.F., Ren, G.M., and Yu, J.Y.: Zircon U-Pb Chronology of Kuanping Rock Group By LA-ICP-MS and its Geological Significance, *Acta Geol. Sin.*, 81, 79-87 (in Chinese with English abstract), 2007.
- 895 He, Y.S., Li, S.G., Hoefs, J., Huang, F., Liu, S.A., and Hou, Z.H.: Post-collisional granitoids from the Dabie orogen: New evidence for partial melting of a thickened continental crust, *Geochim. Cosmochim. Acta*, 75, 3815-3838, <https://doi.org/10.1016/j.gca.2011.04.011>, 2011.
- Hou, Q.Y., Zhao, Z.D., Zhang, H.F., Zhang, B.R., Zhang, L., and Chen, Y.L.: Discussion on the tectonic affinity of ancient oceanic mantle in Western Qinling-Songpan continental tectonic node, China: From elemental and Sr-Nd-Pb isotopic evidences, *Acta Petrol. Sin.*, 22, 2901-2909 (in Chinese with English abstract), 2006.
- 900 Houseman, G.A. and Molnar, P.: Gravitational (Rayleigh-Taylor) instability of a layer with non-linear viscosity and convective thinning of continental lithosphere, *Geophys. J. Int.*, 128, 125-150, <https://doi.org/10.1111/j.1365-246X.1997.tb04075.x>, 1997.
- 905 King, P.L., White, A.J.R., Chappell, B.W., and Allen, C.M.: Characterization and origin of aluminous A-type granites from the Lachlan fold belt, southeastern Australia, *J. Petrol.*, 38, 371-391, <https://doi.org/10.1093/petroj/38.3.371>, 1997.
- Li, H.K., Geng, J.Z., Hao, S., Zhang, Y.Q., and Li, H.M.: A study on the determination of the isotope age of zircon U-Pb by laser ablation multi-receiver plasma mass spectrometry (LA-MC-ICPMS). *Acta Mineralogica Sinica* 29, 600-601 (in Chinese), <https://doi.org/10.16461/j.cnki.1000-4734.2009.s1.014>, 2009.

- 910 Li, S., Wang, T., Wilde, S.A., and Tong, Y.: Evolution, source and tectonic significance of Early Mesozoic granitoid magmatism in the Central Asian Orogenic Belt (central segment), Earth-Sci. Rev., 126: 206–234, <https://doi.org/10.1016/j.earscirev.2013.06.001>, 2013.
- Li, S.Z., Yang, Z., Zhao, S.J., Li, X.Y., Suo, Y.H., Guo, L.L., Yu, S., Dai, L.M., Li, S.J., and Mou, D.L.: Global Early Paleozoic Orogens (II): Subduction-Accretionary-Type Orogeny, Journal of Jilin University (Earth Science Edition), 46, 968-1004 (in Chinese with English abstract), <https://doi.org/10.13278/j.cnki.jjuese.201604102>, 2016b.
- 915 Li, S.Z., Jahn, B.M., Zhao, S.J., Dai, L.M., Li, X.Y., Suo, Y.H., Guo, L.L., Wang, Y.M., Liu, X.C., Lan, H.Y., Zhou, Z.Z., Zheng, Q.L., and Wang, P.C.: Triassic southeastward subduction of North China Block to South China Block: Insights from new geological, geophysical and geochemical data, Earth-Sci. Rev., 166, 270-285, <https://doi.org/10.1016/j.earscirev.2017.01.009>, 2017.
- 920 Li, S.Z., Zhao, S.J., Liu, X., Cao, H.H., Yu, S., Li, X.Y., Somerville, I., Yu, S.Y., and Suo, Y.H.: Closure of the Proto-Tethys Ocean and Early Paleozoic amalgamation of microcontinental blocks in East Asia, Earth-Sci. Rev., 186, 37-75, <https://doi.org/10.1016/j.earscirev.2017.01.011>, 2018b.
- Li, S.Z., Suo, Y.H., Li, X.Y., Liu, B., Dai, L.M., Wang, G.Z., Zhou, J., Li, Y., Liu, Y.M., Cao, X.Z., Somerville, I., Mu, D.L., Zhao, S.J., Liu, J.P., Meng, F., Zhen, L.B., Zhao, L.T., Zhu, J.J., Yu, S.Y., Liu, Y.J., and Zhang, G.W.: Microplate tectonics: new insights from micro-blocks in the global oceans, continental margins and deep mantle, Earth-Sci. Rev., 185, 1029-1064, <https://doi.org/10.1016/j.earscirev.2018.09.005>, 2018c.
- 925 Li, W.C., Chen, R.X., Zheng, Y.F., Hu, Z.: Dehydration and anatexis of UHP metagranite during continental collision in the Sulu orogen, J. Metamorphic. Geol., 32, 915–936, <https://doi.org/10.1111/jmg.12100>, 2014.
- Li, W.Y., Li, S.G., Pei, X.Z., and Zhang, G.W.: Geochemistry and zircon SHRIMP U-Pb ages of the Guanzizhen ophiolite complex, the Western Qinling orogen, China, Acta Petrol. Sin., 23, 2836-2844 (in Chinese with English abstract), 2007.
- 930 Li, W.Y.: Geochronology and Geochemistry of the Ophiolites and Island–arc–type Igneous Rocks in the Western Qinling Orogeny and the Eastern Kunlun Orogeny: Implications for the Evolution of the Tethyan Ocean, Ph.D. thesis, University of Science and Technology of China (in Chinese with English abstract), 174 pp., 2008.
- Li, W.Y., Zhang, Z.W., Wang, Y.L., Zhang, J.W., You, M.X., Zhang, Z.B., and Nanka, O.: Tectonic Transformation of Proto- and Paleo-Tethys and the Metallization of Magmatic Ni-Cu-Co Sulfide Deposits in Kunlun Orogen, Northwest China, Journal of Earth Sciences and Environment, 44, 1-19 (in Chinese with English abstract), <https://doi.org/10.19814/j.jese.2021.08033>, 2022.
- 935 Li, X.Y., Li, S.Z., Yu, S.Y., Santosh, M., Zhao, S.J., Guo, X.Y., Cao, H.H., Wang, Y.M., and Huang, Z.B.: Early Paleozoic arc-back-arc system in the southeastern margin of the north Qilian Orogen, China: Constraints from geochronology, and whole-rock elemental and Sr-Nd-Pb-Hf isotopic geochemistry of volcanic suites, Gondwana Res., 59, 9-26, <https://doi.org/10.1016/j.gr.2018.03.008>, 2018a.
- 940 Liu, C.J.: Study on Material Composition, Tectonic Evolution and Transformation at the Conjunction of Qinling and North Qilian Orogen, M.S. thesis, Chang'an University (in Chinese with English abstract), 120 pp., 2013.

- 945 Liu, L., Liao, X.Y., Zhang, C.L., Chen, D.L., Gong, X.K., and Kang, L.: Multimetamorphic timings of HP-UHP rocks in the North Qinling their geological implications, Acta Petrol. Sin., 29, 1634–1656 (in Chinese with English abstract), 2013.
- Liu, L., Liao, X.Y., Wang, Y.W., Wang, C., Santosh, M., Yang, M., Zhang, C.L., and Chen, D.L.: Early Paleozoic tectonic evolution of the North Qinling Orogenic Belt in Central China: Insights on continental deep subduction and multiphase exhumation, Earth-Sci. Rev., 159, 58-81, <https://doi.org/10.1016/j.earscirev.2016.05.005>, 2016.
- 950 Liu, L., Chen, D.L., Zhang, J.F., Kang, L., Yang, W.Q., Liao, X.Y., and Ma, T.: The Minimum Stable Pressure and Geological Significants of Supersilic Garnet in Continental Felsic Rocks: Constraints from HT-HP Experiments, Acta Geol. Sin., (English Edition) 94, 32-33, <https://doi.org/10.1111/1755-6724.14447>, 2020.
- Liu, Q., Wu, Y.B., Wang, H., Gao, S., Qin, Z.W., Liu, X.C., Yang, S.H., and Gong, H.J. Zircon U-Pb ages and Hf isotope compositions of migmatites from the North Qinling terrane and their geological implications, J. Metamorph. Geol., 32, 177-193, <https://doi.org/10.1111/jmg.12065>, 2014.
- 955 Liu, X.C., Wu, Y.B., Gao, S., Wang, H., Zheng, J.P., Hu, Z.C., Zhou, L., and Yang, S.H.: Record of multiple stage channelized fluid and melt activities in deeply subducted slab from zircon U-Pb age and Hf-O isotope compositions, Geochim. Cosmochim. Acta, 144, 1-24, <https://doi.org/10.1016/j.gca.2014.08.016>, 2014.
- Liu, Y.S., Gao, S., Hu, Z.C., Gao, C.G., Zong, K.Q., and Wang, D.B.: Continental andoceanic crust recycling induced melt-peridotite interactions in the Trans North China orogen: U-Pb dating, Hf isotopes and trace elements in zircons from mantle xenoliths, J. Petrol., 51, 537-571, <https://doi.org/10.1093/petrology/egp082>, 2010.
- 960 Ludwig, K.R.: Isoplot v.3.75: A Geochronological Toolkit for Microsoft Excel, Berkeley Geochronology Center Special Publication, 2012.
- Ma, X.X., Shu, L.S., and Meert, J.G.: Early Permian slab breakoff in the Chinese Tianshan belt inferred from the post-collisional granitoids, Gondwana Res., 27, 228-243. <http://dx.doi.org/10.1016/j.gr.2013.09.018>, 2015.
- 965 Mao, X.H., Zhang, J.X., Yu, S.Y., Li, Y.S., Yu, X.X., and Lu, Z.L.: Early Paleozoic granulite-facies metamorphism and anatexis in the northern West Qinling orogen: Monazite and zircon U-Pb geochronological constraints, Science China: Earth Sciences, 60, 943-957, <https://doi.org/10.1007/s11430-016-9029-7>, 2017.
- Mao, X.H, Zhang, J.X., Yu, S.Y., Li, Y.S., Yu, X.X., Lu, Z.L., and Zhou, G.S.: Metamorphism of Qinling Complex in Northern West Qinling Orogen: Petrology, Phase Equilibria Modelling of Paragneiss and Their Geological Implication, Earth Sci., 43, 278-295 (in Chinese with English abstract), <https://doi.org/10.3799/dqkx.2018.017>, 2018.
- 970 Maniar, P.D. and Piccoli, P.M.: Tectonic discrimination of granitoids, Geol. Soc. Am. Bull., 101, 635-643, [https://doi.org/10.1130/0016-7606\(1989\)101<0635:TDOG>2.3.CO;2](https://doi.org/10.1130/0016-7606(1989)101<0635:TDOG>2.3.CO;2), 1989.
- ~~Mark, B.A., Song, S.G., Wang, C., Zeng, R.Y., and Wen, T.: An oblique subduction model for closure of the Proto-Tethys and Palaeo-Tethys oceans and creation of the Central China Orogenic Belt, Earth Sci. Rev., 240, 104385, <https://doi.org/10.1016/j.earscirev.2023.104385>, 2023.~~
- 975 Mattern, F. and Schneider, W.: Suturing of the Proto- and Paleo-Tethys oceans in the western Kunlun (Xinjiang, China), J. Asian Earth Sci., 18, 637-650, [https://doi.org/10.1016/S1367-9120\(00\)00011-0](https://doi.org/10.1016/S1367-9120(00)00011-0), 2000.

- McKerrow, W.S., Niocaill, C.M., and Dewey, J.F.: The Caledonian Orogeny redefined, *J. Geol. Soc. London.*, 157, 1149-1154, <https://doi.org/10.1144/jgs.157.6.1149>, 2000.
- 980 McDonough, W.F. and Sun, S.S.: The composition of the Earth, *Chem. Geol.*, 120, 223-253, [https://doi.org/10.1016/0009-2541\(94\)00140-4](https://doi.org/10.1016/0009-2541(94)00140-4), 1995.
- Merino, E., Villaseca, C., Orejana, D., and Jeffries, T.: Gahnite, chrysoberyl and beryl co-occurrence as accessory minerals in a highly evolved peraluminous pluton: The Belvís de Monroy leucogranite (Cáceres, Spain), *Lithos*, 179, 137-156, <http://dx.doi.org/10.1016/j.lithos.2013.08.004>, 2013.
- 985 Middlemost, E.A.K.: Naming materials in the magma/igneous rock system, *Earth-Sci. Rev.*, 37, 215-224, [https://doi.org/10.1016/0012-8252\(94\)90029-9](https://doi.org/10.1016/0012-8252(94)90029-9), 1994.
- Miller, C.F.: Are strongly peraluminous magmas derived from pelilicsedimentary sources?, *The Journal of Geology*, 93, 673-689, <https://doi.org/10.1086/628995>, 1985.
- Mushkin, A., Navon, O., Halicz, L., Hartmann, G., and Stein, M.: The Petrogenesis of A-type Magmas from the Amram Massif, Southern Israel, *J. Petrol.*, 44, 815-832, <https://doi.org/10.1093/petrology/44.5.815>, 2003.
- 990 Pan, G.T., Wang, L.Q., Geng, Q.R., Yin, F.G., Wang, B.D., Wang, D.B., Peng, Z.M., and Ren, F.: Space-time structure of the Bangonghu-Shuanghu-Nujiang-Changning-Menglian Mega-suture zone: A discussion on geology and evolution of the Tethys Ocean, *Sedimentary Geology and Tethyan Geology*, 40, 1-19 (in Chinese with English abstract), <https://doi.org/10.19826/j.cnki.1009-3850.2020.07001>, 2020.
- 995 Patiño Douce, A.E. and Beard, J.S.: Dehydration-melting of Biotite Gneiss and Quartz Amphibolite from 3 to 15 kbar, *J. Petrol.*, 36, 707-738, 1995.
- Patiño Douce, A.E.: Generation of metaluminous A-type granites by low-pressure melting of calc-alkaline granitoids, *Geology*, 25, 743-746, [https://doi.org/10.1130/0091-7613\(1997\)025<0743:GOMATG>2.3.CO;2](https://doi.org/10.1130/0091-7613(1997)025<0743:GOMATG>2.3.CO;2), 1997.
- Peccerillo, A. and Taylor, S.R.: Geochemistry of eocene calc-alkaline volcanic rocks from the Kastamonu area, Northern Turkey, *Contrib. Mineral. Petrol.*, 58, 63-81, <https://doi.org/10.1007/BF00384745>, 1976.
- 1000 Pei, X.Z., Ding, S.P., Li, Y., Su, C.Q., Li, Y.J., Chen, S.E., Yang, J.X., Zhang, J., Zhao, X., and Guo, J.F.: Regional geological report of the Peoples Republic of China: Tianshui map (I48C002003), the Institute of Geological Survey of Chang'an University, Map, 2004a. (In Chinese).
- Pei, X.Z., Ding, S.P., Hu, B., Li, Y., Zhang, G.W., and Guo J.F.: Definition of the Guanzizhen ophiolite in Tianshui area western Qinling, and its geological significance, *Geological Bulletin of China*, 23, 1202-1208 (in Chinese with English abstract), 2004.
- 1005 Pei, X.Z., Li, Z.C., Ding, S.P., Li, Y., Hu, B., and Liu, H.B.: Geochemical characteristics and zircon U-Pb ages of island-arc basic igneous complexes in the Tianshui area, West Qinling, *Geology in China*, 32, 5-16 (in Chinese with English abstract), 2005.
- 1010 Pei, X.Z., Liu, H.B., Ding, S.P., Li, Z.C., Hu, B., Sun, R.Q., and Hou, Y.H.: Geochemical characteristics and tectonic significance of the meta-volcanic rocks in the Liziyuan Group from Tianshui area, western Qinling Orogen,

Geotectonica et Metallogenia, 30, 193-205 (in Chinese with English abstract),
<https://doi.org/10.16539/j.ddgzyckx.2006.02.010>, 2006.

- 1015 Pei, X.Z., Ding, S.P., Li, Z.C., Liu, Z.Q., Li, G.Y., Li, R.B., Wang, F., and Li, F.J.: LA-ICP-MS Zircon U-Pb Dating of the
Gabbro from the Guanzizhen Ophiolite in the Northern Margin of the Western Qinling and Its Geological Significance,
Acta Geol. Sin., 81, 1550-1561 (in Chinese with English abstract), 2007a.
- Pei, X.Z., Liu, Z.Q., Ding, S.P., Li, Z.C., Li, G.Y., Li, R.B., Wang, F., and Li, F.J.: Zircon LA-ICP-MS U-Pb Dating of the
Gabbro from the Baihua Igneous Complex in Tianshui Area, Eastern Gansu, and Its Geological Significance, Advances
in Earth Science, 22, 818-827 (in Chinese with English abstract), 2007b.
- 1020 Pei, X.Z., Ding, S.P., Li, Z.C., Liu, Z.Q., Li, R.B., Feng, J.Y., Sun, Y., Zhang, Y.F., Liu, Z.G., Zhang, X.F., Chen, G.C., and
Chen, Y.X.: Early Paleozoic Tianshui-Wushan Tectonic zone of the Northern Margin of West Qinling and its Tectonic
Evolution, Acta Geol. Sin., 83, 1547-1564 (in Chinese with English abstract), 2009.
- Peng, Y.B., Yu, S.Y., Zhang, J.X., Li, Y.S., Li, S.Z., and Lv, P.: Building a continental arc section: Constraints from
Paleozoic granulite-facies metamorphism, anatexis, and magmatism in the northern margin of the Qilian Block,
1025 northern Tibet Plateau, Geol. Soc. Am. Bull., 134, 1301-1318, <https://doi.org/10.1130/B36100.1>, 2021.
- Pearce, J.A., Harris, N.B.W., and Tindle, A.G.: Trace element discrimination diagrams for the tectonic interpretation of
granitic rocks, J. Petrol., 25, 956-983, <https://doi.org/10.1093/petrology/25.4.956>, 1984.
- Petford, N. and Atherton, M.: Na-rich Partial Melts from Newly Underplated Basaltic Crust: the Cordillera Blanca Batholith,
Peru, J. Petrol., 37, 1491-1521, <https://doi.org/10.1093/petrology/37.6.1491>, 1996.
- 1030 Qin, J.F., Lai, S.C., Zhang, Z.H., Zhang, Z.Z., Long, X.P., and Zhu, R.Z.: Crustal growth and evolution in convergent margin:
Evidence from three Paleozoic granitic pulses in the junction zone between Qinling and Qilian orogenic belt, Lithos,
434-435, 106938, <https://doi.org/10.1016/j.lithos.2022.106938>, 2022.
- Qiu, J.S., Xiao, E., Hu, J., Xu, X.S., Jiang, S.Y., and Li, Z.: Petrogenesis of highly fractionated I-type granites in the coastal
area of northeastern Fujian Province: Constraints from zircon U-Pb geochronology, geochemistry and Nd-Hf isotopes,
1035 Acta Petrol. Sin., 24, 2468-2484 (in Chinese with English abstract), 2008.
- Rapp, R.P. and Watson, E.B.: Dehydration melting of meta-basalt at 8-32 kbar: Implications for continental growth and crust
mantle recycling, J. Petrol., 36, 891-931, <https://doi.org/10.1093/petrology/36.4.891>, 1995.
- Raumer, J.F. and Stampfli, G.M.: The birth of the Rheic Ocean-Early Palaeozoic subsidence patterns and subsequent
tectonic plate scenarios, Tectonophysics, 461, 9-20, <https://doi.org/10.1016/j.tecto.2008.04.012>, 2008.
- 1040 Ren, J.S., Zhu, J.B., Li, C., and Liu, R.Y.: Is the Qinling Orogen an Indosinian Collisional Orogenic Belt?, Earth Science, 44,
1476-1486 (in Chinese with English abstract), 2019.
- Ren, L., Liang, H.Y., Bao, Z.W., Zhang, J., Li, K.X., and Huang, W.T.: The petrogenesis of early Paleozoic high-Ba-Sr
intrusions in the North Qinling terrane, China, and tectonic implications, Lithos, 314, 534-550,
<https://doi.org/10.1016/j.lithos.2018.06.027>, 2018.

- 1045 Ren, L., Liang, H.Y., Bao, Z.W., and Huang, W.T.: Early Paleozoic magmatic “flare-ups” in western Qinling orogeny, China: New insights into the convergence history of the North and South China Blocks at the northern margin of Gondwana, *Lithos*, 380-381, 105833, <https://doi.org/10.1016/j.lithos.2020.105833>, 2021.
- Rudnick, R.L.: Making continental crust, *Nature*, 378, 571-578, <https://doi.org/10.1038/378571a0>, 1995.
- 1050 [Sandiford, M. and Powell, R.: Deep crustal metamorphism during continental extension: Modern and ancient examples, *Earth Planet. Sci. Lett.*, 79, 151-158, \[https://doi.org/10.1016/0012-821X\\(86\\)90048-8\]\(https://doi.org/10.1016/0012-821X\(86\)90048-8\), 1986.](#)
- Siégl, C., Bryan, S.E., Allen, C.M., and Gust, D.A.: Use and abuse of zircon-based thermometers: a critical review and a recommended approach to identify antecrystic zircons, *Earth-Sci. Rev.*, 176, 87-116, <https://doi.org/10.1016/J.EARSCIREV.2017.08.011>, 2018.
- Skjerlie, K.P. and Johnston, A.D.: Vapor-absent melting at 10 kbar of a biotite- and amphibole- bearing tonalitic gneiss: Implications for the generation of A-type granites, *Geology*, 20, 263-266, [https://doi.org/10.1130/0091-7613\(1992\)020<0263:VAMAKO>2.3.CO;2](https://doi.org/10.1130/0091-7613(1992)020<0263:VAMAKO>2.3.CO;2), 1992.
- 1055 Song, S.G., Niu, Y.L., Su, L., and Xia, X.H.: Tectonics of the North Qilian Orogen, NW China, *Gondwana Res.*, 23, 1378-1401, <https://doi.org/10.1016/j.gr.2012.02.004>, 2013.
- 1060 [Song, S.G., Niu, Y.L., Su, L., Wei, C.J., and Zhang, L.F.: Adakitic \(tonalitic-trondhjemitic\) magmas resulting from eclogite decompression and dehydration melting during exhumation in response to continental collision, *Geochim. Cosmochim. Acta*, 130, 42–62, <https://doi.org/10.1016/j.gca.2014.01.008>, 2014.](#)
- [Song, S.G., Wang, M.J., Wang, C., and Niu, Y.L.: Magmatism during continental collision, subduction, exhumation and mountain collapse in collisional orogenic belts and continental net growth: A perspective, *Sci. China: Earth Sci.*, 58, 1284-1304, <https://doi.org/10.1007/s11430-015-5102-x>, 2015.](#)
- 1065 Song, S.G., Wu, Z.Z., Yang, L.M., Su, L., Xia, X.H., Wang, C., Dong, J.L., Zhou, C.A., and Bi, H.Z.: Ophiolite belts and evolution of the Proto-Tethys Ocean in the Qilian Orogen, *Acta Petrol. Sin.*, 35, 2948-2970 (in Chinese with English abstract), 2019.
- Song, Z.G., Jia, Q.Z., Zhang, Z.T., and Zhang, Y.: The Early Palaeozoic volcanic rock series and its interconnection relationship between the North Qinling and the North Qilian Orogens, *Northwest Geoscience*, 12, 1-82 (in Chinese with English abstract), 1991.
- 1070 Stampfli, G.M. and Borel, G.D.: A plate tectonic model for the Paleozoic and Mesozoic constrained by dynamic plate boundaries and restored synthetic oceanic isochrons, *Earth Planet. Sc. Lett.*, 196, 17-33, [http://dx.doi.org/10.1016/S0012-821X\(01\)00588-X](http://dx.doi.org/10.1016/S0012-821X(01)00588-X), 2002.
- Streckeisen, A.: To each plutonic rock its proper name, *Earth-Sci. Rev.*, 12, 1-33, [https://doi.org/10.1016/0012-8252\(76\)90052-0](https://doi.org/10.1016/0012-8252(76)90052-0), 1976.
- 1075 [Su, L., Song, S.G., Sun, B., Zhou, D.W., and Hao, J.Y.: The SHRIMP zircon U-Pb ages of the garnet pyroxenite and Fushui complex from Songshugou area and its constrain on the tectonic evolution of Qinling Orogenic belt, *Chin. Sci. Bull.*, 49, 1209-1211 \(in Chinese with English abstract\), 2004.](#)

- 1080 Sun M.S. and Dong H.B.: Re-discussion of the stratigraphic sequence of the Caotangou Group and its age, *Geology of Shaanxi*, 13, 22-30 (in Chinese with English abstract), 1995.
- Sun, S.S. and McDonough, W.F.: Chemical and isotopic systematics of oceanic basalts: Implications for mantle composition and processes, *Geol. Soc. Lond. Spec. Publ.*, 42, 313-345, <https://doi.org/10.1144/GSL.SP.1989.042.01.19>, 1989.
- Sylvester, P.J.: Post-collisional alkaline granites, *The Journal of Geology*, 97, 261-280, <https://doi.org/10.1086/629302>, 1989.
- 1085 Tang, Y., Chen, D.L., Ren, Y.F., Wang, H.J.: Discovery of Early Paleozoic eclogite-facies metamorphic rocks in the western part of North Qinling Orogen and its geological significance, *Acta Petrol. Sin.*, 38, 585-597, <https://doi.org/10.18654/1000-0569/2022.03.01.2022>.
- Taylor, S.R. and McLennan, S.M.: The continental crust: its composition and evolution. Blackwell Scientific Publications, 1985.
- 1090 Thompson, A.B. and England, P.C.: Pressure-Temperature-Time Paths of Regional Metamorphism II. Their Inference and Interpretation using Mineral Assemblages in Metamorphic Rocks, *J. Petrol.*, 25, 929-955, <https://doi.org/10.1093/petrology/25.4.929>, 1984.
- Torsvik, T.H. and Rehnstrom, E.F.: The Tornquist Sea and Baltica-Avalonia docking, *Tectonophysics*, 362, 67-82, [https://doi.org/10.1016/s0040-1951\(02\)00631-5](https://doi.org/10.1016/s0040-1951(02)00631-5), 2003.
- Torsvik, T.H.: Earth history: A journey in time and space from base to top, *Tectonophysics*, 760, 297-313, 1095 <https://doi.org/10.1016/j.tecto.2018.09.009>, 2019.
- Turnbull, R., Tulloch, A., Ramezani, J., and Jongens, R.: Extension-facilitated pulsed S-I-A-type “flare-up” magmatism at 370 Ma along the southeast Gondwana margin in New Zealand: Insights from U-Pb geochronology and geochemistry, *Geol. Soc. Am. Bull.*, 128, 1500-1520, <https://doi.org/10.1130/B31426.1>, 2016.
- 1100 Turpin, L., Cuney, M., Friedrich, M., Bouchez, J.L., and Aubertin, M.: Meta-igneous origin of Hercynian peraluminous granites in N.W. French Massif Central: implications for crustal history reconstructions, *Contrib. Mineral. Petrol.*, 104, 163-172, <https://doi.org/10.1007/BF00306440>, 1990.
- Turner, S.P., Foden, J.D., and Morrison, R.S.: Derivation of some A-type magmas by fractionation of basaltic magma: An example from the Padthaway Ridge, South Australia, *Lithos*, 28, 151-179, [https://doi.org/10.1016/0024-4937\(92\)90029-X](https://doi.org/10.1016/0024-4937(92)90029-X), 1992.
- 1105 van de Zedde, D.M.A. and Wortel, M.J.R.: Shallow slab detachment as a transient source of heat at midlithospheric depths, *Tectonics*, 20, 868-882, <https://doi.org/10.1029/2001TC900018>, 2001.
- van Staal, C.R., Barr, S.M., and Murphy, J.B.: Provenance and tectonice volution of Ganderia: Constraints on the evolution of the Iapetus and Rheic oceans, *Geology*, 40, 987-990, <https://doi.org/10.1130/G33302.1>, 2012.
- 1110 Watson, E.B. and Harrison, T.M.: Zircon Saturation Revisited: Temperature and Composition Effects in a Variety of Crustal Magma Types, *Earth Planet. Sc. Lett*, 64, 295-304, [https://doi.org/10.1016/0012-821X\(83\)90211-X](https://doi.org/10.1016/0012-821X(83)90211-X), 1983.

- Wang, B.D., Wang, L.Q., Zhou, D.Q., Wang, D.B., Yu, Y.P., Yan, G.C., and Wu, Z.: Co-Shuanghu-Changning-Menglian suture zone: The boundary between Gondwanaland and Pan-Cathaysia mainland in the Qinghai-Tibet Plateau, Geological Bulletin of China, 40, 1783-1798 (in Chinese with English abstract), 2021.
- 1115 Wang, H., Wu, Y.B., Gao, S., Zhang, H.F., Liu, X.C., Gong, H.J., Peng, M., Wang, J., and Yuan, H.L.: Silurian granulite-facies metamorphism, and coeval magmatism and crustal growth in the Tongbai orogen, central China, Lithos, 125, 249-271, <https://doi.org/10.1016/j.lithos.2013.07.021>, 2011.
- Wang, H. and Wu, Y.B.: Early Paleozoic HP-UHP metamorphism of the Qinling orogen, Chin. Sci. Bull., 58, 2124-2131 (in Chinese), 2013.
- 1120 Wang, H.L., He, S.P., Chen, J.L., Xu, X.Y., Sun, Y., and Diwu, C.R.: LA-ICPMS Dating of Zircon U-Pb and Tectonic Significance of Honghuapu Subduction-Related Intrusions in the West Segment of Northern Qinling Mountains, Geoscience, 20, 536-544 (in Chinese with English abstract), 2006.
- Wang, H.L., He, S.P., Chen, J.L., Xu, X.Y., Sun, Y., Diwu, C.R., and Li, H.P.: LA-ICPMS zircon U-Pb dating of the Hudian gneissic monzogranite in the western segment of the North Qinling and its geological significance, Geology in China, 34, 17-25 (in Chinese with English abstract), 2007.
- 1125 Wang, H.L., Xu, X.Y., Chen, J.L., Sun, Y., Li, W.Z., Li, P., Li, T., and Zhang, H.: Dating and Geochemical Characteristics of the Yanwan Paleozoic Collisional Intrusion in the West Segment of Northern Qinling Mts, Acta Geol. Sin., 83, 353-364 (in Chinese with English abstract), 2009.
- Wang, J., Zhang, H.F., Xu, W.C., and Cai, H.M.: Petrogenesis of Granites from Dangchuan Area in West Qinling Orogenic Belt and Its Tectonic Implication, Earth Science, 33, 474-486 (in Chinese with English abstract), 2008.
- 1130 Wang, M.J., Song, S.G., Niu, Y.L., and Su, L.: Post-collisional magmatism: Consequences of UHPM terrane exhumation and orogen collapse, N. Qaidam UHPM belt, NW China, Lithos, 210-211, 181-198, <https://doi.org/10.1016/j.lithos.2014.10.006>, 2014.
- Wang, T., Wang, X.X., Tian, W., Zhang, C.L., Li, W.P., and Li, S.: North Qinling Paleozoic granite associations and their variation in space and time: Implications for orogenic processes in the orogens of central China, Scientia Sinica (Terrae), 39, 949-971 (in Chinese with English abstract), 2009.
- 1135 Wang, X.X., Wang, T., and Zhang, C.L.: Neoproterozoic, Paleozoic, and Mesozoic granitoid magmatism in the Qinling Orogen, China: Constraints on orogenic process, J. Asian Earth Sci., 72, 129-151, <https://doi.org/10.1016/j.jseaes.2012.11.037>, 2013.
- Wang, X.X., Wang, T., and Zhang, C.L.: Granitoid magmatism in the Qinling orogen, central China and its bearing on orogenic evolution, Scientia Sinica (Terrae), 45, 1109-1125 (in Chinese with English abstract), 2015.
- 1140 Wang, X.S., Bi, X.W., Leng, C.B., Zhong, H., Tang, H.F., Chen, Y.W., Yin, G.H., Huang, D.Z., and Zhou, M.F.: Geochronology and geochemistry of Late Cretaceous igneous intrusions and Mo-Cu-(W) mineralization in the southern Yidun Arc, SW China: Implications for metallogenesis and geodynamic setting, Ore Geol. Rev., 61, 73-95, <https://doi.org/10.1016/j.oregeorev.2014.01.006>, 2014.

- 1145 Wang, Y.C.: Geological characteristics and Tectonic significance of Caledonian collision-post collision type granite at the conjunction of Qinling and Qilian, M.S. thesis, Chang'an University (in Chinese with English abstract), 96 pp., 2013.
- Weyer, S., Muenker, C., and Mezger, K.: Nb/Ta, Zr/Hf and REE in the depleted mantle: implications for the differentiation history of the crust-mantle system, *Earth Planet. Sc. Lett.*, 205, 309-324, [https://doi.org/10.1016/S0012-821X\(02\)01059-2](https://doi.org/10.1016/S0012-821X(02)01059-2), 2003.
- 1150 Whalen, J.B., Currie, K.L., and Chappell, B.W.: A-type granites: Geochemical characteristics, discrimination and petrogenesis, *Contrib. Mineral. Petrol.*, 95, 407-419, <https://doi.org/10.1007/BF00402202>, 1987.
- Wilson, M.: *Igneous Petrogenesis: A Global Tectonic Approach*, London: Unwin Hyman, 1-466, <https://doi.org/10.1007/978-1-4020-6788-4>, 1989.
- Wu, C.L., Yao, S.Z., Yang, J.S., Zeng, L.S., Chen, S.Y., Li, H.B., Qi, X.X., Joseph, L.W., and Frank, K.M.: Double subduction of the Early Paleozoic North Qilian oceanic plate: Evidence from granites in the central segment of North Qilian, NW China, *Geology in China*, 33, 1197-1208 (in Chinese with English abstract), 2006.
- 1155 Wu, C.L., Gao, Y.H., Li, Z.L., Lei, M., Qin, H.P., Li, M.Z., Liu, C.H., Frost, R.B., Robinson, P.T., and Wooden, J.L.: Zircon SHRIMP U-Pb dating of granites from Dulan and the chronological framework of the North Qaidam UHP belt, NW China, *Sci. China: Earth Sci.*, 57, 2945-2965, <https://doi.org/10.1007/s11430-014-4958-5>, 2014.
- 1160 Wu, F.Y., Jahn, B.M., Wilde, S.A., Lo, C.H., Yui, T.F., Lin, Q., Ge, W.C., and Sun, D.Y.: Highly fractionated -type granites in NE China (I): geochronology and petrogenesis, *Lithos*, 66, 241-273, [https://doi.org/10.1016/S0024-4937\(02\)00222-0](https://doi.org/10.1016/S0024-4937(02)00222-0), 2003.
- Wu, F.Y., Li, X.H., Yang, J.H., and Zheng, Y.F.: Discussions on the petrogenesis of granites, *Acta Petrol. Sin.*, 23, 1217-1238 (in Chinese with English abstract), 2007.
- 1165 Wu, F.Y., Liu, X.C., Ji, W.Q., Wang, J.M., and Yang, L.: Highly fractionated granites: Recognition and research, *Scientia Sinica (Terrae)*, 47, 745-765 (in Chinese with English abstract), 2017.
- Wu, F.Y., Wan, B., Zhao, L., Xiao, W.J., and Zhu, R.X.: Tethyan geodynamics, *Acta Petrol. Sin.*, 36, 1627-1674 (in Chinese with English abstract), 2020.
- 1170 Wu, Y.B. and Zheng, Y.F.: Tectonic evolution of a composite collision orogen: An overview on the Qinling-Tongbai-Hong'an-Dabie-Sulu orogenic belt in central China, *Gondwana Res.*, 23, 1402-1428, <https://doi.org/10.1016/j.gr.2012.09.007>, 2013.
- Wyborn, D., Turner, B.S., and Chappell, B.W.: The Boggy Plain Supersuite: A distinctive belt of I-type igneous rocks of potential economic significance in the Lachlan Fold Belt, *Aust. J. Earth Sci.*, 34, 21-43, <https://doi.org/10.1080/08120098708729392>, 1987.
- 1175 Xia, L.Q., Xia, Z.C., and Xu, X.Y.: Early Palaeozoic Mid-Ocean Ridge-Ocean Island and Back-Arc Basin Volcanism in the North Qilian Mountains, *Acta Geol. Sin.*, 301-312 (in Chinese with English abstract), <https://doi.org/10.19762/j.cnki.dizhixuebao.1998.04.002>, 1998.

- Xiao, X.C., Wang, J., Su, L., and Song, S.G.: A further discussion of the Kūda ophiolite, West Kunlun, and its tectonic significance, *Geological Bulletin of China*, 72, 745-750 (in Chinese with English abstract), 2003.
- 1180 [Xiang, H., Zhong, Z.Q., Li, Y., Qi, M., Zhou, H.W., Zhang, L., Zhang, Z.M., and Santosh, M.: Sapphirine-bearing granulites from the Tongbai orogen, China: Petrology, phase equilibria, zircon U-Pb geochronology and implications for Paleozoic ultrahigh temperature metamorphism, *Lithos*, 208-209, 446-461, <https://doi.org/10.1016/j.lithos.2014.08.017>, 2014.](#)
- 1185 Xie, S.X., Ren, L., Zhang, J., Bao, Z.W., and Liang, H.Y.: Petrogenesis of the Caotangou Group Rhyolite in the Western Part of North Qinling Terrane and its Geological Implication, *Geotectonica et Metallogenia*, 44, 1239-1257 (in Chinese with English abstract), <https://doi.org/10.16539/j.ddgzyckx.2020.06.014>, 2020.
- Xin, Y.J. and Huang, D.Z.: Early Paleozoic arc-back-arc system evolution in the junction of the Qinling and Qilian Orogens: Geochemical constraints from ca. 445–430 Ma magmatic rocks in the Tianshui area, *Int. Geol. Rev.*, 65, 442-467, <https://doi.org/10.1080/00206814.2022.2051084>, 2023.
- 1190 Xu, T., Guo, Q.M., Chen, Q.M., Wang, Q., Zhang, S.H., Zuo, W.Q., and Ren, C.Z.: Age and Geochemical Features of the Early Silurian Jiguanya Highly Fractionated Syn-collisional Granites in the Baoji Area of North Qinling Orogen and Their Geological Significance, *Bulletin of Geological Science and Technology*, 37, 79-87 (in Chinese with English abstract), <https://doi.org/10.19509/j.cnki.dzkq.2018.0111>, 2018.
- 1195 Xu, X.C., Pei, X.Z., Liu, C.J., Li, R.B., Li, Z.C., Wei, B., Wang, Y.Y., Liu, T.J., Ren, H.Z., Chen, W.N., and Chen, Y.X.: Geochemical characteristics of the Yinaigou Early Paleozoic Caotangou Group volcanic rocks in Tianshui of West Qinling Mountains and their geological significance, *Geology in China*, 41, 851-865 (in Chinese with English abstract), 2014.
- 1200 Xu, X.Y., He, S.P., Wang, H.L., Zhang, E.P., Chen, J.L., and Sun, J.M.: Tectonic Framework of North Qinling Mountain and North Qilian Mountain Conjunction Area in Early Paleozoic: A Study of the Evidences from Strata and Tectonic-Magmatic Events, *Northwestern Geology*, 41, 1-21 (in Chinese with English abstract), 2008.
- [Xu, X.Y., Chen, J.L., and Yan, Z.: Geological Map of Qinling and Adjacent Areas \(1:500,000\), Xi'an Map Publishing House, Map, 2012. \(In Chinese\).](#)
- Xu, Y.J., Du, Y.S., Cawood, P.A., Guo, H., Hu, H., and An, Z.H.: Detrital zircon record of continental collision: Assembly of the Qilian Orogen, China, *Sediment. Geol.*, 230, 35-45, <https://doi.org/10.1016/j.sedgeo.2010.06.020>, 2010a.
- 1205 Xu Y.J., Du Y.S., Cawood, P.A., and Yang J.H.: Provenance record of a foreland basin: Detrital zircon U-Pb ages from Devonian strata in the North Qilian Orogenic Belt, China, *Tectonophysics*, 495, 337-347, <https://doi.org/10.1016/j.tecto.2010.10.001>, 2010b.
- 1210 Xu, Z.Q., Yang, J.S., Li, H.B., and Yao, J.X.: The Early Palaeozoic Terrene Framework and the Formation of the High-Pressure (HP) and Ultra-High Pressure (UHP) Metamorphic Belts at the Central Orogenic Belt (COB), *Acta Geol. Sin.*, 80, 1793-1806 (in Chinese with English abstract), 2006.

Xu, Z.Q., Zheng, B.H., and Wang, Q.: From accretion to collision: situation and outlook, Acta Geol. Sin., 95, 75-97, <https://doi.org/10.19762/j.cnki.dizhixuebao.2021050>, 2021.

Xue, F., Kröner, A., Reischmann, T., and Lerch, F.: Palaeozoic pre- and post-collision calc-alkaline magmatism in the Qinling orogenic belt, central China, as documented by zircon ages on granitoid rocks, J. Geol. Soc. London., 153, 409-417, <https://doi.org/10.1144/gsjgs.153.3.0409>, 1996.

Yan, Q.R., Wang, Z.Q., Chen, J.L., Yan, Z., Wang, T., Li, Q.G., Jiang, C.F., and Zhang, Z.Q.: Tectonic setting and SHRIMP age of volcanic rocks in the Xieyuguan and Caotangou Groups: implications for the North Qinling Orogenic Belt, Acta Geol. Sin., 81, 488-500+583-584 (in Chinese with English abstract), 2007.

Yang, J.H., Wu, F.Y., Chung, S.L., Wilde, S.A., and Chu, M.F.: A hybrid origin for the Qianshan A-type granite, northeast China: Geochemical and Sr–Nd–Hf isotopic evidence, Lithos, 89, 89-106, <https://doi.org/10.1016/j.lithos.2005.10.002>, 2006.

Yang, J.S., Xu, Z.Q., Pei, X.Z., Shi, R.D., Wu, C.L., Zhang, J.X., Li, H.B., Meng, F.C., and Rong, H.: Discovery of Diamond in North Qinling: Evidence for a Giant UHPM Belt across Central China and Recognition of Paleozoic and Mesozoic Dual Deep Subduction between North China and Yangtze Plates, Acta Geol. Sin., 76, 484-495 (in Chinese with English abstract), 2002.

Yang, L.M., Song, S.G., Allen, M.B., Su, L., Dong, J.L., and Wang, C.: Oceanic accretionary belt in the West Qinling Orogen: links between the Qinling and Qilian orogens, China, Gondwana Res., 64, 137-162, <https://doi.org/10.1016/j.gr.2018.06.009>, 2018a.

Yang, L.M., Song, S.G., Su, L., Li, Y.G., Dong, J.L., and Li, Q.Y.: Discovery of Liziyuan high-pressure granulite in Tianshui region: Evidence for Late Devonian orogeny in West Qinling Orogen, Acta Petrol. Sin., 34, 2425-2440 (in Chinese with English abstract), 2018b.

Yang, Z., Dong, Y.P., Liu, X.M., and Zhang, J.H.: LA-ICP-MS zircon U-Pb dating of gabbro in the Guanzizhen ophiolite, Tianshui, West Qinling, China, Geological Bulletin of China, 25, 1321-1325 (in Chinese with English abstract), 2006.

Yu, S.Y., Zhang, J.X., Gong, J.H., and Li, Y.S.: Qaidam Mountains Research on HP granulite-facies metamorphism and anatexis: A case study of Dulan area in the North, Acta Petrol. Sin., 29, 2061-2072 (in Chinese with English abstract), 2013.

Zeng, W., Sun, F.Y., Zhou, H.Y., Wang, J.Y., Li, Z.D., Chen, J.Q., Bi, J.H., and Cui, Y.R.: Cassiterite U-Pb Age, Geochemistry and Their Geological Significances of Rare Metal Pegmatites in Guanpo Area, North Qinling, China, Earth Science, 48, 2851-2871 (in Chinese with English abstract), 2023.

Zhai, X.M., Day, H.W., Hacker, B.R., and You, Z.D.: Paleozoic metamorphism in the Qinling orogen, Tongbai Mountains, central China, Geology, 26, 371-374, [https://doi.org/10.1130/0091-7613\(1998\)026<0371:PMITQO>2.3.CO;2](https://doi.org/10.1130/0091-7613(1998)026<0371:PMITQO>2.3.CO;2), 1998.

Zhang, C.L., Yu, H.F., Shen, J.L., Dong, Y.G., Ye, H.M., and Guo, K.Y.: Zircon SHRIMP Age Determination of the Giant-crystal Gabbro and Basalt in Kūda, West Kunlun: Dismembering of the Kuda Ophiolite, Geological Review, 50, 639-643 (in Chinese with English abstract), <https://doi.org/10.16509/j.georeview.2004.06.013>, 2004.

- 1245 [Zhang, C.L., Liu, L., Wang, T., Wang, X.X., Li, L., Gong Q.F., and Li, X.F.: Granitic magmatism related to early Paleozoic continental collision in the North Qinling belt, Chin. Sci. Bull., 58, <https://doi.org/10.1007/s11434-013-6064-z>, 2013.](#)
- Zhang, G.W., Zhang, B.R., Yuan, X.C., and Xiao, Q.H. et al.: Qinling Orogenic Belt and Continental Dynamics, Beijing: Science Press, 1-855 (in Chinese), 2001.
- Zhang, G.W., Guo, A.L., Dong, Y.P., and Yao, A.P.: Rethinking of the Qinling Orogen, Journal of Geomechanics, 25, 746-768 (in Chinese with English abstract), 2019.
- 1250 [Zhang, H.R. and Hou, Z.Q.: Pattern and Process of Continent-Continent Collision Orogeny: A Case Study of the Tethys Collisional Orogen, Acta Geol. Sin., 89, 1539-1559 \(in Chinese with English abstract\), 2015.](#)
- [Zhang, J.X., Yu, S.Y., Meng, F.C., and Li, J.P.: Paired high-pressure granulite and eclogite in collision orogens and their geodynamic implications, Acta Petrol. Sin., 25, 2050-2066 \(in Chinese with English abstract\), 2009.](#)
- 1255 Zhang, J.X., Yu, S.Y., and Meng, F.C.: Ployphase Early Paleozoic metamorphism in the northern Qinling orogenic belt, Acta Petrol. Sin., 27, 1179-1190 (in Chinese with English abstract), 2011.
- Zhang, J.X., Yu, S.Y., Li, S.Z., Li, Y.S., Yu, X.X., Lin, Y.H., and Mao, X.H.: Subduction, accretion and closure of Proto-Tethyan Ocean: Early Paleozoic accretion/collision orogeny in the Altun-Qilian-North Qaidam orogenic system, Acta Petrol. Sin., 31, 3531-3554 (in Chinese with English abstract), 2015.
- 1260 Zhao, G.C., Wang, Y.J., Huang, B.C., Dong, Y.P., Li, S.Z., Zhang, G.W., and Yu, S.: Geological reconstructions of the East Asian blocks: From the breakup of Rodinia to the assembly of Pangea, Earth-Sci. Rev., 186, 262-286, <https://doi.org/10.1016/j.earscirev.2018.10.003>, 2018.
- [Zheng, Y.F., Zhao, Z.F., and Chen, Y.X.: Continental subduction channel processes: Plate interface interaction during continental collision, Chin. Sci. Bull., 58, 4371-4377, <https://doi.org/10.1007/s11434-013-6066-x>, 2013.](#)
- 1265 [Zheng, Y.F., Chen, Y.X., Dai, L.Q., and Zhao, Z.F.: Developing plate tectonics theory from oceanic subduction zones to collisional orogens, Sci. China: Earth Sci., 58, 1045–1069, <https://doi.org/10.1007/s11430-015-5097-3>, 2015.](#)
- [Zheng, Y.F., Chen, Y.X., Chen, R.X, and Dai, L.Q.: Tectonic evolution of convergent plate margins and its geological effects, Sci. China: Earth Sci., 65, 1247-1276, <https://doi.org/10.1007/s11430-022-9947-6>, 2022.](#)
- Zhu, J.C., Rao, B., Xiong, X.L., Li, F.C., and Zhang, P.H.: Comparison and genetic interpretation of Li-F rich, rare-metal bearing granitic rocks, Geochimica, 31, 141-152 (in Chinese with English abstract), <https://doi.org/10.19700/j.0379-1726.2002.02.005>, 2002.
- 1270 [Zhu, R.X., Zhao, P., and Zhao L.: Tectonic evolution and geodynamics of the Neo-Tethys Ocean, Sci. China: Earth Sci., 65, 1-24, <https://doi.org/10.1007/s11430-021-9845-7>, 2022.](#)
- Zhu, T., Dong, Y.P., Wang, W., Xu, J.G., Ma, H.Y., and Cha, L.: The Geochemical Characteristics and Tectonic Setting of Volcanics in Caotangou Group, Northwestern Geology, 41, 59-66 (in Chinese with English abstract), 2008.
- 1275