

External and internal drivers behind the formation, vegetation succession, and carbon balance of subarctic fen margin.

Teemu Juselius-Rajamäki¹, Sanna Piilo¹, Susanna Salminen-Paatero², Emilia Tuomaala¹, Tarmo Virtanen¹, Atte Korhola¹, Anna Autio³, Hannu Marttila³, Pertti Ala-Aho³, Annalea Lohila^{4,5}, Minna Väiliranta¹

¹University of Helsinki, Ecosystem and Environmental Research Program

²University of Helsinki, Department of Chemistry, Radiochemistry

³University of Oulu, Water, Energy and Environmental Engineering Research Unit

⁴Finnish Meteorological Institute, Climate System Research Unit, P.O. Box 503, 00101 Helsinki, Finland

⁵Institute for Atmospheric and Earth System Research/Physics, University of Helsinki, P.O. Box 64, 00014, Finland

Correspondence to: Teemu Juselius-Rajamäki (teemu.juseliuss@helsinki.fi)

Abstract. Peatlands are the most dense terrestrial carbon storage and recent studies have shown that the northern peatlands have continued to expand to new areas to this day. However, depending on the vegetation and hydrological regime in the newly initiated areas, the climate forcing may vary. If these new areas develop as wet, fen-type peatlands with high methane emissions they would initially have a warming effect on the climate. On the other hand, if the development starts as dry bog-type peatlands, these new peatland areas would likely act as a strong carbon sink from early on. However, although some research has concentrated on the expansion of the new northern peatland areas, there remains a significant lack of studies on the successional development of the newly initiated peatland frontiers. In this research, we combined palaeoecological, remote sensing and hydrological modeling methods to study the expansion and successional pathway dynamics in a subarctic fen margin in Finnish Lapland and discussed possible implications for carbon balance of these margin peatland areas. Our results showed that the studied peatland margins had started to develop ca. 2000 years ago and have continued to expand thereafter, and this expansion has occurred in non-linear fashion. In addition, the wet fen-type vegetation persisted in the studied margin for majority of the development history and only the dryer conditions after the Little Ice Age instigated the fen-to-bog transition. However, a notable part of the fen margins in the Lompolovuoma and Lompolojänkkä basins has remained as a wet fen-types, and their persistence was likely caused by the hydrological conditions in the peatland and in the surrounding catchment. Our findings show a large variation in the peatland expansion and succession dynamics even within a single peatland basin. Although changes in climate conditions had initiated the fen-to-bog process in some margins, some had remained in the wet, fen stage showing resilience to allogenic forcings. Thus, when estimating the peatland carbon stocks, and predicting the future trajectories for peatland development, this heterogeneity should be taken into account to avoid errors caused by over-simplification of peatland lateral expansion dynamics.

1 Introduction

After peatland initiation through a primary peat formation, infilling (terrestrialization), or paludification, peatland area is increased by lateral expansion – the most important process of forming new peatland areas in modern climate

(Ruppel et al., 2013). In raised mires, these new peatland margin areas have been generally described as moist minerotrophic fens and spruce swamps (Howie and Meerveld, 2011; Rydin and Jeglum, 2013), while in aapa mires (patterned fens) the margins vary from dry ombrotrophic bogs to wetter lush swamps (Laitinen et al., 2005, 2007). However, although the current vegetation in aapa mire margins has been described in a standard peatland literature, there is an obvious lack of studies on the long-term successional development of these transitional ecotones between peatlands and the surrounding mineral land. A recent study with main focus on aapa mire region of Finland showed that the northern peatlands are still expanding (Juselius-Rajamäki et al., 2023), and whether these newly forming peatland areas initiate and develop as moist fens or dryer bog-types can markedly affect the climate forcing of this recent lateral expansion.

Lateral expansion process is driven by both allogenic and autogenic factors. For instance, forest fire or other disturbance in areas adjacent to a peatland decreases the evapotranspiration and causes rising water table that enables peatland expansion (Kuhry and Turunen, 2006). Similarly, waterlogging may be caused by autogenic development of adjacent peatlands. As the peat accumulates vertically, the surface and groundwater flow pathways are directed towards the margins of peat mound (Autio et al., 2023), creating suitable conditions for new peat formation (Korhola, 1996; Rydin and Jeglum, 2013). On the other hand, drainage ditches located in the mire margins can prevent natural discharge to peatlands blocking the lateral expansion (Sallinen et al., 2019), while high-intensity fires can destroy peat layers setting back the advance of peatland margins (Kuhry, 1994; Simard et al., 2007). Also, climate affects the lateral expansion of peatlands, and for example, during warm and dry climate phase between 8000 – 5000 Before Present (BP) expansion of peatlands slowed down, while wet and humid climate from 5000 to 3000 BP promoted lateral peatland expansion (Korhola, 1994, 1995; Ruppel et al., 2013).

The development of vegetation communities in the newly initiated peatland margins vary according to the non-linear successional trajectory and is driven, particularly by seasonal water availability, and consequently transportation of essential ions (Goud et al., 2018). Depending on topography, surface flow may control the first appearance of vegetation communities. Later groundwater seepage, point-scale or as wider seepage front, transports moisture and dissolved elements for established plants. Compared to raised mires that have grown vertically above the surrounding marginal areas, and often the entire landscape (Howie and Meerveld, 2011; Rydin and Jeglum, 2013), the secondary peatland development pattern over the margins is more complex for aapa mires, because the shape of the peatland varies from flat to concave (Seppä, 2002) and formation of new peatland habitats is dependent of water supplies from snowmelt (Sallinen et al., 2023) and dilution of ion concentrations (pH-levels). Newly established habitat types may range from ombrotrophic bog-types to minerotrophic swamps and fens (Foster and King, 1984; Laitinen et al., 2005, 2007; Mäkilä and Moisanen, 2007; Ruuhijärvi, 1983). However, mechanisms, such as surface water hydraulic forcing, which create different types of margins, are currently poorly understood.

Differences in local hydrology mirrored in the current vegetation communities suggest opposite climatic feedback mechanisms for the peatland centers and marginal areas. The overall climatic effect of peatlands is and has been

strongly controlled by the balance between sequestration of carbon dioxide (CO₂), and release of methane (CH₄) (Frolking and Roulet, 2007). Methane is produced in anoxic conditions and released into the atmosphere via vegetation, ebullition or by diffusion (Lai, 2009; Rydin and Jeglum, 2013). However, in areas where the acrotelm i.e., the oxic and biologically active layer of the peat, is thick most of the methane is oxidated to carbon dioxide (Lai, 2009). Thus, in the peatland margins where dry bog-type vegetation communities dominate, the climate forcing is most likely negative, i.e, cooling impact on climate, due to the continuous uptake of CO₂ and low CH₄ emissions. However, if the water table depth becomes too deep, accelerated decomposition can turn these locations to carbon sources due to increased CO₂ emissions that negate the decrease in CH₄ emissions (Evans et al., 2021). On the other hand, in wet fen-type margins high methane emissions have an opposite effect on short timescales, further amplified by graminoid vegetation communities (Bubier et al., 1993; Juutinen et al., 2013; Kou et al., 2022; Ward et al., 2013).

Often, the interest of (palaeo)peatland researchers has been in the deepest and oldest part of a peatland while the development of peatland margins, i.e., young areas, has attracted less consideration (Korhola et al., 2010; Ruppel et al., 2013). Only recently the focus has turned to peatland margins and peat profile sampling has been extended to the peatland-upland ecotones (Juselius-Rajamäki et al., 2023; Lacourse et al., 2019; Le Stum-Boivin et al., 2019; Mathijssen et al., 2014, 2016, 2017; Peregon et al., 2009; Schaffhauser et al., 2017). Even these studies have focused more on the expansion dynamics of the peatlands, while the vegetation succession of the marginal areas has deserved lesser consideration. As the past vegetation communities can be used to ascertain climate feedback, the knowledge of vegetation succession in peatland margins can be used to better understand how lateral expansion has affected the past climates and helps us to predict the effects of lateral expansion for future climate change.

Here, we studied the expansion and successional pathways of peatland margins in a subarctic fen, Lompolovuoma, located in Finnish Lapland using a novel approach combining palaeoecological, remote sensing and hydrological modeling methods. To have a wider understanding of development and diversity of plant communities in aapa mire margins, we used additional comparable peat profile data from three other peatlands from northern Finland as well as detailed remote sensing-based vegetation and land-cover classification (Räsänen et al., 2021) from Lompolovuoma fen margins. The aim of the research was to study when did the development of the mire margins occur and what was the vegetation in the mire margins during the different stages of this development. In addition, we studied how did the local and external factors contributed to the mire margin development. The results of our study give an insight into aapa mire margin succession patterns, their relation to hydrology, and gives us basic understanding of the peatland climate feedback and carbon balance related to peatland lateral expansion in subarctic areas.

2 Methods and materials

2.1 Study sites

The Lompolovuoma study site is a subarctic fen located in the municipality of Muonio in Finnish Lapland (67° 59' 42" N, 24° 12' E, Fig. 1a). The site belongs to the northern aapa mire zone with more continental climate, shorter

growing season, and more profound frost effects than on the aapa mires located further south (Ruuhijärvi, 1983). The mean annual temperature in the study site is 0.4 °C (2003-2019) and the mean annual precipitation 647 mm (2008–2019) (Marttila et al., 2021).

We studied the margins of a sub-basin in a larger fen complex that comprises of several elongated, north-south aligned fen areas. The vegetation in the central areas of the study site is dominated by typical wet fen taxa, such as various *Carex* species and flark *Sphagnum* species. Strings are mainly occurring in the southern parts of the basin. A stream runs across the peatland basin from south to north towards Lake Pallasjärvi.

Vegetation communities in the studied peatland margins resemble raised pine bog habitats in the south with low hummocks and narrow lawn areas (Laine et al., 2018). The ground layer consists of *Sphagnum fuscum* and *Sphagnum angustifolium*, and *Cladonia* sp. lichens also occurred. In the field layer *Eriophorum vaginatum*, *Rubus chamaemorus* and various dwarf shrubs such as *Empetrum* sp., *Andromeda polifolia* and *Vaccinium vitis-idaea* was found. In addition, stunted *Pinus sylvestris* grow on the hummocks.

To expand our understanding of vegetation succession in aapa mire margins, we used three additional short profiles collected from aapa mires elsewhere in Finland: Syysjärvi, Salamajärvi and Patvinsuo (Fig. 1a). These profiles enabled comparison between different local and geographic settings across Finland. For a full description of the study sites, field sampling, and laboratory analysis for supplementary sites, see Juselius-Rajamäki et al. (2023).




Figure 1. a) Location of the main study site Lompolovuoma marked with an orange star while locations of the comparison sites are marked with blue circles. Borderline separating aapa and raised mire complex areas in Finland is presented. 1. b) The study location within Lompolovuoma basin shows the study transect samples in blue circles, and the long cores in orange circles. For the transect samples the sample code indicates the sample location within the transect: and Sm for the sample closest to the mire margin, Si for the sample in the middle of the transect and Sp for the sample closest to the peatland center. Mire margins are shown with a red line. Aerial image from National Land Survey of Finland, taken in 2023. This figure contains data from National Land Survey of Finland NLS Aerial photographs database.

2.2 Field sampling

The field sampling for the study was conducted during the summer and the autumn of 2022. To study the lateral expansion dynamics, we sampled a total of three transects coded as T followed by transect number with three peat core samples coded as S followed by core location indicator: p for sample located closest to the peatland centre, i for the intermediate sample, and m for the sample located closest to the margin. Each transect ran from the edge of the peatland towards the centre. The transects were placed on the “bog-type” margin with variable width, which led variable lengths in our transects. All profiles were collected from lawn-microform with similar types of vegetation. The first sampling location for each transect was at the extreme end of the “bog-type” margin next to the central fen area. The other samples were collected along the transect from locations fulfilling the above criteria, and in such manner that entire length of “bog-type” -margin at the transect location was covered. We established two transects from the east edge and one transect from the north edge of the fen sub-basin (Fig. 1b). The peat cores were taken with a box corer ($7 \times 4 \times 65$ cm) down to mineral subsoil. To reconstruct Holocene peatland initiation, in addition to the peat cores sampled from the mire margin, four long cores were collected from two different locations of the central part of the study basin: two of the long cores, a and b, were respectively located close to each other and are representing replicates (Fig. 1b). These samples were collected using a Russian peat corer (3×50 cm). The profiles were described and classified in the field, and the length of the profile was measured. The location of each sampling point was recorded using Trimble R8 GPS device with ± 0.05 m accuracy and the distance between each transect sampling point was measured using a tape measurer. After sampling, the peat cores were carefully wrapped in plastic to avoid any contamination and transported to University of Helsinki premises. The samples were stored in a cold room prior to further analysis.

2.3 Laboratory analysis

The short profiles were cut into 1 cm subsamples and from these subsamples, dry bulk density (BD, g/cm^3) and sediment organic matter (OM) based on the loss on ignition method (LOI) were determined (Heiri et al., 2001). We used LOI values to differentiate between the mineral subsoil and the peat. We defined peat initiation depth based on the first layer where $\text{LOI} \geq 70\%$ (Korhola, 1994). In addition, we analyzed the C/N-content as following: 4 cm interval (transect 1 and 3) and 5 cm interval (transect 2) using LECO TruSpec micro-Elemental Determinator. For the long profiles, contact layer between limnotelmatic *Equisetum* peat and fen peat, without visible *Equisetum* remains, was first determined in the field and then confirmed using a stereomicroscope.

To reconstruct past changes in vegetation, plant sub-macrofossil analysis for each short peat profile was conducted at 4 cm intervals and when prominent changes occurred the interval was increased to every second cm. The percentage

proportion of each peat forming vegetation type of a total sample volume (100 %) was analyzed from 5 cm³ peat samples that were gently rinsed under running water in a 100 µm sieve. The residue was analyzed under a stereomicroscope following Välranta et al. (2007) and Mauquoy et al. (2014). For example, seeds and leaves were counted in exact numbers and the percentage of unidentified organic material (UOM) estimated for highly decomposed organic remains that had lost their microscopical characteristics. A compound light microscope was used for higher taxonomic level identification. Software Tilia (Grimm, 1991) and C2 (Juggins, 2007) were used to create diagrams.

To study the lateral expansion and succession dynamics of the fen margins, we applied AMS radiocarbon (¹⁴C) determinations to date the basal peat of each short profile and the depths corresponding to the major regime shifts in vegetation e.g., first occurrence of the *Sphagnum* mosses overlying sedge-dominated peat and the shift to *Sphagnum* dominance. For the long profiles we dated the shift from limnotelmatic *Equisetum* peat to fen peat to gain understanding of long-term development of the Lompolovuoma fen. Terrestrial plant remains and/or charcoal were prioritized for ¹⁴C analyses over bulk peat samples (Quirk et al., 2022). However, in three cases regarding the short cores, the peat was highly decomposed and bulk peat had to be used (Table 1). In addition, bulk peat was used as material for the AMS dating of the long cores. Rootlets were carefully removed from the bulk peat samples. Samples were dated in Poznan Radiocarbon laboratory (Poznan, Poland). We calibrated ¹⁴C BP ages against the INTCAL 20 NH calibration curve (Reimer et al., 2020) and modern dates (pMC % modern carbon) by using the Bomb21 NH1 calibration curve (Hua et al., 2022). Finally, calibrated ages were rounded to the nearest 5 years.

For the comparison profiles, radiocarbon dating results were acquired from Juselius-Rajamäki et al. (2023). In addition, radiolead (²¹⁰Pb) dating was performed for the comparison profiles at the Department of Chemistry, University of Helsinki. The separation method used for ²¹⁰Po was a combination of several previously published methods (Ali et al., 2008; Flynn, 1968; Kauranen and Miettinen, 1966; Sanderson, 2016). Dried peat samples were digested with concentrated acids HNO₃ and HCl. ²⁰⁹Po tracer spike was added to the samples at the beginning of the analysis to monitor the yield loss. After digestion, the samples were evaporated to dryness, dissolved into a dilute HCl solution, filtered, and transferred into deposition vessels made from PTFE. Ascorbic acid was added to reduce interfering impurities, e.g. Fe, in the samples. ²¹⁰Po was deposited spontaneously onto a silver disc in the deposition vessel using a heated water bath (65–75 °C) with constant stirring for 2.5–3 hours. The activity concentration of ²¹⁰Po was measured from the silver disc with a PIPS (passivated implanted planar silicon) detector. The activity concentration of ²¹⁰Pb in the samples was obtained via equilibrium of ²¹⁰Po and ²¹⁰Pb in the samples.

2.4 Age-depth models

Age-depth models with ¹⁴C ages were done using Bacon package ver. 3.2.0 (Blaauw and Christen, 2011) in R software ver. 4.3.1 (R Core Team, 2023). We assumed different peat accumulation rates for different vegetation community stages, and these were acquired from the literature representing similar vegetation communities and geographic locations (Granlund et al., 2022; Mäkilä et al., 2001; Mathijssen et al., 2014; Rydin and Jeglum, 2013; Zhang et al.,

2020). We used these accumulation rates as a prior value for the age-depth models for corresponding vegetation community stages. After the initial model run, if the model fit was not satisfied (Blaauw and Christen, 2011), the prior values were altered to ensure the model fit. Boundaries were set for the profiles based on vegetation community shifts, and different accumulation rates were calculated for different plant communities. For the profiles with both ^{14}C and ^{210}Pb ages, e.g., (SyJ T1Sm, SJ T3Sm, and PS T1Sm), we used Plum package ver. 0.4.0 (Aquino-López et al., 2018) in R software ver. 4.3.1 (R Core Team, 2023). For the comparison peat profiles, the same prior accumulation rates were used as for the Lompolovuoma study site. Again, to accommodate for the individual peat profile characteristics, the rates were modified to ensure age-depth model fit. The individual age-depth models containing the accumulation rates and used boundaries are presented in supplementary data (Fig. S1-S12).

2.5 Lateral expansion rates

Lateral expansion rates (cm/year) were calculated between adjacent peat sections in each transect. The rates were calculated by dividing the horizontal distance between adjacent dated profiles (cm) with the difference of the basal ages, respectively (years). Mean calibrated ages from the age-depth model were used.

2.6 Current vegetation community analysis

We used field and remote sensing-based land cover type data from Räsänen et al. (2021), where the methodology is described in detail, to estimate the proportion of vegetation communities in the peatland margins. Land cover classification was based on field verification data collected in summer 2019, and multisource remote sensed data. Classification was conducted in two steps: first 4 channel 0.5 m pixel sized aerial image from summer 2018 was segmented, and then for these segments (mean size 50 m²) values were calculated from several Lidar, Planetscope and Sentinel images from years 2018 and 2019, and these were classified using random forest classification. Final land cover product had 16 classes, and the overall classification accuracy was 76%. Here, we used a simplified classification based on ombrotrophic – minerotrophic gradient to describe habitat conditions and related vegetation community. In addition, tree-covered fens were separated from open fens. Applied vegetation communities are: “bog”-type (referring to dry conditions), “fen”-type (referring to wet conditions), and tree-covered fens (referring to forested peatland) and these enable comparison with the remote sensing data. These were combined from the land cover type classes with similar ecological characteristics: dwarf shrub pine bogs and dwarf shrub bogs as the bogs, tall sedge fens and flarks as the fens and paludified spruce, birch, and mixed forests as the tree-covered fens. We delineated our study basin Lompolovuoma and adjacent Lompolojänkä basin based on the land cover dataset in ArcGis Pro ver. 3.1.0 (ESRI, 2023) and calculated the proportion of each land cover type for the whole peatland area and for the peatland margins. For the peatland margins, we chose a 25 meter distance from the peatland forest border to represent the marginal peatland area. This distance prevented any overlap of the marginal areas even in the narrowest parts of the peatland and allowed non-biased analysis of the marginal peatland types irrelevant to the topography or vegetation on site.

2.7 Hydrological analyses

To study the hydrological drivers behind the development of divergent peatland types at the fen margins detected in our vegetation coverage analysis, we used the fully integrated physically based-hydrogeological model HydroGeoSphere (Aquanty, 2015). The model allows explicit simulation of water exchange between groundwater and surface water and can be parameterized using physical properties of peat and mineral soils. The high spatial resolution of the model makes it suitable to estimate water fluxes at the scale of vegetation inventories and remote sensing data. This model has been previously implemented for the Pallaslompolo catchment, and the full methodology for this hydrogeological model is described in Autio et al. (2023). Due to the original study boundaries, this model only covers Lompolojänkää sub-basin. In this study, we (1) investigated the resulting hydrological conditions in terms of groundwater-surface water exchange flux and (2) compared the impact of the current (baseline) and the drier climate in terms of water table elevation (Helama et al., 2017) .

In (1), we investigated the prevailing groundwater-surface water exchange fluxes of the transient model run averaged over the summer of 2017 within each peatland type. For (2), we studied the effect of drier climate conditions by comparing the outputs of the steady-state simulations for the current climate with the effective rainfall P_{eff} equal to 385 mm (average for 2016–2018) and the drier climate of P_{eff} equal to 250 mm. The value of 250 mm is within the measured range that varied between 170 mm and 574 mm in 2008–2018 but represents a significantly lower value than the measured long-term mean of 358 mm for the years (2008–2018). Due to the variable density of the model computing mesh, the model output was first plotted in the postprocessing visualisation software Tecplot 360 EX 2022 R2, which accommodates value interpolation over element size. The variables were divided into separate bins according to magnitude, hereafter referred to as contour groups showing spatial variation in model output. The resulting raster image was imported to GIS mapping software (ESRI, 2023), georeferenced and clipped according to the defined peatland margins for each peatland type. The areas of each contour group were then calculated respectively for each peatland type.

3 Results

3.1 Peat initiation and spatial development of the peatland margins

In transect 1, the oldest basal date ca. 2230 cal. BP was dated from the peat profile closest to the mire center (T1Sp) (Table 1). For the intermediate profile (T1Si) the basal age was ca. 1185 cal. BP and for the profile next to the forest (T1Sm) basal age was ca. 990 cal. BP. In transect 2, the oldest basal age found in the intermediate profile (T2Si) was 1930 cal. BP while younger basal ages of 1025 cal. BP and 390 cal. BP were found for the T3Sp and T3Sm, respectively (Table 1). Oldest basal age in transect 3 was 1390 cal. BP in the intermediate sample T2Si (Table 1) while the basal age in the sample closest to the mire center (T3Sp) was 1225 cal. BP and in the peatland margin (T3Sm) 765 cal. BP.

Long core (LC) dating results suggest that a shift from limnotelmatic peat to fen peat occurred ca. 6300 cal. BP at earliest and around 4000 cal. BP at latest (Table 1). This change occurred earlier in the northern part of the sub-basin (LC1a ca. 6290 cal. BP and LC1b ca. 6360 cal. BP). In the southern part, this shift occurred ca. 4865 cal. BP for LC2a and ca. 4365 cal. BP for LC2b.

Table 1. Peat profile description. Coring location describes the location of the sampling across the transects with “Margin” being located closest to the mire-forest boundary, and “Peatland” closest to the mire center. Sample type describes the location within the profile with “Basal” representing the contact layer between peat and mineral subsoil, “Sphagnum occurrence” indicating the first occurrence of Sphagnum mosses and “Sphagnum dominance” the first layer with clear Sphagnum-dominance. Sample description indicates material used in ¹⁴C analyses. Age (cal. BP) with 95 % confidence interval show calibrated median age with 95.4 % confidence intervals.

Laboratory code	Core code	Sample location	Sample type	Depth (cm)	Dated material	¹⁴ C Age (BP)	±	pMC	±2	Age (cal. BP) with 95.4 % confidence interval
Poz-162912	T1Sm	Margin	<i>Sphagnum</i> occurrence	7-8	<i>Sphagnum</i> and feather moss leaves and stems			103.46	0.33	-60 (-5 – -65)
Poz-162911	T1Sm	Margin	Basal	19-20	Bulk	1085	30			990 (1060 – 930)
Poz-162914	T1Si	Intermediate	<i>Sphagnum</i> dominance	27-28	<i>Sphagnum</i> moss leaves and stems, woody	315	30			390 (460 – 305)
Poz-162913	T1Si	Intermediate	Basal	30-31	Woody	1250	50			1185 (1285 – 1065)
Poz-165854	T1Sp	Peatland	<i>Sphagnum</i> dominance	21-22	<i>Sphagnum</i> moss leaves and stems			121.63	0.35	-35 (-5 – -35)
Poz-162924	T1Sp	Peatland	<i>Sphagnum</i> occurrence	36-37	<i>Sphagnum</i> moss leaves and stems, woody	845	30			740 (790 – 685)
Poz-162925	T1Sp	Peatland	Basal	40-41	Woody, charred wood	2210	30			2230 (2325 – 2125)
Poz-162917	T2Sm	Margin	<i>Sphagnum</i> dominance	25-26	<i>Sphagnum</i> moss leaves and stems, woody	85	30			115 (260 – 25)
Poz-162916	T2Sm	Margin	Basal	29-30	Shrub leaves, woody, bulk	320	35			390 (470 – 305)
Poz-165855	T2Si	Intermediate	<i>Sphagnum</i> dominance	32-33	<i>Sphagnum</i> moss leaves and stems	75	30			115 (260 – 30)
Poz-162920	T2Si	Intermediate	<i>Sphagnum</i> occurrence	41-42	<i>Sphagnum</i> and feather moss leaves and stems	570	70			590 (665 – 505)
Poz-162918	T2Si	Intermediate	Basal	48-49	Woody	1995	30			1930 (1995 – 1835)
Poz-162922	T2Sp	Peatland	<i>Sphagnum</i> dominance	35-36	<i>Sphagnum</i> and feather moss leaves and stems	150	30			145 (285 – 50...)
Poz-162921	T2Sp	Peatland	Basal	46-47	Woody	1140	30			1025 (1175 – 960)
Poz-165856	T3Sm	Margin	<i>Sphagnum</i> dominance	8-9	<i>Sphagnum</i> moss leaves and stems, woody			107.25	0.33	-55 (-5 – -60)
Poz-162880	T3Sm	Margin	Basal and <i>Sphagnum</i> occurrence	19-20	Woody, charred wood	870	30			765 (905 – 690)
Poz-165857	T3Si	Intermediate	<i>Sphagnum</i> dominance	14-15	Bulk with majority (>95%) of <i>Sphagnum</i> , woody			109.35	0.34	-50 (-5 – -55)
Poz-162619	T3Si	Intermediate	Basal and <i>Sphagnum</i> occurrence	32-33	Woody	1520	30			1390 (1515 – 1315)
Poz-165859	T3Sp	Peatland	<i>Sphagnum</i> dominance	13-14	Bulk with majority			135.14	0.35	-25 (-25 – -30)

					(>95 %) of Sphagnum					
Poz-162923	T3Sp	Peatland	<i>Sphagnum</i> occurrence	21-22	Woody	105	30			110 (270 – 15)
Poz-162882	T3Sp	Peatland	Basal	40-41	Woody	1290	30			1225 (1290 – 1175)
Poz-165876	LC1a	Fen lawn	Fen peat	127-129	Bulk peat with roots removed	5490	40			6290 (6395 – 6200)
Poz-165959	LC1b	Fen lawn	Fen peat	123-124	Bulk peat with roots removed	5595	35			6360 (6445 – 6300)
Poz-165085	LC2a	Fen lawn	Fen peat	180-181	Bulk peat with roots removed	4305	35			4865 (4965 – 4830)
Poz-165086	LC2b	Fen lawn	Fen peat	189-191	Bulk peat with roots removed	3930	35			4365 (4515 – 4245)

3.2 Peat properties

A shift from mineral layer to organic layer was sharp in all profiles and in the upper parts of the profiles, the loss on ignition (LOI, %) values varied only slightly (Fig. 3, 4, 5). In transect 3, the mineral material has intruded into the peat at depths of 14 cm (T3Sp) and 23 cm (T3Si). Compared to LOI (%), more fluctuations were visible in bulk density (BD, g/cm³) values (Fig. 3, 4, 5). Above the sharp mineral subsoil – peat contact, the BD decreased towards the surface with the lowest values found at the top of peat profiles. A stepwise decrease in BD occurred in peat profiles T1Sp, T2Si, T2Sp and T3Sm while a gradual decrease in BD values was observed in other profiles. Carbon content (%) above the mineral subsoil contact varied only slightly along the peat profiles (Fig. 3, 4, 5) and highest nitrogen contents along the peat profiles were found in the layers closest to the mineral subsoil and the surface (Fig. 3, 4, 5).

3.3 Fossil plant communities and succession of the peatland margins

Three main vegetation stages were identified in the Lompolovuoma margin peat profiles (Fig 2a-c, Fig 3-5). The first and oldest stage consisted of the remnants of cyperaceous and ericaceous vegetation (C-E), but lacked brown mosses usually associated with calcareous fens. This phase was characterized by a high proportion of unidentified organic matter (UOM), indicating a high level of humidification. The second stage contained remains of mixed *Cyperaceae-Ericaceae-Sphagnum* (C-E-S) vegetation. The transition from stage 1 to 2 occurred gradually in some peat profiles while sometimes shift was abrupt. In this transition *Sphagnum* sect. *Acutifolia* started to replace cyperaceous vegetation. In the transects 1 and 3 the high level of decomposition prevented species-level identification of *Sphagnum* mosses in early C-E-S stage. However, in transect 2, the C-E stage was directly overlain by *Sphagnum fuscum*. In the final *Sphagnum-Ericaceae* stage (S-E), the plant community is dominated by *Sphagnum* mosses, and the cyperaceous vegetation is nearly or completely missing. *Sphagnum* species consists of *Sphagnum fuscum*, *S. capillifolium*, *S. russowii*, and *S. angustifolium*. A varying amount (%) of ericaceous vegetation is usually mixed with the sphagna. Varying amounts of forest bryophytes, such as *Pleurozium schreberi* is also detected through the peat layers. In addition, in the marginal profiles of transect 1, the mire vegetation was replaced twice by forest vegetation, and similar replacement occurred once in the margin of transect 2. Macrofossil data is presented in Fig. A2, A3, and A4.




Figure 2a-c. Transect profiles. The figures show the vegetation community stages: C-E (*Cyperaceae-Ericaceae*), C-E-S (*Cyperaceae-Ericaceae-Sphagnum*), and S-E (*SphagnumEricaceae*) as well as the forest community stages in the margins of T1 and T2. In addition, the onset of each stage at the location peat profiles is shown with ages (CE = Common Era) derived from the age-depth model. The ratios between the x and y axes varies between the illustrations. The vegetation is presented to give a rough impression of real-life conditions in the study transects locations and is not in the true scale.

At the onset of peat development in the mire margins, the C-E vegetation community dominated (Fig. 2a-c, Fig. 3-5). In transects 1 and 2 this layer was thin, only up to 4 cm in transect 1 and from 5 cm to 9 cm in transect 2. In the transect 3 the C-E layer was markedly thicker, 16 cm in T3Si and 18 cm in T3Sp. Duration of the C-E stage was highly

variable: in transect 1 the C-E stage lasted between ca. 250 (T1Sm) and 600 years (T1Sp). In transect 2 C-E stage lasted between ca. 200 (T2Sm) and 1000 (T2Si) years. In transect 3 the C-E stage was missing from the profile closest to the mire margin (T3Sm), and *Sphagnum* mosses established directly on top of the mineral subsoil. The duration of the C-E stage in T3Si was ca. 1250 years and in T3Sp ca. 1050 years.

The C-E stage ended asynchronously across Lompolovuoma mire margin and in most of the cases the C-E stage was followed by the mixed C-E-S stage where sphagna started to colonize the margins. The establishment of sphagna marking the start of the C-E-S occurred between ca. 670 and 970 CE in transect 1, between ca. 1020 and 1720 CE in transect 2, and between ca. 1720 and 1770 CE in transect 3. No C-E-S stage was detected in samples T1Sm and T2Sm. Instead, the vegetation shifted towards a mix of ericaceous vegetation, *Pleurozium schreberi* and *Dicranum* sp. Suggesting turn to dryer conditions. In T3Sm, the C-E-S stage occurred directly over the mineral subsoil.


On contrary to asynchronous shift from C-E stage to C-E-S stage, the change to ombrotrophic vegetation community (S-E) with high proportion of sphagna appeared nearly simultaneously across all studied margins. This stage started between ca. 1870 and 1970 in all peat sections in transects 2 and 3 and similarly also in T1Sp. Only in T1Sm (1770 CE) and T1Si (1720 CE) the shift to S-E vegetation community stage occurred earlier. Currently S-E vegetation type is predominant across the transects.

A comparable successional pathway as in Lompolovuoma was detected from Syysjärvi study site in eastern Lapland (Fig. A1). A 1 cm thick ericaceous vegetation layer overlaid mineral soil, and this community was shortly replaced by a 2 cm thick C-E layer similar to the results found in Lompolovuoma. These stages lasted only ca. 15 years, respectively, after which C-E-S stage with some sphagna took over ca. 1970 CE. Above 3 cm thick C-E-S stage, the S-E stage mostly comprised by *Sphagnum capillifolium* that took over in ca. 1980 and has persisted ever since.


Different successional pathways were found from Salamajärvi and Patvinsuo peatland sites (Fig. A1). In Salamajärvi, there was no evidence of cyperaceous vegetation. Rather, the peat layers comprising of ericaceous vegetation with a small amount of *Sphagnum* mosses initiated directly on mineral subsoil in ca. 1830 CE in the margin of the Salamajärvi peatland. Afterwards, proportion of sphagna gradually started to increase and *Sphagnum* mosses became dominant ca. 1950 CE. Currently, *Sphagnum capillifolium* is the dominating moss species.

When peat formation started in Patvinsuo margin (Fig. A1) ca. 1850 CE, the initial vegetation consisted of C-E-S vegetation. At first, proportion of *Sphagnum* mosses started to increase, and ca. 1915 CE those were the dominant taxa. However, between ca. 1915 and 1950 CE *Sphagnum* mosses together with remains of *Cyperaceae* nearly disappeared and mostly ericaceous vegetation remained and supplemented by the presence of *Cenococcum sclerotia* that suggest dry mire margin conditions. However, towards present, the amount of *Sphagnum* mosses again increased and currently they form most of the coring site vegetation, with *Sphagnum russowii* being the most common species.

Transect 1



Transect 2



Transect 3




Figure 5. Fossil plant records (left) including undetected organic matter (UOM) and loss on ignition (LOI), bulk density (BD) and carbon and nitrogen contents and C/N ratio (right) for transect 3. Proportion of vegetation type and LOI in percentages (%), unit for bulk density is g/cm³. Carbon content (%), nitrogen content (%), and C:N ratio is available for profile T3Sp only.

3.4 Lateral expansion rates and vertical peat increment

The average rate of lateral expansion between dated peat profiles varied from 0.53 cm/year (T3Si to T3Sm) to 5.23 cm/year (T1Si to T1Sm). The median lateral expansion rate for all transects was 2.25 cm/year with interquartile range of 1.72 – 2.90 cm/year.

3.5 Vegetation community cover analysis

The total area of Lompolovuoma and Lompolojännkä peatland basin is 141.2 ha, of which 34 % is classified as a bog-type, 46 % as fen-type, and 21 % as tree-covered fen (Table 2). The area 25 meters from the peatland border is in total 43.9 ha and covers 31 % of the total peatland area. In these marginal areas, bog type constituted 44 %, fen-type 23 %, and tree-covered fens 33 % of the mire margin area (Fig. 6, Table 2). In Lompolovuoma basin, where our study transects were located, the coverage of bog-type in the peatland margin is 54 % while in adjacent peatland basin, Lompolojännkä, bog-type is covering smaller area, 35 %. On the contrary, higher coverage of fen-type is found in the margins of the Lompolojännkä (26 %) than in Lompolovuoma (20 %). Similarly, larger areas were covered by tree-covered fens in Lompolojännkä (39 %) than in Lompolovuoma (26 %).




Figure 6. Peatland margin vegetation communities. The area 25 meters from the peatland margin is divided into bog-type (brown), fen-type (yellow), and tree-covered fen type (green) in Lompolovuoma study basin and adjacent Lompolojänkki basin. In addition, location of the study transect peat cores (blue circles) and long cores (orange circles) are shown. This figure contains data from National Land Survey of Finland NLS Aerial photographs database.

Table 2. The vegetation class coverage and peatland area. Table shows the total area of the Lompolovuoma and Lompolojänkkä peatland basins, and proportion of 3 vegetation community classes in the peatland basins: Bog-type, fen-type, and tree-covered fen type. In addition, the total area of the 25 meter margin, and proportions of the vegetation community classes is shown. In the final 2 columns, the proportion of the vegetation community classes is shown individually for Lompolovuoma and Lompolojänkkä basins.

Vegetation class	Peatland		Peatland margins			
	Total area, ha	Total area, %	Margin area, ha	Margin area, %	Lompolo-vuoma	Lompolo-jänkkä
Bog-type	47.4	34 %	19.2	44 %	54 %	35 %
Fen-type	64.8	46 %	10.3	23 %	20 %	26 %
Tree-covered fen	29.0	21 %	14.4	33 %	26 %	39 %
Total area	141.2	100 %	43.9	100 %	100 %	100 %

3.6 Hydrological analyses

The simulated groundwater – surface water (GW-SW) exchange patterns for the current climatic and groundwater table (GWT) elevation change are shown in Fig. 7a and 7b, respectively. The calculated areas by contour group and peatland vegetation group are presented in Table 3 for the GW-SW exchange fluxes and in Table 4 for the changes in terms of GWT elevation.

In terms of exchange flux, the areas classified as fens indicate the dominance of the GW exfiltration over infiltration processes in the simulations. In contrast, the bog areas indicate more balance between infiltration and exfiltration processes with a slight prevalence of the infiltration area. The areas classified as treed fens show the dominance of infiltration. However, ~30 % of the total treed fen area is in the vicinity of the ditch network (the rightmost part of the peatland system), which impacted the peatland vegetation as indicated by aerial photos (National Land Survey of Finland, 2023). After excluding the drained areas from treed fens, the GW-SW exfiltration pattern is more balanced with a slight prevalence of exfiltration.

In terms of groundwater table elevation changes, the simulated drier climatic conditions have a mild impact on the areas classified as fens, with 59 % of the water table decreasing by less than 1cm and 89 % by less than 5 cm. In contrast, the areas classified as bog are more susceptible to GWT changes. They are characterised by significantly less extent of the areas with mild (less than 1 cm and 5 cm) table decrease (only 28 % and 70 % respectively) and a significant portion (30 %) with a substantial decline (more than 5 cm). The treed fen areas, excluding ditches, suggest that the water table decrease would be variable, with more GWT reduction than in the case of open fens but lower than in the case of bogs.




Figure 7a. The GW-SW exchange flux patterns from Lompolojankkä sub-basin averaged for summer 2017 representing the current climate conditions. Positive flux values indicate the locations of groundwater exfiltration and infiltration towards groundwater. 7b. The groundwater table elevation changes result from the drier climate conditions. Negative values indicate the groundwater level decrease and positive increase.

Table 3. Exchange flux contour areas by vegetation type. Treed fens (ditches excluded) exclude the open drainage areas.

Exchange flux (m/s)		Area (m ² / %)			
Min value	Max value	Fen	Bog	Treed fens	Treed fen (ditches excluded)
	<-1E-6	0 / 0 %	18 / 0 %	0 / 0 %	0 / 0 %
-1E-06	-1E-07	1471 / 2 %	2818 / 4 %	3130 / 4 %	2360 / 4 %
-1E-07	-1E-08	18822 / 30 %	34813 / 47 %	46693 / 55 %	26342 / 45 %
-1E-08	0.00	2936 / 5 %	3662 / 5 %	2549 / 3 %	2256 / 4 %
0.00	1E-08	3067 / 5 %	3603 / 5 %	2661 / 3 %	2388 / 4 %
1E-08	1E-07	23583 / 38 %	19196 / 26 %	12452 / 15 %	10989 / 19 %
1E-07	1E-06	12529 / 20 %	9859 / 13 %	16696 / 20 %	14106 / 24 %
1E-6>		83 / 0 %	85 / 0 %	555 / 1 %	552 / 1 %

Total negative flux	23229 / 37 %	41311 / 56 %	52373 / 62 %	30957 / 52 %
Total positive flux	39262 / 63 %	32743 / 44 %	32363 / 38 %	28035 / 48 %
Total flux	62491	74053	84736	58993

Table 4. Water table difference between baseline and drier climates by contour areas and vegetation type. Treed (ditches excluded) exclude the open drainage areas.

Water table decrease from the baseline model (m)		Area (m ² / %)			
Min value	Max value	Fen	Bog	Treed fens	Treed fen (ditches excluded)
-0.2	-0.5	15 / 0 %	642 / 1 %	8084 / 10 %	921 / 2 %
-0.1	-0.2	1496 / 2 %	7855 / 11 %	11539 / 14 %	4043 / 7 %
-0.05	-0.1	5531 / 9 %	13195 / 18 %	11111 / 13 %	6479 / 11 %
-0.01	-0.05	18608 / 30 %	31348 / 42 %	35656 / 42 %	31314 / 53 %
0	-0.01	36846 / 59 %	21040 / 28 %	18428 / 22 %	16317 / 28 %
Total		62495	74080	84818	59074

4. Discussion

We studied lateral expansion and vegetation succession of peatland margins in a subarctic Lompolovuoma fen in Finnish Lapland. Our results show that the studied margins in Lompolovuoma started to develop ca. 2000 years ago and the youngest basal age of 390 cal. BP was located still a few meters from the current forest edge. Peat initiation in the margins occurred in several independent loci that only later coalesced into a continuous peatland. The initial wet *Cyperaceae* – *Ericaceae* marginal communities inhabited the fen margins over the time periods reaching from a few centuries to a millennium and the following establishment of *Sphagnum* moss dominated communities was asynchronous. Starting from the end of the 1800th century, these margins shifted to a climax bog plant community characterized by hummock sphagna and ericaceous vegetation. This change occurred mostly over a remarkably short time, in a few decades. However, our data also showed that forest vegetation had on several occasions intruded to the already established peatland, suggesting an on-going “power-struggle” between peatland and forest ecosystems. The marginal “bog-type” vegetation currently covers roughly 50 % of the margins in the Lompolovuoma sub-basin, while in the adjacent sub-basin of Lompolojänkkä, only 35 % have reached the ombrotrophic stage. Our hydrological GW-SW model indicates that in the “fen-type” margins high water-tables are sustained even during dry climatic conditions, showing a resistance-potential to fen-to-bog transition.

4.1 Non-linear development of peatland margins in Lompolovuoma fen

The formation of Lompolovuoma peat margins investigated here began ca. 2200 years ago. This differs from the central part of the mire, where limnotelmatic Equisetum peat found at the bottom the long peat profiles suggest that peat formation initially occurred over water body. Similar to the results by Juselius-Rajamäki et al. (2023), these data

contradict the traditional perception that peatland expansion has ceased or markedly slowed down during the last 2000 years in Fennoscandia (Ruuhijärvi, 1983; Sjörs, 1983). Rather, the current findings suggest this presumption is due to under-representation of studies and sample collection from the mire marginal areas rather than an actual ceasing of lateral expansion (Kuhry and Turunen, 2006; Ruppel et al., 2013). In transects 2 and 3, the expansion of new peat surfaces occurred from individual miniature loci evidenced by the oldest basal ages found from the middle of the profiles, while in transect 1 the oldest basal age was acquired for the profile closest to the main mire (Fig. 2a-c). However, the basal age and the basal elevation of the T1Sp matches closely to the age and elevation of the oldest bottom age of transect 2, suggesting a relatively simultaneous initiation process.

The basal ages from the studied transects show that after the initial peat formation, the individually formed peat patches spread both downhill towards the main mire area, and uphill towards the adjacent forest. Only later, separate peat patches were connected to main mire basin. Such convergence of the multiple smaller loci to a single peatland mass has been reported both during the early Holocene (Almquist-Jacobson and Foster, 1995; Korhola, 1992, 1994; Mathijssen et al., 2014, 2017) and for more recently developed mire margins (Juselius-Rajamäki et al., 2023). However, the mechanisms behind the development of individual peat patches and the later convergence have received only little attention and remain unresolved (Noble et al., 1984).

In Lompolovuoma, peat initiation occurred in steep slopes on average exceeding 0.5° , a threshold known to restrict peat formation in more continental regions where availability of water is not excessive (Almquist-Jacobson and Foster, 1995; Loisel et al., 2013; Zhao et al., 2014). Thus, in the past, suitable conditions promoting the initiation of individual peat patches must have existed. The peat patches may have started to form in small topographical depressions that, although initially well-drained, may become impervious due to deposition of organic or fine inorganic matter, formation of hardpans in the Spodosol layer, or deposition of ash due to forest fires, creating favourable conditions for peat formation (Klinger, 1996; Le Stum-Boivin et al., 2019; Mallik et al., 1984; Noble et al., 1984; Rydin and Jeglum, 2013). No full-scale subsoil topography measurements were conducted, but field survey data did not reveal any clear depressions underlying any of the oldest peat profiles. Another scenario is, that under sufficiently humid conditions the peat formation began directly on the steep slopes, as suggested for southern Finland peatlands (Korhola, 1996). Climate reconstructions suggest wet climate phase prevailed in Lapland between 2500 and 2000 BP (Eronen et al., 1999; Luoto and Nevalainen, 2015), which may have promoted peat formation even in a relative steep slope, such as presented here.

The vertical growth of peat as a driving mechanism for lateral expansion has been traditionally linked to raised mires (Foster and Wright, 1990). However, although the shape of the Lompolovuoma surface has remained concave, the low hydraulic conductivity of saturated peat (Ingram, 1978; Rydin and Jeglum, 2013) combined with the large amounts of waters flowing from surrounding uphill areas, especially during the snow-melt period (Autio et al., 2023) could nevertheless cause flooding in suitable locations even if these locations were separated from the main mire body. Similarly, previous studies have shown that although no elevated mire centre exists, significant lateral expansion of

peatland has occurred (Almquist-Jacobson and Foster, 1995; Korhola, 1994, 1996; Korhola et al., 2010; Mathijssen et al., 2017), suggesting that even on flat or concave shaped peatland basins peat accumulation can lead to redistribution of waters towards mire margins. Low-severity fires in adjacent forests are also known to promote peatland lateral expansion, as the reduced tree-cover decreases evapotranspiration and promotes colonization of *Sphagnum* due to increased light availability (Le Stum-Boivin et al., 2019; Novenko et al., 2021). However, in our basal layers only a single charred wood piece used for dating was found, while microscopic analysis of the basal layers did not reveal any charcoal (Fig. A2, A3, A4). Thus, forest fires did not likely play an important role in the peat initiation in question.

4.2 Autogenic and allogenic drivers behind the plant community succession

The initial *Cyperaceae-Ericaceae*-dominated stage found in our study site is commonly present in the basal layers of the peatland margins in Finland (Juselius-Rajamäki et al., 2023; Mathijssen et al., 2017). On the other hand, many studies have shown that sphagna is frequently found in the first stages of the paludification process (Le Stum-Boivin et al., 2019; Noble et al., 1984; Rydin and Jeglum, 2013). This variation can also be seen in our comparison profiles, as the margin of Syysjärvi site shows similar development as in Lompolovuoma, while in the more southern Salamajärvi and Patvinsuo *Sphagnum* mosses were already present during the initial paludification (Fig. A1). The lack of *Sphagnum* mosses in Lompolovuoma margin during the peatland initiation is likely explained by the hydrological conditions. At the onset of the peatland expansion, the water table was likely fluctuating, as shown by the presence of both forest mosses and mycorrhizal fungi *Cenococcum geophilum* (van Geel, 1978) linked to dry conditions, and discovered cyperaceous vegetation, for example *Carex limosa* (Fig A3, A4) usually referring to relatively wet hydrological regime (Visser et al., 2000). *Sphagnum* mosses require constantly humid conditions for colonization (Fenton et al., 2007; Sundberg and Rydin, 2002), and even though they can tolerate limited periods of desiccation (Hájek and Vicharová, 2014), the prolonged fluctuating water sources in margins likely prevented early colonization by *Sphagnum* mosses. Only after the gradual development of mire conditions proper in the margins, was the spread of the peat mosses possible.

After the initial C-E stage, colonization of sphagna occurred asynchronously between 670 and 1770 CE. This gradual transition towards mixed *Cyperaceae-Ericaceae-Sphagnum* vegetation was likely moulded by autogenic development as changes driven by allogenic forcing would occur over large areas within a relatively short time span rather than over a millennium, as discussed in Välranta et al. (2017). This conclusion is supported by the fact that no evidence of forest fires was found in the peat profiles. Similarly, no such contemporary climate event has been detected which could promote large scale changes in vegetation and simultaneous spatial colonization of sphagna (Hanhijärvi et al., 2013; Linderholm et al., 2018; Luoto and Nevalainen, 2015). The comparison profiles from Patvinsuo and Salamajärvi also show gradual increase in the *Sphagnum* mosses, albeit at much shorter time scale than witnessed in Lompolovuoma, while in Syysjärvi the shift to *Sphagnum* moss dominance was extremely rapid (Fig. A1).

Although the decomposition of the bottom-most layers of peat prevented complete species-level identification of cyperaceous vegetation, increasing number of *Eriophorum vaginatum* remains were found in layers preceding the *Sphagnum* colonization (Fig. A2, A3, A4). Like *Sphagnum* mosses, tussock-forming cyperaceous vegetation may act as ‘ecosystem engineers’ (Palozzi and Lindo, 2017; Välranta et al., 2017) and the importance of *Eriophorum vaginatum* facilitating the fen-to-bog transition has been recognized in various studies (Hughes, 2000; Hughes and Dumayne-Peaty, 2002; Välranta et al., 2017). These species can alter local conditions, such as hydrology and acidity (Hughes, 2000; Hughes and Dumayne-Peaty, 2002) and produce litter highly resistant to decay, thus promoting peat accumulation (Wein, 1973). This accumulation process can be further amplified by presence of ericaceous vegetation (Hughes, 2000). Although in the studied margins the accumulation of the peat during C-E stage was modest, elevated surface combined with increased acidity seems to have been sufficient to create conditions suitable for establishment of *Sphagnum* species found in the studied margins, likely protecting them from alkaline waters and complete inundation known to impede colonization sphagna (Granath et al., 2010; Ruuhijärvi, 1983; Sallantausta, 2006).

After the colonization, *Sphagnum* mosses accelerate the change in local conditions (Rydin and Jeglum, 2013), increasing their competitiveness against other mire vegetation and leading to ombrotrophication. In some cases, this change can occur rapidly (Tahvanainen, 2011) and synchronously over wide area (Loisel and Bunsen, 2020), while also more gradual changes have been observed (Välranta et al., 2017). In Lompolovuoma, initially, the abundance of sphagna remained low after the first establishment, but a more dramatic change occurred towards the end of the 19th century, when *Sphagnum* mosses started dominating the margin plant communities, and most of the cyperaceous vegetation disappeared, leading to the current S-E vegetation stage. This change coincided with the end of the ‘Little Ice Age’ (LIA), when humid and cool climate conditions were followed by increasingly warm temperatures (Hanhijärvi et al., 2013). Similar post-LIA fen to bog shifts have been reported in previous studies where data is captured from central parts of the peatland (Granlund et al., 2022; Kolari et al., 2022; Loisel and Yu, 2013; Magnan et al., 2018; Piilo et al., 2019; Primeau and Garneau, 2021; Robitaille et al., 2021), while our results show similar recent changes occurring in the margins. Current results are supported by a study from adjacent Lompolojänkää basin showing a similar kind of recent vegetation shift in the margins (Kuuri-Riutta et al., 2024) and those of our comparison profile from Syysjärvi (Fig. A1). Thus, although aapa mires are generally described as having wet central parts and dryer margins, our results show that dryer margins supporting sphagna may have formed rather recently.

Although these recent fen-to-bog transitions have occurred during dry climatic conditions post-LIA, similar shift has also occurred during wet climate phases (Välranta et al., 2017), as the only requirement for the process is the separation of the peat surface from the groundwater supply (Hughes, 2000; Hughes and Barber, 2003). During wet climatic conditions, the accumulation of peat is promoted, rather high water-table levels are maintained, and the fen-to-bog transition leads to a bog pool and lawn communities (Hughes and Barber, 2003). On the other hand, dry climate conditions decrease the water-table, that enables the species with tolerance towards drought or fluctuating water-tables to out-compete other species (Hughes and Barber, 2004). In Lompolovuoma margins, hummock-forming *Sphagnum* species, especially *Sphagnum fuscum*, increased markedly during the ultimate shift to ombrotrophic bog conditions.

The final fen-to-bog transition in the studied mire margins appears to be caused by the drier and warmer climate, as only sporadic presence of non-hummock *Spaghna* was detected in the peat profiles (Fig. A2, A3, A4). Moreover, the most marginal peat profiles in transect 1 and transect 2, as well as in comparison profile from Patvinsuo, show that the peatland vegetation has been replaced completely by forest vegetation on several occasions. This suggests that peatland expansion may be reversed at least temporary.

However, based on the remote sensing data, similar ombrotrophication has not occurred across all margins in Lompolovuoma and adjacent Lompolojänkä basins. The ombrotrophic S-E stage can currently be found roughly in 50 % of the margins of the Lompolovuoma basin while this stage has been reached only in ca. 35 % in adjacent Lompolojänkä. Similarly, the central part of adjacent Lompolojänkä basin has shown no evidence of fen-to-bog transition (Mathijssen et al., 2014), but transition is ongoing in the margins (Kuuri-Riutta et al., 2024). Thus, it appears that for the transition from fen to bog to occur, certain prerequisites and conditions must be met. Our hydrological model, based on the Lompolojänkä basin, showed that while marginal fens were generally ground-water recipients, the bog-type vegetation acted preferentially as surface water infiltration areas. By decreasing the effective precipitation in the hydrological model to mimic dryer conditions, the highest levels of water table drawdown were found in the current “bog-type” margins, marking these locations more likely to suffer drying conditions. Although both the analysis of vegetation cover (Räsänen et al., 2021) and hydrological model (Autio et al., 2023) contain some degree of uncertainty, the application of the hydrological model over the marginal peatland types supports our hypothesis of drop in ground-water levels as a likely cause for the final shift towards ombrotrophic climax stage.

4.3 Implication for carbon balance and future trajectories of vegetation succession in aapa mire margins

It has been shown that in fen conditions climate forcing from peatland complex can remain positive (e.g., climate warming effect) for most of the development history due to high methane emissions and only after continuous carbon uptake and expansion of bog vegetation the climate forcing turns negative (Korhola et al, 1996; Mathijssen et al., 2017, 2022). For example, in adjacent Lompolojänkä basin the modelled climate warming effect persisted up to 2000 years (Mathijssen et al., 2014). Thus, it is likely that during the initial minerotrophic C-E stage lasting between 150 and 1250 years, the mire margin had a climate warming effect. Afterwards, a shift to decay-resistant *Sphagnum* vegetation, lower water table leading to reduced methane emissions and continuous carbon uptake would likely have the same effect. Decrease of the cyperaceous vegetation especially during the last ca. 100 years would have reduced the methane emissions even further (Bubier et al., 1993; Ward et al., 2013). Although our study did not include carbon balance calculations, the shift towards bog community on the studied margins suggest that under current conditions, the margins would likely proceed to have a climate cooling effect. However, drying trend detected in the European peatlands (Swindles et al., 2019) could also turn these locations to carbon sources, if sufficient moisture conditions are not retained (Zhang et al., 2020).

As this study and studies by Juselius-Rajamäki et al. (2023) and Kuuri-Riutta et al. (2024) show, new peatland areas are currently widely being formed in the mire margins all over subarctic and boreal zone under natural conditions. However, in many places this development has been blocked by the ditching of mire margins (Sallinen et al., 2019), while the widespread drying of peatland surfaces during the last ca. 300 years may suggest that detrimental climatic conditions for lateral expansion are forming (Swindles et al., 2019). In addition, as revealed by this study, the succession of mire margins even in the same peatland can differ, with some margins retaining their initial wet minerotrophic characteristics, while others develop to ombrotrophic bogs. Due to the opposite climate forcing, the effect of this recent mire expansion on the climate depends on the scope of different peatland types across new mire margins and their later development. The knowledge on the developing peatland margins and their plant community succession still remains scarce. As the lateral expansion of peatlands has had a significant effect on atmospheric greenhouse gas concentrations in the past (Korhola et al., 2010; Peng et al., 2024), we suggest that more studies across the northern peatland margins are needed to reveal the wider effect of this recent lateral peatland expansion on the global carbon budgets.

5 Conclusions

Our research shows that the studied mire margin in Lompolovuoma basin has continued to increase in area since ca. 2000 BP, but this development has not progressed linearly. Rather, the current mire margin has formed from several individual loci and via patches that have merged as the local hydrology has transformed suitable for peat formation. After the initial wet “fen-type” conditions, that persisted for markedly long period, colonization by *Sphagnum* mosses, the change to current “bog-type” conditions represents a remarkable swift shift. This change was driven by dryer climatic conditions following the LIA as shown by our hydrological model. However, not all margins in Lompolonvuoma and Lompolojänkää basins have shifted to “bog-type” communities suggesting that wetter “fen-types” are at least partially resistant to hydrologically driven regime shifts. This study shows that even on the basin-scale, peatland margins are highly heterogeneous systems, and this should be taken into account when assessing the effects of past and future lateral expansion trend on the peatland area and peatland carbon dynamics.

Comparison profiles






Figure A1. Fossil plant records (left) including undetected organic material (UOM), and loss on ignition (LOI) and bulk density (BD) for supplementary profiles. Proportion of vegetation type and LOI in percentages (%), unit for bulk density is g/cm³

Lompolovuoma transect 1

LVT1Sm



LVT1Si



LVT1Sp




Figure A2. Macrofossil data for peat profiles in transect 1. Unrecognized organic matter (UOM), and plant species or species group are presented as well as remains of mites, *Cenococcum*, and mineral content.

Figure A3. Macrofossil data for peat profiles in transect 2. Unrecognized organic matter (UOM), and plant species or species group are presented as well as remains of mites, *Cenococcum*, and mineral content.

Lompolovuoma transect 3




Figure A4. Macrofossil data for peat profiles in transect 3. Unrecognized organic matter (UOM), and plant species or species group are presented as well as remains of mites, *Cenococcum*, and mineral content.

7 Data availability statement

The data for peat properties, peat core locations and age-depth models are available on the figshare: <https://doi.org/10.6084/m9.figshare.25941493.v1>

8 Author contributions

T.-J-R and M.V. conceived the idea for the article. T.J-R, E.T., M.V., A.A., H.M., and P.A-A. collected the field data. T-J.R. and S.P. performed the macrofossil analysis. T.J-R. and S.S-P. conducted the ²¹⁰Pb-analysis. T.J-R. conducted the spatial analysis. A.A., H.M., and P.A-A. conducted hydrological modelling. T-J.R. created the initial draft for the manuscript. All authors contributed to the drafts and gave the final approval for publication.

9 Conflict of interest statement

The authors declare that they have no conflict of interest.

10 Acknowledgements and financial support

T.J-R. was funded by Tellervo ja Juuso Walden foundation, M.V., S.P. and T.V. received funding from - Research council of Finland project 338631 and 349193. Groundwater modelling and GPR dataset as part of University of Oulu activities were supported by Research Council of Finland ACWI project (project nro 316349), Freshwater Competence Centre (FWCC) and DIWA-flagship. We acknowledge the support from the Ministry of Transport and Communication through ICOS Finland and from the WetHorizons project (Horizon Europe GAP-101056848).

11 References

- Ali, A. A., Ghaleb, B., Garneau, M., Asnong, H., and Loisel, J.: Recent peat accumulation rates in minerotrophic peatlands of the Bay James region, Eastern Canada, inferred by ²¹⁰Pb and ¹³⁷Cs radiometric techniques, *Appl. Radiat. Isotopes*, 66(10), 1350–1358, <https://doi.org/10.1016/j.apradiso.2008.02.091>, 2008.
- Almquist-Jacobson, H., and Foster, D. R.: Toward an Integrated Model for Raised-Bog Development: Theory and Field Evidence, *Ecology*, 76(8), 2503–2516. 1995.
- Aquanty, *HydroGeoSphere Reference Manual*, 2015.
- Aquino-López, M. A., Blaauw, M., Christen, J. A., and Sanderson, N. K.: Bayesian Analysis of ²¹⁰Pb Dating, *J. Agr. Biol. Envir. S.*, 23(3), 317–333, <https://doi.org/10.1007/s13253-018-0328-7>, 2018.
- Autio, A., Ala-Aho, P., Rossi, P. M., Ronkanen, A. K., Aurela, M., Lohila, A., Korpelainen, P., Kumpula, T., Klöve, B., and Marttila, H.: Groundwater exfiltration pattern determination in the sub-arctic catchment using thermal imaging, stable water isotopes and fully-integrated groundwater-surface water modelling, *J. Hydrol.*, 626, 130342, <https://doi.org/10.1016/j.jhydrol.2023.130342>, 2023.

Blaauw, M., and Christen, J. A.: Flexible paleoclimate age-depth models using an autoregressive gamma process, *Bayesian Anal.*, 6(3), <https://doi.org/10.1214/11-ba618>, 2011.

Bubier, J., Costello, A., Moore, T. R., Roulet, N. T., and Savage, K.: Microtopography and methane flux in boreal peatlands, northern Ontario, Canada, *Can. J. Botany*, 71(8), 1056–1063, <https://doi.org/10.1139/b93-122>, 1993.

Eronen, M., Lindholm, M., Saastamoinen, S., and Zetterberg, P.: Variable Holocene climate, treeline dynamics and changes in natural environments in northern Finnish Lapland. *Chemosphere*, 1(4), 377–387, [https://doi.org/10.1016/S1465-9972\(99\)00042-2](https://doi.org/10.1016/S1465-9972(99)00042-2), 1999.

ESRI: ArcGIS Pro (3.1), 2023.

Evans, C. D., Peacock, M., Baird, A. J., Artz, R. R. E., Burden, A., Callaghan, N., Chapman, P. J., Cooper, H. M., Coyle, M., Craig, E., Cumming, A., Dixon, S., Gauci, V., Grayson, R. P., Helfter, C., Heppell, C. M., Holden, J., Jones, D. L., Kaduk, J., Levy, P., Matthews, R., McNamara, N. P., Misselbrook, T., Oakley, S., Page, S. E., Rayment, M., Ridley, L. M., Stanley, K. M., Williamson, J. L., Worrall, F. and Morrison, R.: Overriding water table control on managed peatland greenhouse gas emissions, *Nature*, 593(7860), 548–552, <https://doi.org/10.1038/s41586-021-03523-1>, 2021.

Fenton, N. J., Béland, C., De Blois, S., and Bergeron, Y.: Sphagnum establishment and expansion in black spruce (*Picea mariana*) boreal forests, *Can. J. Botany*, 85(1), 43–50, <https://doi.org/10.1139/B06-148>, 2007.

Flynn, W. W.: The determination of low levels of Polonium-210 in environmental materials, *Anal. Chim. Acta*, 43, 221–227, 1968.

Foster, D. R., and King, G. A.: Landscape Features, Vegetation and Developmental History of a Patterned Fen in South-Eastern Labrador, Canada, *J. Ecol.*, 72(1), 115–143, <https://doi.org/10.2307/2260009>, 1984.

Foster, D. R., and Wright, H. E.: Role of ecosystem development and climate change in bog formation in central Sweden, *Ecology*, 71(2), 450–463, <https://doi.org/10.2307/1940300>, 1990.

Frolking, S., and Roulet, N. T.: Holocene radiative forcing impact of northern peatland carbon accumulation and methane emissions, *Glob. Change Biol.*, 13(5), 1079–1088, <https://doi.org/10.1111/j.1365-2486.2007.01339.x>, 2007.

Goud, E. M., Watt, C., and Moore, T. R.: Plant community composition along a peatland margin follows alternate successional pathways after hydrologic disturbance. *Acta Oecol.*, 91, 65–72, <https://doi.org/10.1016/j.actao.2018.06.006>, 2018.

Granath, G., Strengbom, J., and Rydin, H.: Rapid ecosystem shifts in peatlands: Linking plant physiology and succession, *Ecology*, 91(10), 3047–3056, <https://doi.org/10.1890/09-2267.1>, 2010.

Granlund, L., Vesakoski, V., Sallinen, A., Kolari, T. H. M., Wolff, F., and Tahvanainen, T.: Recent Lateral Expansion of Sphagnum Bogs Over Central Fen Areas of Boreal Aapa Mire Complexes, *Ecosystems*, 25(7), 1455–1475, <https://doi.org/10.1007/s10021-021-00726-5>, 2022.

Grimm, E. C.: TILIA and TILIAGRAPH Software, 1991.

Hájek, T., and Vicherová, E.: Desiccation tolerance of Sphagnum revisited: A puzzle resolved, *Plant Biol.*, 16(4), 765–773, <https://doi.org/10.1111/plb.12126>, 2014.

- Hanhijärvi, S., Tingley, M. P., and Korhola, A.: Pairwise comparisons to reconstruct mean temperature in the Arctic Atlantic Region over the last 2,000 years, *Clim. Dynam.*, 41, 2039–2060, <https://doi.org/10.1007/s00382-013-1701-4>, 2013.
- Heiri, O., Lotter, A. F., and Lemcke, G.: Loss on ignition as a method for estimating organic and carbonate content in sediments: Reproducibility and comparability of results, *J. Paleolimnol.*, 25(1), 101–110, <https://doi.org/10.1023/A:1008119611481>, 2001.
- Helama, S., Jones, P. D., and Briffa, K. R.: Dark Ages Cold Period: A literature review and directions for future research, *Holocene*, 27(10), 1600–1606, <https://doi.org/10.1177/0959683617693898>, 2017.
- Howie, S. A., and Meerveld, I. T. Van.: The essential role of the lag in raised bog function and restoration: A review, *Wetlands*, 31(3), 613–622, <https://doi.org/10.1007/s13157-011-0168-5>, 2011.
- Hua, Q., Turnbull, J. C., Santos, G. M., Rakowski, A. Z., Ancapichún, S., De Pol-Holz, R., Hammer, S., Lehman, S. J., Levin, I., Miller, J. B., Palmer, J. G., and Turney, C. S. M.: Atmospheric Radiocarbon for the Period 1950–2019, *Radiocarbon*, 64(4), 723–745, <https://doi.org/10.1017/RDC.2021.95>, 2022.
- Hughes, P. D. M.: A reappraisal of the mechanisms leading to ombrotrophy in British raised mires, *Ecol. Lett.*, 3, 7–9, <https://doi.org/10.1046/j.1461-0248.2000.00118.x>, 2000.
- Hughes, P. D. M., and Barber, K. E.: Mire development across the fen-bog transition on the Teifi floodplain at Tregaron Bog, Ceredigion, Wales, and a comparison with 13 other raised bogs, *J. Ecol.*, 91(2), 253–264, <https://doi.org/10.1046/j.1365-2745.2003.00762.x>, 2003.
- Hughes, P. D. M., and Barber, K. E.: Contrasting pathways to ombrotrophy in three raised bogs from Ireland and Cumbria, England, *Holocene*, 14(1), 65–77, <https://doi.org/10.1191/0959683604hl690rp>, 2004.
- Hughes, P. D. M., and Dumayne-Peaty, L.: Testing Theories of Mire Development Using Multiple Successions at Crymlyn Bog, West Glamorgan, South Wales, UK, *J. Ecol.*, 90(3), 456–471, 2002.
- Ingram, H. A. P.: Soil Layers in Mires: Function and Terminology, *J. Soil Sci.*, 29(2), 224–227, <https://doi.org/10.1111/j.1365-2389.1978.tb02053.x>, 1978.
- Juggins, S.: User Guide: C2 Software for ecological and palaeoecological data analysis and visualisation, User guide Version 1.5 (vols. 1–73), University of Newcastle, <http://www.staff.ncl.ac.uk/stephen.juggins>, 2007.
- Juselius-Rajamäki, T., Välianta, M., and Korhola, A.: The ongoing lateral expansion of peatlands in Finland, *Glob. Change Biol.*, 29, 7173–7191, <https://doi.org/10.1111/gcb.16988>, 2023.
- Juutinen, S., Välianta, M., Kuutti, V., Laine, A. M., Virtanen, T., Seppä, H., Weckström, J., and Tuittila, E. S.: Short-term and long-term carbon dynamics in a northern peatland-stream-lake continuum: A catchment approach, *J. Geophys. Res.-Biogeo.*, 118, 171–183, <https://doi.org/10.1002/jgrg.20028>, 2013.
- Kauranen, P., and Miettinen, J. K.: 210Po and 210Pb in environmental samples in Finland, in: Radioecological concentration processes, Proceedings of an International Symposium Held in Stockholm 25–29 April, 1966.
- Klinger, L. F.: Coupling of Soils and Vegetation in Peatland Succession, *Arctic Alpine Res.*, 28(3), 380–387, 1996.
- Kolari, T. H. M., Sallinen, A., Wolff, F., Kumpula, T., Tolonen, K., and Tahvanainen, T.: Ongoing Fen–Bog Transition in a Boreal Aapa Mire Inferred from Repeated Field Sampling, Aerial Images, and Landsat Data, *Ecosystems*, 25, 1166–1188, <https://doi.org/10.1007/s10021-021-00708-7>, 2022.

- Korhola, A.: Mire Induction, ecosystem dynamics and lateral expansion on raised bogs in the southern coastal area of Finland, *Fennia*, 170(2), 25–94, 1992.
- Korhola, A.: Radiocarbon Evidence for Rates of Lateral Expansion in Raised Mires in Southern Finland, *Quaternary Res.*, 42, 299–307, <https://doi.org/10.1006/qres.1994.1080>, 1994.
- Korhola, A.: Holocene climatic variations in southern Finland reconstructed from peat-initiation data, *Holocene*, 5, 43–58, <https://doi.org/10.1177/095968369500500106>, 1995.
- Korhola, A.: Initiation of a sloping mire complex in southwestern Finland: Autogenic versus allogenic controls, *Écoscience*, 3(2), 216–222, <https://doi.org/10.1080/11956860.1996.11682334>, 1996.
- Korhola, A., Alm, J., Tolonen, J., Turunen, J., and Jungner, H.: Three-dimensional reconstruction of carbon accumulation and CH₄ emission during nine millenia in a raised mire, *J. Quaternary Sci.*, 11(2), 161–165., 1996.
- Korhola, A., Ruppel, M., Seppä, H., Välranta, M., Virtanen, T., and Weckström, J.: The importance of northern peatland expansion to the late-Holocene rise of atmospheric methane, *Quaternary Sci. Rev.*, 29(5–6), 611–617, <https://doi.org/10.1016/j.quascirev.2009.12.010>, 2010.
- Kou, D., Virtanen, T., Treat, C. C., Tuovinen, J. P., Räsänen, A., Juutinen, S., Mikola, J., Aurela, M., Heiskanen, L., Heikkilä, M., Weckström, J., Juselius, T., Piilo, S. R., Deng, J., Zhang, Y., Chaudhary, N., Huang, C., Välranta, M., Biasi, C., Liu, X., Guo, M., Zhuang, Q., Korhola, A. and Shurpali, N. J.: Peatland Heterogeneity Impacts on Regional Carbon Flux and Its Radiative Effect Within a Boreal Landscape, *J. Geophys. Res.-Biogeo.*, 127(9), <https://doi.org/10.1029/2021JG006774>, 2022.
- Kuhry, P.: The Role of Fire in the Development of Sphagnum-Dominated Peatlands in Western Boreal Canada, *J. Ecol.*, 82(4), 899–910, 1994.
- Kuhry, P., and Turunen, J.: The Postglacial Development of Boreal and Subarctic Peatlands, in: *Boreal peatland ecosystems. Ecological studies*, vol. 188, edited by: R. K. Wieder, D. Vitt, and R. B. Jackson, Springer, Berlin Heidelberg, Germany, 25–46, https://doi.org/10.1007/978-3-540-31913-9_3, 2006.
- Kuuri-Riutta, O., Pilkama, E., Salminen-Paatero, S., Vögeli, C., Mitchell, E. A. D., Lohila, A., Tuittila, E. S., and Välranta, M.: Recent hummock establishment in the margin of a subarctic fen, Finnish Lapland, *Boreas*, <https://doi.org/10.1111/bor.12651>, 2024.
- Lacourse, T., Adeleye, M. A., and Stewart, J. R.: Peatland formation , succession and carbon accumulation at a mid-elevation poor fen in Pacific Canada, *The Holocene*, 29(11), 1694–1707, <https://doi.org/10.1177/0959683619862041>, 2019.
- Lai, D. Y. F.: Methane Dynamics in Northern Peatlands: A Review, *Pedosphere*, 19(4), 409–421, [https://doi.org/10.1016/S1002-0160\(09\)00003-4](https://doi.org/10.1016/S1002-0160(09)00003-4), 2009.
- Laine, J., Vasander, H., Hotanen, J.-P., Nousiainen, H., Saarinen, M., and Penttilä, T.: *Suotyyppit ja turvekankaat – kasvupaikkaopas*, Tapio, 2018.
- Laitinen, J., Rehell, S., and Huttunen, A.: Vegetation-related hydrotopographic and hydrologic classification for aapa mires (Hirvisuo, Finland), *Ann. Bot. Fenn.*, 42(2), 107–121, <https://www.jstor.org/stable/23726854>, 2005.
- Laitinen, J., Rehell, S., Huttunen, A., Tahvanainen, T., Heikkilä, R., and Lindholm, T.: Mire systems in Finland - Special view to aapa mires and their water-flow pattern, *Suo*, 58(1), 1–26, 2007.

- Le Stum-Boivin, É., Magnan, G., Garneau, M., Fenton, N. J., Grondin, P., and Bergeron, Y.: Spatiotemporal evolution of paludification associated with autogenic and allogenic factors in the black spruce-moss boreal forest of Québec, Canada, *Quaternary Res.*, 91(2), 520–532, <https://doi.org/10.1017/qua.2018.101>, 2019.
- Linderholm, H. W., Nicolle, M., Francus, P., Gajewski, K., Helama, S., Korhola, A., Solomina, O., Yu, Z., Zhang, P., D’Andrea, W. J., Debret, M., Divine, D. V., Gunnarson, B. E., Loader, N. J., Massei, N., Seftigen, K., Thomas, E. K., Werner, J., Andersson, S., Berntsson, A., Luoto, T. P., Nevalainen, L., Saarni, S. and Väliranta, M.: Arctic hydroclimate variability during the last 2000 years: Current understanding and research challenges, *Clim. Past*, 14(4), 473–514, <https://doi.org/10.5194/cp-14-473-2018>, 2018.
- Loisel, J., and Bunsen, M.: Abrupt Fen-Bog Transition Across Southern Patagonia: Timing, Causes, and Impacts on Carbon Sequestration, *Front. Ecol. Evol.*, 8(August), 1–19, <https://doi.org/10.3389/fevo.2020.00273>, 2020.
- Loisel, J., and Yu, Z.: Recent acceleration of carbon accumulation in a boreal peatland, south central Alaska, *J. Geophys. Res.-Biogeo.*, 118(1), 41–53, <https://doi.org/10.1029/2012JG001978>, 2013.
- Loisel, J., Yu, Z., Parsekian, A., Nolan, J., and Slater, L.: Quantifying landscape morphology influence on peatland lateral expansion using ground-penetrating radar (GPR) and peat core analysis, *J. Geophys. Res.-Biogeo.*, 118(2), 373–384, <https://doi.org/10.1002/jgrg.20029>, 2013.
- Luoto, T. P., and Nevalainen, L.: Late Holocene precipitation and temperature changes in Northern Europe linked with North Atlantic forcing, *Clim. Res.*, 66, 37–48, <https://doi.org/10.3354/cr01331>, 2015.
- Magnan, G., van Bellen, S., Davies, L., Froese, D., Garneau, M., Mullan-Boudreua, G., Zacccone, C., and Shotyk, W.: Impact of the Little Ice Age cooling and 20th century climate change on peatland vegetation dynamics in central and northern Alberta using a multi-proxy approach and high-resolution peat chronologies, *Quaternary Sci. Rev.*, 185, 230–243, 2018.
- Mäkilä, M., and Moisanen, M.: Holocene lateral expansion and carbon accumulation of Luovuoma, a northern fen in Finnish Lapland, *Boreas*, 36(2), 198–210, <https://doi.org/10.1080/03009480600994460>, 2007.
- Mäkilä, M., Saarnisto, M., and Kankainen, T.: Aapa mires as a carbon sink and source during the Holocene, *J. Ecol.*, 89(4), 589–599, <https://doi.org/10.1046/j.0022-0477.2001.00586.x>, 2001.
- Mallik, A. U., Gimingham, C. H., and Rahman, A. A.: Ecological Effects of Heather Burning: I. Water Infiltration, Moisture Retention and Porosity of Surface Soil, *J. Ecol.*, 72(3), 767–776, <https://doi.org/10.2307/2259530>, 1984.
- Marttila, H., Lohila, A., Ala-Aho, P., Noor, K., Welker, J. M., Croghan, D., Mustonen, K., Meriö, L. J., Autio, A., Muhic, F., Bailey, H., Aurela, M., Vuorenmaa, J., Penttilä, T., Hyöky, V., Klein, E., Kuzmin, A., Korpelainen, P., Kumpula, T., Rauhala, A. and Kløve, B.: Subarctic catchment water storage and carbon cycling – Leading the way for future studies using integrated datasets at Pallas, Finland, *Hydrol. Process.*, 35(9), <https://doi.org/10.1002/hyp.14350>, 2021.
- Mathijssen, P. J. H., Kähkölä, N., Tuovinen, J. P., Lohila, A., Minkkinen, K., Laurila, T., and Väliranta, M.: Lateral expansion and carbon exchange of a boreal peatland in Finland resulting in 7000 years of positive radiative forcing, *J. Geophys. Res.-Biogeo.*, 122(3), 562–577, <https://doi.org/10.1002/2016JG003749>, 2017.

Mathijssen, P. J. H., Tuovinen, J. P., Lohila, A., Aurela, M., Juutinen, S., Laurila, T., Niemelä, E., Tuittila, E. S., and Väiranta, M. Development, carbon accumulation, and radiative forcing of a subarctic fen over the Holocene, *Holocene*, 24(9), 1156–1166, <https://doi.org/10.1177/0959683614538072>, 2014.

Mathijssen, P. J. H., Tuovinen, J. P., Lohila, A., Väiranta, M., and Tuittila, E. S.: Identifying main uncertainties in estimating past and present radiative forcing of peatlands, *Glob. Change Biol.*, 28(13), 4069–4084, <https://doi.org/10.1111/gcb.16189>, 2022.

Mathijssen, P. J. H., Väiranta, M., Korrensalo, A., Alekseychik, P., Vesala, T., Rinne, J., and Tuittila, E. S.: Reconstruction of Holocene carbon dynamics in a large boreal peatland complex, southern Finland, *Quaternary Sci. Rev.*, 142, 1–15, <https://doi.org/10.1016/j.quascirev.2016.04.013>, 2016.

Mauquoy, D., Hughes, P. D. M., Mauquoy, D., Hughes, P. D. M., and Van Geel, B.: A protocol for plant macrofossil analysis of peat deposits, *Mires Peat*, 7, 1-5, 2010.

National Land Survey of Finland.: *Aerial photo V4134*, <https://asiointi.maanmittauslaitos.fi/karttapaiikka/tiedostopalvelu/ortoilmakuva>, 2023.

Noble, M., Lawrence, D., and Streveler, G.: Sphagnum Invasion beneath an Evergreen Forest Canopy in Southeastern Alaska, *The Bryologist*, 87(2), 119–127, 1984.

Novenko, E. Y., Mazei, N. G., Kupriyanov, D. A., Kusilman, M. V., and Olchev, A. V.: Peatland initiation in Central European Russia during the Holocene: Effect of climate conditions and fires, *Holocene*, 31(4), 545–555, <https://doi.org/10.1177/0959683620981709>, 2021.

Palozzi, J. E., and Lindo, Z.: Boreal peat properties link to plant functional traits of ecosystem engineers, *Plant Soil*, 418(1–2), 277–291, <https://doi.org/10.1007/s11104-017-3291-0>, 2017.

Peng, H., Nijp, Jelmer, J., Ratcliffe, J. L., Li, C., Hong, B., Lidberg, W., Zeng, M., Mauquoy, D., Bishop, K., and Nilsson, M. B.: Climatic controls on the dynamic lateral expansion of northern peatlands and its potential implication for the ‘anomalous’ atmospheric CH₄ rise since the mid-Holocene, *Sci. Total Environ.*, 908(7), 168450, <https://doi.org/10.1016/j.scitotenv.2023.168450>, 2024.

Peregon, A., Uchida, M., and Yamagata, Y.: Lateral extension in Sphagnum mires along the southern margin of the boreal region, Western Siberia, *Environ. Res. Lett.*, 4(4), <https://doi.org/10.1088/1748-9326/4/4/045028>, 2009.

Piilo, S. R., Zhang, H., Garneau, M., Gallego-Sala, A., Amesbury, M. J., and Väiranta, M. M.: Recent peat and carbon accumulation following the Little Ice Age in northwestern Québec, Canada, *Environ. Res. Lett.*, 14(7), <https://doi.org/10.1088/1748-9326/ab11ec>, 2019.

Primeau, G., and Garneau, M.: Carbon accumulation in peatlands along a boreal to subarctic transect in eastern Canada, *Holocene*, 31(5), 858–869, <https://doi.org/10.1177/0959683620988031>, 2021.

Quik, C., Palstra, S. W. L., van Beek, R., van der Velde, Y., Candel, J. H. J., van der Linden, M., Kubiak-Martens, L., Swindles, G. T., Makaske, B., and Wallinga, J.: Dating basal peat: The geochronology of peat initiation revisited, *Quat. Geochronol.*, 72(March), 101278, <https://doi.org/10.1016/j.quageo.2022.101278>, 2022.

R Core Team.: *R: A language and environment for statistical computing* (4.2.2), R Foundation for Statistical Computing, 2023.

- Räsänen, A., Manninen, T., Korkiakoski, M., Lohila, A., and Virtanen, T.: Predicting catchment-scale methane fluxes with multi-source remote sensing, *Landscape Ecol.*, 36(4), 1177–1195, <https://doi.org/10.1007/s10980-021-01194-x>, 2021.
- Reimer, P. J., Austin, W. E. N., Bard, E., Bayliss, A., Blackwell, P. G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Hajdas, I., Heaton, T. J., Hogg, A. G., Hughen, K. A., Kromer, B., Manning, S. W., Muscheler, R., Palmer, J. G., Pearson, C., van der Plicht, J., Weimer, R. W., Richards, D. A., Scott, M. E., Southon, J. R., Turney, C. S. M., Wacker, L., Adolphi, F., Büntgen, U., Capano, M., Fahrni, S. M., Fogtmann-Schulz, A., Friedrich, R., Köhler, P., Kudsk, S., Miyake, F., Olsen, J., Reinig, F., Sakamoto, M., Sookdeo, A. and Talamo, S.: The IntCal20 Northern Hemisphere Radiocarbon Age Calibration Curve (0–55 cal kBP), *Radiocarbon*, 62(4), 725–757, <https://doi.org/10.1017/RDC.2020.41>, 2020.
- Robitaille, M., Garneau, M., van Bellen, S., and Sanderson, N. K.: Long-term and recent ecohydrological dynamics of patterned peatlands in north-central Quebec (Canada), *Holocene*, 31(5), 844–857, <https://doi.org/10.1177/0959683620988051>, 2021.
- Ruppel, M., Välijärvi, M., Virtanen, T., and Korhola, A.: Postglacial spatiotemporal peatland initiation and lateral expansion dynamics in North America and northern Europe, *Holocene*, 23(11), 1596–1606, <https://doi.org/10.1177/0959683613499053>, 2013.
- Ruuhijärvi, R.: Finnish mire types and their regional distribution, in: *Ecosystems of the world (Vol. 4B)*, edited by: A. J. P. Gore, Elsevier, Amsterdam, The Netherlands, 47–67. 1983.
- Rydin, H., and Jeglum, J. K. (Eds.): *The Biology of Peatlands*, Oxford University Press, New York, United States of America, <https://doi.org/10.1093/acprof:osobl/9780199602995.001.0001>, 2013.
- Sallantausta, T.: Mire ecohydrology in Finland, in: *Finland – Land of Mires*, edited by T. Lindholm and R. Heikkilä, Finnish Environmental Institute, Vammalan kirjapaino Oy, Vammala, Finland, 105–108, 2006.
- Sallinen, A., Akanegbu, J., Marttila, H., and Tahvanainen, T.: Recent and future hydrological trends of aapa mires across the boreal climate gradient, *J. Hydrol.*, 617, 129022, <https://doi.org/10.1016/j.jhydrol.2022.129022>, 2023.
- Sallinen, A., Tuominen, S., Kumpula, T., and Tahvanainen, T.: Undrained peatland areas disturbed by surrounding drainage: A large scale GIS analysis in Finland with a special focus on Aapa mires, *Mires Peat*, 24, 1–22, <https://doi.org/10.19189/MaP.2018.AJB.391>, 2019.
- Sanderson, N. K.: *Patterns and Drivers of Recent Peatland Carbon Accumulation in Northeastern Canada*, University of Exeter, UK, 2016.
- Schaffhauser, A., Payette, S., Garneau, M., and Robert, É. C.: Soil paludification and Sphagnum bog initiation: the influence of indurated podzolic soil and fire, *Boreas*, 46(3), 428–441, <https://doi.org/10.1111/bor.12200>, 2017.
- Seppä, H.: Mires of Finland: Regional and local controls of vegetation, landforms, and long-term dynamics, *Fennia*, 180(1–2), 43–60, 2002.
- Simard, M., Lecomte, N., Bergeron, Y., Bernier, P. Y., and Paré, D.: Forest productivity decline caused by successional paludification of boreal soils, *Ecol. Appl.*, 17(6), 1619–1637, <https://doi.org/10.1890/06-1795.1>, 2007.

- Sjörs, H.: Mires of Sweden, in: *Ecosystems of the world 4B*, edited by: Gore, A. J. P., Elsevier, Amsterdam, The Netherlands, 69–94, 1983.
- Sundberg, S., and Rydin, H.: Habitat requirements for establishment of *Sphagnum* from spores, *J. Ecol.*, 90(2), 268–278, <https://doi.org/10.1046/j.1365-2745.2001.00653.x>, 2002.
- Swindles, G. T., Morris, P. J., Mullan, D. J., Payne, R. J., Roland, T. P., Amesbury, M. J., Lamentowicz, M., Turner, T. E., Gallego-Sala, A., Sim, T., Barr, I. D., Blaauw, M., Blundell, A., Chambers, F. M., Charman, D. J., Feurdean, A., Galloway, J. M., Galka, M., Green, S. M., Kajukalo, K., Karofeld, E., Korhola, A., Lamentowicz, L., Langdon, P., Marcisz, K., Mauquoy, D., Mazei, Y. A., McKeown, M. M., Mitchell, E. A. D., Novenko, E., Plunkett, G., Roe, H. M., Schoning, K., Sillasoo, Ü., Tsyganov, A. N., van der Linden, M., Väliranta, M. and Warner, B.: Widespread drying of European peatlands in recent centuries, *Nat. Geosci.*, 12(11), 922–928, <https://doi.org/10.1038/s41561-019-0462-z>, 2019.
- Tahvanainen, T.: Abrupt ombrotrophication of a boreal aapa mire triggered by hydrological disturbance in the catchment, *J. Ecol.*, 99(2), 404–415, <https://doi.org/10.1111/j.1365-2745.2010.01778.x>, 2011.
- Väliranta, M., Korhola, A., Seppä, H., Tuittila, E. S., Sarmaja-Korjonen, K., Laine, J., and Alm, J.: High-resolution reconstruction of wetness dynamics in a southern boreal raised bog, Finland, during the late Holocene: A quantitative approach, *Holocene*, 17(8), 1093–1107, <https://doi.org/10.1177/0959683607082550>, 2007.
- Väliranta, M., Salojärvi, N., Vuorsalo, A., Juutinen, S., Korhola, A., Luoto, M., and Tuittila, E. S.: Holocene fen–bog transitions, current status in Finland and future perspectives, *Holocene*, 27(5), 752–764, <https://doi.org/10.1177/0959683616670471>, 2017.
- van Geel, B.: A Palaeoecological study of Holocene peat bog sections in Germany and the Netherlands, *Rev. Palaeobot. Palyno.*, 25, 1–120, <https://doi.org/10.2307/1216527>, 1978.
- Visser, E. J. W., Bogemann, G. M., Van de Steeg, H. M., Pierik, R., and Blom, C. W. P. M.: Flooding tolerance of *Carex* species in relation to field distribution and aerenchyma formation, *New Phytol.*, 148(1), 93–103, <https://doi.org/10.1046/j.1469-8137.2000.00742.x>, 2000.
- Ward, S. E., Ostle, N. J., Oakley, S., Quirk, H., Henrys, P. A., and Bardgett, R. D.: Warming effects on greenhouse gas fluxes in peatlands are modulated by vegetation composition, *Ecol. Lett.*, 16(10), 1285–1293, <https://doi.org/10.1111/ele.12167>, 2013.
- Wein, R. W.: *Eriophorum vaginatum* L, *J. Ecol.*, 61(2), 601–615, 1973.
- Zhang, H., Väliranta, M., Piilo, S., Amesbury, M. J., Aquino-López, M. A., Roland, T. P., Salminen-Paatero, S., Paatero, J., Lohila, A., and Tuittila, E. S.: Decreased carbon accumulation feedback driven by climate-induced drying of two southern boreal bogs over recent centuries, *Glob. Change Biol.*, 26(4), 2435–2448, <https://doi.org/10.1111/gcb.15005>, 2020.
- Zhao, Y., Tang, Y., Yu, Z., Li, H., Yang, B., Zhao, W., Li, F., and Li, Q.: Holocene peatland initiation, lateral expansion, and carbon dynamics in the Zoige Basin of the eastern Tibetan Plateau, *Holocene*, 24(9), 1137–1145, <https://doi.org/10.1177/0959683614538077>, 2014.