



Finely-resolved along-track wave attenuation estimates in the Antarctic marginal ice zone from ICESat-2

Joey J. Voermans¹, Alexander D. Fraser^{2,3}, Jill Brouwer^{2,3,*}, Michael H. Meylan⁴, Qingxiang Liu^{5,1}, and Alexander V. Babanin¹

¹Department of Infrastructure Engineering, University of Melbourne, Parkville, Victoria, Australia

²Australian Antarctic Program Partnership, Institute for Marine and Antarctic Studies, University of Tasmania, nipaluna / Hobart, 7001 Tasmania, Australia

³Institute for Marine and Antarctic Studies, University of Tasmania, nipaluna / Hobart, 7001 Tasmania, Australia

⁴School of Information and Physical Sciences, University of Newcastle, Callaghan, 2308 NSW, Australia

⁵Frontier Science Center for Deep Ocean Multispheres and Earth System (FDOMES) and Physical Oceanography Laboratory, Ocean University of China, Qingdao, China

*Now at NCM Engineering and Technology, CSIRO, Australia

Correspondence: Joey J. Voermans (jvoermans@unimelb.edu.au)

Abstract. Advances in our modelling capacity of wave-ice interactions are hindered by the limited availability of wave observations in sea ice and, specifically, under a broad range of wave and sea ice conditions. Satellite remote sensing provides opportunities to vastly expand the observational dataset of waves in sea ice and the study of wave-ice interactions. Specifically, Brouwer et al. (2022) demonstrated a clear reduction of observed wave energy into the Antarctic Marginal Ice Zone (MIZ) as derived from ICESat-2 observations. Here, we build upon the work of Brouwer et al. (2022) to estimate the wave attenuation rate in the Antarctic MIZ under a wide variety of sea ice conditions. Overall statistics of the observations reveal a linear increase in the wave attenuation rate with relative distance into the MIZ, implying that the wave energy in the MIZ scales as $\sim \exp(\beta x^2 \dots)$, where β is a frequency-dependent attenuation coefficient. Attenuation rates are well-sorted with wave frequency, where highest attenuation rates are observed for the shortest waves. We find that both the magnitude and frequency dependence of the ICESat-2 estimated wave attenuation rates are consistent with in situ observations. We further highlight that the misalignment between the incident wave direction and the measurement transect, and the inhomogeneity of the ice pack may lead to significant local fluctuations and negative values in the estimated wave attenuation rate when evaluating individual transects. The strong dependence of the overall statistics of the wave attenuation rate on the wave frequency and the relative distance into the MIZ alone provides significant opportunities in modelling wave-ice interactions in the Antarctic environment at global and climate scales, as it does not depend on system variables that are not straightforward to measure, retrieve or simulate at such large scales.



1 Introduction

The Antarctic Marginal Ice Zone (MIZ), the region separating the Southern Ocean from the consolidated pack ice, represents an interface of intense air-sea-ice interactions (e.g., Häkkinen, 1986; Weeks, 2010; Squire, 2020; Bennetts et al., 2024). Ocean waves are a critical mediator between the ocean, ice and atmosphere in the MIZ due to their capacity to break the ice (e.g., Kohout and Meylan, 2008; Dumont et al., 2011; Voermans et al., 2020), and, in doing so, waves can rapidly alter the sea ice morphology, its mobility, and, as a consequence, modulate the air-sea-ice fluxes in the MIZ (Williams et al., 2013; Collins III et al., 2015; Boutin et al., 2020; Li et al., 2021) such as the lateral melt of floes and thus the seasonal cycle during ice meltback (Bateson et al., 2022).

The spatial extent to which waves can impact the ice cover, and thus the associated MIZ dynamics, is in large part determined by the amount of wave energy that propagates through the ice, which can, under certain conditions, propagate over a thousand kilometers into the ice pack (Nose et al., 2023; Squire et al., 2009). Wave energy attenuates through various scattering and wave dissipation mechanisms which largely depend on the properties of the sea ice and wave field, including sea ice thickness, concentration, floe size distribution, sea ice mechanical and material properties, wave frequency and wave steepness (for and overview see Shen, 2019; Squire, 2020; Rogers et al., 2021; Thomson, 2022, and references therein). To model waves in sea ice, a considerable number of theories and parameterizations have been established to estimate the wave attenuation rate (e.g., Liu et al., 2020, their Table 1). While the majority of these have been calibrated and/or validated using experimental observations, uncertainty persists under what sea ice and wave conditions the various physical processes that underpin these models may be valid or of importance (Shen, 2019; Voermans et al., 2021; Shen, 2022).

Our understanding and capacity to model the wave attenuation rate in terms of system variables is, in part, hindered by the characteristic length scales associated with the wave-ice interactions mechanisms, the numerical models and sea ice observations. Operational forecasting models and wave observations typically operate or are retrieved at scales of $O(1 - 10 \text{ km})$. Yet, sea ice conditions and wave-ice physics can vary strongly across such distances, where, for example, individual floe dimensions can vary from $O(\text{m})$ to $O(\text{km})$ (Toyota et al., 2006), and ice-ocean roughness scales can vary from $O(\text{mm})$ for smooth surfaces to $O(\text{m})$ for ridges. In most cases, MIZ consists of a mixture of different types of ice and open water conditions where each combination may lead to completely different behaviors of waves in ice (Herman, 2024). This not only makes it challenging as to how the dynamics at small scales need to be captured at much larger scales, whether it be in the form of an effective or average sea ice property, but also how such properties can be realistically derived or observed at much larger scales for usage in numerical simulations. In particular, products of only sea ice concentration and sea ice thickness are currently routinely generated in the Antarctic. Given the many variables required to describe sea ice, this often leads to the necessary assumption that all other sea ice variables in our models are homogeneous, ultimately representing a major source of uncertainty.

A straightforward approach to improve our modeling capacity of wave-ice interactions is to retrieve considerably more data under a diverse range of wave and sea ice conditions in an attempt to identify trends with system variables (Rogers et al., 2021; Montiel et al., 2022; Rabault et al., 2023). Observations of the wave attenuation rate in sea ice can be obtained by measuring the difference in wave energy between two locations, typically using wave-ice buoys deployed on the ice or by any other



motion-recording sensor. Traditionally, it is assumed that wave energy decays exponentially with distance into the MIZ (e.g., Wadhams et al. (1988), although questions have been raised on the a priori assumption of its validity; e.g., see Squire (2018); Herman (2024)):

$$55 \quad E(f, x) = E(f, 0) \exp(-\alpha x) \quad (1)$$

where $E(f, x)$ is the wave energy spectrum, which varies with wave frequency f and the distance into the MIZ x at a rate given by the wave attenuation rate α . The number of estimates of the wave attenuation rate from wave observations has increased drastically in recent years due the rapid progresses in the development of low-cost wave-ice buoys, as they are able to capture the wave field characteristics $E(f, x)$ at high temporal resolution and at high accuracy (Kohout et al., 2015; Rabault et al., 60 2022; Kodaira et al., 2024; Womack et al., 2024). The main restriction of in-situ deployments is, however, that the spatial coverage tends to be extremely sparse due to the logistical challenges in deploying instrumentation in the remote and harsh Antarctic MIZ. Thus, even though the studies associated with such deployments may provide high temporal detail of wave-ice interactions, the complexity and diversity of sea ice conditions during such deployments restricts our general understanding of wave-ice interactions in general and their applicability at much larger scales.

65 Satellite remote sensing may provide critical large spatial coverage that is difficult to achieve with in-situ instruments alone, albeit, at the cost of reduced temporal and frequency resolution. Synthetic Aperture Radar (SAR) imagery was used by Arduin et al. (2017) to retrieve wave field properties in the MIZ, and by Stopa et al. (2018) to derive wave attenuation estimates. Horvat et al. (2020) and Collard et al. (2022) used laser altimeter observations from ICESat-2 to identify waves in sea ice, and was used by Brouwer et al. (2022) and Hell and Horvat (2024) to derive 1D and 2D wave spectra, respectively. While no direct 70 estimates of the wave attenuation rate α were provided, the ICESat-2 derived data of Brouwer et al. (2022) showed a clear reduction of the spectral wave energy into the Antarctic MIZ (i.e. see Figs. B2 and C2 in Brouwer et al., 2022). This highlights the potential of using ICESat-2 observations to estimate the wave attenuation rate in sea ice across large distances. In this study, our objective is to estimate the wave attenuation rate from ICESat-2 altimeter measurements in an attempt to identify trends in the wave attenuation rate across the Antarctic MIZ and under a wide range of sea ice conditions.

75 2 Methods

For the estimation of the wave attenuation rate in the MIZ we make use of the processed ICESat-2 data (Fraser et al., 2024) as derived by Brouwer et al. (2022), which is derived from the Level 3 sea ice height product (ATL07, version 2; Kwok et al. (2021)), from the National Snow and Ice Data Center (NSIDC; <https://nsidc.org/data/atl07>, last access: 5 March 2020). The transects included in this dataset were subjected to extensive quality control measures, including the selection of transects 80 with low cloud coverage and manual discrimination between swell-dominated and ice-structure-dominated contributions to the observed height. The data include 320 transects covering February, May, September and December of 2019, representing times of minimum extent, rapid autumn advance, maximum extent and rapid summer retreat, respectively (Eayrs et al., 2019). The data consist of surface height measurements of the three high-power laser beams crossing the Antarctic MIZ in a predominantly



north-south direction, resampled to a regular horizontal spacing of 8 m (i.e., oversampling the ATL07 mean segment length of
85 ~ 15 m). In addition, we use estimates of the MIZ width x_{MIZ} based on the depth of wave penetration into the MIZ Brouwer
et al. (2022). Specifically, we use the median value of the MIZ distance determined in four wavelengths (165, 239, 345 and
498 m). The reader is referred to Brouwer et al. (2022) for further details on the dataset Fraser et al. (2024) and initial quality
control procedures.

Here, the wave energy density $E(k, x)$, where k is the wave number, was estimated for section lengths L along the transect
90 using Welch's method with Hamming windowing, 50% overlap and segment lengths of 128 sample points. While there are
potential pathways to estimate the wave direction, we assume here that the impact of misalignment angle between wave
propagation and the ICESat-2 transect are of minor influence, a common assumption in wave-ice interaction studies and one
which is more suitable to the Antarctic than the Arctic, given the circumpolar sea ice morphology of the former. That is, we
assume that the apparent wave number k_a is equal to the wave number $k = k_a / \cos(\Delta\theta)$ with $\Delta\theta = 0$. The choice of L is
95 somewhat arbitrary. Large L reduces uncertainty in estimates of $E(k, x)$ whilst potential inhomogeneity of the sea ice cover
requires a relatively small section length. In Fig. 1(a,b) we compare the wave energy estimates for section lengths L of 2048,
8192 m and 16384 m for two transects. For quality control purposes, we estimate the signal-to-noise ratio (eSNR, where 'e'
denotes 'estimated') as $\sqrt{\int_{k_i}^{k_{i+1}} E(k_i, x) \Delta k} / \epsilon$, where we assume an accuracy of the ICESat-2 measurements in sea ice as
 $\epsilon = 2$ cm, following the study of Neumann et al. (2019). It is noted, however, that this is not a robust estimate of the eSNR
100 as it depends on the resolution of the spectral density estimate Δk . We use a value of $eSNR = 1$ as a cut-off threshold. For
some transects, the energy in some wave number bins increases again towards the end of the MIZ, which is most likely due to
a strong increase of the ice-structure contribution to the surface height fluctuations after most wave energy has been attenuated
(as mentioned in Brouwer et al., 2022). For this reason, we choose the absolute minimum of wave energy within the MIZ as a
cut-off instead for these transects.

105 To estimate the apparent attenuation rate α , we make use of the commonly adopted assumption that wave energy decays
exponentially with distance into the ice pack (Eq. 1). When surface elevation measurements are available at different x (i.e.
from an altimeter transect), α can be readily obtained from:

$$\alpha = \frac{-\ln(E(f, x_2)/E(f, x_1))}{\Delta x} \quad (2)$$

with $\Delta x = (x_2 - x_1) \cos(\Delta\theta)$, where x_1 being the position closest to the ice edge. While there are different methods for
110 estimating the wave direction, such as from satellite observations (Hell and Horvat, 2024; Collard et al., 2022) or wave model
hindcast (Stopa et al., 2016; Liu et al., 2021), we assume here that the waves are well aligned with the direction of the ICESat-
2 measurements, i.e., $\Delta\theta = 0$, rather than attempting to apply such corrections. We will discuss this assumption later on.
Naturally, this means that the attenuation rates estimated in this study are likely an underestimation.

Following Eq. 2, the wave attenuation rates corresponding to the wave energy of transects shown in Fig. 1(a,b) are shown in
115 Fig. 1(c,d) for different L and for $\Delta x = 16$ km. As $L = 8192$ m significantly reduces the fluctuations in E and α , we choose
 $L = 8192$ m as the section length, as opposed to a larger length, to restrict uncertainty associated with sea ice inhomogeneity
on our results. Further, in this study we restrict our focus to the wave energy at wave lengths of $\lambda = 128, 227, 341$ and 512 m,




Figure 1. Wave energy (a,b) and apparent wave attenuation rates (c,d) estimated for two transects for $T = 12$ s, and for different section lengths $L = 2948$ m, 8192 m and 16384 m. Parts of transect with $eSNR < 1$ are marked in grey. Dotted and dashed vertical lines depict the inner MIZ boundary as determined by Brouwer et al. (2022) corresponding to $T = 12$ s and a broad range of frequencies, respectively.

corresponding to wave periods of roughly $T = 9, 12, 15$ and 18 s in deep water assuming the linear dispersion relationship for open-water waves.

120 3 Results

3.1 Attenuation examples

The wave energy and attenuation rate of an example transect are shown in Fig. 2 for spectral bins corresponding to $T = 9, 12, 15$ and 18 s. A Sentinel-1 image of the sea ice conditions a day after the ICESat-2 measurements is provided in Fig. 2a, showing consolidated pack ice along the majority of the transect, with the exception of the initial part of the MIZ which looks like sea ice with very low concentration. Aside from minor spatial fluctuations in wave energy, an overall decrease of the spectral wave energy is observed from the ice edge into the MIZ for all wave periods (Fig. 2b). The attenuation rate estimated for this transect is well sorted by wave frequency, showing strongest attenuation for the shortest waves (Fig. 2c). The early cutoff of α for $T = 9$ s is a consequence of the high attenuation rate and the $eSNR$ threshold imposed. The attenuation rate for




Figure 2. An example ICESat-2 transect (a) on 2019-12-26 with estimated spectral wave energy (b) and apparent wave attenuation rate (c). The MIZ, as derived by Brouwer et al. (2022), along the transect is given by the black line (a), which is 61.5 km wide. Black circles denote fractions of the MIZ width, from 0.0 to 1.0, with the larger circles corresponding to $x/x_{MIZ} = 0, 0.5$ and 1. The Sentinel-1 image is from 2019-12-27, i.e., a day after the ICESat-2 transect measurements. The gray lines in (b) represent observations removed after quality control.

$T = 18$ s declines gently before briefly becoming slightly negative halfway the MIZ. We note that the overall magnitudes of the apparent attenuation rates are comparable to those observed by Kohout et al. (2020), Voermans et al. (2021) and Montiel et al. (2022), and slightly larger than Meylan et al. (2014) and Rogers et al. (2021).

Another example transect is shown in Fig. 3, illustrating contrasting trends in the estimates of the apparent attenuation rates compared to Fig. 2. For this transect, fluctuations and larger scale trends in the estimated wave energy are of sufficient magnitude to lead to slightly negative estimates of the apparent wave attenuation rate for all wave periods (Fig. 3c). For $x/x_{MIZ} < 0.5$, α does not show any correlation with wave frequency. This changes for $x/x_{MIZ} > 0.5$ where α increases steeply and appears well sorted again with wave frequency. From the Sentinel-1 imagery it is unclear why this sudden change occurred around $x/x_{MIZ} = 0.5$. While a strong gradient in Sentinel-1 backscatter can be observed between $0.1 < x/x_{MIZ} < 0.2$ (Fig. 3a), the location of this gradient does not appear to coincide with the position of rapid change in α . Imagery on the days before and after the transect measurements do not suggest great change in location of this strong gradient in backscatter.




Figure 3. An example ICESat-2 transect (a) on 2019-05-24 with estimated spectral wave energy (b) and apparent wave attenuation rate (c). The MIZ width along the transect is given by the black line (a), which is 137.5 km wide, on top of a Sentinel-1 image from 2019-05-24 (i.e., the same day as the ICESat-2 transect). The gray lines in (b) represent observations removed after quality control.

140 3.2 Sensitivity of attenuation rate to the wave field direction

While it is tempting to interpret small details of α for each transect in terms of spatial changes in sea ice conditions and/or wave physics, it remains uncertain whether this is realistic. For example, while the occasional negative attenuation rates observed in Fig. 3 would imply an increase in wave energy, potentially due to wind-input or non-linear wave interactions (e.g. Li et al., 2017), they may also simply be methodological artifacts due to a low and/or variable signal-to-noise ratio (Thomson et al., 2021) or non-stationarity of the incoming wave field (Voermans et al., 2023). Here, however, we suggest that some transient fluctuations and negative attenuation rates observed in our estimates may be caused by the assumption of homogeneous sea ice combined with the assumption that the waves propagate along the same direction as the measurement transect.

To illustrate this, we take another example transect (Fig. 4a) where sea ice conditions are highly inhomogeneous, most likely due to ocean eddy-ice interactions (e.g., Manucharyan and Thompson, 2017). We then take the Sentinel-1 backscatter as a crude proxy of the local attenuation rate $\alpha \approx \alpha_p$, where we assume that in open water (dark = low backscatter) the attenuation is very low, and in ice covered regions (bright = high backscatter) the attenuation rate is much higher (see colorbar in Fig. 4). Of course, this is by no means an accurate depiction of the true attenuation rate and is merely used here to illustrate the




Figure 4. Estimated impact of misalignment angle between incident wave direction and ICESat-2 transect for example transect on 01-May-2019: (a) proxy-attenuation rate α_p based on Sentinel-1 backscatter (note, for illustrative purposes only); (b) estimated wave energy along the transect based on different incident wave direction. The observed wave energy for $T = 12$ s for this transect is shown in black.

impact of wave direction on the interpretation of the estimated wave energy along a transect. We then estimate the wave energy along the transect for different wave directions given the spatial variability of α_p and assuming an exponential decay of wave energy with x , i.e., Eq. 1. With such a simple model, wave energy traveling in the same direction as the transect (in this case at 6° relative to north, we use the ‘coming from’ convention here) decreases monotonically along the transect (Fig. 4b). However, if waves would approach from either 15° or 325° relative to north, fluctuations in wave energy along the transect start to appear. If Δx is smaller than the length scale of such fluctuations, a negative apparent attenuation rate will be found. In case of 325° , this leads to a very large fluctuation as waves traveling towards the transect at around $x/x_{MIZ} = 0.4$ experience low attenuation ice conditions when coming from the Southern Ocean in contrast to waves arriving at neighboring points. In other words, waves arriving at different points along the transect are experiencing completely different attenuation through differing sea ice conditions. In this specific case, this leads to a threefold increase in the observed wave energy between $x/x_{MIZ} = 0.2$ and 0.4 along the transect. While the exact location of the local maximum and maximum in wave energy between the ICESat-2 estimates and our simple model are slightly off, the difference in magnitude between them are fairly similar. (We note that the interpretation of the sharp increase in α around $x/x_{MIZ} = 0.5$ for the example transect shown in Fig. 3 can be readily explained by the potential misalignment between the incident wave angle and transect as well, see Appendix A.) The uncertainty associated with the misalignment angle and sea ice inhomogeneity also means that even when the wave direction is known, correction of wave attenuation estimates for misalignment does not necessarily provide a more reliable estimate of the wave attenuation rates, nor does it mean that negative attenuation rates should be disregarded a priori.




Figure 5. The variability of the mean apparent attenuation rate $\bar{\alpha}$ for $T = 12$ s against (a) relative distance into the MIZ, (b) sea ice concentration, (c) sea ice thickness, and (d) wave energy, using all tracks in the Fraser et al. (2024) dataset. Light gray shading corresponds to the 25th and 75th percentiles of the dataset; in (a) the dark gray shading corresponds to the 95% confidence interval of the mean (bootstrap sampling, 1000 samples with replacements). For bins with less than 30 observations, the data are shown by a dashed line, see (a) and (d).

170 3.3 Overall attenuation statistics

The combination of sea ice inhomogeneity and measurement misalignment introduces complications in the interpretation of α as estimated for individual transects. However, given that there is no correlation between the ICESat-2 tracks and local sea ice and wave conditions, we may expect that for some transects the resulting fluctuations in α are either an underestimation or overestimation of the true wave attenuation rate. This would mean that the average apparent attenuation rate $\bar{\alpha}$ of all transects could still provide a reasonable depiction of the mean wave attenuation rate across the MIZ. In Fig. 5a the mean apparent wave attenuation rate $\bar{\alpha}$, averaged over all transects in the dataset, is shown for $T = 12$ s. Here, we removed transects that contained extreme outliers in excess of three standard deviations from the median to reduce noise in the estimates of $\bar{\alpha}$ which reduced the size of the dataset by 13%. The observations suggest a linear increase in $\bar{\alpha}$ from about 0 to 10^{-4} across the MIZ (when plotted on logarithmic scales, it approaches $\bar{\alpha} = 10^{-5}$ at $x = 0$, not shown). While the scatter is reasonably large, noting that the light gray area depicts the 25% and 75% percentiles, this is not necessarily surprising as some transects may present strong fluctuations, and can even become negative (e.g., see Fig. 3). Moreover, local sea ice conditions may vary significantly between



transects or with respect to the ‘average’ sea ice conditions in the MIZ. Trends at the end and start of the MIZ are considered unreliable as few observations are available here. For $x/x_{MIZ} = 0.9$, there is a tendency for a flattening and decrease in $\bar{\alpha}$. However, we expect this to be a methodological artifact caused by a low eSNR and sea ice morphology, i.e., near the inner
185 limit of the MIZ the wave energy is considerably smaller such that sea ice morphology starts to contaminate the wave energy estimates.

In Fig. 5(b) and (c) the mean attenuation rate is compared against AMSR2-derived sea ice concentration (Spreen et al., 2008), downloaded from <https://data.seaice.uni-bremen.de/amr2/> (last access: 22 February 2024), and SMOS sea ice thickness (Huntemann et al., 2014), downloaded from <https://data.seaice.uni-bremen.de/smos/> (last access: 22 February 2024),
190 respectively. While we observe modest trends of $\bar{\alpha}$ with sea ice properties, they are much weaker than the trend with distance into the MIZ. Obviously, sea ice thickness, ice concentration and floe size distribution are strongly correlated with x , and we thus suspect that these are nevertheless responsible for the increase in $\bar{\alpha}$ when scaled with the distance into the MIZ. In Fig. 5d we compare the attenuation rate against wave energy. In general, no trend can be observed with wave energy, except when E is very small. This is however caused by a bias in spatial correlation with distance into the MIZ (red line), i.e., observations of low
195 wave energy are more likely to occur deep into the MIZ, as opposed to high wave energy observations. A similar conclusion can be made when evaluating the decay rate of the total wave energy in the MIZ (see Appendix B).

Trends of the mean wave attenuation rate for $T = 9, 15$ and 18 s are consistent with those of $T = 12$ s (Fig. 6), with all indicating that $\bar{\alpha}$ increases linearly with distance into the MIZ. The apparent attenuation rates are significantly higher for $T = 9$ s compared to the longer wave periods. While no clear sorting can be observed for $x/x_{MIZ} < 0.5$ for $T = 12, 15$ and
200 18 s, gradual separation in terms of frequency can be observed towards the inner MIZ. Some overlap of the 95% confidence bounds of $\bar{\alpha}$, however, persists across the MIZ for the current dataset.

3.4 Seasonality and frequency-dependence of attenuation

To assess whether $\bar{\alpha}$ varies between seasons, the mean attenuation rate for $T = 12$ s was determined for the months February, May, September and December (Fig. 7) as well. While scattered, it appears that the attenuation rate in the months of December
205 and February increase initially much faster with distance into the MIZ, reaching a maximum already about halfway the MIZ. Such a difference would be consistent with the expected wave climate at the ice edge of the Antarctic MIZ during the winter and summer, where wave climate is milder in summer and the sea ice is retreated from the high-energy Southern Ocean (Young et al., 2020). This is expected to lead to stronger wave-induced sea ice break-up in winter, which is expected to reduce the wave attenuation rates in the MIZ in May and September. In addition, we may expect that sea ice mechanical and material properties
210 are different during the seasons which could potentially lead to a change in the apparent wave attenuation rate (e.g., sea ice is thicker in summertime; Worby et al. (2008)). However, some caution is required in the interpretation of the monthly averages shown as the number of observations in February and December are very small. Thus, more data are required to confirm any seasonality of $\bar{\alpha}$.

To evaluate the frequency dependence of $\bar{\alpha}$, the mean apparent attenuation rate between $0.1 < x/x_{MIZ} < 0.6$ is replotted
215 in Fig. 8 (corresponding to the region where estimates are available for all four evaluated frequencies). For $0.08 < f < 0.11$,




Figure 6. (a) The variability of the mean apparent attenuation rate $\bar{\alpha}$ for $T = 9, 12, 15$ and 18 s across the MIZ. The number of observations at each x is given in (b). The shaded areas correspond to the 95% confidence interval of the mean (bootstrap sampling, 1000 samples with replacements).

the attenuation rate $\bar{\alpha}$ seems to scale with a power between 2 and 3, while a considerably flatter trend may be observed for $0.08 < f$. The former is similar to the parameterization of Meylan et al. (2014), whilst the latter appears consistent with the field observations of Montiel et al. (2022) who observe significant flattening for frequencies below about 0.06 Hz. However, as the confidence limits of $\bar{\alpha}$ overlap for $T = 12, 15$ and 18 s, and the range of f for which observations of $\bar{\alpha}$ are available is very limited, more data are required to ascertain such power law.

4 Discussion & Conclusions

The ICESat-2 observations of Fraser et al. (2024) provide a unique dataset of waves in sea ice obtained across a diverse range of Antarctic sea ice conditions (Brouwer et al., 2022). Averaging all estimates of the wave attenuation rate along the transects reveals a strong correspondence between $\bar{\alpha}$ and the relative position within the MIZ. Specifically, we find that $\bar{\alpha}$ increases linearly with distance from the ice edge. The trend with distance into the MIZ is much clearer than sea ice concentration or ice thickness alone. The change of $\bar{\alpha}$ is expected to be a natural consequence of changing sea ice conditions with x/x_{MIZ} where, on average, one may expect increases in sea ice thickness, ice concentration, and floe sizes moving into the MIZ. Additionally,




Figure 7. (a) The variability of the monthly mean apparent attenuation rate $\bar{\alpha}$ for $T = 12$ s across the MIZ for February, May, September and December. The number of observations at each x is given in (b) noting that the observations in February and December are very limited.

variability may be expected with material and mechanical properties of sea ice as well, although no clear observations are available on their trends with x/x_{MIZ} .

230 While the wave attenuation rates estimated using ICESat-2 measurements compare well to the rates observed by others through in-situ experiments (Meylan et al., 2014; Kohout et al., 2020; Voermans et al., 2021; Rogers et al., 2021; Montiel et al., 2022), direct validation of the ICESat-2 derived wave spectra is still required to provide greater certainty of their accuracy. Particularly, such observations are ideally made within a few kilometers from the ICESat-2 tracks to avoid impacts of sea ice inhomogeneity, and should provide further insights into the impact of sea ice variance on biases and aliasing of wave spectra estimates (Hell and Horvat, 2024). Comparison against observations of sea floor cables is attractive in particular as it provides additional details on the spatial variability of sea ice on wave attenuation (Smith et al., 2023). Impact of a misalignment angle between the incident wave direction and the ICESat-2 transects remains a factor of uncertainty on the geometric correction of α , the apparent wave number, the assumption of sea ice homogeneity and the shape of the ice edge in the area immediately adjacent to the transect. Specifically, we have shown here that the inhomogeneity and misalignment angle between the incident wave direction and measurement transect could lead to major fluctuations in the observed wave energy, which in turn leads to strong fluctuations in estimates of the wave attenuation rates including negative rates. Large in-situ buoy deployments should be able to provide further insights in the influence thereof on estimates of α . Lastly, our study assumes that the incoming

235

240




Figure 8. Frequency dependence of $\bar{\alpha}$ across the MIZ for $0.1 < x/x_{MIZ} < 0.62$. Comparison against the empirical model of Meylan et al. (2014) is provided.

wave field is stationary and homogeneous, whereas preexisting spatial variance in wave energy can readily be observed by, for example, wave-current interactions (e.g., see also Fig. 4).

245 While the correspondence observed between $\bar{\alpha}$ and x/x_{MIZ} is empirical, it may provide alternative approaches to model wave attenuation in global models if the MIZ width is a known variable. A linear relationship of the form $\bar{\alpha} = \beta x/x_{MIZ} + \gamma$, would imply that wave energy decays into the Antarctic MIZ as:

$$E(f, x) = E(f, 0) \exp(-\beta x^2 \dots) \quad (3)$$

250 where β is a wave attenuation rate coefficient which varies with frequency. Obviously, such a model ignores the presence of strong local variability of wave-ice conditions, and such an approach is therefore unlikely to model waves in an accurate way at small spatial and short time domains (Herman, 2024). Nevertheless, at global and climate scales, such detail may not necessarily be required and the model may provide a major advantage compared to current physics based models which typically rely on system variables that are not straightforward to measure, retrieve or simulate at global scales.

Data availability. The processed ICESat-2 data underlying this manuscript are freely available at https://data.aad.gov.au/metadata/AAS_4528_ICESat-2-wave-attenuation-tracks or <http://dx.doi.org/doi:10.26179/q9pe-w283>. AMSR2-derived sea ice concentration and SMOS derived thin sea ice thickness were obtained from the University of Bremen (<https://seaice.uni-bremen.de/>).




Figure A1. Impact of sea ice inhomogeneity and misalignment between incident wave direction and ICESat-2 transect (an expansion of the region shown in Fig. 3). (a) Sentinel-1 image from 2019-05-24, (b) wave direction from hindcasts.

Appendix A: Misalignment angle, example transect Fig. 3

In the example transect shown in Fig. 3 a steep increase in the estimated wave attenuation rate α at around $x/x_{MIZ} = 0.5$ was observed. Notably, this steep increase cannot be linked straightforwardly to an obvious change in sea ice conditions based on visual inspection of the corresponding Sentinel-1 image. We consider here the misalignment angle of the incident wave direction and the transect and the inhomogeneity of the sea ice as a likely reason of the steep increase in α . In Fig. A1 a larger spatial area of the sea ice conditions is shown, revealing strong variability in the ice edge position. Particularly, the region of sea ice surrounding $67^{\circ}\text{S } 90^{\circ}\text{E}$ is expected to lead to a sharp drop in wave energy around $x/x_{MIZ} \approx 0.5$ if the incident wave direction is approximately 345° (relative to north). That is, waves arriving along the transect at $x/x_{MIZ} < 0.5$ will have crossed a significantly shorter distance through sea ice in comparison to the wave energy arriving at $x/x_{MIZ} > 0.5$. Wave directional data just north of the observation site ($65.75^{\circ}\text{S } 89^{\circ}\text{E}$) obtained from ERA5 Reanalysis and WaveWatchIII hindcast Liu et al. (2021) suggests an incident wave direction of around 345° is well within the range of possibility.

Appendix B: Attenuation rate of significant wave height

Although the wave attenuation rate is frequency dependent, the rapid attenuation of wave energy at higher frequencies means that the spectrum becomes narrower deeper into the MIZ. In such a case, one may look at the attenuation rate of the significant wave height α_H instead, which has the advantage of being more robustly measured than the wave spectrum but at the cost of




Figure B1. Attenuation rate of significant wave height (a), wave height gradient (b), mean sea ice concentration (c) and relative position in the MIZ (d) against significant wave height for all transects.

the error made by ignoring the frequency dependence (Kohout et al., 2020):

$$\frac{dH}{dx} = -\frac{\alpha_H}{2} H_s \tag{B1}$$

where $H_s = 4\sqrt{m_0}$ is the significant wave height and m_0 is the first order moment of the wave spectrum. While the dataset suggests that the wave attenuation rate of H_s decays strongly with H_s (Fig. B1a), this is in large part biased by the strong correlation between H_s and the relative position within the MIZ (Fig. B1c,d) (see also Kohout et al., 2020). That is, low H_s is predominantly observed deep into the MIZ where sea ice concentration and sea ice thickness are relatively high, whereas larger H_s tends to be observed near the ice edge where sea ice concentration and sea ice thickness tends to be low.

Author contributions. Conceptualization: ADF. Methodology: JJV. Formal analysis: JJV, ADF. Data interpretation: all authors. Writing - Original Draft: JJV. Writing - Review & Editing: all authors.

Competing interests. The contact author has declared that neither they nor their co-authors have any competing interests



Acknowledgements. This project is supported through the Australian Government as part of the Antarctic Science Collaboration Initiative program and contributes to Project 6 (Sea Ice) of the Australian Antarctic Program Partnership (project ID ASCI000002). ADF is supported by the Australian Research Council through a Future Fellowship (FT230100234). ADF acknowledges the generous support of the Harris Charitable Trust through the Antarctic Science Foundation. QL acknowledges the National Natural Science Foundation of China (42106012), the Shandong Provincial Natural Science Fund for Excellent Young Scientists Fund Program (Overseas) (2023HWYQ-056), the Taishan Scholars Program (tsqz20221111) and the Fundamental Research Funds for the Central Universities (202441007). AVB acknowledges support of the Centre for Disaster Management and Public Safety (CDMPS). Authors acknowledge the usage of EO Browser, Sinergise Solutions d.o.o., a Planet Labs company, in retrieving Sentinel-1 imagery. Authors would like to thank Tripp Collins for feedback on this study.



References

- Ardhuin, F., Stopa, J., Chapron, B., Collard, F., Smith, M., Thomson, J., Doble, M., Blomquist, B., Persson, O., Collins III, C. O., et al.: Measuring ocean waves in sea ice using SAR imagery: A quasi-deterministic approach evaluated with Sentinel-1 and in situ data, *Remote Sensing of Environment*, 189, 211–222, 2017.
- 295 Bateson, A. W., Feltham, D. L., Schröder, D., Wang, Y., Hwang, B., Ridley, J. K., and Aksenov, Y.: Sea ice floe size: its impact on pan-Arctic and local ice mass and required model complexity, *The Cryosphere*, 16, 2565–2593, 2022.
- Bennetts, L. G., Shakespeare, C. J., Vreugdenhil, C. A., Foppert, A., Gayen, B., Meyer, A., Morrison, A. K., Padman, L., Phillips, H. E., Stevens, C. L., Toffoli, A., C., C. N., Kiss, A. E., Kraitzman, N., Malyarenko, A., McConnochie, C. D., Meucci, A., Montiel, F., Neme, J., Nikurashin, M., Patel, R. S., Peng, J.-P., Rayson, M., Rosevear, M. G., Sohail, T., Spence, P., and J. S. G.: Closing the loops on Southern
- 300 Ocean dynamics: From the circumpolar current to ice shelves and from bottom mixing to surface waves, *ESS Open Archive*, 2024.
- Boutin, G., Lique, C., Ardhuin, F., Rousset, C., Talandier, C., Accensi, M., and Girard-Ardhuin, F.: Towards a coupled model to investigate wave–sea ice interactions in the Arctic marginal ice zone, *The Cryosphere*, 14, 709–735, 2020.
- Brouwer, J., Fraser, A. D., Murphy, D. J., Wongpan, P., Alberello, A., Kohout, A., Horvat, C., Wotherspoon, S., Massom, R. A., Cartwright, J., et al.: Altimetric observation of wave attenuation through the Antarctic marginal ice zone using ICESat-2, *The Cryosphere*, 16, 2325–
- 305 2353, 2022.
- Collard, F., Marié, L., Nouguier, F., Kleinherenbrink, M., Ehlers, F., and Ardhuin, F.: Wind-Wave Attenuation in Arctic Sea Ice: A Discussion of Remote Sensing Capabilities, *Journal of Geophysical Research: Oceans*, 127, e2022JC018654, 2022.
- Collins III, C. O., Rogers, W. E., Marchenko, A., and Babanin, A. V.: In situ measurements of an energetic wave event in the Arctic marginal ice zone, *Geophysical Research Letters*, 42, 1863–1870, 2015.
- 310 Dumont, D., Kohout, A., and Bertino, L.: A wave-based model for the marginal ice zone including a floe breaking parameterization, *Journal of Geophysical Research: Oceans*, 116, 2011.
- Eayrs, C., Holland, D., Francis, D., Wagner, T., Kumar, R., and Li, X.: Understanding the seasonal cycle of Antarctic sea ice extent in the context of longer-term variability, *Reviews of Geophysics*, 57, 1037–1064, 2019.
- Fraser, A., Brouwer, J., and Wongpan, P.: Altimetric observation of wave attenuation through the Antarctic marginal ice zone using ICESat-2; Dataset, Tech. rep., Australian Antarctic Data Centre, <http://dx.doi.org/doi:10.26179/q9pe-w283>, Accessed: 2023-08-24, 2024.
- 315 Häkkinen, S.: Coupled ice-ocean dynamics in the marginal ice zones: Upwelling/downwelling and eddy generation, *Journal of Geophysical Research: Oceans*, 91, 819–832, 1986.
- Hell, M. C. and Horvat, C.: A method for constructing directional surface wave spectra from ICESat-2 altimetry, *The Cryosphere*, 18, 341–361, 2024.
- 320 Herman, A.: From apparent attenuation towards physics-based source terms—a perspective on spectral wave modeling in ice-covered seas, *Frontiers in Marine Science*, 11, 1413 116, 2024.
- Horvat, C., Blanchard-Wigglesworth, E., and Petty, A.: Observing waves in sea ice with ICESat-2, *Geophysical Research Letters*, 47, e2020GL087629, 2020.
- Huntemann, M., Heygster, G., Kaleschke, L., Krumpfen, T., Mäkynen, M., and Drusch, M.: Empirical sea ice thickness retrieval during the
- 325 freeze-up period from SMOS high incident angle observations, *The Cryosphere*, 8, 439–451, 2014.



- Kodaira, T., Katsuno, T., Nose, T., Itoh, M., Rabault, J., Hoppmann, M., Kimizuka, M., and Waseda, T.: An affordable and customizable wave buoy for the study of wave-ice interactions: design concept and results from field deployments, *Coastal Engineering Journal*, 66, 74–88, 2024.
- Kohout, A. L. and Meylan, M. H.: An elastic plate model for wave attenuation and ice floe breaking in the marginal ice zone, *Journal of Geophysical Research: Oceans*, 113, 2008.
- 330 Kohout, A. L., Penrose, B., Penrose, S., and Williams, M. J.: A device for measuring wave-induced motion of ice floes in the Antarctic marginal ice zone, *Annals of Glaciology*, 56, 415–424, 2015.
- Kohout, A. L., Smith, M., Roach, L. A., Williams, G., Montiel, F., and Williams, M. J.: Observations of exponential wave attenuation in Antarctic sea ice during the PIPERS campaign, *Annals of Glaciology*, 61, 196–209, 2020.
- 335 Kwok, R., Petty, A. A., Cunningham, G., Markus, T., Hancock, D., Ivanoff, A., Wimert, J., Bagnardi, M., Kurtz, N., and ICESat 2 Science Team: ATLAS/ICESat-2 L3A sea ice height, version 2, Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center, 20, <https://doi.org/10.5067/ATLAS/ATL07.005>, 2021.
- Li, J., Kohout, A. L., Doble, M. J., Wadhams, P., Guan, C., and Shen, H. H.: Rollover of apparent wave attenuation in ice covered seas, *Journal of Geophysical Research: Oceans*, 122, 8557–8566, 2017.
- 340 Li, J., Babanin, A. V., Liu, Q., Voermans, J. J., Heil, P., and Tang, Y.: Effects of Wave-Induced Sea Ice Break-Up and Mixing in a High-Resolution Coupled Ice-Ocean Model, *Journal of Marine Science and Engineering*, 9, 365, 2021.
- Liu, Q., Rogers, W. E., Babanin, A., Li, J., and Guan, C.: Spectral modeling of ice-induced wave decay, *Journal of Physical Oceanography*, 50, 1583–1604, 2020.
- Liu, Q., Babanin, A. V., Rogers, W. E., Zieger, S., Young, I. R., Bidlot, J.-R., Durrant, T., Ewans, K., Guan, C., Kirezci, C., et al.: Global wave hindcasts using the observation-based source terms: Description and validation, *Journal of Advances in Modeling Earth Systems*, 13, e2021MS002493, 2021.
- 345 Manucharyan, G. E. and Thompson, A. F.: Submesoscale sea ice-ocean interactions in marginal ice zones, *Journal of Geophysical Research: Oceans*, 122, 9455–9475, 2017.
- Meylan, M. H., Bennetts, L. G., and Kohout, A. L.: In situ measurements and analysis of ocean waves in the Antarctic marginal ice zone, *Geophysical Research Letters*, 41, 5046–5051, 2014.
- 350 Montiel, F., Kohout, A. L., and Roach, L. A.: Physical drivers of ocean wave attenuation in the marginal ice zone, *Journal of Physical Oceanography*, 52, 889–906, 2022.
- Neumann, T. A., Martino, A. J., Markus, T., Bae, S., Bock, M. R., Brenner, A. C., Brunt, K. M., Cavanaugh, J., Fernandes, S. T., Hancock, D. W., et al.: The Ice, Cloud, and Land Elevation Satellite–2 Mission: A global geolocated photon product derived from the advanced topographic laser altimeter system, *Remote sensing of environment*, 233, 111325, 2019.
- 355 Nose, T., Katsuno, T., Waseda, T., Ushio, S., Rabault, J., Kodaira, T., and Voermans, J.: Observation of wave propagation over 1,000 km into Antarctica winter pack ice, *Coastal Engineering Journal*, pp. 1–17, 2023.
- Rabault, J., Nose, T., Hope, G., Müller, M., Breivik, Ø., Voermans, J., Hole, L. R., Bohlinger, P., Waseda, T., Kodaira, T., et al.: Openmetbuoy-v2021: An easy-to-build, affordable, customizable, open-source instrument for oceanographic measurements of drift and waves in sea ice and the open ocean, *Geosciences*, 12, 110, 2022.
- 360 Rabault, J., Müller, M., Voermans, J., Brazhnikov, D., Turnbull, I., Marchenko, A., Biuw, M., Nose, T., Waseda, T., Johansson, M., et al.: A dataset of direct observations of sea ice drift and waves in ice, *Scientific Data*, 10, 251, 2023.



- Rogers, W. E., Meylan, M. H., and Kohout, A. L.: Estimates of spectral wave attenuation in Antarctic sea ice, using model/data inversion, *Cold Regions Science and Technology*, 182, 103–198, 2021.
- 365 Shen, H. H.: Modelling ocean waves in ice-covered seas, *Applied Ocean Research*, 83, 30–36, 2019.
- Shen, H. H.: Wave-ice interaction models and experimental observations, in: *Proceedings of the IUTAM Symposium held at Aalto University, Espoo, Finland, 3-9 June 2019*, pp. 183–200, Springer, 2022.
- Smith, M. M., Thomson, J., Baker, M. G., Abbott, R. E., and Davis, J.: Observations of ocean surface wave attenuation in sea ice using seafloor cables, *Geophysical Research Letters*, 50, e2023GL105243, 2023.
- 370 Spreen, G., Kaleschke, L., and Heygster, G.: Sea ice remote sensing using AMSR-E 89-GHz channels, *Journal of Geophysical Research: Oceans*, 113, 2008.
- Squire, V. A.: A fresh look at how ocean waves and sea ice interact, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 376, 20170342, 2018.
- Squire, V. A.: Ocean wave interactions with sea ice: a reappraisal, *Annual Review of Fluid Mechanics*, 52, 37–60, 2020.
- 375 Squire, V. A., Vaughan, G. L., and Bennetts, L. G.: Ocean surface wave evolution in the Arctic Basin, *Geophysical Research Letters*, 36, 2009.
- Stopa, J. E., Ardhuin, F., and Girard-Ardhuin, F.: Wave climate in the Arctic 1992–2014: Seasonality and trends, *The Cryosphere*, 10, 1605–1629, 2016.
- Stopa, J. E., Sutherland, P., and Ardhuin, F.: Strong and highly variable push of ocean waves on Southern Ocean sea ice, *Proceedings of the*
380 *National Academy of Sciences*, 115, 5861–5865, 2018.
- Thomson, J.: Wave propagation in the marginal ice zone: connections and feedback mechanisms within the air–ice–ocean system, *Philosophical Transactions of the Royal Society A*, 380, 20210251, 2022.
- Thomson, J., Hošeková, L., Meylan, M. H., Kohout, A. L., and Kumar, N.: Spurious rollover of wave attenuation rates in sea ice caused by noise in field measurements, *Journal of Geophysical Research: Oceans*, 126, e2020JC016606, 2021.
- 385 Toyota, T., Takatsuji, S., and Nakayama, M.: Characteristics of sea ice floe size distribution in the seasonal ice zone, *Geophysical research letters*, 33, 2006.
- Voermans, J. J., Rabault, J., Filchuk, K., Ryzhov, I., Heil, P., Marchenko, A., Collins III, C. O., Dabboor, M., Sutherland, G., and Babanin, A. V.: Experimental evidence for a universal threshold characterizing wave-induced sea ice break-up, *The Cryosphere*, 14, 4265–4278, 2020.
- 390 Voermans, J. J., Liu, Q., Marchenko, A., Rabault, J., Filchuk, K., Ryzhov, I., Heil, P., Waseda, T., Nose, T., Kodaira, T., et al.: Wave dispersion and dissipation in landfast ice: comparison of observations against models, *The Cryosphere*, 15, 5557–5575, 2021.
- Voermans, J. J., Xu, X., and Babanin, A. V.: Validity of the wave stationarity assumption on estimates of wave attenuation in sea ice: toward a method for wave–ice attenuation observations at global scales, *Journal of Glaciology*, 69, 803–810, 2023.
- Wadhams, P., Squire, V. A., Goodman, D. J., Cowan, A. M., and Moore, S. C.: The attenuation rates of ocean waves in the marginal ice zone,
395 *Journal of Geophysical Research: Oceans*, 93, 6799–6818, 1988.
- Weeks, W.: *On sea ice*, University of Alaska Press, 2010.
- Williams, T. D., Bennetts, L. G., Squire, V. A., Dumont, D., and Bertino, L.: Wave–ice interactions in the marginal ice zone. Part 2: Numerical implementation and sensitivity studies along 1D transects of the ocean surface, *Ocean Modelling*, 71, 92–101, 2013.
- Womack, A., Alberello, A., de Vos, M., Toffoli, A., Verrinder, R., and Vichi, M.: A contrast in sea ice drift and deformation between winter
400 and spring of 2019 in the Antarctic marginal ice zone, *The Cryosphere*, 18, 205–229, 2024.

<https://doi.org/10.5194/egusphere-2024-2104>

Preprint. Discussion started: 6 August 2024

© Author(s) 2024. CC BY 4.0 License.



Worby, A. P., Geiger, C. A., Paget, M. J., Van Woert, M. L., Ackley, S. F., and DeLiberty, T. L.: Thickness distribution of Antarctic sea ice, *Journal of Geophysical Research: Oceans*, 113, <https://doi.org/10.1029/2007jc004254>, 2008.

Young, I. R., Fontaine, E., Liu, Q., and Babanin, A. V.: The wave climate of the Southern Ocean, *Journal of Physical Oceanography*, 50, 1417–1433, 2020.