

Age of air from ACE-FTS measurements of sulfur hexafluoride

Laura N. Saunders¹, Kaley A. Walker¹, Gabriele P. Stiller², Thomas von Clarmann^{2,†}, Florian Haenel², Hella Garny^{3,4}, Harald Bönisch², Chris D. Boone⁵, Ariana E. Castillo⁶, Andreas Engel⁷, Johannes C. Laube⁸, Marianna Linz^{6,9}, Felix Ploeger^{8,10}, David A. Plummer¹¹, Eric A. Ray^{12,13}, and Patrick E. Sheese¹

¹Department of Physics, University of Toronto, Toronto, Canada

²Institute of Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

³Deutsches Zentrum für Luft- und Raumfahrt (DLR), Institut für Physik der Atmosphäre, Oberpfaffenhofen, Germany

⁴Ludwig-Maximilians-University Munich, Meteorological Institute, Munich, Germany

⁵Department of Chemistry, University of Waterloo, Waterloo, Canada

⁶Department of Earth and Planetary Sciences, Harvard University, Cambridge, USA

⁷Institute for Atmospheric and Environmental Science, Goethe University Frankfurt, Frankfurt, Germany

⁸Institute of Climate and Energy Systems, Stratosphere (ICE-4), Forschungszentrum Jülich, Jülich, Germany

⁹School of Engineering and Applied Sciences, Harvard University, Cambridge, USA

¹⁰Institute for Atmospheric and Environmental Research, University of Wuppertal, Wuppertal, Germany

¹¹Climate Research Division, Environment and Climate Change Canada, Montreal, Canada

¹²Chemical Sciences Division, Earth Systems Research Laboratory, NOAA, Boulder, USA

¹³Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder, Boulder, USA

[†]deceased

Correspondence: Kaley A. Walker (kaley.walker@utoronto.ca)

Abstract. Climate models predict that the Brewer-Dobson Circulation (BDC) will accelerate due to tropospheric warming, leading to a redistribution of trace gases and, consequently, to a change of the radiative properties of the atmosphere. Changes in the BDC are diagnosed by the so-called “age of air”, that is, the time since air in the stratosphere exited the troposphere. These changes can be derived from a long-term observation-based record of long-lived trace gases with increasing concentration in the troposphere, such as sulfur hexafluoride (SF₆). The Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS) provides the longest available continuous time series of vertically-resolved SF₆ measurements, spanning 2004 to the present. In this study, a new age of air product is derived from the ACE-FTS SF₆ dataset. The ACE-FTS product is in good agreement with other observation-based age of air datasets and shows the expected global distribution of age of air values. Age of air from a chemistry climate model is evaluated and the linear trend of observation-based age of air is calculated in twelve regions within the lower stratospheric midlatitudes (14-20 km, 40-70°) in each hemisphere. All trends are negative and significant to two standard deviations. The mean trend in the northern hemisphere (southern hemisphere) is -0.10 ± 0.01 (-0.11 ± 0.02) years per decade. This is therefore the first observation-based age of air trend study to suggest an acceleration of the shallow branch of the BDC, which transports air poleward in the lower stratosphere, in both hemispheres.

1 Introduction

15 While it is well-known that increased greenhouse gas concentrations are causing a warming of the troposphere and a cooling of the stratosphere, so far, few observational datasets constrain how this affects the Brewer-Dobson Circulation (BDC). The BDC is characterized by poleward motion in the stratosphere through two distinct branches (Plumb, 2002; Birner and Bönisch, 2011). The shallow branch transports air from the tropics to higher latitudes in the lower stratosphere (below ~ 20 km), with transit timescales on the order of months to a couple years. The deep branch brings air poleward during the winter at higher
20 altitudes, with transit timescales on the order of more than two years. Model studies suggest that both branches of the BDC are accelerating (e.g., Butchart, 2014, and references therein), but this has been difficult to confirm with measurements.

The speed of the BDC cannot be measured directly, so it must be inferred from other quantities. One way to do this is to estimate the age of air, that is, the time that has passed since air left the troposphere (Waugh and Hall, 2002). Since air typically enters the stratosphere through tropical upwelling and is then subjected to the BDC as it moves poleward, this definition of
25 age of air is a useful indicator of BDC-related mean transport time. At a particular location, younger ages correspond with shorter transit times, while older ages correspond with longer transit times. It is also necessary to address the fact that any finite region under consideration (that is, a macroscopic “air parcel”) is made up of air of different ages. The distribution of these ages is referred to as the age spectrum (Hall and Plumb, 1994), which can be modelled as an inverse Gaussian function with a mean (i.e., the mean age of stratospheric air) and a width. Mixing between air parcels leads to wider age spectra, as it
30 increases the number of different pathways leading to a specific location. A decrease in the mean age of air at higher latitudes, with no change in the tropics, indicates a decrease in the mean transport time from the tropics to higher latitudes, and therefore an acceleration of the BDC (e.g. Neu and Plumb, 1999; Linz et al., 2016, 2017). This motivates the need for trend detection in age of air derived from observations.

Age of air can be estimated from long-lived trace gases that have increasing concentrations in the troposphere, also known
35 as “clock” tracers. Once an air parcel containing such a gas enters the stratosphere, the volume mixing ratio (VMR) does not change due to the absence of chemical sinks (hence the long lifetime), and the concentration is therefore an indicator of when the parcel left the troposphere. CO_2 as well as fluorocarbons and hydrofluorocarbons are some commonly used clock tracers, but one of the most often used is SF_6 , which according to surface measurements (e.g., Levin et al., 2010; Rigby et al., 2010; Simmonds et al., 2020) has been steadily increasing in the troposphere since the mid-20th century due to industrial activity.
40 It has effectively no seasonal cycle or stratospheric sinks, which complicate the use of CO_2 and CFCs, respectively, but SF_6 is destroyed in the mesosphere via photolysis and electron-capture reactions (Morris et al., 1995; Reddmann et al., 2001; Totterdill et al., 2015).

There are several aircraft- and balloon-borne in situ measurements of SF_6 available from the last three decades that have been used to infer age of air (e.g., Volk et al., 1997; Engel et al., 2002). Those that were used in this study are highlighted here.
45 Harnisch et al. (1996) calculated age using cryosampler measurements of SF_6 from the AIRE 93 and SESAME 95 balloon flights over France (44° N) in September 1993 and Sweden (68° N) in March 1995, respectively. Andrews et al. (2001) derived age of air from SF_6 measured by the LACE instrument on board three balloon flights over the USA (35° N and 65° N) and

Brazil (7° S) throughout 1996 and 1997. For that study, age of air was also derived using NASA ER-2 aircraft-based SF₆ measurements taken by the ACATS-IV instrument between 1992 and 1997. The former two studies also derived age of air using CO₂ measurements taken during the same campaigns, and Boering et al. (1996) did so using CO₂ measurements from these ER-2 flights. More recently, Leedham Elvidge et al. (2018) and Adcock et al. (2021) calculated age of air using SF₆ measurements based on whole air samples taken during M55 Geophysica high-altitude aircraft flights. The former used one campaign over Germany in October-November 2009 (48-54° N) alongside two campaigns over Sweden (62-77° N) in January-February 2010 and December 2011. The latter involved one campaign over Greece (33-41° N) in September 2016 and another over Nepal (21-29° N) in July-August 2017. Both studies also determined an “averaged age of air” using measurements of multiple other clock tracers besides SF₆ taken during the same campaigns.

The only vertically-resolved global age of air dataset currently available is derived from MIPAS (Michelson Interferometer for Passive Atmospheric Sounding) measurements of SF₆ and spans the 2002-2012 period (Stiller et al., 2008, 2012; Haenel et al., 2015). This spatially and temporally dense satellite dataset has been invaluable in age of air studies, but the relatively short time series makes trend detection challenging. Hardiman et al. (2017) used model simulations to determine that natural variability could mask changes in the BDC for a time series shorter than 12 years, like that available from MIPAS. This motivates the development of a new satellite-based age of air dataset using ACE-FTS (Atmospheric Chemistry Experiment Fourier Transform Spectrometer), which provides the longest available time series of vertically-resolved global SF₆ measurements. In this study, ACE-FTS retrievals of SF₆ were used to calculate age of air for the February 2004-February 2021 period.

Several studies have used existing measurement-based age of air datasets as well as reanalysis data to evaluate model predictions regarding the acceleration of the BDC. Engel et al. (2009) compiled age of air derived from balloon-borne cryosampler measurements of CO₂ and SF₆ in the northern midlatitude stratosphere (24-35 km) from 1975-2005 into a single time series. Over this 30-year period, a slight positive but statistically insignificant change was detected. Additional balloon data taken in 2015 and 2016 were added to this time series and resulted in an even smaller and still statistically insignificant positive trend (Engel et al., 2017). Bönisch et al. (2011) identified a change in the slopes of O₃-N₂O correlations using aircraft-based in situ measurements taken between 1979 and 2009. This was linked to increased upwelling in the tropics, a potential sign of acceleration of the BDC. Within the same study, the observation was corroborated with Japanese 25-year Reanalysis data (Onogi et al., 2007), which showed a 1 month per decade decrease in transit times into the midlatitude lower stratosphere, suggesting a slight acceleration of the shallow branch. However, the same reanalysis data indicated that the deep branch did not accelerate over this time period. Diallo et al. (2012) also used reanalysis data, from ERA-Interim, to calculate the trend in age of air for the 1989-2010 period. A significant negative trend of 3.5-6 months per decade was found in the southern (and northern equatorward of 40° N) lower stratosphere, but no clear trend was detected in the mid-stratosphere. This also suggests that the shallow branch of the BDC is accelerating while there is no clear trend in the deep branch. Ray et al. (2014) used an idealized model constrained by balloon-borne in situ measurements of SF₆ and CO₂ to assess variability in northern hemisphere age of air and identified a negative trend in the lowest 5 km of the stratosphere, but a positive trend at higher altitudes. Haenel et al. (2015) found similar results in trends based on the MIPAS age of air dataset: negative trends in the lowermost tropical and southern midlatitude stratosphere but positive trends in the northern midlatitudes and at both poles, all on the order of a few

months to a year per decade. The persistent finding across these studies is that, while there is no clear signal (in observations or reanalyses) that the BDC is accelerating as a whole, the shallow branch in the southern hemisphere does appear to have
85 accelerated.

This study details the development of the ACE-FTS age of air dataset as follows. Section 2 describes the ACE-FTS and MIPAS SF₆ datasets, which are compared with each other and with in situ SF₆ measurements in Section 3. Section 4 provides an explanation of how age of air was calculated from ACE-FTS SF₆ measurements, taking into account the complication arising from the sink due to photolysis in the mesosphere. The ACE-FTS age of air results are presented in Section 5. These
90 include an exploration of the effects of the mesospheric sink correction, comparisons with age of air previously derived from in situ and newly derived from MIPAS measurements, a comparison with age of air output from a specified dynamics (nudged) simulation with the Canadian Middle Atmosphere Model (CMAM39), and the calculation of trends in age of air for the 2004-2021 period.

2 Satellite datasets

95 The focus of this study is mean age of stratospheric air (hereafter referred to as simply “age of air”) based on SF₆ measurements from ACE-FTS, for which age of air is calculated for the first time. An updated age of air product based on MIPAS data is also presented, since the details of the method for age of air calculation have evolved since Stiller et al. (2012) and Haenel et al. (2015).

2.1 ACE-FTS

100 ACE-FTS is a high spectral resolution (0.02 cm^{-1}) infrared Fourier-transform spectrometer on board the satellite SCISAT that has been taking measurements since February 2004 (Bernath et al., 2005). It is a solar occultation instrument; the concentrations of atmospheric gases are determined by fitting their known absorption spectra to the spectra obtained during each sunrise or sunset. The retrieval algorithms for versions 3 and 4 are described in detail in Boone et al. (2013) and Boone et al. (2020), respectively. During each occultation, ACE-FTS takes measurements at various altitudes, supplying a profile with 2-3 km
105 vertical resolution which is provided on a 1 km grid. From an orbital inclination of 74°, ACE-FTS measurements reach up to 85° in each hemisphere. It has been making up to 30 occultations per day (2 per orbit) since early 2004, compiling a dataset that contains over one hundred thousand profiles from 6 km up to 150 km, depending on the gas, concentrated at higher latitudes (Mahieu et al., 2008). Due to the nature of solar occultation measurements, ACE-FTS has limited sampling. It takes three months for ACE-FTS to obtain global coverage and the measurements are not distributed evenly in latitude throughout each
110 month. This must be taken into consideration when comparing ACE-FTS with other instruments or with models, usually by subsampling the larger dataset and finding coincident measurements.

Despite the relative sparsity of ACE-FTS measurements when compared with other satellite instruments, its orbital pattern and retrieval method offer several advantages for this study. The first is that solar occultation measurements have a higher signal-to-noise ratio than measurements of atmospheric emission, which is particularly useful for a gas like SF₆ with such

115 a low VMR (i.e., a few parts per trillion). Additionally, the deep branch of the BDC is imprinted in downwelling over polar regions, so age of air at higher latitudes is of interest for inferring changes related to this. The ACE-FTS sampling pattern is suited for this purpose as nearly 50% of profiles are taken poleward of 60° . Most importantly, the 17-year time series of consistent SF_6 measurements is unmatched by previous datasets. The lack of measurement density does present a challenge, but the advantages of using ACE-FTS for age of air studies far outweigh this difficulty.

120 Here, two versions of ACE-FTS were considered: v3.5/3.6 (Feb 2004-Feb 2021) and v4.1/4.2 (Feb 2004-present). v3.5/3.6 was included because the retrieval update for v4 resulted in an unphysical feature in SF_6 between 18 and 30 km. The same issue has been observed in the current ACE-FTS dataset, v5.2. The ACE-FTS data are accompanied by a set of quality flags that indicate whether a measurement is recommended for use. The flags for v4 are provided separately by Sheese and Walker (2020) and for v3, the flags are included in the ACE-FTS VMR files. These quality flags were developed by Sheese et al. (2015) and they were employed as recommended, keeping only measurements with a flag of 0 and removing profiles that contained any measurements with a flag of 4 or 5. There is a drift in v3 known to affect ozone and temperature profiles, and potentially others, that was corrected for in v4 (Boone et al., 2020).

2.2 MIPAS

MIPAS was a Fourier transform spectrometer on the Envisat satellite that measured infrared limb emission spectra from July 2002 to April 2012 (Fischer and Oelhaf, 1996; Fischer et al., 2008). Technical problems in 2004 caused a reduction in spectral resolution from 0.025 cm^{-1} to 0.0625 cm^{-1} (unapodized), but a 20% increase in the number of measured profiles, starting in January 2005. Envisat was in a Sun-synchronous orbit with an inclination of 98.55° and an altitude of 800 km, completing nearly 15 revolutions per day and about 100 scans per orbit, amounting to approximately 1500 profiles per day. Scans were taken at 10 AM or 10 PM local solar time (LST) and covered 87° S - 89° N . The profiles are provided on a uniform altitude grid from 6 to 70 km. The SF_6 retrievals are done by the Institut für Meteorologie und Klimaforschung (IMK) together with the Instituto de Astrofísica de Andalucía (IAA) and are described in von Clarmann et al. (2003), with the changes made for the reduced resolution period explained in von Clarmann et al. (2009). In this study, the lower resolution SF_6 measurements from version 5 (V5R) (Haenel et al., 2015) were used to maximize temporal coverage while using a single dataset. Specifically, V5R_222 and V5R_223 were used because they are the most recent SF_6 products that were retrieved using the same spectroscopic parameters (Varanasi et al., 1994) as versions 3 and 4 of ACE-FTS. As recommended by the instrument team, all SF_6 measurements with a visibility flag of 0 and averaging kernel diagonal less than 0.03 were removed.

SF_6 and derived age of air from MIPAS were both considered for comparison purposes. To account for the difference in measurement density between MIPAS and ACE, the comparisons between the two were done using both the full MIPAS V5R dataset and a subsampled version that only contains measurements which are coincident with an ACE-FTS measurement. MIPAS V5R SF_6 measurements were sampled onto the ACE-FTS grid by, for each ACE-FTS measurement, finding the closest MIPAS measurement that is within 12 h and 1000 km and on the same side of the polar vortex edge (i.e. both are inside, both are outside, or both are inside the edge). The polar vortex edge is defined as any point where the absolute value of scaled potential vorticity (sPV; e.g., Dunkerton and Delisi, 1986; Manney and Zurek, 1993; Manney et al., 1994) is between $1.2 \times 10^{-4} \text{ s}^{-1}$

and $1.6 \times 10^{-4} \text{ s}^{-1}$. Lower sPV values are considered to be outside the vortex while higher values are considered to be inside the vortex. For this step, each altitude was considered separately, meaning that one profile could include some measurements inside the vortex and some measurements outside the vortex. sPV values at the ACE-FTS and MIPAS measurement locations were taken from the Jet Propulsion Laboratory Derived Meteorological Products (JPL DMPs, Manney et al., 2007), which are based on NASA’s Global Modeling and Assimilation Office’s MERRA-2 (Modern Era Retrospective analysis for Research and Applications) (Gelaro et al., 2017). The generous 1000 km coincidence criterion for distance was chosen to maximize the number of coincidences; using stricter coincidence criteria did not significantly improve the comparisons. Approximately half the ACE-FTS measurements over the relevant time period had a coincident MIPAS measurement.

3 SF₆ comparisons

Prior to this study, ACE-FTS SF₆ measurements (version 3.0) had only been compared with output from the SLIMCAT 3D chemical transport model, with good agreement found at all altitudes between 12 and 33 km (Brown et al., 2011). Here, a comparison of mean profiles from ACE-FTS v3.5/3.6 and v4.1/4.2 for the tropics (20° S-20° N), the extratropics (20° up to polar vortex edge, $|\text{sPV}| < 1.2 \times 10^{-4} \text{ s}^{-1}$), and the polar vortex ($|\text{sPV}| > 1.6 \times 10^{-4} \text{ s}^{-1}$) are shown in Fig. 1. The same occultations were used from each version for these comparisons. The error bars represent one standard error of the mean at each altitude. The v4.1/4.2 profiles, shown in dark blue (solid), have a decrease in SF₆ near 18 km accompanied by a slight increase just above. This artifact is a result of how altitudes are determined in the retrieval (private communication, C. Boone, May 2024). The previous version, v3.5/3.6, which is shown in light blue (solid), does not have this feature and was therefore chosen for this study. It should be noted that the decrease in SF₆ below about 14 km is caused by the fact that the altitude of the lowest retrieval varies over the course of the year. Therefore, the low-altitude bins can be missing measurements from certain times of year, resulting in this feature. Hereafter, “ACE-FTS” refers to v3.5/3.6. Comparisons with other SF₆ datasets are presented in the following section.

3.1 Comparisons with MIPAS and in situ data

Figure 1 also shows profile comparisons of coincident ACE-FTS and MIPAS measurements averaged over 2005-2012 as light blue and orange dashed lines, respectively. In all regions, particularly outside the polar vortex, SF₆ decreases more rapidly with height in the stratosphere in the MIPAS measurements. The difference between the mean profiles is greatest in the extratropics, where ACE-FTS VMRs are greater than MIPAS by up to about 10%. Within the polar vortex, SF₆ decreases to about 4.5 pptv at 30 km in both datasets, but the vertical gradients between 20 and 30 km differ. The MIPAS profile decreases more quickly from 20-25 km, but the ACE-FTS profile decreases more quickly from 25-30 km until the profiles meet again near 30 km. The largest differences, about 10%, are at 24-25 km. It is also noteworthy there is a difference of about 1 pptv between the ACE-FTS profile from the full dataset (2004-2021) and that for only measurements that are coincident with MIPAS (2005-2012). NOAA Marine Boundary Layer (MBL) reference SF₆ (Global Monitoring Laboratory, 2023) increased from about 6.5 pptv to 7.5 pptv between 2008 and 2012 (the midpoints of the two time periods), which accounts for the discrepancy. This reinforces




Figure 1. Mean profiles of SF₆ retrievals from ACE-FTS v3.5/3.6 (light blue), ACE-FTS v4.1/4.2 (dark blue), and coincident measurements from ACE-FTS v3.5/3.6 and MIPAS V5R (dashed light blue and dashed orange, respectively) for (a) the tropics (20° S-20° N), (b) the northern extratropics (north of 20° N and absolute sPV less than $1.2 \times 10^{-4} \text{ s}^{-1}$), (c) the northern polar vortex (absolute sPV greater than $1.6 \times 10^{-4} \text{ s}^{-1}$), (d) the southern extratropics (south of 20° S and absolute sPV less than $1.2 \times 10^{-4} \text{ s}^{-1}$), and (e) the southern polar vortex (absolute sPV greater than $1.6 \times 10^{-4} \text{ s}^{-1}$). Error bars represent the standard error of the mean.

the importance of using coincident measurements when comparing observations from two different instruments; comparing the full ACE-FTS mission with the full MIPAS mission would introduce significant bias due to the two different time periods. The mean profile from the full MIPAS mission is similar to that from the sampled MIPAS dataset, and is not shown because it makes the comparison between the mean coincident profiles more difficult to visualize.

185 ACE-FTS and MIPAS SF₆ were also compared with balloon-borne profiles from the BONBON cryosampler (e.g., Engel et al., 2006; Laube et al., 2010). There were three flights available during the ACE-FTS mission: two over Brazil near 5° S on 8 June and 25 June 2005 (B42 and B43), and one over Sweden near 67° N on 10 March 2009 (B45). Following Kolonjari et al. (2024), where ACE-FTS measurements were compared with B42 and B43, the coincidence criteria for these comparisons were quite broad: within 90 days and within 5° latitude, with no restriction on longitude. This ensured that there would be
 190 a sufficient number of coincidences even though ACE-FTS profiles are sparse in the tropics, particularly in June when the satellite's beta angle is high. Using these criteria, the B42 and B43 measurements were each coincident with 20-35 ACE-FTS

measurements and up to over 2000 MIPAS measurements, depending on the altitude. The B45 measurements had over 400 coincident ACE-FTS measurements and over 8000 coincident MIPAS measurements.




Figure 2. SF_6 profiles from three BONBON balloon flights (black triangles) on (a) 8 June 2005 (B42) and (b) 25 June 2005 (B43), both over Brazil near 5° S, and on (c) 10 October 2009 over Sweden near 68° N (B45). Also shown are means of SF_6 profiles from ACE-FTS (blue circles) and MIPAS (orange squares) that are within 90 days and 5° latitude of each BONBON profile. The error bars represent one standard deviation.

The SF_6 values from each flight were compared with the means of coincident ACE-FTS or MIPAS measurements, shown in Fig. 2. The error bars represent one standard deviation. The agreement between all three datasets is excellent; the VMRs are consistent everywhere except near 21–23 km for flight B45 in Kiruna, where the balloon measurements are lower than the ACE-FTS measurements by about 15%. This flight intersected with the polar vortex, which could cause a sharp decrease in SF_6 like that seen in this altitude region. Because of the necessarily broad definitions for coincidence, it is not expected that the satellite data show this isolated dynamical feature. Despite this, the comparison was included due to the limited availability of balloon-based SF_6 profiles for reference during the ACE-FTS measurement period.

Overall, there is good consistency between the ACE-FTS SF_6 profiles and the MIPAS and BONBON datasets. ACE-FTS and MIPAS agree within 10% in all latitude regions, and coincident profiles from both datasets agree within one standard deviation with the BONBON profiles.

4 Methodology for deriving age of air

205 The convolution method described in Garny et al. (2024b), slightly updated since Stiller et al. (2012) and Haenel et al. (2015), was used to infer age of air from SF_6 measurements. An overview of the method is provided here. For consistent comparisons, MIPAS age of air was recalculated for this study using the updated method. To reduce noise, the satellite instrument datasets were first zonally averaged by month, separately for each year (i.e., each bin corresponds to one month in one year). For most of this study, 10° latitude bins (centered on 85°S , 75°S , ..., 85°N) and 3 km altitude bins (centred on 9.5, 12.5, ..., 210 30.5 km) were used. In addition, a dataset with 5° latitude bins and one 2 km altitude bin from 19-21 km was also created for comparison purposes. There is also an age of air product provided using 10° equivalent latitude bins based on the equivalent latitude provided as part of the JPL DMPs. Similarly to the ACE-FTS climatologies described in Koo et al. (2017), at least five measurements were required in each bin (defined by month, year, latitude, and altitude). The mean SF_6 value in each bin was then used to calculate mean age of air; essentially, each bin was treated as an “air parcel”.

215 4.1 Tropospheric SF_6 reference time series

Since age of air is roughly defined as the time since an air parcel left the troposphere, a tropospheric SF_6 time series is required as a reference to match the parcel’s SF_6 VMR with a tropospheric value and time. Specifically, a tropical tropospheric reference time series is needed as it is assumed that most air enters the stratosphere in the tropics. Such a time series is only available from surface measurements, so the age of air calculation includes a small correction to account for the difference between air 220 at the surface and air at the tropopause (see below). The time series used here is made from a combination of datasets. Of most importance are the tropical NOAA MBL reference data, which were used from mid-1995 onward. For an accurate calculation, the reference time series should extend back about 15 years from the earliest stratospheric SF_6 value to allow for the detection of old air. In previous studies (e.g., Stiller et al., 2008, 2012; Haenel et al., 2015), the time series was linearly extrapolated to obtain values for earlier years. Here, the time series was extended back to 1989 using scaled monthly data from Cape Grim in 225 Tasmania (Ray et al., 2014).

4.2 Surface correction

The age of air calculation uses surface SF_6 measurements as a tropospheric reference, but to find the time since entering the stratosphere, the stratospheric SF_6 VMRs should be compared with VMRs at the tropical tropopause. While SF_6 concentrations are not expected to decrease significantly from the surface to the tropopause since the transit timescale is on the order of weeks 230 or months at most, a more accurate age can be calculated if the satellite data are corrected to account for this difference. Further, any measurement bias between the ground-based data and the satellite observations can be detected and corrected for by this method.

This was done by finding a bias correction that shifts the tropical tropospheric ACE-FTS values to the reference time series values and applying it to all ACE-FTS measurements. The tropics were defined to be those profiles between 17.5°S and 235 17.5°N , for consistency with Stiller et al. (2012) and to avoid influence from the extratropics. The troposphere was defined

as those measurements that are at least 2 km below the WMO tropopause, which is another JPL DMP calculated for each ACE-FTS profile based on the algorithm described in Reichler et al. (2003). This was to reduce the influence of troposphere-stratosphere mixing in the upper troposphere and therefore ensure that only the well-mixed portion of the troposphere was included. This correction also accounts for instrument bias and for the possibility of a satellite measurement detecting both tropospheric and stratospheric air due to coarser vertical resolution. If the satellite VMRs are too low relative to the reference time series, for example, then ages calculated based on the reference time series will be too high. Ensuring agreement between the two datasets in the troposphere reduces the influence of these biases.

The monthly tropical tropospheric SF₆ means from ACE-FTS are shown as blue circles in the first panel of Fig. 3 along with the reference time series (black line). There is good agreement, mostly within $\pm 5\%$, but after 2014, the majority of ACE-FTS measurements fall below the reference time series. The differences could be due to changes in measurement density, which increased over time, or the known drift in v3.5/3.6 retrievals. This discrepancy further motivates a correction to the ACE-FTS measurements. The relative differences between monthly mean tropical tropospheric SF₆ from ACE-FTS and the surface monthly means are shown in the second panel of Fig. 3. A cubic least squares fit, shown by the red line, was used to find the relative bias r_t for each month of the time series. Linear and quadratic fits resulted in noticeable structure in the fit residuals, so a cubic fit was chosen. The final panel in Fig. 3 shows the reference time series and the corrected ACE-FTS tropical tropospheric values (blue squares). The corrected time series was calculated by dividing each VMR in the time series by $1+r_t$. The corrected ACE-FTS values no longer drift relative to the reference time series after 2014. The correction was applied to all ACE-FTS SF₆ climatological means based on month and year but not based on latitude or altitude, since latitude and altitude dependence cannot be determined without similarly reliable reference data for other regions, which are not available. This implies that the correction is understood to address a drift in ACE-FTS measurements that appears in all measurements at the same time, independent of their location. The correction is negative throughout most of the time period, meaning that the stratospheric ACE-FTS values were mostly increased. Therefore, the surface correction in this case will result in a younger age of air, by at most a few months.

A similar method was applied to the MIPAS dataset. The correction was linear and there was a mean positive bias of 1.8%, which was removed by dividing the VMR in each bin by 1.018.

4.3 Age of air calculation

If an air parcel were not subject to any mixing, then the amount of SF₆ within it could simply be matched to the reference time series and the time lag would correspond to the parcel's exact age. However, this is never a practical assumption, so it is necessary to consider mixing and treat measured (and modelled) air parcels with an age spectrum. The age spectrum was approximated using an inverse Gaussian function (Waugh and Hall, 2002):

$$G(\Gamma, \Delta, t) = \frac{\Gamma^{3/2}}{2\Delta\sqrt{\pi t^3}} \exp\left(\frac{-\Gamma^3(t/\Gamma - 1)^2}{4\Delta^2 t}\right), \quad (1)$$




Figure 3. (a) Monthly tropical tropospheric SF_6 time series from the NOAA MBL reference data (black line) and ACE-FTS (blue circles, see text for definition of tropics and troposphere). (b) Relative difference between the monthly ACE-FTS SF_6 values and the reference time series (red squares), with a cubic fit to the data (red line). (c) Same as (a) but with the surface correction applied to the ACE-FTS values (blue squares).

where t is the transit time, Γ is the mean age and Δ is the width of the distribution. Prior studies assumed that the “ratio of moments”, $w = \Delta^2/\Gamma$, was a constant value of 0.7 years as in Waugh and Hall (2002). For this study, the ratio of moments for each latitude and altitude was taken from a model study based on a Chemical Lagrangian Model of the Stratosphere (CLaMS) run (Ploeger et al., 2021) that used winds from ERA-Interim and an exponential fit correction to fit the tail of the spectrum. This ratio of moments is assumed to be constant in time, which is consistent with the use of an inverse Gaussian for the age spectrum. For more details on this approach, see Garny et al. (2024b). Generally, this parameter is the source of considerable uncertainty as w values are much larger for ERA5 than for ERA-Interim, for example. ERA-Interim was chosen because of

its close agreement to the observed circulation based on age of air (Linz et al., 2017). The w value varies from about 0.5 to 1.5 years, with the highest values being found in the extratropical lower stratosphere.

Given that each air parcel has an age distribution rather than one single age, the goal of this calculation is to find the mean of the age spectrum for each parcel (i.e., bin). This requires determining which age spectrum, that is, which Γ value, would yield the bin's mean SF_6 VMR (denoted $[\text{SF}_6]_{\text{measured}}$, following the formalism in Stiller et al. (2012)). Since it is not analytically possible to determine the age spectrum based on $[\text{SF}_6]_{\text{measured}}$ directly, the inverse problem must be solved instead. That is, given a mean age Γ and a distribution width Δ , the goal is to find the stratospheric entry time and the corresponding tropospheric SF_6 value. This is essentially the same as treating the age spectrum as a probability density function to find an expectation value for $[\text{SF}_6]_{\text{measured}}$. Specifically, the following integral is evaluated:

$$[\text{SF}_6]_{\text{measured}} = \int_0^{\infty} \tilde{G}(\Gamma, \Delta, t) [\text{SF}_6]_{\text{trop}}(t_{\text{meas}} - t) dt, \quad (2)$$

where $\tilde{G}$ is the normalized age spectrum, t is the transport time, and t_{meas} is the time of the stratospheric measurement. $[\text{SF}_6]_{\text{trop}}(t)$, the tropospheric reference time series (see Section 4.1), is evaluated at $t_{\text{meas}} - t$ because the stratospheric SF_6 VMR should correspond to the tropospheric value from t years before the measurement was taken.

The inverse problem is then solved iteratively. The first guess Γ_0 is taken to be the time lag between a stratospheric SF_6 measurement and the time at which that SF_6 value was found in the troposphere ($t_{\text{trop}, 0}$) according to the reference time series. Γ_0 is then used in Equation 2 along with the distribution width (using $\Delta_0 = \sqrt{w\Gamma_0}$) for the air parcel's latitude and altitude to obtain the expected stratospheric SF_6 value, denoted hereafter by $[\text{SF}_6]_{\text{modelled}}$. A new guess for the tropospheric time ($t_{\text{trop}, \text{new}}$) is then calculated using Newton's method:

$$t_{\text{trop}, \text{new}} = t_{\text{trop}, 0} + \frac{[\text{SF}_6]_{\text{measured}} - [\text{SF}_6]_{\text{modelled}}}{d[\text{SF}_6]_{\text{trop}}(t_{\text{trop}, 0})/dt}. \quad (3)$$

The new guess for the age Γ_{new} is then taken to be $t_{\text{meas}} - t_{\text{trop}, \text{new}}$. This process is repeated, using $4\Gamma_{\text{new}}$ as the upper bound on the integral with 1000 intervals, until the difference between $[\text{SF}_6]_{\text{modelled}}$ and $[\text{SF}_6]_{\text{measured}}$ is below 0.01 pptv, which is well below the standard error of the mean SF_6 value in each bin and typically allows convergence within five iterations.

The uncertainty on the age of air in each bin was estimated in two ways. The first was by propagating the standard deviation of the SF_6 VMR through a simplified age of air calculation. That is, the upper bound on the age was taken to be the lag time between the average SF_6 VMR plus the standard deviation and the equivalent point on the tropospheric reference curve. The lower bound was estimated the same way but using the SF_6 VMR minus the standard deviation. The "standard deviation of the age of air" was taken to be half of the range between these two bounds. The same calculation was also done using the standard error of the mean SF_6 VMR to determine the "standard error of the mean age of air". Both of these uncertainties are provided in the data files.

4.4 Mesospheric sink correction

The calculation described above assumes that SF_6 has no chemical sources or sinks in an air parcel observed in the stratosphere. However, some air in the stratosphere has travelled through the mesosphere and therefore been depleted of SF_6 . This is especially true at higher altitudes and within the polar vortex. In these regions, SF_6 has a lower VMR for a reason other than older age of air, which the above calculation does not account for. Garny et al. (2024a) have established a correction scheme to address this bias in the “apparent” (i.e., uncorrected) age and determine the “ideal” (i.e., sink-corrected) age. Garny et al. (2024a) showed that the apparent age $\tilde{\Gamma}$ is approximately related to the ideal age Γ by the following:

$$\tilde{\Gamma} = \Gamma \left(1 + \frac{F_t(t_{\text{meas}})}{\tau_{\text{eff}}} \right), \quad (4)$$

where τ_{eff} is the effective lifetime of stratospheric SF_6 , t_{meas} is the stratospheric measurement time, $F_t(t_{\text{meas}})$ is the ratio of the reference tropospheric mixing ratio to its time derivative evaluated at $t_{\text{meas}} - \Gamma_0$, and Γ_0 is the linearization point for the Taylor expansion of the reference time series $[\text{SF}_6]_{\text{trop}}(t_{\text{meas}})$ (see Garny et al. (2024a) for details). As recommended by Garny et al. (2024a), Γ_0 was set to 4 years and τ_{eff} was calculated using $\tau_{\text{eff}} = \alpha \exp(\beta \Gamma)$, where α and β are 5.21×10^2 and -3.81×10^{-1} , respectively. $F_t(t)$ was calculated using the reference time series described in Section 4.1, with the gradient in the denominator calculated over a five year period.

For each apparent age value at time $t_{\text{meas},i}$, that is, the age derived for each bin in the ACE-FTS and MIPAS SF_6 climatologies, $F_t(t_{\text{meas}})$ was determined for the relevant year and Equation 4 was solved numerically to find the ideal age. The uncertainty on sink-corrected age of air was estimated by applying the sink correction scheme to the age plus the estimated uncertainty and to the age minus the estimated uncertainty, and using half the range of these two values. This was done for both the standard deviation and the standard error of the mean. It is important to note that the sink correction only depends explicitly on time and not on altitude or latitude. However, there is implicit altitude and latitude dependence since the ideal age depends on the apparent age, and the apparent age changes with altitude and latitude. This is not a comprehensive treatment of the spatial dependence, as 2-year-old air in the tropics has less mesospheric influence than 2-year-old air in the poles, for example. In addition, the sink correction is so far only derived from model data, and there is uncertainty in how well the model simulates the sink. In particular, the sink correction cannot completely remove the bias for air older than about five years (Garny et al., 2024a).

5 Results and discussion

The multi-year zonal mean apparent age of air derived from ACE-FTS (i.e., without the sink correction) is shown in Fig. 4. The annual means from 2004 through 2021 were averaged together so that each year was weighted equally. As expected, age increases with height and with latitude following the flow of the BDC. The apparent age is older in the southern hemisphere at high altitudes because the polar vortex is stronger and SF_6 -poor air descending from the mesosphere is kept isolated from younger midlatitude air. In the remainder of this section, the ACE-FTS derived age of air is investigated in more detail through an exploration into the effects of the mesospheric sink correction and comparisons with age of air derived from MIPAS and

335 from other observational datasets. Finally, applications of the ACE-FTS derived age of air product are demonstrated through a comparison with CMAM39 and an exploration of the long-term trend in age of air.




Figure 4. Multi-year zonal mean apparent age of air based on the ACE-FTS SF₆ dataset (February 2004-February 2021).

5.1 Effect of the mesospheric sink correction

This section explores the effect of the mesospheric sink correction described in Section 4.4 on age of air derived from both ACE-FTS and MIPAS. Figure 5 shows the zonal mean ages derived from ACE-FTS for each season with (middle row) and without (top row) the sink correction, alongside the difference (bottom row). The absolute difference uses the uncorrected age as the reference, so negative values correspond to a decrease in age as a result of the sink correction. The influence of mesospheric air can be seen throughout the seasons at higher altitudes and latitudes. Air appears oldest in the upper stratosphere at high southern latitudes during austral winter and spring where subsidence within the polar vortex is strongest. A similar aging effect can be seen in the northern polar vortex during boreal winter, but to a lesser extent due to increased mixing with the midlatitudes. The absolute differences between the uncorrected and corrected zonal means are below 2 years throughout most of the stratosphere, but are quite large in these polar regions with high subsidence. In the southern polar vortex, the sink correction decreases the derived age by up to 6 years.

The sink correction has a similar effect on MIPAS ages, as shown in Fig. 6. Note that here, the full MIPAS dataset has been used. Similarly to ACE-FTS, the effect of the sink correction is amplified at high latitudes. The correction has a larger effect on MIPAS than on ACE-FTS at midlatitudes between 20 and 30 km. This makes sense given that the MIPAS ages in these regions are older than those from ACE-FTS, particularly above 25 km where the sink correction decreases the age by up to 3 years even in the lower midlatitudes.




Figure 5. (a) Multi-year zonal mean age of air from the full ACE-FTS v3.5/3.6 dataset (February 2004-February 2021). Each column represents one season. (b) Same as (a) but with the sink-correction applied. (c) Absolute differences between the sink-corrected and uncorrected ACE-FTS ages (corrected minus uncorrected).

For both datasets, even with the sink correction, the air at high latitudes and altitudes is significantly older than in other regions. This difference could be at least partially explained by the fact that the sink correction is known to undercorrect for air older than about five years (Garny et al., 2024a). Further validation of the SF_6 sink correction beyond the model-based analysis of Garny et al. (2024a) would require the comparison against age of air estimates from other clock tracers, which are not available so far and so this is left for future studies. However, one such comparison has been done for vortex air from the SOLVE campaign for the northern hemisphere, and CO_2 ages reached about 6 years between 25-32 km (Ray et al., 2017). This is only from one flight, however, and is not necessarily representative.




Figure 6. Same as Fig. 5 but for the full MIPAS V5R dataset (2005-2012).

360 5.2 Comparison with MIPAS

For the comparisons shown here, only coincident ACE-FTS and MIPAS measurements were considered, so the number of measurements included in each bin is reduced compared to when the full datasets are used.

Figure 7 shows the multi-year seasonal zonal mean uncorrected age of air for ACE-FTS (top row) and MIPAS (middle row), and the differences between them (bottom row). The general patterns in the zonal means are the same. Age of air increases with
 365 both height and latitude, and is high inside the polar vortex, partially due to the influence of mesospheric air (only relevant for uncorrected age) and partially due to the longer transit times to these locations. The agreement is also good, within 6 months, in the mid- to high latitudes at below about 20 km, which is an important region when investigating the BDC. This is the endpoint of the shallow branch of the BDC, so if the satellite-based ages are consistent here, then they can be used to investigate changes in the circulation with more certainty. MIPAS shows older air than ACE-FTS almost everywhere except near the tropopause
 370 and above 29 km poleward of 60° S. This is not surprising given that MIPAS SF₆ VMRs are smaller than those from ACE-FTS

above 20 km, particularly outside the polar vortex, with a stronger vertical gradient such that the inter-instrument difference becomes larger with altitude (see Section 3). Below 20 km, the ACE-FTS and MIPAS ages are generally within a year of each other except in the tropics from 17–20 km, where MIPAS air is up to about 2 years older due to the reduced measured SF_6 values in this region (see Fig. 1).




Figure 7. Multi-year (2005–2012) zonal mean uncorrected age of air derived from coincident row (a) ACE-FTS and row (b) MIPAS measurements. Row (c) shows the absolute differences (MIPAS–ACE-FTS). Each column corresponds to one season.

375 The comparisons between the sink-corrected ACE-FTS and MIPAS ages are shown in Fig. 8. The main conclusions are similar, though the difference in age at higher altitudes is reduced because of MIPAS ages being more affected by the sink correction than those from ACE-FTS. This is simply because the sink correction affects older ages more than younger ages, and MIPAS ages are older due to having smaller SF_6 VMRs compared to ACE-FTS. The two datasets have better agreement with the sink correction having been applied, especially in the mid- to high latitudes above 20 km where the difference between
380 the ages decreased from up to 6 years to typically less than 4 years. The tropics are still quite different above 25 km where the

difference is still about 3-4 years, however this is not surprising as the sink correction does not have as much of an effect on younger air.




Figure 8. Same as Fig. 7 but with the sink correction applied to both datasets.

The ACE-FTS and MIPAS ages show similar patterns, though the MIPAS ages are up to three years (more than 50%) older at higher altitudes and latitudes with the mesospheric sink correction applied. It is clear that the difference in vertical gradient of measured SF_6 between the two instruments impacts the derived age of air. Future studies should include comparisons to datasets with ages derived using other clock tracers where ACE-FTS and MIPAS have better agreement.

5.3 In situ comparisons

In situ measurements are a particularly useful reference for comparison. Here, ages calculated from several balloon and aircraft campaigns from 1993 to 2017 (discussed in Section 1) are compared with the ACE-FTS age of air dataset. Both the sink-corrected and uncorrected ACE-FTS ages are included in these comparisons since the in situ ages use both SF_6 and other

Table 1. Overview of the in situ datasets used for comparison with the ACE-FTS age of air product. The table is divided into campaigns that occurred during the ACE-FTS mission (top) and those that occurred prior to the mission (bottom). The former are from aircraft campaigns while the latter are balloon campaigns.

Aircraft campaign	Date	Location	Latitude	Altitude (km)	Gas(es)	Reference
OB09	Oct. 30, 2009	Oberpfaffenhofen, Germany	48-54° N	10-20 km	SF ₆ , C ₃ F ₈ ,	Leedham Elvidge et al. (2018)
	Nov. 4, 2009				HFC-125, HFC-227ea	
K2010	Jan. 20, 2010	Kiruna, Sweden	62-77° N	9-19 km	SF ₆ , C ₂ F ₆ ,	Leedham Elvidge et al. (2018)
	Feb. 2 2010				C ₃ F ₈ , CHF ₃ ,	
K2011	Dec. 11/16, 2011	Kiruna, Sweden	62-77° N	9-19 km	HFC-125, HFC-227ea	Leedham Elvidge et al. (2018)
					SF ₆ , CF ₄ ,	
					C ₂ F ₆ , CHF ₃ ,	
AMO-16	Sep. 1/6, 2016	Kalamata, Greece	33-41° N	10-20 km	HFC-227ea	Adcock et al. (2021)
AMO-17	Jul. 27/29/31, 2017	Kathmandu, Nepal	21-29° N	10-20 km	SF ₆ , C ₂ F ₆ , HFC-125	Adcock et al. (2021)
	Aug. 2/4/6, 2017				SF ₆ , C ₂ F ₆ , HFC-125	
Balloon campaign	Date	Location	Latitude	Altitude (km)	Gas(es)	Reference
AIRE 93	Sep. 30, 1993	Aire sur l'Adour, France	44° N	6-34 km	SF ₆	Harnisch et al. (1996)
SESAME 95	Mar. 7, 1995	Kiruna, Sweden	68° N	6-34 km	SF ₆	Harnisch et al. (1996)
OMS	Sep. 21, 1996	Fort Sumner, USA	35° N	9-26	CO ₂	Andrews et al. (2001)
					SF ₆	Ray et al. (1999)
OMS	Feb. 14, 1997	Juazeiro do Norte, Brazil	7° S	9-32 km	CO ₂	Andrews et al. (2001)
	Nov. 11/20, 1997				SF ₆	Ray et al. (1999)
OMS	Jun. 30, 1997	Fairbanks, USA	65° N	9-32 km	CO ₂	Andrews et al. (2001)
					SF ₆	Ray et al. (1999)

tracers such as CO₂. The uncorrected ACE-FTS ages are compared with in situ ages based on SF₆, which are also uncorrected, while the sink-corrected ACE-FTS ages are compared with those from other tracers with no or very small mesospheric sinks.

Table 1 summarizes the details of the aircraft and balloon campaigns used for comparison. For each aircraft flight, age of air has been derived from SF₆ along with several fluorine-containing compounds (see references in Table 1 for age calculation used for each flight). As such, two types of age are shown in Fig. 9: that derived from SF₆ (grey triangles, top row) and the averaged age derived from all measured clock tracers (black circles, bottom row). The uncorrected ACE-FTS ages are shown in the top row, while the sink-corrected ages are shown in the bottom row. The nearest bin (or in some cases, bins) in space and time from the ACE-FTS age dataset was (were) selected for comparison, and each bin is shown as one point with error bars representing one standard deviation of the age of air (as described above). Where multiple bins with the same altitude were used, the points are separated vertically on the plot for better visibility, but the bins are all centred on the same altitude (9.5, 12.5, 15.5, or 18.5 km).




Figure 9. Comparisons between the ACE-FTS age of air dataset and ages based on aircraft in situ measurements taken during the ACE-FTS mission. Each column corresponds to one campaign (a) OB09, (b) K2010, (c) K2011, (d) AMO-16, and (e) AMO-17. See Table 1 for further details. Black dots: mean age of air based on all clock tracers included in the campaign. Grey triangles: age of air based on SF_6 . Open coloured circles: mean age of air for one bin in the ACE-FTS dataset (top: uncorrected, bottom: sink-corrected). Points for ACE-FTS bins centred on the same altitude are separated vertically for visibility. Error bars show the standard deviation of the age in each bin.

In nearly all cases, the ACE-FTS ages agree with the aircraft data (i.e., the aircraft data are within one standard deviation of the satellite data). One exception are the two ACE-FTS bins at 12.5 km in October and November 2009 for both the uncorrected and the sink-corrected ages. The age in both latitude bins is 1-2 years older (more than one standard deviation) than the nearest aircraft age at 13 km. This is related to the unrealistic decrease in ACE-FTS SF_6 below about 14 km, which biases the derived age older. In the case of the November 2009 bin centred on 55° N at 15.5 km, the ACE-FTS age is also more than one standard deviation from the nearest aircraft age. This can likely be explained by sampling differences; the aircraft age is based on a measurement taken at 48° N and the ACE-FTS bin does not include this latitude. Conversely, the ACE-FTS age in the bin

centred on 45° N does include 48° N and is consistent with the aircraft age. Sampling differences can also explain why a few
410 aircraft ages also fall outside one standard deviation from the ACE-FTS ages in December 2011. The ACE-FTS bin is from
January 2012 while the aircraft data were taken in mid-December 2011. There are no cases where the sink-corrected ages
are consistent while the uncorrected ages are not, and vice versa. This was expected, since the uncorrected ages were only
compared with similarly uncorrected age derived from SF₆ and the sink-corrected ages were only compared with ages based
on other clock tracers.

415 The ACE-FTS ages were also compared with the age derived from five balloon campaigns prior to the beginning of the
satellite mission. Here, only the time of year was considered (as opposed to the exact month and year), so a balloon campaign
from September 1993, for example, was compared with all Septembers of the ACE-FTS mission averaged together. This
introduced a temporal bias, especially considering the fact that age of air is potentially changing over time. As such, these
comparisons, which are shown in Fig. 10, are more useful for looking at the vertical structure and tentatively comparing age
420 of air between two time periods. For these comparisons, the sink-corrected ACE-FTS ages are shown on the same plots as the
uncorrected ages, but as filled markers rather than open markers. The error bars represent the standard deviation of the mean
age of air in the bins that were averaged together across all years. The sink-corrected ages are best compared to the in situ ages
derived from CO₂, so they are not included in comparisons with flights without CO₂ ages (September 1993 and March 1995).
Two times of year were considered for the comparison with the 1997 campaign in Brazil because it included flights in both
425 February and November. As before, the two ACE-FTS bins are vertically separated for visibility, but are still centred on the
same altitude.

The agreement between the ACE-FTS and balloon ages varies by location. A common feature in the in situ ages, visible in
the profiles from AIRE 93 (September 1993), SESAME 95 (March 1995), and OMS 1996 (September 1996), is a constant age
above about 20-25 km. The ACE-FTS profiles do not have this feature. Aside from this difference, the ACE-FTS ages agree
430 with the AIRE 93 (September 1993) ages below 30 km. At 30.5 km, however, the ACE-FTS age is nearly 4 years (100%) older.
The ACE-FTS ages are consistently 1-2 years older than the SESAME 95 (March 1995) ages. This is consistent with a change to
the northern deep branch of the BDC between 1995 and the mid-2000s, but could also be due to interannual variability. For the
OMS flight over the southern USA (September 1996), the uncorrected ages are consistently 1-2 years younger than the in situ
ages derived from SF₆, except at 30.5 km. The sink-corrected ages agree with the in situ CO₂ ages below 20 km, but are up to
435 one year younger above this point. The differences at lower levels (below about 25 km) could be consistent with an acceleration
of the shallow branch of the BDC. The ACE-FTS ages are mostly consistent with those derived from measurements over Brazil
(February and November 1997), though the variability is high. The uncorrected ACE-FTS ages mostly agree with in situ SF₆
ages from the flight over Alaska (June 1997) within a few percent. The SF₆ in situ ages are variable, and Andrews et al. (2001)
attribute this to remnants of the polar vortex containing mesospheric air. The sink-corrected ACE-FTS ages agree with the
440 CO₂ in situ ages within a few percent everywhere except 20-25 km, where the ACE-FTS ages are slightly (up to about 1 year)
younger. This is also consistent with a BDC acceleration, but the difference is small relative to those seen for SESAME 95 and
OMS 1996, and the presence of the polar vortex significantly complicates the comparison. Overall, the significant variability
in age over time inhibits trend interpretation from these limited in situ comparisons.




Figure 10. Comparisons between the ACE-FTS age of air dataset and ages based on balloon in situ measurements taken prior to the ACE-FTS mission. Each panel corresponds to one in situ campaign (a) AIRE 93, (b) SESAME 95, (c)-(e) OMS. See Table 1 for further details. Grey triangles: age of air based on SF₆. Grey squares: mean age of air based on all clock tracers included in the campaign. Open coloured circles: average uncorrected age of air for a set of bins (e.g., all Septembers at 45°N) in the ACE-FTS dataset. Filled coloured circles: average sink-corrected age of air for a set of bins in the ACE-FTS dataset. Points for ACE-FTS averages centred on the same altitude are separated vertically for visibility. Error bars correspond to one standard deviation of the mean ages from the bins that were averaged together.

The zonal mean ages derived from the full ACE-FTS and MIPAS datasets were also compared with a latitudinal cross-section of ages based on aircraft measurements of SF₆ and CO₂ taken between 1992 to 1997 (Boering et al., 1996; Elkins et al., 1996), shown in Fig. 11. It is important to note that the in situ ages derived from SF₆ have not been sink-corrected, so they are equivalent to the uncorrected ages from ACE-FTS and MIPAS (solid lines). As before, the sink-corrected ACE-FTS and MIPAS ages (dashed lines) should be more similar to the in situ ages derived from CO₂. The error bars on the ages based on CO₂ indicate 1 σ uncertainty. The in situ measurements were all taken between 19.5 and 20.5 km, so the ACE-FTS and MIPAS ages for this comparison were calculated using the SF₆ climatologies with the 19-21 km altitude bin. Additionally, these ages were calculated for 5° latitude bins to increase the resolution of the latitudinal cross-section.

Figure 11 shows that uncorrected ACE-FTS ages agree better with the in situ ages derived from SF₆ than the uncorrected MIPAS ages in the tropics and high latitudes, but they are slightly too young (by less than 1 year) in the midlatitudes. With the sink correction applied, ACE-FTS ages are still younger than those from MIPAS and the aircraft data, but they are still mostly consistent with the ages based on CO₂, except between about 20° S and 40° S and near 60° in both hemispheres. It is possible that the differences between the aircraft and ACE-FTS altitude and latitude sampling result in ACE-FTS ages being




Figure 11. Age of air as a function of latitude, based on in situ measurements of SF_6 (black triangles), in situ measurements of CO_2 (grey triangles with 1σ error bars), ACE-FTS (blue lines), and MIPAS (orange lines). The dashed lines represent the sink-corrected ages from ACE-FTS and MIPAS. The in situ measurements are aircraft measurements taken between 1992 and 1997 at 19–21 km. The ACE-FTS and MIPAS ages are based on the full datasets, with all measurements taken between 19 and 21 km averaged into 5° latitude bins.

younger. In particular, the aircraft data could show more distinct mixing barriers than are visible in the satellite data, which average over more time and area. The difference could also be due to the time periods covered by the two datasets (2004–2021 for ACE-FTS vs. 1993–1997 for the aircraft data). If so, this result would indicate that age of air is decreasing over time in the midlatitudes, which could be consistent with an acceleration of the shallow branch of the BDC. A conclusion cannot yet be
460 drawn regarding whether the apparent change in age of air over time is due to biases between the different SF_6 measurements or an actual decrease, so further analysis is required.

5.4 Comparison with CMAM39

CMAM39 is a chemistry-climate model based on the extended version of the Canadian Centre for Climate Modelling and
465 Analysis third-generation Atmospheric General Circulation Model (Scinocca et al., 2008) and includes an extensive description of stratospheric chemistry (de Grandpré et al., 2000; Jonsson et al., 2004). The meteorological fields (temperature and horizontal winds) in the specified dynamics version are “nudged” toward ERA-Interim reanalysis (Dee et al., 2011) up to 1 hPa

with a relaxation time constant of 24 hours. The nudging constrains the large-scale dynamics to follow the reanalysis, allowing for a direct comparison between the model and measurements, though nudging has also been shown to modify certain aspects of the BDC as compared with a free-running simulation (Chrysanthou et al., 2019). The modelled age of air is based on an idealised SF₆ tracer, meaning that it is not subject to the mesospheric sink. Therefore, the sink-corrected ACE-FTS ages were used for this comparison, which spans the 2004-2018 period where ACE-FTS and CMAM39 overlap. To account for sampling differences, CMAM39 has been subsampled onto the ACE-FTS times and locations using an updated version of the advanced method described in Kolonjari et al. (2018). Here, the modelled values were also linearly interpolated in time instead of using a nearest neighbour approach.

The comparisons for each season are shown in Fig. 12, where the ACE-FTS ages are shown in the top row, the CMAM39 ages are shown in the middle row, and the differences (CMAM39–ACE-FTS) are shown in the bottom row. ACE-FTS and CMAM39 agree within 6 months below about 24 km, and it is noteworthy that the modelled ages are mostly younger than the observed ages. Kolonjari et al. (2018) found evidence that the BDC in CMAM30, the predecessor to CMAM39 that was also nudged to ERA-Interim, was too fast, which would result in younger stratospheric air and be consistent with these results. It was also found that at pressures less than 150 hPa, the age in CMAM30 was up to one year older than the age in the free-running version, suggesting that nudging affects the strength of the BDC and tracer transport. The largest differences between the ACE-FTS and CMAM39 ages are at high altitudes and latitudes, particularly in the winter and spring where ACE-FTS air is 3 years older in the southern hemisphere. This is not necessarily entirely due to an issue in the model; ACE-FTS ages were also shown to be older than in situ ages at higher altitudes in Fig. 10. It is also possible that the sink correction does not fully account for the high age bias in ages derived from SF₆ measurements. This could be investigated using comparisons between CMAM39 and ages based on other clock tracers.

5.5 Long-term age of air trend

One of the main advantages of the ACE-FTS age of air product is the long 17-year time series that version 3.5/3.6 provides. While the MIPAS product has excellent global coverage and density, the mission lasted only about a decade, which makes it difficult to detect a long-term trend in a climate system impacted by multi-year oscillations such as the Quasi-Biennial Oscillation (QBO), the El Niño Southern Oscillation (ENSO), and the solar cycle. On the other hand, the relative sparsity of ACE-FTS measurements also causes some difficulty in accounting for these oscillations. Each 10° latitude band does not have an ACE-FTS age value for every month as it does for MIPAS, and fitting a time series with gaps is more challenging. Given these limitations, the aim of this work is to investigate whether significant long-term changes in age of air can be detected from ACE-FTS, despite the sparse sampling. The focus is on regions where ACE-FTS has the highest density of successful SF₆ retrievals: the lower stratosphere (14-20 km) between 40 and 70° latitude in each hemisphere. While the profiles at these latitudes do extend higher into the stratosphere, the number of measurements in each bin is reduced at higher altitudes, increasing the standard error of the zonal mean and therefore the uncertainty in the age, which makes trend detection more difficult. Age of air in the lower stratosphere is largely influenced by the shallow branch of the BDC, so changes in this region




Figure 12. Multi-year (2004-2018) zonal mean age of air derived from row (a) ACE-FTS and row (b) CMAM39 subsampled to the ACE-FTS measurement grid, with the absolute differences (CMAM39-ACE, row (c)). Each column corresponds to one season.

can indicate whether the shallow branch might be changing. A trend was calculated based on the monthly values for each (10° , 3 km) bin within these two regions.

To calculate the linear trend of each age of air time series, a similar approach to the one presented in Stiller et al. (2012) and Haenel et al. (2015) was used. Each sink-corrected time series was fitted using the following function:

$$\Gamma(t) = a + bt + c_1 \text{qbo}_1(t) + d_1 \text{qbo}_2(t) + c_2 \text{enso}(t) + c_3 \text{sol}_1(t) + d_3 \text{sol}_2(t) + \sum_{n=4}^5 \left(c_n \sin \frac{2\pi t}{l_n} + d_n \cos \frac{2\pi t}{l_n} \right), \quad (5)$$

where t is time in years. The QBO and ENSO variations are included via the proxies qbo_1 , qbo_2 , and enso . Previous analyses of longer time series of age of air from MERRA2 reanalysis data and of merged observed water vapour (SWOOSH) have also

found decadal oscillations, the causes of which are not completely clear (Strahan et al., 2020; Tao et al., 2023). In order to account for these, an oscillation term with a period of about 10 years was included. In the numerical approach, the solar cycle was used for this with proxies sol_1 and sol_2 . There is some justification for this given the link between the solar cycle and the velocity of tropical upwelling (Kodera and Shibata, 2006), but this is not sufficient evidence to assume that the decadal oscillation is driven by the solar cycle alone. The final sum term represents the annual and semi-annual cycles, where $l_4 = 1$ year and $l_5 = 0.5$ years. Other harmonic oscillations were omitted because they resulted in an overfit of the sparse ACE-FTS time series and did not significantly affect the linear term, b . Several versions of this function were tested with different combinations of proxies for the QBO, ENSO, and the decadal variation, and it was determined that including all of them produced the best results. That is, the fit residuals had the least structure when more natural variability was accounted for. To allow for temporal offsets between the QBO time series and the age response at different locations, two orthogonal terms were included. The same holds for the decadal variation. This model differs from that used in Haenel et al. (2015), which used eight different harmonics and did not consider the lower frequency oscillations (ENSO and the decadal variation). The latter two were included here because the length of the time series is nearly doubled and more than one period for each oscillation is covered by the time series. The qbo_1 and qbo_2 vectors are monthly orthogonal normalized Singapore winds at 30 and 50 hPa, respectively (Freie Universität Berlin, 2022). The ENSO proxy is the Multivariate ENSO Index (NOAA, 2022). The sol_1 term is the monthly adjusted solar flux at 10.7 cm as determined from daily measurements taken in Penticton, Canada at the Dominion Radio Astrophysical Observatory provided by NOAA (2018) from February to September 2004 and by Space Weather Canada (2022) from October 2004 to February 2021. The orthogonal term, sol_2 , is from the same dataset but shifted by 2 years and 9 months, that is, by one quarter of the solar period of approximately 11 years. As mentioned above, the so-called solar cycle term is included to account for any decadal variation, and is not necessarily believed to have solar variability as its cause. Any variation longer than approximately 10 years would not make sense to be fitted to a 17-year time series. With these proxies, the ACE-FTS age of air data were fit with Equation 5 using the method described in von Clarmann et al. (2010) for irregularly spaced and correlated data. This method uses Gaussian error propagation of the standard error of the monthly zonal mean to calculate the uncertainty on the linear trend. Uncertainties due to the sink correction are not considered.

The fitted curves for the northern and southern hemisphere, with the linear component plotted separately (black dashed line), are shown in Figures 13 and 14, respectively. For the most part, the fit represents the data fairly well; the residuals did not show noticeable structure. The clearest feature in all bins is the annual cycle, which appears to be mostly well-captured though perhaps with a slightly underestimated amplitude at lower latitudes. Similarly to what was observed by Stiller et al. (2012), the amplitude of the seasonal cycle is larger at higher altitudes.

As an example, the contributions of each type of interannual variability are summarized for 65° N (14-17 km) in Fig. 15. This summary is shown for the other 11 bins in the Supplementary Information. The QBO, ENSO, decadal variation, and semiannual cycles each have similar amplitudes of 1-2 months, with the decadal variation being of particular interest as it is noticeable in the time series shown in Fig. 13 at 55 and 65° N (14-17 km). It appears as a low-frequency oscillation causing an increase in age early in the time series up to around 2010, a decrease until 2016, and an increase afterwards. The solar cycle proxy seems to account for this oscillation well; in every bin, the fit residuals are smaller and have less structure when




Figure 13. Monthly age of air derived from the ACE-FTS dataset (blue squares), with the full fit described in the text (black solid line) and its linear component (black dashed line). Each panel (a) through (f) corresponds to one northern hemispheric bin in the lower stratosphere, indicated in the figure.

it is included in the fit. However, even if the time series is fit without the solar cycle term in Eq. 5, the linear trend remains qualitatively the same (i.e., negative) and is significantly different from zero to two standard deviations. Therefore the finding that age of air decreases over the full observation period is independent of the assumption of a solar cycle. The mean trend in the midlatitude lower stratosphere is -0.10 ± 0.01 years per decade in the northern hemisphere and -0.11 ± 0.02 years per decade in the southern hemisphere, where the uncertainty is the standard error of the mean. These findings are consistent with an acceleration of the shallow branch of the BDC in both hemispheres, if tropical ages are not changing, and are therefore in agreement with the results in Ray et al. (2014).

These trends are not consistent with what Haenel et al. (2015) found for this region, which was that the trends in the northern hemisphere region were positive but insignificant and most trend values in the southern hemisphere were negative, but also insignificant except for 40-60° N from approximately 18-20 km. There are several reasons that the trends would not agree. As




Figure 14. Same as Fig. 13 but for the southern hemisphere.

shown in Section 5.2, there are discrepancies between the MIPAS and ACE-FTS age of air datasets. Furthermore, Haenel et al. (2015) did not apply a sink correction to the MIPAS ages, and a sink correction will change the trend (Loeffel et al., 2022).

555 The regression model used here was also altered through the addition of the decadal variation and ENSO and the removal of several harmonics. Another important difference is that the MIPAS trends were calculated for 2002-2012, while the ACE-FTS trends cover the 2004-2021 time period. To explore the impacts of these differences, the trends in the ACE-FTS age of air were calculated over the period that overlapped with the MIPAS mission, January 2005-April 2012. Since only eight years of data were included, ENSO and the decadal variation were omitted from the regression, making the fit more similar to that used for

560 MIPAS data in Haenel et al. (2015), though with fewer harmonics as the ACE-FTS data are not as dense. Additionally, the ACE-FTS time series here start in 2005 rather than in 2002 like the time series that were analyzed in Haenel et al. (2015). The resulting fits for the northern and southern hemispheres are shown in Figures 16 and 17, respectively. In all cases, the fits appear to represent the data well (based on the lack of structure in the residuals), but the linear term is positive and larger in magnitude than for the fits over the longer time series. This is not unexpected; the fits to the long-term data suggested that the




Figure 15. (a) The contribution (in years) of each type of interannual variability to the fitted age of air for 60-70°N, 14-17 km. (b) The total contribution of the 12- and 6-month seasonal cycles.

age was increasing up until 2010 due to the decadal variation, which covers most of the MIPAS mission. Here, however, the decadal variation is not included in the fit so this increase is attributed purely to a long-term increase in the age of air. The same fits were attempted with the decadal variation included, and the linear term remained positive in about half of the bins as the short time series obscures the difference between variability due to the decadal variation and that due to a real trend.

The trends from Figs. 13 through 17 are summarized in Fig. 18, with the full ACE-FTS trends shown in the top panel and the trends for the truncated period (2005-2012) shown in the middle panel. In addition, the linear trends for the MIPAS ages are shown in the bottom panel for comparison with the ACE-FTS trends for the same period. The MIPAS trends were also calculated without including ENSO or the solar cycle proxies. These trends are not equivalent to those in Haenel et al. (2015), which are mostly consistent with zero in these regions, because they include two harmonics instead of eight, ENSO and the solar cycle are not included, MIPAS data from 2002-2004 are not included, and the trends are based on ages produced using the slightly modified age calculation method. The crosses show where the MIPAS and shortened ACE-FTS trends agree within two standard deviations. The trends are consistent within half of the bins. The differences could be partially explained by sampling differences between the instruments; using only coincident measurements resulted in too few profiles to achieve a




Figure 16. Same as Fig. 13 but for only January 2005 to April 2012.

reasonable fit. Qualitatively, the two datasets agree that the trend is positive for the 2005-2012 period throughout the regions under consideration.

580 While rigorous trend estimation remains challenging, valuable conclusions can be drawn from the analysis presented in this section. The comparison between the 2005-2012 and the 2004-2021 ACE-FTS trends shows that the linear term is sensitive to the length of the time series. This is expected due to the high variance in the age of air datasets; it is difficult to separate natural variability from a long-term trend, especially a small one, with a short time series. The 2004-2021 time series revealed a decadal variation signal in the time series, which appeared as a positive trend for the MIPAS mission period in both the ACE-FTS and MIPAS datasets. This is an example of how natural variability can explain short-term trends, and that these trends
 585 therefore should not be extrapolated beyond the time period covered by the dataset from which they were inferred. This further demonstrates the importance of a long age of air time series. A similar phenomenon was observed by Strahan et al. (2020) in the MIPAS age of air data, who showed that the MIPAS time series happens to have been taken during one of the largest oscillations in age of air on record and thus must not be over-interpreted as representative of longer timescales. The trends in




Figure 17. Same as Fig. 14 but for only January 2005 to April 2012.

the ACE-FTS age of air data are consistent with an acceleration of the shallow branch of the BDC in both hemispheres, if there is no change in tropical age. This is consistent with the decrease in the shallow branch transit time found by Bönisch et al. (2011) over the 1979-2009 period. The decrease in age of air in the 60-70° N bins and the 60-70° S bin at 15.5 km would imply that the deep branch is also accelerating if tropical ages are constant. A denser age of air time series at even higher latitudes would be useful for corroborating whether ages there are decreasing.

6 Summary

An age of air dataset was derived from the 17-year ACE-FTS v3.5/3.6 SF₆ time series. The dataset exhibited the expected qualitative pattern based on air motion due to the BDC: air was shown to be older at higher latitudes and altitudes. A correction was applied to account for the influence of the mesospheric SF₆ sink on the calculated age of air; this was found to reduce the derived age and was strongest in polar regions where mesospheric air enters the stratosphere. The ACE-FTS ages displayed the




Figure 18. Zonal distribution of the linear trends for (a) age derived from ACE-FTS over the 2004-2021 period, (b) age derived from ACE-FTS over the 2005-2012 period, and (c) age derived from MIPAS over the 2005-2012 period. The crosses indicate where the ACE-FTS 2005-2012 and MIPAS 2005-2012 trends are within two standard deviations of each other.

600 same behaviour as the ages derived from MIPAS, but MIPAS showed significantly older air at higher latitudes and altitudes. The differences were reduced with the sink correction applied to both datasets, but the ages still differed by up to roughly 3 years. However, ACE-FTS age was mostly consistent with ages derived from in situ aircraft measurements, with a few exceptions that could be explained by location sampling differences between the two instruments or by proximity to the tropopause or the polar vortex edge. Comparisons with age of air derived from balloon measurements of SF_6 and CO_2 from before the ACE-FTS mission are consistent with an acceleration of the shallow branch of the BDC, and possibly a slowing of the deep branch
605 FTS mission are consistent with an acceleration of the shallow branch of the BDC, and possibly a slowing of the deep branch between 1995 and the mid-2000s, though this was based on comparisons with single balloon flights. Both of these indications would be consistent with previously determined trends in age of air derived from observations or from reanalysis data.

Two applications of the ACE-FTS age of air dataset were demonstrated. First, the sink-corrected ages were compared with the age product from CMAM39. It was found that the modelled age was too young compared with the ACE-FTS age, except
610 near the tropical tropopause, which is consistent with previous studies suggesting that the BDC is too fast in the model. The ACE-FTS age of air time series was also used for more detailed examination of the trend through regression analysis. While there remain several limitations to this analysis, the results corroborate what has been found in several other studies. Since ACE-FTS uses solar occultation, it has a high signal-to-noise ratio compared to other remote sensing instruments, but the measurements are spatially sparse. This means that there are significantly fewer measurements in each bin in the age dataset and
615 not every month has a corresponding age in a given latitude bin. While the time series are longer than what has been available for age of air until now on a global scale, these attributes make it nevertheless difficult to account for natural variability. In this study it was determined that despite this, including oscillations such as the QBO, ENSO, and a decadal variation approximated by the solar cycle, using respective proxy data, improved the regression analysis. The linear trend was calculated for bins in the mid- to high latitudes (40-70°) in the lower stratosphere (14-20 km) because these bins have the most data points and are a
620 meaningful location for examining the BDC. A negative linear trend significant to two standard deviations was found in every bin that was considered; the mean trend was -0.10 ± 0.01 years per decade in the northern hemisphere and -0.11 ± 0.02 years per decade in the southern hemisphere. These results are consistent with an acceleration of the shallow branch of the BDC. This was previously seen by Bönisch et al. (2011) for the 1979-2009 period and Haenel et al. (2015) for the 2002-2012 period in parts of southern hemisphere, however without statistical significance. Due to the inclusion of the 60-70° bins, the results may
625 also indicate an acceleration of the deep branch of the BDC, which has not previously been seen with observational data. These results agree, at least qualitatively, with climate model predictions of changes in atmospheric circulation. However, they are prone to uncertainty from natural variability, and longer data records are required for increasing reliability of trend estimates.

Data availability. The age of air datasets are available at <https://doi.org/10.5683/SP3/5AC1F0>. The ACE-FTS Level 2 data used in this study can be obtained via the ACE-FTS website (registration required, <https://database.scisat.ca/>). The MIPAS SF₆ data can be obtained
630 from <https://www.imk-asf.kit.edu/english/308.php>. The CMAM39 data are available through anonymous ftp at [crd-data-donnees-rtc.ec.gc.ca/pub/CCCMA/dplummer/CMAM39-SD_6hr](ftp://crd-data-donnees-rtc.ec.gc.ca/pub/CCCMA/dplummer/CMAM39-SD_6hr).

Author contributions. This study was designed by KAW, GPS, and TvC. LNS performed the analysis and wrote the manuscript, building on prior work from GPS, TVC, and FH. HG developed the sink correction scheme. CDB and PES provided their expertise on ACE-FTS; AE and JCL on the balloon and aircraft data; DAP on the CMAM39 output; and EAR on the SF₆ reference time series. ML and FP provided the
635 ratio of moments based on the CLaMS run. AC tested the age of air dataset. Valuable discussions were provided by all authors throughout the study, as were comments on the manuscript.

Competing interests. At least one co-author is an editor of Atmospheric Chemistry and Physics.

Acknowledgements. This project was funded by grants from the Canadian Space Agency (CSA) and the Natural Sciences and Engineering Research Council of Canada (NSERC). The Atmospheric Chemistry Experiment (ACE) is a Canadian-led mission mainly supported by the
640 CSA. The development of the CMAM39 dataset was funded by the CSA. We thank Peter Bernath for his leadership of the ACE mission. We thank Ted Shepherd, Dylan Jones, John Scinocca, and David Plummer for their leadership and support of the CMAM39-SD project. AE and FP acknowledge funding from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – TRR 301 – Project-ID 428312742.

645 We also wish to acknowledge the significant contribution of Thomas von Clarmann, who passed away just as this work was being completed. He provided excellent guidance and encouragement throughout this study and is greatly missed by his collaborators and friends.

References

- Adcock, K. E., Fraser, P. J., Hall, B. D., Langenfelds, R. L., Lee, G., Montzka, S. A., Oram, D. E., Röckmann, T., Stroh, F., Sturges, W. T., Vogel, B., and Laube, J. C.: Aircraft-Based Observations of Ozone-Depleting Substances in the Upper Troposphere and Lower Stratosphere in and Above the Asian Summer Monsoon, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033137, <https://doi.org/10.1029/2020JD033137>, 2021.
- Andrews, A. E., Boering, K. A., Daube, B. C., Wofsy, S. C., Loewenstein, M., Jost, H., Podolske, J. R., Webster, C. R., Herman, R. L., Scott, D. C., Flesch, G. J., Moyer, E. J., Elkins, J. W., Dutton, G. S., Hurst, D. F., Moore, F. L., Ray, E. A., Romashkin, P. A., and Strahan, S. E.: Mean ages of stratospheric air derived from in situ observations of CO₂, CH₄, and N₂O, *Journal of Geophysical Research: Atmospheres*, 106, 32 295–32 314, <https://doi.org/10.1029/2001JD000465>, 2001.
- Bernath, P. F. et al.: Atmospheric Chemistry Experiment (ACE): Mission overview, *Geophysical Research Letters*, 32, L15S01, <https://doi.org/10.1029/2005GL022386>, 2005.
- Birner, T. and Bönisch, H.: Residual circulation trajectories and transit times into the extratropical lowermost stratosphere, *Atmospheric Chemistry and Physics*, 11, 817–827, <https://doi.org/10.5194/acp-11-817-2011>, 2011.
- Boering, K. A., Wofsy, S. C., Daube, B. C., Schneider, H. R., Loewenstein, M., Podolske, J. R., and Conway, T. J.: Stratospheric Mean Ages and Transport Rates from Observations of Carbon Dioxide and Nitrous Oxide, *Science*, 274, 1340–1343, 1996.
- Bönisch, H., Engel, A., Birner, T., Hoor, P., Tarasick, D. W., and Ray, E. A.: On the structural changes in the Brewer-Dobson circulation after 2000, *Atmospheric Chemistry and Physics*, 11, 3937–3948, <https://doi.org/10.5194/acp-11-3937-2011>, 2011.
- Boone, C., Bernath, P., Cok, D., Jones, S., and Steffen, J.: Version 4 retrievals for the Atmospheric Chemistry Experiment Fourier transform spectrometer (ACE-FTS) and imagers, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 247, 106 939, <https://doi.org/10.1016/j.jqsrt.2020.106939>, 2020.
- Boone, C. D., Walker, K. A., and Bernath, P. F.: Version 3 retrievals for the Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS), *ACE at 10: A Solar Occultation Anthology*, Deepak Publishing, 2013.
- Brown, A. T., Chipperfield, M. P., Boone, C., Wilson, C., Walker, K. A., and Bernath, P. F.: Trends in atmospheric halogen containing gases since 2004, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 112, 2552–2566, 2011.
- Butchart, N.: The Brewer-Dobson circulation, *Reviews of Geophysics*, 52, 157–184, <https://doi.org/10.1002/2013RG000448>, 2014.
- Chrysanthou, A., Maycock, A. C., Chipperfield, M. P., Dhomse, S., Garny, H., Kinnison, D., Akiyoshi, H., Deushi, M., Garcia, R. R., Jöckel, P., Kirner, O., Pitari, G., Plummer, D. A., Revell, L., Rozanov, E., Stenke, A., Tanaka, T. Y., Visioni, D., and Yamashita, Y.: The effect of atmospheric nudging on the stratospheric residual circulation in chemistry–climate models, *Atmospheric Chemistry and Physics*, 19, 11 559–11 586, <https://doi.org/10.5194/acp-19-11559-2019>, 2019.
- de Grandpré, J., Beagley, S. R., Fomichev, V. I., Griffioen, E., McConnell, J. C., Medvedev, A. S., and Shepherd, T. G.: Ozone climatology using interactive chemistry: Results from the Canadian Middle Atmosphere Model, *Journal of Geophysical Research: Atmospheres*, 105, 26 475–26 491, <https://doi.org/10.1029/2000JD900427>, 2000.
- Diallo, M., Legras, B., and Chédin, A.: Age of stratospheric air in the ERA-Interim, *Atmospheric Chemistry and Physics*, 12, 12 133–12 154, <https://doi.org/10.5194/acp-12-12133-2012>, 2012.
- Dunkerton, T. J. and Delisi, D. P.: Evolution of potential vorticity in the winter stratosphere of January–February 1979, *Journal of Geophysical Research: Atmospheres*, 91, 1199–1208, <https://doi.org/10.1029/JD091iD01p01199>, 1986.

- Elkins, J. W., Fahey, D. W., Gilligan, J. M., Dutton, G. S., Baring, T. J., Volk, C. M., Dunn, R. E., Myers, R. C., Montzka, S. A., Wamsley, P. R., Hayden, A. H., Butler, J. H., Thompson, T. M., Swanson, T. H., Dlugokencky, E. J., Novelli, P. C., Hurst, D. F., Lobert, J. M., Ciciora, S. J., McLaughlin, R. J., Thompson, T. L., Winkler, R. H., Fraser, P. J., Steele, L. P., and Lucarelli, M. P.: Airborne gas chromatograph for in situ measurements of long-lived species in the upper troposphere and lower stratosphere, *Geophysical Research Letters*, 23, 347–350, <https://doi.org/10.1029/96GL00244>, 1996.
- Engel, A., Strunk, M., Müller, M., Haase, H.-P., Poss, C., Levin, I., and Schmidt, U.: The temporal development of total chlorine in the high latitude stratosphere based on reference distributions of mean age derived from CO₂ and SF₆, *J. Geophys. Res.*, 107, 4136, <https://doi.org/10.1029/2001JD000584>, 2002.
- Engel, A., Möbius, T., Haase, H.-P., Bönisch, H., Wetter, T., Schmidt, U., Levin, I., Reddmann, T., Oelhaf, H., Wetzel, G., Grunow, K., Huret, N., and Pirre, M.: Observation of mesospheric air inside the Arctic stratospheric polar vortex in early 2003, *Atmospheric Chemistry and Physics*, 6(1), 267–282, <https://doi.org/10.5194/acp-6-267-2006>, 2006.
- Engel, A., Moore, F., Hurst, D., Möbius, T., Bönisch, H., Schmidt, U., Heinz, R., Levin, I., Atlas, E., Aoki, S., Nakazawa, T., Sugawara, S., Elkins, J., Schauffler, S., Andrews, A., and Boering, K.: Age of stratospheric air unchanged within uncertainties over the past 30 years, *Nature Geoscience*, 2, 28–31, <https://doi.org/10.1038/ngeo388>, 2009.
- Engel, A., Bönisch, H., Ullrich, M., Sitals, R., Membrive, O., Danis, F., and Crevoisier, C.: Mean age of stratospheric air derived from AirCore observations, *Atmospheric Chemistry and Physics*, 17, 6825–6838, <https://doi.org/10.5194/acp-17-6825-2017>, 2017.
- Fischer, H. and Oelhaf, H.: Remote sensing of vertical profiles of atmospheric trace constituents with MIPAS limb-emission spectrometers, *Applied Optics*, 35, 2787–2796, <https://doi.org/10.1364/AO.35.002787>, 1996.
- Fischer, H. et al.: MIPAS: an instrument for atmospheric and climate research, *Atmospheric Chemistry and Physics*, 8, 2151–2188, <https://doi.org/10.5194/acp-8-2151-2008>, 2008.
- Freie Universität Berlin: The Quasi-Biennial-Oscillation (QBO) Data Serie, <https://www.geo.fu-berlin.de/en/met/ag/strat/produkte/qbo/index.html>, accessed on 28 Nov 2022, 2022.
- Garny, H., Eichinger, R., Laube, J. C., Ray, E. A., Stiller, G. P., Bönisch, H., Saunders, L., and Linz, M.: Correction of stratospheric age of air (AoA) derived from sulfur hexafluoride (SF₆) for the effect of chemical sinks, *Atmospheric Chemistry and Physics*, 24, 4193–4215, <https://doi.org/10.5194/acp-24-4193-2024>, 2024a.
- Garny, H., Ploeger, F., Abalos, M., Bönisch, H., Castillo, A., von Clarmann, T., Diallo, M., Engel, A., Laube, J., Linz, M., Neu, J., Podglajen, A., Ray, E., Rivoire, L., Saunders, L., Stiller, G., Voet, F., Wagenhäuser, T., and Walker, K.: Age of stratospheric air: Progress on processes, observations and long-term trends, *Reviews of Geophysics*, 62, e2023RG000 832, <https://doi.org/10.1029/2023RG000832>, 2024b.
- Gelaro, R., McCarty, W., Suárez, M. J., Todling, R., Molod, A., Takacs, L., Randles, C. A., Darmenov, A., Bosilovich, M. G., Reichle, R., Wargan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A. M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J. E., Partyka, G., Pawson, S., Putman, W., Rienecker, M., Schubert, S. D., Sienkiewicz, M., and Zhao, B.: The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2), *Journal of Climate*, 30, 5419 – 5454, <https://doi.org/10.1175/JCLI-D-16-0758.1>, 2017.
- Global Monitoring Laboratory: NOAA Greenhouse Gas Marine Boundary Layer Reference, <https://gml.noaa.gov/ccgg/mbll/>, accessed on 10 July 2023, 2023.
- Haenel, F. J., Stiller, G. P., von Clarmann, T., Funke, B., Eckert, E., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., and Reddmann, T.: Reassessment of MIPAS age of air trends and variability, *Atmospheric Chemistry and Physics*, 15, 13 161–13 176, <https://doi.org/10.5194/acp-15-13161-2015>, 2015.

- Hall, T. M. and Plumb, R. A.: Age as a diagnostic of stratospheric transport, *Journal of Geophysical Research*, 99, 1059–1070, <https://doi.org/10.1029/93JD03192>, 1994.
- Hardiman, S. C., Lin, P., Scaife, A. A., Dunstone, N. J., and Ren, H.-L.: The influence of dynamical variability on the observed Brewer-
725 Dobson circulation trend, *Geophysical Research Letters*, 44, 2885–2892, <https://doi.org/10.1002/2017GL072706>, 2017.
- Harnisch, J., Borchers, R., Fabian, P., and Maiss, M.: Tropospheric trends for CF₄ and C₂F₆ since 1982 derived from SF₆ dated stratospheric air, *Geophysical Research Letters*, 23, 1099–1102, <https://doi.org/https://doi.org/10.1029/96GL01198>, 1996.
- Jonsson, A. I., de Grandpré, J., Fomichev, V. I., McConnell, J. C., and Beagley, S. R.: Doubled CO₂-induced cooling in the middle atmosphere: Photochemical analysis of the ozone radiative feedback, *Journal of Geophysical Research: Atmospheres*, 109, D24 103,
730 <https://doi.org/10.1029/2004JD005093>, 2004.
- Kodera, K. and Shibata, K.: Solar influence on the tropical stratosphere and troposphere in the northern summer, *Geophysical Research Letters*, 33, L19 704, <https://doi.org/10.1029/2006GL026659>, 2006.
- Kolonjari, F., Plummer, D. A., Walker, K. A., Boone, C. D., Elkins, J. W., Hegglin, M. I., Manney, G. L., Moore, F. L., Pendlebury, D., Ray, E. A., Rosenlof, K. H., and Stiller, G. P.: Assessing stratospheric transport in the CMAM30 simulations using ACE-FTS measurements,
735 *Atmospheric Chemistry and Physics*, 18, 6801–6828, <https://doi.org/10.5194/acp-18-6801-2018>, 2018.
- Kolonjari, F., Sheese, P. E., Walker, K. A., Boone, C. D., Plummer, D. A., Engel, A., Montzka, S. A., Oram, D. E., Schuck, T., Stiller, G. P., and Toon, G. C.: Validation of Atmospheric Chemistry Experiment Fourier Transform Spectrometer (ACE-FTS) chlorodi-
fluoromethane (HCFC-22) in the upper troposphere and lower stratosphere, *Atmospheric Measurement Techniques*, 17, 2429–2449,
<https://doi.org/10.5194/amt-17-2429-2024>, 2024.
- 740 Koo, J.-H., Walker, K. A., Jones, A., Sheese, P. E., Boone, C. D., Bernath, P. F., and Manney, G. L.: Global climatology based on the ACE-FTS version 3.5 dataset: Addition of mesospheric levels and carbon-containing species in the UTLS, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 186, 52–62, 2017.
- Laube, J. C., Engel, A., Bönisch, H., Möbius, T., Sturges, W. T., Braß, M., and Röckmann, T.: Fractional release factors of long-lived halo-
generated organic compounds in the tropical stratosphere, *Atmospheric Chemistry and Physics*, 10, 1093–1103, <https://doi.org/10.5194/acp-10-1093-2010>, 2010.
745
- Leedham Elvidge, E., Bönisch, H., Brenninkmeijer, C. A. M., Engel, A., Fraser, P. J., Gallacher, E., Langenfelds, R., Mühle, J., Oram, D. E., Ray, E. A., Ridley, A. R., Röckmann, T., Sturges, W. T., Weiss, R. F., and Laube, J. C.: Evaluation of stratospheric age of air from CF₄, C₂F₆, C₃F₈, CHF₃, HFC-125, HFC-227ea and SF₆; implications for the calculations of halocarbon lifetimes, fractional release factors and ozone depletion potentials, *Atmospheric Chemistry and Physics*, 18, 3369–3385, <https://doi.org/10.5194/acp-18-3369-2018>, 2018.
- 750 Levin, I., Naegler, T., Heinz, R., Osusko, D., Cuevas, E., Engel, A., Ilmberger, J., Langenfelds, R. L., Neininger, B., Rohden, C. v., Steele, L. P., Weller, R., Worthy, D. E., and Zimov, S. A.: The global SF₆ source inferred from long-term high precision atmospheric measurements and its comparison with emission inventories, *Atmospheric Chemistry and Physics*, 10, 2655–2662, <https://doi.org/10.5194/acp-10-2655-2010>, 2010.
- Linz, M., Plumb, R. A., Gerber, E. P., and Sheshadri, A.: The Relationship between Age of Air and the Diabatic Circulation of the Strato-
755 sphere, *Journal of the Atmospheric Sciences*, 73, 4507–4518, <https://doi.org/10.1175/JAS-D-16-0125.1>, 2016.
- Linz, M., Plumb, R. A., Gerber, E. P., Haenel, F. J., Stiller, G., Kinnison, D. E., Ming, A., and Neu, J. L.: The strength of the meridional overturning circulation of the stratosphere, *Nature Geoscience*, 10, 663–667, 2017.

- Loeffel, S., Eichinger, R., Garny, H., Reddmann, T., Fritsch, F., Versick, S., Stiller, G., and Haenel, F.: The impact of sulfur hexafluoride (SF_6) sinks on age of air climatologies and trends, *Atmospheric Chemistry and Physics*, 22, 1175–1193, <https://doi.org/10.5194/acp-22-1175-2022>, 2022.
- Mahieu, E., Duchatelet, P., Demoulin, P., Walker, K. A., Dupuy, E., Froidevaux, L., Randall, C., Catoire, V., Strong, K., Boone, C. D., Bernath, P. F., Blavier, J.-F., Blumenstock, T., Coffey, M., De Mazière, M., Griffith, D., Hannigan, J., Hase, F., Jones, N., Jucks, K. W., Kagawa, A., Kasai, Y., Mebarki, Y., Mikuteit, S., Nassar, R., Notholt, J., Rinsland, C. P., Robert, C., Schrems, O., Senten, C., Smale, D., Taylor, J., Tétard, C., Toon, G. C., Warneke, T., Wood, S. W., Zander, R., and Servais, C.: Validation of ACE-FTS v2.2 measurements of HCl , HF , CCl_3F and CCl_2F_2 using space-, balloon- and ground-based instrument observations, *Atmospheric Chemistry and Physics*, 8, 6199–6221, <https://doi.org/10.5194/acp-8-6199-2008>, 2008.
- Manney, G. L. and Zurek, R. W.: Interhemispheric comparison of the development of the stratospheric polar vortex during fall: A 3-dimensional perspective for 1991–1992, *Geophysical Research Letters*, 20, 1275–1278, <https://doi.org/10.1029/93GL01111>, 1993.
- Manney, G. L., Zurek, R. W., O’Neill, A., and Swinbank, R.: On the Motion of Air through the Stratospheric Polar Vortex, *Journal of Atmospheric Sciences*, 51, 2973–2994, [https://doi.org/10.1175/1520-0469\(1994\)051<2973:OTMOAT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<2973:OTMOAT>2.0.CO;2), 1994.
- Manney, G. L., Daffer, W. H., Zawodny, J. M., Bernath, P. F., Hoppel, K. W., Walker, K. A., Knosp, B. W., Boone, C., Remsberg, E. E., Santee, M. L., Harvey, V. L., Pawson, S., Jackson, D. R., Deaver, L., McElroy, C. T., McLinden, C. A., Drummond, J. R., Pumphrey, H. C., Lambert, A., Schwartz, M. J., Froidevaux, L., McLeod, S., Takacs, L. L., Suarez, M. J., Trepte, C. R., Cuddy, D. C., Livesey, N. J., Harwood, R. S., and Waters, J. W.: Solar occultation satellite data and derived meteorological products: Sampling issues and comparisons with Aura Microwave Limb Sounder, *Journal of Geophysical Research: Atmospheres*, 112, D24S50, <https://doi.org/10.1029/2007JD008709>, 2007.
- Morris, R. A., Miller, T. M., Viggiano, A. A., Paulson, J. F., Solomon, S., and Reid, G.: Effects of electron and ion reactions on atmospheric lifetimes of fully fluorinated compounds, *Journal of Geophysical Research*, 100, 1287–1294, <https://doi.org/10.1029/94JD02399>, 1995.
- Neu, J. L. and Plumb, R. A.: Age of air in a “leaky pipe” model of stratospheric transport, *Journal of Geophysical Research: Atmospheres*, 104, 19 243–19 255, <https://doi.org/10.1029/1999JD900251>, 1999.
- NOAA: Monthly Adjusted Solar Flux, https://www.ngdc.noaa.gov/stp/space-weather/solar-data/solar-features/solar-radio/noontime-flux/penticton/penticton_adjusted/listings/listing_drao_noontime-flux-adjusted_monthly.txt, accessed on 21 Dec 2022, 2018.
- NOAA: Multivariate ENSO Index Version 2, <https://psl.noaa.gov/enso/mei/>, accessed on 29 Nov 2022, 2022.
- Onogi, K., Tsutsui, J., Koide, H., Sakamoto, M., Kobayashi, S., Hatsushika, H., Matsumoto, T., Yamakazi, N., Kamahori, H., Takahashi, K., Kadokura, S., Wada, K., Kato, K., Oyama, R., Ose, T., Mannoji, N., and Taira, R.: The JRA-25 Reanalysis, *Journal of the Meteorological Society of Japan. Ser. II*, 85, 369–432, <https://doi.org/10.2151/jmsj.85.369>, 2007.
- Ploeger, F., Diallo, M., Charlesworth, E., Konopka, P., Legras, B., Laube, J. C., Grooß, J.-U., Günther, G., Engel, A., and Riese, M.: The stratospheric Brewer–Dobson circulation inferred from age of air in the ERA5 reanalysis, *Atmospheric Chemistry and Physics*, 21, 8393–8412, <https://doi.org/10.5194/acp-21-8393-2021>, 2021.
- Plumb, R.: Stratospheric Transport, *Journal of the Meteorological Society of Japan*, 80, 793–809, <https://doi.org/10.2151/jmsj.80.793>, 2002.
- Ray, E. A., Moore, F. L., Elkins, J. W., Dutton, G. S., Fahey, D. W., Vömel, H., Oltmans, S. J., and Rosenlof, K. H.: Transport into the Northern Hemisphere lowermost stratosphere revealed by in situ tracer measurements, *Journal of Geophysical Research*, 104, 26 565–26 580, <https://doi.org/10.1029/1999JD900323>, 1999.

- Ray, E. A., Moore, F. L., Rosenlof, K. H., Davis, S. M., Sweeney, C., Tans, P., Wang, T., Elkins, J. W., Bönnisch, H., Engel, A., Sugawara, S., Nakazawa, T., and Aoki, S.: Improving stratospheric transport trend analysis based on SF₆ and CO₂ measurements, *Journal of Geophysical Research: Atmospheres*, 119, 14 110–14 128, <https://doi.org/10.1002/2014JD021802>, 2014.
- Ray, E. A., Moore, F. L., Elkins, J. W., Rosenlof, K. H., Laube, J. C., Röckmann, T., Marsh, D. R., and Andrews, A. E.: Quantification of the SF₆ lifetime based on mesospheric loss measured in the stratospheric polar vortex, *Journal of Geophysical Research: Atmospheres*, <https://doi.org/10.1002/2016JD026198>, 2016JD026198, 2017.
- Reddmann, T., Ruhnke, R., and Kouker, W.: Three-dimensional model simulations of SF₆ with mesospheric chemistry, *Journal of Geophysical Research*, 106, 14 525–14 537, <https://doi.org/10.1029/2000JD900700>, 2001.
- Reichler, T., Dameris, M., and Sausen, R.: Determining the tropopause height from gridded data, *Geophysical Research Letters*, 30, 2042, <https://doi.org/10.1029/2003GL018240>, 2003.
- Rigby, M., Mühle, J., Miller, B. R., Prinn, R. G., Krummel, P. B., Steele, L. P., Fraser, P. J., Salameh, P. K., Harth, C. M., Weiss, R. F., Grealley, B. R., O'Doherty, S., Simmonds, P. G., Vollmer, M. K., Reimann, S., Kim, J., Kim, K.-R., Wang, H. J., Olivier, J. G. J., Dlugokencky, E. J., Dutton, G. S., Hall, B. D., and Elkins, J. W.: History of atmospheric SF₆ from 1973 to 2008, *Atmospheric Chemistry and Physics*, 10, 10 305–10 320, <https://doi.org/10.5194/acp-10-10305-2010>, 2010.
- Scinocca, J. F., McFarlane, N. A., Lazare, M., Li, J., and Plummer, D.: Technical Note: The CCCma third generation AGCM and its extension into the middle atmosphere, *Atmospheric Chemistry and Physics*, 8, 7055–7074, <https://doi.org/10.5194/acp-8-7055-2008>, 2008.
- Sheese, P. and Walker, K.: Data Quality Flags for ACE-FTS Level 2 Version 4.1/4.2 Data Set, V32, <https://doi.org/10.5683/SP2/BC4ATC>, accessed 17 April 2024, 2020.
- Sheese, P. E., Boone, C. D., and Walker, K. A.: Detecting physically unrealistic outliers in ACE-FTS atmospheric measurements, *Atmospheric Measurement Techniques*, 8, 741–750, <https://doi.org/10.5194/amt-8-741-2015>, 2015.
- Simmonds, P. G., Rigby, M., Manning, A. J., Park, S., Stanley, K. M., McCulloch, A., Henne, S., Graziosi, F., Maione, M., Arduini, J., Reimann, S., Vollmer, M. K., Mühle, J., O'Doherty, S., Young, D., Krummel, P. B., Fraser, P. J., Weiss, R. F., Salameh, P. K., Harth, C. M., Park, M.-K., Park, H., Arnold, T., Rennick, C., Steele, L. P., Mitrevski, B., Wang, R. H. J., and Prinn, R. G.: The increasing atmospheric burden of the greenhouse gas sulfur hexafluoride (SF₆), *Atmospheric Chemistry and Physics*, 20, 7271–7290, <https://doi.org/10.5194/acp-20-7271-2020>, 2020.
- Space Weather Canada: Monthly averages of solar 10.7 cm flux, <https://spaceweather.gc.ca/forecast-prevision/solar-solaire/solarflux/sx-5-mavg-en.php>, accessed on 21 Dec 2022, 2022.
- Stiller, G. P., von Clarmann, T., Höpfner, M., Glatthor, N., Grabowski, U., Kellmann, S., Kleinert, A., Linden, A., Milz, M., Reddmann, T., Steck, T., Fischer, H., Funke, B., López-Puertas, M., and Engel, A.: Global distribution of mean age of stratospheric air from MIPAS SF₆ measurements, *Atmospheric Chemistry and Physics*, 8, 677–695, <https://doi.org/10.5194/acp-8-677-2008>, 2008.
- Stiller, G. P., von Clarmann, T., Haenel, F., Funke, B., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., Lossow, S., and López-Puertas, M.: Observed temporal evolution of global mean age of stratospheric air for the 2002 to 2010 period, *Atmospheric Chemistry and Physics*, 12, 3311–3331, <https://doi.org/10.5194/acp-12-3311-2012>, 2012.
- Strahan, S. E. et al.: Observed Hemispheric Asymmetry in Stratospheric Transport Trends From 1994 to 2018, *Geophysical Research Letters*, 47, e2020GL088 567, <https://doi.org/10.1029/2020GL088567>, 2020.
- Tao, M., Konopka, P., Wright, J., Liu, Y., Bian, J., Davis, S., Jia, Y., and Ploeger, F.: Multi-decadal variability controls short-term stratospheric water vapor trends, *Communications Earth Environment*, 4, 441, <https://doi.org/10.1038/s43247-023-01094-9>, 2023.

- Totterdill, A., Kovács, T., Gómez Martín, J. C., Feng, W., and Plane, J. M. C.: Mesospheric Removal of Very Long-Lived Greenhouse Gases SF₆ and CFC-115 by Metal Reactions, Lyman- α Photolysis, and Electron Attachment, *The Journal of Physical Chemistry A*, 119, 2016–2025, <https://doi.org/10.1021/jp5123344>, 2015.
- Varanasi, P., Li, Z., Nemtchinov, V., and Cherukuri, A.: Spectral absorption-coefficient data on HCFC-22 and SF₆ for remote-sensing applications, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 52, 323–332, [https://doi.org/10.1016/0022-4073\(94\)90162-7](https://doi.org/10.1016/0022-4073(94)90162-7), 1994.
- Volk, C. M., Elkins, J. W., Fahey, D. W., Dutton, G. S., Gilligan, J. M., Loewenstein, M., Podolske, J. R., Chan, K. R., and Gunson, M. R.: Evaluation of source gas lifetimes from stratospheric observations, *Journal of Geophysical Research: Atmospheres*, 102, 25 543–25 564, <https://doi.org/10.1029/97JD02215>, 1997.
- 835 von Clarmann, T., Glatthor, N., Grabowski, U., Höpfner, M., Kellmann, S., Kiefer, M., Linden, A., Tsidu, G. M., Milz, M., Steck, T., Stiller, G. P., Wang, D. Y., Fischer, H., Funke, B., Gil-López, S., and López-Puertas, M.: Retrieval of temperature and tangent altitude pointing from limb emission spectra recorded from space by the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS), *Journal of Geophysical Research: Atmospheres*, 108, 4746, <https://doi.org/10.1029/2003JD003602>, 2003.
- 845 von Clarmann, T., Höpfner, M., Kellmann, S., Linden, A., Chauhan, S., Funke, B., Grabowski, U., Glatthor, N., Kiefer, M., Schieferdecker, T., Stiller, G. P., and Versick, S.: Retrieval of temperature, H₂O, O₃, HNO₃, CH₄, N₂O, ClONO₂ and ClO from MIPAS reduced resolution nominal mode limb emission measurements, *Atmospheric Measurement Techniques*, 2, 159–175, <https://doi.org/10.5194/amt-2-159-2009>, 2009.
- von Clarmann, T., Stiller, G., Grabowski, U., Eckert, E., and Orphal, J.: Technical Note: Trend estimation from irregularly sampled, correlated data, *Atmospheric Chemistry and Physics*, 10, 6737–6747, <https://doi.org/10.5194/acp-10-6737-2010>, 2010.
- 850 Waugh, D. and Hall, T.: Age of stratospheric air: theory, observations, and models, *Reviews of Geophysics*, 40, 1010, <https://doi.org/10.1029/2000RG000101>, 2002.