

# Source apportionment and ecotoxicity of PM<sub>2.5</sub> pollution events in a Major Southern Hemisphere Megacity: influence of a biofuel impacted fleet and biomass burning

Guilherme Martins Pereira<sup>1,2</sup>, Leonardo Yoshiaki Kamigauti<sup>1</sup>, Rubens Fabio Pereira<sup>1</sup>, Djacinto Monteiro dos Santos<sup>3</sup>, Thayná da Silva Santos<sup>2</sup>, José Vinicius Martins<sup>4</sup>, Célia Alves<sup>5</sup>, Cátia Gonçalves<sup>5,a</sup>, Ismael Casotti Rienda<sup>5</sup>, Nora Kováts<sup>6</sup>, Thiago Nogueira<sup>7</sup>, Luciana Rizzo<sup>3</sup>, Paulo Artaxo<sup>3</sup>, Regina Maura de Miranda<sup>8</sup>, Marcia Akemi Yamasoe<sup>1</sup>, Edmilson Dias de Freitas<sup>1</sup>, Pérola de Castro Vasconcellos<sup>2</sup>, Maria de Fatima Andrade<sup>1</sup>

<sup>1</sup>Department of Atmospheric Sciences, Institute of Astronomy, Geophysics and Atmospheric Sciences, University of São Paulo, 05508-090, São Paulo, Brazil.

<sup>2</sup>Department of Chemistry, Institute of Chemistry, University of São Paulo, 05508-000, São Paulo, Brazil.

<sup>3</sup>Department of Applied Physics, Institute of Physics, University of São Paulo, 05508-090, São Paulo, Brazil.

<sup>4</sup>Department of Mineralogy and Geotectonics, Institute of Geosciences, University of São Paulo, 05508-080, São Paulo, Brazil.

<sup>5</sup>Center for Environmental and Marine Studies, Department of Environment and Planning, University of Aveiro, 3810-193 Aveiro, Portugal

<sup>6</sup>Center of Natural Environmental Sciences, University of Pannonia, Egyetem str. 10, 8200 Veszprém, Hungary

<sup>7</sup>Department of Environmental Health, School of Public Health, University of São Paulo, 01246-904, São Paulo, Brazil.

<sup>8</sup>School of Arts, Sciences and Humanities, University of São Paulo, 03828-000, São Paulo, Brazil.

<sup>a</sup>Current address: Department of Physics, University of León, Campus de Vegazana, 24071 León, Spain.

*Correspondence to:* Guilherme Martins Pereira (guilherme.martins.pereira@usp.br)

**Abstract.** The Metropolitan Area of São Paulo (MASP) in Brazil has reduced its vehicular emissions in the last decades. However, it is still affected by air pollution events, mainly in the winter, characterized as a dry season. The chemical composition of fine particulate matter (PM<sub>2.5</sub>) was studied in the MASP during a 100-day dry period in 2019. PM<sub>2.5</sub> samples underwent an extensive chemical characterization (including inorganic and organic species), and submicrometer particle number size distributions were simultaneously monitored. PM<sub>2.5</sub> concentrations exceeded the new World Health Organization's daily guidelines on 75% of sampling days, emphasizing the need for strengthening local regulations. Source apportionment (Positive Matrix Factorization, PMF5.0) was performed, and the sources related to vehicular emissions remain relevant (over 40% of PM<sub>2.5</sub>). A high contribution of biomass burning was observed, reaching 25% of PM<sub>2.5</sub> mass and correlated with sample ecotoxicity. This input was associated with north and northwest winds, suggesting other emerging sources besides sugarcane burning (forest fires and sugarcane bagasse power plants). A mixed factor of road dust and vehicular emissions increased throughout the campaign was related to stronger winds, suggesting a significant resuspension. The sulfate secondary formation was related to humid conditions. Additionally, monitoring size particle distribution allowed the observation of particle growth on days impacted by secondary formation. The results pointed out that control measures of high

34 PM<sub>2.5</sub> events should include the control of emerging biomass burning sources in addition to stricter rules concerning vehicular  
35 emissions.

## 36 1 Introduction

37 According to the World Health Organization (WHO), a large part of the world's population lives in places where the  
38 recommended air quality standards are not achieved, and this includes the Metropolitan Area of São Paulo (MASP), Brazil  
39 (WHO, 2021; CETESB 2023), where thousands of tons of pollutants are released into the atmosphere every year. Natural and  
40 anthropogenic sources produce atmospheric pollutants. The MASP has more than 21 million inhabitants and around 7 million  
41 vehicles. In 2022, in the MASP, vehicles were responsible for 96% of CO emissions, 70% of HC, 60% of NO<sub>x</sub>, 8% of SO<sub>2</sub>,  
42 and 37% of fine mode PM (CETESB, 2023). Among the pollutants, particulate matter is extensively studied, as it harms human  
43 health and is associated with cardiovascular diseases and cancer (Cohen et al., 2017). Furthermore, particulate matter has  
44 climatic effects, through the absorption and scattering of solar radiation and indirectly by affecting cloud microphysics (Li et  
45 al., 2022; Pöschl, 2005).

46 Particles are emitted by different natural sources, such as volcanic eruptions, resuspension, and erosion of soils and  
47 vegetation, while the dominant anthropogenic sources are vehicles, industrial activities, and biomass burning. The formation  
48 of secondary particles is quite significant in some cities, such as São Paulo, where sulfate and nitrate concentrations are high  
49 and secondary particles comprise half of the fine mode PM mass (Andrade et al., 2012; CETESB et al., 2023). Particulate  
50 matter is a complex mixture, including inorganic species (water-soluble ions and element oxides) and carbonaceous species,  
51 such as elemental carbon (EC, associated with soot), and organic carbon (OC) associated with organic compounds, some of  
52 them with toxic properties, such as polycyclic aromatic hydrocarbons (PAHs) (Ravindra et al., 2008; Pöschl, 2005). The fine  
53 fraction of particulate matter (PM<sub>2.5</sub>) includes particles with aerodynamic diameters equal or lower than 2.5 µm that can  
54 penetrate the respiratory system, reaching the alveoli. Smaller particles than 100 nm are called ultrafine particles and can reach  
55 other organs through the lung vasculature (Schraufnagel, 2020; Kumar et al., 2014).

56 The vehicle fuel profile in the MASP is unique compared to other metropolises worldwide, with a significant proportion  
57 of biofuels, especially in light duty vehicles (Andrade et al., 2017). The fleet has been running in gasoline and bioethanol  
58 blends (gasohol, 73% of gasoline, 27%, ethanol), hydrated ethanol (5% of water), diesel and biodiesel blends (9% of biodiesel  
59 produced from soybean) (Pereira et al., 2023a,b). The peak levels of ethanol use were reached at the end of 2018 and were  
60 comparable to those of gasohol (Pereira et al., 2023a). Despite the exponential increase in the vehicle fleet since the 1980's,  
61 particulate pollutants have been reduced in the last decades (Andrade et al., 2017). Sugarcane biomass burning in the  
62 countryside also has been reduced due to control measures (Valente and Laurini, 2021). In addition to the large quantity of  
63 pollutants generated locally, increasing events of transport of biomass burning from the Amazon and Central parts of Brazil  
64 have been observed in the MASP at the end of the decade (Pereira et al., 2021; Miranda et al., 2017), occurring mainly in the  
65 dry period (July-October) (Vieira et al., 2023). In 2014 and 2015, biomass-burning sources were apportioned and found to  
66 explain nearly one-fourth of particulate matter in the dry season (Pereira et al., 2017b; Emygdio et al., 2018). Local biomass  
67 burning was also relevant, including the burning of charcoal and wood in barbecue and pizzerias and waste burning (Kumar  
68 et al., 2016). In the eastern region of the city, an area with high population density, roads with heavy vehicle traffic, highly  
69 sealed, and many industries, the contribution in 2019 was 9.9% light vehicles, 42% heavy vehicles, and 47.3% soil dust and  
70 local sources such as waste burning and industries (Vieira et al., 2023). A rise in the contribution of non-exhaust sources is  
71 expected in the next few years due to control measures focused on vehicle exhaust.


72 The São Paulo State Environmental Company (CETESB) monitors legislated pollutants at 26 air quality stations  
73 throughout the MASP. Despite the efforts of local governments to improve air quality in the region, in 2019, they all recorded  
74 concentrations of PM<sub>2.5</sub> above the value recommended by the WHO, reaching critical levels in the drier months (April to  
75 September). Considering the aspects mentioned before, it is crucial to periodically update the chemical characterization of

76 particulate matter since Brazilian environmental agencies do not carry out such extensive monitoring. It can be said that particle  
77 sources in São Paulo always need to be studied and identified, but this depends on good quality chemical composition data.  
78 The composition of aerosols varies with the changing fleet and the policies concerning the burning of biomass, furthermore,  
79 they are also influenced by the changing climate and meteorological conditions. In recent years, dryer weather conditions in  
80 the winter have favored forest and crop fires in central Brazil and the accumulation of pollutants (Souto-Oliveira et al., 2023;  
81 Pereira et al., 2021). This study aims to thoroughly characterize PM<sub>2.5</sub> chemical composition and toxicity in the MASP,  
82 identifying the relative contribution of emission sources. Associations with weather conditions and case studies of air pollution  
83 events were also investigated. The adoption of receptor models was performed to study the emission sources of PM<sub>2.5</sub>, and  
84 these results were associated with particle size distributions.

## 85 2 Materials and methods

### 86 2.1 Sampling of PM<sub>2.5</sub> and particle number size distributions monitoring

87 High-volume samplers (Energética, Brazil) (1.13 m<sup>3</sup> min<sup>-1</sup> flow) were employed to collect PM<sub>2.5</sub> on the rooftop of the Institute  
88 of Chemistry building at the University of São Paulo, in the city of São Paulo, Brazil (23°33'53"S, 46°43'32"W), located in a  
89 green area, and more than 2 km away from a busy expressway (Marginal Pinheiros) (Figure 1). The sampling took place  
90 between June 04 and September 12, 2019, in a pre-lockdown polluted period, starting six months before the pandemic. During  
91 this campaign, a dark precipitation event (August 19) was observed, an unprecedented event that was termed “black rain”  
92 following biomass burning episodes in central South America (Pereira et al., 2021). Ninety-nine samples were collected for  
93 24 hours (starting at 9 AM, local time). Quartz fiber filters (20 cm × 25 cm, Whatman, UK) were employed to collect PM<sub>2.5</sub>.  
94 These filters were decontaminated before sampling by heating at 600 °C for 6 h. Before and after sampling, the filters were  
95 weighed in a microbalance (controlled temperature and humidity for equilibrium: T = 25 °C and RH = 50%). Then, the filters  
96 were wrapped into a laminated sheet and stored at 5 °C to avoid volatilization and reactions of the analytes before the analysis  
97 (de Oliveira Alves et al., 2015; Souza et al., 2014).



99  
100 **Figure 1:** Sampling site location (marked with red cross) (© Google Maps). Heavily trafficked expressways are  
101 indicated with the red dashed line.

102 Particle number size distributions (PNSD) were monitored using a Scanning Mobility Particle Sizer (SMPS 3081, TSI  
103 Inc) in association with a Condensation Particle Counter (CPC 3010, TSI Inc). The system provided particle number size  
104 distributions in the size range from 10 to 450 nm every 2 min. PNSD measurements were averaged to match the filter sampling

periods. Average distributions were also calculated for periods of interest in the dataset. The SMPS data were collected on the rooftop of the Institute of Astronomy, Geophysics, and Atmospheric Sciences of the University of São Paulo, 500 meters from the Institute of Chemistry. PNSD measurements were taken between June and September 2019, totalizing 48 non-consecutive sampling days during the austral winter.

## 2.2 Analytical procedures

Distinct chemical composition analyses were performed, thus, pieces of the filters were sent to each laboratory. At the Institute of Chemistry (University of São Paulo), water-soluble ions (WSI) and PAHs were determined. The extraction of WSI from filters (4 cm<sup>2</sup>) was performed with 10 mL of deionized water (Milli-Q, Merck Millipore, USA) under sonication for 15 min. Then, extracts were filtered with syringe filters (Millex-GV, 0.22 µm, PVDF), and cations and anions were quantified in an ion chromatograph (Modules 819, 830, 833, 818, and 820, Metrohm, Switzerland). Anions F<sup>-</sup>, Cl<sup>-</sup>, NO<sub>2</sub><sup>-</sup>, Br<sup>-</sup>, NO<sub>3</sub><sup>-</sup>, PO<sub>4</sub><sup>3-</sup>, SO<sub>4</sub><sup>2-</sup>, C<sub>2</sub>O<sub>4</sub><sup>2-</sup>, HCO<sub>2</sub><sup>-</sup>, and C<sub>4</sub>H<sub>2</sub>O<sub>4</sub><sup>2-</sup>, and cations Ca<sup>2+</sup>, Mg<sup>2+</sup>, K<sup>+</sup>, NH<sub>4</sub><sup>+</sup>, and Na<sup>+</sup> were quantified. Pereira et al. (2023a) describe the adopted columns and eluents. Fluka (Switzerland) analyte standards were used. Recoveries fell between 80-120% and were obtained by adding known concentrations of standards to blank filters and are reported in Table S1.

Punches of approximately 20 cm<sup>2</sup> from the filters underwent extraction using ultrasonic baths with 80 mL of dichloromethane for three cycles of twenty minutes, as described by Pereira et al. (2017a). The extracts were concentrated via rotary evaporation under low pressure (at 35°C for approximately 30 minutes) and fractionated on a chromatographic column packed with 1.5 g silica gel. The methodology described in Vasconcellos et al. (2010) was modified and adopted in this study, involving three elutions: the first fraction carried alkanes (10 mL of hexane), while the second (9.6 mL of hexane + 5.4 mL of toluene), and the third (7.5 mL of hexane and 7.5 mL of dichloromethane) carried PAHs and their derivatives. The second and third fractions were combined to enhance recovery. The fractionated extracts were concentrated by rotary evaporation under reduced pressure (at 35°C for approximately 15 minutes), filtered, stored in 2 mL vials, dried, and reconstituted with 0.5 µL of hexane. Samples were diluted prior to analysis by gas chromatography coupled with mass spectrometry (GC/MS, Agilent, GC 7820A, and MS 5975), using an Agilent VF-5ms column (stationary phase, 30 m × 0.250 mm × 0.25 µm), with helium as the carrier gas (99.97% purity and flow rate of 1.0 mL min<sup>-1</sup>). The following species were determined: phenanthrene (Phe), anthracene (Ant), fluoranthene (Flt), pyrene (Pyr), retene (Ret), benzo(a)anthracene (BaA), chrysene (Chr), benzo(b)fluoranthene (BbF), benzo(k)fluoranthene (BkF), benzo(e)pyrene (BeP), benzo(a)pyrene (BaP), indene(1,2,3-c,d)pyrene (InP), dibenzo(a,h)anthracene (DBA) and benzo(g,h,i)pyrene (BPe) and coronene (Cor). Recovery was assessed by adding known quantities of PAHs mix to filters containing 15 mg of the certified material (Urban Dust SRM 1649b, NIST, USA). Most species presented values between 80 and 120%, averaging 100% (Table S1).

Carbonaceous species (OC and EC) in the quartz fiber filters were determined through thermal-optical analysis using a *Sunset Laboratory Inc.* carbon analyzer at the Institute of Physics of the University of São Paulo. The EUSAAR-2 temperature protocol (Cavalli et al., 2010) was employed, and the transmittance-based pyrolysis correction was applied, similar to previous studies conducted in the MASP (Monteiro dos Santos et al., 2016). Temperature-resolved carbon fractions are obtained, with increasing temperatures: four for organic fractions (OC1 to OC4, from higher to lower volatility temperatures), four for elemental carbon (EC1 to EC4), and the organic pyrolyzed carbon (PC) is monitored. Total OC is the sum of OC1, OC2, OC3, OC4, and PC, while total EC is the sum of EC1, EC2, EC3, and EC4, subtracted by PC. The first stage in helium medium is carried out under the following conditions of temperature and time steps: 200 °C for 120s (OC1), 300 °C for 150s (OC2), 450 °C for 180s (OC3) and 650 °C for 180s (OC4). The second stage, under He-O<sub>2</sub> medium, the four steps are: 500 °C for 120s (EC1), 550 °C for 120s (EC2), 700 °C for 70s (EC3) and 850 °C for 80s (EC4). Secondary organic carbon (SOC) was estimated considering the 5<sup>th</sup> percentile of the OC/EC ratios, similar as in Monteiro dos Santos et al. (2016). Organic matter (OM) was calculated as 1.6 × OC, as adopted for urban aerosols (Turpin and Lim, 2001).

Quartz fiber filters were subjected to acid digestion using a microwave digester oven (CEM MDS-2000, USA) at the Institute of Geosciences of the University of São Paulo. A strip of 20 cm × 2.5 cm of the filter was submitted to digestion in a closed vessel (PFA) with a mixture of HNO<sub>3</sub> and H<sub>2</sub>O (10 and 15 mL, respectively), then the volume was adjusted to 50 mL. Major and trace elements were quantified by inductively coupled plasma mass spectroscopy (ICP-MS, model iCAP Q, Thermo Fisher Scientific, USA). The percent recoveries and detection limits were previously reported for all the determined analytes and reported in Pereira et al. (2023a).

At the Center for Environmental and Marine Studies (CESAM) of the University of Aveiro (Portugal), sugar species were extracted from the filters with ultrapure Milli-Q water in ultrasonic agitation, as described by Oduber et al. (2021). After the extraction, the solutions were filtered with syringe filters (PTFE; 0.2 µm) and transferred to vials for liquid chromatography analysis with amperometric detection. Sugar compounds such as levoglucosan (Lev), mannosan (Man), galactosan (Gal), mannitol (Mnt), arabitol (Ara), and xylitol (Xyl) were determined with a Thermo Scientific Dionex™ ICS-5000 equipped with an anion-exchange analytical column (CarboPac® PA-1; 2 × 250 mm). Multi-step gradient conditions were adopted, with ultrapure Milli-Q water and two solutions of NaOH (200 mM and 5 mM). Recoveries were reported by Caseiro et al (2007).

### 2.3 Benzo(a)pyrene equivalent indexes

Benzo(a)pyrene equivalent indexes were calculated to evaluate toxicity parameters due to exposure to PAHs. Equations 1 and 2 were employed to calculate the BaP<sub>TEQ</sub> and BaP<sub>MEQ</sub> by multiplying each species' concentrations by their toxic and mutagenic equivalency factors (TEF and MEF), as reported in de Oliveira Alves et al. (2020). The PAH carcinogenicity equivalent index (BaP<sub>Eq</sub>) was also calculated by applying Equation 3, incorporating specific PAH concentrations as adopted by Yassaa et al. (2001) and Cecinato (1997).

$$\text{BaP}_{\text{TEQ}} = ([\text{BaA}] \times 0.1) + ([\text{Chr}] \times 0.01) + ([\text{BbF}] \times 0.1) + ([\text{BkF}] \times 0.1) + ([\text{BaP}] \times 1) + ([\text{InP}] \times 0.1) + ([\text{DBA}] \times 5) + ([\text{BPe}] \times 0.01) \quad (1)$$

$$\text{BaP}_{\text{MEQ}} = ([\text{BaA}] \times 0.082) + ([\text{Chr}] \times 0.017) + ([\text{BbF}] \times 0.25) + ([\text{BkF}] \times 0.11) + ([\text{BaP}] \times 1) + ([\text{InP}] \times 0.31) + ([\text{DBA}] \times 0.29) + ([\text{BPe}] \times 0.19) \quad (2)$$

$$\text{BaP}_{\text{Eq}} = ([\text{BaA}] \times 0.06) + ([\text{BbF}] \times 0.07) + ([\text{BkF}] \times 0.07) + ([\text{BaP}] \times 1) + ([\text{DBA}] \times 0.6) + ([\text{InP}] \times 0.08) \quad (3)$$

### 2.4 Data treatment and Positive Matrix Factorization

Pearson coefficients were obtained to estimate the correlation between different variables (Jamovi software), and *r* was considered significant when *p* < 0.05. Correlations between 0.0 and 0.3 were considered negligible, between 0.3 and 0.5 as weak, between 0.5 and 0.7 as moderate, between 0.7 and 0.9 as strong, and between 0.9 and 1.0, as very strong (Khan et al., 2018). To evaluate equal and unequal variances, the Mann–Whitney U test was also employed (*p* < 0.05). Polar plots were constructed with the mass concentrations as functions of wind speed and direction. They are obtained by a function in OpenAir package (RStudio). Diagnostic ratios were performed between two (or more) chemical species concentrations: OC/EC, EC/Cu, PC/CTot, Fe/Ca<sup>2+</sup>, Cu/Sb, Cu/Zn, Fe/Cu, Sn/Sb, La/Ce, V/Ni, Pb/Cu, Cu/Ca<sup>2+</sup>, BaP/(BaP+BeP), Flt/(Flt+Pir), InP/(InP+BPe), BaA/(BaA+Cri), LMW-PAHs/HMW-PAHs, Pyr/BaP, Pyr/BbF, Flt/BbF, NO<sub>3</sub><sup>-</sup>/EC, SO<sub>4</sub><sup>2-</sup>/NO<sub>3</sub><sup>-</sup>, SO<sub>4</sub><sup>2-</sup>/EC, K<sup>+</sup>/Lev, Lev/Man, SO<sub>4</sub><sup>2-</sup>/Zn, and NO<sub>3</sub><sup>-</sup>/Zn.

The ISORROPIA model calculates the composition and phase state of the water-soluble inorganic aerosol in thermodynamic equilibrium with gas phase precursors, simulating the process of dissolution of atmospheric particles and the ion formation (Bian et al., 2014; Fountoukis and Nenes, 2007; Li et al., 2014; Vieira-Filho et al., 2016a). In this study, the ISORROPIA II was applied. It can solve two types of problems: (i) forward (or "closed system") and (ii) reverse (or "open system"). The reverse problem approach was adopted in this study using aerosol phase concentrations of NH<sub>3</sub> (ammonium ion), H<sub>2</sub>SO<sub>4</sub>, Na<sup>+</sup>, Ca<sup>2+</sup>, K<sup>+</sup>, Mg<sup>2+</sup>, HCl, and HNO<sub>3</sub> to estimate water content, salt aerosol concentrations, and gaseous precursors aerosols. The aerosol-atmosphere system was considered thermodynamically stable (with precipitation of salts).

The enrichment factor (EF) is an approximation employed to identify the degree to which an element in the aerosol is enriched or depleted (da Rocha et al., 2012). EF was calculated using aluminum as a soil tracer, according to crustal element abundances described in the Appendix of Lee (1999). Elements with EF below 10 were considered to be of crustal origin (not enriched), and elements with EF above 10 were considered to be of non-crustal origin (anomalously enriched) (Pereira et al., 2007). Equation 4 is adopted to calculate EF, where  $C_{Xp}$  and  $C_{Alp}$  are the concentrations of elements X and Al in the sample, and  $C_{Xc}$  and  $C_{Alc}$  are their average concentrations in the Earth's crustal material:

$$EF = \frac{\frac{C_{Xp}}{C_{Alp}}}{\frac{C_{Xc}}{C_{Alc}}} \quad (4)$$

The positive matrix factorization (PMF) receptor model was applied to the datasets (Paatero and Tapper, 1994), including ninety-four samples. The USEPA PMF5.0 software was used. The variable classification followed the established definitions: “bad” when the signal-to-noise ratio (S/N) was below 0.2, “weak” with S/N between 0.2 and 2, and “strong” when S/N was above 2 (Lang et al., 2015). The number of samples below the detection limit (Amato et al., 2016; Contini et al., 2016; Paatero and Hopke, 2003) and the thermal stability of the species (Pereira et al., 2017b) were also considered. Weak variables had their uncertainty three-fold increased, and bad variables were excluded from the model (Norris et al., 2014). Twenty-two species were considered as strong ( $\text{NH}_4^+$ ,  $\text{K}^+$ ,  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ , V, Mn, Ni, Zn, As, Rb, Cd, Sb, Pb, Lev, OC1, OC2, OC3, OC4, PC, EC1, EC2, and EC3), six species were classified as weak ( $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ , Cu, EC4, LMW-PAHs, and HMW-PAHs) and the concentrations of particulate matter were defined as a total variable (weak). Concentrations below the detection limits (DL) were substituted by half of the DL value. Uncertainties were calculated following the procedure by Norris et al. (2014). To evaluate the number of factors, Q robust (QR) was compared to Q theoretical value (QT) as in Pereira et al. (2017b). The final model was obtained with an additional 7% modeling uncertainty. Solutions with three to eight factors were tested, and a final solution with five factors was taken as the best result, with factors that could be well interpreted. Bootstrap mapping (BS) and displacement of factor analysis (DISP) were used to analyze factor solutions, confirming a 5-factor solution as the most feasible. Most of the markers used to identify the sources were within the BS interquartile ranges (box) and mapping ranged from 87%-100%, furthermore, no DISP swaps were observed.

## 2.5 Backward air mass trajectories

For specific pollution events, backward air mass trajectories spanning 48 hours were generated using the HYSPLIT model (Draxler and Rolph, 2003) via the READY (Real-time Environmental Applications and Display System) platform provided by NOAA (National Oceanic and Atmospheric Administration). Trajectory frequencies were calculated to illustrate the origin of air masses reaching the MASP in selected polluted periods, starting at 9:00 AM (local time). The trajectories were calculated at a height level of 500 and 3000 meters above ground level (AGL), based on GDAS meteorological fields with 1 degree resolution.

## 2.6 Ecotoxicity assays


The ecotoxicity of  $\text{PM}_{2.5}$  was screened by the kinetic version of the *Aliivibrio fischeri* bioluminescence-based assay (Kováts et al., 2021). This bioassay mimics the respiratory metabolism of biological systems resulting from exposure to particulate matter. This assay provides an easy-to-quantify endpoint to assess the presence of toxic substances. Inhibition of the bacteria's metabolism by toxic substances is demonstrated by the attenuation of its natural light emittance. Two 17 mm diameter filter sections were cut and ground in an agate mortar. The samples were then transferred to 4 mL vials, to which 2 mL of ultra-pure water was added. The suspensions were prepared with continuous agitation. The manufacturer's reconstitution solution was used to rehydrate the lyophilized bacteria (strain NRRL-B-11177, from Lange Co.), which were then stabilized for 35 min at

12°C. For each sample, serial dilutions in 2% NaCl were prepared in 96-well plates. After adding the bacterial suspensions to the samples, the bioluminescence intensity was continuously read for the first 30 seconds by a Luminoskan Ascent Luminometer (Thermo Scientific). The bioluminescence was reread after 30 min of contact. The Ascent Software (Aboatox Co., Finland) was employed to calculate the EC50 (concentration that causes 50% inhibition of bioluminescence compared to the control). Depending on their toxic units (TU = 100/EC50%), samples were cataloged as non-toxic (TU < 1), toxic (1 < TU < 10), very toxic (10 < TU < 100), or extremely toxic (TU > 100).

### 3 Results and discussions

#### 3.1 PM<sub>2.5</sub> chemical composition and general trend

The concentrations of particulate matter presented a wide variation. PM<sub>2.5</sub> concentrations ranged from 7 to 47 µg m<sup>-3</sup>, averaging 24 µg m<sup>-3</sup> (Figure 2a) and exceeding the new World Health Organization's (WHO) daily recommendations on 75% of sampling days (15 µg m<sup>-3</sup>), but none of national and local limits were surpassed (CONAMA, 2018; WHO, 2021). The period was characterized by low precipitation, the variation of meteorological conditions (temperature, relative humidity, pressure and irradiance are described in the SI (Figure S1). In terms of mass fractions, the most abundant elements observed in the 2019 intensive campaign were K > Al > Cu > Fe (Table 1). The species mentioned are linked to dust resuspension, vehicular sources, and biomass burning. Potassium (K) is attributed to soil resuspension and biomass burning, aluminum (Al) is related to soil resuspension, and iron (Fe) and copper (Cu) are associated with vehicular sources in the MASP (Brito et al., 2013; Pereira et al., 2017b, 2023a, 2023b). Biomass burning tracer levoglucosan (Lev) increased in specific periods (Figure 2b) and will be further discussed. As observed in the 2014 intensive campaign, secondarily formed ions were the most abundant: NO<sub>3</sub><sup>-</sup> > SO<sub>4</sub><sup>2-</sup> > NH<sub>4</sub><sup>+</sup> (Figure 2c and 2d). However, the profile has changed slightly, showing a greater abundance of nitrate over sulfate in most of the period, unlike the intensive campaigns of 2008, 2010, 2013, and 2014 (Table S2), when sulfate was predominant in PM (Pereira et al., 2019, 2017a,b, Souza et al., 2014). The total mass of the most common element oxides was estimated (Al<sub>2</sub>O<sub>3</sub>, SiO<sub>2</sub>, TiO<sub>2</sub>, MnO, and Fe<sub>2</sub>O<sub>3</sub>). Since Si was not determined in this study, SiO<sub>2</sub> was estimated based on the Al content (three times the concentrations of Al<sub>2</sub>O<sub>3</sub>) (Alves et al., 2018). The sum of these oxides can be used to estimate the proportion of crustal species in the PM<sub>2.5</sub> (Almeida et al., 2006) and accounted for 7%.



**Figure 2:** Daily variation in the concentrations of (a) PM<sub>2.5</sub>, OM, EC, and (b) Lev and Lev/Man ratio, (c) fractions of OC and (d) EC, (e) ionic composition, and (f) mass balance. In (b), "Dry periods" are marked with red squares (daily minimum RH < 50%), and the dark rain event is marked with the purple box.



**Table 1:** Median, average, minimum, and maximum concentrations of different species and diagnostic ratios<sup>1</sup>.

	Med.	Ave.	Min.	Max.		Med.	Ave.	Min.	Max.
(µg m <sup>-3</sup> )					(Cont.)				
PM <sub>2.5</sub>	23.7	24.5	6.9	47.2	Cu	93	136	13	931
OC	6.3	7.2	1.2	18.6	Zn	39	49	4.0	250
EC	1.8	2.3	0.3	6.7	As	1.2	1.2	0.3	2.9
SOC	2.7	3.4	0	10.6	Se	1.1	1.4	0.1	5.7
(ng m <sup>-3</sup> )					Rb	1.8	2.2	0.1	6.2
Na <sup>+</sup>	143	166	5	549	Sr	4.6	5.2	0.3	15.4
NH <sub>4</sub> <sup>+</sup>	737	954	61	4241	Mo	1.6	4.9	0.1	31.2
K <sup>+</sup>	247	268	6	686	Ag	0.27	0.26	0.01	0.85
Ca <sup>2+</sup>	101	128	17	517	Cd	0.49	0.66	0.03	2.69
Mg <sup>2+</sup>	44	43	6	95	Sn	5.5	8.2	0.2	30.7
Cl <sup>-</sup>	174	321	14	1505	Sb	3.8	5.7	0.3	27.9
NO <sub>3</sub> <sup>-</sup>	2115	2531	134	10711	Ba	14	16	4	54
PO <sub>4</sub> <sup>3-</sup>	320	318	36	784	La	0.25	0.29	0.01	1.47
SO <sub>4</sub> <sup>2-</sup>	1651	1942	69	6139	Ce	0.23	0.26	0.01	2.09
C <sub>2</sub> O <sub>4</sub> <sup>2-</sup>	158	152	12	319	Tl	0.06	0.07	0.00	0.23
(ng m <sup>-3</sup> )					Pb	11	14	1	38
Lev	315	380	38	1437	Bi	0.16	0.2	0.01	0.9
Man	23	29	7	123	U	0.23	0.25	0.06	0.69
Gal	13	16	7	41	<b>Ratios</b>				
Xyl	6	8	2	75	OC/EC	-	3.1	1.1	10.2
(ng m <sup>-3</sup> )					EC/Cu	-	17.3	2.2	146.5
Al	180	208	11	1142	Fe/Ca <sup>2+</sup>	-	0.8	0.0	7.8
K	271	286	16	757	Cu/Sb	-	23.6	3.6	543.1
Ti	7.5	8.9	1.4	141	Cu/Zn	-	2.8	0.2	23.1
V	1.2	1.7	0.1	8.0	Fe/Cu	-	0.7	0.0	6.8
Cr	1.7	3.4	0.0	20.7	La/Ce	-	1.1	0.5	16.7
Mn	6	6.9	0.6	25	V/Ni	-	0.8	0.1	2.8
Fe	64	99	2	674	SO <sub>4</sub> <sup>2-</sup> /NO <sub>3</sub> <sup>-</sup>	-	0.8	0.0	4.9
Co	0.07	0.08	0.00	0.33	K <sup>+</sup> /Lev	-	0.7	0.0	4.7
Ni	1.5	2.0	0.2	27	Lev/Man	-	13	6.7	58

A predominance of organic matter was observed in PM<sub>2.5</sub>, accounting for a mass fraction of 47%, on average, followed by NO<sub>3</sub><sup>-</sup> and EC (nearly 10%) (Figure 2c). OM and EC represented a significant fraction of PM<sub>2.5</sub>, over 50% on average. Similarly, in the Eastern part of São Paulo (2019-2020), organic matter (OM) corresponded to nearly 40% of PM<sub>2.5</sub> (Vieira et

<sup>1</sup> Average diagnostic ratios were calculated with the average species' concentrations.


al., 2023). Previous studies (2008 and 2014) pointed to higher proportions of EC in PM<sub>2.5</sub> (Table S2). The elemental carbon fraction is reducing worldwide (Chow et al., 2022a, Yamagami et al., 2019) (more details are presented in the SI) and a reduction trend related to lower vehicular emissions of this species (Pereira et al., 2023a, 2023b). The emission limits following a control program policy were upgraded in the early 2010s, similar to Euro 5 and Euro 4, for heavy-duty (HDVs) and light-duty vehicles (LDVs), respectively (Pacheco et al., 2017) and more recently in 2022 (CETESB, 2022). The control policies are associated with reducing carbonaceous species, especially EC (Pereira et al., 2023a,b). Furthermore, adopting biofuels (e.g., ethanol and biodiesel blends) reduces these pollutants' emissions (de Abrantes et al., 2009). The relative reduction of sulfate levels, together with EC, is observed worldwide, as reported in the SI. Following a similar trend to the observed in other Latin American metropolises (Gómez-Peláez et al., 2020), the precursor SO<sub>2</sub> reduced relatively more than NO<sub>x</sub> in the MASP in the last decades (Andrade et al., 2017), and its concentrations were eight-fold lower in 2019 compared to 2000 (Figure S2) (CETESB, 2020). The transport sector became the dominant SO<sub>2</sub> source in the MASP through the 1980s to early 2000s after the reduction of industrial emissions following the adoption of electrical boilers (Kumar et al., 2016). Since the early 2010s, S10 diesel, and S50 gasoline have been adopted to control vehicular emissions, although older trucks were allowed to use S500 diesel (CETESB, 2015). From 2024, a resolution will establish new national specifications for road-use diesel oils, discontinuing S500 fuels totally (MME, 2024). Due to these reductions in the MASP, sulfate vehicular emissions were estimated to be fivefold lower than those from stationary sources in 2019 (CETESB, 2020).

The OC and EC fractions are shown in Figures 2e and 2f, respectively. A predominance of OC2 and OC4 was observed among the organic carbon fractions and EC1 and EC2 among the elemental carbon fractions. The relatively low proportion of EC3 suggests that this site is less affected by HDV emissions, since previous experiments in the MASP have shown that EC3 predominated in a diesel-powered HDV-impacted tunnel (Pereira et al., 2023b; Monteiro dos Santos et al., 2016). In the tunnel studies, the OC2 fraction dominated the emissions and had already been observed in urban areas of São Paulo in 2013 (Monteiro dos Santos et al., 2016). OC1 is the most volatile fraction, which may explain its relatively lower proportion in the particulate matter. Pyrolyzed carbon (PC) represented 16% of OC. Previously, PC represented 20-30% of total OC at the university and downtown sites, accounting for less than 10% of OC in a street canyon (Monteiro dos Santos et al., 2016). This fraction encompasses oxygenated components and links to water-soluble organic carbon (WSOC), secondary formation, and primary sources, including biomass burning (Zhu et al., 2014; Pio et al., 2007; Yu et al., 2002). More recently, it has been connected to HDV emissions, possibly due to the adoption of biodiesel (Pereira et al., 2023b).

Levoglucosan, a monosaccharide anhydride widely used as a biomass-burning tracer, presented the highest average concentration (380 ng m<sup>-3</sup>) among isomer sugars, followed by mannosan (29 ng m<sup>-3</sup>). Compared to a previous intensive campaign (2014), the average K<sup>+</sup> level reduced from 809 ng m<sup>-3</sup> to 268 ng m<sup>-3</sup> (almost three-fold), while levoglucosan presented a similar level (509 ng m<sup>-3</sup>) (Pereira et al., 2017). Since potassium levels are high in crop-burning emissions (Chow et al., 2022a), the significant reduction in concentrations may be due to the lower influence of sugarcane burning since control policies have contributed to the decrease in the number of fires (Valente and Laurini, 2021). Galactosan was present in less than half of the samples, averaging 16 ng m<sup>-3</sup>. It is usually observed in lower proportions than mannosan in biomass burning emissions (Bhattacharai et al., 2019). It is possible to observe in the trajectory cluster a predominance of air masses coming from the northwest (Figure 3a and 3b), suggesting the importance of the transport of biomass burning aerosols from rural areas of São Paulo state and central region of Brazil (nearly 50%) during this period and a low influence of trajectories passing through the ocean (nearly 20%). The highest levoglucosan concentrations were observed on September 10 and 11 (1437 and 1119 ng m<sup>-3</sup>) in a dry period that preceded a cold front. Relatively higher concentrations were also observed for K<sup>+</sup> on both days (543 and 555 ng m<sup>-3</sup>). Back trajectories frequencies arriving at 500 m for these two days point to the typical influence of air masses from the north and northwest of São Paulo state (Figure 3c and 3d), as observed in previous campaigns (Pereira et al., 2017a, 2017b). When a height above the boundary layer is considered (3000 m), it is possible to see a frequency of trajectories passing

302 through areas in the country's central region. The states of Minas Gerais and Goiás, located north of São Paulo state, also  
 303 presented many fires from September 10 to 12 (Figure S3) (INPE, 2019).

304




305

306 **Figure 3:** Backward trajectory clusters for the sampling period, arriving at (a) 500 m and (b) 3000 m, and trajectory  
 307 frequencies for September 10 and 11, 2019, arriving at (c) 500 m and (d) 3000 m.

308 Xylitol was the most detected polyol, with an average concentration of  $8 \text{ ng m}^{-3}$ . This species is mainly found in  
 309 biological material, soil biota, and biomass-burning smoke (Marynowski and Simoneit, 2022; Gonçalves et al., 2021; Caseiro  
 310 et al., 2007). Gonçalves et al. (2021) found a relation between this polyol and biomass-fuelled heating and reported an increase  
 311 from  $2.95 \text{ ng m}^{-3}$  in summer to  $19.9 \text{ ng m}^{-3}$  in winter, in an urban background site in Portugal. Although, much lower  
 312 concentrations were observed in an urban site near sugarcane plantation areas in the countryside of São Paulo (below  $2 \text{ ng m}^{-3}$ )  
 313 (Carvalho et al., 2023). The trajectories were obtained for a period with high xylitol, reaching  $75 \text{ ng m}^{-3}$  on the day August  
 314 30 (Figure S4) and point to the influence of air masses from the north and northwest of São Paulo state, corroborating a biomass  
 315 burning contribution (Figure S5). Relatively high xylitol was also observed in September 10 and 11 (above  $25 \text{ ng m}^{-3}$ ), when  
 316 the highest levoglucosan concentrations were observed. Other polyols were detected in a few samples, as discussed in SI.  
 317 Arabitrol and mannitol levels tend to increase in the wet season, contrary to what was observed for biomass-burning tracers in  
 318 a medium-sized city in the São Paulo state (Carvalho et al., 2023). The study period is expected to have lower pollen-related  
 319 emissions since these spore emissions are enhanced under higher temperatures and humidity (Marynowski and Simoneit,  
 320 2022).

321 The total concentration of PAHs in  $\text{PM}_{2.5}$  ranged from  $1.1$  to  $37.3 \text{ ng m}^{-3}$ , with an average of  $10.1 \text{ ng m}^{-3}$ , higher than  
 322 the one observed in the 2019 extensive campaign (including a wet period), of  $4.7 \text{ ng m}^{-3}$  (Serafeim et al., 2023). The most

abundant compounds observed in the present study were BbF, BeP, and BPe (Figure 4 and Table S3), with BbF being potentially carcinogenic and related to emissions from gasoline-powered vehicles (Ravindra et al., 2008). The predominance of BbF is often observed in São Paulo. This compound is emitted in smaller amounts by LDVs than BaP (Pereira et al., 2023a). However, it is less influenced by chemical decomposition and more persistent (Aubin and Farant, 2000), and together with DBA, presents the most extended residence times in the atmosphere (Keyte et al., 2013).



**Figure 4:** Box plots of polycyclic aromatic hydrocarbon concentrations. EEA recommended annual limit for BaP is indicated by the dashed red line.

Benzo(a)pyrene averaged  $0.6 \text{ ng m}^{-3}$  (ranging from  $0.04$  to  $2.39 \text{ ng m}^{-3}$ ) and surpassed the annual limit recommended by the European Environment Agency (EEA;  $1 \text{ ng m}^{-3}$ ) in 16% of the samples (Ravindra et al., 2008). However, half of the samples exceeded this value if the BaP toxicity equivalent index ( $\text{BaP}_{\text{TEQ}}$ ) is considered (average of  $1.9 \text{ ng m}^{-3}$  and range of  $0.06$ - $8.64 \text{ ng m}^{-3}$ ). The benzo(a)pyrene-equivalent index ( $\text{BaP}_{\text{Eq}}$ ), calculated according to Yassaa et al. (2001), ranged between  $0.04$  and  $4.08 \text{ ng m}^{-3}$ , with an average of  $0.9 \text{ ng m}^{-3}$ , lower than those determined for samples collected in winter in 2010, 2012, 2013 and 2014 in São Paulo (ranging from  $1.1$  to  $3.4 \text{ ng m}^{-3}$ ) (Pereira et al., 2017b, 2017a). This reduction may be attributed to emission regulations and adopting biofuels, such as ethanol and biodiesel, which can lower HMW-PAH emissions (de Abrantes et al., 2009; Pereira et al., 2023a). This trend follows a reduction in PAH emissions in recent decades in developed countries due to the establishment of new regulations (Shen et al., 2011).  $\text{BaP}_{\text{MEQ}}$  and  $\text{BaP}_{\text{TEQ}}$  tended to increase with northern and northwestern winds, more associated with drier weather, stable conditions, and biomass-burning-related aerosols (Figure S6). The opposite was observed with S and SE winds when cold fronts and sea breezes were registered.

### 3.2 Diagnostic ratios and polar plots


The OC/EC ratios in the present study ranged from 1.1 to 10.2, with an average of 3.1 (the monthly averages varied from 3.0 to 3.3) (Figure S7a). In previous tunnel and roadside campaigns in the MASP, ratios lower than 1 were observed, increasing to nearly 2 in more background green areas, where there is increased biogenic influence and secondary formation (Brito et al., 2013; Monteiro dos Santos et al., 2016; Pereira et al., 2023b). The distance to main roads can increase the ratios and values between 1.8 and 3.7 are found in background sites (Amato et al., 2016). However, biomass burning can also increase this ratio reaching over 4.1-14.5 (Watson et al., 2001; Wu et al., 2018). It is noteworthy that the values obtained in the present study

349 were mostly higher than those found in the 2014, 2013, 2010, and 2008 intensive campaigns in the same sampling site ( $1.2\text{--}$   
350  $2.3$ )<sup>2</sup>, in similar dry periods, however in shorter monitoring periods (Pereira et al., 2019, Pereira et al., 2017a,b; Souza et al.,  
351 2014). The ratio increase between these years may indicate a more significant influence of biofuel consumption and reduction  
352 of EC emissions (Dawidowski et al., 2024; Pereira et al., 2023a), since the ethanol sales in the state of SP in 2019 were 75%  
353 higher than in 2013 (MME, 2023). Furthermore, an enhanced contribution from secondary organic carbon (SOC) formation  
354 may also contribute to this. SOC was estimated to reach nearly half of OC, with a maximum of 83% (Table 1). The formation  
355 of SOC is enhanced under higher concentrations of oxidants such as ozone, which was observed in the MASP after 2006  
356 (Andrade et al., 2017). During the lockdown (2020), the OC/EC ratio increased (average of 5.7), following lower primary  
357 vehicular emissions and higher relative contribution of secondary processes (Mbengue et al., 2023; Meng et al., 2021), which  
358 was observed in the MASP after 2006 (Andrade et al., 2017). In Santiago, Chile, there is a trend of increasing photochemical  
359 activity and oxidizing capacity in the atmosphere (2001-2018), leading to an increase in the secondary fraction of aerosols  
360 (Menares et al., 2020). The enhancement of SOC under a more oxidative atmosphere and/or higher temperatures requires  
361 further investigation. During the lockdown (2020), the OC/EC ratio increased (average of 5.7), following lower primary  
362 vehicular emissions and higher relative contribution of secondary processes (Farias et al., in review).

363 To investigate associations between wind direction and diagnostic ratios, polar plots were developed (Figure 5). The  
364 OC/EC ratios appeared to increase with relatively stronger SE winds (Figure 5a). These winds can also transport biogenic  
365 organic carbon from forested areas near the coast and air pollutants from coastal petrochemical and harbour areas.  
366 Furthermore, humidity can favor the partition of more polar SOC, as enhanced RH can lead to water uptake by hygroscopic  
367 submicron particles (Satsangi et al., 2021). In SE parts of MASP, more influenced by forests, less volatile oxidized organic  
368 aerosols were predominant and associated with SOC, derived from VOCs from both biogenic and vehicular sources (Monteiro  
369 dos Santos et al., 2021).

---

<sup>2</sup> Ratios (OC/EC and Lev/Man) were obtained for PM<sub>10</sub> in 2010 and 2013, with the assumption that most of these species are found in the fine fraction in São Paulo.



**Figure 5:** Polar plots considering the diagnostic ratios as a function of wind speed and direction for the ratios OC/EC (a),  $\text{SO}_4^{2-}/\text{NO}_3^-$  (b), Lev/Man (c), Cu/Zn (d), EC/Cu (e), V/Ni (e), LMW-PAHs/HMW-PAHs (g), InP/(InP+BPe) (h), and BaP/(BaP+BeP) (i). Distance from the center is related to wind speed and ratios increase from blue to red.

The average  $\text{SO}_4^{2-}/\text{NO}_3^-$  in this study was 0.8 (0.04-4.6), lower than the 1.6 (from July to December) of the 2019 extensive study (Serafeim et al., 2023). The difference can be attributed to the lower temperatures in the present study (from June to September), which were less favorable to the volatilization of particulate nitrate (Tang et al., 2016). In 2014, higher ratios were observed, with 1.2 in the intensive period and 2.2 in the extensive period (Pereira et al., 2017b). The ratio in the 2008 intensive campaign was higher than in 2014 (1.8) (Souza et al., 2014). Besides the ambient temperature, the reduction in this ratio can be associated with the relatively higher reduction of sulfur dioxide levels, if compared to nitrogen oxides (Sect. 3.1). The ratio  $\text{SO}_4^{2-}/\text{NO}_3^-$  appeared to increase with SE and E winds (Figure 5b), suggesting that the production of sulfate was favored by humid and cloudier conditions associated with these air masses and/or a possible influence of sulfur dioxide emitting sources in this area (e.g., industries and HDVs). Sulfate particles can grow with condensation by adding sulfate and water in the aerosol droplet process (Guo et al., 2010). The SE part of MASP is influenced by industrial sources and high sulfate levels associated with locally emitted  $\text{SO}_2$  (Monteiro dos Santos et al., 2021).

Galactosan and mannosan are products of hemicellulose thermal decomposition, while levoglucosan is formed during the combustion of cellulose (Simoneit, 2002). Since the amount of cellulose and hemicellulose varies with biomass type, with hardwood containing relatively higher amounts of cellulose, the Lev/Man ratio can distinguish the smoke from different biofuels (Li et al., 2021; Zhu et al., 2015). Typical ranges of 15–25 and 3–10 have been documented for hardwood and softwood burning, respectively (Li et al., 2021). The average Lev/Man in the present study was 13 (6.7 - 58.0), slightly higher than averages observed from 2014, 2013, and 2010 (ranging from 8 to 12)<sup>1</sup> (Pereira et al., 2017a,b, 2019). Previous ratios were



392 closer to the values found in chamber sugarcane burnings (Lev/Man = 10) and areas impacted by this type of smoke plumes  
393 (Lev/Man = 9) (Hall et al., 2012; Urban et al., 2014). In an agroindustrial region in northern São Paulo state, a change in the  
394 Lev/Man ratio was observed in 2020 (Lev/Man = 19) (Scaramboni, 2023), this increase was attributed to a change in the  
395 biomass burning profile, with reduced sugarcane straw burning and increased use of sugarcane bagasse in power plants in the  
396 last decade (MME, 2024), alongside the contribution of forest fires and other agricultural residue burnings. During high  
397 levoglucosan events at the end of the 2019 campaign, under lower relative humidity (Figure 2), the Lev/Man ratios exceeded  
398 40 and suggested the influence of different types of biomasses. Furthermore, in areas impacted by forest fires in the Amazon,  
399 likely related to hardwood, the ratios ranged from 15 to 24 (Decesari et al., 2005; Graham, 2002), while in central areas of  
400 Brazil, burning grass in the *cerrado* (tropical savanna biome) resulted in low amounts of mannosan, as this biomass contains  
401 less mannose compared to hardwood (Scaramboni et al., 2024). Before the dark precipitation event registered on August 19,  
402 associated with smoke transported from areas in central Brazil and the Amazon (Pereira et al., 2021), Lev/Man values also  
403 reached values above 20.

404 The median  $K^+/Lev$  ratio was 0.7, considerably lower than that observed in the 2014 dry period (1.6) (Pereira et al.,  
405 2017a). Lower  $K^+/Lev$  ratios, typically below 1, have been found in emissions from wood stove combustion and forest fires  
406 (Caseiro et al., 2009). Jung et al. (2014) reported similar  $K^+/Lev$  ratios for hardwood and softwood burning (0.04 and 0.02,  
407 respectively)<sup>2</sup>, but this ratio was higher for crop burning (averaging 1.9)<sup>2</sup>. Urban et al. (2012) documented an average ratio of  
408 around 4 for fine sugarcane-burning particles<sup>3</sup>, justifying this value with the enrichment of  $K^+$  in the leaves and the inefficient  
409 formation of levoglucosan during the flaming phase. The reduction in the proportion of  $K^+/Lev$  in the present study may be  
410 associated with reducing sugarcane burning after regulation. Some  $K^+/Lev$  peaks were observed on days with very low Lev  
411 concentrations (06/17, 07/27, 07/28, 08/06, and 09/01) (Figure S7b), suggesting the influence of soil-related  $K^+$ .

412 The Lev/Man ratio increased with stronger winds, whether from S/SE or N (Figure 5c), denoting the influence of  
413 emissions from burning various types of biomasses. If the polar plot is obtained with Conditional Probability Function (CPF)  
414 probability and Non-parametric Wind Regression (NWR), the increase with stronger N winds is prominent (Figure S8a and  
415 S8b). The lowest ratios were found with low-speed SE winds, probably due to wood burning in some restaurants in that area.  
416 Brazilian pizzerias traditionally use eucalypt logs in wood stoves. However, recently, briquettes have been adopted (Lima,  
417 2015). Sun et al. (2019) reported Lev/Man ratios in the range of 17-19 for different types of briquettes. Some authors found  
418 Lev/Man ratios mostly below 10 for woodstove emissions from different types of wood logs, while briquettes presented ratios  
419 of 29.7 in hot start and 1.4 in cold start conditions (Gonçalves et al., 2011).

420 The diagnostic ratios EC/Cu, Fe/Cu, Cu/Zn, and Cu/Sb were associated with the proportion of LDVs and HDVs in  
421 tunnels in the MASP (Pereira et al., 2023b). In previous studies, EC was more related to HDV emissions, while Cu was linked  
422 to LDV emissions (Brito et al., 2013). EC/Cu ratios in the present varied between 2 and 146, and the average ratio was 17. It  
423 was nearer the values observed for LDVs (5-8), rather than the ratios associated with a more HDV-impacted fleet (80-189).  
424 Sb has been described as a tracer of abrasion sources, including brake wear (Thorpe and Harrison, 2008). Cu is also present in  
425 Brazilian exhaust emissions from ethanol/gasohol vehicles (Brito et al., 2013). The present average Cu/Sb ratio was 24 (4-  
426 543), higher than the median values observed in tunnels (11-14), which may be attributed to the distance of the sampling site  
427 from a high-traffic area and a lower contribution from brake abrasion sources. The average Cu/Zn was 2.8 (0.2-23.1), falling  
428 between the median value observed for LDV (6.2) and that documented for HDV+LDV-impacted tunnels (0.9). The Cu/Zn  
429 and Cu/Sb ratios increased with southern weak winds (Figures 5d and S8c), suggesting an enrichment in copper and  
430 predominance of LDV emissions in this area, possibly associated with the traffic of vehicles in the nearby residential  
431 neighborhood. On the other hand, the EC/Cu and Fe/Cu ratios increased with N winds (Figures 5e and S8d), which may  
432 indicate an influence of HDVs passing in the expressway located north of the campus.

---

<sup>3</sup> These authors presented the ratios as Lev/ $K^+$ .

The average V/Ni ratio was 0.8 (0.1-2.8). Higher values are found for residual fuel oil combustion range from 1 to 3 (Johnson et al., 2014). This ratio increased with strong E/NE/SE winds (Figure 5f), suggesting an influence from oil burning in industries located in these areas (Vieira et al., 2023; Souto-Oliveira et al., 2021). The La/Ce ratio, similarly to the latter, was favored by S/SE winds (Figure S9h). Higher ratios (4.3) were observed for fluidized-bed catalytic cracking (FCC) during petroleum refining, while a lower value (0.7) was documented for automobile catalyst emissions (Kulkarni et al., 2006). A petrochemical complex in the southeastern area of the MASP may explain this influence (Caumo et al., 2022; Gioia et al., 2017). Previous studies have observed that cold fronts may be associated with isotopic fingerprints from the Cubatão petrochemical industrial area outside the MASP (Souto-Oliveira et al., 2018).

Diagnostic ratios between PAHs for the analyzed samples were calculated and shown in Table S3. BaP/(BaP+BeP) values lower than 0.5 indicate that the analyzed particles were aged, as BaP undergoes photolysis more quickly than BeP (Tobiszewski and Namieśnik, 2012). The median was equal to 0.32, suggesting a predominance of aged particles. In three samples (collected on 27/07, 03/08, and 28/08), this ratio was close to 0.5, indicating fresh emissions. The Flt/(Flt+Pir) median was 0.48. According to De La Torre-Roche et al. (2009), values of this ratio ranging from 0.4 to 0.5 are characteristic of the burning of fossil fuels. InP/(InP+BPe) median ratio was equal to 0.48, similar to what was observed in 2014 by Pereira et al. (2017b). Values between 0.2 and 0.5 are associated with emissions from the burning of fossil fuels (Yunker et al., 2002). The BaA/(BaA+Cri) presented an average of 0.38, falling near the observed for vehicular sources (0.2-0.35) (Akyüz and Cabuk, 2010).

ΣLMW-PAHs/ΣHMW-PAHs median ratio was 0.35, within the range reported for pyrogenic (<1) emissions by some authors (Zhang et al., 2008). However, this ratio was previously found to vary with the proportions of LDVs and HDVs (0.7 and 7.5, respectively) (Pereira et al., 2023b) and with ambient temperature (Tobiszewski and Namieśnik, 2012). The ratios Pyr/BaP, Pyr/BbF, and Flt/BbF were associated with the proportion of LDVs and HDVs, increasing with the latter (Pereira et al., 2023b). Nevertheless, the ratios can be influenced by the volatilization of Pyr and Flt in warmer conditions and the photodegradation of BaP. The values were similar to those observed for gasoline emissions (near or below 1).


ΣLMW-PAHs/ΣHMW-PAHs and Pyr/BaP ratios appeared to be affected by winds coming from NW (Figures 5g and S8e), with a mixed biomass burning and vehicular influence, and by cooler winds from the S, which may favor the condensation of LMW-PAHs in the particulate phase (Ravindra et al., 2008). Temperature was the meteorological parameter that most affects total and individual PAHs, with a more substantial influence on LMW-PAHs (Amarillo and Carreras, 2016). InP/(InP+BPe) ratios approached the values (above 0.5) found for grass, wood, and coal combustion with NW winds (Figure 5h) (Yunker et al., 2002). It was not possible to observe the same for Flt/(Flt+Pyr) (Figure S8f), which increased with S and SE winds, suggesting that it may be linked with the cooler/cloudier conditions and the shorter photochemical residence time of Pyr compared to Flt (Keyte et al., 2013). The highest BaP/(BaP+BeP) values were registered for lower wind speeds (Figure 5i), suggesting fresh local emissions (Tobiszewski and Namieśnik, 2012), although it also increased with stronger NW winds.

### 3.3 Source apportionment (PMF)

The factor profiles obtained by Positive Matrix Factorization are shown in Figure 6. A five factor solution was obtained, attributed to: biomass burning (BB), secondary formation (SF) and industries (IN), and two associated with vehicular emissions (VE1 and VE2). Factor 1 (BB) was characterized by biomass-burning-related species such as levoglucosan,  $K^+$ , and some carbonaceous species (Bhattarai et al., 2019; Simoneit, 2002). However, the presence of Cd, Sb, and Pb suggests a mixture with vehicular sources (Thorpe and Harrison, 2008) or that these species may be linked with waste burning (La Colla et al., 2021). This factor was also characterized by a significant contribution from pyrolyzed carbon, a fraction of water-soluble organic carbon (WSOC), often associated with biomass burning and SOA (Yu et al., 2002; Zhu et al., 2014). Recently, it has been observed that the biomass burning factor can mix with secondary organic carbon contribution in a Korean industrial area (Han et al., 2023). Among elemental carbon fractions, EC1 was the most abundant. Some elements, such as Rb, also appeared



475 in the biomass burning profile. This species has been described as a component of some types of soil by Calvo et al. (2013),  
 476 wood-burning emissions (Fine et al., 2001), and biomass burning aerosols in the Amazon (Artaxo et al., 1994). Recently,  
 477 water-soluble rubidium in fine particulate matter has been assigned to wood-burning emissions and considered an alternative  
 478 biomass-burning tracer (Massimi et al., 2020). This factor increased with winds coming from N and NW, corroborating the  
 479 influence of the transport of biomass-burning aerosols from the countryside and forested areas in northern and central Brazil.  
 480 It decreases with S and SW winds, suggesting a reduction with cold fronts and sea breezes (Sánchez-Ccoyllo et al., 2002). The  
 481 results agree with Souto-Oliveira et al., (2023), that showed the impact of the transport of wildfires to the MASP during fine  
 482 particulate matter exceedance events in 2020.



483 **Figure 6:** Profiles identified with the PMF receptor model and polar plots (a-e). Model reference runs (Ref. run) are  
 484 marked in red, bootstrap runs (BS) and standard deviation (sd) in grey.  
 485  
 486

Factor 2 (SF) was highly loaded with secondary formation species, mainly  $\text{NH}_4^+$  and  $\text{SO}_4^{2-}$ . This process likely involves the oxidation of sulfur dioxide, the formation of sulfuric acid, neutralization with ammonia, and the production of ammonium sulfate ( $(\text{NH}_4)_2\text{SO}_4$ ) (Han et al., 2023; Ianniello et al., 2011). Furthermore, this factor also presented loadings of some primary emission species, such as vanadium. Serafeim et al. (2023) also observed high loadings for Ni and V in the secondary-related factor in the MASP. These species originate mainly from industrial emissions (Calvo et al., 2013). Previously, it has been observed with the WRF-Chem model that secondary formation explains 20 to 30% of  $\text{PM}_{2.5}$  in the MASP (Vara-Vela et al., 2016), increasing in the summertime (Pereira et al., 2017b). The SF factor was previously attributed to vehicular-related emissions in a previous study in 2014 (Pereira et al., 2017). However, with the reduction of the emissions of  $\text{SO}_2$  emissions by vehicular sources (CETESB, 2020), the contribution of industries has become relevant (Section 3.1). However, most of the industries are located on the outskirts, whereas vehicular emissions are more concentrated in the city's central areas. Thus, the apportionment of secondary sulfate formation can be a subject for further studies.

Factor 3 showed high loads for V, Ni, and nitrate. Earlier studies in the MASP have associated Ni and V with residual oil burning and industrial sources (Andrade et al., 1994; Castanho and Artaxo, 2001). Heavy oil combustion in industrial boilers also emits nickel and zinc, especially in the ultrafine fraction (smaller than  $0.1 \mu\text{m}$ ) (Jang et al., 2007; Linak et al., 2004). Factor 3 (IN) increased with the same wind direction as Factor 2 (SF), which suggests that it may be partially related to secondary formation, possibly due to emissions of precursor gases, such as  $\text{NO}_x$  and  $\text{SO}_2$ , or associated with the same air masses. Secondary aerosol formation can mislead the separation of factors in PMF, as observed by Faisal et al. (2024).


Factor 4 (VE1) was loaded with soil resuspension constituents, such as  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ , as well as with species previously related to vehicular emission in MASP, such as As and Cu, carbonaceous species (OC1, OC2, OC3, OC4, EC3, and EC4), LMW-PAHs and HMW-PAHs (Pereira et al., 2017b, 2023a, 2023b). In the extensive 2019 campaign, an independent road dust factor was observed, representing 32% of  $\text{PM}_{2.5}$  (Serafeim et al., 2023). The association with construction-related calcium, found in concrete material, can also be considered (Bourotte et al., 2006). EC3 and EC4 were abundant in this factor. EC3 was anteriorly related to HDV emissions in MASP, while EC4 was found in an urban canyon. Furthermore, OC2, OC3, and OC4 were found in LDV emissions in similar proportions (Monteiro dos Santos et al., 2016; Pereira et al., 2023a). Levoglucosan was also observed in a smaller proportion in this factor. As observed in 2014, this factor overlaps with factor 1 (Pereira et al., 2017b), as they increase with the same wind direction. However, the contribution of factor 4 increases with NW stronger winds, corroborating the resuspension of road dust as the main source. Strong NW winds are often attributed to prefrontal conditions (Ribeiro et al., 2018). Another aspect contributing to the overlap is that this anhydrosugar is also found in the  $\text{PM}_{10}$  from the wear between tires and pavements, given that wheel rubbers have cellulose in their composition (Alves et al., 2020). However,  $\text{PM}_{2.5}$ -bound levoglucosan is emitted in much smaller proportions by this non-exhaust vehicle emission source compared to biomass burning (Bhattarai et al., 2019). The “hybrid” factor may occur because the aerosol in the urban atmosphere, comprising mainly vehicular-related species, road dust, and biomass burning emissions, arrives mixed at this semi-background site. Additionally, it is challenging to differentiate HDV from LDV emissions in this site (Pereira et al., 2017a). The increase in time resolution for PMF can reduce mixing in source profiles, decreasing sample variability (Wang et al., 2018). This can be achieved through real-time techniques (e.g., Aerosol Chemical Speciation Monitor) (Monteiro dos Santos et al., 2021). Proportionally, non-exhaust emissions, such as road dust, will increase in the future (Thorpe and Harrison, 2008) since exhaust emissions are on a downward trend due to increasingly stricter regulations and improved treatment systems.

Factor 5 (VE2) presented several species of vehicular origin, such as Mn, Ni, Cu, Zn, Sb, and the potentially toxic Cd and Pb (Brito et al., 2013; Pereira et al., 2023a), in addition to OC and EC fractions, especially EC3, the latter being more associated with HDVs (Pereira et al., 2023b). This vehicular factor increased with lower wind speed, suggesting a local contribution. The site is located next to an avenue with a constant flow of buses and LDVs during the day and is near a busy expressway. Despite the lower proportion, the emissions of buses and trucks contributed to almost half of black carbon in the

530 MASP (Brito et al., 2018). In the SEM study performed by Bourotte et al. (2006), particles rich in carbon, Cu, and Zn were  
 531 identified and associated with incomplete fossil fuel burning. Zinc is also enriched in tire wear particles, as zinc oxide is added  
 532 to the tires in the vulcanization process (Thorpe and Harrison, 2008). Copper was previously assigned to LDV exhaust  
 533 emissions and to the corrosion of the internal parts of the MASP fleet engines (Ferreira da Silva et al., 2010). It was also  
 534 pointed out as a brake dust particle tracer (Thorpe and Harrison, 2008).

535 Figure 7 shows the contributions of each type of source to  $PM_{2.5}$  and their variation between sampling days. Biomass  
 536 burning accounted for 25% of  $PM_{2.5}$ . This contribution was slightly higher than that documented in the intensive campaign of  
 537 2014 (18.3%). In the extensive study performed in 2019 (including dry and wet seasons), the biomass burning factor  
 538 represented 13% of ambient  $PM_{2.5}$ , a similar share to that observed in the 2014 extensive period (Pereira et al., 2017b; Serafeim  
 539 et al., 2023), and lower than the one of the intensive campaign between June and September of the present study, a period  
 540 typically more affected by biomass burning (Pereira et al., 2017a). In a major city located in central Brazil (Cuiabá), biomass  
 541 burning was relevant and represented one third of  $PM_{10}$  (Parra et al., 2024). Several peaks of the biomass burning factor  
 542 coincided with lower RH values. An increase in the contribution of this factor was observed two days before the sky-darkening  
 543 phenomenon at MASP on August 19, when dark precipitation was recorded, attributed to pollutants emitted by biomass burning  
 544 (Pereira et al., 2021). Overall, crop burning and wildfires can explain a significant share of  $PM_{2.5}$  in this period. However,  
 545 regional biomass burning profile is changing, with reduced sugarcane straw burning, increased contribution of wildfires, and  
 546 sugarcane bagasse use as fuel, as discussed in Section 3.2. The study of Knorr et al. (2017) predicts that, in the context of  
 547 climate change, during the fire seasons, wildfire-related  $PM_{2.5}$  levels in many regions (including parts of South America) can  
 548 increase to dangerous levels, even with accentuated reductions of other anthropogenic emissions. In the last decade, wildfire  
 549 smoke has influenced  $PM_{2.5}$  trends across much of the US, including major cities, complicating efforts to control this pollutant.  
 550 (Buchholz et al., 2022). In temperate regions, the high contribution of biomass burning to  $PM_{2.5}$  in the winter still can be  
 551 attributed to residential heating (Nava et al., 2020). In the temperate metropolis Beijing (China), biomass and coal burning can  
 552 represent nearly half of  $PM_{2.5}$  in winter, where these sources are used for heating and cooking in nearby rural areas (Srivastava  
 553 et al., 2021). Due to regulations, open-field burning in China has become less frequent. However, biomass burning for home  
 554 heating in northern China can affect downwind areas such as South Korea (Cheong et al., 2024).

555



556


557

558 **Figure 7:** Daily factor contribution (a) and  $PM_{2.5}$  contributions (b) identified with the PMF receptor model.

559

560 The vehicular-related sources, such as local LDVs and HDVs (12%), vehicular emissions plus road dust (29%), represent  
 561 a significant share of particulate matter in the city (over 40%). Vehicular sources were observed as dominant in 2013, 2014  
 562 and 2015 (Souto-Oliveira et al., 2021; Emygdio et al., 2018; Pereira et al., 2017a). Previous studies were performed in other  
 563 Brazilian cities, where vehicular emissions ranged from 3% to nearly half of  $PM_{2.5}$ . In coastal sites (Rio de Janeiro, Vitória,

and Recife), sea spray can be relevant (Galvão et al., 2019; Justo et al., 2023; dos Santos et al., 2014). Despite the entrance of sea breezes on a few occasions, it was not possible to identify this source in the present study. As discussed in Section 3.1, few trajectories passed by the ocean. The factor VE1 increased throughout the sampling period and dominated  $PM_{2.5}$  at the end of August and the beginning of September, indicating greater dust resuspension at the end of winter as a result of the long dry spell and higher wind speeds (Figure S9). In another source apportionment study performed in the east region of São Paulo in 2019, four factors were found to contribute to  $PM_{2.5}$ : heavy-duty vehicles (42%), light-duty vehicles (9.9%), soil and local particles (38.7%) and local sources (8.6%) (Vieira et al., 2023), with a higher share of vehicular sources than the present study and an increase of soil particles in the dry season. According to Serafeim et al. (2023), vehicular emissions and biomass burning were associated with enhanced oxidative potential. Secondary formation of aerosol from oxide precursors, which can be emitted from vehicular and industrial sources, represented 21% of  $PM_{2.5}$ . Contrastingly, secondary aerosol was predominant in São Paulo during the 2020 lockdown (nearly 40%), followed by biomass burning (30%), with a relative reduction of the contribution of vehicular sources (Farias et al., in review). Industrial sources accounted for only 13.6%, a similar level to that observed in 2014 (over 10%). A relatively low impact of industrial emissions is felt in this part of the city, as nowadays, industries are located on the outskirts of the MASP and the countryside (Kumar et al., 2016) and no large industries are found near the sampling site. Industrial sources can be relevant in other Brazilian cities, pelletizing and steel industry represented more than 60% of  $PM_{2.5}$  in Vitória (Galvão et al., 2019). The contribution of this factor, combined with the frequencies of the trajectories, suggested the influence from areas to the east and northeast, including parts of MASP and other neighboring metropolitan areas with a high concentration of industries. The Simplified Quantitative Transport Bias Analysis (SQTBA) approach was adopted, it recognises the plume dispersion process, considering it as part of back trajectory analysis (Figure 8).



**Figure 8:** (a) Trajectory analysis for the contribution of factor 3, and (b) participation of industries for municipalities in São Paulo state (industrial transformation value), adapted from SEADE (2019).

### 3.4 Correlations between PMF results and other variables

The factors obtained with the PMF receptor model were correlated with other variables: meteorological data (temperature, relative humidity, and wind) collected from the local station, concentrations of chemical species, diagnostic ratios and results obtained from the ISORROPIA thermodynamic model (modeled solid and liquid inorganic aerosol, water content,  $NaNO_{3(s)}$ ,  $Na_2SO_{4(s)}$ ,  $NaHSO_{4(s)}$ ,  $NaCl_{(s)}$ ,  $NH_4Cl_{(s)}$ ,  $NH_4NO_{3(s)}$ ,  $(NH_4)_2SO_{4(s)}$ ,  $NH_4HSO_{4(s)}$ ,  $CaSO_{4(s)}$ ,  $Ca(NO_3)_2_{(s)}$ ,  $CaCl_{2(s)}$ ,  $K_2SO_{4(s)}$ ,  $KHSO_{4(s)}$ ,  $KNO_{3(s)}$ ,  $KCl_{(s)}$ ,  $MgSO_{4(s)}$ ,  $Mg(NO_3)_2_{(s)}$ ,  $MgCl_{2(s)}$ ,  $H^+_{(aq)}$ ,  $Na^+_{(aq)}$ ,  $NH_4^+_{(aq)}$ ,  $Cl^-_{(aq)}$ ,  $NO_3^-_{(aq)}$ ,  $SO_4^{2-}_{(aq)}$ ,  $HSO_4^-_{(aq)}$ ,  $Ca^{2+}_{(aq)}$ ,  $K^+_{(aq)}$  and  $Mg^{2+}_{(aq)}$  (Table S4 and Figure S10). Overall, the components were negatively correlated with RH and positively with temperature, which typically indicates dry air masses and unfavorable conditions for the dispersion of atmospheric pollutants (Sánchez-Ccoyllo et al., 2002), more discussions are presented in the SI.

596 The biomass burning factor correlated with BPe and InP, species typically associated with vehicular emissions  
 597 (Ravindra et al., 2008). However, the InP/(BPe+InP) polar plots suggested an influence of biomass burning with northwest  
 598 strong winds (Sect. 3.2). The low correlation of retene with BB and levoglucosan suggests that this species may be related to  
 599 a local biomass-burning source or the influence of gas-particle partition of this semi-volatile species (Ravindra et al., 2008).  
 600 Chloride was moderately correlated with BB and VE2 factors ( $r > 0.5$ ), strongly correlated with mannosan and EC1 ( $r > 0.7$ ),  
 601 and weakly correlated with levoglucosan, suggesting a different biomass burning profile, such as wood burning in restaurants  
 602 or biomass burning associated with waste (Kumar et al., 2015). Xylitol was weakly correlated with BB and VE1 ( $r > 0.3$ ), it  
 603 also presented a moderate negative correlation with RH ( $r \sim -0.3$ ), suggesting an increase under drier conditions. Notably, it  
 604 presented weaker correlations with potassium than Lev, suggesting a biomass-burning origin less associated with crop burning.  
 605 The association of xylitol with biomass burning was observed in other studies (Clemente et al., 2024; Gonçalves et al., 2021).

606 The secondary aerosol formation factor was moderately correlated with the modeled liquid inorganic aerosol and water  
 607 content of the aerosol, in addition to the modeled secondary species  $\text{NH}_4^+(\text{aq})$ ,  $\text{SO}_4^{2-}(\text{aq})$ , and  $\text{HSO}_4^-(\text{aq})$  ( $r > 0.6$ ). Correlation with  
 608 modeled water content suggests that the secondary formation pathway in São Paulo depends on humidity.  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$   
 609 formation in clouds and fog droplets is attributed mainly due to the heterogeneous aqueous transformation of  $\text{SO}_2$  and  $\text{NO}_x$   
 610 under high relative humidity (RH) during haze episodes in Beijing (Huang et al., 2016). Among the ratios, the secondary  
 611 formation factor presented weak to moderate correlations with  $\text{NO}_3^-/\text{EC}$ ,  $\text{SO}_4^{2-}/\text{EC}$ , and  $\text{NO}_3^-/\text{Zn}$  ( $r > 0.4$ ), suggesting that they  
 612 can be helpful in identifying secondary formation events, an opposite trend to that observed for the BB, VE1, and VE2 factors.

613 The IN factor presented moderate correlations with the liquid inorganic aerosol and water content in the aerosol,  
 614 possibly associated with more humid air masses coming from the east, as particles can grow in the aerosol droplet process  
 615 (Guo et al., 2010). Notably, it was correlated with aerosol acidity ( $\text{H}^+(\text{aq})$ ), perhaps due to gaseous oxides that can undergo  
 616 secondary reactions and produce acidic species.  $\text{H}^+(\text{aq})$  was also moderately correlated with nitrate, suggesting an influence of  
 617 nitric acid (Ianniello et al., 2011). Selenium was moderately associated with the IN factor. This species was previously  
 618 attributed to industrial sources (Vieira et al., 2023). Sn was moderately correlated with IN ( $r \sim 0.5$ ) and strongly correlated with  
 619 Tl. The IN factor presented a weak correlation with the V/Ni ratio ( $r \sim 0.5$ ), denoting the contribution from the burning of crude  
 620 oil (Johnson et al., 2014). IN was also moderately correlated with  $\text{NO}_3^-/\text{EC}$ , suggesting a connection with secondary formation.  
 621 There was a weak correlation between SF and IN, pointing to a common origin between both, maybe due to the emission of  
 622 gaseous precursors and secondary formation.

623 The PMF factor associated with vehicular exhaust and dust resuspension (VE1) presented weak correlations with most  
 624 species. The factor presented moderate correlations with the modeled mass of solid inorganic aerosol ( $r \sim 0.6$ ). This factor also  
 625 displayed a moderate positive correlation with temperature ( $r > 0.6$ ) and negative with relative humidity and pressure ( $r < -$   
 626  $0.5$ ), suggesting an increase of this factor during prefrontal conditions. Among the factors, VE2 and BB presented higher  
 627 correlations with the toxicity equivalent indexes BaP-TEQ and BaP-MEQ ( $r > 0.5$ ), suggesting a contribution of these sources  
 628 to carcinogenicity and mutagenicity. The correlations of VE2 were higher for BaA and HMW-PAHs such as Per, InP, BPe,  
 629 and Cor ( $r > 0.5$ ), which are found in LDV emissions (Pereira et al., 2023a). La was moderately associated with VE2, and  
 630 strongly with Mn, Co, and Ce. La and Ce derive mainly from catalyst emissions (Kulkarni et al., 2006). Fe presented strong  
 631 correlations with VE2, and Ti, Mn, and Ce. These species are linked to vehicular and soil-related sources (Pereira et al., 2017a,  
 632 Brito et al., 2013). Co, a component of tire debris (Thorpe and Harrison, 2008), was moderately correlated with VE2 and  
 633 strongly correlated with La, Tl, and Mn. VE2 presented moderate correlations with  $\text{Fe}/\text{Ca}^{2+}$  ( $r \sim 0.6$ ), which may indicate  
 634 enrichment of anthropogenic iron (Pereira et al., 2023a).

635 Some species were less source-specific.  $\text{PO}_4^{3-}$  presented no correlations with PMF factors and was moderately  
 636 correlated with  $\text{Na}^+$  and  $\text{Ca}^{2+}$  ( $r > 0.45$ ). Oxalate was not correlated with specific factors. It presented moderate correlations  
 637 with  $\text{NH}_4^+$ ,  $\text{K}^+$ ,  $\text{NO}_3^-$ ,  $\text{SO}_4^{2-}$ , As, Rb, Sb, Tl, Pb, and OC ( $r > 0.5$ ), suggesting multiple sources, as previously observed, such as  
 638 vehicle exhaust, biomass burning, and biogenic activity, in addition to secondary formation (Guo et al., 2010).  $\text{Na}^+$ , as



previously observed in the MASP (Vieira-Filho et al., 2016b), was not specific for any source and presented a weak correlation with some other species ( $\text{Ca}^{2+}$ ,  $\text{Cl}^-$ ,  $\text{PO}_4^{3-}$ , As, Rb, Sr, and Cd) ( $r > 0.3$ ), without strong correlation with  $\text{Cl}^-$ , which can be explained by the relatively low influence of sea spray in this site in the studied period (Section 3.1). Aluminium displayed low correlations with the PMF factors and was strongly correlated with Ti ( $r > 0.7$ ) and moderately with Ce ( $r > 0.6$ ), pointing to an origin in crustal materials (Hetem and Andrade, 2016). Ti was also strongly correlated with Ce. Both species presented low enrichment, suggesting a mineral origin, apart from vehicular emissions. Cr was not source-related but presented a relatively high correlation with Ni ( $r \sim 0.5$ ). Both species were previously associated with industrial sources (Bourotte et al., 2011; Castanho and Artaxo, 2001). Arsenic, a highly toxic element, showed relatively moderate correlations with BB, IN, and VE2 and strong with Rb, Ag, and Cd. Arsenic has multiple sources, including industries (Calvo et al., 2013).


### **3.5 Particle size distributions: associations with aerosol source apportionment and meteorological scenarios**

Submicrometer particle number size distributions provided information about the relative contribution of different particle size modes and their temporal evolution. Total particle number concentrations averaged  $1.03 \pm 0.59 \cdot 10^4 \text{ cm}^{-3}$ , with a mean geometric diameter of  $51 \pm 16.1 \text{ nm}$ . Size distributions were dominated by the Aitken mode (mean geometric diameter range 50-100 nm) and the accumulation mode (diameter above 100 nm), respectively, accounting for 46% and 32% of particle number concentrations, on average. The nucleation mode (mean geometric diameter below 30 nm) was less frequent, contributing to an average of 21% of the particle number concentration on average (Table S5). Events of new particle formation and growth were infrequent, possibly due to the relatively high particle loading and condensational sink. Nearly 70% of the monitored particles had diameters below 100 nm, classified as ultrafine particles, which can penetrate the extensive area of the lungs and reach other organs through the lung vasculature (Schraufnagel, 2020). The predominance of Aitken mode particles suggests the influence of relatively fresh particles compared to the typically aged accumulation mode particles. A previous study in the MASP reported associations between inorganic species and the Aitken mode particles, whereas aged oxygenated organic particles were mostly associated with the accumulation mode (Monteiro dos Santos et al., 2021).

The size distribution measurements were averaged to match the filter sampling periods, providing a link with the aerosol chemical composition measurements and source apportionment. No significant differences were observed in the particle size distributions in days with a predominance of vehicular emissions or biomass burning. Although the aerosol chemical composition differed under the influence of different emission sources (Section 3.3), the particle size distributions were typically dominated by the Aitken mode, suggesting the presence of relatively fresh particles. However, there was a clear difference in the size distributions measured under high aerosol loadings ( $\text{PM}_{2.5} \geq 15 \mu\text{g m}^{-3}$ , WHO guideline), with an increase of Aitken and accumulation mode particles, especially at the end of the day (Figure S11). The atmospheric conditions associated with high PM concentration (low boundary layer height, weak ventilation, absence of precipitation, and clear sky conditions) likely favored secondary aerosol production and particle growth by condensation in pre-existing particles (Sánchez-Ccoyllo and Andrade, 2002; Santos et al., 2018), which might explain the increase of larger particles (Monteiro dos Santos et al., 2021).

Four days were selected within the study period to assess the dynamics of particle size distribution in high temporal resolution, and the influence of different meteorological scenarios. Figures 9a and 9b show the evolution of size distributions in two days with a strong contribution of secondary aerosol formation, according to the source apportionment analysis (Figure 7). New particle formation events were observed these days, with sub 30 nm particles emerging in the morning, followed by a consistent growth towards the Aitken mode until the mid-afternoon, with growth rates of  $5.1$  and  $3.3 \text{ nm h}^{-1}$ , respectively. The observed growth may be associated with the condensation of secondary organic and inorganic aerosol species. Previous studies

in São Paulo indicated a large fraction of secondary inorganic species in the Aitken mode, likely produced from the reaction of nitric acid and ammonia (Carbone et al., 2013; Monteiro dos Santos et al., 2021). Another interesting feature was the entrance of the sea breeze around 4 PM local time on August 15<sup>th</sup>, marked by a dashed line in Figure 9a. The sea breeze clearly affected the measured size distributions, leading to a fast particle concentration decrease, from  $1.8 \cdot 10^4$  to  $1.0 \cdot 10^4 \text{ cm}^{-3}$ , associated with a mean geometric diameter increase from  $44.2 \pm 2.0$  to  $47.0 \pm 2.0 \text{ nm}$ . The observed changes in the size distribution may be explained by the transport of cleaner and humid air from the coast. In addition to oceanic air masses, aged particles from industrial emissions near the coast may also be transported to the MASP during sea breeze events.




**Figure 9:** Submicrometer particle number size distributions for (a) August 15, (b) 29, (c) 13, and (d) 19. The x-axis represents the timeline (local time), the y-axis refers to particle diameters and the colors represent the number of particles, normalized by diameter bins.

On the other hand, Figures 9c and 9d show examples of the influence of cold fronts on particle size distributions. The prefrontal period is typically characterized by stagnant atmospheric conditions that favor the occurrence of high  $\text{PM}_{2.5}$  concentrations. The emergence of the cold front on the morning of August 19<sup>th</sup> resulted in increased ventilation and precipitation (Figure S12), leading to a strong decrease in the particle number concentrations, from  $2.8 \cdot 10^4$  to  $0.5 \cdot 10^4 \text{ cm}^{-3}$ . A decrease in the geometric mean diameter was simultaneously observed, from  $94.7 \pm 2.0$  to  $43.8 \pm 2.0 \text{ nm}$ , indicating that the particle removal by rain and winds was more efficient for accumulation mode particles. Overall, Figure 9 provides examples of sub-daily variability in aerosol properties, which cannot be discerned in the aerosol filter samples.

### 3.6 Chemical and source characteristics of pollution event days

The sampling days were separated between high pollution days (with  $\text{PM}_{2.5}$  above the WHO guideline of  $15 \mu\text{g m}^{-3}$ ) and low pollution days (below). These polluted periods were associated with relatively higher average temperature and lower humidity ( $19^\circ\text{C}$  and 74%, compared to  $15^\circ\text{C}$  and 83%). During these pollution events,  $\text{PM}_{2.5}$  increased by 171%, and carbonaceous species (OM+EC) represented a higher mass fraction (nearly 60%) (Figure 10). In polluted periods, the accumulation processes can lead to the formation of organic coatings on black carbon particles (Monteiro dos Santos et al., 2021). On the other hand,

oxides, other WSI, and sulfate accounted for a higher fraction in non-event days. The unidentified fraction, which can be attributed to unmeasured components such as carbonate or water (Pereira et al., 2017a), was also slightly higher in non-event days. In absolute numbers, all PMF factors increased on event days. However, the BB factor contribution increased significantly on polluted days, rising from 4 to 27% of the PM<sub>2.5</sub> mass.



**Figure 10:** Chemical mass balance for clean (PM<sub>2.5</sub> < 15 µg m<sup>-3</sup>) (a) and polluted days (PM<sub>2.5</sub> ≥ 15 µg m<sup>-3</sup>) (b), and PMF source contributions for both periods (c).

OC, EC, K<sup>+</sup>, NO<sub>3</sub><sup>-</sup>, V, Mn, Fe, Co, Zn, Rb, Cd, Sn, Sb, Ce, Tl, Pb, and Lev presented a significantly higher increase than that observed for PM<sub>2.5</sub> in these polluted periods (Table S6 and Figure S13), suggesting an accumulation of these species during these events ( $p < 0.05$ ). In percentage terms, all OC and EC fractions increased more than PM<sub>2.5</sub> ( $p < 0.05$ ), except OC1, EC3, and EC4. Proportionally, retene increased less than PM<sub>2.5</sub>, suggesting that it may be linked to a local biomass-burning source, such as wood burning in restaurants (Andrade et al., 2017; Kumar et al., 2016). Mannosan followed the same trend. The toxicity indexes BaP-TEQ and BaP-MEQ presented significant increases ( $p < 0.05$ ), caused mainly by the rise in HMW-PAHs due to the accumulation of fossil-fuel and biomass burning emitted aerosols (Vasconcellos et al., 2010). Furthermore, there appears to have been an accumulation of other toxic species, such as potentially toxic elements (PTE) such as Cd, Sb, and Pb, from anthropogenic sources (Pereira et al., 2023b; Thorpe and Harrison, 2008). However, the toxicity unity (TU) did not increase as much (~50%). Phosphate was the only species to decrease, although with  $p > 0.05$ . It is often associated with dust and farming activities (Yuan et al., 2008; Allen et al., 2010). In the present study, the most likely origin was in the soil.


Levoglucosan concentration tripled during the polluted days (423 and 140 ng m<sup>-3</sup>, respectively). Ionic potassium quadrupled these days (330 and 82 ng m<sup>-3</sup>), suggesting a higher impact of crop burning (Chow et al., 2022a). However, levoglucosan represented similar portions of OM on both occasions (nearly 3%). Xylitol and mannosan presented a lower increase on polluted days ( $p < 0.05$ ), suggesting that they are associated with other types of biomass burning. In 2020, it was observed that during smoke plume events in the MASP, PM<sub>2.5</sub> surpassed 25 µg m<sup>-3</sup> at 99% of the air quality monitoring stations. These days, the plumes are mainly associated with sugarcane burning and often with the contribution of wildfires in the Amazon and Pantanal biomes (Souto-Oliveira et al., 2023). In that study, the authors were able to differentiate CO<sub>2</sub> from local



wood burning (4-7%) from that from remote forest wildfires (4-26%). In recent years, the influence of remote forest burning has occurred, likely related to natural seasonal variations and climate change, with lower humidity and precipitation and increased fires (Goss et al., 2020, Abram et al., 2021). The influence of biomass burning in the metropolitan area often occurs above the boundary layer. However, these plumes from remote regions can still influence the concentration and composition of PM<sub>2.5</sub> at the surface (Souto-Oliveira et al., 2023).

Enrichment factors for the elements were above 10 for Cr, Cu, Zn, As, Se, Mo, Ag, Cd, Sn, Sb, Ba, Tl, Pb, Bi, and U (Figure 11), suggesting a strong anthropogenic character. Overall, there was an increase in the enrichment factors of metals in the polluted period, five times higher for Fe and similar for U, suggesting that these are mostly soil-bound in the clean period. Despite the very low concentrations, Se, Sb, and Bi presented the highest EFs (near and above 10,000), indicating a very high anthropogenic character. Selenium was previously attributed to anthropogenic sources such as fossil fuel combustion, burning of garbage, tires and paper, coal combustion, oil, and glass industries (Mehdi et al., 2013), and previously attributed to industrial sources in the eastern part of the MASP (Vieira et al., 2023). Antimony is associated with non-exhaust emissions (brake wear) (Thorpe and Harrison, 2008) and associated with non-ferrous metal industries (Calvo et al., 2013). Bismuth is related to anthropogenic sources, such as fossil fuel combustion, metallurgy, and refuse incineration (Ferrari et al., 2000). Other species presented low EFs, being attributed to geogenic sources: Ti, Fe, and Co (EFs near and below 1). Often associated with catalyst emissions (Kulkarni et al., 2006), La and Ce displayed relatively low enrichment factors (below 5). Some species exhibited geogenic character in the clean periods and a more anthropogenic character in the polluted period, with K, V, Ni, and Rb approaching an EF of 10.

748



749 **Figure 11:** Enrichment factors (EF) for clean, polluted and total period.

### 750 3.7 Ecotoxicity assays

Ecotoxicity tests of aqueous particulate matter extracts were performed with the bacteria *Aliivibrio fischeri*. All samples in this study were classified as toxic, with toxic unity (TU) values ranging from 1.7 to 7.1, averaging 3.7. The highest values were obtained in periods impacted by biomass burning (Figure S14). The TU unit showed moderate correlations with levoglucosan, water-soluble K<sup>+</sup>, and the modeled aqueous K<sup>+</sup> ( $r > 0.5$ ), which may be associated with biomass burning aerosols and water-soluble organic species (not quantified in the study) (Urban et al., 2012). Among the factors obtained by receptor modeling, biomass burning had the highest correlation with TU (Table S5), indicating that water-soluble components trigger the toxicity of particles emitted by this source. The vehicular-exhaust factor exhibited the second highest correlation with TU, although weak. In addition, the TU values presented moderate correlations with elements known to be toxic and associated with vehicular sources such as Pb ( $r \sim 0.5$ ). The correlation was moderate ( $r > 0.5$ ) with OC (higher for fractions OC2, OC3, and OC4, less volatile and more oxidized) and EC (higher for fractions EC1 and EC2). Significant correlations between the bioluminescence inhibition responses and the contributions of biomass burning and traffic to particulate matter concentrations were also reported for Coimbra, Portugal (Alves et al., 2021). As observed in the present study, toxicity was statistically

correlated with OC, EC, anhydrosugars, and elements from exhaust and non-exhaust emissions. Since biomass burning and traffic emissions may elicit acute toxic effects, adopting source-specific preventive and remedial measures is necessary.

#### 4 Summary and conclusions

Fine particulate matter (PM<sub>2.5</sub>) was collected in a 100-day dry period in 2019, covering the period from June to early September, when several pollution events were observed, surpassing the WHO daily limit of 15 µg m<sup>-3</sup> in 75% of the days. Chemical characterization was obtained, including water-soluble ions, elements, carbonaceous species, anhydrosugars, and polycyclic aromatic hydrocarbons. Additionally, the size distribution of particles (SMPS) was monitored simultaneously. PM<sub>2.5</sub> levels were, however, lower than in previous studies. The lower sulfate-to-nitrate ratios suggested a decrease in sulfur oxide levels. A higher contribution of organic matter to particulate matter indicated an increase in the secondary formation of organic species and a reduction of elemental carbon emissions by vehicles. However, a further study should be performed to statistically assess this trend over the last decades. The impact of oxidant atmosphere and higher temperatures on the formation of secondary organic carbon and the effect of industrial emissions on sulfate production require further investigation.

As for biomass burning, typically observed in the dry period, there was a decrease in the contribution of sugarcane straw burning, since there was a change in the K<sup>+</sup>/Lev and Lev/Man ratios and relatively lower potassium levels. However, the long-range transport of plumes from forest fires and agricultural burning in regions north and northwest of MASP remains a significant source of PM<sub>2.5</sub> in this period, as the average concentration of levoglucosan remained high. Other authors have observed a further increase in the influence of bagasse-burning plants and the intrusion of aerosols from forest fires originating in central and northern Brazil. This last phenomenon is related to climate change and has increased in the last decades (longer dry periods). The increase in the Lev/Man ratio with stronger winds suggests the contribution of different types of biomasses. Correlations with other species and the PMF receptor model suggested Rb as a reliable biomass-burning tracer. PAHs remain a concern due to increasing concentrations and equivalent toxicity values exceeding 1 ng m<sup>-3</sup> in half of the samples. However, the levels are lower than those observed in previous studies. The vehicular-related species BbF remained an abundant PAH, suggesting it is a persistent constituent. Furthermore, PAH diagnostic ratios fell within the range observed for vehicular emissions. The increased concentrations of these pollutants are likely related to the lower dispersion in this season. Additionally, increasing V/Ni and La/Ce ratios with east and southeast winds suggested a contribution of aerosols from petrochemical industrial areas, which can occur with meteorological conditions characterized by cold fronts and sea breezes.

The PMF receptor model was applied to assess the PM<sub>2.5</sub> sources and a five-factor solution was obtained (biomass burning, secondary formation, industrial, vehicular+road dust, and local vehicular). A high contribution of biomass burning, associated with north and northwest winds, was observed, reaching one-fourth of the particulate matter. Considering the previous source apportionment studies, sources related to vehicular emissions are still dominant (40% of PM<sub>2.5</sub>). A mixed factor of road dust and vehicular emissions increased throughout the campaign, suggesting a more significant influence of resuspension at the end of the winter. Relatively lower, an industrial contribution was observed, increasing with northeast winds that pass through industrial areas of MASP. The PMF solution showed overlapping contributions in some factors, which may be related to the low temporal resolution of sampling and the fact that emissions from various sources mix before reaching the semi-background receptor site. Enhancing the time resolution in future investigations may help the identification of more sources.

Two particle formation events were identified by SMPS in days with pronounced secondary formation and happened before the arrival of the sea breeze. The sulfate secondary formation was related to humid conditions, as suggested by correlations between the contribution of secondary formation and the aerosol's modeled water content (ISORROPIA). During days of pollution events (PM<sub>2.5</sub> > 15 µg m<sup>-3</sup>), carbonaceous species represented a higher fraction of particulate matter, while sulfate's contribution was reduced. An accumulation of PAHs and toxic species such as Cd, Sb, and Pb on these days represents a health concern. These pollution events were associated with a relative increase in the contribution of the biomass burning

factor, whose emissions, added to pollutants emitted and formed locally, contribute to the degradation of air quality in the dry season. Throughout the sampling campaign, all samples were classified as ecotoxic. The ecotoxicity correlated with the biomass burning factor, highlighting the importance of regulating this source for air quality control. These results indicate that controlling PM<sub>2.5</sub> exceedance events should include regulating emerging biomass burning sources and stricter rules concerning vehicular emissions.

#### **Data availability**

The datasets will be available upon request after publishing.

#### **Author contributions**

GMP performed part of the analyses (OC, EC, elements, and water-soluble ions), the data treatment and led the manuscript writing. MFA managed the project, provided infrastructure, designed the study and revised the manuscript. PCV provided infrastructure, designed the study and revised the manuscript. LYK performed the data treatment. RFB monitored the particle size distribution and contributed with the writing. DMS helped with the OC and EC determination and revised the manuscript. TSS performed the PAHs extraction and characterization. CG characterized the anhydrosugars. ICR performed the ecotoxicity essays and helped with the revisions. CA, NK, LR, PA, and EDF provided infrastructure and revised the manuscript. TN, RMM, and MAY revised the paper.

#### **Competing interests**

The authors declare that they have no conflict of interest.

#### **Acknowledgements**

These results were part of the “Chemical and toxicological SOURCE PROFiling” of particulate matter in urban air project (SOPRO, Grant no. POCI-01-0145-FEDER-029574), supported by the Portuguese Foundation for Science and Technology from the Ministry of Science, Technology, and Higher Education (FCT/MCTES), and by the São Paulo Research Foundation (FAPESP; Grant #2018/07848-9). SOPRO project was also funded by FEDER, through COMPETE2020—Programa Operacional Competitividade e Internacionalização (POCI), and by national funds (OE), through FCT/MCTES. The authors also acknowledge FAPESP for Grants #2016/18438-0 (Metroclima project, for research resources) and #2019/01316-8 (URBESP project, scholarship), and National Research Council for the grants (CNPq 301503/2018-4). The financial support to CESAM by FCT/MCTES (UIDP/50017/2020 + UIDB/50017/2020 + LA/P/0094/2020), through national funds, is also acknowledged. Meteorological data (irradiance, atmospheric pressure, temperature and relative humidity) were kindly provided by the IAG-USP meteorological station and the Climatology and Biogeography Laboratory (Department of Geography - FFLCH - USP).

#### **References**

- Abram, N.J., Henley, B.J., Sen Gupta, A., Lippmann, T.J.R., Clarke, H., Dowdy, A.J., Sharples, J.J., Nolan, R.H., Zhang, T., Wooster, M.J., Wurtzel, J.B., Meissner, K.J., Pitman, A.J., Ukkola, A.M., Murphy, B.P., Tapper, N.J., Boer, M.M.: Connections of climate change and variability to large and extreme forest fires in southeast Australia, *Commun. Earth Environ.*, 2, 1-17, <https://doi.org/10.1038/s43247-020-00065-8>, 2021.
- de Abrantes, R., Vicente de Assunção, J., Pesquero, C. R., Bruns, R. E. and Nóbrega, R. P.: Emission of polycyclic aromatic hydrocarbons from gasohol and ethanol vehicles, *Atmos. Environ.*, 43(3), 648–654, <https://doi.org/10.1016/j.atmosenv.2008.10.014>, 2009.
- Akyüz, M. and Cabuk, H.: Gas-particle partitioning and seasonal variation of polycyclic aromatic hydrocarbons in the atmosphere of Zonguldak, Turkey., *Sci. Total Environ.*, 408(22), 5550–5558, <https://doi.org/10.1016/j.scitotenv.2010.07.063>,

2010.

Allen, A.G., Cardoso, A.A., Wiatr, A.G., Machado, C.M.D., Paterlini, W.C., and Baker, J.: Influence of intensive agriculture on dry deposition of aerosol nutrients, *J. Braz. Chem. Soc.*, 21, 1, <https://doi.org/10.1590/S0103-50532010000100014>, 2010.

Almeida, S. M., Pio, C. A., Freitas, M. C., Reis, M. A. and Trancoso, M. A.: Approaching PM<sub>2.5</sub> and PM<sub>2.5-10</sub> source apportionment by mass balance analysis, principal component analysis and particle size distribution., *Sci. Total Environ.*, 368(2–3), 663–674, <https://doi.org/10.1016/j.scitotenv.2006.03.031>, 2006.

Alves, C., Rienda, I. C., Vicente, A., Vicente, E., Gonçalves, C., Candeias, C., Rocha, F., Lucarelli, F., Pazzi, G., Kováts, N., Hubai, K., Pio, C. and Tchepel, O.: Morphological properties, chemical composition, cancer risks and toxicological potential of airborne particles from traffic and urban background sites, *Atmos. Res.*, 264, 105837, <https://doi.org/10.1016/j.atmosres.2021.105837>, 2021.

Alves, C. A., Evtyugina, M., Vicente, A. M. P., Vicente, E. D., Nunes, T. V., Silva, P. M. A., Duarte, M. A. C., Pio, C. A., Amato, F. and Querol, X.: Chemical profiling of PM<sub>10</sub> from urban road dust., *Sci. Total Environ.*, 634, 41–51, <https://doi.org/10.1016/j.scitotenv.2018.03.338>, 2018.

Alves, C. A., Vicente, A. M. P., Calvo, A. I., Baumgardner, D., Amato, F., Querol, X., Pio, C. and Gustafsson, M.: Physical and chemical properties of non-exhaust particles generated from wear between pavements and tyres, *Atmos. Environ.*, 224, 117252, <https://doi.org/10.1016/j.atmosenv.2019.117252>, 2020.

Amarillo, A. C. and Carreras, H.: Quantifying the influence of meteorological variables on particle-bound PAHs in urban environments, *Atmos. Pollut. Res.*, 7(4), 597–602, <https://doi.org/10.1016/j.apr.2016.02.006>, 2016.

Amato, F., Alastuey, A., Karanasiou, A., Lucarelli, F., Nava, S., Calzolari, G., Severi, M., Becagli, S., Gianelle, V. L., Colombi, C., Alves, C., Custódio, D., Nunes, T., Cerqueira, M., Pio, C., Eleftheriadis, K., Diapouli, E., Reche, C., Minguillón, M. C., Manousakas, M.-I. and Querol, X.: AIRUSE-LIFE+: a harmonized PM speciation and source apportionment in five southern European cities, *Atmospheric Chemistry and Physics*, 16(5), 3289–3309, <https://doi.org/10.5194/acp-16-3289-2016>, 2016.

Andrade, F., Orsini, C. and Maenhaut, W.: Relation between aerosol sources and meteorological parameters for inhalable atmospheric particles in Sao Paulo City, Brazil, *Atmos. Environ.*, 28(14), 2307–2315, [https://doi.org/10.1016/1352-2310\(94\)90484-7](https://doi.org/10.1016/1352-2310(94)90484-7), 1994.

Andrade, M.F., Kumar, P., de Freitas, E. D., Ynoue, R. Y., Martins, J., Martins, L. D., Nogueira, T., Perez-Martinez, P., de Miranda, R. M., Albuquerque, T., Gonçalves, F. L. T., Oyama, B. and Zhang, Y.: Air quality in the megacity of São Paulo: Evolution over the last 30 years and future perspectives, *Atmos. Environ.*, 159, 66–82, <https://doi.org/10.1016/j.atmosenv.2017.03.051>, 2017.

Andrade, M.F., de Miranda, R.M., Fornaro, A., Kerr, A., Oyama, B., de Andre, P.A., Saldiva, P. Vehicle emissions and PM<sub>2.5</sub> mass concentrations in six Brazilian cities. *Air Qual. Atmos. Health*, 5, 79–88, <https://doi.org/10.1007/s11869-010-0104-5>, 2012.

Artaxo, P., Gerab, F., Yamasoe, M. A. and Martins, J. V.: Fine mode aerosol composition at three long-term atmospheric monitoring sites in the Amazon Basin, *J. Geophys. Res.*, 99(D11), 22857, <https://doi.org/10.1029/94JD01023>, 1994.

Aubin, S. and Farant, J. P.: Benzo[b]fluoranthene, a potential alternative to benzo[a]pyrene as an indicator of exposure to airborne PAHs in the vicinity of Söderberg aluminum smelters., *J. Air Waste Manag. Assoc.*, 50(12), 2093–2101, <https://doi.org/10.1080/10473289.2000.10464236>, 2000.

Bhattacharai, H., Saikawa, E., Wan, X., Zhu, H., Ram, K., Gao, S., Kang, S., Zhang, Q., Zhang, Y., Wu, G., Wang, X., Kawamura, K., Fu, P. and Cong, Z.: Levoglucosan as a tracer of biomass burning: Recent progress and perspectives, *Atmos. Res.*, 220, 20–33, <https://doi.org/10.1016/j.atmosres.2019.01.004>, 2019.

Bian, Y. X., Zhao, C. S., Ma, N., Chen, J. and Xu, W. Y.: A study of aerosol liquid water content based on hygroscopicity measurements at high relative humidity in the North China Plain, *Atmospheric Chemistry and Physics*, 14(12), 6417–6426, <https://doi.org/10.5194/acp-14-6417-2014>, 2014.

Bourotte, C., Forti, M. C., Melfi, A. J. and Lucas, Y.: Morphology and solutes content of atmospheric particles in an urban and a natural area of São Paulo state, Brazil, *Water Air Soil Pollut.*, 170(1–4), 301–316, <https://doi.org/10.1007/s11270-005-9001-1>, 2006.

Bourotte, C. L. M., Sánchez-Ccoyllo, O. R., Forti, M. C. and Melfi, A. J.: Chemical composition of atmospheric particulate matter soluble fraction and meteorological variables in São Paulo state, Brazil, *Rev. bras. meteorol.*, 26(3), 419–432, <https://doi.org/10.1590/S0102-77862011000300008>, 2011.

897 Brito, J., Rizzo, L. V., Herckes, P., Vasconcellos, P. C., Caumo, S. E. S., Fornaro, A., Ynoue, R. Y., Artaxo, P. and Andrade,  
898 M. F.: Physical–chemical characterisation of the particulate matter inside two road tunnels in the São Paulo Metropolitan Area,  
899 *Atmospheric Chemistry and Physics*, 13(24), 12199–12213, <https://doi.org/10.5194/acp-13-12199-2013>, 2013.

900 Brito, J., Carbone, S., A. Monteiro dos Santos, D., Dominutti, P., de Oliveira Alves, N., Rizzo, L. V., Artaxo, P.: Disentangling  
901 vehicular emission impact on urban air pollution using ethanol as a tracer. *Scientific Reports*, 8, 10679,  
902 <https://doi.org/10.1038/s41598-018-29138-7>, 2018.

903 Buchholz, R.R., Park, M., Worden, H.M., Tang, W., Edwards, D.P., Gaubert, B., Deeter, M.N., Sullivan, T., Ru, M., Chin, M.,  
904 Levy, R.C., Zheng, B., Magzamen, S.: New seasonal pattern of pollution emerges from changing North American wildfires.  
905 *Nat Commun* 13, 2043. <https://doi.org/10.1038/s41467-022-29623-8>, 2022.

906 Calvo, A. I., Alves, C., Castro, A., Pont, V., Vicente, A. M. and Fraile, R.: Research on aerosol sources and chemical  
907 composition: Past, current and emerging issues, *Atmos. Res.*, 120–121, 1–28, <https://doi.org/10.1016/j.atmosres.2012.09.021>,  
908 2013.

909 Carbone, S., Saarikoski, S., Frey, A., Reyes, F., Reyes, P., Castillo, M., Gramsch, E., Oyola, P., Jayne, J., Worsnop, D. R. and  
910 Hillamo, R.: Chemical characterization of submicron aerosol particles in Santiago de Chile, *Aerosol Air Qual. Res.*, 13(2),  
911 462–473, <https://doi.org/10.4209/aaqr.2012.10.0261>, 2013.

912 Carvalho, J.S., do Nascimento, R.K.S., J Cintra, J.V.F.R.F., da Rosa, N.L.C., Grosseli, G.M., Fadini, P.S., Urban, R.C.: Source  
913 apportionment and health impact assessment of atmospheric particulate matter in the city of São Carlos, Brazil, *Chemosphere*,  
914 326, 138450, <https://doi.org/10.1016/j.chemosphere.2023.138450>, 2023.

915 Caseiro, A., Marr, I.L., Claeys, M., Kasper-Giebl, A., Puxbaum, H., Pio, C.: Determination of saccharides in atmospheric  
916 aerosol using anion-exchange high-performance liquid chromatography and pulsed-amperometric detection, *J. Chromatogr.*  
917 A, 1171, 37–45, <https://doi.org/10.1016/j.chroma.2007.09.038>, 2007.

918 Caseiro, A., Bauer, H., Schmidl, C., Pio, C. A. and Puxbaum, H.: Wood burning impact on PM<sub>10</sub> in three Austrian regions,  
919 *Atmos. Environ.*, 43(13), 2186–2195, <https://doi.org/10.1016/j.atmosenv.2009.01.012>, 2009.

920 Castanho, A. D. A. and Artaxo, P.: Wintertime and summertime São Paulo aerosol source apportionment study, *Atmos.*  
921 *Environ.*, 35(29), 4889–4902, [https://doi.org/10.1016/S1352-2310\(01\)00357-0](https://doi.org/10.1016/S1352-2310(01)00357-0), 2001.

922 Caumo, S., Traub, A., Evans, G. and de Castro Vasconcellos, P.: Health risk assessment in atmosphere near a petrochemical  
923 industrial complex: Measuring oxidative potential and oxidative burden, *Atmos. Pollut. Res.*, 13(7), 101457,  
924 <https://doi.org/10.1016/j.apr.2022.101457>, 2022.

925 Cavalli, F., Viana, M., Yttri, K. E., Genberg, J. and Putaud, J. P.: Toward a standardised thermal-optical protocol for measuring  
926 atmospheric organic and elemental carbon: the EUSAAR protocol, *Atmos. Meas. Tech.*, 3(1), 79–89,  
927 <https://doi.org/10.5194/amt-3-79-2010>, 2010.

928 Cecinato, A.: Polynuclear aromatic hydrocarbons (PAH), benz(a)pyrene (BaPY) and nitrated-PAH (NPAH) in suspended  
929 particulate matter. *Ann Chim* 87, 483–496, 1997.

930 CETESB: Companhia de Tecnologia do Saneamento Ambiental: Relatório de qualidade do ar no Estado de São Paulo 2014,  
931 Report of air quality in the São Paulo State 2014, São Paulo, Brazil, available at: [http://ar.cetesb.sp.gov.br/publicacoes-](http://ar.cetesb.sp.gov.br/publicacoes-relatorios/)  
932 [relatorios/](http://ar.cetesb.sp.gov.br/publicacoes-relatorios/) (last access: 1 February 2024), 2015.

933 CETESB: Companhia de Tecnologia do Saneamento Ambiental: Relatório de qualidade do ar no Estado de São Paulo 2019,  
934 Report of air quality in the São Paulo State 2019, São Paulo, Brazil, available at: [http://ar.cetesb.sp.gov.br/publicacoes-](http://ar.cetesb.sp.gov.br/publicacoes-relatorios/)  
935 [relatorios/](http://ar.cetesb.sp.gov.br/publicacoes-relatorios/), 2020.

936 CETESB: Companhia de Tecnologia do Saneamento Ambiental: Relatório de qualidade do ar no Estado de São Paulo 2022,  
937 Report of air quality in the São Paulo State 2022, São Paulo, Brazil, available at: [http://ar.cetesb.sp.gov.br/publicacoes-](http://ar.cetesb.sp.gov.br/publicacoes-relatorios/)  
938 [relatorios/](http://ar.cetesb.sp.gov.br/publicacoes-relatorios/) (last access: 1 February 2024), 2023.

939 CETESB: Proconve – Programa de Controle da Poluição do Ar por Veículos Automotores - Companhia Ambiental do Estado  
940 de São Paulo, <https://cetesb.sp.gov.br/veicular/proconve/>, last access: 1 April 2022, 2022.

941 Cheong, Y., Kim, T., Ryu, J. et al. Source apportionment of PM<sub>2.5</sub> using DN-PMF in three megacities in South Korea. *Air*  
942 *Qual Atmos Health* 17, 2579–2599. <https://doi.org/10.1007/s11869-024-01584-5>, 2024.

943 Chow, W. S., Liao, K., Huang, X. H. H., Leung, K. F., Lau, A. K. H. and Yu, J. Z.: Measurement report: The 10-year trend of  
944 PM<sub>2.5</sub> major components and source tracers from 2008 to 2017 in an urban site of Hong Kong, China, *Atmospheric Chemistry*  
945 *and Physics*, 22(17), 11557–11577, <https://doi.org/10.5194/acp-22-11557-2022>, 2022a.

946 Chow, W. S., Huang, X. H. H., Leung, K. F., Huang, L., Wu, X. and Yu, J. Z.: Molecular and elemental marker-based source  
 947 apportionment of fine particulate matter at six sites in Hong Kong, China., *Sci. Total Environ.*, 813, 152652,  
 948 <https://doi.org/10.1016/j.scitotenv.2021.152652>, 2022b.

949 Clemente, Á., Yubero, E., Nicolás, J.F., Crespo, J. and Galindo N.: Organic tracers in fine and coarse aerosols at an urban  
 950 Mediterranean site: contribution of biomass burning and biogenic emissions. *Environ Sci Pollut Res* 31, 25216–25226  
 951 <https://doi.org/10.1007/s11356-024-32789-x> , 2024.

952 Cohen, A.J., Brauer, M., Burnett, R., Anderson, H.R., Frostad, J., Estep, K., Balakrishnan, K., Brunekreef, B., Dandona, L.,  
 953 Dandona, R., Feigin, V., Freedman, G., Hubbell, B., Jobling, A., Kan, H., Knibbs, L., Liu, Y., Martin, R., Morawska, L., Pope,  
 954 C.A. 3rd, Shin, H., Straif, K., Shaddick, G., Thomas, M., van Dingenen, R., van Donkelaar, A., Vos, T., Murray, C.J.L.,  
 955 Forouzanfar, M.H.: Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an  
 956 analysis of data from the Global Burden of Diseases Study 2015, *The Lancet*, 389(10082), 1907–1918, 2017. Available at:  
 957 [https://doi.org/10.1016/S0140-6736\(17\)30505-6](https://doi.org/10.1016/S0140-6736(17)30505-6).

958 La Colla, N. S., Botté, S.E., and Marcovecchio, J.E.: Atmospheric particulate pollution in South American megacities.,  
 959 *Environ. Rev.* 29, 3, <https://doi.org/10.1139/er-2020-0105>, 2021.

960 CONAMA: Conselho Nacional de MEio Ambiente - Resolução No. 491, 19 De Novembro de 2018,  
 961 <http://www2.mma.gov.br/port/conama/legiabre.cfm?codlegi=740>, last access: 12 December 2018, 2018.

962 Contini, D., Cesari, D., Conte, M. and Donato, A.: Application of PMF and CMB receptor models for the evaluation of the  
 963 contribution of a large coal-fired power plant to PM<sub>10</sub> concentrations., *Sci. Total Environ.*, 560–561, 131–140,  
 964 <https://doi.org/10.1016/j.scitotenv.2016.04.031>, 2016.

965 Dawidowski, L., Constantin, J.G., Murillo, J.H., Gómez-Marín, M., Nogueira, T., Jiménez, S.B., Díaz-Suárez, V., Victorica,  
 966 F.B., Lichtig, P., Resquin, M.D., Vargas-Rojas, M., Murillo-Hernández, J., Correa, J.A.V., Andrade, M.F., dos Santos, D.M.,  
 967 Maldonado, J.F., Aldape, F., Abreu, L.F., Manousakas, M.I. Carbonaceous fraction in PM<sub>2.5</sub> of six Latin American cities:  
 968 Seasonal variations, sources and secondary organic carbon contribution, *Sci. Total Environ*, 948, 174630,  
 969 <https://doi.org/10.1016/j.scitotenv.2024.174630>, 2024.

970 Decesari, S., Fuzzi, S., Facchini, M. C., Mircea, M., Emblico, L., Cavalli, F., Maenhaut, W., Chi, X., Schkolnik, G., Falkovich,  
 971 A., Rudich, Y., Claeys, M., Pashynska, V., Vas, G., Kourtev, I., Vermeylen, R., Hoffer, A., Andreae, M. O., Tagliavini, E.,  
 972 Moretti, F. and Artaxo, P.: Characterization of the organic composition of aerosols from Rondônia, Brazil, during the LBA-  
 973 SMOCC 2002 experiment and its representation through model compounds, , <https://doi.org/10.5194/acpd-5-5687-2005>, 2005.

974 De La Torre-Roche, R. J., Lee, W.-Y. and Campos-Díaz, S. I.: Soil-borne polycyclic aromatic hydrocarbons in El Paso, Texas:  
 975 analysis of a potential problem in the United States/Mexico border region., *J. Hazard. Mater.*, 163(2–3), 946–958,  
 976 <https://doi.org/10.1016/j.jhazmat.2008.07.089>, 2009.

977 Draxler, R. and Rolph, G.: HYSPLIT (Hybrid Single-Particle Lagrangian Integrated Trajectory) model, NOAA Air Resour.  
 978 Lab., Silver Spring, MD., 2003.

979 Emygdio, A.P.M., Andrade, M.F., Gonçalves, F.L.T., Engling, G., Zanetti, R.H.S., Kumar, P. Biomarkers as indicators of  
 980 fungal biomass in the atmosphere of São Paulo, Brazil. *Sci. Total Environ.*, 612, 809–821,  
 981 <https://doi.org/10.1016/j.scitotenv.2017.08.153>, 2018.

982 Faisal, M., Ali, U., Kumar, A., Hazarika, N., Singh, V. and Kumar, M.: Festive fireworks in Delhi: A major source of elemental  
 983 aerosols established through dispersion normalized PMF in a multiyear study, *Atmos. Environ.*, 323, 120394,  
 984 <https://doi.org/10.1016/j.atmosenv.2024.120394>, 2024.

985 Ferrari, C.P., Hong, S., Van de Velde, K., Boutron, C.F., Rudnev, S.N., Bolshov, M., Chisholm, W., Rosman, K.J.R.: Natural  
 986 and anthropogenic bismuth in Central Greenland, *Atmos. Environ.*, 34, 941–948, [https://doi.org/10.1016/S1352-2310\(99\)00257-5](https://doi.org/10.1016/S1352-2310(99)00257-5), 2000.

988 Ferreira da Silva, M., Vicente de Assunção, J., de Fátima Andrade, M. and Pesquero, C. R.: Characterization of metal and  
 989 trace element contents of particulate matter (PM<sub>10</sub>) emitted by vehicles running on Brazilian fuels-hydrated ethanol and  
 990 gasoline with 22% of anhydrous ethanol., *J. Toxicol. Environ. Health Part A*, 73(13–14), 901–909,  
 991 <https://doi.org/10.1080/15287391003744849>, 2010.

992 Fine, P. M., Cass, G. R. and Simoneit, B. R.: Chemical characterization of fine particle emissions from fireplace combustion  
 993 of woods grown in the northeastern United States., *Environ. Sci. Technol.*, 35(13), 2665–2675,  
 994 <https://doi.org/10.1021/es001466k>, 2001.

995 Fountoukis, C. and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for K<sup>+</sup>–Ca<sup>2+</sup>–

996  $\text{Mg}^{2+}\text{-NH}_4^+\text{-Na}^+\text{-SO}_4^{2-}\text{-NO}_3^-\text{-Cl}^-\text{-H}_2\text{O}$  aerosols, *Atmospheric Chemistry and Physics*, 7(17), 4639–4659,  
997 <https://doi.org/10.5194/acp-7-4639-2007>, 2007.

998 Galvão, E.S., Reis, N.C., Lima, A.T., Stuetz, R.M., Orlando, M.T.A., Santos, J.M.: Use of inorganic and organic markers  
999 associated with their directionality for the apportionment of highly correlated sources of particulate matter, *Sci. Total Environ.*,  
1000 651, 1332–1343, <https://doi.org/10.1016/j.scitotenv.2018.09.263>, 2019. Gioia, S. M. C. L., Babinski, M., Weiss, D. J., Spiro,  
1001 B., Kerr, A. A. F. S., Veríssimo, T. G., Ruiz, I. and Prates, J. C. M.: An isotopic study of atmospheric lead in a megacity after  
1002 phasing out of leaded gasoline, *Atmos. Environ.*, 149, 70–83, <https://doi.org/10.1016/j.atmosenv.2016.10.049>, 2017.

1003 Gómez Peláez, L. M., Santos, J. M., de Almeida Albuquerque, T. T., Reis, N. C., Andreão, W. L. and de Fátima Andrade, M.:  
1004 Air quality status and trends over large cities in South America, *Environmental Science & Policy*, 114, 422–435,  
1005 <https://doi.org/10.1016/j.envsci.2020.09.009>, 2020.

1006 Gonçalves, C., Alves, C., Fernandes, A.P., Monteiro, C., Tarelho, L., Evtugina, M., Pio, C.: Organic compounds in  $\text{PM}_{2.5}$   
1007 emitted from fireplace and woodstove combustion of typical Portuguese wood species, *Atmos. Environ.*, 45, 4533–4545,  
1008 <https://doi.org/10.1016/j.atmosenv.2011.05.071>, 2011.

1009 Gonçalves, C., Rienda, I.C., Pina, N., Gama, C., Nunes, T., Tchepel, O., Alves, C.:  $\text{PM}_{10}$ -Bound Sugars: Chemical  
1010 Composition, Sources and Seasonal Variations. *Atmosphere*, 12, 194, <https://doi.org/10.3390/atmos12020194>, 2021.

1011 Goss, M., Swain, D.L., Abatzoglou, J.T., Sarhadi, A., Kolden, C.A., Williams, A.P., Diffenbaugh, N. S. Climate change is  
1012 increasing the likelihood of extreme autumn wildfire conditions across California *Environ. Res. Lett.*, 15, 9,  
1013 <https://doi.org/10.1088/1748-9326/ab83a7>, 2020.

1014 Graham, B.: Water-soluble organic compounds in biomass burning aerosols over Amazonia. Characterization by NMR and  
1015 GC-MS, *J. Geophys. Res.*, 107(D20), 8047, <https://doi.org/10.1029/2001JD000336>, 2002.

1016 Guo, S., Hu, M., Wang, Z. B., Slanina, J. and Zhao, Y. L.: Size-resolved aerosol water-soluble ionic compositions in the  
1017 summer of Beijing: implication of regional secondary formation, *Atmospheric Chemistry and Physics*, 10(3), 947–959,  
1018 <https://doi.org/10.5194/acp-10-947-2010>, 2010.

1019 Hall, D., Wu, C.-Y., Hsu, Y.-M., Stormer, J., Engling, G., Capeto, K., Wang, J., Brown, S., Li, H.-W. and Yu, K.-M.: PAHs,  
1020 carbonyls, VOCs and  $\text{PM}_{2.5}$  emission factors for pre-harvest burning of Florida sugarcane, *Atmos. Environ.*, 55, 164–172,  
1021 <https://doi.org/10.1016/j.atmosenv.2012.03.034>, 2012.

1022 Han, Y.-S., Eun, D.-M., Lee, G., Gong, S.Y., Youn, J.-S., Enhancement of  $\text{PM}_{2.5}$  source appointment in a large industrial city  
1023 of Korea by applying the elemental carbon tracer method for positive matrix factorization (PMF) model, *Atmospheric Pollution*  
1024 *Research*, Volume 14, 101910, <https://doi.org/10.1016/j.apr.2023.101910>, 2023.

1025 Hetem, I.G., and Andrade, M.F.: Characterization of fine particulate matter emitted from the resuspension of road and  
1026 pavement dust in the Metropolitan Area of São Paulo, Brazil, *Atmos.*, 7, 31, <https://doi.org/10.3390/atmos7030031>, 2016.

1027 Huang, X., Liu, Z., Zhang, J., Wen, T., Ji, D. and Wang, Y.: Seasonal variation and secondary formation of size-segregated  
1028 aerosol water-soluble inorganic ions during pollution episodes in Beijing, *Atmos. Res.*, 168, 70–79,  
1029 <https://doi.org/10.1016/j.atmosres.2015.08.021>, 2016.

1030 IAG-USP. Boletim Climatológico Anual da Estação Meteorológica do IAG/USP. Seção Técnica de Serviços Meteorológicos  
1031 – Instituto de Astronomia, Geofísica e Ciências Atmosféricas da Universidade de São Paulo – vol. 22, 2019, São Paulo,  
1032 IAG/USP, 2023

1033 Ianniello, A., Spataro, F., Esposito, G., Allegrini, I., Hu, M., and Zhu, T.: Chemical characteristics of inorganic ammonium  
1034 salts in  $\text{PM}_{2.5}$  in the atmosphere of Beijing (China), *Atmos. Chem. Phys.*, 11(21), 10803–10822., <https://doi.org/10.5194/acp-11-10803-2011>, 2011.

1036 INPE: INPE (Instituto Nacional de Pesquisas Espaciais) – Portal do Monitoramento de Queimadas, available at:  
1037 <https://queimadas.dgi.inpe.br/queimadas/> (last access: 1 February 2024), 2019.

1038 Jang, H.-N., Seo, Y.-C., Lee, J.-H., Hwang, K.-W., Yoo, J.-I., Sok, C.-H. and Kim, S.-H.: Formation of fine particles enriched  
1039 by V and Ni from heavy oil combustion: Anthropogenic sources and drop-tube furnace experiments, *Atmos. Environ.*, 41(5),  
1040 1053–1063, <https://doi.org/10.1016/j.atmosenv.2006.09.011>, 2007.

1041 Johnson, G. R., Juwono, A. M., Friend, A. J., Cheung, H.-C., Stelcer, E., Cohen, D., Ayoko, G. A. and Morawska, L.: Relating  
1042 urban airborne particle concentrations to shipping using carbon based elemental emission ratios, *Atmos. Environ.*, 95, 525–  
1043 536, <https://doi.org/10.1016/j.atmosenv.2014.07.003>, 2014.

1044 Jung, J., Lee, S., Kim, H., Kim, D., Lee, H. and Oh, S.: Quantitative determination of the biomass-burning contribution to  
1045 atmospheric carbonaceous aerosols in Daejeon, Korea, during the rice-harvest period, *Atmos. Environ.*, 89, 642–650,  
1046 <https://doi.org/10.1016/j.atmosenv.2014.03.010>, 2014.

1047 Justo, E.P.S., Quijano, M.F.C., Beringui, K., Ventura, L.B., Pereira, G.M., Vasconcellos, P.C., Gioda, A.: Assessment of the  
1048 impact of the bus fleet and transportation infrastructure works on the air quality in Rio de Janeiro (Olympic Games 2016). *Air*  
1049 *Qual. Atmos. Health.*, 16, 289–309, <https://doi.org/10.1007/s11869-022-01275-z>, 2023.

1050 Keyte, I.J., Harrison, R.M., and  
1051 Lammel, G.: Chemical reactivity and long-range transport potential of polycyclic aromatic hydrocarbons – a review, *Chem.*  
*Soc. Rev.*, 42, 9333–9391, <https://doi.org/10.1039/C3CS60147A>, 2013.

1052 Khan, Z. Y., Kettler, J., Khwaja, H. A., Naqvi, I. I., Malik, A. and Stone, E. A.: Organic aerosol characterization and source  
1053 identification in Karachi, Pakistan, *Aerosol Air Qual. Res.*, 18(10), 2550–2564, <https://doi.org/10.4209/aaqr.2017.12.0579>,  
1054 2018.

1055 Knorr, W., Dentener, F., Lamarque, J.-F., Jiang, L., Arneth, A., 2017. Wildfire air pollution hazard during the 21st century.  
1056 *Atmos. Chem. Phys.* 17, 9223–9236. <https://doi.org/10.5194/acp-17-9223-2017>, 2017.

1057 Kováts, N., Hubai, K., Sainnokhoi, T.-A., Hoffer, A. and Teke, G.: Ecotoxicity testing of airborne particulate matter-  
1058 comparison of sample preparation techniques for the *Vibrio fischeri* assay., *Environ. Geochem. Health*, 43(11), 4367–4378,  
1059 <https://doi.org/10.1007/s10653-021-00927-w>, 2021.

1060 Kulkarni, P., Chellam, S. and Fraser, M. P.: Lanthanum and lanthanides in atmospheric fine particles and their apportionment  
1061 to refinery and petrochemical operations in Houston, TX, *Atmos. Environ.*, 40(3), 508–520,  
1062 <https://doi.org/10.1016/j.atmosenv.2005.09.063>, 2006.

1063 Kumar, P., de Fatima Andrade, M., Ynoue, R. Y., Fornaro, A., de Freitas, E. D., Martins, J., Martins, L. D., Albuquerque, T.,  
1064 Zhang, Y. and Morawska, L.: New directions: From biofuels to wood stoves: The modern and ancient air quality challenges  
1065 in the megacity of São Paulo, *Atmos. Environ.*, 140, 364–369, <https://doi.org/10.1016/j.atmosenv.2016.05.059>, 2016.

1066 Kumar, S., Aggarwal, S. G., Gupta, P. K. and Kawamura, K.: Investigation of the tracers for plastic-enriched waste burning  
1067 aerosols, *Atmos. Environ.*, 108, 49–58, <https://doi.org/10.1016/j.atmosenv.2015.02.066>, 2015.

1068 Kumar, P., Morawska, L., Birmili, W., Paasonen, P., Hu, M., Kulmala, M., Harrison, R. M., Norford, L., and Britter, R.: Ultra  
1069 fine particles in cities, *Environ. Int.*, 66, 1–10, <https://doi.org/10.1016/j.envint.2014.01.013>, 2014.

1070 Lang, Y.-H., Li, G., Wang, X.-M., and Peng, P.: Combination of Unmix and PMF receptor model to apportion the potential  
1071 sources and contributions of PAHs in wetland soils from Jiaozhou Bay, China, *Mar. Pollut. Bull.*, 90, 129–134,  
1072 <https://doi.org/10.1016/j.marpolbul.2014.11.009>, 2015

1073 Lee, J. D.: *Concise Inorganic Chemistry*, 5th Edn., Wiley, 1070 pp., 1999.

1074 Lima, F.D.M. *Quantificação e caracterização físico-química do material particulado fino (MP<sub>2,5</sub>): queima de biomassa em*  
1075 *fornos de pizzaria na cidade de São Paulo*. Master thesis - University of São Paulo, 2015.

1076 Linak, W. P., Miller, C. A., Wood, J. P., Shinagawa, T., Yoo, J.-I., Santoianni, D. A., King, C. J., Wendt, J. O. L. and Seo, Y.-  
1077 C.: High temperature interactions between residual oil ash and dispersed kaolinite powders, *Aerosol Science and Technology*,  
1078 38(9), 900–913, <https://doi.org/10.1080/027868290500805>, 2004.

1079 Li, Q.-F., Wang-Li, L., Shah, S. B., Jayanty, R. K. M. and Bloomfield, P.: Ammonia concentrations and modeling of inorganic  
1080 particulate matter in the vicinity of an egg production facility in Southeastern USA., *Environ. Sci. Pollut. Res. Int.*, 21(6),  
1081 4675–4685, <https://doi.org/10.1007/s11356-013-2417-z>, 2014.

1082 Li, W., Ge, P., Chen, M., Tang, J., Cao, M., Cui, Y., Hu, K. and Nie, D.: Tracers from Biomass Burning Emissions and  
1083 Identification of Biomass Burning, *Atmosphere*, 12(11), 1401, <https://doi.org/10.3390/atmos12111401>, 2021.

1084 Li, J., Carlson, B.E., Yung, Y.L., Lv, D., Hansen, J., Penner, J.E., Liao, H., Ramaswamy, V., Kahn, R.A., Zhang, P., Dubovik,  
1085 O., Ding, A., Lacis, A.A., Zhang, L., Dong, Y.: Scattering and absorbing aerosols in the climate system. *Nat. Rev. Earth*  
1086 *Environ.* 3, 363–379, <https://doi.org/10.1038/s43017-022-00296-7>, 2022.

1087 Massimi, L., Simonetti, G., Buiarelli, F., Di Filippo, P., Pomata, D., Riccardi, C., Ristorini, M., Astolfi, M.L., Canepari, S.:  
1088 Spatial distribution of levoglucosan and alternative biomass burning tracers in atmospheric aerosols, in an urban and industrial  
1089 hot-spot of Central Italy, *Atmos. Res.*, 239, 104904, <https://doi.org/10.1016/j.atmosres.2020.104904>, 2020.

1090 Marynowski, L., and Simoneit, B.R.T.: Saccharides in atmospheric particulate and sedimentary organic matter: status overview  
1091 and future perspectives, *Chemosphere*, 288, 132376, <https://doi.org/10.1016/j.chemosphere.2021.132376>, 2022.



1092 Mbengue, S., Vodička, P., Komínková, K., Zíková, N., Schwarz, J., Prokeš, R., Suchánková, L., Julaha, K., Ondráček, J.,  
1093 Holoubek, I. and Ždimal, V.: Different approaches to explore the impact of COVID-19 lockdowns on carbonaceous aerosols  
1094 at a European rural background site., *Sci. Total Environ.*, 892, 164527, <https://doi.org/10.1016/j.scitotenv.2023.164527>, 2023.

1095 Mehdi, Y., Hornick, J.-L., Istasse, L. and Dufrasne, I.: Selenium in the environment, metabolism and involvement in body  
1096 functions., *Molecules*, 18(3), 3292–3311, <https://doi.org/10.3390/molecules18033292>, 2013.

1097 Menares, C., Gallardo, L., Kanakidou, M., Seguel, R., Huneeus, N.: Increasing trends (2001–2018) in photochemical activity  
1098 and secondary aerosols in Santiago, Chile. *Tellus B: Chem. Phys. Meteorol.*, 72(1), 1–18.  
1099 <https://doi.org/10.1080/16000889.2020.1821512>, 2020.

1100 Meng, J., Li, Z., Zhou, R., Chen, M., Li, Y., Yi, Y., Ding, Z., Li, H., Yan, L., Hou, Z. and Wang, G.: Enhanced photochemical  
1101 formation of secondary organic aerosols during the COVID-19 lockdown in Northern China., *Sci. Total Environ.*, 758, 143709,  
1102 <https://doi.org/10.1016/j.scitotenv.2020.143709>, 2021.

1103 de Miranda, R.M., Lopes, F., do Rosário, N.É., Yamasoe, M.A., Landulfo, E., and Andrade, M.F.: The relationship between  
1104 aerosol particles chemical composition and optical properties to identify the biomass burning contribution to fine particles  
1105 concentration: a case study for São Paulo city, Brazil., *Environ. Monit. Assess.* 189, 6, [https://doi.org/10.1007/s10661-016-](https://doi.org/10.1007/s10661-016-5659-7)  
1106 5659-7, 2017.

1107 MME, Ministério de Minas e Energia (Ministry of Mines and Energy - Brazil). National Agency of Petroleum, Natural Gas  
1108 and Biofuels - Brazilian Statistical Yearbook of Petroleum, Natural Gas and Biofuels, 2023, available at:  
1109 [https://www.gov.br/anp/pt-br/centrais-de-contendo/publicacoes/anuario-estatistico/arquivos-anuario-estatistico-](https://www.gov.br/anp/pt-br/centrais-de-contendo/publicacoes/anuario-estatistico/arquivos-anuario-estatistico-2023/anuario-2023.pdf/view)  
1110 2023/anuario-2023.pdf/view, (last access: 15 January 2024) 2023.

1111 MME, Ministério de Minas e Energia (Ministry of Mines and Energy - Brazil). National Agency of Petroleum, Natural Gas  
1112 and Biofuels - RESOLUÇÃO ANP Nº 968, DE 30 DE ABRIL DE 2024 (ANP RESOLUTION No. 968, OF APRIL 30, 2024)  
1113 available at: <https://www.in.gov.br/en/web/dou/-/resolucao-anp-n-968-de-30-de-abril-de-2024-557405632>, (last access: 01  
1114 July 2024), 2024.

1115 Monteiro dos Santos, D., Rizzo, L. V., Carbone, S., Schlag, P. and Artaxo, P.: Physical and chemical properties of urban  
1116 aerosols in São Paulo, Brazil: links between composition and size distribution of submicron particles, *Atmospheric Chemistry*  
1117 *and Physics*, 21(11), 8761–8773, <https://doi.org/10.5194/acp-21-8761-2021>, 2021.

1118 Monteiro dos Santos, D. A., Brito, J. F., Godoy, J. M. and Artaxo, P.: Ambient concentrations and insights on organic and  
1119 elemental carbon dynamics in São Paulo, Brazil, *Atmos. Environ.*, 144, 226–233,  
1120 <https://doi.org/10.1016/j.atmosenv.2016.08.081>, 2016.

1121 Nava, S., Calzolari, G., Chiari, M., Giannoni, M., Giardi, F., Becagli, S., Severi, M., Traversi, R., Lucarelli, F.: Source  
1122 Apportionment of PM<sub>2.5</sub> in Florence (Italy) by PMF Analysis of Aerosol Composition Records. *Atmosphere*, 11, 484.  
1123 <https://doi.org/10.3390/atmos11050484>, 2020.

1124 Norris, G., Duvall, R., Brown, S. and Bai, S.: EPA Positive Matrix Factorization (PMF) 5.0 Fundamentals and User Guide,  
1125 2014.

1126 Oduber, F., Calvo, A. I., Castro, A., Alves, C., Blanco-Alegre, C., Fernández-González, D., Barata, J., Calzolari, G., Nava, S.,  
1127 Lucarelli, F., Nunes, T., Rodríguez, A., Vega-Maray, A. M., Valencia-Barrera, R. M. and Fraile, R.: One-year study of airborne  
1128 sugar compounds: Cross-interpretation with other chemical species and meteorological conditions, *Atmos. Res.*, 251, 105417,  
1129 <https://doi.org/10.1016/j.atmosres.2020.105417>, 2021.

1130 de Oliveira Alves, N., Brito, J., Caumo, S., Arana, A., de Souza Hacon, S., Artaxo, P., Hillamo, R., Teinilä, K., Batistuzzo de  
1131 Medeiros, S. R. and de Castro Vasconcellos, P.: Biomass burning in the Amazon region: Aerosol source apportionment and  
1132 associated health risk assessment, *Atmos. Environ.*, 120, 277–285, <https://doi.org/10.1016/j.atmosenv.2015.08.059>, 2015.

1133 de Oliveira Alves, N., Martins Pereira, G., Di Domenico, M., Costanzo, G., Benevenuto, S., de Oliveira Fonoff, A. M., de  
1134 Souza Xavier Costa, N., Ribeiro Júnior, G., Satoru Kajitani, G., Cestari Moreno, N., Fotoran, W., Iannicelli Torres, J., de  
1135 Andrade, J. B., Matera Veras, M., Artaxo, P., Menck, C. F. M., de Castro Vasconcellos, P., Saldiva, P.: Inflammation response,  
1136 oxidative stress and DNA damage caused by urban air pollution exposure increase in the lack of DNA repair XPC protein,  
1137 *Environ. Int.*, 145, 106150, <https://doi.org/10.1016/j.envint.2020.106150>, 2020.

1138 Paatero, P. and Hopke, P. K.: Discarding or downweighting high-noise variables in factor analytic models, *Anal. Chim. Acta*,  
1139 490(1–2), 277–289, [https://doi.org/10.1016/S0003-2670\(02\)01643-4](https://doi.org/10.1016/S0003-2670(02)01643-4), 2003.

1140 Paatero, P. and Tapper, U.: Positive matrix factorization: A non-negative factor model with optimal utilization of error  
1141 estimates of data values, *Environmetrics*, 5(2), 111–126, <https://doi.org/10.1002/env.3170050203>, 1994.

1142 Pacheco, M. T., Parmigiani, M. M. M., de Fatima Andrade, M., Morawska, L. and Kumar, P.: A review of emissions and  
 1143 concentrations of particulate matter in the three major metropolitan areas of Brazil, *J. Transp. Health*, 4, 53–72,  
 1144 <https://doi.org/10.1016/j.jth.2017.01.008>, 2017.

1145 Parra, Y.J., Pereira, G.M., Custódio, D., de Figueiredo, S.B., Alves C., Vasconcellos P.C.: Aerosol characterization in a  
 1146 Central-West site of Brazil: influence of farming activities and toxicity. *Air Qual Atmos Health* 17, 599–620.  
 1147 <https://doi.org/10.1007/s11869-023-01467-1>, 2024.

1148 Pereira, P. A. de P., Lopes, W. A., Carvalho, L. S., da Rocha, G. O., Carvalho, N. De, Loyola, J., Quiterio, S. L., Escalera, V.,  
 1149 Arbilla, G., and de Andrade, J. B.: Atmospheric concentrations and dry deposition fluxes of particulate trace metals in Salvador,  
 1150 Bahia, Brazil, *Atmos. Environ.*, 41, 7837–7850, <https://doi.org/10.1016/j.atmosenv.2007.06.013>, 2007.

1151 Pereira, G. M., De Oliveira Alves, N., Caumo, S. E. S., Soares, S., Teinilä, K., Custódio, D., Hillamo, R., Alves, C. and  
 1152 Vasconcellos, P. C.: Chemical composition of aerosol in São Paulo, Brazil: influence of the transport of pollutants, *Air Qual.*  
 1153 *Atmos. Health*, 10(4), 457–468, <https://doi.org/10.1007/s11869-016-0437-9>, 2017a.

1154 Pereira, G. M., Teinilä, K., Custódio, D., Gomes Santos, A., Xian, H., Hillamo, R., Alves, C. A., Bittencourt de Andrade, J.,  
 1155 Olímpio da Rocha, G., Kumar, P., Balasubramanian, R., Andrade, M. de F. and de Castro Vasconcellos, P.: Particulate  
 1156 pollutants in the Brazilian city of São Paulo: 1-year investigation for the chemical composition and source apportionment,  
 1157 *Atmospheric Chemistry and Physics*, 17(19), 11943–11969, <https://doi.org/10.5194/acp-17-11943-2017>, 2017b.

1158 Pereira, G.M., Oraggio, B., Teinilä, K., Custódio, D., Huang, X., Hillamo, R., Alves, C.A., Balasubramanian, R., Rojas, N.Y.,  
 1159 Sanchez-Ccoyllo, O.R., and Vasconcellos, P.C.: A comparative chemical study of PM<sub>10</sub> in three Latin American cities: Lima,  
 1160 Medellín, and São Paulo. *Air Qual. Atmos. Health*, 12, 1141–1152, <https://doi.org/10.1007/s11869-019-00735-3>, 2019.

1161 Pereira, G. M., da Silva Caumo, S. E., Grandis, A., do Nascimento, E. Q. M., Correia, A. L., de Melo Jorge Barbosa, H.,  
 1162 Marcondes, M. A., Buckeridge, M. S. and de Castro Vasconcellos, P.: Physical and chemical characterization of the 2019  
 1163 “black rain” event in the Metropolitan Area of São Paulo, Brazil, *Atmos. Environ.*, 248, 118229,  
 1164 <https://doi.org/10.1016/j.atmosenv.2021.118229>, 2021.

1165 Pereira, G. M., Kamigauti, L. Y., Nogueira, T., Gavidia-Calderón, M. E., Monteiro Dos Santos, D., Evtugina, M., Alves, C.,  
 1166 Vasconcellos, P. de C., Freitas, E. D. and Andrade, M. de F.: Emission factors for a biofuel impacted fleet in South America’s  
 1167 largest metropolitan area., *Environ. Pollut.*, 331(Pt 2), 121826, <https://doi.org/10.1016/j.envpol.2023.121826>, 2023a.

1168 Pereira, G. M., Nogueira, T., Kamigauti, L. Y., Monteiro Dos Santos, D., Nascimento, E. Q. M., Martins, J. V., Vicente, A.,  
 1169 Artaxo, P., Alves, C., de Castro Vasconcellos, P. and de Fatima Andrade, M.: Particulate matter fingerprints in biofuel  
 1170 impacted tunnels in South America’s largest metropolitan area., *Sci. Total Environ.*, 856(Pt 2), 159006,  
 1171 <https://doi.org/10.1016/j.scitotenv.2022.159006>, 2023b.

1172 Pio, C. A., Legrand, M., Oliveira, T., Afonso, J., Santos, C., Caseiro, A., Fialho, P., Barata, F., Puxbaum, H., Sanchez-Ochoa,  
 1173 A., Kasper-Giebl, A., Gelencsér, A., Preunkert, S. and Schock, M.: Climatology of aerosol composition (organic versus  
 1174 inorganic) at nonurban sites on a west-east transect across Europe, *J. Geophys. Res.*, 112(D23),  
 1175 <https://doi.org/10.1029/2006JD008038>, 2007.

1176 Pöschl, U.: Atmospheric aerosols: composition, transformation, climate and health effects., *Angew. Chem. Int. Ed*, 44(46),  
 1177 7520–7540, <https://doi.org/10.1002/anie.200501122>, 2005.

1178 Ravindra, K., Sokhi, R. and Vangrieken, R.: Atmospheric polycyclic aromatic hydrocarbons: Source attribution, emission  
 1179 factors and regulation, *Atmos. Environ.*, 42(13), 2895–2921, <https://doi.org/10.1016/j.atmosenv.2007.12.010>, 2008.

1180 Ribeiro, F.N.D., de Oliveira, A.P., Soares, J., de Miranda, R.M., Barlage, M., Chen, F.: Effect of sea breeze propagation on  
 1181 the urban boundary layer of the metropolitan region of Sao Paulo, Brazil, *Atmos. Res.* 214, 174–188,  
 1182 <https://doi.org/10.1016/j.atmosres.2018.07.015>, 2018.

1183 da Rocha, G. O., Vasconcellos, P. de C., Ávila, S. G., Souza, D. Z., Reis, E. A. O., Oliveira, P. V., and Sanchez-Ccoyllo, O.:  
 1184 Seasonal distribution of airborne trace elements and water-soluble ions in São Paulo Megacity, Brazil, *J. Braz. Chem. Soc.*,  
 1185 23, 1915–1924, <https://doi.org/10.1590/S0103-50532012005000062>, 2012.

1186 Sánchez-Ccoyllo, O. R. and Andrade, M.F.: The influence of meteorological conditions on the behavior of pollutants  
 1187 concentrations in São Paulo, Brazil., *Environ. Pollut.*, 116(2), 257–263, [https://doi.org/10.1016/s0269-7491\(01\)00129-4](https://doi.org/10.1016/s0269-7491(01)00129-4), 2002.

1188 dos Santos, L.H.M., Kerr, A.A.F.S., Veríssimo, T.G., Andrade, M.F., de Miranda, R.M., Fornaro, A., and Saldiva, P.: Analysis  
 1189 of atmospheric aerosol (PM<sub>2.5</sub>) in Recife city, Brazil. *J. Air Waste Manag. Assoc.*, 64(5), 519–528.  
 1190 <https://doi.org/10.1080/10962247.2013.854282>, 2014.

1191 Santos, T. C. dos, Reboita, M. S. and Carvalho, V. S. B.: Investigação da Relação entre Variáveis Atmosféricas e a  
1192 Concentração de  $MP_{10}$  E  $O_3$  no Estado de São Paulo, Rev. Bras. Meteorol., 33(4), 631–645, [https://doi.org/10.1590/0102-](https://doi.org/10.1590/0102-7786334006)  
1193 7786334006, 2018.

1194 Satsangi, A., Mangal, A., Agarwal, A., Lakhani, A., and Kumari, K. M.: Variation of carbonaceous aerosols and water soluble  
1195 inorganic ions during winter haze in two consecutive years. Atmos. Pollut. Res., 12(3), 242–251,  
1196 <https://doi.org/10.1016/j.apr.2020.12.011>, 2021.

1197 Scaramboni, C. Impacto da queima de biomassa na concentração de compostos policíclicos aromáticos e na toxicidade *in vitro*  
1198 do material particulado atmosférico de Ribeirão Preto-SP. Doctoral thesis - University of São Paulo, 2023.

1199 Scaramboni, C., Farias, C.N., Vasconcellos, P.C., Levi, M., Sadiktsis, I., Pozza, S.A., Umbuzeiro, G.A., Watanabe, T.,  
1200 Rodrigues, P.C.O., Grandis, A., Pagliuso, D., Buckeridge, M.S., Campos, M.L.A.M., Kippler, M. Dreij, K., Galvão, M.F.O.:  
1201 Characterization of cross-continental  $PM_{2.5}$ : Insights into emissions and chemical composition, Atmos. Res. 305, 107423,  
1202 <https://doi.org/10.1016/j.atmosres.2024.107423>, 2024.

1203 Schraufnagel, D. E.: The health effects of ultrafine particles., Exp. Mol. Med., 52(3), 311–317, [https://doi.org/10.1038/s12276-](https://doi.org/10.1038/s12276-020-0403-3)  
1204 020-0403-3, 2020.

1205 SEADE, Fundação Sistema Estadual de Análise de Dados - São Paulo State Data Analysis System Foundation, available at:  
1206 [https://www.seade.gov.br/wp-content/uploads/2021/09/MapaIndustria\\_abril2019.pdf](https://www.seade.gov.br/wp-content/uploads/2021/09/MapaIndustria_abril2019.pdf), (last access: 15 January 2024), 2019.

1207 Serafeim, E., Besis, A., Kouras, A., Farias, C. N., Yera, A. B., Pereira, G. M., Samara, C. and de Castro Vasconcellos, P.:  
1208 Oxidative potential of ambient  $PM_{2.5}$  from São Paulo, Brazil: Variations, associations with chemical components and source  
1209 apportionment, Atmos. Environ., 298, 119593, <https://doi.org/10.1016/j.atmosenv.2023.119593>, 2023.

1210 Shen, H., Tao, S., Wang, R., Wang, B., Shen, G., Li, W., Su, S., Huang, Y., Wang, X., Liu, W., Li, B. and Sun, K.: Global  
1211 time trends in PAH emissions from motor vehicles., Atmos. Environ., 45(12), <https://doi.org/10.1016/j.atmosenv.2011.01.054>,  
1212 2011.

1213 Simoneit, B. R. T.: Biomass burning — a review of organic tracers for smoke from incomplete combustion, Applied  
1214 Geochemistry, 17(3), 129–162, [https://doi.org/10.1016/S0883-2927\(01\)00061-0](https://doi.org/10.1016/S0883-2927(01)00061-0), 2002.

1215 Souto-Oliveira, C. E., Babinski, M., Araújo, D.F., and Andrade, M.F.: Multi-isotopic fingerprints (Pb, Zn, Cu) applied for  
1216 urban aerosol source apportionment and discrimination, Sci. Total Environ., 626, 1350-1366,  
1217 <https://doi.org/10.1016/j.scitotenv.2018.01.192>, 2018.

1218 Souto-Oliveira, C. E., Kamigauti, L. Y., Andrade, M. F., and Babinski, M.: Improving Source Apportionment of Urban Aerosol  
1219 Using Multi-Isotopic Fingerprints (MIF) and Positive Matrix Factorization (PMF): Cross-Validation and New Insights. Front.  
1220 Environ. Sci. 9, 623915, <https://doi.org/10.3389/fenvs.2021.623915>, 2021.

1221 Souto-Oliveira, C. E., Marques, M. T. A., Nogueira, T., Lopes, F. J. S., Medeiros, J. A. G., Medeiros, I. M. M. A., Moreira,  
1222 G. A., da Silva Dias, P. L., Landulfo, E. and Andrade, M. de F.: Impact of extreme wildfires from the Brazilian Forests and  
1223 sugarcane burning on the air quality of the biggest megacity on South America., Sci. Total Environ., 888, 163439,  
1224 <https://doi.org/10.1016/j.scitotenv.2023.163439>, 2023.

1225 Souza, D. Z., Vasconcellos, P. C., Lee, H., Aurela, M., Saarnio, K., Teinilä, K. and Hillamo, R.: Composition of  $PM_{2.5}$  and  
1226  $PM_{10}$  collected at urban sites in Brazil, Aerosol Air Qual. Res., 14(1), 168–176, <https://doi.org/10.4209/aaqr.2013.03.0071>,  
1227 2014.

1228 Srivastava, D., Xu, J., Vu, T. V., Liu, D., Li, L., Fu, P., Hou, S., Moreno Palmerola, N., Shi, Z., and Harrison, R. M.: Insight  
1229 into  $PM_{2.5}$  sources by applying positive matrix factorization (PMF) at urban and rural sites of Beijing, Atmos. Chem. Phys.,  
1230 21, 14703–14724, <https://doi.org/10.5194/acp-21-14703-2021>, 2021.

1231 Sun, J., Shen, Z., Zhang, Y., Zhang, Q., Lei, Y., Huang, Y., Niu, X., Xu, H., Cao, J., Ho, S.S.H. and Li, X.: Characterization  
1232 of  $PM_{2.5}$  source profiles from typical biomass burning of maize straw, wheat straw, wood branch, and their processed products  
1233 (briquette and charcoal) in China, Atmos. Environ. 205, 36-45, <https://doi.org/10.1016/j.atmosenv.2019.02.038>, 2019.

1234 Tang, X., Zhang, X., Ci, Z., Guo, J. and Wang, J.: Speciation of the major inorganic salts in atmospheric aerosols of Beijing,  
1235 China: Measurements and comparison with model, Atmos. Environ., 133, 123–134,  
1236 <https://doi.org/10.1016/j.atmosenv.2016.03.013>, 2016.

1237 Thorpe, A. and Harrison, R. M.: Sources and properties of non-exhaust particulate matter from road traffic: A review, Sci.  
1238 Total Environ., 400(1–3), 270–282, <https://doi.org/10.1016/j.scitotenv.2008.06.007>, 2008.

1239 Tobiszewski, M. and Namieśnik, J.: PAH diagnostic ratios for the identification of pollution emission sources., Environ.

1240 Pollut., 162, 110–119, <https://doi.org/10.1016/j.envpol.2011.10.025>, 2012.

1241 Turpin, B.J., Lim, H.: Species contributions to PM<sub>2.5</sub> mass concentrations: revisiting common assumptions for estimating  
1242 organic mass, *Aerosol Sci. Technol.*, 35, 602–610, 2001.

1243 Urban, R. C., Lima-Souza, M., Caetano-Silva, L., Queiroz, M. E. C., Nogueira, R. F. P., Allen, A. G., Cardoso, A. A., Held,  
1244 G. and Campos, M. L. A. M.: Use of levoglucosan, potassium, and water-soluble organic carbon to characterize the origins of  
1245 biomass-burning aerosols, *Atmos. Environ.*, 61, 562–569, <https://doi.org/10.1016/j.atmosenv.2012.07.082>, 2012.

1246 Urban, R. C., Alves, C. A., Allen, A. G., Cardoso, A. A., Queiroz, M. E. C. and Campos, M. L. A. M.: Sugar markers in aerosol  
1247 particles from an agro-industrial region in Brazil, *Atmos. Environ.*, 90, 106–112,  
1248 <https://doi.org/10.1016/j.atmosenv.2014.03.034>, 2014.

1249 Valente, F. and Laurini, M.: Pre-harvest sugarcane burning: A statistical analysis of the environmental impacts of a regulatory  
1250 change in the energy sector, *Cleaner Engineering and Technology*, 4, 100255, <https://doi.org/10.1016/j.clet.2021.100255>,  
1251 2021.

1252 Vara-Vela, A., Andrade, M. F., Kumar, P., Ynoue, R. Y. and Muñoz, A. G.: Impact of vehicular emissions on the formation  
1253 of fine particles in the Sao Paulo Metropolitan Area: a numerical study with the WRF-Chem model, *Atmospheric Chemistry  
1254 and Physics*, 16(2), 777–797, <https://doi.org/10.5194/acp-16-777-2016>, 2016.

1255 Vasconcellos, P. C., Souza, D. Z., Sanchez-Ccoyllo, O., Bustillos, J. O. V, Lee, H., Santos, F. C., Nascimento, K. H., Araújo,  
1256 M. P., Saarnio, K., Teinilä, K., and Hillamo, R.: Determination of anthropogenic and biogenic compounds on atmospheric  
1257 aerosol collected in urban, biomass burning and forest areas in São Paulo, Brazil, *Sci. Total Environ.*, 408, 5836–5844,  
1258 <https://doi.org/10.1016/j.scitotenv.2010.08.012>, 2010.

1259 Vieira-Filho, M. S., Ito, D. T., Pedrotti, J. J., Coelho, L. H. G. and Fornaro, A.: Gas-phase ammonia and water-soluble ions in  
1260 particulate matter analysis in an urban vehicular tunnel., *Environ. Sci. Pollut. Res. Int.*, 23(19), 19876–19886,  
1261 <https://doi.org/10.1007/s11356-016-7177-0>, 2016a.

1262 Vieira-Filho, M.S., Pedrotti, J. J., and Fornaro, A.: Water-soluble ions species of size-resolved aerosols: Implications for the  
1263 atmospheric acidity in São Paulo megacity, Brazil, *Atmos. Res.*, 181, 281–287,  
1264 <https://doi.org/10.1016/j.atmosres.2016.07.006>, 2016b.

1265 Vieira, E. V. R., do Rosario, N. E., Yamasoe, M. A., Morais, F. G., Martinez, P. J. P., Landulfo, E. and Maura de Miranda, R.:  
1266 Chemical characterization and optical properties of the aerosol in São Paulo, Brazil, *Atmosphere*, 14(9), 1460,  
1267 <https://doi.org/10.3390/atmos14091460>, 2023.

1268 Watson, J.G., Chow, J.C., Houck, J.E.: PM<sub>2.5</sub> chemical source profiles for vehicle exhaust, vegetative burning, geological  
1269 material, and coal burning in Northwestern Colorado during 1995, *Chemosphere*, 43, 1141–1151,  
1270 [https://doi.org/10.1016/S0045-6535\(00\)00171-5](https://doi.org/10.1016/S0045-6535(00)00171-5), 2001.

1271 Wang Q.Q., L.P. Qiao, M. Zhou, S.H. Zhu, S. Griffith, L. Li, Yu. J.Z.: Source Apportionment of PM<sub>2.5</sub> Using Hourly  
1272 Measurements of Elemental Tracers and Major Constituents in an Urban Environment: investigation of Time-Resolution  
1273 Influence *J. Geophys. Res.-Atmos.*, 123, 5284–5300, <https://doi.org/10.1029/2017JD027877>, 2018.

1274 WHO, World Health Organization: WHO global air quality guidelines: Particulate matter (PM<sub>2.5</sub> and PM<sub>10</sub>), ozone, nitrogen  
1275 dioxide, sulfur dioxide and carbon monoxide, Geneva., 2021.

1276 Wu, X., Vu, T. V., Shi, Z., Harrison, R. M., Liu, D. and Cen, K.: Characterization and source apportionment of carbonaceous  
1277 PM<sub>2.5</sub> particles in China - A review, *Atmos. Environ.*, 189, 187–212, <https://doi.org/10.1016/j.atmosenv.2018.06.025>, 2018.

1278 Yamagami, M., Ikemori, F., Nakashima, H., Hisatsune, K. and Osada, K.: Decreasing trend of elemental carbon concentration  
1279 with changes in major sources at Mega city Nagoya, Central Japan, *Atmos. Environ.*, 199, 155–163,  
1280 <https://doi.org/10.1016/j.atmosenv.2018.11.014>, 2019.

1281 Yassaa, N., Meklati, B.Y., Cecinato, A., and Marino, F.: Organic aerosols in urban and waste landfill of Algiers metropolitan  
1282 area: occurrence and sources, *Environ. Sci. Technol.* 35, 306–311, <https://doi.org/10.1021/es991316d>, 2001.

1283 Yunker, M. B., Macdonald, R. W., Vingarzan, R., Mitchell, R. H., Goyette, D. and Sylvestre, S.: PAHs in the Fraser River  
1284 basin: a critical appraisal of PAH ratios as indicators of PAH source and composition, *Org. Geochem.*, 33(4), 489–515,  
1285 [https://doi.org/10.1016/S0146-6380\(02\)00002-5](https://doi.org/10.1016/S0146-6380(02)00002-5), 2002.

1286 Yu, J. Z., Xu, J. and Yang, H.: Charring characteristics of atmospheric organic particulate matter in thermal analysis., *Environ.  
1287 Sci. Technol.*, 36(4), 754–761, <https://doi.org/10.1021/es015540q>, 2002.

1288 Yuan, H., Zhuang, G., Li, J., Wang, Z., Li, J.: Mixing of mineral with pollution aerosols in dust season in Beijing: Revealed

1289 by source apportionment study, *Atmo. Environ.* 42, 2141-2157, ISSN 1352-2310,  
1290 <https://doi.org/10.1016/j.atmosenv.2007.11.048>, 2008.

1291 Zhang, W., Zhang, S., Wan, C., Yue, D., Ye, Y. and Wang, X.: Source diagnostics of polycyclic aromatic hydrocarbons in  
1292 urban road runoff, dust, rain and canopy throughfall., *Environ. Pollut.*, 153(3), 594–601,  
1293 <https://doi.org/10.1016/j.envpol.2007.09.004>, 2008.

1294 Zhu, C.-S., Cao, J.-J., Tsai, C.-J., Shen, Z.-X., Han, Y.-M., Liu, S.-X. and Zhao, Z.-Z.: Comparison and implications of PM<sub>2.5</sub>  
1295 carbon fractions in different environments., *Sci. Total Environ.*, 466–467, 203–209,  
1296 <https://doi.org/10.1016/j.scitotenv.2013.07.029>, 2014.

1297 Zhu, C., Kawamura, K. and Kunwar, B.: Effect of biomass burning over the western North Pacific Rim: wintertime maxima  
1298 of anhydrosugars in ambient aerosols from Okinawa, *Atmospheric Chemistry and Physics*, 15(4), 1959–1973,  
1299 <https://doi.org/10.5194/acp-15-1959-2015>, 2015.