

Relationships between the concentration of particulate organic nitrogen and the inherent optical properties of seawater in oceanic surface waters

5 Alain Fumenia¹, Hubert Loisel¹, Rick A. Reynolds², and Dariusz Stramski²

¹Laboratoire d'Océanologie et de Géosciences, Université du Littoral Côte d'Opale, Université Lille, CNRS, IRD, UMR 8187, LOG, Wimereux, France

10 ²Marine Physical Laboratory, Scripps Institution of Oceanography, University of California San Diego, La Jolla, California 92093-0238, USA

Correspondence to: Alain Fumenia (alain.fumenia@univ-littoral.fr)

Contents of this file

Tables S1 to S5

15 The supporting information provides the best-fit coefficients and their standard deviation of the relationships between PON and IOPs for the open-ocean dataset (Table S1) and the whole dataset (Table S2 and S3). In addition, the supporting information includes the slope (S), intercept (I), and statistical metrics of the relationships between PON derived from IOP-based algorithms and measured PON for the open-ocean dataset (Table S4) and the whole dataset (Table S5).

20

25

Table S1. Best-fit coefficients of the relationships between PON (mg m^{-3}) and IOPs(λ) (m^{-1}) for the open-ocean dataset (OOD). R^2 is the coefficient of determination and N the number of samples.

$\text{PON} = A \text{ IOP}(\lambda)^B$	A	B	R^2	N
$\text{PON} = A b_{\text{bp}}(442)^B$	245739.14 ± 260832.07	1.51 ± 0.10	0.75	70
$\text{PON} = A b_{\text{bp}}(510)^B$	206544.76 ± 196112.79	1.44 ± 0.09	0.80	63
$\text{PON} = A b_{\text{bp}}(555)^B$	105514.11 ± 72139.11	1.31 ± 0.07	0.84	74
$\text{PON} = A a_{\text{p}}(412)^B$	246.10 ± 54.57	0.73 ± 0.03	0.87	71
$\text{PON} = A a_{\text{p}}(442)^B$	229.69 ± 50.62	0.74 ± 0.03	0.86	71
$\text{PON} = A a_{\text{p}}(490)^B$	284.23 ± 61.70	0.71 ± 0.03	0.88	71
$\text{PON} = A a_{\text{p}}(510)^B$	356.71 ± 79.86	0.69 ± 0.03	0.89	71
$\text{PON} = A a_{\text{p}}(555)^B$	468.23 ± 127.04	0.62 ± 0.03	0.86	69
$\text{PON} = A a_{\text{p}}(670)^B$	265.76 ± 66.94	0.56 ± 0.03	0.84	71
$\text{PON} = A a_{\text{ph}}(412)^B$	223.93 ± 57.02	0.65 ± 0.04	0.82	71
$\text{PON} = A a_{\text{ph}}(442)^B$	215.76 ± 54.12	0.68 ± 0.04	0.82	71
$\text{PON} = A a_{\text{ph}}(490)^B$	259.53 ± 65.26	0.66 ± 0.03	0.84	71
$\text{PON} = A a_{\text{ph}}(510)^B$	310.04 ± 81.39	0.62 ± 0.03	0.84	71
$\text{PON} = A a_{\text{ph}}(555)^B$	309.01 ± 92.82	0.49 ± 0.03	0.80	69
$\text{PON} = A a_{\text{ph}}(670)^B$	249.45 ± 66.11	0.53 ± 0.03	0.82	71
$\text{PON} = A a_{\text{d}}(412)^B$	1195.9 ± 601.35	0.85 ± 0.06	0.74	71
$\text{PON} = A a_{\text{d}}(442)^B$	1264.5 ± 693.80	0.82 ± 0.06	0.71	71
$\text{PON} = A a_{\text{d}}(490)^B$	1212.0 ± 756.50	0.75 ± 0.07	0.65	71
$\text{PON} = A a_{\text{d}}(510)^B$	1109.51 ± 711.43	0.72 ± 0.07	0.63	71
$\text{PON} = A a_{\text{d}}(555)^B$	878.97 ± 617.02	0.64 ± 0.07	0.56	71
$\text{PON} = A a_{\text{d}}(670)^B$	251.02 ± 185.26	0.41 ± 0.06	0.38	71

Table S2. Best-fit coefficients of the relationships between PON (mg m⁻³) and $b_{bp}(\lambda)$ (m⁻¹) for the whole dataset (WD). R^2 is the coefficient of determination and N the number of samples.

	a_0	a_1	a_2	a_3	R^2	N
$\log_{10}(\text{PON}) = a_0 + a_1(\log_{10}(b_{bp}(442))) + a_2(\log_{10}(b_{bp}(442)))^2 + a_3(\log_{10}(b_{bp}(442)))^3$	2.21	0.57	0.34	0.10	0.60	289
$\log_{10}(\text{PON}) = a_0 + a_1(\log_{10}(b_{bp}(510))) + a_2(\log_{10}(b_{bp}(510)))^2 + a_3(\log_{10}(b_{bp}(510)))^3$	2.63	1.40	0.74	0.15	0.62	279
$\log_{10}(\text{PON}) = a_0 + a_1(\log_{10}(b_{bp}(555))) + a_2(\log_{10}(b_{bp}(555)))^2 + a_3(\log_{10}(b_{bp}(555)))^3$	2.62	1.32	0.68	0.14	0.63	284

Table S3. Best-fit coefficients of the relationships between PON (mg m⁻³) and the spectral absorption coefficients (m⁻¹) of different seawater particulate constituents for the whole dataset (WD). *R*² is the coefficient of determination and *N* the number of samples.

$\text{PON} = A \text{ IOP}a(\lambda)^B$	<i>A</i>	<i>B</i>	<i>R</i> ²	<i>N</i>
$\text{PON} = A a_p(412)^B$	134.53 ± 8.32	0.62 ± 0.02	0.80	392
$\text{PON} = A a_p(442)^B$	152.42 ± 9.50	0.65 ± 0.02	0.82	392
$\text{PON} = A a_p(490)^B$	221.56 ± 16.02	0.67 ± 0.02	0.82	392
$\text{PON} = A a_p(510)^B$	254.27 ± 19.62	0.64 ± 0.02	0.82	392
$\text{PON} = A a_p(555)^B$	326.86 ± 30.76	0.59 ± 0.02	0.79	389
$\text{PON} = A a_p(670)^B$	278.87 ± 21.81	0.59 ± 0.01	0.83	392
$\text{PON} = A a_{ph}(412)^B$	209.99 ± 16.29	0.63 ± 0.02	0.79	392
$\text{PON} = A a_{ph}(442)^B$	220.73 ± 16.96	0.67 ± 0.02	0.80	392
$\text{PON} = A a_{ph}(490)^B$	323.53 ± 27.80	0.69 ± 0.02	0.82	392
$\text{PON} = A a_{ph}(510)^B$	359.60 ± 31.47	0.64 ± 0.02	0.82	392
$\text{PON} = A a_{ph}(555)^B$	363.03 ± 35.65	0.52 ± 0.01	0.79	389
$\text{PON} = A a_{ph}(670)^B$	273.66 ± 21.93	0.55 ± 0.01	0.82	392
$\text{PON} = A a_d(412)^B$	184.38 ± 16.10	0.55 ± 0.02	0.73	392
$\text{PON} = A a_d(442)^B$	223.29 ± 21.18	0.55 ± 0.02	0.73	392
$\text{PON} = A a_d(490)^B$	303.97 ± 34.60	0.56 ± 0.02	0.70	392
$\text{PON} = A a_d(510)^B$	351.94 ± 42.41	0.57 ± 0.02	0.70	392
$\text{PON} = A a_d(555)^B$	504.51 ± 72.94	0.58 ± 0.02	0.68	392
$\text{PON} = A a_d(670)^B$	787.80 ± 160.15	0.56 ± 0.02	0.58	392

45

50

55 Table S4. Slope (S), intercept (I), and statistical metrics of the relationships between PON derived from IOP-based algorithms and measured PON for the open-ocean dataset (OOD). $RMSD$ is the Root Mean Square Deviation (mg m^{-3}), MdB the Median Bias (mg m^{-3}), MdR the Median Ratio, $MdAPD$ the Median Absolute Percentage Difference (%), $MdSA$ the Median Symmetric Accuracy (%), and N the number of samples.

	S	I	R	$RMSD$	MdB	MdR	$MdAPD$	$MdSA$	N
$b_{bp}(442)$	1.06	-0.07	0.87	6.35	0.34	1.01	28.56	35.41	70
$b_{bp}(510)$	1.04	-0.04	0.90	4.30	0.56	1.07	20.11	22.02	63
$b_{bp}(555)$	1.02	-0.03	0.92	6.17	0.28	1.05	23.68	28.30	74
$a_p(412)$	0.98	0.02	0.93	4.57	0.06	1.01	18.08	19.84	71
$a_p(442)$	0.98	0.02	0.93	4.81	0.31	1.04	18.29	21.50	71
$a_p(490)$	0.98	0.04	0.94	4.46	0.31	1.03	17.99	19.88	71
$a_p(510)$	0.98	0.02	0.94	4.39	0.10	1.03	18.91	19.56	71
$a_p(555)$	0.97	0.03	0.93	4.79	-0.09	0.98	20.21	24.51	69
$a_p(665)$	0.95	0.05	0.92	5.35	-0.10	0.98	18.76	21.18	71
$a_{ph}(412)$	0.96	0.05	0.91	4.63	0.22	1.04	17.64	18.69	71
$a_{ph}(442)$	0.96	0.04	0.91	4.85	0.28	1.04	18.81	18.88	71
$a_{ph}(490)$	0.96	0.03	0.92	4.48	0.22	1.01	18.81	19.56	71
$a_{ph}(510)$	0.96	0.04	0.92	4.32	0.25	1.03	17.42	18.65	71
$a_{ph}(555)$	0.93	0.07	0.90	4.55	0.42	1.04	17.66	17.66	69
$a_{ph}(665)$	0.94	0.06	0.91	5.19	0.20	1.04	19.45	19.80	71
$a_d(412)$	0.97	0.03	0.86	7.14	-0.04	0.99	30.44	34.20	71
$a_d(442)$	0.96	0.04	0.84	7.60	0.22	1.03	31.98	36.90	71
$a_d(490)$	0.94	0.06	0.81	8.35	0.30	1.07	35.83	42.91	71
$a_d(510)$	0.92	0.08	0.79	8.58	0.46	1.07	37.21	40.62	71
$a_d(555)$	0.87	0.13	0.75	9.20	0.63	1.10	38.90	54.67	69
$a_d(665)$	0.63	0.36	0.62	10.83	0.72	1.12	47.09	58.05	71

60

65

Table S5. Slope (S), intercept (I), and statistical metrics of the relationships between PON derived from IOP-based algorithms and measured PON for the whole dataset (WD). $RMSD$ is the Root Mean Square Deviation (mg m^{-3}), MdB the Median Bias (mg m^{-3}), MdR the Median Ratio, $MdAPD$ the Median Absolute Percentage Difference (%), $MdSA$ the Median Symmetric Accuracy (%), and N the number of samples.

	S	I	R	$RMSD$	MdB	MdR	$MdAPD$	$MdSA$	N
$b_{bp}(442)$	0.80	0.28	0.83	29.85	0.23	1.02	36.99	47.87	289
$b_{bp}(510)$	0.80	0.27	0.83	28.91	0.65	1.04	34.50	38.79	279
$b_{bp}(555)$	0.82	0.24	0.85	28.18	0.41	1.03	35.88	43.75	284
$a_p(412)$	0.95	0.08	0.89	27.52	-0.48	0.98	30.93	38.94	392
$a_p(442)$	0.96	0.06	0.90	25.36	0.31	1.02	30.47	36.88	392
$a_p(490)$	0.96	0.06	0.91	25.72	0.25	1.02	29.17	34.87	392
$a_p(510)$	0.96	0.06	0.91	26.09	-0.11	0.99	28.49	33.47	392
$a_p(555)$	0.94	0.09	0.89	27.72	0.24	1.02	30.14	37.01	389
$a_p(665)$	0.95	0.07	0.91	23.27	0.08	1.01	25.28	31.62	392
$a_{ph}(412)$	0.95	0.08	0.89	24.77	0.05	1.00	30.30	37.58	392
$a_{ph}(442)$	0.96	0.07	0.90	23.35	0.37	1.03	29.37	35.70	392
$a_{ph}(490)$	0.96	0.06	0.90	23.77	0.33	1.03	29.29	33.57	392
$a_{ph}(510)$	0.96	0.06	0.91	23.80	0.42	1.03	27.33	33.77	392
$a_{ph}(555)$	0.93	0.11	0.89	26.82	0.61	1.03	28.28	33.94	389
$a_{ph}(665)$	0.94	0.08	0.90	23.76	-0.10	0.99	24.91	30.20	392
$a_d(412)$	0.91	0.13	0.85	32.77	0.41	1.02	33.53	43.15	392
$a_d(442)$	0.91	0.13	0.85	32.56	0.62	1.03	34.10	41.24	392
$a_d(490)$	0.90	0.15	0.84	33.19	0.22	1.01	34.32	41.30	392
$a_d(510)$	0.90	0.14	0.84	33.21	0.23	1.01	34.05	42.22	392
$a_d(555)$	0.90	0.15	0.82	33.88	-0.08	0.99	37.67	44.05	389
$a_d(665)$	0.85	0.23	0.76	35.41	0.46	1.03	39.90	48.77	392