

# Size-resolved process understanding of stratospheric sulfate aerosol following the Pinatubo eruption

Allen Hu<sup>1,\*</sup>, Ziming Ke<sup>2,\*</sup>, Xiaohong Liu<sup>1</sup>, Benjamin Wagman<sup>3</sup>, Hunter Brown<sup>3</sup>, Zheng Lu<sup>1</sup>, Mingxuan Wu<sup>4</sup>, Hailong Wang<sup>4</sup>, Qi Tang<sup>2</sup>, Diana Bull<sup>3</sup>, and Kara Peterson<sup>3</sup>, and Shaocheng Xie<sup>2</sup>

<sup>1</sup>Department of Atmospheric Sciences, Texas A&M, College Station TX, 77843, United States

<sup>2</sup>Lawrence Livermore National Laboratory, Livermore CA, 94550, United States

<sup>3</sup>Sandia National Laboratories, Albuquerque NM, 87185, United States

<sup>4</sup>Pacific Northwest National Laboratory, Richland WA, 99352, United States

\* These authors contributed equally to this work.

Correspondence to: Xiaohong Liu ([xiaohong.liu@tamu.edu](mailto:xiaohong.liu@tamu.edu))

**Abstract.** Stratospheric sulfate aerosol produced by volcanic eruptions plays important roles in atmospheric chemistry and the global radiative balance of the atmosphere. The simulation of stratospheric sulfate concentrations and optical properties is highly dependent on the chemistry scheme and microphysical treatment. In this work, we implemented a sophisticated gas-phase chemistry scheme (full chemistry, FC) and a 5-mode version of the Modal Aerosol Module with Prognostic Stratospheric Aerosol (MAM5-PSA) for the interactive treatment of stratospheric sulfate aerosol in the Department of Energy's Energy Exascale Earth System Model version 2 (E3SMv2) model to better simulate the chemistry-aerosol feedback following the Pinatubo eruption, and to compare it against a simulation using simplified chemistry (SC) and the default 4-mode version of the Modal Aerosol Module (MAM4). MAM5-PSA experiments were found to better capture the stratospheric sulfate burden from the eruption of the volcano to the end of 1992 as compared to the High-resolution Infrared Sounder (HIRS) observations, and the formation of sulfate in MAM5-PSA with FC (with an additional OH replenishment reaction) was significantly faster than in MAM4 with FC. Analyses of microphysical processes indicate that more sulfate aerosol mass was generated in total in FC experiments than in SC experiments. MAM5-PSA performs better than MAM4 in simulation of aerosol optical depth (AOD); AOD anomalies from the MAM5-PSA experiment have better agreement with observations. The simulated largest changes in global mean net radiative flux at the top of the atmosphere following the eruption were about  $-3 \text{ W/m}^2$  in MAM5-PSA experiments and roughly  $-1.5 \text{ W/m}^2$  in MAM4 experiments.

## 1 Introduction

Explosive volcanic eruptions can inject material and gas-phase aerosol precursors into the stratosphere, affecting atmospheric chemistry and radiation and often leading to significant global-scale cooling at the surface (Yang and Schlesinger, 2002; Parker et al., 1996). Many major volcanic eruptions have been associated with dramatic drops in temperature, famine and even the absence of a summer season (Robock, 2000), though Mt. Pinatubo was the first major eruption to be observed with modern instruments. Background stratospheric aerosol during volcanically quiescent periods have optical depths of less than 0.01 and a total stratospheric burden of roughly 0.1 Tg (Sheng et al., 2015; Deshler, 2008), and mainly consist of sulfate formed through the oxidation of sulfur dioxide ( $\text{SO}_2$ ) and carbonyl sulfide (OCS) emitted at the Earth's surface (Sheng et al., 2015). These background aerosols are often overshadowed by sulfate aerosols produced by volcanic eruptions, which are formed from the oxidation of  $\text{SO}_2$  into sulfuric acid and subsequent nucleation and condensation to form sulfate aerosol in the stratosphere (Mills et al., 2016). These particles have a lifespan on the order of years, so it is rare for the stratosphere to be unperturbed by volcanic emissions (Seinfeld and Pandis, 2016). Eventually, these stratospheric aerosols are removed by entering the troposphere through sedimentation processes (Kremser et al., 2016).

The eruption of interest in this study is the Mt. Pinatubo eruption on June 15, 1991. It was one of the most powerful volcanic eruptions in recent decades and occurred after the onset of various modern observation techniques. The amount of  $\text{SO}_2$  injected into the atmosphere was likely in the range of 15 to 18 Tg (Carn, 2022; Neely III, 2016; Aubry et al., 2021). A significant cooling of approximately 0.5 degree at the Earth's surface was observed afterwards due to

the strong scattering of solar shortwave radiation by sulfate aerosol (Parker et al., 1996). In recent years interest in geoengineering as a countermeasure against the effects of climate change has been growing (Crutzen, 2006), with a concept that is analogous to that of a volcanic eruption; sulfate aerosol, formed from SO<sub>2</sub> that is deliberately injected into the stratosphere, could lead to significant temperature decreases at the Earth's surface. It is therefore essential to examine volcanic eruptions for better modeling of future eruptions and better understanding of the effects of geoengineering.

Previous studies have shown that the impacts of volcanic aerosol injections on climate are dependent on the latitude, aerosol type, and season of the injection of volcanic aerosols. Tilmes et al. (2017) examined the aerosol loadings and climate responses associated with hypothetical SO<sub>2</sub> injections at different heights and latitudes using the Whole Atmosphere Community Climate Model (WACCM), finding that continuous high-altitude injections (above the equator, 15°S and 15°N, 30°S and 30°N respectively) at 5 km above the tropopause produced 50% higher aerosol burdens than equivalent injections at 1 km above the tropopause due to a longer sedimentation path, although low altitude injections transported more efficiently from the tropics to the midlatitudes, leading to similar temperature decreases (Tilmes et al., 2017). Injections at 15° N and 15° S were found to transport more efficiently to middle and high latitudes and produce higher global mean AOD than injections at the equator due to the influence from the Brewer-Dobson circulation and the polar vortex (Tilmes et al., 2017). Injections at the 30° and 50° latitudes do not efficiently transport towards the tropics. Laakso et al. (2017) investigated the climatic effects of seasonally varying sulfate injections in the Max Planck Institute's Earth System Model (MPI-ESM). They found that, relative to the experiment with stationary injections above the equator, equivalent injections at the latitudinal position of the maximum solar radiation (for example, at 20-40°N in April and at 20-40°S in October) would lead to 15% stronger radiative forcing outside 20°S-20°N but 27% weaker forcing within this area.

Previous studies regarding stratospheric aerosols utilize different methods to simulate aerosol properties and microphysics. The simplest and most direct approach is to prescribe stratospheric aerosol properties or burdens using climatology derived from observations. For example, Zhuo et al. (2021) produced volcanic forcing using the Easy Volcanic Aerosol (EVA) module, which directly generated stratospheric aerosol optical properties for the given volcanic emissions. More sophisticated methods primarily include bulk schemes, modal schemes, and sectional schemes (Liu, 2023). Bulk schemes are the simplest, where aerosols species are not divided into bins or modes, and properties such as size distribution are prescribed. CNRM-ESM2-1 (an Earth system model developed by the Centre National de Recherches Météorologiques) – one of the models used in Tilmes et al. (2022) for a model intercomparison project for geoengineering – prescribed stratospheric aerosol size distributions with no interactive aerosol microphysics. Gao et al. (2023) used the GFDL Earth System Model version 4.1 (GFDL ESM4.1) to prognostically simulate stratospheric sulfate aerosol concentrations for volcanic eruptions but aerosol size was prescribed, with a sulfate dry effective radius of 0.166 µm, 0.25 µm, 0.4 µm or 1 µm in different sensitivity experiments. Modal schemes typically divide the aerosol population according to the modes conventionally used to describe aerosol size distributions (i.e., Aitken mode, accumulation mode, coarse mode, and nucleation mode in some schemes), with each

mode having its own prescribed standard deviations of log-normal distributions while mass and number concentrations of aerosols in each mode are predicted (Mills et al., 2016; Vioni et al., 2023). Mills et al. (2016) used WACCM's three mode version of the Modal Aerosol Module (MAM3) to simulate the Pinatubo eruption by altering the parameters (e.g., standard deviation) of the coarse mode, but in doing so also influenced the simulation of unrelated coarse mode aerosols such as sea salt and dust, as the aerosols in each mode are treated as internal mixtures rather than being separated by species. Brown et al. (2024) simulated the Pinatubo eruption in E3SMv2 using a four mode version MAM (MAM4), with modifications to the default MAM4 used in E3SMv2. Bin schemes (or sectional schemes) divide aerosols into more categories than modal schemes, providing greater resolution with the drawback of additional computational cost. Vattioni et al. (2019) used a sectional aerosol module which was capable of handling aerosols in 40 different size bins to better represent accumulation-mode  $\text{H}_2\text{SO}_4$  of stratospheric aerosol and their direct injection into the atmosphere as opposed to injections of  $\text{SO}_2$ . Laakso et al. (2022) simulated sulfur injection in the stratosphere using the Sectional Aerosol module for Large Scale Applications (SALSA) which utilized 10 size bins.

Recent studies of volcanic eruptions or stratospheric sulfate related to geoengineering have utilized different approaches with regards to stratospheric chemistry. Zhuo et al. (2021) omitted chemistry of sulfate formation entirely, with  $\text{SO}_2$  emissions directly being converted into aerosol optical properties through empirical calculations. Some studies used a simplified chemistry scheme. In the SALSA1 scheme introduced in Kokkola et al. (2008) and implemented in Bergman et al. (2012), prescribed hydroxyl radical (OH) concentrations were used for stratospheric chemistry. Kleinschmitt et al. (2018) used a prescribed  $\text{SO}_2$  to  $\text{H}_2\text{SO}_4$  conversion rate in the Sectional Stratospheric Sulfate Aerosol (S3A) module to simulate  $\text{SO}_2$  injections. Kleinschmitt et al. (2017) reported a doubling of  $\text{SO}_2$  lifetime in the stratosphere when OH is prognostically calculated for major injections of  $\text{SO}_2$ , such as the Pinatubo eruption, but did not quantitatively compare scenarios with OH prescribed or not prescribed. Studies that used WACCM, such as Mills et al. (2016) and Vioni et al. (2019), considered the key chemical reactions within the stratosphere with a prognostic treatment of OH for the oxidation of  $\text{SO}_2$  to form sulfate aerosol.

This paper builds upon the work of Mills et al. (2016) and Brown et al. (2024) by adding a unique stratospheric sulfate coarse mode and by considering the full chemistry in E3SM, to simulate the burden, AOD, and radiative effect of volcanic aerosols. Our aim is to (1) examine the differences in simulated sulfate aerosol burden after the Pinatubo eruption with and without the changes in MAM; (2) examine the differences in sulfate between using E3SM's default "simple" chemistry and a sophisticated "full" chemistry treatment; and (3) quantify the microphysical processes involved in sulfate chemistry and modal aerosol. It is worth noting that changes in stratospheric ozone following volcanic eruptions play a significant role in both stratospheric chemistry and the impacts on temperature, but it is not the focus for this paper.

The model set-up, including aerosol module, chemistry settings, emissions, nudging, and observational data are described in Section 2. Section 3.1 presents the simulated spatial distribution and total burden of stratospheric sulfate

aerosols following the Pinatubo eruption, and Section 3.2 engages in process analyses of the sulfate aerosols and discusses their growth processes in the different experiments. Section 4 presents the conclusion of our research.

## **2 Methodology**


### **2.1 Model overview**

The model utilized in this study is E3SMv2, which runs roughly twice as fast relative to E3SMv1. The model contains further changes to the dynamical core, the dynamical grid and parameterization column grid, and atmospheric physics and chemistry (Golaz et al., 2022). By default, E3SMv2 treats aerosols through an enhanced version of MAM4 with all aerosol types appearing in the coarse mode (Liu et al., 2016), in which the four lognormal size modes represent the Aitken mode, accumulation mode, coarse mode and primary carbon mode respectively (the primary carbon mode is not directly relevant for our study of stratospheric sulfate aerosol). The model also simulates aerosol microphysical processes relevant to stratospheric sulfate, including condensation, nucleation, coagulation, and water uptake. The experiments are run under the “ne30pg2” resolution, with a grid spacing of approximately 110 km for dynamics and about 165 km for physics. Vertically, it has 72 layers of varying thicknesses, with the top at roughly 60 km altitude (Golaz et al., 2022). The emissions and wet and dry removal of aerosols were described in Wang et al. (2020).

In E3SMv2 MAM4, the coarse mode was primarily intended for coarse sea salt and dust, as well as other aerosol species resuspended from raindrop evaporation, with a geometric standard deviation of 1.8, and stratospheric sulfate aerosol cannot enter the coarse mode through renaming (a process in which aerosol particles that grow larger than a given threshold via condensation and coagulation are transferred from one mode to another, rather than staying in the original mode and instead increasing in number to conserve mass) (Liu et al., 2016). In this study, we utilize a new five-mode version of the Modal Aerosol Module with prognostic stratospheric aerosol (MAM5-PSA) to represent aerosol processes. MAM5-PSA is similar to MAM4 but has a fifth mode added to specifically represent stratospheric coarse-mode sulfate aerosol. Renaming is allowed from the accumulation mode into stratospheric coarse mode in MAM5-PSA. This fifth mode has a smaller geometric standard deviation (1.2) for aerosol size aiming to more accurately reflect the gravitational settling velocity of volcanic sulfate aerosol from the atmosphere (Mills et al., 2016). The stratosphere coarse mode is separate from the original coarse mode in MAM4 so that the changes to better represent volcanic aerosol properties do not affect other coarse mode aerosols such as dust and sea salt. This is an improvement upon Mills et al. (2016) and Brown et al. (2024) in which volcanic sulfate aerosols were not separated from other coarse mode aerosols.

To understand the mechanisms of simulated sulfate aerosol in MAM5-PSA, we carry out process analyses for (a) NUCL (nucleation), (b) COAG (coagulation) and (c) RNMAa (renaming from Aitken mode to accumulation mode), and (d) RNMAsc (renaming from accumulation mode to stratospheric coarse mode) (Fig. 1(a)). Condensation processes for each of the modes have also been included, with the positive values representing the transition of mass

from the gaseous phase to the aerosol. We improve the E3SMv2 to output these tendencies in 2D after the 3D tendencies are vertically integrated.



**Figure 1: (a) Flowchart depicting the aerosol modes and analyzed microphysical processes for the formation of stratospheric sulfate in our experiments. The parameters for the lognormal size distribution of the accumulation mode and stratospheric coarse mode are shown.  $\sigma$  represents the geometric standard deviation of aerosol number concentration (a smaller standard deviation corresponds to a longer atmospheric lifespan), and  $D_g$ ,  $D_{g\_hi}$ , and  $D_{g\_lo}$  represent the geometric dry mean diameter and its upper and lower limits, respectively.  $D_g$  values are not static within the model and the values given can be seen as a reference value. Renaming is a process within E3SM where aerosols that grow sufficiently large can transfer from one mode to another. (b) Chart that depicts the differences in the full chemistry (FC) between MAM4 and MAM5-PSA. In MAM4FC, the oxidation of  $\text{SO}_2$  by OH radicals directly produces  $\text{H}_2\text{SO}_4$  with no byproducts. In MAM5FC, this reaction instead produces  $\text{HO}_2$  and  $\text{SO}_3$ .  $\text{SO}_3$  eventually converts into sulfuric acid, while  $\text{HO}_2$  is crucial in the replenishment of OH concentrations.**

## 2.2 Chemical mechanisms

By default, E3SMv2 uses a simplified set of chemical species and reactions (hereafter referred to as “simple chemistry” or “SC”) that omits many less important species and reactions in order to save computational costs. Tracer species in the model have prognostically evolving concentrations and include  $\text{H}_2\text{O}_2$ , dimethyl sulfide (DMS), sulfuric acid ( $\text{H}_2\text{SO}_4$ ),  $\text{SO}_2$ , secondary organic aerosol precursor gases, and aerosols. Prescribed species such as oxygen, ozone, OH, and  $\text{HO}_2$  have fixed, predetermined concentrations. Photolysis is restricted to  $\text{H}_2\text{O}_2$  (and is simplified to produce no product); beyond that, there are only six other reactions:





To better simulate the stratospheric chemistry following a volcanic eruption – albeit at greater computational cost (roughly twice that of SC) – a “full chemistry” (FC) setup is added through replacing the chemistry preprocessor. It is similar to the Model of Ozone and Related chemical Tracers (MOZART) chemistry scheme (Emmons et al., 2010) used in WACCM (Mills et al. (2016); the original work for E3SM is based on Wu et al. (2022), with some relevant stratospheric gas phase reactions added in. This scheme can handle many organic compounds that are ignored in the simple chemistry, containing 16 photolysis reactions and 41 stratospheric reactions that influence the concentrations of many radical species in the stratosphere. Most importantly, OH radicals (as well as other oxidants) are prognostically calculated rather than using prescribed values. This allows the model to represent the localized depletion of these species due to the injection of large amounts of SO<sub>2</sub>, bottlenecking aerosol formation rates.

With the definitions of SC and FC described above, we consider combinations of MAM4SC, MAM4FC, MAM5SC, and MAM5FC in this study (Table 1). One further change is made for MAM5FC: for the oxidation of SO<sub>2</sub> to form sulfuric acid, the reaction in MAM4FC is  $SO_2 + OH \rightarrow H_2SO_4$ , whereas the reactions in MAM5FC are



HO<sub>2</sub> participates in the replenishment of OH radicals in the stratosphere in MAM5FC, a process that is significantly slower in MAM4FC due to the above reaction not producing HO<sub>2</sub> (Fig. 1(b)).

### 2.3 Experimental set-up

In order to examine the differences between MAM4/MAM5-PSA and simple/full chemistry in E3SMv2’s simulation of Pinatubo, one experiment is performed for each permutation. Coupled Model Intercomparison Project Phase 6 (CMIP6) Diagnosis, Evaluation and Characterization of Klima (DECK) emissions files are used with the exception of volcanic emissions, which consists of VolcanEESM emissions regridded and merged into CMIP6 SO<sub>2</sub> emissions (Brown et al., 2024). Pinatubo eruptions are assumed to have occurred on June 15 1991, with 10 Tg of SO<sub>2</sub> evenly emitted between 18 and 20 km altitude, identical to Mills et al. (2016). Sea ice and sea surface temperatures are prescribed according to CMIP6 DECK datasets. Each simulation starts from January 1, 1991, with more than six months as spin up time. Only the meridional and zonal winds were nudged to MERRA-2 reanalysis leaving the temperature and humidity to dynamically respond.

**Table 1: List of numerical simulations.**

Experiment name	Aerosol module	Chemistry
-----------------	----------------	-----------

MAM4SC	MAM4	Simple chemistry
MAM4FC	MAM4	Full chemistry
MAM5SC	MAM5-PSA	Simple chemistry
MAM5FC	MAM5-PSA	Full chemistry (with alterations)

## 2.4 Observational data


AVHRR monthly observations of AOD for the years of 1989-1993 were used for model evaluation. Observed AOD values at 600 nm were downloaded from <https://www.ncei.noaa.gov/data/avhrr-aerosol-optical-thickness/access> (last access: June 5 2024) (Zhao, 2022). The wavelength of observation is different from E3SMv2 which outputs AOD at 550 nm wavelength, but the discrepancy caused by the difference in wavelength can be considered negligible. The website provides retrieved monthly AOD with a spatial resolution of  $0.1^\circ \times 0.1^\circ$ , calculated using AVHRR's daily orbital observation of top-of-atmosphere reflectance over the oceans. E3SM output AOD that is used to compare with AVHRR results is masked according to the pixels that were successfully observed by AVHRR and regridded to be the same resolution. The Global Space-based Stratospheric Aerosol Climatology (GloSSAC) is a global and gap-free data set of zonally averaged optical properties of stratospheric aerosols (focused on aerosol extinction coefficient at 525 and 1020 nm) from 1976–2018 (Thomason et al., 2018). Here, we use the updated version 2 from Kovilakam et al. (2020). Sulfate burdens derived from HIRS observations according to Baran and Foot (1994) were used for model comparison. Data from the balloon-borne University of Wyoming optical particle counters (WOPC) (Deshler et al., 1993; Deshler et al., 2003) is converted from the original particle number size distributions to particle volume size distributions for comparisons with model outputs.

## 3 Results

### 3.1 Simulated concentrations of stratospheric sulfate

Figure 2 presents the simulated horizontal distributions of sulfate aerosol concentrations at 53 hPa at 3, 9 and 15 days after the eruption of Pinatubo respectively for the four experiments. The overall distribution pattern is similar between all four experiments, with sulfate aerosol spreading westwards from Mt. Pinatubo in the Philippines following the eruption because of the Quasi-Biennial Oscillation. After 3 days, the sulfate reaches India; after 9 days, it passes over Africa and reaches the eastern coast of the Atlantic Ocean; after 15 days, it reaches Central America. Within this time period, northward and southward transport is both relatively weak, and the sulfate mostly remains between  $30^\circ\text{S}$  and  $30^\circ\text{N}$  (excluding background sulfates present prior to the eruption).






**Figure 2: Simulated distributions of sulfate aerosol concentrations (kg/kg) at 53 hPa for days 3, 9 and 15 after the eruption of Pinatubo, respectively, for experiments MAM4SC (first row), MAM4FC (second row), MAM5SC (third row), and MAM5FC (bottom row).**

For both MAM4 and MAM5-PSA simulations, sulfate concentrations are generally higher in SC than in FC. In these experiments, sulfate formation is not hindered by a lack of OH because OH concentration is prescribed and not depleted. On June 18, 3 days after the eruption, there is a substantial difference in sulfate concentrations near the origin of the plume. In MAM4SC and MAM5SC, the concentrations reach above  $10^{-6}$  kg/kg. There is no replenishment mechanism for OH radical in MAM4FC, hindering sulfate formation, so the peak value in MAM4FC is below  $10^{-8}$  kg/kg. In MAM5FC, OH can be depleted but a replenishment mechanism is also present, and the highest concentration is between  $10^{-8}$  kg/kg and  $10^{-7}$  kg/kg. These differences between SC and FC weaken over time as the plume spreads and encounters more OH to participate in the oxidation of  $\text{SO}_2$  in FC. On June 30, the peak concentration in MAM4FC is between  $10^{-8}$  and  $10^{-7}$  kg/kg, while in the other three experiments it is between  $10^{-7}$  and  $10^{-6}$  kg/kg.

Figure 3 depicts the vertical profiles of sulfate concentrations for the Aitken and accumulation modes (or the sum of accumulation and stratospheric coarse modes for MAM5-PSA, as MAM4 cannot rename accumulation mode into coarse mode) in each of the four experiments. In all four experiments, the initial sulfate burden before July consists of mostly Aitken mode aerosols with peak concentrations on the order of  $10^{-9}$  kg/kg. Sulfate aerosol either leaves the Aitken mode or is removed from the stratosphere within the span of several weeks in SC experiments, while sulfate persists past October in FC runs. Soon after July 1991, accumulation mode sulfate in MAM4 and accumulation mode plus stratospheric coarse mode sulfate in MAM5-PSA become dominant until they diminish back to background levels, with peak concentrations on the scale of  $10^{-7}$  kg/kg.



**Figure 3: Vertical profiles of sulfate concentrations in the four experiments. The plots are divided by modes (Aitken mode in top panel with units of  $10^{-9}$  kg/kg, accumulation mode or the sum of accumulation mode and stratospheric coarse mode for MAM5-PSA in bottom panel with units of  $10^{-7}$  kg/kg). The plots for the Aitken mode aerosols span the year 1991 as the lifetime of the aerosols are within several months. The plots for the accumulation mode and stratospheric coarse mode aerosols span the years 1991 to 1993. The concentrations are averaged across longitude and latitude within the 30°S to 30°N band to remove interfering signals. Note that the top and bottom panels span different time periods.**


There are significant differences in concentrations and locations of sulfate between SC and FC. In the SC experiments, both the earlier Aitken mode sulfate and the later accumulation and stratospheric coarse mode sulfate remain at a higher altitude than the corresponding FC experiments. The highest concentrations occur between 10 and 30 hPa altitudes for SC, and between 30 and 70 hPa altitudes for FC, which may be explained by the higher  $\text{SO}_2$  concentrations at 30-70 hPa in FC relative to SC, the smaller negative difference in OH concentrations at 30-70 hPa in FC relative to SC, and the higher specific humidity at 30-70 hPa in FC compared to SC (Supplementary Figs. 1-4). For this same reason, the Aitken mode in FC generally has higher concentrations than SC after the eruption (up to  $2.6 \times 10^{-9}$  kg/kg in FC compared to less than  $1.0 \times 10^{-9}$  kg/kg in SC) in both the MAM4 and MAM5-PSA simulations. For the accumulation

and stratospheric coarse modes, SC generally has higher concentration than FC several weeks after the eruption (up to  $1.0 \times 10^{-7}$  kg/kg and  $1.6 \times 10^{-7}$  kg/kg for MAM4SC and MAM5SC respectively compared to less than  $1.0 \times 10^{-7}$  kg/kg in FC).

The main differences between MAM4 and MAM5-PSA are: (1) the inability of sulfate to progress past the accumulation mode in MAM4, and (2) the relatively short lifespan of the accumulation mode aerosols in MAM4 compared to the stratospheric coarse mode aerosols in MAM5-PSA. The shorter lifespan is a result of the reduced geometric standard deviation of the stratospheric coarse mode (1.2, versus 1.6 for MAM4's accumulation mode (as described in Fig. 1). A smaller geometric standard deviation for stratospheric coarse mode in MAM5-PSA means that there are fewer super-coarse aerosols within the population. As super-coarse aerosols sediment more quickly, a smaller geometric standard deviation (1.2) leads to a longer lifespan for sulfate. In MAM4, constraining sulfate to the accumulation mode leads to aerosols, as well as the larger geometric standard deviation leads to a significant number of aerosols being super-coarse and quickly removed. Therefore, sulfate concentrations return to background level after roughly one year in MAM4 (as was also seen in E3SMv2-PA in Brown et al. (2024)) but this takes multiple years in MAM5-PSA.

### 3.2 Simulated burdens of stratospheric sulfate

Figure 4(a) shows the time series of the simulated stratospheric burden of sulfate between  $80^\circ$  S and  $80^\circ$  N, as well as the HIRS monthly observational data for stratospheric sulfate burden for the same latitude bands taken from Baran and Foot (1994). Following the eruption of Mt. Pinatubo on June 15, 1991, sulfate burdens in all four model experiments rise rapidly, reach their peak value in November 1991, then begin to diminish over time. MAM4SC, MAM5SC and MAM5FC have about the same peak sulfate burden of about 5.5 Tg, occurring at roughly the same time in November 1991. The rate of increase for the sulfate burden, as well as the eventual peak sulfate burden, is the lowest in MAM4FC (about 4.8 Tg sulfur on November 30, 1991) due to OH concentrations in the atmosphere being depleted by the extremely large amount of  $\text{SO}_2$  produced by the eruption and being unable to replenish quickly. Starting in January 1992, the stratospheric burdens in MAM4 drop off more quickly compared to MAM5-PSA due to the reduced geometric standard deviation in MAM5-PSA's stratospheric coarse mode, so MAM4's burden drops off excessively quickly compared to the HIRS observation whereas MAM5FC agrees closely with HIRS until the end of 1992. The reduced geometric standard deviation corresponds to a smaller proportion of extremely large aerosols that are removed more quickly from the stratosphere, due to the settling velocity being roughly proportional to the second power of the aerosol diameter (Seinfeld and Pandis, 2016). The burden in MAM5-PSA continues to decrease almost linearly in the following years but does not return to background levels until 1995 (not pictured). For both MAM4 and MAM5-PSA, the full chemistry experiments have stratospheric sulfate burdens that decline more quickly after reaching peak values in November 1991 than the corresponding simple chemistry experiment.




**Figure 4:** (a) Comparisons of simulated stratospheric sulfate burdens (Tg S) between 80° S and 80° N from the four experiments with HIRS observations for years of 1991-1995. Black line depicts the estimates made from HIRS data in Baran and Foot (1994), with the black bars representing the margin of error. (b), (c), and (d) are similar comparisons for the Aitken mode, accumulation mode and stratospheric coarse mode respectively (renaming from accumulation mode into the coarse mode does not exist in MAM4).

Figures 4(b), 4(c) and 4(d) respectively show the variations in stratospheric sulfate burdens of Aitken mode, accumulation mode and stratospheric coarse mode. The results used in these plots are three-day averages on every third day. In MAM4, most of the sulfate burden exists in the form of the accumulation mode, as renaming into the coarse mode is not available. Around November 15, 1991 (the time of peak stratospheric sulfate burden), Aitken mode and accumulation mode contribute, respectively, 0.1% and 99.9% (0.3% and 99.7%) to total burden in MAM4SC (MAM4FC). For MAM5-PSA, the accumulation mode peaks briefly, but is soon almost entirely converted into the stratospheric coarse mode via renaming. Similar to the accumulation mode for MAM4, in both MAM5FC and MAM5SC the stratospheric coarse mode makes up more than 98% of the stratospheric sulfate burden on November 15, 1991, with about two-thirds of the remainder being in the accumulation mode and one-third in the Aitken mode.

It is worth noting that the peak values for the different aerosol modes do not occur concurrently. For the Aitken mode, all four experiments peak for the first time in late June and July (0.04 Tg S on June 30 in MAM4SC, 0.15 Tg S on July 30 in MAM4FC, 0.03 Tg S on June 30 in MAM5SC, and 0.1 Tg S on July 12 in MAM5FC). The results here agree with those in Fig. 2, where aerosol formation is faster in SC compared to FC, and MAM5FC is faster than MAM4FC due to replenishment of OH radicals. Several months after the Pinatubo eruption, there is again an increase in Aitken mode burden due to the eruption of Mount Hudson between August and October. The Mount Spurr and Lascar eruptions of 1992 and 1993 respectively are also visible here.

For the accumulation mode, the peak burden of 5.5 Tg S (4.8 Tg S) occurs on November 15 (November 30) in MAM4SC (MAM4FC). Much like for Aitken mode, the peak value for MAM4FC occurs later than MAM4SC as a result of the slower start in terms of sulfate formation. For MAM5-PSA, accumulation mode stratospheric sulfates are mostly transitory in nature, quickly renaming into the stratospheric coarse mode. The accumulation mode of MAM5SC and MAM5FC has small peak values of 0.4-0.8 Tg S during July 9-24. The stratospheric coarse mode burdens peak around November 15 for both MAM5-PSA experiments at around 5.5 Tg S.

Figure 5 shows the stratospheric burdens of sulfate for different modes and latitudinal regions of 30 °N-80 °N, 30 °S-30 °N, and 80 °S-30 °S. As for Fig. 4, for all regions the vast majority of the stratospheric sulfate burden exists in the accumulation mode for MAM4 and in the stratospheric coarse mode for MAM5-PSA. It should be noted that the total stratospheric sulfate burden exhibits different changes with time for MAM5-PSA. Between 30 °S and 30 °N, the total stratospheric burden in all experiments declines the fastest compared to the other two regions due to both gravitational settling of sulfate and its poleward transport due to the Brewer-Dobson circulation. Between 30 °N-80 °N, the stratospheric sulfate burden peaks in February and March of 1992, as opposed to October 1991 in 30 °S-30 °N, due to the time required for aerosol transport northward via the Brewer-Dobson circulation. The stratospheric sulfate burden has a second peak in January 1993 at 30 °S-30 °N and 30 °S-80 °S corresponding to the Lascar eruption, though this does not counteract the overall decline over time. Between 30 °S-80 °S, the stratospheric sulfate burden peaks in November 1991 for both MAM5-PSA experiments, and afterwards remains relatively steady. For MAM5-PSA the stratospheric sulfate burden appears to oscillate annually in the southern hemisphere, with high values around October-November and low values around April of 1992 and 1993.




**Figure 5: Simulated stratospheric sulfate burden (in Tg S) divided by mode and latitudinal region. The vertical lines represent the Pinatubo, Cerro Hudson, Spurr and Lascar eruptions respectively.**

It can also be seen in Fig. 5 that the relative latitudinal contribution to the total stratospheric sulfate burden changes over time. The burden between 30 °S and 30 °N constitutes about two-thirds or more of the stratospheric sulfate burden in the months after the Pinatubo eruption. Starting from April of 1992, the stratospheric burden of 1.7-2.1 Tg S between 30 °N-80 °N is similar in magnitude to the stratospheric burden between 30 °S and 30 °N in all four experiments. Starting from 1993, the stratospheric burden south of 30 °S in MAM5-PSA begins to make up a much larger portion of the total between 80 °S and 80 °N (about 0.8 Tg S out of the total of about 4.1 Tg S), due to the stratospheric coarse mode burdens not decreasing. This accounts for the slower decline of the total stratospheric burden in MAM5-PSA as seen in Fig. 4(a).

Since the independent stratospheric coarse mode is newly added in this study, Figure 6 depicts the distribution of MAM5-PSA aerosol volume particle size distribution, as compared with WOPC observations (Deshler et al., 1993; Deshler et al., 2003). MAM4 has been omitted as it does not include coarse mode sulfate. It can be seen that with the exception of December 30, 1991, MAM5FC generally better matches WOPC observations in coarse mode mass than MAM5SC. However, MAM5FC coarse mode size is biased towards being smaller than the observations. The

relatively smaller coarse mode aerosols may also help to explain the excessively long lifespan later seen past 1992 in Figs. 4 and 5.



**Figure 6: Comparisons of stratospheric aerosol size distributions from MAM5SC and MAM5FC with observations from WOPC for 1991–1993. WOPC launches are samples taken from the 18 km measurements and matched to the nearest model height and grid cell over Laramie, Wyoming (41.3° N, 105° W). The vertical dashed lines with longer dashes represent the accumulation mode’s geometric mean diameter, while the vertical lines with shorter dashes represent the coarse mode’s geometric mean diameter. The triangles represent the effective diameters calculated from the size distribution.**


### 3.3 Process analyses

In order to better understand the differences between the experiments and how the aerosols transition between the different modes, the tendencies (i.e., the rate of mass change into or out of a certain mode) between 80 °S and 80 °N are plotted in Fig. 7. Here it is assumed that these rates apply only to sulfate and other aerosol contributions are negligible within the stratosphere. The tendencies are integrated over the three dimensions of: all longitudes, the above-mentioned latitude range, and vertically above the tropopause.

The magnitude of the tendency for the formation of Aitken mode aerosols through the nucleation of gaseous  $\text{H}_2\text{SO}_4$  (NUCL) is generally higher in the FC experiments compared to the SC experiments, which explains the higher Aitken mode sulfate concentrations in FC than in SC as shown in Fig. 3. For MAM4FC, the peak value occurred on July 24 and it does not return to background levels until early December 1991. MAM5FC peaks on July 3 and returns to background levels after the middle of August 1991. In both MAM4SC and MAM5SC, the rate peaks on June 24 and returns to background levels by the start of August 1991. Due to the higher OH concentrations in the SC experiments,

the nucleation process begins and ends earlier in the SC experiments compared to FC. Considering the cumulative mass changes depicted in Fig. 7(b), the curves for the SC experiments flatten out earlier than those for FC (i.e., the microphysical process in SC is mostly concluded). For MAM4SC (MAM5SC) the curve flattens out by July 15, 1991 (July 6, 1991), with later, less significant processes contributing to a 39.4% (63.8%) increase by the end of 1993. For MAM4FC (MAM5FC) the curve flattens out by October 19, 1991 (September 16, 1991), with approximately a 9.3% (10.8%) increase by the end of 1993. Note that there was some interference from the Mount Hudson eruption in August 1991, as well as Spurr and Lascar. The total magnitude of the nucleation process is also different between the experiments, with MAM4SC, MAM4FC, MAM5SC and MAM5FC reaching roughly 0.3, 1.3, 0.1 and 0.6 Tg respectively by the end of 1993.





**Figure 7:** The relevant tendencies for each mode in each experiment in the stratosphere between 80°S and 80°N. See Figure 1(a) for description of processes. Positive values represent gained mass from the associated microphysical process from the perspective of the aerosol mode in question. NUCI represents the aerosol mass gain for Aitken mode due to nucleation (i.e., aerosol formation), COAG represents Aitken mode mass loss due to coagulation into the accumulation mode, RNMa represents Aitken mode mass loss due to renaming into the accumulation mode, RNMa represents the gain in mass for stratospheric coarse mode in MAM5-PSA due to renaming from the accumulation mode. The left column plots represent the tendency values over time, while the right column represents the cumulative mass change due to the associated microphysical process (i.e. tendency integrated over time). The bottom three rows represent the condensation tendencies for each mode, with the positive value representing mass transition from gaseous phase to the aerosol. The vertical lines represent the Pinatubo, Cerro Hudson, Spurr and Lascar eruptions respectively.

Figs. 7(c)(d) show the aerosol mass loss rates (negative values) of the Aitken mode due to coagulation into the accumulation mode (COAG). Generally, the peak values occur some time (0 to 9 days) after that of the nucleation process, as coagulation becomes stronger when Aitken mode aerosol concentrations are high and accumulation mode aerosols become available. This happens sooner for the SC experiments than FC (June 27 for MAM4SC vs. August 2 for MAM4FC, June 24 for MAM5SC vs. July 6 for MAM5FC). For the integrated values (Fig. 7(d)), much like in the nucleation process, MAM4 has stronger coagulation than MAM5-PSA, and FC has stronger coagulation than SC. The curves also flatten slightly later than for the nucleation process, about 0 to 9 days after the corresponding curve for nucleation as described above. The total magnitude from largest to smallest is still MAM4FC, MAM5FC, MAM4SC and MAM5SC, at 4 Tg, 2.2 Tg, 1.2 Tg and 0.4 Tg, respectively.

Figs. 7(e)(f) show the aerosol mass loss rate of the Aitken mode due to renaming into the accumulation mode (RNMaa). This occurs concurrently with the coagulation process, though the peak occurs slightly earlier as it only requires the presence of Aitken mode aerosols and not accumulation mode aerosols. MAM4SC, MAM4FC, MAM5SC, and MAM5FC peak on June 18, July 21, June 27, and July 12, respectively. Of particular note is that the response to the Cerro Hudson eruption appears to be much stronger here than in the other processes, and that the difference between MAM4 and MAM5-PSA is small for both FC and SC.

Figs. 7(g)(h) represent the aerosol mass gain rate of the stratospheric coarse mode due to renaming from the accumulation mode (RNMasc). This step occurred after most of the above processes had passed their peaks and accumulation mode aerosols had grown considerably in size. MAM5SC peaked on July 9 while MAM5FC peaked on July 24. Unlike the other processes, in both MAM5SC and MAM5FC the curve in Fig. 7(h) does not flatten, and the renaming process does not completely conclude at the end of January 1993 (explained above in Fig. 5 discussion).


Figs. 7(i)-(n) represent the condensation processes for each of the three modes. In general, this accounts for the changes in mass within each mode, though the mass of each mode is still not strictly conserved due to exchanges between the troposphere and stratosphere. It can be seen that FC has stronger condensation tendencies than SC in the Aitken mode due to the larger amount of Aitken mode aerosols seen in Figure 5. Accumulation mode tendencies are stronger for MAM4 than MAM5-PSA likewise due to very small accumulation mode concentrations in MAM5-PSA. MAM5SC has a much stronger condensation tendency than MAM5FC for the stratospheric coarse mode, which appears to counterbalance the differences in RNMasc, ultimately leading to relatively close stratospheric coarse mode burdens between the two experiments. Similarly, MAM4SC has stronger condensation tendency for the accumulation mode than MAM4FC throughout the span of the experiment. This suggests that in SC,  $\text{H}_2\text{SO}_4$  within the stratosphere tends to directly contribute to accumulation/stratospheric coarse mode mass through condensation, whereas in FC the  $\text{H}_2\text{SO}_4$  contributes a larger portion to stratospheric burden through nucleation and condensation onto the Aitken mode, which eventually grow to accumulation/coarse mode through renaming and coagulation (though the majority of condensation still occurs in the final accumulation/stratospheric coarse mode).

Overall, these processes are delayed in FC compared to SC, though the delay is smaller in MAM5FC compared to MAM5SC due to OH replenishment. The full chemistry experiments also have stronger nucleation, coagulation and renaming processes with larger changes in aerosol mass due to these processes compared to the corresponding simple chemistry experiments. In conjunction with the stratospheric burden plotted in Fig. 4, this suggests that removal processes are also stronger in the full chemistry experiments. MAM4 also generally has stronger nucleation leading to higher Aitken mode concentrations, and therefore also has stronger coagulation processes than MAM5-PSA.

We have also examined the microphysical processes divided by latitudinal regions (Supplementary Figs. 5 and 6). The same patterns as above apply between 30 °S and 30 °N. Above 30 °N and below 30 °S the same signal from the Pinatubo is still present, though slightly delayed due to the time that it took for the aerosol to transport poleward. Signals from other eruptions (Spurr and Lascar) are also present. The main difference is in RNMasc. Above 30 °N and below 30 °S, renaming into the stratospheric coarse mode for MAM5-PSA continues long after the Pinatubo eruption, contributing to the long lifespan of stratospheric sulfate in MAM5-PSA as depicted in Figures 4 and 5. This did not occur between 30 °N and 30 °S, where the stratospheric burden also declines more rapidly as seen in Fig. 5. It is also worth noting that the same pattern does not exist for the other tendencies, and only RNMasc continues to a significant degree past 1992. The condensation tendencies are also not evenly divided across different latitudinal regions, with 30 °S-80 °S and 30 °N-80 °N having stronger condensation tendencies for MAM4FC in the Aitken and accumulation modes, and for MAM5FC in the Aitken and stratospheric coarse modes. Notable increases in condensation can be seen during the Spurr and Lascar eruptions, though it may also be due to Aitken mode aerosols from the original Pinatubo eruption being transported away from the equator and continuing their growth elsewhere.


### 3.4 Simulated AOD and net solar flux

Figure 8 depicts simulated AOD from the four experiments and AVHRR observations from 1991-1993, normalized to the corresponding data from 1989 (i.e., the monthly averages during 1991-1993 subtracted by the corresponding monthly average in 1989 for the same pixel), and masked according to the observed area by AVHRR. The AVHRR AOD anomaly had a maximum value in the range of 0.4-0.45 between 10 °N and 10 °S starting from July 1991, corresponding to the Pinatubo eruption. The four simulations do not have as high of an AOD anomaly for the Pinatubo eruption, with an anomaly of 0.25-0.3 for MAM4SC, 0.15-0.2 for MAM4FC, and 0.35-0.4 for both MAM5SC and MAM5FC experiments. In this respect, MAM5-PSA better captures the AOD anomaly due to the Pinatubo eruption. The simulated AOD maximums occurs slightly north of the equator, to the north of the AOD maximum in the AVHRR observation.



**Figure 8: Latitude vs. time plots of model simulated AOD and AVHRR observed AOD. Both were normalized to 1989 AOD output or observations respectively, to eliminate contributions from the troposphere and background stratospheric AOD. Model output was masked to correspond to areas that AVHRR had observation data for. Results were averaged across longitude.**


The four experiments capture well the northward transport of the AOD anomaly in the latter half of 1991 and the first half of 1992, also matching the generally higher stratospheric sulfate burden above 30 °N compared to below 30 °S (Fig. 5). In all four experiments and the AVHRR observations, the AOD anomaly diminished over time following the Pinatubo eruption, returning towards 0 starting from 1993. This decay is weaker in the MAM5-PSA experiments compared to MAM4 and the AVHRR results, with lingering positive AOD anomalies of 0.05-0.1 throughout 1993, matching the longer lifespan of the stratospheric sulfate in MAM5-PSA as seen in Fig. 4.



**Figure 9: Comparison of AOD anomaly between simulations and observations. The left panel shows the time evolution of monthly AOD anomaly values from simulations and AVHRR and GloSSAC observations, while the right panel shows the time evolution of normalized AOD anomaly.**


We have evaluated the characteristics of the time evolution of simulated global AOD anomaly and its e-folding time against observational data. The time evolution of global monthly AOD anomaly, averaged between 60°S–60°N, is depicted in the left panel of Fig. 9. The AVHRR and the Global Space-based Stratospheric Aerosol Climatology (GloSSAC) were two satellite-based AOD observations that provided global coverage during the Mt. Pinatubo eruption. GloSSAC offered zonal values with a latitudinal resolution of 5° and uniform spatio-temporal coverage up to the year 1994. However, due to instrument saturation for optical depths around 0.15, GloSSAC was less accurate in the center of tropical clouds during the first months after the Mt. Pinatubo eruption. Conversely, AVHRR could only measure stratospheric AOD values larger than 0.01, making it less reliable when AOD values were low (Russell et al., 1996; Quaglia et al., 2023). To calculate AOD anomaly values, background levels (from June 1989 to May 1991) were removed from both observations and simulations. The MAM5FC and MAM5SC simulations produced a reasonable global mean AOD anomaly peak of around 0.15 in November 1991, when the simulated sulfate aerosol burden peaked (Fig. 4). These peaks were higher than the GloSSAC peak (0.12) but lower than the AVHRR peak (0.19). The MAM4FC and MAM4SC simulations produced AOD anomaly peaks (0.11) that were smaller than GloSSAC, indicating an apparent underestimation of AOD anomaly strength. The MAMFC simulation showed a faster AOD anomaly decay than MAM5SC, with values aligning closely with GloSSAC in 1993. In contrast, the MAM4FC and MAM4SC simulations consistently underestimated AOD anomaly compared to GloSSAC from October 1991 to December 1993.

The e-folding time, calculated as the time between the maximum and the 1/e value, was 13 months in AVHRR and 23 months in GloSSAC. The difference between the two measurements is due to the nature of these instruments. The simulated e-folding time in MAM4FC and MAM4SC was 10 months, while MAM5FC and MAM5SC had e-folding times of 19 months and 25 months, respectively. Compared to a previous study (11 to 15 months, Quaglia et al. (2023)), the MAM5FC e-folding time is larger but remains within a reasonable range.



**Figure 10: Latitude vs. time plots of the top-of-model net radiative flux difference between the four experiments and corresponding model simulations with emissions from volcanic eruptions shut off. Units in  $W/m^2$ .**

Figure 10 depicts the differences in the top-of-model net solar flux in the four experiments with and without the emissions from volcanic eruptions (negative values representing cooling from the volcanic aerosol). The location of the maximum of the net solar flux difference agrees well with the location of the AOD anomalies in Fig. 8. The maximum of the net solar flux is simulated to be about  $-6 W/m^2$  for MAM4SC,  $-4 W/m^2$  for MAM4FC, and  $-10 W/m^2$  for both MAM5-PSA experiments. Considering the more accurate AOD simulation in MAM5-PSA, it is expected that MAM5-PSA also better captures climate effects from the Pinatubo eruption. The time variations of the global mean top-of-model net radiative flux shows that, during October-November of 1991, MAM4 experiments produce weaker peak values of around  $-1.5 W/m^2$  while MAM5-PSA produces peak values of roughly  $-3.0 W/m^2$  (Fig. 11). The latter results agree closely with the those simulated by CESM experiments in Mills et al. (2017) as well as the E3SMv2 modified MAM4 results from Brown et al. (2024), where the peak values are also roughly  $-3.0 W/m^2$ .



**Figure 11: Global mean net radiative flux in the four experiments, where negative values represent a net upward flux.**

#### 4 Conclusions and discussion

In this work, we have described an implementation of a 5-mode version of the Modal Aerosol Module (MAM5-PSA) used in E3SMv2 as well as a more complex “full chemistry” set-up, resolving issues in simulating the coarse mode for stratospheric sulfate seen in previous studies. The intention is to provide more accurate simulations of the chemical and microphysical processes within the stratosphere following the Pinatubo eruption, and to compare it against a simulation using simplified chemistry and the default MAM4 aerosol module. Full chemistry in MAM5-PSA includes the addition of a series of chemical reactions which allow the simulation of OH radical replenishment compared to that in MAM4. MAM5-PSA adds a stratospheric coarse mode that specifically represents the microphysical properties of stratospheric sulfate. In particular, the geometric standard deviation is smaller, contributing to a longer lifespan for stratospheric sulfate aerosol. Corresponding to these changes, we carried out four experiments: MAM4SC, MAM4FC, MAM5SC and MAM5FC to simulate the burdens, AOD, and radiative effect of volcanic aerosols following the Pinatubo eruption.

These changes led to large differences in both the temporal variations and the spatial distributions of sulfate concentrations. Directly after the eruption, the SC experiments had the fastest rise in sulfate concentrations, as FC was limited by OH required to oxidize  $\text{SO}_2$ . MAM5FC was significantly faster than MAM4FC however due to the OH radical replenishment. Vertically, sulfate distributions were generally at lower altitudes in FC compared to SC, caused by the differences in  $\text{SO}_2$ , OH, and specific humidity within the stratosphere. With respect to the stratospheric sulfate burden, MAM5FC agreed best with the HIRS observations from the eruption of the volcano to the end of 1992. Starting in 1993, stratospheric sulfate burdens in all four of the experiments began to diminish, but the process was slower in MAM5-PSA because of the stronger deposition in MAM4 relative to MAM5-PSA. The slow removal of stratospheric sulfate burden in MAM5-PSA occurs mostly in the high latitudes but is faster near the equator. Compared to WOPC observations, MAM5-PSA has comparable mass in the coarse mode, but coarse mode particles are smaller,

which may be responsible for the longer lifespan of stratospheric sulfate after the start of 1993. We assessed the time evolution and e-folding time of simulated global AOD anomaly against satellite observations (AVHRR and GloSSAC). The MAM5FC and MAM5SC simulations generally produced reasonable AOD peaks and decay patterns, while MAM4FC and MAM4SC tended to underestimate AOD strength and overestimate the decay rate.

We further analyzed the tendencies associated with the microphysical processes related to the growth of stratospheric sulfate: nucleation (NUCL), coagulation (COAG), condensation (AitkenCond, AccumCond, StratoCoarseCond), and renaming (RNMaa and RN Masc). In general, all of these processes are stronger in FC than in SC. For condensation, FC is stronger than SC in the Aitken mode due to the higher concentrations of Aitken mode sulfate, and MAM4 is stronger than MAM5-PSA in the accumulation mode again due to higher concentrations. It should be noted that the differences in integrated tendencies between the experiments were often much larger than the differences in stratospheric sulfate burdens between the experiments, which suggested that sedimentation was faster in FC than in SC to counterbalance the stronger aerosol production suggested by the stronger tendencies. In terms of timing, the tendencies generally initiate and conclude faster in SC compared to FC due to not having limitations for OH radicals, which correspond well with the occurrence of peaks of sulfate burden. Overall, in terms of accurately portraying the retention of sulfate aerosols in the stratosphere, MAM5-PSA provides a substantial improvement over MAM4 during 1992.

We compared the model output AOD against AVHRR observations to assess overall performance. Immediately after the eruption, MAM5-PSA performs better than MAM4 by producing AOD anomalies closer to AVHRR. In all four experiments, the northward transport of aerosols due to the Brewer-Dobson circulation was accurately captured, but the AOD related to the Cerro Hudson and Spurr eruptions was not distinct compared to AVHRR. These eruptions have a much lower injection height than Pinatubo (11-16 km and 9-14.5 km altitude respectively) and a much smaller emission of SO<sub>2</sub> which may explain the lack of a signal. In terms of the changes in global mean net radiative flux, the simulated largest global mean values following the eruption were roughly -1.5 W/m<sup>2</sup> during October-November of 1991 in MAM4 experiments and about -3 W/m<sup>2</sup> in MAM5-PSA experiments. MAM5-PSA results were much closer to the number seen in Mills et al. (2017) and Brown et al. (2024), also roughly -3 W/m<sup>2</sup>.

A number of factors contribute to the uncertainty of our results. One important factor is the known dry bias in E3SM's stratosphere (Christiane Jablonowski, personal communication, 2024), since water vapor plays an important role in the HO<sub>x</sub> cycle relevant to the initial sulfate formation, as well as later sulfate growth and sedimentation. Another issue is the presence of minor volcanic eruptions such as Cerro Hudson, Spurr and Lascar, which produced much less SO<sub>2</sub> than Pinatubo, but still significant enough to leave a distinguishable signal following Pinatubo's eruption. A third factor is the interaction between the Pinatubo eruption and stratospheric ozone. Volcanic eruptions are known to affect stratospheric ozone concentrations (Peng et al., 2023) directly and indirectly through factors such as the decreases in tropospheric temperatures, modified circulation and the changes in the stratospheric chemical composition (Dhomse et al., 2015). The capabilities of E3SM in simulating stratospheric ozone in the unique circumstances of a volcanic



eruption are not well studied. Further understanding of these factors would greatly aid in the accurate simulation of Pinatubo and other volcanic eruptions.

#### **Code and data availability.**

AVHRR observational data was downloaded from <https://www.ncei.noaa.gov/data/avhrr-aerosol-optical-thickness/access>. The VolcanEESM emissions files used in this study can be found at <https://svn-ccsm-inputdata.cgd.ucar.edu/trunk/inputdata/atm/cam/chem/stratvolc/>. The source code and run scripts used in this study can be found at <https://zenodo.org/doi/10.5281/zenodo.12734295>.

#### **Author contributions.**

Conceptualization: AH, XL and ZK. Data curation: AH, ZK and BW. Formal analysis: AH. Funding acquisition: XL, DB and KP. Investigation: AH. Methodology: AH, XL and ZK. Project administration: XL, HW, SX, DB and KP. Resources: XL, DB and KP. Software: AH implemented MAM5 into E3SMv2 with assistance from ZK, MW, BW and ZL. Original MAM5 implementation in E3SMv1 by ZK. ZK, MW, HW and QT enabled and improved the chemistry scheme that “full chemistry” is based on. Supervision: XL, DB and KP. Validation: AH. Visualization: AH. Writing – original draft preparation: AH. Writing – review and editing: all authors.

#### **Competing interests.**

Some authors are members of the editorial board of Atmospheric Chemistry and Physics.

#### **Disclaimer.**

#### **Acknowledgements.**

XL, AH, BW, HB, DB, and KP acknowledge the support of this work by the Laboratory Directed Research and Development program at Sandia National Laboratories, a multimission laboratory managed and operated by National Technology and Engineering Solutions of Sandia, LLC., a wholly owned subsidiary of Honeywell International, Inc., for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA-0003525. XL, ZK, ZL, MW, HW, QT, and SX acknowledge the support by the U.S. Department of Energy (DOE), Office of Science, Office of Biological and Environmental Research, Earth and Environmental System Modeling program as part of the Energy Exascale Earth System Model (E3SM) project. The work at the Lawrence Livermore National Laboratory (LLNL) was performed under the auspices of the U.S. DOE by LLNL under Contract DE-AC52-07NA27344. The Pacific Northwest National Laboratory (PNNL) is operated for DOE by Battelle Memorial Institute under contract DE-AC05-76RLO1830.

## References

- Aubry, T. J., Engwell, S., Bonadonna, C., Carazzo, G., Scollo, S., Van Eaton, A. R., Taylor, I. A., Jessop, D., Eycheenne, J., Gouhier, M., Mastin, L. G., Wallace, K. L., Biass, S., Bursik, M., Grainger, R. G., Jellinek, A. M., and Schmidt, A.: The Independent Volcanic Eruption Source Parameter Archive (IVESPA, version 1.0): A new observational database to support explosive eruptive column model validation and development, *Journal of Volcanology and Geothermal Research*, 417, 107295, <https://doi.org/10.1016/j.jvolgeores.2021.107295>, 2021.
- Baran, A. and Foot, J.: New application of the operational sounder HIRS in determining a climatology of sulphuric acid aerosol from the Pinatubo eruption, *Journal of Geophysical Research: Atmospheres*, 99, 25673–25679, 1994.
- Bergman, T., Kerminen, V. M., Korhonen, H., Lehtinen, K. J., Makkonen, R., Arola, A., Mielonen, T., Romakkaniemi, S., Kulmala, M., and Kokkola, H.: Evaluation of the sectional aerosol microphysics module SALSA implementation in ECHAM5-HAM aerosol-climate model, *Geosci. Model Dev.*, 5, 845–868, 10.5194/gmd-5-845-2012, 2012.
- Brown, H. Y., Wagman, B., Bull, D., Peterson, K., Hillman, B., Liu, X., Ke, Z., and Lin, L.: Validating a microphysical prognostic stratospheric aerosol implementation in E3SMv2 using observations after the Mount Pinatubo eruption, *Geosci. Model Dev.*, 17, 5087–5121, 10.5194/gmd-17-5087-2024, 2024.
- Carn, S.: Multi-Satellite Volcanic Sulfur Dioxide L4 Long-Term Global Database V4 [dataset], 10.5067/MEASURES/SO2/DATA405, 2022.
- Crutzen, P. J.: Albedo enhancement by stratospheric sulfur injections: a contribution to resolve a policy dilemma?, *Climatic change*, 77, 211, 2006.
- Deshler, T.: A review of global stratospheric aerosol: Measurements, importance, life cycle, and local stratospheric aerosol, *Atmospheric Research*, 90, 223–232, 2008.
- Deshler, T., Johnson, B. J., and Rozier, W. R.: Balloonborne measurements of Pinatubo aerosol during 1991 and 1992 at 41 N: Vertical profiles, size distribution, and volatility, *Geophysical Research Letters*, 20, 1435–1438, 1993.
- Deshler, T., Hervig, M., Hofmann, D., Rosen, J., and Liley, J.: Thirty years of in situ stratospheric aerosol size distribution measurements from Laramie, Wyoming (41 N), using balloon-borne instruments, *Journal of Geophysical Research: Atmospheres*, 108, 2003.
- Dhomse, S. S., Chipperfield, M. P., Feng, W., Hossaini, R., Mann, G. W., and Santee, M. L.: Revisiting the hemispheric asymmetry in midlatitude ozone changes following the Mount Pinatubo eruption: A 3-D model study, *Geophysical Research Letters*, 42, 3038–3047, <https://doi.org/10.1002/2015GL063052>, 2015.
- Emmons, L. K., Walters, S., Hess, P. G., Lamarque, J.-F., Pfister, G. G., Fillmore, D., Granier, C., Guenther, A., Kinnison, D., and Laepple, T.: Description and evaluation of the Model for Ozone and Related chemical Tracers, version 4 (MOZART-4), *Geoscientific Model Development*, 3, 43–67, 2010.
- Gao, C. Y., Naik, V., Horowitz, L. W., Ginoux, P., Paulot, F., Dunne, J., Mills, M., Aquila, V., and Colarco, P.: Volcanic drivers of stratospheric sulfur in GFDL ESM4, *Journal of Advances in Modeling Earth Systems*, 15, e2022MS003532, 2023.
- Golaz, J. C., Van Roekel, L. P., Zheng, X., Roberts, A. F., Wolfe, J. D., Lin, W., Bradley, A. M., Tang, Q., Maltrud, M. E., and Forsyth, R. M.: The DOE E3SM Model Version 2: Overview of the physical model and initial model evaluation, *Journal of Advances in Modeling Earth Systems*, 14, e2022MS003156, 2022.
- Kleinschmitt, C., Boucher, O., and Platt, U.: Sensitivity of the radiative forcing by stratospheric sulfur geoengineering to the amount and strategy of the SO<sub>2</sub> injection studied with the LMDZ-S3A model, *Atmos. Chem. Phys.*, 18, 2769–2786, 10.5194/acp-18-2769-2018, 2018.
- Kleinschmitt, C., Boucher, O., Bekki, S., Lott, F., and Platt, U.: The Sectional Stratospheric Sulfate Aerosol module (S3A-v1) within the LMDZ general circulation model: description and evaluation against stratospheric aerosol observations, *Geosci. Model Dev.*, 10, 3359–3378, 10.5194/gmd-10-3359-2017, 2017.
- Kokkola, H., Korhonen, H., Lehtinen, K. E. J., Makkonen, R., Asmi, A., Järvenoja, S., Anttila, T., Partanen, A. I., Kulmala, M., Järvinen, H., Laaksonen, A., and Kerminen, V. M.: SALSA – a Sectional Aerosol module for Large Scale Applications, *Atmos. Chem. Phys.*, 8, 2469–2483, 10.5194/acp-8-2469-2008, 2008.
- Kremser, S., Thomason, L. W., von Hobe, M., Hermann, M., Deshler, T., Timmreck, C., Toohey, M., Stenke, A., Schwarz, J. P., and Weigel, R.: Stratospheric aerosol—Observations, processes, and impact on climate, *Reviews of Geophysics*, 54, 278–335, 2016.
- Laakso, A., Korhonen, H., Romakkaniemi, S., and Kokkola, H.: Radiative and climate effects of stratospheric sulfur geoengineering using seasonally varying injection areas, *Atmos. Chem. Phys.*, 17, 6957–6974, 10.5194/acp-17-6957-2017, 2017.

- Laakso, A., Niemeier, U., Vioni, D., Tilmes, S., and Kokkola, H.: Dependency of the impacts of geoengineering on the stratospheric sulfur injection strategy—Part 1: Intercomparison of modal and sectional aerosol modules, *Atmospheric Chemistry and Physics*, 22, 93-118, 2022.
- Liu, X.: Aerosols and Climate Effects, in: *Fast Processes in Large-Scale Atmospheric Models*, Wiley, 53-86, <https://doi.org/10.1002/9781119529019.ch3>, 2023.
- Liu, X., Ma, P.-L., Wang, H., Tilmes, S., Singh, B., Easter, R., Ghan, S., and Rasch, P.: Description and evaluation of a new four-mode version of the Modal Aerosol Module (MAM4) within version 5.3 of the Community Atmosphere Model, *Geoscientific Model Development*, 9, 505-522, 2016.
- Mills, M. J., Richter, J. H., Tilmes, S., Kravitz, B., MacMartin, D. G., Glanville, A. A., Tribbia, J. J., Lamarque, J. F., Vitt, F., and Schmidt, A.: Radiative and chemical response to interactive stratospheric sulfate aerosols in fully coupled CESM1 (WACCM), *Journal of Geophysical Research: Atmospheres*, 122, 13,061-013,078, 2017.
- Mills, M. J., Schmidt, A., Easter, R., Solomon, S., Kinnison, D. E., Ghan, S. J., Neely III, R. R., Marsh, D. R., Conley, A., and Bardeen, C. G.: Global volcanic aerosol properties derived from emissions, 1990–2014, using CESM1 (WACCM), *Journal of Geophysical Research: Atmospheres*, 121, 2332-2348, 2016.
- Neely III, R. R. S., A.: VolcanEESM: Global volcanic sulphur dioxide (SO<sub>2</sub>) emissions database from 1850 to present - Version 1.0 [dataset], 10.5285/76ebdc0b-0eed-4f70-b89e-55e606bcd568, 2016.
- Parker, D., Wilson, H., Jones, P. D., Christy, J., and Folland, C. K.: The impact of Mount Pinatubo on world-wide temperatures, *International Journal of Climatology: A Journal of the Royal Meteorological Society*, 16, 487-497, 1996.
- Peng, Y., Yu, P., Portmann, R. W., Rosenlof, K. H., Zhang, J., Liu, C.-C., Li, J., and Tian, W.: Perturbation of Tropical Stratospheric Ozone Through Homogeneous and Heterogeneous Chemistry Due To Pinatubo, *Geophysical Research Letters*, 50, e2023GL103773, <https://doi.org/10.1029/2023GL103773>, 2023.
- Quaglia, I., Timmreck, C., Niemeier, U., Vioni, D., Pitari, G., Brodowsky, C., Brühl, C., Dhomse, S. S., Franke, H., and Laakso, A.: Interactive stratospheric aerosol models' response to different amounts and altitudes of SO<sub>2</sub> injection during the 1991 Pinatubo eruption, *Atmospheric Chemistry and Physics*, 23, 921-948, 2023.
- Robock, A.: Volcanic eruptions and climate, *Reviews of geophysics*, 38, 191-219, 2000.
- Russell, P. B., Livingston, J. M., Pueschel, R. F., Bauman, J. J., Pollack, J. B., Brooks, S. L., Hamill, P., Thomason, L. W., Stowe, L. L., Deshler, T., Dutton, E. G., and Bergstrom, R. W.: Global to microscale evolution of the Pinatubo volcanic aerosol derived from diverse measurements and analyses, *Journal of Geophysical Research: Atmospheres*, 101, 18745-18763, <https://doi.org/10.1029/96JD01162>, 1996.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, Wiley, 2016.
- Sheng, J. X., Weisenstein, D. K., Luo, B. P., Rozanov, E., Stenke, A., Anet, J., Bingemer, H., and Peter, T.: Global atmospheric sulfur budget under volcanically quiescent conditions: Aerosol-chemistry-climate model predictions and validation, *Journal of Geophysical Research: Atmospheres*, 120, 256-276, 2015.
- Tilmes, S., Richter, J. H., Mills, M. J., Kravitz, B., MacMartin, D. G., Vitt, F., Tribbia, J. J., and Lamarque, J. F.: Sensitivity of aerosol distribution and climate response to stratospheric SO<sub>2</sub> injection locations, *Journal of Geophysical Research: Atmospheres*, 122, 12,591-512,615, 2017.
- Tilmes, S., Vioni, D., Jones, A., Haywood, J., Séférian, R., Nabat, P., Boucher, O., Bednarz, E. M., and Niemeier, U.: Stratospheric ozone response to sulfate aerosol and solar dimming climate interventions based on the G6 Geoengineering Model Intercomparison Project (GeoMIP) simulations, 2022.
- Vattioni, S., Weisenstein, D., Keith, D., Feinberg, A., Peter, T., and Stenke, A.: Exploring accumulation-mode H<sub>2</sub>SO<sub>4</sub> versus SO<sub>2</sub> stratospheric sulfate geoengineering in a sectional aerosol–chemistry–climate model, *Atmospheric Chemistry and Physics*, 19, 4877-4897, 2019.
- Vioni, D., Bednarz, E. M., Lee, W. R., Kravitz, B., Jones, A., Haywood, J. M., and MacMartin, D. G.: Climate response to off-equatorial stratospheric sulfur injections in three Earth system models—Part 1: Experimental protocols and surface changes, *Atmospheric Chemistry and Physics*, 23, 663-685, 2023.
- Vioni, D., MacMartin, D. G., Kravitz, B., Tilmes, S., Mills, M. J., Richter, J. H., and Boudreau, M. P.: Seasonal injection strategies for stratospheric aerosol geoengineering, *Geophysical Research Letters*, 46, 7790-7799, 2019.
- Wang, H., Easter, R. C., Zhang, R., Ma, P. L., Singh, B., Zhang, K., Ganguly, D., Rasch, P. J., Burrows, S. M., and Ghan, S. J.: Aerosols in the E3SM Version 1: New developments and their impacts on radiative forcing, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001851, 2020.
- Wu, M., Wang, H., Easter, R. C., Lu, Z., Liu, X., Singh, B., Ma, P.-L., Tang, Q., Zaveri, R. A., Ke, Z., Zhang, R., Emmons, L. K., Tilmes, S., Dibb, J. E., Zheng, X., Xie, S., and Leung, L. R.: Development and Evaluation of E3SM-MOSAIC: Spatial Distributions and Radiative Effects of Nitrate Aerosol, *Journal of Advances in Modeling Earth Systems*, 14, e2022MS003157, <https://doi.org/10.1029/2022MS003157>, 2022.

Yang, F. and Schlesinger, M. E.: On the surface and atmospheric temperature changes following the 1991 Pinatubo volcanic eruption: A GCM study, *Journal of Geophysical Research: Atmospheres*, 107, ACL 8-1-ACL 8-14, 2002.

Zhao, X. a. N. C. P.: NOAA Climate Data Record (CDR) of AVHRR Daily and Monthly Aerosol Optical Thickness (AOT) over Global Oceans (Monthly subset) (4.0) [dataset], 10.25921/w3zj-4y48, 2022.

Zhuo, Z., Kirchner, I., Pfahl, S., and Cubasch, U.: Climate impact of volcanic eruptions: the sensitivity to eruption season and latitude in MPI-ESM ensemble experiments, *Atmospheric Chemistry and Physics*, 21, 13425-13442, 2021.