

Control of spatio-temporal variability of ocean nutrients in the East Australian Current

Megan Jeffers¹, Christopher C. Chapman², Bernadette M. Sloyan², and Helen Bostock¹

¹School of the Environment, Faculty of Science, The University of Queensland, Australia

²CSIRO Environment, Hobart Marine Laboratories, Hobart, Australia

Correspondence: Megan Jeffers (m.jeffers@uq.edu.au)

Abstract. The East Australian Current (EAC), the South Pacific's southward flowing western boundary current, dominates the marine environment of the east coast of Australia. Upwelling of deep EAC nutrient rich water into the oligotrophic surface waters is very important for maintaining upper ocean productivity. However, the role of EAC dynamics in upper ocean nutrient variability and resulting productivity is poorly understood. In this study we use physical and biogeochemical data collected 5 from 2012–2022 to improve understanding of the variability of the nutrients in the upper water column at $\sim 27^{\circ}$ S, a subtropical region strongly influenced by [the](#) EAC.

The 10-year data set shows that there is a seasonal increase in nutrient concentrations in the upper water column (0–200 m) in the [Austral](#)~~austral~~ spring (September–November) and autumn (March–May), and a minimum in winter (June–August). We also find that the nutrient concentrations in the upper water column are influenced by the position of the EAC jet. We 10 modes of variability in the EAC's position are identified: an inshore mode with [the](#) jet flowing along the continental slope and; an offshore mode with the current core detached from the continental slope and flowing over the adjacent abyssal plain. The position of the EAC jet influences the location of upwelling of nutrient-rich water at depth (>200 m). For the EAC inshore mode, cooler, nutrient-rich waters are restricted to the area of the continental shelf and slope that is inshore of the EAC. The offshore mode exhibits a wider distribution of nutrient-rich waters over both the inshore shelf and slope and, [the](#) offshore 15 abyssal Tasman Sea. Our analysis highlights the important interactions between [nutrient concentrations and distribution](#) and the highly variable EAC [and the distribution of high nutrient waters](#), which has implications for primary production, fisheries, and the biological carbon pump.

1 Introduction

The East Australian Current (EAC) is the western boundary current (WBC) of the South Pacific subtropical gyre, and plays 20 a significant role in the southwest Pacific Ocean circulation and nutrient distribution. The EAC transports warm water from the Coral Sea into the [more temperate climate of the](#) Tasman Sea (Archer et al., 2017a; Sloyan et al., 2016). The upper EAC waters (0–200 m) are oligotrophic (low nutrient concentration), however below 200 m the EAC transports nutrient rich water southwards ([Suthers et al., 2011; Chiswell et al., 2015](#)).

The EAC is fed by the South Equatorial Current via several westward flowing jets between 15°-S and 23°-S. The North 25 Vanuatu and the North Caledonian jets flow into the northern Coral Sea at ~15 °-S and ~19°-S, respectively, providing inflowing waters that are warm and salty, with low oxygen and nutrient concentrations (Kessler and Cravatte, 2013). At these latitudes the EAC has a relatively weak southward transport of between 6–8 Sv (Ganachaud et al., 2014; Kessler and Cravatte, 2013). Around 23°-S, the addition of the sub-surface South Caledonian Jet strengthens the EAC flow to a mean southward transport of 24.6 Sv and also increases the subsurface oxygen concentrations (Kessler and Cravatte, 2013; Sloyan et al., 2016). 30 This marks the start of the EAC jet “intensification zone” that extends southwards until the separation latitude at approximately 31–34°-S (Oke et al., 2019; Ridgway and Dunn, 2003). At the separation latitude, the EAC bifurcates into two eddy-dominated flow paths, with one proceeding southward along the eastern Australian coast to Tasmania known as the south EAC extension, and the other flowing across the Tasman Sea to New Zealand as the Tasman Front, or east EAC extension (Ganachaud et al., 2014; Godfrey et al., 1980; Oke et al., 2019; Ridgway and Dunn, 2003; Sutton and Bowen, 2014). **However,** 35 **recent work has shown that the** **The** latitude of the separation point is significantly influenced by changes in the EAC further upstream (Li et al., 2021) (Cetina-Heredia et al., 2014; Li et al., 2021), emphasising the importance of understanding the EAC jet **further upstream** in this intensification zone.

The EAC exhibits seasonal changes in its physical flow. Several observational-based studies have revealed a seasonal change of approximately 9 Sv in the volume transport of the EAC, with the maximum occurring in austral summer and minimum in 40 winter (Archer et al., 2017a; Godfrey et al., 1980; Ridgway and Godfrey, 1997; Sloyan et al., 2024). Temperature and salinity also show a seasonal cycle when considering the surface waters (potential density $\rho_0 > 26.2 \text{ kg/m}^3$), which are warmer and fresher during the summer (Oke et al., 2019; Sloyan et al., 2016). Additionally, at 29°S, Everett et al. (2014) identified a seasonal cycle in dissolved nutrients with increased dissolved nitrate and silicate concentrations in austral winter and spring (June - September).

Between the latitudes of 25–30°-S the EAC, whilst maintaining a more coherent jet-like flow, is highly dynamic due to 45 interactions with mesoscale eddies (Archer et al., 2017a; Oke et al., 2019) and larger scale forcing (Sloyan and O’Kane, 2015; Bull et al., 2020). This meandering of the jet can cause significant changes to the vertical structure of the current and southward transport (Roughan et al., 2022b; Sloyan et al., 2016). Previous work at ~27°-S identified a meander in the EAC jet which shifted the position of the main poleward (southward) flow laterally, on and off the continental slope (Sloyan 50 et al., 2016). **Similar variability and jet meandering has been observed in other WBCs. The Kuroshio Current experiences variability in its core position and strength, and experiences meandering and interactions with bathymetry and mesoscale eddies** (Ebuchi and Hanawa, 2003; Kawabe, 2005, 1995; Waseda et al., 2003). **The Florida Current is the upstream portion of the Gulf Stream, and has a very similar jet structure to the EAC (Archer et al., 2018), and exhibits a similar meandering behaviour, with meanders occurring on a time-scale of 3–30 days (Archer et al., 2017b). Other WBCs also experience meandering, including the Gulf Stream (Mao et al., 2023), Agulhas Current (Goschen et al., 2015), and the Brazil Current (Da Silveira et al., 2008).**

Additionally, Such current variability has been linked to biogeochemical variability in the EAC and other WBC systems. For example, in the EAC exhibits seasonal changes in its physical flow. Several observational-based studies have revealed a seasonal change of approximately 9 Sv in the volume transport of the EAC, with Chapman et al. (2024) showed observations

of nutrient injection into the surface layers during meandering, and that the vertical velocities that drove this were catalysed by the interaction of EAC with mesoscale eddies. Similarly, in the Kuroshio, the large meander mode results in an uplift of the nutricline, which increases nutrient availability in the near surface waters (Hayashida et al., 2023). In the upstream region of the Kuroshio, increased speeds of the current causes an uplift of nutrients onshore (Chen et al., 2022). Additionally, during periods where the Kuroshio sits closer to the coast, it interacts with bathymetry which causes strong vertical mixing and uplift of nutrients to the continental shelf (Durán Gómez and Nagai, 2022). Mesoscale variability in the Florida Current is linked to the upwelling of cool, nutrient rich waters between the shelf break and the offshore meander (Fiechter and Mooers, 2007; Kourafalou and Kang, 2015). Upwelling has also been linked to the movement of jet meanders in the Agulhas Current, with upwelling occurring when the maximum occurring in austral summer and minimum in winter (Archer et al., 2017a; Godfrey et al., 1980; Ridgway and Godfrey, 1997). Temperature and salinity also show a seasonal cycle when considering the surface waters (potential density $\rho_0 > 26.2 \text{ kg/m}^3$), which are warmer and fresher during the summer (Oke et al., 2019) current shifts onto the continental shelf, or shifts offshore (Goschen et al., 2015). It is clear that meandering of WBCs upstream of their separation points is not unique to the EAC, and the current meandering can result in upwelling, which has been observed in several WBC systems.

Upwelling of deeper EAC nutrient rich water into the oligotrophic surface waters influences the primary production patterns, plankton community composition, and nutrient utilization strategies of marine organisms along the east coast of Australia (Everett et al., 2014; Hassler et al., 2011). At 29° S, Everett et al. (2014) identified a seasonal cycle in dissolved nutrients with increased dissolved nitrate and silicate concentrations in Austral winter and spring (June–September). Previous work has shown that the EAC separation point is characterized by upwelling, resulting in an enhanced nutrient supply and increased biological activity, contributing to more productive fisheries (Everett et al., 2011; Figueira and Booth, 2010; Hassler et al., 2011; Olson, 2001; Suthers et al., 2023). The upwelling of the nutrients to the surface layer in this region has been attributed to various factors, including wind forcing, EAC divergence, and the presence of eddies (Everett et al., 2011; Gibbs et al., 1998; Godfrey et al., 1980; Rochford, 1975; Roughan and Middleton, 2002, 2004). Some studies attribute the increased nutrient supply to the higher velocity of the current induced by topographic forcing (Blackburn and Cresswell, 1993; Boland and Church, 1981; Oke and Middleton, 2000), while other studies attributed the increased nutrient concentrations to the action of the EAC separating from the coast (Tranter et al., 1986). At the separation point, it has been suggested that shifts in the position of the EAC are associated with periods of increased nutrient upwelling (Rochford, 1975), resulting in the transport of nutrient-rich waters onto the continental shelf and coastal areas (Everett et al., 2014; Oke and Middleton, 2001; Roughan and Middleton, 2004). Roughan and Middleton (2002) find that when the EAC shifted towards the coast, nutrient upwelling to the surface occurred alongside uplifted isotherms. This upwelling supports considerable productivity in coastal and shelf regions in an otherwise relatively low productivity region (Nieblas et al., 2009; Roughan et al., 2022a; Schaeffer et al., 2013). While there is evidence that EAC dynamics influence nutrient supply to the euphotic upper ocean (Chapman et al., 2024), our understanding of nutrient dynamics within the EAC is limited, largely due to a lack of data (Everett et al., 2011, 2014; Hassler et al., 2011; McGillicuddy Jr, 2016; Oke and Middleton, 2001; Rocha et al., 2019; Schaeffer et al., 2016).

In this study we examine the spatial and temporal variability of nutrients from the continental shelf to the off-shore region at $\sim 27^\circ \text{ S}$. We use 10-years of nutrient bottle data collected between 2012–2022 (Sloyan et al., 2016, 2024). We examine

the seasonality of the nutrients and the role of the position of the EAC, relative to the continental shelf and slope, in influencing
95 the distribution of nutrients in the upper water column and find that there is a seasonal increase in nutrient concentrations in the upper water column (0–200 m) in the Austral spring (September–November) and autumn (March–May), and a minimum in winter (June–August). It is also clear that the nutrient concentrations in the upper water column are influenced by the bimodal distribution of the EAC jet position. As the position of the EAC jet shifts longitudinally, it can be broadly categorised into two modes: the inshore mode and offshore mode. When the EAC jet is in its inshore mode—jet located on the continental slope—
100 cooler, nutrient-rich waters are restricted to an area inshore of the EAC jet. The offshore EAC jet mode—EAC jet located over the abyssal plain—is associated with a wider distribution of nutrient-rich waters that extends from the continental shelf, across the slope and into the offshore waters of the adjacent abyssal plain. Our analysis highlights the important interactions between nutrient concentrations and distribution and the highly variable EAC, which has implications for primary production, fisheries, and the biological carbon pump. Understanding the dynamical implications of the position of EAC on nutrient distribution
105 is essential for elucidating the broader implications of the EAC's role on marine ecosystems and fisheries. This long-term dataset offers a valuable insight into the oceanography, biogeochemical cycling, and the EAC's dynamical influence on the surface and mixed layer oceanography, biogeochemical cycling, and nutrient concentrations.

2 Methods

2.1 Data

110 2.1.1 The EAC Moorings

To capture the behaviour of the EAC in the intensification zone where it has a defined jet-like structure, a “picket-fence” mooring array was established by CSIRO and the Australian Integrated Marine Observing System (IMOS) at approximately 27°S, offshore of Brisbane, Australia (Sloyan et al., 2024). The mooring array was operational from 2012 to 2022, except for a 22-month period between 2013 and 2015. The EAC mooring array extends from 153.5°E to 155.5°E, covering a depth range
115 of approximately 60 m to 5000 m (Figure 1; Sloyan et al. (2024)). The moorings were equipped with Acoustic Doppler current profiling (ADCP) instruments that provided a vertical profile of horizontal current velocity and discrete temperature and salinity instruments at various vertical intervals. The data used in this study is the daily mooring velocity data product with missing or bad velocities “filled” using a machine learning approach (Sloyan et al. (2023, 2024); <https://doi.org/10.25919/10h0-yf37>; Sloyan et al. (2023, 2024)).

120 2.1.2 Conductivity, Temperature, Depth and Hydrochemical Data

For the assessment of the EAC's physical and biogeochemical properties, we utilize 136 utilise 162 Conductivity, Temperature, Depth (CTD) and Niskin bottle profiles collected during research voyages (Figure 1, Table 1). The CTD Of the 162 CTD stations, nutrient samples were collected on 136. CTD profiles provide measurements of temperature, salinity, and dissolved oxygen at 1 dbar pressure intervals. Nutrients were measured on water samples. Water samples for nutrient analyses were




Figure 1. Location of CTD data in context of broader surroundings. Panel (a) shows 20-year time mean (Jan 2000 – Dec 2019) of surface geostrophic currents from IMOS OceanCurrent. The location of the EAC mooring array is indicated by the red line. The black star shows the location of Brisbane, Australia and hydrochemistry NC=New Caledonia. The black square outlines the region panel (b). Panel (b) shows the location of CTD stations (red dots) and EAC array moorings (black yellow triangles) used in this study. The bathymetric bold white line shows the position of the 154.2°E line used for defining EAC modes. The bathymetric map of the study area is also shown. The with the contour interval (black line) is every 500 m from 50500–4000 m water depth.

125 collected by Niskin bottles at discrete depths that span the entire water column. These water samples were analysed for nitrate, phosphate, and silicate (and nitrite and ammonia that are not used here). No voyages were conducted during Austral austral summer (November–March) in the region to avoid cyclone season, however. However all months from April to November were sampled, providing observations for Austral austral autumn, winter, and spring seasons over the 10-year period. All CTD data is available from the CSIRO Data Trawler (<https://www.cmar.csiro.au/data/trawler/>).

130 Although the CTD data are full depth (depths >4500 m over the abyssal plain), we use data from the upper 200 m of the water column, as this depth range spans the EAC jet core (Sloyan et al., 2024). This depth range also includes a significant phytoplankton production, which is at a maximum in the Coral and Tasman Sea at a range of depths between

Table 1. List of voyages from which the CTD and nutrient data were used for this study. Including voyage ID, dates, research vessel and number of CTD ~~casts~~ deployed and with hydrochemistry.

Voyage ID	Voyage Dates	Vessel	No. Deployments	CTD casts	No. CTD
SS2012_V01	20 th April – 29 th April 2012	Southern Surveyor RV <i>Southern Surveyor</i>	<u>15</u>		
SS2013_V05	20 th August – 2 nd September 2013	Southern Surveyor RV <i>Southern Surveyor</i>	25		
IN2015_V02	15 th May – 26 th May 2015	RV <i>Investigator</i>	<u>17</u>		
IN2016_V04	30 th August – 23 rd September 2015	RV <i>Investigator</i>	<u>4</u>		
IN2016_V06	28 th October – 13 th November 2016	RV <i>Investigator</i>	12		
IN2018_V03	29 th April – 10 th May 2018	RV <i>Investigator</i>	<u>14</u>		
IN2019 IN2019_V05	9 th September – 29 th September 2019	RV <i>Investigator</i>	<u>26</u>		
IN2021_V03	7 th May – 2 nd June 2021	RV <i>Investigator</i>	<u>28</u>		
IN2022_V06	14 th July – 28 th July 2022	RV <i>Investigator</i>	<u>21</u>		

40–100 m ~~depth~~ (Ellwood et al., 2013). In the ~~southwest~~ Southwest Pacific plankton tend to be nitrate and phosphate limited (Ellwood et al., 2013; Ustic et al., 2021) as the southwest (Ellwood et al., 2013; Ustic et al., 2021; Hassler et al., 2011; Doblin et al., 2013). The Southwest Pacific contains a low dissolved phosphorus region centred around 28°S (Martiny et al., 2019). However, $\text{NO}_3^-:\text{PO}_4^{3-}$ ratios show that nitrate is still the primary limiting nutrient (Hassler et al., 2011), particularly in the top 200 m (Doblin et al., 2016). Silicate is also limited a key nutrient in this region, although as siliceous diatoms dominate the phytoplankton community (Ellwood et al., 2013; Eriksen et al., 2019). (Eriksen et al., 2019; Thompson et al., 2009). However, like nitrate and phosphate, silicate is also limited in this region, and is experiencing a decline (Ellwood et al., 2013; Thompson et al., 2009).

140

2.2 Methods Analysis

We collate the 10-years of ship-based hydrographic data and analyse the physical (temperature and salinity) and biogeochemical (oxygen and nutrients) properties between 0–200 m ~~dbar~~ across the longitudinal extent of the EAC mooring array. Mixed Layer depth (MLD) was calculated using density-based procedures developed by Holte and Talley (2009). This method uses an algorithm to choose the MLD from a suite of possible MLDs which are calculated using multiple threshold (difference from the surface) and gradient (where the depth-gradient exceeds a criteria) criteria. The threshold value of temperature and density for both the difference and gradient methods are 0.2°C and 0.03 kg m⁻³, and 0.005°C dbar⁻¹ and 0.0005 kg m⁻³ dbar⁻¹, respectively.

145

The hydrographic data are grouped by ~~season and the austral seasons of autumn (March–May), winter (June–August), and spring (September–November), and~~ by mode position of the EAC jet (Figure 2). For each day that a CTD station was occupied, we classified the EAC into an “inshore mode” or an “offshore mode” ~~using a qualitative description of the velocity profiles, based on the meridional velocities velocity profiles~~ and position of the EAC core from the mooring data. The ~~inshore mode~~




Figure 2. All-Full 10 years of hydrographic data points (coloured dots and lines) and grid (black lines) used for interpolated property sections. Distribution of nutrient data (a) grouped by season (austral autumn (March–May), winter (June–August), and spring (September–November)), and (b) EAC mode, and distribution of CTD data (c) grouped by seasons, and (d) and EAC mode.

was defined when the strongest southward flow was between 40–100 m (approximately the EAC core) and adjacent to, or “attached” to the continental slope; westward of approximately EAC is considered to be in an inshore mode when the EAC core is located westward of 154.2° E. The inshore mode is similar E (approximately over the continental slope, Figure 1 (b), bold white line), otherwise it is considered an offshore mode. Comparing the velocity structure of the “inshore” and “offshore” modes with Sloyan et al. (2016), we note that the inshore mode essentially corresponds to the climatological position of the EAC in this area southward velocities, which we have confirmed by taking composite averages of the meridional velocity across all days where a CTD station was occupied and where time steps are classified as inshore mode (Figure 3). The offshore




Figure 3. Panel (a) shows the cumulative sum of days observed for the inshore (blue) and offshore (purple) modes, considering only when a CTD station was occupied. Meridional velocity (m/s ; positive is equatorward) plotted for the upper 500 m of the water column from the daily EAC mooring data product for (ab) mean over all days with a CTD station. A one-day example for (c) inshore mode, and (bd) a one-day offshore mode. Also plotted is mean example of the EAC jet for all days classified as (e) inshore mode, and (ef) a one-day mean example of the EAC jet offshore mode. Velocity is plotted for the upper 500 m of the water column.

160 mode was defined when the strongest southward flow was “detached” from the continental slope, eastward of approximately 154.2° E and found over the abyssal plain or at the eastern edge of the mooring array. The inshore and offshore modes approximately match the mode 1 and mode (b) and (e) respectively). For the time period considered here, 57% of samples are categorised as belonging to the inshore mode. In contrast, the “offshore mode” approximately matches mode 2 velocities from Sloyan et al. (2016) from Sloyan et al. (2016) and diverges from the climatological position of the EAC.

165 After grouping the hydrographic profiles into by season and EAC jet position, we interpolate the temperature, salinity, oxygen and nutrient data to a 10 dbar vertical and 0.1° longitude grid (Figure 2) to produce property sections for each season

and (Figures 4 and 6) and EAC mode (Figures 2, 4, and 68 and 9). To test for statistical significance, a Monte Carlo simulation was run. For each ~~parameter~~variable, a random subset of data points were selected and used to make an interpolated pressure-longitude transect. This was repeated 1000 times and sections were constructed by selecting only the <5th and >95th percentile observations for each grid cell from the Monte Carlo. Stippling was added to all seasonal and mode sections for areas of statistical significance at the 95% confidence interval.

3 Results

3.1 Seasonality

We will first focus on the seasonal variability of the physical and biogeochemical variables in the EAC during austral autumn (March–May), winter (June–August), and spring (September–November).

The upper 200 m of the water column shows in-situ temperature ranges between \sim 15–25°C, practical salinity ranges between \sim 35–35.8 and ~~oxyen~~oxygen ranges between 160–230 $\mu\text{mol/L}$ for all seasons (Figure 4). In general, water is warmest, freshest and most oxygenated in the upper \sim 50 m and becomes cooler and reduced in oxygen with depth. The surface waters (0–100 mbar) are warmest in autumn, cooler during the winter period, and begin to rise slightly in temperature during spring. During both austral autumn and spring, there is a sharper gradient in both temperature, salinity, and oxygen between the surface and 200 mbar depth across the transect compared to winter. Oxygen concentrations above the mixed layer are lowest in autumn, increase in winter, and reach a maximum in spring. Below the mixed layer, the lowest oxygen concentrations are found near the western boundary and highest oxygen concentrations along the eastern edge of the section. Salinity is highly variable, with the near surface (above the mixed layer) having a minimum in autumn, increasing through winter, and reaching a maximum in spring. The subsurface salinity (50–150 mbar) has highest values in spring. The winter seasonal changes in temperature, salinity and oxygen have little ~~statistical significance~~significant divergence from the mean state, however, the seasonal extremes of warm, fresh, ~~oxygen poor~~low oxygen waters in autumn, and cool, saline, more oxygenated waters in spring are statistically significant, particularly in the near-surface.

There ~~was is~~ a slight seasonal cycle in MLD. ~~There was a moderate mixed layer in Autumn (mean MLD \sim 36.08 m in autumn)~~ (interquartile range (IQR) of 21.75–47.75 m, and 90th percentile 57.75 m), followed by a deepening in winter (mean MLD of 43.65 m, IQR of 16.50–62.75 m and 90th percentile of 85.80 m), and shoaling in spring (mean MLD of 25.40 m, IQR of 15.13–29.38 m and 90th percentile of 41.35 m).

The Temperature, Salinity, Oxygen (TS-O) diagrams show that the surface waters ($\rho_\theta < 26.5$, $\rho_\theta < 26.4$) are warmer with slightly lower salinity in autumn, cooling and freshening in winter, and increasing in temperature and salinity again in spring (Figure 5). Surface waters are generally oxygen-rich, with oxygen decreasing in the subsurface waters. There is a slight increase in oxygen in the surface waters in winter and spring, likely due to slight cooling in the temperature. All seasons show anomalous low oxygen concentration within the surface layer, these are particularly evident in ~~Autumn~~autumn (Figure 5).




Figure 4. Vertical sections of (a), (b), (c) *in-situ* temperature ($^{\circ}\text{C}$), (d), (e), (f) *practical* salinity, and (g), (h), (i) *oxygen* ($\mu\text{mol/L}$) for the upper 200 m dbar compiled from the 10 years (2012–2022) of data for austral autumn (March, April, May, left column), winter (June, July, August, middle column) and spring (September, October, November, right column). Solid lines *shows* mean MLD and dashed lines show the 90th percentile MLD averaged across longitude. Stippling indicates *statistically significant departures from the average obtained using all data* *statistical significance* following the procedure described in the *Data and Methods* *Analysis* section (*section 2.2*). Black shading represents the seafloor.

Nutrient concentrations are generally low at the surface (0–50 m dbar) and increase with depth (Figure 6). The ratio of nitrate to phosphate for 0–100 m dbar and 100–200 m dbar is 13.19 and 15.11, respectively. These values approximately agree with the Redfield ratio, particularly between 100–200 m dbar. There is evidence *for* *of* a subtle seasonal cycle in nutrients *when grouping together all data* in the upper 200 m dbar of the water column, *although at these depths the changes between seasons*




Figure 5. T-S-O Temperature-Salinity-Oxygen (TS-O) diagrams for austral (a) autumn, (c) winter, and (e) spring with oxygen concentrations coloured. Temperature-Oxygen (T-O) diagrams for austral (b) autumn (yellow), (d) winter (blue) and (f) spring (green). All Other CTD stations data are shown in (light grey dots) in each figure, and potential density isopycnals (kg m^{-3}) are shown in light grey shown as black dotted lines on T-S-O TS-O diagrams.

are subtle (as shown in Figure 7 (a)–(c)). Within each season, there is a large degree of variability. In general, autumn and spring have similar IQR, and winter has the smallest IQR for all nutrients sampled. This suggests winter has the lowest nutrient

Table 2. Number of days sampled in each season and mode. The percentage of days categorised in each mode for every season is shown in brackets, as well as total percentage number of days sampled in each mode and season.

Mode	Autumn	Winter	Spring	Total
Inshore	16 (38%)	17 (81%)	27 (64%)	60 (57%)
Offshore	26 (62%)	4 (19%)	15 (36%)	45 (43%)
Total	42 (40%)	21 (20%)	42 (40%)	105

205 concentrations available to the surface waters compared to the other seasons. Changes are broadly consistent across the three different nutrients, particularly in the variance and IQR. Although broadly similar, there are differences in the mean nutrient concentrations between the three nutrients with austral season. There is little change to ~~nitrato~~ the nitrate mean, although there is an increased number of observations with high concentration in autumn ~~and spring~~ compared to winter ~~and spring~~ (Figure 7 (a)). Phosphate experiences a slight increase ~~each season in the seasonal mean~~ from autumn through to spring (Figure 7 (b)).

210 Silicate has the largest mean in ~~winter and lowest in spring~~ autumn and winter and the lowest mean in spring (Figure 7 (c)).

The nutrients also exhibit a seasonal change with the longitude. The western edge of the section (westward of approximately 154° ~~E~~) has a similar distribution of nutrients throughout the water column during autumn and winter and a slight decrease in nutrient concentration during spring (Figure 6). The eastern section of the transect, particularly eastward of ~154.5° E, surface waters between 50–200 ~~m dbar~~ are more nutrient-rich during autumn, with reduced nutrient concentration during winter, and 215 show an increase in nutrient concentration again during spring, although with more variability (Figure 6).

3.2 Response of the ocean environment to EAC variability

Here, we present the physical and biogeochemical characteristics sorted by EAC inshore or offshore mode. The inshore mode (Figure 3 (b)–(c) and (e)) approximately follows the climatological position of the EAC, where meridional flow between 40–100 m is ~~approximately generally~~ $>0.4 \text{ m/s}$ in the poleward/southward direction and sits ~~on to over~~ the continental slope, with 220 the majority of the ~~core current~~ sitting westward of 154.2° E. The inshore mode has southward velocity over the continental slope and northward velocity further offshore at the eastern edge of the mooring array. In contrast, the EAC “offshore” mode (Figure 3 (e)–(d) and (f)) has an EAC jet that is displaced eastward from the continental slope and found to the east of ~154.3° E. In this “offshore” mode there is evidence for northward flow occurring inshore at depths below ~100 m. Both the inshore mode and the offshore mode occurred during all the seasons sampled, however, winter was dominated by ~~the~~ inshore mode, 225 whereas autumn and spring had a more even distribution of both modes (Table 2). ~~Additionally, all voyages experienced a “switch” between the two EAC modes, except for the 2016 voyages, during which only the inshore mode was observed.~~

Similarly to the seasonally grouped sections, ~~sections grouped into~~ EAC jet “inshore” or “offshore” mode sections ~~have~~ the warmest, freshest and most oxygenated water ~~is~~ found in the upper ~50 ~~m dbar~~, and sections cool and have reduced in oxygen concentrations with depth (Figure 8). ~~As mentioned~~ The TS-O diagram (not shown) reveals little discernible difference 230 between the two modes, as the modes have similar temperature and oxygen ranges.




Figure 6. Vertical sections of ((a), (b), (c)) **Nitrate** ($\mu\text{mol/L}$) **nitrate**, ((d), (e), (f)) **Phosphate** ($\mu\text{mol/L}$) **phosphate**, and ((g), (h), (i)) **Silicate** ($\mu\text{mol/L}$) **silicate** for the upper 200 m dbar compiled from the 10-years (2012–2022) of data for austral autumn (March, April, May, left column), winter (June, July, August, middle column) and spring (September, October, November, right column). Solid lines **shows** mean MLD and dashed lines show the 90th percentile MLD averaged across longitude. Stippling indicates **statistically significant departures from the average obtained using all data** **statistical significance** following the procedure described in the **Data and Methods** **Analysis** section (section 2.2). Black shading represents the seafloor.

As mentioned, the velocity profile of the inshore mode is similar to the climatological mean velocity profile of the EAC, thus, there **is** **are only** a few significant changes to the properties of the water column. The offshore mode is cooler **and more oxygenated** than the inshore mode across most of the section. However, for the western portion, **over the continental slope and** below the mixed layer, we find that the **off-shore** **offshore** mode is warmer, more saline, and more oxygenated than the




Figure 7. Box plots of nutrient concentrations across (a), (b), (c) the three seasons (Austral autumn, winter, and spring) and (d), (e), (f) the two modes sampled. Coloured dots are data points. Coloured boxes represent interquartile range (IQR), with the line inside the box representing the median (50th percentile). The whiskers (vertical black line) represents the range that contains 1.5 times the IQR above or below the first and third quartile.

235 corresponding locations of the inshore mode. These offshore mode property differences are generally significant for the portion of the section transect that is west of the EAC core. There ~~was~~is no difference in mean MLD between the two modes (35.77 m vs 34.61 m), or in variability variability of MLD (IQR of 17.75–48.50 m vs 15.38–48.75 m). The 90th percentiles were 63.00 m vs 65.10 m.

240 **Violin plots of nutrient concentrations across (a, b, c) the three seasons (Austral autumn (A), winter (W), and spring (S) and (d, e, f) the two modes sampled.**




Figure 8. Vertical sections of (a), (b) in-situ temperature ($^{\circ}\text{C}$), (c), (d) practical salinity, and (e), (f) oxygen ($\mu\text{mol/L}$) for the upper 200m dbar compiled from the 10 years (2012–2022) of data for the two EAC jet modes — inshore mode (left column), and offshore mode (right column). Solid lines shows mean MLD and dashed lines show the percentile MLD averaged across longitude. Dashed lines show the 0.6, 0.4, 0.25, 0.2, and 0.15 m/s contours of mean southward velocity for the corresponding EAC mode. Stippling indicates statistically significant departures from the average obtained using all data statistical significance following the procedure described in the Data and Methods Analysis section (section 2.2). Black shading represents the seafloor.

The TS-O diagram (not shown) reveal little discernible difference between the two modes. In general, the surface waters are more oxygen-rich with oxygen decreasing in the subsurface waters. The two modes have a similar temperature and oxygen ranges.

When considering changes in nutrient concentrations for all data ~~from across the top in the upper~~ 200 m-dbar of the water column ~~across all longitudes combined~~, there is ~~not a no~~ clear change in ~~mean~~ nutrient concentrations between the two modes, although the offshore mode has a larger IQR for all three nutrients sampled (Figure 7 d-f)(d)-(f)). The larger IQR in the offshore mode results from all nutrients having greater variability, reflecting a larger amount of samples with high nutrient concentration. This suggests a greater availability of nutrients in the upper 200 dbar when compared with the inshore mode. Similar to the changes across the three seasons, changes across the two modes are similar between the three nutrients. ~~Each mode has similar range and distribution of nutrient concentration with depth for all three nutrients sampled. However~~ Additionally, the differences in nutrient concentrations ~~become more clear~~ between the two modes are on the same order of magnitude as seasonal changes.

Differences in nutrient concentrations become clearer when considering the effect of depth and longitude (Figure 9).

The nutrient sections based on EAC jet location show that generally nutrient poor water is found above the mixed layer during the inshore mode, and the highest nutrient concentrations occur at the western edge of the section at a pressure of approximately 150 m-depth dbar (Figure 9). For the offshore mode, waters are generally more nutrient-rich. Whilst there are not widespread statistically significant increases in nutrient concentrations between the two modes and the mean state, there is evidence for an increase in the concentrations of nitrate and phosphate above and near the mixed layer during the offshore mode when compared to the inshore mode.

260 4 Discussion

In this study, we use physical and biogeochemical data collected over a 10-year period to determine an offshore seasonal cycle in nutrient concentrations, and reveal the effect of the varying position of the EAC on the availability of nutrients to the surface waters in the upstream EAC system. The results suggest that ~~nitrate~~there is a seasonal cycle in biogeochemical properties. Nitrate, phosphate, and silicate peak in concentration in autumn and spring below the mixed layer, but these are associated with different oxygen concentrations. There is a low-oxygen high nutrient peak in autumn and a higher-oxygen and nutrient peak in early spring. Previous observations from coastal sites have found a spring peak in nutrients at 29°S (Everett et al., 2014) and also at the North Stradbroke Island National Reference Site ~~(at 27°S; (Butler et al., 2020))~~S (Butler et al., 2020). They also suggested an early winter minimum in silicate. Additionally, Rocha et al. (2019) found that the nutricline deepens by approximately 50 m during the summer, reducing availability of nutrients to the surface waters.

270 While we ~~also find observe~~ a seasonal nutrient signal, we ~~show an interesting impact of the position also highlight the influence~~ of the EAC core's position relative to the continental shelf/slope. The ~~different modes in the EAC jet did not alter the average properties of the water column, but they did impact the position of the EAC affects the~~ vertical distribution of nutrients in the upper water column, and longitudinally and their longitudinal gradients along the transect line. For both modes, ~~we see evidence of upwelling, with of~~. For the inshore mode, cooler, fresher, oxygen poor, and nutrient-rich water ~~being found~~ closer to the surface. For the inshore mode, this upwelling signature occurs on the western (inshore) edge of the property sections (west of 154°E), ~~whereas for the trapped between the EAC and continental slope. Whilst there is a statistical increase~~




Figure 9. Vertical sections of (a), (b) Nitrate (umol/L)nitrate, (c), (d) Phosphate (umol/L)phosphate, and (e), (f) Silicate (umol/L)silicate for the upper 200m dbar compiled from the 10-years (2012–2022) of data for the two EAC jet modes — inshore mode (left column), and offshore mode (right column). Solid lines shows show mean MLD and dashed lines show the percentile MLD averaged across longitude. Dashed lines show the 0.6, 0.4, 0.25, 0.2, and 0.15 m/s contours of mean southward velocity for the corresponding EAC mode. Stippling indicates statistically significant departures from the average obtained using all data statistical significance following the procedure described in the Data and Methods Analysis section (section 2.2). Black shading represents the seafloor.

in nutrient concentrations, due to the depth (150 dbar) and presence of the EAC jet, these nutrients would be inaccessible to the surface layer. For the offshore mode, the upwelling signature is found off-shore there is an uplift of nutrients across the transect, particularly offshore at $\sim 154.7^{\circ}$ –E. Additionally, while the MLD is similar between the two modes, meaning that




Figure 10. The EAC velocity anomaly of the first two modes of the Complex Empirical Orthogonal Function (CEO) analysis. For the leading two modes, the spatial maps of (a), (d) phase ($-\pi - \pi$) and (b), (e) amplitude and (c), (f) reconstructed velocity anomaly (m/s) are shown. The inshore mode approximately matches the CEO mode 1 (a), (b), (c), which accounts for 65.4% of velocity anomaly. The CEO mode 2 (d), (e), (f) fits the offshore mode, and accounts for 13.3% of velocity anomaly. The reconstructed velocity anomaly for each mode is determined from the spatial amplitude and phase maps.

280 ~~upwelling in the offshore mode results in higher concentrations of nutrients there is a shoaling of the nutricline and an increase in nutrients closer to and entering the mixed layer during the while the EAC is in the~~ offshore mode compared to the inshore mode. The effect of the EAC core location highlights a potential mechanism for episodic nutrient supply to the surface layer, as both the inshore mode and offshore mode were observed in all three seasons. A complex empirical orthogonal function (CEO) analysis of the southward velocity component shows that the onshore and offshore EAC modes explain greater than 285 78% of the velocity variance with periods of 120-60 days (Figure 10). The first two CEO modes explain the wavering of the EAC across the mooring array, with the first mode showing the EAC located near the continental shelf, flowing over the abyssal plain, accounting for 65.4% of the velocity anomaly. The second mode shows an EAC detached from the continental shelf and accounts for 13.3% of the velocity anomaly. This highlights the variability in the EAC, separate from seasonality and large scale dynamics.

290 ~~The upwelling that occurs-~~

~~The distribution of nutrients that occur~~ in the two EAC jet modes may be connected to different drivers. The encroachment of the EAC towards the continental shelf can cause upwelling along the slope (Schaeffer et al., 2013). Additionally, increasing EAC speeds has previously been linked to upwelling (Archer et al., 2017a; Oke and Middleton, 2001; Roughan and Middleton,

2002). In the Kuroshio, increased speeds where the current sits close to the shelf in its upstream region causes a similar uplift of nutrients onshore (Chen et al., 2022). As such, the inshore mode of the EAC – characterised by high speeds and a position close to the continental shelf – may still experience some upwelling of nutrient rich waters, however this upwelling will be limited to the near slope/shelf area. This is compared to the offshore mode, where upwelling could be linked to eddy interactions and sub-mesoscale circulations (e.g., Roughan et al., 2017; Roughan and Middleton, 2004; Suthers et al., 2011), meaning upwelling can occur over a broader longitudinal range, trapped by the EAC position.

300 The drivers of the shift in the position of the EAC core, related to the offshore mode, are yet to be determined. Early studies implied seasonality plays a role in jet meandering (Ridgway and Godfrey, 1997), while more recent studies found seasonality plays no role in the core position (Archer et al., 2017a). It may be related to the energy of the EAC system, as a higher energy jet separates further north and creates larger scale meandering displays different dynamics to a low energy jet (Li et al., 2021). However, none of the previous studies have not been able to attribute EAC wavering or meandering to one clear cause (Archer 305 et al., 2017a; Bowen et al., 2005; Sloyan et al., 2016). Despite this, the effects of mesoscale eddy interactions are heavily implicated, as there are strong interactions between eddies and the mean flow path of the EAC (Li and Roughan, 2023).

Similar variability has been observed in other WBCs, for example the Gulf Stream (Mao et al., 2023), the Florida Current (Archer et al., 2018), Agulhas Current (Goschen et al., 2015), and the Brazil Current (Da Silveira et al., 2008). The Kuroshio Current also experiences variability in its core position and strength, and experiences meandering and interactions with 310 bathymetry and mesoscale eddies (Ebuehi and Hanawa, 2003; Kawabe, 2005, 1995; Waseda et al., 2003). South of Japan, the Kuroshio Current follows three main flow paths, flow following the coastline, flow that separates from the coast resulting in offshore flow, and finally, a large meander mode that greatly changes the current position (Kawabe, 1985, 1995). Additionally, in the upstream region of the Kuroshio near Taiwan, it has been found that increased speed of the Kuroshio current causes an increased uplift of nutrients onshore (Chen et al., 2022). This is similar to the inshore position In the process of classifying 315 EAC modes for this study, it was noted that the offshore mode is often associated with a cyclonic eddy sitting on the inshore side of the EAC, where there are faster speeds near the continental shelf and upwelling of nutrients along the slope. Additionally, periods where the Kuroshio sits closer to the coast, it interacts with bathymetry which causes strong vertical mixing and uplift of nutrients to the continental shelf (Durán Gómez and Nagai, 2022). The Florida Current is upstream of the Gulf Stream, and has a very similar jet structure to the EAC when considering jet coordinates (Archer et al., 2018), 320 and exhibits a similar meandering behaviour, with meanders occurring on a time-scale of 3–30 days (Archer et al., 2017b). Mesoscale variability in the Florida Current is mainly linked to two mechanisms, frontal eddies and current meandering, both of which are linked to the upwelling of cool, nutrient rich waters between the shelf break and the offshore meander (Fiechter and Mooers, 2007; Kourafalou and Kang, 2012). Upwelling has also been linked to the movement of jet meanders in the Agulhas Current, with upwelling occurring when the current shifts onto the continental shelf, or shifts offshore (Goschen et al., 2015) 325. It is clear that meandering of the current upstream of the separation point is not unique to the EAC, and the dynamics of WBC meandering, including the identification of upwelling, has been observed in several WBC systems. However, we still lack a detailed understanding of the impact of current meandering on the biogeochemistry of the upper ocean and continental shelf, similar to what was observed by Chapman et al. (2024). As such, upwelling is likely linked to oceanic internal processes, such

as eddy interactions and sub-mesoscale circulations (e.g., Roughan et al., 2017; Roughan and Middleton, 2004; Suthers et al., 2011; Chapman et al., 2019) rather than direct control by winds. Topographic interaction could play a role, particularly during the inshore mode, however, further investigation is needed.

Previous work has found existing relationships between ENSO and EAC transport (Holbrook et al., 2011). Though we do not present the results here, we found that the position of the EAC and El Niño–Southern Oscillation (ENSO) Index (SOI), using a linear model, to be highly correlated. Periods of El Niño (positive SOI) showed increased occurrences of the EAC offshore mode nutricline (the maximum absolute vertical gradient in nutrient concentrations) during the offshore mode compared to the inshore mode (Figure 11). This effect is observable in Figure 9 as the inshore mode has a low vertical gradient across the majority of the transect, compared to the offshore mode, which has a sharper gradient in nutrients observable within the top 200 dbar. This effect of a shoaling nutricline was also observed in the Kuroshio “large meander” mode, which combined with winter convective mixing, made nutrients accessible to the surface waters (Hayashida et al., 2023). However, a full analysis of the connection between ENSO and EAC current position could not be conducted due to the limited duration of the dataset. It is worth noting that even when considering other long datasets, only a handful of strong ENSO events have occurred in conjunction with comprehensive oceanographic and mooring data; therefore there is limited data to robustly test these hypotheses. So, while there is some evidence for a link between EAC meandering and ENSO, longer timeseries data is needed to understand the mechanisms of such a connection. This area within the upstream EAC does not experience strong winter convective mixing and is characterised by relatively small seasonal changes in MLD. We found a maximum mean MLD of 40 m in winter, similar to what has previously been observed (Sobral et al., 2024; Condie and Dunn, 2006). Previous studies found minimum MLDs in summer of 20 m (Sobral et al., 2024) and 30 m (Condie and Dunn, 2006). This indicates that the MLD changes by only 10–20 m throughout the year. As such, convective mixing has low potential for replenishing mixed layer nutrients. Instead, we find that the properties of the water column are primarily being influenced by EAC variability.

The Southwest Pacific surface waters are oligotrophic, and plankton in this area are nitrate and phosphate limited (Ellwood et al., 2013; Ustick et al., 2021). The upwelling of nutrient rich waters, observed in this study, can cause plankton blooms (Silsbe and Malkin, 2016) and has implications for ecosystem function (Brander et al., 2003; Hays et al., 2005; McGillicuddy Jr, 2016; Richardson and Schoeman, 2004). For example, data from a single voyage (IN2019_V05 from Table 1) showed uplift of deeper waters and upwelling velocities along the strongly tilted isopycnals which form the inshore flank of the EAC (Chapman et al., 2024). When the EAC was in the offshore position, this uplift was associated with surface phytoplankton blooms and increased zooplankton biomass (Chapman et al., 2024). Whilst we have observed mechanisms which lead to an increase of nutrients in the upper water column, exploring biological responses to such nutrient changes is out of the scope of this paper. Future work should explore measurements of the deep chlorophyll-a maximum or the plankton communities observed during these research voyages to determine the productivity response to nutrient upwelling. These relationships further with sampling and high-resolution biogeochemical models. The improved understanding of biogeochemical dynamics and the associated plankton response will improve our ability to understand how marine ecosystems may respond to current meandering future variability in the EAC system.




Figure 11. Nutricline depth (dots) for CTD casts for (a) inshore and (b) offshore mode. Nutricline depth was calculated as the maximum vertical gradient of full-depth bottle samples for nitrate. Blue line is polynomial robust line of fit for calculated nutricline depths. A lowess line of fit was also plotted and agreed well with the robust line of fit. Dashed black lines show the 0.6, 0.4, 0.25, 0.2, and 0.15 m/s contours of mean southward velocity for each corresponding mode. Bold red line shows mean mixed layer depth and dashed red line shows 90th percentile mixed layer depth. Grey shading represents the seafloor.

5 Conclusions

This study systematically assesses the nutrient variability in the EAC intensification region, and is one of the few studies
365 focusing on waters beyond the coastal and shelf environment. We find seasonal variability in the temperature, salinity, oxygen, and nutrients of the EAC, and importantly we identify the influence of EAC jet meandering on the ~~location of the upwelled nutrients. Furthermore, we categorized uplift of nutrients. We categorised~~ the EAC into two distinct modes based on the position of the current relative to the continental shelf and slope — the inshore and offshore modes. The ~~inshore mode is generally low in nutrients, with high nutrient waters trapped and concentrated between the EAC jet the continental slope. The~~ offshore mode,
370 when the EAC jet sits over the abyssal plain, exhibits ~~a shoaled nutricline, with~~ nutrient-enriched waters ~~and extending further offshore. Meanwhile the inshore mode shows upwelling of nutrients along the edge of~~ reaching the mixed layer in the offshore region. Whilst we cannot attribute these changes to any one previous hypothesis of EAC variability, we see that the position and
375 strength of the continental shelf. ~~Understanding what drives the position of EAC influences the supply of nutrients to the upper ocean, likely due to oceanic processes such as eddy interactions. Understanding the EAC and the nutrient dynamics response of nutrients to the movement of the EAC jet~~ contributes to our knowledge of the EAC's role in supporting and shaping the biological communities and productivity within this region.

Author contributions. BS and CC planned and led voyages and supplied data. MJ undertook the analysis and prepared the original paper draft with contributions from CC, BS and HB. All authors contributed to preparation of the final published paper.

Competing interests. At least one of the (co-)authors is a member of the editorial board of Ocean Science.

380 *Acknowledgements.* We acknowledge the use of the CSIRO Marine National Facility (<https://ror.org/01mae9353>) and grant of sea time on RV ~~Investigator~~ Investigator in undertaking this research. ~~CSIRO Environment and IMOS funded the collection of the EAC mooring array. IMOS is enabled by the National Collaborative Research Infrastructure Strategy (NCRIS). CSIRO Environment funds were used to create the EAC data products. The authors would also like to acknowledge the hydro-chemistry teams from all voyages that undertook nutrient analysis, and the ship's crew, officers and MNF support staff for their support during various field campaigns.~~

Archer, M., Keating, S., Roughan, M., Johns, W., Lumpkin, R., Beron-Vera, F., and Shay, L.: The kinematic similarity of two western boundary currents revealed by sustained high-resolution observations, *Geophysical research letters*, 45, 6176–6185, 2018.

Archer, M. R., Roughan, M., Keating, S. R., and Schaeffer, A.: On the variability of the East Australian Current: Jet structure, meandering, and influence on shelf circulation, *Journal of Geophysical Research: Oceans*, 122, 8464–8481, 2017a.

390 Archer, M. R., Shay, L. K., and Johns, W. E.: The surface velocity structure of the Florida Current in a jet coordinate frame, *Journal of Geophysical Research: Oceans*, 122, 9189–9208, 2017b.

Blackburn, S. and Cresswell, G.: A coccolithophorid bloom in Jervis Bay, Australia, *Marine and Freshwater Research*, 44, 253–260, 1993.

Boland, F. and Church, J.: The East Australian Current 1978, *Deep Sea Research Part A. Oceanographic Research Papers*, 28, 937–957, 1981.

395 Bowen, M. M., Wilkin, J. L., and Emery, W. J.: Variability and forcing of the East Australian Current, *Journal of Geophysical Research: Oceans*, 110, 2005.

Brander, K., Dickson, R., and Edwards, M.: Use of Continuous Plankton Recorder information in support of marine management: applications in fisheries, environmental protection, and in the study of ecosystem response to environmental change, *Progress in Oceanography*, 58, 175–191, 2003.

400 Bull, C. Y. S., Kiss, A. E., Gupta, A. S., Jourdain, N. C., Argüeso, D., Di Luca, A., and Sérazin, G.: Regional versus remote atmosphere-ocean drivers of the rapid projected intensification of the East Australian Current, *Journal of Geophysical Research: Oceans*, 125, e2019JC015889, <https://doi.org/10.1029/2019JC015889>, 2020.

Butler, E., Skuza, M., and Lønborg, C.: Spatial and temporal trends in concentrations of nutrients, 2020.

Cetina-Heredia, P., Roughan, M., van Sebille, E., and Coleman, M. A.: Long-term trends in the East Australian Current separation latitude and 405 eddy driven transport, *Journal of Geophysical Research: Oceans*, 119, 4351–4366, <https://doi.org/10.1002/2014JC010071>, 2014.

Chapman, C. C., Sloyan, B. M., Schaeffer, A., Suthers, I. M., and Pitt, K. A.: Offshore plankton blooms via mesoscale and sub-mesoscale interactions with a western boundary current, *Journal of Geophysical Research: Oceans*, 129, e2023JC020547, 2024.

Chen, C.-C., Lu, C.-Y., Jan, S., Hsieh, C.-h., and Chung, C.-C.: Effects of the coastal uplift on the Kuroshio ecosystem, Eastern Taiwan, the 410 western boundary current of the North Pacific Ocean, *Frontiers in Marine Science*, 9, 796 187, 2022.

Chiswell, S. M., Bostock, H. C., Sutton, P. J., and Williams, M. J.: Physical oceanography of the deep seas around New Zealand: a review, *New Zealand Journal of Marine and Freshwater Research*, 49, 286–317, <https://doi.org/10.1080/00288330.2014.992918>, 2015.

Condie, S. A. and Dunn, J. R.: Seasonal characteristics of the surface mixed layer in the Australasian region: implications for primary production regimes and biogeography, *Marine and Freshwater Research*, 57, 569–590, 2006.

415 Da Silveira, I., Lima, J., Schmidt, A., Ceccopieri, W., Sartori, A., Franscisco, C., and Fontes, R.: Is the meander growth in the Brazil Current system off Southeast Brazil due to baroclinic instability?, *Dynamics of Atmospheres and Oceans*, 45, 187–207, 2008.

Doblin, M. A., Petrou, K., Sinutok, S., Seymour, J. R., Messer, L. F., Brown, M. V., Norman, L., Everett, J. D., McInnes, A. S., Ralph, P. J., et al.: Nutrient uplift in a cyclonic eddy increases diversity, primary productivity and iron demand of microbial communities relative to a western boundary current, *PeerJ*, 4, e1973, 2016.

420 Durán Gómez, G. S. and Nagai, T.: Elevated Nutrient Supply Caused by the Approaching Kuroshio to the Southern Coast of Japan, *Frontiers in Marine Science*, 9, 842 155, 2022.

Ebuchi, N. and Hanawa, K.: Influence of mesoscale eddies on variations of the Kuroshio path south of Japan, *Journal of oceanography*, 59, 25–36, 2003.

Ellwood, M. J., Law, C. S., Hall, J., Woodward, E. M. S., Strzepek, R., Kuparinen, J., Thompson, K., Pickmere, S., Sutton, P., and Boyd, P. W.: Relationships between nutrient stocks and inventories and phytoplankton physiological status along an oligotrophic meridional transect in the Tasman Sea, *Deep Sea Research Part I: Oceanographic Research Papers*, 72, 102–120, 2013.

Eriksen, R. S., Davies, C. H., Bonham, P., Coman, F. E., Edgar, S., McEnnulty, F. R., McLeod, D., Miller, M. J., Rochester, W., Slotwinski, A., et al.: Australia's long-term plankton observations: the integrated marine observing system national reference station network, *Frontiers in Marine Science*, p. 161, 2019.

430 Everett, J., Baird, M., and Suthers, I.: Three-dimensional structure of a swarm of the salp *Thalia democratica* within a cold-core eddy off southeast Australia, *Journal of Geophysical Research: Oceans*, 116, 2011.

Everett, J. D., Baird, M. E., Roughan, M., Suthers, I. M., and Doblin, M. A.: Relative impact of seasonal and oceanographic drivers on surface chlorophyll a along a Western Boundary Current, *Progress in Oceanography*, 120, 340–351, 2014.

Fiechter, J. and Mooers, C.: Primary production associated with the Florida Current along the East Florida Shelf: Weekly to seasonal variability from mesoscale-resolution biophysical simulations, *Journal of Geophysical Research: Oceans*, 112, 2007.

435 Figueira, W. F. and Booth, D. J.: Increasing ocean temperatures allow tropical fishes to survive overwinter in temperate waters, *Global Change Biology*, 16, 506–516, 2010.

Ganachaud, A., Cravatte, S., Melet, A., Schiller, A., Holbrook, N., Sloyan, B., Widlansky, M., Bowen, M., Verron, J., Wiles, P., et al.: The Southwest Pacific Ocean circulation and climate experiment (SPICE), *Journal of Geophysical Research: Oceans*, 119, 7660–7686, 2014.

440 Gibbs, M. T., Middleton, J. H., and Marchesiello, P.: Baroclinic response of Sydney shelf waters to local wind and deep ocean forcing, *Journal of Physical Oceanography*, 28, 178–190, 1998.

Godfrey, J., Cresswell, G., Golding, T., Pearce, A., and Boyd, R.: The separation of the East Australian Current, *Journal of Physical Oceanography*, 10, 430–440, 1980.

Goschen, W., Bornman, T., Deyzel, S., and Schumann, E.: Coastal upwelling on the far eastern Agulhas Bank associated with large meanders 445 in the Agulhas Current, *Continental Shelf Research*, 101, 34–46, 2015.

Hassler, C., Djajadikarta, J., Doblin, M., Everett, J., and Thompson, P.: Characterisation of water masses and phytoplankton nutrient limitation in the East Australian Current separation zone during spring 2008, *Deep Sea Research Part II: Topical Studies in Oceanography*, 58, 664–677, 2011.

450 Hayashida, H., Kiss, A. E., Miyama, T., Miyazawa, Y., and Yasunaka, S.: Anomalous Nutricline Drives Marked Biogeochemical Contrasts During the Kuroshio Large Meander, *Journal of Geophysical Research: Oceans*, 128, e2023JC019697, 2023.

Hays, G. C., Richardson, A. J., and Robinson, C.: Climate change and marine plankton, *Trends in ecology & evolution*, 20, 337–344, 2005.

Holbrook, N. J., Goodwin, I. D., McGregor, S., Molina, E., and Power, S. B.: ENSO to multi-decadal time scale changes in East Australian Current transports and Fort Denison sea level: Oceanic Rossby waves as the connecting mechanism, *Deep Sea Research Part II: Topical Studies in Oceanography*, 58, 547–558, 2011.

455 Holte, J. and Talley, L.: A new algorithm for finding mixed layer depths with applications to Argo data and Subantarctic Mode Water formation, *Journal of Atmospheric and Oceanic Technology*, 26, 1920–1939, 2009.

Kawabe, M.: Sea level variations at the Izu Islands and typical stable paths of the Kuroshio, *Journal of the Oceanographical Society of Japan*, 41, 307–326, 1985.

Kawabe, M.: Variations of current path, velocity, and volume transport of the Kuroshio in relation with the large meander, *Journal of physical oceanography*, 25, 3103–3117, 1995.

Kawabe, M.: Variations of the Kuroshio in the southern region of Japan: Conditions for large meander of the Kuroshio, *Journal of oceanography*, 61, 529–537, 2005.

Kessler, W. S. and Cravatte, S.: Mean circulation of the Coral Sea, *Journal of Geophysical Research: Oceans*, 118, 6385–6410, 2013.

Kourafalou, V. H. and Kang, H.: Florida Current meandering and evolution of cyclonic eddies along the Florida Keys Reef Tract: Are they interconnected?, *Journal of Geophysical Research: Oceans*, 117, 2012.

465 Li, J. and Roughan, M.: Energetics of Eddy–Mean Flow Interactions in the East Australian Current System, *Journal of Physical Oceanography*, 53, 595–612, 2023.

Li, J., Roughan, M., and Kerry, C.: Dynamics of interannual eddy kinetic energy modulations in a western boundary current, *Geophysical Research Letters*, 48, e2021GL094 115, 2021.

470 Mao, S., He, R., Bane, J., Gawarkiewicz, G., and Todd, R. E.: A data-assimilative modeling investigation of Gulf Stream variability, *Deep Sea Research Part II: Topical Studies in Oceanography*, 211, 105 319, 2023.

Martiny, A. C., Lomas, M. W., Fu, W., Boyd, P. W., Chen, Y.-l. L., Cutter, G. A., Ellwood, M. J., Furuya, K., Hashihama, F., Kanda, J., et al.: Biogeochemical controls of surface ocean phosphate, *Science Advances*, 5, eaax0341, 2019.

McGillicuddy Jr, D. J.: Mechanisms of physical-biological-biogeochemical interaction at the oceanic mesoscale, *Annual Review of Marine Science*, 8, 125–159, 2016.

475 Nieblas, A.-E., Sloyan, B. M., Hobday, A. J., Coleman, R., and Richardsone, A. J.: Variability of biological production in low wind-forced regional upwelling systems: A case study off southeastern Australia, *Limnology and Oceanography*, 54, 1548–1558, 2009.

Oke, P. R. and Middleton, J. H.: Topographically induced upwelling off eastern Australia, *Journal of Physical Oceanography*, 30, 512–531, 2000.

480 Oke, P. R. and Middleton, J. H.: Nutrient enrichment off Port Stephens: The role of the East Australian Current, *Continental Shelf Research*, 21, 587–606, 2001.

Oke, P. R., Roughan, M., Cetina-Heredia, P., Pilo, G. S., Ridgway, K. R., Rykova, T., Archer, M. R., Coleman, R. C., Kerry, C. G., Rocha, C., et al.: Revisiting the circulation of the East Australian Current: Its path, separation, and eddy field, *Progress in Oceanography*, 176, 102 139, 2019.

485 Olson, D. B.: Biophysical dynamics of western transition zones: A preliminary synthesis, *Fisheries Oceanography*, 10, 133–150, 2001.

Richardson, A. J. and Schoeman, D. S.: Climate impact on plankton ecosystems in the Northeast Atlantic, *Science*, 305, 1609–1612, 2004.

Ridgway, K. and Dunn, J.: Mesoscale structure of the mean East Australian Current System and its relationship with topography, *Progress in oceanography*, 56, 189–222, 2003.

Ridgway, K. and Godfrey, J.: Seasonal cycle of the East Australian Current, *Journal of Geophysical Research: Oceans*, 102, 22 921–22 936, 490 1997.

Rocha, C., Edwards, C. A., Roughan, M., Cetina-Heredia, P., and Kerry, C.: A high-resolution biogeochemical model (ROMS 3.4+ bio_Fennel) of the East Australian Current System, *Geoscientific Model Development*, 12, 441–456, 2019.

Rochford, D.: Nutrient enrichment of east Australian coastal waters. II. Laurieton upwelling, *Marine and Freshwater Research*, 26, 233–243, 1975.

495 Roughan, M. and Middleton, J. H.: A comparison of observed upwelling mechanisms off the east coast of Australia, *Continental Shelf Research*, 22, 2551–2572, 2002.

Roughan, M. and Middleton, J. H.: On the East Australian Current: Variability, encroachment, and upwelling, *Journal of Geophysical Research: Oceans*, 109, 2004.

Roughan, M., Keating, S., Schaeffer, A., Cetina Heredia, P., Rocha, C., Griffin, D., Robertson, R., and Suthers, I.: A tale of two eddies: The 500 biophysical characteristics of two contrasting cyclonic eddies in the East Australian Current System, *Journal of Geophysical Research: Oceans*, 122, 2494–2518, 2017.

Roughan, M., Cetina-Heredia, P., Ribbat, N., and Suthers, I. M.: Shelf transport pathways adjacent to the East Australian Current reveal 505 sources of productivity for coastal reefs, *Frontiers in Marine Science*, 8, 789 687, 2022a.

Roughan, M., Hemming, M., Schaeffer, A., Austin, T., Beggs, H., Chen, M., Feng, M., Galibert, G., Holden, C., Hughes, D., et al.: Multi- 510 decadal ocean temperature time-series and climatologies from Australia's long-term National Reference Stations, *Scientific Data*, 9, 157, 2022b.

Schaeffer, A., Roughan, M., and Morris, B. D.: Cross-shelf dynamics in a western boundary current regime: Implications for upwelling, *Journal of Physical Oceanography*, 43, 1042–1059, 2013.

Schaeffer, A., Roughan, M., Jones, E. M., and White, D.: Physical and biogeochemical spatial scales of variability in the East Australian 515 Current separation from shelf glider measurements, *Biogeosciences*, 13, 1967–1975, 2016.

Silsbe, G. M. and Malkin, S. Y.: Where light and nutrients collide: The global distribution and activity of subsurface chlorophyll maximum layers, *Aquatic microbial ecology and biogeochemistry: A dual perspective*, pp. 141–152, 2016.

Sloyan, B. M. and O'Kane, T. J.: Drivers of decadal variability in the Tasman Sea, *Journal of Geophysical Research: Oceans*, 120, 3193–3210, 520 <https://doi.org/https://doi.org/10.1002/2014JC010550>, 2015.

Sloyan, B. M., Ridgway, K. R., and Cowley, R.: The East Australian Current and property transport at 27°S from 2012 to 2013, *Journal of Physical Oceanography*, 46, 993–1008, 2016.

Sloyan, B. M., Chapman, C. C., Cowley, R., and A, C. A.: Application of machine learning techniques to ocean mooring time series data, *Journal of Atmospheric and Oceanic Technology*, 40, 241–260, 2023.

Sloyan, B. M., Cowley, R., and Chapman, C. C.: East Australian Current velocity, temperature and salinity data products, *Scientific Data*, 11, 525 10, 2024.

Sobral, F., Roughan, M., Malan, N., and Li, J.: Mean-state and seasonal variability in temperature structure and heat transport in the East Australian Current System from a multi-decadal regional ocean model, *Journal of Geophysical Research: Oceans*, 129, e2023JC020438, 2024.

Suthers, I. M., Young, J. W., Baird, M. E., Roughan, M., Everett, J. D., Brassington, G. B., Byrne, M., Condie, S. A., Hartog, J. R., Hassler, 530 C. S., et al.: The strengthening East Australian Current, its eddies and biological effects—an introduction and overview, 2011.

Suthers, I. M., Schaeffer, A., Archer, M., Roughan, M., Griffin, D. A., Chapman, C. C., Sloyan, B. M., and Everett, J. D.: Frontal eddies provide an oceanographic triad for favorable larval fish habitat, *Limnology and Oceanography*, 2023.

Sutton, P. J. and Bowen, M.: Flows in the Tasman front south of Norfolk island, *Journal of Geophysical Research: Oceans*, 119, 3041–3053, 535 2014.

Thompson, P., Baird, M., Ingleton, T., and Doblin, M.: Long-term changes in temperate Australian coastal waters: implications for phytoplankton, *Marine Ecology Progress Series*, 394, 1–19, 2009.

Tranter, D. J., Carpenter, D. J., and Leech, G. S.: The coastal enrichment effect of the East Australian Current eddy field, *Deep Sea Research Part A. Oceanographic Research Papers*, 33, 1705–1728, 1986.

Ustick, L. J., Larkin, A. A., Garcia, C. A., Garcia, N. S., Brock, M. L., Lee, J. A., Wiseman, N. A., Moore, J. K., and Martiny, A. C.:
535 Metagenomic analysis reveals global-scale patterns of ocean nutrient limitation, *Science*, 372, 287–291, 2021.
Waseda, T., Mitsudera, H., Taguchi, B., and Yoshikawa, Y.: On the eddy–Kuroshio interaction: Meander formation process, *Journal of
Geophysical Research: Oceans*, 108, 2003.