

Observed bottom warming in the East Siberian Sea driven by the intensified vertical mixing

Xiaoyu Wang^{1, 2, 3}, Longjiang Mu³, and Xianyao Chen^{1, 2, 3,*}

¹ Frontier Science Center for Deep Ocean Multispheres and Earth System, Ocean University of China, Qingdao, China

² Physical Oceanography Laboratory, Ocean University of China, Qingdao, China

³ Laoshan Laboratory, Qingdao, China

Correspondence to: Xianyao Chen (chenxy@ouc.edu.cn)

Abstract. The East Siberian Sea (ESS) features the broadest continental shelf on Earth and contains nearly 80% of the world's subsea permafrost. A persistent cold bottom layer, with temperatures at freezing point, inhibits the upward transport of heat, thus preventing the thawing of permafrost and subsequent methane release from sediments. However, in early September 2016, we observed an unprecedented warming of over 3°C at the seabed, approximately 46 meters deep in the ESS, following a relatively moderate Arctic cyclone. We attribute this notable bottom warming to enhanced wave-induced vertical mixing, which facilitates uniform stirring of the Arctic marginal seas and allows surface heat to reach the bottom layer. As sea ice continues to retreat in the Arctic continental shelf, wind-driven waves have increased space to develop. Consequently, even moderate cyclones can trigger substantial vertical mixing, a phenomenon not previously documented. Given the accelerated warming of the Arctic and the rapid decline of sea ice, we anticipate that more open water will foster the growth of larger wind-driven waves and intensified vertical mixing, leading to greater heat influx to the bottom layers of Arctic shelves in the future.

1 Introduction

The East Siberian Arctic shelf includes the Laptev Sea, the East Siberian Sea, and the Russian side of the Chukchi Sea. It is the broadest ($\sim 2.1 \times 10^6$ km 2) and shallowest (mean depth < 50 m) shelf in the world ocean, with nearly 80% of existing subsea permafrost and up to 50% methane flux of the global coastal seas area (Shakhova et al., 2019; Sayedi et al., 2020). In summer, the seafloor of the East Siberian Arctic shelf is generally covered by cold bottom waters (< 1.0 °C) with several to tens of meters thick (Dmitrenko et al., 2010; Bauch & Cherniavskaya, 2018; Wang et al., 2021). This cold layer acts as a barrier to reduce the methane emissions from sediments by about half (Ferré et al., 2020; Altuna et al., 2021).

However, recent observations show that the near-bottom water over the Laptev Sea mid-shelf has exhibited pronounced warming due to the increasing downward heat transport during the retreat of the surface sea ice since mid-1980s (Janout & Lenn, 2014; Janout et al., 2016). In the later summer of 2007, the wind-induced vertical mixing caused a dramatic and abrupt near-bottom warming of about 3.0°C on the Laptev Sea mid-shelf (30-m water depth) (Hölemann et al., 2011).

The warming impacts on bottom water temperature anomaly remained for up to 3 months after the wind event, leading to the

删除了：The East Siberian Sea has the broadest continental shelf on Earth and nearly 80% of the subsea permafrost worldwide. There exists a cold layer with the temperature at freezing points around -1.5°C above the sea floor of the shelf, preventing the heat transport from above to melt the permafrost and release the methane from sediments. However, we observed a worrying warming trend at the seafloor caused by enhanced vertical mixing in the shelf of the East Siberian Sea. In an ice-reduced Arctic continental shelf, even a moderate cyclone can result in the rapid growth of high wave to stir the marginal sea uniformly, which is not observed before. The intensified mixing can transport enormous heat downward, leading to a remarkable warming of more than 3°C at the bottom. As the Arctic is experiencing accelerated warming and the sea ice is rapidly retreating, the East Siberian Sea will undoubtedly suffer more extreme heatwaves, which might cause unpredictable climate impacts on the Arctic biochemical processes and greenhouse gas emission.

删除了：CH₄

删除了： and temperatures close to the freezing point

warm bottom waters during the winter (Kraineva et al., 2019). A series of year-round oceanographic moorings at a deeper
50 location (40-m water depth) in the Laptev Sea recorded near-bottom temperatures reaching 2.0°C during 2012/2013 (Janout
et al., 2016). Turbulent mixing is critical in determining vertical heat transfer in the water column. The strength of vertical
mixing on the East Siberian Arctic shelf is dominated by wind forcing, tide-/shear-induced mixing, surface stress from ice
motion, and buoyancy loss during ice formation. When the sea ice declines dramatically at the end of the summer melt
season, wind forcing and tides become dominant roles (Janout & Lenn, 2014). Previous studies showed that cyclone-induced
55 strong wind can cause a mixing layer deepening of about 5 - 10 m (Long & Perrie, 2012; Peng et al., 2021). This slight
deepening ~~can only~~ cause significant impacts on the ~~thermohaline bottom~~ conditions on the inner shelf (water depth < 30 m).

The East Siberian Sea (ESS) has a much broader shelf than the Laptev Sea. However, ~~for the lack of in-situ~~ observations, few ~~studies focus on the~~ bottom warming and related physical processes in the ESS. In the summer of 2016,
60 the sea ice within the ESS sector decreased significantly, leaving a nearly ice-free shelf by early September (Figure 1). ~~On~~
~~September 11th, a uniformly mixed water column on the mid-shelf of the ESS was observed with the bottom water~~
~~temperature reaching nearly 3.0°C.~~ It is the first observation of such remarkable vertical mixing event in the ESS during the
period with the most intensified upper stratification over the course of a year. This paper reports this extreme mixing event
and the related processes. We also present the long-term vertical mixing intensification during the sea ice retreat and its
thermal impacts on the bottom layer in the ESS in the recent decade.

65 ~~The paper is structured as follows. Section 2 provides a data and methods description. Section 3 presents the process~~
~~of an extreme vertical mixing event firstly observed in the ESS and its influences on the heat budget of shelf waters.~~
~~Discussion and conclusions are given in Section 4.~~

2 Data and Methods

2.1 In-situ observations

70 The cruise (LA77), using the vessel R/V Akademik M.A. Lavrentyev, carried out two hydrographic sections across the
ESS. The cross-shelf observations started at station LA77-17 on September 2, and ended at station LA77-40 on September
15, 2016 (Figure 1b). Twenty-three temperature-salinity profiles were obtained using an SBE-911 plus Conductivity,
Temperature, and Depth (CTD) sensor with temperature and salinity accuracies of 0.001°C and 0.003, respectively. ~~The~~
~~sensor was calibrated at the Sea-bird facilities in Seattle before and after the cruise, and the CTD casts were processed~~
75 ~~according to Sea-bird processing procedures.~~ Nine temperature-salinity profiles were obtained using expendable CTD
(XCTD) with temperature and salinity accuracies of 0.02°C and 0.04, respectively. ~~The in-situ CTD and XCTD data are~~
~~available at <https://doi.org/10.5281/zenodo.4507584>.~~ The wind speed at each station was recorded by a mechanical
anemometer mounted at about 10 m high.

删除了: is intense enough to

删除了: thermal

删除了: of the bottom layer

删除了: due to

删除了: observations and

删除了: During this period, fieldwork in the ESS found that the
whole water column on the mid-shelf was uniformly mixed and the
bottom water temperature reaching nearly 3.0°C....

删除了: s

设置了格式: 字体: (中文) + 中文正文(宋体), (中文) 中文(中国)

2.2 Reanalysis product and satellite-derived data

We use the hourly and monthly atmospheric reanalysis products provided by the European Center for Medium-Range Weather Forecasts (ECMWF) (Hersbach et al., 2020). The spatial resolution of meteorological and surface heat flux datasets is $0.25^\circ \times 0.25^\circ$. That of the ocean wave datasets is $0.5^\circ \times 0.5^\circ$.

删除了：， and that of the ocean wave datasets is $0.5^\circ \times 0.5^\circ$.

We use the Global Ocean Physics Reanalysis (GLORYS12V1) product with an eddy-resolving (1/12° horizontal resolution and 50 vertical levels) resolution to evaluate the warming trends of bottom waters on the shelf bottom (Lellouche, et al., 2021). The model component is the Nucleus for European Modelling of the Ocean (NEMO) platform driven at surface by ECMWF ERA-Interim then ERA5 reanalysis. Along track altimeter data (Sea Level Anomaly), satellite sea surface temperature, sea ice concentration and in-situ temperature and salinity profiles are jointly assimilated. Moreover, a 3D-VAR scheme provides a correction for the slowly-evolving large-scale biases in temperature and salinity. This reanalysis has good representation of Arctic sea ice concentration and water temperature, and it is widely used in the analysis of Arctic hydrographic structure changes (Hall et al., 2021; Hudson et al., 2024; Ivanov et al., 2024).

The sea surface temperature (SST used in the analysis) is from the Optimally Interpolated SST daily product v5.1 with a spatial resolution of $0.25^\circ \times 0.25^\circ$, supported by Remote Sensing Systems sponsored by National Oceanographic Partnership Program (NOPP) and the NASA Earth Science Physical Oceanography Program.

2.3 Mixed layer depth, wavelength of surface wave, and ocean heat content

The mixed layer depth (MLD) is usually between 15-25 m on the Arctic shelves, and its change can be used to demonstrate the regionally diapycnal mixing strength (Peralta-Ferriz and Woodgate, 2015). The method to determine the MLD is according to the maximum depth at which the density is within a certain threshold of the shallowest measured density. Due to complex hydrographic conditions around the Arctic Ocean, there are various thresholds from 0.01 to 0.25 kg/m³ (Timmermans et al., 2012; Peralta-Ferriz and Woodgate, 2015). In this study, we use a threshold value of 0.05 kg/m³ to detect the MLD. Note that the other threshold density values would not make a difference in determining the extreme mixing event.

To evaluate how an extreme diapycnal-mixing event may impact the vertical heat budget on the central shelf of the ESS, the wavelength of surface wave, and the ocean heat content, are considered. The downward maximum mixing depth caused by wind-induced surface waves is roughly consistent with its own wavelength (Qiao et al., 2010), so we can infer the maximum mixing depth of surface waves according to its wavelength size. The ocean heat content is used to examine to what extent the surface ocean heat being transported to the bottom layer by an extreme diapycnal mixing

The surface wavelength λ (meter) is calculated using equations derived from the dispersion relation of regular waves (Holthuijsen, 2007), as follows:

删除了：We use the datasets of a Global Ocean Physics Reanalysis product (GLORYS12V1), with a spatial resolution of $1/12^\circ \times 1/12^\circ$, to investigate the variations of bottom water temperature and mixed layer depth. The GLORYS12V1, provided by the Copernicus Marine Environment Monitoring Service, is a data-assimilated eddy-resolving global reanalysis based mainly on a real-time global forecasting system. In-situ observations, remotely sensed SST, sea ice concentration, and along-track altimeter data are jointly assimilated. The Copernicus Marine Environment Monitoring Service also provides surface velocities derived from absolute dynamic topography....

删除了：defining mixed layer depth (MLD)

删除了：rich vertical structures caused by water advection and the vertical mixing...

设置了格式：非突出显示

135

$$\begin{aligned}\lambda &= \frac{gT^2}{2\pi} && \text{when } d/\lambda \geq 0.5, \\ \lambda &= \frac{gT^2}{2\pi} \tanh\left(\frac{2\pi d}{\lambda}\right) && \text{when } 0.05 \leq d/\lambda < 0.5, \\ \lambda &= \sqrt{gd} \times T && \text{when } d/\lambda \leq 0.05,\end{aligned}$$

where d is the water depth in meters, T the wave period in seconds, and g the acceleration of gravity, 9.8 m/s^2 .

Ocean heat content Q (Joule) is calculated for the surface and bottom layers as follows:

删除了:

140 where ρ is average water density ($\sim 1026 \text{ kg/m}^3$), C_p seawater heat capacity ($\sim 4200 \text{ J/(kg}\cdot\text{C)}$, θ and θ_{fz} the potential temperature and freezing point ($^{\circ}\text{C}$) of each layer, respectively, and z_1 and z_2 the depth range of each layer. [The surface layer is the MLD while the bottom layer is from the lower boundary of MLD to the seafloor.](#)

2.4 Days with heavy winds and high waves

145 Energy induced by heavy winds and high ocean waves are primary factors that enhance the diapycnal mixing in the upper ocean. We determine the days with heavy winds and high wave following three criteria: First, the local (in one grid) hourly wind speed is higher than 13 m/s , and its cumulative duration within one day is more than 6 hours; Second, the grids with the wind speed higher than the criteria will occupy at least 20% of the total area of the studied region ($155\text{--}174.5^{\circ}\text{E}$, $71.5\text{--}74.5^{\circ}\text{N}$); Third, for the days with high waves, its hourly significant wave height must be higher than 2.4 m .




Figure 1: Observations in the East Siberian Sea (ESS).

(a) The path of the Arctic cyclone from September 9 to 12, 2016. The white area indicates the ocean covered by sea ice. (b) Location of stations (blue triangles) in the ESS. The red triangle marks the station LA77-35. Dots show the trajectory of the

删除了: Observation station locations

155 cyclone center. The red dashed line marks the area with wind speed exceeding 12 m/s. (c) The temperature along the section from LA77-17 to LA77-27 and then from LA77-28 to LA77-40. The horizontal coordinate is time. The black square 160 rectangles denote the mixed layer depth at each station, and the red triangle marks the location of station LA77-35. The red rectangular box denotes where the significant bottom water warming occurs. The inset figure shows the temperature profiles of all stations, in which, the red line denotes the temperature profile of the station LA77-35. (d) Same as (c) but for the salinity. (e) Variation of wind speeds during the observation period. Black dots denote the 6-hour wind speeds derived from the hourly ERA5 reanalysis data according to the vessel's GPS positions. The red asterisks denote in-situ observations of wind speeds from an onboard anemometer.

删除了: blue

删除了: marks

删除了: 

3 Results

3.1 A record deepening of the surface mixed layer in Summer

Observations in September 2016 showed that the water column on the ESS shelf has a two-layer structure: the surface mixed layer and the bottom layer separated by the seasonal pycnocline ([Figure 1c and 1d](#)). Temperatures in the surface mixed layer were between 2–5°C, and the salinity was lower than 31.0, while the temperatures in the bottom layer dropped below -1°C, and the salinity ranged between 31.0–34.0. The mixed layer depth was only 10–20 m on the mid-shelf, and about 30 m near the slope. However, observation at LA77-35 showed that the mixed layer deepened to the sea floor, resulting in a uniform water column of 45 m thick ([Figure 1c and 1d](#)). At the same time, the bottom water temperature increased to about 2°C, above the freezing point. Because station LA77-35 was far from the influences of Atlantic Water and coastal river plumes, we deduce that the unusual bottom warming at LA77-35 was due to intense diapycnal mixing from the surface layer.

From September 9 to 12, 2016, a synoptic-scale Arctic cyclone moved southward across the ESS shelf ([Figure 1a](#)) with open water about 1100 km long along its path. When its center moved to the mid-shelf on September 11, the cyclone caused strong winds in the western ESS. Station LA77-35 was right in the center of heaviest wind, where the observed wind speed exceeded 14 m/s on September 11 ([Figure 1b and 1e](#)). The CTD profile of LA77-35 was obtained on September 12 when the sea condition calmed down. Hence the observed ocean state at LA77-35 represented the final influences of the cyclone on the mid-shelf. The extreme deepening of the mixed layer at LA77-35 was driven by the windy weather condition. The intense diapycnal mixing completely eroded the stratification on the shelf, resulting in a record deepening of the mixed layer in the Pacific sector of the Arctic Ocean during summertime.

3.2 Vertical mixing intensified by the growing waves in open water

The sea level pressure at the center of the cyclone was 1012–1024 hPa ([Figure 2](#)), indicating it was just a moderate cyclone, compared with the minimum observed sea level pressures during two Arctic cyclones in August 2016 (967 hPa) and August 2008 (976 hPa), respectively (Long & Perrie, 2012; Peng et al., 2021). The resulting maximum wind speed was also lower than the previous two. However, the deepened mixed layer observed in September 2016 in the ESS reached about 25 m depth, much larger than the 5–10 m deepening reported by the previous studies ([Long & Perrie, 2012; Peng et al., 2021](#)). The main difference between the cyclone we experienced during September 9–12, 2016, and the reported ones was that its path was along the entirely open waters, allowing the growing of the wave and the continuous wind energy input into the ocean.

At 00:00 UTC on September 11, when the cyclone center was still in the Arctic basin, the significant wave height (using wave height for short in the following text) near the continental slope was about 1.5 m ([Figure 2a](#)). Half a day later

when the cyclone center moved southward onto the outer shelf, the maximum wave height gradually grew to about 2.0 m. When the cyclone landed on September 12, nearly all the wave heights on the western ESS shelf were above 2.0 m. The 200 maximum wave height exceeded 3.0 m, and its location was close to station LA77-35 (Figure 2c).

The turbulent mixing induced by wave breaking is mainly confined within the near-surface zone with the depth scale of wave heights (Soloviev & Lukas, 2003; Sulisz et al., 2015). To examine whether the wave-induced mixing can reach the seafloor of the mid-shelf, we calculated the wavelength change during the cyclone's passing (Figure 2d-f). Before September 205 11, the wave length in the mid-shelf region is around 20-30 m. However, after the cyclone passed, the wavelength increased to about 70 m, which is long enough to stir the whole water column with 30-60 m depth by enhanced shearing and wave breaking. The well-developed wave enhanced the turbulent mixing in the mid-shelf, significantly deepening the mixed layer, and fully eroding the summer ocean stratification from surface to bottom. Therefore, although the cyclone we experienced was limited in intensity, its wind energy input to the ocean is maximal accumulation along the large areas of open waters from north to south.

210




Figure 2: Distribution of significant wave height (top panel) and the wave length (bottom panel) every 12 hours during September 11-12.

(a-c) The black contours represent sea level pressure in units of hPa. The arrows represent the 10 m wind field. The white 215 area denotes sea ice extent, and the red asterisk marks the location of station LA-77-35. (d-f) Arrows indicate the wave direction. Thin black lines in (d-f) indicate 30 m, 50 m, and 100 m isobath, respectively.

删除了: On September 12,

删除了: in the mid-shelf region with 30 to 60 m depths

删除了: s

删除了:

删除了: From September 11 to 12.

删除了: .

删除了: Same as top but for the distribution of wavelength.

225 **3.3 Heat Transfer to the bottom of mid-shelf by intensified vertical mixing**

Observations showed that the mixed layer depth in the ESS in early September was only 15 - 25 m, and water temperatures in the bottom layer were below -1°C (Figure 1c). However, the wind-induced diapycnal mixing at station LA77-35 caused a remarkable warming of about 3°C near the seafloor of the mid-shelf. This intensified downward mixing process transferred about 1.9×10^8 J/m² of heat from the surface to the bottom layer, equivalent to a monthly mean heat flux around 73 W/m², an order of magnitude higher than the net solar radiation reaching the bottom during summertime.

During September 10-11, the SST at station LA77-35 decreased by about 1.2°C (Figure 3a). This rapid decrease in temperature was associated with the enhanced diapycnal mixing caused by the cyclone, as well as the enhanced surface heat flux from the ocean to the atmosphere. According to the ERA5 Reanalysis data, the net shortwave radiation decreased to about 40 W/m² in this region (positive value means heat input to the ocean), the sensible and latent heat fluxes at station 235 LA77-35 increased to -40 W/m² and -50 W/m², respectively (Figure 3b). The net surface heat flux surface was about -73 W/m², which can result in the SST decrease of about 0.15-0.19°C within two days, if assuming a mixed layer depth of 15-20 m before the cyclone. The SST at station LA77-35 decreased by more than 1.0°C, and the area around the station with the same cooling rate was about 14,000 km², suggesting the area with the significant surface cooling most likely experienced the same-extent bottom warming induced by intensified vertical mixing, as observed at LA77-35.

240 Note that the reduction of the SST from September 10 to 12 should be more than 2°C under such strong diapycnal mixing of about 45m in depth. But the SST at station LA77-35 only decreased by about 1.2°C. We infer that the additional heat is from the advection of the warming coastal water in the west of the ESS to the mid-shelf (Figure 3e). The heat advection by coastal water only explains about 30-40% of the surface and the mixed layer temperature variability, the vertical heat transport enhanced by the vertical mixing still dominants the bottom warming in this area.

245

删除了: It should be noted that the net surface heat flux, about 45 W/m² during September 10 to 12 can lead to more than 2°C reduction of the SST, if assuming the mixed layer thickness is about 20 m, whereas the SST at station LA77-35 only decreased by about

删除了: But t




Figure 3: Sea surface temperature (SST), air-sea energy flux, and the surface geostrophic current in the ESS.

(a) Changes of mean SST in the region around station LA77-35 (74.375°N , 157.125°E); (b) Same as (a) but for the mean air-sea energy fluxes. Positive value means heat input into the ocean. Black dots denote the net surface energy flux. (c) Difference of the SST between September 10 and 12. The red dot marks the location of station LA77-35. (d) Distribution of mean surface energy balance between September 10 to 12. (e) Same as (d) but for the mean surface geostrophic current and SST during the observation period.

删除了: asterisk

删除了: (derived from sea surface height)

260 3.4 Intensification of the wind-induced mixing on the ESS shelf

We further investigate the long-term trend of wind-induced mixing in the ESS during the recent decade. The key parameters include the sea ice extent, the day with heavy winds and high waves, the mixed layer depth, and the bottom water temperature (Figure 4). The summer sea ice extent became lower than $0.5 \times 10^6 \text{ km}^2$ since 2007, and remained at similar level since then (Figure 4a). Simultaneously, the number of days with heavy winds and high waves gradually increased (Figure

4b). All of the top five years with the most prevailing windy processes occur in recent decade. Especially, in September 2016 and 2017, heavy winds and high waves prevailed in the ESS for more than a week (Figure 4b). As a result, the mean mixed layer depth in September exhibits significant deepening trend from about 11 m to nearly 15 m during last two decades (Figure 4c), and the bottom temperature reaches nearly 0°C during 2015 and 2016, indicating an increasing influence of wind-induced synoptic processes on vertical mixing (Figure 4d). This intensified vertical mixing may result from a synergistic effect of wave-induced mixing (Mellor 2008; Qiao et al., 2010), wind-induced inertial oscillations (Lenn et al., 2011), the alignment of wind and current directions (Burchard and Rippeth, 2009; Lincoln et al., 2015), or the propagation of shelf waves carrying signals from distant storms (Schulz et al., 2021).




Figure 4: Variability of the sea ice extent, number of days with heavy winds and high waves, SST, and the bottom water temperature in the ESS during 2001-2021.

(a) Variation of sea ice extent in July, August, September (thick black line), and October in 2016. **(b)** Number of days with heavy winds and high waves in September. **(c)** SST and the mixed layer depth in the ESS in summer. **(d)** Same as (c) but for the bottom water temperature.

4 Discussion and conclusions

285 Deep water mixing can rapidly ventilate bubble-transported and dissolved methane from the bottom layer, producing high emission rates to the atmosphere (Shakhova et al., 2014; James et al., 2016). Previous studies reported that summer cyclones can deepen the surface mixed layer of the Arctic Ocean by 5 - 10 m, still away from the seafloor (Long & Perrie, 2012; Peng et al., 2021). Here we show an event with a thickened mixed layer to the seafloor with a 25 m depth in the ESS. This recorded-deepening of the mixed layer in the Pacific sector results from an Arctic cyclone moving onto the shelf from 290 open waters with nearly a thousand ~~kilometres~~ from north to south. Over such large open water area, the wave can grow sufficiently and become strong enough to strengthen the vertical mixing of the mid-shelf ocean. The intensified mixing transports a large amount of heat from the upper layer straightly to the seafloor, leading to remarkable warming of about 3°C in the bottom layer. Our observational study provides an important supplement and extension of previous knowledge of wind-induced bottom water warming in the Laptev Sea shelf (Hölemann et al., 2011; Janout et al., 2016; Kraineva et al., 295 2019). Furthermore, based on the in-situ observations, we estimate that the area of bottom water warming on the mid-shelf can at least reach about 14,000 km².

300 The Arctic has warmed nearly 2-4 times faster than the global average since 1979 (Serreze & Barry, 2011; Rantanen et al., 2022). At the same time, the summer sea ice covering the Arctic shelves also decreased dramatically and remained at a low level since 2007 (Park et al., 2020; Liu et al., 2021). Lengthening summers, reduced sea ice extent, and increasing solar 305 heating are pushing more heat into the ocean (Angelopoulos et al., 2020; Dong et al., 2022). With the increasing trends of the open-water area, heavy wind days, and high wave days during summertime since 2012, more vigorous wind-induced mixing and the enhanced bottom warming will happen in the ESS. Now the average bottom water temperature in summer is still below 0°C. However, with the further reduction of sea ice extent and increasing heat content in the mixed layer in the future, the bottom water temperature will undoubtedly become warmer than the freezing point, which is very serious for the bottom 310 permafrost and the methane buried in the sediments. Previous studies report that temperatures of the surface sediments in the ESS remain at about -1.5 to -1.8°C (Chuvalin et al., 2021; 2022), slightly colder than bottom water temperatures in summer. It indicates that the heat entering the bottom layer by wind-induced mixing will continue to enter the sediments, eventually causing the sediment temperature to deviate from the freezing point, which will accelerate subsea methane emissions (Sultan et al., 2020; Wild et al., 2022). Quantifying the impact of these processes is beyond the scope of this paper, but it deserves attention in the future.

315 Observations in the Laptev Sea revealed that the near-bottom warming could remain for at least 2-3 months when the sea ice refreezes again (Janout et al., 2016). This means that the heat stored in the bottom layer during summertime can partly return to the surface layer to melt ice. Unfortunately, the lack of long-term mooring observations prevents us from evaluating how many proportions of heat can reemerge to melt sea ice in the following winter in the ESS. Analyzing and quantifying the vertical distribution and transport of ocean heat on the Arctic shelves will improve the understanding of the

删除了: kilometers

删除了: Kraineva et al., 2019

impacts of the Arctic Ocean warming on regional climate changes. To address this question, long-term mooring systems are urgently needed to monitor the ocean-seabed thermal processes in the ESS.

320 Competing interests

The contact author has declared that none of the authors has any competing interests.

Acknowledgments

This study is supported by the National Key R&D Program of China under Grant 2019YFA0607000, the Chinese Natural Science Foundation under Grants 42276248 and 42176235. The authors would like to thank the European Center for Medium-Range Weather Forecasts for access to the ERA5 reanalysis data, the National Oceanographic Partnership Program (NOPP) and the NASA Earth Science Physical Oceanography Program for access to the Optimally Interpolated SST data, the Copernicus Marine Environment Monitoring Service (CMEMS) for access to the Global Ocean Physics Reanalysis product and the surface geostrophic current data.

The in-situ observation data and the detail introductions are available at <https://doi.org/10.5281/zenodo.4507584>.

330 The ERA5 hourly and monthly data are available at <https://doi.org/10.24381/cds.adbb2d47> and <https://doi.org/10.24381/cds.fl17050d7>.

The Microwave Optimally Interpolated SST data are produced by Remote Sensing Systems and are available at <https://www.remss.com/measurements/sea-surface-temperature/>.

335 The Global Ocean Physics Reanalysis product is available at <https://doi.org/10.48670/moi-00021>, and the surface geostrophic current data are available at <https://doi.org/10.48670/moi-00148>.

References

Angelopoulos, M., Overduin, P. P., Miesner, F., Grigoriev, M. N., & Vasiliev, A. A. Recent advances in the study of Arctic submarine permafrost. *Permafrost and Periglacial Processes*, 31(3), 442-453. <https://doi.org/10.1002/ppp.2061>, 2020.

Bauch, D., & Cherniavskaya, E. Water Mass Classification on a Highly Variable Arctic Shelf Region: Origin of Laptev Sea 340 Water Masses and Implications for the Nutrient Budget. *Journal of Geophysical Research: Oceans*, 123(3), 1896-1906. <https://doi.org/10.1002/2017JC013524>, 2018.

Burchard, H. & Rippeth, T.P., Generation of bulk shear spikes in shallow stratified tidal seas. *Journal of Physical Oceanography*, 39(4), pp.969-985, 2009.

345 Chuvilin, E., Bukhanov, B., Grebenkin, S., Tumskoy, V., Shakhova, N., Dudarev, O., Semiletov, I., & Spasennykh, M. [Thermal properties of sediments in the East Siberian Arctic Seas: A case study in the Buor-Khaya Bay. *Marine and Petroleum Geology*, 123, 104672. https://doi.org/10.1016/j.marpetgeo.2020.104672, 2021.](https://doi.org/10.1016/j.marpetgeo.2020.104672)

350 Chuvilin, E., Bukhanov, B., Yurchenko, A., Davletshina, D., Shakhova, N., Spivak, E., Rusakov, V., Dudarev, O., Khaustova, N., Tikhonova, A., Gustafsson, O., Tesi, T., Martens, J., Jakobsson, M., Spasennykh, M., & Semiletov, I. [In-situ temperatures and thermal properties of the East Siberian Arctic shelf sediments: Key input for understanding the dynamics of subsea permafrost. *Marine and Petroleum Geology*, 138, 105550. https://doi.org/10.1016/j.marpetgeo.2022.105550, 2022.](https://doi.org/10.1016/j.marpetgeo.2022.105550)

355 Dmitrenko, I. A., Kirillov, S. A., Tremblay, L. B., Kassens, H., Anisimov, O. A., Lavrov, S. A., Razumov, S. O., & Grigoriev, M. N. [Recent changes in shelf hydrography in the Siberian Arctic: Potential for subsea permafrost instability. *Journal of Geophysical Research: Oceans*, 116\(C10\) https://doi.org/10.1029/2011JC007218, 2011.](https://doi.org/10.1029/2011JC007218)

360 Dong, J., Shi, X., Gong, X., Astakhov, A. S., Hu, L., Liu, X., Yang, G., Wang, Y., Vasilenko, Y., Qiao, S., Bosin, A., & Lohmann, G. [Enhanced Arctic sea ice melting controlled by larger heat discharge of mid-Holocene rivers. *Nature Communications*, 13\(1\), 5368. http://doi.org/10.1038/s41038-022-33106-1, 2022.](https://doi.org/10.1038/s41038-022-33106-1)

El Bani Altuna, N., Rasmussen, T. L., Ezat, M. M., Vadakkepuliyambatta, S., Groeneveld, J., & Greaves, M. [Deglacial bottom water warming intensified Arctic methane seepage in the NW Barents Sea. *Communications Earth & Environment*, 2\(1\), 188. http://doi.org/10.1038/s43247-021-00264-x, 2021.](https://doi.org/10.1038/s43247-021-00264-x)

Ferré, B., Jansson, P. G., Moser, M., Serov, P., Portnov, A., Graves, C. A., Panieri, G., Gründger, F., Berndt, C., Lehmann, M. F., & Niemann, H. [Reduced methane seepage from Arctic sediments during cold bottom-water conditions. *Nature Geoscience*, 13\(2\), 144-148. http://doi.org/10.1038/s41561-019-0515-3, 2020.](https://doi.org/10.1038/s41561-019-0515-3)

Hall, S. B., Subrahmanyam, B., Nyadjro, E. S., and Samuels, A.: [Surface freshwater fluxes in the arctic and subarctic seas during contrasting years of high and low summer sea ice extent. *Remote Sens.*, 13, 1570. https://doi.org/10.3390/rs13081570, 2021.](https://doi.org/10.3390/rs13081570)

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., & Thépaut, J. The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999-2049. https://doi.org/10.1002/qj.3803, 2020.

Hölemann, J. A., Kirillov, S., Klagge, T., Novikhin, A., Kassens, H., & Timokhov, L. [Near-bottom water warming in the Laptev Sea in response to atmospheric and sea-ice conditions in 2007. *Polar Research*, http://doi.org/10.3402/polar.v30i0.6425, 2011.](https://doi.org/10.3402/polar.v30i0.6425)

375 Holthuijsen, L. H. [Waves in oceanic and coastal waters. Cambridge university press, https://doi.org/10.1017/CBO9780511618536, 2007.](https://doi.org/10.1017/CBO9780511618536)

Hudson, P. A., Martin, A. C. H., Josey, S. A., Marzocchi, A., & Angeloudis, A. Drivers of Laptev Sea interannual variability in salinity and temperature. *Ocean Science*, 20(2), 341-367. <http://doi.org/10.5194/os-20-341-2024>, 2024

380 Ivanov, V. V., Danshina, A. V., Smirnov, A. V., & Filchuk, K. V. Transformation of the atlantic water between svalbard and Franz Joseph Land in the late winter 2018–2019. *Deep Sea Research Part I: Oceanographic Research Papers*, 206, 104280. <https://doi.org/10.1016/j.dsr.2024.104280>, 2024

James, R. H., Bousquet, P., Bussmann, I., Haeckel, M., Kipfer, R., Leifer, I., Niemann, H., Ostrovsky, I., Piskozub, J., Rehder, G., Treude, T., Vielstädte, L., & Greinert, J. Effects of climate change on methane emissions from seafloor sediments in the Arctic Ocean: A review. *Limnology and Oceanography*, 61(S1), S283-S299. <https://doi.org/10.1002/lno.10307>, 2016.

385 Janout, M. A., & Lenn, Y. Semidiurnal Tides on the Laptev Sea Shelf with Implications for Shear and Vertical Mixing. *Journal of Physical Oceanography*, 44(1), 202-219. <http://doi.org/10.1175/JPO-D-12-0240.1>, 2014.

Janout, M., Hölemann, J., Juhls, B., Krumpen, T., Rabe, B., Bauch, D., Wegner, C., Kassens, H., & Timokhov, L. Episodic 390 warming of near-bottom waters under the Arctic sea ice on the central Laptev Sea shelf. *Geophysical Research Letter*, 43(1), 264-272. <https://doi.org/10.1002/2015GL066565>, 2016.

Kraineva, M., Golubeva, E., & Platov, G. Simulation of the near-bottom water warming in the Laptev Sea in 2007–2008. *Bull. Nov. Comp. Center Num. Model. in Atmosph.*, 17, 21-30, 2019.

395 Lellouche, J.-M., Greiner, E., Romain, B.-B., Gilles, G., Angélique, M., Marie, D., Clément, B., Mathieu, H., Olivier, L. G., Charly, R., Tony, C., Charles-Emmanuel, T., Florent, G., Giovanni, R., Mounir, B., Yann, D., and Pierre-Yves, L. T.: The Copernicus Global 1/12° Oceanic and Sea Ice GLORYS12 Reanalysis, *Front. Earth Sci.*, 9, 698876, <https://doi.org/10.3389/feart.2021.698876>, 2021.

Lenn, Y.-D., T. P. Rippeth, C. P. Old, S. Bacon, I. Polyakov, V. Ivanov, & J. Hölemann, Intermittent intense turbulent mixing under ice in the Laptev Sea continental shelf. *Journal of Physical Oceanography*, 41, 531-547, 2011.

400 Lincoln, B. J., T. P. Rippeth, & J. H. Simpson, Surface mixed layer deepening through wind shear alignment in a seasonally stratified shallow sea, *Journal of Geophysical Research*, 121, 6021–6034, doi:10.1002/2015JC011382, 2015.

Liu, Z., Risi, C., Codron, F., He, X., Poulsen, C. J., Wei, Z., Chen, D., Li, S., & Bowen, G. J. Acceleration of western Arctic sea ice loss linked to the Pacific North American pattern. *Nature Communications*, 12(1), 1519. <http://doi.org/10.1038/s41467-021-21830-z>, 2021.

405 Long, Z., & Perrie, W. Air-sea interactions during an Arctic storm. *Journal of Geophysical Research: Atmospheres*, 117(D15) <https://doi.org/10.1029/2011JD016985>, 2012.

Mellor, G. L., The three dimensional, current and surface wave equations: a revision. *Journal of physical oceanography*, 33, 1978-1989, 2008.

Park, H., Watanabe, E., Kim, Y., Polyakov, I., Oshima, K., Zhang, X., Kimball, J. S., & Yang, D. Increasing riverine heat 410 influx triggers Arctic sea ice decline and oceanic and atmospheric warming. *Science Advances*, 6(45), e4699. <http://doi.org/doi:10.1126/sciadv.abc4699>, 2020.

Peng, L., Zhang, X., Kim, J., Cho, K., Kim, B., Wang, Z., & Tang, H. Role of Intense Arctic Storm in Accelerating Summer Sea Ice Melt: An In Situ Observational Study. *Geophysical Research Letter*, 48(8), e2021GL092714G. <https://doi.org/10.1029/2021GL092714>, 2021.

415 Peralta-Ferriz, C., & Woodgate, R. A. Seasonal and interannual variability of pan-Arctic surface mixed layer properties from 1979 to 2012 from hydrographic data, and the dominance of stratification for multiyear mixed layer depth shoaling. *Progress in Oceanography*, 134, 19-53. <https://doi.org/10.1016/j.pocean.2014.12.005>, 2015.

Qiao, F., Yuan, Y., Ezer, T., Xia, C., Yang, Y., Lü, X., & Song, Z., A three-dimensional surface wave-ocean circulation coupled model and its initial testing. *Ocean Dynamics*, 60, 1339–1355, 2010.

420 Rantanen, M., Karpechko, A. Y., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., & Laaksonen, A. The Arctic has warmed nearly four times faster than the globe since 1979. *Communications Earth & Environment*, 3(1), 168. <http://doi.org/10.1038/s43247-022-00498-3>, 2022.

Sayedi, S. S., Abbott, B. W., Thornton, B. F., Frederick, J. M., Vonk, J. E., Overduin, P., Schädel, C., Schuur, E. A. G., Bourbonnais, A., Demidov, N., Gavrilov, A., He, S., Hugelius, G., Jakobsson, M., Jones, M. C., Joung, D., Kraev, G., 425 Macdonald, R. W., David McGuire, A., Mu, C., O Regan, M., Schreiner, K. M., Stranne, C., Pizhankova, E., Vasiliev, A., Westermann, S., Zarnetske, J. P., Zhang, T., Ghandehari, M., Baeumler, S., Brown, B. C., & Frei, R. J. Subsea permafrost carbon stocks and climate change sensitivity estimated by expert assessment. *Environmental Research Letters*, 15(12), 124075. <http://doi.org/10.1088/1748-9326/abcc29>, 2020.

Schulz, K., Büttner, S., Rogge, A., Janout, M., Hölemann, J., & Rippeth, T. P., Turbulent mixing and the formation of an 430 intermediate nepheloid layer above the Siberian continental shelf break. *Geophysical Research Letters*, 48, e2021GL09298, <https://doi.org/10.1029/2021GL092988>, 2021.

Serreze, M. C., & Barry, R. G. Processes and impacts of Arctic amplification: A research synthesis. *Global and Planetary Change*, 77(1), 85-96. <https://doi.org/10.1016/j.gloplacha.2011.03.004>, 2011.

435 Shakhova, N., Semiletov, I., & Chuvalin, E. Understanding the Permafrost-Hydrate System and Associated Methane Releases in the East Siberian Arctic Shelf. *Geosciences*, 9(6), 251, 2019.

Shakhova, N., Semiletov, I., Leifer, I., Sergienko, V., Salyuk, A., Kosmach, D., Chernykh, D., Stubbs, C., Nicolsky, D., Tumskoy, V., & Gustafsson, Ö. Ebullition and storm-induced methane release from the East Siberian Arctic Shelf. *Nature Geoscience*, 7(1), 64-70. <http://doi.org/10.1038/ngeo2007>, 2014.

Soloviev, A., & Lukas, R. Observation of wave-enhanced turbulence in the near-surface layer of the ocean during TOGA 440 COARE. *Deep Sea Research Part I: Oceanographic Research Papers*, 50(3), 371-395. [https://doi.org/10.1016/S0967-0637\(03\)00004-9](https://doi.org/10.1016/S0967-0637(03)00004-9), 2003.

Sulisz, W., Paprota, M., & Wang, J. Theoretical and Experimental Investigations of Wave-Induced Vertical Mixing. *Mathematical Problems in Engineering*, 950849. <http://doi.org/10.1155/2015/950849>, 2015.

Sultan, N., Plaza-Faverola, A., Vadakkepuliyambatta, S., Buentz, S., & Knies, J. Impact of tides and sea-level on deep-sea 445 Arctic methane emissions. *Nature Communications*, 11(1), 5087. <http://doi.org/10.1038/s41467-020-18899-3>, 2020.

Timmermans, M., Cole, S., & Toole, J. Horizontal Density Structure and Restratification of the Arctic Ocean Surface Layer. *Journal of Physical Oceanography*, 42(4), 659-668. <http://doi.org/10.1175/JPO-D-11-0125.1>, 2012.

Wang, X., Zhao, J., Lobanov, V. B., Kaplunenko, D., Rudykh, Y. N., He, Y., & Chen, X. Distribution and Transport of Water Masses in the East Siberian Sea and Their Impacts on the Arctic Halocline. *Journal of Geophysical Research: Oceans*, 126(8), e2020J-e16523J. <https://doi.org/10.1029/2020JC016523>, 2021.

Wild, B., Shakhova, N., Dudarev, O., Ruban, A., Kosmach, D., Tumskoy, V., Tesi, T., Grimm, H., Nybom, I., Matsubara, F., Alexanderson, H., Jakobsson, M., Mazurov, A., Semiletov, I., & Gustafsson, Ö. Organic matter composition and greenhouse gas production of thawing subsea permafrost in the Laptev Sea. *Nature Communications*, 13(1), 5057. <http://doi.org/10.1038/s41467-022-32696-0>, 2022.

删除了: Angelopoulos, M., Overduin, P. P., Miesner, F., Grigoriev, M. N., & Vasiliev, A. A. Recent advances in the study of Arctic submarine permafrost. *Permafrost and Periglacial Processes*, 31(3), 442-453. <https://doi.org/10.1002/ppp.2061>, 2020.⁴⁵⁰
Bauch, D., & Cherniavskaya, E. Water Mass Classification on a Highly Variable Arctic Shelf Region: Origin of Laptev Sea Water Masses and Implications for the Nutrient Budget. *Journal of Geophysical Research: Oceans*, 123(3), 1896-1906. <https://doi.org/10.1002/2017JC013524>, 2018.⁴
Chuvilin, E., Bukanov, B., Grebenkin, S., Tumskoy, V., Shakova, N., Dudarev, O., Semiletov, I., & Spasennykh, M. Thermal properties of sediments in the East Siberian Arctic Seas: A case study the Buor-Khayn Bay. *Marine and Petroleum Geology*, 123, 104672. <https://doi.org/10.1016/j.marpetgeo.2020.104672>, 2021.⁴
Chuvilin, E., Bukanov, B., Yurchenko, A., Davletshina, D., Shakova, N., Spivak, E., Rusakov, V., Dudarev, O., Khastova, N., Tikhonova, A., Gustafsson, O., Tesi, T., Martens, J., Jakobsson, M., Spasennykh, M., & Semiletov, I. In-situ temperatures and thermal properties of the East Siberian Arctic shelf sediments: Key input for understanding the dynamics of subsea permafrost. *Marine and Petroleum Geology*, 138, 105550. <https://doi.org/10.1016/j.marpetgeo.2022.105550>, 2022.⁴
Dmitrenko, I. A., Kirillov, S. A., Tremblay, L. B., Kassens, H., Anisimov, O. A., Lavrov, S. A., Razumov, S. O., & Grigoriev, M. N. Recent changes in shelf hydrography in the Siberian Arctic: Potential for subsea permafrost instability. *Journal of Geophysical Research: Oceans*, 116(C10) <https://doi.org/10.1029/2011JC007218>, 2011.⁴
Dong, J., Shi, X., Gong, X., Astakhov, A. S., Hu, L., Liu, X., Yang, G., Wang, Y., Vasilenko, Y., Qiao, S., Bosin, A., & Lohmann, G. Enhanced Arctic sea ice melting controlled by larger heat discharge of mid-Holocene rivers. *Nature Communications*, 13(1), 5368. <http://doi.org/10.1038/s41467-022-33106-1>, 2022.⁴
El Bani Altuna, N., Rasmussen, T. L., Ezat, M. M., Vadakkepuliyambatta, S., Groeneveld, J., & Greaves, M. Deglacial bottom water warming intensified Arctic methane seepage in the NW Barents Sea. *Communications Earth & Environment*, 2(1), 188. <http://doi.org/10.1038/s43247-021-00264-x>, 2021.⁴
Ferré, B., Jansson, P. G., Moser, M., Serov, P., Portnov, A., Graves, C. A., Panieri, G., Gründger, F., Berndt, C., Lehmann, M. F., & Niemann, H. Reduced methane seepage from Arctic sediments during cold bottom-water conditions. *Nature Geoscience*, 13(2), 144-148. <http://doi.org/10.1038/s41561-019-0515-3>, 2020.⁴
Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellán, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimes, P., Hogan, R. J., Holm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P., Rozum, I., Vamborg, F., Villaume, S., & Thépaut, J. The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146(730), 1999-2049. <https://doi.org/10.1002/qj.3803>, 2020.⁴
Hölemann, J. A., Kirillov, S., Klagge, T., Novikhin, A., Kassens, H., & Timokhov, L. Near-bottom water warming the Laptev Sea in response to atmospheric and sea-ice conditions in 2007. *Polar Research*, <http://doi.org/10.3402/polar.v30.06425>, 2011.⁴
Holthuijsen, L. H. *Waves in oceanic and coastal waters*. Cambridge university press, <https://doi.org/10.1017/CBO9780511618536>, 2007.⁴
James, R. H., Bousquet, P., Bussmann, I., Haeckel, M., Kipfer, R., Leifer, I., Niemann, H., Ostrovsky, I., Piskozub, J., Rehder, G., Treude, T., Vielstädte, L., & Greinert, J. Effects of climate ch[...]

第 16 页: [1] 删除了

Bus Stal

2024/10/28 16:40:00