

1 **Widespread increase in discharge from West Antarctic Peninsula** 2 **glaciers since 2018**

3 Benjamin J. Davison¹, Anna E. Hogg¹, Carlos Moffat², Michael P. Meredith³, Benjamin, J. Wallis¹

4 ¹School of Earth and Environment, University of Leeds, Leeds, United Kingdom

5 ²School of Marine Science and Policy, University of Delaware, Newark, DE, USA

6 ³British Antarctic Survey, Cambridge, United Kingdom

7 *Correspondence to:* Benjamin J. Davison (b.davison@leeds.ac.uk)

8 **Abstract.** Many glaciers on the Antarctic Peninsula have retreated and accelerated in recent decades. Here we show that there
9 was a widespread, quasi-synchronous and sustained increase in grounding line discharge from glaciers on the west coast of the
10 Antarctic Peninsula since 2018. Overall, west Antarctic Peninsula discharge trends increased by over a factor of three, from
11 50 Mt yr⁻² during 2017 to 2020 up to 160 Mt yr⁻² in the years following, leading to a 7.4 % increase in grounding line discharge
12 since 2017. The acceleration in discharge was concentrated at glaciers connected to deep, cross-shelf troughs hosting warm
13 ocean waters, and the acceleration occurred during a period of anomalously high subsurface water temperatures on the
14 continental shelf. Given that many of the affected glaciers have retreated over the past several decades in response to ocean
15 warming, thereby highlighting their sensitivity to ocean forcing, we argue that the recent period of anomalously warm water
16 was likely a key driver of the observed acceleration. However, the acceleration also occurred during a time of anomalously
17 high atmospheric temperatures and glacier surface runoff, which could have contributed to speed-up by directly increasing
18 basal water pressure and, by invigorating near-glacier ocean circulation, increasing submarine melt rates. The spatial pattern
19 of glacier acceleration therefore provides an indication of glaciers that are exposed to warm ocean water at depth and/or have
20 active surface-to-bed hydrological connections; however, many stages in the chain of events leading to glacier acceleration,
21 and how that response is affected by glacier-specific factors, remain insufficiently understood. Both atmospheric and ocean
22 temperatures in this region and its surroundings are likely to increase further in the coming decades, therefore there is a pressing
23 need to improve our understanding of recent changes in Antarctic Peninsula glacier dynamics in response atmospheric and
24 oceanic changes in order to improve projections of their behaviour over the coming century.

25 **1 Introduction**

26 The Antarctic Peninsula (AP) hosts over 800 tidewater glaciers, which collectively hold an ice mass equivalent to 69±5 mm
27 of global sea level rise (Huss and Farinotti, 2014). Substantial changes in glacier and ice shelf area have occurred across the
28 AP since the mid-20th century (Cook and Vaughan, 2010; Doake and Vaughan, 1991; Rott et al., 1996). Many studies have
29 focused on changes to AP ice shelves, including the retreat of Wordie Ice Shelf from 1966 to 1989 (Doake and Vaughan, 1991;
30 Vaughan and Doake, 1996), Prince Gustav Ice Shelf during 1989 to 1995 (Cooper, 1997), Larsen-A in 1995 (Rott et al., 1996),

31 Larsen-B in 2002 (Rack and Rott, 2004; Scambos et al., 2003) and Wilkins Ice Shelf in 2008 (Braun et al., 2009). These
32 changes in ice shelf area have generally been attributed to rising surface air temperatures, leading to extensive melt ponding,
33 hydrofracture and rapid successive calving of elongate icebergs parallel to the ice shelf edge (Scambos et al., 2009). Glacier
34 acceleration and thinning has followed the collapse of these ice shelves due to loss of ice shelf buttressing – the Larsen-B
35 tributary glaciers have become a heavily researched example of this response (Rignot et al., 2004; Scambos et al., 2004; Wuite
36 et al., 2015; Rott et al., 2018; Seehaus et al., 2018). Although the well-documented initial acceleration and subsequent
37 deceleration of those glaciers was substantial, measurements of AP mass change over recent decades remain uncertain because
38 of very large uncertainties in bed elevation and surface mass balance (Rignot et al., 2019; Gardner et al., 2018; Hansen et al.,
39 2021; Rott et al., 2018), though recent efforts to downscale regional climate model output has led to significant improvements
40 (Noël et al., 2023).

41 Outside of ice shelf tributary glaciers, tidewater glaciers on the AP have received less research attention. The majority of such
42 glaciers on the west coast have retreated since at least the 1980s (Cook et al., 2005; Cook and Vaughan, 2010; Cook et al.,
43 2014), seemingly in response to increased flow of relatively warm ($> 1^{\circ}\text{C}$) Circumpolar Deep Water (CDW) onto the
44 continental shelf south of Bransfield Strait (Cook et al., 2016). Glaciers in the southwest AP draining into the George VI Ice
45 Shelf and Bellingshausen Sea have accelerated (Hogg et al., 2017) and thinned (Wouters et al., 2015) since the late-2000s. In
46 addition to these long-term changes in area, speed and thickness, many glaciers along the west AP coast appear to undergo
47 seasonal changes in ice velocity (Wallis et al., 2023b; Boxall et al., 2022), which may be driven by changes in surface and
48 upper-layer ocean temperature, surface-derived meltwater flow at the ice-bed interface, changes in sea ice coverage or some
49 combination thereof. Pulses of meltwater supply to the ice-bed interface, caused by rapid supraglacial lake drainage or extreme
50 melt events, may cause some glaciers on the AP to undergo rapid, short-lived accelerations (Tuckett et al., 2019) but, insofar
51 as they do occur, they remain challenging to detect (Rott et al., 2020).

52 More recently, a large and sustained acceleration and retreat of Cadman Glacier on the west AP has been documented (Wallis
53 et al., 2023a). This acceleration and retreat began in 2018 during a period of anomalously high subsurface ocean temperatures
54 on the continental shelf, due to an incursion of warm CDW. Whilst the glaciers immediately adjacent to Cadman Glacier were
55 protected from this incursion of warm CDW by shallow sills, many glaciers on the west AP will not have such protective sills,
56 raising the possibility of a more widespread response of glaciers on the west AP. Identifying and attributing such a response
57 is important because understanding drivers of grounded ice speed change is informative for interpreting present-day glacier
58 mass changes and for reducing uncertainties in projections of future glacier mass change. In this study, we examine changes
59 in ice speed, grounding line discharge, terminus positions and ocean temperature along a substantial section of the west AP
60 (Figure 1) during this period of anomalously high atmospheric and subsurface ocean temperature.




Figure 1. Study area overview. (a) April 2014 to April 2024 mean ice speed and bathymetry (Morlighem et al., 2020) of the Antarctic Peninsula. Routinely repeated Conductivity-Temperature-Depth (CTD) stations from the Palmer Long-Term Ecological Research programme shown by numbered and coloured dots. The crossed dots indicate CTD stations acquired since 2009. Glacier drainage basins (Cook et al., 2014) are outlined in black and the inset shows basins Hp-I and West Graham Land outlined in red. (b) Ice speed change between the periods 2017/04/01 to 2020/09/01 and 2020/04/01 to 2023/09/01, as a percentage of the long-term average speed. DB in (b) indicates Darbel Bay.

61 2 Methods

62 2.1 Grounding line discharge

63 Grounding line discharge is the rate of mass flowing across the glacier grounding line towards the sea. In the case of tidewater
 64 glaciers with relatively stable termini, it approximates the calving flux. We use the dataset of Davison et al. (2023), which
 65 provides monthly-average grounding line discharge through 16 flux gates located between 3 and 6 km upstream of the
 66 MEaSURES grounding line (Mouginot et al., 2017); readers are referred to Davison et al. (2023) for full methodological details.
 67 For the purposes of this study, we use the ‘FrankenBed’ version of the discharge dataset, which uses a 100x100 m bedrock
 68 grid for the Antarctic Peninsula (Huss and Farinotti, 2014), removes firn air content using the Institute for Marine and
 69 Atmospheric Research Utrecht Firn Densification Model (Veldhuijsen et al., 2022) and accounts for changes in surface
 70 elevation over time using time-dependent polynomial fits to observed surface elevation changes posted on a 5x5 km grid at

71 quarterly intervals (Shepherd et al., 2019). The correction for firm air content affects the total grounding line discharge through
72 each basin but has no impact on the trends in grounding line discharge. The correction for changes in surface elevation results
73 in an overall 1 % decrease in grounding line discharge from 1996 to 2021, and thus is not expected to significantly affect
74 grounding line discharge trends at the majority of glaciers examined here. Some glaciers on the west AP, such as Cadman
75 Glacier, have undergone substantial thinning in recent years (Wallis et al., 2023a), and those changes are included in this
76 dataset. During the study period (2017 to 2023), all the discharge estimates are calculated using 100x100 m velocity estimates
77 derived from intensity tracking of Sentinel-1 6- and 12-day image pairs, making them particularly suitable for resolving
78 changes in speed on the relatively narrow outlet glaciers of the AP. The discharge dataset includes discharge time-series for
79 all glacier basins on the AP as defined by Cook et al. (2014). In this study, we restrict our analysis to 569 glaciers in the west
80 AP, which we define as basins whose centre coordinate falls within West Graham Land or basin Hp-I, as defined by Mouginot
81 et al. (2017) (Figure 1).

82 **2.2 Discharge change point**

83 For each tidewater glacier basin on the west AP, we used change point analysis to identify the single most substantial change
84 in grounding line discharge linear trends since 2017. Change points were defined as the time at which the linear discharge
85 trends before and after the change point differ the most. To reduce aliasing seasonal discharge variability, we excluded change
86 points falling within 20 months (25 %) of the beginning or end of the study period. For all basins, we calculated the linear
87 discharge trend before and after the identified change point to highlight glaciers that underwent a trend acceleration or even a
88 trend reversal. Although we calculate a change point for all glaciers, we note that not all glaciers underwent a significant
89 change in discharge. To identify glaciers with a significant acceleration, we isolated basins where the discharge trend during
90 the second period was positive, at least 50 % greater than during the first period and where the P-value of the trend during the
91 second period was less than 0.1 – we chose not to restrict our analysis just to basins with more significant trends (e.g. $P < 0.05$)
92 because of the short time periods over which trends were calculated. We then further tested for the sensitivity of the timing of
93 the change point, by incrementing the change point in one-month intervals for three months either side of the initial change
94 point. Only glaciers for which each of the above conditions were met using all seven change points were considered to have
95 undergone a significant, sustained discharge trend change that was not sensitive to seasonal variability. Throughout this study,
96 we present discharge trends and trend changes for all glaciers identified as having undergone a significant and sustained
97 discharge trend change, focusing on the timing and spatial distribution of those changes with respect to changes in atmospheric
98 and oceanic conditions. Furthermore, ten of those glaciers with the strongest changes in discharge trend (locations in Figure
99 1) were selected for detailed examination and for demonstration of the discharge trend changes.

100 **2.3 Terminus positions**

101 For each of the ten example glaciers, we measured interannual changes in glacier terminus position by delineating termini in
102 all available cloud-free Sentinel-2 imagery between February and May each year from 2016 to 2023. Higher frequency

103 measurements show that there is seasonal terminus advance and retreat along the west AP, with the most advanced positions
104 generally occurring at the end of the austral winter and the most retreated positions occurring at the end of summer (Wallis et
105 al., 2023b). By focusing on Sentinel-2 imagery from February to May, our measurements approximate the seasonally most
106 retreated position whilst avoiding the difficulties posed by low radar backscatter during the melt events and by Digital
107 Elevation Model artefacts that can affect Sentinel-1 Ground Range Detected imagery in this area of steep topography. We
108 perform the terminus delineations in the Google Earth Engine Digitisation Tool (GEEDiT), and use the multi-centrelines
109 method in the Margin Change Quantification Tool (MaQiT) to calculate width-averaged terminus position change for each
110 glacier (Lea, 2018). When calculating width-averaged terminus position change, we only include sections of the terminus
111 delineated at every measurement epoch.

112 **2.4 Atmospheric and ocean temperature change**

113 We extract daily 2 m atmospheric temperatures over the west AP from 1979 through 2023 from ERA5 reanalyses (Hersbach
114 et al., 2020) and calculate daily anomalies relative to the 1979-2008 daily climatology. We calculate ocean temperature
115 anomalies along five Conductivity-Temperature-Depth (CTD) sections occupied during the Palmer Long-Term Ecological
116 Research (LTER) programme (Smith et al., 1995). The Palmer LTER CTD dataset provides quasi-annual snapshots of
117 conservative ocean temperature, typically during January, along transects from beyond the continental shelf break to near the
118 west AP coastline. For this study, we selected the five transects occupied most frequently (locations in Figure 1), each separated
119 by approximately 100 km, extending from Marguerite Bay in the south to Palmer Basin in the north. In 2009, the Palmer-
120 LTER programme extended its sampling grid latitudinally but reduced its cross-shore resolution (Figure 1). Here, we calculate
121 conservative temperature anomalies during each cruise relative to the 1999-2008 mean for each transect, during which time
122 the programme was still using the high-resolution grid. We also examine daily runoff time-series from 5x5 km resolution
123 RACMO2.3p2 (van Wessem et al., 2018).

124 **3 Results**

125 **3.1 Acceleration of grounding line discharge**

126 We observe widespread changes in speed on the AP between the April 2017 to September 2020 and April 2020 to September
127 2023 periods (Figure 1b; Figure 2). Many tidewater glaciers draining the west AP accelerated by 5 to 20 % since April 2017,
128 leading to an overall 7 Gt yr^{-1} (7.4 %) increase in west AP grounding line discharge. This acceleration was most pronounced
129 in the fast-flowing trunks of the larger outlet glaciers and was clearest at Montgolfier Glacier, Niepce Glacier, Luke Glacier,
130 Comrie Glacier and Wilkinson Murphy Glacier, where speeds increased by over 20 % (Figure 2). At some glaciers, such as
131 Blanchard Glacier and Montgolfier Glacier, we observe deceleration in the shear margins and around high elevation ice falls
132 (Figure 2b, c), which we hypothesise is due to shear margin weakening and dynamic thinning, respectively.




Figure 2. Speed change of selected glaciers between the periods 2017/04/01 to 2020/09/01 and 2020/04/01 to 2023/09/01. (a) Whitecloud, (b) Blanchard, (c) Montgolfier, (d) Moser, (e) Bolton, (f) Niepce, (g) Leay, (h) Wiggins, (i) Bussey, (j) Luke, (k) Comrie and, (l) Wilkinson Murphy. The background is the 15x15 m Landsat Image Mosaic of Antarctica (Bindshadler et al., 2008).

133 Throughout the observation period, grounding line discharge increased at 177 basins on the west AP, such that it was
 134 significantly correlated with time ($R^2 \geq 0.5$ and $P < 0.05$), whilst 49 basins underwent an overall decrease in grounding line
 135 discharge. For some basins, the discharge increase is relatively steady and is part of a longer-term trend – these glaciers are
 136 not the focus of this study. We instead focus on glaciers that underwent a notable increase in linear discharge trends between
 137 2018 and 2021 (Figures 3 to 5). To illustrate these linear trend increases, grounding line discharge at Wilkinson Murphy
 138 Glacier remained steady at 2017 levels, with fluctuations of magnitude less than 5 % from 2017 to June 2020, after which
 139 discharge increased at a rate of 3.4 % yr^{-1} to a maximum around 10 % greater than 2017 levels (Figure 3j). Similarly, the
 140 positive trends in discharge at Montgolfier Glacier, Niepce Glacier and Luke Glacier all increased by more than a factor of
 141 five between May 2021 and January 2022 (Figure 3b, d, h). Some glaciers, such as Moser Glacier, Leay Glacier and Bussey




Figure 3. Grounding line discharge change at selected glaciers. In each panel, grounding line discharge change (relative to the 2017 mean) and associated error are shown as black dots and grey shading. The timing of the change in discharge trend is shown by the dashed line with the date labelled. The linear trends before and after the change point are shown in magenta and green respectively. The red shading indicates the austral summer (December through February).

142 Glacier transitioned from a period of weakly declining discharge to very strongly increasing discharge during this broad period
 143 of acceleration (Figure 3c, e, g).

144 These large increases in linear discharge trends are widespread along the west AP (Figures 4 and 5). Overall, 97 of the 569
 145 glaciers on the west AP exhibited a 50 % or greater increase in linear discharge trend. Of those 97 glaciers, 42 were insensitive
 146 to the timing of the discharge change point within a 7-month period. In comparison, only 7 glaciers underwent a significant
 147 decrease in discharge trend when calculated using the same methods. There is a clear spatial pattern to these increases in linear
 148 discharge trends: the majority of glaciers north of Blanchard Glacier and south of Wilkinson Murphy Glacier generally had
 149 little change in discharge trend since 2017. The majority of glaciers that underwent a significant increase in discharge were
 150 located in the central west AP, between Blanchard and Wilkinson Murphy glaciers. Within the central west AP, there appears
 151 to be some clustering to the discharge changes. Some areas, such as Darbel Bay (location in Figure 1), host several glaciers
 152 that appear to have little change in discharge. In the case of Darbel Bay, the bathymetry is shallow (<100 m based on




Figure 4. Overview of discharge trend changes. (a) Linear trend in discharge from January 2017 to the change point for each basin on the west coast of the Antarctic Peninsula. (b) Linear trend in discharge from the change point to July 2023. (c) The change in discharge trend before and after the change point, with positive values indicating a trend increase. (d) The timing of the discharge trend change. Basins with significant trends ($P < 0.05$; a,b) or significant trend increases (see text for details; c,d) are outlined in black.

153 BedMachine v3; Morlighem et al., 2020), limiting the transport of warm CDW to the coast. However, other ‘low responders’
 154 do not always coincide with areas of shallow bathymetry and sometimes have responsive neighbouring glaciers. As in Wallis
 155 et al. (2023a), these cases may reflect the presence of shallow bathymetric sills not captured by BedMachine v3, which would
 156 act as barriers to incursions of warm water below the sill depth (Bao and Moffat, 2024).

157 There is broad consistency in the timing of discharge trend changes amongst west AP glaciers (Figures 4 to 6). A vast majority
 158 of glaciers with significant discharge trend increases began to accelerate between the November 2020 and November 2021
 159 (Figure 5d and 6), though there is spread around this period (Figures 3, 4d and 5d). Prior to the change point for each glacier,
 160 there was a range of discharge trends, with some glaciers decelerating, accelerating or remaining approximately steady with
 161 less discharge than in 2023 (Figures 5 and 6). Since the 2020/2021 austral summer, however, there has been a widespread,
 162 quasi-synchronous acceleration of glaciers along a large section of the central west AP, leading to peak discharge at or near
 163 the end of our observations in 2023 (Figure 6).




Figure 5. Violin plot overview of discharge trend changes. (a) Linear trend in discharge from January 2017 to the change point for each basin on the west coast of the Antarctic Peninsula. (b) Linear trend in discharge from the change point to July 2023. (c) The change in discharge trend before and after the change point, with positive values indicating a trend increase. (d) The timing of the discharge trend change. Basins with significant trend increases (see text for details) are labelled ‘Sig’. Note the change in y-axis scale between panel (a) and panel (b).

164 3.2 Terminus position change

165 We examined changes in terminus position at the end of the austral summer from 2016 to 2023 at our 10 example glaciers.
 166 Perhaps surprisingly, inter-annual terminus position changes at 7 of the 10 selected glaciers is negligible or not discernible
 167 from seasonal fluctuations in terminus position (not shown). Bussey Glacier exhibited modest but clear retreat of just 20 m on
 168 average and by 150 m on its true left margin (Figure 7). Wiggins Glacier experienced slightly greater retreat of over 100 m
 169 averaged across the width of the terminus and by approximately 240 m at the most affected section (Figure 6). Wilkinson
 170 Murphy Glacier retreated by 1 km on average since 2017 and by over 1.5 km across much of its fast-flowing centre (Figure
 171 7). The timing of terminus position changes at these glaciers broadly coincides with the observed changes in grounding line
 172 discharge, with the majority of retreat occurring since 2019.

173 3.3 Ocean temperature change

174 The conservative temperature anomalies from the Palmer LTER CTD transects (locations in Figure 1) clearly show a warming
 175 trend on the west AP continental shelf below 100 m from 1993 to 2021, and a cooling trend above 100 m (Figures 8 and 9).
 176 The significant linear trends in water temperature across all transects range from 0.02 °C dec⁻¹ to 0.21 °C dec⁻¹. Of particular
 177 relevance to this study, from 2018 to 2021 there was a positive temperature anomaly at 100 to 200 m depth that built to a peak




Figure 6. Discharge change across the west Antarctic Peninsula. (a) Frequency-density of normalized discharge time-series. Only west AP basins with a trend increase of more than 50 % (N=97) were included to illustrate the synchronicity of the acceleration. (b) Grounding line discharge change (relative to the 2017 mean) of West Graham Land and associated error are shown as black dots and grey shading. The dashed line shows the timing of the change in discharge trend. The magenta and green lines show the linear trends before and after the change point. The red shading indicates the austral summer (December through February).

178 of over 1°C above the long-term average in December 2021, with an anomaly maximum around 100 m depth (Figures 8 and
 179 9). There is variability superimposed on these trends; for example, there was a period of more rapid warming below 100 m
 180 during the 1990s. In addition, the summers of 2013 through to 2017 were generally cooler than the summers of 2007 through
 181 to 2009 along transect 200 (Figure 8). These patterns are well-documented by several other publications (e.g. Cook et al., 2016;
 182 Martinson et al., 2008) and the warm periods are associated with sea ice coverage changes and wind-driven CDW warming
 183 and shoaling within the Antarctic Circumpolar Current (Moffat and Meredith, 2018; Schmidtko et al., 2014), allowing more
 184 and warmer CDW to access the continental shelf.

185 4. Discussion

186 Many glaciers on the west AP have been retreating over recent decades (Cook et al., 2005). This retreat appears to have a
 187 strong latitudinal pattern, with southern glaciers retreating faster, driven by a long-term increase in subsurface ocean
 188 temperatures (Cook et al., 2016; Meredith and King, 2005), caused in turn by warming, shoaling and greater penetration of
 189 CDW onto the continental shelf (Moffat and Meredith, 2018). In addition, many of the west AP glaciers are clearly responsive
 190 to shorter-term changes in ocean temperature and, possibly, surface melt supply, resulting in seasonal changes in ice velocity
 191 and terminus position (Wallis et al., 2023b; Boxall et al., 2022). Therefore, it is reasonable to assume that the west AP glaciers
 192 could be responsive to multi-year anomalies in subsurface ocean temperature and/or meltwater supply. Our observations reveal
 193 a widespread, quasi-synchronous and sustained increase in grounding line discharge across the west AP, centred around the
 194 austral summer of 2021 (Figures 3 to 6). The response is concentrated in the central west AP, where warm CDW accesses the
 195 glaciers via deep, cross-shelf troughs in the continental shelf. The majority of glaciers further north, which are not exposed to




Figure 7. Overview of terminus position changes at four of the selected glaciers. The left column (a, c, e) show width-averaged terminus position change relative to the first measurement. The right column (b, d, f) illustrates the location of the terminus at each measurement time, overlaid on the 15x15 m Landsat Image Mosaic of Antarctica (Bindschadler et al., 2008). The units in (b), (d) and (f) are 1000 km and the projection is South Polar Stereographic (EPSG 3031).

196 CDW, exhibit muted or no change in grounding line discharge trends (Fig 4c). There is variability in the timing and magnitude
 197 of glacier response along the coast, which will be governed by individual glacier geometry (Seehaus et al., 2018), proximal
 198 fjord bathymetry (Wallis et al., 2023a; Bao and Moffat, 2024) as well as the competition between distinct processes (e.g. cross-
 199 shelf transport and modification of CDW vs transport of cold water from the Weddell Sea around the tip of the Peninsula)
 200 setting the subsurface ocean temperature (Moffat and Meredith, 2018; Venables et al., 2017). In places, this results in very
 201 different responses between neighbouring glaciers and, for some glaciers, a continuation of their longer-term discharge trends
 202 (Figure 4).

203 The widespread, quasi-synchronous and sustained nature of the discharge change points to a regional, sustained forcing. The
 204 hydrographic observations show that there was a widespread and coherent increase in subsurface ocean temperatures on the
 205 continental shelf from 2018 onwards, centred at 100 to 200 m depth and extending to the ocean bed on the continental shelf
 206 (Figures 8 and 9). We do not have observations from the waters immediately adjacent to any of the west AP tidewater glaciers,




Figure 8. Conservative temperature anomalies relative to the 1999–2008 mean along transect 200. The vertical grey dashed lines indicate individual cast locations – note that the panel outlines obscure casts at the transect endpoints. The dark grey shading is topography from BedMachine v3 (Morlighem et al., 2020) and the Antarctic Peninsula coast is on the right.

207 so we do not have direct evidence that the anomalously warm waters came into contact with the tidewater glaciers and elevated
 208 submarine melt rates. However, the Palmer LTER data indicate that anomalously warm modified CDW was present across the
 209 continental shelf south of Bransfield Strait during the 2018 to 2021 period, including in the deep, glacially-carved troughs that
 210 connect the shelf edge to the west AP glaciers (Cook et al., 2016; Arndt et al., 2013; Couto et al., 2017). In addition, diverse
 211 local CTD measurements along the west AP have documented the presence of CDW in immediate proximity to glacier termini
 212 in the same region (Meredith et al., 2022; Venables et al., 2023), demonstrating that CDW does penetrate to parts of the coast.




Figure 9. Conservative temperature anomaly time-series. Each panel illustrates time-series of conservative temperature anomalies within the given depths for each transect. The robust linear trends in temperature for each transect are quoted and significant trends ($P < 0.05$) are in bold.

213 It is therefore highly likely that the anomalously warm water present on the continental shelf from 2018 to at least 2021 came
 214 into widespread contact with the west AP glaciers south of Bransfield Strait.

215 Assuming that this contact did happen and that there was no commensurate drop in current velocity at the ice-ocean interface,
 216 we would expect terminus submarine melt rates to increase. Glacier terminus depths along the west AP are poorly mapped,
 217 but the available data indicate that many glaciers are several hundred metres thick at the terminus (Cook et al., 2016; Arndt et
 218 al., 2013). Glaciers with grounding lines deeper than 100 m would be exposed to the anomalously warm CDW during each
 219 austral summer since 2018, likely leading to enhanced undercutting. The temperature anomalies were greatest around 100 to
 220 200 m depth; therefore, the enhancement of undercutting would lead to more pronounced quasi-linear or step-like undercuts
 221 for glaciers shallower than 200 m and parabolic undercuts for more deeply grounded glaciers. Comparable undercut profiles
 222 have been observed at glaciers in Greenland in the presence of similar vertical temperature profiles (Fried et al., 2015; Rignot
 223 et al., 2015).

224 The majority of theoretical and numerical perspectives (Slater et al., 2021; Ma and Bassis, 2019; Benn et al., 2017; Krug et
 225 al., 2015; O’Leary and Christoffersen, 2013) suggest that such profiles of undercutting can amplify calving, leading to retreat
 226 and glacier acceleration. We observe retreat at just three of our ten example glaciers, only one of which (Wilkinson Murphy
 227 Glacier) was very substantial. We do not have terminus position measurements at the tens of other west AP glaciers that
 228 accelerated since the austral summer of 2020/2021. In the absence of terminus retreat, more rapid submarine melting must be
 229 balanced by faster ice velocities (Krug et al., 2015), such that the position of the calving front becomes a function of the
 230 velocity and thickness of the upstream ice, rather than the driver of upstream ice velocity changes (Benn et al., 2007).

231 If enhanced submarine melting were the primary driver of the glacier acceleration, then the spatial pattern of glacier
 232 acceleration provides information about the pathways by which the warm water accessed the west AP coastline. Most of the

233 glaciers that accelerated were located between Adelaide Island and Anvers Island, where several deep troughs provide a direct
234 pathway across the shelf along which CDW intrusions can access the central west AP (Cook et al., 2016; Arndt et al., 2013;
235 Couto et al., 2017). Some glaciers, such as Blanchard Glacier, located further north, where CDW influence on deep water
236 temperatures is at least seasonal (Wang et al., 2022), also accelerated. Such instances likely reflect the convoluted topographic
237 routes that dissect the west AP shelf and the competition between CDW and Weddell Sea waters on deep water temperatures,
238 among other processes. The majority of the northern-most glaciers along the West AP, which drain into Bransfield Strait and
239 are not exposed to warm CDW, showed weak or no acceleration. In addition, we observe acceleration at some glaciers that,
240 according to bathymetry products (Morlighem et al., 2020), are grounded in shallow water. For example, Luke Glacier and
241 Comrie Glacier (locations in Figure 1) are essentially land-terminating in BedMachine v3 yet are several hundred metres thick
242 in an independent thickness product (Huss and Farinotti, 2014). These and other similar sites may therefore indicate regions
243 to target in future bathymetric mapping efforts, or at least for improvement in future bed topographic assimilation efforts.

244 At most depths along the central west AP continental shelf, the conservative temperature anomalies since 2018 were similar
245 to, or slightly larger than, other warm periods in the late-2000s (Figure 9), so it is possible that ocean forcing alone was not
246 sufficient to drive the observed acceleration. In addition to warming ocean waters, ERA5 atmospheric temperatures over the
247 west AP have been anomalously high persistently since 2016 (Figure 10a). There were record high atmospheric temperatures
248 over the AP in February 2020 and 2022 (Gorodetskaya et al., 2023; Francelino et al., 2021). These heatwaves caused record-
249 high levels of snowmelt and rainfall (Gorodetskaya et al., 2023) that in turn led to extreme melt ponding, for example on the
250 George VI and Larsen-C ice shelves in 2020 (Banwell et al., 2021; Bevan et al., 2020). Output from RACMO2.3p2 (van
251 Wessem et al., 2018) - a 5.5 km regional climate model - shows that there is a modest amount of runoff (i.e. snowmelt that
252 does not refreeze in the firn) from the west AP (Figure 10b). The presence of plumes along the west AP coastline (Rodrigo et
253 al., 2016) provide strong evidence that at least some of this surface-derived meltwater and runoff does reach the ice-bed
254 interface and is discharged at the grounding line. Theoretical perspectives (e.g. Jenkins, 2011; Slater et al., 2016) and numerous
255 observational and modelling studies from other regions (e.g. Jackson et al., 2017; Sutherland et al., 2019; Straneo et al., 2011;
256 Carroll et al., 2016) show that the turbulent mixing and entrainment induced by subglacial discharge-driven plumes increases
257 glacier submarine melt rates. The RACMO2.3p2 runoff data indicate that runoff was much higher during February 2020 and
258 2021 than during the preceding years; this would drive more vigorous plumes and faster submarine melt rates, potentially
259 amplifying the effect of the observed warmer subsurface waters (Slater and Straneo, 2022).

260 In addition to modifying submarine melt rates, surface-to-bed meltwater injection could directly increase glacier speeds by
261 increasing basal water storage and by transiently increasing basal water pressure and basal sliding rates. There is some evidence
262 from Sentinel-1 ice velocity estimates supporting the relevance of this process on the AP over weekly to seasonal time-scales,
263 based on the co-occurrence of periods of elevated speed with periods of meltwater availability inferred from regional climate
264 model output (Tuckett et al., 2019; Wallis et al., 2023b; Boxall et al., 2022). However, care must be taken to avoid aliasing
265 apparent velocity changes caused by melt-induced changes in radar penetration depth (Rott et al., 2020). There is a large body




Figure 10. Atmospheric conditions over the west Antarctic Peninsula. (a) 2 m atmospheric temperature anomalies relative to the 1979–2008 daily climatology over the west AP from ERA5 reanalysis. The anomalies are smoothed with a 90-moving window. (b) Modelled runoff from a 5x5 km run of RACMO2.3p2, integrated over the west AP, from 2016 to 2023. Daily runoff is plotted grey and 30-day smoothed runoff in black. Panel (a) was plotted using the anomaly function in MATLAB (Greene, 2024).

266 of evidence that meltwater-induced accelerations on other ice masses generally have little impact on annual ice displacement,
 267 because of meltwater-induced subglacial drainage mechanisms that result in compensatory periods of slower ice flow (e.g.
 268 Sole et al., 2013). On the AP, there are insufficient observations of meltwater-induced ice flow variations to determine whether
 269 similar compensatory subglacial drainage mechanisms also operate there. It is possible that the combination of moderately
 270 thick, fast-flowing ice, low meltwater supply, thick snowpack and potentially extensive firn aquifers (Van Wessem et al., 2021)
 271 may result in qualitatively different meltwater-induced ice velocity changes compared to those observed elsewhere. In addition,
 272 the extreme meltwater production in 2020 and 2022 may have reduced firn pore space, allowing more surface-derived
 273 meltwater to penetrate to the ice-bed interface in subsequent, lower melt years. Further satellite observations and field-based
 274 studies are required to characterise the surface-to-bed hydrological drainage systems and the mechanisms through which they
 275 affect ice flow on the AP.

276 The widespread increase in grounding line discharge of the west AP presented in this study has implications for glacier mass
 277 balance. Although the glaciers on the AP are small compared to their neighbours in parts of West Antarctica, they are changing
 278 rapidly such that AP contributed 14 % of Antarctica’s total mass loss from 1992 to 2020 (Otosaka et al., 2023). Previous work
 279 has linked warming subsurface ocean waters to widespread glacier retreat along the west AP (Cook et al., 2016) and more
 280 recent work has further shown an ocean-driven ice tongue collapse and acceleration of Cadman Glacier on the west AP (Wallis
 281 et al., 2023a). The observations presented in this study build on this understanding by showing a widespread, quasi-
 282 synchronous acceleration of grounding line discharge along the west AP linked to a period of anomalously high air and
 283 subsurface ocean temperatures. Unless surface mass balance increased commensurately, this recent acceleration of west AP

284 glaciers will accelerate the rate of west AP mass loss, contributing to faster rates of sea level rise. In addition, the increase in
285 grounding line discharge constitutes an increased solid freshwater input to the Bellingshausen Sea, which numerical modelling
286 suggests can increase ocean heat transport to West Antarctic ice shelves, potentially leading to faster submarine melt rates
287 (Flexas et al., 2022).

288 **5. Conclusions**

289 During the past half-century, tidewater glaciers on the west coast of the Antarctic Peninsula have retreated in response to rising
290 subsurface ocean temperatures and they remain responsive to seasonal changes in atmospheric and ocean temperatures. This
291 study identifies a widespread, quasi-synchronous and sustained increase in grounding line discharge of many glaciers along
292 the west coast of the Antarctic Peninsula around the 2020/2021 austral summer. In many cases, grounding line discharge trends
293 more than doubled and led to 5 to 20 % increases in grounding line discharge over a 2.5 year period. The acceleration of
294 grounding line discharge occurred at a time of anomalously high, though not exceptional, subsurface ocean temperatures on
295 the continental shelf, which would have increased terminus submarine melt rates and could have driven the observed glacier
296 acceleration. The co-occurrence of record-high air temperatures and surface melting may have contributed to the glacier
297 acceleration by increasing surface-to-bed meltwater delivery, potentially amplifying submarine melt rates and directly
298 increasing glacier sliding speeds. In the absence of *in-situ* observations on the glacier surface and in the waters immediately
299 adjacent to glacier calving fronts, there remain many uncertainties regarding the chain of events leading to this period of glacier
300 acceleration, but we are hopeful that future campaigns to improve seafloor mapping, acquire near-glacier hydrographic
301 measurements and to measure glacier velocity *in-situ* will provide important new understanding of the processes driving
302 changes in ice flow on the Antarctic Peninsula. Nevertheless, it is clear that the recent period of anomalous atmospheric and
303 ocean temperatures have, together or in isolation, driven a widespread and sustained acceleration of many west AP glaciers.
304 Given that the atmosphere and ocean in the region are projected to warm further in the coming decades, we recommend further
305 research in this area to improve understanding of glacier response to changing environmental conditions across the Antarctic
306 Peninsula.

307

308 *Data availability.* The grounding line discharge dataset are available on Zenodo (<https://zenodo.org/records/10417864>). The
309 Palmer LTER dataset were compiled for a previous study and made available on Zenodo
310 (<https://zenodo.org/records/10009821>). BedMachine v3 (Morlighem et al., 2020) is available from:
311 <https://nsidc.org/data/nsidc-0756/versions/3>. The Antarctica Peninsula basin shapefiles (Cook et al., 2014) are available from:
312 <http://add.scar.org/>. The Landsat Image Mosaic of Antarctica (Bindschadler et al., 2008) is available from:
313 <https://lima.usgs.gov/fullcontinent.php>. ERA5 reanalysis (Hersbach et al., 2020) is available from:
314 <https://cds.climate.copernicus.eu/cdsapp#!/dataset/reanalysis-era5-single-levels>.

315

316

317 *Author contributions.* BJD conceived the study, performed the analysis and wrote the manuscript. BJW and CM compiled the
318 raw CTD data into a format more amenable for analysis. All authors discussed the results and implications, and contributed to
319 the manuscript preparation.

320

321 *Competing interests.* The contact author has declared that none of the authors have any competing interests.

322

323 *Acknowledgements.* We are grateful to the creators of the open-access satellite imagery, datasets and tools used in this study,
324 and to the crews of all cruises involved in producing the Palmer LTER dataset. We particularly thank Rich Iannuzzi and
325 Michael Cappola for the initial quality control of the LTER dataset. Data processing was undertaken on ARC3 and ARC4, part
326 of the high-performance computing facilities at the University of Leeds, UK.

327

328 *Financial support.* BJD and AEH are supported by ESA through the Polar+ Ice Shelves project (ESA-IPL-POE-EF-cb-LE-
329 2019-834) and the SO-ICE project (ESA AO/1-10461/20/I-NB), by NERC via the DeCadeS project (NE/T012757/1) and by
330 the UK EO Climate Information Service (NE/X019071/1). BJW is supported by the Panorama NERC Doctoral Training
331 Partnership under grant NE/S007458/1. The Palmer LTER program is supported by NSF-OPP Grant #2026045.

332 **References**

333 Arndt, J. E., Schenke, H. W., Jakobsson, M., Nitsche, F. O., Buys, G., Goleby, B., Rebesco, M., Bohoyo, F., Hong, J., Black,
334 J., Greku, R., Udintsev, G., Barrios, F., Reynoso-Peralta, W., Taisei, M., and Wigley, R.: The international bathymetric chart
335 of the Southern Ocean (IBCSO) version 1.0-A new bathymetric compilation covering circum-Antarctic waters, *Geophys. Res.*
336 *Lett.*, 40, 3111–3117, <https://doi.org/10.1002/grl.50413>, 2013.

337 Banwell, A. F., Tri Datta, R., Dell, R. L., Moussavi, M., Brucker, L., Picard, G., Shuman, C. A., and Stevens, L. A.: The 32-
338 year record-high surface melt in 2019/2020 on the northern George VI Ice Shelf, Antarctic Peninsula, *Cryosphere*, 15, 909–
339 925, <https://doi.org/10.5194/tc-15-909-2021>, 2021.

340 Bao, W. and Moffat, C.: Impact of shallow sills on heat transport and stratification regimes in proglacial fjords, *Cryosph.*, 18,
341 187–203, <https://doi.org/10.5194/tc-18-187-2024>, 2024.

342 Benn, D. I., Warren, C. R., and Mottram, R. H.: Calving processes and the dynamics of calving glaciers, *Earth-Science Rev.*,
343 82, 143–179, <https://doi.org/10.1016/j.earscirev.2007.02.002>, 2007.

344 Benn, D. I., Aström, J., Zwinger, T., Todd, J., Nick, F. M., Cook, S., Hulton, N. R. J., and Luckman, A.: Melt-under-cutting
345 and buoyancy-driven calving from tidewater glaciers: New insights from discrete element and continuum model simulations,
346 *J. Glaciol.*, 63, 691–702, <https://doi.org/10.1017/jog.2017.41>, 2017.

347 Bevan, S., Luckman, A., Hendon, H., and Wang, G.: The 2020 Larsen C Ice Shelf surface melt is a 40-year record high,

348 Cryosphere, 14, 3551–3564, <https://doi.org/10.5194/tc-14-3551-2020>, 2020.

349 Bindschadler, R., Vornberger, P., Fleming, A., Fox, A., Mullins, J., Binnie, D., Paulsen, S. J., Granneman, B., and Gorodetzky,
350 D.: The Landsat Image Mosaic of Antarctica, *Remote Sens. Environ.*, 112, 4214–4226,
351 <https://doi.org/10.1016/j.rse.2008.07.006>, 2008.

352 Boxall, K., Christie, F. D. W., Willis, I. C., Wuite, J., and Nagler, T.: Seasonal land-ice-flow variability in the Antarctic
353 Peninsula, *Cryosphere*, 16, 3907–3932, <https://doi.org/10.5194/tc-16-3907-2022>, 2022.

354 Braun, M., Humbert, A., and Moll, A.: Changes of Wilkins Ice Shelf over the past 15 years and inferences on its stability,
355 *Cryosphere*, 3, 41–56, <https://doi.org/10.5194/tc-3-41-2009>, 2009.

356 Carroll, D., Sutherland, D. A., Hudson, B., Moon, T., Catania, G. A., Shroyer, E. L., Nash, J. D., Bartholomaeus, T. C., Felikson,
357 D., Stearns, L. A., Noël, B. P. Y., and van den Broeke, M. R.: The impact of glacier geometry on meltwater plume structure
358 and submarine melt in Greenland fjords, *Geophys. Res. Lett.*, 43, 9739–9748, <https://doi.org/10.1002/2016GL070170>, 2016.

359 Cook, A. J. and Vaughan, D. G.: Overview of areal changes of the ice shelves on the Antarctic Peninsula over the past 50
360 years, *Cryosphere*, 4, 77–98, <https://doi.org/10.5194/tc-4-77-2010>, 2010.

361 Cook, A. J., Fox, A. J., Vaughan, D. G., and Ferrigno, J. G.: Retreating glacier fronts on the Antarctic Peninsula over the past
362 half-century, *Science (80-.)*, 308, 541–544, <https://doi.org/10.1126/science.1104235>, 2005.

363 Cook, A. J., Vaughan, D. G., Luckman, A. J., and Murray, T.: A new Antarctic Peninsula glacier basin inventory and observed
364 area changes since the 1940s, *Antarct. Sci.*, 26, 614–624, <https://doi.org/10.1017/S0954102014000200>, 2014.

365 Cook, A. J., Holland, P. R., Meredith, M. P., Murray, T., Luckman, A., and Vaughan, D. G.: Ocean forcing of glacier retreat
366 in the western Antarctic Peninsula, *Science (80-.)*, 353, 283–286, <https://doi.org/10.1126/science.aae0017>, 2016.

367 Cooper, A. P. R.: Historical observations of Prince Gustav ice shelf, *Polar Rec. (Gr. Brit.)*, 33, 285–294,
368 <https://doi.org/10.1017/S0032247400025389>, 1997.

369 Couto, N., Martinson, D. G., Kohut, J., and Schofield, O.: Distribution of Upper Circumpolar Deep Water on the warming
370 continental shelf of the West Antarctic Peninsula, *J. Geophys. Res. Ocean.*, 122, 5306–5315, [https://doi.org/10.1002/](https://doi.org/10.1002/2017JC012840)
371 [2017JC012840](https://doi.org/10.1002/2017JC012840), 2017.

372 Davison, B. J., Hogg, A. E., Slater, T., and Rigby, R.: Antarctic Ice Sheet grounding line discharge from 1996 through 2023,
373 *Earth Syst. Sci. Data Discuss.*, <https://doi.org/10.5194/essd-2023-448>, 2023.

374 Doake, C. S. M. and Vaughan, D. G.: Rapid disintegration of the Wordie Ice Shelf in response to atmospheric warming, *Nature*,
375 350, 328–330, <https://doi.org/10.1038/350328a0>, 1991.

376 Flexas, M. M., Thompson, A. F., Schodlok, M. P., Zhang, H., and Speer, K.: Antarctic Peninsula warming triggers enhanced
377 basal melt rates throughout West Antarctica, *Sci. Adv.*, 8, 1–12, <https://doi.org/10.1126/sciadv.abj9134>, 2022.

378 Francelino, M. R., Schaefer, C., de Los Milagros Skansi, M., Colwell, S., Bromwich, D. H., Jones, P., King, J. C., Lazzara, M.
379 A., Renwick, J., Solomon, S., Brunet, M., and Cerveny, R. S.: WMO evaluation of two extreme high temperatures occurring
380 in February 2020 for the antarctic peninsula region, *Bull. Am. Meteorol. Soc.*, 102, E2053–E2061,
381 <https://doi.org/10.1175/BAMS-D-21-0040.1>, 2021.

382 Fried, M. J., Catania, G. A., Bartholomaus, T. C., Duncan, D., Davis, M., Stearns, L. A., Nash, J., Shroyer, E., and Sutherland,
383 D.: Distributed subglacial discharge drives significant submarine melt at a Greenland tidewater glacier, *Geophys. Res. Lett.*,
384 42, 9328–9336, <https://doi.org/10.1002/2015GL065806>, 2015.

385 Gardner, A. S., Moholdt, G., Scambos, T., Fahnestock, M., Ligtenberg, S., Van Den Broeke, M., and Nilsson, J.: Increased
386 West Antarctic and unchanged East Antarctic ice discharge over the last 7 years, *Cryosphere*, 12, 521–547,
387 <https://doi.org/10.5194/tc-12-521-2018>, 2018.

388 Gorodetskaya, I. V., Durán-alarcón, C., González-herrero, S., Clem, K. R., Zou, X., Rowe, P., Imazio, P. R., Campos, D.,
389 Santos, C. L., Dutrievoz, N., and Wille, J. D.: Record-high Antarctic Peninsula temperatures and surface melt in February
390 2022: a compound event with an intense atmospheric river, *npj Clim. Atmos. Sci.*, 6, <https://doi.org/10.1038/s41612-023->
391 00529-6, 2023.

392 Greene, C. A.: Chad Greene (2024). anomaly (<https://www.mathworks.com/matlabcentral/fileexchange/61327-anomaly>),
393 MATLAB Central File Exchange. Chad Greene (2024). anomaly
394 (<https://www.mathworks.com/matlabcentral/fileexchange/61327-anomaly>), MATLAB Central File Exchange., 2024.

395 Hansen, N., Langen, P. L., Boberg, F., Forsberg, R., Simonsen, S. B., Thejll, P., Vandecrux, B., and Mottram, R.: Downscaled
396 surface mass balance in Antarctica: Impacts of subsurface processes and large-scale atmospheric circulation, *Cryosphere*, 15,
397 4315–4333, <https://doi.org/10.5194/tc-15-4315-2021>, 2021.

398 Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers,
399 D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., De Chiara,
400 G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L.,
401 Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Rosnay, P.,
402 Rozum, I., Vamborg, F., Villaume, S., and Thépaut, J. N.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146, 1999–
403 2049, <https://doi.org/10.1002/qj.3803>, 2020.

404 Hogg, A. E., Shepherd, A., Cornford, S. L., Briggs, K. H., Gourmelen, N., Graham, J. A., Joughin, I., Mouginot, J., Nagler,
405 T., Payne, A. J., Rignot, E., and Wuite, J.: Increased ice flow in Western Palmer Land linked to ocean melting, *Geophys. Res.*
406 *Lett.*, 44, 4159–4167, <https://doi.org/10.1002/2016GL072110>, 2017.

407 Huss, M. and Farinotti, D.: A high-resolution bedrock map for the Antarctic Peninsula, *Cryosphere*, 8, 1261–1273,
408 <https://doi.org/10.5194/tc-8-1261-2014>, 2014.

409 Jackson, R. H., Shroyer, E. L., Nash, J. D., Sutherland, D. A., Carroll, D., Fried, M. J., Catania, G. A., Bartholomaus, T. C.,
410 and Stearns, L. A.: Near-glacier surveying of a subglacial discharge plume: Implications for plume parameterizations,
411 *Geophys. Res. Lett.*, 44, 6886–6894, <https://doi.org/10.1002/2017GL073602>, 2017.

412 Jenkins, A.: Convection-Driven Melting near the Grounding Lines of Ice Shelves and Tidewater Glaciers, *J. Phys. Oceanogr.*,
413 41, 2279–2294, <https://doi.org/10.1175/JPO-D-11-03.1>, 2011.

414 Krug, J., Durand, G., Gagliardini, O., and Weiss, J.: Modelling the impact of submarine frontal melting and ice mélange on
415 glacier dynamics, *Cryosph. Discuss.*, 9, 183–221, <https://doi.org/10.5194/tcd-9-183-2015>, 2015.

416 Lea, J. M.: The Google Earth Engine Digitisation Tool (GEEDiT) and the Margin change Quantification Tool (MaQiT) –
417 simple tools for the rapid mapping and quantification of changing Earth surface margins, *Earth Surf. Dyn.*, 551–561,
418 <https://doi.org/10.5194/esurf-6-551-2018>, 2018.

419 Ma, Y. and Bassis, J. N.: The Effect of Submarine Melting on Calving From Marine Terminating Glaciers, *J. Geophys. Res.*
420 *Earth Surf.*, 124, 334–346, <https://doi.org/10.1029/2018JF004820>, 2019.

421 Martinson, D. G., Stammerjohn, S. E., Iannuzzi, R. A., Smith, R. C., and Vernet, M.: Western Antarctic Peninsula physical
422 oceanography and spatio-temporal variability, *Deep. Res. Part II Top. Stud. Oceanogr.*, 55, 1964–1987,
423 <https://doi.org/10.1016/j.dsr2.2008.04.038>, 2008.

424 Meredith, M. P. and King, J. C.: Rapid climate change in the ocean west of the Antarctic Peninsula during the second half of
425 the 20th century, *Geophys. Res. Lett.*, 32, 1–5, <https://doi.org/10.1029/2005GL024042>, 2005.

426 Meredith, M. P., Inall, M. E., Alexander Brearley, J., Ehmen, T., Sheen, K., Munday, D., Cook, A., Retallick, K., Van
427 Landeghem, K., Gerrish, L., Annett, A., Carvalho, F., Jones, R., Naveira Garabato, A. C., Bull, C. Y. S., Wallis, B. J., Hogg,
428 A. E., and Scourse, J.: Internal tsunamigenesis and ocean mixing driven by glacier calving in Antarctica, *Sci. Adv.*, 8, 1–11,
429 <https://doi.org/10.1126/sciadv.add0720>, 2022.

430 Moffat, C. and Meredith, M.: Shelf-ocean exchange and hydrography west of the Antarctic Peninsula: A review, *Philos. Trans.*
431 *R. Soc. A Math. Phys. Eng. Sci.*, 376, <https://doi.org/10.1098/rsta.2017.0164>, 2018.

432 Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R.,
433 Fretwell, P., Goel, V., Greenbaum, J. S., Gudmundsson, H., Guo, J., Helm, V., Hofstede, C., Howat, I., Humbert, A., Jokat,
434 W., Karlsson, N. B., Lee, W. S., Matsuoka, K., Millan, R., Mouginot, J., Paden, J., Pattyn, F., Roberts, J., Rosier, S., Ruppel,
435 A., Seroussi, H., Smith, E. C., Steinhage, D., Sun, B., Broeke, M. R. van den, Ommen, T. D. van, Wessem, M. van, and Young,
436 D. A.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nat. Geosci.*, 13,
437 132–137, <https://doi.org/10.1038/s41561-019-0510-8>, 2020.

438 Mouginot, J., Scheuchl, B., and Rignot, E.: MEaSUREs Antarctic Boundaries for IPY 2007-2009 from Satellite Radar, Version
439 2, Boulder, Color. USA. NASA Natl. Snow Ice Data Cent. Distrib. Act. Arch. Cent., <https://doi.org/10.5067/AXE4121732AD>,
440 2017.

441 Noël, B., van Wessem, J. M., Wouters, B., Trusel, L., Lhermitte, S., and van den Broeke, M. R.: Higher Antarctic ice sheet
442 accumulation and surface melt rates revealed at 2 km resolution, *Nat. Commun.*, 14, 7949, [https://doi.org/10.1038/s41467-](https://doi.org/10.1038/s41467-023-43584-6)
443 023-43584-6, 2023.

444 O’Leary, M. and Christoffersen, P.: Calving on tidewater glaciers amplified by submarine frontal melting, *Cryosphere*, 7, 119–
445 128, <https://doi.org/10.5194/tc-7-119-2013>, 2013.

446 Ootaka, I. N., Shepherd, A., Ivins, E. R., Schlegel, N. J., Amory, C., Van Den Broeke, M. R., Horwath, M., Joughin, I., King,
447 M. D., Krinner, G., Nowicki, S., Payne, A. J., Rignot, E., Scambos, T., Simon, K. M., Smith, B. E., Sørensen, L. S., Velicogna,
448 I., Whitehouse, P. L., Geruo, A., Agosta, C., Ahlstrøm, A. P., Blazquez, A., Colgan, W., Engdahl, M. E., Fettweis, X., Forsberg,
449 R., Gallée, H., Gardner, A., Gilbert, L., Gourmelen, N., Groh, A., Gunter, B. C., Harig, C., Helm, V., Khan, S. A., Kittel, C.,

450 Konrad, H., Langen, P. L., Lecavalier, B. S., Liang, C. C., Loomis, B. D., McMillan, M., Melini, D., Mernild, S. H., Mottram,
451 R., Mouginit, J., Nilsson, J., Noël, B., Pattle, M. E., Peltier, W. R., Pie, N., Roca, M., Sasgen, I., Save, H. V., Seo, K. W.,
452 Scheuchl, B., Schrama, E. J. O., Schröder, L., Simonsen, S. B., Slater, T., Spada, G., Sutterley, T. C., Vishwakarma, B. D.,
453 Van Wessem, J. M., Wiese, D., Van Der Wal, W., and Wouters, B.: Mass balance of the Greenland and Antarctic ice sheets
454 from 1992 to 2020, *Earth Syst. Sci. Data*, 15, 1597–1616, <https://doi.org/10.5194/essd-15-1597-2023>, 2023.

455 Rack, W. and Rott, H.: Pattern of retreat and disintegration of the Larsen B ice shelf, Antarctic Peninsula, *Ann. Glaciol.*, 39,
456 505–510, <https://doi.org/10.3189/172756404781814005>, 2004.

457 Rignot, E., Casassa, G., Gogineni, P., Krabill, W., Rivera, A., and Thomas, R.: Accelerated ice discharge from the Antarctic
458 Peninsula following the collapse of Larsen B ice shelf, *Geophys. Res. Lett.*, 31, 2–5, <https://doi.org/10.1029/2004GL020697>,
459 2004.

460 Rignot, E., Fenty, I., Xu, Y., Cai, C., and Kemp, C.: Undercutting of marine-terminating glaciers in West Greenland, *Geophys.*
461 *Res. Lett.*, 42, 5909–5917, <https://doi.org/10.1002/2015GL064236>, 2015.

462 Rignot, E., Mouginit, J., Scheuchl, B., Van Den Broeke, M., Van Wessem, M. J., and Morlighem, M.: Four decades of
463 Antarctic ice sheet mass balance from 1979–2017, *Proc. Natl. Acad. Sci. U. S. A.*, 116, 1095–1103,
464 <https://doi.org/10.1073/pnas.1812883116>, 2019.

465 Rodrigo, C., Giglio, S., and Varas, A.: Glacier sediment plumes in small bays on the Danco Coast, Antarctic Peninsula, *Antarct.*
466 *Sci.*, 28, 395–404, <https://doi.org/10.1017/S0954102016000237>, 2016.

467 Rott, H., Skvarca, P., and Nagler, T.: Rapid collapse of northern Larsen Ice Shelf, *Antarctica*, *Science* (80-.), 271, 788–792,
468 <https://doi.org/10.1126/science.271.5250.788>, 1996.

469 Rott, H., Abdel Jaber, W., Wuite, J., Scheiblaue, S., Floricioiu, D., Van Wessem, J. M., Nagler, T., Miranda, N., and Van Den
470 Broeke, M. R.: Changing pattern of ice flow and mass balance for glaciers discharging into the Larsen A and B embayments,
471 *Antarctic Peninsula*, 2011 to 2016, *Cryosphere*, 12, 1273–1291, <https://doi.org/10.5194/tc-12-1273-2018>, 2018.

472 Rott, H., Wuite, J., De Rydt, J., Gudmundsson, G. H., Floricioiu, D., and Rack, W.: Impact of marine processes on flow
473 dynamics of northern Antarctic Peninsula outlet glaciers, *Nat. Commun.*, 11, 10–12, <https://doi.org/10.1038/s41467-020->
474 16658-y, 2020.

475 Scambos, T., Fricker, H. A., Liu, C. C., Bohlander, J., Fastook, J., Sargent, A., Massom, R., and Wu, A. M.: Ice shelf
476 disintegration by plate bending and hydro-fracture: Satellite observations and model results of the 2008 Wilkins ice shelf
477 break-ups, *Earth Planet. Sci. Lett.*, 280, 51–60, <https://doi.org/10.1016/j.epsl.2008.12.027>, 2009.

478 Scambos, T. A., Hulbe, C., and Fahnestock, M.: Climate-induced ice shelf disintegration in the Antarctic Peninsula, in:
479 *Antarctic Peninsula Climate Variability: Historical and Paleoenvironmental Perspectives*, vol. 79, 79–92, 2003.

480 Scambos, T. A., Bohlander, J. A., Shuman, C. A., and Skvarca, P.: Glacier acceleration and thinning after ice shelf collapse in
481 the Larsen B embayment, *Antarctica*, *Geophys. Res. Lett.*, 31, 2001–2004, <https://doi.org/10.1029/2004GL020670>, 2004.

482 Schmidtko, S., Heywood, K. J., Thompson, A. F., and Aoki, S.: Multidecadal warming of Antarctic waters, *Science* (80-.),
483 346, 1227–1231, <https://doi.org/10.1126/science.1256117>, 2014.

484 Seehaus, T., Cook, A., Silva, A. B., and Braun, M. H.: Changes in glacier dynamics at the northern Antarctic Peninsula since
485 1985, *Cryosph.*, 12, 577–594, <https://doi.org/10.5194/tc-12-577-2018>, 2018.

486 Shepherd, A., Gilbert, L., Muir, A. S., Konrad, H., McMillan, M., Slater, T., Briggs, K. H., Sundal, A. V., Hogg, A. E., and
487 Engdahl, M. E.: Trends in Antarctic Ice Sheet Elevation and Mass, *Geophys. Res. Lett.*, 46, 8174–8183,
488 <https://doi.org/10.1029/2019GL082182>, 2019.

489 Slater, D. A. and Straneo, F.: Submarine melting of glaciers in Greenland amplified by atmospheric warming, *Nat. Geosci.*,
490 15, 794–799, <https://doi.org/10.1038/s41561-022-01035-9>, 2022.

491 Slater, D. A., Goldberg, D. N., Nienow, P. W., and Cowton, T. R.: Scalings for Submarine Melting at Tidewater Glaciers from
492 Buoyant Plume Theory, *J. Phys. Oceanogr.*, 46, 1839–1855, <https://doi.org/10.1175/JPO-D-15-0132.1>, 2016.

493 Slater, D. A., Benn, D. I., Cowton, T. R., Bassis, J. N., and Todd, J. A.: Calving Multiplier Effect Controlled by Melt Undercut
494 Geometry, *J. Geophys. Res. Earth Surf.*, 126, 1–17, <https://doi.org/10.1029/2021JF006191>, 2021.

495 Smith, R., Baker, K., Fraser, W., Hofmann, E., Karl, D., Klink, J., Quentin, L., Prezelin, B., Ross, R., Trivelpiece, W., and
496 Vernet, M.: The Palmer LTER: A Long-Term Ecological Research Program at Palmer Station, Antarctica, *Oceanography*, 8,
497 77–86, <https://doi.org/10.5670/oceanog.1995.01>, 1995.

498 Sole, A., Nienow, P., Bartholomew, I., Mair, D., Cowton, T., Tedstone, A., and King, M. A.: Winter motion mediates dynamic
499 response of the Greenland Ice Sheet to warmer summers, *Geophys. Res. Lett.*, 40, 3940–3944,
500 <https://doi.org/10.1002/grl.50764>, 2013.

501 Straneo, F., Curry, R. G., Sutherland, D. a., Hamilton, G. S., Cenedese, C., Våge, K., and Stearns, L. a.: Impact of fjord
502 dynamics and glacial runoff on the circulation near Helheim Glacier, *Nat. Geosci.*, 4, 322–327,
503 <https://doi.org/10.1038/ngeo1109>, 2011.

504 Sutherland, D. A., Jackson, R. H., Kienholz, C., Amundson, J. M., Dryer, W. P., Duncan, D., Eidam, E. F., Motyka, R. J., and
505 Nash, J. D.: Direct observations of submarine melt and subsurface geometry at a tidewater glacier, *Science (80-.)*, 365, 369–
506 374, <https://doi.org/10.1126/science.aax3528>, 2019.

507 Tuckett, P. A., Ely, J. C., Sole, A. J., Livingstone, S. J., Davison, B. J., van Wessem, M. J., and Howard, J.: Rapid accelerations
508 of Antarctic Peninsula outlet glaciers driven by surface melt, *Nat. Commun.*, 10, [https://doi.org/10.1038/s41467-019-12039-](https://doi.org/10.1038/s41467-019-12039-2)
509 2, 2019.

510 Vaughan, D. and Doake, C. S. M.: Recent retreat of ice shelves on the Antarctic Peninsula, *Nature*, 379, 328–331, 1996.

511 Veldhuijsen, Sanne, B. M., van de Berg, W. J., Brils, M., Munneke, P. K., and van den Broeke, M. R.: Characteristics of the
512 contemporary Antarctic firm layer simulated with IMAU-FDM v1.2A (1979-2020), *Cryosph. Discuss.*,
513 <https://doi.org/10.5194/tc-2022-118>, 2022.

514 Venables, H., Meredith, M. P., Hendry, K. R., ten Hoopen, P., Peat, H., Chapman, A., Beaumont, J., Piper, R., Miller, A. J.,
515 Mann, P., Rossetti, H., Massey, A., Souster, T., Reeves, S., Fenton, M., Heiser, S., Pountney, S., Reed, S., Waring, Z., Clark,
516 M., Bolton, E., Mathews, R., London, H., Clement, A., Stuart, E., Reichardt, A., Brandon, M., Leng, M., Arrowsmith, C.,
517 Annett, A., Henley, S. F., and Clarke, A.: Sustained year-round oceanographic measurements from Rothera Research Station,

518 Antarctica, 1997–2017, *Sci. Data*, 10, 1–13, <https://doi.org/10.1038/s41597-023-02172-5>, 2023.

519 Venables, H. J., Meredith, M. P., and Brearley, J. A.: Modification of deep waters in Marguerite Bay, western Antarctic
520 Peninsula, caused by topographic overflows, *Deep. Res. Part II Top. Stud. Oceanogr.*, 139, 9–17,
521 <https://doi.org/10.1016/j.dsr2.2016.09.005>, 2017.

522 Wallis, B. J., Hogg, A. E., Meredith, M. P., Close, R., Hardy, D., McMillan, M., Wuite, J., Nagler, T., and Moffat, C.: Ocean
523 warming drives rapid dynamic activation of marine-terminating glacier on the west Antarctic Peninsula, *Nat. Commun.*, 14,
524 <https://doi.org/10.1038/s41467-023-42970-4>, 2023a.

525 Wallis, B. J., Hogg, A. E., van Wessem, J. M., Davison, B. J., and van den Broeke, M. R.: Widespread seasonal speed-up of
526 west Antarctic Peninsula glaciers from 2014 to 2021, *Nat. Geosci.*, 16, 231–237, [https://doi.org/10.1038/s41561-023-01131-](https://doi.org/10.1038/s41561-023-01131-4)
527 4, 2023b.

528 Wang, X., Moffat, C., Dinniman, M. S., Klinck, J. M., Sutherland, D. A., and Aguiar-González, B.: Variability and Dynamics
529 of Along-Shore Exchange on the West Antarctic Peninsula (WAP) Continental Shelf, *J. Geophys. Res. Ocean.*, 127,
530 e2021JC017645, <https://doi.org/10.1029/e2021JC017645>, 2022.

531 van Wessem, J. M., van de Berg, W. J., Noël, B. P. Y., van Meijgaard, E., Birnbaum, G., Jakobs, C. L., Krüger, K., Lenaerts,
532 J. T. M., Lhermitte, S., Ligtenberg, S. R. M., Medley, B., Reijmer, C. H., van Tricht, K., Trusel, L. D., van Ulf, L. H., Wouters,
533 B., Wuite, J., and van den Broeke, M. R.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2,
534 part 2: Antarctica (1979–2016), *Cryosph.*, 12, 1479–1498, <https://doi.org/10.5194/tc-12-1479-2018>, 2018.

535 Van Wessem, J. M., Steger, C. R., Wever, N., and Van Den Broeke, M. R.: An exploratory modelling study of perennial firn
536 aquifers in the Antarctic Peninsula for the period 1979–2016, *Cryosphere*, 15, 695–714, [https://doi.org/10.5194/tc-15-695-](https://doi.org/10.5194/tc-15-695-2021)
537 2021, 2021.

538 Wouters, B., Martin-Español, A., Helm, V., Flament, T., Van Wessem, J. M., Ligtenberg, S. R. M., Van Den Broeke, M. R.,
539 and Bamber, J. L.: Dynamic thinning of glaciers on the Southern Antarctic Peninsula, *Science (80-.)*, 348, 899–903,
540 <https://doi.org/10.1126/science.aaa5727>, 2015.

541 Wuite, J., Rott, H., Hetzenecker, M., Floricioiu, D., De Rydt, J., Gudmundsson, G. H., Nagler, T., and Kern, M.: Evolution of
542 surface velocities and ice discharge of Larsen B outlet glaciers from 1995 to 2013, *Cryosphere*, 9, 957–969,
543 <https://doi.org/10.5194/tc-9-957-2015>, 2015.

544