

Modelling Flood Losses to Microbusinesses in Ho Chi Minh City, Vietnam

Anna Buch^{1,2,3}, Dominik Paprotny², Kasra Rafiezadeh Shahi^{3,4}, Heidi Kreibich³, Nivedita Sairam³

¹~~Institute of Geography, Heidelberg~~ University of Heidelberg, GIScience Research Group, Heidelberg, 69117, Germany

²Potsdam Institute for Climate Impact Research (PIK), Transformation Pathways, Potsdam, 14473, Germany

³German Research Centre for Geosciences (GFZ), Section 4.4 Hydrology, Potsdam, 14473, Germany

⁴~~Potsdam Institute for Climate Impact Research (PIK), Earth System Analysis, Potsdam, 14473, Germany~~

Correspondence to: Anna Buch (anna.buch@uni-heidelberg.de), Nivedita Sairam (~~nivedita~~nivedita.sairam@gfz-potsdam.de)

Abstract. Microbusinesses are important sources of livelihood for low- and middle-income households. In Ho Chi Minh City (HCMC), Vietnam, many microbusinesses are set up ~~in~~on the ground floor of residential houses susceptible to urban floods. Increasing flood risk in HCMC threatens the financial resources of microbusinesses by damaging business contents and causing business interruption. Since flood loss estimations are rarely conducted at object-level resolution and are often focused on households or large companies, the losses suffered by microbusinesses are often overlooked. This study aims to derive the drivers of flood losses [%] in microbusinesses by applying a Conditional Random Forest to survey data (~~relative~~ content losses: n=317; ~~relative~~ business interruption losses: n=361) collected from microbusinesses in HCMC. ~~The variability of the losses to -business~~ content losses and ~~due to~~ business interruption were adequately explained by the revenues of the businesses from monthly sales, age of the building where the business is established and water depth in the building during the flood event. Based on the identified drivers, probabilistic loss models (non-parametric Bayesian Networks) were developed using a combination of data-driven and expert-based model formulation. The models estimated the flood losses for HCMC's microbusinesses with a mean absolute error of 3.8 % for content losses and 18.7 % for business interruption losses. The Bayesian Network model for business interruption performed with a similar predictive performance when it was regionally transferred and applied to comparable survey data from another Vietnamese city, Can Tho ~~City~~. The flood loss models introduced in this study make it possible to derive flood risk metrics specific to microbusinesses to support adaptation decision making and risk transfer mechanisms.

Plain Language Summary. Many households in Vietnam depend on revenues from microbusinesses (shop-houses). However, losses caused by regular flooding to the microbusinesses are not modelled. Business turnover, building age and water depth are found to be the main drivers of flood losses to microbusinesses. We built and validated probabilistic models (non-parametric Bayesian Networks) that estimate flood losses to microbusinesses. The results help in flood risk management and adaption decision making for microbusinesses.

1 Introduction

Comprehensive risk management requires empirical evidence on drivers of risk and assessment of potential impacts. The lack of information on vulnerability of certain economic sectors or social groups, and their often-limited participation in local risk management, in turn foster a lack of awareness among decision-makers leading to biased risk management strategies. As impacts of climate change become more severe, comprehensive risk management that protects society as a whole is imperative - in particular the vulnerable and under-represented groups. However, it is often not plausible in low- and middle-income countries due to poor data availability. An example of a vulnerable economic sector in a society with a high flood risk, explored in this study, are microbusinesses in Ho Chi Minh City (HCMC), Vietnam. [These micro-sized companies are quite common across South-east \(SE\) and South \(S\) Asia; a description of their operations and economic relevance in regard to Vietnam's urban areas are provided in Sect. 2.1.](#)

HCMC is one of the world's most exposed cities to flood risk under current and future conditions (Hallegatte et al. 2013; Seussolini et al. 2017). Similar to other Vietnamese metropolises, HCMC lies in the great delta areas in the south of the country. These flat, riverine and coastal regions experience regular flooding in particular during the rainy season. In HCMC, these regular floods are often the result of compound events caused by the simultaneous occurrence of high tides, heavy rainfall and high flows of the Saigon and Dong Nai rivers and their tributaries (Tran 2014; Thuy et al. 2019). Other large cities in the delta areas of South Vietnam also experience regular urban flooding, for instance, Can Tho City in the Mekong Delta (abbreviated as Can Tho). Urban floods in Can Tho are predominantly fluvial in nature, such as a major flood event in 2011. Despite the ongoing efforts to improve protection and adaptation measures on private and municipal levels, climate change and the ongoing growth of these important economic centres increase their risk to urban flooding (Güneralp et al. 2015; Rentschler et al. 2022). The existing infrastructure and adaptation measures in these cities are unable to counterbalance the new risks caused by intensified flood events and ongoing urban pressure (e.g. Bouwer 2011; Jha et al. 2012; Formetta & Feyen 2019; Kreibich et al. 2022).

We define microbusinesses, including household businesses, according to the definition of the World Bank, as very small businesses with less than ten employees. However, this general definition for microbusinesses needs to be adapted to the regional context of South-East (SE) and South (S) Asia. Microbusinesses in these countries tend to employ usually less than three people. In most cases, they are located on the ground floor of a building with residences on the upper floors, commonly called shophouses in Vietnamese cities. Microbusinesses provide an important source of income for unemployed family members and people with limited opportunities on the labour market, likewise migrant workers and people who received less possibilities of schooling (Samantha 2018). Together with the operations of small and medium-sized companies (SMEs), microbusinesses drive the rapid economic developments of many SE Asian states since the last decades (Trinh & Thanh 2017). According to Vietnam's economic census of 2017, it is estimated that around 75 % of all enterprises are microbusinesses in the country (General Statistics Office 2018). Vietnam's microbusinesses engage around 11 % of all employees (General Statistics Office 2018) and the density of microbusinesses is particularly high in economic centres such as HCMC and other

65 delta cities like Can Tho. The economic importance of HCMC becomes evident when the region's contribution to Vietnam's
total economic output is considered — the HCMC region accounts for approximately 40 % of the national GDP (General
Statistics Office 2018). These values highlight the relevance of microbusinesses for Vietnam's local and national economy.
Microbusinesses are particularly vulnerable to the negative consequences of regular flooding due to their limited financial
resources and inadequate support by local authorities and the government (JETRO 2017; Leitold et al. 2020). As a
70 consequence, microbusinesses often rely on their neighbouring network to cope with flooding (Chinh et al. 2016; 2017; Leitold
& Revilla Diez 2019; Leitold et al. 2021). Bank loans or microcredits are less common due to the usually rather low credit
rating (Patankar 2019). In terms of flood losses this means that repair measures and other business investments are often
directly financed by the savings of the microbusiness owner. Insufficient or missing flood insurance policies can further
exacerbate the situation of flood affected businesses (KPMG 2016; Patankar 2019). Besides temporal decline in revenues,
75 repair costs or poor future prospects, worse case impacts may include business closures or unemployment among business
owners and their employees (Jha et al. 2012).

In addition to the largely studied structural damages, the commercial sector, specifically, microbusinesses also suffers directly
from economic loss of business content (e.g. inventory, goods, equipment) and due to business interruption. The latter refers
to the decline in business revenues due to interrupted business operations of flood-affected businesses during a reference period
80 such as the flood month or period of flooding (Meyer et al., 2013; Chinh et al., 2016). However, our definition of interruption
losses does not consider long-term losses or impacts on businesses outside the flood zone. The literature on commercial losses
often focuses on companies of various sizes in Europe or the US and these studies indicate that indirect losses represent a
significant share of flood consequences (e.g. Hallegatte, 2008; Merz et al., 2010; Koks and Thissen, 2016; Sieg et al., 2019;
and Tsinda et al., 2019). Since the business structures and available resources for larger firms differ considerably from those
85 of small- and micro-sized companies (JETRO 2017; Leitold and Diez, 2019), the state-of-the-art approaches for commercial
flood loss modelling are not generalizable to Vietnam's microbusinesses. However, the better the drivers of flood losses for a
specific sector are understood, the more informed loss assessments can be made and investments towards flood adaptation
improved (Sieg et al., 2017).

Modelling flood losses in low- and middle-income countries is often hampered by the lack of comprehensive and open-source
90 data, which necessitates reliance on primary data collection campaign. The lack of information on flood losses among
microbusinesses is explained by the fact that they mainly operate in the informal sector, which makes it difficult to record and
thus to estimate their flood losses (Garschagen, 2015; Rand and Tarp, 2020). Despite these limitations, only some studies
have analysed and modelled content losses to micro-, small and medium-sized businesses and SMEs companies in South (SE)-
and South-East (SE)-Asia (Chinh et al., 2016; Wijayanti et al., 2017; Samantha, 2018). To the authors knowledge, there is no
95 existing analysis elucidating the drivers of flood losses in microbusinesses in the context of low- and middle-income countries.
However, the identification of the loss drivers is crucial to develop meaningful flood loss models that capture the role of the
drivers in influencing losses (Rözer et al., 2019). The heterogeneity in flood loss processes at the object-level necessitates the
development of multi-variable, probabilistic approaches capable of capturing non-linear effects (Schröter et al., 2014; Vogel

et al., 2014; Rözer et al., 2019; Paprotny et al., 2020; [Paprotny et al., 2021](#); Rafiezadeh Shahi et al., 2024). The absence of such probabilistic loss models in the context of microbusinesses impedes quantification and inclusion of uncertainties for adaptation decision making. Furthermore, multivariate flood loss models are rarely evaluated under conditions other than those under which they were developed, consequently their applicability for spatial/temporal transfers remains unknown (Apel et al., 2009; Gerl et al., 2014; Ootegem et al., 2017; Vogel et al., 2018; Amadio et al., 2019). Our study aims to address these limitations in the state-of-the-art flood loss modelling approaches for microbusinesses in the context of low- and middle-income countries by deriving empirical evidence on the drivers of flood losses ~~to in~~ microbusinesses in HCMC; calibrating and validating ~~a~~ process-based Bayesian Network (BN) models for HCMC that predict content and business interruption losses; and evaluating the transferability of the ~~BN~~ models by applying them on comparable data from a different city (Can Tho). The manuscript is organized ~~into as follows~~ [the following sections shown \(see, Supplementary Information Fig. S1\)](#): Section 2 ~~explains~~ [comprises an overview of the research domain and](#) the empirical survey datasets used in the study, Sect. 3 the methodology implemented including feature selection and the development of ~~the~~ probabilistic flood loss models, Sect. 4 presents and discusses the results of this study, followed by conclusions in Sect. 5.

2 ~~Case studies~~ and ~~d~~Data — ~~Post-flood survey of microbusinesses~~

2.1 ~~Case studies~~

[HCMC is one of the world's most exposed cities to flood risk under current and future conditions \(Hallegatte et al., 2013; Scussolini et al., 2017\). Similar to other SE-Asian metropolises, HCMC lies in a river delta area close to the coast. These densely populated, flat, riverine and coastal regions experience regular flooding in particular during the rainy season \(Garschagen, 2015; Tierolf et al., 2021; Nguyen et al., 2021\). In HCMC, these regular floods are often the result of compound events caused by the simultaneous occurrence of high tides, heavy rainfall and high flows of the Saigon and Dong Nai rivers and their tributaries \(Tran, 2014; Thuy et al., 2019\). Other large cities in the delta areas of South Vietnam also experience regular urban flooding, for instance, Can Tho City in the Mekong Delta \(abbreviated as Can Tho\). Urban floods in Can Tho are predominantly fluvial in nature, such as a major flood event in 2011. Despite the ongoing efforts to improve protection and adaptation measures on private and municipal levels, climate change and the ongoing growth of these important economic centres increase their risk to urban flooding \(Güneralp et al., 2015; Rentschler et al., 2022\). The existing infrastructure and adaptation measures in these cities are unable to counterbalance the new risks caused by intensified flood events and ongoing urban pressure \(e.g. Bouwer, 2011; Bloch et al., 2012; Fornetta and Feyen, 2019; Kreibich et al., 2022\). The overview map in Fig. 1 illustrates the locations of both cities \(HCMC, Can Tho\) used in the case studies of this paper. We define microbusinesses, including household-businesses, according to the definition of the World Bank, as very small businesses with less than ten employees. However, this general definition for microbusinesses needs to be adapted to the regional context of SE- and SE-Asia. Microbusinesses in these countries tend to employ usually less than three people. In most cases, microbusinesses are located on the ground floor of a building with residences on the upper floors, commonly called](#)

Formatiert: Überschrift 1, AbstandVor: 0 Pt.

shophouses in Vietnamese cities. Microbusinesses provide an important source of income for unemployed family members and people with limited opportunities on the labour market, likewise migrant workers and people who received less possibilities of schooling (Samantha, 2018). Together with the operations of small and medium-sized companies (SMEs), microbusinesses drive the rapid economic developments of many SE-Asian states since the last decades (Trinh and Thanh, 2017). According to Vietnam's economic census of 2017, it is estimated that around 75 % of all enterprises are microbusinesses in the country (General Statistics Office, 2018). Vietnam's microbusinesses engage around 11 % of all employees (General Statistics Office, 2018) and the density of microbusinesses is particularly high in economic centres such as HCMC and other delta-cities like Can Tho. The economic importance of HCMC becomes evident when the region's contribution to Vietnam's total economic output is considered - the HCMC region accounts for approximately 40 % of the national GDP (General Statistics Office, 2018). These values highlight the relevance of microbusinesses for Vietnam's local and national economy. The microbusiness owners are particularly vulnerable to the negative consequences of regular flooding due to their limited financial resources and inadequate support by local authorities and the government (Leitold et al., 2020). As a consequence, the owners often rely on their neighbouring network to cope with flooding (Chinh et al., 2016; Chinh et al., 2017; Leitold and Revilla Diez, 2019; Leitold et al., 2021). Bank loans or microcredits are less common due to the usually rather low credit rating (Patankar, 2019). In terms of flood losses this means that repair measures and other business investments are often directly financed by the savings of the microbusiness owner. Insufficient or missing flood insurance policies can further exacerbate the situation of flood-affected businesses (KPMG, 2016; Patankar, 2019). Besides temporal decline in revenues, repair costs or poor future prospects, worse case impacts may include business closures or unemployment among business owners and their employees (Bloch et al., 2012).

~~The flood loss models for microbusinesses are built using empirical data from HCMC and the transferability of the models is evaluated using empirical data from Can Tho. Both datasets are based on in-person structured surveys undertaken with flood-affected microbusinesses. The owner or the manager of the microbusiness was asked to respond to the survey.~~

2.2 Data – post-flood survey of microbusinesses

~~The flood loss models for microbusinesses are built using empirical data from HCMC and the transferability of the models is evaluated using empirical data from Can Tho. Both datasets are based on in-person structured surveys undertaken with flood-affected microbusinesses. The owner or the manager of the microbusiness was asked to respond to the survey. They were informed about the project, how their responses would be used, and that they could leave the survey at any time. No personal or health-related information was collected in either survey. The data is stored and handled exclusively within the German Research Centre for Geosciences (GFZ) in compliance with data privacy and data protection regulations.~~

The workflow of this study presents in the first main step the preparation of the survey datasets (see, Supplementary Information Fig. S1); the respective key aspects are described in detail in the following subsections and in the Supplementary Information, Sect. S1.

2.2.1 Ho Chi Minh City

165 The survey at HCMC was conducted during September-October 2020 and collected responses of 250 microbusinesses which experienced flooding between 2010 and the time of the survey (2020). The majority of microbusinesses surveyed in HCMC are shops or retailers (76 %) mostly selling groceries or other everyday objects. Around 17 % are services, such as restaurants or for reparations, and only 7 % produce consumer goods or processes raw materials. The presented shares of the business sectors in the HCMC survey are representative for entire Vietnam (General Statistics Office, 2018).

170 In order to achieve a reasonable representation of HCMC, we selected the districts with the most frequent flood risk and heterogeneity in socio-economic conditions. Within each district, the shophouses were chosen randomly. The sample size in each district was not chosen based on statistical considerations, but on recommendation from local experts.

175 The interviewees could respond to questions on two flood events – the most severe and the most recent event. However, not every interviewee provided information for both events, which leads to a number of 397 loss records – which leads to 397 loss records in the HCMC dataset. Each record in this dataset comprises information about one or two types of flood losses experienced during an event. In detail, 361 samples of the loss records contain information about business interruption losses reported as relative values (e.g. reduced sales and production), while a similar sample size comprises flood losses to business content but reported as monetary values (e.g. to furniture, electrical devices, stored products and vehicles). The conversion of the latter loss type to relative scales, reduced the their to 317 samples (relative content loss) – number of content loss samples.

180 by using exposure information about the value of business content as exposure information (see, Supplementary Information Sect. S1), as described in the Supplementary Information (Section 4). Consequently, the sample size referring to relative content losses (n=317) is smaller than for relative interruption losses (n=361). Hereinafter both types of relative flood losses are referred to as flood loss variables (Table 1).

185 Figure 1 visualizes the approximate locations of the microbusinesses surveyed in HCMC. However, their exact geolocations are not shown to protect the anonymity of the interviewees. Furthermore, the map shows that the surveyed microbusinesses are often located near to an open channel or tributary river.

190 The questionnaire covers aspects relating to the economic flood losses of microbusinesses and their potential drivers, i.e. flood characteristics, building conditions, business characteristics, undertaken emergency and precautionary measures, the respondent's risk perception and their socio-economic profile. The majority of microbusinesses surveyed in HCMC are shops or retailers (76 %) mostly selling groceries or other everyday objects. Around 17 % are services, such as restaurants or for reparations, and only 7 % produce consumer goods or processes raw materials. The presented shares of the business sectors in the HCMC survey are representative for entire Vietnam (General Statistics Office 2018).

Formatiert: Überschrift 3

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman), Englisch (VereingteStaaten)

hat formatiert: Nicht Hervorheben

hat formatiert: Nicht Hervorheben

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman), Schriftfarbe: Automatisch




Figure 1: Location of surveyed microbusinesses affected by floods between 2010 and 2020 in Ho Chi Minh City (HCMC). The geolocation of one microbusiness was highly inaccurate and thus it is not shown in the map. The overview map on the lower right side shows the location of both study areas (HCMC and Can Tho) in Southeast Asia.

2.2.2 Can Tho

Between August and December 2011 severe flooding affected several districts of Can Tho causing damages to various economic sectors. The survey was undertaken in January-February 2012 and received responses from 373 microbusinesses out of which 313 furnished information on losses to business content and due to business interruption on business interruption losses as well as content losses. The questionnaire was is comparable to the survey undertaken in HCMC, except that each interviewee reported only about the most severe flood event during 2011 and provided information about the value of their business content. The latter was used to calculate relative content losses. Furthermore, the microbusinesses' locations were not queried. All other pre-processing steps were the same as for the HCMC data.

The value distributions of common variables queried by the HCMC and Can Tho survey is are shown in the Supplementary Information, Figure S12. They consisted of about 88 questions covering topics including flood hazard characteristics of the

- hat formatiert: Schriftart: 9 Pt., Fett, Schriftfarbe: Automatisch, Englisch (Vereinigtes Königreich)
- Formatiert: Zeilenabstand: einfach
- hat formatiert: Englisch (Vereinigtes Königreich), Muster: Transparent
- hat formatiert: Muster: Transparent
- hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman), 10 Pt., Nicht Kursiv
- hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman), Nicht Kursiv
- hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman), 10 Pt., Nicht Kursiv

210 2011 flood event, flood preparedness, warning and emergency measures, flood losses to business contents and losses due to
business interruption, risk perceptions, and the business and socio-economic characteristics of the respondents. Compared to
the HCMC survey, the Can Tho dataset includes fewer microbusinesses operating in the trading sector (46 %). Consequently,
more respondents provide services (45 %) or belong to Can Tho's manufacturing sector (9 %). Details on the pre-processing
of the Can Tho survey data are provided in the Supplementary Information, Sect. (Section S1).
In order to derive the drivers of flood losses and to develop the loss models, the 14 pre-processed candidate predictors from
the HCMC dataset are used (Table 1) are used, while the data from Can Tho is used to assess the models' transferability.

Formatiert: Leerraum zwischen asiatischem und westlichem
Text nicht anpassen, Leerraum zwischen asiatischem Text und
Zahlen nicht anpassen

Table 1: Candidate predictors and response variables of flood losses from HCMC

Candidate predictor	Abbreviation	Value range [Mean, Median]	Explanation
Water depth [cm]	water depth	1 – 150 [34; 30]	Water depth [cm] refers to the measured flood water level above the ground floor of the shophouse
Inundation duration [h]	inundation duration	0.2 – 240 [11; 3]	Duration [h] of flood inundation of the shophouse
Contamination (indicator)	contamination	0: no visible 1: light 2: heavier [x; 1]	Type of visible contamination of the flood water
Flow velocity [m/s]	flow velocity	0.1 – 0.5 (calm – turbulent) [x; 0.3]	Flow velocity [m/s] of flood water on the street
Structural precautionary measures (indicator)	structural measures	0.0 – 1.0 [0.2; 0.0]	Ratio between number of implemented measures and number of possible measures. These measures are often implemented during major renovations or building constructions. They comprise the usage of water-resistant building material and the elevation of the building or parts of it.
Non-structural precautionary measures (indicator)	nNon-structural measures	0.0 – 1.0 [0.4; 0.3]	Ratio of number of implemented measures and number of possible measures. These measures need to be purchased before the flood event. They are quite affordable compared to structural measures and comprise wet-proofing of valuables, installation of the electricity control system at a higher level, acquisition of mobile water-barriers and pumping equipment.
Emergency measures (indicator)	emergency measures	0.0 – 1.0 [0.4; 0.5]	Ratio of number of implemented measures and number of possible measures. These measures can be applied shortly before or during the flood event. They comprise saving of documents, relocation of furniture, vehicles or products., usage of sandbags and sealing of doors and windows.
Building age [(years)]	building age	0 – 100 [20; 18]	The age of the shophouses at the time of flooding [(years)]
Building area [(sqm)]	building area	12 – 850 [87; 74]	Building footprint of the shophouse [(sqm)]
Flood experience [n]	flood experience	3 – 151 [82; 76]	Number of experienced floodings between 2010 and 2020 [(n)]
Flood resilience (indicator)	flood resilience	0 – 5 (weak – strong) [x; 3]	Interviewee's appraisal of support by authorities or the neighbourhood

Formatierte Tabelle

Number of employees [number]		<u>no. employees</u>	1 – 9 [x; 2]	Number of employees -[x]
Average monthly income [Euro 2020]		<u>mtlhy. income</u>	18 – 3314 [430; 295]	Available monthly income [Euro 2020] of the interviewee, in most cases the owner of the microbusiness
Average monthly sale [Euro 2020]		<u>mtlhy. sales</u>	92 – 2762 [370; 276]	Averaged revenue from monthly sales and production [Euro 2020] . The variable is representative for the value and quantity of goods and products hold by an individual microbusiness, i.e. it reflects the business size and type.
Flood loss variables				
Relative business interruption loss [%]		<u>rbred</u>	0 – 100 [18.2; 10.0]	Decline in revenues due interrupted business operations (e.g. reduced production and sales) during the flood event [%]. The decline is relative to the potential revenue that would be generated without the flood. 0 % represents no business interruption; 100 % a complete business downtime during the flood event.
Relative content loss [%]	Chance of content loss	<u>chance of rclose</u>	0, 1 (zero-loss, loss) [x; 0]	Chance of flood losses to business content. 0 represents the absence of content loss (zero-loss case); 1 the occurrence of content loss (loss case).
	Degree of content loss [%]	<u>degree of rclose</u>	0.2 – 93.5 [12.3; 4.0]	Flood losses relative to the value of business content, only loss cases [%]; Values close to 0 % represent minor flood losses to business content; 100 % the entire loss of business content.

3 Methodology

Our approach for modelling flood impacts specific to microbusinesses consists of two components (see, Supplementary Information Fig. S1). First, we identify the drivers of content and interruption losses to HCMC’s microbusinesses based on the set of candidate predictors (Table 1). For this feature selection, a variant of Random Forest – Conditional Random Forest – was chosen which since it provides a feature importance method not biased towards correlated predictors (see, Sect. 3.1.4). Second, we calibrate probabilistic loss models – non-parametric Bayesian Networks – specific to microbusinesses based on the identified drivers (see, Sect. 3.2).

Since more than half of the businesses in both cities reported no or only marginal content losses (see, Supplementary Information Fig. S34.a and Fig. S4.b), we model the chance of loss to business content separately from the degree of loss. The former represents the absence or presence of content loss to microbusinesses and is binary (absence/presence), while the latter represents the severity of experienced loss and is a continuous value (0, 100]. Since only a majority 40 % reported zero occurred interruption losses, the aspects of business interruption loss (chance and degree of interruption loss) were not considered separately (see, Supplementary Information Fig. S23). The predictive performances of the Machine Learning (ML)-model used for feature selection and the flood loss models were assessed by the Mean Absolute Error (MAE), Root Mean Square Error (RMSE), Mean Bias Error (MBE) and Symmetric Mean Absolute Percentage Error (SMAPE). The MAE metric is chosen due to its outlier robustness as a selection criterion for the cross-validation of the ML-based models (Chicco et al., 2021). The equations of the performance metrics are listed in the Supplementary Information, in Table S1.

Formatiert: Abstand Nach: 12 Pt.

3.1 Feature selection

3.1.1 Conditional Random Forest

The candidate predictors for flood losses presented in Table 1 exhibit a moderate to high degree of multicollinearity, for instance, the flood-related features are strongly correlated to each other. For this reason, Conditional Inference Trees were applied to account for these correlations during feature selection. The ML model utilised for feature selection is Conditional Inference Trees, Conditional Inference Trees which were initially introduced by Hothorn et al. (2006) and extended by Strobl et al. (2007) to an ensemble of trees, a so-called Conditional Inference Random Forest (CRF). Each tree is grown only by a subset of features, which were identified before as significant based on their p-values (Hothorn et al., 2006). By this approach predictive features are identified, despite their potential collinearity to other candidate predictors. The choice of an unbiased version of the permutation-based feature importance method – namely Conditional Permutation Importance (CPI) - further reduces the chance of biased importance scores for correlated features (CPI; Debeer & Strobl, 2020). The CPI accounts on linear and non-linear interactions of correlated predictors using a Chi-square test (Debeer & Strobl, 2020). Though the CPI is a measure well suited for the feature selection from CRF models (Levshina, 2020), the method is rather computationally expensive, but applicable for the presented approach due to the rather small sets of training samples.

For each of the flood loss variables a CRF model was trained and evaluated via nested cross-validation. The training and evaluation of the CRF model was done via nested cross-validation. Nested cross-validation is a state-of-the-art technique for determining an unbiased generalisation ability of a model (Krstajic et al., 2014). It is particularly recommended for relatively small datasets (Brill, 2022; Liu et al., 2022). Repeated 10 inner folds were used for hyperparameter tuning and 10 outer folds for performance evaluation of the estimators. Of these 10 evaluated estimators, the estimator with the best performance (smallest MAE-score) was used for feature selection, i.e. for identifying the drivers for the degree of content loss and relative interruption loss to microbusinesses.

3.2 Probabilistic flood loss models for microbusinesses

3.2.1 Probabilistic Logistic Regression

The chance of content loss, as one component of relative content loss, is modeled using a probabilistic logistic regression model, applied on the candidate predictors from Table 1. To prevent model overfitting, probabilistic logistic regression incorporates L1 and L2 regularization, which effectively manage multicollinearity in the feature space. The model returns the probability of assigning a microbusiness to either zero-loss or loss category. However, the sample sizes between both categories are imbalanced (see, Supplementary Information Fig. S34.a). To overcome this imbalance, the logistic regression model was trained on a weighted sample of zero-loss and loss cases. Similar as to the CRF (see, Sect. 3.1.1), the logistic regression model was also trained and evaluated by nested cross-validation consisting of 10 inner and 10 outer folds. However, we used

all validated classifiers for modelling the chance of content loss; rather than a single classifier due to their rather moderate predictive performance.

3.2.2 Bayesian Network

Bayesian Networks (BNs) are probabilistic, graphical models with many applications to flood loss modelling (Vogel et al., 2014; Wagenaar et al., 2018; Rözer et al., 2019; Paprotny et al., 2020; Paprotny et al. 2021; Rafieezadeh Shahi et al., 2024).

They are better in regional transfer settingsable compared to than other ML-based models, such as regularized linear regressions, du~~since BNs that they can be applied on incomplete information.~~ Furthermore, ~~t~~They have the benefit of explicitly representing the dependency structures, quantifying uncertainty and the possibility of including expert knowledge alongside data. In more detail, the dependency structure of a BN represents (assumed) causal relations between variables, these dependencies can be set based on knowledge or logical conclusions.

In this study, non-parametric Bayesian Networks, were chosen for modelling the degree of content loss and for modelling the relative business interruption loss. As the term “non-parametric” indicates, this type of Bayesian Network does not rely on prior assumptions about the distribution of the data (Du ~~&~~and Swamy, 2019). Non-parametric BNs were first introduced by Kurowicka ~~and~~ Cooke (2006) and later extended by Hanea et al. (2006; 2015). They rather make use of the ranks of the empirical data which is favourable in terms of the varying distributions of flood losses ~~and its their~~ potential drivers. These drivers are used to construct the graphs of the BNs. Confirmed by the Cramer-von Mises measure for the single variable-pairs of the BN graphs, the joint distributions of the variables are represented by Gaussian copulas.

The constructed flood loss models are calibrated and internally validated on the flood losses reported in HCMC. The performance of the single Bayesian Network model for relative interruption loss was determined by 5-fold cross-validation, while the performance of the modelling approach used for relative content loss was assessed by calculating the prediction bias directly between the reported losses and their probabilistic estimates.

The transferability of these models is assessed based on their performance in predicting flood losses in Can Tho. The performance of the models for each prediction task is benchmarked against the performance of a reference Random Forest (RF) model (Chinh et al., 2017).

4. Results and discussion

The ~~results-section is~~are structured as follows. Firstly, the performance metrics of the CRF model are reported and the most important flood loss drivers for microbusinesses are derived and discussed briefly (see, Sect. 4.1). Subsequently, the identified drivers are used to construct the Bayesian Network flood loss models (see, Sect. 4.2). The ~~loss models -flood loss models-~~are validated (see, Sect. 4.3) ~~and their~~. Finally, the transferability ~~of the models~~ to other delta-cities is tested using the survey data from Can Tho as a case study (see, Sect. 4.4). Finally, the model uncertainties and the limitations of the proposed approach are discussed (see, Sect. 4.5).


hat formatiert: Schriftart: 10 Pt.

hat formatiert: Schriftart: 10 Pt.

hat formatiert: Schriftart: 10 Pt.

4.1 Drivers of flood losses to microbusinesses

The cross-validation of the CRF model shows that all its estimators, validated on the outer folds of the nested cross-validation, have similar moderate performances in predicting the degree of content losses and the relative interruption losses. Furthermore, the similar sets of hyperparameter values across the validated estimators show that the applied ML-algorithm is suitable for both prediction tasks. The prediction of the degree of content losses results in an averaged MAE of 12.8 %, RMSE of 18.4 %, MBE of -0.2 % and SMAPE of 51.4 %, while the prediction of relative interruption losses leads to an averaged MAE of 17.5 %, RMSE of 22.6 %, MBE of 0.3 % and SMAPE of 59.9 %. However, high SMAPE scores are caused by less severe cases of content loss being overestimated, while moderate and severe loss cases are often underestimated by the estimators. The same applies to the prediction of interruption losses. Revenues returned from business operations (mthly. sales) are the most influencing factor for the severity (degree) of loss to business content, while the number of applied emergency measures has the greatest impact on interruption losses. Further main drivers for the degree of content loss and relative interruption loss are the age of the shophouse (building age), hydrological variables and the monthly income (Fig. 2.a and Fig. 2.bure4).






Figure 21: Feature importance of the best-performed Conditional Random Forest (CRF) estimator for predicting (a) the degree of losses to business content and (b) for relative losses due to business interruption losses to microbusinesses in HCMC. Only the five most predictive features are shown.

The identified drivers of flood losses to microbusinesses in HCMC differ partly from those of less flood-experienced companies in high-income countries. The companies' flood experience, its number of employees and the building area are identified in this research as less important loss factors for microbusinesses but were identified as relevant for larger companies in Europe. For instance, in Germany, the company's flood experience, size (number of employees) and the building area were identified as relevant for larger companies (e.g. Kreibich et al., 2007 (flood experience); Sieg et al., 2017 (employees- content loss); Sultana et al., 2018 (employees- interruption loss); Schoppa et al., 2020 (building area)). However, these factors were not found to be critical in the case of HCMC. Of these factors, the missing impact of flood experience could be explained by HCMC's regular floodings which lead to a high level of adaptive behaviour across the residents (Harish et al., 2023).

4.2 Bayesian Network flood loss models

The graph of the non-parametric Bayesian Network for estimating the degree of loss to business content consists of six nodes, the graph for relative business interruption loss of five nodes; the structures of the graphs are visualised in the Figures 23 and Fig. 4. The first parent node of each BN graph was set based on the strongest unconditional rank correlation between a predictor and the flood-target-loss variable for (degree of content loss, and relative interruption loss), respectively. This highest unconditional correlation coefficient exists for both constructed BNs for the variable-pair of water depth inside the building to the corresponding flood loss variable (Spearman's rank coefficient value, rho, for degree of content loss: 0.37; for relative interruption loss: 0.24). However, in the feature space for relative interruption losses, exists an equally strong correlation exists between to the target and the indicator of emergency measures. This feature was identified by the CRF model as the most predictable-predictive for estimating relative interruption losses (Fig. 2.b); but was considered unimportant predictor during the conditionalization of the BN. However, the conditionalization of the BN with emergency measures showed that

Formatiert: Abstand Nach: 12 Pt., Rahmen: Oben: (Kein Rahmen), Unten: (Kein Rahmen), Links: (Kein Rahmen), Rechts: (Kein Rahmen), Zwischen: (Kein Rahmen), Tabstopps: Nicht an 4,75 cm

information about emergency measures becomes unimportant when adding further features to the BN graph, so the corresponding BN graph for relative interruption loss was constructed without it (Fig. 4). The variables for the remaining parent nodes (2. – 5. parent node) were selected based on the strongest conditional ranking correlation using. During this process, the CRF ranking was used as a guideline so that the most important drivers were tested first as to prioritize the testing of potential parent nodes.




Figure 23: Structure of the Bayesian Network for predicting the degree of loss to business content loss (degree of rcloss). The values represent the rank correlation coefficients between the variables (rho).

Formatiert: Zeilenabstand: einfach

The predictors of flood losses and their assumed dependencies in the BN graphs are presented in the following:

- The degree of losses to business content and relative interruption losses, correlate with water depth in the shophouse (water depth). It is the predictor with the strongest rank correlation to both flood loss types (rho: 0.34 in Fig. 3, rho: 0.23 in Fig. 4) and was also previously identified as a relevant predictor by the CRF model. Rising water levels in the building directly increase the potential damage to low-lying goods, equipment and machinery (Kreibich et al., 2010; Chinh et al., 2015; Sieg et al., 2017). Apart from (non-)structural damages, the flooding of business premises itself or indirectly through power outages potentially leads to business interruptions (Kreibich et al., 2009; Sultana et al., 2018).
- High flow velocities (flow velocity) on the streets are associated in the BN graphs with more severe business interruptions as indicated by a correlation coefficient of 0.23 (Fig. 4) but are not important for modelling the degree of content losses. Business activities are potentially affected when high velocities hamper the transportation, such as by relocated objects


blocking streets, or damage infrastructure, such as the energy systems (Jha-Bloch et al., 2012). Additionally, flow velocities have a direct effect on the water level in buildings by pressing water through openings in windows or doors, as also expressed in the BN graph for relative interruption losses of Fig. 4 (rho: 0.36). However, the missing impact of flow velocity on the degree of content loss is explained by the high level of preparedness of HCMC's residents, such as the relocation of vehicles before potential flooding (Chinh et al., 2016), whereas business activities, especially those of shops and small retailers, cannot or can only partially be relocated to other premises.

- Age of the shophouse (**building age**) and degree of content loss have a negative relationship in the BN graph (rho: -0.22 in Fig. 3). The majority of shophouses in the HCMC samples were built in the last 30 years before the flood event, i.e. mainly between the 1980s and late 2000s. These "newer" shophouses reported the most severe content losses and can be explained by the strong urban pressure in these decades. The findings are confirmed by Downes and Storch (2014), Chinh et al. (2015) and Nguyen et al. (2016), who highlight that "newer" buildings in HCMC are more flood-exposed than "older" ones.
- The revenues from business operations (**mtbly. sales**) are positively correlated with the degree of content loss in the respective BN graph as shown in Fig. 3 (rho: 0.29), but only a weak positive correlation exists to relative interruption losses. Monthly sales are seen as an indicator for the microbusiness size and its type of business content. Higher sales have effect on both exposure and vulnerability, as sale volume reflects a heterogeneity among companies in type of contents (Schoppa et al., 2020). The variable of monthly sales has a negative correlation with the indicator for non-structural precautionary measures in the graph for degree of content loss (rho: -0.18 in Fig. 3). This is theoretically explained by the connections within the data: businesses with limited revenues are more likely to acquire non-structural measures before the flood event, as loss of contents would have existential consequences for small retailers compared to more prosperous businesses.
- As shown by the BN graph in Fig. 3 reduces the implementation of non-structural precautionary measures (**non-structural measures**) reduce the severity (degree) of content losses in microbusinesses (rho: -0.23 in Fig. 3), though they-such measures are not relevant for modelling relative interruption losses. The impact of precautionary measures on reducing commercial content losses is well studied (Kreibich et al., 2007; Kreibich et al., 2010; Chinh et al., 2016; Sieg et al., 2017; Schoppa et al., 2020). Non-structural measures usually prevent water from infiltrating into the building, but not in all cases. For instance, Chinh et al. (2016) found that in Can Tho flood water can also come from the sewage system and thus bypass implemented precautionary measures. Consequently, there is no link with water depth in our model due to weak correlation between water depth and non-structural measures.
- The implementation of structural precautionary measures (**structural measures**) has mitigating effects on the severity of content and interruption losses in microbusinesses (rho: -0.19 in Fig. 3, rho: -0.11 in Fig. 4). The moderate dependencies in the BN graphs are in line with the findings of various studies, which highlight the usage of structural measures as an efficient individual precautionary measure (Scussolini et al., 2017; Trinh and Thanh, 2017; Du et al., 2020; Harish et al., 2023). The efficiency of these measures is represented in the BN graphs indirectly by lower water levels in the

390

shophouses and directly in the flood loss variables, e.g. in elevated buildings, there is less chance that flood water will enter the building through overloaded drainage systems.

- A higher number of employees (**no. employees**) is linked with lower interruption losses in the respective BN graph ([rho: -0.13 in Fig. 4](#)). Despite its rather weak negative rank correlation it improves the predictive accuracy of the BN model. The number of employees refers to the availability to human resources on which the business owner can draw on, which in turn affects the possibility to keep the business running during the flood event, [for example, e.g.](#) by relocating important business processes. [The findings of Sultana et al. \(2018\) confirm the number of employees as an important predictor of interruption losses in German companies, though they identify a positive association which is contrary to the findings of this study.](#)



395


Figure 34: Structure of the Bayesian Network for predicting relative business interruption losses (rbred). The values represent the rank correlation coefficients between the variables (rho).

Formatiert: Zeilenabstand: Mehrere 1,15 ze

4.3 Flood loss model validation

4.3.1 Relative content loss

At the first glance, the modelling approach consisting of logistic regression and Bayesian Network seems to perform quite well when predicting relative content losses (MAE: of 3.8 %, RMSE: of 12.3 %). It marginally underestimates losses (MBE: -2.4 %) and has a remarkable low SMAPE of 16.3 % indicating a good level of preciseness of the estimates. The mean value of the modelled relative content losses is of similar magnitude as the observed loss ratios (observed mean: 4.7 %, predicted mean: 4.6 %), as shown also by the clustering of the data points in the lower value range in Fig. 54.a. However, the figure also shows—illustrates—that more severe losses to business content are consistently underestimated by the models.



Formatiert: AbstandNach: 12 Pt., Rahmen: Oben: (Kein Rahmen), Unten: (Kein Rahmen), Links: (Kein Rahmen), Rechts: (Kein Rahmen), Zwischen: (Kein Rahmen)


Figure 45: Scatterplots of observed and modelled relative content losses (rcloss) to HCMC's microbusinesses for (a) the combination of logistic regression (LR) and Bayesian Network (BN) and (b) the reference Random Forest (RF) model used for benchmarking. The grey points represent the observations of zero-loss. The ML-based classifiers assigned to most cases an absence of content loss (zero-loss), thus only one grey point seems to be visualised in Figure 45.a).

The general well predictive performance of the modelling approach is caused by the usually low probability values for the chance of content loss. Having a critical look to the predicted probabilities of chance of content loss, it becomes clear that the observed small prediction bias is caused by the circumstance that the logistic regression estimated instances of chance of content loss usually as zero-loss cases, and thus assigns low probability of losses to most predictor combinations combinations (see, the high share of cases predicted as zero-losses in the left half of Fig. 6.a). (see, centre histogram of Figure 5.a). The high-large numbershare of observations of content loss wrongly predicted as zero-losses further illustrates this (see, False Positives in the lower left corner of Fig. 65.b); only 25 % of the experienced content losses (loss cases) are correctly predicted by the ML-classifiers (see, True Negatives in the lower right corner of Fig. 56.b).


As a consequence, the estimates for relative loss to business content are mostly reduced by more than half as soon as they are multiplied with the predicted probabilities for chance of loss. In particular, the estimates of severe cases of content loss are reduced in their magnitudes. Furthermore, the ML-based classifiers could hardly distinguish between cases with an absence of loss (zero-loss) and small loss fractions (near zero-loss), which further deteriorated the-their calibration and performance-of the-classifiers.

In comparison to the modelling approach, the reference Random Forest model (Chinh et al., 2017) less-does not captures reported cases of zero-loss as such. This is shown when comparing the predicted values of zero-loss cases from the modelling approach (see, grey dots in Fig. 45.a) with the ones from the reference RF model (see, grey dots in Fig. 45.b). However, the general predictive performance is only marginally worse (Table 2). The cross-validated RF-estimators have on average similar magnitudes in the RMSE (12.4 %) and MBE (1.3 %) as the modelling approach, but higher MAE (7.2 %) and SMAPE (78.9 %).


430




435



Distributions of observed and predicted chance of loss



(a)



(b)

Figure 65: (a) Distribution of observed (either 0.0 or 1.0) and predicted probabilities of for chance of content loss from the ML-based classifiers. A vertical dashed line separates the observed and predicted cases of zero-loss from the observed and predicted loss cases, and (b) The corresponding confusion matrix for chance of content loss. The values in front of the brackets are the sample numbers; values in the brackets the sample numbers normalized over the observations.

Formatiert: AbstandNach: 12 Pt., Rahmen: Oben: (Kein Rahmen), Unten: (Kein Rahmen), Links: (Kein Rahmen), Rechts: (Kein Rahmen), Zwischen: (Kein Rahmen)

Formatiert: AbstandNach: 0 Pt.


4.3.2 Relative interruption loss

The cross-validation of the BN model for relative interruption losses results in an averaged MAE of 18.7 %, RMSE of 24.5 %, MBE of 0.17 % and SMAPE of 61.9 %. The modelled mean value in the interruption losses is almost equal to the observed mean of around 18.5 %, yet the variation in the observations is not well represented in the model estimates, as visualised in

440

Figure 6.a. Nearly all reported cases of interruption loss are predicted by the BN with loss fractions between 10 % and 40 %. This is much narrower compared to the variation seen in the reported loss ratios ranging between 0 % to 100 % decrease in business revenues. Additionally, the figure shows that more severe cases of interruption loss are underestimated by the BN, despite their rather frequent occurrence.

445 The reference RF model results in similar high prediction errors as the BN (Table 2). They particularly overestimate cases of zero- and near zero-loss and underestimate severe loss cases (Figure 76.a and Figure 76.b).



Formatiert: Tabstopps: 1,27 cm, Links + 2,54 cm, Links + 3,81 cm, Links + 5,08 cm, Links + 6,35 cm, Links + 7,62 cm, Links + 8,89 cm, Links + 10,16 cm, Links + 11,43 cm, Links + 12,7 cm, Links + 15,04 cm, Links

Figure 76: Scatterplots for observed and modelled relative interruption losses (rbred) to HCMC’s microbusinesses for (a) the Bayesian Network and (b) the reference Random Forest model used for benchmarking. The grey points represent the observations of zero-loss, i.e. the absence of interruption loss.

Formatiert: Abstand Nach: 0 Pt.

4.4 Transferability of the flood loss models

In order to test the transfer capability, In this subsection the results of the transferability of the flood loss models are presented and discussed. The interruption loss model calibrated on microbusinesses in HCMC was applied to predict interruption losses in Can Tho using a comparable survey dataset. The same procedure was applied to the models for content losses. However, the transferred logistic regression model was not able to capture the variation for chance of loss in the Can Tho samples. Thus, the study only presents the results of the transferred Bayesian Network model for interruption losses and the corresponding reference Random Forest model. Cumulative distribution functions (CDFs) were chosen to visualise the main aspects of their model performances.

The generalisation ability of the Bayesian Network BN model on the Can Tho samples results in similar prediction errors than during training on the HCMC samples, except for the SMAPE score. The transfer of the BN leads to a MAE of 17.9 %, RMSE of 23.5 %, MBE of 0.2 % and SMAPE of 23.2 %. The error scores show that the model’s capacity to estimate interruption losses remains unchanged when transferred to Can Tho, in contrast to the transferability of the reference Random RFForest model which resulted in a degraded performance (Table 2). These findings are reflected by cumulative distribution functions (CDFs) in Fig. 8.a and Fig. 8.b. The CDFs for the BN in Figure 7.a and for the reference RF model in Figure 7.b reflect these findings. The cumulative distributions shown in this study represent the change in the predictive accuracy of a model due to regional transfer. In other words, the CDFs provide insight into the extent to which a transferred flood loss model suffers from the different information contained in the Can Tho samples. The CDFs are shown in their normalized version to facilitate comparability despite different sample sizes.

The probability of the BN to predict a Can Tho sample precisely (prediction bias < ±10 %) remains unchanged (Figure 87.a) but drops for the reference RF model from around 45 % (HCMC samples) to 25 % (Can Tho samples) (Figure 78.b). The In most cases the reference RF model underestimates the reported interruption losses in nearly 90 % of the Can Tho samples, but in only 30 % of the HCMC samples (Fig. 8.b), as shown when comparing the MBE scores between the transferred reference RF model and the transferred BN in Table 2.

These findings presented results of this subsection show that the reference RF model is less transferable than the BN, despite both models performing similarly well in their calibration site (i.e., HCMC).

Formatiert: Abstand Nach: 12 Pt.

hat formatiert: Schriftart: Fett




Figure 78: Cumulative distribution function (CDF, normalized) of prediction errors for modelling business interruption losses (rbred) in HCMC and in the transfer region, Can Tho. (a) CDF of the Bayesian Network model performance; (b) CDF of the reference Random Forest model performance. The CDF for the reference RF model is cut by 50 % as no larger prediction errors exist.

Transfer experiments on (Bayesian Network) flood loss models have ~~shown~~ highlighted that model transfer usually leads to a stagnation or drop in the model's performance, in particular, when the new conditions differ remarkably from the calibration region (Schröter et al., 2014; Wagenaar et al., 2018). However, there was no drop in the performance of the BN model performance when transferred across regions. a drop in model performance when regional transferred could not be observed in this study due to very similar local conditions between the calibration (HCMC) and application transfer site (Can Tho).

hat formatiert: Schriftart: 9 Pt.

Formatiert: Abstand Nach: 0 Pt.

hat formatiert: Schriftart: 9 Pt.

These local conditions are reflected in the similar predictor ranges and distributions of both survey datasets (see, Supplementary Information Fig. S4-2). Additionally, the high heterogeneity in the HCMC samples, in particular, in the hydrological, building- and business-related predictors, has the potential to increase the model robustness for new study sites (Wagenaar et al., 2018).

Table 2: Model validation of flood loss models in HCMC and in the transfer region (Can Tho). The different sample sizes are due to the differences in the number of cases reported and the way in which incomplete samples are treated in the models. rcloss: relative loss to business content, rbred: relative loss due to business interruption, LR: probabilistic logistic regression, BN: Bayesian Network, RF: reference Random Forest, x: not applicable

	Model validation [sample size]	MAE [%]	RMSE [%]	MBE [%]	SMAPE [%]
HCMC					
<i>rcloss</i>	LR + BN [284]	3.8	12.3	-2.4	16.3
	RF [284]	7.2	12.4	1.3	78.9
<i>rbred</i>	BN [361]	18.7	24.5	0.17	61.9
	RF [314]	16.4	21.8	1.7	58.6
Can Tho (transfer region)					
<i>rcloss</i>	LR + BN [266]	x	x	x	x
	RF [266]	13.5	19.6	0.8	75.0
<i>rbred</i>	BN [313]	17.9	23.5	0.2	23.2
	RF [267]	25.7	32.6	-23.5	41.1

4.5 **Applicability, Limitations and uncertainties**

Reliable flood loss models are essential tools for stakeholders and practitioners across multiple sectors, including insurance, urban planning, flood risk management, and climate adaptation decision-making. The flood loss models presented in this study specifically address the economic impacts of flooding on microbusinesses in Vietnamese cities, focusing on business interruptions and content losses. To our knowledge, this type of economic sector is underrepresented in risk management and the proposed models can advance decision making with a focus on this sector. By representing key drivers of loss as a graph structures, the models offers an interpretable and transparent framework for understanding how various factors contribute to flood-related damages. The models are based on a non-parametric Bayesian Networks, which enables probabilistic estimation of flood losses while explicitly quantifying uncertainty in both data and model formulation. This feature makes the models particularly robust, allowing for transparent assessment of risk and greater confidence in the results. Unlike traditional deterministic models, the Bayesian approach ensures flexibility in handling incomplete or uncertain data, which is a common challenge in flood loss estimation. The combination of an interpretable model structure and transparent uncertainty

quantification opens the door for operationalizing this modelling approach in practical settings. It provides stakeholders with a clearer understanding of how flood losses are calculated, promoting trust and facilitating decision-making. Furthermore, the model's ability to function effectively even with missing or limited data enhances its transferability to similar geographic regions and contexts. This adaptability is particularly valuable for expanding its application in data-scarce environments or in rapidly urbanizing areas where flood risks are evolving.

Despite these advantages, the results have indicated high uncertainty in reconstructing flood losses from survey data. One possible further analysis would be comparing the model estimates with other studies. However, comparability is limited by the fact that in contrast to our the flood losses were determined at object-level modelling, state-of-the-art while flood loss modelling in low- and middle-income countries is mainly carried out on meso- or macroscale (Booij, 2004; Aerts et al., 2020; Tierolf et al., 2021), with commercial losses reported only in absolute values (Wijayanti et al., 2017; Patankar, 2019; Tsinda et al., 2019) and often without validation (Ke et al., 2012; Patankar and Patwardhan, 2015; Yang et al., 2016).

We did not observe an increase in model uncertainty regional transfer for the Bayesian Network model for interruption losses due to the regional transfer did not affect the width in model uncertainties. Furthermore, the mean values of the empirical flood-interruption losses are for both regions (HCMC, Can Tho) response variables within the uncertainty ranges (within 95 % confidence interval), independent from the region of application. However, as seen above, before the majority of interruption-related losses are underestimated by the flood loss models remarkably.

The example of regional transfer illustrates the potential of non-parametric, continuous Bayesian Network models compared to a Random Forest models. However, since the transfer capability was validated for only one case study, there is a need to models presented here need to be calibrated and validate the models under further local and temporal conditions, to truly estimate specific flood impacts on microbusinesses in new regions.

5. Conclusions

We proposed a first approach to estimate flood losses to microbusinesses by combining expert knowledge with survey data of flood-affected microbusinesses from HCMC and Can Tho in Vietnam. A Conditional Random Forest model was applied to obtain the main drivers of content and interruption losses from a set of heterogeneous samples and potential predictors which are partly correlated to each other. The identified drivers were used to calibrate knowledge-based probabilistic loss models consisting of non-parametric, continuous Bayesian Networks and logistic regression. The findings of this study indicate that information on business revenues from monthly sales and production, building age, and hydrological characteristics of the flood is crucial in estimating content and interruption losses for microbusinesses. The probabilistic flood loss models were calibrated and validated against reported flood losses in HCMC and in a transfer case study of Can Tho, performed in the calibration region (HCMC) with a MAE of 3.8 % for relative content losses and 18.7 % for relative interruption losses. The transfer of the interruption model to another delta city (Can Tho) illustrates the generally good transferability of probabilistic models. However, the calibration of the developed models under further local and temporal

~~conditions would improve their applicability for decision makers. The study resulted in interpretable and transferrable probabilistic flood loss models for predicting content losses and business interruption losses to microbusinesses. In addition, the developed loss-models are openly provided and integrating them to flood risk assessments will has the potential to advance risk management decision making with a focus on microbusinesses.~~

550 *Data and Code Availability.* The survey data will be made openly available in the HOWAS21 database (<https://howas21.gfz-potsdam.de/>) after an embargo of three years after the end of the project (in 2027). The data can be accessed in the meantime from the authors. Source code (python) is openly available at <https://github.com/A-Buch/flood-loss-models-4-HCMC/tree/microbusiness-paper>. The Bayesian Network flood loss models are created with the PyBanshee toolbox (Koot et al., 2023, https://github.com/mike-mendoza/py_banshee.git); the Conditional Random Forest models ~~s are is~~ based on the R package partykit (Hothorn ~~and~~ Zeileis, 2015, <https://www.jmlr.org/papers/v16/hothorn15a.html>).

Author contributions. Conceptualisation: AB, DP, NS; Data curation, formal analysis and visualization: AB; Methodology: AB, DP, KRS, HK, NS; Supervision: DP, NS; Writing – original draft: AB; Writing – review & editing: AB, DP, KRS, HK, NS

560 *Competing interests.* ~~Co-author Heidi Kreibich is member of the editorial board of Natural Hazards and Earth System Sciences. Some authors are members of the editorial board of NHESS.~~

Funding. The data collection was undertaken by project DECIDER (grant nos. 01LZ1703G and 01LZ1703A) and WISDOM II (“Water related Information System for a Sustainable development of the Mekong Delta”) funded by the BMBF (German Ministry for Education and Research). Nivedita Sairam is funded by project HI-Clif (grant no. 01LN2209A) funded by the BMBF.

References

570 Aerts, J. P. M., Uhlemann-Elmer, S., Eilander, D., and Ward, P. J.: Comparison of estimates of global flood models for flood hazard and exposed gross domestic product: a China case study, Nat. Hazard Earth Sys., 20, 3245–3260, <https://doi.org/10.5194/nhess-20-3245-2020>, 2020.

Amadio, M., Scorzini, A. R., Carisi, F., Essenfelder, A. H., Domeneghetti, A., Mysiak, J., and Castellarin, A.: Testing empirical and synthetic flood damage models: the case of Italy, Nat. Hazard Earth Sys., 19, 661–678, <https://doi.org/10.5194/nhess-19-661-2019>, 2019.

hat formatiert: Schriftfarbe:Automatisch

hat formatiert: Absatz-Standardschriftart,Schriftart: (Standard)Times New Roman, Schriftfarbe:Automatisch

hat formatiert: Schriftfarbe:Automatisch

hat formatiert: Überschrift 1 Zchn, Schriftart: 10 Pt., Kursiv, Englisch (VereinigtesKönigreich)

hat formatiert: Schriftart: 10 Pt., Nicht Fett, Englisch (VereinigtesKönigreich)

Formatiert: Überschrift 1, Zeilenabstand: 1,5 Zeilen

hat formatiert: Schriftart: (Standard)Times New Roman

Formatiert: Standard, Zeilenabstand: einfach

hat formatiert: Schriftart: Nicht Kursiv

Formatiert: Standard

hat formatiert: Schriftart: Nicht Kursiv

hat formatiert: Schriftart: (Standard)+Überschriften(Times New Roman)

Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: 1,5 Zeilen

hat formatiert: Schriftart: (Standard)+Überschriften(Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften(Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften(Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften(Times New Roman)

Apel, H., Aronica, G. T., Kreibich, H., and Thieken, A. H.: Flood risk analyses—how detailed do we need to be?, Nat. Hazards, 49, 79–98, <https://doi.org/10.1007/s11069-008-9277-8> <https://doi.org/10.1007/s11069-008-9277-8>, 2009.

575 Bloch, R., Jha, A. K., and Lamond, J.: Cities and Flooding: A Guide to Integrated Urban Flood Risk Management for the 21st Century, The World Bank, <https://doi.org/10.1596/978-0-8213-8866-2>, 2012.

Booij, M. J.: Flood Damage Assessment and Modelling in the Red River basin in Vietnam, International Workshop on Flood Controls Decision Support Systems (FLOCODS), in: International Workshop on Flood Controls Decision Support Systems (FLOCODS), Hanoi, Vietnam, 10-12 February 2004, 1-14, available at: <https://www.utwente.nl/en/et/cem/research/wem/people-attachments/booij/publications/44.pdf> <https://www.utwente.nl/en/et/cem/research/wem/people-attachments/booij/publications/44.pdf> (last access: 20 January 2025), 2004.

580 Bouwer, L. M.: Have Disaster Losses Increased Due to Anthropogenic Climate Change?, B. Am. Meteorol. Soc., 92, 39–46, <https://doi.org/10.1175/2010BAMS3092.1> <https://doi.org/10.1175/2010BAMS3092.1>, 2011.

585 Brill, F. A.: Applications of machine learning and open geospatial data in flood risk modelling, PhD Thesis, University Potsdam, <https://doi.org/10.25932/PUBLISHUP-55594> <https://doi.org/10.25932/PUBLISHUP-55594>, 2022.

Chicco, D., Warrens, M. J., and Jurman, G.: The coefficient of determination R-squared is more informative than SMAPE, MAE, MAPE, MSE and RMSE in regression analysis evaluation, PeerJ., 7:e623, <https://doi.org/10.7717/peerj-es.623> <https://doi.org/10.7717/peerj-es.623>, 2021.

590 Chinh, D. T., Gain, A., Dung, N. V., Haase, D., and Kreibich, H.: Multi-Variate Analyses of Flood Loss in Can Tho City, Mekong Delta, Water, 8, 6, <https://doi.org/10.3390/w8010006> <https://doi.org/10.3390/w8010006>, 2015.

Chinh, D. T., Bubeck, P., Dung, N. V., and Kreibich, H.: The 2011 flood event in the Mekong Delta: preparedness, response, damage and recovery of private households and small businesses, Disasters, 40, 753–778, <https://doi.org/10.1111/disa.12171> <https://doi.org/10.1111/disa.12171>, 2016.

595 Chinh, D. T., Dung, N. V., Gain, A., and Kreibich, H.: Flood Loss Models and Risk Analysis for Private Households in Can Tho City, Vietnam, Water, 9, 313, <https://doi.org/10.3390/w9050313> <https://doi.org/10.3390/w9050313>, 2017.

Debeer, D. and Strobl, C.: Conditional permutation importance revisited, BMC Bioinformatics, 21, 307, <https://doi.org/10.1186/s12859-020-03622-2> <https://doi.org/10.1186/s12859-020-03622-2>, 2020.

Downes, N. K. and Storch, H.: Current Constraints and Future Directions for Risk Adapted Land-Use Planning Practices in the High-Density Asian Setting of Ho Chi Minh City. Planning Practice & Research, 29, 220–237, <https://doi.org/10.1080/02697459.2014.929835>, 2014.

600 Downes, N. K., Storch, H., Schmidt, M., Nguyen, T. C. V., Dinh, L. C., Tran, T. N., and Hoa, L. T.: Understanding Ho Chi Minh City's Urban Structures for Urban Land-Use Monitoring and Risk Adapted Land-Use Planning, in: Sustainable Ho Chi Minh City: Climate Policies for Emerging Mega Cities, edited by: Katzschner, A., Waibel, M., Schwede, D., Katzschner, L., Schmidt, M., and Storch, H., Springer, Cham, 89–116, https://doi.org/10.1007/978-3-319-04615-0_6, 2016.

605

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

Formatiert: Abstand Nach: 0 Pt.

Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: 1,5 Zeilen

- Du, K.-L. and Swamy, M. N. S.: Neural Networks and Statistical Learning, Springer London, London, <https://doi.org/10.1007/978-1-4471-7452-3><https://doi.org/10.1007/978-1-4471-7452-3>, 2019.
- Du, S., Scussolini, P., Ward, P. J., Zhang, M., Wen, J., Wang, L., Koks, E., Diaz-Loaiza, A., Gao, J., Ke, Q., and Aerts, J. C. J. H.: Hard or soft flood adaptation? Advantages of a hybrid strategy for Shanghai, *Global Environ. Chang.*, 61, 102037, <https://doi.org/10.1016/j.gloenvcha.2020.102037><https://doi.org/10.1016/j.gloenvcha.2020.102037>, 2020.
- Formetta, G. and Feyen, L.: Empirical evidence of declining global vulnerability to climate-related hazards, *Global Environ. Chang.*, 57, 101920, <https://doi.org/10.1016/j.gloenvcha.2019.05.004><https://doi.org/10.1016/j.gloenvcha.2019.05.004>, 2019.
- GADM: Download GADM data (version 4.1). GADM Maps and Data, https://gadm.org/download_country.html (last access: 20 January 2025), 2022.
- Garschagen, M.: Risky Change? Vietnam's Urban Flood Risk Governance between Climate Dynamics and Transformation, *Pac. Aff.*, 88, 599–621, <https://doi.org/10.5509/2015883599><https://doi.org/10.5509/2015883599>, 2015.
- General Statistics Office: Results of the 2017 economic census, Statistical Publishing House, [available at: https://www.gso.gov.vn/en/data-and-statistics/2019/03/results-of-the-2017-economic-census/](https://www.gso.gov.vn/en/data-and-statistics/2019/03/results-of-the-2017-economic-census/)<https://www.gso.gov.vn/en/data-and-statistics/2019/03/results-of-the-2017-economic-census/> (last access: 20 January 2025), 2018.
- Gerl, T., Bochow, M., and Kreibich, H.: Flood Damage Modeling on the Basis of Urban Structure Mapping Using High-Resolution Remote Sensing Data, *Water*, 6, 2367–2393, <https://doi.org/10.3390/w6082367><https://doi.org/10.3390/w6082367>, 2014.
- Güneralp, B., Güneralp, I., and Liu, Y.: Changing global patterns of urban exposure to flood and drought hazards, *Global Environ. Chang.*, 31, 217–225, <https://doi.org/10.1016/j.gloenvcha.2015.01.002><https://doi.org/10.1016/j.gloenvcha.2015.01.002>, 2015.
- Hallegatte, S.: An Adaptive Regional Input-Output Model and its Application to the Assessment of the Economic Cost of Katrina, *Risk Anal.*, 28, 779–799, <https://doi.org/10.1111/j.1539-6924.2008.01046.x><https://doi.org/10.1111/j.1539-6924.2008.01046.x>, 2008.
- Hallegatte, S., Green, C., Nicholls, R. J., and Corfee-Morlot, J.: Future flood losses in major coastal cities, *Nat. Clim. Change*, 3, 802–806, <https://doi.org/10.1038/nclimate1979><https://doi.org/10.1038/nclimate1979>, 2013.
- Hanea, A. M., Kurowicka, D., and Cooke, D.: Hybrid Method for Quantifying and Analyzing Bayesian Belief Nets, *Qual. Reliab. Eng. Int.*, 22, 709–729, <https://doi.org/10.1002/qre.808><https://doi.org/10.1002/qre.808>, 2006.
- Hanea, A. M., Morales Napoles, O., and Ababei, D.: Non-parametric Bayesian networks: Improving theory and reviewing applications, *Reliab. Eng. Syst. Safe.*, 144, 265–284, <https://doi.org/10.1016/j.res.2015.07.027><https://doi.org/10.1016/j.res.2015.07.027>, 2015.
- Harish, T. V., Sairam, N., Yang, L. E., Garschagen, M., and Kreibich, H.: Identifying the drivers of private flood precautionary measures in Ho Chi Minh City, Vietnam, *Nat. Hazard. Earth Sys.*, 23, 1125–1138, <https://doi.org/10.5194/nhess-23-1125-2023><https://doi.org/10.5194/nhess-23-1125-2023>, 2023.

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman), Englisch (Vereinigtes Königreich)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman), Englisch (Vereinigtes Königreich)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman), Englisch (Vereinigtes Königreich)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: 1,5 Zeilen

font formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

font formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

... hat formatiert

hat formatiert	...
hat formatiert	

hat formatiert

hat formatiert

hat formatiert

hat formatiert

hat formatiert

hat formatiert

hat formatiert

hat formatiert	...
hat formatiert	...

hat formatiert

640 Hothorn, T. and Zeileis, A.: partykit: A Modular Toolkit for Recursive Partytioning in R, J. Mach. Learn. Res., 16, 3905–3909,
available at: <https://www.jmlr.org/papers/volume16/hothorn15a/hothorn15a.pdf><https://www.jmlr.org/papers/volume16/hothorn15a/hothorn15a.pdf> (last access: 20 January), 2015.

Hothorn, T., Hornik, K., and Zeileis, A.: Unbiased Recursive Partitioning: A Conditional Inference Framework, Journal of
645 Computational and Graphical Statistics, 15, 651–674,
<https://doi.org/10.1198/106186006X133933><https://doi.org/10.1198/106186006X133933>, 2006.

Bloch, R., Jha, A. K., and Lamond, J.: Cities and Flooding: A Guide to Integrated Urban Flood Risk Management for the 21st
Century, The World Bank, <https://doi.org/10.1596/978-0-8213-8866-2>, 2012.

Ke, Q., Jonkman, S. N., Van Gelder, P. H. A. J. M., and Rijcken, T.: Flood damage estimation for downtown Shanghai –
650 sensitivity analysis, in: Conference of the international society for integrated disaster risk management, Beijing, China, 7-9
September 2012, 260–273, available at: <https://research.tudelft.nl/en/publications/flood-damage-estimation-for-downtown-shanghai-sensitivity-analysis><https://research.tudelft.nl/en/publications/flood-damage-estimation-for-downtown-shanghai-sensitivity-analysis> (last access: 20 January 2025), 2012.

Koks, E. E. and Thissen, M.: A Multiregional Impact Assessment Model for disaster analysis, Econ. Syst. Res., 28, 429–449,
655 <https://doi.org/10.1080/09535314.2016.1232701><https://doi.org/10.1080/09535314.2016.1232701>, 2016.

Koot, P., Mendoza-Lugo, M. A., Paprotny, D., Morales-Nápoles, O., Ragno, E., and Worm, D. T. H.: PyBanshee version (1.0):
A Python implementation of the MATLAB toolbox BANSHEE for Non-Parametric Bayesian Networks with updated features,
SoftwareX, 21, 101279, <https://doi.org/10.1016/j.softx.2022.101279><https://doi.org/10.1016/j.softx.2022.101279>, 2023.

KPMG: Preparing MSMEs for effective disaster management, KPMG, available at:
660 <https://assets.kpmg.com/content/dam/kpmg/pdf/2016/05/Disaster-Management-Preparedness-SME.pdf><https://assets.kpmg.com/content/dam/kpmg/pdf/2016/05/Disaster-Management-Preparedness-SME.pdf> (last access:
20 January 2025), 2016.

Kreibich, H., Müller, M., Thielen, A. H., and Merz, B.: Flood precaution of companies and their ability to cope with the flood
in August 2002 in Saxony, Germany, Water Resour. Res., 43, W03408,
665 <https://doi.org/10.1029/2005WR004691><https://doi.org/10.1029/2005WR004691>, 2007.

Kreibich, H., Piroth, K., Seifert, I., Maiwald, H., Kunert, U., Schwarz, J., Merz, B., and Thielen, A. H.: Is flow velocity a
significant parameter in flood damage modelling?, Nat. Hazard. Earth Sys., 9, 1679–1692, <https://doi.org/10.5194/nhess-9-1679-2009><https://doi.org/10.5194/nhess-9-1679-2009>, 2009.

Kreibich, H., Seifert, I., Merz, B., and Thielen, A. H.: Development of FLEMOcs – a new model for the estimation of flood
670 losses in the commercial sector, Hydrolog. Sci. J., 55, 1302–1314,
<https://doi.org/10.1080/02626667.2010.529815><https://doi.org/10.1080/02626667.2010.529815>, 2010.

Kreibich, H., Van Loon, A. F., Schröter, K., Ward, P. J., Mazzoleni, M., Sairam, N., Abeshu, G. W., Agafonova, S.,
AghaKouchak, A., Aksoy, H., Alvarez-Garretón, C., Aznar, B., Balkhi, L., Barendrecht, M. H., Biancamaria, S., Bos-

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

Burgering, L., Bradley, C., Budiyo, Y., Buytaert, W., Capewell, L., Carlson, H., Cavus, Y., Couasnon, A., Coxon, G.,
675 Daliakopoulos, I., De Ruiter, M. C., Delus, C., Erfurt, M., Esposito, G., François, D., Frappart, F., Freer, J., Frolova, N., Gain,
A. K., Grillakis, M., Grima, J. O., Guzmán, D. A., Huning, L. S., Ionita, M., Kharlamov, M., Khoi, D. N., Kieboom, N.,
Kireeva, M., Koutroulis, A., Lavado-Casimiro, W., Li, H.-Y., LLasat, M. C., Macdonald, D., Mård, J., Mathew-Richards, H.,
McKenzie, A., Mejia, A., Mendiando, E. M., Mens, M., Mobini, S., Mohor, G. S., Nagavciuc, V., Ngo-Duc, T., Thao Nguyen
Huynh, T., Nhi, P. T. T., Petrucci, O., Nguyen, H. Q., Quintana-Seguí, P., Razavi, S., Ridolfi, E., Riegel, J., Sadik, M. S.,
680 Savelli, E., Sazonov, A., Sharma, S., Sørensen, J., Arguello Souza, F. A., Stahl, K., Steinhausen, M., Stoelzle, M., Szalińska,
W., Tang, Q., Tian, F., Tokarczyk, T., Tovar, C., Tran, T. V. T., Van Huijgevoort, M. H. J., Van Vliet, M. T. H., Vorogushyn,
S., Wagener, T., Wang, Y., Wendt, D. E., Wickham, E., Yang, L., Zambrano-Bigiarini, M., Blöschl, G., and Di Baldassarre,
G.: The challenge of unprecedented floods and droughts in risk management, *Nature*, 608, 80–86,
<https://doi.org/10.1038/s41586-022-04917-5><https://doi.org/10.1038/s41586-022-04917-5>, 2022.

Krstajic, D., Buturovic, L. J., Leahy, D. E., and Thomas, S.: Cross-validation pitfalls when selecting and assessing regression
685 and classification models, *J. Cheminformatics*, 6, 10, <https://doi.org/10.1186/1758-2946-6-10>[https://doi.org/10.1186/1758-](https://doi.org/10.1186/1758-2946-6-10)
[2946-6-10](https://doi.org/10.1186/1758-2946-6-10), 2014.

Kurowicka, D. and Cooke, R.: Uncertainty Analysis with High Dimensional Dependence Modelling, 1st ed., Wiley Ser. Prob.
Stat., <https://doi.org/10.1002/0470863072><https://doi.org/10.1002/0470863072>, 2006.

690 Leitold, R. and Revilla Diez, J.: Exposure of manufacturing firms to future sea level rise in Ho Chi Minh City, Vietnam, *J.-*
Maps, 15, 13–20, <https://doi.org/10.1080/17445647.2018.1548385><https://doi.org/10.1080/17445647.2018.1548385>, 2019.

Leitold, R., Revilla Diez, J., and Tran, V.: Are we expecting too much from the private sector in flood adaptation? Scenario-
based field experiments with small- and medium-sized firms in Ho Chi Minh City, Vietnam, *Climatic Change*, 163, 359–378,
<https://doi.org/10.1007/s10584-020-02888-y><https://doi.org/10.1007/s10584-020-02888-y>, 2020.

695 Leitold, R., Garschagen, M., Tran, V., and Revilla Diez, J.: Flood risk reduction and climate change adaptation of
manufacturing firms: Global knowledge gaps and lessons from Ho Chi Minh City, *Int. J. Disast. Risk Re.*, 61, 102351,
<https://doi.org/10.1016/j.ijdrr.2021.102351><https://doi.org/10.1016/j.ijdrr.2021.102351>, 2021.

Levshina, N.: Conditional Inference Trees and Random Forests, in: A Practical Handbook of Corpus Linguistics, edited by:
Paquot, M. and Gries, S. Th., Cham: Springer, Cham, —611–643, [https://doi.org/10.1007/978-3-030-46216-](https://doi.org/10.1007/978-3-030-46216-1_25)
700 [1-25](https://doi.org/10.1007/978-3-030-46216-1_25)https://doi.org/10.1007/978-3-030-46216-1_25, 2020.

Liu, X., Kounadi, O., and Zurita-Milla, R.: Incorporating Spatial Autocorrelation in Machine Learning Models Using Spatial
Lag and Eigenvector Spatial Filtering Features, *ISPRS Int. J. Geo-Inf.*, 11, 242,
<https://doi.org/10.3390/ijgi11040242><https://doi.org/10.3390/ijgi11040242>, 2022.

Merz, B., Kreibich, H., Schwarze, R., and Thieken, A.: Review article “Assessment of economic flood damage,” *Nat. Hazard.*
705 *Earth Sys.*, 10, 1697–1724, <https://doi.org/10.5194/nhess-10-1697-2010><https://doi.org/10.5194/nhess-10-1697-2010>, 2010.

Meyer, V., Becker, N., Markantonis, V., Schwarze, R., Van Den Bergh, J. C. J. M., Bouwer, L. M., Bubeck, P., Ciavola, P.,
Genovese, E., Green, C., Hallegatte, S., Kreibich, H., Lequeux, Q., Logar, I., Papyrakis, E., Pfurtscheller, C., Poussin, J.,

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) +Überschriften (Times New Roman)

Przyluski, V., Thieken, A. H., and Viavattene, C.: Review article: Assessing the costs of natural hazards – state of the art and knowledge gaps, *Nat. Hazard. Earth Sys.*, 13, 1351–1373, <https://doi.org/10.5194/nhess-13-1351-2013>, 2013.

Nguyen, T. B., Samsura, D. A. A., Van Der Krabben, E., and Le, A.-D.: Saigon-Ho Chi Minh City, *Cities*, 50, 16–27, <https://doi.org/10.1016/j.cities.2015.08.007>, 2016.

Nguyen, M. T., Sebesvari, Z., Souvignet, M., Bachofer, F., Braun, A., Garschagen, M., Schinkel, U., Yang, L. E., Nguyen, L. H. K., Hochschild, V., Assmann, A., and Hagenlocher, M.: Understanding and assessing flood risk in Vietnam: Current status, persisting gaps, and future directions, *Journal of Flood Risk Management*, 14, e12689, <https://doi.org/10.1111/jfr3.12689>, 2021.

Ootegem, L. V., Herck, K. V., Creten, T., Verhofstadt, E., Foresti, L., Goudenhoofd, E., Reyniers, M., Delobbe, L., Tuyls, D. M., and Willems, P.: Exploring the potential of multivariate depth-damage and rainfall-damage models, *J. Flood Risk Management*, 11, 916–929, <https://doi.org/10.1111/jfr3.12284>, 2017.

Paprotny, D., Kreibich, H., Morales-Nápoles, O., Terefenko, P., and Schröter, K.: Estimating exposure of residential assets to natural hazards in Europe using open data, *Nat. Hazard. Earth Sys.*, 20, 323–343, <https://doi.org/10.5194/nhess-20-323-2020>, 2020.

Paprotny, D., Kreibich, H., Morales-Nápoles, O., Wagenaar, D., Castellarin, A., Carisi, F., Bertin, X., Merz, B., and Schröter, K.: A probabilistic approach to estimating residential losses from different flood types, *Nat. Hazards*, 105, 2569–2601, <https://doi.org/10.1007/s11069-020-04413-x>, 2021.

Patankar, A.: Impacts of Natural Disasters on Households and Small Businesses in India, *Asian Development Bank Economics Working Paper Series No. 603*, <https://doi.org/10.2139/ssrn.3590902>, 2019.

Patankar, A. and Patwardhan, A.: Estimating the uninsured losses due to extreme weather events and implications for informal sector vulnerability: a case study of Mumbai, India, *Nat. Hazards*, 80, 285–310, <https://doi.org/10.1007/s11069-015-1968-3>, 2015.

Rafiezadeh Shahi, K., Sairam, N., Schoppa, L., Sang, L. T., Tan, D. L. H., and Kreibich, H.: BN-FLEMOA: A Bayesian Network-based Flood Loss Estimation Model for Adaptation Planning in Ho Chi Minh City, Vietnam, *ESS Open Archive [preprint]*, <https://doi.org/10.22541/essoar.172081523.38063336/v1>, 12 July 2024.

Rand, J. and Tarp, F.: *Micro, Small, and Medium Enterprises in Vietnam*, 1st ed., Oxford University Press, Oxford, <https://doi.org/10.1093/oso/9780198851189.003.0001>, 2020.

Rentschler, J., Salhab, M., and Jafino, B. A.: Flood exposure and poverty in 188 countries, *Nat. Commun.*, 13, 3527, <https://doi.org/10.1038/s41467-022-30727-4>, 2022.

Rözer, V., Kreibich, H., Schröter, K., Müller, M., Sairam, N., Doss-Gollin, J., Lall, U., and Merz, B.: Probabilistic Models Significantly Reduce Uncertainty in Hurricane Harvey Pluvial Flood Loss Estimates, *Earth's Future*, 7, 384–394, <https://doi.org/10.1029/2018EF001074>, 2019.

- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- Formatiert: Abstand Vor: 0 Pt., Nach: 0 Pt., Zeilenabstand: 1,5 Zeilen
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- Formatiert: Abstand Nach: 0 Pt., Zeilenabstand: 1,5 Zeilen
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)
- hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)

Samantha, G.: The Impact of Natural Disasters on Micro, Small and Medium Enterprises (MSMEs): A Case Study on 2016 Flood Event in Western Sri Lanka, *Procedia Engineer.*, 212, 744–751, <https://doi.org/10.1016/j.proeng.2018.01.096>, 2018.

745 Schoppa, L., Sieg, T., Vogel, K., Zöller, G., and Kreibich, H.: Probabilistic Flood Loss Models for Companies, *Water Resour. Res.*, 56, e2020WR027649, <https://doi.org/10.1029/2020wr027649>, 2020.

Schröter, K., Kreibich, H., Vogel, K., Riggelsen, C., Scherbaum, F., and Merz, B.: How useful are complex flood damage models?, *Water Resour. Res.*, 50, 3378–3395, <https://doi.org/10.1002/2013wr014396>, 2014.

750 Scussolini, P., Tran, T. V. T., Koks, E., Diaz-Loaiza, A., Ho, P. L., and Lasage, R.: Adaptation to Sea Level Rise: A Multidisciplinary Analysis for Ho Chi Minh City, Vietnam, *Water Resour. Res.*, 53, 10841–10857, <https://doi.org/10.1002/2017wr021344>, 2017.

Sieg, T., Vogel, K., Merz, B., and Kreibich, H.: Tree-based flood damage modeling of companies: Damage processes and model performance, *Water Resour. Res.*, 53, 6050–6068, <https://doi.org/10.1002/2017wr020784>, 2017.

755 Sieg, T., Schinko, T., Vogel, K., Mechler, R., Merz, B., and Kreibich, H.: Integrated assessment of short-term direct and indirect economic flood impacts including uncertainty quantification, *PLOS ONE*, 14, e0212932, <https://doi.org/10.1371/journal.pone.0212932>, 2019.

Strobl, C., Boulesteix, A.-L., Zeileis, A., and Hothorn, T.: Bias in random forest variable importance measures: Illustrations, sources and a solution, *BMC Bioinformatics*, 8, 25, <https://doi.org/10.1186/1471-2105-8-25>, 2007.

760 Sultana, Z., Sieg, T., Kellermann, P., Müller, M., and Kreibich, H.: Assessment of Business Interruption of Flood-Affected Companies Using Random Forests, *Water*, 10, 1049, <https://doi.org/10.3390/w10081049>, 2018.

765 Thuy, N. B., Tien, T. Q., Wettre, C., and Hole, L. R.: Monsoon-Induced Surge during High Tides at the Southeast Coast of Vietnam: A Numerical Modeling Study, *Geosciences*, 9, 72, <https://doi.org/10.3390/geosciences9020072>, 2019.

Tierolf, L., De Moel, H., and Van Vliet, J.: Modeling urban development and its exposure to river flood risk in Southeast Asia, *Comp. Environ. Urban*, 87, 101620, <https://doi.org/10.1016/j.compenvurbsys.2021.101620>, 2021.

770 Tran, T. N.: Improvement of flood risk assessment under climate change in Ho Chi Minh City with GIS applications, PhD Thesis, University of Technology in Cottbus - Senftenberg, 186 pp., available at: https://opus4.kobv.de/opus4-btu/files/3078/Tran_Thong_Nhat_Thesis_2014.pdf (last access: 20 January 2025), 2014.

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard)+Überschriften (Times New Roman)

775 Trinh, P. T. T. and Thanh, N. D.: Development Characteristics of SME Sector in Vietnam: Evidence from the Vietnam Enterprise Census 2006-2015 (Working Paper WP-18), working paper, VEPR [Viet Nam Institute for Economic and Policy Research, supported by the Friedrich Naumann Foundation for Freedom], Hanoi, Vietnam, ~~available at:~~ http://vepr.org.vn/upload/533/20171222/EN_VEPR%20WP%2018.pdf~~http://vepr.org.vn/upload/533/20171222/EN_VEPR%20WP%2018.pdf~~ (last access: 20 January 2025), 2017.

780 Tsinda, A., Kind, C., Hess, J. S., Mugsiha, R., and Bizoza, A. R.: Estimating damage costs of flooding on small- and medium-sized enterprises in Kigali, Rwanda, Jambá Journal of Disaster Risk Studies, 11, a755, <https://doi.org/10.4102/jamba.v11i1.755>~~https://doi.org/10.4102/jamba.v11i1.755~~, 2019.

U.S. Geological Survey: Earth Explorer, <https://earthexplorer.usgs.gov/> (last access: 20 January 2025), n.d.

Vogel, K., Riggelsen, C., Scherbaum, F., Schröter, K., Kreibich, H., and Merz, B.: Challenges for Bayesian network learning in a flood damage assessment application, in: Safety, Reliability, Risk and Life-Cycle Performance of Structures and Infrastructures, edited by: Deodatis, G., Ellingwood, B. R., Frangopol, Dan M, D. M., CRC Press, 3123–3130, <https://doi.org/10.1201/b16387-452>~~https://doi.org/10.1201/b16387-452~~, 2014.

785 Vogel, K., Weise, L., Schröter, K., and Thieken, A. H.: Identifying Driving Factors in Flood-Damaging Processes Using Graphical Models, Water Resour. Res., 54, 8864–8889, <https://doi.org/10.1029/2018wr022858>~~https://doi.org/10.1029/2018wr022858~~, 2018.

790 Wagenaar, D., Lüdtke, S., Schröter, K., Bouwer, L. M., and Kreibich, H.: Regional and Temporal Transferability of Multivariable Flood Damage Models, Water Resour. Res., 54, 3688–3703, <https://doi.org/10.1029/2017wr022233>~~https://doi.org/10.1029/2017wr022233~~, 2018.

Wijayanti, P., Zhu, X., Hellegers, P., Budiyo, Y., and Van Ierland, E. C.: Estimation of river flood damages in Jakarta, Indonesia, Nat. Hazards, 86, 1059–1079, <https://doi.org/10.1007/s11069-016-2730-1>, 2017.

795 Yang, L., Kajitani, Y., Tatano, H., and Jiang, X.: A methodology for estimating business interruption loss caused by flood disasters: insights from business surveys after Tokai Heavy Rain in Japan, Nat. Hazards, 84, 411–430, <https://doi.org/10.1007/s11069-016-2534-3>, 2016.

hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)

hat formatiert: Schriftart: (Standard) + Überschriften (Times New Roman)

Formatiert: Einzug: Links: 0 cm, Erste Zeile: 0 cm