

Review article: Understanding the placement of fire emissions from the Brazilian Cerrado biome in the global carbon budget Fire emissions in the Brazilian Cerrado: dynamics, estimates, management and their role in the global carbon budget

5 Renata Moura da Veiga¹, Celso von Randow¹, Chantelle Burton², Douglas I. Kelley³, Manoel Cardoso¹, Fabiano Morelli¹

¹National Institute for Space Research (INPE), Avenida dos Astronautas 1758, 12227-010, Sao Jose dos Campos, SP, Brazil

²Met Office Hadley Centre, FitzRoy Road, Exeter, Devon, EX1 3PB, United Kingdom

10 ³UK Centre for Ecology and Hydrology, Maclean Building, Bensor Lane, Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, United Kingdom

Correspondence to: Renata Moura da Veiga (rmvrenata@gmail.com)

Abstract. Estimating fire emissions in the Brazilian Cerrado requires ~~a comprehensive approach that integrates fire parameters, mitigation strategies and policies, observational data, fire and vegetation modelling techniques, and policy.~~ Despite the Cerrado's significant ~~Although high quantities of global contribution to global~~ fire emissions ~~come from the Cerrado,~~ research in this area is still overlooked when compared to other savanna countries. Of 77 papers systematically reviewed papers, 54% address fire dynamics, management and policy. While these are key drivers to provide a holistic understanding of fire emissions, linking them to estimates is challenging, especially due to the difficulty in valuing the qualitative aspects of fire. ~~This study systematically reviewed 69 papers on fire emissions from the Cerrado. The aim his review was examines how fire dynamics, management and policy shape emissions and identify opportunities to better integrate these fields to better~~ integrate these fields for a more comprehensive understanding of fire emissions in the Cerrado. ~~to provide insights into the placement of the Cerrado in the global carbon budget and identify the research gaps that would support improved estimation of the biome's carbon balance. Our review finds that, in the Cerrado, studies often focus on quantifying fire dynamics parameters and emissions, and that a holistic approach is required to estimate fire emissions, which is hindered due to the difficulty in valuing the qualitative aspects of fire.~~ Most papers integrate different methodological techniques to achieve their ~~results.~~ Evidence suggests a rise in growing interest in ~~understanding~~ fire emissions in the Cerrado, reflected in the rising ~~increased~~ number of studies ~~overthroughtout~~ the years. More research is required to understand the aspects of fire dynamics in the Cerrado, how these reflect fire emissions ~~locally and globally~~ and potential mitigation activities that could help reduce the Cerrado's contribution to the global carbon budget. This could be achieved by including fire management representation in land surface models and using observational data to constrain and assess their utility.

1 Introduction

The Brazilian Cerrado biome (hereafter referred to as “the Cerrado”) covers approximately 2 million km² in Central Brazil. It is the second largest biome in Brazil, covering almost 24% of the country (IBGE, 2004). 48.66% (965,783 km²) of the Cerrado is covered by natural vegetation (MapBiomass, 2022). ~~We define natural areas as those covered by natural vegetation of the Cerrado, where anthropogenic land uses do not occur (e.g. pasture and agriculture). According to this criteria, 48.66% (965,783 km²) of the Cerrado is covered by natural vegetation (MapBiomass, 2022).~~ The Cerrado is one of the species-rich tropical savannas in the world (Klink and Machado, 2005; Schmidt and Eloy, 2020; Simon et al., 2009), and it is heavily influenced by fire (Fidelis, 2020; Pivello, 2011). Fire is the most critical factor in maintaining its health and biodiversity (Fidelis, 2020; Franke et al., 2018; Pivello, 2011), and the wide variety of ecosystems contained within it (Franke et al., 2018). Fire is also a major driver of carbon emissions, which influences the Cerrado’s overall contribution to the global carbon budget.

Fires in the Cerrado release substantial amounts of carbon into the atmosphere, primarily in the form of carbon dioxide (CO₂), but also as other greenhouse gases such as carbon monoxide (CO) and methane (CH₄) – CO₂ and CO combined account for 95% of the carbon emitted during biomass burning (Ward and Hardy, 1991). CO₂ and CO are both involved in atmospheric chemistry and in the greenhouse effect in different ways. Savanna burning dominates the emission of CO through incomplete combustion due to limited oxygen (Ehhalt et al., 2001). Similarly, CO₂ is released during complete combustion of biomass burning (Prentice et al., 2001). CO₂ is a major greenhouse gas, and it is crucial in absorbing and trapping infrared radiation in the atmosphere. However, the increased concentration of CO₂ in the atmosphere has warmed the Earth in alarming amounts (IPCC, 2022). Thus, understanding the emission of carbon, especially CO₂, during the combustion process is important to recognize the impact of these gases on fire emissions, especially in fire-prone settings like the Cerrado.

Beyond immediate emissions, fire participates in many complex interactions in the carbon cycle, influencing carbon balance over time ~~also influences carbon balance over time~~. For example, post-fire recovery critically shapes the Cerrado's long-term carbon balance (Burton et al. 2024; Gomes et al., 2020b; 2024; Hamilton et al., 2024). If vegetation fully regenerates to its pre-fire state, there is no net effect on atmospheric CO₂ levels over time. ~~However, even in this scenario, fires can influence other greenhouse gases and aerosols.~~ Alternatively, if fire activity decreases and vegetation accumulates, the landscape may shift to a net carbon sink. Conversely, if fires reduce long-term vegetation cover, the Cerrado could become a sustained carbon source, as observed globally (Burton et al., 2024; Jones et al. 2024). In fact, Gomes et al. (2024) indicate that fires in the Cerrado have acted as a source of carbon emissions to the atmosphere over the past decades.

During biomass burning, a large amount of carbon gases is released into the atmosphere. These emissions are mainly in the form of carbon dioxide (CO₂), carbon monoxide (CO), and methane (CH₄) – CO₂ and CO combined account for 95% of the carbon emitted during biomass burning (Ward and Hardy, 1991). CO₂ and CO are both involved in atmospheric chemistry and the greenhouse effect in different ways. CO is recognized as a major indirect greenhouse gas, meaning that it does not

absorb enough terrestrial infrared radiation to be considered a direct greenhouse gas, but it influences the concentration of other direct greenhouse gases, such as CH₄ and tropospheric ozone, through atmospheric chemistry (Ehhalt et al., 2001).

Savanna burning dominates the emission of CO through incomplete combustion due to limited oxygen (Ehhalt et al., 2001). Similarly, CO₂ is released during complete combustion of biomass burning (Prentice et al., 2001). CO₂ is a major greenhouse gas, and it is crucial in absorbing and trapping infrared radiation in the atmosphere, causing the greenhouse effect. However, the increased concentration of CO₂ in the atmosphere has intensified the greenhouse effect and warmed the Earth in alarming amounts. Thus, understanding the emission of CO and CO₂ during the combustion process is important to recognize the impact of these gases in fire emissions, especially in fire-prone settings like the Cerrado. Due to their importance, the studies captured by this review often report emissions in terms of carbon released by fire, including all the carbon components emitted during biomass burning, or in terms of CO₂ alone, due to its impact on the greenhouse effect.

Its emissions and influence on vegetation growth, cover and structure make fire a major driver of the Cerrado's carbon balance. The Cerrado's fires are potentially responsible for more than 30% (about 0.13 PgC year⁻¹) of Brazil's total fire emissions (da Silva Junior et al., 2020). As a comparison, the The Cerrado also accounts for about 14% of Brazil's emission from land use and land cover change (SEEG, 2023). Brazil is the highest emitter in the world in this category (Friedlingstein et al., 2023), contributing with up to 0.4 PgC year⁻¹ (Rosan et al., 2021). Thus, understanding the Cerrado's fire-driven carbon emissions is critical for accurately quantifying national greenhouse gas contributions, assessing Brazil's progress toward climate commitments, such as its Nationally Determined Contributions (NDCs), and evaluating its role in the global carbon budget. Accurate quantification of these emissions helps evaluate carbon sources and sinks, greenhouse gas contributions, and the impact of land management and climate policies at both national and global scales.

In the Cerrado, the extensive range of vegetation assemblages influences fire through variations in fuel type and structure across the region, as well as microclimate conditions (Flannigan et al., 2009; Gomes et al., 2020a). The Cerrado comprises a mosaic of vegetation formations, or physiognomies, that range from grasslands (such as *Campo Limpo* and *Campo Sujo*) to savannas (such as *Cerrado Ralo*, *Cerrado Típico*, and *Cerrado Denso*) to forest formations (such as *Cerradão*) (Ribeiro and Walter, 2008). The open formations (grasslands and savannas) are dominated by grasses and herbs, or fine fuel load, with few shrubs and trees (Ribeiro and Walter, 2008). Open formations have higher wind speed, higher air temperature, lower relative humidity, and lower fuel moisture than forests (Hoffmann et al., 2012).

The Cerrado presents two distinct seasons: the rainy season (November–March), characterized by biomass accumulation and higher fuel moisture, and the dry season (April–October), when the accumulated biomass available for burning becomes highly flammable, and fire can rapidly spread (Gomes et al., 2020b; Hoffmann et al., 2012; Silva et al., 2021). Intense wildfires in the Cerrado tend to occur in the late dry season (LDS; August–October) (Moura et al., 2019; Silva et al., 2021), when fire emissions are expected to be higher.

In the Cerrado, higher temperatures and reduced precipitation are now more common due to climate change (Hofman et al., 2021), which also change its fire regimes. Arruda et al. (2024) observed a shift in the peak of fire activity, with the highest concentration of fire events changing from September to October between 1985 and 2022. Fire events are also

becoming more frequent and intense (Gomes et al., 2024; Oliveira et al., 2022; Pivello et al., 2021). The IPCC AR6 WGI/WGII (IPCC, 2021, 2022) and the UNEP “Spreading like Wildfire” report (UNEP, 2022) warn that climate change increases drought conditions, which can aggravate heatwaves, increasing the risk of fire occurrence and the intensity and frequency of extreme events, such as wildfires. This happens because the combination of extreme weather events that occur simultaneously, or compound events, can amplify their effects (dos Santos et al., 2024). Drought-heatwaves episodes and extreme fire events intensified by climate change also impact hydrological processes, including precipitation and evaporation trends, groundwater recharge and soil infiltration capacity (Klink et al., 2020; Libonati et al., 2022). This is particularly important because the Cerrado region supports aquifers that supply major hydrographic basins in the whole country (Klink et al., 2020).

~~national implications for climate policies and international commitments (da Silva Junior et al., 2020; Pivello et al., 2021). The Cerrado's role in Brazil's overall emissions profile is, therefore, critical, with fires contributing to a substantial share of the country's fire emissions, which has national implications for climate policies and international commitments (da Silva Junior et al., 2020; Pivello et al., 2021).~~

The Cerrado's extensive range of vegetation assemblages, in turn, influences fire through variations in fuel type and structure across the region, as well as microclimate conditions (Flannigan et al., 2009; Gomes et al., 2020a). The Cerrado comprises a mosaic of vegetation formations, or physiognomies, that range from grasslands (such as *Campo Limpo* and *Campo Sujo*) to savannas (such as *Cerrado Ralo*, *Cerrado Típico*, and *Cerrado Denso*) to forest formations (such as *Cerradão*) (Ribeiro and Walter, 2008). The open formations (grasslands and savannas) are dominated by grasses and herbs, or fine fuel load, with few shrubs and trees (Ribeiro and Walter, 2008). Open formations have higher wind speed, higher air temperature, lower relative humidity, and lower fuel moisture than forests (Hoffmann et al., 2012). This combination of fuel and climate results in more intense and faster fires in grasslands and savannas than in forest areas in the Cerrado (Hoffmann et al., 2012).

The Cerrado presents two distinct seasons: the rainy season (November–March), characterized by biomass accumulation and higher fuel moisture, and the dry season (April–October), when the accumulated biomass available for burning becomes highly flammable, and fire can rapidly spread (Gomes et al., 2020b; Hoffmann et al., 2012; Silva et al., 2021). Although natural fires ignited by lightning occur in the wet season (Batista et al., 2018; Ramos Neto and Pivello, 2000), many fires result from anthropogenic activities during the dry season (Coutinho, 1990; Eloy et al., 2019; Pivello, 2011). Intense wildfires in the Cerrado tend to occur in the late dry season (LDS; August–October) (Moura et al., 2019; Silva et al., 2021), when fire emissions are expected to be higher. Therefore, fire occurrence in the Cerrado is also seasonal.

Fire participates in many complex interactions in the carbon cycle, from releasing carbon to benefiting ecosystems trajectories (Hamilton et al., 2024). Fire and climate regulate one another and can be in a positive feedback loop—climate and humans can influence fire patterns, and fire can influence climate by releasing carbon (Bowman et al., 2009). In the Cerrado, higher temperatures and reduced precipitation are now more common due to climate change, which also changes its fire regimes, with fire events becoming increasingly common (Gomes et al., 2020b; Hofmann et al., 2021).

The IPCC AR6 WGI/WGII (IPCC, 2021, 2022) and the UNEP “Spreading like Wildfire” report (UNEP, 2022) warn that climate change increases drought conditions, which can aggravate heatwaves, increasing the risk of fire occurrence and

the intensity and frequency of extreme events, such as wildfires. This happens because the combination of extreme weather events that occur simultaneously, or compound events, can amplify their effects (dos Santos et al., 2024). For example, the year 2020 was marked by compound drought-heatwave episodes, which favored fire activity and the increase in burned area in the Cerrado (Libonati et al., 2022; dos Santos et al., 2024).

135 Drought-heatwave episodes and extreme fire events intensified by climate change also impact hydrological processes, including precipitation and evaporation trends, groundwater recharge and soil infiltration capacity (Klink et al., 2020; Libonati et al., 2022). This is particularly important because the Cerrado region supports aquifers that supply major hydrographic basins in the whole country (Klink et al., 2020).

140 Beyond immediate emissions, fire also influences carbon balance over time. For example, post-fire recovery critically shapes the Cerrado's long-term carbon balance (Burton et al., 2024; Gomes et al., 2020b; Hamilton et al., 2024). If vegetation fully regenerates to its pre-fire state, there is no net effect on atmospheric CO₂ levels over time. However, even in this scenario, fires can influence other greenhouse gases and aerosols. Alternatively, if fire activity decreases and vegetation accumulates, the landscape may shift to a net carbon sink. Conversely, if fires reduce long-term vegetation cover, the Cerrado could become a sustained carbon source, as observed globally (Burton et al., 2024).

145 During biomass burning, a large amount of carbon gases is released into the atmosphere. These emissions are mainly in the form of carbon dioxide (CO₂), carbon monoxide (CO), and methane (CH₄). CO₂ and CO combined account for 95% of the carbon emitted during biomass burning (Ward and Hardy, 1991). CO₂ and CO are both involved in atmospheric chemistry and the greenhouse effect in different ways. CO is recognized as a major indirect greenhouse gas, meaning that it does not absorb enough terrestrial infrared radiation to be considered a direct greenhouse gas, but it influences the concentration of other direct greenhouse gases, such as CH₄ and tropospheric ozone, through atmospheric chemistry (Ehhalt et al., 2001).

150 Savanna burning dominates the emission of CO through incomplete combustion due to limited oxygen (Ehhalt et al., 2001). Similarly, CO₂ is released during complete combustion of biomass burning (Prentice et al., 2001). CO₂ is a major greenhouse gas, and it is crucial in absorbing and trapping infrared radiation in the atmosphere, causing the greenhouse effect. However, the increased concentration of CO₂ in the atmosphere has intensified the greenhouse effect and warmed the Earth in alarming amounts. Thus, understanding the emission of CO and CO₂ during the combustion process is important to recognize the impact of these gases in fire emissions, especially in fire-prone settings like the Cerrado. Due to their importance, the studies captured by this review often report emissions in terms of carbon released by fire, including all the carbon components emitted during biomass burning, or in terms of CO₂ alone, due to its impact on the greenhouse effect.

160 From 1997–2016, savanna fires from Southern Hemisphere South America, which the Cerrado dominates, accounted for 6.36% of the global carbon from fires annually (Van Der Werf et al., 2017). This contribution is substantial, as it highlights the Cerrado's role as one of the world's major fire-emitting ecosystems. To put this into perspective, savanna fires from the Australia and New Zealand region, which refer to the Australian savanna, account for 4.55% of the global carbon from fire emissions emitted each year for the same period (Van Der Werf et al., 2017). Compared with the Cerrado, a relatively high

number of fire studies were performed in Australia. Da Veiga and Nikolakis (2022) counted 64 papers from Australia and 29 from Brazil when documenting the interaction between fire management and carbon programs worldwide.

Fire studies in Australia have important benefits for understanding fire impacts in that region, and acts as an example to other savanna environments. Those include performing estimates of the release of GHG into the atmosphere and designing methods that can help reduce these emissions and providing insights into the interaction between actors in integrated fire management (IFM) practices (da Veiga and Nikolakis, 2022; Moura et al., 2019; Russell Smith et al., 2009). The cultural, socio-economic and ecological aspects of fire are crucial to execute and evaluate IFM activities (Myers, 2006). IFM integrates traditional knowledge and its connection with fire, and Australia is a leader in documenting these (da Veiga and Nikolakis, 2022). Measuring the social and cultural dimensions of fire presents significant challenges, and often is excluded from fire emission estimates in the Cerrado.

Natural activities to avoid the release of GHG or to increase carbon storage by letting vegetation recover, termed natural climate solutions (NCS), have emerged as a possibility to reduce emissions (Griscom et al., 2020). Tropical NCS, like improved fire management, could mitigate 6.56 petagrams of CO₂ annually (Pg CO₂ year⁻¹) between 2030 and 2050 (Griscom et al., 2020). Management activities, which include fire management, hold 26% of NCS' mitigation potential in Asia, Africa and Latin America. Lipsett-Moore et al. (2018) found that prescribed savanna burning, a typical fire management activity, could reduce 75% of emissions from LDS fires in South America. However, it is worth noting that some uncertainties may affect the estimates of fire management potential (Griscom et al., 2020). For example, limitations associated with spatial resolution and with expanding the details of local results to a broader scale.

In this context, understanding fire dynamic parameters provides grounding for assessing fire emissions in the Cerrado, and the interaction between these is essential for uncovering the factors that influence the Cerrado's role in the global carbon cycle and the broader implications for national and international policy. While much of the fire dynamic parameters literature emphasizes the importance of these parameters to estimate fire emissions (e.g. Arruda et al., 2024; Batista et al., 2018; Pereira et al., 2021), few studies aim to directly link these two components.

Linking fire dynamics to estimated emissions also guides mitigation by identifying aspects for potential intervention. For example, natural activities to avoid the release of GHG or to increase carbon storage by letting vegetation recover, termed natural climate solutions (NCS), have emerged as a possibility to reduce emissions (Griscom et al., 2020). Tropical NCS, like improved fire management, could mitigate 6.56 petagrams of CO₂ annually (Pg CO₂ year⁻¹) between 2030 and 2050 (Griscom et al., 2020). Management activities, which include fire management, hold 26% of NCS' mitigation potential in Asia, Africa and Latin America. Lipsett-Moore et al. (2018) found that prescribed savanna burning, a typical fire management activity, could reduce 75% of emissions from LDS fires in South America. However, it is worth noting that some uncertainties may affect the estimates of fire management potential (Griscom et al., 2020), such as limitations associated with spatial resolution and with expanding the details of local results to a broader scale.

Thus, it is essential to account for fire emissions and recognize mitigation mechanisms worldwide. This is particularly important/notably relevant for the Cerrado due to its intrinsic connection to fire. This review examines how fire

carbon emissions in the Cerrado are estimated, particularly what are the fire dynamic parameters used to estimate emissions, what fire carbon emissions estimates have been published, and their implications on fire management and policy. Due to the importance of carbon in the global carbon budget, this systematic review focuses on emissions in terms of carbon released by fire, including all the carbon components emitted during biomass burning, or in terms of CO₂ alone, due to its impact on the greenhouse effect.

Thus, ~~this systematic literature review synthesizes published material on fire emissions in areas of the Cerrado aims that do not explicitly include anthropogenic land uses; to gain a comprehensive understanding of the Cerrado's fire emissions within the global carbon budget by evaluating how fire parameters can inform emission estimates and mitigation strategies. Particularly, we with~~ aims to: (a) outline current emissions estimates, specially ~~fiably~~ carbon-CO₂, in regions that encompass the Cerrado or are limited to it; (b) ~~describe or~~ fire dynamic factors that ~~help~~ support these estimates; ~~in regions that encompass Cerrado or are limited to it; (c)~~ understand how these estimates fit the carbon budget; ~~(e) understand how measurements are done, how to use different, none standardized measurement, and what observational products are available;~~ (d) identify ~~potential~~ mitigation ~~opportunities~~ strategies in the biome and their link to fire-driven carbon fluxes; and (e) identify research gaps. This will support improvements for future fire emission estimation, ~~and~~ provide insights into the placement of the Cerrado in the global carbon balance and assist fire policies.

2 Methods

According to Moher et al. (2009, 1), a systematic review is “a review of a clearly formulated question that uses systematic and explicit methods to identify, select, and critically appraise relevant research, and to collect and analyze data from the studies that are included in the review”. It is based on rigorous criteria and a well-established and reproducible methodological approach to evaluate and synthesize the state of understanding of a specific topic (Cronin et al., 2008; Foo et al., 2021; Moher et al., 2015). We conducted this systematic review according to the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) Statement (Moher et al., 2009), and the practical guide on conducting systematic searches by Foo et al. (2021). PRISMA was designed to guide authors in transparently reporting on the review process (Moher et al., 2009).

We followed the four main steps outlined by Foo et al. (2021) to conduct this systematic review: (1) decide on a review question; (2) execute search; (3) initial literature screening; and (4) full-text screening. We also combined the literature review steps outlined by Cronin et al. (2008), which includes selecting a review topic, searching the literature, gathering, reading, analyzing the literature, and writing the review. After establishing our research question as “How compiling published material on fire emissions in natural areas of the Cerrado ~~that do not explicitly include anthropogenic land uses~~ can provide a holistic ~~better~~ understanding of ~~the placement of these emissionstheir role~~ in the global carbon budget?”, we combined keywords in English with the Boolean operators AND and OR in the Google Scholar database: fire AND emissions AND (Cerrado OR “Brazilian Savanna” OR (Savanna AND Brazil)). Keyword searches revealed a total of 3,7617 records.

230 We applied four inclusionary criteria to identify relevant literature: papers had to be (1) published in peer-reviewed journals with an impact factor greater than 1; (2) encompass the Cerrado biome; (3) be published after 2003, ~~for a two-decade period (2003–2022);~~ and (4) be conducted in areas that do not explicitly include anthropogenic land uses, ~~here referred to as “natural areas”.~~ We define natural areas as those covered by natural vegetation of the Cerrado, where anthropogenic land uses do not occur (e.g. pasture and agriculture). According to this criteria, 48.66% (965,783 km²) of the Cerrado is covered by ~~natural vegetation (MapBiomass, 2022).~~ While we acknowledge the role of anthropogenic fires and the importance of further research to integrate fires in natural and anthropogenic areas to provide a complete analysis of fire emissions in the Cerrado, we focus on non-anthropogenic areas, or natural areas, to provide a clearer ecological perspective on fire emissions in the Cerrado and their implications for the global carbon budget. Anthropogenic burning, as agricultural and pastoral, are better documented and accounted for, while natural fire regimes in the Cerrado remain understudied, particularly in terms of their

235 contribution to the global carbon budget. Thus, identifying the key drivers of fire emissions in natural landscapes provides a strong basis for improving emissions estimates, understanding fire-climate feedbacks, and assessing long-term ecosystem resilience in the Cerrado. Because papers found by this review often do not specify the land use of their study area, we have not included papers that explicitly document fire occurring in anthropized areas, ~~as a proxy for documenting existing literature on natural areas of the Cerrado.~~

240

245 To improve the assessment of our research question, we have also incorporated in our review papers that don't focus only on the Cerrado, but rather include it as part of the analysis. Although we explicitly limited the region to the Cerrado in our keyword search, we often encountered papers that include the Cerrado, but are not limited to it. Including these papers in our review allows for a complete analysis of fire emissions in the Cerrado, from global to local scales.

~~Despite its constraints, the impact factor was included as an indicator of scientific quality (Andersen et al., 2006; da~~

250 ~~Veiga and Nikolakis, 2022; Ketcham and Crawford, 2007).~~ The search starts in 2003 because fire emissions estimation is highly dependent on satellite data, much of it from MODIS (Moderate-Resolution Imaging Spectroradiometer) satellite burn products, and there is greater certainty in these data from 2003. ~~(see Andela et al., 2019; Silva et al., 2019).~~ The influence of MODIS data on our understanding of global fires can be seen in fire-vegetation modelling. There is broader disagreement between models on simulated burned area before the MODIS era, and much higher agreement during the satellite period


255 (Kloster and Lasslop, 2017; Hantson et al., 2020; Rabin et al., 2017). ~~We then conducted the search for a two-decade period, covering research from 2003 to 2022.~~

~~Empirical studies provide new information based on observation or experiments. Review and perspective papers analyze previously published studies by evaluating existing literature or expressing opinions on a specific topic. Although review and perspective papers do not focus on bringing original research, they supply the current knowledge of a specific topic and highlight pertinent published literature (Cronin et al., 2008). These were included in this systematic literature review to contribute to a more complete analysis of fire emissions in the Cerrado.~~

260

The criteria led to the initial screening of 10990 papers. Although we used keywords to conduct our review, the searches still returned papers not in English, or that did not mention fire emissions. 3224 papers were excluded due to being

265 duplicates, not in English, not mentioning fire emissions, or explicitly analyzing fire in anthropized lands. We full-text screened
the remaining ~~77~~⁶⁹ papers to confirm they met all the eligibility criteria. Figure 1 demonstrates the systematic literature review
process through the PRISMA diagram.






Figure 1: Adapted PRISMA flow diagram demonstrating the systematic literature review process divided into three steps: identification of potential papers through searched terms in the Google Scholar database, and exclusion of papers based on the four criteria established for this research; screening of the papers selected and exclusion of papers with the reported reasons; and inclusion of papers in this literature review.

Once we had selected the papers, we adopted a systematic scheme to review, analyze, and synthesize them. We retrieved the following information from the 7769 papers: title, year of publication, authors, journal, study design, area of study, what is measured, and how it is measured, and institutions of authors. ~~The 69 papers selected for this literature review are available in the Supplement.~~ Review and perspective papers were included in this systematic literature review to contribute to a more complete analysis of fire emissions in the Cerrado. Review and perspective papers analyze previously published studies by evaluating existing literature or expressing opinions on a specific topic, while empirical studies provide new

information based on observation or experiments. Although they do not focus on bringing original research, they supply the current knowledge of a specific topic and highlight pertinent published literature (Cronin et al., 2008).


The 77 papers selected for this literature review are available in the Supplement. After selecting the papers, we also wrote a summary of each study paper to simplify the understanding of its content and build a solid basis for writing the review (Cronin et al., 2008). From this, we framed the review by dividing the literature into themes. This approach has proven to be an efficient way to conduct and discuss the results from a systematic literature review in da Veiga and Nikolakis (2022).

We classified the reviewed papers based on (a) location range, from global to local scales: global, tropical region, South America, Brazil and Cerrado; (b) topic of search, where we identified three topics in the literature: fire dynamics parameters, fire emission estimates, and fire management and policy; and (c) study design, divided into empirical, review, and perspective. These classifications allowed a deep understanding of the general scope of current published literature on fire emissions in the Cerrado, including the purpose of the studies. This, in turn, enabled the identification of key trends for future research, which are outlined in this review. We divide our results into four sections, first summarizing the overall trends in current publications, and then highlighting the key findings from each of the three literature topics identified through our review process.

3 Results

3.1 Systematic literature review process

We reviewed 7769 papers and incorporated them into the literature review process. From 2003–2022, the number of papers published has increased since 2003 (Fig. 2), indicating a rise in interest in understanding fire emissions in the Cerrado and a rise in papers on fire sciences being published.



360




Figure 2: Number of papers published per year, since 2003, included in the literature review, with trendline (p-value ≤ 0.005); Number of papers published per year from the 69 papers included in this literature review, from 2003 to 2022.

There is an statistically significant increasing trend increasing tendency ($p \leq 0.005$) in the number of papers published throughout the time series, with a sharp drop in publications in the year 2022. but the year 2022 did not follow the growth trend shown in Fig. 2. This sharp drop in publications could indicate a gap in publications this year or a limitation of our research method that could not capture publications in 2022. It could also indicate a shifted focus away from the Cerrado studies due to political or financial constraints to encourage scientific studies in the region, or due to a shifted focus towards other regions of Brazil. For example, papers about fire dynamics and emissions in the Pantanal and in the Amazon rainforest were published in 2022 (see Barbosa et al., 2022; Dutra et al., 2022; Menezes et al., 2022; Silva et al., 2022; Walker et al., 2022). Papers published in 2022 related to fire dynamics and emissions in the Pantanal and in the Amazon show fire as a consequence of the compound impact of land use and climate in these regions (Barbosa et al., 2022; Silva et al., 2022; Walker et al., 2022).

Although we explicitly limited the region to the Cerrado in our keyword search, we often encountered papers that include the Cerrado, but are not limited to it. Including these papers in our review allows for a complete analysis of emissions in the Cerrado, from global to local scales. 2826 papers focused on the Cerrado, while 252 provide a global scale analysis, 96 relate to Brazil, 9 to South America, and 6 to the Tropical region. These numbers suggest that most papers are either restricted to the Cerrado, or provide a broad global analysis, with fewer papers in between, as shown in Fig. 3.

Papers not restricted to the Cerrado provide a broader perspective of emissions, encompassing the biome instead of being limited to it. Often papers referred to one limited region were expanded to represent a broader area. For example, within the South American Biomass Burning Analysis (SAMBBA) Project in 2012, Hodgson et al. (2018) used airborne flights in Tocantins, a Brazilian state dominated by the Cerrado, to represent fire emissions from smoke sampling for the whole Cerrado

395 biome. Conversely, Mistry et al. (2019) use the example of Brazil and Venezuela to illustrate how fire management can support emission reduction in South America.

400 As per the study design, we identified ~~6051~~ empirical papers (784%), ~~145~~ review papers (1822%), and 3 perspective papers (4%). ~~Empirical studies provide new information based on observation or experiments. Review and perspective papers analyze previously published studies by evaluating existing literature or expressing opinions on a specific topic. Although review and perspective papers do not focus on bringing original research, they supply the current knowledge of a specific topic and highlight pertinent published literature (Cronin et al., 2008). These were included in this systematic literature review to contribute to a more complete analysis of fire emissions in the Cerrado.~~ We further connect the study design with the coverage of the study area in Fig. 3.





Figure 3: Number of papers per study design and per coverage of study area in both percentage (chart) and

~~absolute~~ **absolute** numbers (data table).

We also observed that international (non-Brazilian) institutions drive most of the research captured by this literature review. We gathered the institution from the first author of each paper, of which 473 are international (612.3%) and 3026 are Brazilian (397.7%). From the Brazilian-led papers, 130 (43.338.5%) are from institutions located within the Cerrado biome area. From the international-led studies, 164 papers (342.6%) from the international-led studies involve authors from Brazilian institutions (Fig. 4), while 14half of the Brazilian-led studies (13 papers, 4750%) include authors from international institutions..




Figure 4: Institutions of the first authors from the papers reviewed. The chart on the left indicates that 473 papers involve first authors from international (non-Brazilian) institutions (Oranges), while 3026 come from Brazilian institutions (Greens). From the

~~international-led papers, 16 involve authors from Brazilian institutions, while 31 do not. From the Brazilian-led papers, 13 are from institutions located within the Cerrado biome, while 17 are not, of which 10 are within the Cerrado region. The chart on the right indicates that, from the international-led papers, 14 involve authors from Brazilian institutions, while 29 do not.~~

By reading the papers and through full-text screening of the content, we identified three topics of search from this systematic literature review process: (1) fire dynamics parameters, where the study aims to measure and evaluate parameters that can further be used to estimate emissions; (2) emission estimates, which include papers that focus on the quantification of fire emissions; (3) fire management and policy, where papers discuss the importance of fire management and fire policy to influence emission rates.

Of the ~~7769~~ papers reviewed, ~~4637~~ relate to fire dynamics parameters used to estimate emissions, ~~430~~ report the amounts of fire emissions, and ~~127~~ report fire management and policy — the total does not round up to ~~7769~~ because ~~2415~~ papers are related to more than one topic. These numbers indicate that many papers are not related to reporting emissions but provide information to support the understanding and estimation of fire emissions, ~~demonstrating a potential to expand the study of GHG emissions from fire in the Cerrado. For example, Santos et al. (2021) use satellite imagery to estimate emissions and parameters, such as burned area and fire intensity, to support the application of prescribed burning in the Cerrado, but actual emissions estimates are not included.~~

~~Of the 43 papers that report fire emissions, 23 focus only on fire emissions, while 18 analyze fire emissions and fire dynamics parameters, 1 focuses on fire emissions and fire management, and 1 on the three topics. From the 23 papers exclusive to fire emissions, only 2 are restricted to the Cerrado, one focusing on net CO₂ (Gomes et al., 2024), and the other on fine particulate matter (Mataveli et al., 2019). The remaining papers include the Cerrado region, but are not limited to it (9 provide a global analysis, 5 relate to the Tropical region, 4 to South America and 3 to Brazil). These numbers demonstrate the potential to expand the study of emissions from fire in the Cerrado.~~

~~Most papers provide new information on the parameters used to estimate fire emissions and on emissions themselves rather than analyzing existing literature. These often handle modelling techniques, satellite observations, in situ observations and even results found in previous studies, reflecting the importance of literature reviews in supporting the development of new data. These tools are often combined to provide a more complete analysis. Modelling and satellite observation are frequently integrated, with satellite observations being data providers to models, or models being evaluated by satellite observations, or even through data assimilation between satellite data and modelling simulations. From our results, 22 empirical papers integrate modelling and satellite data, 4 integrate modelling and literature review, 4 integrates satellite and in situ observations, and 1 integrates modelling, satellite data, literature review and in situ observations. Figure 5 shows the methodological techniques used across study areas in empirical papers.~~




Figure 5: Treemap of the methodological techniques used across study areas in empirical papers. The numbers present the number of studies of each methodological technique within each study area. The study areas Global, Tropical region, South America and Brazil are regions that include results for the Cerrado. 26 papers combine different techniques and are double-counted.

To our knowledge, and according to our search criteria, this is the first systematic literature review to provide an overview of fire emissions in the Cerrado. -Of the 3026 papers specific to the Cerrado, 4 were classified as review papers, and these are mostly related to the parameters associated with fire emissions, fire behavior, and fire ecology (see Arruda et al., 2018; Bustamante et al., 2012; Gomes et al., 2018; Gomes et al., 2020a). We further synthesize the main findings of the three topics from the systematic literature review process.

3.2 Fire dynamics parameters to estimate fire emissions

Papers under ‘fire dynamic parameters’ encompass 464% of the studies reviewed, underscoring the importance of variables like burned area, fuel characteristics, combustion completeness, combustion efficiency and emission factor in fire emissions research. These parameters directly influence emission estimates, with their combination playing key roles in determining carbon emissions from fires. By examining these variables within the specific ecological and climatic context of the Cerrado, we gain insights into how fire behavior and emissions in this biome interact.

~~The area burned, typically measured via satellite or ground surveys, is one of the primary parameters for estimating emissions (Libonati et al., 2015; Mangeon et al., 2016; Silva et al. 2021). Coupled with the available biomass for burning and~~

its characteristics — which depend on vegetation type, density, moisture and seasonal growth patterns — these elements set the stage for potential emissions. Fire intensity, driven by conditions such as dry weather, strong winds, and fuel accumulation, influences combustion efficiency. High intensity fires tend to consume more fuel, resulting in higher combustion efficiency and more complete combustion. This reduces emissions of pollutants such as carbon monoxide and particulate matter but increases emissions of carbon dioxide. In contrast, incomplete combustion results in higher emissions of pollutants such as particulate matter and carbon monoxide and produces pyrogenic carbon, which may persist in soils over long periods. Combustion completeness further influences the amount of biomass converted to carbon and released into the atmosphere. Together, these parameters allow for the estimation of emissions based on the combination of burned area, fuel load, and combustion completeness.

The prevalence of studies on these fire dynamics parameters reflects ~~both~~ the accessibility of these variables and indicates the importance to link a gap in linking fire dynamics directly to emission, with studies often highlighting the potential applicability of their research in fire emission estimation (e.g. Libonati et al., 2015; Pereira Junior et al., 2014). This focus on fire dynamics provides some of the most current information available, yet it suggests a need for more research to correlate fill the gaps in understanding the chain from fire drivers to emissions. We further discuss the fire dynamics parameters found in the literature review process.

3.2.1 Burned area and fuel characteristics

Burned area detection in the Cerrado has mainly been measured via the satellites Aqua and Terra/MODIS, and Landsat/TM (Libonati et al., 2015; ~~Mataveli et al., 2019~~; Oliveira et al., 2021; Pereira et al., 2021), and is one of the primary parameters for estimating emissions. The Global Fire Emissions Database version 5 (GFED5) primarily relies on MODIS burned area products to provide global burned area (Chen et al., 2023). GFED5 estimates 21.35 Mha year⁻¹ (2001–2020) for burned area in open savannas of the Southern Hemisphere of South America (SHSA), where the Cerrado is located. GFED5 burned area results are more aligned with data from higher-resolution satellite sensors (Chen et al., 2023), indicating enhancement compared to previous versions, and a potential to improve fire emissions estimates derived from GFED5 (Chen et al., 2023).

Libonati et al. (2015) developed a regional algorithm using MODIS data to increase the accuracy of estimations of burned area in the Cerrado, demonstrating that regional products more accurately capture vegetation diversity. Also, to capture the variety of fire dynamics throughout the biome, Silva et al. (2021) map fire characteristics for the 19 ecoregions of the Cerrado, including the patterns and trends of burned area using MODIS data. Results show a great variation of size of burned area in the Cerrado, with large areas detected in the boundaries with other biomes (Silva et al., 2021). Using Aqua and Terra/MODIS data and the algorithm developed by Libonati et al. (2015), the National Institute for Space Research (INPE, in Portuguese) in Brazil has estimated that 16.2% of the Cerrado was burned in 2007 (the equivalent of 329,138km²), a critical year in terms of wildfires in Brazil, and 3.6% in 2009 (74,085km²), a year with low overall rates of burned area in the country.

To increase the accuracy of estimations of burned area in the Cerrado, Libonati et al. (2015) developed a regional algorithm to capture the local characteristics of the biome, considering the complexity of plant diversity and climate.

Also, to capture the diversity of fire dynamics throughout the biome, Silva et al. (2021) propose dividing the Cerrado into 19 ecoregions based on the patterns and trends of fire intensity, seasonality, frequency, extent and sear size. Results show a great variation of fire intensity and size of burned area in the Cerrado, with large areas detected in the boundaries with other biomes (Silva et al., 2021). Using Aqua and Terra/MODIS data and the algorithm developed by Libonati et al. (2015), the National Institute for Space Research (INPE, in Portuguese) in Brazil has estimated that 6.8% of the Cerrado was burned in 2020 (the equivalent of 139,644km²), a critical year in terms of wildfires in Brazil, and 3.6% in 2009 (74,353km²), a year with low overall rates of burned area in the country.

The year 2020 was a significant year in terms of burned area in the Cerrado due to a combination of factors (Pivello et al., 2021). 2020 was a drought year in the biome, intensified by prolonged dry season and heatwave (Hofmann et al., 2021; Libonati et al., 2022; dos Santos et al., 2024). This compound drought heatwave episode aggravated fire activity in the Cerrado (Libonati et al., 2022; dos Santos et al., 2024). Although no estimates were found correlating the compound event of 2020 with fire emissions, it is expected that the drought heatwave episode led to increased fire emissions due to the increased fire activity and burned area that occurred in that year.

The extent of area burned is highly connected to the fuel characteristics, including fuel type (vegetation components), moisture (amount of water content), and load, and these are essential to determine the intensity and occurrence of fire, together with climatic conditions (Alvarado et al., 2017; Gomes et al., 2020b; Haas et al., 2022). Fuel type also influences the monthly peak in fire of each vegetation type, due to climatic conditions. Grassy material tends to burn first because of its high flammability, with the fire peak from grassland happening in August (Arruda et al., 2024). In comparison, savannas tend to encounter more fires in September and October (Arruda et al., 2024). These impact the extent of area burned in each vegetation type. From 1985 to 2022, 85% of burned area in the Cerrado affected native vegetation, with savannas accounting for 54%, and grasslands, 11% (Arruda et al., 2024).

Fuel load in the Cerrado is highly connected to the seasonal variation in precipitation (Oliveira et al., 2021). For example, drought was the strongest predictor of burned areas made by Alvarado et al. (2017). Rainfall in the wet season allows fuel to build up, determining the fuel load available for burning. In contrast, rainfall in the dry season determines the moisture content of the accumulated fuel and, thus, the probability of burning (Alvarado et al., 2017). Fuel load increases with vegetation density (Oliveira et al., 2021), but grasslands have more biomass available for burning than forests due to the fine fuel load in each physiognomy. Approximately 95% of the biomass of the herbs and grass layers is available for burning in the Cerrado. In comparison, only 0.01% of the shrubs and tree layers are available due to the high quantity of woody material and low quantity of grassy material (Gomes et al., 2020b).

The proportion of influence between climate and fuel on the resultant burned area and emissions varies and is still being discussed. In a global context, burned area is likely to increase worldwide due to climate change (Burton et al., 2024). In South America an increase in temperature and decrease in moisture availability is expected to drive an increase in burned

area, including in the Cerrado (Burton et al., 2021). Similarly, Silva et al. (2019) predicts increased burned area in the Cerrado under all IPCC Representative Concentration Pathways (RPC) scenarios. The RCP 2.6, the scenario most closely aligned with the United Nation's 1.5 °C target, could lead to an increase in burned area up to 22% by 2050 in the Cerrado.

studies suggest that burned area is more sensitive to climate than to fuel availability (Bistinas et al., 2014; Haas et al., 2022; Kelley et al., 2019). From the global analysis of Kelley et al. (2019), we can infer that this trend is also observed in the Cerrado, where moisture is a greater limiting factor to burned area predictions.

3.2.2 Combustion efficiency, combustion completeness and emission factor

Combustion efficiency, combustion completeness and emission factor have been identified in the literature as Another important parameters identified in the literature to estimate fire emissions. Combustion completeness, which refers to the amount of biomass converted to gas, aerosols and particulates during the combustion process and released into the atmosphere (Carvalho Jr. et al., 1998). Similarly, combustion efficiency identifies “the percentage of carbon released during combustion of biomass fuels in the chemical form of carbon dioxide” (Ward and Hardy, 1991, 117–118). Combustion efficiency is often measured by the amount of CO₂ emitted divided by the amount of CO₂ and CO emissions combined, termed modified combustion efficiency (see Andreae, 2019; Hodgson et al., 2018; Vernooij et al., 2021). MCE tends to be higher in open savannas (Vernooij et al., 2023). Values above 0.9 tend to characterize fires in a flaming stage, and these are predominant in the Cerrado due to the dry fine fuel that is likely to rapidly burn (Hodgson et al., 2018).

Using airborne sensors, modified combustion efficiency has been reported to be 0.94±0.02 in a flight above Tocantins in 2012 (Hodgson et al., 2018). Vernooij et al. (2021) used a UAV-based approach (uncrewed aerial vehicle) to sample smoke from grassland and savanna formations in 2017–2018, in a Protected Area also in Tocantins (*Estação Ecológica Serra Geral do Tocantins, EESGT*), to evaluate the seasonal burning differences. The authors concluded that late dry season fires (LDS; after July 1; more intense fires) have slightly higher modified combustion efficiency when compared to early dry season fires (EDS; before July 1; more mild fires): 0.963 and 0.957, respectively. In further studies, Vernooij et al. (2023) discuss that the MCE found in Vernooij et al. (2021) from measurements in EESGT may have underestimated the MCE in other parts of the Cerrado when analysing MCE in a biome-scale. These values are considered high and are consistent with other savannas in the world. Global MCE for savannas and grasslands averages 0.94 (Andreae, 2019), while MCE in the African and in the Australian savannas have been reported as 0.938±0.019 and 0.86–0.99, respectively (Hodgson et al., 2018).

Gomes et al. (2020b) developed a model to simulate the relationship between fire frequency, plant biomass, and fire emissions in the Cerrado based on microclimate variables (wind speed, average air temperature, relative air humidity, and vapor pressure deficit) and aboveground biomass of grasses, herbs, shrubs, and trees. Burning of biomass has proven to be higher in open formations due to fuel characteristics; when analyzing the percentage of biomass burned about the total biomass, or the combustion factor, the BEFIRE model (Behavior and Effect of Fire) has estimated a combustion factor of 87% for grasslands, 86% for savannas, and 39% for forests in the Cerrado.

590 The emission factor (EF) is an essential component that contributes to estimating fire emission, and it refers to the ratio of a particular gas released divided by the fuel consumed, expressed in grams of that specific gas per kilogram of dry matter (Palacios-Orueta et al., 2005).- EF depends on the type and moisture content of the fuel consumed, amount of fuel burned, amount and concentration of the emitted gas, meteorological conditions during the fire (Palacios-Orueta et al., 2005; Vernooij et al., 2023). Thus, EF is highly variable across studies and often the source of uncertainty in fire emission estimates.

595 In fact, Andreae (2019) argues that the emission factors of CO₂ and CO from forest fires are the main source of uncertainty in relation to the impact of vegetation fires on the global carbon cycle.

Andreae (2019) and Hodgson et al. (2018) argue that local values of EF can be used to improve estimates.-Similarly, Vernooij et al. (2023) proposes incorporating dynamic EF to improve representation of savanna’s temporal and spatial variabilities, better capturing its fire regime and resultant emissions. When comparing the early (EDS) and late dry season (LDS) in savannas worldwide, including the Cerrado, Vernooij et al. (2023) finds that fuel moisture content and relative humidity were lower in the LDS compared to the EDS, increasing fuel consumption and fire intensity over the dry season, and resulting in higher EFs in the LDS.-The values of EF found from recent studies are summarized in Table 1.

600

Table 1: Emission factors values from areas that include the Cerrado found by this review.

Study	Study area	Emission Factor	Value (g kg ⁻¹)	
Van Der Werf et al. (2017)	Global — savannas	EF _{CO₂}	1,686	
		EF _{CO}	63	
		EF _{CH₄}	1.94	
<u>Andreae (2019)</u>	<u>Global – savannas</u>	<u>EF_{CO₂}</u>	<u>1,660</u>	610
		<u>EF_{CO}</u>	<u>69</u>	
		<u>EF_{CH₄}</u>	<u>2.7</u>	
<u>Vernooij et al. (2023)</u>	<u>Global – savannas</u>	<u>EF_{CO₂}</u>	<u>1,685</u>	
		<u>EF_{CO}</u>	<u>64</u>	
		<u>EF_{CH₄}</u>	<u>1.85</u>	
Hodgson et al. (2018)	Cerrado	EF _{CO₂}	1,711 ± 175	615
		EF _{CO}	74 ± 8	
		EF _{CH₄}	2.23 ± 0.23	
Vernooij et al. (2021)	Cerrado	EF _{CO₂}	1,664	
		EF _{CO}	48	
		EF _{CH₄}	0.78	

620 3.2.3 Fire behavior and intensity

Fire behavior is limited by fuel characteristics, availability and microclimate conditions. Fire behavior is often analyzed in terms of fire intensity (Gomes et al., 2020a; Silva et al., 2021), fire spread (Gomes et al., 2020a), heat released (Gomes et al., 2020a), fuel consumption (Andela et al., 2016), and fire return interval (Gomes et al., 2020b; Pereira Junior et al., 2014). Fire intensity here is defined as the rate at which fire releases heat energy (DeBano et al., 1998). For the Cerrado,
625 this means that fire intensity follows a seasonal pattern, increasing in the dry season months (Silva et al., 2021), and that it also is highly influenced by the vegetation type, increasing from forests to savannas and grasslands, where fine fuel consumption is higher (Gomes et al., 2020a). Silva et al. (2021) indicates ~~a larger~~ higher values of fire intensity in the end of the dry season during the dry months in the Cerrado, for 2001–2019, with more than 90% of burned area happening between June and October, when fuel moisture is lower and fuel availability for burning is higher.

630 Fire intensity can also be measured through the fire radiative power (FRP). FRP is the instantaneous amount of energy released by fire in the combustion process (Wooster, 2002). FRP often derives from MODIS data, and it relates to the intensity of fire and to the amount of biomass being consumed (Wooster, 2002). Although FRP has been used to provide estimates of fire intensity, Sperling et al. (2020) states that FRP from MODIS is underestimated for the Cerrado. Through FRP, Silva et al. (2021) estimates fire intensity in the Cerrado, with high values (FRP > 63.7 MW) found in the border with other biomes.
635 Continuous observations of FRP can be integrated over time, resulting in the Fire Radiative Energy (FRE) (Van der Werf et al., 2017). FRE represents the total amount of energy released by fire during the combustion process, and it relates to the total amount of biomass consumed by fire, being directly related to fire emissions (Wooster, 2002).

FRP data was also combined by Andela et al. (2016) to derive fuel consumption estimates, which depends on combustion completeness and amount of fuel available for burning. For the savannas in South America, where the Cerrado is
640 included, high values of fuel consumption were estimated compared to other savannas (Andela et al., 2016). For Andela et al. (2016), fuel consumption is partly driven by fire return periods, with grasses favoring short fire return periods, and thus resulting in low fuel build-up rates and lower fuel consumption. Pereira Junior et al. (2014) modelled fire return interval for a Protected Area in the Cerrado from 1997–2008), finding fire return intervals of three to six years, depending on the vegetation type. Shorter fire return periods (biennial) have been reported for other areas of the Cerrado (see Batista et al, 2018; Gomes et
645 al., 2020b), typically related to the fine fuel load.

Fire spread in Cerrado is connected to the amount of fine fuel load available for burning and moisture (Gomes et al., 2020a; Silva et al., 2021). Fires in the Cerrado spread from savannas and grasslands to forests due to higher fine fuel load available for burning, and the same pattern is observed for heat released (Gomes et al., 2020a), which is also higher in savannas and grasslands than in forests. These reflect combustion completeness, combustion factor and carbon
650 the emission rates of the different physiognomies in the biome (Gomes et al., 2020a).

Together, these findings emphasize the critical role that specific fire dynamics parameters play in shaping fire emissions within the Cerrado. To estimate carbon emissions from fire, researchers utilize various methodologies and data

sources to quantify the carbon released during combustion. By understanding how burned area, fuel load, combustion completeness, combustion efficiency and emission factor vary across the landscape and seasons, researchers can better estimate fire emissions and their contribution to global greenhouse gas levels budget. These insights lay the foundation for targeted fire management strategies that consider both ecological and climatic factors unique to the Cerrado. In the next sections, we discuss how current research brings this information together to estimate fire emission in the Cerrado.

Satellite observations also play a pivotal role in measuring fire intensity, and this includes providing direct measurements of fire intensity by quantifying the radiative energy emitted during the carbon oxidation of combustion. This instantaneous measure is termed Fire Radiative Power (FRP) (Van Der Werf et al., 2017). FRP considers temperature, area affected by fire, and emissivity rates and often uses MODIS active fires data as inputs (Ichoku et al., 2008; Mataveli et al., 2019; Vermote et al., 2009). When multiple FRP observations are available, it is possible to estimate the quantity of energy radiated during the combustion process, termed Fire Radiative Energy (FRE) (Vermote et al., 2009). The parameters combined are used to estimate fire emissions worldwide, including the Cerrado. The variables identified in this review as predominant factors associated with fire emissions estimates in the Cerrado are summarized in Fig. 65.




Figure 65: Variables associated with estimating fire emissions in the Cerrado found in the literature. The Cerrados's physiognomies, separated into forests, savannas and grasslands, increase in fine fuel load and decrease in fuel moisture from forests to grasslands. Microclimatic conditions also change across the physiognomies, with increasing wind speed and air temperature, and decreasing relative humidity from forests to grasslands. The Cerrado's seasonality is divided into wet and dry seasons. The wet season is characterized by high precipitation, lightning ignitions and accumulated biomass, whereas the dry season is characterized by low precipitation, anthropogenic ignitions and flammable biomass. Fuel characteristics (square boxes), climatic conditions (circle boxes)

and ignition (hexagon boxes) interact (dashed lines) to determine the Cerrado's fire behavior. Two aspects of fire behavior are presented (numbers 1 and 2): 1) fire spreads from forests to grasslands; 2) fire intensity increases in the dry season. The Cerrado's physiognomies, seasonality and fire behavior together drive the size of burned area, resulting in fire emissions (solid lines). High-intensity fires typically consume more fuel, leading to higher combustion efficiency and more complete combustion. Combustion completeness then affects the proportion of biomass converted into carbon and released into the atmosphere, also resulting in fire emissions (solid lines). The image representing the Cerrado's physiognomies was adapted from the Brazilian Agricultural Research Corporation (Embrapa, 2024).

~~Together, these findings emphasize the critical role that specific fire dynamics parameters play in shaping fire emissions within the Cerrado. To estimate carbon emissions from fire, researchers utilize various methodologies and data sources to quantify the carbon released during combustion. By understanding how burned area, fuel load, combustion completeness, combustion efficiency and emission factor vary across the landscape and seasons, researchers can better estimate fire emissions and their contribution to global greenhouse gas levels. These insights lay the foundation for targeted fire management strategies that consider both ecological and climatic factors unique to the Cerrado. In the next sections, we discuss how current research brings this information together to estimate fire emission in the Cerrado.~~

3.3 Estimated fire **carbon** emissions in the Cerrado

Papers under 'emission estimates' account for 43% (43 papers) of the papers reviewed, with 23 papers focusing on estimating emissions alone, while the remaining papers include a combination of emissions estimates, fire dynamics parameters and/or fire management and policy. Given the importance of carbon gases released into the atmosphere during combustion, mainly in the form of carbon dioxide (CO₂) (Ward and Hardy, 1991), to the global carbon budget, this section will synthesize the main findings related to fire carbon emissions from the Cerrado found in the literature review process, notwithstanding the impact of non-carbon gases emitted during biomass burning.

The amount of GHG emitted to the atmosphere by fires is typically inferred by models implemented at different scales ranging from local to global analyses – of the 43 papers under 'emission estimates', 25 handle modelling techniques. Global and regional analyses tend to be less detailed, as they usually focus on capturing absolute emissions and on studying general aspects of large areas through a coarse resolution, and these are necessary to assess the impact of emissions on the global carbon balance (Palacios-Orueta et al., 2005; Rabin et al., 2018).

~~Two global emissions datasets were often used in the papers reviewed to develop and evaluate models of fire occurrence and effects, and these are GFED and Global Fire Assimilation System (GFAS). Both rely on MODIS products to estimate emissions. GFED quantifies fire emissions estimates are globally, and estimations are based on MODIS burned area products and on the Carnegie-Ames-Stanford Approach (CASA) model (Van Der Werf et al., 2017). GFAS estimates fire emissions globally by using a conversion factor that links MODIS-derived FRP observations to combustion rates, resulting in the fuel consumption, which is then combined with emissions factors to estimate fire emissions (Kaiser et al., 2012).~~

For example, Burton et al. (2021) uses the Interactive Fire and Emission Algorithm for Natural Environments (INFERNO) integrated into the Joint UK Land Environment Simulator (JULES) to evaluate current and future fire emissions in South America, and results are compared to both GFED and GFAS. Pereira et al. (2016) estimates fire emissions also in

South America using the Brazilian Biomass Burning Emission Model with FRP assimilation (3BEM FRP), and results are also compared to GFAS.

From 1997 to 2016, GFED fire emissions averaged 2.2 PgC year⁻¹ globally (Van Der Werf et al., 2017). In the SHSA region (Southern Hemisphere South America), where the Cerrado is located, fire emissions averaged 0.291 PgC year⁻¹, of which 49.3% were from savanna fires (0.14 PgC year⁻¹). Considering these values, savanna fires from SHSA, which broadly include the Cerrado, account for 6.36% of annual global total carbon emissions from fires. GFAS does not provide subregional analysis, but carbon emissions for South America averaged 0.349 PgC year⁻¹ (2003–2008), while GFED version 3.1 averaged 0.299 PgC year⁻¹ (Kaiser et al., 2012).

~~Version 4s of GFED also includes small burned area detection to improve its results, and small burned area detection in GFED4s relies on MODIS burned area product, on active fire from MODIS, and on surface reflectance observations (Van Der Werf et al., 2017). According to Van Der Werf et al. (2017), GFED values could be improved using FRP and FRE to reduce uncertainty from small burning area detection.~~

~~Pan et al. (2020) compare global datasets of biomass burning emissions, including the Global Fire Emission Dataset (GFED). GFED quantifies fire emissions globally, and estimations are based on MODIS burned area products and on the Carnegie Ames Stanford Approach (CASA) model (Van Der Werf et al., 2017). Version 4s of GFED also includes small burned area detection to improve its results, and small burned area detection in GFED4s relies on MODIS burned area product, on active fire from MODIS, and on surface reflectance observations (Van Der Werf et al., 2017). According to Van Der Werf et al. (2017), GFED values could be improved using FRP and FRE to reduce uncertainty from small burning area detection.~~

~~GFED was extensively used to develop and evaluate models of fire occurrence and effects (e.g. Hantson et al., 2020; Kelley et al., 2013; Lasslop et al., 2020; Rabin et al., 2018). In one application for South America, for example, the Interactive Fire and Emission Algorithm for Natural Environments (INFERNO) (Mangeon et al., 2016; Burton et al., 2019) was integrated into the Joint UK Land Environment Simulator (JULES) (Clark et al., 2011) and used to evaluate current and future fire emissions in that region (Burton et al., 2021). Other models developed for South America are based on FRP to estimate emissions, such as the Global Fire Assimilation System (GFAS) (Kaiser et al., 2012) and the 3BEM with FRP (3BEM FRP) (Pereira et al., 2016).~~

Conversely, local studies include the complexity and diversity of a limited region, which is essential to capture changes in fire dynamics and to assess the components that influence emissions at smaller scales (Palacios-Orueta et al., 2005). They might be extrapolated to represent more extensive areas. ~~For example, Santos et al. (2021) used satellite-derived products to analyze the influence of prescribed burning in the dynamics of fire in two Indigenous Territories in the Cerrado, from which attributes for the whole biome can be inferred.~~ Although there are uncertainties with both global and local scales, the use of remote-sensing techniques contributes to the accuracy of emission estimations, and it is the core of much recent research regarding fire occurrence and emissions worldwide (Lasslop et al., 2019).

This literature review identified 4 papers dedicated to estimating fire carbon emissions from the Cerrado fires, being one exclusively focused on estimating fire-related carbon emissions from the Cerrado (Gomes et al., 2024), while the other

studies either include the Cerrado, but are not restricted to it (da Silva Junior et al., 2020), include emissions as well as fire dynamics parameters (Gomes et al., 2020a; Oliveira et al., 2021). Yet, some studies that estimate fire emissions in the Cerrado are not focused on carbon emissions, but on aerosol (e.g. Hodgson et al., 2018; Pereira et al., 2009; 2022), fine particulate matter (PM_{2.5}) (e.g. Mataveli et al., 2019; 2023; Santos et al., 2021) and nitrogen compounds (e.g. Pope et al., 2020).

Gomes et al. (2024) uses satellite-derived burned area maps, surface fuel material maps, combustion factors, and emissions factors to estimate net CO₂ emissions from the Cerrado fires, among other greenhouse gases. From 1985 to 2020, the Cerrado savanna emitted approximately 2,227,964 Gg of CO₂ (0.018 PgC year⁻¹) and removed approximately 1,495,725 Gg of CO₂ (0.011 PgC year⁻¹) due to biomass regrowth, resulting in a net CO₂ emission of 859,701 Gg (0.007 PgC year⁻¹) (Gomes et al., 2024).

From GFED4s and MODIS data, da Silva Junior et al. (2020) estimated carbon emissions for the Brazilian biomes between 1999 and 2018. Brazilian biomes produced 8.09 PgC of fire emissions (equivalent to 0.40 PgC year⁻¹). By analyzing all the Brazilian biomes, da Silva Junior et al. (2020) put fire emissions in the Cerrado into a national perspective, where it contributes 32.04% of total fire emissions (about 0.13 PgC year⁻¹), similar to the values found by Van Der Werf et al. (2017) for the savanna fire emissions in the SHSA region. According to da Silva Junior et al. (2020), the Cerrado is a major contributor to Brazil's fire emissions.

Gomes et al. (2020a) modelled carbon emissions associated with fine fuel consumption, finding 0.230 kg m⁻² for grassland, 0.210 kg m⁻² for savanna, and 0.053 kg m⁻² for forests, and concluding that fine fuel load was the main predictor of the amount of carbon released through fire. When considering different scenarios (moderate, medium, and extreme) for fine fuel available for burning, wind speed, and vapor pressure deficit using the BEFIRE model, Gomes et al. (2020b) showed that carbon emissions from fine fuel consumption increased with the intensity of the scenario (0.19 kg m⁻² for moderate, 0.23 kg m⁻² for medium, and 0.26 kg m⁻² for extreme). Because the model only considers fine fuel, which is more abundant in grasslands due to the presence of grasses, followed by savannas and forests, carbon emissions decrease with the increase of woody biomass. These simulations confirm that fire-associated emissions depend on the vegetation type (Gomes et al., 2020a, 2020b).

Oliveira et al. (2021) modelled fire emissions across the Cerrado by estimating fuel loads through remote sensing data over four years (2015–2018). Results averaged 0.066 ± 0.013 Pg CO₂ year⁻¹ (0.018 ± 0.00354 PgC year⁻¹). When accounting for regrowth uptake, net emission was 0.015 ± 0.004 Pg CO₂ year⁻¹ (0.00487 ± 9.65 10⁻⁴ PgC year⁻¹). Oliveira et al. (2021) consider these values low and suggest incorporating a more detailed vegetation map and the burning intensity of different fuel types to improve their estimates. The values of fire carbon emissions restricted to the Cerrado reported in this section are summarized in Table 2.

From 1997 to 2016, GFED fire emissions averaged 2.2 PgC year⁻¹ globally (Van Der Werf et al., 2017). In the SHSA region (Southern Hemisphere South America), where the Cerrado is located, fire emissions averaged 0.291 PgC year⁻¹, of which 49.3% were from savanna fires (0.14 PgC year⁻¹). Considering these values, savanna fires from SHSA, which broadly include the Cerrado, account for 6.36% of annual global total carbon emissions from fires. As a comparison, the Australian savanna encompassed in the AUST region (Australia and New Zealand) is the region's largest source of fire emissions,

accounting for 86.3% of the 0.116 PgC year⁻¹ emitted in AUST. This means that AUST savanna fires contribute 4.55% of annual global carbon emissions from fires. The values reported in this section are summarized in Table 2. From GFED4s and MODIS data, da Silva Junior et al. (2020) estimated carbon emissions for the Brazilian biomes between 1999 and 2018. Brazilian biomes produced 8.09 PgC of fire emissions (equivalent to 0.40 PgC year⁻¹). By analyzing all the Brazilian biomes, da Silva Junior et al. (2020) put fire emissions in the Cerrado into a national perspective, where it contributes 32.04% of total fire emissions (about 0.13 PgC year⁻¹), similar to the values found by Van Der Werf et al. (2017) for the savanna fire emissions in the SHSA region. The Cerrado presented significant emissions in the dry season (da Silva Junior et al., 2020). According to da Silva Junior et al. (2020), the Cerrado is a major contributor to Brazil's fire emissions. Using the BEFIRE model, Gomes et al. (2020a) estimated carbon emissions from fine fuel consumption of 0.230 kg m⁻² for grassland, 0.210 kg m⁻² for savanna, and 0.053 kg m⁻² for forests. When considering different scenarios (moderate, medium, and extreme) for fine fuel available for burning, wind speed, and vapor pressure deficit, the study showed that carbon emissions increased with the intensity of the scenario (0.19 kg m⁻² for moderate, 0.23 kg m⁻² for medium, and 0.26 kg m⁻² for extreme), with herbs and grasses accounting for 80% of the total emissions, and trees and shrubs for the other 20% (Gomes et al., 2020b). Because the model only considers fine fuel, which is more abundant in grasslands, followed by savannas and forests (Ribeiro and Walter, 2008), carbon emissions decrease with the increase of woody biomass.

Table 2: Values found in the literature for fire carbon emissions in the Cerrado. Units are: petagrams of carbon per year (PgC year⁻¹) and kilogram per square meter (kg m⁻²).

Study	Study area	Value	Units	Observation
da Silva Junior et al. (2020)	Brazil Cerrado	0.4 0.13	Pg C year ⁻¹ Pg C year ⁻¹	Based on MODIS data and GFED4s.
Gomes, Miranda, Soares-Filho, et al. (2020a)	Cerrado – Grassland Cerrado – Savanna Cerrado – Forest	0.23 0.21 0.053	Kg m ⁻² Kg m ⁻² Kg m ⁻²	The study only considers fine fuel, and it is based on eight experimental burn studies across Cerrado.
Oliveira et al. (2021)	Cerrado	0.018 ± 0.00354	Pg C year ⁻¹	Includes results from a bayesian model developed by the authors based on remote sensing imagery, as in Franke et al. (2018).
Gomes et al. (2024)	Cerrado – Savanna	0.018	Pg C year ⁻¹	Focuses only on the savanna formation of the Cerrado.

840 The difference in values, and even in the units used to report results (Table 2), indicates the complexity of estimating fire emissions, especially in a diverse region such as the Cerrado. The estimation of fire emissions relies on multiple variables, each quantified by different methodologies and databases. Also, the procedure used to extrapolate measurements and estimate emissions on broader scales differs among studies. When combined, these variabilities result in uncertainties associated with estimating fire emissions, often reported by the studies synthesized in this review ([e.g. Andreae, 2019; Oliveira et al., 2021; Vernooij et al., 2023](#)). ~~For this reason, Houghton (2003) states that comparing results from contrasting approaches can be misleading if the processes behind them are poorly understood.~~

~~Reducing uncertainties is also critical to generating more accurate estimates of fire emissions. Within the REgional Carbon Cycle Assessment and Processes (RECCAP) scope, the Global Carbon Project explains that contrasting results can help evaluate the models' strengths, weaknesses, and uncertainties. This could lead to a more holistic understanding of the processes involved in fire emissions, further improving estimates and providing better grounding for future projections.~~

~~The papers reviewed have shown that input data uncertainty affects output accuracy. In the Cerrado studies, for example, uncertainties regarding the accuracy of spatial patterns of physiognomies and climatic seasonality throughout the biome (i.e., length of dry and wet seasons and rainfall amount) impact the absolute estimates of fire emissions and future projections. Some changes that can be made to improve carbon accounting from fire in the Cerrado include: acknowledging the heterogeneity of the biome, especially regarding climatic seasonality and fuel characteristics (Bustamante et al., 2018; Gomes et al., 2018; Oliveira et al., 2021); incorporating location-specific algorithms that capture the diversity of the biome (Libonati et al., 2015; Oliveira et al., 2021); better accounting of carbon stocks existent in the biome (Bustamante et al., 2018); and accounting for other carbon pools, such as soils and belowground, which are large components of carbon in the Cerrado's physiognomies (Bustamante et al., 2012). These will improve understanding of fire emissions in the Cerrado and their placement in the global carbon balance.~~

3.4 The influence of fire management and policy in estimating fire emissions in the Cerrado

In synthesizing the literature on fire emission in the Cerrado, we identified [128% \(12 papers\)](#) of papers focused on fire management and policy, [most all \(8\)](#) under the 'review' and 'perspective' categories. This indicates that fire management and policy are important in understanding fire dynamics in the Cerrado. Still, papers that address these do not usually bring new information based on observation or experiments but tend to synthesize or opine on existing literature. For example, this review captured [only one study that quantifies](#) ~~no studies quantifying~~ the amount of fire emissions mitigated by fire management in the Cerrado ([Franke et al. \(2024\)](#)), probably due to the difficulty in quantifying the social and cultural aspects of fire, which are intrinsic to fire management and policy. Estimating the influence of humans on fire emissions is a complex task, which is also reflected in the lack of equations and algorithms to reproduce fire management strategies in land surface models. [This complexity emphasizes](#) ~~at makes sense, given~~ all factors that need to be considered beyond quantifying the amount of GHG emitted to the atmosphere.

Fire exclusion policies arose in Brazil as a reaction to the misuse of fire for pasture management and deforestation, especially in the 20th century (Durigan and Ratter, 2016). ~~As a result, t~~These policies dominated for decades in the Cerrado (Durigan and Ratter, 2016), also due to the misbelief that fire harms the biome, resulting in fuel accumulation and changes in fire season and regime (Pivello, 2011; Pivello et al., 2021). Consequently, extensive LDS wildfires replaced small patchy burns (~~Fidelis, 2020;~~ Moura et al., 2019; Pivello, 2011). A shift towards recognizing fire as essential to maintaining the Cerrado's diversity and ecosystem services led to a change in the Federal Legislation in 2012 (Federal Law 12,651/2012) to explicitly allow the use of fire fire-management for ecologicaleconservation purposes in fire-prone settings (Durigan, 2020; Pivello et al., 2021). ~~(Federal Law 12,651/2012). Further, the -acknowledgment of controlled fires an essential socio-ecological component~~ in fire-dependent ecosystems led to the Brazilian Integrated Fire Management Policy (Pivello et al., 2021), sanctioned in 2024 (Federal Law 14,944/2024).

The changes in policy to recognize the importance of fire to the Cerrado is also reflected in the increased numbers of Protected Areas under fire management – the Pilot Integrated Fire Management project, named the Cerrado-Jalapão, started in three Protected Areas. The Cerrado-Jalapão project improved the understanding of fire dynamics in the biome (Durigan and Ratter, 2016). In 2024, Franke et al. (2024) identified 36 Protected Areas (PAs) undergoing Integrated Fire Management (IFM) activities. Similarly, Santos et al. (2021) documented an increase in the number and spatial distribution of prescribed burning in PAs within the Cerrado from 2015–2018. Prescribed burning is a common activity in fire management used to reduce fuel load in fire-prone settings around the world (Santos et al., 2021).

~~In 2014, the Pilot Integrated Fire Management project emerged in protected areas of the Cerrado with conservation purposes based on the Australian model (Schmidt et al., 2018), named the Cerrado Jalapão project. Cerrado Jalapão encouraged a further shift away from fire suppression policies, allowed scientific experiments to report the ecological benefits of fire in the Cerrado, and improved the understanding of fire dynamics in the biome (Durigan and Ratter, 2016; Schmidt and Eloy, 2020).~~

Despite the increasing recognition of fire's importance to the Cerrado and the subsequent expansion of fire management operations, as well as the global the-relevance of fire management in reducing emissions (Moura et al., 2019)to fire emissions, this review identified only one study that quantifies ~~did not identify studies quantifying~~ the reduction in fire carbon emissions achieved through fire management in the Cerrado. Franke et al. (2024) show an emission abatement of 26,677 tCO₂e year⁻¹ (2014 – 2019) in specific protected areas, and a reduction potential of more than 1,085 tCO₂e year⁻¹ (2014 – 2019) when the result is scaled-up to all protected areas in the Cerrado. The reduction in emissions from prescribed burning is due to lower fuel consumption and combustion factors of early dry season fires when compared to mid/late dry season fires (Franke et al., 2024). These values are considered conservative due to analyzing mainly fine fuels, and Franke et al. (2024) argue that estimations could be improved by using high-resolution data that would allow the identification of small-scala fires.

~~However, we~~This literature review also documents -found studies that estimate fire regime characteristics of fire management activities, such as early dry season burning, important for further emission estimates – Mistry et al. (2019) suggests reduced burned area and late dry season emissions from the Integrated Fire Management in the Cerradoparameters

of prescribed burning. Batista et al. (2018) compares two areas within the Canastra National Park, one under fire suppression and the other under fire management. The area under fire management presented less burned area annually (2000–2015), with a higher proportion of areas burned in the EDS. Similarly, Franke et al. (2024) identified increased EDS and decreased LDS burned area extents as a result of fire management. Over 2014–2019, Franke et al. (2024) estimated an average of 20.4% EDS burned area in Protected Areas (PAs) under Integrated Fire Management (IFM), against 4.7% in PAs that do not apply IFM. Conversely, middle to late dry season burned area averaged 76.5% in IFM PAs, while that average reaches 87.1% in non-IFM PAs. Santos et al. (2021) also documents reduced burned area in the late dry season due to fire management, which led to reduced fire intensity and reduced extreme wildfires.

Assessing fire emissions in the Cerrado is particularly complex due to the region's ecological diversity and the interplay of fire dynamics, policy, and cultural practices. Fire management remains challenged by the lack of understanding of fire dynamics parameters from early and late dry season burning (Mistry et al., 2019; Pivello et al. 2021), as estimated biomass, combustion completeness and burned area, and these affect fire emissions (Mistry et al., 2019). Understanding the role of these aspects in estimating fire emissions contributes to the development of consistent fire policies in Brazil (Durigan, 2020; Durigan and Ratter, 2016; Moura et al., 2019), which influence Brazil's national and international commitments to carbon emission reductions (Pivello et al., 2021).

Prescribed burning is a common activity under fire management, where fire under controlled conditions is seen as a technical and ecological tool to control fuel load, provide vegetation patchiness, and avoid late dry season intense wildfires while maintaining the ecosystem integrity (Myerspan, 2006; Schmidt et al., 2018). Prescribed burning is usually done at the end of the wet season or early dry season, and it reduces fire intensity and can result in less fire spread (dos Santos et al., 2021; Schmidt et al., 2018), suggesting a reduction in emissions from fire management activities.

Dos Santos et al. (2021) have shown that LDS burns have higher combustion factor, heat released, and fire intensity when compared to EDS burns. Fire management has reduced LDS area burned by 40–57% in the three PAs encompassed in the Cerrado Jalapão project during the first three years (2014–2016) of implementation (Schmidt et al., 2018). In the Canastra National Park in Brazil, areas under fire management also presented less annual area burned (Batista et al., 2018).

~~These reaffirm the potential of management activities to reduce emissions, as shown in other savanna countries. In northern Australia, more specifically in the WALFA (West Arnhem Land Fire Abatement) area, a region recognized as a reference for integrating fire studies with traditional knowledge, EDS burns emit 48% of what is emitted in the LDS (Russell-Smith et al., 2009). The WALFA project applies EDS burns to reduce LDS burns, and during its first 7 years of implementation, GHG emissions have reduced more than 37% when compared to the pre-project 10-year emissions baseline (Russell-Smith et al., 2013). Similarly, Khatun, Corbera, and Ball (2017) suggest that, in the Tanzanian miombo, EDS burns could avoid carbon emissions and enhance carbon uptake by approximately 10 tC ha⁻¹ in a 20-year period. Studies in Mozambique and Botswana explore the potential of EDS burns to reduce emissions in southern African savannas (Russell-Smith et al., 2021).~~

Assessing fire becomes even more challenging when dealing with a diverse region such as the Cerrado. From reviewing the literature, essential aspects that influence the estimation of emissions emerged, such as fire dynamics, ecology,

940 policy, and culture (see Arruda et al., 2018; Durigan and Ratter, 2016; Pivello et al., 2021). Understanding the role of these aspects in estimating fire emissions contributes to the development of consistent fire policies in Brazil (Arruda et al., 2018; Bustamante et al., 2018; Durigan and Ratter, 2016; Gomes et al., 2018), which influence Brazil's national and international commitments to carbon emission reductions (da Silva Junior et al., 2020; Pivello et al., 2021).

4 Discussion

945 By analysing existing literature on fire emissions in the Cerrado, we identified the main topics that allow for a ~~This systematic review allows for a~~ broad and holistic ~~etter~~ understanding of ~~the role of these emissions~~ Cerrado's placement in the carbon budget on regional, national and global scales, ~~and these include not only fire carbon emission estimates, but also fire dynamic parameters and fire management and policy.~~ -In synthesizing literature, we ~~observed~~ identified a growing interest in ~~understanding~~ fire ~~sciencedynamics~~ in the Cerrado from multiple perspectives, as shown in the number of papers published annually. Pivello et al. (2021) also observes an increase of studies on fire. This is probably due to an increased recognition of fire's importance in the global carbon balance and the increased number of alarming fire events reported in recent years ~~because of~~ due to climate change ~~and intense human activities~~ (Burton et al., 2024; Hofmann et al. 2021; Oliveira et al., 2022; ~~Loehman et al. 2014~~).

955 The year 2022 did not follow the growth trend, which could indicate a gap in publications this year or a limitation of our research method that could not capture publications in 2022. The following years tend to increase publications compared to 2022. The decrease in 2022 could also indicate a shifted focus away from the Cerrado studies due to political or financial constraints to encourage scientific studies in the region, or due to a shifted focus towards other regions of Brazil. Pereira et al. (2024) indicate an increase in papers published about fires in the Pantanal after the 2020 megafire in the biome. Papers related to fire dynamics and emissions in the Pantanal published in 2022 show fire as a consequence of the compound impact of land use and climate in these regions, and similar findings were reported for the Amazon rainforest in papers also published in 2022 (see Barbosa et al., 2022; Dutra et al., 2022; Menezes et al., 2022; Silva et al., 2022; Walker et al., 2022).

960 -We identified that papers often encompass a global and regional emissions analysis, with local (Cerrado) analysis accounting for ~~about~~ 368% of papers included in this review. The remaining 642% of papers represent global and regional (Brazil, South America, Tropical region) analysis, often understanding the role of emissions ~~from a holistic perspective on a broad scale~~ and providing insights into the influence of local emissions on the global carbon balance. Additionally, ~~we found that many most of the~~ papers covered in this systematic literature review ~~are is led driven~~ by non-Brazilian institutions ~~and/or do not include~~ lack authors ~~from associated with~~ Brazilian institutions. In fact, fire emissions studies in the Cerrado are rarely led by institutions within the region. -Of all the papers included, only 10 involve first authors from institutions located within the Cerrado region. This ~~highlights indicates an~~ opportunitiesy to enhance collaboration between Brazilian and non-Brazilian institutions, and ~~even a potential to increase~~ stronger partnership between different regions within Brazil.

970

975 The papers reviewed have shown that input data uncertainty affects output accuracy. In the Cerrado studies, for example, uncertainties regarding the accuracy of spatial patterns of physiognomies and climatic seasonality throughout the biome (i.e., length of dry and wet seasons and rainfall amount) impact the absolute estimates of fire emissions and future projections. Some changes that can be made to improve carbon accounting from fire in the Cerrado include: acknowledging the heterogeneity of the biome, especially regarding climatic seasonality and fuel characteristics (Gomes et al., 2018; Oliveira et al., 2021); incorporating location-specific algorithms and datasets that improves the representation of the Cerrado (Libonati et al., 2015; Mataveli et al., 2024; Oliveira et al., 2021); and accounting for other carbon pools, such as soils and belowground, which are large components of carbon in the Cerrado's physiognomies (Bustamante et al., 2012). These will improve understanding of fire emissions in the Cerrado and their placement in the global carbon balance.

980 Examining fire carbon emissions reveals that local emissions reflect the global carbon budget. A key factor in carbon balance analysis is vegetation regrowth, since a significant portion of the CO₂ emitted by fire is sequestered during post-fire biomass recovery (Andreae, 2019; Van Der Werf et al., 2017; Gomes et al., 2024). This literature review identified only one study that quantifies the net CO₂ emissions from the Cerrado fires from 1985–2020 (Gomes et al., 2024). Vegetation regrowth removed 63.5% of the CO₂ emitted, indicating that fire in the Cerrado has been a source of carbon to the atmosphere in recent
985 decades (Gomes et al., 2024). For a shorter time series (2015–2018), Oliveira et al. (2021) also found the Cerrado fires to be a net emitter of CO₂. Further research is needed to enhance the understanding of the long-term carbon balance of Cerrado fires. This literature review contributes by providing an overview of published studies on fire emissions in the region.

990 Compiling literature on fire emissions in the Cerrado has revealed several studies that focus on fire dynamics, management, and policy rather than estimating emissions, with studies often stating their relevance in providing insights into emission estimates and the importance of the Cerrado's fire emission in the global carbon balance. This indicates that estimating fire emissions requires a holistic approach. For example, including the perspectives of fire culture, ecology and policy within emissions is essential given the importance of fire to the biome. Fire culture refers to the interaction between humans and fire, encompassing the factors that drive societies to use it (see Pivello et al., 2021). The use of fire, shaped by cultural traditions and socioeconomic conditions, can influence the extent of burned areas and the resulting fire emissions.
995 Traditional communities, for instance, may occasionally use fire on a small scale (Pivello et al., 2021).

1000 Coupled with the available biomass for burning and its characteristics — which depend on vegetation type, density, moisture and seasonal growth patterns — burned area, fuel characteristics, combustion completeness, combustion efficiency and emission factor set the stage for potential emissions. Fire intensity, driven by conditions such as dry weather, strong winds, and fuel accumulation, influences combustion efficiency. High-intensity fires tend to consume more fuel, resulting in higher combustion efficiency and more complete combustion. This reduces emissions of pollutants such as carbon monoxide and particulate matter but increases emissions of carbon dioxide. In contrast, incomplete combustion results in higher emissions of pollutants such as particulate matter and carbon monoxide and produces pyrogenic carbon, which may persist in soils over long periods. Combustion completeness further influences the amount of biomass converted to carbon and released into the

atmosphere. Together, these parameters allow for the estimation of emissions based on the combination of burned area, fuel load, and combustion completeness.

Thus, this review indicates a critical need to develop interdisciplinary studies to bridge fire policies and fire emissions in the Cerrado. Understanding fire dynamics, including the opportunities for mitigating emissions from fire activities, is essential for recognizing fire's role in achieving global environmental and climate targets. For instance, Martin (2019) identifies United Nations Sustainable Development Goals that are related to fire and land management, as goals 3 (good health and well-being), 13 (climate action), and 15 (life on land). These impact the 2015 Paris Agreement target to limit warming to 1.5 °C by 2100. In fact, da Silva Junior et al. (2020) find that fire activities undermine Brazil's emission reduction targets under the Paris Agreement. The Paris Agreement outlines commitments for climate actions and acknowledges the importance of mitigation and removal actions, where fire management can play an important role. If great effort is put into mitigating emissions and removing carbon, the 1.5 °C could be achieved, although ambitious, with Brazil holding the highest mitigation potential in the land sector (Roe et al., 2019). Together with other countries, improved forest management – which includes fire management – in Brazil could be able to increase carbon removal by 40 GtCO₂ by 2050 (Roe et al., 2019).

Studies that discuss the cultural and socioeconomic aspects of fire often do not discuss them from the emissions' perspective. Despite the difficulty in quantifying the social and cultural aspects of fire, these, the lack of inclusion of these aspects in fire emission estimates could also be due to the shift towards recognizing fire as essential to the Cerrado being recent, especially when compared to other fire-prone settings. For example, the WALFA (West Arnhem Land Fire Abatement) project in northern Australia became entirely active in 2005 (Russell-Smith et al., 2013), where traditional people, scientists and governmental institutions collaborate to reduce fire emissions through fire management activities (Russell-Smith et al., 2013). Meanwhile, the Pilot Integrated Fire Management project in the Cerrado started in 2014 (Schmidt et al., 2018).

Compared with the Cerrado, a relatively high number of fire studies were performed in Australia. Da Veiga and Nikolakis (2022) counted 64 papers from Australia and 29 from Brazil when documenting the interaction between fire management and carbon programs worldwide. The Australian studies, especially in the WALFA region, act as an example to other savanna environments, including to the Cerrado's Pilot Integrated Fire Management project (Schmidt et al., 2018). Meanwhile, the Pilot Integrated Fire Management project in the Cerrado started in 2014 (Schmidt et al., 2018). The WALFA project reaffirms the potential of management activities to reduce emissions in savanna countries. EDS burns in the region emit 48% of what is emitted in the LDS (Russell-Smith et al., 2009). Similarly, Khatun, Corbera, and Ball (2017) suggest that, in the Tanzanian miombo, EDS burns could avoid carbon emissions and enhance carbon uptake. Studies in Mozambique and Botswana explore the potential of EDS burns to reduce emissions in southern African savannas (Russell-Smith et al., 2021).

Climate change increasingly affects fires, and adaptation and mitigation activities are essential to limit these effects (Burton et al., 2024). Direct human impacts may offset the effects of climate change in fire worldwide (Burton et al., 2024), especially in fire-prone environments, and this indicates an opportunity to investigate the potential of fire management to mitigate emissions in the Cerrado, and to understand fire emissions in the biome. The lack of data availability and accessibility

has been acknowledged in the literature as a gap in research regarding fire dynamics in the Cerrado, primarily due to its heterogeneity (Bustamante et al., 2018; Gomes et al., 2018; Oliveira et al., 2021). Developing more specific algorithms able to capture the climatic seasonality and fuel diversity of the Cerrado, understanding the above and belowground carbon pools of each physiognomy by in situ evaluations and satellite-derived approaches, and incorporating these into fire emissions estimates in the Cerrado could improve carbon measurements in the biome.

Consequently, this would improve fire emission estimates in the Cerrado. Pathways towards this improvement include connecting observational information with modeling and a better assessment and quantification of the impact of qualitative aspects in fire estimates. Examples of how these can be achieved are by valuing prescribed burning emissions and including these in fire modeling, representing fire management in land surface models, using in situ observations to assess models' utility and as input data to modeling, and incorporating the ecological, social and cultural aspects of fire in fire emission estimates. These could address uncertainty and improve models' accuracy, thus providing better accounting of fire emissions in the Cerrado and worldwide.

These reaffirm the potential of management activities to reduce emissions, as shown in other savanna countries. In northern Australia, more specifically in the WALFA (West Arnhem Land Fire Abatement) area, a region recognized as a reference for integrating fire studies with traditional knowledge, EDS burns emit 48% of what is emitted in the LDS (Russell-Smith et al., 2009). The WALFA project applies EDS burns to reduce LDS burns, and during its first 7 years of implementation, GHG emissions have reduced more than 37% when compared to the pre-project 10-year emissions baseline (Russell-Smith et al., 2013). Similarly, Khatun, Corbera, and Ball (2017) suggest that, in the Tanzanian miombo, EDS burns could avoid carbon emissions and enhance carbon uptake by approximately 10 tC ha⁻¹ in a 20-year period. Studies in Mozambique and Botswana explore the potential of EDS burns to reduce emissions in southern African savannas (Russell-Smith et al., 2021).

~~Most papers provide new information on the parameters used to estimate fire emissions and on emissions themselves rather than analyzing existing literature. These often handle modelling techniques, satellite observations, on-site observations and even results found in previous studies, reflecting the importance of literature reviews in supporting the development of new data. These tools are often combined to provide a more complete analysis. Modelling and satellite observation are frequently integrated, with satellite observations being data providers to models, or models being evaluated by satellite observations, or even through data assimilation between satellite data and modelling simulations. From our results, 13 empirical papers integrate modelling and satellite data, 2 integrate modelling and literature review, 1 integrates satellite and on-site observations, and none directly integrate modelling and on-site observations. Figure 6 shows the methodological techniques used across study areas in empirical papers.~~




Figure 6: Treemap of the methodological techniques used across study areas in empirical papers. The numbers present the number of studies of each methodological technique within each study area. The study areas Global, Tropical region, South America and Brazil are regions that include results for the Cerrado. Some papers combine different techniques and are double-counted.

Therefore, estimating local fire emissions is reflected in the global carbon budget. A vital factor to consider when analyzing fire emissions is vegetation regrowth since much of the emitted CO_2 is sequestered during the recovery of the post-fire ecosystem (Andreae, 2019; Van Der Werf et al., 2017). In savanna environments such as the Cerrado, fine fuel load, responsible for 95% of fuel available for burning in the biome, can regain its initial biomass 15 months after a fire event (Gomes et al., 2020b). However, it is uncertain if biomass recovery in the Cerrado fully offsets fire emissions, especially under changing fire regimes, and better accounting for vegetation regrowth is needed to understand if fire dynamics are a net CO_2 source or sink to the atmosphere in the long term and what are the mitigation opportunities to assess this issue.

Our research question is “How compiling published material on fire emissions in natural areas of the Cerrado ~~that do not explicitly include anthropogenic land uses~~ can provide a holistic better understanding of their role ~~placement of these emissions~~ in the global carbon budget?”. Analyzing published papers on fire emissions in these areas in the Cerrado provides valuable insights into its role in the carbon balance. This includes understanding the parameters used to estimate emissions, quantifying the amount of carbon, especially CO_2 , released into the atmosphere by fires, and identifying important aspects of

fire dynamics that are sources of uncertainty or are not considered in fire emission estimates. These are summarized in Table 3.

Table 3. Parameters included in current studies and parameters to be considered for future studies to estimate fire emissions in the Cerrado captured by this review.

Parameters included in current studies in the Cerrado	Parameters to be considered for future fire emission estimates in the Cerrado	1105
Biomass burned	Belowground and soil carbon pools	
Burned area	Fire culture	
Combustion efficiency	Fire ecology	
Combustion factor	Fire policy	
Emission factor	Location-specific algorithms	1110
Fuel characteristics	Socioeconomic aspects of fire	

~~Aiming at compiling literature on fire emissions in the Cerrado has led to several papers that do not explicitly estimate fire emissions themselves, but rather discuss fire dynamics and parameters used to estimate emissions. This indicates that there is a gap in the literature regarding fire emissions estimates in the Cerrado. However, studies have indicated that fires in the Cerrado play an important role in the global carbon balance. For example, Van Der Werf et al. (2017) found that savanna fire emissions from the Southern Hemisphere South America region, which includes the Cerrado, averaged 0.14 PgC year⁻¹ over 20 years, accounting for more than 6% of global fire emissions per year. Similarly, and from a national perspective, da Silva Junior et al. (2020) have shown Cerrado fires contribute more than 32% of the Brazilian total fire emissions (about 0.13 PgC year⁻¹ over the 20 years).~~

~~Our review also indicates that published literature fails to analyze fire emissions from a holistic approach in the Cerrado. Including the perspectives of fire culture, ecology and policy within emissions is essential given the importance of fire to the biome. However, studies that discuss these aspects often do not discuss them from the emissions' perspective. Despite the difficulty in quantifying the social and cultural aspects of fire, the lack of inclusion of these in fire emission estimates could also be due to the shift towards recognizing fire as essential to the Cerrado being recent, especially when compared to other fire prone settings. For example, the WALFA project in northern Australia became entirely active in 2005 (Russell Smith et al., 2013), where traditional people, scientists and governmental institutions collaborate to reduce fire emissions through fire management activities (Russell Smith et al., 2013). Meanwhile, the Pilot Integrated Fire Management project in the Cerrado started in 2014 (Schmidt et al., 2018).~~

~~Thus, this review indicates a critical need to develop interdisciplinary studies to bridge fire policies and fire emissions in the Cerrado. Understanding fire dynamics, including the opportunities for mitigating emissions from fire activities, is essential for recognizing fire's role in achieving global environmental and climate targets. For instance, Martin (2019) identifies United Nations Sustainable Development Goals that are related to fire and land management, as goals 3 (good health and well-being), 13 (climate action), and 15 (life on land). These impact the 2015 Paris Agreement target to limit warming to 1.5 °C by~~

2100. The Paris Agreement outlines commitments for climate actions and acknowledges the importance of mitigation and removal actions, where fire management can play an important role. The 1.5 °C target is ambitious, yet achievable if great effort is put into mitigating emissions and removing carbon, with Brazil holding the highest mitigation potential in the land sector (Roe et al., 2019). Together with other countries, improved forest management—which includes fire management—in Brazil could be able to increase carbon removal by 40 GtCO₂ by 2050 (Roe et al., 2019).

Climate change increasingly affects fires, and adaptation and mitigation activities are essential to limit these effects (Burton et al., 2024). Direct human impacts may offset the effects of climate change in fire worldwide (Burton et al., 2024), especially in fire-prone environments, and this is an opportunity to investigate the potential of fire management to mitigate emissions in the Cerrado, and to understand fire emissions in the biome. Pathways towards improving fire emissions in the Cerrado include connecting observational information with modeling and a better assessment and quantification of the impact of qualitative aspects in fire estimates. Examples of how these can be achieved are by valuing prescribed burning emissions and including these in fire modeling, representing fire management in land surface models, using on-site observations to assess models' utility and as input data to modeling, and incorporating non-carbon aspects of fire in fire emission estimates, such as the ecological, social and cultural aspects. These could address uncertainty and improve models' accuracy, thus providing better accounting of fire emissions in the Cerrado and worldwide.

This systematic literature review presents an overall assessment of published literature on fire emissions in the Cerrado to understand its placement in the carbon budget, considering the criteria used to narrow our search. Including Portuguese as a research language and consulting other search database platforms, such as Web of Science, could have resulted in more papers included. The “grey literature” was not in the scope of our research method but could potentially result in more findings. “Grey literature” is defined as “*unpublished research (e.g., dissertations, conference abstracts, preprints or unpublished datasets), or those published outside of traditional academic publishing (e.g., governmental reports)*” (Foo et al., 2021, 1711). However, to our knowledge, this is the first systematic literature review to present a general and holistic scope of fire emissions in the Cerrado.

Finally, the challenges in estimating fire emissions worldwide lie not only in measuring carbon directly released, but also in capturing the nuanced effects of fire on ecosystems and the broader Earth system (Hamilton et al., 2024). Continued and diversified research is needed to improve the understanding of fire dynamics in the Cerrado and how these reflect fire emissions locally and globally. These will assess current knowledge gaps regarding fire emission estimates in the Cerrado by enhancing the understanding of the role of emissions from the biome in the global carbon budget and potential mitigation activities in achieving global environmental and climatic goals, and by providing a better grounding for future projections.

5 Conclusion

This systematic literature review synthesized ~~69-77~~ peer-reviewed papers, from local to global scales, according to a set of criteria to understand fire emissions in the Cerrado and their ~~the placement of fire emissions in the Cerrado~~ in the global

carbon budget, from a holistic perspective. Based on our knowledge and search criteria, this is the first systematic literature review to provide an overview an integrated understanding of fire emissions in the Cerrado, where fire dynamics, management and policy emerge as crucial for estimating fire emissions. By providing necessary context on the drivers of emissions, their variability, and potential mitigation strategies, we highlight how insights from fire behavior and fire regimes are crucial for emissions estimation-

Although our research question and keyword search focus on fire emissions, the literature review process often revealed papers on fire dynamics that are drivers to emissions estimates, indicating a gap in connecting fire dynamics and emissions research in the Cerrado. ~~Thus, w~~We then examine how fire dynamics shape emissions and identify where further integration of these fields could improve our holistic understanding of fire emissions in the Cerrado. This is possible due to a clearer understanding of the variables considered when estimating emissions, a deeper comprehension of the published values estimated, and the identification of important aspects of fire regimes dynamics that influence fire policy and mitigation strategies still need to be considered.

Fire management and policy emerged as key themes in our literature search because they play a crucial role in shaping fire regimes and, consequently, fire emissions in the Cerrado. While there is limited empirical research quantifying the direct impact of fire management on emissions in this region, discussing management strategies is essential to fully assessing the factors that influence fire-driven carbon fluxes. Understanding how management affects fire regimes provides a pathway to improving emissions estimates and identifying potential mitigation strategies.

~~Thus, ese demonstrate that~~ this review demonstrates that understanding the placement of fire emissions in the global carbon budget requires estimating fire emissions is complex and requires a holistic approach that draws together disciplines across fire science, especially in a distinct environment such as the Cerrado. The complexity of estimating fire emissions, and estimating emission outreach measuring carbon, and. ~~Thus,~~ this review highlights the urgent need for interdisciplinary studies to connect fire parameters with fire emissions, which then influence fire policies and the achievement of climatic commitments. Continued research is needed to fully understand and quantify the influence of fire dynamics and mitigation strategies onand emissions in the Cerrado, ~~and we suggest pathways to achieving improvements.~~

~~In assessing documented literature, we identified three prominent topics when searching about fire emissions in the Cerrado: fire dynamics parameters, emission estimates, and fire management and policy. These demonstrate that estimating fire emissions is complex and requires a holistic approach that draws together disciplines across fire science, especially in a distinct environment such as the Cerrado, and estimating emission outreach measuring carbon. Thus, this review highlights the urgent need for interdisciplinary studies to connect fire parameters with fire emissions, which then influence fire policies and the achievement of climatic commitments. This complexity is reflected in the divergence of values and units of carbon emissions estimates and in the multiple parameters used to estimate emissions.~~

~~This review demonstrates that papers fail to report on fire emissions themselves, with fire dynamics and parameters used to estimate emissions in the Cerrado often being the focus of published literature. However, we found that reviewing literature on fire emissions from the Brazilian Cerrado biome contributes to the understanding of the placement of these~~

~~emissions in the global carbon budget. This is possible due to a clearer understanding of the variables considered when estimating emissions, a deeper comprehension of the published values estimated, and the identification of important aspects of fire dynamics that still need to be considered. These are typically of qualitative nature.~~

1205 **Code availability:** No codes were used in this literature review.

Data availability: The list of papers reviewed is available in the Supplement.

Authors' contribution: RMV conceptualized the study and prepared the original draft. RMV, CvR, CB, DIK, MC and FM
1210 interpreted and analyzed data. RMV, CvR, CB, DIK and MC substantially revised and edited the study.

Competing interests: The authors declare that they have no conflict of interest.

Acknowledgments: RMV thanks the São Paulo Research Foundation (FAPESP) for grants 2020/06470-2 and 2022/13322-5.
1215 CvR acknowledges FAPESP grant 2017/22269-2 and CNPq grant 314780/2020-3. CB was funded by the Met Office Climate Science for Service Partnership (CSSP) Brazil project which is supported by the Department for Science, Innovation & Technology (DSIT). DIK was supported by the Natural Environment Research Council as part of the NC-International programme [NE/X006247/1] delivering National Capability. MC acknowledges the support from the São Paulo Research Foundation (FAPESP, Process 2015/50122-0).

1220 **References**

Alvarado, S. T., Fornazari, T., Cóstola, A., Morellato, L. P. C., and Silva, T. S. F.: Drivers of fire occurrence in a mountainous Brazilian cerrado savanna: Tracking long-term fire regimes using remote sensing, *Ecol. Indic.*, 78, 270–281, <https://doi.org/10.1016/j.ecolind.2017.02.037>, 2017.

~~Andela, N., Morton, D. C., Giglio, L., Paugam, R., Chen, Y., Hantson, S., Van Der Werf, G. R., and Anderson, J. T.: The
1225 Global Fire Atlas of individual fire size, duration, speed and direction, *Earth Syst. Sci. Dat.*, 11, 529–552,
<https://doi.org/10.5194/essd-11-529-2019>, 2019.~~

Andersen, J., Belmont, J., and Cho, C. T.: Journal impact factor in the era of expanding literature, *J. Microbiol. Immunol. Infect.*, 39, 436–443, 2006.

Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmos. Chem. Phys.*,
1230 19, 8523–8546, <https://doi.org/10.5194/acp-19-8523-2019>, 2019.

- Arruda, F. V., de Sousa, D. G., Teresa, F. B., do Prado, V. H. M., da Cunha, H. F., and Izzo, T. J.: Trends and gaps of the scientific literature about the effects of fire on Brazilian Cerrado, *Biota Neotrop.*, 18, 1–6, <https://doi.org/10.1590/1676-0611-bn-2017-0426>, 2018.
- 235 [Arruda, V. L. S., Alencar, A. A. C., de Carvalho Junior, O. A., Ribeiro, F. de F., de Arruda, F. V., Conciani, D. E., da Silva, W. V., Shimbo, J. Z.: Assessing four decades of fire behavior dynamics in the Cerrado biome \(1985 to 2022\), *Fire Ecol.*, 20, <https://doi.org/10.1186/s42408-024-00298-4>, 2024.](#)
- Barbosa, M. L. F., Haddad, I., da Silva, A. L. N., da Silva, G. M., da Veiga, R. M., Hoffmann, T. B., de Souza, A. R., Dalagnol, R., Streher, A. S., Pereira, F. R. S., Aragão, L. E. O.C., Anderson, L. O., and Poulter, B.: Compound impact of land use and extreme climate on the 2020 fire record of the Brazilian Pantanal, *Glob. Ecol. Biogeogr.*, 1–16, 1240 <https://doi.org/10.1111/geb.13563>, 2022.
- Batista, E. K. L., Russell-Smith, J., França, H., and Figueira, J. E. C.: An evaluation of contemporary savanna fire regimes in the Canastra National Park, Brazil: Outcomes of fire suppression policies, *J. Environ. Manag.*, 205, 40–49, <https://doi.org/10.1016/j.jenvman.2017.09.053>, 2018.
- 245 ~~Bistinas, I., Harrison, S. P., Prentice, I. C., and Pereira, J. M. C.: Causal relationships versus emergent patterns in the global controls of fire frequency, 11, 5087–5101, <https://doi.org/10.5194/bg-11-5087-2014>, 2014.~~
- ~~Bowman, D. M. J. S., Baleh, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A., D’Antonio, C. M., DeFries, R. S., Doyle, J. C., Harrison, S. P., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Marston, J. B., Moritz, M. A., Prentice, I. C., Roos, C. I., Scott, A. C., Swetnam, T. W., Van der Werf, G. R., and Pyne, S. J.: Fire in the earth system, *Science*, 324, 481–484, <https://doi.org/10.1126/science.1163886>, 2009.~~
- 1250 Brazilian Agricultural Research Corporation (Embrapa, in Portuguese): Cerrado Biome, <https://www.embrapa.br/cerrados/colecao-entomologica/bioma-cerrado>, last access: 16 May 2024.
- Brazilian Statistics Institute (IBGE, in Portuguese): Territory, <https://brasilemsintese.ibge.gov.br/territorio>, last access: 02 February 2024, –2004.
- 255 ~~Burton, C., Betts, R., Cardoso, M., Feldpausch, R. T., Harper, A., Jones, C. D., Kelley, D. I., Robertson, E., and Wiltshire, A.: Representation of fire, land use change and vegetation dynamics in the Joint UK Land Environment Simulator vn4.9 (JULES), *Geosci. Model Dev.*, 12, 179–193, <https://doi.org/10.5194/gmd-12-179-2019>, 2019.~~
- Burton, C., Kelley, D. I., Jones, C. D., Betts, R. A., Cardoso, M., and Anderson, L.: South American fires and their impacts on ecosystems increase with continued emissions, *Clim. resil. Sustain.*, 1–15, <https://doi.org/10.1002/cli2.8>, 2021.
- 1260 Burton, C., Lampe, S., Kelley, D. I., Thiery, W., Hantson, S., Christidis, N., Gudmundsson, L., Forrest, M., Burke, E., Chang, J., Huang, H., Ito, A., Kou-Giesbrecht, S., Lasslop, G., Li, W., Nieradzick, L., Li, F., Chen, Y., Randerson, J., Reyer, P. O. C., and Mengel, M.: Global burned area increasingly explained by climate change, *Nat. Clim. Change*, 14, 1186–1192, 2024.
- Bustamante, M. M. C., Silva, J. S. O., Cantinho, R. Z., Shimbo, J. Z., Oliveira, P. V. C., Santos, M. M. O., Ometto, J. P. H. B., Cruz, M. R., Mello, T. R. B., Godiva, D., and Nobre, C. A.: Engagement of scientific community and transparency in C

- accounting: the Brazilian case for anthropogenic greenhouse gas emissions from land use, land-use change and forestry, Environ. Res. Lett., 13, 055005, <https://doi.org/10.1088/1748-9326/aabb37>, 2018.
- Bustamante, M., Nardoto, G., Pinto, A., Resende, J., Takahashi, F., and Vieira, L.: Potential impacts of climate change on biogeochemical functioning of Cerrado ecosystems, Braz. J. Biol., 72, 655–671, <https://doi.org/10.1590/S1519-69842012000400005>, 2012.
- Carvalho Jr., J. A., Higuchi, N., Araújo, T. M., and Santos, J. C.: Combustion completeness in a rainforest clearing experiment in Manaus, Brazil, J. Geophys. Res., 103, D11, 13195–13199, 1998.
- ~~Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C., and Cox, P. M.: The Joint UK Land Environment Simulator (JULES), model description — Part 2: Carbon fluxes and vegetation dynamics, Geosci. Model Dev., 4, 701–722, <https://doi.org/10.5194/gmd-4-701-2011>, 2011.~~
- ~~Coutinho, L. M.: Fire in the Ecology of the Brazilian Cerrado, in: Fire in the Tropical Biota, 82–105, https://doi.org/10.1007/978-3-642-75395-4_6, 1990.~~
- ~~Chen, Y., Hall, J., Van Wees, D., Andela, N., Hantson, S., Giglio, L., Van Der Werf, G. R., Morton, D. C., Randerson, J. T.: Multi-decadal trends and variability in burned area from the fifth version of the Global Fire Emissions Database (GFED5), Earth Syst. Sci. Data, 15, 5227–5259, <https://doi.org/10.5194/essd-15-5227-2023>, 2023.~~
- Cronin, P., Ryan, F., and Coughian, M.: Undertaking a literature review: a step-by-step approach, Br. J. Community Nurs., 17, 2008.
- da Silva Junior, C. A., Teodoro, P. E., Delgado, R. C., Teodoro, L. P. R., Lima, M., Pantaleão, A. A., Baio, F. H. R., de Azevedo, G. B., Azevedo, G. T. O. S., Capristo-Silva, G. F., Arvor, D., and Facco, C. U: Persistent fire foci in all biomes undermine the Paris Agreement in Brazil, Sci. Rep., 10, 16246, <https://doi.org/10.1038/s41598-020-72571-w>, 2020.
- da Veiga, R. M., and Nikolakis, W: Fire Management and Carbon Programs: A Systematic Literature Review and Case Study Analysis, Soc. Natur. Resour., 35, 896–913, <https://doi.org/10.1080/08941920.2022.2053618>, 2022.
- ~~DeBano, L.F., Neary, D.G., Ffolliott, P.F.: Fire Effects on Ecosystems, John Wiley & Sons, New York, 1998.~~
- ~~dos Santos, A. C., Montenegro, S. da R., Ferreira, M. C., Barradas, A. C. S., and Schmidt, I. B.: Managing fires in a changing world: Fuel and weather determine fire behavior and safety in the neotropical savannas, J. Environ. Manage., 289, <https://doi.org/10.1016/j.jenvman.2021.112508>, 2021.~~
- dos Santos, D. M., de Oliveira, A. M., Duarte, E. S. F., Rodrigues, J. A., Menezes, L. S., Albuquerque, R., Roque, F. de O., Peres, L. F., Hoelzemann, J. J., and Libonati, R.: Compound dry-hot-fire events connecting Central and Southeastern South America: an unapparent and deadly ripple effect, Nat. Hazards, 1, 1–13, <https://doi.org/10.1038/s44304-024-00031-w>, 2024.
- ~~Durigan, G.: Zero-fire: Not possible nor desirable in the Cerrado of Brazil. Flora, 268, <https://doi.org/10.1016/j.flora.2020.151612>, 2020.~~
- Durigan, G., and Ratter, J. A.: The need for a consistent fire policy for Cerrado conservation, J. Appl. Ecol., 53, 11–15, <https://doi.org/10.1111/1365-2664.12559>, 2016.

- Dutra, D. J., Anderson, L. O., Fearnside, P. M., Graça, P. M. L. de A., Yanai, A. M., Dalagnol, R., Burton, C., Jones, C., Betts, R., and Aragão, L. E. O. C.: Fire Dynamics in an Emerging Deforestation Frontier in Southwestern Amazonia, Brazil, *Fire*, 6, 21–24, <https://doi.org/10.3390/fire6010002>, 2022.
- Ehhalt, D., Prather, M., Dentener, F., Derwent, R., Dlugokencky, E., Holland, E., Isaksen, I., Katima, J., Kirchhoff, V., Matson, O., Midgley, P., and Wang, M.: Atmospheric Chemistry and Greenhouse Gases, in: *Climate Change 2001: The Scientific Basis, Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Houghton, J. T., Ding, Y., Griggs, D. J., Noguer, M., van der Linden, P. J., Dai, X., Maskell, K., and Johnson, C. A., Cambridge University Press, Cambridge, United Kingdom and New York, USA, 239–287, ISBN: 0521014956, 2001.
- ~~Eloy, L., A. Bilbao, B., Mistry, J., and Schmidt, I. B.: From fire suppression to fire management: Advances and resistances to changes in fire policy in the savannas of Brazil and Venezuela, *Geogr. J.*, 185, 10–22, <https://doi.org/10.1111/geoj.12245>, 2019.~~
- Fidelis, A.: Is fire always the “bad guy”?, *Flora*, 268, <https://doi.org/10.1016/j.flora.2020.151611>, 2020.
- ~~Fidelis, A., Alvarado, S., Barradas, A., and Pivello, V.: The Year 2017: Megafires and Management in the Cerrado: *Fire*, 1, 1–49, <https://doi.org/10.3390/fire1030049>, 2018.~~
- Flannigan, M. D., Krawchuk, M. A., De Groot, W. J., Wotton, B. M., and Gowman, L. M.: Implications of changing climate for global wildland fire: *Int. J. Wildland Fire*, 18, 483–507, <https://doi.org/10.1071/WF08187>, 2009.
- Foo, Y. Z., O’Dea, R. E., Koricheva, J., Nakagawa, S., and Lagisz, M.: A practical guide to question formation, systematic searching and study screening for literature reviews in ecology and evolution, *Methods Ecol. Evol.*, 12, 1705–1720, <https://doi.org/10.1111/2041-210X.13654>, 2021.
- Franke, J., Barradas, A. C. S., Borges, M. A., Menezes Costa, M., Dias, P. A., Hoffmann, A. A., Orozco Filho, J. C., Melchiori, A. E., and Siegert, F.: Fuel load mapping in the Brazilian Cerrado in support of integrated fire management, *Remote Sens. Environ.*, 217, 221–232, <https://doi.org/10.1016/j.rse.2018.08.018>, 2018.
- Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le Quéré, C., Luijkx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp, L., Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T.-T.-T., Chevallier, F., Chini, L. P., Cronin, M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato, E., Keeling, R. F., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X., Lefèvre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G., Morgan, E. J., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O’Brien, K. M., Olsen, A., Omar, A. M., Ono, T., Paulsen, M., Pierrot, D., Pocock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C., Rosan, T. M., Schwinger, J., Séférian, R., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tans, P. P., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., van Ooijen, E.,

- Wanninkhof, R., Watanabe, M., Wimart-Rousseau, C., Yang, D., Yang, X., Yuan, W., Yue, X., Zaehle, S., Zeng, J., and Zheng, B.: Global Carbon Budget 2023, *Earth Syst. Sci. Data*, 15, 5301–5369, <https://doi.org/10.5194/essd-15-5301-2023>, 2023.
- Gomes, L., Miranda, H. S., & Bustamante, M. M. C.: How can we advance the knowledge on the behavior and effects of fire in the Cerrado biome?: *For. Ecol. Manag.*, 417, 281–290, <https://doi.org/10.1016/j.foreco.2018.02.032>, 2018.
- Gomes, L., Miranda, H. S., Silvério, D. V., and Bustamante, M. M. C.: Effects and behaviour of experimental fires in grasslands, savannas, and forests of the Brazilian Cerrado, *For. Ecol. Manag.*, 458, <https://doi.org/10.1016/j.foreco.2019.117804>, 2020a.
- Gomes, L., Miranda, H. S., Soares-Filho, B., Rodrigues, L., Oliveira, U., and Bustamante, M. M. C.: Responses of Plant Biomass in the Brazilian Savanna to Frequent Fires, *Front. for. glob. change.*, 3, 1–11, <https://doi.org/10.3389/ffgc.2020.507710>, 2020b.
- Gomes, L., Schüller, J., Silva, C., Alencar, A., Zimbres, B., Arruda, V., Silva, W. V. da, Souza, E., Shimbo, J., Marimon, B. S., Lenza, E., Fagg, C. W., Miranda, S., Morandi, P. S., Marimon-Junior, B. H., Bustamante, M.: Impacts of Fire Frequency on Net CO₂ Emissions in the Cerrado Savanna Vegetation, *Fire*, 7, <https://doi.org/10.3390/fire7080280>, 2024.
- Greenhouse Gas Emissions and Removals Estimation System (SEEG, in Portuguese): seeg.eco.br, last access: 14 November 2024, 2023.
- Griscom, B. W., Busch, J., Cook-Patton, S. C., Ellis, P. W., Funk, J., Leavitt, S. M., Lomax, G., Turner, W. R., Chapman, M., Engelmann, J., Gurwick, N. P., Landis, E., Lawrence, D., Malhi, Y., Schindler Murray, L., Navarrete, D., Roe, S., Scull, S., Smith, P., Streck, C., Walker, W. S., and Worthington, T.: National mitigation potential from natural climate solutions in the tropics. *Phil. Trans. R. Soc. B*, 375, <https://doi.org/10.1098/rstb.2019.0126>, 2020.
- ~~Haas, O., Prentice, I. C., and Harrison, S. P.: Global environmental controls on wildfire burned area, size, and intensity, *Environ. Res. Lett.*, 17, <https://doi.org/10.1088/1748-9326/ac6a69>, 2022.~~
- Hamilton, D. S., Kelley, D. I., Perron, M. M. G., Lloret, J., Burton, C., Bergas-Masso, E., Liguori-Bills, N., Barkley, A. E., Buchholtz, R., Diez, S., Dintwe, K., Forkel, M., Hall, J., Hantson, S., Hayman, G., Hebden, S., Jones, M. W., Kulkarni, C., Nowell, B., McCarty, J. L., Santín, C., Schneider, S. R., Shuman, J. K., Thoreson, J., Plummer, S., Poulter, B., Vannière, B.: Igniting Progress: Outcomes from the FLARE workshop and three challenges for the future of transdisciplinary fire science, <https://doi.org/10.5281/zenodo.12634068>, 2024.
- Hantson, S., Kelley, D. I., Arneth, A., Harrison, S. P., Archibald, S., Bachelet, D., Forrest, M., Hickler, T., Lasslop, G., Li, F., Mangeon, S., Melton, J. R., Nieradzik, L., Rabin, S. S., Prentice, I. C., Sheehan, T., Sitch, S., Teckentrup, L., Voulgarakis, A., and Yue, C.: Quantitative assessment of fire and vegetation properties in simulations with fire-enabled vegetation models from the Fire Model Intercomparison Project, *Geosci. Model Dev.*, 13, 3299–3318. <https://doi.org/10.5194/gmd-13-3299-2020>, 2020.
- Hodgson, A. K., Morgan, W. T., O’Shea, S., Bauguutte, S., Allan, J. D., Darbyshire, E., Flynn, M. J., Liu, D., Lee, J., Johnson, B., Haywood, J. M., Longo, K. M., Artaxo, P. E., and Coe, H.: Near-field emission profiling of tropical forest and Cerrado fires in Brazil during SAMBBA 2012, *Atmos. Chem. Phys.*, 18, 5619–5638, <https://doi.org/10.5194/acp-18-5619-2018>, 2018.

- Hoffmann, W. A., Jaconis, S. Y., Mckinley, K. L., Geiger, E. L., Gotsch, S. G., and Franco, A. C.: Fuels or microclimate? Understanding the drivers of fire feedbacks at savanna-forest boundaries, *Austral Ecol.*, 37, 634–643, <https://doi.org/10.1111/j.1442-9993.2011.02324.x>, 2012.
- Hofmann, G. S., Cardoso, M. F., Alves, R. J. V., Weber, E. J., Barbosa, A. A., de Toledo, P. M., Pontual, F. B., Salles, L. de O., Hasenack, H., Cordeiro, J. L. P., Aquino, F. E., and de Oliveira, L. F. B.: The Brazilian Cerrado is becoming hotter and drier, *Glob. Change Biol.*, 27, 4060–4073, <https://doi.org/10.1111/gcb.15712>, 2021.
- ~~Houghton, R. A.: Why are estimates of the terrestrial carbon balance so different?, *Glob. Change Biol.*, 9, 500–509, <https://doi.org/10.1046/j.1365-2486.2003.00620.x>, 2003.~~
- ~~Ichoku, C., Giglio, L., Wooster, M. J., and Remer, L. A.: Global characterization of biomass burning patterns using satellite measurements of fire radiative energy, *Remote Sens. Environ.*, 112, 2950–2962, <https://doi.org/10.1016/j.rse.2008.02.009>, 2008.~~
- Intergovernmental Panel on Climate Change (IPCC): Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change. Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, A.L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J.B.R., Maycock, T.K., Waterfield, T. Yelekçi, O., Yu, R., and Zhou, B. (eds.), Cambridge, UK and New York, USA: Cambridge University Press, <https://doi.org/10.1017/9781009157896>, 2021.
- Intergovernmental Panel on Climate Change (IPCC): Impacts, Adaptation, and Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Pörtner, H-O, Roberts, D.C., Tignor, M.M.B., Poloczanska, E., Mintenbeck, K., Alegría, A., Craig, M., Langsdorf, S., Löschke, S., Möller, V., Okem, A., and Rama B. (eds.), Cambridge, UK and New York, USA: Cambridge University Press, <https://doi.org/10.1017/9781009325844>, 2022.
- Kaiser, J. W., Heil, A., Andreae, M. O., Benedetti, A., Chubarova, N., Jones, L., Morcrette, J.-J., Razinger, M., Schultz, M. G., Suttie, M., and van der Werf, G. R: Biomass burning emissions estimated with a global fire assimilation system based on observed fire radiative power, *Biogeosciences*, 9, 527–554, <https://doi.org/10.5194/bg-9-527-2012>, 2012.
- ~~Kelley, D. I., Prentice, I. C., Harrison, S. P., Wang, H., Simard, M., Fisher, J. B., and Willis, K. O.: A comprehensive benchmarking system for evaluating global vegetation models, *Biogeosciences*, 10, 3313–3340, <https://doi.org/10.5194/bg-10-3313-2013>, 2013.~~
- ~~Kelley, D. I., Bistinas, I., Whitley, R., Burton, C., Matthews, T. R., and Dong, N.: How contemporary bioclimatic and human controls change global fire regimes, *Nat. Clim. Change*, 9, 690–696, <https://doi.org/10.1038/s41558-019-0540-7>, 2019.~~
- Ketcham, C. M., and Crawford, J. M.: The impact of review articles. *Lab. Invest.*, 87, 1174–1185, <https://doi.org/10.1038/labinvest.3700688>, 2007.
- Khatun, K., Corbera, E., and Ball, S.: Fire is REDD+: offsetting carbon through early burning activities in south-eastern Tanzania, *Oryx*, 51, 43–52, <https://doi.org/10.1017/S0030605316000090>, 2017.
- Klink, C. A., and Machado, R. B.: Conservation of the Brazilian Cerrado, *Conserv. Biol.*, 19, 707–713, <https://doi.org/10.1111/j.1523-1739.2005.00702.x>, 2005.

- 1400 Klink, C. A., Sato, M. N., Cordeiro, G. G., and Ramos, M. I. M.: The role of vegetation on the dynamics of water and fire in the cerrado ecosystems: Implications for management and conservation. *Plants*, 9, 12, 1–27, <https://doi.org/10.3390/plants9121803>, 2020.
- Kloster, S., and Lasslop, G.: Historical and future fire occurrence (1850 to 2100) simulated in CMIP5 Earth System Models, *Global Planet. Change*, 150, 58–69, <https://doi.org/10.1016/J.GLOPLACHA.2016.12.017>, 2017.
- 1405 Lasslop, G., Coppola, A. I., Voulgarakis, A., Yue, C., and Veraverbeke, S.: Influence of Fire on the Carbon Cycle and Climate, *Curr. Clim. Change Rep.*, 5, 112–123, <https://doi.org/10.1007/s40641-019-00128-9>, 2019.
- ~~Lasslop, G., Hantson, S., Harrison, S. P., Bachelet, D., Burton, C., Forkel, M., Forrest, M., Li, F., Melton, J. R., Yue, C., Archibald, S., Scheiter, S., Armeth, A., Hickler, T., and Sitch, S.: Global ecosystems and fire: Multi-model assessment of fire-induced tree cover and carbon storage reduction, *Glob. Change Biol.*, 26, 5027–5041, <https://doi.org/10.1111/gcb.15160>, 2020.~~
- 1410 Libonati, R., DaCamara, C., Setzer, A., Morelli, F., and Melchiori, A.: An Algorithm for Burned Area Detection in the Brazilian Cerrado Using 4 μm MODIS Imagery, *Remote Sens.*, 7, 15782–15803, <https://doi.org/10.3390/rs71115782>, 2015.
- Libonati, R., Geirinhas, J. L., Silva, P. S., dos Santos, D. M., Rodrigues, J. A., Russo, A., Peres, L. F., Narcizo, L., Gomes, M. E. R., Rodrigues, A. P., DaCamara, C. C., Pereira, J. M. C., and Trigo, R. M.: Drought–heatwave nexus in Brazil and related
- 1415 impacts on health and fires: A comprehensive review. *Ann. N Y Acad Sci.*, 1517, 44–62, <https://doi.org/10.1111/nyas.14887>, 2022.
- Lipsett-Moore, G. J., Wolff, N. H., and Game, E. T.: Emissions mitigation opportunities for savanna countries from early dry season fire management, *Nat. Commun.*, 9, 2247, <https://doi.org/10.1038/s41467-018-04687-7>, 2018.
- ~~Loehman, R. A., Reinhardt, E., and Riley, K. L.: Wildland fire emissions, carbon, and climate: Seeing the forest and the trees –A cross-scale assessment of wildfire and carbon dynamics in fire-prone, forested ecosystems, *For. Ecol. Manag.*, 317, 9–19, <https://doi.org/10.1016/j.foreco.2013.04.014>, 2014.~~
- ~~Mangeon, S., Voulgarakis, A., Gilham, R., Harper, A., Sitch, S., and Folberth, G.: INFERNO: A fire and emissions scheme for the UK Met Office’s Unified Model, *Geosci. Model Dev.*, 9, 2685–2700, <https://doi.org/10.5194/gmd-9-2685-2016>, 2016.~~
- ~~Martin, D. A.: Linking fire and the United Nations Sustainable Development Goals, *Sci. Total Environ.*, 662, 547–558, <https://doi.org/10.1016/j.scitotenv.2018.12.393>, 2019.~~
- 1425 MapBiomas: Collection 8.0 of the Annual Land Use Land Cover Maps of Brazil, <https://plataforma.brasil.mapbiomas.org/>, 2022.
- Mataveli, G. A. V., Silva, M. E. S., França, D. de A., Brunsell, N. A., de Oliveira, G., Cardozo, F. da S., Bertani, G., and Pereira, G.: Characterization and Trends of Fine Particulate Matter (PM_{2.5}) Fire Emissions in the Brazilian Cerrado during
- 1430 2002–2017. *Remote Sens.*, 11, 2254. <https://doi.org/10.3390/rs11192254>, 2019.
- Mataveli, G., Pereira, G., Sanchez, A., de Oliveira, G., Jones, M. W., Freitas, S. R., Aragão, L. E. O. C: Updated Land Use and Land Cover Information Improves Biomass Burning Emission Estimates, *Fire*, 6, 1–15, <https://doi.org/10.3390/fire6110426>, 2023.

- Menezes, L. S., de Oliveira, A. M., Santos, F. L. M., Russo, A., de Souza, R. A. F., Roque, F. O., and Libonati, R.: Lightning patterns in the Pantanal: Untangling natural and anthropogenic-induced wildfires, *Sci. Total Environ.*, 820, <https://doi.org/10.1016/j.scitotenv.2022.153021>, 2022.
- Mistry, J., Schmidt, I. B., Eloy, L., and Bilbao, B.: New perspectives in fire management in South American savannas: The importance of intercultural governance, *Ambio*, 48, 172–179, <https://doi.org/10.1007/s13280-018-1054-7>, 2019.
- Moher, D., Liberati, A., Tetzlaff, J., and Altman, D. G.: Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement, *BMJ*, 339, 332–336, <https://doi.org/10.1136/bmj.b2535>, 2009.
- Moher, D., Shamseer, L., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., Shekelle, P., Stewart, L. A., and Group, P.-P.: Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement, *Syst. Rev.*, 4, 1–9, 2015.
- Moura, L. C., Scariot, A. O., Schmidt, I. B., Beatty, R., and Russell-Smith, J.: The legacy of colonial fire management policies on traditional livelihoods and ecological sustainability in savannas: Impacts, consequences, new directions, *J Environ. Manage.*, 232, 600–606, <https://doi.org/10.1016/j.jenvman.2018.11.057>, 2019.
- ~~Myers, R. L.: Living with Fire Sustaining Ecosystems & Livelihoods Through Integrated Fire Management Global Fire Initiative. <http://nature.org/fire>, 2006.~~
- Oliveira, U., Soares-Filho, B., Bustamante, M., Gomes, L., Ometto, J. P., Rajão, R.: Determinants of Fire Impact in the Brazilian Biomes. *Front. For. Glob. Change*, 1–12, <https://doi.org/10.3389/ffgc.2022.735017>, 2022.
- Oliveira, U., Soares-Filho, B., Costa, W. L. C., Gomes, L., Bustamante, M., and Miranda, H.: Modeling fuel loads dynamics and fire spread probability in the Brazilian Cerrado, *For. Ecol. Manage.*, 482, <https://doi.org/10.1016/j.foreco.2020.118889>, 2021.
- Palacios-Orueta, A., Chuvieco, E., Parra, A., and Carmona-Moreno, C.: Biomass burning emissions: A review of models using remote-sensing data, *Environ. Monit. Assess.*, 104, 189–209, <https://doi.org/10.1007/s10661-005-1611-y>, 2005.
- ~~Pan, X., Ichoku, C., Chin, M., Bian, H., Darmenov, A., Colarco, P., Ellison, L., Kuesera, T., Da Silva, A., Wang, J., Oda, T., and Cui, G.: Six global biomass burning emission datasets: Intercomparison and application in one global aerosol model, *Atmos. Chem. Phys.*, 20, 969–994, <https://doi.org/10.5194/acp-20-969-2020>, 2020.~~
- Pereira, A. M. M., Oliveira, M. da R., Bao, F., Souza, E. B. de, Pott, A., Escobar, A. C. de S., Carvalho, S. S. de, Damasceno-Júnior, G. A.: Changes, trends, and gaps in research dynamics after the megafires in the Pantanal. *Environ. Dev. Sustain.*, <https://doi.org/10.1007/s10668-024-05081-8>, 2024.
- Pereira, G., Freitas, S. R., Moraes, E. C., Ferreira, N. J., Shimabukuro, Y. E., Rao, V. B., Longo, K. M.: Estimating trace gas and aerosol emissions over South America: Relationship between fire radiative energy released and aerosol optical depth observations, *Atmos. Environ.*, 43, 6388–6397, <https://doi.org/10.1016/j.atmosenv.2009.09.013>, 2009.
- Pereira, G., Longo, K. M., Freitas, S. R., Mataveli, G., Oliveira, V. J., Santos, P. R., Rodrigues, L. F., Cardozo, F. S.: Improving the south America wildfires smoke estimates: Integration of polar-orbiting and geostationary satellite fire products in the Brazilian biomass burning emission model (3BEM). *Atmos. Environ.*, 273, <https://doi.org/10.1016/j.atmosenv.2022.118954>,

2022.

- Pereira, G., Siqueira, R., Rosário, N. E., Longo, K. L., Freitas, S. R., Cardozo, F. S., Kaiser, J. W., and Wooster, M. J.:
1470 Assessment of fire emission inventories during the South American Biomass Burning Analysis (SAMBBA) experiment,
Atmos. Chem. Phys., 16, 6961–6975, <https://doi.org/10.5194/acp-16-6961-2016>, 2016.
- Pereira Junior, A. C., Oliveira, S. L. J., Pereira, J. M. C., Turkman, M. A. A.: Modelling Fire Frequency in a Cerrado Savanna
Protected Area, PLoS ONE, 9, e102380. <https://doi.org/10.1371/journal.pone.0102380>, 2014.
- Pivello, V. R.: The use of fire in the cerrado and Amazonian rainforests of Brazil: Past and present, Fire Ecol., 7, 24–39,
1475 <https://doi.org/10.4996/fireecology.0701024>, 2011.
- Pivello, V. R., Vieira, I., Christianini, A. V., Ribeiro, D. B., da Silva Menezes, L., Berlinck, C. N., Melo, F. P. L., Marengo, J.
A., Tornquist, C. G., Tomas, W. M., and Overbeck, G. E.: Understanding Brazil’s catastrophic fires: Causes, consequences
and policy needed to prevent future tragedies, Perspect. Ecol.Conserv., <https://doi.org/10.1016/j.pecon.2021.06.005>, 2021.
- Pope, R. J., Arnold, S. R., Chipperfield, M. P., Reddington, C. L. S., Butt, E. W., Keslake, T. D., Feng, W., Latter, B. G.,
1480 Kerridge, B. J., Siddans, R., Rizzo, L., Artaxo, P., Sadiq, M., Tai, A. P. K.: Substantial Increases in Eastern Amazon and
Cerrado Biomass Burning-Sourced Tropospheric Ozone. Geophys. Res. Lett., 47(3), 1–10,
<https://doi.org/10.1029/2019GL084143>, 2020.
- Prentice, I. C., Farquhar, G. D., Fasham, M. J. R., Goulden, M. L., Heimann, M., Jaramillo, V. J., Kheshgi, H. S., Le Quéré,
C., Scholes, R. J., and Wallace, D. W. R. The carbon cycle and atmospheric carbon dioxide, in: Climate Change 2001: The
1485 Scientific Basis, Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate
Change, edited by: Houghton, J. T., Ding, Y., Griggs, D. J., Noguer, M., Linden, P. J. V. D., Dai, X., Maskell, K., and Johnson
C.A., Cambridge University Press, Cambridge, United Kingdom and New York, USA, 239–287, ISBN: 0521014956, 2001.
- Rabin, S. S., Melton, J. R., Lasslop, G., Bachelet, D., Forrest, M., Hantson, S., Kaplan, J. O., Li, F., Mangeon, S., Ward, D.
S., Yue, C., Arora, V. K., Hickler, T., Kloster, S., Knorr, W., Nieradzick, L., Spessa, A., Folberth, G. A., Sheehan, T.,
1490 Voulgarakis, A., Kelley, D. I., Prentice, I. C., Sitch, S., Harrison, S., and Arneth, A.: The Fire Modeling Intercomparison
Project (FireMIP), phase 1: experimental and analytical protocols with detailed model descriptions, Geosci. Model Dev., 10,
1175–1197, <https://doi.org/10.5194/gmd-10-1175-2017>, 2017.
- Rabin, S. S., Ward, D. S., Malyshev, S. L., Magi, B. I., Shevliakova, E., and Pacala, S. W.: A fire model with distinct crop,
pasture, and non-agricultural burning: Use of new data and a model-fitting algorithm for FINAL.1, Geosci. Model Dev., 11,
1495 815–842, <https://doi.org/10.5194/gmd-11-815-2018>, 2018.
- Ramos Neto, M. B., and Pivello, V. R.: Lightning fires in a Brazilian Savanna National Park: Rethinking management
strategies, Environ. Manag., 26, 675–684, <https://doi.org/10.1007/s002670010124>, 2000.
- Ribeiro, J. F., and Walter, B. M. T.: As principais fitofisionomias do bioma Cerrado, in: Cerrado: Ecologia e flora, 151–212,
2008.
- 1500 Roe, S., Streck, C., Obersteiner, M., Frank, S., Griscom, B., Drouet, L., Fricko, O., Gusti, M., Harris, N., Hasegawa, T.,
Hausfather, Z., Havlík, P., House, J., Nabuurs, G. J., Popp, A., Sánchez, M. J. S., Sanderman, J., Smith, P., Stehfest, E., and

- Lawrence, D.: Contribution of the land sector to a 1.5 °C world. *Nat. Clim. Change*, 9, 11, 817–828, <https://doi.org/10.1038/s41558-019-0591-9>, 2019.
- Rosan, T. M., Klein Goldewijk, K., Ganzenmüller, R., O'Sullivan, M., Pongratz, J., Mercado, L. M., Aragao, L. E. O. C., Heinrich, V., Randow, C. V., Wiltshire, A., Tubiello, F. N., Bastos, A., Friedlingstein, P., and Sitch, S.: A multi-data
1505 assessment of land use and land cover emissions from Brazil during 2000–2019, *Environ. Res. Lett.*, 16, 074004, <https://doi.org/10.1088/1748-9326/ac08c3>, 2021.
- Russell-Smith, J., Cook, G. D., Cooke, P. M., Edwards, A. C., Lendrum, M., Meyer, C. (Mick), and Whitehead, P. J.: Managing fire regimes in north Australian savannas: Applying Aboriginal approaches to contemporary global problems, *Front. Ecol.*
1510 *Environ.*, 11, e55–e63. <https://doi.org/10.1890/120251>, 2013.
- Russell-Smith, J., Murphy, B. P., Meyer, C. P., Cook, G. D., Maier, S., Edwards, A. C., Schatz, J., and Brocklehurst, P.: Improving estimates of savanna burning emissions for greenhouse accounting in northern Australia: limitations, challenges, applications, *Inter. J. Wildland Fire*, 18, 1–18, <https://doi.org/10.1071/WF08009>, 2009.
- Russell-Smith, J., Yates, C., Vernooij, R., Eames, T., van der Werf, G., Ribeiro, N., Edwards, A., Beatty, R., Lekoko, O., Mafoko, J., Monagle, C., and Johnston, S.: Opportunities and challenges for savanna burning emissions abatement in southern
1515 Africa, *J. Environ. Manage.*, 288, 1–17, <https://doi.org/10.1016/j.jenvman.2021.112414>, 2021.
- Santos, F. L. M., Nogueira, J., de Souza, R. A. F., Falleiro, R. M., Schmidt, I. B., and Libonati, R.: Prescribed burning reduces large, high-intensity wildfires and emissions in the Brazilian savanna, *Fire*, 4, 1–21, <https://doi.org/10.3390/fire4030056>, 2021.
- Schmidt, I. B., Moura, L. C., Ferreira, M. C., Eloy, L., Sampaio, A. B., Dias, P. A., and Berlinck, C. N.: Fire management in
1520 the Brazilian savanna: First steps and the way forward, *J. Appl. Ecol.*, 55, <https://doi.org/10.1111/1365-2664.13118>, 2018.
- Schmidt, I. B., and Eloy, L.: Fire regime in the Brazilian Savanna: Recent changes, policy and management, *Flora*, 268, <https://doi.org/10.1016/j.flora.2020.151613>, 2020.
- Silva, P. S., Bastos, A., Libonati, R., Rodrigues, J. A., and DaCamara, C. C.: Impacts of the 1.5 °C global warming target on future burned area in the Brazilian Cerrado, *For. Ecol. Manage.*, 446, 193–203, <https://doi.org/10.1016/j.foreco.2019.05.047>,
1525 2019.
- ~~Silva, P. S., Geirinhas, J. L., Lapere, R., Laura, W., Cassain, D., Alegria, A., and Campbell, J.: Heatwaves and fire in Pantanal: Historical and future perspectives from CORDEX CORE, J. Environ. Manage., 323, <https://doi.org/10.1016/j.jenvman.2022.116193>, 2022.]~~
- Silva, P. S., Nogueira, J., Rodrigues, J. A., Santos, F. L. M., Pereira, J. M. C., DaCamara, C. C., Daldegan, G. A., Pereira, A. A., Peres, L. F., Schmidt, I. B., and Libonati, R.: Putting fire on the map of Brazilian savanna ecoregions, *J. Environ. Manag.*,
1530 296, <https://doi.org/10.1016/J.JENVMAN.2021.113098>, 2021.
- Simon, M. F., Grether, R., de Queiroz, L. P., Skema, C., Pennington, R. T., and Hughes, C. E.: Recent assembly of the Cerrado, a neotropical plant diversity hotspot, by in situ evolution of adaptations to fire, *Proc. Natl. Acad. Sci.*, 106, 20359–20364, <https://doi.org/10.1073/pnas.0903410106>, 2009.

- 1535 United Nations Environment Programme (UNEP): Spreading like Wildfire – The Rising Threat of Extraordinary Landscape Fires. A UNEP Rapid Response Assessment, Sullivan, A., Baker, E., & Kurvits T. (eds.), Nairobi, 2022.
- Van Der Werf, G. R., Randerson, J. T., Giglio, L., Van Leeuwen, T. T., Chen, Y., Rogers, B. M., Mu, M., Van Marle, M. J. E., Morton, D. C., Collatz, G. J., Yokelson, R. J., and Kasibhatla, P. S.: Global fire emissions estimates during 1997-2016, *Earth Syst. Sci. Data*, 9, 697–720, <https://doi.org/10.5194/essd-9-697-2017>, 2017.
- 1540 ~~Vermote, E., Ellicott, E., Dubovik, O., Lapyonok, T., Chin, M., Giglio, L., and Roberts, G. J.: An approach to estimate global biomass burning emissions of organic and black carbon from MODIS fire radiative power, *J. Geophys. Res.*, 114, <https://doi.org/10.1029/2008JD011188>, 2009.~~
- Vernooij, R., Eames, T., Russell-Smith, J., Yates, C., Beatty, R., Evans, J., Edwards, A., Ribeiro, N., Wooster, M., Strydom, T., Giongo, M. V., Borges, M. A., Menezes Costa, M., Barradas, A. C. S., Van Wees, D., Van Der Werf, G. R.: Dynamic
- 1545 savanna burning emission factors based on satellite data using a machine learning approach, *Earth Syst. Dyn.*, 14(5), 1039–1064, <https://doi.org/10.5194/esd-14-1039-2023>, 2023.
- Vernooij, R., Giongo, M., Borges, M. A., Costa, M. M., Barradas, A. C. S., and van der Werf, G. R.: Intraseasonal variability of greenhouse gas emission factors from biomass burning in the Brazilian Cerrado, *Biogeosciences*, 18, 1375–1393, <https://doi.org/10.5194/bg-18-1375-2021>, 2021.
- 1550 Walker, K., Flores-Anderson, A., Villa, L., Griffin, R., Finer, M., and Herndon, K: An analysis of fire dynamics in and around indigenous territories and protected areas in a Brazilian agricultural frontier, *Environ. Res. Lett.*, 17, 8, <https://doi.org/10.1088/1748-9326/ac8237>, 2022.
- Ward, D. E., and Hardy, C. C.: Smoke emissions from wildland fires, *Environ. Int.*, 17, 117–134, [https://doi.org/10.1016/0160-4120\(91\)90095-8](https://doi.org/10.1016/0160-4120(91)90095-8), 1991.
- 1555 Wooster, M. J.: Small-scale experimental testing of fire radiative energy for quantifying mass combusted in natural vegetation fires, *Geophys. Res. Lett.*, 29(21), 21–24, <https://doi.org/10.1029/2002GL015487>, 2002.