

Modeling actinic flux and photolysis frequencies in dense biomass burning plumes

Jan-Lukas Tirpitz¹, Santo Fedele Colosimo¹, Nathaniel Brockway², Robert Spurr³, Matt Christi⁴, Samuel Hall⁵, Kirk Ullmann⁵, Johnathan Hair⁶, Taylor Shingler⁶, Rodney Weber⁷, Jack Dibb⁸, Richard Moore⁶, Elizabeth Wiggins⁶, Vijay Natraj⁹, Nicolas Theys¹⁰, and Jochen Stutz¹

¹Department of Atmospheric and Oceanic Sciences, UCLA, Los Angeles, CA, USA

²University of Alaska, Fairbanks, AK, USA

³RT Solutions Inc., Cambridge, MA, USA

⁴Fort Collins, CO, USA

⁵National Center for Atmospheric Research, Atmospheric Chemistry Observations and Modeling Laboratory, Boulder, Colorado, USA

⁶National Aeronautics and Space Administration, Langley Research Center, Hampton, Virginia, USA

⁷School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA, USA

⁸Earth Systems Research Center, University of New Hampshire, Durham, NH, USA

⁹Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

¹⁰Royal Belgian Institute for Space Aeronomy, Brussels, Belgium

Correspondence: Jan-Lukas Tirpitz (jltirpitz@atmos.ucla.edu)

Abstract. Biomass-burning (BB) affects air quality and climate by releasing large amounts of gaseous and particulate pollutants into the atmosphere. Photochemical processing during daylight transforms these emissions, influencing their overall environmental impact. Accurately quantifying the photochemical drivers, namely actinic flux and photolysis frequencies, is crucial to constrain this chemistry. However, the complex radiative transfer within BB plumes presents a significant challenge for both direct observations and numerical models.

This study introduces an expanded version of the 1D VLIDORT-QS radiative transfer (RT) model, named VLIDORT for PhotoChemistry (VPC). VPC is designed for photochemical and remote sensing applications, particularly in BB plumes and other complex scenarios. To validate VPC and investigate photochemical conditions within BB plumes, the model was used to simulate spatial distributions of actinic fluxes and photolysis frequencies for the Shady wildfire (Idaho, US, 2019), based on plume composition data from the NOAA/NASA FIREX-AQ (Fire Influence on Regional to Global Environments and Air Quality) campaign.

Comparison between modeling results and observations by the UCAR CAFS (Charged-coupled device Actinic Flux Spectroradiometer) yield a modeling accuracy of 10 - 20%. Systematic biases between model and observations are within 2%, indicating that the uncertainties are most likely due to variability in the input data caused by the inhomogeneity of the plume as well as 3D RT effects not captured in the model. Random uncertainties are largest in the ultra-violet (UV) spectral range, where they are dominated by uncertainties in the plume particle size distribution and brown carbon (BrC) absorptive properties.

The modeled actinic fluxes show a decrease from the plume top to bottom of the plume with a strong spectral dependence caused by BrC absorption, which darkens the plume towards shorter wavelengths. In the visible (Vis) spectral range, actinic

fluxes above the plume are enhanced by up to 60%. In contrast, in the UV, actinic fluxes above the plume are not affected or
20 even reduced by up to 10%. Strong reductions exceeding an order of magnitude in and below the plume occur for both spectral
ranges but are more pronounced in the UV.

Copyright statement. TEXT

1 Introduction

Biomass-burning (BB) constitutes a major source of particulate and gaseous atmospheric pollutants with significant impact on
25 air quality and climate (Crutzen and Andreae, 1990; Bond et al., 2013; Klimont et al., 2017). As a consequence of climate
change, natural BB events are expected to increase in frequency and intensity, thereby gaining even further significance in
the coming decades (Jaffe et al., 2020; McClure and Jaffe, 2018; O’Dell et al., 2019). Particles emitted from BB primarily
consist of black carbon (BC) and organic carbon (OC) (Reid et al., 2005; Levin et al., 2010) while emitted gases include
carbon monoxide, nitrogen oxides, a plethora of volatile organic compounds, and greenhouse gases such as CO₂, CH₄ and
30 N₂O (Andreae, 2019). During daytime, both BB particles and gases undergo photochemical processing in the plume, leading
to the removal and transformation of emitted gases, formation of secondary pollutants, and changes in particle properties (Hand
et al., 2010; Forrister et al., 2015; Sumlin et al., 2017; Peng et al., 2021; Xu et al., 2021; Liu et al., 2021; Hennigan et al., 2012;
Cappa et al., 2020; Kiland et al., 2023). The main driver for these processes is photochemistry inside the plume, which is often
optically thick and highly inhomogeneous. Consequently, the accurate quantification of the photochemical drivers is crucial to
35 assess the impact of biomass-burning emissions on the environment.

Understanding and predicting the photochemistry in a given environment, requires knowledge of the speed at which the
involved reactions proceed. For a photochemical reaction



initiated by a photon $h\nu$, the reaction rate

$$40 \frac{d[A]}{dt} = -J[A] \tag{2}$$

is determined by the photolysis frequency J (expressed in Hz), which can be calculated according to

$$J = \int \sigma(\lambda)\Phi(\lambda)F(\lambda)d\lambda. \tag{3}$$

The absorption cross-section $\sigma(\lambda)$ of A describes the probability of photons being absorbed, while the quantum yield $\Phi(\lambda)$
is the probability for absorbed photons to undergo the reaction. $\sigma(\lambda)$ and $\Phi(\lambda)$ are typically known from laboratory measure-
45 ments. The actinic flux $F(\lambda)$ describes the spherically integrated, directionally independent number of photons at wavelength
 λ . $F(\lambda)$ is controlled by environmental factors, such as altitude, solar position, cloudiness, aerosols and Earth surface proper-

ties. All of these conditions are temporally and spatially variable. Consequently, $F(\lambda)$ is highly variable and its determination can be challenging.

Direct measurements of $F(\lambda)$ can be performed with high accuracy using instruments with direction-insensitive reception optics, and spectrometers that cover the required spectral and dynamic range (e.g. Junkermann et al., 1989; Kraus and Hofzumahaus, 1998; Shetter and Müller, 1999; Bais et al., 2003). However, such ~~measurements can only be performed in-situ and are thus~~ measurements are limited in spatio-temporal coverage and are not performed on all aircraft missions studying trace gases or aerosols. Another, often complementary, approach is to simulate actinic fluxes using radiative transport (RT) models. With such models, continuous spatial distributions of actinic fluxes can be derived. RT models need to be constrained with detailed information on the atmospheric conditions, in particular on the spatial distribution and properties of clouds and aerosols. In chemical transport models the inputs for the built-in RT models are driven by the chemical transport simulation itself. For the interpretation of field experiments and observations, the compilation of adequate RT model input data is more challenging.

Over the past 50 years, both direct measurements of actinic fluxes and RT simulations have been applied to obtain a complete picture of the radiative conditions for clear sky conditions (e.g. Turco, 1975; Meier et al., 1997; Shetter et al., 2002; Volz-Thomas et al., 1996; Hofzumahaus et al., 2002; Wagner et al., 2011) and above, below, and inside clouds (e.g. Thompson, 1984; Madronich, 1987; Junkermann, 1994; Kelley et al., 1995; Lefer et al., 2003; Tie et al., 2003; Liu et al., 2006; Neu et al., 2007; Ryu et al., 2017). In contrast, for BB plumes, corresponding studies rely on observations only, whereas the modeling of actinic fluxes and photolysis frequencies and their spatial distribution has received less attention. For instance, many current experimental BB plume chemistry studies are based on directly measured actinic fluxes (Decker et al., 2021; Liao et al., 2021) or irradiances (e.g. Lindsay et al., 2022), or on the deduction of actinic fluxes from concentration measurements of gases involved in well-known proxy reactions (Peng et al., 2021). Such measurements are however limited in spatio-temporal coverage as they can only provide information at a single location and time. For atmospheric chemistry studies in BB plumes, a more complete picture is desirable, as the high spatial variability of actinic fluxes and photolysis frequencies significantly impacts plume processing (e.g. Decker et al., 2021; Palm et al., 2021). Continuous distributions of actinic flux and photolysis frequencies ~~in BB plumes with the~~ with the required accuracy and spatial resolution ~~needed for atmospheric chemistry studies~~ can only be inferred using RT models ~~(e.g. Decker et al., 2021; Palm et al., 2021)~~. Trentmann et al. (2003a; 2003b) performed such RT modeling, but only for a synthetic BB plume, created with a three-dimensional chemical transport model. To our knowledge, similar studies for real plume scenarios do not currently exist. This is not surprising, as modeling is typically challenged by a lack of information to constrain the RT model, arising from the complex and variable nature of such plumes, in particular when they are young (hours old) and optically dense. The RT in BB plumes is dominated by the aerosol, for which the spatial distributions and optical properties are often not well-known. A peculiarity in BB plumes is the optically active fraction of OC, typically referred to as "brown carbon" (BrC), which becomes strongly absorbing towards shorter wavelengths (Laskin et al., 2015). It dominates the RT in the ultra-violet (UV) spectral range, where photochemistry is most responsive. Its concentrations and optical properties are difficult to assess, as they depend on the burned fuel, burning phase and smoke age (Forrister et al., 2015; Laskin et al., 2015; Sumlin et al., 2018; Zeng et al., 2022; Shetty et al., 2023).

In addition, challenges arise from the modeling side. ~~Simulation of horizontal radiative transport effects~~, In the presence of optically thick clouds or plumes, horizontal transport effects such as side illumination and shadowing ~~require~~ can significantly impact the radiation field, particularly when spatial scales on the order of the cloud size or smaller have to be resolved (Stephens and Platt, 1987; Mayer, 2009; Wagner et al., 2023). Considering such effects requires three-dimensional RT models (e.g. Mayer, 2009; Deutschmann et al., 2011), which are computationally expensive. Most models, including the model used in this study, therefore assume a horizontally homogeneous atmosphere; they solve the radiative transfer equation in one dimension and are thus referred to as "1D RT models". This assumption strongly increases efficiency but can introduce errors due to the inability to simulate RT in the horizontal dimension. Investigations of these effects have been performed based on synthetic data (Trentmann, 2003) showing the significance of shadowing. 1D RT models, and to a certain extent 3D-models, often suffer from the incomplete knowledge and simplified description of the input data, particularly in complex environments where atmospheric inhomogeneities can introduce uncertainties in the model initialization. These inhomogeneity effects and challenges with horizontal RT in real BB plume scenarios have not been well studied. There is thus a need to better understand how well 1D RT models can describe actinic fluxes and photolysis frequencies in these plumes.

The accuracy of modeled actinic fluxes and photolysis frequencies can only be ensured by comparison with established models or with measurements. While such comparisons have been conducted extensively for well-defined atmospheric scenarios (e.g. Barnard et al., 2004; Kelley et al., 1995; Castro et al., 1997; Volz-Thomas et al., 1996; Dickerson et al., 1997; Kazantzidis et al., 2001; Vuilleumier et al., 2001; Balis et al., 2002; Shetter et al., 2003; Hofzumahaus et al., 2004), they have not been performed for the complex radiative conditions encountered in optically dense BB plumes.

In the present study we derive continuous spatial distributions of actinic flux and photolysis frequencies in a real and optically dense BB plume, by means of RT simulations. The key objectives of the study are:

1. Introduce and validate a recently developed radiative transfer model (VPC) for photochemical and remote sensing applications in BB plumes.
2. Assess the viability and limitations of actinic flux and photolysis modeling in BB plumes, using 1D RT models and state-of-the-art plume composition measurements.
3. Provide insights into spatial actinic flux and photolysis frequency distributions in BB plumes.

The VPC model (Section 2) is an extended version of the VLIDORT-QS model (Spurr, 2006), a variant of the widely-used VLIDORT RTM. VPC can calculate actinic fluxes based on a range of input parameters. We constrain VPC using recent airborne BB plume composition measurements performed during the FIREX-AQ campaign in 2019 (Section 3), to simulate 2D distributions of actinic flux and photolysis frequencies over 20 plume cross-sections (Section 4). We compare these modeling results to FIREX-AQ actinic flux and photolysis frequency measurements to validate the VPC results (Section 5) and to quantify systematic and random uncertainties. We investigate spectral and environmental dependencies, and perform model sensitivity studies to identify critical factors that limit the modeling accuracy. In addition, we investigate selected model results to provide deeper insight into the spatial and spectral distribution of actinic fluxes and photolysis frequencies in BB plumes (Section 6).

115 Potential implications and future studies in BB plume radiative transport, chemical modeling and remote sensing applications are discussed at the end of this paper (Section 6).

2 Model description

VLIDORT for photochemistry (VPC), is built around the Quasi-Spherical Vector Linearized Discrete Ordinate Radiative Transfer (VLIDORT-QS) code (Section 2.1). The VLIDORT family of RT codes are widely used in the trace gas remote sensing
120 community (Spurr, 2006; Spurr and Christi, 2019). These codes have been tested for many different atmospheric conditions and are thus well-suited for the study and interpretation of field observations. While there are other well-established and efficient RT models for the calculations of photolysis frequencies, such as the Tropospheric UV and Visible (TUV) radiation model ~~fortran code (Madronich and Floecke, 1999)~~ Fortran code (Madronich and Floecke, 1999) or Fast-J (Wild et al., 2000), VPC particularly facilitates the interpretation of field observations of BB plumes. To allow the study of dense BB plumes, VPC has
125 to meet the following requirements: (1) a suitable and flexible representation of BB plume particles in the model, (2) efficient retrieval and sensitivity analyses common in remote sensing studies, (3) accurate RT results for a wide variety of geometries, including high solar zenith angles (SZA) and limb viewing directions, (4) simulation of the light's ~~polarisation~~ polarization state, allowing for retrieval of aerosol information from polarimetric remote sensing data (which will be explored in future studies) and (5) the simultaneous calculation of radiances and actinic fluxes, useful for applications where actinic fluxes are de-
130 duced from radiance observations, or where photochemistry and remote sensing are closely linked (e.g. for plume composition measurements from satellite).




Figure 1. Schematic of the VPC model.

An overview of the model is depicted in Fig. 1. On the input side, VPC features the ~~Aerosol module, which can account for multiple types of aerosol~~ so-called "aerosol module", designed to describe complex aerosol mixtures in a flexible way (Section 2.2.2: the module allows usage of an arbitrary number of aerosol types, each with individual properties that are described either
135 via bulk optical or microphysical parameters (see Section 2.2.2 for details)). On the output side, a module has been added to convert the RT-simulated actinic flux spectra to photolysis frequencies for various chemical reactions (Section 2.4). VLIDORT-QS, the photolysis frequency module and parts of the aerosol module are implemented in Fortran, but have been embedded

in object-oriented Python wrappers, to obtain a structured and modular user interface, including useful functions for data pre- and post-processing as well as visualization. The package supports not only combined use but also stand-alone individual use
140 of the Aerosol, VLIDORT-QS and Photolysis frequency modules. The current model default configuration is optimized for tropospheric applications in the wavelength range between 295 and 650 nm. Extension to stratospheric applications requires adding additional absorbers (e.g. O₂ absorption in the deep UV).

2.1 VLIDORT-QS radiative transport model

The core of VPC is the VLIDORT-QS radiative transfer code, which is designed for a spherically curved atmosphere. Single
145 scattering (or first-order radiative transfer) is treated exactly for a spherically curved medium, while multiple scattering radiation fields along a given line-of-sight are calculated using a modified version of the standard VLIDORT model (Spurr, 2006) working in plane-parallel scattering mode. A detailed description of the VLIDORT-QS model, including validations against 3D Monte-Carlo RT models, can be found in Spurr et al. (2022).

Since the original work on VLIDORT-QS was reported, we have added a number of additional features to the model.
150 Most important for the present study, is the ability to generate directionally integrated fluxes (actinic fluxes and irradiances) at all segment boundaries. The model also features a complete linearization scheme, i.e. in addition to the radiation field itself VLIDORT-QS will generate a complete range of analytically-derived radiance or Stokes-vector Jacobians (weighting or sensitivity functions) with respect to any atmospheric parameter, including aerosol loading and optical properties in BB plumes. This capability makes VLIDORT-QS suitable for retrieval and sensitivity analyses common in remote sensing studies,
155 and adds this ability to observations of actinic fluxes. Finally a number of performance enhancements have been made in order to speed up the calculations; in particular the code has been made thread-safe for use in parallel-computing environments such as OpenMP.

2.2 Model inputs

Table 1 provides an overview on the most relevant model input parameters, which are discussed in more detail in the following
160 subsections.

2.2.1 Vertical grid

In VPC the atmosphere is represented by a finite number of stacked optically homogeneous layers. Common layer numbers are on the order of 100 over an altitude range from zero to about 70 km ASL. Layer thicknesses typically increase with altitude from about 100m at the surface to several km at the top of the atmosphere (e.g. Madronich and Flocke, 1999). Per default,
165 heights h_l for each layer boundary $l = \{0, 1, \dots, l_{\max}\}$ are defined via $h_l = c_1(c_2^l - 1)$, resulting in an exponential grid with increasing layer thickness, but VPC also allows for user defined arbitrary vertical grids.

Table 1. Overview of VPC input parameters. Index l indicates parameters defined individually for each atmospheric model layer. λ indicates dependence on wavelength (see Section 2.5 for further details). Aerosol parameters are defined for each aerosol type $a = \{1, 2, 3, \dots\}$. Parameters with an index $m = \{1, 2, 3\}$ need to be specified for each mode of the trimodal PSD.

Subgroup	Symbol	Description	See also
Microphysical aerosol module	$r_{m,a}$	PSD mode median radii	
	$\sigma_{m,a}$	PSD modal widths	
	$f_{m,a}$	Fraction of particles residing in first two PSD modes ($m = \{1, 2\}$). Third mode fraction $f_3 = 1 - f_1 - f_2$ results from PSD normalisation <u>normalization</u> .	
	$n_a(\lambda)$	Particle real refractive index	
	$\kappa_a(\lambda)$	Particle imaginary refractive index	
Optical aerosol module	$\omega_a(\lambda)$	Aerosol single scattering albedo (SSA)	
	$g_a(\lambda)$	Asymmetry parameter for Henyey-Greenstein scattering phase function	
	$\alpha_{i,a}^{(E)}$	wavelength-dependence coefficients for aerosol extinction coefficient E	Eq. 4
Profile information	h_l	Model vertical grid (altitudes above ground of atmospheric layer boundaries)	
	p_l	Pressure	
	T_l	Temperature	
	$E_{l,a}(\lambda_0)$	Aerosol extinction coefficients at reference wavelength λ_0	
	$c_l^{(s)}$	Number concentration of variable number of trace gases $s = \{\text{O}_3, \text{NO}_2, \text{SO}_2, \dots\}$	
Surface	$A(\lambda)$	Lambertian surface albedo	
Other	λ_w	List of simulated wavelengths	
	$\Delta\lambda_{FWHM}$	Spectral averaging	Section 2.5
		Solar zenith angle	
		Viewing geometry* (viewing elevation, relative azimuth, observer altitude)	

* Of relevance for radiance calculations only.

2.2.2 Aerosol module

In the RT simulation performed with VLIDORT-QS, particles are described by a single set of effective bulk aerosol optical properties, namely the extinction coefficient, single scattering albedo (SSA) and scattering phase matrix. The purpose of the aerosol module is to pre-calculate these quantities for mixtures of aerosol types with different properties. The module allows to include an arbitrary number of aerosol types in the RT simulation. As described in detail in the following paragraphs, for each aerosol type the user chooses between two available aerosol models, ~~provide~~provides aerosol properties as required by the respective model, and defines the aerosol vertical distribution.

The two available aerosol models are in the following referred to as *microphysical* and *optical* aerosol model respectively (Fig. 1). The *microphysical aerosol model* describes aerosol in terms of a particle size distribution (PSD) and complex particle refractive index (real part n and imaginary part κ). These are fed to an integrated Mie aerosol model (Spurr et al., 2012) to derive exact bulk optical properties, including the exact scattering phase matrices, assuming spherical particles. The PSD is tri-modal, with up to three log-normal distributions. The (normalized) number of particles residing in each mode is described by the modal fractions f_m , with $f_1 + f_2 + f_3 = 1$. ~~In the~~ In the current VPC version, the Mie model assumes homogeneous particles, i.e. without coatings. Internally or externally mixed aerosols are realized by defining a single internally mixed aerosol type or multiple externally mixed aerosol types with individual properties, respectively. In the presented study we make use of the latter approach (Section 4). In the future, the microphysical model will be expanded to account for coated or non-spherical particles (see e.g. Kattawar and Hood, 1976; Borghese et al., 1979; Mackowski and Mishchenko, 1996; Muinonen et al., 1996).

For some applications, e.g. remote sensing retrievals with limited information in the measurements, it is useful to stay in the optical domain and skip the additional layer of complexity added by a microphysical aerosol representation. This is the purpose of the *optical aerosol model*, wherein aerosol is described by the wavelength-dependence coefficients $\alpha_{i,a}^{(E)}$ (Eq. 4) and the SSA $\omega_a(\lambda)$. The scattering phase function is defined via the scattering asymmetry parameter (AP) $g_a(\lambda)$, assuming a Henyey-Greenstein formalism (Henyey and Greenstein, 1941).

The properties of each aerosol type are constant with altitude. Altitude dependencies are realized via superposition of multiple aerosol types with different vertical profiles. To simplify the interpretation of field observations in VPC, aerosol amounts are expressed in terms of the aerosol extinction coefficient $E_{l,a}(\lambda_0)$ (extinction per km). Extinction coefficients are often provided by remote sensing observation. They are fed to the model for a specific reference wavelength λ_0 , for each model layer l and each aerosol type a .

2.2.3 Trace gas absorption

For the radiative transport simulation, VPC includes literature cross-sections for O_3 , (Brion et al., 1998) O_4 , (Thalman and Volkamer, 2013), H_2O (Rothman et al., 2010; Lampel et al., 2015) and NO_2 (Vandaele et al., 1998). Users can provide vertical concentration profiles for each of these gases to account for their absorption. VPC also facilitates the implementation of additional gases.

2.2.4 Atmospheric boundaries

For the extraterrestrial solar irradiance at the top of the atmosphere (TOA), VPC users can choose to either use the spectrum by Chance and Kurucz (2010) or by Coddington et al. (2023). Surface properties in VLIDORT-QS can be defined via a Lambertian equivalent reflectance (LER) or providing the full bidirectional reflectance distribution function (BRDF). In the current VPC framework only the wavelength-dependent LER is implemented. If required, BRDF functionality might be added in the future.

2.3 Wavelength-dependent parameters

205 Some model input parameters depend on wavelength, such as refractive indices, albedos, extinction coefficient and asymmetry parameter. In this section we refer to them by $x(\lambda)$. For each wavelength-dependent parameter, the model offers two approaches:

1. Providing independent values for a set of discrete wavelengths.
2. Providing a single value x_0 for a reference wavelength λ_0 and coefficients α_i for an Angstrom wavelength-dependence
210 of arbitrary order i_{max} .

In the second case, parameter values for each wavelength are internally calculated assuming:

$$\ln\left(\frac{x}{x_0}\right) = \sum_{i=1}^{i_{max}} \alpha_i \left(\ln \frac{\lambda_0}{\lambda}\right)^i \quad (4)$$

This ~~parameterisation~~ parameterization represents an extended version of the classic Ångström wavelength-dependence (Ångström, 1929). In fact, for $i_{max} = 1$, Eq. 4 can be rearranged to obtain the well-known dependence:

$$215 \quad \frac{x}{x_0} = \left(\frac{\lambda}{\lambda_0}\right)^{-\alpha_1}, \quad (5)$$

Increasing i_{max} to two, adds a quadratic term to Eq. 4, effectively accounting for a log-linear wavelength-dependence of the Ångström parameter, similar to ~~parameterisations~~ parameterizations proposed by King and Byrne (1976); Kaufman (1993) or Schuster et al. (2006). Increasing i_{max} adds higher order wavelength-dependencies. The ~~parameterisation~~ parameterization was chosen because of its general applicability. We found that Eq. 4 can successfully capture wavelength-dependencies not
220 only of extinction but also other parameters, such as SSA, AP and aerosol refractive indexes measured during FIREX-AQ (see Supplement S1). In contrast to the conventional Ångström dependence, Eq. 4 can also describe the strong non-log-linear wavelength-dependence of BrC absorption in BB plumes.

As ~~decribed~~ described in Section 2.2.2, aerosol vertical profiles are provided in terms of extinction coefficient $E(\lambda_0)$ at a reference wavelength λ_0 . The wavelength-dependence of $E(\lambda)$ is either derived from internal Mie model calculations, when
225 using the microphysical aerosol model, or according to Eq. 4, when using the optical aerosol model. In the case of the SSA ω , Eq. 4 was found to better describe (i.e. with lower i_{max} values) the single scattering co-albedo $1 - \omega$ instead of ω itself. Thus, in the optical aerosol model Eq. 4 is applied to $1 - \omega$.

2.4 Model outputs

The main outputs of a RT simulation with VLIDORT-QS are actinic flux spectra $F_l(\lambda)$ at each layer boundary of the model's
230 vertical grid. Separate spectra are provided for three contributions to $F_l(\lambda)$:

1. Direct solar beam, attenuated by atmospheric extinction

2. Downwelling diffuse radiation, describing the amount of scattered photons incident from the hemisphere above the observer

3. Upwelling diffuse radiation, describing the amount of scattered photons incident from the hemisphere below the observer

235 This separation is useful as it allows for deeper analysis of the results, also considering that real measurements are typically performed separately for upwelling (diffuse only) and downwelling (direct + diffuse) actinic fluxes.

In addition, radiance spectra for a prescribed viewing geometry are calculated in the same simulation process (see Table 1). VLIDORT-QS and the Mie model feature an analytical linearisation scheme (see e.g. Spurr et al., 2022), which can provide Jacobians (sensitivities) of the outputs with respect to any input parameter with high computational efficiency.

240 For the conversion of actinic flux spectra to photolysis frequencies (performed by the "PF-module", see Fig. 1), we isolated and adapted a corresponding module from TUV (version 5.4) (Madronich and Flocke, 1999). It reads and prepares all required cross-sections and quantum yields to calculate photolysis frequencies for 113 photochemical reactions based on Eq. 3. Using this established code as basis ensures consistency with the atmospheric chemistry community. We made a few adaptations: we added missing data on the N_2O_5 absorption cross section for the 340 to 410 nm range, and we adapted ClNO_2 cross section
245 and quantum yield based on the 2015 JPL recommendations (Burkholder et al., 2019). Furthermore, we extended the module by including 14 additional reactions for halogen chemistry.

The TUV module was also used to process the measured actinic flux spectra from FIREX-AQ (see Section 3.3). Note that for the present study, only 48 photolysis reactions relevant to the spectral range of the measurements were considered. An overview of all of the reactions available is provided in Supplement S5.

250 2.5 Spectral averaging and interpolation

RT simulations can be performed for user-defined sets of wavelengths. Ideally, actinic flux spectra are calculated line-by-line, at a resolution resolving even narrow solar Fraunhofer and atmospheric absorption-lines (on the order of few pm in the UV-Vis). However, using very small wavelength intervals is inefficient, ~~while the~~. On the other hand, subsampling the wavelength range decreases the accuracy of the simulation ~~suffers if too few wavelength are chosen~~ (e.g. madronichNumericalIntegrationErrors1990).

255 A number of steps have therefore been taken to make ~~the model~~ VPC flexible and more efficient. The resolution of the originally highly-resolved ($\Delta\lambda \approx 0.01$ nm) literature spectra used by the model (Section 2.2.3 and 2.2.4) can be reduced prior to simulation by Gaussian smoothing. ~~Although this is not physically correct (ideally, smoothing is applied to the line-by-line modeled spectrum after simulation), this~~, i.e. convolution of the spectra with a Gaussian kernel of defined width, typically on the order of 1 nm FWHM (full width at half maximum). This option is useful for efficient calculation of outputs averaged over
260 ~~broader wavelength intervals~~ few nm wavelength intervals, at the cost of relatively small ~~introduceed errors~~ errors, introduced by the commutation of RT modeling and spectral smoothing. We found $< 1\%$ ($< 3\%$) ~~error in~~ errors in the photolysis frequencies for a smoothing kernel of 1 nm (2 nm) FWHM and typical atmospheric scenarios.

To improve efficiency further, a novel spectral interpolation approach was developed; this allows reproduction of high resolution ($\Delta\lambda \approx 1$ nm) actinic flux spectra from simulations at a few suitable wavelengths. Several models use approaches for

265 efficient photolysis rate calculation by reducing the number of full radiative transport calculations to a few wavelengths (<10) (e.g. Landgraf and Crutzen, 1998; Williams et al., 2006; Wild et al., 2000; Madronich and Flocke, 1999). They are however limited in accuracy ($\approx 10\%$ in photolysis frequencies in the UV) and optimized for applications in clean atmospheres. Our approach uses more wavelengths (10 to 30, depending on SZA and atmospheric conditions) and is thus slower but more accurate (< 2% error in photolysis frequencies) and it provides high resolution spectra as an intermediate product. Furthermore, it also
 270 works for the particular conditions encountered in this study, including the presence of the strongly wavelength-dependent BrC absorption and high SZAs.

Our approach is inspired by the passive Differential Optical Absorption Spectroscopy (DOAS) measurement technique (Platt and Stutz, 2008). The passive DOAS approach measures radiances of scattered skylight, but the basic concepts described in the following hold for both radiances and actinic fluxes. Most of the structures observed in an actinic flux spectrum $F(\lambda)$ are
 275 solar Fraunhofer lines already present in the extraterrestrial solar irradiance spectrum I_0 . The actual atmospheric signal can be isolated and conveniently described by the actinic flux optical depth (AFOD)

$$\tau(\lambda) = -\ln\left(\frac{F(\lambda)}{I_0(\lambda)}\right). \quad (6)$$

We make use of the fact that only trace gas absorption introduces significant narrow-band features in $\tau(\lambda)$, whereas scattering and aerosol absorption impose a spectrally smooth signal. Therefore, just as for passive DOAS radiance OD spectra, AFOD
 280 spectra can be approximated by:

$$\tau(\lambda) = \sum_{i=0}^{n_p} P_i \lambda^i + \sum_s S_s(\lambda) \cdot \sigma_s(\lambda, T_s). \quad (7)$$

Here, the first term represents a polynomial that accounts for the spectrally smooth scattering and aerosol absorption signals. The second term accounts for optical depth contributions of each trace gas species s with absorptions strong enough to significantly influence actinic fluxes. ~~In this study and in most other cases, this is only ozone.~~ $\sigma^{(s)}(\lambda, T_s)$ are the trace gas absorption
 285 cross-sections from literature, which can depend on the gas temperature T_s . $S^{(s)}(\lambda)$ is the observed slant column density (SCD), which is the gas' concentration integrated along the effective light path. The term "effective" is used here to indicate that contributing photons travel along an infinite number of light paths, each with a distinct probability. These probabilities change with wavelength, as do the effective lightpath lengths and the SCD. Following Pukite et al. (2010), we ~~parameterise~~
parameterize the SCD wavelength-dependence by

$$290 \quad S(\lambda) = \sum_{j=0}^{n_\lambda} S_{\lambda,j} \lambda^j + \sum_{k=1}^{n_\sigma} S_{\sigma,k} \sigma(\lambda), \quad (8)$$

where we omitted the index s for readability. The first sum is a polynomial, with coefficients S_λ . It describes the SCD including a spectrally smooth wavelength-dependence that arises from changes in RT due to scattering and aerosol absorption. S_σ accounts for changes in effective light path due to strong narrow-band gas absorption, which favours short light path lengths through the absorbing gas.

295 If in our case O_3 is the only absorbing gas of relevance. Then, Eq. 7 and Eq. 8 describe any actinic flux spectrum based on $n_p + n_\lambda + n_\sigma + 1$ free parameters, including T_s . These parameters can be inferred by fitting Eq. 7 to simulated AFODs at a few suitable wavelengths. From the retrieved parameter values, the full high resolution spectrum can be reconstructed.

The optimization of the interpolation settings applied for this study, as well as an accuracy assessment, are described in detail in Supplement S2. In short, we use simulations at 23 wavelengths to reproduce high resolution actinic flux spectra
300 ($\Delta\lambda = 2\text{ nm}$ from 298 to 640 nm). Typical differences to the exact line-by-line calculated spectra were assessed for a wide range of conditions, yielding a root-mean-square deviation (RMSD) of 0.3%. Maximum differences are observed in the UV and for SZAs $> 80^\circ$ but never exceed 7%. Resulting errors in photolysis frequencies are even smaller, with a RMSD of 0.2% and a maximum of 1.8%. We therefore conclude that errors introduced by the interpolation are negligible compared to other sources of uncertainty encountered in the present study.

305 In the future, more investigations, similar to those in Supplement S2, can be performed to further enhance efficiency and make the interpolation generally applicable for other spectral ranges, spectral resolution and conditions.

3 Measurement data

The present study is based on data from the 2019 NOAA/NASA FIREX-AQ measurement campaign (Warneke et al., 2023). In the course of this 6-week airborne campaign, more than 90 BB plumes were sampled in-situ as well as remotely from the
310 NASA DC-8 research aircraft. For our purpose, the FIREX-AQ observations offer a unique opportunity, as they provide a view of plume composition and radiative conditions of unprecedented comprehensiveness and accuracy. An overview of all of the instruments aboard the aircraft and the raw data is available in the NASA FIREX-AQ data archive (Aknan and Chen, 2023). This section provides basic information on the flight selected for our analysis and instrumentation relevant for our study.

3.1 Shady Fire Overview

315 Our case study focuses on measurements from the "Shady Fire" on July 25, 2019 in Idaho ~~because the~~, which we chose for various reasons. Compared to other fires, the data coverage is high. The multiple-hour-long flight included three plume overflights as well as 20 plume transects during daylight and clear sky conditions. Most instruments successfully collected data throughout the entire flight. The burned area over the sampling period was small ($\approx 2\text{ km}^2$) and the burned fuel was homogeneous. The plume was large with an extent of about ~~$10 \times 1 \times 100\text{ km}$~~ $10\text{ km} \times 1\text{ km} \times 100\text{ km}$ ($W \times H \times L$), ensuring
320 good spatial sampling despite the high aircraft speed ($\approx 150\text{ m/s}$) and justifying the 1D model assumption of a ~~horizontal~~ horizontally homogeneous atmosphere. Observed solar azimuth angles (SAAs) were between 250° and 290° , and the Sun is therefore almost aligned with the plume axis (270° azimuthal orientation) over the entire flight (Fig. 2), which is a favourable configuration for avoiding horizontal radiative transport effects (Section 6). ~~Compared to other fires, the data coverage is very high, allowing for a detailed analysis and validation of the VPC model. The multiple-hour long flight included three plume overflights as well as 20 plume transects with most instruments collecting data.~~ Wind conditions were very stable. At the sampling altitude (4200 to 5200 m MSL), transect-averaged wind speed and wind direction over the entire flight were $9 \pm 3\text{ m/s}$
325




Figure 2. Upper panel shows Shady fire geolocation and a part of the flight track (see red box in Fig. 3), including an overflight along the plume axis and following transects. Approximate location and movement of the Sun during the flight are indicated on the left. Lower left shows a Nadir infrared image of the burning area (Aknan and Chen, 2023), recorded from the aircraft at the end of the second overflight.




Figure 3. The Shady fire sampling flight profile. Top panel shows aerosol extinction profiles from the lidar. The flight track is embedded and color coded based on in-situ aerosol extinction measurements. Lower panel shows the lidar aerosol optical depth (AOD) at 532 nm and solar elevation angle. Grey shaded areas indicate the evaluated transects. The red box indicates the data which is visualized on the map in Fig. 2.

and $(270 \pm 12)^\circ$. At the same time, a constantly low relative humidity of $(32 \pm 5)\%$ prevented excessive hygroscopic growth of the aerosol particles. All in all, the Shady Fire represents a particularly favorable case. Even though it might not represent typical conditions, it is an ideal starting point for our purposes, as model validation and error analysis occur in a comparably controlled environment. Future applications of the presented modeling approach to other plumes under less favorable conditions are discussed in Section 7.

Fig. 2 and Fig. 3 provide a general overview of the conditions, fire location and flight path. Plume overflights were performed against the wind along the plume axis and were followed by plume transects ~~with sequentially increasing distance to at progressively increasing distances from~~ the fire.

335 Actinic flux and photolysis simulations were performed for each of the transects labeled in Fig.3. Some transects were excluded due to gaps in crucial measurement data. Data from overflights were used only to infer the aerosol lidar ratio (see Section 4.2).

3.2 Measurements of plume properties

Measurements of plume geometry, composition and aerosol properties were used to constrain the VPC model.

340 A combined High Spectral Resolution Lidar (HSRL) and an ozone Differential Absorption Lidar called DIAL-HSRL (Hair et al., 2008; Browell, 1989) measured aerosol backscatter coefficient at 532 nm at a temporal resolution of 10 s (translating into a horizontal spatial resolution of about 1.5 km) and vertical resolution of 30 m during FIREX-AQ. Measurements were performed simultaneously in the upward and downward directions. Profiles range from near the surface to about 8 km altitude, unless parts of the atmosphere are shielded by optically opaque plume layers. Although the backscatter coefficient is
345 determined from a ratio of two channels with the same viewing geometry, for this system, the profiles contain a vertical gap of about 250 m (see also Figure 3) extending above and below the aircraft due to changes in the geometrical overlap of the telescope and spatially dependent gains of the two photo-detectors that are used for the backscatter coefficient measurements (Wandinger and Ansmann, 2002; Simeonov et al., 1999). The lidar independently provides aerosol extinction profiles, however, at a reduced temporal and vertical resolution (~~60s and 100m~~ 60 s and 100 m) and with an increased geometrical overlap
350 (~~> 2km around the aircraft Hair et al., 2008~~) (> 2 km around the aircraft Hair et al., 2008).

A TSI model 3340 laser aerosol spectrometer measured aerosol size distributions at 1 s temporal resolution, ~~at aircraft cabin temperature and dry humidity. The data are~~ reported for standard temperature and pressure and dry humidity. The measured size range covered particle radii between 50 nm and 2.5 microns. For the flight segments of relevance, the average relative humidity was $(32 \pm 5)\%$. Hygroscopic scattering enhancements for BB aerosol at such humidities are reported to be on the order of 1% and below (Kotchenruther and Hobbs, 1998; Chang et al., 2023). PSD and other particle properties measured under dry conditions are therefore expected to be representative for ambient conditions. TSI Nephelometers, a Radiance Research Particle Soot Absorption Photometer (PSAP) and the NOAA Aerosol Optical Properties Suite (AOP, Langridge et al., 2011; Lack et al., 2012) were used to determine the in-situ aerosol absorption, scattering and extinction coefficients at a temporal
360 resolution of 1 s and various wavelengths. Measurements at 405, 532, and 664 nm wavelength were performed under dry and humidified conditions to determine the hygroscopicity factor, which was then used to scale the measurements to ambient humidity. Measurements at 450, 550, 532 and 700 nm were performed at dry conditions only. A comparison of data for dry and ambient conditions did not indicate hygroscopic growth effects exceeding other measurement uncertainties.

BrC particle absorption between 300 and 700 nm was investigated with a technique in the following referred to as SAEB (spectral analysis of extracted BrC chromophores). For each plume transect particles were gathered on separate filters, which
365 were later extracted with water and afterwards methanol (Liu et al., 2014, 2015; Zeng et al., 2020, 2022). While most organic

material dissolves in either of the solvents, insoluble black carbon remainders can be removed from the solution using pore filters. Spectral analysis of the remaining BrC chromophore solution in a Liquid Waveguide Capillary Cell (LWCC) then provides BrC absorption spectra, which we used to infer the wavelength-dependence of the BrC imaginary refractive index (see Section 4.2).

370 Carbon monoxide (CO) in-situ measurements from the DACOM instrument (Sachse et al., 1987) were used as plume indicator and for interpolation of gaps in aerosol extinction data (Section 4.2). O₃ was measured with a nitric oxide chemiluminescence monitor. The DC-8 aircraft's meteorology and navigation systems provided temperature, pressure, humidity, wind speed, wind direction, geolocation, altitude ASL, ground speed, and radar measured altitude above ground at 1 s temporal resolution.

3.3 Measurements of actinic fluxes and photolysis frequencies

375 For the validation of our modeling results, we use actinic flux spectra measured on the aircraft with Charged-coupled device Actinic Flux Spectroradiometers (CAFS). CAFS instruments measure in-situ down- and up-welling radiation and combine them to provide 4π steradian actinic flux density spectra from 298 to 640 nm. (Shetter and Müller, 1999; Hall et al., 2018). The sampling resolution is ≈ 0.8 nm with a FWHM of 1.7 nm at 297 nm. The absolute spectral sensitivity of the instruments was determined in the laboratory with 1000 W NIST-traceable tungsten-halogen lamps with a wavelength-dependent uncertainty of 3 to 5%. During deployments, spectral sensitivity drift was assessed with secondary calibration lamps while wavelength assignment was monitored with Hg line sources and comparisons to spectral features in the extraterrestrial flux. The optical collectors were characterized for angular and azimuthal response and the effective planar receptor distance. For FIREX-AQ, upgraded electronics and cooling improved the signal to noise allowing for 1 Hz acquisition. From the measured actinic flux, photolysis frequencies are calculated for 48 atmospheric trace gases (listed in Supplement S5), using the same module from the TUV model as VPC (Section 2.4).

380
385

In the following analysis, CAFS actinic fluxes are shown in conjunction with a confidence interval, consisting of two uncertainty contributions: a limit of detection of 6×10^{10} photons $s^{-1} cm^2 nm^{-1}$, estimated from noise analysis under low light conditions, and a span error of 5%, arising from uncertainties in the instrument's calibration.

4 Model setup

390 To initialize our model, the measurements introduced in Section 3.2 were preprocessed and combined with literature and satellite data as described in the following subsections. An overview of the input data is provided in Table 2. We use a vertical grid of 87 layers extending from ground to 62 km above ground level (AGL). The layer thickness was set to 100 m between zero and 6 km (approximately covering the altitude range of the flight), 250 m between 6 and 9 km, 3000 m between 9 and 42 km, and 5000 m between 42 and 62 km altitude AGL. The spectrum by Chance and Kurucz (2010) was used for the extraterrestrial solar irradiance. All literature spectra were spectrally smoothed prior to simulation ($\Delta\lambda = 2$ nm) as described in Section 2.5, which was found to approximately match the resolution of the CAFS instrument. Full simulations were performed on an irregular wavelength grid with 23 nodes (see Supplement S2), with denser sampling towards UV wavelengths. The grid

395

was optimized for the interpolation approach described in Section 2.5 and Supplement S2. The final actinic flux spectra were calculated from these simulations using the aforementioned interpolation approach with a spectral sampling interval of 0.2 nm. For the VLIDORT scattering RT solver, the number of discrete-ordinate streams in the polar half-hemisphere was set to 8. In addition, the "delta-M scaling" ansatz was used to deal with sharply-peaked forward scattering characteristic of aerosols.

Spatial distributions (2D plume cross-sections) of actinic flux and photolysis frequencies are modeled for each of the 20 transects highlighted in the lower panel of Fig. 3. The horizontal resolution of the modeled distributions is determined by the speed of the aircraft and the temporal resolution at which input data is available. As indicated in Table 2, different model input parameters are updated at different ~~intervals~~ temporal intervals, depending on the parameter's data availability, variability and its relevance for the RT:

1. 10 seconds: the temporal resolution of the model simulations is ultimately limited by the resolution of the lidar backscatter profile measurements (10 s). All model input data available at shorter time spans are therefore averaged to at least 10 s intervals prior to simulation, which corresponds to an approximate horizontal resolution of 1.5 km.
2. Per transect: ~~in general BB-plumes exhibit high temporal and spatial variability, but many properties show little (or at least undetectable) variation over individual transects for some parameters (e.g. particle filter measurements) only transect average observations exist. Other parameters (e.g. PSD, particle refractive indices and optical black carbon fraction). Some observations are thus averaged per transect.)~~ appeared to be constant over individual transects within the measurement uncertainty. Those were averaged to reduce measurement noise.
3. Fixed: for some parameters constant values are used for the entire flight, e.g. for those taken from literature or in the case of scarce data coverage.

4.1 Background aerosol

Two aerosol types are used to represent tropospheric and stratospheric background aerosol, both types based on the microphysical model of the VPC aerosol module (Section 2.2.2). For stratospheric background aerosol, extinction profiles were taken from the SAGE II database (SAGE Science Team, 2012). The size distribution was adapted from Wrana et al. (2021). For the particle refractive index we assume a mixture of 75 % H_2SO_4 and 25 % of water (Levoni et al., 1997), yielding a value of $1.43 + i \cdot 10^{-8}$, based on Palmer and Williams (1975) and Segelstein (1981). We use the same value for all wavelengths. For tropospheric background aerosol, the extinction profile is inferred by averaging outside-plume observations of the lidar during the flight. Similarly, outside-plume measurements from the laser aerosol spectrometer were averaged to obtain the size distribution. For the refractive index we use a constant value of $1.53 + i \cdot 0.007$, as reported for clean continental air by Levoni et al. (1997).

Table 2. Overview of the data used to constrain the model for actinic flux and photolysis frequency simulations in the Shady fire plume.

	Parameter	Data source description	Update interval
Tropospheric background aerosol	Size distribution	Laser aerosol spectrometer measurements outside plume	Fixed
	Refractive index	Levoni, 1997	Fixed
	Extinction profile	Lidar extinction profiles outside plume	Fixed
Stratospheric background aerosol	Size distribution	Wrana, 2021	Fixed
	Refractive index	Levoni, 1997	Fixed
	Extinction profile	SAGE II database	Fixed
Brown carbon (BrC) aerosol	Size distribution	Laser aerosol spectrometer	Per transect
	Real refractive index	Sumlin et al. (2018)	Per transect
	Imag. refractive index	Wavelength-dependence from spectroscopy of particle solvent extracts Magnitude retrieved from SSA from Nephelometer, PSAP and AOP	Fixed Per transect
	Extinction profile	Combined lidar backscatter profiles, Nephelometer extinction and DACOM carbon monoxide measurements	10 s
Black carbon (BC) aerosol	Size distribution	Laser aerosol spectrometer	Per transect
	Refractive index	OPAC Hess, 1998	Fixed
	Extinction profile	Same approach as for brown carbon	10 s
Other	Optical black carbon fraction	Retrieved from SSA from Nephelometer, PSAP and AOP	Per transect
	Surface albedo	TROPOMI LER (Tilstra et al., 2023)	10 s
	O ₃ profile	Troposphere: Chemoluminescence monitor Stratosphere: Std. Atmosphere, OMI (Bhartia, 2012) column	Per transect
	Pressure profile	Std. atmosphere, scaled with in-situ	10 s
	Temperature profile	Std. atmosphere, troposphere scaled with in-situ	10 s

4.2 Plume aerosol

In the plume, we consider the two optically dominant particle compounds: black carbon (BC) and brown carbon **BrC** (BrC). BC and BrC are each represented by their own aerosol type in the model. Both types use the microphysical model of the VPC aerosol module (Section 2.2.2), and we assume particles to be spherical and internally homogeneous. The results presented in this study indicate that the plume bulk optical properties can be sufficiently reproduced using this simplified approach, even though real morphology and mixing states of BB particles can be much more complex (see e.g. Hand et al., 2010; Liu et al., 2021). Particle properties are assumed to be altitude-independent, as measurements performed at different altitudes with respect to the plume center did not show significant changes.

435 PSD parameters were inferred for each transect by averaging corresponding laser aerosol spectrometer data and fitting a bimodal log-normal distribution as accepted by the microphysical aerosol model (Section 2.2.2). We assume the same PSD for BrC and BC, since the laser aerosol spectrometer cannot distinguish between the aerosol types.

The refractive index for BC was taken from the OPAC database (Hess et al., 1998). We used a fixed value of $1.7 + i \cdot 0.46$ for all wavelengths. The real part of the BrC refractive index was set to a fixed value of 1.53, following Sumlin et al. (2018).
440 The imaginary refractive index κ_{BrC} was inferred from SAEB measurements. The recorded SAEB spectra (Section 3.2) reflect the wavelength-dependence of the BrC material absorption coefficient α_{BrC} which can be converted to an unscaled imaginary refractive index using the relation $\kappa_{\text{BrC}} \propto \alpha_{\text{BrC}} \cdot \lambda$ (Bohren and Huffman, 1998). With the PSD and refractive index information, we set up the microphysical aerosol model for BC and BrC mixtures to simulate bulk overall SSAs. We fit these SSAs to the ones observed by the Nephelometer, PSAP and AOP measurements to retrieve the BC to BrC ratio and the magnitude of
445 κ_{BrC} for each plume transect. To describe the BC to BrC ratio we introduce the "optical BC fraction" which represents the contribution of BC to the total extinction of BC and BrC at a prescribed reference wavelength λ_0 . It is used below to calculate separate vertical profiles for BrC and BC. Average retrieved values for κ_{BrC} and the optical BC fraction at 300 nm are on the order of 0.05 and 0.08, respectively. A typical BrC imaginary refractive index obtained this way is shown in Fig. S1. The retrieved values reproduce the plume bulk optical properties well (Section 5), but it should be noted that their physical meaning
450 is limited due to the simplifications implicit in the aerosol modeling approach.

Aerosol extinction profiles were inferred from lidar, Nephelometer, and DACOM data. Due to the limitations in the lidar aerosol extinction profiles (Section 3.2) the lidar aerosol backscatter profiles (B) had to be used to achieve suitable spatial resolution and vertical coverage. Conversion to extinction (E) was achieved by investigating aerosol lidar ratios $S = E/B$ observed during plume overflights (see Fig. 3). S was found to depend primarily on the smoke age t (see Fig. S3). This dependence was
455 parameterized by fitting a second order polynomial $S(t)$, used to calculate the extinction profiles $E = S(t_{\text{transect}}) \cdot B$ for each transect. The vertical gap arising from the lidar's non-overlap region around the aircraft was filled using in-situ measured extinction from the Nephelometer and linear interpolation. Smaller temporal gaps in the Nephelometer data were filled using carbon monoxide data from the DACOM instrument, applying a conversion factor inferred from Nephelometer-DACOM-correlations in the same transect (typical Pearson r correlation > 0.99). Missing data above or below the plume was either zero-padded
460 or extrapolated assuming a Gaussian plume shape. Separation into BC and BrC contributions was made based on the current transect's optical BC fraction.

4.3 Surface reflectance

Surface reflectances are taken from the TROPOMI database for monthly Lambertian-equivalent reflectivity (LER, Tilstra et al., 2023). The database has a spatial resolution of $0.125^\circ \times 0.125^\circ$ ($10 \text{ km} \times 14 \text{ km}$ for the Shady fire location), and its 39 spectral
465 channels cover a wavelength range from about 330 to 2300 nm. Below 330 nm, we assume a linear decrease to zero at 250 nm. We consider these approximations as reasonable, since the surface albedo is generally low with minimal impact on the total actinic flux.

4.4 Trace gases

The only trace gas taken into account for our study was O_3 . Based on in-situ measurements performed on the aircraft, we estimated the absorption of other trace gases to be small ($< 3\%$ in actinic flux spectra) and omitted them for simplicity. The O_3 tropospheric profile was inferred from O_3 in-situ measurements on the aircraft during ascent and descent at take-off and landing. For the stratosphere we assumed the 1976 US Standard Atmosphere, scaled such that the O_3 total column matched with OMI satellite observations (Bhartia, 2012).

5 Results

We ran the VPC model for the 20 transects indicated in Fig. 3. Simulation runs were performed at 10 s temporal resolution (corresponding to a horizontal spatial resolution of ≈ 1.5 km) and 100 m vertical resolution at flight altitude, roughly matching the resolution of the lidar observations (Section 3). For each run, the model was constrained as described in Section 4, based on the plume composition and vertical distribution observed at the respective time and location. In total, 350 model runs (10 to 30 runs per transect) were performed. Each run provides vertical profiles for the actinic flux and photolysis frequencies.

For the investigation of photolysis frequencies, we will focus on four important atmospheric reactions with different spectral sensitivity:







Figure 4 illustrates the spectral sensitivity by showing action spectra (actinic flux spectrum \times reactant absorption cross-section \times quantum yield) for each reaction, assuming typical modeled actinic flux spectra with and without the plume present. The action spectra weighted average wavelengths for the four reactions are approximately 310, 360, 375, and 550 nm.

5.1 General features of the BB plume environment

It is instructive to start with a discussion of general features of the BB plume environment. Fig. 5 and Fig. 6 show combined overviews of modeling results and observations for four selected transects (5, 8, 14 and 20) covering different situations. Similar plots for all 20 transects can be found in Supplement S6. To provide a picture over the full wavelength range, Fig. 7 shows measured and modeled spectra for three selected model runs (as highlighted by the black rectangles in Fig. 5) with the aircraft flying within, above and below the plume.

The underlying data for the selected transects were sampled at different times of the flight (compare transect labels in Fig. 3) and at different distances to the fire (20, 40, 100 and 60 km, respectively). As indicated by the 2D distributions of aerosol extinction coefficients (Fig. 5, A-C, and 6, A), the sampled plume cross sections have a horizontal (vertical) extent of




Figure 4. Spectral sensitivity of the four photochemical reactions considered in the comparison. Shown are action spectra (actinic flux \times reactant absorption cross-section \times quantum yield) assuming a typical tropospheric Rayleigh atmosphere actinic flux (dotted lines) or dense plume (AOD \approx 3 at 532 nm) actinic flux spectrum (solid lines).

10 to 15 km (1 to 2 km) and are located about 2 to 3 km above the ground. Plume density and shape vary strongly between different transects. In a young and dense plume, extinction coefficients and AODs of up to 6 km^{-1} and > 3 (both at 532 nm) are encountered, respectively (Fig. 5, A). Around the vertical middle of the plume, this leads to reductions in actinic flux (Fig. 5, G) and photolysis frequencies (Fig. 5, J) of more than an order of magnitude compared to the clean atmosphere outside the plume. The reduction is most pronounced in the UV and UV-centered photolysis frequencies, i.e. the actinic flux at 340 nm is strongly reduced (Fig. 5, G), and is near zero at wavelengths below 340 nm (Fig. 7, A). The aerosol model results identify the increasing BrC absorption as the main reason (Fig. S1), rendering the plume darker and more opaque at the same time. BrC approximately triples plume aerosol extinction and reduces the SSA from 0.9 to 0.75, when comparing 532 to 300 nm. The radiative conditions above, within and below the plume, are very different and, depending on the wavelength, can lead to both enhancement or reduction in actinic flux and photolysis frequencies, in comparison to the situation with a clean atmosphere. During transect 8, the relative height of the aircraft with respect to the plume varies (Fig. 5, B). We see enhancements in actinic flux and photolysis frequencies (Fig. 5, panel H and panel K), particularly towards Vis wavelengths, when the aircraft is above or in the upper region of the plume, and reduction in both UV and Vis when the aircraft moves deeper into or below the plume. This behaviour is also apparent in the example spectra (Fig. 7) as well as the actinic flux vertical profiles (Fig. 5, panels D-F and 6, panel B), and will be investigated in more detail in Section 6.

Transect 14 represents a plume older than >3 h. Older plumes are typically wider, more ragged, and optically thinner (Fig. 5, panel B, panel C). Accordingly, the reductions in actinic flux and photolysis frequencies are less pronounced (Fig. 5, panel I, panel L).

5.2 Comparison of model and measurement

In this section we compare the modeled actinic fluxes and photolysis frequencies to the ones observed by the CAFS. The comparison serves to validate the VPC model and to assess the accuracy of modeling results. While the model provides full vertical profiles of actinic flux and photolysis frequencies, the comparison is limited to a single altitude per model run, namely the altitude of the aircraft, where the respective CAFS measurement was performed. However, the dataset includes




Figure 5. Overview of modeled and observed (CAFS) actinic fluxes and photolysis frequencies for three example transects. The first and second rows show plume cross-sections of measured aerosol extinction (532nm) and modeled actinic flux, respectively. The observer is looking downwind. Solar relative azimuth angles (RAA) are given with respect to the viewing direction: 0° , 90° or 180° indicate sun in front, to the right or in the back of observer, respectively. Altitudes are given with respect to mean sea level (MSL). The third row shows time series of in-situ actinic fluxes at three wavelengths, normalized to TOA irradiance. Black rectangles indicate those observations discussed in more detail throughout the paper (see also Fig. 13, Fig. 7, and Fig. 12). The fourth row shows in-situ photolysis frequencies for three example reactions. For reference, dashed lines indicate Rayleigh atmosphere actinic fluxes and photolysis frequencies, created by re-running the model without the plume present.

measurements at different relative altitudes with respect to the plume. We therefore expect the average agreement of model and measurements to be representative for the entire modeled distributions.

5.2.1 Comparison of actinic fluxes

The qualitative agreement of model and measurement can already be seen in the combined plots of actinic flux time series in Fig. 5, panels G-I and 6, panel C. The model reproduces major reductions and enhancements in the time series very well. Occasional outliers typically occur at the plume edges, particularly late in the evening (see highlighted model runs in Fig. 6, panel C), and are discussed later (Section 6).




Figure 7. Modeled and measured (CAFS) actinic flux spectra for three example observations (as highlighted in Fig. 5). To put the flux magnitudes into perspective, "Rayleigh atmosphere" spectra were calculated by re-running the model without plume aerosols. Lower panels show relative differences between model and measurements. The modeled spectra were interpolated between simulation nodes following the procedure described in Section 2.5. Model errors were estimated using the sensitivity studies described in Section 5.2.3 and do not consider errors from horizontal radiative transport effects.

errors increasing with aerosol signal (e.g. errors in the aerosol representation or properties) and twilight conditions. This enables a more direct analysis of the RT effects inside the plume.




Figure 8. Correlation plots of measured (CAFS) and modeled (VPC) actinic flux optical density (AFOD, see Eq. 6). Data points in the plot are limited to the same wavelengths (340, 440 and 550 nm) as shown in Fig. 5. For the difference histogram and linear regression results on the right, the whole dataset was considered.

540 To quantify the overall agreement of model and observations, we performed a correlation analysis on the entire dataset (Fig. 8), including all 20 transects, all wavelengths, and also locations outside the plume. CAFS observations - with an original




Figure 9. Agreement of modeled and measured actinic flux optical density (AFOD) in dependence of wavelength and plume total AOD. AODs were calculated from lidar observations at 532nm. Boxes span 25st to 75st percentile. Whiskers span 10th to 90th percentile. Grey shaded areas show average CAFS measurement uncertainty.

temporal resolution of 1 s - were averaged over each of the modeled 10 s time intervals. We applied two filters: (1) we only considered data for which the CAFS signal-to-noise ratio is higher than 5 (removes < 2% of the data), and (2) we subsequently ignored the highest first percentile of differences to remove the most severe outliers, which are mostly associated with inhomogeneities in the plume not captured by the model. Based on this data, we obtain a Pearson correlation coefficient of 0.98 between model and measurements. The RMSD in AFOD is 0.17, which corresponds to a 17% deviation in the actinic flux. Slope (0.98) and intercept (-0.004) from the linear regression analysis indicate no significant systematic differences. As described in Section 2.4 and 3.3, both CAFS and VPC provide upwelling and downwelling actinic flux contributions separately. Correlation analysis results for each of the contributions are provided in Supplement S4. The upwelling part exhibits larger deviations (RMSD of 24%) than those for the downwelling part (RMSD of 17%), which is most likely due to imperfect representation of surface properties, surface illumination (e.g. shadowing by the plume) and the assumption of constant ground elevation in the model. However, the impact of this increased deviation is limited, since in the presented data, upwelling radiation contributes on average only $\approx 20\%$ to the total actinic flux.

To obtain a more differentiated picture, we investigated the dependence of model-measurement differences on wavelength, plume AOD and solar geometry by binning of the data (Fig. 9 and 10). For the AOD binning, we used the AODs at 532 nm observed by the lidar. Overall, we found this AOD to be a good proxy to identify those observations significantly affected by the presence of the plume, even though the impact of the plume signal also depends on the aircraft's altitude with respect to the plume. We find a clear increase in model-measurement difference towards short wavelengths (Fig. 9, A and also Fig. 7) and high AODs (Fig. 9, B). For wavelengths > 400 nm and low AODs, RMSDs of 0.05 to 0.1 are observed. In contrast, for wavelengths around ≈ 300 nm and high AODs (≈ 4 at 532 nm and ≈ 10 at 300 nm), RMSDs increase up to 0.4 (Fig. 9, C). At high AODs, Fig. 9 B indicates a general systematic underestimation. Only observations above and in the upper plume contribute to this data bin, since observations below do not meet the CAFS signal-to-noise filtering criterion mentioned before.

For intermediate AODs ($1 < \text{AOD} < 3$ at 532nm), including all wavelengths, the RMSD is 0.18. We consider this value to be representative for the typical in-plume model accuracy.

565 For the dependence on the solar geometry, we chose three segments of the flight (transects 1, 2 and 3; transects 6, 7 and 8; transects 18, 19, 20) with different solar geometries but otherwise similar conditions (Fig. 10, A). During the first and third segment, the Sun illuminates the plume from the side (albeit at a small relative azimuth angle of $\approx 15^\circ$) with respect to the plume axis), at low ($\approx 50^\circ$) and high SZA ($\approx 80^\circ$), respectively. During the second segment, the sun is approximately aligned with the plume axis at a moderate SZA ($\approx 60^\circ$). During side illumination, in particular at high SZAs, model-measurement differences increase and model results underestimate actinic fluxes (Fig. 10, B). Compared to the second segment, the RMSD for the third segment almost doubles (from 0.13 to 0.24). As discussed in detail in Section 6, this might be caused by the 1D model assumption of a horizontally homogeneous atmosphere.




Figure 10. Agreement of model and measurement for three segments of the measurement flight with different solar geometries, as illustrated in Panel A. RAAs were calculated with respect to the plume axis, i. e. the prevalent wind direction. Boxes span 25th to 75th percentile. Whiskers span 10th to 90th percentile. Grey shaded areas show average CAFS measurement uncertainty.




Figure 11. Agreement of modeled and measured photolysis frequencies for the four example reactions with different spectral sensitivity. Instead of root-mean-squares of absolute differences, we show root-mean-square values of relative differences here (RMSRD), for easier comparison of different reactions.

5.2.2 Comparison of photolysis frequencies

From a chemical point of view, it is interesting to assess how the model-measurement differences in actinic fluxes propagate into the photolysis frequencies. We therefore performed correlation and regression analyses similar to those in Fig. 8 for the modeled and measured photolysis frequencies of the 48 tropospheric photochemical reactions already mentioned in Section 2.4. Again, the first percentile of data with highest differences was omitted. Results for the four reactions introduced at the beginning of Section 5 are shown in Fig. 11. The results for the remaining reactions are listed in Supplement S2. Note that, instead of the root-mean-square of absolute deviations (RMSDs), we calculate the root-mean-square of relative deviations (RMSRD) here, according to:

$$\text{RMSRD} = \sqrt{\frac{1}{N} \sum_{i=1}^N \left(\frac{y_i - x_i}{x_i} \right)^2} \quad (9)$$

with x_i and y_i being measured and observed photolysis frequencies, respectively. This facilitates the comparison between different reactions.

As mentioned before, CAFS and VPC use the same photolysis frequency code, quantum yield and absorption cross-section data. The conversion from actinic fluxes to photolysis frequencies therefore does not introduce additional differences between model and measurement; photolysis frequencies simply reflect the actinic flux differences, weighted by the action spectrum of the corresponding reaction (Fig. 4). Accordingly, systematic differences (slope and offset of the linear regressions) and RMSRDs for the photolysis frequencies are similar to those obtained for actinic fluxes and follow the patterns already seen in panel A of Fig. 9. Larger differences are found for UV-driven reactions like the O_3 photolysis (Fig. 11, A). Smaller differences are observed for Vis-driven reactions, especially the NO_3 photolysis (Fig. 11, D), which also benefits from integration over a broad spectral range (see action spectrum in Fig. 4). For all 48 reactions, Pearson correlation coefficients are > 0.96 , slopes

are between 0.96 and 1.04 and intercepts range between -5% and $+2\%$ of the average observed photolysis frequency for the respective reaction (Supplement S2). RMSRDs are between 0.12 and 0.21, with an average of 0.15.

5.2.3 Model sensitivity to input parameters

595 To better understand the origin of the model-measurement differences and identify critical factors, we performed model sensitivity studies, i.e. we investigated the response of the modeling results to variations in the input parameters.

Table 3. Parameter variation magnitudes

Group	Parameter	Variation
PSD	Fine mode median radius r_1	10%
	Coarse mode median radius r_2	10%
	Fine mode width σ_1	0.1
	Coarse mode width σ_2	0.1
	Modal fraction f	0.1
Refractive index	Real part n	0.1
	Imaginary part κ	20%
Other	Ozone column	5%
	Surface LER	100%
	Aerosol extinction	10%
	Temperature	5%
	Pressure	2%
	Optical BC fraction	0.05

The variations applied to the model inputs are listed in Table 3. Their magnitudes correspond to the approximate uncertainties in the respective parameter. Variations of PSD parameters, pressure and temperature were estimated based on specifications of the corresponding FIREX-AQ instruments, but also considering noise, variability and limited coverage of the measurements. 600 Refractive index and optical BC fraction variations are based on the variability reported in the literature (Sarpong et al., 2020; Lack et al., 2012; Andreae and Gelencser, 2006) and from uncertainties in SSA measurements, propagated through the Mie model fit described in Section 4.2. O₃ column and surface LER variations are based on reported uncertainties in the corresponding satellite measurements (Shavrina et al., 2007; Balis et al., 2007; Kroon et al., 2008; Frith et al., 2020; Tilstra et al., 2023). Aerosol extinction uncertainty was assumed to be dominated by uncertainties in the lidar ratio, which we 605 calculated from the scattering in Fig. S3. The listed variations were applied to the three simulated cases highlighted in Fig. 5 and shown in Fig. 7.

The resulting differences in the modeling results (Fig. 12), including their spectral dependencies, exhibit very similar magnitudes and patterns as those differences between model and measurements reported in Section 5.2.1. Most parameters induce a




Figure 12. Response of actinic flux spectra (panel A, B and C) and photolysis frequencies (panel D, E and F) to the parameter variations defined in Table 3, for the three example spectra presented in Fig. 7. Parameters have been grouped here for readability by quadratically adding their propagated impact. Legend in panel A applies to the entire figure.

spectrally smooth difference in the actinic flux, which increases significantly towards the UV (Fig. 12, A and B). Furthermore, the differences increase with the plume AOD, i.e. they are largest (with a total of up to 60% in the UV) inside the optically dense plume of transect 5 (Fig. 12, A), and become smaller (up to 40%) for the thinner plume encountered during transect 8 (Fig. 12, C). Smallest differences are found above the plume (Fig. 12, B). Here, the actinic flux is dominated by the contribution from direct solar radiation (see also Fig. 13), which is not affected by plume aerosols, but prone to uncertainties in the stratospheric O₃ column below 315 nm.

In and below the plume, uncertainties in the UV are dominated by particle properties such as refractive indices and the PSD. Investigating the contributions of separate parameters and aerosol types (not shown in Fig. 12) provides further insights: as expected from the generally low optical BC fraction (10% at around 500 nm) and the strong increase of BrC absorption towards the UV, these uncertainties are dominated by the BrC properties, while BC parameter variations contribute less than 20% to the total uncertainty below 350 nm.

In the Vis, uncertainties are generally lower than those in the UV, since plume optical thickness decreases with wavelength. The uncertainty is dominated by the optical BC fraction, which is approximately proportional to the plume extinction. The contribution from BrC becomes negligible towards longer wavelengths.

6 Discussion

The major features in the observed actinic flux and photolysis frequency time series (Fig. 5 and 6), as well as in the actinic flux spectra (Fig. 7), are well reproduced by the model. RMSDs over the entire dataset are on the order of 10% to 20%, in both actinic fluxes and photolysis frequencies. Systematic differences in our comparison are on the order of a few percent, as

illustrated by the regression slope of 0.98 and negligible intercept between measured and modeled actinic fluxes (Fig. 8) and slopes between 0.99 to 1.02 in the measured and modeled photolysis rates (Fig. 11). These differences are smaller than the uncertainty in the CAFS measurements (Fig. 8, 9, 11).

630 Generally we find a surprisingly good model-measurement agreement, considering the complexity, heterogeneity and variability of BB plumes and the large variations in actinic flux and photolysis frequencies over more than an order of magnitude on short spatial and temporal scales. To put our results into perspective: similar airborne (Kelley et al., 1995; Volz-Thomas et al., 1996) and ground-based (Barnard et al., 2004; Castro et al., 1997; Dickerson et al., 1997; Kazantzidis et al., 2001; Vuilleumier et al., 2001; Balis et al., 2002; Shetter et al., 2003; Hofzumahaus et al., 2004) model-measurement comparisons
635 for less polluted and nearly homogeneous clear-sky atmospheres typically achieve agreements on the order of 10 %.

Besides the relatively small CAFS measurement error, model-measurement differences arise for two major reasons: Errors in the model input parameters and simplifications taken in the modeling approach.

Errors in model input parameters partly arise from instrumental measurement errors and uncertainties in the literature data that were used to constrain the model (Section 4). Furthermore, the limited spatio-temporal coverage and resolution of this data
640 does not fully encompass real variations in the inhomogeneous plume, thereby adding considerable uncertainty to the model constraints. Non-linear relations between plume properties and actinic fluxes might also hinder exact simulations when using averaged parameters.

Model simplifications are the representation of aerosol (as spherical particulates and externally mixed) and the application of a 1D model, which cannot account for horizontal RT effects. The latter are caused by the horizontal inhomogeneities of
645 the scenario. Such effects comprise for instance side-illumination and shadowing, especially at the edges of the plumes. But also on smaller scales and within the plume, horizontal RT can affect the results. For instance, the fact that the lidar aerosol extinction profile was measured in zenith direction while the RT calculation were performed for non-zero SZA can introduce inconsistencies in the aerosol extinction profiles.

For our sensitivity studies (Section 5.2.3), we estimated the uncertainties in the model input parameters, considering both
650 instrumental errors and uncertainties due to limited coverage and resolution of the measurements. Propagating these uncertainties through the model, yields model uncertainties very similar to the observed model-measurement differences, not only in magnitude but also in the wavelength and AOD dependence (compare Fig. 9 and 12). We conclude that, at least as average behavior over the dataset, the model-measurement differences are dominated by uncertainties in the model input. Hence, even with comprehensive state-of-the-art measurements as performed during FIREX-AQ, the information on the plume environment
655 is still the limiting factor for accurate modeling of actinic fluxes and photolysis frequencies in BB plumes. Most critical is the uncertainty in brown carbon aerosol properties (refractive indices and PSD), which dominates the model uncertainties in the UV (Fig. 12), where photochemistry is most sensitive. BrC properties are difficult to assess as they are highly variable and the mechanisms affecting the absorption's spectral dependence are not yet fully understood (e.g. Laskin et al., 2015; Shetty et al., 2023). A recent comparison study revealed inconsistencies between different approaches to derive BrC optical properties from
660 FIREX-AQ data (Zeng et al., 2022), presumably in part due to insoluble BrC components that the SAEB technique (Section 3) cannot detect (Liu et al., 2013; Shetty et al., 2019; Chakrabarty et al., 2023). In our study, this effect is likely reduced,

since we combine the SAEB observations with direct in-situ optical measurements of the SSA (Section 4.2). However, the absence of SSA observations in the UV and the inhomogeneity of the plume hinders an accurate consideration of BrC in the modeling process. We propose that further investigation of the discrepancies between in-situ extractive BrC sampling and remotely-sensed actinic flux or radiance spectra may shed light on this issue and ultimately better constrain radiative transport and photochemistry in BB plumes.

Considering that most of the model-measurement differences can be explained by uncertainties in the model input, simplifications in the model are unlikely to limit the modeling accuracy under typical conditions. However, horizontal radiative transport effects might become problematic for specific cases. These effects - in particular side-illumination and shadowing - are expected to be most pronounced when the SAA is not aligned with the plume axis, i.e. the Sun illuminates the plume from the side (first and third sketch in Fig. 10, A) and when SZAs are large. Indeed, we observe a dependence in model-measurement differences, matching these expectations (Fig. 10, B). Furthermore, horizontal radiative transport effects are likely to occur at the plume edges, where horizontal inhomogeneity is large. An extreme case, where all three conditions are fulfilled, are the plume edges of transect 20. The actinic flux time series indicates strong under- and over-estimations (black boxes in Fig. 6, C), which can be explained by plume side-illumination and shadowing, respectively. It should be noted that the Shady Fire dataset is favourable in this context, since the sun is almost aligned with the plume axis over the entire flight (Section 3). Plumes under less favourable geometries might lead to larger horizontal radiative transport effects. On the other hand, corrections, for example based on additional measurements of direct solar AOD, might reduce these effects in the future (Várnai and Davies, 1999).

With the confidence that we can model actinic fluxes and photolysis frequencies accurately, it is worth discussing the modeled spatial distributions of actinic fluxes and photolysis frequencies. Figure 13 shows actinic flux vertical profiles for Case 1 during transect 5 (black box in Fig. 5, panel G). We separated contributions from direct, diffuse downwelling and diffuse upwelling flux and investigated two wavelengths close to the lower and upper end of the photochemically relevant wavelength range (320 and 600 nm, in panel A and panel B, respectively). For the wavelengths in between these values, the profiles were found to transition steadily into each other.

Below the plume, the contribution from direct sunlight is considerably reduced in the UV and Vis, leading to a net reduction in the actinic fluxes, despite the enhancement of the diffusive radiation fields (e.g. the Vis downwelling contribution in Fig. 13, panel B). Above the plume, the behaviour is very different for UV and Vis. In the Vis (Fig. 13, panel B), more light is scattered upwards, as the plume albedo is larger than the typically dark Earth surface ($LER \lesssim 10\%$). This leads to an enhancement of the Vis diffuse upwelling flux on the order of 20% for the example in Fig. 13, panel B, and up to 60% over other parts of the Shady fire. The enhancement decreases only slightly with altitude for the infinitely horizontally extended plume in our 1D model. This behaviour is very similar to the one reported for cloud layers, which are known to have substantially ($> 100\%$) increased (decreased) actinic fluxes and photolysis frequencies above (below) the clouds (e.g. van Weele and Duhnkerke, 1993; Lefer et al., 2003).

This behavior changes significantly in the UV. While upwelling radiation from clouds is similar in the UV and Vis spectral range, actinic fluxes above BB plumes remain similar to those in a pure Rayleigh atmosphere. This is due to increased darkening




Figure 13. Detailed plot of the simulated actinic flux vertical profile, for the case 1 in transect 5 (black box in Fig. 5, panel G), in the UV (left) and Vis (right). Colored areas indicate separate contributions from direct as well as diffuse upwelling and downwelling radiation. The thin black line indicates the plume extinction profile in arbitrary units. Note the different altitude scales of upper and lower panels. To put fluxes into perspective, dashed lines show results for a Rayleigh atmosphere without plume.

of the plume by BrC absorption. For wavelengths below 320nm, actinic fluxes above the plume are even reduced by a few percent (light gray shading above the plume in Fig. 14, panel B), indicating that the plume appears even darker than the combined reflectance of surface and atmosphere in a clean atmosphere. Within and below the plume the shape of the profile is similar for UV and Vis but reductions are much stronger in the UV.

For chemical studies, the spatial distribution of photolysis frequencies in the plume is of most interest. Figure 14 shows such distributions, as obtained from the VPC model based on data from transect 5. The distributions mostly reflect the findings on the actinic flux vertical profiles. Photolysis frequencies for UV-driven reactions such as the O_3 photolysis (Fig. 14, B) remain unchanged (or are slightly lower) above the plume, and drop to near-zero towards the center or bottom of the plume. For Vis-driven reactions like the NO_3 photolysis (Fig. 14, D), photolysis frequencies are enhanced by up to 40% above the plume, while the reduction within and below the plume by up to 60% is much weaker than that in the UV. The ultimate impact of these spatial and spectral features on the plume processing should be investigated with high resolution chemical models in the future.




Figure 14. 2D distributions of three photolysis frequencies (Panel B,C,D) for the transect 5 plume cross-section. To illustrate the situation, panel A shows the plume shape in terms of aerosol extinction (same as Fig. 5, panel A). "Enhancement factors" represent the ratio of modeled photolysis frequencies with and without plume.

7 Conclusions

710 We have introduced and validated VPC, a VLIDORT-based quasi-spherical 1D RT model ~~optimized for RT and remote sensing applications in BB plumes. The model, which is constrained,~~ VPC can calculate radiances, radiative fluxes and photolysis frequencies for a wide range of atmospheric conditions, including high loads of complex aerosol mixtures as they occur in BB or other plumes. VPC also efficiently calculates Jacobians of the simulated quantities with respect to the input parameters, facilitating its use as a forward model in remote sensing retrievals or similar inversion problems.

715 We have constrained the model by a comprehensive set of aerosol measurements performed during FIREX-AQ ~~, calculates actinic flux and calculated actinic fluxes~~ and photolysis frequencies in BB plumes with an accuracy of 10 – 20% compared to direct measurements.

Previous model-measurement comparisons in clean atmospheres have found differences of < 10%. Considering the highly complex and inhomogeneous RT environment in dense BB plumes, the agreement between VPC and the observations is remarkably good. The average difference of actinic fluxes over the entire dataset is 17%, with larger values ($\approx 40\%$) at $\approx 300\text{nm}$ and for large plume AODs (≈ 10 at 300nm). For the resulting photolysis frequencies, RMSRDs range between 13% and 21%, with lower (higher) errors for Vis- (UV-) driven reactions. A further analysis of the comparison results show very small systematic differences on the order of $\approx \pm 2\%$ between the model and the observations. Our sensitivity studies suggest that most of the RMSD can be explained by the uncertainties in the model input data, mostly arising from the limited

725 spatio-temporal coverage and resolution of the corresponding measurements. In the UV, the model error is dominated by the uncertainty in the properties of the strongly absorbing BrC. These uncertainties can stem from the variability of refractive

indexes as well as size distribution inside the plume. In the Vis, the amount and properties of black carbon dominate the model error. Due to the decrease of both strong BrC absorption and plume optical density towards the Vis, the model error in the Vis is lower by about a factor of 3 compared to the error in the UV. Given these findings, and considering that photochemistry is particularly sensitive to the UV spectral range, more research on BrC optical properties in BB plumes is needed to better constrain radiative transport and photochemistry in BB plumes.

The model-measurement comparison also provides insights into the potential limitations of the 1D approximation in the RT simulations. We observe a systematic increase in the model-measurement difference by a factor of two for solar geometries with high SZA and when the SAA is not aligned with the plume direction. In addition, occasional outliers at the plume edges, where horizontal gradients of environmental parameters are largest, can be identified. Both these considerations indicate that, under certain conditions, 3D RT effects can have significant impact.

~~Previous model-measurement comparisons in clean atmospheres have found differences of $< 10\%$. Considering the highly complex and inhomogeneous RT environment in dense BB plumes, the agreement between VPC and the observations is remarkably good.~~

The model results also ~~provided~~ provide insights on how photochemistry is affected within and in proximity of BB plumes. Despite the absorption of BC, BB plumes appear bright in the nadir at visible wavelength compared to the typical Earth surface. Accordingly, the additional upward reflected light leads to enhancements of actinic fluxes in the Vis above the BB plume of up to $\approx 60\%$ compared to fluxes in a Rayleigh atmosphere. This enhancement can reach the upper troposphere and the stratosphere, and resembles the impact of clouds on actinic fluxes, reported in previous studies. However, the increase in BrC absorption significantly darkens the BB plume towards the UV. Consequently, the observed above-plume enhancement gradually disappears towards shorter wavelengths and actinic fluxes can even be reduced compared to fluxes in a Rayleigh atmosphere for wavelengths $\lesssim 310$ nm. Inside the plume, actinic fluxes steadily decrease from top to bottom and remain approximately constant in the atmosphere below. The decrease is more pronounced in the UV than in the Vis.

Our results show ~~very~~ good agreement between observations and measurements in the Shady fire. However, our results also highlight the challenges in describing actinic fluxes in complex dense BB plumes. The Shady fire ~~was~~ has been selected for its favorable conditions, such as the large plume size, clear sky conditions ~~during the flight,~~ comprehensive sampling, and the alignment of the sun with the plume axis ~~which reduce plume inhomogeneities and 3D radiative transport effects~~. An expansion of our analysis to ~~the entire other~~ FIREX-AQ dataset of over 90 fires plumes would allow assessment of actinic fluxes and RT modeling challenges in a wider variety of ~~biomass burning (BB) plumes.~~ BB plumes. From the 90 fires sampled during FIREX-AQ, about five other fires provide similarly favorable conditions. Prevailing challenges for the study of other fires are scarce sampling and the presence of clouds. Identifying the most critical unknowns from additional sensitivity studies might be of great value in the future to constrain actinic fluxes in and around BB plumes with less measurements and efforts.

The availability of a linearized RT model, such as VLIDORT-QS, together with our sensitivity results also open the opportunity to use actinic flux and other remote sensing observations to study aerosol optical properties. The increasing disagreement of measured and modeled actinic fluxes towards lower wavelength strongly implies that the BrC optical properties from the in-

situ SAEB observations may not fully represent the true properties of the plume aerosol. The next application for VLIDORT-QS is to retrieve BrC aerosol optical properties from the measured actinic flux spectra.

765 Further applications are conceivable in the context of chemical transport modeling (CTM). To improve computational efficiency, CTMs typically run at relatively coarse spatial resolution (several km) and resort to simplifications (e.g. 1D-RT modeling) and parameterizations to account for photochemical processes. Detailed modeling based on accurate measurements of the atmospheric state, similar as presented in this study, can help to understand magnitudes and origins of the uncertainties introduced into CTMs by these approximations.

770 Adding the facility to calculate actinic fluxes to VLIDORT-QS allows us to improve trace gas remote sensing retrievals of BB plumes. These retrievals depend on the trace gas vertical concentrations profiles, which for many of the target gases, such as O₃, NO₂, HCHO, and HONO, depend on the vertical profiles of the respective photolysis frequencies. By combining actinic flux profiles from VLIDORT-QS with a ~~chemical transport model~~CTM, self-consistent retrievals of trace gases should be possible.

Code and data availability. The raw data from the FIREX-AQ campaign is available on the "NASA Airborne Science Data for Atmospheric Composition" database: <https://www-air.larc.nasa.gov/cgi-bin/ArcView/firexaq>. The processed model input data and simulation results are available at DOI 10.5281/zenodo.12802619. The latest version of the VPC model is on a private Github repository. Access is provided by the authors on request.

A1

Author contributions. JLT performed the RT simulations, conducted the comparison and wrote the first draft of the manuscript. JLT, SFC and NB developed the VPC framework. RS and MC expanded the VLIDORT-QS Fortran code and provided support for its use in the VPC framework. SH and KU collected and processed the CAFS data and advised on its assimilation in the study. JH and TS collected and processed the lidar profiles and advised on their assimilation in the study. RW and JD collected and processed the SAEB data and advised on its assimilation in the study. RM collected and processed data from the TSI Laser Aerosol Spectrometer, the PSAP and the Nephelometer. VN and NT provided advice on RT, aerosol property and satellite remote-sensing aspects of the work. JS initiated the investigations and contributed to all project activities as the primary supervisor. All authors contributed to the manuscript.

785 *Competing interests.* The authors declare that they have no competing interests.

Acknowledgements. UCLA's research was funded by NOAA Grant: NA17OAR4310005 (Remote Sensing of Radical Precursor Chemistry in Biomass Burning Plumes) and, NASA Grant number 80NSSC21K1447 (Advancing UV/Vis remote sensing of biomass burning plumes:

Table A1. Abbreviations

AFOD	Actinic flux optical depth
AGL	Above ground level
AOD	Aerosol optical depth
BB	Biomass burning
BC	Black carbon
BrC	Brown carbon
CAFS	Charged-coupled device Actinic Flux Spectro-radiometer
<u>CTM</u>	<u>Chemical transport model</u>
<u>FWHM</u>	<u>Full width at half maximum</u>
MSL	Mean sea level
PSD	Particle size distribution
RAA	Solar relative azimuth angle
RMSD	Root-mean-square deviation
RMSRD	Root-mean-square of relative deviation
RT	Radiative transfer
SAA	Solar azimuth angle
SAEB	Spectral analysis of extracted BrC chromophores
SSA	Single scattering albedo
SZA	Solar zenith angle
UV	Ultra-violet (spectral range)
Vis	Visible (spectral range)
VPC	VLIDORT for photochemistry

Brown carbon, actinic flux, and trace gas retrievals). RW was supported by NASA award No. 80NSSC18K0662. JD was supported by NASA grant 80NSSC18K0631. SH and KU are supported by the NSF National Center for Atmospheric Research, which is a major facility sponsored by the U.S. National Science Foundation under Cooperative Agreement No. 1852977. Their research was funded by NASA Award Nos. 80NSSC18K0638 and 80NSSC21K1446. Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with NASA (grant No. 80NM0018D0004). We acknowledge the following FIREX-AQ PIs and participants for providing their data: Glenn Diskin (DACOM), Nick Wagner (NOAA AOP) and Elizabeth Wiggins (NASA LARGE). We acknowledge the free use of the TROPOMI surface DLER database provided through the Sentinel-5p+ Innovation project of the European Space Agency (ESA). The TROPOMI surface DLER database was created by the Royal Netherlands Meteorological Institute (KNMI).

References

- Aknan, A. and Chen, G.: NASA Airborne Science Data for Atmospheric Composition, <https://www-air.larc.nasa.gov/missions/firex-aq/>, 2023.
- 800 Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmospheric Chemistry and Physics*, 19, 8523–8546, <https://doi.org/10.5194/acp-19-8523-2019>, 2019.
- Andreae, M. O. and Gelencser, A.: Black carbon or brown carbon? The nature of light-absorbing carbonaceous aerosols, *Atmos. Chem. Phys.*, 2006.
- Bais, A. F., Madronich, S., Crawford, J., Hall, S. R., Mayer, B., van Weele, M., Lenoble, J., Calvert, J. G., Cantrell, C. A., Shetter, R. E., Hofzumahaus, A., Koepke, P., Monks, P. S., Frost, G., McKenzie, R., Krotkov, N., Kylling, A., Swartz, W. H., Lloyd, S., Pfister, G., 805 Martin, T. J., Roeth, E.-P., Griffioen, E., Ruggaber, A., Krol, M., Kraus, A., Edwards, G. D., Mueller, M., Lefer, B. L., Johnston, P., Schwander, H., Flittner, D., Gardiner, B. G., Barrick, J., and Schmitt, R.: International Photolysis Frequency Measurement and Model Intercomparison (IPMMI): Spectral actinic solar flux measurements and modeling, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002891>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2002JD002891>, 2003.
- Balis, D., Kroon, M., Koukouli, M. E., Brinkma, E. J., Labow, G., Veeffkind, J. P., and McPeters, R. D.: Validation of Ozone Monitoring Instrument total ozone column measurements using Brewer and Dobson spectrophotometer ground-based observations, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2007JD008796>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2007JD008796>, 2007.
- 810 Balis, D. S., Zerefos, C. S., Kourtidis, K., Bais, A. F., Hofzumahaus, A., Kraus, A., Schmitt, R., Blumthaler, M., and Gobbi, G. P.: Measurements and modeling of photolysis rates during the Photochemical Activity and Ultraviolet Radiation (PAUR) II campaign, *Journal of Geophysical Research: Atmospheres*, 107, PAU 5–1–PAU 5–12, <https://doi.org/10.1029/2000JD000136>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2000JD000136>, 2002.
- Barnard, J. C., Chapman, E. G., Fast, J. D., Schmelzer, J. R., Slusser, J. R., and Shetter, R. E.: An evaluation of the FAST-J photolysis algorithm for predicting nitrogen dioxide photolysis rates under clear and cloudy sky conditions, *Atmospheric Environment*, 38, 3393–3403, <https://doi.org/10.1016/j.atmosenv.2004.03.034>, 2004.
- 820 Bhartia, P. K.: OMI/Aura Ozone(O3) Total Column 1-Orbit L2 Swath 13x24 km, <https://doi.org/10.5067/AURA/OMI/DATA2024>, 2012.
- Bohren, C. F. and Huffman, D. R.: *Absorption and Scattering of Light by Small Particles*, Wiley, 1 edn., <https://doi.org/10.1002/9783527618156>, 1998.
- Bond, T. C., Doherty, S. J., Fahey, D. W., Forster, P. M., Berntsen, T., DeAngelo, B. J., Flanner, M. G., Ghan, S., Kärcher, B., Koch, D., Kinne, S., Kondo, Y., Quinn, P. K., Sarofim, M. C., Schultz, M. G., Schulz, M., Venkataraman, C., Zhang, H., Zhang, S., Bellouin, N., 825 Guttikunda, S. K., Hopke, P. K., Jacobson, M. Z., Kaiser, J. W., Klimont, Z., Lohmann, U., Schwarz, J. P., Shindell, D., Storelvmo, T., Warren, S. G., and Zender, C. S.: Bounding the role of black carbon in the climate system: A scientific assessment: BLACK CARBON IN THE CLIMATE SYSTEM, *Journal of Geophysical Research: Atmospheres*, 118, 5380–5552, <https://doi.org/10.1002/jgrd.50171>, 2013.
- Borghese, F., Denti, P., Toscano, G., and Sindoni, O. I.: Electromagnetic scattering by a cluster of spheres, *Applied Optics*, 18, 116–120, <https://doi.org/10.1364/AO.18.000116>, publisher: Optica Publishing Group, 1979.
- 830 Brion, J., Chakir, A., Charbonnier, J., Daumont, D., Parisse, C., and Malicet, J.: Absorption Spectra Measurements for the Ozone Molecule in the 350–830 nm Region, *Journal of Atmospheric Chemistry*, 30, 291–299, <https://doi.org/10.1023/A:1006036924364>, 1998.

- Browell, E.: Differential absorption lidar sensing of ozone, *Proceedings of the IEEE*, 77, 419–432, <https://doi.org/10.1109/5.24128>, conference Name: Proceedings of the IEEE, 1989.
- Burkholder, J. B., Sander, S. P., Abatt, J. R., Barker, J. R., Cappa, C., Crouse, J. D., Dibble, T. S., Huie, R. E., Kolb, C., Kurylo, M. J.,
835 Orkin, V. L., Percival, C. J., Wilmouth, D. M., and Wine, P. H.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Evaluation No. 19, Tech. Rep. JPL Publication 19-5, Jet Propulsion Laboratory, Pasadena, <http://jpldataeval.jpl.nasa.gov>, 2019.
- Cappa, C. D., Lim, C. Y., Hagan, D. H., Coggon, M., Koss, A., Sekimoto, K., de Gouw, J., Onasch, T. B., Warneke, C., and Kroll, J. H.:
840 Biomass-burning-derived particles from a wide variety of fuels – Part 2: Effects of photochemical aging on particle optical and chemical properties, *Atmospheric Chemistry and Physics*, 20, 8511–8532, <https://doi.org/10.5194/acp-20-8511-2020>, publisher: Copernicus GmbH, 2020.
- Castro, T., Ruiz-Suárez, L. G., Ruiz-Suárez, J. C., Molina, M. J., and Montero, M.: Sensitivity analysis of a UV radiation transfer model and experimental photolysis rates of NO₂ in the atmosphere of Mexico City, *Atmospheric Environment*, 31, 609–620, [https://doi.org/10.1016/S1352-2310\(96\)00199-9](https://doi.org/10.1016/S1352-2310(96)00199-9), 1997.
- Chakrabarty, R. K., Shetty, N. J., Thind, A. S., Beeler, P., Sumlin, B. J., Zhang, C., Liu, P., Idrobo, J. C., Adachi, K., Wagner, N. L.,
845 Schwarz, J. P., Ahern, A., Sedlacek, A. J., Lambe, A., Daube, C., Lyu, M., Liu, C., Herndon, S., Onasch, T. B., and Mishra, R.: Shortwave absorption by wildfire smoke dominated by dark brown carbon, *Nature Geoscience*, 16, 683–688, <https://doi.org/10.1038/s41561-023-01237-9>, number: 8 Publisher: Nature Publishing Group, 2023.
- Chance, K. and Kurucz, R.: An improved high-resolution solar reference spectrum for earth’s atmosphere measurements in the ultraviolet, visible, and near infrared, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 111, 1289–1295,
850 <https://doi.org/10.1016/j.jqsrt.2010.01.036>, 2010.
- Chang, K.-E., Hsiao, T.-C., Tsay, S.-C., Lin, T.-H., Griffith, S. M., Liu, C.-Y., and Chou, C. C. K.: Embedded information of aerosol type, hygroscopicity and scattering enhancement factor revealed by the relationship between PM_{2.5} and aerosol optical depth, *Science of The Total Environment*, 867, 161 471, <https://doi.org/10.1016/j.scitotenv.2023.161471>, 2023.
- Coddington, O. M., Richard, E. C., Harber, D., Pilewskie, P., Woods, T. N., Snow, M., Chance, K., Liu, X., and Sun, K.: Version 2 of
855 the TSIS-1 Hybrid Solar Reference Spectrum and Extension to the Full Spectrum, *Earth and Space Science*, 10, e2022EA002 637, <https://doi.org/10.1029/2022EA002637>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2022EA002637>, 2023.
- Crutzen, P. J. and Andreae, M. O.: Biomass Burning in the Tropics: Impact on Atmospheric Chemistry and Biogeochemical Cycles, *Science*, 250, 1669–1678, <https://doi.org/10.1126/science.250.4988.1669>, publisher: American Association for the Advancement of Science, 1990.
- Decker, Z. C. J., Robinson, M. A., Barsanti, K. C., Bourgeois, I., Coggon, M. M., DiGangi, J. P., Diskin, G. S., Flocke, F. M., Franchin, A.,
860 Fredrickson, C. D., Gkatzelis, G. I., Hall, S. R., Halliday, H., Holmes, C. D., Huey, L. G., Lee, Y. R., Lindaas, J., Middlebrook, A. M., Montzka, D. D., Moore, R., Neuman, J. A., Nowak, J. B., Palm, B. B., Peischl, J., Piel, F., Rickly, P. S., Rollins, A. W., Ryerson, T. B., Schwantes, R. H., Sekimoto, K., Thornhill, L., Thornton, J. A., Tyndall, G. S., Ullmann, K., Van Rooy, P., Veres, P. R., Warneke, C., Washenfelder, R. A., Weinheimer, A. J., Wiggins, E., Winstead, E., Wisthaler, A., Womack, C., and Brown, S. S.: Nighttime and daytime dark oxidation chemistry in wildfire plumes: an observation and model analysis of FIREX-AQ aircraft data, *Atmospheric Chemistry and*
865 *Physics*, 21, 16 293–16 317, <https://doi.org/10.5194/acp-21-16293-2021>, 2021.
- Deutschmann, T., Beirle, S., Frieß, U., Grzegorski, M., Kern, C., Kritten, L., Platt, U., Prados-Román, C., Pukı̄te, J., Wagner, T., Werner, B., and Pfeilsticker, K.: The Monte Carlo atmospheric radiative transfer model McArtim: Introduction and validation of Jacobians and 3D features, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 112, 1119–1137, <https://doi.org/10.1016/j.jqsrt.2010.12.009>, 2011.

- Dickerson, R. R., Kondragunta, S., Stenchikov, G., Civerolo, K. L., Doddridge, B. G., and Holben, B. N.: The Impact of Aerosols on Solar Ultraviolet Radiation and Photochemical Smog, *Science*, 278, 827–830, <https://doi.org/10.1126/science.278.5339.827>, publisher: American Association for the Advancement of Science, 1997.
- Forrister, H., Liu, J., Scheuer, E., Dibb, J., Ziemba, L., Thornhill, K. L., Anderson, B., Diskin, G., Perring, A. E., Schwarz, J. P., Campuzano-Jost, P., Day, D. A., Palm, B. B., Jimenez, J. L., Nenes, A., and Weber, R. J.: Evolution of brown carbon in wildfire plumes, *Geophysical Research Letters*, 42, 4623–4630, <https://doi.org/10.1002/2015GL063897>, 2015.
- Frith, S. M., Bhartia, P. K., Oman, L. D., Kramarova, N. A., McPeters, R. D., and Labow, G. J.: Model-based climatology of diurnal variability in stratospheric ozone as a data analysis tool, *Atmospheric Measurement Techniques*, 13, 2733–2749, <https://doi.org/10.5194/amt-13-2733-2020>, publisher: Copernicus GmbH, 2020.
- Hair, J. W., Hostetler, C. A., Cook, A. L., Harper, D. B., Ferrare, R. A., Mack, T. L., Welch, W., Izquierdo, L. R., and Hovis, F. E.: Airborne High Spectral Resolution Lidar for profiling aerosol optical properties, *Applied Optics*, 47, 6734, <https://doi.org/10.1364/AO.47.006734>, 2008.
- Hall, S. R., Ullmann, K., Prather, M. J., Flynn, C. M., Murray, L. T., Fiore, A. M., Correa, G., Strode, S. A., Steenrod, S. D., Lamarque, J.-F., Guth, J., Josse, B., Flemming, J., Huijnen, V., Abraham, N. L., and Archibald, A. T.: Cloud impacts on photochemistry: building a climatology of photolysis rates from the Atmospheric Tomography mission, *Atmospheric Chemistry and Physics*, 18, 16 809–16 828, <https://doi.org/10.5194/acp-18-16809-2018>, publisher: Copernicus GmbH, 2018.
- Hand, J. L., Day, D. E., McMeeking, G. M., Levin, E. J. T., Carrico, C. M., Kreidenweis, S. M., Malm, W. C., Laskin, A., and Desyaterik, Y.: Measured and modeled humidification factors of fresh smoke particles from biomass burning: role of inorganic constituents, *Atmospheric Chemistry and Physics*, 10, 6179–6194, <https://doi.org/10.5194/acp-10-6179-2010>, 2010.
- Hennigan, C. J., Westervelt, D. M., Riipinen, I., Engelhart, G. J., Lee, T., Collett, J. L., Pandis, S. N., Adams, P. J., and Robinson, A. L.: New particle formation and growth in biomass burning plumes: An important source of cloud condensation nuclei, *Geophysical Research Letters*, 39, 2012GL050 930, <https://doi.org/10.1029/2012GL050930>, 2012.
- Heney, L. C. and Greenstein, J. L.: Diffuse radiation in the Galaxy, *The Astrophysical Journal*, 93, 70, <https://doi.org/10.1086/144246>, 1941.
- Hess, M., Koepke, P., and Schult, I.: Optical Properties of Aerosols and Clouds: The Software Package OPAC, *Bulletin of the American Meteorological Society*, 79, 831–844, [https://doi.org/10.1175/1520-0477\(1998\)079<0831:OPOAAC>2.0.CO;2](https://doi.org/10.1175/1520-0477(1998)079<0831:OPOAAC>2.0.CO;2), publisher: American Meteorological Society Section: Bulletin of the American Meteorological Society, 1998.
- Hofzumahaus, A., Kraus, A., Kylling, A., and Zerefos, C. S.: Solar actinic radiation (280–420 nm) in the cloud-free troposphere between ground and 12 km altitude: Measurements and model results, *Journal of Geophysical Research: Atmospheres*, 107, <https://doi.org/10.1029/2001JD900142>, 2002.
- Hofzumahaus, A., Lefer, B. L., Monks, P. S., Hall, S. R., Kylling, A., Mayer, B., Shetter, R. E., Junkermann, W., Bais, A., Calvert, J. G., Cantrell, C. A., Madronich, S., Edwards, G. D., Kraus, A., Müller, M., Bohn, B., Schmitt, R., Johnston, P., McKenzie, R., Frost, G. J., Griffioen, E., Krol, M., Martin, T., Pfister, G., Röth, E. P., Ruggaber, A., Swartz, W. H., Lloyd, S. A., and Van Weele, M.: Photolysis frequency of O₃ to O(1D): Measurements and modeling during the International Photolysis Frequency Measurement and Modeling Intercomparison (IPMMI), *Journal of Geophysical Research: Atmospheres*, 109, <https://doi.org/10.1029/2003JD004333>, <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2003JD004333>, 2004.

- 905 Jaffe, D. A., O'Neill, S. M., Larkin, N. K., Holder, A. L., Peterson, D. L., Halofsky, J. E., and Rappold, A. G.: Wildfire and prescribed burning impacts on air quality in the United States, *Journal of the Air & Waste Management Association* (1995), 70, 583–615, <https://doi.org/10.1080/10962247.2020.1749731>, 2020.
- Junkermann, W.: Measurements of the $J(O^1D)$ actinic flux within and above stratiform clouds and above snow surfaces, *Geophysical Research Letters*, 21, 793–796, <https://doi.org/10.1029/93GL03498>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/93GL03498>,
910 1994.
- Junkermann, W., Platt, U., and Volz-Thomas, A.: A photoelectric detector for the measurement of photolysis frequencies of ozone and other atmospheric molecules, *Journal of Atmospheric Chemistry*, 8, 203–227, <https://doi.org/10.1007/BF00051494>, 1989.
- Kattawar, G. W. and Hood, D. A.: Electromagnetic scattering from a spherical polydispersion of coated spheres, *Applied Optics*, 15, 1996–1999, <https://doi.org/10.1364/AO.15.001996>, publisher: Optica Publishing Group, 1976.
- 915 Kaufman, Y. J.: Aerosol optical thickness and atmospheric path radiance, *Journal of Geophysical Research: Atmospheres*, 98, 2677–2692, <https://doi.org/10.1029/92JD02427>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/92JD02427>, 1993.
- Kazantzidis, A., Balis, D. S., Bais, A. F., Kazadzis, S., Galani, E., Kosmidis, E., and Blumthaler, M.: Comparison of Model Calculations with Spectral UV Measurements during the SUSPEN Campaign: The Effect of Aerosols, *Journal of the Atmospheric Sciences*, 58, 1529–1539, [https://doi.org/10.1175/1520-0469\(2001\)058<1529:COMCWS>2.0.CO;2](https://doi.org/10.1175/1520-0469(2001)058<1529:COMCWS>2.0.CO;2), publisher: American Meteorological Society Section: Journal
920 of the Atmospheric Sciences, 2001.
- Kelley, P., Dickerson, R. R., Luke, W. T., and Kok, G. L.: Rate of NO₂ photolysis from the surface to 7.6 km altitude in clear-sky and clouds, *Geophysical Research Letters*, 22, 2621–2624, <https://doi.org/10.1029/95GL02563>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/95GL02563>, 1995.
- Kiland, K. J., Mahrt, F., Peng, L., Nikkho, S., Zaks, J., Crescenzo, G. V., and Bertram, A. K.: Secondary Organic Aerosol from Biomass Burning Phenolics Could Increase Brown Carbon Lifetimes, Seed Ice Clouds, and Transport Pollutants, <https://doi.org/10.26434/chemrxiv-2023-n4qvm>, 2023.
- 925 King, M. D. and Byrne, D. M.: A Method for Inferring Total Ozone Content from the Spectral Variation of Total Optical Depth Obtained with a Solar Radiometer, *Journal of the Atmospheric Sciences*, 33, 2242–2251, [https://doi.org/10.1175/1520-0469\(1976\)033<2242:AMFITO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033<2242:AMFITO>2.0.CO;2), publisher: American Meteorological Society Section: Journal of the Atmospheric Sciences,
930 1976.
- Klimont, Z., Kupiainen, K., Heyes, C., Purohit, P., Cofala, J., Rafaj, P., Borcken-Kleefeld, J., and Schöpp, W.: Global anthropogenic emissions of particulate matter including black carbon, *Atmospheric Chemistry and Physics*, 17, 8681–8723, <https://doi.org/10.5194/acp-17-8681-2017>, 2017.
- Kotchenruther, R. A. and Hobbs, P. V.: Humidification factors of aerosols from biomass burning in Brazil, *Journal of Geophysical Research: Atmospheres*, 103, 32 081–32 089, <https://doi.org/10.1029/98JD00340>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/98JD00340>, 1998.
- 935 Kraus, A. and Hofzumahaus, A.: Field Measurements of Atmospheric Photolysis Frequencies for O₃, NO₂, HCHO, CH₃CHO, H₂O₂, and HONO by UV Spectroradiometry, in: *Atmospheric Measurements during POPCORN — Characterisation of the Photochemistry over a Rural Area*, edited by Rudolph, J. and Koppmann, R., pp. 161–180, Springer Netherlands, Dordrecht, https://doi.org/10.1007/978-94-017-0813-5_8, 1998.
940

- Kroon, M., Veefkind, J. P., Sneep, M., McPeters, R. D., Bhartia, P. K., and Levelt, P. F.: Comparing OMI-TOMS and OMI-DOAS total ozone column data, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD008798>, <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2007JD008798>, 2008.
- 945 Lack, D. A., Richardson, M. S., Law, D., Langridge, J. M., Cappa, C. D., McLaughlin, R. J., and Murphy, D. M.: Aircraft Instrument for Comprehensive Characterization of Aerosol Optical Properties, Part 2: Black and Brown Carbon Absorption and Absorption Enhancement Measured with Photo Acoustic Spectroscopy, *Aerosol Science and Technology*, 46, 555–568, <https://doi.org/10.1080/02786826.2011.645955>, 2012.
- Lampel, J., Pöhler, D., Tschritter, J., Frieß, U., and Platt, U.: On the relative absorption strengths of water vapour in the blue wavelength range, *Atmospheric Measurement Techniques*, 8, 4329–4346, <https://doi.org/10.5194/amt-8-4329-2015>, 2015.
- 950 Landgraf, J. and Crutzen, P. J.: An Efficient Method for Online Calculations of Photolysis and Heating Rates, *Journal of the Atmospheric Sciences*, 55, 863–878, [https://doi.org/10.1175/1520-0469\(1998\)055<0863:AEMFOC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1998)055<0863:AEMFOC>2.0.CO;2), 1998.
- Langridge, J. M., Richardson, M. S., Lack, D., Law, D., and Murphy, D. M.: Aircraft Instrument for Comprehensive Characterization of Aerosol Optical Properties, Part I: Wavelength-Dependent Optical Extinction and Its Relative Humidity Dependence Measured Using Cavity Ringdown Spectroscopy, *Aerosol Science and Technology*, 45, 1305–1318, <https://doi.org/10.1080/02786826.2011.592745>, 2011.
- 955 Laskin, A., Laskin, J., and Nizkorodov, S. A.: Chemistry of Atmospheric Brown Carbon, *Chemical Reviews*, 115, 4335–4382, <https://doi.org/10.1021/cr5006167>, publisher: American Chemical Society, 2015.
- Lefer, B. L., Shetter, R. E., Hall, S. R., Crawford, J. H., and Olson, J. R.: Impact of clouds and aerosols on photolysis frequencies and photochemistry during TRACE-P: 1. Analysis using radiative transfer and photochemical box models, *Journal of Geophysical Research: Atmospheres*, 108, 2002JD003 171, <https://doi.org/10.1029/2002JD003171>, 2003.
- 960 Levin, E. J. T., McMeeking, G. R., Carrico, C. M., Mack, L. E., Kreidenweis, S. M., Wold, C. E., Moosmüller, H., Arnott, W. P., Hao, W. M., Collett Jr., J. L., and Malm, W. C.: Biomass burning smoke aerosol properties measured during Fire Laboratory at Missoula Experiments (FLAME), *Journal of Geophysical Research: Atmospheres*, 115, <https://doi.org/10.1029/2009JD013601>, <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2009JD013601>, 2010.
- Levoni, C., Cervino, M., Guzzi, R., and Torricella, F.: Atmospheric aerosol optical properties: a database of radiative characteristics for different components and classes, *Applied Optics*, 36, 8031–8041, <https://doi.org/10.1364/ao.36.008031>, 1997.
- 965 Liao, J., Wolfe, G. M., Hannun, R. A., St. Clair, J. M., Hanisco, T. F., Gilman, J. B., Lamplugh, A., Selimovic, V., Diskin, G. S., Nowak, J. B., Halliday, H. S., DiGangi, J. P., Hall, S. R., Ullmann, K., Holmes, C. D., Fite, C. H., Agastra, A., Ryerson, T. B., Peischl, J., Bourgeois, I., Warneke, C., Coggon, M. M., Gkatzelis, G. I., Sekimoto, K., Fried, A., Richter, D., Weibring, P., Apel, E. C., Hornbrook, R. S., Brown, S. S., Womack, C. C., Robinson, M. A., Washenfelder, R. A., Veres, P. R., and Neuman, J. A.: Formaldehyde evolution in US wildfire
- 970 plumes during the Fire Influence on Regional to Global Environments and Air Quality experiment (FIREX-AQ), *Atmospheric Chemistry and Physics*, 21, 18 319–18 331, <https://doi.org/10.5194/acp-21-18319-2021>, 2021.
- Lindsay, A. J., Anderson, D. C., Wernis, R. A., Liang, Y., Goldstein, A. H., Herndon, S. C., Roscioli, J. R., Dyroff, C., Fortner, E. C., Croteau, P. L., Majluf, F., Krechmer, J. E., Yacovitch, T. I., Knighton, W. B., and Wood, E. C.: Ground-based investigation of HO_x and ozone chemistry in biomass burning plumes in rural Idaho, *Atmospheric Chemistry and Physics*, 22, 4909–4928, <https://doi.org/10.5194/acp-22-4909-2022>, publisher: Copernicus GmbH, 2022.
- 975 Liu, D., Li, S., Hu, D., Kong, S., Cheng, Y., Wu, Y., Ding, S., Hu, K., Zheng, S., Yan, Q., Zheng, H., Zhao, D., Tian, P., Ye, J., Huang, M., and Ding, D.: Evolution of Aerosol Optical Properties from Wood Smoke in Real Atmosphere Influenced by Burning Phase and Solar

- Radiation, *Environmental Science & Technology*, 55, 5677–5688, <https://doi.org/10.1021/acs.est.0c07569>, publisher: American Chemical Society, 2021.
- 980 Liu, H., Crawford, J. H., Pierce, R. B., Norris, P., Platnick, S. E., Chen, G., Logan, J. A., Yantosca, R. M., Evans, M. J., Kittaka, C., Feng, Y., and Tie, X.: Radiative effect of clouds on tropospheric chemistry in a global three-dimensional chemical transport model, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006403>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2005JD006403>, 2006.
- Liu, J., Bergin, M., Guo, H., King, L., Kotra, N., Edgerton, E., and Weber, R. J.: Size-resolved measurements of brown carbon in water and
985 methanol extracts and estimates of their contribution to ambient fine-particle light absorption, *Atmospheric Chemistry and Physics*, 13, 12 389–12 404, <https://doi.org/10.5194/acp-13-12389-2013>, publisher: Copernicus GmbH, 2013.
- Liu, J., Scheuer, E., Dibb, J., Ziemba, L. D., Thornhill, K. L., Anderson, B. E., Wisthaler, A., Mikoviny, T., Devi, J. J., Bergin, M., and Weber, R. J.: Brown carbon in the continental troposphere, *Geophysical Research Letters*, 41, 2191–2195, <https://doi.org/10.1002/2013GL058976>, 2014.
- 990 Liu, J., Scheuer, E., Dibb, J., Diskin, G. S., Ziemba, L. D., Thornhill, K. L., Anderson, B. E., Wisthaler, A., Mikoviny, T., Devi, J. J., Bergin, M., Perring, A. E., Markovic, M. Z., Schwarz, J. P., Campuzano-Jost, P., Day, D. A., Jimenez, J. L., and Weber, R. J.: Brown carbon aerosol in the North American continental troposphere: sources, abundance, and radiative forcing, *Atmospheric Chemistry and Physics*, 15, 7841–7858, <https://doi.org/10.5194/acp-15-7841-2015>, publisher: Copernicus GmbH, 2015.
- Mackowski, D. W. and Mishchenko, M. I.: Calculation of the T matrix and the scattering matrix for ensembles of spheres, *JOSA A*, 13,
995 2266–2278, <https://doi.org/10.1364/JOSAA.13.002266>, publisher: Optica Publishing Group, 1996.
- Madronich, S.: Photodissociation in the atmosphere: 1. Actinic flux and the effects of ground reflections and clouds, *Journal of Geophysical Research: Atmospheres*, 92, 9740–9752, <https://doi.org/10.1029/JD092iD08p09740>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/JD092iD08p09740>, 1987.
- Madronich, S. and Flocke, S.: The Role of Solar Radiation in Atmospheric Chemistry, in: *Environmental Photochemistry*, edited by
1000 Hutzinger, O. and Boule, P., vol. 2 / 2L, pp. 1–26, Springer Berlin Heidelberg, Berlin, Heidelberg, https://doi.org/10.1007/978-3-540-69044-3_1, series Title: The Handbook of Environmental Chemistry, 1999.
- Mayer, B.: Radiative transfer in the cloudy atmosphere, *The European Physical Journal Conferences*, 1, 75–99, <https://doi.org/10.1140/epjconf/e2009-00912-1>, 2009.
- McClure, C. D. and Jaffe, D. A.: US particulate matter air quality improves except in wildfire-prone areas, *Proceedings of the National
1005 Academy of Sciences*, 115, 7901–7906, <https://doi.org/10.1073/pnas.1804353115>, publisher: Proceedings of the National Academy of Sciences, 2018.
- Meier, R. R., Anderson, G. P., Cantrell, C. A., Hall, L. A., Lean, J., Minschwaner, K., Shetter, R. E., Shettle, E. P., and Stamnes, K.: Actinic radiation in the terrestrial atmosphere, *Journal of Atmospheric and Solar-Terrestrial Physics*, 59, 2111–2157, [https://doi.org/10.1016/S1364-6826\(97\)00047-3](https://doi.org/10.1016/S1364-6826(97)00047-3), 1997.
- 1010 Muinonen, K., Nousiainen, T., Fast, P., Lumme, K., and Peltoniemi, J. I.: Light scattering by Gaussian random particles: Ray optics approximation, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 55, 577–601, [https://doi.org/10.1016/0022-4073\(96\)00003-9](https://doi.org/10.1016/0022-4073(96)00003-9), 1996.
- Neu, J. L., Prather, M. J., and Penner, J. E.: Global atmospheric chemistry: Integrating over fractional cloud cover, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD008007>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2006JD008007>, 2007.
- 1015

- O'Dell, K., Ford, B., Fischer, E. V., and Pierce, J. R.: Contribution of Wildland-Fire Smoke to US PM_{2.5} and Its Influence on Recent Trends, *Environmental Science & Technology*, 53, 1797–1804, <https://doi.org/10.1021/acs.est.8b05430>, 2019.
- 1020 Palm, B. B., Peng, Q., Hall, S. R., Ullmann, K., Campos, T. L., Weinheimer, A., Montzka, D., Tyndall, G., Permar, W., Hu, L., Flocke, F., Fischer, E. V., and Thornton, J. A.: Spatially Resolved Photochemistry Impacts Emissions Estimates in Fresh Wildfire Plumes, *Geophysical Research Letters*, 48, e2021GL095443, <https://doi.org/10.1029/2021GL095443>, 2021.
- Palmer, K. F. and Williams, D.: Optical constants of sulfuric Acid; application to the clouds of venus?, *Applied Optics*, 14, 208–219, <https://doi.org/10.1364/AO.14.000208>, 1975.
- 1025 Peng, Q., Palm, B. B., Fredrickson, C. D., Lee, B. H., Hall, S. R., Ullmann, K., Campos, T., Weinheimer, A. J., Apel, E. C., Flocke, F., Permar, W., Hu, L., Garofalo, L. A., Pothier, M. A., Farmer, D. K., Ku, I.-T., Sullivan, A. P., Collett, J. L., Fischer, E., and Thornton, J. A.: Observations and Modeling of NO_x Photochemistry and Fate in Fresh Wildfire Plumes, *ACS Earth and Space Chemistry*, 5, 2652–2667, <https://doi.org/10.1021/acsearthspacechem.1c00086>, 2021.
- Platt, U. and Stutz, J.: Differential Optical Absorption Spectroscopy, *Physics of Earth and Space Environments*, Springer Berlin Heidelberg, Berlin, Heidelberg, <https://doi.org/10.1007/978-3-540-75776-4>, 2008.
- 1030 Puķīte, J., Kühn, S., Deutschmann, T., Platt, U., and Wagner, T.: Extending differential optical absorption spectroscopy for limb measurements in the UV, *Atmospheric Measurement Techniques*, 3, 631–653, <https://doi.org/10.5194/amt-3-631-2010>, 2010.
- Reid, J. S., Koppmann, R., Eck, T. F., and Eleuterio, D. P.: A review of biomass burning emissions part II: intensive physical properties of biomass burning particles, *Atmospheric Chemistry and Physics*, 5, 799–825, <https://doi.org/10.5194/acp-5-799-2005>, publisher: Copernicus GmbH, 2005.
- 1035 Rothman, L. S., Gordon, I. E., Barber, R. J., Dothe, H., Gamache, R. R., Goldman, A., Perevalov, V. I., Tashkun, S. A., and Tennyson, J.: HITEMP, the high-temperature molecular spectroscopic database, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 111, 2139–2150, <https://doi.org/10.1016/j.jqsrt.2010.05.001>, 2010.
- Ryu, Y.-H., Hodzic, A., Descombes, G., Hall, S., Minnis, P., Spangenberg, D., Ullmann, K., and Madronich, S.: Improved modeling of cloudy-sky actinic flux using satellite cloud retrievals, *Geophysical Research Letters*, 44, 1592–1600, <https://doi.org/10.1002/2016GL071892>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/2016GL071892](https://onlinelibrary.wiley.com/doi/pdf/10.1002/2016GL071892), 2017.
- 1040 Sachse, G. W., Hill, G. F., Wade, L. O., and Perry, M. G.: Fast-response, high-precision carbon monoxide sensor using a tunable diode laser absorption technique, *Journal of Geophysical Research: Atmospheres*, 92, 2071–2081, <https://doi.org/10.1029/JD092iD02p02071>, 1987.
- SAGE Science Team: SAGE II/ERBS Level 2 Solar Event Species Profiles Native Binary file - Version 7.0, https://doi.org/10.5067/ERBS/SAGEII/SOLAR_BINARY_L2-V7.0, 2012.
- 1045 Sarpong, E., Smith, D., Pokhrel, R., Fiddler, M. N., and Bililign, S.: Refractive Indices of Biomass Burning Aerosols Obtained from African Biomass Fuels Using RDG Approximation, *Atmosphere*, 11, 62, <https://doi.org/10.3390/atmos11010062>, number: 1 Publisher: Multidisciplinary Digital Publishing Institute, 2020.
- Schuster, G. L., Dubovik, O., and Holben, B. N.: Angstrom exponent and bimodal aerosol size distributions, *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006328>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1029/2005JD006328](https://onlinelibrary.wiley.com/doi/pdf/10.1029/2005JD006328), 2006.
- 1050 Segelstein, D. J.: The complex refractive index of water, Thesis, University of Missouri–Kansas City, <https://mospace.umsystem.edu/xmlui/handle/10355/11599>, accepted: 2011-09-06T20:38:49Z, 1981.

- Shavrina, A. V., Pavlenko, Y. V., Veles, A., Syniavskiy, I., and Kroon, M.: Ozone columns obtained by ground-based remote sensing in Kiev for Aura Ozone Measuring Instrument validation, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2007JD008787>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2007JD008787>, 2007.
- 1055 Shetter, R. E. and Müller, M.: Photolysis frequency measurements using actinic flux spectroradiometry during the PEM-Tropics mission: Instrumentation description and some results, *Journal of Geophysical Research: Atmospheres*, 104, 5647–5661, <https://doi.org/10.1029/98JD01381>, 1999.
- Shetter, R. E., Cinquini, L., Lefer, B. L., Hall, S. R., and Madronich, S.: Comparison of airborne measured and calculated spectral actinic flux and derived photolysis frequencies during the PEM Tropics B mission, *Journal of Geophysical Research: Atmospheres*, 107, PEM 6–1–PEM 6–12, <https://doi.org/10.1029/2001JD001320>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2001JD001320>, 2002.
- 1060 Shetter, R. E., Junkermann, W., Swartz, W. H., Frost, G. J., Crawford, J. H., Lefer, B. L., Barrick, J. D., Hall, S. R., Hofzumahaus, A., Bais, A., Calvert, J. G., Cantrell, C. A., Madronich, S., Müller, M., Kraus, A., Monks, P. S., Edwards, G. D., McKenzie, R., Johnston, P., Schmitt, R., Griffioen, E., Krol, M., Kylling, A., Dickerson, R. R., Lloyd, S. A., Martin, T., Gardiner, B., Mayer, B., Pfister, G., Röth, E. P., Koepke, P., Ruggaber, A., Schwander, H., and van Weele, M.: Photolysis frequency of NO₂: Measurement and modeling during the International Photolysis Frequency Measurement and Modeling Intercomparison (IPMMI), *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2002JD002932>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2002JD002932>, 2003.
- Shetty, N., Liu, P., Liang, Y., Sumlin, B., Daube, C., Herndon, S., Goldstein, A. H., and Chakrabarty, R. K.: Brown carbon absorptivity in fresh wildfire smoke: associations with volatility and chemical compound groups, *Environmental Science: Atmospheres*, 3, 1262–1271, <https://doi.org/10.1039/D3EA00067B>, publisher: RSC, 2023.
- 1070 Shetty, N. J., Pandey, A., Baker, S., Hao, W. M., and Chakrabarty, R. K.: Measuring light absorption by freshly emitted organic aerosols: optical artifacts in traditional solvent-extraction-based methods, *Atmospheric Chemistry and Physics*, 19, 8817–8830, <https://doi.org/10.5194/acp-19-8817-2019>, publisher: Copernicus GmbH, 2019.
- Simeonov, V., Larcheveque, G., Quaglia, P., Bergh, H. v. d., and Calpini, B.: Influence of the photomultiplier tube spatial uniformity on lidar signals, *Applied Optics*, 38, 5186–5190, <https://doi.org/10.1364/AO.38.005186>, publisher: Optica Publishing Group, 1999.
- 1075 Spurr, R. and Christi, M.: The LIDORT and VLIDORT Linearized Scalar and Vector Discrete Ordinate Radiative Transfer Models: Updates in the Last 10 Years, in: *Springer Series in Light Scattering: Volume 3: Radiative Transfer and Light Scattering*, edited by Kokhanovsky, A., pp. 1–62, Springer International Publishing, Cham, https://doi.org/10.1007/978-3-030-03445-0_1, 2019.
- Spurr, R., Wang, J., Zeng, J., and Mishchenko, M. I.: Linearized T-matrix and Mie scattering computations, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 113, 425–439, <https://doi.org/10.1016/j.jqsrt.2011.11.014>, 2012.
- 1080 Spurr, R., Natraj, V., Colosimo, S., Stutz, J., Christi, M., and Korkin, S.: VLIDORT-QS: A quasi-spherical vector radiative transfer model, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 291, 108 341, <https://doi.org/10.1016/j.jqsrt.2022.108341>, 2022.
- Spurr, R. J.: VLIDORT: A linearized pseudo-spherical vector discrete ordinate radiative transfer code for forward model and retrieval studies in multilayer multiple scattering media, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 102, 316–342, <https://doi.org/10.1016/j.jqsrt.2006.05.005>, 2006.
- 1085 Stephens, G. L. and Platt, C. M. R.: Aircraft Observations of the Radiative and Microphysical Properties of Stratocumulus and Cumulus Cloud Fields, *Journal of Applied Meteorology and Climatology*, 26, 1243–1269, [https://doi.org/10.1175/1520-0450\(1987\)026<1243:AOTRA>2.0.CO;2](https://doi.org/10.1175/1520-0450(1987)026<1243:AOTRA>2.0.CO;2), publisher: American Meteorological Society Section: *Journal of Applied Meteorology and Climatology*, 1987.

- Sumlin, B. J., Pandey, A., Walker, M. J., Pattison, R. S., Williams, B. J., and Chakrabarty, R. K.: Atmospheric Photooxidation Diminishes Light Absorption by Primary Brown Carbon Aerosol from Biomass Burning, *Environmental Science & Technology Letters*, 4, 540–545, <https://doi.org/10.1021/acs.estlett.7b00393>, publisher: American Chemical Society, 2017.
- Sumlin, B. J., Heinson, Y. W., Shetty, N., Pandey, A., Pattison, R. S., Baker, S., Hao, W. M., and Chakrabarty, R. K.: UV–Vis–IR spectral complex refractive indices and optical properties of brown carbon aerosol from biomass burning, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 206, 392–398, <https://doi.org/10.1016/j.jqsrt.2017.12.009>, 2018.
- Thalman, R. and Volkamer, R.: Temperature dependent absorption cross-sections of O₂–O₂ collision pairs between 340 and 630 nm and at atmospherically relevant pressure, *Physical Chemistry Chemical Physics*, 15, 15 371, <https://doi.org/10.1039/c3cp50968k>, 2013.
- Thompson, A. M.: The effect of clouds on photolysis rates and ozone formation in the unpolluted troposphere, *Journal of Geophysical Research: Atmospheres*, 89, 1341–1349, <https://doi.org/10.1029/JD089iD01p01341>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/JD089iD01p01341>, 1984.
- Tie, X., Madronich, S., Walters, S., Zhang, R., Rasch, P., and Collins, W.: Effect of clouds on photolysis and oxidants in the troposphere, *Journal of Geophysical Research: Atmospheres*, 108, <https://doi.org/10.1029/2003JD003659>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2003JD003659>, 2003.
- Tilstra, L. G., De Graaf, M., Trees, V., Litvinov, P., Dubovik, O., and Stammes, P.: A directional surface reflectance climatology determined from TROPOMI observations, preprint, Others (Wind, Precipitation, Temperature, etc.)/Remote Sensing/Data Processing and Information Retrieval, <https://doi.org/10.5194/amt-2023-222>, 2023.
- Trentmann, J.: Three-dimensional solar radiation effects on the actinic flux field in a biomass-burning plume, *Journal of Geophysical Research*, 108, 4558, <https://doi.org/10.1029/2003JD003422>, 2003.
- Trentmann, J., Andreae, M. O., and Graf, H.: Chemical processes in a young biomass-burning plume, *Journal of Geophysical Research: Atmospheres*, 108, 2003JD003 732, <https://doi.org/10.1029/2003JD003732>, 2003.
- Turco, R. P.: Photodissociation rates in the atmosphere below 100km, *Geophysical surveys*, 2, 153–192, <https://doi.org/10.1007/BF01447907>, 1975.
- van Weele, M. and Duynkerke, P. G.: Effect of clouds on the photodissociation of NO₂: Observations and modelling, *Journal of Atmospheric Chemistry*, 16, 231–255, <https://doi.org/10.1007/BF00696898>, 1993.
- Vandaele, A., Hermans, C., Simon, P., Carleer, M., Colin, R., Fally, S., Mérianne, M., Jenouvrier, A., and Coquart, B.: Measurements of the NO₂ absorption cross-section from 42 000 cm⁻¹ to 10 000 cm⁻¹ (238–1000 nm) at 220 K and 294 K, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 59, 171–184, [https://doi.org/10.1016/S0022-4073\(97\)00168-4](https://doi.org/10.1016/S0022-4073(97)00168-4), 1998.
- Volz-Thomas, A., Lerner, A., Pätz, H.-W., Schultz, M., McKenna, D. S., Schmitt, R., Madronich, S., and Röth, E. P.: Airborne measurements of the photolysis frequency of NO₂, *Journal of Geophysical Research: Atmospheres*, 101, 18 613–18 627, <https://doi.org/10.1029/96JD01375>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/96JD01375>, 1996.
- Vuilleumier, L., Bamer, J. T., Harley, R. A., and Brown, N. J.: Evaluation of nitrogen dioxide photolysis rates in an urban area using data from the 1997 Southern California Ozone Study, *Atmospheric Environment*, 35, 6525–6537, [https://doi.org/10.1016/S1352-2310\(01\)00379-X](https://doi.org/10.1016/S1352-2310(01)00379-X), 2001.
- Várnai, T. and Davies, R.: Effects of Cloud Heterogeneities on Shortwave Radiation: Comparison of Cloud-Top Variability and Internal Heterogeneity, *Journal of the Atmospheric Sciences*, 56, 4206–4224, [https://doi.org/10.1175/1520-0469\(1999\)056<4206:EOCHOS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1999)056<4206:EOCHOS>2.0.CO;2), 1999.

- Wagner, J., Angelini, F., Blumthaler, M., Fitzka, M., Gobbi, G., Kift, R., Kreuter, A., Rieder, H., Simic, S., Webb, A., and Weihs, P.: Investigation of the 3-D actinic flux field in mountainous terrain, *Atmospheric Research*, 102, 300–310, <https://doi.org/10.1016/j.atmosres.2011.07.008>, 2011.
- 1130 Wagner, T., Warnach, S., Beirle, S., Bobrowski, N., Jost, A., Pukite, J., and Theys, N.: Investigation of three-dimensional radiative transfer effects for UV–Vis satellite and ground-based observations of volcanic plumes, *Atmospheric Measurement Techniques*, 16, 1609–1662, <https://doi.org/10.5194/amt-16-1609-2023>, publisher: Copernicus GmbH, 2023.
- Wandinger, U. and Ansmann, A.: Experimental determination of the lidar overlap profile with Raman lidar, *Applied Optics*, 41, 511–514, <https://doi.org/10.1364/AO.41.000511>, publisher: Optica Publishing Group, 2002.
- 1135 Warneke, C., Schwarz, J. P., Dibb, J., Kalashnikova, O., Frost, G., Al-Saad, J., Brown, S. S., Brewer, W. A., Soja, A., Seidel, F. C., Washenfelder, R. A., Wiggins, E. B., Moore, R. H., Anderson, B. E., Jordan, C., Yacovitch, T. I., Herndon, S. C., Liu, S., Kuwayama, T., Jaffe, D., Johnston, N., Selimovic, V., Yokelson, R., Giles, D. M., Holben, B. N., Goloub, P., Popovici, I., Trainer, M., Kumar, A., Pierce, R. B., Fahey, D., Roberts, J., Gargulinski, E. M., Peterson, D. A., Ye, X., Thapa, L. H., Saide, P. E., Fite, C. H., Holmes, C. D., Wang, S., Coggon, M. M., Decker, Z. C. J., Stockwell, C. E., Xu, L., Gkatzelis, G., Aikin, K., Lefer, B., Kaspari, J., Griffin, D., Zeng, L., Weber, R., Hastings, M., Chai, J., Wolfe, G. M., Hanisco, T. F., Liao, J., Campuzano Jost, P., Guo, H., Jimenez, J. L., Crawford, J., and The FIREX-AQ Science
- 1140 Team: Fire Influence on Regional to Global Environments and Air Quality (FIREX-AQ), *Journal of Geophysical Research: Atmospheres*, 128, e2022JD037758, <https://doi.org/10.1029/2022JD037758>, 2023.
- Wild, O., Zhu, X., and Prather, M. J.: Fast-J: Accurate Simulation of In- and Below-Cloud Photolysis in Tropospheric Chemical Models, *Journal of Atmospheric Chemistry*, 37, 245–282, <https://doi.org/10.1023/A:1006415919030>, 2000.
- 1145 Williams, J. E., Landgraf, J., Bregman, A., and Walter, H. H.: A modified band approach for the accurate calculation of on-line photolysis rates in stratospheric-tropospheric Chemical Transport Models, *Atmospheric Chemistry and Physics*, 6, 4137–4161, <https://doi.org/10.5194/acp-6-4137-2006>, 2006.
- Wrana, F., Von Savigny, C., Zalach, J., and Thomason, L. W.: Retrieval of stratospheric aerosol size distribution parameters using satellite solar occultation measurements at three wavelengths, *Atmospheric Measurement Techniques*, 14, 2345–2357, <https://doi.org/10.5194/amt-14-2345-2021>, 2021.
- 1150 Xu, L., Crouse, J. D., Vasquez, K. T., Allen, H., Wennberg, P. O., Bourgeois, I., Brown, S. S., Campuzano-Jost, P., Coggon, M. M., Crawford, J. H., DiGangi, J. P., Diskin, G. S., Fried, A., Gargulinski, E. M., Gilman, J. B., Gkatzelis, G. I., Guo, H., Hair, J. W., Hall, S. R., Halliday, H. A., Hanisco, T. F., Hannun, R. A., Holmes, C. D., Huey, L. G., Jimenez, J. L., Lamplugh, A., Lee, Y. R., Liao, J., Lindaas, J., Neuman, J. A., Nowak, J. B., Peischl, J., Peterson, D. A., Piel, F., Richter, D., Rickly, P. S., Robinson, M. A., Rollins, A. W., Ryerson, T. B., Sekimoto, K., Selimovic, V., Shingler, T., Soja, A. J., St. Clair, J. M., Tanner, D. J., Ullmann, K., Veres, P. R., Walega, J., Warneke, C.,
- 1155 Washenfelder, R. A., Weibring, P., Wisthaler, A., Wolfe, G. M., Womack, C. C., and Yokelson, R. J.: Ozone chemistry in western U.S. wildfire plumes, *Science Advances*, 7, eabl3648, <https://doi.org/10.1126/sciadv.abl3648>, 2021.
- Zeng, L., Zhang, A., Wang, Y., Wagner, N. L., Katich, J. M., Schwarz, J. P., Schill, G. P., Brock, C., Froyd, K. D., Murphy, D. M., Williamson, C. J., Kupc, A., Scheuer, E., Dibb, J., and Weber, R. J.: Global Measurements of Brown Carbon and Estimated Direct Radiative Effects, *Geophysical Research Letters*, 47, e2020GL088747, <https://doi.org/10.1029/2020GL088747>, <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020GL088747>, 2020.
- 1160 Zeng, L., Dibb, J., Scheuer, E., Katich, J. M., Schwarz, J. P., Bourgeois, I., Peischl, J., Ryerson, T., Warneke, C., Perring, A. E., Diskin, G. S., DiGangi, J. P., Nowak, J. B., Moore, R. H., Wiggins, E. B., Pagonis, D., Guo, H., Campuzano-Jost, P., Jimenez, J. L., Xu, L., and

- Weber, R. J.: Characteristics and evolution of brown carbon in western United States wildfires, *Atmospheric Chemistry and Physics*, 22, 8009–8036, <https://doi.org/10.5194/acp-22-8009-2022>, publisher: Copernicus GmbH, 2022.
- 1165 Ångström, A.: On the Atmospheric Transmission of Sun Radiation and on Dust in the Air, *Geografiska Annaler*, 11, 156–166, <https://doi.org/10.2307/519399>, publisher: [Wiley, Swedish Society for Anthropology and Geography], 1929.