

Warm-phase Microphysical Evolution in Large-Eddy Simulations of Tropical Cumulus Congestus: Evaluating Drop Size Distribution Evolution using Polarimetry Retrievals, In Situ Measurements, and a Thermal-Based Framework

McKenna W. Stanford^{1,2,†}, Ann M. Fridlind², Andrew S. Ackerman², Bastiaan van Diedenhoven³, Qian Xiao⁴, Jian Wang⁴, Toshihisa Matsui^{5,6}, Daniel Hernandez-Deckers⁷, and Paul Lawson⁸

¹Center for Climate Systems Research, Columbia University, New York, NY, USA

²NASA Goddard Institute for Space Studies, New York, NY, USA

³SRON Netherlands Institute for Space Research, Utrecht, the Netherlands

⁴Center for Aerosol Science and Engineering, Department of Energy, Environmental and Chemical Engineering, Washington University in St. Louis, Saint Louis, MO, USA

⁵Mesoscale Atmospheric Processes Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD, USA

⁶Earth System Science Interdisciplinary Center – ESSIC, University of Maryland, College Park, MD, USA

⁷Grupo de Investigación en Ciencias Atmosféricas, Departamento de Geociencias, Universidad Nacional de Colombia, Bogotá, Colombia

⁸Stratton Park Engineering Company, Inc., Boulder, CO, USA

[†]Current affiliation: Atmospheric, Climate, and Earth Sciences Division, Pacific Northwest National Laboratory, Richland, WA, USA

Correspondence: McKenna W. Stanford (mckenna.stanford@pnnl.gov)

Abstract.

Owing to uncertainties in convective microphysics processes, improving parameterizations in Earth system models (ESMs) can benefit from observationally constrained cases suitable for scaling between cloud-resolving models and ESMs. We propose a benchmark large-eddy simulation (LES) cumulus congestus case study from the NASA Cloud, Aerosol, and Monsoon Processes Philippines Experiment (CAMP²Ex) for evaluating and improving ESMs in single-column model (SCM) mode. We seek observational constraints using novel polarimetric retrievals and in situ cloud microphysics measurements. Simulations using bulk and bin microphysics initialized with observed aerosol profiles are compared to cloud-top retrievals of cloud droplet effective radius (R_{eff}), effective variance (ν_{eff}), and number concentration (N_d) from the airborne Research Scanning Polarimeter (RSP). Both schemes reproduce characteristics of cloud-top N_d and R_{eff} that increase and decrease with altitude, respectively. Cloud-top N_d is low-biased relative to RSP retrievals in both schemes, potentially due to limitations in both simulations and retrieval assumptions. Cloud-top R_{eff} is low-biased in the bulk scheme but reasonably reproduced by the bin scheme. Profiles of N_d and R_{eff} are sensitive to the collision-coalescence process and the vertical variation of aerosol size distribution. Comparison of simulated and in situ droplet size distributions (DSDs) show that, to first order, integrated moments are always sensitive to sizes $< \sim 30 \mu\text{m}$, and can also be sensitive to larger sizes if the DSDs are sufficiently broad, with implications for the assumed maximum observed size retrieved by the RSP. The bin scheme captures the observed extended tail of the DSD while

the bulk scheme is unable due to parametric constraints. Differences in expected relationships between in situ measurements of cloud cores and cloud-top retrievals by RSP demonstrate difficulty in well constraining the case presented herein. Finally, a thermal-tracking framework demonstrates that the dilution of N_d throughout a thermal's lifetime is heavily determined by collision-coalescence and the height-varying aerosol distribution, and that in the absence of these, the impact of entrainment on diluting N_d is largely offset by secondary activation. Implications for evaluating warm-phase convective microphysics schemes in ESMs and translating results to global, space-based polarimetry platforms are discussed.

1 Introduction

Cumulus congestus clouds play an important role in the global water and energy budget. In the tropics, they represent the intermediate mode of the trimodal tropical convection distribution in between shallow trade-wind cumuli and deep convection (Johnson et al., 1999). Definitions vary in the literature, but cumulus congestus generally have cloud top heights (CTHs) between 4 and 8 km that are either stabilized around the 0 °C level (so-called "terminal congestus") or penetrate the 0 °C level with sustained vertical growth ("transient congestus"). Wall et al. (2013) showed using 5 years of CloudSat profiles that congestus contribute up to 12% of the total cloud population in the tropics and up to 18% of all clouds with tops lower than 8 km over regions such as the Amazon, central Africa, and the maritime continent. During the Tropical Ocean Global Atmosphere Coupled Ocean Atmosphere Response Experiment (TOGA-COARE) field campaign (Webster and Lukas, 1992) conducted over the western Pacific warm pool, Johnson et al. (1999) concluded that 57% of precipitating convective clouds were identified as congestus and contributed to 28% of total convective rainfall. Transient congestus, which were shown to account for ~ 30-40% of congestus clouds observed by CloudSat by Luo et al. (2009), are also important for promoting growth to deeper convective clouds (Kuang and Bretherton, 2006; Waite and Khouider, 2010; Hohenegger and Stevens, 2013). Understanding their dynamical and microphysical evolution is therefore crucial for developing cumulus and convection parameterizations for large-scale models that account for their contributions to global precipitation and role in redistributing heat, momentum, aerosol, and moisture throughout the troposphere.

Mechanistically, congestus dynamics and microphysics are intricately linked. Cumulus clouds are composed of numerous thermals with relatively short lifetimes (3-5 minutes; Hernandez-Deckers and Sherwood, 2016, 2018; Matsui et al., 2024) that successively rise to the thermodynamic neutral buoyancy level, unless their ascent is precluded by the effects of dry-air entrainment. Supersaturation in these thermals acts as the primary source of condensation. However, thermals generate toroidal circulations that enhance cloud dilution on the inflow branch via entrainment of relatively dry environmental air, therefore additionally acting to evaporate condensate (e.g., Lasher-Trapp et al., 2005; Moser and Lasher-Trapp, 2017; Morrison et al., 2020; Peters et al., 2020; Pardo et al., 2020). Chandrakar et al. (2021) used a detailed Lagrangian microphysics model to simulate relatively shallow cumulus (CTHs < 5 km) and found that entrainment of aerosols by thermal circulations also played a significant role in secondary activation (activation of cloud droplets above cloud base). Entrainment and secondary activation are two mechanisms that contribute to the broadening of drop size distributions (DSDs) with altitude as a thermal rises, in addition to the collision-coalescence process and condensational growth. DSD broadening with height has been evaluated

extensively in large-eddy simulations (LES) of cumulus (e.g., Lasher-Trapp et al., 2005; Cooper, 1989; Grabowski and Abade,
50 2017; Morrison et al., 2018; Chandrakar et al., 2021), is supported by theory (Cooper, 1989), and has been documented
observationally in cumulus clouds (Warner, 1969; Manton, 1979; Lawson et al., 2015, 2017, 2022). Broadening mechanisms
additionally impact ice production through ice multiplication processes. Laboratory studies of secondary ice production by a
drop-shattering process indicate more numerous tiny splinters emitted during multiplication and more frequent fragmentation
with increased size of the frozen drop (e.g., Phillips et al., 2018; Lauber et al., 2018). Because DSD broadening is generally
55 coincident with the production of larger drop sizes that reach the drizzle size regime, the initiation of ice multiplication once the
drops are lofted may be dependent on the efficiency of the broadening mechanism. Indeed, Lawson et al. (2015, 2017, 2022)
showed observational evidence of fractured frozen drops and spicules, which are indicative of a drop-shattering event (Keinert
et al., 2020), in aircraft measurements of tropical cumulus congestus where DSD broadening with height was also observed.
On the other hand, copious ice production was not observed in a high-based congestus case sampled over the United Arab
60 Emirates (UAE) shown in Morrison et al. (2022) where DSDs never showed substantial precipitation-sized drops.

In this work we evaluate the microphysical evolution of a tropical cumulus congestus case study using LES and available
observational constraints. Here we focus only on the liquid phase and the DSD evolution using a thermal-based framework.
Future work will build on this foundation, extending evaluation to the ice phase and exploring the role of ice multiplication. The
selected case was observed during the NASA Cloud, Aerosol, and Monsoon Processes Philippines Experiment (CAMP²Ex)
65 aircraft-based field campaign (Reid et al., 2023) and is representative of a field of cumulus congestus with growing tops that
eventually realize an organized structure with cloud tops reaching ~ -15 °C. Simulations are performed using both bulk and
bin microphysics schemes and are constrained in a number of ways. First, observed aerosol particle size distributions (PSDs)
are used as input to represent trimodal lognormal distributions that vary with height. Second, large-scale thermodynamic and
vertical motion conditions are harvested from a nested mesoscale simulation. Third, cloud-top drop number concentration (N_d),
70 effective radius (R_{eff}), and effective variance (ν_{eff} , a measure of the DSD spectral width) are constrained to the extent possible
using retrievals from the airborne Research Scanning Polarimeter (RSP; Cairns et al., 1999) and in-situ microphysics measure-
ments. The multi-angle, multi-wavelength RSP measures total and polarized reflectance at cloud top and allows retrieval of R_{eff}
and ν_{eff} using the sharply-defined cloud-bow at scattering angles in the rainbow region of the visible spectrum. The retrieval of
both R_{eff} and ν_{eff} gives sufficient information about the DSD to retrieve N_d with relatively few assumptions (Sinclair et al., 2019)
75 that verges from bi-spectral approaches where ν_{eff} is considered a constant. Importantly, more RSP retrievals were available
during the CAMP²Ex campaign than in any prior campaign. To this end, we evaluate the utility of spatiotemporally expansive
RSP retrievals to supplement in situ aircraft transects in representing the warm-phase microphysical evolution of congestus,
which to our knowledge is the first such study to do so. Using airborne polarimetry retrievals for model evaluation is particularly
relevant with the recent successful launch of the Plankton, Aerosol, Cloud, Ocean Ecosystem (PACE) satellite (Werdell et al.,
80 2019), which includes two multi-angle polarimeters that have significantly coarser footprints relative to the airborne platform.
Understanding the capabilities and limitations of these instruments and will be crucial for extrapolating airborne platforms and
high-resolution models to space-based platforms and global models. Finally, we incorporate a thermal-tracking framework at

high temporal frequency (Hernandez-Deckers and Sherwood, 2016, 2018) to isolate microphysical processes occurring within thermals and their contribution and control over the evolving DSD.

85 Cloud-top N_d and R_{eff} retrievals and in situ DSD measurements during this CAMP²Ex case study indicate agreement with past studies in which DSDs broaden with altitude, N_d decreases with height, and R_{eff} increases with height. Sensitivity tests focus on two processes that exert potentially leading controls on such basic and widely observed profile features: (1) the efficiency of collision-coalescence and its parameterization in different warm-rain formulations and (2) the height-variation of aerosol PSDs. The thermal-tracking framework then examines the role of entrainment and mixing in modulating these
90 profiles. Morrison et al. (2022) similarly evaluated bin microphysics LES of a high-based congestus case from the UAE and compared it to in situ aircraft observations. They also examined the effects on DSD evolution of collision-coalescence and entrainment processes, as well as secondary activation, aerosol loading, and giant cloud condensation nuclei (CCN). Their simulations indicated that warm-rain generation was more controlled by the sub-cloud aerosol distributions than by activation of entrained aerosol and that dilution effects on warm-rain formation was limited under such conditions. Notably, the case they
95 studied lacked development of large precipitation-sized drops whereas here we investigate a case that produced substantial precipitation-sized drops. Chandrakar et al. (2024) also analyzed a CAMP²Ex congestus case using Lagrangian particle-based microphysics and found the inclusion of turbulent enhancement of collision coalescence to be essential in representing DSD broadening and precipitation formation, with little influence by giant CCN. Herein we also consider the turbulent enhancement of collision coalescence in the bin microphysics scheme, but omit giant CCN in part for lack of observational constraints on
100 the vertical profile and lack of evidence as a dominant control on profile features.

An additional key component of this work is to develop an LES case study for evaluating physics schemes in single-column model (SCM) versions of large-scale models, for example as has been demonstrated for subtropical marine stratocumulus-to-cumulus transitions (Sandu and Stevens, 2011; Neggers, 2015; Neggers et al., 2017), Arctic mixed-phase boundary layer clouds (Klein et al., 2009; Fridlind and Ackerman, 2017), and the currently ongoing cold air outbreak LES-SCM intercom-
105 parison project (Juliano et al., 2022). While LES intercomparison projects have been successfully carried out for case studies of shallow trade cumulus (Siebesma et al., 2003), precipitating trade cumulus (VanZanten et al., 2011), continental cumulus (Vogelmann et al., 2015; Endo et al., 2015; Lin et al., 2015), and tropical deep convection (Fridlind et al., 2012; Varble et al., 2011), considerably less attention has been paid to the congestus regime until recently. For example, a precipitating cumulus congestus case from the CAMP²Ex campaign was presented for LES and cloud-resolving model intercomparison studies at
110 the 11th International Cloud Modeling Workshop (ICMW) in Seoul, South Korea in 2024. However, a key difference between the ICMW set-up and that presented herein is the former’s use of spatially patterned surface heat fluxes as a convective forcing mechanism, which is not straightforward to replicate in typical SCM setups. In serving as an LES benchmark case for SCM simulations, we mean to indicate that (1) the meteorological and aerosol set-up is suitable for initializing and forcing the two model types identically, (2) the simulated conditions reproduce basic cloud macroscopic features observed (e.g., cloud
115 top height), and (3) the degree to which simulations statistically reproduce various measurements of cloud microphysical features has been established to the degree possible (e.g., R_{eff}). To this end, we provide observational constraints on the profiles of environmental aerosol, thermodynamics and large-scale vertical motion, and statistically robust polarimetric retrievals that sup-

plement sparse in situ measurements of warm-cloud microphysics. Finally, using the thermal-based framework, we investigate implications for developing convective microphysics in large-scale models that utilize spectral convection parameterizations with varying entrainment rates.

The remainder of the article is structured as follows. Observations including environmental aerosol and in situ cloud measurements, RSP retrievals, and a description of the case study are provided in Section 2. The LES setup and experimental design are given in Section 3 and results are presented in Section 4. Finally, a discussion of using this case as a benchmark for tropical congestus SCM studies, potential implications for convective microphysics parameterization development, and translation to space-based polarimetry platforms are provided in Section 5, followed by conclusions in Section 6.

2 Observations

Observations were obtained during the National Aeronautics and Space Administration (NASA) Cloud, Aerosol, and Monsoon Processes Philippines Experiment (CAMP²Ex; Reid et al., 2023). CAMP²Ex was an aircraft-based field campaign held from 25 August - 5 October 2019 and based out of the Clark International Airport, Philippines. Aircraft included NASA's P-3B and the Stratton Park Engineering Company's (SPEC) Learjet 35, which carried out 17 and 13 science flights, respectively. The goal of CAMP²Ex was to characterize the interaction of clouds, aerosol, and radiation in the monsoon system of Southeast Asia's maritime continent. Flights sampled a range of cloud conditions from shallow cumulus to deep convective systems. Here, we focus on a cumulus congestus event that occurred on 25 September 2019.

2.1 Aerosols: Fast Integrated Mobility Spectrometer (FIMS)

Aerosol size distributions with electrical mobility diameter (diameter derived from measuring the electrical mobility of a particle and thus the drag force that is a function of the particle shape and orientation) ranging from 10 to 600 nm were measured at 1 Hz resolution by the Fast Integrated Mobility Spectrometer (FIMS; Wang et al., 2017b, a, 2018; Kulkarni and Wang, 2006) onboard the NASA P-3B aircraft. FIMS operates by simultaneously detecting particles of different sizes based on the displacement of charged particles in an electric field. Calibration was performed before and after the CAMP²Ex campaign. Calibrations of sizing accuracy and detection efficiency followed the procedure described in Wang et al. (2017a). Aerosol PSDs ranging from 10 to 600 nm were derived in 30 size bins with a bin width ($\Delta \log_{10} D_p$) equal to 0.061, where D_p is the single particle diameter, using the inversion technique of Wang et al. (2018).

Aerosol representation in our numerical simulations follows the methodology of Fridlind et al. (2017). Out-of-cloud FIMS measurements were composited over altitude ranges (Fig. 1b-c) and fitted to a distribution by minimizing the sum of squared residuals as follows:

$$\text{Residual} = \sum (F_i - \text{Obs}_i)^2 \quad (1)$$

where F_i is the fitted concentration and Obs_i is the FIMS concentration for size bin i . Three lognormal modes are then derived with a lognormal distribution for each mode represented by:

$$dN/d\ln D_p = \frac{N}{\sqrt{2\pi\ln(\sigma_g)}} \exp\left[-\frac{\ln^2(D_p/D_g)}{2\ln^2(\sigma_g)}\right] \quad (2)$$

where D_g is the geometric mean particle diameter and σ_g is the geometric standard deviation. Aerosol number concentrations (N_a) vary with height at 0.5-km increments (Fig. 1a) while D_g and σ_g are held constant with height. D_g and σ_g of the three modes were derived from fitting the average PSD of 0-3 km to a trimodal distribution by optimizing the residual defined in Eq. 1. The realized trimodal D_g are 27, 62, and 153 nm, which roughly correspond to nucleation, Aitken, and accumulation modes, respectively. The hygroscopicity parameter (κ) is derived from time-averaged aerosol mass spectrometer (AMS) measurements for this research flight. Since the AMS does not provide size-resolved chemical composition, κ is assumed constant for all three modes. However, ongoing work to derive size-resolved κ based on CCN measurements suggests little variability of κ with supersaturation (i.e., size). We assume that aerosols consist of only $(\text{NH}_4)_2\text{SO}_4$ as the inorganic component with $\kappa = 0.53$ and an organic species with $\kappa = 0.1$ (Petters and Kreidenweis, 2007). The mass fractions of both components are converted to volume fractions using densities of 1.77 g cm^{-3} for $(\text{NH}_4)_2\text{SO}_4$ and 1.4 g cm^{-3} for average organic components (Hallquist et al., 2009). Consequently, the AMS-based κ using Equation (7) of Petters and Kreidenweis (2007) is 0.4. This derivation assumes that aerosols are internally mixed. Characterizing aerosol mixing state was not possible during CAMP²Ex. However, back-trajectories for this flight show exclusively fetches over open ocean, suggesting that the aerosols were minimally influenced by pollution sources (e.g., biomass burning smoke and anthropogenic sources from the metro Manila region). Furthermore, Xu et al. (2021) suggest that the internal mixing assumption in clean marine aerosol environments did not induce any significant error in a CCN closure study.

2.2 Clouds: Research Scanning Polarimeter (RSP) retrievals

Retrievals of cloud-top N_d , R_{eff} , and ν_{eff} are performed using the airborne Research Scanning Polarimeter (RSP; Cairns et al., 1999). The RSP makes total intensity and polarimetric measurements in nine spectral bands in the visible/near infrared and shortwave infrared. Onboard the P-3B aircraft, the RSP scans a given point on a cloud from multiple viewing angles (Alexandrov et al., 2012a). The CTH is retrieved using a multi-angle parallax method (Sinclair et al., 2017). Single-scattered light between scattering angles of 135° and 165° describe the sharply-defined cloud-bow in polarized reflectance that is used to retrieve the cloud-top R_{eff} and ν_{eff} of the DSD (Alexandrov et al., 2012a). Cloud optical depth is retrieved from near-nadir reflectance measurements at a wavelength of 865 nm. Further details of RSP retrievals are provided by, e.g., Sinclair et al. (2017, 2020, 2021).

The analytical expression for R_{eff} is defined following Hansen and Travis (1974) as:

$$R_{\text{eff}} = \frac{\int_{r_1}^{r_2} \pi r^3 \frac{dn}{dr} dr}{\int_{r_1}^{r_2} \pi r^2 \frac{dn}{dr} dr} = \frac{\langle r^3 \rangle}{\langle r^2 \rangle} \quad (3)$$




Figure 1. (a) Profiles of aerosol number concentration (N_a) for three lognormal modes with height-invariant geometric mean particle diameter (D_p) and geometric standard deviation (σ_g) listed above the panel. (b)-(d) Lognormal aerosol size distributions measured by the fast integrated mobility spectrometer (FIMS, dashed grey), the three derived modes (blue, light blue, and red), and the trimodal fit for all modes (black) for three example altitude ranges that span sub-cloud to upper entrainment environments (0-0.5 km, 2.5-3 km, and 5.5-6 km).

where r is drop radius, $\frac{dn}{dr}dr$ is the number of particles per unit volume between radii of r and $r + dr$, r_1 and r_2 are the upper and lower size limits of a give size distribution, respectively, and $\langle r^n \rangle = \int_{r_1}^{r_2} r^n \frac{dn}{dr} dr$ is the n^{th} moment of the DSD. The effective variance is is defined as:

$$\nu_{\text{eff}} = \frac{\int_{r_1}^{r_2} (r - R_{\text{eff}})^2 \pi r^2 \frac{dn}{dr} dr}{R_{\text{eff}}^2 \int_{r_1}^{r_2} \pi r^2 \frac{dn}{dr} dr} = \frac{\langle r^4 \rangle \langle r^2 \rangle}{\langle r^3 \rangle^2} - 1 \quad (4)$$

and is a dimensionless measure of the width of the size distribution. Droplet number concentrations at cloud top are derived from the RSP-retrieved R_{eff} and ν_{eff} following, e.g., Grosvenor et al. (2018) and Sinclair et al. (2019), as follows:

$$N_d = \frac{\sqrt{5}}{2\pi k} \left(\frac{f_{\text{ad}} c_w \tau_c}{Q_{\text{ext}} \rho_w R_{\text{eff}}^5} \right)^{\frac{1}{2}} \quad (5)$$

where f_{ad} is the subadiabatic factor, τ_c is the cloud optical depth, c_w is the adiabatic condensation rate (e.g., Brenguier et al., 2000; Painemal and Zuidema, 2011), Q_{ext} is a unitless extinction efficiency factor, ρ_w is the bulk density of liquid water (1000 kg m⁻³), and k is a parameter that relates ν_{eff} to R_{eff} following:

$$k = \left(\frac{R_v}{R_{\text{eff}}} \right)^3 = (1 - \nu_{\text{eff}})(1 - 2\nu_{\text{eff}}) \quad (6)$$

where R_v is the volume-mean droplet radius (Grosvenor et al., 2018). The k parameter has historically been used by the satellite community and is considered constant for bi-spectral retrievals, though the parameter's natural variability has been documented via aircraft observations (e.g. Painemal and Zuidema, 2011). This illustrates a distinct advantage of the polarimeter's ability to retrieve ν_{eff} and subsequently implement the k parameter into Eq. 5. Here, k , R_{eff} , and τ_c are retrieved by the RSP, Q_{ext} is assumed to be 2 for Mie scattering, c_w is calculated via dropsondes from the CAMP²Ex flight and decreases with altitude, and f_{ad} is assumed to be 0.8 following in situ profiles from the NASA North Atlantic Aerosols and Marine Ecosystems Study (Behrenfeld et al., 2019; Alexandrov et al., 2018). Grosvenor et al. (2018) conclude that different observations suggest f_{ad} of 0.66 ± 0.22 , which encompasses the value chosen here. However, since N_d scales with $f_{\text{ad}}^{\frac{1}{2}}$, using 0.66 versus 0.8 would decrease N_d by only a factor of 0.9. RSP retrievals are considered to represent ~ 1 optical depth below cloud top (Miller et al., 2018; Alexandrov et al., 2018). Importantly, RSP retrievals are inherently truncated at a size limit that is dependent on the size distribution being observed (see Fig. 6 in Reid et al., 2023). Here, we assume this lower size limit be a diameter (radius) of 200 (100) μm . Using in situ cloud microphysics (discussed next), we explore the implications of using RSP retrievals at this limit and discuss relevant sensitivities.

2.3 Clouds: In situ cloud microphysics measurements

In situ cloud probes are used to evaluate DSD characteristics within cloud cores. Probes were instrumented on both aircraft but only Learjet measurements are used herein. These include the fast forward scattering spectrometer probe (FFSSP; O'Connor et al., 2008; Lawson et al., 2017), the Nevzorov liquid and total water content device (Korolev et al., 1998), the 10- μm channel 2D-stereo (2D-S) optical array probe (Lawson et al., 2006), and the high-volume precipitation spectrometer probe (HVPS;

200 Lawson et al., 1998). In situ probes are used to evaluate similarities with RSP retrievals and to provide continuous size distributions that extend beyond the limits of the RSP. Further discussion of combining these instruments to produce continuous size distributions is provided in Appendix D.

2.4 Case Description

The event on 25 September 2019 corresponded to Research Flight 14 (RF14) for the P-3B and RF12 for the SPEC Learjet from
205 ~ 0300 UTC to 0900 UTC. The 0°C level during this case was ~ 5 km above ground level (AGL). The P-3B aircraft initially sampled cumulus clusters with CTHs ~ 4 - 5 km, as demonstrated in Fig. 2, which shows a 30-km overpass of a cumulus congestus cluster during this case along with time-height cross sections of radar reflectivity from the W-band 3rd Generation Airborne Precipitation Radar (APR3) and a time-series of RSP-retrieved cloud-top N_d , R_{eff} , and ν_{eff} . Cloud-top heights over ocean were likely limited by a prominent dry layer ~ 4 - 4.5 km AGL where a dewpoint depression of $\sim 20^\circ\text{C}$ was evident
210 in dropsondes (see Fig. 3b). However, the strength of imposed large-scale vertical motion additionally modulated maximum CTHs in simulations, which is connected to the thermodynamic profile by shifting the vertical structure of relative humidity.

The P-3B mostly sampled at a cruising altitude of ~ 7 km AGL. The overpass shown in Fig. 2 was over ocean and during the earlier portion of the flight. During the latter portion of the flight (after ~ 0600 UTC), the P-3B sampled a more vigorous cluster of convection near the southern end of Catanduanes Island (see location in Fig. 4). Continental surface fluxes and
215 enhanced terrain (up to ~ 1.5 km above sea level) from the island are presumed to have invigorated convection along with continued moistening from the cloud system situated to the north of the island (not shown). Importantly, sampling of clouds over Catanduanes Island by the P-3B were mostly in-situ, which since the RSP requires the plane to fly above cloud top, means the RSP was unable to sample the tops of the most vigorous clouds. Therefore, forthcoming RSP retrievals are shown mostly for cumulus congestus clusters over ocean with CTHs warmer than 0°C before the aircraft sampled the more vigorous
220 convection over the island.

The Learjet also performed a number of cloud penetrations prior to 0600 UTC, after which it continuously sampled in-cloud transects at higher altitudes over the southern portion of Catanduanes Island (Fig. 4). While sampling the island convection, the Learjet indeed sampled CTHs reaching up to 7 km AGL where ice formation proceeded. In-situ images from a cloud particle imager (CPI; Lawson et al., 2001; Woods et al., 2018) showed indications of ice multiplication by means of fractured drops and
225 spicules. Ice multiplication for this event, including the development of a drop shattering parameterization based on laboratory data, is explored in a subsequent study.

3 Model Setup and Experimental Design

Here we describe the LES model setup and experimental design. The LES is initialized with horizontally uniform thermodynamic and large-scale vertical motion (w_{LS}) profiles. While thermodynamic profiles can be obtained from dropsondes, aircraft,
230 and/or radiosondes released nearby, w_{LS} is more difficult to quantify observationally. Prior work indicated the importance of including w_{LS} as LES forcing in order to realize observed conditions. In the absence of observational data, we use mesoscale




Figure 2. Time series of (a) profiles of W-band radar reflectivity from the 3rd Generation Airborne Precipitation Radar (APR3) and CTHs retrieved from the Research Scanning Polarimeter (RSP), (b) the RSP-retrieved cloud-top drop number concentration (N_d), (c) effective radius (R_{eff}), and (d) effective variance (ν_{eff}).

simulations by the NASA Unified Weather Research and Forecasting (NU-WRF; Peters-Lidard et al., 2015) model to estimate thermodynamic and dynamic forcing.

3.1 Harvesting Thermodynamic and Dynamic Forcing from WRF

235 Specific details of the NU-WRF simulation set-up are provided in Appendix A, and only a brief description is provided here. We employ a nested set-up with an outer domain with a horizontal mesh of 3 km and an inner domain with a horizontal mesh of 600 m. The inner 600-m mesh domain is shown in Fig. 4 which encompasses the relevant portions of the P-3B and SPEC Learjet flights. Thermodynamic profiles (temperature and humidity) are averaged over this domain and shown in Fig. 3b. Two dropsondes from the flight are also shown in Fig. 3b, where dropsonde #1 was released in an ambient environment far from
240 active convection and dropsonde #2 was released in an ambient but near-convection environment. Regardless, both dropsondes agree well with the profile derived from the NU-WRF domain-average.

Various sectors of the domain shown in Fig. 4 were evaluated to derive w_{LS} . While there was considerable variability across the domain and time, spatial and temporal averaging (across hours that corresponded to flight times, $\sim$ 0300-0900 UTC) yielded a characteristic profile that is shown in Fig. 3a, with positive vertical motion in the lowest 5 km and subsidence between 5 km
245 and 10 km. The subsidence between 5 and 10 km conceptually seems consistent with the extreme dry layer represented in the sounding $\sim$ 5 km. As a result, we proceed with the idealized w_{LS} profile provided in Fig. 3a. We note that while we do not consider this w_{LS} profile to be very well-constrained with respect to observations, we do consider it to be plausible and sufficient for representing the large-scale vertical motion in the presented simulations. Furthermore, because a foundational objective of this study is to establish a cumulus congestus case study that can be evaluated in SCM simulations for convective
250 microphysics development and evaluation in large-scale models, it is pertinent to provide a measure of w_{LS} to also force the SCM. Appendix B provides further details of the importance of including w_{LS} to force the LES dynamic conditions.

3.2 Large-Eddy Simulations

Large-eddy simulations are performed using the Distributed Hydrodynamic Aerosol and Radiative Modeling for Atmospheres (DHARMA; Stevens and Bretherton, 1996; Stevens et al., 2002; Ackerman et al., 2000) model with doubly periodic boundary
255 conditions and a horizontal mesh at 100 m. The domain size is 19.2 x 19.2 km and spans 20 km vertically. Vertical grid spacing increases linearly from $\sim$ 20-100 m in the lowest 1 km, constant at 100 m between 1 and 15 km, and coarsens above 15 km. A larger domain size of 38.4 x 38.4 km did not qualitatively change results shown here. A dynamic Smagorinsky turbulence model from Kirkpatrick et al. (2006) is used for parameterizing subgrid fluxes, and ocean surface fluxes follow the bulk aerodynamic formula of Zeng et al. (1998). Radiation is neglected for simplicity because the impacts of radiative
260 heating and cooling on microphysical processes are considered to be second-order relative to congestus dynamics. Simulations are integrated for 12 h and use a dynamics time step of 1 second. Although ice particles were identified at temperatures below 0 °C when the aircrafts sampled invigorated convection over the southern portion of Catanduanes Island, we neglect ice here to narrow focus on warm-phase microphysical processes. The inclusion of ice will be discussed in a follow-up study, but sensitivity tests including ice indicate this does not significantly affect the warm-phase microphysical evolution described




Figure 3. Profiles of (a) large-scale vertical motion (w_{LS}) and (b) thermodynamic profiles shown on a skew T – log P diagram from the NU-WRF-derived sounding used to initialize the LES and two dropsondes released in the ambient (#1) and near-convection (#2) environments by the P-3B aircraft, as included in the legend.




Figure 4. Inner NU-WRF domain used for harvesting large-scale vertical motion and thermodynamics to initialize the DHARMA simulations (blue box) and flight paths of the SPEC Learjet and the NASA P-3B aircraft.

265 here. Simulations are performed using both bulk and bin (size-resolved) microphysics schemes. An overview of all sensitivity experiments is provided in Table 1.

3.2.1 Bulk Microphysics

Bulk microphysics follow a substantially modified version of the two-moment scheme of Morrison et al. (2005, 2009) using only cloud and rain hydrometeor species, with the addition of three aerosol modes. The control simulation (CNTL) and all but
270 one sensitivity simulation use the droplet autoconversion and self-collection parameterization of Seifert and Beheng (2001). Rain accretion, self-collection, breakup, and fall speed follow Seifert (2008), and the gamma size distribution for rain uses a shape parameter of 3. Sensitivity to the warm-rain formulation is tested using the Khairoutdinov and Kogan (2000) scheme for autoconversion and accretion and employing an exponential size distribution to emulate the parameterization most commonly employed in bulk microphysics models (experiment named KK). Aerosol is activated using Köhler theory following
275 Abdul-Razzak and Ghan (2000) for multiple aerosol modes. This method derives a maximum supersaturation (equal to the critical supersaturation of the smallest activated particles) as a function of dimensionless parameters that include solute effects, curvature effects, and PSD lognormal distribution properties. Here, we use a prognostic supersaturation value after microphysical relaxation that follows from Morrison and Grabowski (2008a). The number concentration of activated and unactivated aerosol are tracked throughout model integration. The sensitivity of N_d and R_{eff} profile modulation to collision-coalescence is
280 examined by turning off autoconversion as the lower extreme (experiment named NO_AC) and by scaling the autoconversion efficiency by a factor of two as an upper extreme (experiment named 2X_AC). The impact of the height-resolved aerosol PSD is explored by fixing each mode of the trimodal lognormal size distributions to the cloud-base value, therefore making the profile constant with height as a number mixing ratio (experiment named FIXED_AERO). The combined sensitivity to autoconversion, collision-coalescence, and the height-resolved aerosol PSD is then explored in the FIXED_AERO_NO_AC
285 experiment. Finally, a resolution sensitivity test is performed with the same physics as CNTL but with a horizontal mesh at 50 m that is discussed in Appendix C.

3.2.2 Bin Microphysics

The bin microphysics scheme is based on the Community Aerosol-Radiation-Microphysics Application (CARMA) code (Ackerman et al., 1995; Jensen et al., 1998). Prognostic species include bin-wise unactivated aerosol number, liquid drop number,
290 and aerosol core mass in liquid drops. Core mass is tracked in a droplet size bin by solving a continuity equation for the total dissolved aerosol mass, enabling calculation of the mean solute effect on droplet growth rate in a manner that conserves total solute mass. For each species, we use a geometric mass grid with a constant ratio of masses between two adjacent bins. The mass ratio ($\mathcal{M}$) is defined as $m_{i+1}/m_i = 2^{1/s}$, where m_i is the drop mass in the i^{th} bin and s is a bin width parameter such that drop mass is doubled every s bins. For liquid drops, $\mathcal{M} = 1.65$ ($s \sim 1.384$) and for aerosols $\mathcal{M} = 1.35$ ($s \sim 2.309$). For
295 spherical particles, the corresponding geometric size grid is $r_{i+1}/r_i = 2^{1/(3s)}$ where r_i is the radius of the i^{th} bin. For aerosols, the mass in the smallest bin corresponds to a diameter of ~ 10 nm and the largest bin corresponds to a diameter of ~ 1.5 μm . For liquid drops, the size bins range from 2 μm to 7.78 mm in diameter. Lee et al. (2021) evaluated the impacts of numerical

broadening in CARMA using the geometric grid compared to a hybrid grid that transitions from linear to geometric bin spacing, with the linear grid intended to limit numerical broadening caused by solving condensation. In parcel simulations, Lee et al. (2021) found that the introduction of turbulent-induced collision enhancement (discussed below) limited the effects of numerical broadening from condensation alone and reduced differences between the two grid choices relative to the absence of turbulent enhancement. In LES of a drizzling marine stratocumulus case study, they also found that the hybrid grid led to smaller drops and delayed onset of surface precipitation that agreed slightly better with radar observations of DSD moments, but that overall differences between simulations were small relative to differences with observations. For simplicity, we retain the geometric grid here, which appears reasonably resolved in comparison with observations (see Fig.10).

Specific details of microphysical processes including condensation, evaporation, and sedimentation are given in Ackerman et al. (1995). As in the bulk simulations, we initialize with a height-varying trimodal aerosol distribution (Fig. 1) with a hygroscopicity parameter of 0.4 for all bins. Activation of unactivated aerosol within a bin occurs when supersaturation exceeds the critical supersaturation calculated using the Köhler equilibrium relations. Upon activation, aerosol number is added to the smallest droplet size bin and aerosol mass is transferred to the corresponding droplet core mass bin. Collision-coalescence is performed using the exponential collection scheme of Bott (2000) to solve the stochastic collection equation with collision efficiencies from Hall (1980). Raindrop breakup follows from Hall (1980) and Low and List (1982). Complete evaporation occurs when the average volume of dissolved aerosol in a droplet bin exceeds the droplet volume. The corresponding droplet core mass concentration is then added back to the aerosol size distribution in a manner that conserves volume and number. Whereas this single-moment bin scheme is superior to the bulk scheme in conservation of aerosol mass and number, the bulk scheme use of prognostic number concentration in three aerosol modes with fixed modal parameters errs on the side of preserving aerosol dispersion. The baseline bin microphysics simulation is named BIN. We perform two additional sensitivity experiments to explore the role of turbulent enhancement of collision-coalescence. In the first experiment (BIN_TURB), the theoretical turbulent collision kernel from Ayala et al. (2008) is incorporated following the implementation described by Lee et al. (2021) that uses the explicitly calculated turbulent kinetic energy dissipation rate (ϵ) from the subgrid-scale (SGS) diffusion scheme and the collision efficiency enhancement from Wang and Grabowski (2009). Chen et al. (2018) found in large-eddy simulations of an Arctic mixed-phase cloud that the modeled turbulent broadening was narrower than observations of Doppler spectral width. They calculated that the total dissipation rate (i.e., numerical plus SGS) was a factor of 6 larger than the SGS dissipation rate and agreed better with observations. Moreover, Chen et al. (2018) revisited the study of Rémillard et al. (2017) who simulated a drizzling marine stratocumulus case and determined that DHARMA's total dissipation rate was ~ 3 times larger than the SGS dissipation rate, demonstrating that the magnitude of the ratio between the total and SGS dissipation rate is strongly case-dependent. Here, following the Chen et al. (2018) approach, we determined that the total mean dissipation rate is ~ 10 times larger than that computed by the SGS scheme, indicating that the more turbulent cloud environment simulated herein experiences substantially greater turbulent broadening than that calculated by the SGS scheme. Therefore, in a second experiment, ϵ is scaled by a factor of 10 (BIN_TURB_10X), which conceptually can be considered the most appropriate parameterization using the bin scheme with turbulent enhancement included.

Table 1. Simulation Details. Line colors devoted to each simulation in forthcoming plots are given in the rightmost column.

Simulation Name	Description	Line Color
CNTL	Setup described in text w/ Seifert and Beheng (2001) warm-rain formulation	Gray
NO_AC	As in CNTL, w/ autoconversion turned off	Red
FIXED_AERO	As in CNTL, w/ aerosol PSD fixed to cloud-base value throughout profile	Orange
FIXED_AERO_NO_AC	As in CNTL, w/ autoconversion turned off and aerosol PSD fixed to cloud-base value throughout profile	Purple
2X_AC	As in CNTL, w/ autoconversion efficiency scaled by a factor of 2	Blue
KK	As in CNTL, w/ Khairoutdinov and Kogan (2000) warm-rain formulation and an exponential size distribution	Green
BIN	As in CNTL, w/ size-resolved bin microphysics	Pink
BIN_TURB	As in BIN, w/ turbulent enhancement of collision-coalescence	Brown
BIN_TURB_10X	As in BIN_TURB, w/ turbulent enhancement scaled by a factor of 10	Light Blue

3.3 Thermal Identification and Tracking Framework

To investigate the role of entrainment in modulating profiles of N_d and to facilitate interpretation of cloud droplet production and evolution within their source elements (i.e., cumulus thermals), a thermal identification and tracking framework is employed. The algorithm used is described in Hernandez-Deckers and Sherwood (2016), based on an early version by Sherwood et al. (2013), and has been used in several recent studies to investigate the interacting roles of microphysics, aerosols, and convective dynamics (e.g., Hernandez-Deckers and Sherwood, 2018; Hernandez-Deckers et al., 2022; Matsui et al., 2024). A thorough description of the algorithm is provided in Hernandez-Deckers et al. (2022) and Matsui et al. (2024) so only a brief description is provided here. Rising volumes of cloudy air, above a minimum vertical velocity (w) threshold (1 m s^{-1}) and condensate threshold (0.01 g kg^{-1}), are tracked at high temporal frequency (1 minute). The minimum lifetime for thermals identified in this study is 3 minutes. Composite thermal statistics are centered around a thermal’s maximum ascent rate (which differs from the maximum w due to inhomogeneity across the thermal’s width resulting from toroidal circulations). The algorithm assumes a spherical shape, which Hernandez-Deckers and Sherwood (2016) showed to be a valid approximation compared to more plume-like structures. In transient convection simulations, Sherwood et al. (2013) and Hernandez-Deckers and Sherwood (2016) showed that thermals are generally short-lived ($\sim 4\text{-}5$ minutes) and rather small (relative to the horizontal mesh we use), which is consistent with thermals detected in the current study (not shown).

4 Results

4.1 Observed Profiles of N_d , R_{eff} , and ν_{eff}

Cloud-top N_d , R_{eff} , and ν_{eff} retrieved by the RSP are composited over temperatures warmer than 0 °C and sorted as a function of CTH (Fig. 5). In situ data from the Learjet FFSSP are also shown in Fig. 5 for points with liquid water content (LWC) > 0.1 g m⁻³. For the in situ data, R_{eff} and ν_{eff} are calculated using Eqs. 3 and 4, respectively, where the upper integration limit is the upper size threshold of the FFSSP ($r = 25 \mu\text{m}$).

Median cloud-top N_d decreases from $\sim 500 \text{ cm}^{-3}$ for CTHs between 0 and 1 km AGL to $\sim 100 \text{ cm}^{-3}$ for CTHs between 4 and 5 km AGL (Fig. 5a). Cloud-top R_{eff} follows an inverse relationship with N_d , increasing from $\sim 5 \mu\text{m}$ for CTHs below 1 km AGL to $\sim 15 \mu\text{m}$ for CTHs between 4 and 5 km AGL. Notably, R_{eff} values of up to $15 \mu\text{m}$ are conceptually aligned with the onset of precipitation at around this altitude (Rosenfeld and Gutman, 1994; Gerber, 1996; Andreae et al., 2004; Freud and Rosenfeld, 2012). Active precipitation in this case is consistent with the radar reflectivity transect shown in Fig. 2a with relatively high reflectivity values (> 20 dBZ) near cloud top and vertically continuous radar echoes reaching the surface. Median cloud-top ν_{eff} increases with increasing CTH in the lowest 2 km, decreases between 2 and 4 km, and increases above 4 km. Finer CTH binning (not shown) indicates a robust increase with CTH above 4 km, implying a broadening of the DSD at altitudes where precipitation starts to form.

In situ data from the FFSSP show smaller N_d relative to RSP for each altitude bin, while FFSSP R_{eff} is slightly larger than the RSP. Below 3 km, FFSSP ν_{eff} is constant (~ 0.075) and smaller than the RSP, but shows a more substantial increase with increasing height at higher altitudes, which is indicative of DSD broadening captured by the in situ measurements. We note that the largest size bin for the FFSSP is $25 \mu\text{m}$ in radius, while we are here assuming that the RSP DSD is truncated at a radius of $100 \mu\text{m}$. While it is possible to merge the FFSSP DSD with the 2D-S10 to get an extended distribution, doing so requires robust statistical averaging for stitching the DSDs together. Here, we chose to retain the 1-Hz native resolution of the FFSSP. This is justified with the knowledge that (1) N_d would be largely insensitive to the inclusion of larger particles, (2) R_{eff} would only increase by including an instrument with larger size bins, and (3) the in situ R_{eff} is already larger than the RSP. We also note that using a smaller LWC threshold of 0.01 g m⁻³ decreases the in situ N_d , shifting it further from the RSP. Despite magnitudinal differences, both instruments show similar trends with increasing CTH.

Discrepancies between RSP and in situ measurements may be due to several factors. At the foundation, RSP retrieves cloud top quantities while in situ measurements in this case include regions in or near the cloud core. The expectation is that N_d should be higher in cloud cores than at cloud top, but the opposite is seen here for reasons that are not readily apparent. For N_d , the RSP retrieval may be limited by assumptions in Eq. 5, which is derived from an adiabatic cloud model (e.g., Grosvenor et al., 2018). This method assumes that N_d is constant with height for a given cloud profile and that LWC increases linearly with height as a constant fraction of its adiabatic value (f_{ad}). The presented RSP-retrieved N_d assumes $f_{\text{ad}} = 0.8$, which may be too high for these cumulus clouds that are subject to substantial entrainment. However, f_{ad} can also be highly variable in addition to its uncertainty. A lower value of f_{ad} would decrease N_d , which scales with the square root of f_{ad} . Moreover, errors in R_{eff} propagate to errors in N_d that scale with the power of -5/2. Therefore, slightly lower RSP R_{eff} relative to in situ is conceptually consistent




Figure 5. Profiles of drop (a) number concentration (N_d), (b) effective radius (R_{eff}), and (c) effective variance (ν_{eff}) as a function of cloud top height (CTH) bins (bin width = 1 km) from the RSP (grays) and as a function of in-situ altitude bins from the FFSSP (greens). Boxes show the interquartile range and whiskers show the 10th and 90th percentiles. The RSP and in situ data are offset from the center of the 1-km bins with RSP on top and in situ on bottom. Circle markers with lines indicate the median and are placed at the center of the height bin for both instruments.

with higher RSP N_d . The in situ measurements, on the other hand, are likely significantly undersampled due to a small number of subjective cloud transects in highly variable cloud environments, and may contain poorly quantified instrumental biases of their own. Nonetheless, we accept these uncertainties here to explore the ability for simulations to produce profiles of N_d and R_{eff} that bound the RSP retrievals and in situ measurements.

385 4.2 Simulated Cloud System Evolution

We represent the simulated cloud-system evolution via time-height series of domain-averaged in-cloud cloud droplet number concentrations (N_c ; Fig. 6), which excludes the rain species in order to better emphasize the evolution of cloud droplets only. Clouds initiate within the first 30 minutes and maintain a cloud base height (CBH) of ~ 0.5 km AGL throughout the entirety of the simulation. Cloud tops grow monotonically from ~ 2 km initially to ~ 6 km at ~ 10 h for CNTL. Time series of rain water path (RWP) are also shown in Fig. 6, where rain onset occurs at ~ 6 h in CNTL. As discussed later, the simulated system is composed of numerous thermals that successively reach higher altitudes. After the onset of precipitation in CNTL, cold pools form such that the system begins to cluster with relatively more vigorous convection (not shown).

Cloud droplet number concentrations decrease with height in CNTL from ~ 400 cm^{-3} near cloud base down to under 50 cm^{-3} near cloud top in the last 3 hours. In comparison, N_c in NO_AC (i.e., no rain formation, Fig. 6d) decreases with height to ~ 150 cm^{-3} near the end of the simulation, indicating the role of collision-coalescence in scavenging N_c in CNTL. The FIXED_AERO simulation (Fig. 6b) realizes higher N_c throughout the entire profile, which results mainly from the 3rd and largest mode of the trimodal N_a distribution (Fig. 1). The largest mode is activated most frequently (not shown), and by forcing a constant profile with a value equal to that at cloud base (~ 650 cm^{-3} ; Fig. 1a) instead of decreasing with altitude, entrainment of these aerosols at high altitudes induces more secondary activation. This also acts to slightly delay higher RWPs relative to CNTL and leads to slightly weaker N_c scavenging. The FIXED_AERO_NO_AC simulation (Fig. 6d) represents the most extreme difference relative to CNTL, with N_c only decreasing by ~ 150 - 200 cm^{-3} between cloud base and cloud top near the end of the simulation, and even more modest decreases at times prior. The KK simulation experiences an earlier onset of precipitation that realizes significantly reduced N_c by hour 12 (Fig. 6d). The KK scheme has a more aggressive accretion process than the Seifert and Beheng (2001) scheme (Stevens and Seifert, 2008) and this experiment uses an exponential size distribution (as opposed to a gamma with a shape parameter of 3 as in all other experiments). The combination of these parameter choices leads to reduced RWP relative to CNTL and larger surface precipitation rates (not shown). Likewise, precipitation onset occurs earlier in 2X_AC (Fig. 6f) when autoconversion is scaled higher, which leads to the most significant scavenging of N_c by precipitation and a sharper decay of the system.

For the purpose of discussing system evolution using the bin scheme in comparison to the bulk scheme, liquid water in the bin scheme is arbitrarily separated using a radius threshold of 25 μm to categorize as cloud and rain species. However, we note that further analysis herein will make no such separation (unless otherwise stated). There are notable differences between the BIN experiment (Fig. 6g) and CNTL. Cloud droplet number concentrations are higher in BIN throughout the entire profile, indicating more efficient aerosol activation in BIN relative to CNTL. Turbulent enhancement of collision-coalescence (BIN_TURB; Fig. 6h) results in slightly earlier precipitation onset that decreases N_c slightly above 4 km near the end of the




Figure 6. Time-height series of in-cloud domain average cloud droplet number concentration (N_c) for the (a) CNTL, (b) FIXED_AERO, (c) KK, (d) NO_AC, (e) FIXED_AERO_NO_AC, (f) 2X_AC, (g) BIN, (h) BIN_TURB, and (i) BIN_TURB_10X simulations. Blue lines (right ordinates) show the rain water path (RWP) time series.

415 simulation. Scaling turbulent enhancement by a factor of 10 (BIN_TURB_10X; Fig. 6i) enhances precipitation further and scavenges smaller drops more efficiently.

4.3 Comparison of Simulations and RSP Retrievals

Simulated CTH is calculated by integrating optical depth (τ) from the top of the domain downward until it exceeds a threshold value of 1 (to match the assumed τ threshold of RSP), where $\tau(z)$ is calculated following, for example, Hansen and Travis
420 (1974) and Stephens (1978):

$$\tau(z) = \int_z^{z+\Delta z} \beta_{\text{ext}} dz = \int_z^{z+\Delta z} \int_0^\infty \pi Q_{\text{ext}} r^2 \frac{dn}{dr} dr dz \quad (7)$$

where β_{ext} is the extinction coefficient, Q_{ext} is the dimensionless extinction efficiency (assumed to be 2), and dz is the height difference across a given level. For the bulk scheme, $\frac{dn}{dr}$ (also used for R_{eff} and ν_{eff}) is reconstructed using the gamma distribution parameters and includes contributions from both cloud and rain species. The range of drop radii is chosen to correspond to the 50 bins used by the bin scheme (see Section 3.2.2). Drop number concentration is the sum of the cloud
425 and rain species in the bulk scheme. Effective radius and variance are calculated following Eqs. 3 and 4, respectively, and are integrated to an upper size limit of $r = 100 \mu\text{m}$ —the assumed size threshold for the RSP. Implications for this truncation size are discussed below.

Cloud-top N_d , R_{eff} , and ν_{eff} are then calculated using an extinction-weighting following:

$$N_d^{\text{top}} = \frac{\int_{z(\tau>0)}^{z(\tau\geq 1)} \beta_{\text{ext}} \int_0^{100\mu\text{m}} \frac{dn}{dr} dr dz}{\int_{z(\tau>0)}^{z(\tau\geq 1)} \beta_{\text{ext}} dz} \quad (8)$$

$$R_{\text{eff}}^{\text{top}} = \frac{\int_{z(\tau>0)}^{z(\tau\geq 1)} \beta_{\text{ext}} \int_0^{100\mu\text{m}} \pi r^3 \frac{dn}{dr} dr dz}{\int_{z(\tau>0)}^{z(\tau\geq 1)} \beta_{\text{ext}} \int_0^{100\mu\text{m}} \pi r^2 \frac{dn}{dr} dr dz} \cdot \frac{\int_{z(\tau>0)}^{z(\tau\geq 1)} \beta_{\text{ext}} dz}{\int_{z(\tau>0)}^{z(\tau\geq 1)} \beta_{\text{ext}} dz}. \quad (9)$$

$$\nu_{\text{eff}}^{\text{top}} = \frac{\int_{z(\tau>0)}^{z(\tau\geq 1)} \beta_{\text{ext}} \int_0^{100\mu\text{m}} (r - R_{\text{eff}})^2 \pi r^2 \frac{dn}{dr} dr dz}{\int_{z(\tau>0)}^{z(\tau\geq 1)} \beta_{\text{ext}} \int_0^{100\mu\text{m}} \pi r^2 \frac{dn}{dr} dr dz} \cdot \frac{\int_{z(\tau>0)}^{z(\tau\geq 1)} \beta_{\text{ext}} dz}{\int_{z(\tau>0)}^{z(\tau\geq 1)} \beta_{\text{ext}} dz}. \quad (10)$$

where R_{eff} in Eq. 9 is the local grid point value integrated to a size limit of $r = 100 \mu\text{m}$. Equations 8-10 represent the
 430 weighting between diffuse cloud top and a typical step increase in optical depth across the layer we define explicitly as cloud
 top (i.e., when accumulated τ exceeds a value of 1 from domain-top), which is consistent with the assumed RSP retrievals
 (e.g., Alexandrov et al., 2012a).

Profile distributions of cloud-top N_d , R_{eff} , and ν_{eff} for the CNTL and BIN_TURB_10X simulations are shown in Fig. 7.
 The latter is chosen as it represents the most appropriate implementation of turbulence-enhanced collision-coalescence with
 435 the DHARMA model; it also exhibits the most efficient precipitation formation among the three bin experiments. Cloud-top
 statistics are accumulated across the last three hours of the simulation to capture the effect of precipitation that was active in
 the observed system (see Fig. 2a), and to roughly represent flight timing that corresponds to initialization time of the NU-WRF
 mesoscale simulation used for DHARMA initial conditions. Median observed in situ values are also shown as green lines,
 along with simulated "in situ" values in orange (i.e., conditioned only on "cloudy" grid points where $\text{LWC} > 0.1 \text{ g m}^{-3}$, as for
 440 the FFSSP). For the bulk scheme, these simulated "in situ" values include only the cloud species in order to mimic observed
 subjective sampling of cloud cores (i.e., minimizing influence from rain-dominant grid points). Similarly for the bin scheme,
 these lines represent only cloudy grid points where the total liquid drop number concentration exceeds 1 cm^{-3} (based on the
 minimum N_d measured by the FFSSP).

For both CNTL and BIN_TURB_10X, simulated cloud-top N_d (Fig. 7a,b) decreases monotonically with increasing CTH
 445 qualitatively similarly to RSP but values are significantly lower than retrieved. Consistent with the expectation for N_d to be
 higher in cloud cores than at cloud top, simulated "in situ" median N_d is higher than simulated cloud-top N_d by up to a factor
 of ~ 2 for CNTL and by up to a factor of 8 for some mid-level altitude bins in BIN_TURB_10X, converging together with "in
 situ" measurements only at the highest altitudes where core regions are not far from cloud tops.

Simulated cloud-top R_{eff} agrees better with observations in general (Fig. 7c,d). Cloud-top R_{eff} in CNTL is slightly smaller
 450 than RSP retrievals at most altitudes whereas BIN_TURB_10X generally agrees well with RSP medians at all altitudes, and
 both schemes appropriately increase cloud-top R_{eff} with increasing CTH. Simulated "in situ" R_{eff} in CNTL is larger than cloud-
 top and also agrees fairly well with FFSSP R_{eff} . These results imply that cloud-top R_{eff} is better constrained by RSP than N_d
 for the case studied here.

Finally, cloud-top ν_{eff} becomes much larger with increasing height above 3 km in both simulations compared to RSP
 455 (Fig. 7e,f), which agrees both qualitatively and quantitatively with the profiles shown by Alexandrov et al. (2020, their Fig. 6).
 By contrast, CNTL produces a constant profile for simulated "in situ" ν_{eff} (cloud species only) ~ 0.075 , which is structurally
 required by using a constant shape parameter in the bulk scheme (i.e., any variability in cloud-top ν_{eff} from the prescribed
 cloud-species value arises due to the coexistence of cloud and rain species in contributing grid cells, which occurs often near
 cloud top). For BIN_TURB_10X, "in situ" ν_{eff} increases slightly with altitude as seen in FFSSP observations but is offset a bit
 460 larger, indicating DSDs broader than observed. "In situ" ν_{eff} in BIN_TURB_10X is still smaller than the cloud-top values since
 this profile conditions grid points to omit precipitation-dominant grid cells, which was not a condition for cloud-top sampling.
 Overall, the differences between simulated cloud-top and "in situ" ν_{eff} imply that cloud-top identification using Eq. 10 is rather
 sensitive to drops in the precipitation size range, even at and near cloud top and to a greater degree at the highest altitudes




Figure 7. Profiles of cloud-top (a,b) drop number concentration, (c,d) effective radius, and (e,f) effective variance for the CNTL simulation (left column) and the BIN_TURB_10X simulation (right column) as a function of cloud top height. Green lines show the median values from FFSSP in situ measurements. Orange lines show the simulated median "in situ" values (conditioned only on cloudy grid points). For cloud top quantities (with robust observational samples), boxes show the interquartile range and whiskers show the 10th and 90th percentiles. Distributions from simulations are across the entire domain for the last 3 h of the simulation. The height bin width is 1 km with the RSP and simulations offset relative to the mid-bin for visibility.

where both N_d and R_{eff} appear reasonably well reproduced. Indeed, decreasing the size cut-off for cloud-top distributions in
 465 Eqs. 8-10 from $r = 100 \mu\text{m}$ to $50 \mu\text{m}$ significantly decreased the cloud-top ν_{eff} values for both CNTL and BIN_TURB_10X,
 but did not have a large impact on cloud-top N_d and R_{eff} (not shown).

Taking the results shown in Fig. 7 as a whole, differences between "in situ" and cloud top sampling of N_d and ν_{eff} emerges
 from this work as a factor that is beyond our scope to resolve. Simulation numerical deficiencies could be to blame, but previous
 work remains limited to date. For example, finite vertical grid spacing (100 m) may not capture a realistic transition of DSDs
 470 from cloud-core to cloud-top (e.g., Sato et al., 2018). However, we did not see any gross overestimation of R_{eff} compared
 with either in situ observations or cloud top retrievals (Sato et al., 2018, cf. their Fig. 11). Related to this, numerical diffusion
 due to vertical advection and spurious evaporation of cloud droplets near cloud edges in these Eulerian schemes may be a
 source for low-biased cloud-top N_d , as documented by Chandrakar et al. (2022), especially for the bin scheme. However, that
 study did not include a cloud-top sampling approach; at the highest altitudes (which are near cloud tops by definition), bin
 475 and Lagrangian N_d largely converge (Chandrakar et al., 2022, cf. their Fig. 12). Comparison of retrieval-dependent cloud-top
 sampling (comparable to RSP and PACE measurements) and in-cloud sampling across bulk, bin, and Lagrangian schemes in a
 future model intercomparison study such as the ICMW CAMP²Ex case would be helpful to establish the roles of microphysics
 and dynamics schemes from both structural and numerical standpoints. We note that results could also depend on the case
 set-up insofar as it captures the progressive lifecycles of multiple cumulus towers and their decaying outflow, reflected in
 480 the RSP retrieval statistics here. In CNTL, a doubling of domain size did not produce quantitatively different results (not
 shown), suggesting that the frequency of cold pool collisions driving low-level updrafts and cloud-base activation is not likely
 a limitation in the current setup. On the other hand, the larger differences between "in situ" and cloud-top N_d and ν_{eff} simulated
 in BIN_TURB_10X versus CNTL at some elevations could be attributable to scheme-dependent structural differences in the
 outflow and decay of early cumulus towers that failed to reach the highest altitudes. We finally note potential limitations on
 485 the observational side, whether it be a breakdown in assumptions of the RSP-retrieved N_d , sparse in situ samplings, or both, as
 discussed in Section 4.1.

Median profiles of cloud-top N_d , R_{eff} , and ν_{eff} for all simulations are shown in Fig. 8, with the inclusion of N_d in units of
 number mixing ratio (kg^{-1}) in Fig. 8b,f. This air density-dependent conversion is performed to control for the impacts of dilu-
 tion by expansion, which Morrison et al. (2022) showed to account for 41% of N_d reduction between 4 and 9 km in a high-based
 490 congestus simulation. In doing so, we isolate the relative impacts on N_d via collision-coalescence and aerosol profile repre-
 sentation using the sensitivity experiments, with any remaining changes in the profile attributed to dilution by entrainment or,
 conversely, subsequent activation of cloud droplets above cloud base. For the bulk simulations (Fig. 8b), cloud-top N_d follows
 an intuitive pattern. The 2X_AC and KK simulations produce a sharper reduction of N_d with height due to more aggressive
 precipitation formation. Conversely, the NO_AC simulation produces a much more gradual decrease due to no precipitation,
 495 while FIXED_AERO produces a shallower slope due to less efficient precipitation owing to entrainment of greater concentra-
 tions of larger (more readily activated) aerosols at higher altitudes. At the extreme, the FIXED_AERO_NO_AC profile shows
 no significant reduction of N_d at all, and an increase with height above 3.5 km on a mass mixing ratio basis. There is a ubiqui-
 tous low bias of simulated cloud-top N_d relative to RSP, as discussed above, including at the highest elevations where CNTL




Figure 8. Profiles of median (a,e) N_d in units of cm^{-3} , (b,f) N_d in units of kg^{-1} , (c,g) R_{eff} , and (d,h) ν_{eff} as a function of cloud top height for the RSP (solid black), FFSSP in situ measurements (dashed black), bulk simulations (top row), and bin simulations (bottom row).

and BIN_TURB_10X cloud-top N_d converge to both observed and simulated "in situ" N_d . Profiles of cloud-top R_{eff} (Fig. 8c) for the sensitivity simulations generally follow an inverse relationship from N_d relative to CNTL. That is, KK and 2X_AC produce larger R_{eff} while NO_AC, FIXED_AERO, and FIXED_AERO_NO_AC produce smaller R_{eff} compared to CNTL. For cloud-top ν_{eff} , the three simulations without autoconversion produce a constant vertical profile because they are limited to the cloud species' fixed shape parameter, while the simulations with more aggressive precipitation formation produce larger ν_{eff} . All of the bin simulations produce lower cloud-top N_d compared to RSP but similar cloud-top R_{eff} , with BIN_TURB_10X agreeing best, consistent with Chandrakar et al. (2024) demonstrating the importance of turbulence-broadening in congestus simulations. Finally, cloud-top ν_{eff} is high-biased in all bin simulations relative to RSP. In the next section, we utilize full in situ size distributions to further evaluate the results shown here.

Table 2. Thermodynamic, kinematic, and microphysical description of four selected cloud passes used to construct composite drop size distributions (DSDs). Cloud passes are defined as contiguous segments with LWC $> 0.1 \text{ g m}^{-3}$. Values listed for DHARMA are for the CNTL simulation and are from the mean DSD conditioned on average vertical velocity (see text). ν_{eff} is discussed in text because of its sensitivity to size thresholding. The maximum vertical velocity for CP 3 is the minimum downdraft speed. Individual DSDs from each instrument and a related discussion is provided in Appendix D.

CP	Transect Length [km]	Sample Size	Temp. [$^{\circ}\text{C}$]			w [m s^{-1}]		LWC [g m^{-3}]		R_{eff} [μm]		$R_{\text{eff}} < 100 \mu\text{m}$ [μm]		N_d [cm^{-3}]	
			Avg.	Min.	Max.	Avg.	Max	Avg.	Max	In Situ	DHARMA	In Situ	DHARMA	In Situ	DHARMA
1	1.9	13	19.41	19.0	19.8	0.76	1.6	0.25	0.5	8.1	8.1	7.4	7.2	348	290
2	12.3	79	17.45	15.0	19.2	1.38	4.8	0.65	1.4	16.7	9.0	8.9	8.4	384	326
3	1.0	6	7.28	5.9	8.9	-2.82	-6.4	1.15	1.8	18.5	17.1	15.2	10.3	128	99
4	1.66	11	1.04	0.3	2.0	4.15	9.2	0.62	0.9	55.2	26.4	19.1	15.9	52	92

4.4 Comparison of Simulated and In Situ DSDs

In situ DSDs are analyzed using a methodology that composites instrument measurements of particle size across contiguous horizontal transects with LWC exceeding 0.1 g m^{-3} measured by the Nevzorov hotwire probe, referred to as "cloud passes". Cloud passes were chosen to span a temperature range from cloud base to the 0°C level with transects long enough to obtain robust sample sizes. A thermodynamic, kinematic, and microphysical summary of the four selected cloud passes is given in Table 2. The first cloud pass (hereafter CP) was performed in a relatively weak updraft a few hundred meters above cloud base at 19.41°C with a transect length of $\sim 2 \text{ km}$. The second CP was a long ascent ($\sim 12 \text{ km}$ horizontally) through a moderate updraft cluster that spanned a temperature range from $15\text{--}19^{\circ}\text{C}$ (mean temperature of 17.45°C). The third CP at 7.28°C was performed $\sim 300 \text{ m}$ below cloud top in a downdraft with a 1-km long transect. Finally, the fourth CP was performed in a relatively strong updraft (maximum vertical velocity, w , of 9.2 m s^{-1}) at 1.04°C with a transect length of $\sim 1.5 \text{ km}$. Composite DSDs are constructed by stitching together the FFSSP, 2D-S10, and HVPS probes following the methodology outlined in Appendix D. For each size bin of a given instrument, the composite DSD is averaged across each 1-Hz sample within the CP before stitching. This methodology yields four continuous size distributions in varying thermodynamic and kinematic environments that are used as an observational target for the simulations. Each in situ sample should be considered as representative of one single CP realization from the simulations, and thus the goal herein is to test whether a given observed CP DSD falls within the range of potentially encountered CP DSDs. Uncertainties associated with performing composite DSDs are further discussed in Appendix D.

For simulations, we focus only on CNTL and BIN_TURB_10X. Simulated CPs are identified at the temperature level of each observed CP using the same LWC thresholding and are selected between hours 9-12. For each cloud pass, the continuous size distribution (including both cloud and rain species for CNTL) is averaged across all grid points within the CP to yield a single sample. Drop size distributions are dynamically constrained by conditioning CPs where the average CP w is within 50% of the observed average CP w . Tests were performed with no conditioning and with conditioning on maximum CP w ,




Figure 9. Drop size distributions (DSD) from the Learjet (light blue), individual cloud passes from CNTL (black), and the mean DSD from CNTL (red) for cloud passes identified at temperatures of (a) 19.41 °C, (b) 17.45 °C, (c) 7.28 °C, and (d) 1.04 °C. Simulated cloud passes are conditioned to be within 50% of the observed cloud-pass-average vertical velocity. Sample sizes represent the number of simulated cloud passes for a given temperature level. Cumulative integrations are shown for (e)-(h) N_d , (i)-(l) R_{eff} , and (m)-(p) v_{eff} . The vertical dashed blue line indicates the assumed threshold for RSP retrievals.

530 yielding negligible qualitative differences but a narrowing of simulated CP DSD range that motivated including the dynamical condition. Drop size distributions for CNTL are shown in Fig. 9a-d (mean DSD in red, individual CP DSDs in transparent black) along with observed DSDs (light blue). The CNTL simulation produces the two prominent size modes at $\sim 10 \mu\text{m}$ (so-called "cloud" mode) and $\sim 0.4\text{-}0.5 \text{ mm}$ ("rain" mode) with varying degrees of accuracy. The most obvious discrepancy evident in all CNTL CP samples is their more pronounced bimodality and a relative dearth of N_d in the size range between $\sim 50\text{-}200$
535 μm (what may be considered a "drizzle" mode), which would be difficult to reproduce in CNTL given the structural separation of two species in the bulk scheme. At the warmer temperatures, the peak of the mean cloud-mode N_d is consistently smaller than observed. Overall, the CNTL simulation appropriately captures the observed broadening of the DSD with decreasing temperature (increasing height) demonstrated in Reid et al. (2023).

To better interpret how DSD structure impacts computations of N_d , R_{eff} , and ν_{eff} , the cumulative integration of these quantities are shown for each cloud pass in the bottom three rows of Fig. 9. A vertical dashed line is placed at a diameter of $200 \mu\text{m}$ to visualize the assumed size threshold for cloud-top retrievals. For all temperatures besides the coldest, the mean simulated DSD produces smaller N_d than observed (Fig. 9e-g), consistent with cloud-top comparisons in Fig. 7a, while the individual DSDs encompass the observed DSD. At the coldest temperature, the mean simulated cumulatively integrated N_d is smaller than observed, but the mean is skewed, evident by the individual DSDs clustering around the observed DSD. In both simulations
545 and observations, the cumulatively integrated N_d is very sensitive to a narrow diameter range that increases from $\sim 5\text{-}20 \mu\text{m}$ at the warmest temperature to $\sim 10\text{-}40 \mu\text{m}$ for the coldest, broadest DSD, completely saturating at larger sizes.

The mean cumulatively integrated R_{eff} is captured very well by CNTL for the two warmest temperatures (Fig. 9i-j), but is lower than observed for the colder temperatures, which is consistent with cloud-top comparisons (Fig. 7c). The individual DSDs again bound the observed DSD, albeit without ever reproducing the observed pattern of increase with diameter at the colder
550 temperatures. Cumulatively integrated R_{eff} relevant to cloud-top sampling is most sensitive to a diameter range between the lower size limit up to $\sim 30 \mu\text{m}$ for the two warmer temperatures. The relatively broader simulated DSDs at colder temperatures show a sensitivity for the "cloud mode" up to $\sim 50 \mu\text{m}$, with an additional abrupt increase for sizes larger than $200 \mu\text{m}$ as the precipitation mode grows. Interestingly, there is little size contribution to R_{eff} within the dearth of N_d in the drizzle size range ($50\text{-}200 \mu\text{m}$) even for the observed DSD, implying that besides for the coldest temperature, the source size region for biases
555 in R_{eff} relative to RSP originate largely within the cloud mode. Therefore, while N_d is always insensitive to truncation at the size threshold for RSP retrievals in this case, integrated R_{eff} can be sensitive to the truncation threshold when the DSDs are sufficiently broad.

Finally, cumulatively integrated ν_{eff} is shown in Fig. 9m-p. Except at the coldest temperature, the mean simulated ν_{eff} is generally larger than observed for diameters below $200 \mu\text{m}$, and the biases originate at relatively small sizes. However, at the
560 two warmest temperatures, the observed ν_{eff} is bounded by the cluster of simulated ν_{eff} within the cloud mode, where the saturated value up to $200 \mu\text{m}$ aligns with that shown for the simulated "in situ" profiles in Fig. 7 (i.e., when the precipitation mode is neglected). At colder temperatures, ν_{eff} is remarkably volatile around the size threshold of $200 \mu\text{m}$ and increases exponentially at larger sizes or even at values smaller than $200 \mu\text{m}$ for the colder temperatures. The demonstrated increase of ν_{eff} with height is consistent with the cloud-top profiles shown in Fig. 7 and implies that cloud-top representations of ν_{eff}




Figure 10. As in Fig. 9, but for the BIN_TURB_10X simulation.

565 that are comparable to RSP could be sensitive to the structure of the individual underlying DSDs if they contain drizzle mode drops.

570 An identical CP analysis is performed for the BIN_TURB_10X simulation (Fig. 10). The bin-scheme DSDs are distinctly more bimodal at warmer temperatures compared to the in situ DSDs, quite similar to the bulk scheme DSDs. A persistent bias across all temperature levels is the bin scheme producing a larger number of cloud droplets smaller than $\sim 10 \mu\text{m}$, particularly at warmer temperatures. We note that this extension toward the smallest size bins is also visible in Lagrangian particle-based simulations of a CAMP²Ex case in Chandrakar et al. (2024) (see their Fig. 3). At the coldest temperature (Fig. 10d), the bin scheme notably produces a more continuous transition from smaller to larger sizes, as in observations, due to the scheme's freedom from parametric constraints. The cumulatively integrated N_d for the bin scheme captures the enhanced influence of sizes below $10 \mu\text{m}$ for the warmest temperatures and saturates at sizes $\sim 20\text{-}30 \mu\text{m}$, as in the bulk scheme and observations.

575 Cumulatively integrated R_{eff} agrees very well with in situ DSDs at warmer temperatures and better represents observed R_{eff} than the bulk scheme at colder temperatures, consistent with cloud-top comparisons. As with the bulk scheme, cumulatively integrated R_{eff} is largely insensitive to the drizzle mode besides for the coldest and broadest DSDs. Finally, cumulatively integrated ν_{eff} shows that DSDs for all temperatures tend to be broader than observed, and that this bias originates in the cloud mode for sizes $< 30 \mu\text{m}$. Similarly to the bulk scheme, ν_{eff} is very sensitive to the size thresholding for which ν_{eff} is
580 calculated, increasing exponentially for diameters $> 200 \mu\text{m}$ at the warmest temperatures and for diameters $> 30 \mu\text{m}$ at the coldest temperatures. This reflects the very large cloud-top ν_{eff} seen at higher altitudes in Fig. 7f when a size threshold of $r = 100 \mu\text{m}$ is used. Overall, in situ DSD evaluation shows that biases in N_d , R_{eff} , and ν_{eff} all originate at rather small sizes below $30 \mu\text{m}$, while R_{eff} and ν_{eff} can be sensitive to larger sizes if the DSD is sufficiently broad. Potential numerical and structural issues with each scheme, along with size thresholds determining what is observed by the RSP, can both contribute to biases of
585 all three variables at cloud top.

4.5 Thermal-based Evaluation

We next present a more objective, process-based investigation of DSD evolution at the source of cloud droplet production—cumulus thermals (hereafter referred to simply as thermals; Hernandez-Deckers et al., 2022; Matsui et al., 2024). While thermal microphysics properties can represent either in-cloud or cloud-top microphysics depending on the locations and lifecycle
590 stages, their successive evolution can provide a source mechanism for droplet activation and the drizzle process, eventually characterizing cloud-top microphysics in convective clouds. Thermals are identified and tracked between hours 9-12 of the simulation for the CNTL and FIXED_AERO_NO_AC simulations only using 1-min simulation output, with the latter simulation chosen to control for the effects of the height-resolved aerosol PSD and collision-coalescence on modulating the N_d profile. The CNTL simulation yielded 278 thermals while the FIXED_AERO_NO_AC simulation yielded 349 thermals. The mode of
595 thermal lifetime is 3 min, which is the minimum lifetime to be considered as a valid thermal, with a near-exponential decrease in thermal lifetime at larger values (not shown), which indicates that the majority of detected thermals are rather short-lived and is consistent with results from Hernandez-Deckers and Sherwood (2016, 2018). Thermals are composited in time relative to their maximum ascent rates (defined as time $t = 0$) and are normalized by their radii (R). The evolution of composite thermals for the CNTL simulation in the X-Z plane between times $t = -3$ min and $t = +3$ min (i.e., 3 time steps prior to and after the
600 central maximum ascent rate) is shown in Fig. 11. Of all thermals for each simulation, at least $\sim 20 \%$ are represented at times $t = -3$ min and $t = +3$ min (or in other words, at least 20% of thermals last 7 minutes or longer). The evolution of thermals in Fig. 11 are displayed by variables of vertical velocity (w), supersaturation (S), cloud water mass mixing ratio (q_c), cloud water number mixing ratio (N_c ; independent of the rain species), and rain water mass mixing ratio (q_r).

Thermals in CNTL realize central w of at least 10 m s^{-1} (Fig. 11a1-a7) and continuous supersaturations at the center of the
605 thermal throughout their lifetime (Fig. 11b1-b7). The evolution of cloud water mass (q_c) shows relatively larger cloud water contents near the beginning of the thermal lifetime, while near the end ($t = +3$ min, Fig. 11c7), there is a shearing of maximum regions of q_c towards the right side of the thermal. Consistent with the cloud-top evaluation, N_c decreases during thermal-centered evolution (Fig. 11d1-d7) and R_{eff} increases (not shown). Supersaturations persist at thermal center (Fig. 11b1-b7) and




Figure 11. Composite thermal evolution (left to right) for the CNTL simulation from 3 minutes prior to maximum ascent rate ($t = -3$ min, leftmost column) to 3 minutes after maximum ascent rate ($t = +3$ min, rightmost column), where maximum ascent rate is centered at $t = 0$ min. Composites are normalized by thermal radius (R) and are displayed in the X - Z plane. Variables are distributed by rows: (a) vertical velocity (w), (b) supersaturation (S), (c) cloud water mass mixing ratio (q_c), (d) cloud water number concentration mixing ratio (N_c), and (e) rain mass mixing ratio (q_r). Gray streamlines are perturbation flow and the black ring represents a normalized $R = 1$. Values in the top right corner of panel are thermal-averaged values for a given time step. The dashed line in supersaturation panels indicates the contour where $S = 0$.




Figure 12. As in Fig. 11 but for the FIXED_AERO_NO_AC simulation, with the exclusion of q_r .

actually increase near the end of thermal lifetime. The evolution of q_r (Fig. 11e1-e7) shows that rain is largely absent in the
610 initial time-step and reaches a maximum at $t = +3$ min with a structure indicating significant unloading of rain.

The same thermal evolution is shown for the FIXED_AERO_NO_AC simulation in Fig. 12, but excluding q_r evolution since autoconversion is neglected. Notably, vertical velocities and supersaturations are weaker in this simulation (Fig. 12a1-a7 and b1-bd7, respectively). The N_c evolution (Fig. 12d1-d7) shows a negligible change of N_c throughout the thermal's lifetime, with the clearest difference being the shift in the axis of maximum N_c due to shear.

615 These thermal evolutions imply that by controlling for the height-resolved aerosol profile as well as the impact of collision-coalescence on N_c throughout a thermal's lifetime, the thermal-averaged N_c is largely uniform during ascent. However, entrainment is expected to be active in this environment, so why does N_c appear to remain constant? This is investigated via vertical profiles of averaged thermal characteristics (Fig. 13) and thermal characteristics as a function of average fractional entrainment rate (ε , Fig. 14; see Hernandez-Deckers and Sherwood, 2016, for a description of how this is explicitly calculated). The CNTL
620 and FIXED_AERO_NO_AC simulations produce similar thermal radii between ~ 400 -600 m (Fig. 13a). Both simulations also show similar ε , which is expected, since ε tends to be inversely proportional to radius (Hernandez-Deckers and Sherwood, 2018). At higher altitudes, FIXED_AERO_NO_AC produces weaker average and maximum w (Fig. 13b,c), which is likely a combination of condensate loading in CNTL as well as the lack of cold pool development in FIXED_AERO_NO_AC prevent-




Figure 13. Profiles of thermal-averaged (a) radius (R), (b) average vertical velocity (w), (c) maximum w , (d) average fractional entrainment rate (ε), (e) cloud water mass mixing ratio (q_c), (f) cloud water number mixing ratio N_c , and (g) cloud water effective radius ($R_{\text{eff},c}$). Profiles are constructed by 0.5-km bins. In (e)-(g), domain-averages for cloudy grid points ($q_c > 0.01 \text{ g kg}^{-1}$) are shown as dotted lines and domain-averages for convective cloudy grid points ($q_c > 0.01 \text{ g kg}^{-1}$ and $w > 1 \text{ m s}^{-1}$) are shown as dashed lines. Shaded regions show the standard deviation from the mean.

ing clustering and organization that is realized in CNTL. Profiles of thermal-averaged q_c , N_c , and $R_{\text{eff},c}$ are shown in Fig. 13e-g, along with profile averages of cloudy grid points ($\text{LWC} > 0.01 \text{ g m}^{-3}$) and convective cloudy grid points ($\text{LWC} > 0.01 \text{ g m}^{-3}$ and $w > 1 \text{ m s}^{-1}$) independent of the thermal framework. No matter the definition, Fig. 13f shows that N_c in FIXED_AERO_NO_AC is consistently constant with height, similar to the relatively constant cloud-top N_d profile presented in Fig. 8b. This suggests that in the absence of a height-varying aerosol profile and collision-coalescence, the impact of entrainment on reducing N_c is offset by secondary droplet activation along cloud edges (i.e., entrainment of aerosols along the inflow branch of toroidal circulations)—a process which has been shown to be a relevant aerosol activation mechanism in prior studies (Morrison et al., 2022; Chandrakar et al., 2021).

Distributing thermal characteristics as a function of ε (Fig. 14) shows that all variables generally decrease with increasing ε except for the strongest ε bins, further implying that the effects of entrainment on diluting N_c in the profiles shown in Fig. 13




Figure 14. Thermal average properties as a function of fractional entrainment rate (ε). ε bin widths are $0.5 \times 10^{-3} \text{ m}^{-1}$. Shaded regions show the standard deviation from the mean and at least 5 thermals are required for a given ε bin.

is offset by secondary activation. The increase of q_c and N_c for FIXED_AERO_NO_AC for the strongest ε may be due to substantial entrainment of aerosol, though the robustness of this signal should be evaluated in more detail in future work. Interestingly, R_{eff} also decreases with increasing ε , consistent with a dominant effect of higher q_c where entrainment is weakest that dilutes substantially at higher rates.

The influence of entrainment on microphysical properties in Figs. 13 and 14 has implications for the development of convective microphysics schemes in large-scale models. In bulk mass-flux based convection parameterizations (e.g., Tiedtke, 1989; Gregory and Rowntree, 1990; Bechtold et al., 2008), entrainment is calculated for a single plume where a cloud ensemble is considered to be impacted equally by a given entrainment profile (e.g., Fig. 13). In spectral parameterizations (e.g. Arakawa and Schubert, 1974; Lin and Arakawa, 1997) that represent a variety of cloud types, each cloud type can experience varying degrees of entrainment (e.g., Fig. 14). Therefore, it is plausible to utilize a thermal-tracking framework with explicitly calculated entrainment as a means for linking convective microphysics to a given model's convection parameterization, which is discussed more in the following section.

5 Discussion

5.1 Choosing a Benchmark LES Cumulus Congestus Case for ESM Evaluation and Improvement

Neggers (2015) showed in an intercomparison of simulated marine stratocumulus-to-cumulus transitions that SCMs can provide a unique fingerprint of their ESM counterparts. For that cloud regime, such a conclusion suggests that the model state, whether it be SCM or ESM, was dominated by the boundary-layer behavior (or "fast physics" relative to the large-scale flow). In a similar manner, the case study presented herein can be used to investigate convective microphysics in ESMs that are driven by the convection scheme, whether diagnostically (see Elsaesser et al., 2017) or prognostically (see Song and Zhang, 2011). In this framework, simple tests regarding the structural implementation of convective microphysics may be explored to determine the leading factors of microphysical shortcomings in the congestus regime. Appropriate thermodynamic forcing, large-scale vertical motion conditions, and aerosol input for this case are provided as supplements for such testing in high-resolution models and SCMs. Furthermore, additional CAMP²Ex cases may be constructed with different aerosol loading and varying convective structures (e.g., Chandrakar et al., 2024). Indeed, Reid et al. (2023, see their Fig. 6f) show the vast range of aerosol conditions measured during CAMP²Ex. Moreover, the role of entrainment in modulating microphysical parameters (see Section 4.5) can be linked to spectral convection parameterizations. As a simple example, the National Aeronautics and Space Administration (NASA) Goddard Institute for Space studies (GISS) ModelE3 (GISS-ModelE3; Cesana et al., 2019, 2021) convection parameterization uses a two-plume model, with one weakly entraining plume and the other strongly entraining, which is not currently aerosol-aware. Thermal-based analysis such as that presented here can estimate the impact of entrainment for each plume on diluting N_d in the associated convective microphysics scheme, in addition to providing an observation-constrained foundation for adding aerosol-aware physics. Of course, results are expected to be case- and regime-specific, and scheme development is expected to be conducted with a range of diverse cases and regimes.

5.2 Translation to Space-based Platforms and Global Simulations

The Hyper-angular Rainbow Polarimeter (HARP-2; Martins et al., 2018) on PACE provides the opportunity to evaluate global distributions of cloud-top microphysical characteristics, extending beyond bi-spectral retrievals such as those provided by the Moderate Resolution Imaging Spectroradiometer (MODIS) on the Aqua and Terra satellites. Fu et al. (2022) performed an intercomparison of cloud-top R_{eff} during CAMP²Ex between MODIS, in situ measurements, and RSP retrievals using both the polarimetric method (as used herein) and its alternative bi-spectral method. They found a persistent high-bias in R_{eff} using bi-spectral methods by 4-7 μm in the median and demonstrated that cloud heterogeneity and 3D radiative transfer effects were at the source of the high-biased bi-spectral R_{eff} retrieval. Potential biases in R_{eff} can have important implications for evaluating ESMs that ubiquitously employ bulk microphysics schemes. For example, R_{eff} is used directly as input to space-based lidar simulators for calculating the particle backscatter coefficient (Chepfer et al., 2008; Cesana et al., 2021). Moreover, the direct use of R_{eff} in ESM radiation schemes will propagate error for comparison with satellite retrievals. Furthermore, just as RSP, HARP-2 has the potential of inferring full drop size distributions (without assuming a gamma shape) using the Rainbow Fourier Transform algorithm (Alexandrov et al., 2012b; Reid et al., 2023; Sinclair et al., 2021), which may provide more detailed information similar to that obtained from the in situ DSDs in this study. In addition to polarimetric cloud retrievals, PACE also provides detailed polarimetric aerosol retrievals (Hasekamp et al., 2019; Gao et al., 2023) which opens the opportunity to study congestus development globally as a function of aerosol loading and type. We note, however, that the polarimeters on PACE will provide retrievals on a spatial scale of about $5.2 \times 5.2 \text{ km}^2$ which is more than an order of magnitude coarser than the RSP observations used in this study. Understanding of the effect of increased footprint size could be advanced by using forward simulation approaches and model output from case studies such as this. On a related note, there is considerable room for improvement in adjusting the comparison of simulated microphysics with polarimetry platforms. For example, Section 4.4 demonstrated the sensitivity of R_{eff} and ν_{eff} to size thresholding assumed for the RSP, and we assumed a τ threshold of 1 to define cloud top. However, the size threshold observed by the RSP is likely dependent on the DSD being observed and its extinction characteristics. Simulators for airborne or spaceborne polarimeters should thus explore methods to determine the relevant (and potentially modular) size thresholds at which extinction saturates the retrieval.

6 Conclusions

We present analyses of a tropical cumulus congestus case study based on large-eddy simulations with observed trimodal aerosol size distribution profiles as input and evaluated against airborne polarimetric retrievals and in situ cloud microphysics measurements. Novel retrievals from the Research Scanning Polarimeter (RSP) provided a statistically robust continuous profile of cloud-top effective radius (R_{eff}) and effective variance (ν_{eff}), which used together allowed the derivation of cloud-top drop number concentration (N_d). Simulations were performed using both bulk and bin microphysics schemes, and the sensitivity of drop size distribution (DSD) evolution to the collision-coalescence process and the height-resolved aerosol profile was quantified. A thermal-tracking analysis framework was then used to objectively describe the microphysical evolution at the source

of cloud droplet production and quantify the effects of dilution via entrainment. The primary conclusions are summarized as follows:

- 700 – Polarimetric retrievals and in situ measurements of N_d and R_{eff} both indicate characteristic profiles of decreasing N_d and increasing R_{eff} with increasing cloud top height (CTH). By contrast, cloud-top ν_{eff} retrievals exhibit a weak minimum near 4 km whereas in situ measurements indicate monotonically increasing values up to the melting level.
- 705 – Both bulk and bin schemes reproduce a decrease of N_d and increase of R_{eff} with increasing CTH. Whereas R_{eff} as a function of height is also quite well reproduced by both schemes, cloud-top N_d is consistently lower than retrieved, "in situ" N_d is consistently higher than measured, and the observational data products do not exhibit the expected relationship (in situ values greater than cloud-top values) that is seen in the simulations. In the bulk scheme, median "in situ" ν_{eff} is uniform with height at a value similar to RSP retrievals following the fixed droplet spectral width in that scheme; in the bin scheme, greater structural freedom enables an increase with height similar to that observed in situ albeit with a consistent high bias. Differences throughout are likely attributable in some part to both uncertainties in the observational data products and potential numerical and structural issues in the simulations.
- 710 – Neglecting collision-coalescence in a sensitivity experiment increases median cloud-top N_d from the control experiment by nearly a factor of 3 circa 5 km, indicating a powerful control on the observed profile of N_d .
- 715 – Neglecting the height-variation of aerosol increases median cloud-top N_d by up to a factor of 2 at CTHs circa 5 km, reflecting the role of thermal entrainment transiting aerosol gradients aloft.
- 720 – Neglecting both collision-coalescence and the height-variation of aerosol produces a cloud-top N_d profile (per kg) that is nearly constant with height below circa 3.5 km and increases with height thereafter, indicating that entrainment alone cannot drive the observed trend.
- 725 – Comparison with in situ cloud pass DSDs shows that both schemes can generally bracket the observed DSDs, which should be considered as only a sparse sample of possible realized DSDs. Compared to in situ DSDs, the bulk scheme cannot capture the observed tail of the distribution, whereas the bin scheme effectively captures the continuum of particle sizes across the drop and drizzle modes (a feature that will be an important foundation for examining ice formation in follow-on work). Both schemes produce cloud modes that appear broader and less peaked than observed.
- 730 – Juxtaposing the cumulatively integrated N_d , R_{eff} , and ν_{eff} dependence on the DSD shape indicates that all integrated moments are, to first order, significantly sensitive to sizes $< 30 \mu\text{m}$. The latter two moments, especially ν_{eff} , can also be sensitive to larger sizes if the DSD is sufficiently broad.
- 735 – Comparing simulated DSDs at cloud top with RSP retrievals can be sensitive to the assumed truncation size, especially for ν_{eff} , and differences between RSP and in situ measurements highlight limitations in strongly constraining the case studied here.

- A thermal-tracking analysis framework demonstrates that neglecting both collision-coalescence and height-variation of aerosol leads to a nearly constant profile of thermal-averaged N_d (per kg) regardless of sampling approach, suggesting that the influence of entrainment on diluting thermal N_d is almost exactly balanced by secondary activation in the absence of aerosol loss processes in this case.

The case study presented here offers a benchmark framework for ESM simulations of tropical cumulus congestus, which are lacking relative to other convective modes that have been the focus of informative LES-SCM intercomparison projects (e.g., Siebesma et al., 2003; Sandu and Stevens, 2011; Neggers, 2015; Neggers et al., 2017; Vogelmann et al., 2015; Endo et al., 2015; Lin et al., 2015). Such a task is aided by supplying large-scale thermodynamics and vertical motion derived from mesoscale simulations of the presented case study as well as trimodal, lognormal aerosol distribution profiles, the combination of which can be identically used to drive SCM versions of ESMs. Use of this case as an observationally evaluated benchmark can contribute to informing development of convective microphysics schemes in large-scale models that currently exist in highly variable states of sophistication. This framework also lends itself to informing future global analysis of warm-phase microphysics evolution via the recently successfully launched space-borne polarimeters on the PACE satellite. In follow-on work with LES, this case offers a natural transition to the evaluation of ice-phase microphysics in tropical congestus, where establishing the realism of simulated drizzle-containing DSDs (e.g., Fig. 10c-d) is an important foundation for simulating secondary ice formation.

Code availability. Python plotting scripts used to make all figures can be found on the NASA Center for Climate Simulation (NCCS) data portal: <https://portal.nccs.nasa.gov/datashare/giss-camp2ex/> and at the following DOI: <Insert Zenodo DOI when review process is completed>. DHARMA source code is maintained at NASA GISS and can be made available upon request. The NASA-Unified WRF (NU-WRF) is maintained at NASA GSFC, and available for public use upon request (<https://nuwrf.gsfc.nasa.gov/>).

Data availability. The NASA Center for Climate Simulation (NCCS) data portal (<https://portal.nccs.nasa.gov/datashare/giss-camp2ex/>) archives the following data: 3D output from the CNTL and BIN_TURB_10X simulations (other simulations can be made available upon request), input aerosol profiles and distribution parameters, thermodynamic and large-scale vertical motion forcing files, NU-WRF namelists needed for reproduction of the mesoscale simulation, files containing time-height series of domain-mean variables for all simulations presented herein, and thermal data files. These files (sans 3D output) are also available at the following DOI for permanent storage: <Insert Zenodo DOI when review process is completed>. Aircraft data from the Research Scanning Polarimeter (RSP), 3rd Generation Advanced Precipitation Radar (APR-3), in situ cloud probes, and Fast Integrated Mobility Spectrometer (FIMS) are available on the CAMP2Ex data repository (<https://www-air.larc.nasa.gov/cgi-bin/ArcView/camp2ex>).

Appendix A: NU-WRF Simulation Setup

A mesoscale simulation used for harvesting large-scale thermodynamic conditions and vertical motion for LES initialization is performed using the NASA Unified Weather Research and Forecasting (NU-WRF; Peters-Lidard et al., 2015) model with the v3.9.1.1 WRF dynamical core. The NU-WRF simulation was initialized with the National Centers for Environmental Prediction (NCEP) Global Data Assimilation System (GDAS) FNL (Final) operational global analysis and forecast dataset with a spatial resolution of $0.25^\circ \times 0.25^\circ$ and a temporal resolution of 6 h (National Centers for Environmental Prediction National Weather Service NOAA U.S. Department of Commerce, 2015). The simulation was integrated for 21 h from 12 UTC on 24 September 2019 through 9 UTC on 25 September. We use a one-way nested grid with an outer domain size of 2400 x 2400 km and a horizontal grid spacing (Δ_h) of 3 km and an inner domain size of 480 x 480 km with $\Delta_h = 600$ m. The outer and inner domains use a time step of 9 and 1.5 s, respectively, and each has 151 stretched vertical levels. We employ the National Severe Storms Laboratory (NSSL) microphysics scheme (Ziegler, 1985; Mansell et al., 2010; Mansell and Ziegler, 2013) with 6 double-moment hydrometeor species including cloud water, rain, cloud ice, snow, graupel, and hail. Trimodal, lognormal aerosol profiles are treated as described for the DHARMA simulations with activation following Abdul-Razzak and Ghan (2000) using the minimum supersaturation from Morrison and Grabowski (2008b). Other physics include the 2017 Goddard radiation package (Matsui et al., 2020), the Noah-MP land surface model (LSM; Niu et al., 2011; Yang et al., 2011), the Mellor-Yamada-Nakanishi-Niino (MYNN) 2.5 level turbulent kinetic energy (TKE) planetary boundary layer (PBL) scheme (Nakanishi and Niino, 2009), and the 2D horizontal Smagorinsky scheme for horizontal subgrid-scale diffusion (Skamarock et al., 2008).

Appendix B: Deriving Large-scale Vertical Motion

Large-scale vertical motion (w_{LS}) was evaluated by averaging vertical velocity across various subdomains of the NU-WRF simulation on a 600-m horizontal mesh (domain shown in Fig. 4). While w_{LS} was rather variable across the domain and time, a characteristic profile of ascent below ~ 5 km and subsidence above was apparent when averaging over a time period of 6 h that corresponded to flight timing (~ 0300 - 900 UTC). The magnitude of ascent in the lower troposphere ultimately modulated the acceleration of growing CTH throughout simulation integration, but a maximum of 2 cm s^{-1} is consistent with subdomain averages. The idealized profile shown in Fig. 4a proved sufficient and appropriate for realizing a simulation similar to the observed system evolution, as determined by the timing of CTH growth from the simulation initialization. As stated in the main text, we do not consider this profile to be well-constrained, but rather plausible, and note that derivations of w_{LS} are essential to appropriately represent the dynamical evolution of at least the congestus case studied here. Indeed, Fig. B1 shows the evolution of domain-mean LWC for the CNTL simulation and a simulation with no imposed w_{LS} . In the simulation without imposed w_{LS} (Fig. B1b), cloud tops initially grow to ~ 2 km and stagnate to a maximum of only ~ 2.5 km at the end of the simulation. Therefore, imposed w_{LS} is essential to realize a simulation similar to the observed system.




Figure B1. Time-height series of domain-mean liquid water content (LWC) for (a) the CNTL simulation and (b) a simulation with the same physics as CNTL but without imposed large-scale vertical motion.

Appendix C: Resolution Sensitivity Test

Resolution sensitivity was investigated by performing a simulation with identical physics to the CNTL simulation (which uses a horizontal mesh of 100 m) but with a horizontal mesh of 50 m. Time-height series of domain-maximum w , domain-mean LWC, and in-cloud domain-mean cloud-species number concentration (N_c) are shown in Fig. C1. Differences between the two resolutions are negligible for each property for the purposes of this study. Cloud-top R_{eff} and N_d were also evaluated relative to RSP (not shown) and the 50-m simulation was found to be indistinguishable from CNTL. This test demonstrates microphysical convergence for the CNTL simulation presented herein.

Appendix D: Composite In Situ DSDs

Composite observed in situ size distributions are constructed by stitching together instruments in various size ranges. These include the fast forward scattering spectrometer probe (FFSSP; 2-50 μm with bin width ranging from 1.5 to 4 μm), the 10 μm 2D-S (Stereo) optical array spectrometer (2D-S10; 10 μm - 3 mm with bin width ranging from 10 to 200 μm), and the high volume precipitation spectrometer (HVPS; 150 μm - 4.5 mm with bin width ranging from 150 μm to 3 mm). The size




Figure C1. Time height series of (a,d) domain-maximum vertical velocity (w), (b,e) domain-mean liquid water content (LWC), and (c,f) in-cloud domain-mean cloud-species droplet number concentration (N_c) for the CNTL simulation with a horizontal mesh of 100 m (top row) and a simulation with a horizontal mesh of 50 m (bottom row).

800 thresholds used to stitch the individual instrument DSDs is chosen based on convergence of overlapping size ranges between the instruments for a given statistical sample. As shown in Fig. D1, there is considerable uncertainty between overlapping sizes of the smallest size bins for a given instrument and the largest size bins for another instrument. For example, drop numbers in the smallest size bins of the 2D-S10 are 2 orders of magnitude lower than measured by the FFSSP at its largest size bins (Fig. D1a). However, convergence is realized at the upper limit of the FFSSP ($50 \mu\text{m}$) such that a continuous distribution
805 emerges. Furthermore, the size threshold is not consistent between different cloud passes and instead depends on the samples measured by the instrument. For example, the size threshold for stitching the 2D-S10 and HVPS is chosen at $250 \mu\text{m}$ for the 19.41°C cloud pass (Fig. D1a) but at $550 \mu\text{m}$ for the 1.04°C cloud pass (Fig. D1d). Ultimately, the selected cloud passes and size thresholds used to create composite DSDs are chosen to create the most realistic transition between instrument size regions while retaining physical characteristics of the DSD shape.




Figure D1. Observed drop size distributions (DSDs) for the 4 selected temperature levels discussed in the text. Size distributions from individual instruments are shown as colored lines and the composite DSDs from stitching together instruments are shown as black lines.

810 *Author contributions.* MWS and AMF conceptualized the study. MWS prepared simulations with the help of ASA and AMF (DHARMA) and TM (NU-WRF). Aerosol input data was prepared by QX and JW. Thermal tracking code and guidance was provided by DHD. BvD prepared RSP data and guided interpretation. PL prepared in situ cloud probe data and guided use. MWS performed formal analysis and was primary author of manuscript. All authors contributed to editing the final manuscript.

Competing interests. AMF is a member of the editorial board of Atmospheric Chemistry and Physics.

815 *Financial support.* This work was supported by the NASA Atmospheric Composition Campaign Data Analysis and Modeling Program, and the NASA Modeling, Analysis, and Prediction Program.

Acknowledgements. Computing support to perform simulations was provided the by NASA Center for Climate Simulations. This work was part of the NASA CAMP2EX campaign (10.5067/Suborbital/CAMP2EX2018/DATA001). We thank Gregory Elsaesser and Kuniaki Inoue for their helpful discussions on convective parameterization, convective microphysics structure, and development pathways. Pacific
820 Northwest National Laboratory is operated by Battelle for the 644 U.S. Department of Energy under Contract DE-AC05-76RLO1830.

References

- Abdul-Razzak, H. and Ghan, S. J.: A parameterization of aerosol activation 2. Multiple aerosol types, *Journal of Geophysical Research Atmospheres*, 105, 6837–6844, <https://doi.org/10.1029/1999JD901161>, 2000.
- Ackerman, A. S., Hobbs, P. V., and Toon, O. B.: A Model for Particle Microphysics, Turbulent Mixing, and Radiative Transfer in the
825 Stratocumulus-Topped Marine Boundary Layer and Comparisons with Measurements, *Journal of Atmospheric Sciences*, 52, 1204 – 1236, [https://doi.org/10.1175/1520-0469\(1995\)052<1204:AMFPMT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1995)052<1204:AMFPMT>2.0.CO;2), 1995.
- Ackerman, A. S., Toon, O. B., Stevens, D. E., Heymsfield, A. J., Ramanathan, V., and Welton, E. J.: Reduction of Tropical Cloudiness by Soot, *Science*, 288, 1042–1047, <https://doi.org/10.1126/SCIENCE.288.5468.1042>, 2000.
- Alexandrov, M. D., Cairns, B., Emde, C., Ackerman, A. S., and van Diedenhoven, B.: Accuracy assessments of cloud droplet size re-
830 trievals from polarized reflectance measurements by the research scanning polarimeter, *Remote Sensing of Environment*, 125, 92–111, <https://doi.org/10.1016/J.RSE.2012.07.012>, 2012a.
- Alexandrov, M. D., Cairns, B., and Mishchenko, M. I.: Rainbow Fourier transform, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 113, 2521–2535, <https://doi.org/10.1016/J.JQSRT.2012.03.025>, 2012b.
- Alexandrov, M. D., Cairns, B., Sinclair, K., Wasilewski, A. P., Ziemba, L., Crosbie, E., Moore, R., Hair, J., Scarino, A. J., Hu, Y., Stammes,
835 S., Shook, M. A., and Chen, G.: Retrievals of cloud droplet size from the research scanning polarimeter data: Validation using in situ measurements, *Remote Sensing of Environment*, 210, 76–95, <https://doi.org/10.1016/J.RSE.2018.03.005>, 2018.
- Alexandrov, M. D., Miller, D. J., Rajapakshe, C., Fridlind, A., van Diedenhoven, B., Cairns, B., Ackerman, A. S., and Zhang, Z.: Vertical profiles of droplet size distributions derived from cloud-side observations by the research scanning polarimeter: Tests on simulated data, *Atmospheric Research*, 239, 104 924, <https://doi.org/10.1016/J.ATMOSRES.2020.104924>, 2020.
- 840 Andreae, M. O., Rosenfeld, D., Artaxo, P., Costa, A. A., Frank, G. P., Longo, K. M., and Silva-Dias, M. A.: Smoking Rain Clouds over the Amazon, *Science*, 303, 1337–1342, https://doi.org/10.1126/SCIENCE.1092779/SUPPL_FILE/ANDREAE.SOM.PDF, 2004.
- Arakawa, A. and Schubert, W. H.: Interaction of a Cumulus Cloud Ensemble with the Large-Scale Environment, Part I, *Journal of the Atmospheric Sciences*, 31, 674–701, [https://doi.org/10.1175/1520-0469\(1974\)031](https://doi.org/10.1175/1520-0469(1974)031), 1974.
- Ayala, O., Rosa, B., and Wang, L. P.: Effects of turbulence on the geometric collision rate of sedimenting droplets. Part 2. Theory and
845 parameterization, *New Journal of Physics*, 10, 075 016, <https://doi.org/10.1088/1367-2630/10/7/075016>, 2008.
- Bechtold, P., Köhler, M., Jung, T., Doblas-Reyes, F., Leutbecher, M., Rodwell, M. J., Vitart, F., and Balsamo, G.: Advances in simulating atmospheric variability with the ECMWF model: From synoptic to decadal time-scales, *Quarterly Journal of the Royal Meteorological Society*, 134, 1337–1351, <https://doi.org/10.1002/QJ.289>, 2008.
- Behrenfeld, M. J., Moore, R. H., Hostetler, C. A., Graff, J., Gaube, P., Russell, L. M., Chen, G., Doney, S. C., Giovannoni, S., Liu, H., Proctor,
850 C., Bolaños, L. M., Baetge, N., Davie-Martin, C., Westberry, T. K., Bates, T. S., Bell, T. G., Bidle, K. D., Boss, E. S., Brooks, S. D., Cairns, B., Carlson, C., Halsey, K., Harvey, E. L., Hu, C., Karp-Boss, L., Kleb, M., Menden-Deuer, S., Morison, F., Quinn, P. K., Scarino, A. J., Anderson, B., Chowdhary, J., Crosbie, E., Ferrare, R., Hair, J. W., Hu, Y., Janz, S., Redemann, J., Saltzman, E., Shook, M., Siegel, D. A., Wisthaler, A., Martin, M. Y., and Ziemba, L.: The North Atlantic Aerosol and Marine Ecosystem Study (NAAMES): Science motive and mission overview, *Frontiers in Marine Science*, 6, 424 783, <https://doi.org/10.3389/FMARS.2019.00122/BIBTEX>, 2019.
- 855 Bott, A.: A Flux Method for the Numerical Solution of the Stochastic Collection Equation: Extension to Two-Dimensional Particle Distributions, *Journal of the Atmospheric Sciences*, 57, 284 – 294, [https://doi.org/10.1175/1520-0469\(2000\)057<0284:AFMFTN>2.0.CO;2](https://doi.org/10.1175/1520-0469(2000)057<0284:AFMFTN>2.0.CO;2), 2000.

Brenguier, J.-L., Pawlowska, H., Schüller, L., Preusker, R., Fischer, J., Fouquart, Y., Brenguier, J.-L., Pawlowska, H., Schüller, L., Preusker, R., Fischer, J., and Fouquart, Y.: Radiative Properties of Boundary Layer Clouds: Droplet Effective Radius versus Number Concentration., JAtS, 57, 803–821, [https://doi.org/10.1175/1520-0469\(2000\)057](https://doi.org/10.1175/1520-0469(2000)057), 2000.

Cairns, B., Russell, E. E., and Travis, L. D.: Research Scanning Polarimeter: calibration and ground-based measurements, <https://doi.org/10.1117/12.366329>, 3754, 186–196, <https://doi.org/10.1117/12.366329>, 1999.

Cesana, G., Del Genio, D. A., Ackerman, S. A., Kelley, M., Elsaesser, G., Fridlind, M. A., Cheng, Y., and Yao, M. S.: Evaluating models response of tropical low clouds to SST forcings using CALIPSO observations, Atmospheric Chemistry and Physics, 19, 2813–2832, <https://doi.org/10.5194/ACP-19-2813-2019>, 2019.

Cesana, G. V., Ackerman, A. S., Fridlind, A. M., Silber, I., and Kelley, M.: Snow Reconciles Observed and Simulated Phase Partitioning and Increases Cloud Feedback, Geophysical Research Letters, 48, e2021GL094 876, <https://doi.org/10.1029/2021GL094876>, 2021.

Chandrakar, K. K., Grabowski, W. W., Morrison, H., and Bryan, G. H.: Impact of Entrainment Mixing and Turbulent Fluctuations on Droplet Size Distributions in a Cumulus Cloud: An Investigation Using Lagrangian Microphysics with a Subgrid-Scale Model, Journal of the Atmospheric Sciences, 78, 2983–3005, <https://doi.org/10.1175/JAS-D-20-0281.1>, 2021.

Chandrakar, K. K., Morrison, H., Grabowski, W. W., and Bryan, G. H.: Comparison of Lagrangian Superdroplet and Eulerian Double-Moment Spectral Microphysics Schemes in Large-Eddy Simulations of an Isolated Cumulus Congestus Cloud, Journal of the Atmospheric Sciences, 79, 1887–1910, <https://doi.org/10.1175/JAS-D-21-0138.1>, 2022.

Chandrakar, K. K., Morrison, H., Grabowski, W. W., and Lawson, R. P.: Are turbulence effects on droplet collision–coalescence a key to understanding observed rain formation in clouds?, Proceedings of the National Academy of Sciences of the United States of America, 121, e2319664 121, https://doi.org/10.1073/PNAS.2319664121/SUPPL_FILE/PNAS.2319664121.SAPP.PDF, 2024.

Chen, Y. S., Verlinde, J., Clothiaux, E. E., Ackerman, A. S., Fridlind, A. M., Chamecki, M., Kollias, P., Kirkpatrick, M. P., Chen, B. C., Yu, G., and Avramov, A.: On the Forward Modeling of Radar Doppler Spectrum Width From LES: Implications for Model Evaluation, Journal of Geophysical Research: Atmospheres, 123, 7444–7461, <https://doi.org/10.1029/2017JD028104>, 2018.

Chepfer, H., Bony, S., Winker, D., Chiriaco, M., Dufresne, J. L., and Sèze, G.: Use of CALIPSO lidar observations to evaluate the cloudiness simulated by a climate model, Geophysical Research Letters, 35, <https://doi.org/10.1029/2008GL034207>, 2008.

Cooper, W. A.: Effects of Variable Droplet Growth Histories on Droplet Size Distributions. Part I: Theory, Journal of Atmospheric Sciences, 46, 1301 – 1311, [https://doi.org/10.1175/1520-0469\(1989\)046<1301:EOVDGH>2.0.CO;2](https://doi.org/10.1175/1520-0469(1989)046<1301:EOVDGH>2.0.CO;2), 1989.

Elsaesser, G. S., Del Genio, A. D., Jiang, J. H., and Lier-Walqui, M. v.: An Improved Convective Ice Parameterization for the NASA GISS Global Climate Model and Impacts on Cloud Ice Simulation, Journal of Climate, 30, 317–336, <https://doi.org/10.1175/JCLI-D-16-0346.1>, 2017.

Endo, S., Fridlind, A. M., Lin, W., Vogelmann, A. M., Toto, T., Ackerman, A., McFarquhar, G. M., Jackson, R. C., Jonsson, H. H., and Liu, Y.: RACORO continental boundary layer cloud investigations: 2. Large-eddy simulations of cumulus clouds and evaluation with in situ and ground-based observations, Journal of Geophysical Research: Atmospheres, 120, 5993–6014, <https://doi.org/10.1002/2014JD022525>, 2015.

Freud, E. and Rosenfeld, D.: Linear relation between convective cloud drop number concentration and depth for rain initiation, Journal of Geophysical Research Atmospheres, 117, <https://doi.org/10.1029/2011JD016457>, 2012.

Fridlind, A. M. and Ackerman, A. S.: Simulations of Arctic Mixed-Phase Boundary Layer Clouds: Advances in Understanding and Outstanding Questions, Mixed-Phase Clouds: Observations and Modeling, pp. 153–183, <https://doi.org/10.1016/B978-0-12-810549-8.00007-6>, 2017.

- Fridlind, A. M., Ackerman, A. S., Chaboureaud, J. P., Fan, J., Grabowski, W. W., Hill, A. A., Jones, T. R., Khaiyer, M. M., Liu, G., Minnis, P., Morrison, H., Nguyen, L., Park, S., Petch, J. C., Pinty, J. P., Schumacher, C., Shipway, B. J., Varble, A. C., Wu, X., Xie, S., and Zhang, M.: A comparison of TWP-ICE observational data with cloud-resolving model results, *Journal of Geophysical Research: Atmospheres*, 117, 5204, <https://doi.org/10.1029/2011JD016595>, 2012.
- 900 Fridlind, A. M., Li, X., Wu, D., Van Lier-Walqui, M., Ackerman, A. S., Tao, W. K., McFarquhar, G. M., Wu, W., Dong, X., Wang, J., Ryzhkov, A., Zhang, P., Poellot, M. R., Neumann, A., and Tomlinson, J. M.: Derivation of aerosol profiles for MC3E convection studies and use in simulations of the 20 May squall line case, *Atmospheric Chemistry and Physics*, 17, 5947–5972, <https://doi.org/10.5194/ACP-17-5947-2017>, 2017.
- Fu, D., Di Girolamo, L., Rauber, R. M., McFarquhar, G. M., Nesbitt, S. W., Loveridge, J., Hong, Y., Van Diedenhoven, B., Cairns, B., Alexandrov, M. D., Lawson, P., Woods, S., Tanelli, S., Schmidt, S., Hostetler, C., and Scarino, A. J.: An evaluation of the liquid cloud droplet effective radius derived from MODIS, airborne remote sensing, and in situ measurements from CAMP2Ex, *Atmospheric Chemistry and Physics*, 22, 8259–8285, <https://doi.org/10.5194/ACP-22-8259-2022>, 2022.
- 905 Gao, M., Franz, B. A., Zhai, P. W., Knobelspiesse, K., Sayer, A. M., Xu, X., Martins, J. V., Cairns, B., Castellanos, P., Fu, G., Hannadige, N., Hasekamp, O., Hu, Y., Ibrahim, A., Patt, F., Puthukkudy, A., and Werdell, P. J.: Simultaneous retrieval of aerosol and ocean properties from PACE HARP2 with uncertainty assessment using cascading neural network radiative transfer models, *Atmospheric Measurement Techniques*, 16, 5863–5881, <https://doi.org/10.5194/AMT-16-5863-2023>, 2023.
- 910 Gerber, H.: Microphysics of Marine Stratocumulus Clouds with Two Drizzle Modes, *Journal of Atmospheric Sciences*, 53, 1649 – 1662, [https://doi.org/10.1175/1520-0469\(1996\)053<1649:MOMSCW>2.0.CO;2](https://doi.org/10.1175/1520-0469(1996)053<1649:MOMSCW>2.0.CO;2), 1996.
- Grabowski, W. W. and Abade, G. C.: Broadening of Cloud Droplet Spectra through Eddy Hopping: Turbulent Adiabatic Parcel Simulations, *Journal of the Atmospheric Sciences*, 74, 1485–1493, <https://doi.org/10.1175/JAS-D-17-0043.1>, 2017.
- 915 Gregory, D. and Rowntree, P. R.: A Mass Flux Convection Scheme with Representation of Cloud Ensemble Characteristics and Stability-Dependent Closure, *Monthly Weather Review*, 118, 1483 – 1506, [https://doi.org/10.1175/1520-0493\(1990\)118<1483:AMFCSW>2.0.CO;2](https://doi.org/10.1175/1520-0493(1990)118<1483:AMFCSW>2.0.CO;2), 1990.
- Grosvenor, D. P., Sourdeval, O., Zuidema, P., Ackerman, A., Alexandrov, M. D., Bennartz, R., Boers, R., Cairns, B., Chiu, J. C., Christensen, M., Deneke, H., Diamond, M., Feingold, G., Fridlind, A., Hünerbein, A., Knist, C., Kollias, P., Marshak, A., McCoy, D., Merk, D., Painemal, D., Rausch, J., Rosenfeld, D., Russchenberg, H., Seifert, P., Sinclair, K., Stier, P., van Diedenhoven, B., Wendisch, M., Werner, F., Wood, R., Zhang, Z., and Quaas, J.: Remote Sensing of Droplet Number Concentration in Warm Clouds: A Review of the Current State of Knowledge and Perspectives, *Reviews of Geophysics*, 56, 409–453, <https://doi.org/10.1029/2017RG000593>, 2018.
- 920 Hall, D.: A detailed microphysical model within a two-dimensional dynamic framework: Model description and preliminary results, *Journal of the Atmospheric Sciences*, pp. 2486–2507, [https://doi.org/10.1175/1520-0469\(1980\)037](https://doi.org/10.1175/1520-0469(1980)037), 1980.
- Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maenhaut, W., McFiggans, G., Mentel, T. F., Monod, A., Prévôt, A. S., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: Current and emerging issues, *Atmospheric Chemistry and Physics*, 9, 5155–5236, <https://doi.org/10.5194/ACP-9-5155-2009>, 2009.
- 930 Hansen, J. E. and Travis, L. D.: Light scattering in planetary atmospheres, *Space Science Reviews*, 16, 527–610, <https://doi.org/10.1007/BF00168069/METRICS>, 1974.

- Hasekamp, O. P., Fu, G., Rusli, S. P., Wu, L., Di Noia, A., Brugh, J. a. d., Landgraf, J., Martijn Smit, J., Rietjens, J., and van Amerongen, A.: Aerosol measurements by SPEXone on the NASA PACE mission: expected retrieval capabilities, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 227, 170–184, <https://doi.org/10.1016/J.QSRT.2019.02.006>, 2019.
- Hernandez-Deckers, D. and Sherwood, S. C.: A Numerical Investigation of Cumulus Thermals, *Journal of the Atmospheric Sciences*, 73, 4117–4136, <https://doi.org/10.1175/JAS-D-15-0385.1>, 2016.
- Hernandez-Deckers, D. and Sherwood, S. C.: On the Role of Entrainment in the Fate of Cumulus Thermals, *Journal of the Atmospheric Sciences*, 75, 3911–3924, <https://doi.org/10.1175/JAS-D-18-0077.1>, 2018.
- 940 Hernandez-Deckers, D., Matsui, T., and Fridlind, A. M.: Updraft dynamics and microphysics: On the added value of the cumulus thermal reference frame in simulations of aerosol-deep convection interactions, *Atmospheric Chemistry and Physics*, 22, 711–724, <https://doi.org/10.5194/ACP-22-711-2022>, 2022.
- Hohenegger, C. and Stevens, B.: Preconditioning Deep Convection with Cumulus Congestus, *Journal of the Atmospheric Sciences*, 70, 448–464, <https://doi.org/10.1175/JAS-D-12-089.1>, 2013.
- 945 Jensen, E. J., Toon, O. B., Tabazadeh, A., Sachse, G. W., Anderson, B. E., Chan, K. R., Twohy, C. W., Gandrud, B., Aulenbach, S. M., Heymsfield, A., Hallett, J., and Gary, B.: Ice nucleation processes in upper tropospheric wave-clouds observed during SUCCESS, *Geophysical Research Letters*, 25, 1363–1366, <https://doi.org/10.1029/98GL00299>, 1998.
- Johnson, R. H., Rickenbach, T. M., Rutledge, S. A., Ciesielski, P. E., and Schubert, W. H.: Trimodal Characteristics of Tropical Convection, *Journal of Climate*, 12, 2397 – 2418, [https://doi.org/10.1175/1520-0442\(1999\)012<2397:TCOTC>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<2397:TCOTC>2.0.CO;2), 1999.
- 950 Juliano, T. W., Tornow, F., Fridlind, A. M., Ackerman, A. S., Geerts, B., Kosović, B., Ovchinnikov, M., Wu, P., and Xue, L.: Evaluating simulated convective clouds during Arctic cold-air outbreaks: A model intercomparison study based on COMBLE, [White Paper], https://www.gewexevents.org/wp-content/uploads/COMBLE_white_paper.pdf, 2022.
- Keinert, A., Spannagel, D., Leisner, T., and Kiselev, A.: Secondary Ice Production upon Freezing of Freely Falling Drizzle Droplets, *Journal of the Atmospheric Sciences*, 77, 2959–2967, <https://doi.org/10.1175/JAS-D-20-0081.1>, 2020.
- 955 Khairoutdinov, M. and Kogan, Y.: A New Cloud Physics Parameterization in a Large-Eddy Simulation Model of Marine Stratocumulus, *Monthly Weather Review*, 128, 229 – 243, [https://doi.org/10.1175/1520-0493\(2000\)128<0229:ANCPPI>2.0.CO;2](https://doi.org/10.1175/1520-0493(2000)128<0229:ANCPPI>2.0.CO;2), 2000.
- Kirkpatrick, M. P., Ackerman, A. S., Stevens, D. E., and Mansour, N. N.: On the Application of the Dynamic Smagorinsky Model to Large-Eddy Simulations of the Cloud-Topped Atmospheric Boundary Layer, *Journal of the Atmospheric Sciences*, 63, 526–546, <https://doi.org/10.1175/JAS3651.1>, 2006.
- 960 Klein, S. A., McCoy, R. B., Morrison, H., Ackerman, A. S., Avramov, A., de Boer, G., Chen, M., Cole, J. N., del Genio, A. D., Falk, M., Foster, M. J., Fridlind, A., Golaz, J. C., Hashino, T., Harrington, J. Y., Hoose, C., Khairoutdinov, M. F., Larson, V. E., Liu, X., Luo, Y., McFarquhar, G. M., Menon, S., Neggers, R. A., Park, S., Poellot, M. R., Schmidt, J. M., Sednev, I., Shipway, B. J., Shupe, M. D., Spangenberg, D. A., Sud, Y. C., Turner, D. D., Veron, D. E., von Salzen, K., Walker, G. K., Wang, Z., Wolf, A. B., Xie, S., Xu, K. M., Yang, F., and Zhang, G.: Intercomparison of model simulations of mixed-phase clouds observed during the ARM Mixed-Phase Arctic Cloud Experiment. I: single-layer cloud, *Quarterly Journal of the Royal Meteorological Society*, 135, 979–1002, <https://doi.org/10.1002/QJ.416>, 2009.
- 965 Korolev, A. V., Strapp, J. W., Isaac, G. A., and Nevzorov, A. N.: The Nevzorov Airborne Hot-Wire LWC–TWC Probe: Principle of Operation and Performance Characteristics, *Journal of Atmospheric and Oceanic Technology*, 15, 1495–1510, [https://doi.org/https://doi.org/10.1175/1520-0426\(1998\)015<1495:TNAHWL>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0426(1998)015<1495:TNAHWL>2.0.CO;2), 1998.

- 970 Kuang, Z. and Bretherton, C. S.: A Mass-Flux Scheme View of a High-Resolution Simulation of a Transition from Shallow to Deep Cumulus Convection, *Journal of the Atmospheric Sciences*, 63, 1895–1909, <https://doi.org/10.1175/JAS3723.1>, 2006.
- Kulkarni, P. and Wang, J.: New fast integrated mobility spectrometer for real-time measurement of aerosol size distribution—I: Concept and theory, *Journal of Aerosol Science*, 37, 1303–1325, <https://doi.org/10.1016/J.JAEROSCI.2006.01.005>, 2006.
- Lasher-Trapp, S. G., Cooper, W. A., and Blyth, A. M.: Broadening of droplet size distributions from entrainment and mixing in a cumulus
975 cloud, *Quarterly Journal of the Royal Meteorological Society*, 131, 195–220, <https://doi.org/10.1256/QJ.03.199>, 2005.
- Lauber, A., Kiselev, A., Pander, T., Handmann, P., and Leisner, T.: Secondary ice formation during freezing of levitated droplets, *Journal of the Atmospheric Sciences*, 75, 2815–2826, <https://doi.org/10.1175/JAS-D-18-0052.1>, 2018.
- Lawson, P., Gurganus, C., Woods, S., and Bruintjes, R.: Aircraft observations of cumulus microphysics ranging from the tropics to midlatitudes: Implications for a "new" secondary ice process, *Journal of the Atmospheric Sciences*, 74, 2899–2920, [https://doi.org/10.1175/JAS-](https://doi.org/10.1175/JAS-D-17-0033.1)
980 [D-17-0033.1](https://doi.org/10.1175/JAS-D-17-0033.1), 2017.
- Lawson, P. R., Woods, S., and Morrison, H.: The Microphysics of Ice and Precipitation Development in Tropical Cumulus Clouds, *Journal of the Atmospheric Sciences*, 72, 2429–2445, <https://doi.org/10.1175/JAS-D-14-0274.1>, 2015.
- Lawson, R. P., Stewart, R. E., and Angus, L. J.: Observations and Numerical Simulations of the Origin and Development of Very Large Snowflakes, *Journal of the Atmospheric Sciences*, 55, 3209 – 3229, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0469(1998)055<3209:OANSOT>2.0.CO;2)
985 [0469\(1998\)055<3209:OANSOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1998)055<3209:OANSOT>2.0.CO;2), 1998.
- Lawson, R. P., Baker, B. A., Schmitt, C. G., and Jensen, T. L.: An overview of microphysical properties of Arctic clouds observed in May and July 1998 during FIRE ACE, *Journal of Geophysical Research: Atmospheres*, 106, 14 989–15 014, <https://doi.org/10.1029/2000JD900789>, 2001.
- Lawson, R. P., O'Connor, D., Zmarzly, P., Weaver, K., Baker, B., Mo, Q., and Jonsson, H.: The 2D-S (Stereo) Probe: Design and Preliminary
990 Tests of a New Airborne, High-Speed, High-Resolution Particle Imaging Probe, *Journal of Atmospheric and Oceanic Technology*, 23, 1462–1477, <https://doi.org/10.1175/JTECH1927.1>, 2006.
- Lawson, R. P., Bruintjes, R., Woods, S., and Gurganus, C.: Coalescence and Secondary Ice Development in Cumulus Congestus Clouds, *Journal of the Atmospheric Sciences*, 79, 953–972, <https://doi.org/10.1175/JAS-D-21-0188.1>, 2022.
- Lee, H., Fridland, A. M., and Ackerman, A. S.: An Evaluation of Size-Resolved Cloud Microphysics Scheme Numerics for Use with Radar
995 Observations. Part II: Condensation and Evaporation, *Journal of the Atmospheric Sciences*, 78, 1629–1645, [https://doi.org/10.1175/JAS-](https://doi.org/10.1175/JAS-D-20-0213.1)
[D-20-0213.1](https://doi.org/10.1175/JAS-D-20-0213.1), 2021.
- Lin, C. and Arakawa, A.: The Macroscopic Entrainment Processes of Simulated Cumulus Ensemble. Part II: Testing the Entraining-Plume Model, *Journal of the Atmospheric Sciences*, 54, 1044 – 1053, [https://doi.org/10.1175/1520-0469\(1997\)054<1044:TMEPOS>2.0.CO;2](https://doi.org/10.1175/1520-0469(1997)054<1044:TMEPOS>2.0.CO;2), 1997.
- 1000 Lin, W., Liu, Y., Vogelmann, A. M., Fridlind, A., Endo, S., Song, H., Feng, S., Toto, T., Li, Z., and Zhang, M.: RACORO continental boundary layer cloud investigations: 3. Separation of parameterization biases single-column model CAM5 simulations of shallow cumulus, *Journal of Geophysical Research: Atmospheres*, 120, 6015–6033, <https://doi.org/10.1002/2014JD022524>, 2015.
- Low, T. B. and List, R.: Collision, Coalescence and Breakup of Raindrops. Part II: Parameterization of Fragment Size Distributions, *Journal of Atmospheric Sciences*, 39, 1607 – 1619, [https://doi.org/10.1175/1520-0469\(1982\)039<1607:CCABOR>2.0.CO;2](https://doi.org/10.1175/1520-0469(1982)039<1607:CCABOR>2.0.CO;2), 1982.
- 1005 Luo, Z., Liu, G. Y., Stephens, G. L., and Johnson, R. H.: Terminal versus transient cumulus congestus: A CloudSat perspective, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2008GL036927>, 2009.

- Mansell, E. R. and Ziegler, C. L.: Aerosol Effects on Simulated Storm Electrification and Precipitation in a Two-Moment Bulk Microphysics Model, *Journal of the Atmospheric Sciences*, 70, 2032–2050, <https://doi.org/10.1175/JAS-D-12-0264.1>, 2013.
- Mansell, E. R., Ziegler, C. L., and Bruning, E. C.: Simulated Electrification of a Small Thunderstorm with Two-Moment Bulk Microphysics, *Journal of the Atmospheric Sciences*, 67, 171–194, <https://doi.org/10.1175/2009JAS2965.1>, 2010.
- Manton, M. J.: On the broadening of a droplet distribution by turbulence near cloud base, *Quarterly Journal of the Royal Meteorological Society*, 105, 899–914, <https://doi.org/10.1002/QJ.49710544613>, 1979.
- Martins, J. V., Fernandez-Borda, R., McBride, B., Remer, L., and Barbosa, H. M.: The harp hyperangular imaging polarimeter and the need for small satellite payloads with high science payoff for earth science remote sensing, *International Geoscience and Remote Sensing Symposium (IGARSS)*, 2018-July, 6304–6307, <https://doi.org/10.1109/IGARSS.2018.8518823>, 2018.
- Matsui, T., Zhang, S. Q., Lang, S. E., Tao, W. K., Ichoku, C., and Peters-Lidard, C. D.: Impact of radiation frequency, precipitation radiative forcing, and radiation column aggregation on convection-permitting West African monsoon simulations, *Climate Dynamics*, 55, 193–213, <https://doi.org/10.1007/S00382-018-4187-2/FIGURES/13>, 2020.
- Matsui, T., Hernandez-Deckers, D., Giangrande, S., Biscaro, T., Fridlind, A., and Braun, S.: Thermal-Driven Graupel Generation Process to Explain Dry-Season Convective Vigor over the Amazon, *EGUsphere* [preprint], 2024.
- Miller, D. J., Zhang, Z., Platnick, S., Ackerman, A. S., Werner, F., Cornet, C., and Knobelspiesse, K.: Comparisons of bispectral and polarimetric retrievals of marine boundary layer cloud microphysics: Case studies using a LES-satellite retrieval simulator, *Atmospheric Measurement Techniques*, 11, 3689–3715, <https://doi.org/10.5194/AMT-11-3689-2018>, 2018.
- Morrison, H. and Grabowski, W. W.: A Novel Approach for Representing Ice Microphysics in Models: Description and Tests Using a Kinematic Framework, *Journal of the Atmospheric Sciences*, 65, 1528–1548, <https://doi.org/10.1175/2007JAS2491.1>, 2008a.
- Morrison, H. and Grabowski, W. W.: Modeling Supersaturation and Subgrid-Scale Mixing with Two-Moment Bulk Warm Microphysics, *Journal of the Atmospheric Sciences*, 65, 792–812, <https://doi.org/10.1175/2007JAS2374.1>, 2008b.
- Morrison, H., Curry, J. A., and Khvorostyanov, V. I.: A New Double-Moment Microphysics Parameterization for Application in Cloud and Climate Models. Part I: Description, *Journal of the Atmospheric Sciences*, 62, 1665–1677, <https://doi.org/10.1175/JAS3446.1>, 2005.
- Morrison, H., Thompson, G., and Tatarskii, V.: Impact of Cloud Microphysics on the Development of Trailing Stratiform Precipitation in a Simulated Squall Line: Comparison of One- and Two-Moment Schemes, *Monthly Weather Review*, 137, 991–1007, <https://doi.org/10.1175/2008MWR2556.1>, 2009.
- Morrison, H., Witte, M., Bryan, G. H., Harrington, J. Y., and Lebo, Z. J.: Broadening of Modeled Cloud Droplet Spectra Using Bin Microphysics in an Eulerian Spatial Domain, *Journal of the Atmospheric Sciences*, 75, 4005–4030, <https://doi.org/10.1175/JAS-D-18-0055.1>, 2018.
- Morrison, H., Peters, J. M., Varble, A. C., Hannah, W. M., and Giangrande, S. E.: Thermal Chains and Entrainment in Cumulus Updrafts. Part I: Theoretical Description, *Journal of the Atmospheric Sciences*, 77, 3637–3660, <https://doi.org/10.1175/JAS-D-19-0243.1>, 2020.
- Morrison, H., Lawson, P., and Chandrakar, K. K.: Observed and Bin Model Simulated Evolution of Drop Size Distributions in High-Based Cumulus Congestus Over the United Arab Emirates, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035711, <https://doi.org/10.1029/2021JD035711>, 2022.
- Moser, D. H. and Lasher-Trapp, S.: The Influence of Successive Thermals on Entrainment and Dilution in a Simulated Cumulus Congestus, *Journal of the Atmospheric Sciences*, 74, 375–392, <https://doi.org/10.1175/JAS-D-16-0144.1>, 2017.
- Nakanishi, M. and Niino, H.: Development of an Improved Turbulence Closure Model for the Atmospheric Boundary Layer, *Journal of the Meteorological Society of Japan. Ser. II*, 87, 895–912, <https://doi.org/10.2151/JMSJ.87.895>, 2009.

- 1045 National Centers for Environmental Prediction National Weather Service NOAA U.S. Department of Commerce: NCEP GDAS/FNL 0.25 Degree Global Tropospheric Analyses and Forecast Grids, <https://doi.org/10.5065/D65Q4T4Z>, 2015.
- Neggers, R. A.: Attributing the behavior of low-level clouds in large-scale models to subgrid-scale parameterizations, *Journal of Advances in Modeling Earth Systems*, 7, 2029–2043, <https://doi.org/10.1002/2015MS000503>, 2015.
- Neggers, R. A., Ackerman, A. S., Angevine, W. M., Bazile, E., Beau, I., Blossey, P. N., Boutle, I. A., de Bruijn, C., Cheng, A., van der
1050 Dussen, J., Fletcher, J., Dal Gesso, S., Jam, A., Kawai, H., Cheedela, S. K., Larson, V. E., Lefebvre, M. P., Lock, A. P., Meyer, N. R., de Roode, S. R., de Rooy, W., Sandu, I., Xiao, H., and Xu, K. M.: Single-Column Model Simulations of Subtropical Marine Boundary-Layer Cloud Transitions Under Weakening Inversions, *Journal of Advances in Modeling Earth Systems*, 9, 2385–2412, <https://doi.org/10.1002/2017MS001064>, 2017.
- Niu, G. Y., Yang, Z. L., Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., Kumar, A., Manning, K., Niyogi, D., Rosero, E., Tewari, M., and Xia,
1055 Y.: The community Noah land surface model with multiparameterization options (Noah-MP): 1. Model description and evaluation with local-scale measurements, *Journal of Geophysical Research: Atmospheres*, 116, 12 109, <https://doi.org/10.1029/2010JD015139>, 2011.
- O'Connor, D., Baker, B., and Lawson, R. P.: Upgrades to the FSSP-100 Electronics, Int. Conf. on Clouds and Precipitation, Cancun, Mexico, IAMAS, http://cabernet.atmosfcu.unam.mx/ICCP-2008/abstracts/Program_on_line/Poster_13/O-Connor_extended.pdf, 2008.
- Painemal, D. and Zuidema, P.: Assessment of MODIS cloud effective radius and optical thickness retrievals over the
1060 Southeast Pacific with VOCALS-REx in situ measurements, *Journal of Geophysical Research: Atmospheres*, 116, 24 206, <https://doi.org/10.1029/2011JD016155>, 2011.
- Pardo, L. H., Morrison, H., Machado, L. A., Harrington, J. Y., and Lebo, Z. J.: Drop Size Distribution Broadening Mechanisms in a Bin Microphysics Eulerian Model, *Journal of the Atmospheric Sciences*, 77, 3249–3273, <https://doi.org/10.1175/JAS-D-20-0099.1>, 2020.
- Peters, J. M., Morrison, H., Varble, A. C., Hannah, W. M., and Giangrande, S. E.: Thermal Chains and Entrainment in Cumulus Updrafts.
1065 Part II: Analysis of Idealized Simulations, *Journal of the Atmospheric Sciences*, 77, 3661–3681, <https://doi.org/10.1175/JAS-D-19-0244.1>, 2020.
- Peters-Lidard, C. D., Kemp, E. M., Matsui, T., Santanello, J. A., Kumar, S. V., Jacob, J. P., Clune, T., Tao, W. K., Chin, M., Hou, A., Case, J. L., Kim, D., Kim, K. M., Lau, W., Liu, Y., Shi, J., Starr, D., Tan, Q., Tao, Z., Zaitchik, B. F., Zavodsky, B., Zhang, S. Q., and Zupanski, M.: Integrated modeling of aerosol, cloud, precipitation and land processes at satellite-resolved scales, *Environmental Modelling & Software*,
1070 67, 149–159, <https://doi.org/10.1016/J.ENVSOFT.2015.01.007>, 2015.
- Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, *Atmospheric Chemistry and Physics*, 7, 1961–1971, <https://doi.org/10.5194/ACP-7-1961-2007>, 2007.
- Phillips, V. T., Patade, S., Gutierrez, J., and Bansemer, A.: Secondary Ice Production by Fragmentation of Freezing Drops: Formulation and Theory, *Journal of the Atmospheric Sciences*, 75, 3031–3070, <https://doi.org/10.1175/JAS-D-17-0190.1>, 2018.
- 1075 Reid, J. S., Maring, H. B., Narisma, G. T., van den Heever, S., Di Girolamo, L., Ferrare, R., Lawson, P., Mace, G. G., Simpas, J. B., Tanelli, S., Ziemba, L., van Didenhoven, B., Bruintjes, R., Bucholtz, A., Cairns, B., Cambaliza, M. O., Chen, G., Diskin, G. S., Flynn, J. H., Hostetler, C. A., Holz, R. E., Lang, T. J., Schmidt, K. S., Smith, G., Sorooshian, A., Thompson, E. J., Thornhill, K. L., Trepte, C., Wang, J., Woods, S., Yoon, S., Alexandrov, M., Alvarez, S., Amiot, C. G., Bennett, J. R., Brooks, M., Burton, S. P., Cayan, E., Chen, H., Collow, A., Crosbie, E., DaSilva, A., DiGangi, J. P., Flagg, D. D., Freeman, S. W., Fu, D., Fukada, E., Hilario, M. R. A., Hong, Y.,
1080 Hristova-Veleva, S. M., Kuehn, R., Kowch, R. S., Leung, G. R., Loveridge, J., Meyer, K., Miller, R. M., Montes, M. J., Moum, J. N., Nenes, T., Nesbitt, S. W., Norgren, M., Nowottnick, E. P., Rauber, R. M., Reid, E. A., Rutledge, S., Schlosser, J. S., Sekiyama, T. T., Shook, M. A., Sokolowsky, G. A., Stamnes, S. A., Tanaka, T. Y., Wasilewski, A., Xian, P., Xiao, Q., Xu, Z., and Zavaleta, J.: The coupling

- between tropical meteorology, aerosol lifecycle, convection, and radiation, during the Cloud, Aerosol and Monsoon Processes Philippines Experiment (CAMP2Ex), *Bulletin of the American Meteorological Society*, -1, <https://doi.org/10.1175/BAMS-D-21-0285.1>, 2023.
- 1085 Rémillard, J., Fridlind, A. M., Ackerman, A. S., Tselioudis, G., Kollias, P., Mechem, D. B., Chandler, H. E., Luke, E., Wood, R., Witte, M. K., Chuang, P. Y., and Ayers, J. K.: Use of Cloud Radar Doppler Spectra to Evaluate Stratocumulus Drizzle Size Distributions in Large-Eddy Simulations with Size-Resolved Microphysics, *Journal of Applied Meteorology and Climatology*, 56, 3263–3283, <https://doi.org/10.1175/JAMC-D-17-0100.1>, 2017.
- Rosenfeld, D. and Gutman, G.: Retrieving microphysical properties near the tops of potential rain clouds by multispectral analysis of AVHRR data, *Atmospheric Research*, 34, 259–283, [https://doi.org/10.1016/0169-8095\(94\)90096-5](https://doi.org/10.1016/0169-8095(94)90096-5), 1994.
- 1090 Sandu, I. and Stevens, B.: On the Factors Modulating the Stratocumulus to Cumulus Transitions, *Journal of the Atmospheric Sciences*, 68, 1865–1881, <https://doi.org/10.1175/2011JAS3614.1>, 2011.
- Sato, Y., Shima, S. i., and Tomita, H.: Numerical Convergence of Shallow Convection Cloud Field Simulations: Comparison Between Double-Moment Eulerian and Particle-Based Lagrangian Microphysics Coupled to the Same Dynamical Core, *Journal of Advances in Modeling Earth Systems*, 10, 1495–1512, <https://doi.org/10.1029/2018MS001285>, 2018.
- 1095 Seifert, A.: On the Parameterization of Evaporation of Raindrops as Simulated by a One-Dimensional Rainshaft Model, *Journal of the Atmospheric Sciences*, 65, 3608–3619, <https://doi.org/10.1175/2008JAS2586.1>, 2008.
- Seifert, A. and Beheng, K. D.: A double-moment parameterization for simulating autoconversion, accretion and selfcollection, *Atmospheric Research*, 59-60, 265–281, [https://doi.org/10.1016/S0169-8095\(01\)00126-0](https://doi.org/10.1016/S0169-8095(01)00126-0), 2001.
- 1100 Sherwood, S. C., Hernández-Deckers, D., Colin, M., and Robinson, F.: Slippery Thermals and the Cumulus Entrainment Paradox, *Journal of the Atmospheric Sciences*, 70, 2426–2442, <https://doi.org/10.1175/JAS-D-12-0220.1>, 2013.
- Siebesma, A. P., Bretherton, C. S., Brown, A., Chlond, A., Cuxart, J., Duynkerke, P. G., Jiang, H., Khairoutdinov, M., Lewellen, D., Moeng, C.-H., Sanchez, E., Stevens, B., and Stevens, D. E.: A Large Eddy Simulation Intercomparison Study of Shallow Cumulus Convection, *Journal of the Atmospheric Sciences*, 60, 1201 – 1219, [https://doi.org/10.1175/1520-0469\(2003\)60<1201:ALESIS>2.0.CO;2](https://doi.org/10.1175/1520-0469(2003)60<1201:ALESIS>2.0.CO;2), 2003.
- 1105 Sinclair, K., Van Diedenhoven, B., Cairns, B., Yorks, J., Wasilewski, A., and McGill, M.: Remote sensing of multiple cloud layer heights using multi-Angular measurements, *Atmospheric Measurement Techniques*, 10, 2361–2375, <https://doi.org/10.5194/AMT-10-2361-2017>, 2017.
- Sinclair, K., van Diedenhoven, B., Cairns, B., Alexandrov, M., Moore, R., Crosbie, E., and Ziemba, L.: Polarimetric retrievals of cloud droplet number concentrations, *Remote Sensing of Environment*, 228, 227–240, <https://doi.org/10.1016/J.RSE.2019.04.008>, 2019.
- 1110 Sinclair, K., van Diedenhoven, B., Cairns, B., Alexandrov, M., Moore, R., Ziemba, L. D., and Crosbie, E.: Observations of Aerosol-Cloud Interactions During the North Atlantic Aerosol and Marine Ecosystem Study, *Geophysical Research Letters*, 47, e2019GL085 851, <https://doi.org/10.1029/2019GL085851>, 2020.
- Sinclair, K., van Diedenhoven, B., Cairns, B., Alexandrov, M., Dzambo, A. M., and L’Ecuyer, T.: Inference of Precipitation in Warm Stratiform Clouds Using Remotely Sensed Observations of the Cloud Top Droplet Size Distribution, *Geophysical Research Letters*, 48, e2021GL092 547, <https://doi.org/10.1029/2021GL092547>, 2021.
- 1115 Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X.-Y., Wang, W., and Powers, J. G.: A Description of the Advanced Research WRF Version 3, NCAR Tech. Note NCAR/TN-475+STR, pp. 1–113, <https://doi.org/10.5065/D68S4MVH>, 2008.
- Song, X. and Zhang, G. J.: Microphysics parameterization for convective clouds in a global climate model: Description and single-column model tests, *Journal of Geophysical Research: Atmospheres*, 116, 2201, <https://doi.org/10.1029/2010JD014833>, 2011.
- 1120

- Stephens, G. L.: Radiation Profiles in Extended Water Clouds. II: Parameterization Schemes, *Journal of Atmospheric Sciences*, 35, 2123 – 2132, [https://doi.org/10.1175/1520-0469\(1978\)035<2123:RPIEWC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1978)035<2123:RPIEWC>2.0.CO;2), 1978.
- Stevens, B. and Seifert, A.: Understanding macrophysical outcomes of microphysical choices in simulations of shallow cumulus convection, *Journal of the Meteorological Society of Japan. Ser. II*, 86A, 143–162, <https://doi.org/10.2151/JMSJ.86A.143>, 2008.
- 1125 Stevens, D. E. and Bretherton, S.: A Forward-in-Time Advection Scheme and Adaptive Multilevel Flow Solver for Nearly Incompressible Atmospheric Flow, *Journal of Computational Physics*, 129, 284–295, <https://doi.org/10.1006/JCPH.1996.0250>, 1996.
- Stevens, D. E., Ackerman, A. S., and Bretherton, C. S.: Effects of Domain Size and Numerical Resolution on the Simulation of Shallow Cumulus Convection, *Journal of the Atmospheric Sciences*, 59, 3285–3301, [https://doi.org/10.1175/1520-0469\(2002\)059](https://doi.org/10.1175/1520-0469(2002)059), 2002.
- Tiedtke, M.: A Comprehensive Mass Flux Scheme for Cumulus Parameterization in Large-Scale Models, *Monthly Weather Review*, 117, 1779 – 1800, [https://doi.org/10.1175/1520-0493\(1989\)117<1779:ACMFSF>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1779:ACMFSF>2.0.CO;2), 1989.
- 1130 VanZanten, M. C., Stevens, B., Nuijens, L., Siebesma, A. P., Ackerman, A. S., Burnet, F., Cheng, A., Couvreux, F., Jiang, H., Khairoutdinov, M., Kogan, Y., Lewellen, D. C., Mechem, D., Nakamura, K., Noda, A., Shipway, B. J., Slawinska, J., Wang, S., and Wyszogrodzki, A.: Controls on precipitation and cloudiness in simulations of trade-wind cumulus as observed during RICO, *Journal of Advances in Modeling Earth Systems*, 3, <https://doi.org/10.1029/2011MS000056>, 2011.
- 1135 Varble, A., Fridlind, A. M., Zipser, E. J., Ackerman, A. S., Chaboureau, J.-P., Fan, J., Hill, A., Mcfarlane, S. A., Pinty, J.-P., Shipway, B., Varble, C. ., Fridlind, A. M., Zipser, E. J., Ackerman, A. S., Chaboureau, J.-P., Fan, J., Hill, A., Mcfarlane, S. A., Pinty, J.-P., and Shipway, B.: Evaluation of cloud-resolving model intercomparison simulations using TWP-ICE observations: Precipitation and cloud structure, *Journal of Geophysical Research: Atmospheres*, 116, 12 206, <https://doi.org/10.1029/2010JD015180>, 2011.
- Vogelmann, A. M., Fridlind, A. M., Toto, T., Endo, S., Lin, W., Wang, J., Feng, S., Zhang, Y., Turner, D. D., Liu, Y., Li, Z., Xie, S., Ackerman, A. S., Zhang, M., and Khairoutdinov, M.: RACORO continental boundary layer cloud investigations: 1. Case study development and ensemble large-scale forcings, *Journal of Geophysical Research: Atmospheres*, 120, 5962–5992, <https://doi.org/10.1002/2014JD022713>, 2015.
- 1140 Waite, M. L. and Khouider, B.: The Deepening of Tropical Convection by Congestus Preconditioning, *Journal of the Atmospheric Sciences*, 67, 2601–2615, <https://doi.org/10.1175/2010JAS3357.1>, 2010.
- 1145 Wall, C., Liu, C., and Zipser, E.: A climatology of tropical congestus using CloudSat, *Journal of Geophysical Research: Atmospheres*, 118, 6478–6492, <https://doi.org/10.1002/JGRD.50455>, 2013.
- Wang, J., Pikridas, M., Pinterich, T., Spielman, S. R., Tsang, T., McMahon, A., and Smith, S.: A Fast Integrated Mobility Spectrometer for rapid measurement of sub-micrometer aerosol size distribution, Part II: Experimental characterization, *Journal of Aerosol Science*, 113, 119–129, <https://doi.org/10.1016/J.JAEROSCI.2017.05.001>, 2017a.
- 1150 Wang, J., Pikridas, M., Spielman, S. R., and Pinterich, T.: A fast integrated mobility spectrometer for rapid measurement of sub-micrometer aerosol size distribution, Part I: Design and model evaluation, *Journal of Aerosol Science*, 108, 44–55, <https://doi.org/10.1016/J.JAEROSCI.2017.02.012>, 2017b.
- Wang, L. P. and Grabowski, W. W.: The role of air turbulence in warm rain initiation, *Atmospheric Science Letters*, 10, 1–8, <https://doi.org/10.1002/ASL.210>, 2009.
- 1155 Wang, Y., Pinterich, T., and Wang, J.: Rapid measurement of sub-micrometer aerosol size distribution using a fast integrated mobility spectrometer, *Journal of Aerosol Science*, 121, 12–20, <https://doi.org/10.1016/J.JAEROSCI.2018.03.006>, 2018.
- Warner, J.: The Microstructure of Cumulus Cloud. Part I. General Features of the Droplet Spectrum, *Journal of Atmospheric Sciences*, 26, 1049 – 1059, [https://doi.org/10.1175/1520-0469\(1969\)026<1049:TMOCCP>2.0.CO;2](https://doi.org/10.1175/1520-0469(1969)026<1049:TMOCCP>2.0.CO;2), 1969.

- Webster, P. J. and Lukas, R.: TOGA COARE: The Coupled Ocean–Atmosphere Response Experiment, *Bulletin of the American Meteorological Society*, 73, 1377 – 1416, [https://doi.org/10.1175/1520-0477\(1992\)073<1377:TCTCOR>2.0.CO;2](https://doi.org/10.1175/1520-0477(1992)073<1377:TCTCOR>2.0.CO;2), 1992.
- 1160 Werdell, P. J., Behrenfeld, M. J., Bontempi, P. S., Boss, E., Cairns, B., Davis, G. T., Franz, B. A., Gliese, U. B., Gorman, E. T., Hasekamp, O., Knobelspiesse, K. D., Mannino, A., Martins, J. V., McClain, C., Meister, G., and Remer, L. A.: The plankton, aerosol, cloud, ocean ecosystem mission status, science, advances, *Bulletin of the American Meteorological Society*, 100, 1775–1794, <https://doi.org/10.1175/BAMS-D-18-0056.1>, 2019.
- 1165 Woods, S., Lawson, R. P., Jensen, E., Bui, T. P., Thornberry, T., Rollins, A., Pfister, L., and Avery, M.: Microphysical Properties of Tropical Tropopause Layer Cirrus, *Journal of Geophysical Research: Atmospheres*, 123, 6053–6069, <https://doi.org/10.1029/2017JD028068>, 2018.
- Xu, W., Fossum, K. N., Ovadnevaite, J., Lin, C., Huang, R. J., O’Dowd, C., and Ceburnis, D.: The impact of aerosol size-dependent hygroscopicity and mixing state on the cloud condensation nuclei potential over the north-east Atlantic, *Atmospheric Chemistry and Physics*, 21, 8655–8675, <https://doi.org/10.5194/ACP-21-8655-2021>, 2021.
- 1170 Yang, Z. L., Niu, G. Y., Mitchell, K. E., Chen, F., Ek, M. B., Barlage, M., Longuevergne, L., Manning, K., Niyogi, D., Tewari, M., and Xia, Y.: The community Noah land surface model with multiparameterization options (Noah-MP): 2. Evaluation over global river basins, *Journal of Geophysical Research: Atmospheres*, 116, 12 110, <https://doi.org/10.1029/2010JD015140>, 2011.
- Zeng, X., Zhao, M., and Dickinson, R. E.: Intercomparison of Bulk Aerodynamic Algorithms for the Computation of Sea Surface Fluxes Using TOGA COARE and TAO Data, *Journal of Climate*, 11, 2628–2644, [https://doi.org/10.1175/1520-0442\(1998\)011<2628:IOBAAF>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<2628:IOBAAF>2.0.CO;2), 1998.
- 1175 Ziegler, C. L.: Retrieval of Thermal and Microphysical Variables in Observed Convective Storms. Part 1: Model Development and Preliminary Testing, *Journal of Atmospheric Sciences*, 42, 1487 – 1509, [https://doi.org/10.1175/1520-0469\(1985\)042<1487:ROTAMV>2.0.CO;2](https://doi.org/10.1175/1520-0469(1985)042<1487:ROTAMV>2.0.CO;2), 1985.