

Recent improvements and maximum covariance analysis of aerosol and cloud properties in the EC-Earth3-AerChem model

Manu Anna Thomas¹, Klaus Wyser¹, Shiyu Wang¹, Marios Chatziparaschos², Paraskevi Georgakaki⁴, Montserrat Costa-Surós², Maria Gonçalves Ageitos^{2,3}, Maria Kanakidou⁶, Carlos Pérez García-Pando^{2,7}, Athanasios Nenes^{4,8}, Twan van Noije⁵, Philippe Le Sager⁵, and Abhay Devasthale¹

¹Swedish Meteorological and Hydrological Institute (SMHI), Folkborgsvägen 17, 60176 Norrköping, Sweden.

²Barcelona Supercomputing Center (BSC), Plaça d'Eusebi Güell, 1-3, Les Corts, 08034 Barcelona, Spain.

³Universitat Politècnica de Catalunya (UPC), C. Jordi Girona 31, 08034 Barcelona, Spain.

⁴Laboratory of Atmospheric Processes and their Impacts (LAPI), School of Architecture, Civil and Environmental Engineering, École Polytechnique Fédérale de Lausanne (EPFL), 1015 Lausanne, Switzerland.

⁵The Royal Netherlands Meteorological Institute (KNMI), Utrechtseweg 297, NL-3731 De Bilt, Netherlands.

⁶University of Crete, Heraklion location, Voutes University campus, Crete, Greece.

⁷Catalan Institution for Research and Advanced Studies (ICREA), Passeig Lluís Companys 23, 08010 Barcelona, Spain.

⁸Center for Studies of Air Quality and Climate Change, Institute of Chemical Engineering Sciences (ICE-HT), Foundation for Research and Technology Hellas (FORTH), 26504 Patras, Greece.

Correspondence: Manu Anna Thomas (Manu.Thomas@smhi.se)

Abstract. Given the importance of aerosols, clouds and their interactions in the climate system, it is imperative that the global Earth system models accurately represent processes associated with them. This is an important prerequisite if we were to narrow the uncertainties in future climate projections. In practice, this means that the continuous model evaluations and improvements grounded in observations are necessary. Numerous studies in the last few decades have shown both the usability and the limitations of utilizing satellite-based observations in understanding and evaluating aerosol-cloud interactions, particularly under varying meteorological and satellite sensor sensitivity paradigms. Furthermore, the vast range of spatio-temporal scales at which aerosol and cloud processes occur adds another dimension to the challenges while evaluating climate models.

In this context, the aim of this study is two-fold. 1) We evaluate the most recent, significant changes in the representation of aerosol and cloud processes implemented in the EC-Earth3-AerChem model in the framework of the EU project FORCeS compared to its previous CMIP6 version (Coupled Model Intercomparison Project Phase 6; <https://pcmdi.llnl.gov/CMIP6/>). We focus particularly on evaluating cloud physical properties and radiative effects, wherever possible, using a satellite simulator. We report overall improvements in EC-Earth3-AerChem model. In particular, the strong warm bias chronically seen over the Southern Ocean is reduced significantly. 2) A statistical, maximum covariance analysis is carried out between aerosol optical depth (AOD) and cloud droplet (CD) effective radius based on the recent EC-Earth3-AerChem/FORCeS simulation to understand to what extent the Twomey effect can manifest itself in the larger spatio-temporal scales. We focus on the three oceanic low-level cloud regimes that are important due to their strong net cooling effect and where pollution outflow from the nearby continent is simultaneously pervasive. We report that the statistical covariability between AOD and CD effective radius is indeed dominantly visible even at the climate scale when the aerosol amount and composition are favourably preconditioned for allowing aerosol-cloud interactions. Despite this strong covariability, our analysis shows a strong cooling/warming in

20 shortwave cloud radiative effects at the top of the atmosphere in our study regions associated with an increase/decrease in CD effective radius. And this cooling/warming can be attributed to the increase/decrease in low cloud fraction, in line with the previous observational studies.

1 Introduction

Aerosols can potentially act as cloud condensation nuclei (CCN) or ice nuclei, influencing cloud formation, structure, and properties. The type, concentration, and size distribution of aerosols impact cloud microphysics, altering droplet or ice crystal size, number concentration, and cloud albedo (Twomey, 1974, 1977; Albrecht, 1989; Ramanathan et al., 2001). Conversely, clouds can also affect aerosols through wet deposition processes and also, indirectly via impact on meteorology and gas phase photochemistry, thereby, influencing their distribution and removal from the atmosphere. This interdependency of aerosol and clouds mediated by local meteorology plays a crucial role in climate regulation, radiative forcing and the distribution of precipitation. However, even after decades of research, a full grasp of the nature of aerosol-cloud interactions remains one of the big challenges and constitutes one of the largest sources of uncertainty in our understanding of climate forcing and feedbacks (Solomon et al., 2007; Quaas et al., 2009; Carslaw et al., 2010; Bellouin et al., 2019).

The vast range of spatio-temporal scales at which aerosol-cloud interactions occur helps us to appreciate why it is so difficult to pin down their role in the climate system. For example, processes like condensation and sublimation occur at nano- to micrometer spatial scales, while the interplay between aerosols and clouds over the major pollution outflow regions occur at much larger spatial scales. The temporal component of aerosol-cloud interactions also stretches from microseconds, to days and weeks of persistent pollution outflow, to the decades of emission policy changes. To account for these wide ranges of spatio-temporal scales, the harmonization and representation in a physically consistent and observationally constrained manner in global climate models is extremely challenging. At the same time, the evaluation of climate models must cover these different spatio-temporal scales. This can help us to assess, among others, at what spatio-temporal scales the aerosol-cloud interactions can distinctly manifest themselves and could even dominate the local variability.

In this context, this study focuses on understanding variability at larger scales. We specifically aim to understand the co-variability between AOD and CD effective radius at the climate scale. We acknowledge that both AOD and CD effective radius can be independently influenced by a number of processes and that CCN number concentration, especially at the cloud base, are better suited to investigate aerosol impact on cloud albedo via changes in droplet radius, as discussed in a number of previous studies (e.g. Quaas et al. (2020) and the references therein). The correspondence between AOD and CCN number concentration has been observed in a number of studies. Andreae (2009) identified a correlation between AOD and CCN concentration by comparing AERONET AOD with CCN measurements. Romakkaniemi et al. (2012) suggested the potential use of AOD as a proxy for CCN concentration. While uncertainties persist, Tang et al. (2014) noted the limitations of AOD as an indicator of CCN, acknowledging variations in CCN concentrations for a given AOD based on regional characteristics and meteorological conditions, such as relative humidity. Despite these complexities, Andreae (2009) observed a high correlation coefficient of

0.98 between AOD and CCN concentration. Using the ground measurements of CCN at the Zeppelin Observatory (78.91°N, 11.89°E), Ahn et al. (2021) derived a CCN–AOD correlation with a coefficient of determination R^2 of 0.59.

We choose AOD and CD effective radius here because these variables are better grounded in observations and can be
55 evaluated more directly using satellite-based observations. Furthermore, we assume that if aerosols indeed are one of the primary drivers of changes in cloud properties, especially under favourable meteorological conditions (Fanourgakis et al., 2019), their co-variability at a much larger, temporal and spatial scales shall visibly manifest itself in a relation between AOD and CD effective radius.

In this study, we first evaluate the recent improvements made in the EC-Earth3-AerChem model using a suite of satellite-
60 based observations and we then employ maximum co-variance analysis on AOD and CD effective radius to explore the intricate interplay between aerosols and clouds. We primarily focus on the oceanic regions that sustain low-level liquid water clouds and where the pollution outflow also occurs at least to some degree. The net cooling effect of low-level liquid water clouds over open oceans is very important and their susceptibility to aerosols has big implications for the energy budget.

Within the framework of the FORCES project (<https://forces-project.eu/>), this study aims to identify and implement the
65 crucial aerosol and cloud processes that could potentially reduce uncertainties in aerosol forcing estimates. This is detailed in Section 2.2. These improvements are being incorporated into the three global models that were previously used for CMIP6 (<https://pcmdi.llnl.gov/CMIP6/>): NorESM2-LM (Seland et al., 2020), MPI-ESM1.2-HAM (Mauritsen et al., 2019; Neubauer et al., 2019) and EC-Earth3-AerChem (van Noije et al., 2021). This paper focuses on the simulations carried out with EC-Earth3-AerChem model.

70 **2 Model, observations and methods**

2.1 EC-Earth3-AerChem model description

The EC-Earth3-AerChem model is based on the EC-Earth3 family of models (Döscher et al., 2022). The atmospheric compo-
nent of the model is based on IFS cycle 36r4 which includes the land surface model H-TESSSEL (Balsamo et al., 2009). The resolution of the atmospheric model is T255L91 with a grid spacing of approximately 80 km and 91 levels in the vertical that
75 extends to 0.01 hPa. The AerChem configuration features the aerosol and chemistry model, Tracer Model version 5 (TM5) (Krol et al., 2005; Huijnen et al., 2010). This chemistry and aerosol module is computationally expensive and is run at a much coarser resolution of $3^\circ \times 2^\circ$ with 34 vertical levels compared to the atmospheric model. The cloud macrophysics and microphysics are part of the atmosphere model and so are run at the same T255L91 resolution. The details of the CMIP6 version of the EC-Earth3-AerChem model are described in van Noije et al. (2021).

80 **2.2 EC-Earth3-AerChem model updates within FORCeS**

The FORCeS project aimed at the improvement of various crucial processes in the global climate models that would have the potential to influence aerosol radiative forcing and climate feedbacks, particularly concerning aerosols and clouds so as to

improve our future climate projections. In this context, the different minerals present in dust aerosols are now explicitly traced and their interaction with specific climate processes has been implemented and the cloud activation scheme was updated. The following paragraphs give a more detailed, but, a brief overview of these updates.

2.2.1 Cloud activation scheme

The formation of cloud droplets from ambient aerosol particles is a crucial process that needs to be accurately represented in global models to predict aerosol-cloud interactions and hence, the indirect effects of aerosols on climate. To achieve this, the modeling community has developed parameterization schemes that can realistically simulate the cloud droplet number concentrations based on ambient supersaturation, aerosol size and number distribution and updraft velocities. The schemes range from simpler schemes such as Abdul-Razzak and Ghan (2000) and Shipway (2015) to more complex schemes such as Barahona et al. (2010) and Morales-Betancourt and Nenes (2014). All parameterizations show a good overall agreement with the parcel model, but exhibit different sensitivities to aerosol perturbations under different meteorological conditions or pollution regimes (Ghan et al., 2011). For example, the Abdul-Razzak and Ghan (2000) scheme simulates low/high maximum supersaturations (S_{max}) and activation at low/high aerosol loads compared to the Morales-Betancourt and Nenes (2014) scheme making it more sensitive to increases in aerosol loads. The Shipway (2015) scheme is closer to the Morales-Betancourt and Nenes (2014) scheme, but tends to overestimate S_{max} and activation in polluted environments. Also, the simpler schemes show larger biases compared to the complex schemes in clean marine regions, particularly over the Southern Ocean. Hence, the current cloud activation scheme of Abdul-Razzak and Ghan (2000) is replaced by the Morales-Betancourt and Nenes (2014) scheme in the FORCeS version of the EC-Earth3-AerChem model. This scheme makes use of the population splitting concept wherein the growing population of droplets is divided into three enabling a more accurate estimate for the number concentration of droplets formed at the time of S_{max} .

2.2.2 Dust mineralogy

EC-Earth3-AerChem calculates dust emission online (van Noije et al., 2021). The emission scheme considers dry paleolakes as preferential dust sources and accounts for seasonal variations in vegetation cover. Saltation of dust particles occurs when the surface wind exceeds a certain threshold velocity, which depends on the soil particle size distribution and soil conditions. Furthermore, snow cover prevents dust emission in the model (Tegen et al., 2002, 2004). Dust is initially emitted into the accumulation and coarse insoluble modes, defined in the M7 aerosol microphysics module (Vignati et al., 2004). Within FORCeS, the dust emission module has been further developed to explicitly incorporate the atmospheric cycle of dust minerals relevant for their climate impacts. These include: (1) iron oxides, which control the absorption of shortwave (SW) radiation by dust aerosols (e.g., Sokolik and Toon, 1999; Di Biagio et al., 2019) (2) quartz and feldspars, which constitute efficient ice-nucleators (e.g., Atkinson et al., 2013; Harrison et al., 2019), and (3) calcite, which intervenes in atmospheric chemistry processes and affects the aerosol pH.

To estimate the emission of the different minerals, we derive the size-distributed mass fraction at emission for each of them considering information on the soil mineralogy of the dust sources (Claquin et al., 1999; Nickovic et al., 2012). To account

for the differences in the mineral's size distribution reported in the soil and those found in the aerosols, we apply an extension of the Brittle Fragmentation Theory for dust emission proposed by Kok (2011). This approach has shown a better agreement with observations (e.g., Perlwitz et al., 2015a, b) than relying exclusively on the soil size fractions. These newly incorporated minerals are then coupled to different processes in the model. The effect of iron oxides in the SW optical properties for dust is accounted for by assuming internal mixtures of iron oxides and other host minerals. A Maxwell Garnett mixing rule is applied to derive online the dust refractive index, considering the volume fraction of hematite and a homogeneous matrix of other minerals. The reference refractive indices are taken from Scanza et al. (2015). The abundance of quartz and K-feldspars is considered to derive the ice nucleating particles, and then coupled to the formation of mixed-phase clouds in the model. In order to estimate the specific fraction of K-feldspar, we assume it to be 35% of the total feldspar originally provided in the soil mineralogy datasets (Atkinson et al., 2013; Chatziparaschos et al., 2023). Finally, the calcite in dust is explicitly linked to the thermodynamic equilibrium of inorganic species, solved in the model with the ISORROPIA-Lite (Kakavas et al., 2022) model (see section 2.2.5).

2.2.3 MPOA source

Marine organic aerosols have been suggested as a relevant source of ice-nucleating particles, particularly in remote marine environments (Wilson et al., 2015). The EC-Earth3-AerChem FORCeS model version includes a new source of Marine Primary Organic Aerosols (MPOA), that considers the partitioning between insoluble marine organics and sea salt. The MPOA emission is calculated as a fraction of the submicron sea salt aerosols dependent on the chlorophyll-a (Chl-a) present in the ocean surface layer (O'Dowd et al., 2008; Vignati et al., 2010). In addition, a coarse mode MPOA is included (Facchini et al., 2008; Myriokefalitakis et al., 2010), which also depends on the online calculated sea salt emission (Gong, 2003). The model uses Chl-a concentration, a monthly averaged product derived from MODIS (Moderate Resolution Imaging Spectroradiometer) satellite observations.

2.2.4 Primary and secondary ice production

In regions such as the remote Arctic/Antarctica where the primary ice nucleating particles (INP) are sparse (Wex et al., 2019; Moore et al., 2024), a considerable source of ice crystals is via secondary ice production (SIP) processes in supercooled clouds (temperatures higher than approximately -10°C) (Järvinen et al., 2022; Järvinen et al., 2023). This process by which ice crystals are generated outnumber those generated by primary INPs (Field et al., 2017). The mechanisms that contribute to the INP formation are not adequately represented in the models resulting in a significant underestimation of the observed ice crystal number concentrations which in turn would impact the radiative forcing estimates (Vergara-Temprado et al., 2018; Zhao et al., 2023). In this project, an aerosol-and-temperature-sensitive parameterization (Costa-Surós et al., in preparation) has been developed to substitute the temperature-based parameterization by Meyers et al. (1992) for estimating the ice crystal number concentrations (ICNC) in mixed phase clouds. Specifically, it considers the primary ice crystal formation by immersion freezing dependent on K-feldspar and quartz (Harrison et al., 2019), deposition nucleation of soot and dust (Ullrich et al., 2017) and immersion freezing of marine organic aerosols (Wilson et al., 2015). The ICNCs formed via SIP are quantified

using the RaFSIP (Random Forest Secondary Ice Production) scheme version 1, as detailed in Georgakaki and Nenes (2024).
150 RaFSIP is a data-driven parameterization developed from a two-year simulation with a 10-km horizontal grid spacing, covering the period from 2016 to 2017 over the pan-Arctic region using the Weather Research and Forecasting (WRF) model (<https://www.mmm.ucar.edu/models/wrf>) with explicit SIP microphysics (Sotiropoulou et al., 2021; Georgakaki et al., 2022). Unlike conventional SIP parameterizations (for e.g., Phillips et al., 2017), the RaFSIP scheme requires only a few input variables, offering a streamlined and easily implementable solution for large-scale models without detailed microphysics schemes,
155 which would not otherwise support the explicit treatment of SIP.

The effect of SIP through collisional break-up, droplet-shattering and Hallett-Mossop rime splintering is described in the model through the introduction of an Ice Enhancement Factor (IEF) - a multiplication factor applied to primary ice production rates. The notion of the IEF is often used in the literature to indicate the prevalence of SIP. This approach shows promise for parameterizing SIP effects in mixed-phase clouds as it allows direct comparison with IEF (ICNCs/INPs) derived from in-situ
160 observations or retrieved from remote-sensing data (for e.g., Wieder et al., 2022).

2.2.5 ISORROPIA-Lite for inorganic aerosols

A new thermodynamic module for inorganic aerosols, ISORROPIA-Lite (Kakavas et al., 2022), is also implemented in the updated EC-Earth3-AerChem model replacing EQSAM (Metzger et al., 2002). ISORROPIA-lite is based on the ISORROPIA-II (Fountoukis and Nenes, 2007) code and it treats the thermodynamics of aerosol containing Ca^{2+} , K^+ , Mg^{2+} , SO_4^{2-} , Na^+ ,
165 NH_4^+ , NO_3^- , Cl^- , H_2O and their equilibrium with gas-phase HNO_3 , NH_3 , HCl and H_2O . Furthermore, the bulk calculation of ammonium nitrate has been replaced by one which distributes these components over the accumulation and coarse soluble modes.

2.2.6 Model tuning

The model undergoes a tuning process after major updates to minimize the radiative imbalance at the top of the atmosphere
170 (TOA) and at the surface (global mean net radiative flux at the TOA, TOA and surface SW and LW flux, cloud radiative effects in LW and SW) in AMIP simulations. This involves adjusting specific parameters with respect to CERES-EBAF data set (see Section 2.3) that is used here as observational reference. In this study, the following three key parameters are adjusted for the tuning of the EC-Earth3-AerChem model:

- the conversion efficiency parameter, RPRCON, which determines the rate of autoconversion of cloud water to rain, is
175 increased from $1.34 \cdot 10^{-3}$ to $1.41 \cdot 10^{-3}$,
- the critical radius for autoconversion is reduced from 8.75 to 7.5 μm and
- the standard deviation of the updraft velocity distribution is reduced by 0.2 m/s (from 0.8 m/s to 0.6 m/s) in the cloud droplet activation process.

Prior to these adjustments, the combined model updates resulted in a TOA net radiation imbalance of more than -3 W/m^2 in comparison to the CMIP6 version of the model which had an imbalance of -0.46 W/m^2 . By fine-tuning the aforementioned parameters, the model is closer to a radiative balance with a net TOA imbalance of -0.7 W/m^2 .

In the following we will refer to the CMIP6 configuration of the EC-Earth3-AerChem model as ECE3-CMIP6 while the configuration including FORCeS updates will be referred to as ECE3-FORCeS.

2.3 Observational datasets

To evaluate the impacts of the model updates detailed in Section 3, we use a suite of satellite-based datasets of cloud and aerosol properties. These are described below.

MODIS-Aqua: Retrievals of cloud properties from the MODIS onboard the Aqua satellite in the framework of NASA's Earth Observing System (EOS) are used. We specifically use Collection 6 Level-2 and Level-3 (MYD06_L2 and MYD08_L3) information on cloud fraction, optical thickness, liquid water path and droplet effective radius. Eighteen years of data from 2003 through 2020 is used.

CERES-EBAF: The radiative flux components from the Clouds and the Earth's Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) dataset are used to evaluate the top of the atmosphere shortwave and longwave fluxes. The latest Edition 4.2 data at Level-3b is used for the analysis (Loeb et al., 2018; Kato et al., 2018). The time period from 2000-2016 is made use of in this study.

CPR-CloudSat: The latest Release 5 of CloudSat Level 2B-CWC-RVOD and 2B-CWC-RO (CloudSat Radar-Only Cloud Water Content) products are used to evaluate the cloud liquid (CLWP) and ice water paths (CIWP) respectively. These products are derived for each radar profile as seen by CloudSat's Cloud Profiling Radar (CPR) for those profiles for which clouds are likely based on the radar profile analysis (https://www.cloudsat.cira.colostate.edu/cloudsat-static/info/dl/2b-cwc-ro/2B-CWC-RO_PDICD.P1_R05.rev0_.pdf). In 2B-CWC-RVOD product, the retrievals of CLWP are constrained using cloud optical depth information from MODIS (<https://www.cloudsat.cira.colostate.edu/data-products/2b-cwc-rvod>). Unlike the passive sensors, the CPR onboard CloudSat can sense the entire cloud column, day and night, thus providing better estimates of CLWP and CIWP. We use the data from 2007 through 2011.

CALIPSO: We further use the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO) Lidar Level-3 Global Energy and Water Cycle Experiment (GEWEX) Cloud, Standard Version 1-00 data product (CAL_LID_L3_GEWEX_Cloud-Standard-V1-00) to obtain information on total cloud fraction and its subdivision into low-, medium- and high-cloud fraction (NASA/LARC/SD/ASDC, 2019). This latest Level-3 version is based on the version 4.20 Level-2 5-km merged layer product for the period 2007-2016.

The climatologies of cloud and aerosol properties from these datasets averaged over their respective time periods are statistically compared with the model simulations in Section 3.

210 2.4 Maximum Covariance analysis: description and methodology

Maximum Covariance Analysis (MCA) (Bretherton et al., 1992; Cherry, 1996) is a statistical technique that extracts coherent patterns in two data sets that explain the maximum fraction of the covariance between them. This means that the analysis identifies the regions where the two fields co-vary to a maximum extent in the different spatial modes. The resulting modes are orthogonal to each other making the patterns complex and difficult to interpret physically. Hence, 'rotated' MCA is employed here to maximize the separation between the patterns by relaxing the orthogonality constraint thereby improving their interpretability. Varimax rotation technique is used in this study. This analysis provides two maps: 1. homogeneous regression map and 2. heterogeneous regression map. The homogeneous patterns are the correlation co-efficients between the input data of one field and scores (see the section below on how scores are calculated) of the same field, whereas the heterogeneous patterns are the correlation co-efficients between the input data of one field and scores of the second field and vice versa. The resulting heterogeneous maps are a typical characteristic of the MCA as they bring out the covariability between the two data sets. The scores are similar to the principal components (PCs) that captures the amplitude and temporal variation or in other words, the variability associated with each spatial pattern (or mode). The squared covariance fraction (SCF) is an invariant quantity even if the modes are modified or transformed in some way. This fraction assesses the relative importance of each mode.

2.4.1 Mathematical background

225 A brief overview of the MC analysis is given below.

Assume two data sets, X and Y, with variables a and b respectively which can be represented as matrices, X(a x n) and Y(b x n), n is the number of observations. Both X and Y are first standardized and detrended.

The scores are defined as projections of X and Y on to the singular vectors of the cross covariance matrix of X and Y. The cross covariance matrix C_{xy} are computed as in the equation below.

230
$$C_{xy} = \left(\frac{1}{n}\right)XY^T$$

where Y^T is the transpose matrix of Y.

A singular value decomposition is performed on the cross covariance matrix as,

$$C_{xy} = UDV^T$$

where U and V are correspondingly the eigen vectors for the two data sets, X and Y and D is a diagonal matrix with singular values.

The scores are then calculated as,

$$A_x = U^T X$$

$$A_y = V^T Y$$

The homogeneous patterns for X and Y fields are defined as,

240 $\text{hom}_X = \text{Corr}(X, A_x)$

$$\text{hom}_Y = \text{Corr}(Y, A_y)$$

Similarly, the heterogeneous patterns for X and Y fields are defined as,

$$\text{het}_X = \text{Corr}(X, A_y)$$

$$\text{het}_Y = \text{Corr}(Y, A_x)$$

245 where 'Corr' refers to the correlation co-efficient between the two fields within the brackets which is calculated as in the equation below. For example,

$$\text{Corr}(X, A_x) = \frac{\sum (X - \bar{X})(A_x - \bar{A}_x)}{\sqrt{\sum (X - \bar{X})^2 \sum (A_x - \bar{A}_x)^2}}$$

where the summation, \sum is the summation over total number of observations.

The squared covariance fraction quantifies the extent to which each mode i accounts for the explained proportion of the total squared covariance and is defined as,

250
$$\text{SCF}_i = \frac{\sigma_i^2}{\sum_{i=1}^m \sigma_i^2}$$

where m is the total number of modes and σ_i is the i^{th} singular value of the covariance matrix.

3 Results-I: Evaluation of the recent improvements in ECE3-FORCeS model

Historical atmosphere-only (AMIP) FORCeS simulations for the period 1980-2020 are used for the analysis. The comparisons
 255 are made with the corresponding CMIP6 simulations that cover a period from 1980 - 2018. Both AMIP simulations are fully compliant with the CMIP6 AMIP protocol, but have been extended beyond the CMIP6 historical period using anthropogenic and biomass burning emissions of ozone and aerosol precursors, greenhouse gas concentrations and other forcings following the CMIP6 protocol for the SSP2-4.5 scenario experiment. SSTs and sea ice fractions are prescribed using the CMIP6 version (1.1.8) of the PCMDI forcing data set ([https://www.wdc-climate.de/ui/cmip6?input=input4MIPs.CMIP6.CMIP.PCMDI.PCMDI-](https://www.wdc-climate.de/ui/cmip6?input=input4MIPs.CMIP6.CMIP.PCMDI.PCMDI-AMIP-1-1-8.ocean.mon.tosbcs.gn.v20220622)
 260 [AMIP-1-1-8.ocean.mon.tosbcs.gn.v20220622](https://www.wdc-climate.de/ui/cmip6?input=input4MIPs.CMIP6.CMIP.PCMDI.PCMDI-AMIP-1-1-8.ocean.mon.tosbcs.gn.v20220622)), which extends out through 2021. The forcing dataset is based on UK MetOffice HadISST and NCEP OI2 (Durack et al., 2022).

To facilitate the evaluation of the model simulated variables against satellite observations in a consistent manner, the COSP (Cloud Feedback Model Intercomparison Project (CFMIP) Observation Simulator Package) (Bodas-Salcedo et al., 2011) that is developed by the CFMIP community is available in EC-Earth3-AerChem. COSP provides tools and algorithms to simulate
 265 what a climate model would "see" from the viewpoint of the satellite that is used. This allows for a more direct comparison with satellite observations (Pincus et al., 2012).

In the following sections, we focus on the evaluation of the ECE3-FORCeS version and provide a comparison against both the ECE-CMIP6 version and observations.




Figure 1. Climatological mean total, low-, middle- and high-cloud fractions respectively in the four columns from (a) CALIPSO observations, (b) COSP-CALIPSO simulations from ECE3-FORCeS and their differences with respect to (c) CALIPSO observations and (d) ECE3-CMIP6 simulations. The values in the brackets refer to the global mean of each category. The observations extend from 83°S - 83°N.

3.1 Evaluation of COSP-simulated cloud fractions

270 In this section, simulated cloud and aerosol from the ECE3-FORCeS version of the model are evaluated against satellite observations and compared against the simulations of ECE3-CMIP6 version of the model. Here, the simulated cloud parameters derived from the COSP simulator of the model are evaluated.

The EC-Earth3-AerChem model outputs the following COSP parameters: CALIPSO-COSP total, low-, middle- and high-cloud fractions, MODIS-COSP total, water and ice cloud fractions. The CALIPSO-COSP simulated climatological mean total and the cloud fractions at three altitudes are presented in Fig. 1 (b). To facilitate the evaluation, CALIPSO observations (a)

275

and the differences of the simulated values with respect to CALIPSO observations (c) and CMIP6 simulations (d) are also shown. The values in brackets correspond to the global mean. The ECE3-FORCeS model simulates the spatial distribution of the total cloud fraction reasonably well compared to the observations, with an overestimation in the polar latitudes. This overestimation stems from the biases which is primarily seen in the low-cloud fraction. The high-cloud fraction, on the other hand, is considerably underestimated globally with respect to the observations particularly over the oceanic regions west of South America, Africa and Australia. This explains the underestimation of the total cloud fraction over the oceanic regions by 20-30% with respect to the observations. The simulated global mean total cloud fraction is 65%, whereas the observed mean fraction is 69%. The aerosol and cloud processes have resulted in an increase in the total cloud fraction in the high latitudes in both hemispheres with respect to the corresponding ECE3-CMIP6 simulations and this increase can be attributed to the increase in low-cloud fraction and partly to an increase in middle-cloud fraction. This increase in low cloud fraction can be attributed to updates in the model, such as the cloud droplet activation that targets liquid water clouds. No significant change is seen in the tropics and mid-latitudes with the model updates.

The ECE3-FORCeS and ECE3-CMIP6 zonal averaged climatological mean total, low-, middle- and high-cloud fractions are evaluated against CALIPSO observations (see Fig. 2 (a) - (d)). The black solid lines in each category of the cloud fraction refer to the observed value. The total cloud fraction in the Southern Hemisphere (SH) high latitudes is much closer to the observations in the ECE3-FORCeS model. It can be clearly seen that the improvements in the ECE3-FORCeS version of the model do not bring a significant change in the cloud fraction over the tropics and mid-latitudes compared to its ECE3-CMIP6 counterpart. However, the total and the low-cloud fractions in ECE3-FORCeS simulations in the Northern Hemisphere (NH) high latitudes tend to deviate even further from the observations compared to those in ECE3-CMIP6 runs.

A comparison of the simulated climatological mean total/water/ice cloud fractions with MODIS observations (Fig. 3) shows that though the model simulates the spatial distribution of total cloud fraction well. However, it fails to reproduce the magnitude globally, particularly, over the tropics and mid latitudes as can be clearly seen in the difference plot (Fig. 3(c)). The model captures the liquid water cloud fraction more realistically and, over the aforementioned regions, there is a slight overestimation of about 10-15% compared to the observations. The global mean values of the water cloud fraction are close to 30% in both observations and the simulation. The ice cloud fraction, on the other hand, is overestimated over the mid and high latitudes in both hemispheres in the model, thereby overestimating the global mean values of the ice cloud fraction by 8%. It can be noted that the simulated ice cloud fraction over the stratocumulus cloud decks is comparable to what is seen by MODIS. This would indicate that the mixed cloud fraction in the model is underestimated which could partly explain the underestimation seen in the simulated total cloud fraction across these regions.

3.2 Evaluation of cloud radiative effects

The cloud radiative effects (CRE) at the TOA is defined as the difference between the TOA all-sky and clear-sky fluxes. The spatial distribution and zonal means of the simulated climatological mean shortwave (SW) and longwave (LW) TOA CRE are evaluated against CERES-EBAF observations and the results are illustrated in Fig. 4 as differences in the simulated values from the observed values. The global means are shown in brackets. The overestimation of the SW cloud forcing in the ECE3-

Zonal mean cloud fraction: CALIPSO obs. vs CALIPSO-COSP simulations




Figure 2. Zonally averaged climatological mean CALIPSO-COSP simulated (a) total, (b) low-, (c) middle- and (d) high-cloud fractions. Solid black lines correspond to the CALIPSO observations. The green dashed lines are the ECE3-CMIP6 simulations and the dotted yellow lines correspond to the ECE3-FORCeS simulations.

310 FORCeS simulations over the Southern ocean and along the equator and over the pollutant outflow regions of Africa, North and South Americas and eastern China is more than halved compared to ECE3-CMIP6 simulations where the biases were more than 40 W/m^2 (Fig. 4, column #1). The global mean biases in the SW CRE is only one-tenth in the ECE3-FORCeS model compared to its previous model version.

On the other hand, the LW CREs are underestimated, particularly along the equatorial belt (Fig. 4, column #2). The slight
 315 positive biases in LW in ECE3-CMIP6 simulations in the stratocumulus regions over the western coast of the continents is reduced in ECE3-FORCeS. However, we overestimate the LW forcing in the NH high latitudes and over the Southern Oceans in the ECE3-FORCeS model. The global mean LW CRE shifted from a negative bias (-0.96) to a slightly more positive bias (0.55). There is a considerable improvement in the SW CRE with the FORCeS model updates which can be seen clearly
 320 in the zonal mean plots shown to the right in Fig. 4. The overestimation in the SW CRE at the TOA is notably reduced by more than 20 W/m^2 over the Southern ocean. In contrast, there is an overestimation in the LW CRE at latitudes $> 50^\circ$ in the FORCeS version, with biases in the range of $3\text{-}5 \text{ W/m}^2$, while in the mid-latitudes (between 50°S and 50°N), there is a slight improvement.




Figure 3. Climatological mean total (left), liquid water (middle) and ice (right) cloud fractions respectively in the three columns from (a) MODIS observations (b) ECE3-FORCeS simulations and (c) their difference with respect to MODIS observations.

3.3 Evaluation of cloud microphysical properties

Measuring cloud water is crucial for investigating microphysical processes and the indirect impacts of aerosols. Cloud water is commonly represented using column-integrated metrics like cloud liquid water path and cloud ice water path. Since the COSP simulator for CloudSat products is not available, we use the standard model output for CLWP and CIWP for comparison with CloudSat retrievals. This comparison could still be reasonable as CloudSat can see through the whole cloudy column. Furthermore, using the MODIS visible optical depth constrained product for CLWP and Radar-Only product for CIWP provide the best reference for cloud water free from the contamination by precipitation signal.

The spatial distribution of the ECE3-FORCeS simulated climatological mean fields against CloudSat observations and the differences from ECE3-CMIP6 simulations are presented in Fig. 5. These maps need to be interpreted with caution as the color bar range varies between model and observations. Despite the model's realistic simulation of spatial distribution, both CLWP and CIWP are markedly underestimated when compared to the actual observations. Following the FORCeS updates, CLWP increased over high- and mid-latitudes in both hemispheres, while decreased over the tropics compared to the ECE3-CMIP6 version of the model. A significant global decrease of up to 3.5 g/m^2 in CIWP is simulated in the ECE3-FORCeS model with regional decreases of up to $10\text{-}15 \text{ g/m}^2$, particularly over the oceans in the NH and SH.

The simulated climatological mean CD effective radius is evaluated against MODIS observations as in Fig. 6. The global mean values are shown in the brackets for the same region (75°S - 75°N) as for the MODIS observations. The magnitude and spatial pattern along the equatorial oceans are very close to the observations. However, the droplets sizes are comparatively

Climatological mean cloud radiative effect @TOA (W/m^2)




Figure 4. (left) Climatological mean CRE (in W/m^2) in SW (left) and LW (right) at the TOA from (a) CERES-EBAF observations, (b) differences of CMIP6 simulations and (c) ECE3-FORCeS simulations from observations. (right) Zonally averaged climatological mean CRE in SW and LW shown as differences from the observations.

340 smaller in the simulations than in the observations. The observed global climatological mean is $14.7 \mu m$ compared to a value of $7 \mu m$ in the ECE3-FORCeS model simulation. The differences in the CD effective radius are in the range of uncertainty of the observations wherein the biases in regional monthly mean MODIS derived values are at least 1 to $10 \mu m$ depending on cloud horizontal heterogeneity and solar zenith angle (Fu et al., 2019). Notably, the latest model updates have led to an increase in CD effective radius, especially over the tropical oceans, as observed in comparison to the ECE3-CMIP6 simulations.

345 4 Results-II: Maximum Covariance Analysis: Covariability between AOD and CD effective radius

4.1 Regions selected for the analysis

In order to investigate the maximum covariance between AOD and CD effective radius, we select the regions based on two prerequisites: a) the regions shall sustain low-level liquid water clouds and b) the regions experience at least some degree of pollution outflow from the continents. Among all the oceanic cloud regimes, these low-level clouds are most likely to
 350 be influenced by aerosols and stand the highest chance of testing the hypothesis that aerosol-cloud covariability can also be visible in the larger spatio-temporal scales. We selected three such regions based on our previous study, Devasthale and Thomas

Climatological mean cloud liquid and ice paths (g/m^2)




Figure 5. Climatological mean cloud liquid water (left) and ice (right) paths (g/m^2) from (a) CloudSat observations, (b) ECE3-FORCeS simulations and (c) their differences with respect to ECE3-CMIP6 simulations. The global mean values are given in the brackets.




Figure 6. Climatological mean CD effective radius (μm) from MODIS observations (left), ECE3-FORCeS simulations (center) and their differences with respect to ECE3-CMIP6 simulations (right). The global mean values are given in brackets.

(2011), which investigated the frequency of occurrence of aerosol-cloud overlap globally in different seasons. They are shown in Fig. 7 by the blue boxes.

The table below provides details about the selected regions, each distinguished by specific types of aerosols. For instance, the Eastern China region is marked by anthropogenic aerosols, persisting throughout the year but with a reduced intensity during

355




Figure 7. Frequency of aerosol-cloud overlap estimated based up on 4 years (May 2006 - June 2010) of CALIPSO satellite observations (Devasthale and Thomas, 2011). The regions selected are marked by the blue boxes.

the June-July-August season due to pollutant removal by wet deposition. The remaining two regions, designated as BB1 and BB2 respectively, for the west coast of Africa and the west coast of South America, are notably influenced by biomass burning aerosols in the free troposphere and marine aerosols in the boundary layer (Bourgeois et al., 2015, 2018). The prevalence of transported biomass burning aerosols is higher during September - November for Africa and June - November for South America during the dry season.

Table 1. Regions selected for the study. 'BB' refers to biomass burning.

Regions	longitude; latitude	time periods
Eastern China	95 E - 150 E; 20 N - 50 N	All months
BB1: Africa	10 W - 20 E; 25 S - 5 S	Sept - Nov
BB2: South America	100 W - 70 W; 15 S - 5 N	Jun - Nov

The climatological mean AOD and low-cloud fraction for the three regions selected for this study are shown in Fig. 8. Among these regions, Eastern China experiences the highest AOD, reaching 0.7, in the area of maximum anthropogenic pollution. In the other two regions, AOD peaks over land with values around 0.25, corresponding to areas where biomass burning occurs. The Andes mountain range, situated to the west of the South American continent, acts as a barrier that partially hinders the transport of pollutants to the oceans. The low-cloud fraction ranges from 0.3 to 0.7 over these regions. It is also to be noted that the regions where the maximum pollution outflow over the oceans occurs and the regions where the maximum frequency of low-level liquid clouds is observed are not the same. The maximum aerosol-cloud overlap in fact occurs over the regions further remote from the individual maxima in AOD and cloud fraction.




Figure 8. Climatological mean AOD (left) and low-cloud fraction (right) over the selected three regions: (a) Eastern China (b) BB1: Africa (blue box) and (c) BB2: South America (blue box).

4.2 MC analysis over Eastern China

370 As explained in Section 2.4, maximum covariance analysis is applied to assess the covariability between AOD and CD effective
radius across Eastern China. Before applying the MCA, both these variables are deseasonalized and detrended. The hetero-
geneous patterns, the patterns that brings out the areas of covariability between the two fields, are presented. Only regions
exhibiting statistical significance at the 95% confidence level are showcased. We present the first three modes derived from the
analysis for Eastern China in Fig. 9 (a) - (c). These three modes collectively explain 66% of the total observed variability. A dis-
375 tinct pattern exhibiting opposite signs between AOD (left) and CD effective radius (right) emerges and each mode captures the
different regions where this inverse covariability is dominant. The opposite signature between these two fields clearly suggests
that the Twomey effect over those regions in the corresponding modes can influence the variability at the large spatio-temporal

380 scales. The first two modes capture the covariability over the outflow regions where the aerosol advection over nearby oceanic low-level clouds is quite frequent (Devasthale and Thomas, 2011), whereas the third mode captures the covariability over the continental source regions. It is intriguing that the covariability between AOD and CD effective radius is not only noticeable in oceanic outflow regions but also extends to land regions. This is mainly due to the fact that different cloud regimes and thermodynamical conditions exist over the land and the oceanic regions in the selected study area. The covariability is especially stronger just off the Eastern Chinese coast in the first two modes, suggesting a noticeable influence of pollution advection on the low-level clouds, possibly dominating over other factors that can also cause covariability.




Figure 9. Statistically significant heterogeneous patterns over Eastern China in the three modes (a)-(c) derived from the maximum covariance analysis.

385 **4.3 MC analysis over BB1: Africa**

The maximum covariance analysis applied to AOD and CD effective radius over this region reveals that only the first mode is the most significant and dominant mode explaining a variability of up to 20%. The remaining modes do not exhibit statistically significant covariability and the SCFs in the subsequent modes are less than 10%. So, here we focus only on the first mode. The covariability seen in this only one mode is nonetheless very significant and noteworthy, as this mode exhibits the regions
390 where the Twomey effect is likely dominant, particularly over the pollutant outflow regions. The heterogeneous pattern in these two fields for this mode is shown in Fig. 10.

As mentioned earlier, this study region is dominated by the seasonal biomass burning in southern Africa that occurs during the months of June through November, transporting enormous amounts of absorbing aerosols across the southeast Atlantic over the extensive stratocumulus decks. This outflow region has therefore been the center of many measurement campaigns
395 and studies focusing on characterizing aerosols and clouds and their interactions (see, for e.g., Redemann et al., 2021). To what extent these aerosols affect the underlying stratocumulus decks is still the topic of intense research. Although the freshly emitted biomass burning aerosols are not ideal condensation nuclei, numerous studies have previously shown that the morphology, composition and size distribution can change considerably during the transport, making them one of the dominant local sources of condensation nuclei, thereby increasing cloud droplet number concentration and changing droplet size (Petters et al., 2009;
400 Lu et al., 2018; Che et al., 2022; Royer et al., 2023).

The significant covariability between AOD and CD effective radius here indeed suggests that as these biomass burning aerosols are advected over the oceanic stratocumulus regions in the model simulation, they undergo aging and mixing with other anthropogenic aerosols as well as marine aerosols, possibly becoming more efficient CCNs. In the ECE3-AerChem model, the open biomass burning emissions are distributed up to 2 kms following the vertical profiles for forest fires as given in Table
405 A1 of van Noije et al. (2014). The accompanying advection of humidity however makes the aerosol-cloud interactions over these regions less predictable as the local meteorology together with the changing aerosol aging, coating and size distribution introduces high internal variability.

4.4 MC analysis over BB2: S. America

Although the frequency of aerosol-cloud overlap is also high in this outflow region, the weakest covariability between AOD
410 and CD effective radius is observed over the ocean compared to previous two study areas as shown in Figure 11. Although the first mode alone accounts for 59% of the total variability, the spatial extent of the statistically significant covariability and its magnitude is limited and weak over the ocean compared to the land. A number of factors that can affect the variability in AOD and clouds need to be considered here. The aerosols need to be convected and advected over much longer distances in the free troposphere crossing the Andes mountain range (Bourgeois et al., 2015, 2018). The high residence time and aging during
415 the transport likely lead to strong changes in aerosol properties and size distribution before they are available as condensation nuclei after the descent over the cooler upwelling waters off the western coast of south America. Furthermore, the large-scale dynamical variability associated with the El Niño Southern Oscillation (ENSO) strongly influences the variability of low-




Figure 10. Statistically significant heterogeneous patterns in mode 1 derived from the MCA over biomass burning (BB1) outflow region of Africa.

level cloudiness in this region. ENSO also likely induces variability in the background marine aerosols through changes in sea surface temperatures and winds. Therefore, the correspondence between AOD and CD effective radius at larger temporal
420 scales is much weaker.




Figure 11. Statistically significant heterogeneous patterns in mode 1 derived from the MCA over biomass burning (BB2) outflow region of South America.

4.5 Implications for cloud radiative effects

Additionally, we examined the implications of this covariability between AOD and CD effective radius for the cloud radiative effects. To investigate this, a composite analysis is carried out based on the PCs of AOD as shown in Fig. 12 as an example for mode 1 over Eastern China.





Figure 12. The monthly principal components of AOD corresponding to the mode 1 of the MCA.

425 Two composites are generated, one for which the PCs of AOD are greater than zero and another where the PCs of AOD
are less than zero. In the composite analysis for PCs of AOD > 0 , we investigated the anomalies of TOA SW CRE, AOD,
CD effective radius, CLWP, CDNC (Cloud droplet number concentrations), specific humidity, and low-cloud cover fraction.
The composite analysis for the PCs of AOD < 0 is exactly the opposite of those obtained for PCs of AOD > 0 . For brevity,
we present the outcomes (Fig. 13 for the three modes specifically over Eastern China for one composite analysis when PCs of
430 AOD > 0 .

In this composite, the anomalies in TOA SW CREs indicate pronounced cooling over the pollution source regions and
the pollutant outflow region in modes 1 and 2, while the opposite is observed in mode 3 as can be seen in Fig. 13. The
significant cooling observed in the first two modes can stem from two processes, either enhanced aerosol-cloud interactions
(Twomey effect) and/or to an increase in low-cloud cover, leading to brighter clouds and increased reflection. Regarding the
435 first hypothesis, the AOD anomalies are in fact lower in this composite together with a corresponding positive anomaly in CD
effective radius, indicative of larger droplets. The cloud droplet number concentration is also lower. This indicates that the
Twomey effect cannot explain the cooling anomalies in the TOA SW CREs. However, a higher low-cloud fraction in tandem
with higher CLWP and specific humidity is seen. This higher low-cloud fraction can adequately account for the marked cooling
in TOA SW CREs. Consequently, the results show that, despite the existence of robust aerosol-cloud interactions at the larger
440 spatio-temporal scales, the TOA SW CREs are predominantly driven by the changes in low-level cloud fraction in these
simulations. Previous studies (Gryspeerd et al., 2019) have shown the inverse relationship between CDNC and CLWP over the
heavily polluted regions based on satellite observations. The results of this composite analysis are in line with those previous
studies, suggesting that CDNC-CLWP control can indeed have larger impact on cloud radiative effects than the Twomey effect.

Similar conclusions can be derived for the first modes over the biomass burning regions of Africa and S. America. The 445 results are shown in the Appendix Figs A1, A2 for Africa and A3, A4 for S. America.






Figure 13. Each panel represents respectively the anomalies of TOA SW cloud radiative effects (W/m^2), AOD, CD effective radius (μm), specific humidity (g/kg), cloud liquid water path (g/m^2), CDNC (cm^{-3}) and low cloud fraction in the three modes over eastern China.

5 Discussions and conclusions

In the framework of the EU project FORCeS, a number of significant improvements in the representation of aerosol and cloud processes were implemented in the ECE3-AerChem model, ECE3-FORCeS. The ECE3-FORCeS includes now a representation of the atmospheric cycle of dust minerals that are relevant for their climate impacts, as well as a new source of marine organic aerosols. As a result, the absorption of radiation by dust in the SW is now dependent on the online calculated abundance of hematite. The model also incorporates an aerosol-sensitive scheme for the estimation of primary ice crystals, which depends on the abundance of quartz, k-feldspars and marine organics for the immersion freezing, and on dust and soot for

deposition nucleation. This primary ice can lead to secondary ice production in supercooled conditions, which is quantified via RaFSIP (Georgakaki and Nenes, 2024). A new warm cloud activation scheme by Morales-Betancourt and Nenes (2014) is also implemented, enabling a more accurate estimate for the number concentration of droplets formed at the time of maximum supersaturation. Finally, a new thermodynamic module for inorganic aerosols, ISORROPIA-LITE, has been also implemented in the model.

These changes and improvements are bound to have an impact on the cloud properties and their radiative effects. Therefore, we used 41 years of historical atmosphere-only (AMIP) simulations for the period 1980-2020 from this recent FORCeS model version to confront them with the satellite-based observations. A suite of satellite sensor retrievals and simulators are used to facilitate the comparison. Due to increased cloud fraction and cloud liquid water path compared to its previous CMIP6 version, the strong warm biases often seen over the Southern ocean is reduced significantly in the FORCeS version of ECE3-AerChem. The bias in the SW TOA CREs is reduced by nearly 50% over this region.

Using these simulations from the ECE3-FORCeS, we further carried out a maximum covariance analysis between AOD and CD effective radius. The aim of this exercise was not to identify any new climate process, but rather to test if the covariability between AOD and CD effective radius could also be seen at the larger spatial and temporal scales. Among the three oceanic regions that were chosen based on the high frequency of aerosol and low-level cloud overlap, the strongest statistically significant inverse covariability between AOD and CD effective radius was observed over the eastern Chinese outflow region. Here, the aerosol loading is generally quite high and the aerosol sources are mostly anthropogenic. The conditions favourable for allowing aerosol-cloud interactions, such as persistence of westerly winds, high humidity and aerosol load in the lower troposphere, favourable aerosol composition and size distribution over the outflow region likely lead to strong inverse covariability between AOD and CD effective radius seen even at larger spatio-temporal scales. This means that the Twomey effect can manifest itself even at a climate scale by dominating the other possible drivers of local variability. Note that the model simulations do not suffer from the limitations often discussed when using satellite data for similar analyses, such as seeing only cloud top information or sampling issues related to simultaneous aerosol and cloud retrievals. Furthermore, the most of the aerosol outflow in this region occurs in the lowermost troposphere and hence AOD could be indeed a good proxy for the cloud condensation nuclei.

In the case of the other two regions in the southeast Atlantic (along the west coast of Africa: BB1) and eastern equatorial Pacific (along the west coast of S. America: BB2), biomass burning is a significant source of aerosols during the months selected for the study. The covariance between AOD and CD effective radius is statistically significant only in the first mode of variability in these regions. However, it can be seen that the covariance is more pronounced over the pollutant outflow region in BB1 and the source region in BB2. This is most likely due to fact that the aerosol vertical distribution and its variability is more disengaged from the variability in the underlying cloud decks due to longer transport distances in BB2 due to the Andes mountain range and the subsequent changes in aerosol composition and size distribution that may not favor the covariability. In such case, the total AOD may indeed not be a proxy for cloud condensation nuclei, as argued by the previous studies (Quaas et al., 2020). Our analysis also show a strong cooling/warming in the TOA SW CREs despite strong covariability (larger droplets with low AOD and vice versa) and can be attributed to stronger/weaker control by the relationship between CDNC

and CLWP. Further investigations are underway to carry out similar study using satellite observations, but the lack of reliable, multidecadal data of AOD, CCN and CDNC is currently making such an analysis challenging.

490 The updates to the EC-Earth3-AerChem model described in this work improve the representation of aerosols and aerosol-cloud interactions. They address previously missing processes, such as secondary ice particles, and improve existing parameterizations, such as, cloud droplet activation. These modifications make the model more realistic and closer to what is observed, but, there are still biases in the cloud microphysical properties. One of the reasons may be that the model was re-tuned using a subset of the parameters identified in the tuning strategy of the CMIP6 version of the model. However, finding a new set of
495 tuning parameters to improve clouds while maintaining radiation balance, cloud forcing, surface temperatures, precipitation patterns, etc., is challenging. A complete re-tuning was beyond the scope of this project. Future model developments aim to reduce biases through new parametrizations for updraft velocity and secondary ice production (RaFSIP v2). The extent to which these changes, along with re-tuning, could mitigate the biases requires further investigation. The goal is to incorporate these improvements that were achieved during the FORCeS project, into the next version of the EC-Earth model which will then be
500 used to contribute to CMIP7, particularly AerChemMIP, to provide a better understanding of the role of various aerosols in the climate and its sensitivity.

Code and data availability. The EC-Earth3 code is available from the EC-Earth development portal for members of the consortium. All code related to CMIP6 forcing is implemented in the component models. The model codes developed at ECMWF, including the atmosphere model IFS, are intellectual property of ECMWF and its member states. Permission to access the EC-Earth3 source code can be requested
505 from the EC-Earth community via the EC-Earth website (<http://www.ec-earth.org/about/contact/>, EC-Earth-Consortium (2023)) and may be granted if a corresponding software license agreement is signed with ECMWF. The repository tag for the version of EC-Earth that is used in this work is "projects/FORCeS". Currently, only European users can be granted access due to license limitations of the atmosphere model. However, the model code is made accessible to both the Editor and the reviewers throughout the review process. The component models NEMO, LPJ-GUESS, TM5, and PISM are not limited by their licenses.

510 All model output data are freely available from any ESGF data node as part of CMIP6 (<https://esgf-data.dkrz.de/search/cmip6-dkrz/>); search for `< model_id >=EC-Earth3-AerChem`, `< experiment_id >=amip`, and `< variant_label >=r1i1p1f1` for ECE3-CMIP6 or `< variant_label >=r1i1p4f1` for ECE3-FORCeS. The python scripts used in generating the figures in the manuscript, the python code for the calculation of maximum covariance analysis and the observational data for model evaluation are archived in Zenodo under <https://doi.org/10.5281/zenodo.10781927>




Figure A1. Anomalies of top of the atmosphere SW CRE (W/m^2), AOD and CD radii (μm) in the three modes over BB1: Africa




Figure A2. Anomalies of specific humidity (g/kg), CLWP (g/m^2) and low cloud fraction in the three modes over BB1: Africa




Figure A3. Anomalies of top of the atmosphere SW CRE (W/m^2), AOD and CD radii (μm) in the three modes over BB2: S. America




Figure A4. Anomalies of specific humidity (g/kg), CLWP (g/m²) and low cloud fraction in the three modes over BB2: S. America

515 *Author contributions.* MT and KW conceptualized the research. MT performed the analysis and drafted the manuscript. SW post-processed the raw model data. AD provided and analyzed the satellite data observations that were used in the analysis. MC, PG, MC-S, MGA, MK, AN and CPG-P were involved in the implementation of the different processes in the ECE3-AerChem model and contributed to the detailed text regarding these as described in Section 3 of the manuscript. KW, TvN and PLS designed and carried out the simulations. All the co-authors contributed to the interpretation of the results and review of the manuscript.

520 *Competing interests.* The authors have no competing interests to declare.

Acknowledgements. The authors acknowledge the funding from the Horizon European Union 2020 Framework Programme (FORCeS project; grant No. 821205). MC-S has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie (grant No. 754433) from H2020-MSCA-COFUND-2016 (STARS). CPG-P, MC, MC-S and MGA acknowledge the funding from the European Research Council under the H2020 Research and Innovation Programme through the Consolidator Grant, FRAGMENT (grant No. 773051), the AXA Research Fund through the AXA Chair on Sand and Dust storms at the Barcelona Supercomputing Center (BSC), and the support of the Department of Research and Universities of the Government of Catalonia to the Research Group Atmospheric Composition (Code: 2021890 SGR 01550). The model simulations and data storage were enabled by resources provided by the National Academic Infrastructure for Supercomputing in Sweden (NAISS) at Linköping University and has been partially funded by the Swedish Research Council (grant No. 2022-06725).

525

530 References

- Abdul-Razzak, H. and Ghan, S. J.: A parameterization of aerosol activation: 2. Multiple aerosol types, *J. Geophys. Res.: Atmos.*, 105, 6837–6844, <https://doi.org/10.1029/1999JD901161>, 2000.
- Ahn, S. H., Yoon, Y., Choi, T., Lee, J., Kim, Y., Lee, B., Ritter, C., Aas, W., Krejci, R., Ström, J., Tunved, P., and Jung, C. H.: Relationship between cloud condensation nuclei (CCN) concentration and aerosol optical depth in the Arctic region, *Atmos. Env.*, 267, 118 748, <https://doi.org/10.1016/j.atmosenv.2021.118748>, 2021.
- 535 Albrecht, B.: Aerosols, cloud microphysics and fractional cloudiness, *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.
- Andreae, M. O.: Correlation between cloud condensation nuclei concentration and aerosol optical thickness in remote and polluted regions, *Atmos. Chem. Phys.*, 9, 543–556, <https://doi.org/10.5194/acp-9-543-2009>, 2009.
- 540 Atkinson, J. D., Murray, B. J., Woodhouse, M. T., Whale, T. F., Baustian, K. J., Carslaw, K. S., Dobbie, S., O’Sullivan, D., and Malkin, T. L.: The importance of feldspar for ice nucleation by mineral dust in mixed-phase clouds, *Nature*, 498, 355–358, <https://doi.org/10.1038/nature12278>, 2013.
- Balsamo, G., Beljaars, A., Scipal, K., Viterbo, P., van den Hurk, B., Hirschi, M., and Betts, A. K.: A revised hydrology for the ECMWF Model: Verification from field site to terrestrial water storage and impact in the integrated forecast system, *J. Hydromet.*, 10, 623 – 643, <https://doi.org/10.1175/2008JHM1068.1>, 2009.
- 545 Barahona, D., West, R. E. L., Stier, P., Romakkaniemi, S., Kokkola, H., and Nenes, A.: Comprehensively accounting for the effect of giant CCN in cloud activation parameterizations, *Atmos. Chem. Phys.*, 10, 2467–2473, <https://doi.org/10.5194/acp-10-2467-2010>, 2010.
- Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A. L., Dufresne, J. L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle, F., Mauritsen, T., McCoy, D. T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding aerosol radiative forcing of climate change, *Rev. Geophys.*, 58(1), <https://doi.org/10.1029/2019rg000660>, 2019.
- Bodas-Salcedo, A., Webb, M. J., Bony, S., Chepfer, H., Dufresne, J.-L., Klein, S. A., Zhang, Y., Marchand, R., Haynes, J. M., Pincus, R., and John, V. O.: COSP: Satellite simulation software for model assessment, *Bull. American Met. Soc.*, 92, 1023 – 1043, <https://doi.org/10.1175/2011BAMS2856.1>, 2011.
- 555 Bourgeois, Q., Ekman, A. M. L., and Krejci, R.: Aerosol transport over the Andes from the Amazon Basin to the remote Pacific Ocean: A multiyear CALIOP assessment, *J. Geophys. Res.: Atmos.*, 120, 8411–8425, <https://doi.org/10.1002/2015JD023254>, 2015.
- Bourgeois, Q., Ekman, A. M. L., Renard, J.-B., Krejci, R., Devasthale, A., Bender, F. A.-M., Riipinen, I., Berthet, G., and Tackett, J. L.: How much of the global aerosol optical depth is found in the boundary layer and free troposphere?, *Atmos. Chem. Phys.*, 18, 7709–7720, <https://doi.org/10.5194/acp-18-7709-2018>, 2018.
- 560 Bretherton, C., Smith, C., and Wallace, J.: An intercomparison of methods for finding coupled patterns in climate data, *J. Clim.*, 5, 541–560, <http://www.jstor.org/stable/26197080>, 1992.
- Carslaw, K. S., Boucher, O., Spracklen, D. V., Mann, G. W., Rae, J. G. L., Woodward, S., and Kulmala, M.: A review of natural aerosol interactions and feedbacks within the Earth system, *Atmos. Chem. Phys.*, 10, 1701–1737, <https://doi.org/doi:10.5194/acp-10-1701-2010>, 2010.
- 565

- Chatziparaschos, M., Daskalakis, N., Myriokefalitakis, S., Kalivitis, N., Nenes, A., Gonçalves Ageitos, M., Costa-Surós, M., Pérez García-Pando, C., Zanolì, M., Vrekoussis, M., and Kanakidou, M.: Role of K-feldspar and quartz in global ice nucleation by mineral dust in mixed-phase clouds, *Atmos. Chem. Phys.*, 23, 1785–1801, <https://doi.org/10.5194/acp-23-1785-2023>, 2023.
- 570 Che, H., Segal-Rozenhaimer, M., Zhang, L., Dang, C., Zuidema, P., Dobracki, A., Sedlacek, A. J., Coe, H., Wu, H., Taylor, J., Zhang, X., Redemann, J., and Haywood, J.: Cloud processing and weeklong ageing affect biomass burning aerosol properties over the south-eastern Atlantic, *Comm. Earth Env.*, 3(1), 182, 10.1038/s43247-022-00517-3, 2022.
- Cherry, S.: Singular value decomposition analysis and canonical correlation analysis, *J. Clim.*, 9, 2003–2009, <https://www.jstor.org/stable/26201389>, 1996.
- Claquin, T., Schulz, M., and Balkanski, Y.: Modeling the mineralogy of atmospheric dust sources, *J. Geophys. Res.: Atmos.*, 104, 22 243–22 256, <https://doi.org/10.1029/1999JD900416>, 1999.
- 575 Devasthale, A. and Thomas, M. A.: A global survey of aerosol-liquid water cloud overlap based on four years of CALIPSO-CALIOP data, *Atmos. Chem. Phys.*, 11, 1143–1154, <https://doi.org/10.5194/acp-11-1143-2011>, 2011.
- Di Biagio, C., Formenti, P., Balkanski, Y., Caponi, L., Cazaunau, M., Panguì, E., Journet, E., Nowak, S., Andreae, M. O., Kandler, K., Saeed, T., Piketh, S., Seibert, D., Williams, E., and Doussin, J.-F.: Complex refractive indices and single-scattering albedo of global dust aerosols in the shortwave spectrum and relationship to size and iron content, *Atmos. Chem. Phys.*, 19, 15 503–15 531, <https://doi.org/10.5194/acp-19-15503-2019>, 2019.
- 580 Döscher, R., Acosta, M., Alessandri, A., Anthoni, P., Arsouze, T., Bergman, T., Bernardello, R., Boussetta, S., Caron, L.-P., Carver, G., Castrillo, M., Catalano, F., Cvijanovic, I., Davini, P., Dekker, E., Doblás-Reyes, F. J., Docquier, D., Echevarria, P., Fladrich, U., Fuentes-Franco, R., Gröger, M., v. Hardenberg, J., Hieronymus, J., Karami, M. P., Keskinen, J.-P., Koenigk, T., Makkonen, R., Massonnet, F., Ménégos, M., Miller, P. A., Moreno-Chamarro, E., Nieradzìk, L., van Noije, T., Nolan, P., O'Donnell, D., Ollinaho, P., van den Oord, G., Ortega, P., Prims, O. T., Ramos, A., Reerink, T., Rousset, C., Ruprich-Robert, Y., Le Sager, P., Schmith, T., Schrödner, R., Serva, F., Sicardi, V., Sloth Madsen, M., Smith, B., Tian, T., Tourigny, E., Uotila, P., Vancoppenolle, M., Wang, S., Wårlind, D., Willén, U., Wyser, K., Yang, S., Yepes-Arbós, X., and Zhang, Q.: The EC-Earth3 Earth system model for the Coupled Model Intercomparison Project 6, *Geosci. Mod. Dev.*, 15, 2973–3020, <https://doi.org/10.5194/gmd-15-2973-2022>, 2022.
- 590 Durack, P. J., Taylor, K. E., Ames, S., Po-Chedley, S., and Mauzey, C.: PCMDI AMIP SST and sea-ice boundary conditions version 1.1.8, <https://doi.org/10.22033/ESGF/input4MIPs.16921>, 2022.
- EC-Earth-Consortium: IPCC DDC: EC-Earth-Consortium EC-Earth3-AerChem model output prepared for CMIP6 AerChemMIP. World Data Center for Climate (WDCC) at DKRZ, <https://doi.org/10.26050/WDCC/AR6.C6ACEEEEEEA>, 2023.
- Facchini, M. C., Decesari, S., Rinaldi, M., Carbone, C., Finessi, E., Mircea, M., Fuzzi, S., Moretti, F., Tagliavini, E., Ceburnis, D., and 595 O'Dowd, C. D.: Important source of marine secondary organic aerosol from biogenic amines, *Env. Sci. Tech.*, 42, 9116–9121, <https://doi.org/10.1021/es8018385>, 2008.
- Fanourgakis, G. S., Kanakidou, M., Nenes, A., Bauer, S. E., Bergman, T., Carslaw, K. S., Grini, A., Hamilton, D. S., Johnson, J. S., Karydis, V. A., Kirkevåg, A., Kodros, J. K., Lohmann, U., Luo, G., Makkonen, R., Matsui, H., Neubauer, D., Pierce, J. R., Schmale, J., Stier, P., Tsigaridis, K., van Noije, T., Wang, H., Watson-Parris, D., Westervelt, D. M., Yang, Y., Yoshioka, M., Daskalakis, N., Decesari, S., 600 Gysel-Beer, M., Kalivitis, N., Liu, X., Mahowald, N. M., Myriokefalitakis, S., Schrödner, R., Sfakianaki, M., Tsimpidi, A. P., Wu, M., and Yu, F.: Evaluation of global simulations of aerosol particle and cloud condensation nuclei number, with implications for cloud droplet formation, *Atmos. Chem. Phys.*, 19, 8591–8617, <https://doi.org/10.5194/acp-19-8591-2019>, 2019.

- Field, P. R., Lawson, R. P., Brown, P. R. A., Lloyd, G., Westbrook, C., Moisseev, D., Miltenberger, A., Nenes, A., Blyth, A., Choulaton, T., Connolly, P., Buehl, J., Crosier, J., Cui, Z., Dearden, C., DeMott, P., Flossmann, A., Heymsfield, A., Huang, Y., Kalesse, H., Kanji, Z. A., Korolev, A., Kirchgaessner, A., Lasher-Trapp, S., Leisner, T., McFarquhar, G., Phillips, V., Stith, J., and Sullivan, S.: Secondary Ice Production: Current state of the science and recommendations for the future, *Met. Monographs*, 58, 7.1 – 7.20, <https://doi.org/10.1175/AMSMONOGRAPHS-D-16-0014.1>, 2017.
- Fountoukis, C. and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for $K^+ - Ca^{2+} - Mg^{2+} - NH_4^+ - Na^+ - SO_4^{2-} - NO_3^- - Cl^- - H_2O$ aerosols, *Atmos. Chem. Phys.*, 7, 4639–4659, <https://doi.org/10.5194/acp-7-4639-2007>, 2007.
- Fu, D., Di Girolamo, L., Liang, L., and Zhao, G.: Regional biases in MODIS marine liquid water cloud drop effective radius deduced through fusion With MISR, *J. Geophys. Res.: Atmos.*, 124, 13 182–13 196, <https://doi.org/10.1029/2019JD031063>, 2019.
- Georgakaki, P. and Nenes, A.: RaFSIP: Parameterizing ice multiplication in models using a machine learning approach, *Journal of Advances in Modeling Earth Systems*, 16, e2023MS003923, <https://doi.org/10.1029/2023MS003923>, 2024.
- Georgakaki, P., Sotiropoulou, G., Vignon, E., Billault-Roux, A.-C., Berne, A., and Nenes, A.: Secondary ice production processes in winter-time alpine mixed-phase clouds, *Atmos. Chem. Phys.*, 22, 1965–1988, <https://doi.org/10.5194/acp-22-1965-2022>, 2022.
- Ghan, S. J., Abdul-Razzak, H., Nenes, A., Ming, Y., Liu, X., Ovchinnikov, M., Shipway, B., Meskhidze, N., Xu, J., and Shi, X.: Droplet nucleation: Physically-based parameterizations and comparative evaluation, *Journal of Advances in Modeling Earth Systems*, 3, <https://doi.org/doi.org/10.1029/2011MS000074>, 2011.
- Gong, S. L.: A parameterization of sea-salt aerosol source function for sub- and super-micron particles, *Global Biogeochem. Cyc.*, 17, <https://doi.org/10.1029/2003GB002079>, 2003.
- Gryspeerd, E., Goren, T., Sourdeval, O., Quaas, J., Mülmenstädt, J., Dipu, S., Unglaub, C., Gettelman, A., and Christensen, M.: Constraining the aerosol influence on cloud liquid water path, *Atmos. Chem. Phys.*, 19, 5331–5347, <https://doi.org/10.5194/acp-19-5331-2019>, 2019.
- Harrison, A. D., Lever, K., Sanchez-Marroquin, A., Holden, M. A., Whale, T. F., Tarn, M. D., McQuaid, J. B., and Murray, B. J.: The ice-nucleating ability of quartz immersed in water and its atmospheric importance compared to K-feldspar, *Atmos. Chem. Phys.*, 19, 11 343–11 361, <https://doi.org/10.5194/acp-19-11343-2019>, 2019.
- Huijnen, V., Williams, J., van Weele, M., van Noije, T., Krol, M., Dentener, F., Segers, A., Houweling, S., Peters, W., de Laat, J., Boersma, F., Bergamaschi, P., van Velthoven, P., Le Sager, P., Eskes, H., Alkemade, F., Scheele, R., Nédélec, P., and Pätz, H.-W.: The global chemistry transport model TM5: description and evaluation of the tropospheric chemistry version 3.0, *Geosci. Mod. Dev.*, 3, 445–473, <https://doi.org/10.5194/gmd-3-445-2010>, 2010.
- Järvinen, E., Nehlert, F., Xu, G., Waitz, F., Mioche, G., Dupuy, R., Jourdan, O., and Schnaiter, M.: Investigating the vertical extent and short-wave radiative effects of the ice phase in Arctic summertime low-level clouds, *Atmos. Chem. Phys.*, 23, 7611–7633, <https://doi.org/10.5194/acp-23-7611-2023>, 2023.
- Järvinen, E., McCluskey, C. S., Waitz, F., Schnaiter, M., Bansemer, A., Bardeen, C. G., Gettelman, A., Heymsfield, A., Stith, J. L., Wu, W., D’Alessandro, J. J., McFarquhar, G. M., Diao, M., Finlon, J. A., Hill, T. C. J., Levin, E. J. T., Moore, K. A., and DeMott, P. J.: Evidence for Secondary Ice Production in Southern Ocean Maritime Boundary Layer Clouds, *J. Geophys. Res.: Atmos.*, 127, e2021JD036411, <https://doi.org/https://doi.org/10.1029/2021JD036411>, 2022.
- Kakavas, S., Pandis, S. N., and Nenes, A.: ISORROPIA-Lite: A comprehensive atmospheric aerosol thermodynamics module for Earth System models, *Tellus B: Chem. Phys. Met.*, 74(1), 1–23, <https://doi.org/10.16993/tellusb.33>, 2022.

- Kato, S., Rose, F. G., Rutan, D. A., Thorsen, T. J., Loeb, N. G., Doelling, D. R., Huang, X., Smith, W. L., Su, W., and Ham, S.-H.: Surface Irradiances of Edition 4.0 Clouds and the Earth's Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) Data Product, *J. Clim.*, 31, 4501 – 4527, <https://doi.org/10.1175/JCLI-D-17-0523.1>, 2018.
- Kok, J. F.: A scaling theory for the size distribution of emitted dust aerosols suggests climate models underestimate the size of the global dust cycle, *Proceedings of the Nat. Academy of Sci.*, 108, 1016–1021, <https://doi.org/10.1073/pnas.1014798108>, 2011.
- Krol, M., Houweling, S., Bregman, B., van den Broek, M., Segers, A., van Velthoven, P., Peters, W., Dentener, F., and Bergamaschi, P.: The two-way nested global chemistry-transport zoom model TM5: algorithm and applications, *Atmos. Chem. Phys.*, 5, 417–432, <https://doi.org/10.5194/acp-5-417-2005>, 2005.
- Loeb, N. G., Doelling, D. R., Wang, H., Su, W., Nguyen, C., Corbett, J. G., Liang, L., Mitrescu, C., Rose, F. G., and Kato, S.: Clouds and the Earth's Radiant Energy System (CERES) Energy Balanced and Filled (EBAF) Top-of-Atmosphere (TOA) Edition-4.0 Data Product, *J. Clim.*, 31, 895 – 918, <https://doi.org/10.1175/JCLI-D-17-0208.1>, 2018.
- Lu, Z., Liu, X., Zhang, Z., Zhao, C., Meyer, K., Rajapakse, C., Wu, C., Yang, Z., and Penner, J. E.: Biomass smoke from southern Africa can significantly enhance the brightness of stratocumulus over the southeastern Atlantic Ocean, *Pro. Nat. Academy Sci.*, 115, 2924–2929, <https://doi.org/10.1073/pnas.1713703115>, 2018.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohengger, C., Ilyina, T., Jahns, T., Jimenez-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J.-S., Tian, F., Voigt, A., Vrese, P., Wieners, K.-H., Wilkenskeld, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and its response to increasing CO₂, *J. Adv. Mod. Earth Sys.*, 11, 998–1038, <https://doi.org/https://doi.org/10.1029/2018MS001400>, 2019.
- Metzger, S., Dentener, F., Pandis, S., and Lelieveld, J.: Gas/aerosol partitioning: 1. A computationally efficient model, *J. Geophys. Res.: Atmos.*, 107, ACH 16–1–ACH 16–24, <https://doi.org/10.1029/2001JD001102>, 2002.
- Meyers, M. P., DeMott, P. J., and Cotton, W. R.: New primary ice-nucleation parameterizations in an explicit cloud model, *J. Applied Met. Clim.*, 31, 708 – 721, [https://doi.org/10.1175/1520-0450\(1992\)031<0708:NPINPI>2.0.CO;2](https://doi.org/10.1175/1520-0450(1992)031<0708:NPINPI>2.0.CO;2), 1992.
- Moore, K. A., Hill, T. C. J., McCluskey, C. S., Twohy, C. H., Rainwater, B., Toohey, D. W., Sanchez, K. J., Kreidenweis, S. M., and DeMott, P. J.: Characterizing Ice Nucleating Particles Over the Southern Ocean Using Simultaneous Aircraft and Ship Observations, *J. Geophys. Res.: Atmos.*, 129, e2023JD039 543, <https://doi.org/https://doi.org/10.1029/2023JD039543>, 2024.
- Morales-Betancourt, R. and Nenes, A.: Droplet activation parameterization: the population-splitting concept revisited, *Geosci. Mod. Dev.*, 7, 2345–2357, <https://doi.org/10.5194/gmd-7-2345-2014>, 2014.
- Myriokefalitakis, S., Vignati, E., Tsigaridis, K., Papadimas, C., Sciare, J., Mihalopoulos, N., Facchini, M. C., Rinaldi, M., Dentener, F. J., Ceburnis, D., Hatzianastasiou, N., O'Dowd, C. D., van Weele, M., and Kanakidou, M.: Global modeling of the oceanic source of organic aerosols., *Adv. Met.*, 939171, <https://doi.org/10.1155/2010/939171>, 2010.
- NASA/LARC/SD/ASDC: CALIPSO Lidar Level 3 Global Energy and Water Cycle Experiment (GEWEX) Cloud, Standard V1-00, https://doi.org/10.5067/CALIOP/CALIPSO/LID_L3_GEWEX_Cloud-Standard-V1-00, 2019.

- Neubauer, D., Ferrachat, S., Siegenthaler-Le Drian, C., Stier, P., Partridge, D. G., Tegen, I., Bey, I., Stanelle, T., Kokkola, H., and Lohmann, U.: The global aerosol–climate model ECHAM6.3–HAM2.3 – Part 2: Cloud evaluation, aerosol radiative forcing, and climate sensitivity, *Geosci. Mod. Dev.*, 12, 3609–3639, <https://doi.org/10.5194/gmd-12-3609-2019>, 2019.
- Nickovic, S., Vukovic, A., Vujadinovic, M., Djurdjevic, V., and Pejanovic, G.: Technical Note: High-resolution mineralogical database of dust-productive soils for atmospheric dust modeling, *Atmos. Chem. Phys.*, 12, 845–855, <https://doi.org/10.5194/acp-12-845-2012>, 2012.
- O’Dowd, C. D., Langmann, B., Varghese, S., Scannell, C., Ceburnis, D., and Facchini, M. C.: A combined organic-inorganic sea-spray source function, *Geophys. Res. Lett.*, 35, <https://doi.org/10.1029/2007GL030331>, 2008.
- Perlwitz, J. P., Pérez García-Pando, C., and Miller, R. L.: Predicting the mineral composition of dust aerosols – Part 1: Representing key processes, *Atmos. Chem. Phys.*, 15, 11 593–11 627, <https://doi.org/10.5194/acp-15-11593-2015>, 2015a.
- 685 Perlwitz, J. P., Pérez García-Pando, C., and Miller, R. L.: Predicting the mineral composition of dust aerosols – Part 2: Model evaluation and identification of key processes with observations, *Atmos. Chem. Phys.*, 15, 11 629–11 652, <https://doi.org/10.5194/acp-15-11629-2015>, 2015b.
- Petters, M. D., Carrico, C. M., Kreidenweis, S. M., Prenni, A. J., DeMott, P. J., Collett Jr., J. L., and Moosmüller, H.: Cloud condensation nucleation activity of biomass burning aerosol, *J. Geophys. Res.: Atmos.*, 114, <https://doi.org/https://doi.org/10.1029/2009JD012353>, 2009.
- 690 Phillips, V. T. J., Yano, J.-I., Formenton, M., Iltoviz, E., Kanawade, V., Kudzotsa, I., Sun, J., Bansemer, A., Detwiler, A. G., Khain, A., and Tessendorf, S. A.: Ice Multiplication by Breakup in Ice–Ice Collisions. Part II: Numerical Simulations, *J. Atmos. Sci.*, 74, 2789 – 2811, <https://doi.org/10.1175/JAS-D-16-0223.1>, 2017.
- Pincus, R., Platnick, S., Ackerman, S. A., Hemler, R. S., and Hofmann, R. J. P.: Reconciling simulated and observed views of clouds: MODIS, ISCCP, and the limits of instrument simulators, *J. Climate*, 25, 4699–4720, <http://www.jstor.org/stable/26192027>, 2012.
- 695 Quaas, J., Ming, Y., Menon, S., Takemura, T., Wang, M., Penner, J. E., Gettelman, A., Lohmann, U., Bellouin, N., Boucher, O., Sayer, A. M., Thomas, G. E., McComiskey, A., Feingold, G., Hoose, C., Kristjánsson, J. E., Liu, X., Balkanski, Y., Donner, L. J., Ginoux, P. A., Stier, P., Grandey, B., Feichter, J., Sednev, I., Bauer, S. E., Koch, D., Grainger, R. G., Kirkevåg, A., Iversen, T., Seland, , Easter, R., Ghan, S. J., Rasch, P. J., Morrison, H., Lamarque, J.-F., Iacono, M. J., Kinne, S., and Schulz, M.: Aerosol indirect effects –general circulation model intercomparison and evaluation with satellite data, *Atmos. Chem. Phys.*, 9, 8697–8717, <https://doi.org/10.5194/acp-9-8697-2009>, 2009.
- 700 Quaas, J., Arola, A., Cairns, B., Christensen, M., Deneke, H., Ekman, A. M. L., Feingold, G., Fridlind, A., Gryspeerdt, E., Hasekamp, O., Li, Z., Lipponen, A., Ma, P.-L., Mülmenstädt, J., Nenes, A., Penner, J. E., Rosenfeld, D., Schrödner, R., Sinclair, K., Sourdeval, O., Stier, P., Tesche, M., van Dierenhoven, B., and Wendisch, M.: Constraining the Twomey effect from satellite observations: issues and perspectives, *Atmos. Chem. Phys.*, 20, 15 079–15 099, <https://doi.org/10.5194/acp-20-15079-2020>, 2020.
- 705 Ramanathan, V., Crutzen, P., Kiehl, J., and Rosenfeld, D.: Aerosols, climate, and the hydrological cycle, *Science*, 294, 2119–2124, <https://doi.org/10.1126/science.1064034>, 2001.
- Redemann, J., Wood, R., Zuidema, P., Doherty, S. J., Luna, B., LeBlanc, S. E., Diamond, M. S., Shinozuka, Y., Chang, I. Y., Ueyama, R., Pfister, L., Ryoo, J.-M., Dobracki, A. N., da Silva, A. M., Longo, K. M., Kacenenbogen, M. S., Flynn, C. J., Pistone, K., Knox, N. M., Piketh, S. J., Haywood, J. M., Formenti, P., Mallet, M., Stier, P., Ackerman, A. S., Bauer, S. E., Fridlind, A. M., Carmichael, G. R., Saide, P. E., Ferrada, G. A., Howell, S. G., Freitag, S., Cairns, B., Holben, B. N., Knobelspiesse, K. D., Tanelli, S., L’Ecuyer, T. S., Dzambo, A. M., Sy, O. O., McFarquhar, G. M., Poellot, M. R., Gupta, S., O’Brien, J. R., Nenes, A., Kacarab, M., Wong, J. P. S., Small-Griswold, J. D., Thornhill, K. L., Noone, D., Podolske, J. R., Schmidt, K. S., Pilewskie, P., Chen, H., Cochrane, S. P., Sedlacek, A. J., Lang, T. J., Stith, E., Segal-Rozenhaimer, M., Ferrare, R. A., Burton, S. P., Hostetler, C. A., Diner, D. J., Seidel, F. C., Platnick, S. E., Myers, J. S.,

- Meyer, K. G., Spangenberg, D. A., Maring, H., and Gao, L.: An overview of the ORACLES (ObseRvations of Aerosols above CLouds and their intERactionS) project: aerosol–cloud–radiation interactions in the southeast Atlantic basin, *Atmos. Chem. Phys.*, 21, 1507–1563, <https://doi.org/10.5194/acp-21-1507-2021>, 2021.
- Romakkaniemi, S., Arola, A., Kokkola, H., Birmili, W., Tuch, T., Kerminen, V.-M., Räisänen, P., Smith, J. N., Korhonen, H., and Laaksonen, A.: Effect of aerosol size distribution changes on AOD, CCN and cloud droplet concentration: Case studies from Erfurt and Melpitz, Germany, *J. Geophys. Res.: Atmos.*, 117, D0702, <https://doi.org/10.1029/2011JD017091>, 2012.
- Royer, H. M., Pöhlker, M. L., Krüger, O., Blades, E., Sealy, P., Lata, N. N., Cheng, Z., China, S., Ault, A. P., Quinn, P. K., Zuidema, P., Pöhlker, C., Pöschl, U., Andreae, M., and Gaston, C. J.: African smoke particles act as cloud condensation nuclei in the wintertime tropical North Atlantic boundary layer over Barbados, *Atmos. Chem. Phys.*, 23, 981–998, <https://doi.org/10.5194/acp-23-981-2023>, 2023.
- Scanza, R. A., Mahowald, N., Ghan, S., Zender, C. S., Kok, J. F., Liu, X., Zhang, Y., and Albani, S.: Modeling dust as component minerals in the Community Atmosphere Model: development of framework and impact on radiative forcing, *Atmos. Chem. Phys.*, 15, 537–561, <https://doi.org/10.5194/acp-15-537-2015>, 2015.
- Seland, Ø., Bentsen, M., Olivié, D., Toniazzo, T., Gjermundsen, A., Graff, L. S., Debernard, J. B., Gupta, A. K., He, Y.-C., Kirkevåg, A., Schwinger, J., Tjiputra, J., Aas, K. S., Bethke, I., Fan, Y., Griesfeller, J., Grini, A., Guo, C., Ilicak, M., Karset, I. H. H., Landgren, O., Liakka, J., Moseid, K. O., Nummelin, A., Spensberger, C., Tang, H., Zhang, Z., Heinze, C., Iversen, T., and Schulz, M.: Overview of the Norwegian Earth System Model (NorESM2) and key climate response of CMIP6 DECK, historical, and scenario simulations, *Geosci. Mod. Dev.*, 13, 6165–6200, <https://doi.org/10.5194/gmd-13-6165-2020>, 2020.
- Shipway, B. J.: Revisiting Twomey’s approximation for peak supersaturation, *Atmos. Chem. Phys.*, 15, 3803–3814, <https://doi.org/10.5194/acp-15-3803-2015>, 2015.
- Sokolik, I. N. and Toon, O. B.: Incorporation of mineralogical composition into models of the radiative properties of mineral aerosol from UV to IR wavelengths, *J. Geophys. Res.: Atmos.*, 104, 9423–9444, <https://doi.org/10.1029/1998JD200048>, 1999.
- Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K., Tignor, M., and Miller, H. L.: IPCC, 2007: Summary for Policy-makers, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.*, Cambridge University Press, Cambridge, UK and New York, NY, USA., 2007.
- Sotiropoulou, G., Vignon, E., Young, G., Morrison, H., O’Shea, S. J., Lachlan-Cope, T., Berne, A., and Nenes, A.: Secondary ice production in summer clouds over the Antarctic coast: an underappreciated process in atmospheric models, *Atmos. Chem. Phys.*, 21, 755–771, <https://doi.org/10.5194/acp-21-755-2021>, 2021.
- Tang, J., Wang, P., Mickley, L. J., Xia, X., Liao, H., Yue, X., Sun, L., and Xia, J.: Positive relationship between liquid cloud droplet effective radius and aerosol optical depth over Eastern China from satellite data, *Atmos. Env.*, 84, 244–253, <https://doi.org/10.1016/j.atmosenv.2013.08.024>, 2014.
- Tegen, I., Harrison, S. P., Kohfeld, K., Prentice, I. C., Coe, M., and Heimann, M.: Impact of vegetation and preferential source areas on global dust aerosol: Results from a model study, *J. Geophys. Res.: Atmos.*, 107, AAC 14–1–AAC 14–27, <https://doi.org/10.1029/2001JD000963>, 2002.
- Tegen, I., Werner, M., Harrison, S. P., and Kohfeld, K. E.: Relative importance of climate and land use in determining present and future global soil dust emission, *Geophys. Res. Lett.*, 31, <https://doi.org/10.1029/2003GL019216>, 2004.
- Twomey, S. A.: Pollution and the planetary albedo, *Atmos. Env.*, 8, 1251–1256, [https://doi.org/10.1016/0004-6981\(74\)90004-3](https://doi.org/10.1016/0004-6981(74)90004-3), 1974.
- Twomey, S. A.: The influence of pollution on the shortwave albedo of clouds, *J. Atmos. Sci.*, 34, 1149–1152, [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2), 1977.

- Ullrich, R., Hoose, C., Möhler, O., Niemand, M., Wagner, R., Höhler, K., Hiranuma, N., Saathoff, H., and Leisner, T.: A new ice nucleation active site parameterization for desert dust and soot, *J. Atmos. Sci.*, 74, 699 – 717, <https://doi.org/10.1175/JAS-D-16-0074.1>, 2017.
- 755 van Noije, T., Bergman, T., Le Sager, P., O'Donnell, D., Makkonen, R., Gonçalves-Ageitos, M., Döschner, R., Fladrich, U., von Hardenberg, J., Keskinen, J.-P., Korhonen, H., Laakso, A., Myriokefalitakis, S., Ollinaho, P., Pérez García-Pando, C., Reerink, T., Schrödner, R., Wyser, K., and Yang, S.: EC-Earth3-AerChem: a global climate model with interactive aerosols and atmospheric chemistry participating in CMIP6, *Geosci. Mod. Dev.*, 14, 5637–5668, <https://doi.org/10.5194/gmd-14-5637-2021>, 2021.
- 760 van Noije, T. P. C., Le Sager, P., Segers, A. J., van Velthoven, P. F. J., Krol, M. C., Hazeleger, W., Williams, A. G., and Chambers, S. D.: Simulation of tropospheric chemistry and aerosols with the climate model EC-Earth, *Geosci. Mod. Dev.*, 7, 2435–2475, <https://doi.org/10.5194/gmd-7-2435-2014>, 2014.
- Vergara-Temprado, J., Miltenberger, A. K., Furtado, K., Grosvenor, D. P., Shipway, B. J., Hill, A. A., Wilkinson, J. M., Field, P. R., Murray, B. J., and Carslaw, K. S.: Strong control of Southern Ocean cloud reflectivity by ice-nucleating particles, *Pro. Nat. Academy Sci.*, 115, 2687–2692, <https://doi.org/10.1073/pnas.1721627115>, 2018.
- Vignati, E., Wilson, J., and Stier, P.: M7: An efficient size-resolved aerosol microphysics module for large-scale aerosol transport models, *J. Geophys. Res.: Atmos.*, 109, <https://doi.org/10.1029/2003JD004485>, 2004.
- 765 Vignati, E., Facchini, M., Rinaldi, M., Scannell, C., Ceburnis, D., Sciare, J., Kanakidou, M., Myriokefalitakis, S., Dentener, F., and O'Dowd, C.: Global scale emission and distribution of sea-spray aerosol: Sea-salt and organic enrichment, *Atmos. Env.*, 44, 670–677, <https://doi.org/10.1016/j.atmosenv.2009.11.013>, 2010.
- 770 Wex, H., Huang, L., Zhang, W., Hung, H., Traversi, R., Becagli, S., Sheesley, R. J., Moffett, C. E., Barrett, T. E., Bossi, R., Skov, H., Hünerbein, A., Lubitz, J., Löffler, M., Linke, O., Hartmann, M., Herenz, P., and Stratmann, F.: Annual variability of ice-nucleating particle concentrations at different Arctic locations, *Atmos. Chem. Phys.*, 19, 5293–5311, <https://doi.org/10.5194/acp-19-5293-2019>, 2019.
- Wieder, J., Ihn, N., Mignani, C., Haarig, M., Bühl, J., Seifert, P., Engemann, R., Ramelli, F., Kanji, Z. A., Lohmann, U., and Henneberger, J.: Retrieving ice-nucleating particle concentration and ice multiplication factors using active remote sensing validated by in situ observations, *Atmos. Chem. Phys.*, 22, 9767–9797, <https://doi.org/10.5194/acp-22-9767-2022>, 2022.
- 775 Wilson, T. W., Ladino, L. A., Alpert, P. A., Breckels, M. N., Brooks, I. M., Browse, J., Burrows, S. M., Carslaw, K. S., Huffman, J. A., Judd, C., Kilhau, W. P., Mason, R. H., McFiggans, G., Miller, L. A., Nájera, J. J., Polishchuk, E., Rae, S., Schiller, C. L., Si, M., Temprado, J. V., Whale, T. F., Wong, J. P., Wurl, O., Yakobi-Hancock, J. D., Abbatt, J. P., Aller, J. Y., Bertram, A. K., Knopf, D. A., and Murray, B. J.: A marine biogenic source of atmospheric ice-nucleating particles., *Nature*, 525(7568):234-8, <https://doi.org/10.1038/nature14986>. PMID: 26354482, 2015.
- 780 Zhao, X., Liu, X., Burrows, S., DeMott, P. J., Diao, M., McFarquhar, G. M., Patade, S., Phillips, V., Roberts, G. C., Sanchez, K. J., Shi, Y., and Zhang, M.: Important Ice Processes Are Missed by the Community Earth System Model in Southern Ocean Mixed-Phase Clouds: Bridging SOCRATES Observations to Model Developments, *J. Geophys. Res.: Atmos.*, 128, e2022JD037513, <https://doi.org/10.1029/2022JD037513>, 2023.