

# Unprecedented Twenty-First Century Glacier Loss on Mt. Hood, Oregon, U.S.A.

Nicolas Bakken-French<sup>1</sup>, Stephen J. Boyer<sup>1</sup>, [B. Clay Southworth<sup>1</sup>](#), Megan Thayne<sup>1</sup>, Dylan H. Rood<sup>2\*</sup>, and Anders E. Carlson<sup>1</sup>

<sup>1</sup>Oregon Glaciers Institute, Corvallis, OR 97330, U.S.A.

<sup>2</sup>Department of Earth Science and Engineering, Imperial College London, London SW7 2AZ, U.K.

Correspondence to: Dylan H. Rood (d.rood@imperial.ac.uk)

**Abstract.** As part of the southern Cascades, Mt. Hood is the tallest and most glaciated peak in Oregon, U.S.A. Despite alpine glaciers being one of the clearest indicators of human-caused climate change, the 21st century behavior of glaciers on Mt. Hood has not been directly documented [at the ground level](#). Here we directly measure changes in Mt. Hood's glacier extents from 2003 to 2023 and find dramatic retreat of all glaciers, with one glacier disappearing, [and another four retreating](#) towards this status. [By 2023, Mt. Hood glaciers lost ~18% of their 2015/2016 area and ~44% of their 1981 area.](#) The seven largest glaciers on the volcano lost [~40% of their area between 2000 and 2023.](#) Comparison to historic records of glacier area back to 1907 shows that this 21st-century retreat is unprecedented [relative to the previous century](#) and has outpaced modeled glacier changes [of 21st century retreat](#). The rate of [area loss](#) over the last 23 years ( $1.7\text{--}1.8\% \text{ year}^{-1}$ ) is [about triple](#) the fastest rate documented in the last century from 1907 to 1946 ( $0.54\text{--}0.58\% \text{ year}^{-1}$ ) and [about six times faster than the 20th century average](#) ( $0.29\text{--}0.33\% \text{ year}^{-1}$ ). [It has also equaled to exceeded by up to 67% the highest simulated rates of glacier recession under a high greenhouse gas emissions scenario.](#) We demonstrate that this century-scale retreat strongly correlates with regional 30-year [mean climate warming of 1.1–1.2°C](#) and [15-year-mean climate warming of 1.3–1.5°C](#) since the early 1900s, but not with regional changes in precipitation. We conclude that Mt. Hood's glaciers are retreating in response to a warming climate and that this recession has accelerated in the 21st century.

## 1 Introduction

The global [loss of mass and retreat](#) of glaciers in the 21st century is accelerating (Zemp et al., 2015; Hugonnet et al., 2021) with century-scale glacier recession attributed to regional climate change in response to anthropogenic greenhouse gas emissions (Marzeion et al., 2014; Roe et al., 2017, 2021). The 2021–2022 hydrological year was the 35th consecutive year of negative annual [mass](#) balance for 50 reference glaciers on six continents (Pelto, 2023; [World Glacier Monitoring Service, 2022](#)). If global average temperature increases by  $1.5^\circ\text{C}$  relative to the pre-industrial period, the limit set forth in the Paris Agreement, roughly half of the world's individual glaciers are projected to disappear [with ~26% loss in global glacier mass](#) (Rounce et al., 2023). Indeed, in the western United States, this deglaciation is already coming to fruition ([Fig. 1A](#)). The Trinity Alps in the State of California ([Fig. 1A](#)) [have lost ~97% of their late 1800s glaciated area, with only ~0.017 km<sup>2</sup> of one glacier still showing evidence of ice flow as of 2015](#) (Garwood et al., 2020). The Olympic Mountains of Washington

Deleted: w

Deleted: ,

Deleted: with

Deleted: two

Deleted: nearing this status, and a third

Deleted: had

Deleted: ~2.8 km<sup>2</sup>, or

Deleted: in the 21st century

Deleted: with respect

Deleted: retreat

Deleted: more

Deleted: than double

Deleted: future

Deleted: average

Deleted: ~

Deleted: , with attendant consequences for water resources

Deleted:

Deleted: 3

Formatted: Superscript

Deleted: became a glacier-free mountain range in the last decade

State (Fig. 1A) lost about half of their late 1800s glacier area by 2015 with 35 glaciers disappearing since 1980 (Fountain et al., 2022). Eleven of the 37 named glaciers in Glacier National Park in the State of Montana (Fig. 1A) ceased to be classified as glaciers as of 2015 with glaciated area in the park declining by ~68% from the 1800s to 2015 (Fagre et al., 2017). Washington State's Mt. Rainier (Fig. 1A), the most glacierized peak in the contiguous United States, has seen its glaciated area decline by ~42% between 1896 and 2021 and had one glacier recently disappear with two more nearing stagnation (Beason et al., 2023). Nearby in the Washington North Cascades (Fig. 1A), 33 glaciers were listed as extinct in the GLIMS Glacier Database as of 2018 (GLIMS and NSIDC, 2018).

Deleted: have lost

Deleted: glaciated




Figure 1: Location map of Oregon (A) and Mt. Hood (B). A. Western United States with mountain ranges mentioned denoted (WA = Washington, MT = Montana, OR = Oregon, CA = California, ID = Idaho, NV = Nevada, UT = Utah, AZ = Arizona); inset shows location with black outline. B. Glacier extents on Mt. Hood with photo locations in Fig. 3 noted. 1907/1901 and 2004 ice extents from Jackson & Fountain (2007); 2015/2016 ice extents from Fountain et al. (2023). 2023 extents from this study. Inset shows figure location (black outline) within Hood River County (area for reanalysis data in Fig. 6). Credits: Esri, U.S. Census Bureau, Earthstar Geographics, Maxar.

Deleted: Location map of United States, Oregon, Hood River County and Mt. Hood. 1907/1901 and 2004 ice extents from


Deleted: Contour interval is 50 m.

70 Despite these ominous predictions and dire observations for glaciers, data on the 21st century behavior of glaciers in the Cascade Range of Oregon are limited. Glaciers in Oregon were first noted in 1870 on Mt. Hood, the tallest and most glacierized peak in Oregon (Fig. 1A, B), only days after the first glaciers in the contiguous United States were recorded on Mt. Shasta in California (O'Connor, 2013). Mt. Hood is an active volcano that reached a height close to its current elevation of 3,425 meters above sea level by 30,000 years ago (U.S.G.S., 2023, <https://www.usgs.gov/volcanoes/mount-hood/science/eruption-history-mount-hood-oregon>). The last major eruption was ~1,500 years ago that created pyroclastic flows and lahars on the southwest side of the volcano upon which Zigzag and Palmer (now gone) glaciers rested (Fig. 1B). Its last eruption occurred in the late 1700s, producing a lava dome south of the summit around which Coalman Glacier flows (Fig. 1B). In the late 1800s, fumaroles formed near this lava dome (Lillquist and Walker, 2006). Mt. Hood has high amounts of rock fall and variations in geothermal heat that could affect ablation, ice flow, and ice extent (e.g., Lundstrom et al., 1993; Lillquist and Walker, 2006; Jackson and Fountain, 2007; Howcutt et al., 2023).

Changes in Mt. Hood's glaciers were monitored by the Research Committee of the Mazama Mountaineering Club from the 1930s to mid-1980s (e.g., Phillips, 1935, 1938; Mason, 1954; Handwith Jr., 1959; Dodge, 1971, 1987). In 1981, the U.S. Geological Survey measured the thickness and area of Mt. Hood's glaciers, estimating their volume (Driedger and Kennard, 1986a, 1986b). However, the extent of Oregon Cascade glaciers on current U.S. Geological Survey and U.S. Forest Service maps are still based on air photos from the 1950s (Fountain et al., 2017). The glaciated area of Mt. Hood was last measured with field observations in 2004 (Jackson and Fountain, 2007) whereas glacier lengths with field observations were last recorded in 2000 (Lillquist and Walker, 2006). Fountain et al. (2023) documented snow and ice areas in the western United States using 2015 and 2016 imagery for Mt. Hood's glaciers but did not test mapped accuracy with field observations. The timing of this imagery also missed what multi-year and longer impacts the 2015 record-low snowpack (Mote et al., 2016) and the June 2021 Pacific Northwest heatwave (Philip et al., 2022; Thompson et al., 2022) would have on Mt. Hood's glaciers (e.g., Pelto, 2018; Pelto et al., 2022).

Regarding climate change impacting Oregon's glaciers, Menounos et al. (2019) documented minimal elevation change for Mt. Hood's glaciers from 2000 to 2018, interpreting from this an approximately neutral mass balance for the first 19 years of the 21st century. Near zero mass balance for 2000-2018 is at odds with the mass changes for global reference glaciers that include four glaciers to the north of Mt. Hood in the state of Washington; these four all had cumulative negative mass balances for the same time period (Pelto, 2023). For the 20th century, Lillquist and Walker (2006) did not find a relationship between glacier-length change on Mt. Hood and temperature or precipitation change over the 20th century. Conversely, climate-glacier modeling simulates the emergence of glacier retreat from natural variability in western North America since the 1960s (Marzeion et al., 2014). As such, recent impacts of human-caused climate change on Mt. Hood's glaciers have been unclear and largely unexplored from direct field observations.

Glacier change on Mt. Hood is of importance to downstream ecosystems and economies (see Robel et al., 2024), such as



- Deleted: ... [1]
- Deleted: recent
- Deleted: (Fig. 1). Glaciers in Oregon were first noted in 1870 ... [2]
- Moved (insertion) [1]
- Deleted: Mt. Hood is an active volcano, with
- Deleted: and topographic relief that influence accumulation,
- Deleted: (e.g., Fig. 2)
- Deleted: glaciers, but
- Deleted: s
- Moved down [2]: Lillquist and Walker (2006) did not find a relationship between glacier-length change on Mt. Hood and temperature or precipitation change over the 20th century. Similarly,
- Deleted: In terms of
- Deleted: A n
- Deleted:
- Moved (insertion) [2]
- Deleted: 3
- Deleted: Similarly, C...imate-glacier modeling also ... [3]
- Deleted: In addition to deciphering the regional impacts of human-caused climate change, gG

orchards and farms in the Hood River Valley (Nolin et al., 2010; U.S. Bureau of Reclamation, 2015; Frans et al., 2016, 2018), unique glacier-fed spring water microbiomes (Miller et al., 2021), and salmon (Pitman et al., 2020; Thieman et al., 2021). The Hood River Valley has 12,000 acres producing seven varieties of pears, 1500 acres producing five varieties of apples, 1500 acres devoted to cherries, and 100 acres of grapes for an expanding wine industry (Columbia Gorge Fruit Growers, 2024, <http://www.cgfg.org/information/crops-information>). Glaciers contribute ~40-70% of the stream flow in the late summer to the upper Hood River where flow is diverted for irrigation and local hydropower (Nolin et al., 2010). As such, future water resource management plans for the Hood River basin included projections of future glacier change on Mt. Hood (U.S. Bureau of Reclamation, 2015; Thieman et al., 2021). White River and Newton-Clark glaciers also contribute late-summer meltwater to a robust orchard industry (Columbia Gorge Fruit Growers, 2024, <http://www.cgfg.org/information/crops-information>). Furthermore, Mt. Hood is also one of the most climbed mountaineering objectives in the world (~10,000 attempts per year), which is becoming more dangerous during a diminishing climbing season with greater rockfall as the volcano has changed in recent years (O'Neil, 2023).

Here we document glacier change on Mt. Hood over the last 20 years. We conducted new repeat photography and field observations of glacier extent in 2003 and 2023 to provide detailed ground-based evidence for glacier change during a period where satellite-based altimetry previously found no significant change in these glaciers. We also utilized a series of field campaigns with satellite imagery to determine the magnitude of glacier area and length change up to mid-September 2023. We place these 21st-century glacier changes in the context of 20th-century glacier changes and test the relationship between ~120 years of glacier-length variability and regional climate change.

## 2 Data and Methods

### 2.1 2003 Frontal GPS Positions and Repeat Photography

In 2003 (28 August to 4 October), the lowest observed actively flowing terminus positions and stagnant-ice positions were measured (latitude-longitude-altitude) by global positioning system (GPS) for every glacier on Mt. Hood (Fig. S1). These were discrete observations, which served as points to assess future minimum changes in terminus elevation. Repeat measurements and comparison to known elevation markers determined an elevation uncertainty of  $\pm 6$  m. Termini were photographed with locations noted (Fig. S1) for future repeat photography (i.e., 2023). All glaciers were described in detail in the field (Supplement). The goal of this initial 2003 field effort was to provide baseline observations and archival photographic evidence from the early 21st century against which future glacier change could be assessed; we subsequently conducted further observations for this assessment 20 years later in 2023.

From 15 to 22 September 2023, repeat photographs were taken at the same locations as the 2003 photographs (Fig. S1) and new field observations were made for all of the glaciers (Supplement). From this repeat mapping (see below), we were able

Deleted: . O

Deleted: depend on Mt. Hood's glacier meltwater, particularly in late summer when glacier meltwater can supply up to ~70% of streamflow (Nolin et al., 2010; U.S. Bureau of Reclamation, 2015;

Deleted: ). U

Deleted: recently described on Mt. Hood depend on glacier-fed spring water (Miller et al., 2021)

Deleted: . Mt. Hood glacier meltwater also maintains cold instream temperatures and late summer streamflow upon which salmon rely

Deleted: To note t

Deleted: For example, t include that

Deleted: for

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Deleted: In addition

Deleted: 's cryosphere

Deleted: field mapping

Deleted: s

Deleted: was used to find

Deleted: , which combined

Deleted: with

Deleted: satellite imagery

Deleted:

Deleted: 2

Deleted: ;

Deleted: repeat

Moved up [1]: Mt. Hood is an active volcano, with high amounts of rock fall and variations in geothermal heat and topographic relief that influence accumulation, ablation, ice flow, and ice extent (e.g., Lundstrom et al., 1993; Lillquist and Walker, 2006; Jackson and Fountain, 2007; Howcutt et al., 2023). For this reason, a

Deleted: Figure 2 shows the location of the repeat photographs.

Deleted: For this reason, a

Deleted: like

Deleted: ten

Deleted: did

Deleted: comparable

to determine the minimum amount the terminus rose in elevation over the last 20 years. We selected photographs to repeat from an archive of photographs taken in 2003. Using the image and field notes regarding the location of the images from 2003, we located the precise site of each selected 2003 photograph. The same frame was maintained using relatively unchanged features such as buttresses, cliffs, and moraines as points of reference.

## 2.2 2023 Glacier Inventory, Mapping and Uncertainty

In contrast to the point measurements in 2003, full glacier extents were mapped in 2023, using an iterative approach over the years 2020 to 2023. Note that we mapped only the extent of actively flowing glacier ice, both debris-free and debris-covered, and excluded stagnated non-flowing debris-free and debris-covered ice (see below for criteria for differentiation and delineation of flowing from stagnant glacier ice). We focused on actively flowing glacier ice, rather than include stagnant ice, to be consistent with prior glacier-extent mapping on Mt. Hood (e.g., Lillquist and Walker, 2006; Jackson and Fountain, 2007; Fountain et al., 2023) and in other mountain ranges of the western United States (e.g., Fagre et al., 2017; Garwood et al., 2020; Fountain et al., 2022; Beason et al., 2023).

Because manual mapping is suggested for glaciers with debris cover (e.g., Paul et al., 2013), we initially manually mapped actively flowing glacier extents using composite Google Earth imagery in the application CalTopo (caltopo.com). As these images span multiple years and are not necessarily recent, we then manually refined ice margins using weekly Sentinel-2 satellite images that were taken shortly before the first snowfall from the summers of 2020, 2021, 2022, and 2023. These images were processed and made available in CalTopo one day after acquisition (see Supplement for a complete list of satellite imagery). In October 2020, September 2021, and September 2023, we iteratively took these ice margins (updated year by year) into the field for direct mapping using CalTopo on a portable GPS enabled tablet. These three years were excellent for field mapping as the length of 2020 summer and the warmth of the 2021 and 2023 summers resulted in extremely low snow cover that facilitated ice-margin identification in regions that would typically have snow cover throughout the summer.

In sequence, glacier margins were first manually mapped from Google Earth imagery, then modified by late summer 2020 Sentinel-2 imagery and finalized by in-the-field ice-margin mapping. Then, the 2020 ice-margin maps were updated with late summer 2021 Sentinel-2 imagery that were subsequently further adjusted and mapped in the field. The 2021 maps were updated with late summer 2022 Sentinel-2 imagery and again with late summer 2023 Sentinel-2 imagery, which were then adjusted and finalized in September of 2023 by in-the-field measurement. All termini in the end-of-summer ablation area were walked, resulting in ~60%, on average, of each ice margin being physically mapped in the field. Notably, this means all debris-covered ice margins were mapped in the field and their delineation was not based solely on satellite imagery. Two glaciers, Zigzag, Glisan, were field mapped entirely.

Deleted: For the 2023 extent of glaciers on Mt. Hood,

Deleted: we initially mapped ice extent using composite Google Earth imagery. As these

Deleted: just

Deleted: -

Deleted: -

Deleted: ,

260 Debris-covered termini are common on Mt. Hood glaciers, and such field mapping of the termini is critical to accurately delineate glacier margins (e.g., Lundstrom et al., 1993; Lillquist and Walker, 2006; Jackson and Fountain, 2007; Ellinger, 2010). We followed prior methods for differentiating actively flowing debris-covered ice from stagnant debris-covered ice on Mt. Hood (e.g., Lillquist and Walker, 2006): an identifiable glacier front, a lobate form of the terminus, and a visible connection of the lobate terminus to debris-free flowing ice. This methodology is consistent with the observation of a convex shape of actively flowing glacier margins by Leonard and Fountain (2003) and Leigh et al. (2019). To these criteria, we added the presence of crevasses observed in the field as another indicator of ice flow whether under debris or in bare ice (Leigh et al., 2019).

275 Manual remote-sensed ice limits compared excellently with direct field observations, so any uncertainty in these field-based glacier extents was difficult to quantify as the values were small. Using only remote sensing, Fountain et al. (2023) had their lowest ice-extent uncertainty for a Mt. Hood glacier at  $\pm 1\%$ , while Jackson and Fountain (2007) reported an uncertainty for their 2000 Mt. Hood glacier areas of  $\pm 0\%$ . Our uncertainty is most likely lower than  $\pm 1\%$ , but to be conservative we assumed a  $\pm 7\%$  uncertainty in the glacier area for 2023. We based this  $\pm 7\%$  uncertainty on the average area uncertainty for the 2004 extents of Mt. Hood glaciers, determined by Jackson and Fountain (2007), which was  $\pm 7\%$ . Our conservative  $\pm 7\%$  is slightly greater than the uncertainties determined from testing the accuracy of remote sensed glacier area from the European Alps and Alaska of 2.6-5.7% (Paul et al., 2013). This means any conclusions we made from our mapped glacier area were likely not dependent on our assumed glacier area uncertainty.

### 2.3 Glacier Area Change and Uncertainty

280 To update glacier change on Mt. Hood for the 21st century, we compare our 2023 glacier extents to the most recent inventories conducted for Mt. Hood. Fountain et al. (2023) produced Mt. Hood glacier areas from remote sensing imagery from either 2015 or 2016; this allowed an assessment of glacier-area change over the last seven to eight years. We also compared our 2023 glacier areas to the 1981 glacier areas for Mt. Hood determined by Driedger & Kennard (1986b) that were based on field mapping and aerial photographs; this afforded a 42-year perspective on glacier change.

285 We assessed rates of glacier change by comparing our 2023-measured area with the glacier area records of Jackson and Fountain (2007) for the time period 1907 (or 1901 in the case of Eliot Glacier) to 2004 for seven glaciers on Mt. Hood: White River, Newton-Clark, Eliot, Coe, Ladd, Sandy, and Reid. Because of the different temporal resolutions of each glacier-area record, we focus on area change between observational years common to all seven glaciers: 1907, 1946, 1972, 2000, and 2023. The one exception is Newton-Clark Glacier, which lacks an aerial extent for 1946. We extrapolated between the measured extent in 1935 and 1956 when the glacier lost  $\sim 0.04 \text{ km}^2$  over these 21 years. To aid comparison between glaciers of different sizes on Mt. Hood and to other glaciers in the Pacific Northwest, we calculated relative changes in glacier area determined as a percent of its prior area. In calculating the rate of change, we propagated through the uncertainty

**Deleted:** These outlines were refined over four years for the summers of 2020, 2021, 2022, and 2023, which allowed for precise determination of margins. In October of 2020, September of 2021, and September of 2023, we mapped glacier extents in the field and checked these glacier limits for all the glaciers on Mt. Hood. Debris-

**Deleted:** from

**Deleted:** and

**Deleted:** Specifically, glacier margins were walked with GPS and debris-covered stagnant ice was separated from debris-covered active ice. In all cases M

**Deleted:** , we found excellent agreement between our remote-sensed ice limits and the field-observed ice limits. As such, any ...

**Deleted:** is

**Deleted:** are

**Deleted:** , which

**Deleted:** is the

**Deleted:**

**Deleted:** (

**Deleted:** ,

**Deleted:** ¶

**Deleted:** place the rate of glacier area loss on

**Deleted:** in the context of the 20th century

**Formatted:** Normal, Line spacing: single, Keep with next

**Deleted:** from remote sensing

**Deleted:** we

**Deleted:** compare

**Deleted:** in comparing

**Deleted:** We focus on the change from 2000 to 2023, rather than 2000 to 2004 and then 2004 to 2023, to make the duration over which the most recent period of change is measured closer to the earlier temporal spacing. In calculating the rate of change, we propagate

320 in the area from Jackson and Fountain (2007) and our 2023 areas. For the 2000 area, we conservatively applied a  $\pm 7\%$   
uncertainty of the 2004 Jackson and Fountain (2007) glacier margins. The propagation of uncertainty was through  
quadrature where the squares of the individual percent uncertainties were summed, and their square root taken.

#### 2.4 Change in Glacier Length and Climate

325 Glacier length changes are commonly used to assess glacier relationships to a changing climate (e.g., Harper, 1993; Haeberli  
and Hoelzle, 1995; Pelto and Hedlund, 2001; Oerlemans, 2005; Lillquist and Walker, 2006; Leclercq and Oerlemans, 2012;  
Leclercq et al., 2014; Pelto, 2016; Roe et al., 2016, 2021; Huston et al., 2021; Hoelzle et al., 2023). Here we compare the  
cumulative change in glacier length for five glaciers on Mt. Hood from 1901 through 2023 to records of regional climate  
change. Lillquist and Walker (2006) measured glacier length change for White River, Newton-Clark, Eliot, Coe, and Ladd  
glaciers from 1901 to 2000. To this record we add the change in length up to 2023. Specifically, we calculate the distance  
330 between the lowest observed actively flowing ice in 2023 (Fig. 1B) and the lowest actively flowing ice in 2000 of Lillquist  
and Walker (2006) along the same flow line they used back to 1901, which was down the center of the glacier. In the case of  
Newton-Clark Glacier, the length was measured in the northern, due-east-facing Newton drainage (Fig. 1B). Changes in  
glacier length were normalized to the 1901 length of the glacier to account for different glacier sizes. Lillquist and Walker  
(2006) did not provide uncertainties in their glacier lengths and reported change at 1 m accuracy. Here, we report our length  
335 changes at 10 m accuracy.

For records of regional climate change, we used U.S. National Oceanic and Atmospheric Administration (NOAA)  
temperature and precipitation reanalysis data for the region of Hood River County (Fig. 1) from 1895 to 2023 (U.S. NOAA,  
2024, <https://www.ncei.noaa.gov/access/monitoring/climate-at-a-glance/>). We chose Hood River County as this is the  
340 smallest geographic region for the reanalysis data that include these five glaciers and would still have direct weather station  
measurements within the geographic domain over the period of comparison. Snow telemetry stations on and around Mt.  
Hood were only installed in the late 1970s, with temperature sensors beginning to work consistently in the late 1980s (U.S.  
National Resources Conservation Service, 2024, [https://www.nrcs.usda.gov/conservation-basics/conservation-by-  
state/oregon/oregon-snow-survey/about-us](https://www.nrcs.usda.gov/conservation-basics/conservation-by-state/oregon/oregon-snow-survey/about-us)).

345 For temperature, we focused on changes in annual May-October and June-September means. May through October is  
generally the period of time in the Oregon Cascades from peak winter snowpack to the first snowfall of the next winter, or  
the maximum length of the ablation season. Since this can vary, we also examined June-September temperature as a shorter  
ablation season. Indeed, Pelto (2018) determined the ablation season to be May-September or June-September for glaciers in  
Washington. For precipitation, we analyzed annual November-April and October-May means. November through April in  
Oregon generally spans the time period from the first winter snow until peak spring snowpack, or the accumulation season.  
350 To account for variability in this season, we also used October through May precipitation.

Deleted: own

Deleted: . In the case of the

Deleted: apply

Deleted: ¶

In the early 1900s, the conical shape of Mt. Hood allowed some glaciers to expand laterally at lower elevations, giving them broad ablation areas relative to narrower accumulation areas (Fig. 1) (Jackson and Fountain, 2007). This could make initial glacier-area responses to a given change in climate greater than later responses to the same magnitude of climate change, as earlier ablation zones were potentially wider than later ablation zones. To account for this possible volcanic geometrical control on the absolute rate of glacier area change over the past ~120 years, we compare the change in glacier area relative to its prior area. ...

Deleted: as a metric

Deleted:

Deleted: for

Deleted: the years 1901, 1928, 1938, 1946, 1959, 1967, 1972, 1979, 1984, 1989, 1995, and 2000. Note that Newton-Clark and White River are missing the year 1928 while Ladd is missing the year 1959; these omissions impact the number of observations used to determine the significance of regressions. Lillquist and Walker (2006) also had observations of length change for 2001 for Eliot and Coe; we exclude this year as it is one year removed from 2000 and only exists for two out of the five glacier records. To this record and

Deleted: and along the same flow line

Deleted: is

Deleted: 2

Deleted: to

Deleted: We compare the cumulative glacier-length change to records of regional climate change. As...

Deleted: metrics

Deleted: records

Deleted: 3

Formatted: Font: Not Bold

Deleted: and

Deleted: temperature

Deleted: fol

Deleted: includes

Deleted: , with the latter being the

Deleted: and

Deleted: precipitation

Deleted: , with the latter being the maximum length of

395 First, we calculated the changes in temperature and precipitation relative to their 1895-1924 means. Second, we produced a 30-year backward-running mean for the yearly changes in temperature and precipitation. We employed a 30-year backward-  
400 running mean because the response time of Pacific Northwest glaciers similar to those on Mt. Hood is 20-30 years (Pelto and Hedlund, 2001). The prior 30-year mean reflects this lagged relationship between climate change and glacier change. We explored the lower limit of the glacier-response times by using a 15-year backward-running mean. While Pelto and Hedlund (2001) found a lower limit to glacier-response time of 20 years, glaciers on Mt. Baker in Washington (Fig. 1A) may respond faster at 11-20 years (Pelto, 2016), which we encompass with our 15-year backward-running mean. Third, the relationship  
410 between the normalized change in glacier length at a given year of measurement was compared to the prior 30-year or 15-year mean change in temperature or precipitation. Correlations were determined through regression, with  $p < 0.05$  indicating significant correlation.

**Deleted:** moving prior-30-year average because this is the U.S. NOAA definition of a climate period. We explore

**Deleted:** We chose Hood River County as this is the smallest geographic region that includes these five glaciers and would still have direct weather station measurements within the geographic domain over the period of comparison. Snow telemetry stations on and around Mt. Hood were only installed in the late 1970s, with temperature sensors beginning to work consistently in the late

**Formatted:** Font: Not Bold

### 3 Results

405 Since 2003, every glacier on Mt. Hood underwent a rise in terminus elevation (Table 1, Fig. 2), with an average rise of ~150 m (observed changes and repeat photographs provided in Supplement). Note that these are inherently minimum estimates of the rises given the point-nature of the 2003 observations. There does not appear to be any pattern in the degree of terminus rise relative to glacier aspect on Mt. Hood (Fig. 2). The greatest terminus rise occurred in the southeast-facing Clark drainage of Newton-Clark Glacier (>300 m) followed by the southwest-facing Zigzag Glacier (>250 m; Fig. 3A, B), the west-facing Sandy Glacier (>240 m; Fig. 3G, H) and the northwest-facing Ladd Glacier (>230 m; Fig. 3E, F). Of the three glaciers with substantial debris cover of their termini, Ladd (>230 m) and Coe (>170 m) (Fig. 3C, D) had terminus rises greater than the average while Eliot's terminus rose >70 m (Table 1, Fig. 2).

**Formatted:** Normal, Line spacing: single, Widow/Orphan control

415 In the last seven to eight years, every glacier on Mt. Hood lost area (Fig. 1B), resulting in  $18 \pm 1\%$  loss in total glacier area (Table 1) relative to 2015/2016 at a rate of  $2.40 \pm 0.02\% \text{ yr}^{-1}$  ( $\text{yr} = \text{year}$ ). Glisan Glacier ceased to show evidence of flow between 2015 (Fountain et al., 2023) and our first field observation in 2020, with a stagnant ice area of  $0.009 \pm 0.0006 \text{ km}^2$  as of 2023. Zigzag, Coalman and Langille glaciers lost >40% of their area since 2015/2016. Ladd Glacier separated from its uppermost accumulation area (Fig. 3E, F) while Coe Glacier separated from its debris-covered terminus (Fig. 3C, D), both due to thinning over ice falls after 2016 (Fountain et al., 2023). A similar separation from its debris-covered terminus also occurred for Eliot Glacier (Fountain et al., 2023). When compared to the 1981 glacier inventory (Driedger and Kennard, 1986b), the glaciated area loss by 2023 for Mt. Hood glaciers was ~44% (Table 1) at a rate of  $\sim 1.0\% \text{ yr}^{-1}$ . The largest  
420 percent area lost since 1981 was the disappearance of south-facing Palmer Glacier (100%), followed by southwest-facing Zigzag Glacier (~87%), and north-facing Ladd Glacier (~74%).

**Formatted:** Heading 3, No widow/orphan control

**Formatted:** Font: Not Bold

**Deleted:**


**Formatted:** Superscript



Glacier (Multiple Termini)	2003 Terminus Elevation (m asl)	2023 Terminus Elevation (m asl)	20-Year Terminus $\Delta$ Elevation (m)	1981 Area km <sup>2</sup>	2015/16 Area km <sup>2</sup> (uncert)	2023 Area km <sup>2</sup> (uncert)	1981-2023 % $\Delta$ Area	2015/16-2023 % $\Delta$ Area (uncert)
Zigzag	2350	2600	250	0.77	0.168 (0.005)	0.101 (0.007)	87	40 (3)
Palmer	gone			0.13			100	
Coalman (E)	3130	3130	0	0.08	0.099 (0.001)	0.057 (0.004)	29	42 (3)
Coalman (W)	3190	3230	40					
White River	2130	2240	110	0.54	0.287 (0.046)	0.271 (0.019)	50	6 (1)
Newton-Clark (N)	2420	2480	60	1.99	1.134 (0.181)	0.975 (0.068)	51	14 (2)
Newton-Clark (S)	2320	2620	300					
Eliot	2050	2120	70	1.68	1.086 (0.174)	0.912 (0.064)	46	16 (3)
Langille (remain)		2520		0.40	0.233 (0.037)	0.138 (0.010)	66	41 (7)
Langille (upper)	2070	gone						
Langille (lower)	1960	gone						
Coe	1920	2090	170	1.24	0.779 (0.125)	0.696 (0.049)	44	11 (2)
Ladd	2100	2330	230	0.90	0.347 (0.055)	0.235 (0.016)	74	32 (6)
Glisan (upper)	2040	gone						
Glisan (lower)	1930	gone	>220		0.041 (0.006)			100
Sandy	1890	2130	240	1.19	0.751 (0.120)	0.631 (0.044)	47	16 (3)
Reid	2240	2350	110	0.75	0.469 (0.005)	0.417 (0.029)	44	11 (1)
Total				9.67	5.39 (0.32)	4.43 (0.12)	44	18 (1)

Table 1: Mt. Hood glacier observations from 2003 and 2023. Change ( $\Delta$ ) in terminus elevation, glacier area in 1981 (Driedger and Kennard, 1986b) and 2015/2016 (Fountain et al., 2023), and area change ( $\Delta$ ) as a percent (%) up to 2023 provided.

Deleted: in



435 Figure 2: Minimum change ( $\Delta$ ) in Mt. Hood glacier termini elevation from 2003 to 2023 relative to glacier position on Mt. Hood; C = Coalman, WR = White River, NC = Newton-Clark with its two termini labeled, and E = Eliot.




Figure 3: Select repeat photographs from Mt. Hood; see Fig. 1B for photo locations. A and B: Zigzag Glacier across the valley (shot location 45.29275°, -121.80044°) repeat photograph from 2003 (A) and 2023 (B). 1. Denotes the reduction in area of Zigzag Glacier to its 2023 limit (dashed white line). 2. Remnant stagnant ice that used to be part of the actively flowing glacier. 3. Western terminus of Coalman Glacier that has retreated but is still visible in 2023 at edge of cloud. 4. Top of White River Glacier that has retracted since 2003. C and D: Langille and Coe Glaciers (shot location 45.39459°, -121.69982°) in 2003 (C) and 2023 (D). 5. Disappearance of remnant ice of Langille Glacier. 6. Loss of an ice fall on Coe Glacier due to terminus retreat to just below the 6 in D. E and F: Ladd Glacier (shot location 45.39718°, -121.70424°) in 2003 (E) and 2023 (F). 7. Separation of Ladd from its uppermost accumulation area. 8. Ice fall that is now the current terminus of Ladd. 9. Retreat of the ice margin where actively flowing ice is now detached stagnant ice. G and H: Sandy Glacier (shot location 45.39477°, -121.73124°) in 2003 (G) and 2023 (H). 10. Accumulation of rock fall on the glacier in the last 20 years. 11. Retreat of glacier ice that is now exposed bedrock. 12. 2023 terminus location and ice caves (H) that were in the centre of the glacier in 2003 (G). 13. 2003 terminus with ice cave (G) that is now stagnant, debris-covered ice (H).

The seven largest glaciers on Mt. Hood (White River, Newton-Clark, Eliot, Coe, Ladd, Sandy, and Reid) that have area records extending back to 1907 (Jackson and Fountain, 2007) lost as a whole  $57\pm 3\%$  of their area since 1907 and  $39.8\pm 1.7\%$  of their area since 2000 (Table 2, Fig. 4). From 2000 to 2023, the rate of area loss was  $1.73\pm 0.17\%$  yr<sup>-1</sup>, about six times faster than the 1907-2000 rate of  $0.31\pm 0.02\%$  yr<sup>-1</sup> (Table 2). Ladd Glacier underwent the greatest rate of retreat in the 21st century, which was associated with the partial loss of its accumulation zone (Table 2; Fig. 3E, F). Conversely, White River Glacier had the greatest 20th century rate of retreat (Table 2), which has been previously attributed to the development of fumaroles by 1907 that reduced its accumulation area (Lillquist and Walker, 2006). Coe Glacier underwent the greatest increase in retreat rate in the 21st century about 12 times faster than its 20th century rate of area loss (Table 2).

Glacier	1907 Area km <sup>2</sup> (uncert)	2000 Area km <sup>2</sup> (uncert)	2023 Area km <sup>2</sup> (uncert)	1907-2023 % Δ Area (uncert)	2000-2023 % Δ Area (uncert)	1907-2000 % yr <sup>-1</sup> (uncert)	2000-2023 % yr <sup>-1</sup> (uncert)
White River	1.04 (0.11)	0.44 (0.03)	0.271 (0.019)	74 (13)	38.4 (3.8)	0.62 (0.08)	1.67 (0.17)
Newton-Clark	2.06 (0.15)	1.54 (0.11)	0.975 (0.068)	53 (10)	36.7 (3.7)	0.27 (0.03)	1.60 (0.16)
Eliot	2.03 (0.16)	1.66 (0.12)	0.912 (0.064)	55 (10)	45.1 (4.5)	0.18 (0.02)	1.55 (0.15)
Coe	1.41 (0.13)	1.21 (0.08)	0.696 (0.049)	51 (12)	42.5 (4.1)	0.15 (0.02)	1.85 (0.18)
Ladd	1.07 (0.11)	0.71 (0.05)	0.235 (0.016)	78 (12)	66.9 (6.6)	0.36 (0.04)	2.94 (0.29)
Sandy	1.61 (0.17)	1.02 (0.07)	0.631 (0.044)	61 (13)	38.1 (3.7)	0.39 (0.05)	1.71 (0.17)
Reid	0.79 (0.13)	0.53 (0.04)	0.417 (0.029)	47 (18)	21.3 (2.2)	0.35 (0.06)	0.90 (0.09)
Seven combined	9.98 (0.37)	7.11 (0.21)	4.282 (0.127)	57 (3)	39.8 (1.7)	0.31 (0.02)	1.73 (0.17)

Table 2: Area for seven largest glaciers on Mt. Hood and their change (Δ) in area and rate of change expressed as percent. 1907 and 2000 areas are from Jackson and Fountain (2007).

Formatted: Caption, Line spacing: single, No widow/orphan control

Formatted: English (UK)

Deleted: Mt. Hood has 12 named glaciers recorded on U.S. federal maps; we show the most recent U.S. Forest Service map (2016) in Fig. 2. We first describe the changes in the 11 remaining glaciers over the last 20 years, Palmer Glacier having ceased to exist prior to this time period. All volcanic age and rock descriptions come from the U.S. Geological Survey Eruption History of Mount Hood, Oregon website (U.S. Geological Survey, 2023). The glacier areas noted for 2015 or 2016 are from Fountain et al. (2023). We then examine potential factors affecting the magnitude of glacier change. We also investigate the change in area with respect to historic changes since 1907 for seven of these glaciers. Lastly, we test the hypothesis that changes in five of these glaciers since 1901 are related to regional climate change.

3.1 Observed Glacier Change in the 21st Century


Figures 3, 5 and 6 show repeat photographs of the glaciers on Mt. Hood, whereas Fig. 4 has photographs of Coalman Glacier from 2003 and 2021. Figure 7 is a summary of the changes in glacier length and terminus elevation from 2003 to 2023. Figure 8 illustrates glacier area for seven of the glaciers and their sum from 1901/1907 to 2023. Our field measurements are provided in Table 1.

Formatted: Heading 3, Line spacing: 1.5 lines, No widow/orphan control

Deleted: that

Deleted: From 2000 to 2023, these glaciers lost a combined  $\sim 2.8$  km<sup>2</sup> ( $7.11\pm 0.21$  km<sup>2</sup> to  $4.3\pm 0.12$  km<sup>2</sup>) or  $\sim 40\%$  of their 2000 area (Fig. 8). When related to their 1907 combined areas, they have lost  $\sim 5.7$  km<sup>2</sup> ( $9.98\pm 0.37$  km<sup>2</sup> to  $4.3\pm 0.12$  km<sup>2</sup>) or  $\sim 60\%$  of their 1907 area. For context, from 1907 to 2000, these glaciers lost  $\sim 2.9$  km<sup>2</sup> ( $9.98\pm 0.37$  km<sup>2</sup> to  $7.11\pm 0.21$  km<sup>2</sup>) or  $\sim 30\%$  of their 1907 area. Between 2000 to 2023, the total area lost by these seven glaciers is  $\sim 2.8$  km<sup>2</sup>.

Formatted: Font: Not Bold



Deleted:





Figure 4: Glacier area from 1907 (or 1901 for Eliot Glacier) to 2004 from Jackson and Fountain (2007), 2015 or 2016 from Fountain et al. (2023), and 2023 (new data), with combined area noted in the lower right. Uncertainty noted by vertical bars; note that the uncertainty on the combined record is smaller than the symbol size. A. White River Glacier. B. Newton-Clark Glacier. C. Eliot Glacier. D. Coe Glacier. E. Ladd Glacier. F. Sandy Glacier. G. Reid Glacier. H. Combined record of seven glaciers.

The 21st century rate of total area loss for these seven glaciers ( $1.73 \pm 0.17 \% \text{ yr}^{-1}$ ) exceeded the prior maximum rate of area loss from 1907-1946 ( $0.56 \pm 0.02 \% \text{ yr}^{-1}$ ) by a factor of about three (Fig. 5H). Five of the seven glaciers had individual 2000-2023 recession rates faster than their 1907-1946 rates (Fig. 5B-F). Reid Glacier's 2000-2023 rate of recession was  $0.90 \pm 0.09 \% \text{ yr}^{-1}$  while its 1907-1946 rate was  $0.84 \pm 0.14 \% \text{ yr}^{-1}$  (Fig. 5G). White River Glacier retreated at  $1.67 \pm 0.17 \% \text{ yr}^{-1}$  2000-2023 whereas its 1907-1946 rate was  $1.41 \pm 0.19 \% \text{ yr}^{-1}$  (Fig. 5A).

- Formatted: Superscript
- Formatted: Normal, Space Before: 12 pt, After: 12 pt
- Formatted: Superscript
- Formatted: Superscript
- Formatted: Superscript
- Formatted: Superscript
- Formatted: Superscript



520 **Figure 5: Area change with respect to preceding ice area expressed as % yr<sup>-1</sup> (yr = year) for seven glaciers calculated from Jackson and Fountain (2007) (1907-2000) and 2023 observations, with combined rate in the lower right. Gray shading shows the uncertainty. A. White River (WR) Glacier. B. Newton-Clark (NC) Glacier. C. Eliot Glacier. D. Coe Glacier. E. Ladd Glacier. F. Sandy Glacier. G. Reid Glacier. H. Combined record of seven glaciers.**

555 Updating the length records for White River, Newton-Clark, Eliot, Coe and Ladd glaciers of Lillquist and Walker (2006) to 2023 found that four of these glaciers had now lost ~35% (White River) to ~42% (Eliot) of their 1901 length while Ladd Glacier had lost ~65% of its 1901 length (Fig. 6C). Regression of relative-length change identified significant ( $p < 0.05$ ) relationships (Table 3) with both 30-year and 15-year backward-running mean-annual, May-October and June-September temperature change relative to the 1895-1924 mean (Fig. 6B). Conversely, no significant relationships (Table 3) were found with either 30-year or 15-year backward-running mean-annual, November-April or October-May precipitation change relative to the 1895-1924 mean (Fig. 6A). Note that Newton-Clark and White River glaciers are missing an observation for

560 1928 while Ladd is missing the year 1959; this impacted the number of observations for determining regression significance.

Formatted: English (UK)

Formatted: Normal, Line spacing: single

Deleted: 3.2 Potential factors in glacier retreat¶

Deleted: All glaciers on Mt. Hood retreated in the last 20 years with a rise in the elevation of their termini. We find a strong relationship between the magnitude of terminus retreat and the degree of terminus-elevation rise (Fig. 7C), with an  $r^2$  of 0.86 ( $p < 0.00$ ). We now examine potential controls that could explain the different magnitude of terminus retreat between the glaciers. We consider glacier aspect, accumulation-area slope, highest point of the accumulation zone, valley shape and potential buttressing, debris cover, and substrate, as well as glacier-specific influences (e.g., fumaroles) and the amount of 20th century retreat. ¶

The degree of terminus retreat and rise in elevation does not correlate with aspect (Fig. 7A, B). There is also no significant relationship between the slope of glacier accumulation area ( $p = 0.28$ ) or the highest elevation of the accumulation area ( $p = 0.33$ ) and the magnitude of terminus retreat. While difficult to quantify buttressing, we do not see a correlative pattern of glaciers flowing over a broad surface with minimal buttressing (Zigzag, Newton-Clark) receding more than glaciers in wide valleys (Sandy, Ladd), narrow valleys (White River, Reid), or very narrow valleys (Coe, Eliot). Debris cover may influence the magnitude of glacier retreat but does not correlate with the magnitude of retreat. The glaciers with the highest debris cover (Eliot, Coe) have retreated around the average of Mt. Hood's glaciers while a third glacier with heavy debris cover (Ladd) has, for other reasons, retreated the most (Figure 8C). Furthermore, substrate type and age (i.e., centuries-old pyroclastic flows versus tens of millennia old andesite or dacite rock) does not appear to be a factor, because glaciers on older rock (Coe, Eliot, Ladd) versus younger volcanic sediment (Zigzag, White River) have similar magnitudes of margin recession. ¶

When comparing adjacent glaciers, topographic and other factors could explain different amounts of retreat. Sandy Glacier retreated ~310 m more than Reid Glacier, which could be due to Sandy having an ice cave system that likely enhances ablation; Reid having greater shading and buttressing; and Reid terminating at a cliff that impeded down-valley extension thereby reducing its initial retreat to a climate perturbation. The existence of fumaroles since at least 1907 certainly have influenced the behavior of Coalman Glacier's two lobes (Lillquist and Walker, 2006), which had the least amount of retreat on Mt. Hood (30 and 110 m). In the absence of this volcanic heat, Coalman Glacier might still be connected to White River and Zigzag glaciers as part of their accumulation zones, and thus not have retreated at all. Coe and Eliot are similar glaciers, but Eliot has more debris cover and shading than Coe, which probably explains Coe's greater retreat. Ladd Glacier's loss of part of its accumulation zone and the westward bend of its ablation zone must have contributed to the largest retreat of any glacier on Mt. Hood in this century. ¶

3.3 Decadal Rate of Area Loss for Seven Glaciers¶

Figure 9 shows the absolute rate of change in aerial extent for these seven glaciers while Fig. 10 illustrates the relative rate of change in aerial extent. From 2000 to 2023, White River Glacier lost area at a rate of 0.066-0.081 km<sup>2</sup> decade<sup>-1</sup>, which is a little less than half the maximum rate of area loss that occurred from 1907 to 1946 of 0.127-0.165 km<sup>2</sup> decade<sup>-1</sup>. However, when assessed as the rate of loss relative to prior extent, the most recent 23 years of retreat at 15.0-18.4 % decade<sup>-1</sup> was probably faster than the 1907-1946 rate of retreat at 12.2-15.9 % decade<sup>-1</sup>. Reid Glacier also retreated at the second fastest rate of 0.043-0.053 km<sup>2</sup> decade<sup>-1</sup> for 2000-2023, which is slightly slower than its 1907-1946 rate of 0.056-0.078 km<sup>2</sup> d... [7]




Figure 6: Hood River County climate change (NOAA) and Mt. Hood glacier length change (Lillquist and Walker, 2006). A. 30-year (solid) and 15-year (dashed) backward-running mean for total annual (dark blue) and November-April (light blue) precipitation relative to 1895-1924 average. B. 30-year (solid) and 15-year (dashed) backward-running mean for mean annual (dark red) and May-October (pink) temperature relative to 1895-1924 average. C. Change in glacier length normalized to 1901 length (Lillquist and Walker, 2006) from 1901 to 2023 for White River (purple), Newton-Clark (green), Eliot (orange), Coe (black), and Ladd (red) Glaciers.

Glacier	Annual T (1901-2023) $r^2, p$	MJJASO T (1901-2023) $r^2, p$	Annual (1901-2023) $r^2, p$
White River	0.63, <0.01	0.67, <0.01	<i>0.10, &gt;0.05</i>
Newton-Clark	0.42, 0.03	0.58, 0.01	<i>0.01, &gt;0.05</i>
Eliot	0.91, <0.01	0.86, <0.01	0.77, <0.01
Coe	0.87, <0.01	0.87, <0.01	0.55, <0.01
Ladd	0.82, <0.01	0.73, <0.01	0.70, <0.01

**Deleted:**  
**Table 2:** Regression  $r^2$  and  $p$  from Fig. 12. Italics indicates non-significant results. An asterisk identifies a significant correlation that is opposite of an expected glacier-climate relationship. We test the influence of our 2023 observations on the statistically significant relationships between glacier length and temperature by conducting the same analysis but for 1901 to 2000. For Eliot, Coe, and Ladd glaciers, the strength of the relationship decreases for annual and May-October temperature but remains statistically significant ( $p = < 0.01$  to  $0.02$ ). In contrast, White River and Newton-Clark glaciers do not have statistically significant relationships with either annual or May-October temperature. A similar test with annual and November-April precipitation yields significant relationships only for Coe Glacier ( $p = 0.01, 0.02$ ), where greater glacier retreat is associated with higher annual ( $p = 0.01$ ) and November-April ( $p = 0.02$ ) precipitation (Fig. 12), which is opposite of the usual relationship between glacier retreat and lower precipitation.

- Deleted: 11
- Deleted: Prior 30-year average
- Deleted: Prior
- Deleted: average
- Deleted:

695


Glacier (30 yr)	Ann T	M-O T	J-S T	Ann P	N-A P	O-M P
White River	<b>0.63</b>	<b>0.67</b>	<b>0.69</b>	0.02	0.02	0.01
Newton-Clark	<b>0.42</b>	<b>0.58</b>	<b>0.46</b>	0.04	0.08	0.02
Eliot	<b>0.93</b>	<b>0.94</b>	<b>0.95</b>	0.07	0.05	0.07
Coe	<b>0.87</b>	<b>0.90</b>	<b>0.90</b>	0.16	0.09	0.16
Ladd	<b>0.82</b>	<b>0.73</b>	<b>0.77</b>	0.06	0.05	0.04
Glacier (15 yr)	Ann T	M-O T	J-S T	Ann P	N-A P	O-M P
White River	<b>0.78</b>	<b>0.76</b>	<b>0.70</b>	0.13	0.24	0.08
Newton-Clark	<b>0.50</b>	<b>0.62</b>	<b>0.50</b>	0.02	0.14	<0.01
Eliot	<b>0.79</b>	<b>0.85</b>	<b>0.86</b>	<0.01	<0.01	0.02
Coe	<b>0.60</b>	<b>0.66</b>	<b>0.72</b>	0.05	0.01	0.10
Ladd	<b>0.73</b>	<b>0.67</b>	<b>0.65</b>	<0.01	0.02	0.01

Table 3: Regression results ( $r^2$ ) for 30-year (top) and 15-year (bottom) backward-running means of temperature (T) and precipitation (P) relative to normalized glacier length. Bold indicates significant results where  $p < 0.05$ . Ann = Annual, M-O = May-October, J-S = June-September, N-A = November-April, O-M = October-May.

#### 4 Discussion

In the 21st century, every glacier on Mt. Hood has lost area, resulting in about 40% glaciated area loss since 2000, with a minimum rise in terminus elevation of ~150 m on average over the last 20 years. This rapid area loss combined with a rise in terminus elevation are indicative of large mass balance losses (e.g., Haerberli and Hoelzle, 1995; Pelto and Brown, 2012; Pelto, 2016; Hoelzle et al., 2023). Such clear glacier retraction and inferred negative mass balance stand at odds with the Menounos et al. (2019) finding of near-neutral mass change for glaciers on Mt. Hood from 2000 to 2018. We note that their study lacked ground truthing of their geodetic measurements in Oregon, which combined with the high and growing debris cover on many of Mt. Hood's glaciers (e.g., Fig. 3H), could explain this disagreement. Furthermore, Florentine et al. (2024) found that not accounting for glacier area change when measuring geodetic mass balance reduces the mass-balance-change signal in western North American glaciers, with greater area loss leading to greater mass-balance bias. Here, our field observations clearly show large-scale glacier loss, consistent with the 21st-century retreat of glaciers around the globe (Zemp et al., 2015) and in agreement with the cumulative mass loss of benchmark glaciers to the north of Mt. Hood in Washington State (O'Neel et al., 2019; Pelto, 2023; World Glacier Monitoring Service, 2022).

The combined retreat for Mt. Hood's seven largest glaciers over the first 23 years of the 21st century stands out as unparalleled since at least the beginning of the 1900s (Fig. 4, 5), similar to the findings of Beason et al. (2023) for glaciers on Mt. Rainier in Washington and agreeing with a shift to more negative annual mass balance for western United States benchmark glaciers since 1990 (O'Neel et al., 2019). We compare Mt. Hood's rate of glacier recession of  $1.73 \pm 0.17 \text{ \% yr}^{-1}$  from 2000 to 2023 (seven largest glaciers; Table 2) and  $2.40 \pm 0.02 \text{ \% yr}^{-1}$  from 2015/2016 to 2023 (all glaciers; Table 1) to other western United States glaciated ranges with comparable records of actively flowing glacier area. To the north in Washington State (Fig. 1A), Mt. Rainier glaciers retreated at  $0.59 \pm 0.05 \text{ \% yr}^{-1}$  from 1994 to 2021 and  $0.69 \pm 0.05 \text{ \% yr}^{-1}$  from



2015 to 2021 (Beason et al., 2023) while glaciers in the Olympic Mountains retreated at  $1.44 \pm 0.02$  %  $\text{yr}^{-1}$  from 1990 to 2015 and  $2.75 \pm 0.04$  %  $\text{yr}^{-1}$  from 2009 to 2015 (Fountain et al., 2022). To the northeast in Montana (Fig. 1A), Glacier National Park glaciers receded at  $-0.77$  %  $\text{yr}^{-1}$  from 1998 to 2015 and  $-0.83$  %  $\text{yr}^{-1}$  from 2005 to 2015 (Fagre et al., 2017). Lastly, to the south in California (Fig. 1A), glaciers in the Trinity Alps retreated at  $3.89 \pm 0.02$  %  $\text{yr}^{-1}$ , between 1994 and 2015, leading to near complete deglaciation of the mountain range by 2015 (Garwood et al., 2020). Thus, the rates of glacier retreat on Mt. Hood in the 21st century are some of the highest rates in the western United States, but have yet to exceed the 2009-2015 rate in the Olympic Mountains, where the rate-change record ended in 2015, or the rate in the Trinity Alps, where the record ended in 2015 with near complete deglaciation.

In the last 20 years, Glisan Glacier on Mt. Hood has retreated to a 2023 area of  $\sim 0.009$   $\text{km}^2$  and ceased to be an actively flowing glacier, joining Palmer Glacier in this former-glacier status (Table 1; Supplement). We find that two other glaciers are nearing this reclassification as former glaciers; Zigzag and Coalman (Supplement). At  $\sim 0.1$   $\text{km}^2$ , Zigzag Glacier was only distinguishable from stagnant ice due to the presence of a few crevasses that have yet to start melting in on themselves (i.e., movement is keeping them open). Coalman Glacier at  $\sim 0.06$   $\text{km}^2$  (Table 1) is now below the area threshold of  $0.1$   $\text{km}^2$  used by the U.S. Geological Survey to distinguish active glaciers (Fagre et al., 2017), but is above the  $0.01$   $\text{km}^2$  threshold used in global inventories (Pfeffer et al., 2014). More importantly, Coalman still showed evidence of flow with one active crevasse. With these two glaciers losing 40-42% of their area in the last seven to eight years (Table 1), we predict that in the near future they will stagnate and become former glaciers. Langille Glacier also lost >40% of its area while Ladd Glacier lost about a third of its area, both in the last seven years (Table 1). If this rapid retreat continues, we predict that Langille and Ladd glaciers will be the next Mt Hood glaciers, after Zigzag and Coalman, to stagnate.

Because of the significant relationship between the change in glacier length for five of Mt. Hood's glaciers and the change in regional temperature up to 2023 (Fig. 6B, C; Table 3), we argue that the retreat of glaciers over the last  $\sim 120$  years reflects regional climate warming (Roe et al., 2016, 2021). We do not find a similar relationship with precipitation (Fig. 6A, C; Table 3). A Mt. Hood glacier-length-temperature relationship agrees with mass-balance-temperature relationships for glaciers to the north in Washington State and for western United States benchmark glaciers. Peltó (2018) found significant relationships between annual mass balance with June-September temperature for 11 glaciers in Washington. O'Neel et al. (2019) noted the lack of a winter-mass-balance trend for the four benchmark glaciers in Washington and Alaska and a positive winter-mass-balance trend for the one glacier in Montana, meaning these glaciers' negative annual-mass-balance trends were being driven by summer temperatures.

We suspect that the lack of a Mt. Hood glacier length-temperature relationship for the 20th century in the Lillquist and Walker (2006) study may partly reflect their methodology. Lillquist and Walker (2006) used a 5-year running average for both temperature and precipitation, shorter than the 11- to 30-year response time for glaciers like those on Mt. Hood (Peltó and Hedlund, 2001; Peltó, 2016), which we approximate with our 30-year and 15-year backward-running means.

**Deleted:** this Olympic Mountains and that the rates on Mt. Hood are yet to exceed ...

**Deleted:** ,

**Deleted:** whose

**Deleted:** c

**Deleted:** status

**Deleted:** is

**Deleted:** of  $0.1$   $\text{km}^2$

**Deleted:** still has one active crevasse

**Formatted:** Superscript

**Deleted:** At  $\sim 0.1$   $\text{km}^2$ , Zigzag Glacier is also near this U.S. Geological Survey glacier demarcation. With these two glaciers

**Deleted:** 30-

**Deleted:** suggest

**Deleted:** and join Palmer and Glisan as former glaciers

**Deleted:** is also retreating rapidly, losing

**Deleted:** Glacier

**Deleted:** reach stagnation on Mt. Hood

**Deleted:** The rate of area loss in the last 23 years is the greatest in the last  $\sim 120$  years for four of the seven glaciers on Mt. Hood, and another two glaciers have retreat rates near their 1907-1946 rates (Fig. 9) despite the potential influence of volcano geometry that would favor earlier faster rates (Fig. 1). Only White River Glacier had a significantly faster absolute rate of area loss from 1907 to 1946 (Fig. 9). This faster rate was likely facilitated by volcanic activity between 1853 and 1907 that separated White River Glacier from Coalman Glacier, with the reduction in accumulation area driving greater White River retreat (Lillquist and Walker, 2006). The sum of the retreat rates of the seven Mt. Hood glaciers over the first 23 years of the 21st century stands out as unparalleled since at least the beginning of the 1900s (Fig. 9). In the 20th century, the maximum absolute and relative rate of area loss for these glaciers occurred 1901-1947. This 1907-1946 interval of retreat corresponds with a warming-drying period in the Pacific Northwest that is attributable to human greenhouse gas emissions (Marvel et al., 2019). Furthermore, the stabilization or even slight advance of Mt. Hood's glaciers 1946-1972 (Fig. 8, 9) coincides with a cooler-wetter period in the Pacific Northwest that was in response to human-aerosol emissions counteracting human-greenhouse gas emissions (Marvel et al., 2019).

**Deleted:** statistically

**Deleted:** Interestingly, we did find significant relationships between Eliot, Coe, and Ladd glacier lengths and prior 30-year average temperatures for the 20th century while ...

**Deleted:** did not.

**Deleted:** We attribute this difference, at least partly, to how the climate data were analyzed.

**Deleted:** , which would mean the glacier's current length is only related to the current year and two prior years of weather, with ... [11]

**Deleted:**



Frans et al. (2016) modeled the evolution of Newton-Clark, Eliot, Coe, and Ladd glaciers from 2000 to the end of the century under intermediate (RCP4.5) and high (RCP8.5) emission scenarios, with 10 simulations per emission scenario. From 2000 to 2025, they simulated a 10-30% loss in glacier area regardless of emission scenario. In contrast, we find that these four glaciers lost 30-50% of their combined area between 2000 and 2023. This means that glacier recession on Mt. Hood in the 21st century has proceeded at a faster rate than simulated for even the highest emission scenario. This also raises concerns over the persistence of Mt. Hood glaciers at all in the 21st century. Frans et al. (2016) modeled continued glacier presence on Mt. Hood in 2100 even under RCP8.5 forcing. However, the observed glacier retreat in the last 20 years has equaled to exceeded by 67% the maximum loss simulated under RCP8.5 emissions for the last 20 years.

We hypothesize that the observed faster retreat of glaciers on Mt. Hood relative to their simulated retreat may, in part, be due to climate-related weather phenomena that had yet to occur at the time of the model forcing of Frans et al. (2016). In 2015, Oregon experienced the lowest spring snowpack ever recorded (Mote et al., 2016), which was followed by the warmest summer ever recorded (U.S. NOAA, 2024, <https://www.ncei.noaa.gov/access/monitoring/climate-at-a-glance/>). Such extreme events are absent from the Frans et al. (2016) forcing and may have significantly impacted glacier mass balance (Pelto, 2018). Likewise, the June 2021 Pacific Northwest heatwave was one of the six most extreme events ever recorded on the planet (Thompson et al., 2022), which would have negatively impacted glacier mass balance (Pelto et al., 2022). In 2020, more forest area burned in the Oregon Cascades than all 36 prior years of record combined (Abatzoglou et al., 2021) while the 2021 heatwave further exacerbating forest fires (Jain et al., 2024). Since 2015, fire-smoke particulates have been rising in Oregon (Burke et al., 2023). The particulates from these fires accumulate on Oregon glaciers, where they reduce albedo and increase ablation (Kaspari et al., 2015; Allgaier et al., 2022). Given the under projection of glacier change on Mt. Hood and given the analogous regional snow-drought and heatwave impacts (Mote et al., 2016; Philip et al., 2022; Thompson et al., 2022) and rise in fire-smoke particulates (Neff et al., 2012; Kaspari et al., 2020) in both Washington and British Columbia, new assessments of recent (e.g., post 2015) glacier retreat in Washington's Olympic, Central Cascade, and North Cascade ranges and in the ranges of British Columbia where similar glacier model simulations have been conducted (Clarke et al., 2015; Frans et al., 2018) could be used to test the accuracy of these model results for the first quarter of the 21st century.

Our finding that the Frans et al. (2016, 2018) model under simulates recent glacier recession on Mt. Hood is of vital import to downstream communities. The U.S. Bureau of Reclamation conducted a Hood River Basin Study that utilized the same model as Frans et al. (2016, 2018). This federal study projected only a 1-4% decline in Mt. Hood glacier area through 2060 (U.S. Bureau of Reclamation, 2015). We argue that this study, which underlies water management plans for the Hood River Basin and reliant agriculture (Thieman et al., 2021), contains overly optimistic simulations of future glacier retreat, and that water resource managers who rely on this federal study are ill-prepared for future late-summer declining stream flow and rising instream temperatures. Thieman et al. (2021) noted that glacier disappearance in the Hood River Basin would be catastrophic for salmon populations. Our findings imply this this could happen sooner than is projected in the current Hood

**Deleted:** However, we did not find a significant relationship for White River and Newton-Clark glaciers with temperature or precipitation over the 20th century. White River's early 20th-century retreat clearly had a non-climatic cause of glacier change from the partial loss of its accumulation zone due to volcanic activity (Lilquist and Walker, 2006). Newton-Clark's main body in the Newton drainage terminated at a cliff for much of the 20th century. This cliff might have limited its down valley extent, reducing its response to a warming climate until the climate warmed sufficiently to drive retreat from the cliff as is now the case. Nevertheless, both glaciers have a significant temperature relationship once the 21st century changes are included. ¶

**Deleted:** area of

**Deleted:** changed from 5.12±0.19 km<sup>2</sup> in 2000 to 2.97±0.11 km<sup>2</sup> in 2023, losinglost

**Deleted:** probably

**Deleted:** over the same time period

**Deleted:** were not included in

**Deleted:** U.S. NOAA, 2023

**Deleted:** se light-absorbing

**Deleted:** this

**Deleted:** along with a

**Deleted:** we suggest that such

**Deleted:** hindsight

**Deleted:** could be conducted for

**Deleted:** glaciers

**Deleted:** all the glaciers of British Columbia

**Deleted:** The rapid glacier retreat in the 21st century that we document here raises questions over the timing of peak meltwater for glaciers on Mt. Hood. Huss and Hock (2018) simulated peak melt water occurring between 2010 and 2020 using the intermediate RCP4.5 emission scenario. Frans et al. (2018) modeled peak melt water for Mt. Hood glaciers as occurring between 2010 and 2050 with a median of 2030 for RCP4.5 and RCP8.5. Note that this timing of peak meltwater is based on the same simulations of Frans et al. (2016) that, with the now-afforded hindsight, under-projected retreat of Mt. Hood glaciers. While our glacier-area-change data do not directly address the timing of peak meltwater, we suggest that the rapid retreat we document here implies that peak meltwater has already occurred on Mt. Hood, similar to the simulations of Huss and Hock (2018). ¶

**Deleted:** with increased meltwater discharge

**Deleted:** , meaning peak water had yet to be reached in these simulations (U.S. Bureau of Reclamation, 2015). We argue that this

**Deleted:** that

**Deleted:** ¶

River Basin plans that have 2100 as the earliest possible year for deglaciation. The cold instream temperatures of Hood River are due to glacier meltwater and afford the river its diversity in salmon, steelhead, and bull trout (Thieman et al., 2021), which deglaciation of the river basin in this century would upset. Rapid glacier recession also imperils unique, and recently discovered, microbiomes on Mt Hood that only exist at glacier-fed springs on the volcano (Miller et al., 2021). As such, glacier-dependent ecosystems on Mt. Hood may be more at risk than previously surmised. However, the most visible impact, and the one affecting the greatest number of people in the near future, will likely be on the orchards of Hood River and Wasco counties whose orchards are the most important component of their economies (Columbia Gorge Fruit Growers, 2024, <http://www.cgfg.org/information/crops-information>).

**Deleted:** . D

**Deleted:** this river diversity as late-summer instream temperatures permanently warm. Rapid glacier recession also imperils unique, and

## 5 Conclusions

Contrary to results from remote-sensing alone, we document from field observations that the glaciers of Mt. Hood retreated, and most likely lost mass, in the 21st century, losing ~40% of their area (seven largest glaciers) in the last 23 years and ~18% of their area (all glaciers) in the last seven to eight years at unprecedented rates with respect to the last 120 years ( $1.73 \pm 0.17$  % yr<sup>-1</sup> and  $2.40 \pm 0.02$  % yr<sup>-1</sup>, respectively). The rate of area loss in the 21st century has proceeded about three times faster than the fastest rate of the 20th century and about six times faster than the average 20th century rate. Two glaciers have ceased to be glaciers (Glisan, Palmer) with another four glaciers rapidly losing area towards this reclassification (Zigzag, Coalman, Langille, Ladd). The century-scale retreat of Mt. Hood glaciers can be attributed to a warming climate, with regional 30-year mean temperature 1.1-1.3°C warmer and 15-year mean temperature 1.3-1.5°C warmer than the 1895-1924 mean. As such, the warming climate of the last 120 years, especially the last 20 years, has significantly impacted the glacier coverage of Mt. Hood, with this recession continuing unless and until the sign of multi-decadal temperature change is reversed.

**Deleted:** We find that the overall recession of glaciers on Mt. Hood in the first two decades of the 21st century is unprecedented with respect to the last 120 years, surpassing in sum the glacier retreat that occurred in the early 1900s. This

**Deleted:** is

**Deleted:** average

**Deleted:** now ~

**Deleted:** than the 1895-1924 average

**Deleted:** On Mt. Hood, two glaciers have ceased to be glaciers (Glisan, Palmer) with another two glaciers nearing this reclassification (Zigzag, Coalman) and a third rapidly losing area towards this threshold (Ladd). ...

**Deleted:**

**Deleted:** and is threatening water resources and dependent ecosystems and economies. Tt

**Deleted:** will

**Deleted:** ue

**Deleted:** unless and

## Data availability

The data referred to in this paper have all been provided within the tables and figures in the main text and supplement.

## Author contributions

The author contributions, following the CRediT authorship guidelines, are as follows: NB-F, SJB, and AEC: conceptualization; NB-F, SJB, WCS, and AEC: methodology; NB-F, SJB, WCS, and AEC: validation; NB-F, SJB, WCS, MT, and AEC: analysis; NB-F, SJB, WCS, MT, DHR, and AEC: investigation; NB-F, SJB, WCS, DHR, and AEC: resources; NB-F, SJB, WCS, MT, and AEC: data curation; AEC: original draft; NB-F, SJB, WCS, MT, DHR, and AEC:

review and editing; NB-F, SJB, WCS, MT, DHR, and AEC: visualization; NB-F, SJB, WCS, MT, DHR, and AEC:  
985 administration; NB-F, SJB, WCS, MT, DHR, and AEC: funding acquisition.

### Competing interests

The authors declare that they have no conflict of interest.

### Acknowledgements

990 We thank the U.S. National Forest Service for permitting research access to the Mt. Hood Wilderness Area and Andrew  
Fountain for sharing the historical area record for Mt. Hood. [Insight from reviewers A. Fountain, M. Pelto and an  
anonymous reviewer along with community comments from F. Paul greatly improved this manuscript.](#) This work was  
supported by a conservation grant from the Mazamas Mountaineering Club. The Oregon Glaciers Institute is a non-profit  
501-c-3 that is supported by private donations.

### References

995 Abatzoglou, J. T., Rupp, D. E., O'Neill, L. W., and Sadegh, M.: Compound Extremes Drive the Western Oregon Wildfires  
of September 2020, *Geophys. Res. Lett.*, 48, e2021GL092520, 2021.

Allgaier, M., Cooper, M. G., Carlson, A. E., Cooley, S. W., Ryan, J. C., and Smith, B. J.: Direct measurement of optical  
properties of glacier ice using a photon-counting diffuse LiDAR, *J. Glaciol.*, 68, 1210-1220, 2022.

1000

Beason, S. R., Kenyon, T. R., Jost, R. P., and Walker, L. J.: Changes in glacier extents and estimated changes in glacial  
volume at Mount Rainier National Park, Washington, USA from 1896 to 2021, *National Resource Report*  
NPS/MORA/NRR-2023/2524, 2023.

1005

Burke, M., Childs, M. L., de la Cuesta, B., Qui, M., Li, J., Gould, C. F., Heft-Neal, S. and Wara, M.: The contribution of  
wildfire to PM<sub>2.5</sub> trends in the USA, *Nature*, 622, 761-766, 2023.

Clarke, G. K. C., Jarosch, A. H., Anslow, F. S., Radić, V., and Menounos, B.: Projected deglaciation of western Canada in  
the twenty-first century, *Nat. Geosci.*, 8, 372-377, 2015.

1010

[Columbia Gorge Fruit Growers: Crops Information, http://www.cgfg.org/information/crops-information, 2024.](http://www.cgfg.org/information/crops-information, 2024)

Dodge, N. A.: The Eliot Glacier: New Methods and Some Interpretations, *Mazama Bulletin*, 53, 25-33, 1971.

1015 Dodge, N. A.: Eliot Glacier: Net Mass Balance, *Mazama Bulletin*, 69, 52-55, 1987.

[Driedger, C. L., and Kennard, P. M.: Glacier volume estimation on Cascade volcanoes: An analysis and comparison with other methods, \*Ann. Glaciol.\*, 8, 59-64, 1986a.](#)

1020 Driedger, C. L., and Kennard, P. M.: Ice Volumes on Cascade Volcanoes: Mount Rainier, Mount Hood, Three Sisters, and Mount Shasta, U.S. Geological Survey Professional Paper 1365, 1986b.

Ellinger, J. R.: The Changing Glaciers of Mt. Hood, Oregon and Mt. Rainier, Washington: Implications for Periglacial Debris Flows M.S. Thesis Oregon State University, 2010.

1025

Fagre, D. B., McKeon, L. A., Dick, K. A., and Fountain, A. G.: Glacier margin time series (1966, 1998, 2005, 2015) of the named glaciers of Glacier National Park, MT, USA, U.S. Geological Survey data release <https://dx.doi.org/10.5066/F7P26WB1>, 2017.

1030 Florentine, C., Sass, L., McNeil, C., Baker, E., and O'Neil, S.: How to handle glacier area change in geodetic mass balance: *J. Glaciol.*, [doi.org/jog.2023.86](https://doi.org/jog.2023.86), 2024.

Fountain, A. G., Glenn, B., and Basagic, H. J.: The geography of glaciers and perennial snowfields in the American West, Arctic, Antarctic, and Alpine [Res.](#), 49, 391-410, 2017.

1035

Fountain, A. G., Glenn, B., and Meneil, C.: Inventory of glaciers and perennial snowfields of the conterminous USA, *Earth Syst. Sci. Data*, 15, 4077-4104, 2023.

Fountain, A. G., Gray, C., Glenn, B., Menounos, B., Pflug, J., and Riedel, J. L.: Glaciers of the Olympic Mountains, Washington – The Past and Future 100 Years, *J. Geophys. Res.*, 127, e2022JF006670, 2022.

1040

Frans, C., Istanbuluoglu, E., Lettenmaier, D. P., Clarke, G., Bohn, T. J., and Stumbaugh, M.: Implications of decadal to century scale glacio-hydrological change for water resources of the Hood River basin, OR, USA, *Hydrological Processes*, 30, 4314-4329, 2016.

1045

Deleted: Research

Deleted: Rsch

Frans, C., Istanbuluoglu, E., Lettenmaier, D. P., Fountain, A. G., and Riedel, J.: Glacier Recession and the Response of Summer Streamflow in the Pacific Northwest United States, 1960-2099, *Water Res. Res.*, 54, 6202-6225, 2018.

Deleted: Rsch

Garwood, J. M., Fountain, A. G., Lindke, K. T., van Hattem, M. G., and Basagic, H. J.: 20th Century Retreat and Recent Drought Accelerated Extinction of Mountain Glaciers and Perennial Snowfields in the Trinity Alps, California, *Northwest Science* 94, 44-61, 2020.

[GLIMS and NSIDC: Global Land Ice Measurements from Space Glacier Database, Compiled and Made Available by the International GLIMS Community and the National Snow and Ice Data Center, doi: 10.7265/N5V98602, 2005 updated 2018.](#)

[Haeberli, W., and Hoelzle, M.: Application of inventory data for estimating characteristics of and regional climate-change effects on mountain glaciers: a pilot study with the European Alps. \*Ann. Glaciol.\*, 21, 206-212, 1995.](#)

Handewith, H. Jr.: Recent Glacier Variations on Mt. Hood, *Mazama Bulletin*, 41, 23-28, 1959.

[Harper, J. T.: Glacier Terminus Fluctuations on Mount Baker, Washington, U.S.A., 1940-1990 and Climatic Variations, \*Arctic and Alpine Res.\*, 25, 332-340, 1993.](#)

[Hoelzle, M., Haeberli, W., Dischl, M., and Peschke, W.: Secular glacier mass balances derived from cumulative glacier length changes, \*Global Planet. Change\*, 36, 295-306, 2023.](#)

Howcutt, S., Spagnolo, M., Rea, B. R., Jaszewski, J., Barr, I., Coppola, D., De Siena, L., Girona, T., Gomez-Patron, A., Mullan, D., and Pritchard, M. E.: Icy Thermometers: Quantifying the impact of volcanic heat on glacier elevation, *Geology*, 51, 1143-1147, 2023.

Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussaillant, I., Brun, F., and Kääb, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592, 726-731, 2021.

[Huston, A., Siler, N., Roe, G. H., Pettit, E., and Steiger, N. J.: Understanding drivers of glacier-length variability over the last millennium, \*The Cryosphere\*, 15, 1645-1662, 2021.](#)

Deleted: ¶

Huss, M., and Hock, R.: Global-scale hydrological response to future glacier mass loss, *Nature Climate Change*, 8, 136-140, 2018.¶

Jackson, K. M., and Fountain, A. G.: Spatial and morphological change on Eliot Glacier, Mount Hood, Oregon, USA, *Ann. Glaciol.*, 46, 222-226, 2007.

Jain, P., Sharma, A. R., Acuna, D. C., Abatzoglou, J. T., and Flannigan, M.: Record-breaking fire weather in North America in 2021 was initiated by the Pacific northwest heat dome, *Communications Earth & Environment*, 5, 202, 2024.

Kaspari, S. D., Pittenger, D., Jenk, T. M., Morgenstern, U., Schwikowski, M., Buening, N., and Stott, L.: Twentieth Century Black Carbon and Dust Deposition on South Cascade Glacier, Washington State, USA, as Reconstructed From a 158-m-Long Ice Core, *J. Geophys. Res.*, 125, e2019JD031126, 2020.

Kaspari, S., Skiles, S. M., Delaney, I., Dixon, D., and Painter, T. H.: Accelerated glacier melt on Snow Dome, Mount Olympus, Washington, USA, due to deposition of black carbon and mineral dust from wildfire, *J. Geophys. Res.*, 120, 2793-2807, 2015.

Lawrence, D. B.: Mt. Hood's Latest Eruption and Glacier Advances, *Mazama Bulletin*, 30, 22-29, 1948.

Leclercq, P. W., and Oerlemans, J.: Global and hemispheric temperature reconstruction from glacier length fluctuations, *Clim. Dynam.*, 38, 1065-1079, 2012.

Leclercq, P. W., Oerlemans, J., Basagic, H. J., Bushueva, I., Cook, A. J., and Le Bris, R.: A data set of worldwide glacier length fluctuations, *The Cryosphere*, 8, 659-672, 2014.

Leigh, J. R., Stokes, C. R., Carr, J. R., Evans, I. S., Andreassen, L. M., and Evans, D. J. A.: Identifying and mapping very small (<0.5 km<sup>2</sup>) mountain glaciers on coarse to high-resolution imagery, *J. Glaciol.*, 65, 873-888, 2019.

Leonard, K. C., and Fountain, A. G.: Map-based methods for estimating glacier equilibrium-line altitudes, *J. Glaciol.*, 49, 329-336, 2003.

Lillquist, K., and Walker, K.: Historical Glacier and Climate Fluctuations at Mount Hood, Oregon, Arctic, Antarctic, and Alpine *Res.*, 38, 399-412, 2006.

Lundstrom, S. C., McCafferty, A. E., and Coe, J. A.: Photogrammetric analysis of 1984-89 surface altitude change of the partially debris-covered Eliot Glacier, Mount Hood, Oregon, U.S.A., *Ann. Glaciol.*, 17, 167-170, 1993.

Marzeion, B., Cogely, J. G., Richter, K., and Parkes, D.: Attribution of global glacier mass loss to anthropogenic and natural causes, *Science*, 345, 919-921, 2014.

Deleted: Rsch

Deleted: sch

Formatted: Superscript

Deleted: Rsch

Deleted: ¶  
Marvel, K., Cook, B. I., Bonfils, C. J. W., Durack, P. J., Smerdon, J. E., and Williams, A. P.: Twentieth-century hydroclimate changes consistent with human influence, *Nature*, 569, 59-65, 2019. ¶

Mason, R. S.: Recent Survey of Coe and Eliot Glaciers, Mazama Bulletin, 36, 37-39, 1954.

1130 Menounos, B., Hugonnet, R., Shean, D., Gardner, A., Howat, I., Berthier, E., Pelto, B., Tennant, C., Shea, J., Noh, M.-J., Brun, F., and Dehecq, A.: Heterogeneous Changes in Western North American Glaciers Linked to Decadal Variability in Zonal Wind Strength, *Geophys. Res. Lett.*, 46, 200-209, 2019.

Deleted: Rsch

Miller, J. B., Frisbee, M. D., Hamilton, T. L., and Murugapiran, S. K.: Recharge from glacial meltwater is critical for alpine springs and their microbiomes, *Environ. Res. Lett.*, 16, 064012, 2021.

Deleted: Rsch

1135 Mote, P. W., Rupp, D. E., Li, S., Sharp, D. J., Otto, F., Uhe, P. F., Xiao, M., Lettenmaier, D. P., Cullen, H., and Allen, M. R.: Perspectives on the causes of exceptionally low 2015 snowpack in the western United States, *Geophys. Res. Lett.*, 43, 10980-10988, 2016.

Deleted: Rsch

1140 Neff, P. D., Steig, E. J., Clark, D. H., McConnell, J. R., Pettit, E. C., and Menounos, B.: Ice-core net snow accumulation and seasonal snow chemistry at a temperate-glacier site: Mount Waddington, southwest British Columbia, Canada, *J. Glaciol.*, 58, 1165-1175, 2012.

Nelson, L. A.: A New Glacier on Mt. Hood, Mazama Bulletin, 6, 67-68, 1924.

1145 Nolin, A. W., Philippe, J., Jefferson, A., and Lewis, S. L.: Present-day and future contributions of glacier runoff to summertime flows in a Pacific Northwest watershed: Implications for water resources, *Water Res. Res.*, 46, W12509, 2010.

Deleted: Rsch

O'Connor, J. E.: Our Vanishing Glaciers, *Oregon Historical Quarterly*, 114, 402-427, 2013.

1150 Oerlemans, J.: Extracting a Climate Signal from 169 Glacier Records, *Science*, 308, 675-677, 2005.

[O'Neil, S., McNeil, C., Sass, L. C., Florentine, C., Baker, E. H., Peitzsch, E., McGrath, D., Fountain, A. G., and Fagre, D.: Reanalysis of the US Geological Survey Benchmark Glaciers: long-term insight into climate forcing of glacier mass balance, \*J. Glaciol.\*, 65, 850-866, 2019.](#)

1155

O'Neil, D.: Changes at Elevation, 1859 *Oregon's Magazine*, 77, 60-67, 2023.

- 1165 [Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S. P., Kononov, V., Le Bris, R., Mölg, N., Nosenko, G., Nuth, C., Pope, A., Racoviteanu, A., Rastner, P., Raup, B., Scharrer, K., Steffen, S., and Winsvold, S.: On the accuracy of glacier outlines derived from remote-sensing data, \*Ann. Glaciol.\*, \*\*54\*\*, 171-182, 2013.](#)
- [Pelto, M.: Climate Driven Retreat of Mount Baker Glaciers and Changing Water Resources, \*Springer Briefs in Climate Studies\*, doi: 10.1007/978-3-319-22605-7, 2016.](#)
- 1170 Pelto, M. S.: How Unusual Was 2015 in the 1984-2015 Period of the North Cascade Glacier Annual Mass Balance?, *Water*, **10**, w10050543, 2018.
- Pelto, M. S.: Alpine Glaciers, *State of the Climate in 2022*, **2**. Global Climate, S43-S44, 2023.
- 1175 [Pelto, M., and Brown, C.: Mass balance loss of Mount Baker, Washington glaciers 1990-2010, \*Hydrol. Process.\*, \*\*26\*\*, 2601-2607, 2012](#)
- [Pelto, M. S., and Hedlund, C.: Terminus behavior and response time of North Cascade glaciers, Washington, U.S.A., \*J. Glaciol.\*, \*\*47\*\*, 497-506, 2001.](#)
- 1180 Pelto, M. S., Dryak, M., Pelto, J., Matthews, T., and Perry, L. B.: Contributions of Glacier Runoff during Heat Waves in the Nooksack River Basin USA, *Water*, **14**, w14071145, 2022.
- 1185 [Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., Kienholz, C., Miles, E. S., Moholdt, G., Mölg, N., Paul, F., Radić, V., Rastner, P., Raup, B. H., Rich, J., Sharp, M. J., and The Randolph Consortium: The Randolph Glacier Inventory: a globally complete inventory of glaciers, \*J. Glaciol.\*, \*\*60\*\*, 537-552, 2014.](#)
- 1190 Philip, S. Y., Kew, S. F., van Oldenborgh, G. J., Anslow, F. S., Seneviratne, S. I., Vautard, R., Coumou, D., Ebi, K. L., Arrighi, J., Singh, R., van Aalst, M., Marghida, C. P., Wehner, M., Yang, W., Li, S., Schumacher, D. L., Hauser, M., Bonnet, R., Luu, L. N., Lehner, F., Gillett, N., Tradowsky, J. S., Vecchi, G. A., Rodell, C., Stull, R. B., Howard, R., and Otto, F. E. L.: Rapid attribution analysis of the extraordinary heat wave on the Pacific coast of the US and Canada in June 2021, *Earth Syst. Dynam.*, **13**, 1689-1713, 2022.
- 1195 Phillips, K.: Recent Changes in Hood's Glaciers, *Mazama Bulletin*, **17** 45-50, 1935.



Phillips, K.: Our Vanishing Glaciers, Mazama Bulletin, 20, 24-41, 1938.

1200 Pitman, K. J., Moore, J. W., Sloat, M. R., Beaudreau, A. H., Bidlack, A. L., Brenner, R. E., Hood, E. W., Pess, G. R.,  
Mantua, N. J., Milner, A. M., Radić, V., Reeves, G. H., Schindler, D. E., and Whited, D. C.: Glacier Retreat and Pacific  
Salmon, *BioScience*, 70, 220-236, 2020.

1205 [Robel, A. A., Ultee, L., Ranganathan, M., and Nash, M.: For whom and by whom is glaciology?. \*J. Glaciol.\*, accepted  
manuscript, doi.org/10.1017/jog.2024.29, 2024.](#)

Roe, G. H., Baker, M. B., and Herla, F.: Centennial glacier retreat as categorical evidence of regional climate change, *Nat.  
Geosci.*, 10, 95-99, 2017.

1210 Roe, G. H., Christian, J. E., and Marzeion, B.: On the attribution of industrial-era glacier mass loss to anthropogenic climate  
change, *The Cryosphere*, 15, 1889-1905, 2021.

Rounce, D. R., Hock, R., Maussion, F., Hugonnet, R., Kochtitzky, W., Huss, M., Berthier, E., Brinkerhoff, D., Compagno,  
L., Copland, L., Farinotti, D., Menounos, B., and McNabb, R. W.: Global glacier change in the 21st century: Every increase  
in temperature matters, *Science*, 379, 78-83, 2023.

1215 Thieman, C., Danielsen, A., Salminen, E., Berge, H., and Ross, K.: Watershed 2040, Hood River Basin Partnership Strategic  
Action Plan, Oregon Watershed Enhancement Board, 2021.

1220 Thomsson, V., Kennedy-Asser, A. T., Vosper, E., Lo, Y. T. E., Huntingford, C., Andrews, O., Collins, M., Hegerl, G. C.,  
and Mitchell, D.: The 2021 western North American heat wave among the most extreme events ever recorded globally, *Sci.  
Adv.*, 8, eabm6860, 2022.

1225 U.S. Bureau of Reclamation: Reclamation Managing Water in the West: Hood River Basin Study,  
<https://www.usbr.gov/pn/studies/hoodriver/index.html>, 2015.

U.S. Geological Survey: Eruption History of Mount Hood, Oregon, [https://www.usgs.gov/volcanoes/mount-  
hood/science/eruption-history-mount-hood-oregon](https://www.usgs.gov/volcanoes/mount-hood/science/eruption-history-mount-hood-oregon), 2023.

1230 U.S. National Oceanic and Atmospheric Administration: Climate at a Glance,  
<https://www.ncei.noaa.gov/access/monitoring/climate-at-a-glance/>, 2024.

[U.S. National Resources Conservation Service: Oregon, Water, About Us, https://www.nrcs.usda.gov/conservation-basics/conservation-by-state/oregon/oregon-snow-survey/about-us, 2024.](https://www.nrcs.usda.gov/conservation-basics/conservation-by-state/oregon/oregon-snow-survey/about-us, 2024)

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

235 [World Glacier Monitoring Service: Reference glacier data, https://wgms.ch/products\\_ref\\_glaciers/, 2022.](https://wgms.ch/products_ref_glaciers/, 2022)

Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S. U., Helzle, M., Paul, F., Haeberli, W., Denzinger, F., Ahlström, A. P., Anderson, B., Bajracharya, S., Baroni, C., Braun, L. N., Cáceres, B. E., Casassa, G., Cobos, G., Dávila, L. R., Delgado Granados, H., Demuth, M. N., Espizua, L., Fischer, A., Fujita, K., Gadek, B., Ghazanfar, A., Hagen, J. O., Holmlund, P., Karimi, N., Li, Z., Pelto, M., Pitte, P., Popovnin, V. V., Portocarrero, C. A., Prinz, R., Sangewar, C. V., Severskiy, I., Sigurdsson, O., Soruco, A., Usabaliyev, R., and Vincent, C.: Historically unprecedented global glacier decline in the early 21st century, *J. Glaciol.*, 61, 745-762, 2015.

<b>Page 3: [1] Deleted</b>	<b>Anders Carlson</b>	<b>15/04/2024 12:09:00</b>
----------------------------	-----------------------	----------------------------

<b>Page 3: [2] Deleted</b>	<b>Anders Carlson</b>	<b>23/04/2024 11:34:00</b>
----------------------------	-----------------------	----------------------------

<b>Page 3: [2] Deleted</b>	<b>Anders Carlson</b>	<b>23/04/2024 11:34:00</b>
----------------------------	-----------------------	----------------------------

<b>Page 3: [3] Deleted</b>	<b>Anders Carlson</b>	<b>23/04/2024 08:02:00</b>
----------------------------	-----------------------	----------------------------

<b>Page 3: [3] Deleted</b>	<b>Anders Carlson</b>	<b>23/04/2024 08:02:00</b>
----------------------------	-----------------------	----------------------------

<b>Page 11: [4] Deleted</b>	<b>Anders Carlson</b>	<b>17/04/2024 14:21:00</b>
-----------------------------	-----------------------	----------------------------

<b>Page 11: [5] Deleted</b>	<b>Anders Carlson</b>	<b>17/04/2024 14:33:00</b>
-----------------------------	-----------------------	----------------------------

<b>Page 11: [6] Deleted</b>	<b>Anders Carlson</b>	<b>19/04/2024 05:32:00</b>
-----------------------------	-----------------------	----------------------------

<b>Page 13: [7] Deleted</b>	<b>Anders Carlson</b>	<b>17/04/2024 14:35:00</b>
-----------------------------	-----------------------	----------------------------

<b>Page 15: [8] Deleted</b>	<b>Anders Carlson</b>	<b>15/04/2024 12:13:00</b>
-----------------------------	-----------------------	----------------------------

<b>Page 15: [9] Deleted</b>	<b>Anders Carlson</b>	<b>12/04/2024 08:56:00</b>
-----------------------------	-----------------------	----------------------------

<b>Page 15: [9] Deleted</b>	<b>Anders Carlson</b>	<b>12/04/2024 08:56:00</b>
-----------------------------	-----------------------	----------------------------

<b>Page 15: [10] Formatted</b>	<b>Anders Carlson</b>	<b>20/04/2024 03:01:00</b>
--------------------------------	-----------------------	----------------------------

Font: Not Bold

▲  
**Page 15: [10] Formatted**

**Anders Carlson**

**20/04/2024 03:01:00**

Font: Not Bold

▲  
**Page 16: [11] Deleted**

**Anders Carlson**

**07/05/2024 08:25:00**

▼

←

▲