

Unprecedented Twenty-First Century Glacier Loss on Mt. Hood, Oregon, U.S.A.

Nicolas Bakken-French¹, Stephen J. Boyer¹, B. Clay Southworth¹, Megan Thayne¹, Dylan H. Rood^{2*}, and Anders E. Carlson¹

¹Oregon Glaciers Institute, Corvallis, OR 97330, U.S.A.

²Department of Earth Science and Engineering, Imperial College London, London SW7 2AZ, U.K.

Correspondence to: Dylan H. Rood (d.rood@imperial.ac.uk)

Abstract. As part of the southern Cascades, Mt. Hood is the tallest and most glacierized peak in Oregon, U.S.A. Despite alpine glaciers being one of the clearest indicators of human-caused climate change, the 21st century behavior of glaciers on Mt. Hood has not been directly documented at the ground level. Here we directly measure changes in the extents of Mt. Hood's glaciers from 2003 to 2023 and find dramatic retreat of all glaciers, with one glacier ~~ceasing to flow (joining another glacier that ceased to flow before 2003)~~ and another ~~three glaciers~~ retreating towards this status. By 2023, Mt. Hood glaciers lost ~~~17%~~ of their 2015/2016 area and ~~~39%~~ of their 1981 area. ~~The rate of area loss from 2015/2016 to 2023 (~2.10 % year⁻¹) was ~2.6 times faster than the rate from 1981 to 2015/2016 (~0.81 % year⁻¹).~~ The seven largest glaciers on the volcano lost ~~~25%~~ of their area between 2000 and 2023. Comparison to historic records of glacier area back to 1907 shows that this 21st-century retreat is unprecedented relative to the previous century. The rate of area loss over the last 23 years (~~~1.07 % year⁻¹~~) ~~was ~1.9 times faster than the~~ fastest rate documented in the last century from 1907 to 1946 (~~~0.56 % year⁻¹~~) and ~~~3.5 times~~ faster than the 20th century average (~~~0.31 % year⁻¹~~). ~~This unprecedented rate of retreat corresponds with regional summer warmth reaching 1.7-1.8°C (2013-2023 average) relative to the~~ early 1900s, but not with regional changes in winter precipitation. We conclude that Mt. Hood's glaciers are retreating in response to a warming climate and that this recession has accelerated in the 21st century.

1 Introduction

The global loss of mass and retreat of glaciers in the 21st century is accelerating (Zemp et al., 2015; Hugonnet et al., 2021) with century-scale glacier recession attributed to regional climate change in response to anthropogenic greenhouse gas emissions (Marzeion et al., 2014; Roe et al., 2017, 2021). The 2021-2022 hydrological year was the 35th consecutive year of negative annual mass balance for 50 reference glaciers on six continents (Pelto, 2023; World Glacier Monitoring Service, 2022). If global average temperature increases by 1.5°C relative to the pre-industrial period, the limit set forth in the Paris Agreement, roughly half of the world's individual glaciers are projected to disappear with ~26% loss in global glacier mass (Rounce et al., 2023). Indeed, in the western United States, this deglaciation is already coming to fruition (Fig. 1A). The Trinity Alps in the State of California (Fig. 1A) have lost ~97% of their late 1800s glacierized area, with only ~0.017 km² of one glacier still showing evidence of ice flow as of 2015 (Garwood et al., 2020). The Olympic Mountains of Washington

- Deleted:** actively flowing
- Deleted:** disappearing
- Deleted:** four
- Deleted:** 18
- Deleted:** 44
- Deleted:** 40
- Deleted:** and has outpaced modeled glacier changes of 21st century retreat
- Deleted:** 1.7-1.8
- Deleted:** is
- Deleted:** about
- Deleted:** triple the
- Deleted:** 4-0.58
- Deleted:** about six
- Deleted:** 0.29-0.33
- Deleted:** It has also equaled or exceeded by up to 67% the highest simulated rates of glacier recession under a high greenhouse gas emissions scenario.
- Deleted:** We demonstrate that this century-scale retreat strongly correlates with regional 30-year-mean climate warming of 1.1-1.2°C and 15-year-mean climate warming of 1.3-1.5°C since

State (Fig. 1A) lost about half of their late 1800s glacier area by 2015 with 35 glaciers disappearing since 1980 (Fountain et al., 2022). Eleven of the 37 named glaciers in Glacier National Park in the State of Montana (Fig. 1A) ceased to be classified as **actively flowing** glaciers as of 2015 with glacierized area in the park declining by ~68% from the 1800s to 2015 (Fagre et al., 2017). Washington State's Mt. Rainier (Fig. 1A), the most glacierized peak in the contiguous United States, has seen its glacierized area decline by ~42% between 1896 and 2021 and had one glacier recently disappear with two more nearing stagnation (Beason et al., 2023). Nearby in the Washington North Cascades (Fig. 1A), 33 glaciers were listed as extinct in the Global Land Ice Measurements from Space (GLIMS) Glacier Database at the National Snow and Ice Data Center (NSIDC) as of 2023 (GLIMS and NSIDC, 2023).




Figure 1: Location map of Oregon (A) and Mt. Hood (B). A. Western United States with mountain ranges mentioned denoted (WA = Washington, MT = Montana, OR = Oregon, CA = California, ID = Idaho, NV = Nevada, UT = Utah, AZ = Arizona); inset shows location with black outline. B. Glacier extents on Mt. Hood with photo locations in Fig. 3 and 6 noted. 1907/1901 and 2004 ice extents from Jackson & Fountain (2007); 2015/2016 ice extents from Fountain et al. (2023). 2023 extents from this study. Inset shows figure location (black outline) within Hood River County (area for reanalysis data in Fig. 7). Credits: Esri, U.S. Census Bureau, Earthstar Geographics, Maxar.

Deleted: 6

70 Despite these ominous predictions and dire observations for glaciers, data on the 21st century behavior of glaciers in the Cascade Range of Oregon are limited. Glaciers in Oregon were first noted in 1870 on Mt. Hood, the tallest and most glacierized peak in Oregon (Fig. 1A, B), only days after the first glaciers in the contiguous United States were recorded on Mt. Shasta in California (O'Connor, 2013). Mt. Hood is an active volcano that reached a height close to its current elevation of 3,425 meters above sea level by 30,000 years ago (U.S.G.S., 2023, <https://www.usgs.gov/volcanoes/mount-hood/science/eruption-history-mount-hood-oregon>). The last major eruption was ~1,500 years ago that created pyroclastic

75 flows and lahars on the southwest side of the volcano upon which Zigzag and Palmer glaciers rested (Fig. 1B). Its last eruption occurred in the late 1700s, producing a lava dome south of the summit around which Coalman Glacier flows (Fig. 1B). In the late 1800s, fumaroles formed near this lava dome (Lillquist and Walker, 2006). Mt. Hood has high amounts of rock fall and variations in geothermal heat that could affect ablation, ice flow, and ice extent (e.g., Lundstrom et al., 1993; Lillquist and Walker, 2006; Jackson and Fountain, 2007; Howcutt et al., 2023).

80 Changes in Mt. Hood's glaciers were monitored by the Research Committee of the Mazama Mountaineering Club from the 1930s to mid-1980s (e.g., Phillips, 1935, 1938; Mason, 1954; Handwith Jr., 1959; Dodge, 1971, 1987). In 1981, the U.S. Geological Survey measured the thickness and area of Mt. Hood's glaciers, estimating their volume (Driedger and Kennard, 1986a, 1986b). However, the extent of Oregon Cascade glaciers on current U.S. Geological Survey and U.S. Forest Service maps are still based on air photos from the 1950s (Fountain et al., 2017). The glacierized area of Mt. Hood was last measured

85 with field observations in 2004 (Jackson and Fountain, 2007). Fountain et al. (2023) documented snow and ice areas in the western United States using 2015 and 2016 imagery for Mt. Hood's glaciers but did not test mapped accuracy with field observations. The timing of this imagery also missed what multi-year and longer impacts the 2015 record-low snowpack (Mote et al., 2016) and the June 2021 Pacific Northwest heatwave (Philip et al., 2022; Thompson et al., 2022) would have on Mt. Hood's glaciers (e.g., Pelto, 2018; Pelto et al., 2022).

90 Regarding climate change impacting Oregon's glaciers, Menounos et al. (2019) documented minimal elevation change for Mt. Hood's glaciers from 2000 to 2018, interpreting from this an approximately neutral mass balance for the first 19 years of the 21st century. Near zero mass balance for 2000-2018 is at odds with the mass changes for global reference glaciers that include four glaciers to the north of Mt. Hood in the state of Washington; these four all had cumulative negative mass balances for the same time period (Pelto, 2023). For the 20th century, Lillquist and Walker (2006) did not find a relationship

95 between glacier-length change on Mt. Hood and temperature or precipitation change over the 20th century. Conversely, climate-glacier modeling simulates the emergence of glacier retreat from natural variability in western North America since the 1960s (Marzeion et al., 2014). As such, recent impacts of human-caused climate change on Mt. Hood's glaciers have been unclear and largely unexplored from direct field observations.

100 Glacier change on Mt. Hood is of importance to downstream ecosystems and economies (see Robel et al., 2024), such as orchards and farms in the Hood River Valley (Nolin et al., 2010; U.S. Bureau of Reclamation, 2015; Frans et al., 2016,

Deleted: (now gone)

Deleted: whereas glacier lengths with field observations were last recorded in 2000 (Lillquist and Walker, 2006). Fountain et al. (2023)

2018), unique glacier-fed spring water microbiomes (Miller et al., 2021), and salmon and steelhead (Pitman et al., 2020; Thieman et al., 2021). The Hood River Valley has ~~50 km²~~ producing seven varieties of pears, ~~6 km²~~ producing five varieties of apples, ~~6 km²~~ devoted to cherries, and ~~0.4 km²~~ of grapes for an expanding wine industry (Columbia Gorge Fruit Growers, 2024, <http://www.cgfg.org/information/crops-information>), along with two hydropower plants with 4.4 megawatt capacity (Farmers Irrigation District, 2024, <https://www.fidhr.org/index.php/en/about-us/hydroelectric>). Glaciers contribute ~40-70% of the stream flow in the late summer to the upper Hood River where flow is diverted for irrigation and local hydropower (Nolin et al., 2010). As such, future water resource management plans for the Hood River basin included projections of future glacier change on Mt. Hood (U.S. Bureau of Reclamation, 2015; Thieman et al., 2021). White River and Newton-Clark glaciers also contribute late-summer meltwater to a robust orchard industry in another county (Columbia Gorge Fruit Growers, 2024, <http://www.cgfg.org/information/crops-information>). Furthermore, Mt. Hood is ~~one of the most~~ climbed mountaineering objectives in the world (~10,000 attempts per year), which is becoming more dangerous during a diminishing climbing season with greater rockfall as the volcano has changed in recent years (O'Neil, 2023).

Here we document glacier change on Mt. Hood over the last 20 years. We conducted new repeat photography and field observations of glacier extent in 2003 and 2023 to provide detailed ground-based evidence for glacier change during a period where satellite-based altimetry previously found no significant change in these glaciers. We also utilized a series of field campaigns with satellite imagery to determine the magnitude of glacier area change up to mid-September 2023. We place these 21st-century glacier changes in the context of 20th-century glacier ~~and regional climate~~ changes.

2 Data and Methods

2.1 2003 and 2023 Flowing-Ice-Margin Positions and Repeat Photography

In 2003 (28 August to 4 October), the lowest observed actively flowing glacier ~~locations~~ and stagnant-ice positions were measured (latitude-longitude-altitude) by global positioning system (GPS) for every glacier on Mt. Hood (Fig. S1). These were discrete observations, which served as points to assess future minimum changes in elevation. Repeat measurements and comparison to known elevation markers determined an elevation uncertainty of ± 6 m. Termini were photographed with locations noted (Fig. S1) for future repeat photography (i.e., 2023). All glaciers were described in detail in the field (Supplement). The goal of this initial 2003 field effort was to provide baseline observations and archival photographic evidence from the early 21st century against which future glacier change could be assessed. ~~We conducted further~~ observations for this assessment 20 years later in 2023. ~~From 15 to 22 September 2023,~~ repeat photographs were taken at the same locations as the 2003 photographs (Fig. S1) and new field observations were made for all of the glaciers (Supplement). ~~We selected photographs to repeat from an archive of photographs taken in 2003. Using the image and field notes for the~~ location of the 2003 images, we located the precise site of each selected 2003 photograph. The same frame was maintained using relatively unchanged features such as buttresses, cliffs, and moraines as points of reference.

Deleted: 12,000

Deleted: acres

Deleted: 1500

Deleted: acres

Formatted: Superscript

Formatted: Superscript

Deleted: 1500

Deleted: acres

Deleted: 100

Deleted: acres

Formatted: Superscript

Formatted: Superscript

Deleted: also

Deleted: and length

Deleted: and test the relationship between ~120 years of glacier-length variability and regional climate change...

Deleted: Frontal GPS

Deleted: -terminus

Deleted: positions

Formatted: No widow/orphan control, Don't keep with next

Deleted: terminus

Deleted: ; w

Deleted: subsequently

Deleted: ¶

Deleted: From this repeat mapping (see below), we were able to determine the minimum amount the terminus rose in elevation over the last 20 years. We selected photographs to repeat from an archive

Deleted: regarding

Deleted: from 2003

160 From 2020 to 2023, the full extent of the actively flowing parts of glaciers were mapped to determine the degree to which
the lowest flowing ice had risen in elevation, which is related to the surface mass balance of the glacier (e.g., Haeberli and
165 Hoelzle, 1995; Hoelzle et al., 2023). Because manual mapping is suggested for glaciers with debris cover (e.g., Paul et al.,
2013), we initially manually mapped actively flowing glacier extents using composite Google Earth imagery in the
application CalTopo (caltopo.com). As these images spanned multiple years and were not necessarily recent, we then
170 manually refined ice margins using weekly Sentinel-2 satellite images that were taken shortly before the first snowfall from
the summers of 2020, 2021, 2022, and 2023. These images were processed and made available in CalTopo one day after
acquisition (see Supplement for a complete list of satellite imagery). The combination of satellite-imagery delineation of
margins with ground-truth mapping relative to the images is a robust, effective technique. In October 2020, September 2021,
and September 2023, we iteratively took these ice margins (updated year by year) into the field for direct mapping using
CalTopo on a portable GPS enabled tablet. We finalized our ice limits with the ESRI World Imagery Basemap from
September 2023.

175 These three years were excellent for field mapping as the length of the 2020 summer and the warmth of the 2021 and 2023
summers resulted in extremely low snow cover that facilitated ice-margin identification in regions that would typically have
snow cover throughout the summer. All termini in the end-of-summer ablation area were walked, resulting in ~60%, on
average, of each ice margin being physically mapped in the field. Notably, this means all debris-covered ice margins were
180 mapped in the field and their delineation was not based solely on satellite imagery. Debris-covered termini are common on
Mt. Hood glaciers, and such field mapping of the termini is critical to accurately delineate glacier margins (e.g., Lundstrom
et al., 1993; Lillquist and Walker, 2006; Jackson and Fountain, 2007; Ellinger, 2010). We followed prior methods for
differentiating actively flowing debris-covered ice from stagnant debris-covered ice on Mt. Hood (e.g., Lillquist and Walker,
2006): an identifiable glacier front, a lobate form of the terminus, and a visible connection of the lobate terminus to debris-
185 free flowing ice. This methodology is consistent with the observation of a convex shape of actively flowing glacier margins
by Leonard and Fountain (2003) and Leigh et al. (2019). To these criteria, we added the presence of crevasses observed in
the field as another indicator of ice flow whether under debris or in bare ice (Leigh et al., 2019). Using these criteria, we
determined and walked this transition from actively flowing to stagnant ice, mapping this demarcation in the field.

2.2 2023 Glacier Inventory and Uncertainty

185 To determine the current glacier extent and compare to past inventories to assess glacier area change, we produced a 2023
glacier inventory for the 12 glaciers of Mt. Hood (Fig. 1B) following UNESCO guidelines that require inclusion of inactive
or stagnant ice (Müller et al., 1977; Cogley et al., 2011). Specifically, we expanded our actively flowing ice areas discussed
above to include stagnant ice (debris-free and debris-covered). We also included separate stagnant-ice bodies and perennial
snowfields that used to be connected to a glacier. Following standard practices (Paul et al., 2013; Leigh et al., 2019), our
190 2023 ice limits were mapped on new high-resolution (0.3 m) Maxar (Vivid) imagery captured on 8 September 2023

Deleted: In contrast to the point measurements in 2003,

Deleted: extent

Deleted: were mapped in 2023, using an iterative approach over the years 2020 to 2023 to determine the deg

Deleted: new

Deleted: Earth imagery

Deleted: . Note that we mapped only the extent of actively flowing glacier ice, both debris-free and debris-covered, and excluded stagnated non-flowing debris-free and debris-covered ice (see below for criteria for differentiation and delineation of flowing from stagnant glacier ice). We focused on actively flowing glacier ice, rather than include stagnant ice, to be consistent with prior glacier-extent mapping on Mt. Hood (e.g., Lillquist and Walker, 2006; Jackson and Fountain, 2007; Fountain et al., 2023) and in other mountain ranges of the western United States (e.g., Fagre et al., 2017; Garwood et al., 2020; Fountain et al., 2022; Beason et al., 2023).

Deleted: ¶

Deleted: , Mapping

Formatted: Normal, Line spacing: single, No widow/orphan control

Deleted: E

210 available through the ESRI World Imagery Basemap
(https://services.arcgisonline.com/ArcGIS/rest/services/World_Imagery/MapServer; Fig. 1B) guided by our field
215 observations from 2020, 2021 and 2023.

Manual remote-sensed ice-limit comparisons with direct field observations yielded extent differences that were small enough
(i.e., less than a meter or two) to make uncertainty difficult to quantify. Using only remote sensing, Fountain et al. (2023)
215 had their lowest ice-extent uncertainty for a Mt. Hood glacier at $\pm 1\%$, while Jackson and Fountain (2007) reported an
uncertainty for their 2000 Mt. Hood glacier areas of $\pm 0\%$. Our uncertainty is most likely lower than $\pm 1\%$, but to be
conservative we assumed a $\pm 7\%$ uncertainty in the glacier area for 2023. We based this $\pm 7\%$ uncertainty on the average area
uncertainty for the 2004 extents of Mt. Hood glaciers determined by Jackson and Fountain (2007), which was $\pm 7\%$. Our
220 conservative $\pm 7\%$ is slightly greater than the uncertainties determined from testing the accuracy of remote sensed glacier
area from the European Alps and Alaska of 2.6-5.7% (Paul et al., 2013). This means any conclusions we made from our
mapped glacier area were likely not dependent on our assumed glacier area uncertainty.

2.3 Glacier Area Change and Uncertainty

We compared our 2023 glacier inventory to the most recent inventories conducted for Mt. Hood. Fountain et al. (2023)
225 produced a Mt. Hood glacier inventory from remote sensing imagery from either 2015 or 2016; this allowed an assessment
of glacier-area change over the last seven to eight years. We also compared our 2023 glacier areas to the 1981 glacier
inventory for Mt. Hood determined by Driedger & Kennard (1986b) that was based on field mapping and aerial photographs;
this afforded a 42-year perspective on glacier change.

We assessed rates of glacier change by comparing our 2023-measured inventory area with the glacier area records of Jackson
and Fountain (2007) for the time period 1907 (or 1901 in the case of Eliot Glacier) to 2004 for seven glaciers on Mt. Hood:
230 White River, Newton-Clark, Eliot, Coe, Ladd, Sandy, and Reid (Fig. 1B). Because of the different temporal resolutions of
each glacier-area record, we calculated area change between observational years common to all seven glaciers: 1907, 1946,
1972, 2000, and 2023. We focused on the changes from 2000 to 2023, rather than 2000 to 2004 and then 2004 to 2023, to
make the duration over which the most recent intervals of change were measured closer to the earlier temporal spacing.
Newton-Clark Glacier lacked an aerial extent for 1946; we extrapolated between the measured extent in 1935 and 1956 when
235 the glacier lost $\sim 0.04 \text{ km}^2$ over these 21 years. To aid comparison between glaciers of different sizes on Mt. Hood and to
other glaciers in the Pacific Northwest, we calculated relative changes in glacier area determined as a percent of its prior
area. In calculating the rate of change, we propagated through the uncertainty in the area from Jackson and Fountain (2007)
and our 2023 areas. For the 2000 area, we conservatively applied a $\pm 7\%$ uncertainty of the 2004 Jackson and Fountain
(2007) glacier margins. The propagation of uncertainty was through quadrature where the squares of the individual percent
240 uncertainties were summed, and their square root taken.

Deleted: imagery

Deleted: from September 2023

Moved up [1]: In contrast to the point measurements in 2003, full glacier extents were mapped in 2023, using an iterative approach over the years 2020 to 2023. Note that we mapped only the extent of actively flowing glacier ice, both debris-free and debris-covered, and excluded stagnated non-flowing debris-free and debris-covered ice (see below for criteria for differentiation and delineation of flowing from stagnant glacier ice). We focused on actively flowing glacier ice, rather than include stagnant ice, to be consistent with prior glacier-extent mapping on Mt. Hood (e.g., Lillquist and Walker, 2006; Jackson and Fountain, 2007; Fountain et al., 2023) and in other mountain ranges of the western United States (e.g., Fagre et al., 2017; Garwood et al., 2020; Fountain et al., 2022; Beason et al., 2023). [1]

Deleted: In measuring glacier dimensions, it is important to be clear on what is considered a glacier and thus included in the glacier extents. Clarke (1987) noted in his review of glacier research since the 1820s that “the most interesting property of glaciers is that they flow”, using the present tense. The U.S. Geological Survey defines a glacier as “a large, perennial accumulation of crystalline ice, snow, rock, sediment, and often water that originates on land and moves down slope under the influence of its own weight and gravity” (U.S. Geological Survey, 2024, <https://www.usgs.gov/faqs/what-glacier>). The NSIDC, which hosts data for GLIMS, defines a glacier as “an accumulation of ice and snow that slowly flows over land” (NSIDC, 2024b, <https://nsidc.org/learn/parts-cryosphere/glaciers/glacier-quick-facts>). However, the NSIDC has another definition of a glacier as: “a mass of ice that originates on land, usually having an area larger than one tenth of a square kilometer; many believe that a glacier must show some type of movement; others believe that a glacier can show evidence of past or present movement” (NSIDC, 2024a, <https://nsidc.org/learn/cryosphere-glossary/glacier>). Therefore, our mapping of only actively flowing glacier ice follows many traditional definitions of what constitutes a glacier, but we recognize that another definition of a glacier exists that includes ice that used to move and is now stagnant. We return to our decision to map only actively flowing glacier ice, rather than include stagnant ice, in the Discussion section. [1]
Because manual mapping is suggested for glaciers with debris cover (e.g., Paul et al., 2013), we initially manually mapped actively flowing glacier extents using composite Google Earth imagery in the application CalTopo (caltopo.com). As these images spanned multiple years and were not necessarily recent, we then manually refined ice margins using weekly Sentinel-2 satellite images that were taken shortly before the first snowfall from the summers (... [1]

Deleted: To update glacier change on Mt. Hood for the 21st century, w...

Deleted: extents

Deleted: areas

Deleted: Note that Fountain et al. (2023) also mapped snowfields (e.g., stagnant ice bodies) and debris-covered stagnant ice bodies in addition to actively flowing glaciers. When determining glacier change, we looked at changes in flowing glacier area for the ice bodies contiguous with those remaining actively flowing in 20 (... [2]

Deleted: areas

Deleted: were

Deleted: of actively flowing glacier ice

380 2.4 Changes in Climate

For records of regional climate change to use in assessing potential climatic causes of glacier change, we used U.S. National Oceanic and Atmospheric Administration (NOAA) temperature and precipitation reanalysis data for the region of Hood River County (Fig. 1B) from 1895 to 2023 (U.S. NOAA, 2024, <https://www.ncei.noaa.gov/access/monitoring/climate-at-a-glance/>). We chose Hood River County as this is the smallest geographic region for the reanalysis data that includes the majority of Mt. Hood and its glaciers (eight of the 12 glaciers with the other four adjacent to the boundary, Fig. 1B) and would still have direct weather station measurements within the geographic domain over the period of comparison. Snow telemetry stations on and around Mt. Hood were only installed in the late 1970s, with temperature sensors beginning to work consistently in the late 1980s (U.S. National Resources Conservation Service, 2024, <https://www.nrcs.usda.gov/conservation-basics/conservation-by-state/oregon/oregon-snow-survey/about-us>).

390 For temperature, we focused on changes in May-October and June-September means. May through October is generally the period of time in the Oregon Cascades from peak winter snowpack to the first snowfall of the next winter, or the maximum length of the ablation season. Since this can vary, we also examined June-September temperature as a shorter ablation season. Pelto (2018) determined the best fit between mass balance and ablation temperature to be May-September or June-September for glaciers in Washington. For precipitation, we analyzed November-April and October-May means. November through April in Oregon generally spans the time period from the first winter snow until peak winter snowpack, or the accumulation season. To account for seasonal variability, we also used October through May precipitation. We calculated the changes in temperature and precipitation relative to their 1895-1924 means. Glaciers in the North Cascades of Washington have 20-30 year response times (Pelto and Hedlund, 2001) while glaciers on the Mt. Baker volcano in Washington (a closer analogue to Mt. Hood) have faster response times down to 11 years (Pelto, 2016) (Fig. 1A). As such, we applied an 11-year running mean to the temperature and precipitation records.

3 Results

Since 2003, the elevation of the lowest actively flowing ice rose for every glacier on Mt. Hood (Table 1, Fig. 2, S1), with an average rise of ~150 m (observed changes and repeat photographs provided in Supplement). Note that these are inherently minimum estimates of the rises given the point-nature of the 2003 observations. There does not appear to be any pattern in the degree of rise relative to glacier aspect on Mt. Hood (Fig. 2). The greatest rise occurred in the southeast-facing Clark drainage of Newton-Clark Glacier (>300 m) followed by the southwest-facing Zigzag Glacier (>250 m; Fig. 3A, B), the west-facing Sandy Glacier (>240 m; Fig. 3E, F) and the northwest-facing Ladd Glacier (>230 m). Of the three glaciers with substantial debris cover of their termini, Ladd (>230 m) and Coe (>170 m) (Fig. 3C, D) had rises greater than the average while Eliot's rose >70 m (Table 1, Fig. 2).

Deleted: in Glacier Length and

Deleted: Glacier length changes are commonly used to assess glacier relationships to a changing climate (e.g., Harper, 1993; Haeblerli and Hoelzle, 1995; Pelto and Hedlund, 2001; Oerlemans, 2005; Lillquist and Walker, 2006; Leclercq and Oerlemans, 2012; Leclercq et al., 2014; Pelto, 2016; Roe et al., 2016, 2021; Huston et al., 2021; Hoelzle et al., 2023). Here we compared the cumulative change in glacier length for five glaciers on Mt. Hood from 1901 through 2023 to records of regional climate change. Lillquist and Walker (2006) measured glacier-length change for White River, Newton-Clark, Eliot, Coe, and Ladd glaciers from 1901 to 2000. To this record we added the change in length up to 2023. Specifically, we calculated the distance between the lowest observed actively flowing ice in 2023 (Fig. 1B) and the lowest actively flowing ice in 2000 of Lillquist and Walker (2006) along the same flow line they used back to 1901, which was down the center of the glacier. In the case of Newton-Clark Glacier, the length was measured in the northern, due-east-facing Newton drainage (Fig. 1B). Changes in glacier length were normalized to the 1901 length of the glacier to account for different glacier sizes. Lillquist and Walker (2006) did not provide uncertainties in their glacier lengths and reported change at 1 m accuracy. Here, we report our length changes at 10 m accuracy.

Deleted: these five glaciers

Deleted: annual,

Deleted: annual,

Deleted: First, w

Deleted: . Second, we

Deleted: produced a 30-year backward-running mean for the yearly changes in temperature and precipitation. We employed a 30-year backward-running mean because the response time of Pacific Northwest glaciers similar to those on Mt. Hood is 20-30 years (...)

Deleted: The prior 30-year mean reflects this lagged relationship between climate change and glacier change. We explored the lower limit of the glacier-response times by using a 15-year backward-running mean. While Pelto and Hedlund (2001) found a lower limit to glacier-response time of 20 years, glaciers on Mt. Baker in Washington (Fig. 1A) may respond faster at 11-20 years (Pelto, 2016), which we encompass with our 15-year backward-running mean. Third, the relationship between the normalized change (... [3])

Formatted: Don't keep with next

Deleted: underwent a rise in terminus elevation for its actively flowing ice (Table 1, Fig. 2

Formatted: No widow/orphan control

Deleted: terminus

Deleted: terminus

Deleted: 3G

Deleted: H

Deleted: ; Fig. 3E, F

Deleted: terminus

Deleted: terminus

Glacier (Multiple Termini)	2003 Lowest Flowing Ice (m asl)	2023 Lowest Flowing Ice (m asl)	20-Year Flowing Ice Δ Elevation (m)	1981 Area km ²	2015/16 Area km ² (uncert)	2023 Area km ² (uncert)	1981-2023 % Δ Area	2015/16-2023 % Δ Area (uncert)
Zigzag	2350	2600	250	0.77	0.351 (0.047)	0.239 (0.017)	69	40 (3)
Palmer	gone			0.13	0.096 (0.024)	0.066 (0.005)	49	
Coalman (E)	3130	3130	0	0.08	0.099 (0.001)	0.052 (0.004)	35	48 (3)
Coalman (W)	3190	3230	40					
White River	2130	2240	110	0.54	0.335 (0.048)	0.257 (0.018)	52	23 (4)
Newton-Clark (N)	2420	2480	60	1.99	1.280 (0.185)	1.050 (0.073)	47	18 (3)
Newton-Clark (S)	2320	2620	300					
Eliot	2050	2120	70	1.68	1.538 (0.174)	1.456 (0.102)	13	5 (1)
Langille		2520		0.40	0.316 (0.040)	0.169 (0.012)	58	47 (7)
Langille (upper)	2070	gone						
Langille (lower)	1960	gone						
Coe	1920	2090	170	1.24	1.045 (0.125)	0.906 (0.063)	27	13 (2)
Ladd	2100	2330	230	0.90	*0.474 (0.056)	0.550 (0.038)	39	
Glisan (upper)	2040	gone			0.082 (0.014)	0.009 (0.001)		89 (17)
Glisan (lower)	1930	gone	220					
Sandy	1890	2130	240	1.19	0.798 (0.121)	0.719 (0.050)	40	10 (2)
Reid	2240	2350	110	0.75	0.469 (0.005)	0.416 (0.029)	45	11 (1)
Total				9.67	6.88 (0.32)	5.85 (0.16)	40	17 (1)

Table 1: Mt. Hood glacier observations from 2003 and 2023. Change (Δ) in elevation for lowest flowing ice, total glacier area in 1981 (Driedger and Kennard, 1986b), 2015/2016 (Fountain et al., 2023) and 2023, and total area change (Δ) as a percent (%) up to 2023 provided. The asterisk notes the 2015/2016 area for Ladd Glacier which excluded buried ice that remained in 2023.




Figure 2: Minimum change (Δ) in the lowest elevation of actively flowing ice from 2003 to 2023 relative to glacier position on Mt. Hood; C = Coalman, WR = White River, NC = Newton-Clark with its two termini labeled, and E = Eliot.

Deleted: actively

Deleted: full

Deleted: and

Formatted: Caption, No widow/orphan control




Figure 3: Select repeat photographs from Mt. Hood; see Fig. 1B for photo locations. A and B: Zigzag Glacier across the valley (shot location 45.29275°, -121.80044°) repeat photograph from 2003 (A) and 2023 (B). 1. Denotes the reduction in area of Zigzag Glacier to its 2023 **actively flowing limit** (dashed white line). 2. Remnant stagnant ice that used to be part of the actively flowing glacier. 3. Western terminus of Coalman Glacier that has retreated but is still visible in 2023 at edge of cloud. 4. Top of White River Glacier that has retracted since 2003. C and D: Langille and Coe Glaciers (shot location 45.39459°, -121.69982°) in 2003 (C) and 2023 (D). 5. **Near-complete disappearance** of remnant ice of Langille Glacier. 6. Loss of an ice fall on Coe Glacier due to terminus retreat to just below the 6 in D. E and F: Sandy Glacier (shot location 45.39477°, -121.73124°) in 2003 (E) and 2023 (F). 7. Accumulation of rock fall on the glacier in the last 20 years. 8. Retreat of glacier ice that is now exposed bedrock. 9. 2023 terminus location and ice caves (F) that were in the **center** of the glacier in 2003 (E). 10. 2003 terminus with ice cave (E) that is now stagnant, debris-covered ice (F).

Deleted: In the last seven to eight years, every glacier on Mt. Hood lost area (Fig. 1B), resulting in 18±1% loss in total glacier area (Table 1) relative to 2015/2016 at a rate of 2.40±0.02 % yr⁻¹ (yr = year). Glisan Glacier ceased to show evidence of flow between 2015 (Fountain et al., 2023) and our first field observation in 2020, with a stagnant ice area of 0.009±0.0006 km² as of 2023. Zigzag, Coalman and Langille glaciers lost >40% of their area since 2015/2016. Ladd Glacier separated from its uppermost accumulation area (Fig. 3E, F) while Coe Glacier separated from its debris-covered terminus (Fig. 3C, D), both due to thinning over ice falls after 2016 (Fountain et al., 2023). A similar separation from its debris-covered terminus also occurred for Eliot Glacier (Fountain et al., 2023). When compared to the 1981 glacier inventory (Driedger and Kennard, 1986b), the glacierized area loss by 2023 for Mt. Hood glaciers was ~44% (Table 1) at a rate of ~1.0 % yr⁻¹. The largest percent area lost since 1981 was the disappearance of south-facing Palmer Glacier (100%), followed by southwest-facing Zigzag Glacier (~87%), and north-facing Ladd Glacier (~74%).

Page Break

... [4]

Deleted: Disappearance

Deleted: Ladd Glacier (shot location 45.39718°, -121.70424°) in 2003 (E) and 2023 (F). 7. Separation of Ladd from its uppermost accumulation area. 8. Ice fall that is now the current termi... [5]

Deleted: G

Deleted: H

Deleted: 10

Deleted: 11

Deleted: 12

Deleted: H

Deleted: centre

Deleted: G

Deleted: 13

Deleted: G

Deleted: H

In the last seven to eight years, every glacier on Mt. Hood lost area (Fig. 1B), resulting in 17±1% loss in glacier area (Table 1) relative to 2015/2016 at 2.27±0.13 % yr⁻¹ (yr = year). This area loss and rate do not include the area changes of Ladd Glacier. While Fountain et al. (2023) mapped Ladd's 2015/2016 area, we found that they excluded a region of buried ice that still existed in 2023, with our 2023 area (0.550±0.038 km²) being greater than the 2015/2016 area (0.474±0.056 km²). Nevertheless, our 2023 area was smaller than the 2004 area (0.67±0.05 km²) of Jackson and Fountain (2007). Glisan Glacier ceased to show evidence of flow between 2015 (Fountain et al., 2023) and our first field observation in 2020, with a stagnant ice area of 0.009±0.0006 km² as of 2023 (Table 1). Zigzag, Coalman and Langille glaciers lost >40% of their area since 2015/2016. Ladd Glacier separated from its uppermost accumulation area (Fig. 3E, F) while Coe Glacier almost completely separated from its debris-covered terminus (Fig. 3C, D), both due to thinning over ice falls after 2016 (Fountain et al., 2023). A similar separation from its debris-covered terminus is occurring for Eliot Glacier (Fountain et al., 2023).

Relative to the 1981 glacier inventory (Driedger and Kennard, 1986b), the glacierized area of Mt. Hood decreased by ~39% by 2023 (Table 1) at a rate of ~0.93 % yr⁻¹. This area change and rate do not include Glisan Glacier as this glacier's area was not included in the 1981 inventory. The largest relative area lost since 1981 was the southwest-facing Zigzag Glacier (~69%), followed by northeast-facing Langille Glacier (~58%), and south-facing White River Glacier (~52%). For the ten glaciers with consistent 1981, 2015/2016 and 2023 areas, which thus excludes Ladd and Glisan, the rate of area change from 1981 to 2015/2016 was 0.81±0.04 % yr⁻¹. From 2015/2016 to 2023, the rate of area loss was 2.10±0.06 % yr⁻¹. Accordingly, glacier retreat accelerated by a factor of ~2.6 between the two periods.

The seven largest glaciers on Mt. Hood (White River, Newton-Clark, Eliot, Coe, Ladd, Sandy, and Reid) that have area records extending back to 1907 (Jackson and Fountain, 2007) lost as a whole 47±2 % of their area since 1907 and 24.7±1.0% of their area since 2000 (Table 2, Fig. 4). From 2000 to 2023, the rate of area loss was 1.07±0.04 % yr⁻¹, ~3.5 times faster than the 1907-2000 rate of 0.31±0.02 % yr⁻¹ (Table 2). South-facing White River Glacier underwent the greatest rate of retreat in the 21st and 20th century, with the former being nearly three times faster than the latter (Table 2). North-facing Coe Glacier underwent the greatest 21st century acceleration in retreat, increasing by more than a factor of seven from 0.15±0.05 % yr⁻¹ to 1.09±0.11 % yr⁻¹ (Table 2).

Formatted: Font: Not Bold

Formatted: Font: Not Bold, Superscript

Formatted: Font: Not Bold

Formatted: Font: Not Bold, Superscript

Formatted: Font: Not Bold

Deleted: 57

Deleted: 3

Deleted: 39

Deleted: 8

Deleted: 7

Deleted: 73

Deleted: 17

Deleted: about six

Deleted: Ladd


Deleted: century

Deleted: , which was associated with the partial loss of its accumulation zone (Table 2; Fig. 3E, F). Conversely, White River Glacier had the greatest 20th century rate of retreat (Table 2), which has been previously attributed to the development of fumaroles by 1907 that reduced its accumulation area (Lillquist and Walker, 2006). Coe Glacier underwent the greatest increase in retreat rate in the 21st century about 12 times faster than its 20th century rate of area loss

Glacier	1907 Area km ² (uncert)	2000 Area km ² (uncert)	2023 Area km ² (uncert)	1907-2023 % Δ Area (uncert)	2000-2023 % Δ Area (uncert)	1907-2000 % yr ⁻¹ (uncert)	2000-2023 % yr ⁻¹ (uncert)
White River	1.04 (0.11)	0.44 (0.03)	0.257 (0.018)	75 (10)	41.7 (4.1)	0.62 (0.08)	1.81 (0.18)
Newton-Clark	2.06 (0.15)	1.54 (0.11)	1.050 (0.073)	49 (5)	31.8 (3.2)	0.27 (0.03)	1.38 (0.14)
Eliot	2.03 (0.16)	1.66 (0.12)	1.456 (0.102)	28 (3)	12.3 (1.2)	0.18 (0.02)	0.53 (0.05)
Coe	1.41 (0.13)	1.21 (0.08)	0.906 (0.063)	36 (4)	25.1 (2.4)	0.15 (0.02)	1.09 (0.11)
Ladd	1.07 (0.11)	0.71 (0.05)	0.550 (0.038)	49 (6)	22.6 (2.2)	0.36 (0.04)	0.98 (0.10)
Sandy	1.61 (0.17)	1.02 (0.07)	0.719 (0.050)	55 (7)	29.6 (2.9)	0.39 (0.05)	1.29 (0.13)
Reid	0.79 (0.13)	0.53 (0.04)	0.416 (0.029)	47 (8)	21.5 (2.2)	0.35 (0.06)	0.94 (0.10)
Seven combined	9.98 (0.37)	7.11 (0.21)	5.352 (0.158)	47 (2)	24.7 (1.0)	0.31 (0.02)	1.07 (0.04)

Glacier	1907 Area km ² (uncert)	2000 Area km ² (uncert)	2023 Area km ² (uncert)
White River	1.04 (0.11)	0.44 (0.03)	0.257 (0.018)
Newton-Clark	2.06 (0.15)	1.54 (0.11)	1.050 (0.073)
Eliot	2.03 (0.16)	1.66 (0.12)	1.456 (0.102)
Coe	1.41 (0.13)	1.21 (0.08)	0.906 (0.063)
Ladd	1.07 (0.11)	0.71 (0.05)	0.550 (0.038)
Sandy	1.61 (0.17)	1.02 (0.07)	0.719 (0.050)
Reid	0.79 (0.13)	0.53 (0.04)	0.416 (0.029)
Deleted: Seven combined	9.98 (0.37)	7.11 (0.21)	5.352 (0.158)

600 Table 2: Area for seven largest glaciers on Mt. Hood and their change (Δ) in area and rate of change expressed as percent. 1907 and 2000 areas are from Jackson and Fountain (2007).



605 Figure 4: Glacier area from 1907 (or 1901 for Eliot Glacier) to 2004 from Jackson and Fountain (2007), 2015 or 2016 from Fountain et al. (2023), and 2023 (new data), with combined area noted in the lower right. Uncertainty noted by vertical bars; note that the uncertainty on the combined record is smaller than the symbol size. A. White River Glacier. B. Newton-Clark Glacier. C. Eliot Glacier. D. Coe Glacier. E. Ladd Glacier. F. Sandy Glacier. G. Reid Glacier. H. Combined record of seven glaciers.

610 The 21st century rate of total area loss for these seven glaciers ($1.07 \pm 0.04 \text{ \% yr}^{-1}$) was 1.9 times faster the prior maximum rate of area loss from 1907-1946 ($0.56 \pm 0.02 \text{ \% yr}^{-1}$) (Fig. 5H). Six of the seven glaciers had individual 2000-2023 recession rates faster than their 1907-1946 rates (Fig. 5B-F). Reid Glacier's 2000-2023 rate of recession was $0.94 \pm 0.10 \text{ \% yr}^{-1}$ while its 1907-1946 rate was $0.84 \pm 0.14 \text{ \% yr}^{-1}$ (Fig. 5G).

- Deleted:** 73
- Deleted:** 17
- Deleted:** exceeded
- Deleted:** by a factor of about three
- Deleted:** Five
- Deleted:** 90
- Deleted:** 09
- Deleted:** White River Glacier retreated at $1.67 \pm 0.17 \text{ \% yr}^{-1}$ 2000-2023 whereas its 1907-1946 rate was $1.41 \pm 0.19 \text{ \% yr}^{-1}$ (Fig. 5A). ...




Figure 5: Area change with respect to preceding ice area expressed as % yr⁻¹ (yr = year) for seven glaciers calculated from Jackson and Fountain (2007) (1907-2000) and 2023 observations, with combined rate in the lower right. Gray shading shows the uncertainty. A. White River (WR) Glacier. B. Newton-Clark (NC) Glacier. C. Eliot Glacier. D. Coe Glacier. E. Ladd Glacier. F. Sandy Glacier. G. Reid Glacier. H. Combined record of seven glaciers.

4 Discussion

In the 21st century, every glacier on Mt. Hood lost area, resulting in the loss of ~25% glacierized area since 2000, with a minimum rise in the elevation of actively flowing termini of ~150 m on average over the last 20 years. This rapid area loss combined with a rise in the lowest elevation for flowing ice are indicative of large mass balance losses (e.g., Haeberli and Hoelzle, 1995; Pelto and Brown, 2012; Pelto, 2016; Hoelzle et al., 2023). Such clear glacier retreat and negative mass balance stand at odds with the Menounos et al. (2019) finding of near-neutral mass change for glaciers on Mt. Hood from 2000 to 2018. We note that their study lacked ground truthing of their geodetic measurements in Oregon, which combined with the high and growing debris cover on many of Mt. Hood's glaciers (e.g., Fig. 3H), could explain this disagreement. Furthermore, Florentine et al. (2024) found that not accounting for glacier area change when measuring geodetic mass balance reduces the mass-balance-change signal in western North American glaciers, with greater area loss leading to greater mass-balance bias. Here, our field observations clearly show large-scale glacier loss, consistent with the 21st-century retreat of glaciers around the globe (Zemp et al., 2015) and in agreement with the cumulative mass loss of benchmark glaciers to the north of Mt. Hood in Washington State (O'Neel et al., 2019; Pelto, 2023; World Glacier Monitoring Service, 2022).

Reconstructions of Ladd Glacier's extent in the 21st century highlight the difficulty that debris cover presents for mapping glacier extent in regions with significant rock fall. The 2015/2016 extent by Fountain et al. (2023), which did not include field checking, excluded a component of the debris-covered ice that we identified in the field in 2003 and that still remained

Formatted: Caption, Space Before: 0 pt, After: 0 pt

Deleted: [X]

Updating the length records for White River, Newton-Clark, Eliot, Coe and Ladd glaciers of Lillquist and Walker (2006) to 2023 (Table S1) found that four of these glaciers had now lost ~35% (White River) to ~42% (Eliot) of their 1901 length while Ladd Glacier...

Deleted: has

Deleted: about

Deleted: 40

Deleted: loss

Deleted: terminus

Deleted: terminus

Formatted: Space Before: 12 pt, After: 12 pt, No widow/orphan control

in 2023 (Fig. 6). Jackson and Fountain (2007) also included this debris covered ice in their 2004 Ladd extent (Fig. 1B), with this study including field checking of ice limits. This demonstrates the importance of field observations when mapping extents of glaciers with high amounts of debris cover.




Figure 6: Ladd Glacier terminus in 2023 (see Fig. 1B for photo location). White dashed lines are the 2023 limit while red dotted lines are the 2015/2016 ice limit of Fountain et al. (2023). 1. Notes ice bodies that Fountain et al. (2023) included in Ladd's 2015/2016 area. 2. Debris covered stagnant ice that Fountain et al. (2023) did not include in the 2015/2016 area of the glacier that remained in 2023 and was included in its 2023 limit.

The combined retreat for Mt. Hood's seven largest glaciers over the first 23 years of the 21st century stands out as unparalleled since at least the beginning of the 1900s (Fig. 4, 5), similar to the findings of Beason et al. (2023) for glaciers on Mt. Rainier in Washington and agreeing with a shift to more negative annual mass balance for western United States benchmark glaciers since 1990 (O'Neel et al., 2019). We compare Mt. Hood's rate of glacier recession of 1.07 ± 0.04 % yr⁻¹ from 2000 to 2023 (seven largest glaciers; Table 2) and 2.27 ± 0.13 % yr⁻¹ from 2015/2016 to 2023 (all glaciers except Ladd; Table 1) to other western United States glacierized ranges with comparable records of actively flowing glacier area. To the north in Washington State (Fig. 1A), Mt. Rainier glaciers retreated at 0.59 ± 0.05 % yr⁻¹ from 1994 to 2021 and 0.69 ± 0.05 % yr⁻¹ from 2015 to 2021 (Beason et al., 2023) while glaciers in the Olympic Mountains retreated at 1.44 ± 0.02 % yr⁻¹ from 1990 to 2015 and 2.75 ± 0.04 % yr⁻¹ from 2009 to 2015 (Fountain et al., 2022). To the northeast in Montana (Fig. 1A), Glacier National Park glaciers receded at ~ 0.77 % yr⁻¹ from 1998 to 2015 and ~ 0.83 % yr⁻¹ from 2005 to 2015 (Fagre et al., 2017). Lastly, to the south in California (Fig. 1A), glaciers in the Trinity Alps retreated at 3.89 ± 0.02 % yr⁻¹, between 1994 and 2015, leading to near complete deglaciation of the mountain range by 2015 (Garwood et al., 2020). Thus, the rates of glacier retreat on Mt. Hood in the 21st century are some of the highest rates in the western United States, but have yet to exceed the 2009-2015 rate in the Olympic Mountains, where the rate-change record ended in 2015, or the rate in the Trinity Alps, where the record ended in 2015 with near complete deglaciation.

Formatted: Caption, Line spacing: single

Deleted: 73

Deleted: 17

Deleted: 2.40±0.02 %

In the last 20 years, Glisan Glacier on Mt. Hood retreated to a 2023 area of $\sim 0.009 \text{ km}^2$ and ceased to be an actively flowing glacier, joining Palmer Glacier in this stagnant-glacier status (Table 1; Supplement). We find that two other glaciers are nearing this reclassification as stagnant glaciers: Zigzag and Coalman (Supplement). With only $\sim 0.1 \text{ km}^2$ of flowing ice remaining, Zigzag Glacier was only distinguishable from stagnant ice due to the presence of a few crevasses that have yet to start melting in on themselves (i.e., movement is keeping them open). Coalman Glacier at $\sim 0.06 \text{ km}^2$ (Table 1) is now below the area threshold of 0.1 km^2 used by the U.S. Geological Survey to distinguish active flowing glaciers (Fagre et al., 2017), but is above the 0.01 km^2 threshold used in global inventories (Pfeffer et al., 2014). More importantly, Coalman still showed evidence of flow with one active crevasse. These two glaciers also lost 40-48% of their area while Langille Glacier lost $\sim 47\%$ of its area in the last seven to eight years (Table 1). We predict that in the near future all three will stagnate if this rapid retreat continues.

Fluctuations in the rate of glacier area change on Mt. Hood generally corresponded with past changes in regional summer temperature and winter precipitation. The 20th century maximum rate of retreat from 1907 to 1946 (Fig. 7C) occurred during a period of warming summer temperatures (Fig. 7B) and reduced winter precipitation (Fig. 7A). Conversely, the interval of overall glacier stability from 1946 to 1972 (Fig. 4H, 7C) with some glaciers advancing (Fig. 4A, 4C, 4D, 4F, 4G) started with summer cooling followed by temperatures below the 1940s maximum concurrent with initial maxima in winter precipitation followed by elevated precipitation relative to the early 1900s. Marvel et al. (2019) previously attributed these changes in Pacific Northwest hydroclimate to anthropogenic forcing, with the initial warmer/drier period forced by rising anthropogenic greenhouse gas emissions and the subsequent cooler/wetter period caused by aerosol emissions counteracting continued greenhouse gas emissions. Renewed glacier retreat from 1972 to 2000 began with reduced winter precipitation and warming summer temperatures, but this pattern then changed in the 1990s where summer temperatures continued to warm along with greater winter precipitation. Glaciers still retreated in the 1990s despite greater winter precipitation (Fig. 4, 7C).

Deleted: We now return to our finding of unprecedented 21st century retreat of glaciers on Mt. Hood based on mapping only actively flowing glacier ice and comparing 2023 areas to equivalent actively flowing glacier areas in 2015/2016, 2000 and earlier years. If we had followed the other interpretation of what constitutes a glacier and included all stagnant ice in the 2023 glacier extents, would retreat in the 21st century still be unprecedented? As a test, we added to the 2015/2016 and 2023 areas of glaciers (Table 1) the area of additional snow/ice bodies noted in the Supplement mapped by Fountain et al. (2023). We conservatively assumed that these snow and ice bodies did not change in area from 2015/2016 to 2023, which was not the case as some of these ice bodies disappeared (see Supplement). With this additional fixed area, the change in glacierized/perennial snow area on Mt. Hood from 2015/2016 to 2023 was $>14\pm 1\%$ versus the change in area for actively flowing glaciers of $18\pm 1\%$. We thus conclude that defining a glacier as a flowing ice body does not greatly change the percentage of glacier area loss in the last seven to eight years.

As another test, we looked at the seven glaciers with records back to 1907 (Fig. 4, 5) (Jackson and Fountain, 2007). We added their peripheral 2015/2016 stagnant buried ice/stagnating ice/perennial snowfield area that Fountain et al. (2023) mapped as separate from the contiguous flowing glacier area to our 2023 glacier areas (Table S2). This additional area was $1.085\pm 0.041 \text{ km}^2$. About 0.718 km^2 ($\sim 66\%$) of this area was in the two buried ice bodies that were explicitly mapped as separate from and not parts of Coe and Eliot glaciers as of 2016 (Fountain et al., 2023). Another 12% of this area was in stagnating ice/perennial snowfields that had separated from Ladd Glacier as of 2016. We then compared these new larger glacier/ice/snow areas to the 2000 areas for these seven glaciers, noting that this is not an equal comparison. First, we again assumed that the areas of these peripheral snow/ice bodies did not change in the last seven to eight years, which is not the case. Second, the 2000 areas of these glaciers only included contiguous actively flowing glacier ice and explicitly did not include such peripheral snow ice/bodies or contiguous stagnant ice (Jackson and Fountain, 2007). With these assumptions, we found that these seven glaciers lost area at $>1.07\pm 0.1 \text{ \% yr}^{-1}$ 2000-2023, which is still unprecedented w... [8]

Deleted: has

Deleted: former-glacier

Deleted: former

Deleted: At

Deleted: ,

Deleted: With t

Deleted: losing

Deleted: 2

Deleted:


Deleted: , we

Deleted: they

Deleted: and become former glaciers.

Deleted: Langille Glacier also lost $>40\%$ of its area while Ladd Glacier lost about a third of its area, both in the last seven years (Table 1). I...

Deleted: , we predict that Langille and Ladd glaciers will be the next Mt Hood glaciers, after Zigzag and Coalman, to stagnate...



850
855
Figure 7: Hood River County climate change and Mt. Hood glacier area change rate. A. 11-year running mean for change in total October-May (light blue) and November-April (blue) precipitation relative to 1895-1924 average. B. 11-year running mean change in May-October (pink) and June-September (orange) temperature relative to 1895-1924 average. C. Rate of glacier area change for the seven glaciers (glac = glaciers) with records back to 1907 (Jackson and Fountain, 2007) (step plot with gray shading showing the uncertainty) and the rate of area change for 10 glaciers (excludes Ladd and Glisan) between 1981 and 2015/2016 and 2015/2016 and 2023 (Driedger and Kennard, 1986b; Fountain et al., 2023) (red crosses where the length is the period of time covered and the height the uncertainty). Note the inverted axis where faster rate of area loss is oriented up.

Deleted: The unprecedented rate of glacier retreat in 21st century (2000-2023) and from 2015/2016 to 2023 (Fig. 6C) corresponds with rising annual and summer temperatures (Fig. 6B). The 2013-2023 When the rate of glacier retreat ...

Formatted: Caption, Space Before: 0 pt, After: 0 pt

The unprecedented rate of glacier retreat in 21st century (2000-2023) and from 2015/2016 to 2023 (Fig. 7C) corresponded with regional summer temperatures warming to maxima (Fig. 7B). The 2013-2023 mean May-October and June-September temperatures were $\sim 1.7^{\circ}\text{C}$ and $\sim 1.8^{\circ}\text{C}$, respectively, warmer than their 1895-1924 means. Indeed, recent summer warmth in the Oregon, Washington and southern British Columbia is unequaled in more than a millennium (Heeter et al., 2023). Conversely, contemporary regional precipitation changes would not explain the 21st century glacier retreat on Mt. Hood, because the 2013-2023 means were around the 1895-1924 means (November-April was slightly below while October-May was slightly above the 1895-1924 means) and the preceding decade had greater amounts of precipitation (Fig. 7A).

As such, we argue that the retreat of Mt. Hood glaciers over the last ~ 120 years reflects regional climate warming (Roe et al., 2016, 2021) and that recent unprecedented retreat reflects the unequaled recent warmth. This agrees with mass-balance-temperature relationships for glaciers to the north in Washington State and for western United States benchmark glaciers. Pelto (2018) found significant relationships between annual mass balance with June-September temperature for 11 glaciers in Washington. O'Neel et al. (2019) noted the lack of a winter-mass-balance trend for the four benchmark glaciers in Washington and Alaska and a positive winter-mass-balance trend for the one glacier in Montana, meaning these glaciers' negative annual-mass-balance trends were being driven by summer temperatures.

5 Conclusions

We documented from field observations combined with remote sensing that the glaciers of Mt. Hood retreated and accordingly lost mass in the 21st century, losing $\sim 25\%$ of their area (seven largest glaciers) in the last 23 years and $\sim 17\%$ of their area (11 of 12 glaciers) in the last seven to eight years at unprecedented rates ($1.07 \pm 0.04\%$ yr^{-1} and $2.27 \pm 0.13\%$ yr^{-1} , respectively) with respect to the last 120 years. This is contrary to geodetic mass balance changes from remote-sensing data alone (Menounos et al., 2019). The rate of area loss in the 21st century has proceeded ~ 1.9 times faster than the fastest rate of the 20th century and ~ 3.5 times faster than the average 20th century rate. Two glaciers have ceased to flow (Glisan, Palmer) with another three glaciers rapidly losing area towards stagnation (Zigzag, Coalman, Langille). This recent rapid retreat of Mt. Hood glaciers can be attributed to a warming climate, with 2013-2023 regional summer means of $1.7\text{-}1.8^{\circ}\text{C}$ warmer than 1895-1924 means. As such, the warming climate of the last 120 years, especially the last 20 years, has significantly impacted the glacier coverage of Mt. Hood, with this recession continuing unless and until the sign of multi-decadal temperature change is reversed.

Data availability

The data referred to in this paper have all been provided within the tables and figures in the main text and supplement or as an asset available on a FAIR-aligned data repository (available at <https://doi.org/XX.XXXX/XXXXXXXXXX-XXXX-XXX>).

Deleted: Because of the significant relationship between the change in glacier length for five of Mt. Hood's glaciers and the change in regional temperature up to 2023 (Fig. 6B, C; Table 3), we argue that the retreat of glaciers over the last ~ 120 years reflects regional climate warming (Roe et al., 2016, 2021). We do not find a similar relationship with precipitation (Fig. 6A, C; Table 3), which does not necessarily mean that precipitation changes are not important but rather that no long-term trend exists for precipitation. A Mt. Hood glacier-length-temperature relationship agrees ...

Deleted: ¶
We suspect that the lack of a Mt. Hood glacier length-temperature relationship for the 20th century in the Lillquist and Walker (2006) study may partly reflect their methodology. Lillquist and Walker (2006) used a 5-year running average for both temperature and precipitation, shorter than the 11- to 30-year response time for glaciers like those on Mt. Hood (Pelto and Hedlund, 2001; Pelto, 2016), which we approximate with our 30-year and 15-year backward-running means. ¶
Frans et al. (2016) modeled the evolution of Newton-Clark, Eliot, Coe, and Ladd glaciers from 2000 to the end of the century under intermediate (RCP4.5) and high (RCP8.5) emission scenarios, with 10 simulations per emission scenario. From 2000 to 2025, they simulated a 10-30% loss in glacier area regardless of emission scenario. In contrast, we find that these four glaciers lost 30-50% of their combined area between 2000 and 2023. This means that glacier recession on Mt. Hood in the 21st century has proceeded ... [9]

Deleted: m

Deleted: ost likely

Deleted: ,

Deleted: 40

Deleted: 18

Deleted: all

Deleted: with respect to the last 120 years

Deleted: 73

Deleted: 17

Deleted: 40

Deleted: 02

Deleted: suggested by

Deleted: about three

Deleted: times faster than

Deleted: about six

Deleted: be glaciers

Deleted: four

Deleted: this reclassification

Deleted: , Ladd

Deleted: e century-scale

Deleted: regional 30-year mean temperature $1.1\text{-}1.3^{\circ}\text{C}$ warmer and 15-year mean temperature $1.3\text{-}1.5\text{...}$

Deleted: the

Author contributions

The author contributions, following the CRediT authorship guidelines, are as follows: NB-F, SJB, and AEC: conceptualization; NB-F, SJB, WCS, [DHR](#), and AEC: methodology; NB-F, SJB, WCS, and AEC: validation; NB-F, SJB, WCS, MT, [DHR](#), and AEC: analysis; NB-F, SJB, WCS, MT, DHR, and AEC: investigation; NB-F, SJB, WCS, DHR, and AEC: resources; NB-F, SJB, WCS, MT, and AEC: data curation; AEC: original draft; NB-F, SJB, WCS, MT, DHR, and AEC: review and editing; NB-F, SJB, WCS, MT, DHR, and AEC: visualization; NB-F, SJB, WCS, MT, DHR, and AEC: administration; NB-F, SJB, WCS, MT, DHR, and AEC: funding acquisition.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

We thank the U.S. National Forest Service for permitting research access to the Mt. Hood Wilderness Area and Andrew Fountain for sharing the historical area record for Mt. Hood. Insight from reviewers A. Fountain, M. Pelto, [F. Paul](#) and an anonymous reviewer greatly improved this manuscript, along with guidance from editor [B. Marzeion](#). This work was supported by a conservation grant from the Mazamas Mountaineering Club. The Oregon Glaciers Institute is a non-profit 501-c-3 that is supported by private donations.

References

[Beason, S. R., Kenyon, T. R., Jost, R. P., and Walker, L. J.:](#) Changes in glacier extents and estimated changes in glacial volume at Mount Rainier National Park, Washington, USA from 1896 to 2021, National Resource Report NPS/MORA/NRR-2023/2524, 2023.

[Cogley, J. G., Hock, R., Rasmussen, L. A., Arendt, A. A., Bauder, A., Braithwaite, R. J., Jansson, P., Kaser, G., Möller, M., Nicholson, L., and Zemp, M.:](#) Glossary of Glacier Mass Balance and Related Terms, IHP-VII Technical Documents in Hydrology No. 86, IACS Contribution No. 2, UNESCO-IHP, 2011.

Columbia Gorge Fruit Growers: Crops Information, <http://www.cgfg.org/information/crops-information>, 2024.

Dodge, N. A.: The Eliot Glacier: New Methods and Some Interpretations, Mazama Bulletin, 53, 25-33, 1971.

Deleted: along with community comments from F. Paul

Deleted: [Abatzoglou, J. T., Rupp, D. E., O'Neill, L. W., and Sadegh, M.:](#) Compound Extremes Drive the Western Oregon Wildfires of September 2020, *Geophys. Res. Lett.*, 48, e2021GL092520, 2021.

[Allgaier, M., Cooper, M. G., Carlson, A. E., Cooley, S. W., Ryan, J. C., and Smith, B. J.:](#) Direct measurement of optical properties of glacier ice using a photon-counting diffuse LiDAR, *J. Glaciol.*, 68, 1210-1220, 2022.

Deleted:

Deleted: [Burke, M., Childs, M. L., de la Cuesta, B., Qui, M., Li, J., Gould, C. F., Heft-Neal, S. and Wara, M.:](#) The contribution of wildfire to PM_{2.5} trends in the USA, *Nature*, 622, 761-766, 2023.

[Clarke, G. K. C.:](#) A short history of scientific investigations on glaciers, *J. Glaciol.*, special issue, 4-24, 1987.

[Clarke, G. K. C., Jarosch, A. H., Anslow, F. S., Radić, V., and Menounos, B.:](#) Projected deglaciation of western Canada in the twenty-first century, *Nat. Geosci.*, 8, 372-377, 2015.

- Dodge, N. A.: Eliot Glacier: Net Mass Balance, *Mazama Bulletin*, 69, 52-55, 1987.
- 1065 Driedger, C. L., and Kennard, P. M.: Glacier volume estimation on Cascade volcanoes: An analysis and comparison with other methods, *Ann. Glaciol.*, 8, 59-64, 1986a.
- Driedger, C. L., and Kennard, P. M.: Ice Volumes on Cascade Volcanoes: Mount Rainier, Mount Hood, Three Sisters, and Mount Shasta, U.S. Geological Survey Professional Paper 1365, 1986b.
- 1070 Ellinger, J. R.: The Changing Glaciers of Mt. Hood, Oregon and Mt. Rainier, Washington: Implications for Periglacial Debris Flows M.S. Thesis Oregon State University, 2010.
- Fagre, D. B., McKeon, L. A., Dick, K. A., and Fountain, A. G.: Glacier margin time series (1966, 1998, 2005, 2015) of the named glaciers of Glacier National Park, MT, USA, U.S. Geological Survey data release
1075 <https://dx.doi.org/10.5066/F7P26WB1>, 2017.
- Farmers Irrigation District, <https://www.fidhr.org/index.php/en/about-us/hydroelectric>, 2024.
- Florentine, C., Sass, L., McNeil, C., Baker, E., and O'Neel, S.: How to handle glacier area change in geodetic mass balance:
1080 *J. Glaciol.*, doi.org/jog.2023.86, 2024.
- Fountain, A. G., Glenn, B., and Basagic, H. J.: The geography of glaciers and perennial snowfields in the American West, Arctic, Antarctic, and Alpine Res., 49, 391-410, 2017.
- 1085 Fountain, A. G., Glenn, B., and McNeil, C.: Inventory of glaciers and perennial snowfields of the conterminous USA, *Earth Syst. Sci. Data*, 15, 4077-4104, 2023.
- Fountain, A. G., Gray, C., Glenn, B., Menounos, B., Pflug, J., and Riedel, J. L.: Glaciers of the Olympic Mountains, Washington – The Past and Future 100 Years, *J. Geophys. Res.*, 127, e2022JF006670, 2022.
- 1090 Frans, C., Istanbuluoglu, E., Lettenmaier, D. P., Clarke, G., Bohn, T. J., and Stumbaugh, M.: Implications of decadal to century scale glacio-hydrological change for water resources of the Hood River basin, OR, USA, *Hydrological Processes*, 30, 4314-4329, 2016.

1095 Frans, C., Istanbuluoglu, E., Lettenmaier, D. P., Fountain, A. G., and Riedel, J.: Glacier Recession and the Response of Summer Streamflow in the Pacific Northwest United States, 1960-2099, *Water Res. Res.*, 54, 6202-6225, 2018.

Garwood, J. M., Fountain, A. G., Lindke, K. T., van Hattem, M. G., and Basagic, H. J.: 20th Century Retreat and Recent Drought Accelerated Extinction of Mountain Glaciers and Perennial Snowfields in the Trinity Alps, California, *Northwest Science* 94, 44-61, 2020.

GLIMS and NSIDC: Global Land Ice Measurements from Space Glacier Database, Compiled and Made Available by the International GLIMS Community and the National Snow and Ice Data Center, doi: 10.7265/N5V98602, 2005 updated 2023.

1105 Haeberli, W., and Hoelzle, M.: Application of inventory data for estimating characteristics of and regional climate-change effects on mountain glaciers: a pilot study with the European Alps, *Ann. Glaciol.*, 21, 206-212, 1995.

Handewith, H. Jr.: Recent Glacier Variations on Mt. Hood, *Mazama Bulletin*, 41, 23-28, 1959.

1110 [Heeter, K. J., Harley, G. L., Abatzoglou, J. T., Anchukaitis, K. J., Cook, E. R., Coulthard, B. L., Dye, L. A., and Homfeld, I. K.: Unprecedented 21st century heat across the Pacific Northwest of North America, *npj climate and atmospheric science*, 6, 5, 2023.](#)

1115 Hoelzle, M., Haeberli, W., Dischl, M., and Peschke, W.: Secular glacier mass balances derived from cumulative glacier length changes, *Global Planet. Change*, 36, 295-306, 2023.

Howcutt, S., Spagnolo, M., Rea, B. R., Jaszewski, J., Barr, I., Coppola, D., De Siena, L., Girona, T., Gomez-Patron, A., Mullan, D., and Pritchard, M. E.: Icy Thermometers: Quantifying the impact of volcanic heat on glacier elevation, *Geology*, 51, 1143-1147, 2023.

1120 Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussaillant, I., Brun, F., and Kääb, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592, 726-731, 2021.

1125 Jackson, K. M., and Fountain, A. G.: Spatial and morphological change on Eliot Glacier, Mount Hood, Oregon, USA, *Ann. Glaciol.*, 46, 222-226, 2007.

Lawrence, D. B.: Mt. Hood's Latest Eruption and Glacier Advances, *Mazama Bulletin*, 30, 22-29, 1948.

Deleted:

Harper, J. T.: Glacier Terminus Fluctuations on Mount Baker, Washington, U.S.A., 1940-1990 and Climatic Variations, Arctic and Alpine Res., 25, 332-340, 1993....

Deleted:

Huston, A., Siler, N., Roe, G. H., Pettit, E., and Steiger, N. J.: Understanding drivers of glacier-length variability over the last millennium, *The Cryosphere*, 15, 1645-1662, 2021.

Deleted:

Jain, P., Sharma, A. R., Acuna, D. C., Abatzoglou, J. T., and Flannigan, M.: Record-breaking fire weather in North America in 2021 was initiated by the Pacific northwest heat dome, *Communications Earth & Environment*, 5, 202, 2024.

Kaspari, S. D., Pittenger, D., Jenk, T. M., Morgenstern, U., Schwikowski, M., Buenning, N., and Stott, L.: Twentieth Century Black Carbon and Dust Deposition on South Cascade Glacier, Washington State, USA, as Reconstructed From a 158-m-Long Ice Core, *J. Geophys. Res.*, 125, e2019JD031126, 2020.

Kaspari, S., Skiles, S. M., Delaney, I., Dixon, D., and Painter, T. H.: Accelerated glacier melt on Snow Dome, Mount Olympus, Washington, USA, due to deposition of black carbon and mineral dust from wildfire, *J. Geophys. Res.*, 120, 2793-2807, 2015.

Deleted:

Leclercq, P. W., and Oerlemans, J.: Global and hemispheric temperature reconstruction from glacier length fluctuations, *Clim. Dynam.*, 38, 1065-1079, 2012.

Leclercq, P. W., Oerlemans, J., Basagic, H. J., Bushueva, I., Cook, A. J., and Le Bris, R.: A data set of worldwide glacier length fluctuations, *The Cryosphere*, 8, 659-672, 2014.

Leigh, J. R., Stokes, C. R., Carr, J. R., Evans, I. S., Andreassen, L. M., and Evans, D. J. A.: Identifying and mapping very small (<0.5 km²) mountain glaciers on coarse to high-resolution imagery, *J. Glaciol.*, 65, 873-888, 2019.

Leonard, K. C., and Fountain, A. G.: Map-based methods for estimating glacier equilibrium-line altitudes, *J. Glaciol.*, 49, 329-336, 2003.

Lillquist, K., and Walker, K.: Historical Glacier and Climate Fluctuations at Mount Hood, Oregon, Arctic, Antarctic, and Alpine Res., 38, 399-412, 2006.

Lundstrom, S. C., McCafferty, A. E., and Coe, J. A.: Photogrammetric analysis of 1984-89 surface altitude change of the partially debris-covered Eliot Glacier, Mount Hood, Oregon, U.S.A., *Ann. Glaciol.*, 17, 167-170, 1993.

[Marvel, K., Cook, B. I., Bonfils, C. J. W., Durack, P. J., Smerdon, J. E., and Williams, A. P.: Twentieth-century hydroclimate changes. Consistent with human influences, *Nature*, 569, 59-65, 2019.](#)

Marzeion, B., Cogely, J. G., Richter, K., and Parkes, D.: Attribution of global glacier mass loss to anthropogenic and natural causes, *Science*, 345, 919-921, 2014.

Mason, R. S.: Recent Survey of Coe and Eliot Glaciers, *Mazama Bulletin*, 36, 37-39, 1954.

Menounos, B., Hugonnet, R., Shean, D., Gardner, A., Howat, I., Berthier, E., Pelto, B., Tennant, C., Shea, J., Noh, M.-J., Brun, F., and Dehecq, A.: Heterogeneous Changes in Western North American Glaciers Linked to Decadal Variability in Zonal Wind Strength, *Geophys. Res. Lett.*, 46, 200-209, 2019.

Miller, J. B., Frisbee, M. D., Hamilton, T. L., and Murugapiran, S. K.: Recharge from glacial meltwater is critical for alpine springs and their microbiomes, *Environ. Res. Lett.*, 16, 064012, 2021.

Mote, P. W., Rupp, D. E., Li, S., Sharp, D. J., Otto, F., Uhe, P. F., Xiao, M., Lettenmaier, D. P., Cullen, H., and Allen, M. R.: Perspectives on the causes of exceptionally low 2015 snowpack in the western United States, *Geophys. Res. Lett.*, 43, 10980-10988, 2016.

[Müller, F., Cafilisch, T., and Müller, G.: Instructions for Compilation and Assemblage of Data for a World Glacier Inventory, Temporary Technical Secretariat for World Glacier Inventory, International Commission on Snow and Ice, 1977.](#)

Deleted: ¶

Deleted: National Snow and Ice Data Center: Glacier, <https://nsidc.org/learn/cryosphere-glossary/glacier>, 2024a. ¶

National Snow and Ice Data Center: What is a glacier, <https://nsidc.org/learn/parts-cryosphere/glaciers/glacier-quick-facts>, 2024b. ¶

Neff, P. D., Steig, E. J., Clark, D. H., McConnell, J. R., Pettit, E. C., and Menounos, B.: Ice-core net snow accumulation and seasonal snow chemistry at a temperate-glacier site: Mount Waddington, southwest British Columbia, Canada, *J. Glaciol.*, 58, 1165-1175, 2012. ¶

Nelson, L. A.: A New Glacier on Mt. Hood, *Mazama Bulletin*, 6, 67-68, 1924.

1210 Nolin, A. W., Phillippe, J., Jefferson, A., and Lewis, S. L.: Present-day and future contributions of glacier runoff to summertime flows in a Pacific Northwest watershed: Implications for water resources, *Water Res. Res.*, 46, W12509, 2010.

O'Connor, J. E.: Our Vanishing Glaciers, *Oregon Historical Quarterly*, 114, 402-427, 2013.

1215 O'Neil, S., McNeil, C., Sass, L. C., Florentine, C., Baker, E. H., Peitzsch, E., McGrath, D., Fountain, A. G., and Fagre, D.: Reanalysis of the US Geological Survey Benchmark Glaciers: long-term insight into climate forcing of glacier mass balance, *J. Glaciol.*, 65, 850-866, 2019.

O'Neil, D.: Changes at Elevation, 1859 *Oregon's Magazine*, 77, 60-67, 2023.

1220 Paul, F., Barrand, N. E., Baumann, S., Berthier, E., Bolch, T., Casey, K., Frey, H., Joshi, S. P., Konovalov, V., Le Bris, R., Mölg, N., Nosenko, G., Nuth, C., Pope, A., Racoviteanu, A., Rastner, P., Raup, B., Scharrer, K., Steffen, S., and Winsvold, S.: On the accuracy of glacier outlines derived from remote-sensing data, *Ann. Glaciol.*, 54, 171-182, 2013.

1225 Pelto, M.: Climate Driven Retreat of Mount Baker Glaciers and Changing Water Resources, *Springer Briefs in Climate Studies*, doi: 10.1007/978-3-319-22605-7, 2016.

Pelto, M. S.: How Unusual Was 2015 in the 1984-2015 Period of the North Cascade Glacier Annual Mass Balance?, *Water*, 10, w10050543, 2018.

1230 Pelto, M. S.: Alpine Glaciers, *State of the Climate in 2022*, 2. *Global Climate*, S43-S44, 2023.

Pelto, M., and Brown, C.: Mass balance loss of Mount Baker, Washington glaciers 1990-2010, *Hydrol. Process.*, 26, 2601-2607, 2012

1235 Pelto, M. S., and Hedlund, C.: Terminus behavior and response time of North Cascade glaciers, Washington, U.S.A., *J. Glaciol.*, 47, 497-506, 2001.

1240 Pelto, M. S., Dryak, M., Pelto, J., Matthews, T., and Perry, L. B.: Contributions of Glacier Runoff during Heat Waves in the Nooksack River Basin USA, *Water*, 14, w14071145, 2022.

Deleted: [Orlemans, J.: Extracting a Climate Signal from 169 Glacier Records, *Science*, 308, 675-677, 2005.](#)

Deleted: [Paterson, W. S. B.: *The Physics of Glaciers*, third edition, 1994.](#)

- Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., Kienholz, C., Miles, E. S., Moholdt, G., Mölg, N., Paul, F., Radić, V., Rastner, P., Raup, B. H., Rich, J., Sharp, M. J., and The Randolph Consortium: The Randolph Glacier Inventory: a globally complete inventory of glaciers, *J. Glaciol.*, 60, 537-552, 2014.
- 1250
- Philip, S. Y., Kew, S. F., van Oldenborgh, G. J., Anslow, F. S., Seneviratne, S. I., Vautard, R., Coumou, D., Ebi, K. L., Arrighi, J., Singh, R., van Aalst, M., Marghidan, C. P., Wehner, M., Yang, W., Li, S., Schumacher, D. L., Hauser, M., Bonnet, R., Luu, L. N., Lehner, F., Gillett, N., Tradowsky, J. S., Vecchi, G. A., Rodell, C., Stull, R. B., Howard, R., and Otto, F. E. L.: Rapid attribution analysis of the extraordinary heat wave on the Pacific coast of the US and Canada in June 2021, *Earth Syst. Dynam.*, 13, 1689-1713, 2022.
- 1255
- Phillips, K.: Recent Changes in Hood's Glaciers, *Mazama Bulletin*, 17 45-50, 1935.
- 1260
- Phillips, K.: Our Vanishing Glaciers, *Mazama Bulletin*, 20, 24-41, 1938.
- Pitman, K. J., Moore, J. W., Sloat, M. R., Beaudreau, A. H., Bidlack, A. L., Brenner, R. E., Hood, E. W., Pess, G. R., Mantua, N. J., Milner, A. M., Radić, V., Reeves, G. H., Schindler, D. E., and Whited, D. C.: Glacier Retreat and Pacific Salmon, *BioScience*, 70, 220-236, 2020.
- 1265
- Robel, A. A., Ultee, L., Ranganathan, M., and Nash, M.: For whom and by whom is glaciology?, *J. Glaciol.*, accepted manuscript, doi.org/10.1017/jog.2024.29, 2024.
- Roe, G. H., Baker, M. B., and Herla, F.: Centennial glacier retreat as categorical evidence of regional climate change, *Nat. Geosci.*, 10, 95-99, 2017.
- 1270
- Roe, G. H., Christian, J. E., and Marzeion, B.: On the attribution of industrial-era glacier mass loss to anthropogenic climate change, *The Cryosphere*, 15, 1889-1905, 2021.
- 1275
- Rounce, D. R., Hock, R., Maussion, F., Hugonnet, R., Kochtitzky, W., Huss, M., Berthier, E., Brinkerhoff, D., Compagno, L., Copland, L., Farinotti, D., Menounos, B., and McNabb, R. W.: Global glacier change in the 21st century: Every increase in temperature matters, *Science*, 379, 78-83, 2023.
- 1280
- Thieman, C., Danielsen, A., Salminen, E., Berge, H., and Ross, K.: Watershed 2040, Hood River Basin Partnership Strategic Action Plan, Oregon Watershed Enhancement Board, 2021.

Thompson, V., Kennedy-Asser, A. T., Vosper, E., Lo, Y. T. E., Huntingford, C., Andrews, O., Collins, M., Hegerl, G. C., and Mitchell, D.: The 2021 western North American heat wave among the most extreme events ever recorded globally, *Sci. Adv.*, 8, eabm6860, 2022.

1285

U.S. Bureau of Reclamation: Reclamation Managing Water in the West: Hood River Basin Study, <https://www.usbr.gov/pn/studies/hoodriver/index.html>, 2015.

1290

U.S. Geological Survey: Eruption History of Mount Hood, Oregon, <https://www.usgs.gov/volcanoes/mount-hood/science/eruption-history-mount-hood-oregon>, 2023.

U.S. National Oceanic and Atmospheric Administration: Climate at a Glance, <https://www.noaa.gov/access/monitoring/climate-at-a-glance/>, 2024.

1295

U.S. National Resources Conservation Service: Oregon, Water, About Us, <https://www.nrcs.usda.gov/conservation-basics/conservation-by-state/oregon/oregon-snow-survey/about-us>, 2024.

World Glacier Monitoring Service: Reference glacier data, https://wgms.ch/products_ref_glaciers/, 2022.

1300

Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S. U., Helzle, M., Paul, F., Haeberli, W., Denzinger, F., Ahlström, A. P., Anderson, B., Bajracharya, S., Baroni, C., Braun, L. N., Cáceres, B. E., Casassa, G., Cobos, G., Dávila, L. R., Delgado Granados, H., Demuth, M. N., Espizua, L., Fischer, A., Fujita, K., Gadek, B., Ghazanfar, A., Hagen, J. O., Holmlund, P., Karimi, N., Li, Z., Pelto, M., Pitte, P., Popovnin, V. V., Portocarrero, C. A., Prinz, R., Sangewar, C. V., Severskiy, I., Sigurdsson, O., Soruco, A., Usabaliev, R., and Vincent, C.: Historically unprecedented global glacier decline in the early

1305

21st century, *J. Glaciol.*, 61, 745-762, 2015.

Deleted:
U.S. Geological Survey: What is a glacier?,
<https://www.usgs.gov/faqs/what-glacier>, 2024.

Page 6: [1] Deleted	Anders Carlson	13/07/2024 00:29:00
----------------------------	-----------------------	----------------------------

x

Page 6: [2] Deleted	Anders Carlson	14/07/2024 08:21:00
----------------------------	-----------------------	----------------------------

v

▲

Page 7: [3] Deleted	Anders Carlson	14/07/2024 08:34:00
----------------------------	-----------------------	----------------------------

v

▲

Page 9: [4] Deleted	Anders Carlson	13/07/2024 00:36:00
----------------------------	-----------------------	----------------------------

x

Page 9: [5] Deleted	Anders Carlson	24/07/2024 08:31:00
----------------------------	-----------------------	----------------------------

v

▲

Page 12: [6] Deleted	Anders Carlson	13/07/2024 00:34:00
-----------------------------	-----------------------	----------------------------

v

▲

Page 12: [7] Deleted	Anders Carlson	13/07/2024 00:34:00
-----------------------------	-----------------------	----------------------------

x

Page 14: [8] Deleted	Anders Carlson	12/07/2024 14:09:00
-----------------------------	-----------------------	----------------------------

x

Page 16: [9] Deleted	Anders Carlson	13/07/2024 00:17:00
-----------------------------	-----------------------	----------------------------

v

▲