

Opinion: The Role of AerChemMIP in Advancing Climate and Air Quality Research

Paul T. Griffiths^{1,2*}, Laura J. Wilcox^{3,*}, Robert J. Allen^{4,*}, Vaishali Naik⁵, Fiona M. O'Connor^{6,7}, Michael Prather⁸, Alex Archibald², Florence Brown⁹, Makoto Deushi¹⁰, William Collins¹¹, Stephanie Fiedler¹², Naga Oshima¹⁰, Lee T. Murray¹³, Bjørn H. Samset¹⁴, Chris Smith^{15,16}, Steven Turnock^{5,17}, Duncan Watson-Parris¹⁸, and Paul J. Young^{19,20}

¹School of Chemistry, Bristol University, UK.

²National Centre for Atmospheric Science, Cambridge University, UK

³National Centre for Atmospheric Science, Department of Meteorology, University of Reading, Reading, UK

⁴Department of Earth and Planetary Sciences, UC Riverside, Riverside, CA, USA

⁵NOAA Geophysical Fluid Dynamics Laboratory, Princeton, NJ, USA

⁶Met Office Hadley Centre, Exeter, UK

⁷Department of Mathematics & Statistics, Global Systems Institute, University of Exeter, UK

⁸Department of Earth System Science University of California, Irvine, US

⁹Institute for Atmospheric and Climate Science, ETH Zurich, Switzerland

¹⁰Department of Atmosphere, Ocean, and Earth System Modeling Research, Meteorological Research Institute, Tsukuba, Japan

¹¹Department of Meteorology, University of Reading, Reading, UK

¹²GEOMAR Helmholtz Centre for Ocean Research Kiel & Faculty of Mathematics and Natural Sciences, Christian-Albrechts-University of Kiel, Germany

¹³Department of Earth and Environmental Sciences, University of Rochester, NY, USA

¹⁴CICERO Center for International Climate Research, Oslo, Norway

¹⁵School of Earth and Environment, University of Leeds, UK

¹⁶International Institute for Applied Systems Analysis, Laxenburg, Austria

¹⁷University of Leeds Met Office Strategic (LUMOS) Research Group, University of Leeds, UK

¹⁸Scripps Institution of Oceanography and Halicioğlu Data Science Institute, UC San Diego, USA

¹⁹JBA Risk Management Ltd, Skipton, UK

²⁰School of Engineering, Newcastle University, Newcastle, UK

*These authors contributed equally to this work.

Correspondence: Paul Griffiths (paul.griffiths@ncas.ac.uk)

Abstract.

The Aerosol Chemistry Model Intercomparison Project (AerChemMIP) was endorsed by the Coupled-Model Intercomparison Project 6 (CMIP6) and was designed to quantify the climate and air quality impacts of aerosols and chemically reactive gases. AerChemMIP provided the first consistent calculation of Effective Radiative Forcing (ERF) for a wide range of forcing agents, which was a vital contribution to the sixth Assessment Report of the Intergovernmental Panel on Climate Change (AR6). It supported the quantification of composition-climate feedback parameters and the climate response to short-lived climate forcers (SLCFs), as well as enabling the future impacts of air pollution mitigation to be identified, and the study of interactions between climate and air quality in a transient simulations. Here we review AerChemMIP in detail, and assess the project against its stated objectives, its contribution to the CMIP6 project, and to the wider scientific efforts designed to

10 understand the role of aerosols and chemistry in the Earth System. We assess the successes of the project, and the remaining challenges and gaps. We conclude with some recommendations that we hope will provide input to planning for future MIPs in this area. In particular, we highlight the necessity of sufficient ensemble size for the attribution of regional climate responses, and the need for coordination across projects to ensure key science questions are addressed.

1 Introduction

15 The goal of the Coupled Model Intercomparison Projects (CMIP), used to inform successive generations of Intergovernmental Panel on Climate Change (IPCC) assessments, is to provide attribution and understanding of climate changes in the past, for the present day, and in future projections. Here we use ‘climate’ as writ large, encompassing the Earth system, particularly the atmospheric composition of greenhouse gases (GHG), chemically active trace gases and aerosols. Model assessment in a multi-model context, core to the CMIPs, is a necessary part of this process, establishing both the accuracy of models in
20 matching observations and their consistency in projecting change, thus enabling confidence in climate actions. An essential element of CMIPs is not just presenting the model spread, but also in providing critical diagnostics so that we can understand the cause of model differences and identify better modelling approaches.

The IPCC released the Sixth Assessment Reports (AR6) over the period 2021-2023 based on and informed by the 6th Phase of CMIP (CMIP6). CMIP6 defined a series of common experiments including, the DECK (Diagnostic, Evaluation and Char-
25 acterization of Klima; Eyring et al., 2016) and historical experiments, standardised input datasets to drive model experiments (input4MIPs), sponsored infrastructure activities, such as input4MIPs and the Earth System Grid Federation (ESGF), and endorsed a host of sub-MIPs focused on specific science questions. The design of the DECK provided experiments for assessing internal model variability (*piControl*), calculating climate sensitivity (*abrupt-4xCO2* and *1pctCO2*), and providing data for comparison with observations (AMIP). Alongside the coupled transient *historical* experiment, these simulations represented
30 the ‘entry card’ for CMIP6 (Eyring et al., 2016).

In this paper, we review the Aerosol and Chemistry MIP (AerChemMIP; Collins et al., 2017), one of the endorsed MIPs of CMIP6, examining its requested experiments, model participation, and the overall contribution to our understanding of the role of short-lived climate forcers (SLCFs) in climate change, including chemistry-aerosol-climate couplings and feedbacks. The criteria for the endorsed MIPs were set by CMIP6 (Meehl et al., 2014; Eyring et al., 2016; Stouffer et al., 2017) such that they
35 needed to address one or more of three broad scientific questions: (Q1) How does the Earth System respond to forcing? (Q2) What are the origins and consequences of systematic model biases? (Q3) How can we assess future climate changes given climate variability, predictability and uncertainties in scenarios? AerChemMIP addressed all of these questions, to varying degrees.

2 AerChemMIP - objectives and protocol design

40 A primary objective for AerChemMIP was to diagnose and document forcings and climate responses to changes in aerosols and GHGs, such as ozone and methane, in the CMIP6 models (Collins et al., 2017). The AerChemMIP experiments enabled the assessment of the role of SLCFs, i.e., aerosols and chemically reactive gases, in historical and future climate change, as well as a more robust quantification of climate sensitivities to chemistry and aerosol changes based on the current generation of comprehensive Earth System Models (ESMs) (Thornhill et al., 2021a). The focus of AerChemMIP on SLCFs was due to
45 the findings of the IPCC 5th Assessment Report (AR5) that SLCFs were the main source of uncertainty in estimates of the effective radiative forcing (ERF). SLCF contributions to anthropogenic forcing include changes in aerosols and their precursors; methane; tropospheric ozone, formed from sunlight, nitrogen oxides (NO_x), carbon monoxide, volatile organic compounds and methane; and stratosphere ozone levels, which are affected by ozone-depleting substances (ODSs). These have contributed significantly to past climate change, with a combined pre-industrial (PI) to present-day (PD) GHG forcing from short-lived
50 GHGs such as methane and tropospheric ozone, estimated in AR5 to be comparable to that of carbon dioxide, and cooling from the direct and indirect effects of aerosol of a similar magnitude.

Taken together, anthropogenic aerosols act to cool the climate, and have offset around a third of GHG-driven warming since 1850 (Szopa et al., 2021a). They have also accounted for the largest component of uncertainty in anthropogenic radiative forcing through successive IPCC assessment reports, reflecting, amongst other issues, diverse approaches to modelling aerosol
55 processes and their interaction with the climate system, and difficulties in using observations to constrain aerosol forcing (Boucher et al., 2013; Bellouin et al., 2020; Forster et al., 2021). CMIP5 brought a significant advance in the representation of aerosol-climate interactions, with two thirds of models including a representation of aerosol-cloud interactions, and more models including interactive aerosol schemes (Wilcox et al., 2013; Ekman, 2014). In CMIP6, a new generation of ESMs came online with structural changes such as two-moment aerosol schemes; online interactive biogenic volatile organic compound
60 (BVOC) emissions, which serve as ozone and aerosol precursors; and ocean biogeochemistry describing e.g., sulfur-containing aerosol precursor species, such as dimethyl sulfide (DMS) with implications for the natural background aerosol state. This change in the background state has the potential to have a large impact on the calculation of PI-to-PD radiative forcing from anthropogenic sulfate aerosol (Carslaw et al., 2013).

Of the reactive gases, methane and ozone were of primary interest for AerChemMIP. Methane is an important GHG, and
65 atmospheric concentrations have increased by a factor of ~2.5 since PI times (e.g., Skeie et al., 2023). It serves as a precursor to tropospheric ozone and stratospheric water vapor, and as a sink for the hydroxyl radical, with a significant impact on tropospheric oxidising capacity and therefore the lifetimes of several climate forcers (e.g., halocarbons, and methane itself). Accurate simulation of atmospheric methane trends from emissions remains a challenge, as there exist a variety of anthropogenic and natural sources such as wetlands, which are capable of responding to climate change, and for this reason, as in
70 CMIP5, many of the CMIP6 models used prescribed model lower boundary conditions for historical and future simulations, produced from integrated assessment models.

Tropospheric ozone is an important SLCF. It is a GHG that has its largest radiative impact in the upper troposphere/lower stratosphere (UT/LS). High surface levels of ozone are associated with adverse effects on human health and vegetation (e.g., Anenberg et al., 2009; Emberson, 2020). It is therefore important to attribute the causes of increases since PI times and understand its evolution in the future with changing climate and precursor emissions. CMIP6 was the first CMIP in which significant number of modelling centres included interactive tropospheric and stratospheric ozone in their flagship models, with whole atmosphere chemical schemes allowing the effect of stratospheric changes on tropospheric chemistry to be attributed (Stevenson et al., 2020; Zeng et al., 2022). Biogenic VOCs serve as ozone precursors, and their emissions are dependent on climate state. All models bar one (MRI-ESM2-0) used in the CMIP6 assessment for tropospheric ozone had some form of online isoprene emissions, with some including more detailed treatments (monoterpene emissions, the effect of carbon dioxide (CO₂) inhibition, and/or a larger suite of online BVOC emissions) (Griffiths et al., 2021).

2.1 AerChemMIP objectives

The AerChemMIP science questions, as set out in Collins et al. (2017) using the nomenclature of near-term climate forcers (NTCFs), subsequently renamed in AR6 to SLCFs, were:

- A1. How have anthropogenic emissions contributed to global radiative forcing and affected regional climate over the historical period?
- A2. How might future policies (on climate, air quality, and land use) affect the abundances of NTCFs and their climate impacts?
- A3. How do uncertainties in historical NTCF emissions affect radiative forcing estimates?
- A4. How important are climate feedbacks to natural NTCF emissions, atmospheric composition, and radiative effects?

As a CMIP6-endorsed MIP, AerChemMIP was designed to build on the DECK and historical experiments, particularly the historical coupled atmosphere-ocean and the atmosphere-only AMIP experiments, and to interface with future scenario simulations coordinated by ScenarioMIP (O'Neill et al., 2016). AerChemMIP experiments were sorted into Tiers according to priority, with the Tiers loosely tied to the science questions: Tier 1 experiments aimed to answer questions A1 and A2, Tier 2 experiments addressed A4 with Tiers 1-3 aiming to answer A3. The AerChemMIP requested additional diagnostics to complement analyses of CMIP and ScenarioMIP experiments. Figure 1 shows the AerChemMIP experimental tiers

In support of its objectives, the AerChemMIP protocol defined historical and future experiments, as well as transient experiments, which are based on the parent CMIP *historical* and ScenarioMIP *SSP3-7.0* experiments, and idealised, so-called 'timeslice' experiments which were forced with SSTs and sea-ice from the corresponding coupled AOGCM and with other boundary conditions appropriate to a single year, used for calculating ERFs.

The calculation of ERFs, addressing A1, was an important focus of AerChemMIP. When the concept of ERF was introduced in AR5 Working Group 1 (WG1), ERFs for historical GHGs, natural forcings, aerosols, and ozone were inferred from coupled transient simulations by removing the effect of the surface temperature response (Forster et al., 2013). ERF contributions from

	Coupled A/O - <u>climate response</u>		Transient experiment, prescribed SST - <u>transient ERF</u>		Idealized timeslice - <u>PI-PD ERF</u>	
	hist-	ssp370-	histSST-	ssp370SST-	piClim-	
T1 (A1-A4)	piNTCF 1950HC	ssp370 lowNTCF	histSST piNTCF 1950HC piCH4	ssp370SST lowNTCF	Control NTCF CH4 HC	
T2 (A3, A4)	piAer		piAer piO3 piN2O	lowAer lowBC lowO3 lowCH4 ssp126Lu	aer BC O3 N2O	
T3 (A3)					NOx VOC SO2 OC NH3 2xDMS 2xfire 2xNOx 2xVOC	

Figure 1. Schematic of AerChemMIP protocol experiments, showing tiers (T), AerChemMIP objectives (A) and species involved

ozone and aerosols were diagnosed in offline calculations (Shindell et al., 2013), based on changes in composition derived in the Atmospheric Chemistry and Climate MIP (ACCMIP; Lamarque et al., 2013). The model configurations participating in ACCMIP were in many cases different from those in CMIP5 in terms of both resolution and complexity of chemistry and aerosols which resulted in ACCMIP not being able to fully describe the forcings in the coupled models. The systematic approach for calculating ERFs due to individual SLCF species in AerChemMIP represents a significant advance for CMIP6 over the CMIP5 approach.

2.2 AerChemMIP timeslice experiments

The AerChemMIP protocol described the simulations needed to calculate PI-to-PD and transient historical ERFs for individual species, such as methane (CH₄) and sulfur dioxide (SO₂). The *piClim-X* experiments in AerChemMIP consisted of a control experiment (*piClim-control*) that had all WMGHG concentrations and SLCF emissions at PI levels, and individual experiments

where one species (or group of species) were changed to PD levels. These simulations were run as 'timeslice' experiments in which an atmosphere-only GCM is forced with SSTs from a corresponding coupled atmosphere-ocean GCM and other boundary conditions appropriate to a single year. The internal model variability (mainly clouds) generates considerable interannual variability in ERFs and the perpetually-repeating boundary conditions allow for repeated realizations of the atmospheric state that both incorporate this internal variability and allow for improved statistics for the estimation of small changes in e.g. TOA radiative forcing. Moreover, these shorter, 30-year timeslice simulations were able to be performed for more forcing agents, with more models, compared to analogous 165-year transient historical simulations. The AOGCM timeslice experiments allow the ERFs to be diagnosed directly, in the absence of slow climate responses and feedbacks, rather than being inferred. It should be noted that some groups found it necessary to extend the *piClim-X* experiments to 45 years to allow stratospheric concentrations of WMGHGs sufficient time to spin up (O'Connor et al., 2021).

AerChemMIP timeslice experiments for natural 2x experiments are analogous to the above, but here the specified PI natural emission flux is doubled, either by scaling the parameterisations in an interactive scheme or the input data files for specified emissions (Collins et al., 2017). For most of these experiments, the emission flux to be doubled is intuitive (e.g., dust emissions are doubled for *piClim-2xdust*). For *piClim-2xfire*, fire-related emissions for several species are doubled, e.g., NO_x, BC, OC, CO, VOCs.

2.3 AerChemMIP historical transient experiments

AerChemMIP coupled-transient-attribution experiments, addressing the role of historical or future emissions changes, were grouped into species categories (*hist-piNTCF*, *-piO3*, *-1950HC*) to keep down computational cost, while the less expensive prescribed SST experiments were performed over a wider range of forcings to quantify preindustrial (PI) to present-day (PD) and historical transient ERF due to changes in the individual forcing agents. This 'everything but' (i.e. every forcing follows the historical trajectory except for a single named forcing held at PI levels) approach meant that any nonlinearities arising due to a warming climate were captured in the AerChemMIP experiments, making them a useful counterpoint to the DAMIP single forcing experiments. These were designed to be used in optimal fingerprinting approaches for detection and attribution, involving a linear regression of historical climate observations onto corresponding models. The DAMIP approach assumed that the climate responses to single forcings can be combined linearly, an assumption that often breaks down at regional scales and can be tested by a comparison between AerChemMIP and DAMIP experiments.

2.4 AerChemMIP ssp370 experiments

Future climate and air pollution was targeted using the Shared Socioeconomic Pathway 3-7.0 (SSP3-7.0, Rao et al., 2017; Riahi et al., 2017) - a 'regional rivalry' scenario without climate policy and weak air pollution mitigation policies. ScenarioMIP contributed the AOGCM coupled *ssp370* experiments (40 models, 379 experiments) as a baseline for understanding the role of aerosols and chemistry in a future climate (O'Neill et al., 2016). AerChemMIP provided complementary *ssp370SST* experiments for transient ERF calculations, and a coupled AOGCM *ssp370-lowNTCF* experiment to attribute the role of aerosols and gaseous emissions, excluding methane, on climate and composition (Allen et al., 2020). For most short-lived precursor

species for air quality and aerosol climate forcing, including NO_x , CO , CH_4 , NMVOCs, SO_2 , BC and OC, the emissions or surface boundary conditions in the *SSP3-7.0* scenario were substantially higher than those in the most extreme warming scenario considered by CMIP6 (*SSP5-8.5*). For the purposes of attributing the impact of emissions changes, this scenario offered the strongest signal and therefore the greatest potential for attribution. For the purposes of policy development, simulations were required until 2055 with most models extending to 2100. Land-use changes were separately assessed via the Land Use MIP (LUMIP; Lawrence et al., 2016) experiment, *ssp370-ssp126Lu*.

To isolate the effects of air pollution mitigation policy, the *SSP3-7.0-lowNTCF* scenario uses the same socio-economic scenario and the same emissions drivers as *SSP3-7.0*, but with ‘strong’ air pollution mitigation policies. In the case of air pollutant species (e.g., SO_2 , BC, OC, NO_x , NH_3 and AVOCs), the emissions factors used in the sustainability pathway *SSP1* were adopted (Gidden et al., 2019). *SSP3-7.0-lowNTCF* was designed so that the reduced NTCF emissions relative to *SSP3-7.0* came only from changes in air quality policy, neglecting changes coming from simultaneous changes in climate policy. The decrease in air pollutant species emissions is due to swift ramping up of end-of-pipe measures for air pollution control, not from the reduction in co-emissions accompanying the CO_2 emissions also seen in *SSP1*. The result was that, under *SSP3-7.0-lowNTCF*, global emissions of all aerosols and gaseous precursors decrease, particularly by mid-century by $\sim 30\text{--}50\%$: reductions comparable to those seen in *SSP2-4.5* e.g. Figure 1 of Wilcox et al. (2023). In contrast, the corresponding emissions under *SSP3-7.0* generally increase (weakly) by mid-century by $\sim 10\%$ (Allen et al., 2021).

Subsequent to the publication of Collins et al. (2017), further experiments were developed to attribute the role of individual SLCFs, such as the coupled AOGCM experiment *ssp370-lowNTCFCH4*, addressing the additional role of methane mitigation (Allen et al., 2021), and *ssp370pdSST* employing future emissions, but prescribing a PD climatology of SSTs and sea ice, to characterise the effect of future climate change on composition (Zanis et al., 2022). For completeness, these experiments are included in Table 1, despite not being assigned a Tier in Collins et al. (2017), using their designation on ES-DOC (<https://search.es-doc.org/>). These experiments were critical for drawing two conclusions relating to SLCFs in the IPCC AR6: 1) changes in future air pollution are more likely to be driven by changes in anthropogenic emissions than climate change; and 2) controls on SLCF emissions, particularly methane, are important for meeting climate goals with simultaneous air quality benefits.

2.5 AerChemMIP and other CMIP6 MIPs

AerChemMIP aimed to draw on the skills and interests of the atmospheric chemistry, aerosol, and radiative forcing communities to address research questions of mutual interest. The community coalesced around semiannual “TriMIP” meetings (reviewed in Fiedler et al. (2024)), involving meetings of the Precipitation Driver and Response Model Intercomparison Project (PDRMIP; Myhre et al., 2017), the Radiative Forcing Model Intercomparison Project (RFMIP; Pincus et al., 2016), and AerChemMIP. These three MIPs shared many interests and scientific goals, but with different foci and approaches (Fiedler et al., 2024).

AerChemMIP was designed to advance the understanding of the role of SLCFs in transient climate change, and the magnitude of the climate response to realistic forcings.

PDRMIP investigated the role of various drivers of climate change for mean and extreme precipitation changes. It included a range of present-day equilibrium experiments with large, idealised perturbations in either emissions or concentrations of well-mixed GHGs (WMGHGs), SLCFs, or natural forcings. The application of such large forcings resulted in clear signals that advanced our physical understanding of the climate response to these forcings. However, information from such idealised experiments can be difficult to apply to the real world, and PDRMIP was complemented by AerChemMIP in this regard which focused on well-defined pathways describing historical and future forcings.

RFMIP aimed to provide both a foundational understanding of the Earth system response to forcing, and described a range of experiments for the quantification of the ERF of individual or groups of forcing agents. Quantifying ERF was a common goal between AerChemMIP and RFMIP, with complementary protocols: firstly, AerChemMIP and RFMIP both included experiments designed to calculate the 2014 vs. 1850 ERF due to SLCFs. There were common control simulations of (*piClim-control* and *piClim-aerO3* in RFMIP and *piClim-control* and *piClim-NTCF* in AerChemMIP). These ERF calculations were based on common definitions of present-day (PD) vs. pre-industrial (PI, 1850) for individual species (concentrations or emissions) and a common protocol of 30-year (perpetually repeating) timeslice experiments driven by fixed monthly climatological (1850) SSTs and sea ice, as recommended by Forster et al. (2016). Secondly, both RFMIP and AerChemMIP protocols defined experiments to diagnose the transient ERF for the historical period (*piClim-histaerO3* for RFMIP, and *histSST* and *histSST-piNTCF* for AerChemMIP). However, while the timeslice experiments for the 2014 vs. 1850 ERF due to SLCFs were identical in RFMIP and AerChemMIP, the experimental designs were different for these transient experiments. For transient experiments, pre-industrial SSTs and sea-ice concentrations were prescribed for RFMIP compared to SSTs and sea-ice concentrations from the historical experiment for AerChemMIP. AerChemMIP also used the ‘everything but’ design, while RFMIP used the single forcing approach taken by the Detection and Attribution MIP (DAMIP; Gillett et al., 2016).

AerChemMIP also aligned with Detection and Attribution MIP (DAMIP; Gillett et al., 2016), that used single forcing experiments in multiple generations of CMIP, to quantify the proportion of an observed trend likely to be driven by a particular forcer. However, the DAMIP approach rests on the assumption that the climate response to individual forcers, or groups of forcers, can be combined linearly to reproduce the total climate response. This assumption doesn’t always hold, especially for regional climate (Stephens et al., 2016; Deng et al., 2020; Aizawa et al., 2022), and even the forcings themselves do not combine linearly (O’Connor et al., 2021). AerChemMIP experiments therefore provide additional experiments to enable assessment of DAMIP analyses.

3 Contributions and successes

At the time of writing, there are 12 published articles in the special issue of *Atmospheric Chemistry and Physics* (Shonk et al., 2020; Stevenson et al., 2020; Zanis et al., 2020; Wilcox et al., 2020; Turnock et al., 2020; Allen et al., 2020; Mortier et al., 2020; Griffiths et al., 2021; Glibb et al., 2021; Thornhill et al., 2021a; O’Connor et al., 2021; Zhang et al., 2021a), with some early synthesis reports also contributed (Forster et al., 2016). While it is difficult to completely separate AerChemMIP from the wider CMIP6 publications, a further 22 articles mention AerChemMIP [Web of Science, 2024-07-26]. Articles highlighted

the role of oxidants (Karset et al., 2018; O'Connor et al., 2021; Griffiths et al., 2021), aerosol processes (Zhang et al., 2021a),
215 and the role of methane (Stevenson et al., 2020; O'Connor et al., 2022).

The number of experiments (37) and the number of simulated model and ensemble years (1,265 for Tier 1, 1,369 for Tier
2, and 270 for Tier 3) made AerChemMIP the largest of the CMIP6-endorsed MIPs. Despite this, there was good uptake by
modelling centres, with up to 19 models performing at least one of the AerChemMIP experiments. Table 1 shows details of
the data available via the ESGF archive in August 2024. Over the historical period, the coupled atmosphere-ocean general
220 circulation model (AOGCM) experiments are well represented with more than ten models performing *hist-piNTCF* and *hist-*
piAer. In general, experiments targeting historical aerosol emissions received the most effort (Figure 1). The average number
of models per experiment was seven, and for coupled hist-AOGCM experiments, there was an average of three ensembles
per experiment. The AerChemMIP research questions were at the cutting edge of the capability of the current generation of
climate models, and participation reflected model capability e.g. models using offline chemistry were necessarily excluded from
225 experiments, such as *hist-piO3*, limiting the number of models running this experiment to five. These five ESMs (UKESM1-0-
LL, MRI-ESM2-0, GISS-E2.1-G, CESM2-WACCM and GFDL-ESM4) with interactive chemistry in fact ran the majority of
the experiments of which they were capable, contributing to all AerChemMIP tiers. The experiment with the lowest number of
participating models was *piClim-NH3*, run only by IPSL-CM6A-LR-INCA and GISS-E2.1-G, which are two of only a small
number of CMIP6 models to feature online ammonia emissions and to treat explicitly the formation of ammonium-containing
230 aerosol. While the absolute number of participating models is a useful metric, the statistics do inevitably reflect the number
of models *capable* of running a given experiment. In this sense, AerChemMIP enjoyed very pleasing success with community
members contributing data to nearly every experiment that their models were capable of running.




Figure 2. The AerChemMIP experiments, and the number of models used to perform them (orange bars), based on the availability of data in the ESGF archive in August 2024. Blue bars show the number of publications using each experiment to date, according to the Web of Science.

EXPT ID	hist-1950HC	hist-piNTCF	histSST	histSST-1950HC	histSST-piCH4	histSST-piNTCF	piClim-CH4	piClim-HC	piClim-NTCF	piClim-control
TIER	1	1	1	1	1	1	1	1	1	1
MODELS	6	11	12	6	8	11	9	6	11	22
Members	19	27	13	6	8	12	10	6	12	32
EXPT ID	hist-piAer	histSST-piAer	histSST-piN2O	histSST-piO3	piClim-2xdust	piClim-2xSS	piClim-BC	piClim-N2O	piClim-O3	piClim-aer
TIER	2	2	2	2	2	2	2	2	2	2
MODELS	10	8	4	4	9	9	11	6	6	19
Members	24	9	4	4	10	10	12	7	6	25
EXPT ID	piClim-2xDMS	piClim-2xNOx	piClim-2xVOC	piClim-2xfire	piClim-NH3	piClim-NOx	piClim-OC	piClim-SO2	piClim-VOC	
TIER	3	3	3	3	3	3	3	3	3	
MODELS	6	4	6	6	2	6	10	11	6	
Members	7	4	7	6	2	6	12	13	6	
EXPT ID	ssp370-lowNTCF	ssp370SST	ssp370SST-lowNTCF	ssp370SST-lowAer	ssp370SST-BC	ssp370SST-lowO3	ssp370SST-lowCH4	ssp370-ssp126Lu	ssp370SST-lowNTCFCH4	
TIER	1	1	1	2	2	2	2	2	2	
MODELS	13	11	8	6	6	3	4	5	6	
Members	40	11	8	6	6	3	4	5	6	
EXPT ID	ssp370pdSST	ssp370-lowNTCFCH4								
TIER	2+	1+								
MODELS	7	6								
Members	7	12								

Table 1. Information on AerChemMIP simulations, including (horizontally displayed) experiment abbreviations (EXPT ID) and (vertically displayed) experiment tier (TIER, i.e., priority), number of models that performed a given experiment (MODELS) and the total number of ensemble members (MEMBERS; i.e., independent realizations).

AerChemMIP within CMIP6 provided several key points of analysis to inform AR6. These included: quantitative understanding of the role of anthropogenic drivers in historical oxidising capacity; an assessment of emissions-based effective radiative forcings for SLCFs, cumulatively and individually; an improved estimate of forcing by ozone-depleting substances using an observational constraint; assessment of climate and air quality impacts due to mitigation of SLCFs (in particular the role of methane); a more robust quantification of non-CO₂ biogeochemical-climate feedbacks; and several evaluations including tropospheric ozone, stratospheric ozone and water vapour, and air quality. The work informed quantification and improved understanding of model biases (e.g., UKESM1 response to ozone-depleting substances (Morgenstern et al., 2020); surface ozone biases (Liu et al., 2022); historical temperature biases (Zhang et al., 2021a)) and the development of emulators (Smith et al., 2024). It continued the use of fixed SST and coupled simulations for quantifying responses on different timescales pioneered in PDRMIP, demonstrated the value of a consistent set of aerosol and gas phase experiments for attribution, and improved links to other projects such as the Chemistry Climate Model Initiative (CCMI) which used the CMIP6 scenarios for its projections of ozone recovery.

3.1 AerChemMIP *piClim* experiments

As shown in Table 1, the -CH₄ and halocarbon (-HC) chemistry experiments have received the most effort. For experiments requiring online chemistry, participating centres were more likely to perform *piClim-X* experiments than the analogous *histSST-piX* experiments.

The ERFs from the *piClim-X* experiments were analysed by Thornhill et al. (2021b) and more recently by Kalisoras et al. (2023) from the online radiation diagnostics. The ERFs were further broken down into instantaneous radiative forcings (IRFs) and adjustment terms due to temperature (tropospheric and stratospheric), water vapour, surface albedo, and clouds using offline radiative kernels. Using this approach, the separation of the ERF due to aerosols into aerosol-radiation interactions (ari) and aerosol-cloud interactions (aci) showed reasonable agreement with the double-call radiation diagnostics for aerosol and cloud forcing based on the methodology of Ghan (2013). Oshima et al. (2020) estimated the present-day ERFs from individual anthropogenic agents comprehensively and suggested the importance of the interactions of aerosols with ice clouds over the tropics and possible important role of black carbon (BC) in Arctic surface warming. A notable result of the analysis of the adjustment terms was that for some models (UKESM1 in particular) there was a significant contribution to the ERFs for ozone precursors (such as CH₄ and anthropogenic NO_x) from changes in cloud radiative effects due to changes in the atmospheric oxidising capacity. O'Connor et al. (2021, 2022) concluded that much of the cloud radiative effect originated through changes in levels of OH and O₃ and associated changes in the gas-phase (rather than aqueous-phase) oxidation of SO₂ and aerosol size distribution. Further interactions between NO_x and ERF would be expected through changes in nitrate aerosol. Nitrate aerosols are treated in GFDL-ESM4, for instance, but not in UKESM1. The role of nitrate aerosol in aerosol-cloud interactions remains largely unexplored in CMIP6 models, and future generations of AerChemMIP may need additional experiments to understand the coupling between NO_x emissions, nitrate aerosol and aerosol-cloud interactions.

AerChemMIP found that NO_x emissions made the largest contribution to the PI-to-PD tropospheric ozone forcing (Thornhill et al., 2021b), whereas ACCMIP had found that the largest contribution was from methane (Stevenson et al., 2013), although

part of this difference in the attribution of the ozone forcing may be related to the difference in NO_x emissions between CMIP5 and CMIP6.

The NO_x emissions used in CMIP6 from the Community Emission Data System (CEDS; Hoesly et al., 2018) are smaller
270 than the CMIP5 estimates until the mid-20th century. This is largely because of explicit representation of the lower NO_x emissions from biomass fuels in early periods, which combusts at lower temperatures as compared to coal. In 1970, CEDS NO_x emissions began to diverge from CMIP5 estimates, generally becoming larger due to waste, transportation, and energy sectors. CEDS emissions remain about 10% larger than those of CMIP5 in 1980 and 1990. Both global estimates increase and start to flatten around 1990. However, CEDS values flatten until 2000 and then increase again, while CMIP5 values decrease
275 from 1990 to 2000. IPCC also noted that differences in modelling protocol may have an effect.

In terms of ozone, many more models included a representation of chemistry in the stratosphere and these whole atmosphere schemes enabled the calculation of ERFs from stratospheric ODSs such as halocarbons and nitrous oxide (N₂O), and a more complete calculation of the ERFs from tropospheric ozone precursors, from which the production of ozone often extends into the lower stratosphere. AerChemMIP was not able to isolate an ERF due solely to ozone changes as the diagnosed ERFs
280 included changes in WMGHG (CH₄, N₂O, ODSs) and impacts on aerosols. In future MIPs, prescribed ozone experiments or methods for isolating ozone radiative effects might be needed (Collins et al., 2024).

The halocarbon ERFs diagnosed in AerChemMIP showed significant reductions compared to that expected from the direct greenhouse effect with a range of -0.18 to 0.32 W m⁻² (Thornhill et al., 2021b). This strong reduction in ERF compared to RF is partly attributed by Morgenstern et al. (2020) to negative cloud responses in the southern hemisphere. Moreover,
285 AerChemMIP models span a wide range of simulated ozone depletion. Morgenstern et al. (2020) found that there was a strong correlation between the modelled historical total ozone column (TOC) change and the halocarbon ERF. The observed TOC trend was used to generate an emergent constraint on the halocarbon ERF of -0.05 to 0.13 W m⁻², a much narrower range than from Thornhill et al. (2021b), but still with a chance of a negative ERF. Despite the uncertainty, largely the result of a large underestimate in a single model's ozone field, the claim that the Montreal Protocol had a positive climate benefit still holds.

As well as radiative diagnostics, the *piClim-X* experiments provide information on how the SLCFs influence atmospheric composition. These include impacts on ozone concentrations and the oxidants that destroy methane and thereby control its lifetime. The effect of N₂O on the methane lifetime (through depletion of tropical upper stratospheric ozone) is larger in the AerChemMIP models than had previously been estimated. This AerChemMIP result led to a slight reduction in the global warming potential (GWP) of N₂O as assessed in IPCC AR6 (Forster et al., 2021) although partially compensated by a positive
295 ozone forcing through the contribution of N₂O to tropospheric and lower stratospheric ozone production.

AerChemMIP also analysed the radiative effects of other emissions relevant to the Earth system: dust, sea salt, DMS, BVOCs, lightning NO_x, and fires. Since emissions of these species are generally prognostic within climate models, rather than prescribed, emissions were doubled compared to the *piClim-control* instead of prescribing 2014 emissions, as was done for anthropogenic emissions. For dust emissions, there was little agreement even on the sign of the ERF (Thornhill et al.,
300 2021a), reflecting the diverse dust emission rates across models, which are largely due to difference in the near-surface wind (Zhao et al., 2022), and diversity in simulated dust properties (Kok et al., 2021). For BVOC emissions, the models agreed that

the negative ERF from increased organic aerosols dominated over the positive contribution from increased ozone production. These experiments were used to derive radiative efficiencies (per mass emitted) for the natural species, and, in combination with the DECK abrupt-4xCO₂ experiment, to calculate climate feedback parameters ($\text{W m}^{-2} \text{K}^{-1}$) due to biogeochemical processes (Thornhill et al., 2021a). The dominant aerosol and chemistry feedbacks were found to be negative.

The quantification of ERFs by species is essential information in attributing climate change to the emissions of different pollutants that can be used by policy-makers to target mitigation of specific emissions. The results from AerChemMIP were used to derive the contributions of emissions of different species to the PI-to-PD forcing (Figure 6.12, Szopa et al., 2021a) and global surface temperature (Figure SPM2c) for IPCC AR6. This work also forms the basis for updates to climate change indicators (Forster et al., 2023).

Figure 3 shows the application of AerChemMIP *piClim-X* experiments in the attribution of drivers of historical climate change. AerChemMIP provided experiments and underpinning data for each species apart from the CO₂ forcing from CO₂ emissions, as indicated by the hatching. These data contribute to Figure SPM.2 of IPCC (2021).

Using *piClim-X* experiments, studies looked at the so-called “fast” circulation responses to aerosols, i.e., the responses that are independent of SST changes (e.g., Amiri-Farahani et al., 2020; Zanis et al., 2020). As expected, the fast PI-to-PD response to all aerosols included continental cooling, especially in the Northern Hemisphere, with the largest cooling over East Asia and India. Interestingly, however, multi-model mean Arctic winter warming occurred (albeit with large inter-model variability), consistent with warm air advection associated with intensification of the Icelandic Low and an anticyclonic anomaly over southeastern Europe (Zanis et al., 2020). The corresponding fast precipitation responses were largest in the tropics and generally associated with a precipitation decrease over continental regions, consistent with weakening of the monsoons of east Asia, Africa and the Americas (Zanis et al., 2020). Amiri-Farahani et al. (2020) used *piClim-2xfire* simulations to investigate the fast atmospheric circulation response to fire emissions, including anomalous ascent and upper-level divergence over the African continent. Previous analyses of idealized PDRMIP (Myhre et al., 2017) simulations have shown the utility of decomposing the climate responses into fast and slow components, particularly for precipitation. For example, the global mean fast precipitation response scales with the change in atmospheric absorption (e.g., due to black carbon) and the slow response scales with the change in surface temperature (Samset et al., 2016; Liu et al., 2018). A similar decomposition has also been used to understand precipitation responses to methane shortwave absorption under both idealized and realistic methane perturbations (Allen et al., 2023, 2024b).




Figure 3. Attributed change in near-surface temperature for 1750–2019 from emitted species. Data is replotted from Figure 6.12 of the IPCC AR6 Working Group 1 (Szopa et al., 2021b; Blichner and Berntsen, 2023). Assessments that were derived directly or indirectly from AerChemMIP experiments are shown unhatched.

3.2 AerChemMIP *histSST* experiments

330 The transient historical prescribed SST (*histSST*) experiments were designed principally to calculate transient ERFs. They were used to attribute historical changes in ERF to individual forcing agents, and in calculations of changes to the Earth's energy budget and integrated radiative forcing. Transient ERFs reveal more detail about historical changes in SLCFs. The magnitude and pattern of forcing can differ markedly from the ERFs calculated from *piClim* experiments, depending on the time period they are calculated for. An example of this is the aerosol radiative forcing, which peaks over North America and

335 Europe in the mid 1980s, and over Asia at the end of the simulation in 2014 (e.g. Kalisoras et al. (2023)). This means that ERFs calculated from *piClim* experiments have a spatial pattern that is more strongly influenced by Asia than most of the historical period, potentially causes issues for attribution if applied to periods before 1980. The *histSST* experiments also found wider application, with experiments such as *histSST-1950HC* and *histSST-piCH₄* allowing attribution of the effect of historical emissions and/or concentrations on atmospheric composition. Stevenson et al. (2020) used these experiments to

340 identify the drivers of hydroxyl (OH) change over the historical period, examining the effect of changing ODSs and ozone precursor emissions on methane levels, via changes to the methane sink, and the strength of methane chemical feedbacks. An analysis of the linearity of the total change in methane over the historical period vs. that in the individual attribution experiments indicated the potential role of climate change, higher global temperatures and the increase in OH derived from increased humidity, but their analysis suggests the utility of a separate experiment to identify the climate-driven, rather than

345 emissions-driven, changes in composition. Subsequently, Zeng et al. (2022) used the *histSST* experiments to determine the role of emissions changes on both stratospheric and tropospheric composition, focusing on the ozone response. In this case, *histSST-piN₂O* was crucial, despite being available in a smaller number of models, for assessing the role of long-lived GHGs, and changes to stratospheric temperatures, on ozone levels. The climate (i.e. CO₂)-driven change in ozone was calculated similar to Stevenson et al. (2020) as the residual between *histSST* and the sum of relevant *histSST* single- or multi-forcing

350 attribution experiments. The analysis by Zeng et al. (2022) contributed to the World Meteorological Organization (WMO) assessment of the recovery of stratospheric ozone [2023]. In CMIP6, O'Connor et al. (2021) identified the need for separate *histSST-piVOC* and *histSST-piNO_x* to disentangle the drivers of e.g., tropospheric ozone change, similar to *piClim-VOC* and *piClim-NO_x* (O'Connor et al., 2021). Experiments such as these are useful in transient experiments where the timing of emissions changes can be used to identify drivers of e.g., ozone production efficiency, against a changing set of emissions.

355 Figure 4 shows analysis of the ozone burdens in various AerChemMIP experiments (Griffiths et al., 2023) for four of the models employing online chemistry. Here the AerChemMIP experiments isolate the response to a consistent perturbation to the anthropogenic emissions of e.g., ozone precursors applied to each model. Data availability prevents a full comparison across each experiment, but there are data available for *histSST-piNTCF*, *histSST-1950HC* and *histSST* experiments for all four models, while three also feature *histSST-piO₃* and *histSST-piCH₄*. It can be seen that GFDL-ESM4 and MRI-ESM2-0

360 show similar responses to historical changes in concentrations of methane and halocarbon species, and that most models except UKESM1 show an increase in ozone due to historical emissions (via the NTCF experiment). While further work is required to

understand the origin of these differences, the figure highlights the usefulness of the idealised *histSST-piX* for understanding the origin of model diversity.




Figure 4. Ozone burdens in *histSST-X* AerChemMIP experiments taken from models using online chemistry. Burdens were calculated using the online tropopause as in Griffiths et al. (2021).

3.3 AerChemMIP *hist* experiments

Fully coupled transient simulations enable the impacts of SLCFs, aerosols, and halocarbons (*hist-piNTCF*, *-piAer*, and *-1950HC*) on surface temperature, the hydrological cycle, and atmospheric and oceanic circulation to be assessed. Although these experiments were Tier 1 and 2, six models performed the simulations in time to be used for the IPCC AR6 report. Ultimately, the modelling centres' contributions increased to ten models for *hist-piAer* and eleven for *hist-piNTCF*, but the ensemble size from participating models remains small, with a majority of centres only providing a single member for these experiments. Unfortunately, this is not sufficient for use in attribution studies of regional climate change, or of changes in the atmospheric or oceanic circulation. Compared to global/hemispheric scales, detecting and attributing a regional climate response is generally more difficult due to a smaller signal to noise (i.e., internal climate variability) ratio, and community uptake of the AerChemMIP simulations has been limited as a result. More ensemble members are available for some of the similar DAMIP experiments, such as *hist-aer*. However, the AerChemMIP experiments have been used to attribute global- and hemispheric-scale climate trends to SLCFs. Zhang et al. (2021a) showed that the common bias of CMIP6 ESMs overestimating the magnitude of mid-twentieth century cooling was primarily due to the higher aerosol burden in these models compared to their physical model counterparts. Using the difference between the historical and *hist-piAer* experiments, Zhang et al. (2021a) confirmed that the bias was driven by high aerosol burdens, rather than high sensitivities to aerosol forcing, as had previously been speculated. In a separate publication, Zhang et al. (2021b) used the same experiments to show that the dominant influence of anthropogenic aerosols on the terrestrial carbon sink is due to the increase in diffuse radiation from an increase in aerosol emissions leading to an increase in photosynthesis, rather than due to the aerosol influence on temperature, precipitation, or the amount of incident shortwave radiation at the surface.

The *hist-piAer* experiments have been used to confirm the main processes behind features of PD climate. Diamond et al. (2022) discuss the current asymmetry between northern hemisphere (NH) and southern hemisphere (SH) albedo, which they find to be a transient feature of global climate. The NH is more reflective in clear skies, but the SH is more cloudy. However, the difference in continental coverage between the hemispheres is offset by the larger extent of Antarctic ice compared to the Arctic,

so that PD asymmetry in clear-sky albedo is dominated by aerosol (confirmed by a comparison between the *historical* and *hist-piAer* experiments). The *hist-piNTCF* experiments have also been useful for investigating where emissions or processes do not play a major role in climate change. DeRepentigny et al. (2020) used the experiments to indicate that SLCFs are not important for the timing of the occurrence of an ice-free Arctic (in CESM2), or the deceleration of the rate of PD ice loss in their model. The initial occurrence of an ice-free Arctic in the near-future is instead primarily controlled by internal variability, while the rate of sea ice decline on longer timescales is determined by CO₂ concentrations. Zeng et al. (2022) use a comparison between the *histSST-piX* experiments and *hist-piNTCF* to confirm that coupling to an interactive ocean has little impact on simulated ozone trends, confirming the utility of fixed SST experiments for studying atmospheric composition. Similarity in methane lifetimes between the *historical* and *histSST* experiments was noted by Stevenson et al. (2020).

The ability to use the difference between the *historical* and *hist-piX* experiments to isolate the role of a set of forcings in climate trends is also useful for model evaluation. Moseid et al. (2020) evaluated global and regional trends in downwelling shortwave radiation at the surface between 1961 and 2014, comparing model output from the *historical* simulation to surface observations. CMIP6 models generally performed well compared to observations over Europe, but poorly over China. Using *hist-piAer*, Moseid et al. (2020) demonstrated that this was due to incorrect SO₂ emission trends over China.

3.4 AerChemMIP *ssp370* experiments

Allen et al. (2021) used the AOGCM *ssp370* and *ssp370-lowNTCF* experiments, together with the additional *ssp370-lowNTCFCH4* experiment, to investigate air quality benefits and climate impacts from SLCF mitigation. Coupled AOGCM experiments allowed the effect of SLCF on surface temperature and precipitation to be derived, finding significant perturbations to the hydrological cycle and highlighting the beneficial role of methane concentration reductions in counteracting the warming and wetting effects from aerosol reductions. Similar work by Shim et al. (2021) focuses on air quality in Asia, as does Li et al. (2022) which focuses on both air quality and climate in Asia. Zanis et al. (2022) used the *ssp370SST* and *ssp370pdSST* experiments to derive the change in surface ozone with increasing temperature, the “ozone-climate penalty,” showing an overall negative relationship between surface ozone and global temperature change, the result of large rates of ozone destruction over marine areas, with increases in ozone over the polluted regions of South and East Asia. Recently, the *ssp370* and related experiments have been used in studies examining the future burden of disease due to changing air quality (Akritidis et al., 2024; Turnock et al., 2023). The *o3ste* diagnostic output, a stratospheric ozone tracer intended to map stratosphere to troposphere exchange, was shown to be useful for identifying the role of circulation changes, particularly increased stratosphere-to-troposphere transport of ozone as the level of ODSs decrease in the future, on future ozone levels. Brown et al. (2022) used *ssp370SST* and *ssp370pdSST* to identify the emissions and chemistry drivers of the ozone-climate penalty in Africa and South America, including lightning NO_x changes, changes in the formation of NO_x reservoirs such as isoprene nitrate and peroxyacetyl nitrate (PAN), and temperature-driven changes in the emissions of BVOCs such as isoprene. A difference in the sign of the ozone-climate penalty was noted, depending on background NO_x levels, and highlighting the need for detailed chemical diagnostics when considering air quality/climate change interactions. The coupled AOGCM experiments were also used to investigate the effects of

SLCF mitigation on the AMOC, showing that future reductions in aerosol and ozone precursors alone induces end-of-century weakening of the AMOC, but this weakening is offset if methane reductions are applied (Hassan et al., 2022).

Figure 5 shows an illustration of how AerChemMIP data may be used for combined air quality/climate co-benefit studies. The figure shows that non-methane NTCF (NMNTCF) mitigation (aerosols and precursor gases only) improves air quality (both particulate matter with a diameter of less than 2.5 μm (PM2.5) and ozone), but at the expense of climate warming. Methane mitigation yields global cooling (a climate benefit) as well as improved ozone-related air quality with minor changes in PM2.5. All-NTCF mitigation (methane as well as aerosols and precursor gases) yields both a climate benefit (cooling, but less than that under methane mitigation) as well as an air quality benefit in terms of both PM2.5 and ozone.




Figure 5. Global annual mean 2090-2099 relative to 2014-2005 mitigation response scatterplots, based on analysis in the work of Allen et al. (2021). Surface temperature [K] versus surface (a) PM2.5 [$\mu\text{g m}^{-3}$] and (b) ozone [ppb] for five AerChemMIP models (as designated in the legend by symbols) and the corresponding multi-model mean (MMM) under non-methane near-term climate forcer mitigation (NMTNCF; black); all-NTCF mitigation (red); and methane-only mitigation (CH4; blue). Error bars represent the 95% confidence interval, estimated as twice the standard error. In the case of multiple realizations (UKESM1-0-LL, MRI-ESM2-0 and GISS-E2-G each performed 3 realizations per simulation), the symbol represents the model mean.

As mentioned above, AerChemMIP contributed to the development of the WGI IPCC AR6, being referenced in the Summary for Policymakers, Chapters 4, 6, 7, and the Atlas, as well as contributing important assessment/evaluation papers that underpin the Report. Furthermore, the AerChemMIP experiments allowed the attribution of historical surface temperature changes to composition changes by comparing the CMIP historical experiment to the AerChemMIP *hist-piAer* experiment, and to attribute radiative forcing to individual components, using 6 models (ESGF now shows near-surface temperature (tas) from 10 models).

The AerChemMIP data contributed to the development of emulators. The contributions to PD ozone forcing from CH₄, NO_x, N₂O, halocarbons, CO and VOCs, and climate, are derived from AerChemMIP *piClim-X* single forced experiments (Thornhill et al., 2021b, a) and the coupled CMIP6 *historical* experiment (Skeie et al., 2020). This relationship was used in AR6 Working Group 1 Chapter 7 to derive the historical ozone forcing time series. AerChemMIP also allowed an evaluation of the sensitivity of methane's chemical lifetime to reactive gases and climate (Thornhill et al., 2021a, b). The *histSST-piAer* experiment, along with RFMIP's *piClim-histaer*, allowed diagnosis of historical aerosol forcing from emissions of SO₂, BC and OC, which was used to construct time series of historical aerosol forcing (both ARI and ACI components) for AR6 (Smith et al., 2021). All of these emissions- and burden-derived relationships are now incorporated into the FaIR reduced-complexity climate model (Smith et al., 2018; Leach et al., 2021). Furthermore, methane's contribution to ozone forcing and the methane lifetime self-feedback factor were used in computation of emissions metrics for methane in AR6 such as GWP (Forster et al., 2021). The *ssp370-lowNTCF* simulations were also used to supplement ScenarioMIP and DAMIP simulations in the training of machine-learning based emulators participating in the ClimateBench benchmark dataset (Watson-Parris et al., 2022).

In summary, AerChemMIP provided significant advances, notably coupled transient experiments and the attribution of the role of SLCFs in radiative forcing and climate changes, enabled keystone analyses of air quality, and identification of the interactions of climate and air quality.

4 AerChemMIP – challenges and gaps

4.1 Timelines

The design of CMIP6, with no hard deadlines for data release, meant that modelling centres were free to deliver data as it became available. However, the IPCC Assessment Report timeline process was again very tight, making preparation for CMIP6 challenging. This was constrained further by the late release of the forcing datasets needed to perform simulations, and the cost of data processing to Climate Model Output Rewriter (CMOR) standards. Data came on-stream over the period 2019-2021, and, across AerChemMIP, only a quarter of the models eventually delivering data had done so by the end of June 2019, with three quarters delivering by the December 2019 submission deadline for papers to be included in AR6. With hindsight, it may have been better for key experiments to have been identified early and for the data to address assessment-relevant scientific questions to be available sooner.

Although the availability of model documentation and description was improved for CMIP6 relative to CMIP5 with the advent of ES-DOC, data for many models was published ahead of model documentation. This meant that it was difficult to use these models in process studies as modelling centres needed to be contacted directly to obtain information, e.g., which parameterization schemes had been used for aerosol microphysics.

4.2 Coordination across MIPs

In its preparation, AerChemMIP assumed extensive collaboration with DECK, the CMIP historical experiments, RFMIP, and ScenarioMIP. The *piControl*, *historical*, and *ssp370* experiments provided baselines for the AerChemMIP experiments, and
465 provided SST and sea-ice fields for experiments where these were prescribed, while complementary experiments for calculating ERFs for WMGHGs, land use, and natural forcing came from RFMIP.

DAMIP and AerChemMIP proposed similar coupled historical experiments, and closer integration with DAMIP's complementary *hist-X* experiment would have been beneficial. At the time of writing, six models have contributed data from both DAMIP *hist-Aer* and AerChemMIP *hist-piAer*. Given the demands on modelling centres, it was perhaps prohibitively costly
470 to ask for both variants (*hist-X* and *hist-piX*) for an attribution experiment, but it would certainly have been preferable to have both variants available. The *hist-piX* experiment design avoids the assumption of linearity that underpins the analysis of *hist-X* experiments. For many of the species considered by AerChemMIP, a degree of nonlinearity is expected in the climate response as the world warms, related to changing reaction rates or changes in cloud distribution and properties. However, for the quantification of the effect of such nonlinearities, these "everything but" style experiments need to be paired with single
475 forcing simulations, requiring better overlap between model participation in DAMIP and AerChemMIP. Diamond et al. (2022) used the AerChemMIP *hist-piAer* as a substitute for missing DAMIP *hist-GHG* experiments, in order to include UKESM1 and MPI-ESM-1-2-HAM in their investigation of delayed eastern equatorial Pacific warming, but without further analysis of the linearity of the response to changing aerosol emissions in a warming world it is difficult to know how sound this assumption is, especially as the degree of linearity is likely to be model-dependent (Simpson et al., 2023).

480 4.3 Coordinated variable request

There was a lack of consistent diagnostics across participating models in CMIP6, even for standard Tier 1 variables. This raised some challenges for analysis of the AerChemMIP data, effectively making a small (6-7 model) ensemble even smaller, and limiting the utility of the experiments. Consistent output variables over all models and all scenarios would have allowed larger ensembles to be used in analysis: models were sometimes rejected from studies for not including (Tier 1) variables of interest
485 (Griffiths et al., 2021).

SSP3-7.0, a pathway involving weak air quality control measures, was chosen as a future baseline by AerChemMIP. From a policy perspective, it is clear that diagnostic data from other SSPs with stronger air quality measures, such as SSP2-4.5 (middle of the road) and SSP1-1.9 (sustainability), would have been useful. AerChemMIP specified priority variables and their domain and frequency for the *historical* and *ssp370* experiments, and it was envisaged that a similar level of detail would be
490 provided by other centres for other ScenarioMIP experiments. Given the pressure on data processing and data archival, some centres omitted the AerChemMIP diagnostics from other ScenarioMIP experiments, prohibiting comparative analysis of e.g., chemical and aerosol processes. In particular, the inclusion of air quality diagnostics in the *ssp245* experiment would have allowed a more direct comparison between CMIP6 studies and those based around the ECLIPSE (Evaluating the Climate and Air Quality Impacts of Short-Lived Pollutants) experiments (Stohl et al., 2015) experiments, which use an RCP4.5 baseline.

495 Improved availability of air quality diagnostics over a wide range of future emission pathways would have enabled better understanding of the future interactions of climate change and air quality, and these diagnostics merit inclusion in a wider variety of scenarios by all centres in future MIP eras. At present, attribution of the drivers of air quality changes in future scenarios other than SSP3-7.0 in a multi-model sense is still lacking, due in part to the lack of data to perform such analyses, although Turnock et al. were able to perform analyses across the various SSPs for surface ozone and PM_{2.5} (Turnock et al., 500 2020).

4.4 Experiment design

AerChemMIP encouraged participating models to include interactive aerosol, and online tropospheric and stratospheric chemistry schemes, which meant that it was only possible for a small subset of CMIP6 models to perform all AerChemMIP experiments (Table 1). The bulk of AerChemMIP data comes from 11 models: BCC-ESM1, CESM2-WACCM, CNRM-ESM 505 (interactive stratosphere only), EC-Earth3-AerChem, GFDL-ESM4, GISS-E2, MIROC6-SPRINTARS, MPI-ESM1.2, MRI-ESM2, NorESM2 and UKESM1, of which MPI-ESM1.2 and NorESM2 use offline chemistry. Although similar to the number (seven) of models participating in the chemistry model intercomparison projects of Hemispheric Transport of Air Pollution (HTAP) Phase 2 and the Air Quality Model Evaluation International Initiative (AQMEII), this represents approximately 15% of the 70 models, and model variants, contributing to CMIP6. Within this small subset, there is large variance between model 510 results, and it becomes difficult to construct, and have faith in, reliable multi-model means. In this situation, identifying outliers also becomes challenging.

In addition to expanding the range of scenarios with AerChemMIP diagnostics, it is useful in each scenario to specify additional complementary experiments, similar to the *ssp370-pdSST* which was used as a complement to *ssp370SST* (Turnock et al., 2022), so as to be able to determine ozone-climate penalties, and to assess the linearity of the climate response. This is 515 similar to the RFMIP *piClim-histaerO3* (historical) experiment, but for the future. Additional experiments to address the role of methane in future climate and air quality are needed to more completely address the role of SLCFs in climate change.

4.5 Ensemble size

A single member of a coupled transient simulation is insufficient to identify where differences between models are due to differences in the response to forcing, as opposed to internal variability or other structural differences between the models. 520 AerChemMIP requested at least three ensemble members per coupled experiment. However, these were not always performed, with some centres submitting only one member per experiment. Unfortunately, this makes it impossible to identify forced responses in transient experiments with realistic forcing, and meant that while these simulations could be used for the AerChemMIP model ensemble as a whole, they could not be used to understand inter-model differences. Small ensemble sizes more generally present difficulties for the analysis of regional climate responses, which are key for AerChemMIP-related issues, such 525 as air quality, and also for climate extremes. Recent work by Fiore et al. (2022) used a 15 member initial condition ensemble to examine the role of ensemble size in simulating atmospheric composition trends and separating forced trends from internal variability, demonstrating that on multi-decadal timescales, the two are comparably important in some regions. Monerie

et al. (2021) examined the role of ensemble size in identifying regional precipitation trends, and concluded that 10 members represent a good balance between regional information and computational expense.

530 4.6 Diagnostics

There is a need for all MIPs to carefully review requested diagnostics for future MIPs to ensure that they are well-designed to address science goals, and that diagnostics are delivered in a standard format. For AerChemMIP, diagnostics relating to the ozone budget and tropopause (e.g., dynamical tropopause) were needed and there was a shortage of output for aerosol optical properties (Fiedler et al., 2024).

535 In the case of tropospheric ozone, ozone production and destruction rates, *o3prod* and *o3loss*, were specified in CMIP6 to cover only a portion of the ozone budget. They are, by definition, insufficient to infer stratospheric ozone input to the troposphere, as the budget closure cannot be guaranteed, making the quantitative equivalence of a residual budget term with stratospheric input impossible. This is especially true as models include additional chemistry, such as tropospheric halogen chemistry, that will strongly influence our interpretation of which reactions should be included in *o3prod* and *o3loss*. Further-
540 more, subsetting some reactions was in practice prone to human error in implementation. The CCMI diagnostic *do3chm*, the tendency due to chemistry, is strongly encouraged for future MIPs. Whereas production (P) and loss (L) terms isolated are preferable, operator tendencies (i.e., net $P - L$) are much more straightforward to code as a diagnostic and are therefore less prone to implementation errors, while still containing valuable information. The increasing adoption of whole-atmosphere on-line chemistry models, which allows a consistent treatment of ozone chemistry and removes the need for boundary conditions
545 or prescribed ozone fields, is a significant advantage.

New diagnostic output of WMO thermal tropopause height/pressure was available in CMIP6, which allowed the separate evaluation of tropospheric and stratospheric ozone. The tropopause introduces strong variations in stratospheric ozone column between models which are less pronounced when an ozonopause is used (as in CMIP5). For the diagnosis of stratosphere-troposphere transport, a diagnostic tropopause is probably more useful, e.g., a potential vorticity based or blended tropical/extra-
550 tropical tropopause definition. Tracer-tracer correlations are also useful here to understand downward transport, and synthetic tracers would add significant value, e.g., *e90* (Abalos et al., 2017; Prather et al., 2011). It may be advantageous to perform whole atmosphere evaluation and assessment in the future.

Given the difficulty in consistently defining the tropopause, it may be wise to consider assessment of model performance against e.g., total column ozone, and Earth Observation (EO) products that target the UT/LS region where the radiative forcing
555 of ozone is largest. Tropospheric ozone burden and column should be treated as less reliable quantities for intercomparison and assessment until such time as reliable diagnostic output is available, and their derivation from EO products is less problematic (Gaudel et al., 2018). It was regrettable that some centres did not provide tropopause output, preventing calculations of burdens and other useful diagnostics.

4.7 Emulators and impact assessments

560 Consideration of the experimental design for the development of flexible and comprehensive emulators, whether a physical reduced complexity model (Smith et al., 2018) or machine learning based (Watson-Parris et al., 2021), requires many similar considerations as for answering the fundamental questions addressed above, but also presents other challenges. For example, the use of idealised, single forcing experiments, to isolate individual contributions to radiative forcing and composition, can be very valuable in this setting either for training or validation. In order to ensure that the emulator is interpolating between
565 simulations rather than extrapolating beyond them (in emissions, concentration or forcing) it is valuable to have data points at the extrema of the relevant space. In this regard, very high emissions scenarios (such as SSP5-8.5), very ambitious scenarios (such as SSP1-1.9) and other corner cases (such as SSP3-7.0-lowNTCF) are extremely useful, regardless of their realism, as is participation in single-forcing experiments by a diversity of models. However, while such scenarios have very different global mean emissions, they do not explore very different spatial distributions of emissions, which are so important for aerosol and
570 other short lived chemical species. Sampling this space is the focus of the Regional Aerosol MIP (RAMIP, Wilcox et al. (2023)) which, with the aid of large ensembles, will ensure robust signals.

While AerChemMIP has shone a light on model diversity in e.g., aerosol forcing, sampling the process uncertainty explicitly within each model would provide valuable insights into the contribution to this spread from structural versus parametric uncertainty (Lee et al., 2011). While sampling full parametric uncertainty requires hundreds of simulations, simple scalings of
575 the aerosol indirect effect, for example, would allow useful determination of the role of such processes in inter-model diversity and persistent discrepancies with observations. Such ensembles can also be incorporated into the emulators described above to more fully capture model uncertainties, and potentially constrain them with energy budget considerations.

5 Planning and designing future aerosol and chemistry MIPs

So far we have considered the success of AerChemMIP's objectives and its support of wider CMIP goals, as well as identifying
580 some remaining areas for future study. In this section, we present some reflections by the AerChemMIP community on future CMIP and AerChemMIP activities, addressing this from the perspective of the the CMIP, sub-MIP and modelling centre level.

5.1 Coordination across MIPs

CMIP is comprised of various specialist sub-MIPs such as AerChemMIP. It is to be expected that new sub-MIPs will arise and evolve over time and may eventually be folded into the standard DECK or core experiments. The recent proposal of a 'Fast
585 Track' for CMIP7, incorporating experiments previously in AerChemMIP/RFMIP exemplifies this. The CMIP project brings essential early-stage planning and coordination for the community, provides crucial oversight and enables cross-cutting activities such as Fast Track, Task Teams for supporting CMIP activities, such as forcing updates, defining standard data requests and designing simulations. From our perspective, an important role for CMIP remains in providing oversight of and coordination between the individual participating MIPs. This coordination across MIPs is vital as each sub-MIP forms a part of the CMIP

590 landscape, and CMIP needs to ensure its underpinning goals and science questions are being addressed. It is also important to ensure that each sub-MIP integrates well with other MIPs, avoiding duplication of effort and enabling the best possible exploitation of modelling efforts.

Coordination is needed in several areas: firstly, by working with the community to define key science questions; secondly, by working with sub-MIPs to define key experiments and required analyses; thirdly, by identifying priority variables, such as the IPCC priority variable list, which is also useful for prioritising data processing and availability; lastly, by coordinating 595 experiments, protocols and diagnostic output across modelling centres to standardise data delivery. CMIP can also coordinate the community review that is critical to ensure experiments meet community needs, address open science questions and achieve buy in from modelling centres e.g., covidMIP where specific science questions motivated quick turnaround.

The task of model evaluation, which aims to build and improve confidence in climate model projections, is an important 600 part of CMIP activities. While AR6 did not in this cycle feature a chapter devoted to the evaluation of climate models, instead moving this work within the individual chapters as required, the importance of assessing and evaluating the components of individual climate models and the overall performance of ESMs remains clear. Future versions of CMIP are expected to continue to assess model performance and to quantify the causes of the diversity in model projections. The preceding phases of CMIP show that the progressive evolution of climate models and model capability necessarily changes what evaluation is 605 possible, how this assessment should be done and modifies the evaluation and assessment requirements and metrics. Effective model evaluation requires the supporting MIPs to be mindful of the progress of the state of the field and it should be expected that the evaluation activities may change further over time. As ESMs become increasingly complex, understanding sources of intermodel diversity requires more effort as there are more processes included and the coupling between them is likely stronger. Additionally, as ESMs evolve the structural differences between models may play an increasing role in driving intermodel 610 differences. As model complexity and the treatment of feedbacks increases, understanding and attributing differences becomes more challenging and more important.

In addition to model evaluation, an important goal of CMIP is to understand the evolution of ESMs. This requires some consistency in experimental design and diagnostic data across the various phases of CMIP. The use of Digital Object Identifiers associated with climate model datasets through CMIP6 is a welcome step forward. In CMIP6, the adoption of CF-compliant 615 formats and the use of CMOR functions to reprocess model output allowed the archiving of consistent output between models and a better interface to evaluation code such as ESMValTool (Schlund et al., 2023). ESMValTool and other model evaluation frameworks such as PCMDI (Lee et al., 2024) provide an important piece of infrastructure for model evaluation, and it would be helpful for the community to standardize further around these tools, as it enables the distribution of model evaluation methods, multi-model comparison and traceability of model performance across MIP eras. In this regard, researchers may wish to port 620 their evaluation scripts from the CMIP6 Github repository to ESMValTool and other standardised evaluation platforms. Use of ESMValTool or similar models during model development would provide a traceable picture of model evolution between release versions.

It is essential for model evaluation, intercomparison, and process studies that a single and, above all, comprehensive source of information on MIPs, models, and simulations is available. The ES-DOC service provided by CMIP (<https://search.es->

625 doc.org/) has proved to be very valuable. The ES-DOC format was initiated in the fifth phase of CMIP as a metadata repository to provide such information (Guilyardi et al., 2013), and was then extended for use in CMIP6 (Balaji et al., 2018; Pascoe et al., 2020). ES-DOC improves our understanding of model data, increases the value of data for use in the future, and, by making earlier work more findable, potentially minimizes the need to re-run models. In AerChemMIP, ES-DOC was used to successfully document new model simulations (e.g., ssp370-lowNTCFCH4; Allen et al. (2021)) that were proposed after the publication of the AerChemMIP protocol paper (Collins et al., 2017). The Errata system (<https://errata.es-doc.org/static/index.html>) also worked well for reporting errors in the simulations. For example, when UKESM1 atmosphere-only simulations were found to have a bug, an issue notice was raised to document that data and the relevant experiments were to be withdrawn from the ESGF and replaced. While ES-DOC also aimed to provide standard information on models, this was less successful. In some cases, model information provided by modelling centres did not appear on ES-DOC. In other cases, the information available was not sufficiently detailed from an AerChemMIP perspective, was available but difficult to find, or incorrect. It would be beneficial, for example, to have standard and more consistent information on the chemistry/aerosol schemes used, and to connect better to the individual model description and evaluation papers. In general, high quality, useful, and perhaps even overly explicit, model description papers are required. Although there were improvements over CMIP5, it was still unfortunately common in AR6 for model description papers to appear after the data had been uploaded to ESGF, leaving a gap in information when preparing multi-model assessments, identifying outliers, and generating high-confidence projections. Where model components were common across several generations of a model, even basic details about this component tended to be omitted from the description paper of the CMIP6 model version. While including such information would make a description paper cumbersome and cause problems with plagiarism checks by journals, such details are important for process studies, and for weighting multi-model ensembles to avoid dominance by a particular model family and closely related models, and would be a valuable addition to ES-DOC.

Improving the search facility (e.g., by component, or by process), making the questionnaire provided to modelling centres less opaque, making the repository straightforward and accessible to correct or update, and better communication with modelling centres and MIPs would all be beneficial. ES-DOC is being re-visited for CMIP7 and the new CMIP International Project Office (IPO) will help to provide a forum for improved communication. Together, these improvements should facilitate multi-model assessments and decrease the burden on centres responding to the questionnaire or to clarifying questions by scientists involved in model evaluation. However, sufficient resources will be required to overcome the technical challenges identified and to fully meet user needs.

To align with the principles of findability, accessibility, interoperability, and reusability (FAIR), it may ultimately be necessary for ESGF or similar repositories to consider archiving source code. It may be helpful if model description papers could feature a minimum, standardised set of information, and it may be necessary to make model descriptions machine-readable or to expand the ES-DOC requirements. Trawling the model literature for intercomparison/process papers is difficult and time-consuming and leads to large amounts of duplicated effort.

5.2 The AerChemMIP project

In preparing for the next phase of an Aerosols and Chemistry MIP, it is expected that the underpinning science questions will be reviewed along with the criteria for participating models, the experiment designs and the diagnostic data request. It is however envisaged that the next phase of CMIP will feature a second phase of AerChemMIP, with the focus remaining on the role of aerosols and chemistry in the Earth system and climate change.

It will be necessary for ScenarioMIP, AerChemMIP and CMIP to coordinate in defining future scenarios and to define trajectories for SLCFs in support of the CMIP science questions. For AerChemMIP, SSP3-7.0 was chosen as the future baseline and provided high signal-to-noise in counterfactual experiments involving strong air quality interventions. Recent work has highlighted that focus on a single scenario may limit the usefulness of climate change projections in impact assessments. This is particularly relevant in the context of SLCFs, as future aerosol/precursor emissions scenarios span a large range, including minimal changes through the entire 21st century (e.g., SSP3-7.0) to rapid reductions over the next three decades (e.g., SSP1-1.9) that are comparable to the growth of emissions over the entire industrial era (e.g., Persad et al., 2023). In addition to these global emissions differences, we also note the importance of large diversity in regional SLCF emissions.

In light of the CMIP7 commitment to halve its carbon emissions relative to CMIP6, the ensemble size and the volume of output diagnostics needs to be considered alongside other sources of experimental cost such as model resolution and the model complexity (Fiedler et al., 2024). In this light, for future MIPs it may be fruitful to revisit the goals of experiments and intended analyses and energy/storage requirements.

For diagnosing ERFs, *piClim-X* timeslice experiments are required, and for understanding climate responses, coupled atmosphere-ocean *hist-piX* experiments should be retained. Transient ERFs can be calculated from the *histSST-piX* experiments. Combined experiments targeting SLCFs are clearly beneficial, although additional single forcing experiments are useful for understanding drivers. Clearly some tradeoff and accommodations need to be made in a MIP addressing both reactive gases and aerosols - *piNTCF* and *piAer* experiments, *piNOx* and *piVOC* experiments have all been useful but any expansion in experiment number should be considered in light of the stated aim to keep the CMIP computational expense to a minimum. Diagnosis of model sensitivities and response to forcings, performed in AerChemMIP with single-forcing experiments, may be possible with an expanded set of diagnostics rather than these dedicated attribution experiments. This should be considered.

It may be necessary to consider experiment design and the participating models in tandem. As noted above, the number of ESMs participating in AerChemMIP using online chemistry may be a concern in the future in light of the decreased participation in CMIP6. In terms of representation of aerosol, there was a wide range of complexity in the treatment of aerosol and aerosol-cloud interactions in CMIP6, but the representation of aerosol processes was generally not a barrier to participation in AerChemMIP. However, only a small number of ESMs featured online chemistry: ACCMIP, for CMIP5, featured 15 atmospheric chemistry models in its assessment of tropospheric ozone, while only five models were able to be used in CMIP6 ozone assessments. The low number of online chemistry models introduced challenges in evaluation and robustly identifying outliers. Future CMIP experiments aiming to address the role of aerosols and reactive trace gases such as AerChemMIP should aim to achieve greater participation, perhaps beyond ESMs, across the modelling community, and to reverse the declining trend in

participation. For the purposes of AerChemMIP, it may be beneficial to consider if models not meeting the DECK entry-card could be included as they are a valuable additional resource for understanding the origins of inter-model diversity. In ACCMIP, chemical transport models (CTMs) were included using time-slice approaches and/or offline meteorology. A future chemistry-focused MIP remains an attractive prospect, and has been the subject of recent discussions (Archibald et al., 2022). This should focus on both PI and PD conditions, with the objective of understanding the sources of model spread in both periods, and in quantifying model skill. In evaluating model skill, observations are essential making the AMIP DECK period of 1979-2015, designed to cover the post-satellite era, the most valuable.

Models of Intermediate Complexity could be useful for longer transient experiments and idealised forcings. As we discuss above, there is a need to include more processes, both to assess the role of processes missing from ESMs and to understand how structural differences impact future predictions. For this purpose, experiments over limited periods of the historical or future periods may be useful, particularly if coupled with higher quality (process-level or time resolution) diagnostics. The value from these experiments would be amplified if all centres/models chose the same period e.g., 2050s. This also reduces additional storage and processing overheads, and makes it easier to re-run models.

CMIP6 showed that evaluation and assessment of coupled ESMs is an increasing challenge, particularly for atmospheric composition. In planning for future phases of AerChemMIP, the presence of ESMs featuring online atmospheric chemistry, the increasing use of online components for natural or biogenic sources of ozone and aerosol precursors is to be expected and encouraged, as they provide more realistic treatment of Earth system feedbacks. Certainly, in CMIP6, interactive descriptions of ocean biogeochemistry, land surface feedbacks and emissions of biogenic species were increasingly common. Moreover, CMIP6 showed that the inter-model range of natural/biogenic sources of ozone and aerosol precursors is now large, and is a key driver in intermodel diversity, particularly in the PI period, both in ozone, sinks for methane such as OH, and, as BVOCs can also oxidize to form secondary organic aerosol (SOA), the large range in BVOC emissions also contributes to intermodel diversity in SOA. Future projections of atmospheric composition using ESMs will depend sensitively on the response of these natural emissions, and other processes, to climate. The AerChemMIP *piClim-2x* experiments which target natural sources of SLCFs will be useful here, and should be expanded as required, for instance if online methane or fire-related emissions become standard. Expanding these types of simulations to coupled transient experiments would be useful and would allow insight on the climate impacts associated with dust, fires, sea salt, etc. in a multi-model context. It will also be essential to perform intercomparison and evaluation of these natural emissions, and the response to climate change of emissions such as NO_x from natural sources such as lightning (Finney et al., 2016; Murray et al., 2013), or wetland methane emissions, will need further investigation, and may merit separate intermodel comparison exercises, such as done for wetland CH₄ emissions (WETCHIMP, Melton et al., 2013).

AerChemMIP provided three tiers of experiments, and various sets of experiments across the period 1850-2100. Practically, it was found that not all experiments required the same effort to set up and process, with variants of experiments, e.g., *piClim-SO2* and *piClim-NOx*, requiring relatively less effort to set up than e.g., *ssp370pdSST* and *ssp370-lowNTCF*, with the longer transient experiments requiring significantly more supervision during execution compared to the time slice experiments. Within centres, experiments motivated by new and targeted science questions clearly received significant additional effort. Incorpor-

rating these ‘bottom-up’ designs of experiments, and making clear how proposed experiments have the ability to address and respond to current and emerging science questions will be beneficial to future MIPs.

It may also be more fruitful to produce mid- to large-ensembles for a smaller number of experiments, than a large number of experiments with a small (<5 member) number of ensembles. As mentioned above, it would be advantageous to develop new MIP experiments in tandem with the design for their diagnostic output required for their analysis, and also to provide criteria for verification of required output. This is a critical point to avoid missing diagnostics or experiments, potentially limiting the usefulness of experiments and reducing the value of potentially costly experiments. As an example, attribution of dynamical responses in AerChemMIP coupled transient experiments is often a challenge due to small ensemble size - as the number of ensemble members increases, it becomes easier to distinguish weak signals and the effects of structural uncertainty from internal model variability. The viable ensemble size for analysis needs to be considered both when designing and performing experiments. AerChemMIP requested three ensembles for each of the coupled transient *hist-piX* experiments, but these were not delivered by all participating models. It is now clear that a larger ensemble size is required to characterise regional climate responses, especially regional precipitation changes (Monerie et al., 2021). With this in mind, MIPs focussed on attribution typically request larger ensembles: DAMIP requested five members per experiment for CMIP6, and the RAMIP requested ten. However, if the focus is on composition and/or forcing, fewer experiments are required, although timeslice experiments longer than 30 years are required for many species.

Given the size of the ensemble in the historical and SSP simulations used as the AerChemMIP baselines, there is an opportunity to expand the AerChemMIP ensemble member size, so that the experiments provide clearer climate information at spatial and temporal scales where internal variability is large, and so that more robust conclusions about the role of model structural uncertainty can be drawn.

To verify the presence of required output, an AerChemMIP variable request, i.e. a list of required diagnostics, may be useful. This should additionally list the analyses that they underpin. For instance, it may be necessary to document a consistent method for generation of PM_{2.5} concentrations, or to specify which species are necessary to be output at high time resolution, such as planetary boundary layer height and dry deposition fluxes, for as full an understanding as possible of future air pollution and its drivers. One option may be to ensure better coordination between ScenarioMIP and AerChemMIP to ensure that the AerChemMIP diagnostic data request is present in all ScenarioMIP experiments, for at least five ensemble members, ideally more.

5.3 Modelling centres

In understanding model evolution, traceability is essential. Finally, we encourage modelling centres to use their model description papers to document the differences/changes with respect to an existing model or model description paper. Understanding how models have changed also requires the availability of codebases to interested researchers.

It is also necessary for modelling centres to document as clearly as possible the origin of variant data. A standard approach for identifying model variants needs to be adopted for CMIP7. This was inconsistent in CMIP6, resulting in model variants being used incorrectly by authors. CMIP naming conventions support the identification of model variants through use of a

physics code in ensemble member names: the 'p1' in 'r1i1p1f1', for example. For CMIP6, there was a burden on the user to establish what these codes meant, and how the data should be treated to take this into account, as conventions differed between centres. For example, p1 and p2 variants of CanESM5 included stochastic perturbations and could be combined into a single CanESM5 ensemble. However, for GISS-E2, the p codes indicate the use of different aerosol and chemistry schemes, and these
765 variants should be treated as different models. When models variants are using different modules, this would be better reflected in the use of different model names rather than different physics versions, which was the widely adopted approach to indicating different model resolution, for example, in CMIP6 (e.g., NorESM2-LM vs. NorESM2-MM).

6 Conclusions

The AerChemMIP Project, endorsed by CMIP6, has led to significant progress in our understanding of the role of aerosols and
770 reactive gases in the climate system, both from a historical perspective and extending out into the future, and has worked well alongside PDRMIP and RFMIP.

The design of AerChemMIP focused on the effect of composition changes. Radiative forcing was calculated using a comparison between perturbation and control atmosphere-only timeslice experiments, a protocol common with RFMIP. The role of historical emissions changes was examined in "all-but-one" transient atmosphere-only attribution experiments allowing radiative
775 forcing to be calculated and the role of the drivers of composition changes to be deduced. Counterfactual coupled transient atmosphere-ocean experiments produced important data on the climate response to historical emissions. Transient experiments investigating the future SSP3-7.0 pathway allowed insight into the role of SLCFs in future climate and air quality.

The resulting literature based on AerChemMIP compares very well against questions A1 ('How have anthropogenic emissions contributed to global radiative forcing and affected regional climate over the historical period?' and A2 ('How might
780 future policies (on climate, air quality and land use) affect the abundances of SLCFs and their climate impacts?') with the *piClim-X* in particular being targeted at A1, and *ssp370* and *ssp370-lowNTCF* experiments targeting A2. The effect on global climate (A1/A2) was well-characterised (e.g., Allen et al., 2020; Thornhill et al., 2021a, b; Allen et al., 2021) but, as we note above, the coupled transient experiments were generally only performed with small ensemble sizes, which limited their scope.

AerChemMIP performed perhaps less well against A3 ('How do uncertainties in historical SLCF emissions affect radiative
785 forcing estimates?'), at least in time for AR6. The original aim in Collins et al. (2017) was to scale the ERFs from all the *piClim* experiments with the emission uncertainty to quantify the contribution from emission uncertainty to the NTCF forcing uncertainty. This task is still feasible and could be identified as a remaining task for future MIPs. Some effort was made to bound the effect of emissions changes through the *ssp370-lowNTCF*. In future MIPs, uncertainties in anthropogenic emissions estimates, which were not provided for CMIP6, would be a welcome addition for this task, although the size of such a task
790 is certainly daunting, particularly in coupled experiments. However, there is work being done to look into the sensitivity of models to uncertainties in emissions (e.g., Booth et al., 2018; Fyfe et al., 2021; Ahsan et al., 2023; Holland et al., 2024).

AerChemMIP performed well against the objectives of question A4 ('A4. How important are climate feedbacks to natural NTCF emissions, atmospheric composition, and radiative effects?') via the *piClim-2x* experiments. However, the use of the

795 DECK *abrupt-4xCO2* and *piControl* simulations to quantify how these natural emissions change with climate needs to be improved. In particular, there is a need to separate the radiative effects of CO₂ from the biophysical effects of CO₂ in any new experiments in a consistent way, due to the impact of future CO₂ on biogeochemical feedbacks (e.g., Arora et al., 2020; Allen et al., 2024a).

800 The AerChemMIP contributions to AR6 were mainly through Chapter 6 ("Short-lived Climate Forcers", Szopa et al., 2021b) and Chapter 7 ("The Earth's Energy Budget, Climate Feedbacks and Climate Sensitivity", Forster et al., 2021). For example, the *piClim-X* experiments were instrumental in being able to construct the emission-based forcing bar chart and (figures 6.12, TS.15) and from these to drive attributions of the historical temperature rise (figures TS.15 and SPM.2). Such contributions came both through the AerChemMIP specific experiments and through the AerChemMIP specific diagnostics. For instance, the historical ozone RF was diagnosed from the historical simulations, using the ozone mixing ratio on model levels: a diagnostic requested by AerChemMIP.

805 AerChemMIP contributed not just to AR6 but to ongoing research efforts. The database of experiments is rich, both in terms of experiments, participating models and the sophistication of the treatment of chemistry and aerosols and their role in the climate system. These data form a part of the climate data landscape and are enabling new analyses to be performed. We hope that this legacy of AerChemMIP also enables future work on the role of aerosols and short-lived reactive gases in the climate system.

Model	Centre	Model components						ES-DOC		References	
		Atmosphere	Aerosol	Chemistry	Ocean	Sea Ice	Land	Aerosol	Chem	Data	Documentation
ACCESS-CM2	CSIRO-ARCCSS	MetUM-HadGEM3-GA7.1	UKCA-GLOMAP-mode		ACCESS-OM2 (GFDL-MOM5)	CICE5.1.2	CABLE2.5	N		Dix et al. (2019)	Bi et al. (2020)
ACCESS-ESM1-5	CSIRO	HadGAM2	CLASSIC (v1.0)		ACCESS-OM2 (GFDL-MOM5)	CICE4.1	CABLE2.4	Y		Ziehn et al. (2019)	Ziehn et al. (2020)
AWI-ESM-1-1-LR	AWI	ECHAM6.3.04p1	MACv2-SP		FESOM 1.4	FESOM 1.4	JSBACH 3.20 with dynamic vegetation	N		Danek et al. (2020)	Sidorenko et al. (2015)
BCC-CSM2-MR	BCC	BCC_AGCM3_MR	MACv2-SP		MOM4	SIS2	BCC_AVIM2	N		Wu et al. (2018)	Wu et al. (2019)
BCC-ESM1	BCC	BCC_AGCM3_LR		BCC-AGCM3-Chem	MOM4	SIS2	BCC_AVIM2	N	N	Zhang et al. (2018)	Wu et al. (2020)
CAMS-CSM1-0	CAMS	ECHAM5_CAMS			MOM4	SIS 1.0	CoLM 1.0	N	N	Rong (2019)	Chen et al. (2019), Rong et al. (2018)
CanESM5	CCCma	CanAM5	Interactive	Specified oxidants	NEMO3.4.1	LIM2	CLASS3.6-CTEM1.2	Y		Swart et al. (2019a)	Swart et al. (2019b)
CAS-ESM2-0	CAS	IAP AGCM 5.0	IAP AACM	IAP AACM	LICOM2.0	CICE4	CoLM	N	N	Chen et al. (2019)	Zhang et al. (2020)
CESM2-FV2	NCAR	CAM6	MAM4	MAM4	POP2	CICE5.1	CLM5	N	N	Danabasoglu (2019a)	Danabasoglu et al. (2020)
CESM2	NCAR	CAM6	MAM4	MAM4	POP2	CICE5.1	CLM5	Y	Y	Danabasoglu (2019d)	Danabasoglu et al. (2020)
CESM2-WACCM-FV2	NCAR	WACCM6	MAM4	MAM4	POP2	CICE5.1	CLM5	N	N	Danabasoglu (2019b)	Danabasoglu et al. (2020)
CESM2-WACCM	NCAR	WACCM6	MAM4	MAM4	POP2	CICE5.1	CLM5	N	N	Danabasoglu (2019c)	Danabasoglu et al. (2020)
CIesm	THU	CIesm-AM	MAM4	trop_mam4	CIesm-OM	CICE4	CIesm-LM	N	N	Huang (2019)	Lin et al. (2020)
CMCC-CM2-HR4	CMCC	CAM4	MACv2-SP		NEMO3.6	CICE4.0	CLM4.5	N		Scoccimarro et al. (2020)	Cherchi et al. (2019)
CMCC-CM2-SR5	CMCC	CAM5.3	MAM3		NEMO3.6	CICE4.0	CLM4.5	Y		Lovato and Peano (2020)	Cherchi et al. (2019)
CNRM-CM6-1	CNRM-CERFACS	Arpege 6.3	Prescribed monthly fields computed by TAC-TIC_v2	OZL_v2	NEMO3.6	Gelato 6.1	Surfex 8.0c	N	N	Voldoire (2018)	Voldoire et al. (2019)
CNRM-ESM2-1	CNRM-CERFACS	Arpege 6.3	TACTIC_v2	REPROBUS-C_v2	NEMO3.6	Gelato 6.1	Surfex 8.0c	Y	Y	Seferian (2018)	S��ferian et al. (2019)
E3SM-1-0	E3SM-Project	EAM v1.0	MAM4	Troposphere specified oxidants for aerosols. Stratosphere linearized interactive ozone (LINOZ v2)	MPAS-Ocean v6.0	MPAS-Seaice v6.0	ELM v1.0	N	N	Bader et al. (2019a)	Golaz et al. (2019)
E3SM-1-1	E3SM-Project	EAM v1.1	MAM4	Troposphere specified oxidants for aerosols. Stratosphere linearized interactive ozone (LINOZ v2)	MPAS-Ocean v6.0	MPAS-Seaice v6.0	ELM v1.1	N	N	Bader et al. (2019b)	Burrows et al. (2020)
EC-Earth3-AerChem	EC-Earth-Consortium	IFS cy36r4	TM5	TM5	NEMO3.6	LIM3	HTESSEL	Y	Y	EC-Earth (2020)	van Noije et al. (2021)
EC-Earth3	EC-Earth-Consortium	IFS cy36r4	MACv2-SP	TM5	NEMO3.6	LIM3	HTESSEL	N	N	EC-Earth (2019b)	D��scher et al. (2021)
EC-Earth3-Veg	EC-Earth-Consortium	IFS cy36r4	MACv2-SP	TM5	NEMO3.6	LIM3	LPJ-GUESS v4	N	N	EC-Earth (2019a)	D��scher et al. (2021)
FGOALS-f3-L	CAS	FAMIL2.2	MACv2-SP		LICOM3.0	CICE4.0	CLM4.0	N		YU (2019)	HE et al. (2020)
FGOALS-g3	CAS	GAMIL3	MACv2-SP		LICOM3.0	CICE4.0	CAS-LSM	N		Li (2019)	Li et al. (2020)
FIO-ESM-2-0	FIO-QLNM	CAM4	Prescribed monthly fields		POP2-W (POP2 coupled with MASNUM surface wave model)	CICE4.0	CLM4.0	N		Song et al. (2019)	Bao et al. (2020)

Continued on next page

GFDL-CM4	NOAA-GFDL	GFDL-AM4.0.1	Interactive	fast chemistry, aerosol only	GFDL-OM4p25 (GFDL-MOM6)	GFDL-SIM4p25 (GFDL-SIS2.0)	GFDL-LM4.0.1	Y	N	Guo et al. (2018)	Held et al. (2019)
GFDL-ESM4	NOAA-GFDL	GFDL-AM4.1	Interactive	GFDL-ATMCHEM4.1	GFDL-OM4p5	GFDL-SIM4p5	GFDL-LM4.1	Y	Y	Krasting et al. (2018)	Dunne et al. (2020)
GISS-E2-1-G-CC	NASA-GISS	GISS-E2.1	Varies with physics version	Varies with physics version	GISS Ocean	GISS SI	GISS LSM	N	N	NASA/GISS (2019a)	Kelley et al. (2020), Miller et al. (2021)
GISS-E2-1-G (p1)	NASA-GISS	GISS-E2.1	'none'	Non-interactive	GISS Ocean	GISS SI	GISS LSM	N		NASA/GISS (2018)	Kelley et al. (2020), Miller et al. (2021)
GISS-E2-1-H (p1)	NASA-GISS	GISS-E2.1	'none'	Non-interactive	HYCOM Ocean	GISS SI	GISS LSM	N		NASA/GISS (2019b)	Kelley et al. (2020), Miller et al. (2021)
GISS-E2-1-G (p3)	NASA-GISS	GISS-E2.1	OMA	GPUCCINI	GISS Ocean	GIS SI	GISS LSM	N	N	NASA/GISS (2018)	Kelley et al. (2020), Miller et al. (2021)
GISS-E2-1-H (p3)	NASA-GISS	GISS-E2.1	OMA	GPUCCINI	HYCOM Ocean	GISS SI	GISS LSM	N	N	NASA/GISS (2019b)	Kelley et al. (2020), Miller et al. (2021)
GISS-E2-1-G (p5)	NASA-GISS	GISS-E2.1	MATRIX	GPUCCINI	GISS Ocean	GISS SI	GISS LSM	N	N	NASA/GISS (2018)	Kelley et al. (2020), Miller et al. (2021)
GISS-E2-1-H (p5)	NASA-GISS	GISS-E2.1	MATRIX	GPUCCINI	HYCOM Ocean	GISS SI	GISS LSM	N	N	NASA/GISS (2019b)	Kelley et al. (2020), Miller et al. (2021)
HadGEM3-GC31-LL	MOHC	MetUM-HadGEM3-GA7.1	UKCA-GLOMAP-mode		NEMO-HadGEM3-GO6.0	CICE-HadGEM3-GS18	JULES-HadGEM3-GL7.1	N	N	Ridley et al. (2019a)	Andrews et al. (2020), Kuhlbrodt et al. (2018)
HadGEM3-GC31-MM	MOHC	MetUM-HadGEM3-GA7.1	UKCA-GLOMAP-mode		NEMO-HadGEM3-GO6.0	CICE-HadGEM3-GS18	JULES-HadGEM3-GL7.1	N	N	Ridley et al. (2019b)	Andrews et al. (2020)
INM-CM4-8	INM	INM-AM4-8	INM-AER1		INM-OM5	INM-ICE1	INM-LND1	N		Volodin et al. (2019a)	Volodin et al. (2010)
INM-CM5-0	INM	INM-AM5-0	INM-AER1		INM-OM5	INM-ICE1	INM-LND1	N		Volodin et al. (2019b)	Volodin et al. (2017), Volodin and Kostykin (2016)
IPSL-CM6A-LR	IPSL	LMDZ	INCA v6 AER		NEMO-OPA	NEMO-LIM3	ORCHIDEE (v2.0)	N		Boucher et al. (2021)	Boucher et al. (2020)
KACE-1-0-G	NIMS-KMA	MetUM-HadGEM3-GA7.1	UKCA-GLOMAP-mode		MOM4p1	CICE-HadGEM3-GS18	JULES-HadGEM3-GL7.1	N		Byun et al. (2019)	Lee et al. (2020a)
KIOST-ESM	KIOST	GFDL-AM2.0	MACv2-SP	Simple carbon aerosol model (emission type)	MOM5.0	SIS	CLM4	N	N	Kim et al. (2019)	Pak et al. (2021)
MIROC6	MIROC	CCSR AGCM	SPRINTARS6.0		COCO4.9	COCO4.9	MATSIRO6.0	Y		Tatebe and Watanabe (2018)	Tatebe et al. (2019)
MIROC-ES2L	MIROC	CCSR AGCM	SPRINTARS6.0		COCO4.9	COCO4.9	MATSIRO6.0 + VISIT-e ver.1.0	N		Hajima et al. (2019)	Hajima et al. (2020)
MPI-ESM-1-2-HAM	HAMMOZ-Consortium	ECHAM6.3	HAM2.3	sulfur chemistry	MPIOM1.63	thermodynamic (Semtner zero-layer) dynamic (Hibler 79) sea ice model	JSBACH 3.20	Y		Neubauer et al. (2019)	Mauritsen et al. (2019)
MPI-ESM1-2-HR	MPI-M	ECHAM6.2	MACv2-SP		MPIOM1.63	thermodynamic (Semtner zero-layer) dynamic (Hibler 79) sea ice model	JSBACH3.20	N		Jungclaus et al. (2019)	Mauritsen et al. (2019), Stevens et al. (2013)
MPI-ESM1-2-LR	MPI-M	ECHAM6.3	MACv2-SP		MPIOM1.63	thermodynamic	JSBACH3.20	N		Wieners et al. (2019)	Wieners et al. (2019), Stevens et al. (2013)

Continued on next page

						(Semtner zero-layer) dynamic (Hibler 79) sea ice model					
MRI-ESM2-0	MRI	MRI-AGCM3.5	MASINGAR mk2r4	MRI-CCM2.1	MRI.COM4.4	MRI.COM4.4	HAL 1.0	Y	Y	Yukimoto et al. (2019)	YUKIMOTO et al. (2019)
NESM3	NUIST	ECHAM v6.3	MACv2-SP		NEMO v3.4	CICE4.1	JSBACH v3.1	N		Cao and Wang (2019)	Cao et al. (2018)
NorCPM1	NCC	CAM-OSLO4.1	OsloAero4.1	OsloChemSimp4.1	MICOM1.1	CICE4	CLM4	Y	Y	Bethke et al. (2019)	Bethke et al. (2021)
NorESM2-LM	NCC	CAM-OSLO	OsloAero	OsloChemSimp	MICOM	CICE	CLM	Y	Y	Seland et al. (2019)	Seland et al. (2020)
NorESM2-MM	NCC	CAM-OSLO	OsloAero	OsloChemSimp	MICOM	CICE	CLM	Y	Y	Bentsen et al. (2019)	Seland et al. (2020)
SAM0-UNICON	SNU	CAM5.3 with UNI- CON	MAM3		POP2	CICE4.0	CLM4.0	Y		Park and Shin (2019)	Park et al. (2019)
TaiESM1	AS-RCEC	TaiAM1	SNAP	SNAP	POP2	CICE4	CLM4.0	N	N	Lee and Liang (2020)	Lee et al. (2020b)
UKESM1-0-LL	MOHC	MetUM- HadGEM3-GA7.1	UKCA-GLOMAP- mode	UKCA-StratTrop	NEMO- HadGEM3- GO6.0	CICE- HadGEM3- GSI8	JULES-ES- 1.0	N	N	Tang et al. (2019)s	Sellar et al. (2019)

Table 2: Model atmospheric, aerosol, chemistry, ocean, sea ice and land-use components of CMIP6 models, with the models contributing multiple coupled transient experiments to AerChemMIP highlighted in **bold**. A full description of the representation of aerosol-radiation interactions, aerosol-cloud interactions, atmospheric chemistry, dust emissions, and biomass burning emissions in the AerChemMIP models can be found in Wilcox. This description uses text extracted from model documentation papers and entries in ES-DOC.

Model	Centre	Model components			Model grid				Emergent properties			Reference
		Atmosphere	Aerosol	Chemistry	nx	ny	nz	Model Top	ECS	ERF_AA	ASR_HD	
ACCESS-CM2	CSIRO-ARCCSS	MetUM-HadGEM3-GA7.1	UKCA-GLOMAP-mode		192	144	85	85km	4.72	-1.09	2.19	Bi et al. (2020)
ACCESS-ESM1-5	CSIRO	HadGAM2	CLASSIC (v1.0)		192	145	38	39km	3.87	-1.15	1.90	Ziehn et al. (2020)
AWI-ESM-1-1-LR	AWI	ECHAM6.3.04p1	MACv2-SP		192	96	47	80km	3.16			Sidorenko et al. (2015)
BCC-CSM2-MR	BCC	BCC_AGCM3_MR	MACv2-SP		320	160	46	1.46hPa	3.04		0.60	Wu et al. (2019)
BCC-ESM1	BCC	BCC_AGCM3_LR		BCC-AGCM3-Chem	128	64	26	2.19hPa	3.26		2.04	Wu et al. (2020)
CAMS-CSM1-0	CAMS	ECHAM5_CAMS			320	160	31	10hPa	2.29		-0.01	Chen et al. (2019), Rong et al. (2018)
CanESM5	CCCma	CanAM5	Interactive	Specified oxidants	128	64	49	1hPa	5.62	-0.85	1.61	Swart et al. (2019b)
CAS-ESM2-0	CAS	IAP AGCM 5.0	IAP AACM	IAP AACM	256	128	35	2.2hPa	3.51		2.51	Zhang et al. (2020)
CESM2-FV2	NCAR	CAM6	MAM4	MAM4	144	96	32	2.25hPa	5.14		2.95	Danabasoglu et al. (2020)
CESM2	NCAR	CAM6	MAM4	MAM4	288	192	32	2.25hPa	5.16	-1.37	2.55	Danabasoglu et al. (2020)
CESM2-WACCM-FV2	NCAR	WACCM6	MAM4	MAM4	144	96	70	4.5E-6 hPa	4.79		2.82	Danabasoglu et al. (2020)
CESM2-WACCM	NCAR	WACCM6	MAM4	MAM4	288	192	70	4.5E-6 hPa	4.75		2.74	Danabasoglu et al. (2020)
CIesm	THU	CIesm-AM	MAM4	trop_mam4	288	192	30	2.255hPa	5.67		0.45	Lin et al. (2020)
CMCC-CM2-HR4	CMCC	CAM4	MACv2-SP		288	192	26	2hPa			0.58	Cherchi et al. (2019)
CMCC-CM2-SR5	CMCC	CAM5.3	MAM3		288	192	30	2.2hPa	3.52		0.48	Cherchi et al. (2019)
CNRM-CM6-1	CNRM-CERFACS	Arpege 6.3	Prescribed monthly fields computed by TACTIC_v2	OZL_v2		128	91	78.4km	4.83	-1.15	1.05	Voldoire et al. (2019)
CNRM-ESM2-1	CNRM-CERFACS	Arpege 6.3	TACTIC_v2	REPROBUS-C_v2	256	128	91	78.4km	4.76	-0.74	1.21	S��f��rian et al. (2019)
E3SM-1-0	E3SM-Project	EAM v1.0	MAM4	Troposphere specified oxidants for aerosols. Stratosphere linearized interactive ozone (LINOZ v2)			72	0.1hPa	5.32	-1.65	3.37	Golaz et al. (2019)
E3SM-1-1	E3SM-Project	EAM v1.1	MAM4	Troposphere specified oxidants for aerosols. interactive ozone (LINOZ v2)			72	0.1hPa			4.04	Burrows et al. (2020)
EC-Earth3-AerChem	EC-Earth-Consortium	IFS cy36r4	TM5	TM5	512	256	91	0.01hPa	3.90	-0.70	3.95	van Noije et al. (2021)
EC-Earth3	EC-Earth-Consortium	IFS cy36r4	MACv2-SP	TM5	512	256	91	0.01hPa		-0.91	2.43	D��scher et al. (2021)
EC-Earth3-Veg	EC-Earth-Consortium	IFS cy36r4	MACv2-SP	TM5	512	256	91	0.01hPa	4.31		-0.37	D��scher et al. (2021)
FGOALS-f3-L	CAS	FAMIL2.2	MACv2-SP		360	180	32	2.16hPa	3		1.43	HE et al. (2020)
FGOALS-g3	CAS	GAMIL3	MACv2-SP		360	180	26	2.19hPa	2.88		0.10	Li et al. (2020)
FIO-ESM-2-0	FIO-QLNM	CAM4	Prescribed monthly fields		192	288	26	2hPa			0.46	Bao et al. (2020)
GFDL-CM4	NOAA-GFDL	GFDL-AM4.0.1	Interactive	fast chemistry, aerosol only	360	180	33	1hPa	5	-0.73	1.81	Held et al. (2019)
GFDL-ESM4	NOAA-GFDL	GFDL-AM4.1	Interactive	GFDL-ATMCHEM4.1	360	180	49	1hPa			1.43	Dunne et al. (2020)
SAM0-UNICON	NASA-GISS	GISS-E2.1	Varies with physics version	Varies with physics version	144	90	40	0.1hPa			1.84	Kelley et al. (2020), Miller et al. (2021)
GISS-E2-1-G (p1)	NASA-GISS	GISS-E2.1	'none'	Non-interactive	144	90	40	0.1hPa	2.72	-1.32	1.76	Kelley et al. (2020), Miller et al. (2021)

Continued on next page

GISS-E2-1-H (p1)	NASA-GISS	GISS-E2.1	'none'	Non-interactive	144	90	40	0.1hPa	3.11		1.68	Kelley et al. (2020), Miller et al. (2021)
GISS-E2-1-G (p3)	NASA-GISS	GISS-E2.1	OMA	GPUCCINI	144	90	40	0.1hPa	2.60	-1.85		Kelley et al. (2020), Miller et al. (2021)
GISS-E2-1-H (p3)	NASA-GISS	GISS-E2.1	OMA	GPUCCINI	144	90	40	0.1hPa	3.10			Kelley et al. (2020), Miller et al. (2021)
GISS-E2-1-G (p5)	NASA-GISS	GISS-E2.1	MATRIX	GPUCCINI	144	90	40	0.1hPa	2.80	-1.36		Kelley et al. (2020), Miller et al. (2021)
GISS-E2-1-H (p5)	NASA-GISS	GISS-E2.1	MATRIX	GPUCCINI	144	90	40	0.1hPa				Kelley et al. (2020), Miller et al. (2021)
HadGEM3-GC31-LL	MOHC	MetUM-HadGEM3-GA7.1	UKCA-GLOMAP-mode		192	144	85	85km	5.55	-1.10	2.47	Andrews et al. (2020), Kuhlbrodt et al. (2018)
HadGEM3-GC31-MM	MOHC	MetUM-HadGEM3-GA7.1	UKCA-GLOMAP-mode		423	324	85	85km	5.42		2.65	Andrews et al. (2020)
INM-CM4-8	INM	INM-AM4-8	INM-AER1		180	120	21	0.01 sigma	1.83			Volodin et al. (2010)
INM-CM5-0	INM	INM-AM5-0	INM-AER1		180	120	73	002 sigma	1.92		0.18	Volodin et al. (2017), Volodin and Kostykin (2016)
IPSL-CM6A-LR	IPSL	LMDZ	INCA v6 AER		144	143	79	80km	4.56	-0.59	0.35	Boucher et al. (2020)
KACE-1-0-G	NIMS-KMA	MetUM-HadGEM3-GA7.1	UKCA-GLOMAP-mode		192	144	85	85km	4.48		1.76	Lee et al. (2020a)
KIOST-ESM	KIOST	GFDL-AM2.0	MACv2-SP	Simple carbon aerosol	192	96	32	2hPa	3.36		0.02	Pak et al. (2021)
				model (emission type)								
MIROC6	MIROC	CCSR AGCM	SPRINTARS6.0		256	128	81	04hPa	2.61	-1.06	0.81	Tatebe et al. (2019)
MIROC-ES2L	MIROC	CCSR AGCM	SPRINTARS6.0		128	64	40	3hPa	2.68		0.94	Hajima et al. (2020)
MPI-ESM-1-2-HAM	HAMMOZ-Consortium	ECHAM6.3	HAM2.3	sulfur chemistry	192	96	47	0.01hPa	2.96		2.23	Mauritsen et al. (2019)
MPI-ESM1-2-HR	MPI-M	ECHAM6.2	MACv2-SP		384	192	95	0.01hPa	3		0.11	Mauritsen et al. (2019), Stevens et al. (2013)
MPI-ESM1-2-LR	MPI-M	ECHAM6.3	MACv2-SP		192	96	47	0.01hPa	3		0.24	Mauritsen et al. (2019), Stevens et al. (2013)
MRI-ESM2-0	MRI	MRI-AGCM3.5	MASINGAR mk2r4	MRI-CCM2.1	320	160	80	0.01hPa	3.15	-1.19	2.35	YUKIMOTO et al. (2019)
NESM3	NUIST	ECHAM v6.3	MACv2-SP		192	96	47	1hPa	4.72		1.45	Cao et al. (2018)
NorCPM1	NCC	CAM-OSLO4.1	OsloAero4.1	OsloChemSimp4.1	144	96	26	2hPa	3.05		2.41	Bethke et al. (2021)
NorESM2-LM	NCC	CAM-OSLO	OsloAero	OsloChemSimp	144	96	32	3mb	2.54	-1.21	2.24	Seland et al. (2020)
NorESM2-MM	NCC	CAM-OSLO	OsloAero	OsloChemSimp	288	192	32	3mb	2.50	-1.26	2.13	Seland et al. (2020)
SAM0-UNICON	SNU	CAM5.3 with UNICON	MAM3		288	192	30	2hPa	3.72	-1.23	1.68	Park et al. (2019)
TaiESM1	AS-RCEC	TaiAM1	SNAP	SNAP	288	192	20	2hPa	4.31		2.79	Lee et al. (2020b)
UKESM1-0-LL	MOHC	MetUM-HadGEM3-GA7.1	UKCA-GLOMAP-mode	UKCA-StratTrop	192	144	85	85km	5.34	-1.11	2.84	Sellar et al. (2019)

Table 3: Model atmospheric, aerosol, and chemistry components of CMIP6 models; their atmospheric grids; and emergent properties; with the models contributing multiple coupled transient experiments to AerChemMIP highlighted in **bold**. Equilibrium Climate Sensitivity (ECS) is taken from Schlund et al. (2020). Effective Radiative Forcing due to anthropogenic aerosol (ERF_AA) was calculated by the authors for the period 2014 vs. 1850 using RFMIP *piClim-aer* and *piClim* simulations. The difference in absorbed solar radiation between the Southern and Northern Hemispheres (ASR_HD) is used as a proxy for aerosol radiative forcing, which can be calculated from historical simulations. Here, the change in this index over the period from 1850 to 1985, when it is most closely related to aerosol forcing, is shown, and is taken from Menary et al. (2020).

810 *Data availability.* All of the data from the CMIP and AerChemMIP simulations analysed in this study have been published on the Earth System Grid Federation. The data for Tables XX and YY are available at Wilcox.

Author contributions. PTG, LJW and RJA led the writing of this manuscript. All authors contributed to the writing and discussion of this paper.

Competing interests. At least one of the (co-)authors is a member of the editorial board of Atmospheric Chemistry and Physics.

815 *Acknowledgements.* PTG and LJW are supported by the Natural Environment Research Council (NERC; grant NE/W004895/1, TerraFIRMA). ATA, PTG, and LJW are additionally supported by the National Centre for Atmospheric Science. RJA acknowledges support from NSF grant AGS-2153486. RJA, BHS, and LJW acknowledge funding by the Research Council of Norway through grant no. 324182 (CATHY). MD and NO were supported by the Environment Research and Technology Development Fund (JPMEERF20202003, JPMEERF20205001 and JPMEERF20232001) of the Environmental Restoration and Conservation Agency Provided by the Ministry of Environment of Japan, the
820 Arctic Challenge for Sustainability II (ArCS II), Program Grant Number JPMXD1420318865, and a grant for the Global Environmental Research Coordination System from the Ministry of the Environment of Japan (MLIT2253). CJS was supported by a NERC/IIASA Collaborative Research Fellowship (NE/T009381/1). STT was funded by the Met Office Climate Science for Service Partnership (CSSP) China project under the International Science Partnerships Fund (ISPF). FMO'C was supported by the European Union's Horizon 2020 project ESM2025 (under grant agreement No 101003536) and the Met Office Hadley Centre Climate Programme funded by DSIT, UK. SF is sup-
825 ported by the German Science Foundation (DFG) with the Collaborative Research Center DETECT (SFB1502, grant no. DFG 450058266) and by HORIZON Europe with the project EXPECT (grant no. 101137656).

References

- Abalos, M., Randel, W. J., Kinnison, D. E., and Garcia, R. R.: Using the Artificial Tracer e90 to Examine Present and Future UTLS Tracer Transport in WACCM, *Journal of the Atmospheric Sciences*, 74, 3383–3403, <https://doi.org/10.1175/JAS-D-17-0135.1>, publisher: American Meteorological Society Section: *Journal of the Atmospheric Sciences*, 2017.
- 830 Ahsan, H., Wang, H., Wu, J., Wu, M., Smith, S. J., Bauer, S., Suchyta, H., Olivié, D., Myhre, G., Matsui, H., Bian, H., Lamarque, J.-F., Carslaw, K., Horowitz, L., Regayre, L., Chin, M., Schulz, M., Skeie, R. B., Takemura, T., and Naik, V.: The Emissions Model Inter-comparison Project (Emissions-MIP): quantifying model sensitivity to emission characteristics, *Atmospheric Chemistry and Physics*, 23, 14 779–14 799, <https://doi.org/10.5194/acp-23-14779-2023>, 2023.
- 835 Aizawa, T., Oshima, N., and Yukimoto, S.: Contributions of Anthropogenic Aerosol Forcing and Multidecadal Internal Variability to Mid-20th Century Arctic Cooling—CMIP6/DAMIP Multimodel Analysis, *Geophysical Research Letters*, 49, e2021GL097093, <https://doi.org/https://doi.org/10.1029/2021GL097093>, e2021GL097093 2021GL097093, 2022.
- Akritidis, D., Bacer, S., Zanis, P., Georgoulas, A. K., Chowdhury, S., Horowitz, L. W., Naik, V., O'Connor, F. M., Keeble, J., Sager, P. L., van Noije, T., Zhou, P., Turnock, S., West, J. J., Lelieveld, J., and Pozzer, A.: Strong increase in mortality attributable to ozone pollution under a climate change and demographic scenario, 19, 024 041, <https://doi.org/10.1088/1748-9326/ad2162>, 2024.
- 840 Allen, R. J., Turnock, S., Nabat, P., Neubauer, D., Lohmann, U., Olivié, D., Oshima, N., Michou, M., Wu, T., Zhang, J., Takemura, T., Schulz, M., Tsigaridis, K., Bauer, S. E., Emmons, L., Horowitz, L., Naik, V., van Noije, T., Bergman, T., Lamarque, J.-F., Zanis, P., Tegen, I., Westervelt, D. M., Le Sager, P., Good, P., Shim, S., O'Connor, F., Akritidis, D., Georgoulas, A. K., Deushi, M., Sentman, L. T., John, J. G., Fujimori, S., and Collins, W. J.: Climate and air quality impacts due to mitigation of non-methane near-term climate forcers, *Atmospheric Chemistry and Physics*, 20, 9641–9663, <https://doi.org/10.5194/acp-20-9641-2020>, publisher: Copernicus GmbH, 2020.
- 845 Allen, R. J., Horowitz, L. W., Naik, V., Oshima, N., O'Connor, F. M., Turnock, S., Shim, S., Le Sager, P., Van Noije, T., Tsigaridis, K., Bauer, S. E., Sentman, L. T., John, J. G., Broderick, C., Deushi, M., Folberth, G. A., Fujimori, S., and Collins, W. J.: Significant climate benefits from near-term climate forcer mitigation in spite of aerosol reductions, *Environmental Research Letters*, <https://doi.org/10.1088/1748-9326/abe06b>, 2021.
- 850 Allen, R. J., Zhao, X., Randles, C. A., Kramer, R. J., Samset, B. H., and Smith, C. J.: Surface warming and wetting due to methane's long-wave radiative effects muted by short-wave absorption, *Nature Geoscience*, 16, 314–320, <https://doi.org/10.1038/s41561-023-01144-z>, 2023.
- Allen, R. J., Gomez, J., Horowitz, L. W., and Shevliakova, E.: Enhanced future vegetation growth with elevated carbon dioxide concentrations could increase fire activity, *Communications Earth and Environment*, 5, 54, <https://doi.org/10.1038/s43247-024-01228-7>, 2024a.
- 855 Allen, R. J., Zhao, X., Randles, C. A., Kramer, R. J., Samset, B. H., and Smith, C. J.: Present-Day Methane Shortwave Absorption Mutes Surface Warming and Wetting Relative to Preindustrial Conditions, *EGUsphere*, 2024, 1–41, <https://doi.org/10.5194/egusphere-2024-872>, 2024b.
- Amiri-Farahani, A., Allen, R. J., Li, K.-F., Nabat, P., and Westervelt, D. M.: A La Niña-Like Climate Response to South African Biomass Burning Aerosol in CESM Simulations, *Journal of Geophysical Research: Atmospheres*, 125, e2019JD031 832, <https://doi.org/10.1029/2019JD031832>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2019JD031832>, 2020.
- 860 Andrews, M. B., Ridley, J. K., Wood, R. A., Andrews, T., Blockley, E. W., Booth, B., Burke, E., Dittus, A. J., Florek, P., Gray, L. J., Haddad, S., Hardiman, S. C., Hermanson, L., Hodson, D., Hogan, E., Jones, G. S., Knight, J. R., Kuhlbrodt, T., Misios, S., Mizieliński, M. S., Ringer, M. A., Robson, J., and Sutton, R. T.: Historical Simulations With HadGEM3-GC3.1 for CMIP6, *Journal of Advances in Modeling*

- Earth Systems, 12, e2019MS001995, <https://doi.org/https://doi.org/10.1029/2019MS001995>, e2019MS001995 10.1029/2019MS001995, 2020.
- Anenberg, S. C., West, J. J., Fiore, A. M., Jaffe, D. A., Prather, M. J., Bergmann, D., Cuvelier, K., Dentener, F. J., Duncan, B. N., Gauss, M., Hess, P., Jonson, J. E., Lupu, A., MacKenzie, I. A., Marmer, E., Park, R. J., Sanderson, M. G., Schultz, M., Shindell, D. T., Szopa, S., Vivanco, M. G., Wild, O., and Zeng, G.: Intercontinental impacts of ozone pollution on human mortality, *Environmental Science & Technology*, 43, 6482–6487, <https://doi.org/10.1021/es900518z>, 2009.
- Archibald, A. T., Collins, W., Evans, M., Griffiths, P., O'Connor, F., Wild, O., and Young, P.: Ace in the hole or a house of cards: Will a DeCK experiment help atmospheric chemistry?, in: *Egusphere*, <https://doi.org/10.5194/egusphere-egu22-9442>, 2022.
- Arora, V. K., Katavouta, A., Williams, R. G., Jones, C. D., Brovkin, V., Friedlingstein, P., Schwinger, J., Bopp, L., Boucher, O., Cadule, P., Chamberlain, M. A., Christian, J. R., Delire, C., Fisher, R. A., Hajima, T., Ilyina, T., Joetzjer, E., Kawamiya, M., Koven, C. D., Krasting, J. P., Law, R. M., Lawrence, D. M., Lenton, A., Lindsay, K., Pongratz, J., Raddatz, T., Séférian, R., Tachiiri, K., Tjiputra, J. F., Wiltshire, A., Wu, T., and Ziehn, T.: Carbon–concentration and carbon–climate feedbacks in CMIP6 models and their comparison to CMIP5 models, *Biogeosciences*, 17, 4173–4222, <https://doi.org/10.5194/bg-17-4173-2020>, 2020.
- Bader, D. C., Leung, R., Taylor, M., and McCoy, R. B.: E3SM-Project E3SM1.0 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.4497>, 2019a.
- Bader, D. C., Leung, R., Taylor, M., and McCoy, R. B.: E3SM-Project E3SM1.1 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.11485>, 2019b.
- Balaji, V., Taylor, K. E., Juckes, M., Lawrence, B. N., Durack, P. J., Lautenschlager, M., Blanton, C., Cinquini, L., Denvil, S., Elkington, M., Guglielmo, F., Guilyardi, E., Hassell, D., Kharin, S., Kindermann, S., Nikonov, S., Radhakrishnan, A., Stockhause, M., Weigel, T., and Williams, D.: Requirements for a global data infrastructure in support of CMIP6, *Geoscientific Model Development*, 11, 3659–3680, <https://doi.org/10.5194/gmd-11-3659-2018>, publisher: Copernicus GmbH, 2018.
- Bao, Y., Song, Z., and Qiao, F.: FIO-ESM Version 2.0: Model Description and Evaluation, *Journal of Geophysical Research: Oceans*, 125, e2019JC016036, <https://doi.org/https://doi.org/10.1029/2019JC016036>, e2019JC016036 2019JC016036, 2020.
- Bellouin, N., Quaas, J., Gryspeerdt, E., Kinne, S., Stier, P., Watson-Parris, D., Boucher, O., Carslaw, K. S., Christensen, M., Daniau, A. L., Dufresne, J. L., Feingold, G., Fiedler, S., Forster, P., Gettelman, A., Haywood, J. M., Lohmann, U., Malavelle, F., Mauritsen, T., McCoy, D. T., Myhre, G., Mülmenstädt, J., Neubauer, D., Possner, A., Rugenstein, M., Sato, Y., Schulz, M., Schwartz, S. E., Sourdeval, O., Storelvmo, T., Toll, V., Winker, D., and Stevens, B.: Bounding global aerosol radiative forcing of climate change, *Reviews of Geophysics*, 58, e2019RG000660, <https://doi.org/10.1029/2019RG000660>, 2020.
- Bentsen, M., Olivie, D. J. L., Seland, y., Toniazzo, T., Gjermundsen, A., Graff, L. S., Debernard, J. B., Gupta, A. K., He, Y., Kirkevåg, A., Schwinger, J., Tjiputra, J., Aas, K. S., Bethke, I., Fan, Y., Griesfeller, J., Grini, A., Guo, C., Ilicak, M., Karset, I. H. H., Landgren, O. A., Liakka, J., Moseid, K. O., Nummelin, A., Spensberger, C., Tang, H., Zhang, Z., Heinze, C., Iversen, T., and Schulz, M.: NCC NorESM2-MM model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.8040>, 2019.
- Bethke, I., Wang, Y., Counillon, F., Kimmritz, M., Fransner, F., Samuelsen, A., Langehaug, H. R., Chiu, P.-G., Bentsen, M., Guo, C., Tjiputra, J., Kirkevåg, A., Olivie, D. J. L., Seland, y., Fan, Y., Lawrence, P., Eldevik, T., and Keenlyside, N.: NCC NorCPM1 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.10894>, 2019.
- Bethke, I., Wang, Y., Counillon, F., Keenlyside, N., Kimmritz, M., Fransner, F., Samuelsen, A., Langehaug, H., Svendsen, L., Chiu, P.-G., Passos, L., Bentsen, M., Guo, C., Gupta, A., Tjiputra, J., Kirkevåg, A., Olivie, D., Seland, Ø., Solsvik Vågane, J., Fan, Y., and Eldevik,

- T.: NorCPM1 and its contribution to CMIP6 DCP, Geoscientific Model Development, 14, 7073–7116, <https://doi.org/10.5194/gmd-14-7073-2021>, 2021.
- Bi, D., Dix, M., Marsland, S., O’Farrell, S., Sullivan, A., Bodman, R., Law, R., Harman, I., Srbinovsky, J., Rashid, H. A., Dobrohotoff, P., Mackallah, C., Yan, H., Hirst, A., Savita, A., Dias, F. B., Woodhouse, M., Fiedler, R., and Heerdegen, A.: Configuration and spin-up of ACCESS-CM2, the new generation Australian Community Climate and Earth System Simulator Coupled Model, *Journal of Southern Hemisphere Earth Systems Science*, 70, 225–251, <https://doi.org/10.1071/ES19040>, 2020.
- Blichner, S.-M. and Berntsen, T.: Chapter 6 of the Working Group I Contribution to the IPCC Sixth Assessment Report - data for Figure 6.12 (v20220815), <https://doi.org/10.5285/8855e410adf547b4afd039a5b88487f4>, 2023.
- Booth, B. B. B., Harris, G. R., Jones, A., Wilcox, L., Hawcroft, M., and Carslaw, K. S.: Comments on “Rethinking the Lower Bound on Aerosol Radiative Forcing”, *Journal of Climate*, 31, 9407 – 9412, <https://doi.org/10.1175/JCLI-D-17-0369.1>, 2018.
- Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M., Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S., Sherwood, S., Stevens, B., and Zhang, X.: Clouds and aerosols, in: *Climate change 2013: The physical science basis. Contribution of working group I to the fifth assessment report of the intergovernmental panel on climate change*, edited by Stocker, T., Qin, D., Plattner, G.-K., Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P., Cambridge University Press, Cambridge, UK and New York, NY, USA, <https://doi.org/10.1017/CBO9781107415324.016>, section: 7 Type: Book section, 2013.
- Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D’Andrea, F., Davini, P., de Lavergne, C., Denvil, S., Deshayes, J., Devilliers, M., Ducharne, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L., Falletti, L., Flavoni, S., Foujols, M.-A., Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez, Lionel, E., Guilyardi, E., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M., Khodri, M., Krinner, G., Lebas, N., Levvasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luyssaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meurdesoif, Y., Mignot, J., Musat, I., Ottlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thiéblemont, R., Traore, A. K., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N.: Presentation and Evaluation of the IPSL-CM6A-LR Climate Model, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002010, <https://doi.org/https://doi.org/10.1029/2019MS002010>, e2019MS002010 10.1029/2019MS002010, 2020.
- Boucher, O., Denvil, S., Levvasseur, G., Cozic, A., Caubel, A., Foujols, M.-A., Meurdesoif, Y., Balkanski, Y., Checa-Garcia, R., Hauglustaine, D., Bekki, S., and Marchand, M.: IPSL IPSL-CM6A-LR-INCA model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.13601>, 2021.
- Brown, F., Folberth, G. A., Sitch, S., Bauer, S., Bauters, M., Boeckx, P., Cheesman, A. W., Deushi, M., Dos Santos Vieira, I., Galy-Lacaux, C., Haywood, J., Keeble, J., Mercado, L. M., O’Connor, F. M., Oshima, N., Tsigaridis, K., and Verbeeck, H.: The ozone–climate penalty over South America and Africa by 2100, *Atmospheric Chemistry and Physics*, 22, 12 331–12 352, <https://doi.org/10.5194/acp-22-12331-2022>, publisher: Copernicus GmbH, 2022.
- Burrows, S. M., Maltrud, M., Yang, X., Zhu, Q., Jeffery, N., Shi, X., Ricciuto, D., Wang, S., Bisht, G., Tang, J., Wolfe, J., Harrop, B. E., Singh, B., Brent, L., Baldwin, S., Zhou, T., Cameron-Smith, P., Keen, N., Collier, N., Xu, M., Hunke, E. C., Elliott, S. M., Turner, A. K., Li, H., Wang, H., Golaz, J.-C., Bond-Lamberty, B., Hoffman, F. M., Riley, W. J., Thornton, P. E., Calvin, K., and Leung, L. R.: The DOE E3SM v1.1 Biogeochemistry Configuration: Description and Simulated Ecosystem-Climate Responses to Historical Changes in Forcing, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001766, <https://doi.org/https://doi.org/10.1029/2019MS001766>, e2019MS001766 10.1029/2019MS001766, 2020.

Byun, Y.-H., Lim, Y.-J., Sung, H. M., Kim, J., Sun, M., and Kim, B.-H.: NIMS-KMA KACE1.0-G model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.8378>, 2019.

Cao, J. and Wang, B.: NUIST NESMv3 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.8769>, 2019.

Cao, J., Wang, B., Yang, Y.-M., Ma, L., Li, J., Sun, B., Bao, Y., He, J., Zhou, X., and Wu, L.: The NUIST Earth System Model (NESM) version 3: description and preliminary evaluation, *Geoscientific Model Development*, 11, 2975–2993, <https://doi.org/10.5194/gmd-11-2975-2018>, 2018.

Carslaw, K. S., Lee, L. A., Reddington, C. L., Pringle, K. J., Rap, A., Forster, P. M., Mann, G. W., Spracklen, D. V., Woodhouse, M. T., Regayre, L. A., and Pierce, J. R.: Large Contribution of Natural Aerosols to Uncertainty in Indirect Forcing, *Nature*, 503, 67–71, <https://doi.org/10.1038/nature12674>, 2013.

Chen, H.-M., Li, J., Su, J.-Z., Hua, L.-J., Rong, X.-Y., Xin, Y.-F., and Zhang, Z.-Q.: Introduction of CAMS-CSM model and its participation in CMIP6, *Advances in Climate Change Research*, 15, 540–544, 2019.

Cherchi, A., Fogli, P. G., Lovato, T., Peano, D., Iovino, D., Gualdi, S., Masina, S., Scoccimarro, E., Materia, S., Bellucci, A., and Navarra, A.: Global Mean Climate and Main Patterns of Variability in the CMCC-CM2 Coupled Model, *Journal of Advances in Modeling Earth Systems*, 11, 185–209, <https://doi.org/https://doi.org/10.1029/2018MS001369>, 2019.

Collins, W. J., Lamarque, J.-F., Schulz, M., Boucher, O., Eyring, V., Hegglin, M. I., Maycock, A., Myhre, G., Prather, M., Shindell, D., and Smith, S. J.: AerChemMIP: quantifying the effects of chemistry and aerosols in CMIP6, *Geoscientific Model Development*, 10, 585–607, <https://doi.org/10.5194/gmd-10-585-2017>, publisher: Copernicus GmbH, 2017.

Collins, W. J., O'Connor, F. M., Barker, C. R., Byrom, R. E., Eastham, S. D., Hodnebrog, Ø., Jöckel, P., Marais, E. A., Mertens, M., Myhre, G., Nützel, M., Olivié, D., Bieltvedt Skeie, R., Stecher, L., Horowitz, L. W., Naik, V., Faluvegi, G., Im, U., Murray, L. T., Shindell, D., Tsigaridis, K., Abraham, N. L., and Keeble, J.: Climate Forcing due to Future Ozone Changes: An intercomparison of metrics and methods, *EGUsphere*, 2024, 1–50, <https://doi.org/10.5194/egusphere-2024-3698>, 2024.

Danabasoglu, G.: NCAR CESM2-FV2 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.11297>, 2019a.

Danabasoglu, G.: NCAR CESM2-WACCM-FV2 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.11298>, 2019b.

Danabasoglu, G.: NCAR CESM2-WACCM model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.10071>, 2019c.

Danabasoglu, G.: NCAR CESM2 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.7627>, 2019d.

Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D. A., DuVivier, A. K., Edwards, J., Emmons, L. K., Fasullo, J., Garcia, R., Gettelman, A., Hannay, C., Holland, M. M., Large, W. G., Lauritzen, P. H., Lawrence, D. M., Lenaerts, J. T. M., Lindsay, K., Lipscomb, W. H., Mills, M. J., Neale, R., Oleson, K. W., Otto-Bliesner, B., Phillips, A. S., Sacks, W., Tilmes, S., van Kampenhout, L., Vertenstein, M., Bertini, A., Dennis, J., Deser, C., Fischer, C., Fox-Kemper, B., Kay, J. E., Kinnison, D., Kushner, P. J., Larson, V. E., Long, M. C., Mickelson, S., Moore, J. K., Nienhouse, E., Polvani, L., Rasch, P. J., and Strand, W. G.: The Community Earth System Model Version 2 (CESM2), *Journal of Advances in Modeling Earth Systems*, 12, e2019MS001916, <https://doi.org/https://doi.org/10.1029/2019MS001916>, e2019MS001916 2019MS001916, 2020.

Danek, C., Shi, X., Stepanek, C., Yang, H., Barbi, D., Hegewald, J., and Lohmann, G.: AWI AWI-ESM1.1LR model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.9328>, 2020.

- Deng, J., Dai, A., and Xu, H.: Nonlinear Climate Responses to Increasing CO₂ and Anthropogenic Aerosols Simulated by CESM1, *Journal of Climate*, 33, 281–301, <https://doi.org/10.1175/JCLI-D-19-0195.1>, publisher: American Meteorological Society Section: Journal of Climate, 2020.
- 980 DeRepentigny, P., Jahn, A., Holland, M. M., and Smith, A.: Arctic Sea Ice in Two Configurations of the CESM2 During the 20th and 21st Centuries, *Journal of Geophysical Research: Oceans*, 125, e2020JC016133, <https://doi.org/10.1029/2020JC016133>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020JC016133>, 2020.
- Diamond, M. S., Gristey, J. J., Kay, J. E., and Feingold, G.: Anthropogenic aerosol and cryosphere changes drive Earth’s strong but transient clear-sky hemispheric albedo asymmetry, *Communications Earth & Environment*, 3, 1–10, <https://doi.org/10.1038/s43247-022-00546-y>, number: 1 Publisher: Nature Publishing Group, 2022.
- 985 Dix, M., Bi, D., Dobrohotoff, P., Fiedler, R., Harman, I., Law, R., Mackallah, C., Marsland, S., O’Farrell, S., Rashid, H., Srbinovsky, J., Sullivan, A., Trenham, C., Vohralik, P., Watterson, I., Williams, G., Woodhouse, M., Bodman, R., Dias, F. B., Domingues, C., Hannah, N., Heerdegen, A., Savita, A., Wales, S., Allen, C., Druken, K., Evans, B., Richards, C., Ridzwan, S. M., Roberts, D., Smillie, J., Snow, K., Ward, M., and Yang, R.: CSIRO-ARCCSS ACCESS-CM2 model output prepared for CMIP6 CMIP historical, 990 <https://doi.org/10.22033/ESGF/CMIP6.4271>, 2019.
- Döscher, R., Acosta, M., Alessandri, A., Anthoni, P., Arneth, A., Arsouze, T., Bergmann, T., Bernadello, R., Bousetta, S., Caron, L.-P., Carver, G., Castrillo, M., Catalano, F., Cvijanovic, I., Davini, P., Dekker, E., Doblas-Reyes, F. J., Docquier, D., Echevarria, P., Fladrich, U., Fuentes-Franco, R., Gröger, M., v. Hardenberg, J., Hieronymus, J., Karami, M. P., Keskinen, J.-P., Koenigk, T., Makkonen, R., Massonnet, F., Ménégos, M., Miller, P. A., Moreno-Chamarro, E., Nieradzik, L., van Noije, T., Nolan, P., O’Donnell, D., Ollinaho, P., van den Oord, 995 G., Ortega, P., Prims, O. T., Ramos, A., Reerink, T., Rousset, C., Ruprich-Robert, Y., Le Sager, P., Schmith, T., Schrödner, R., Serva, F., Sicardi, V., Sloth Madsen, M., Smith, B., Tian, T., Tourigny, E., Uotila, P., Vancoppenolle, M., Wang, S., Wårlind, D., Willén, U., Wyser, K., Yang, S., Yepes-Arbós, X., and Zhang, Q.: The EC-Earth3 Earth System Model for the Climate Model Intercomparison Project 6, *Geoscientific Model Development Discussions*, 2021, 1–90, <https://doi.org/10.5194/gmd-2020-446>, 2021.
- Dunne, J. P., Horowitz, L. W., Adcroft, A. J., Ginoux, P., Held, I. M., John, J. G., Krasting, J. P., Malyshev, S., Naik, V., Paulot, F., Shevliakova, 1000 E., Stock, C. A., Zadeh, N., Balaji, V., Blanton, C., Dunne, K. A., Dupuis, C., Durachta, J., Dussin, R., Gauthier, P. P. G., Griffies, S. M., Guo, H., Hallberg, R. W., Harrison, M., He, J., Hurlin, W., McHugh, C., Menzel, R., Milly, P. C. D., Nikonov, S., Paynter, D. J., Ploshay, J., Radhakrishnan, A., Rand, K., Reichl, B. G., Robinson, T., Schwarzkopf, D. M., Sentman, L. T., Underwood, S., Vahlenkamp, H., Winton, M., Wittenberg, A. T., Wyman, B., Zeng, Y., and Zhao, M.: The GFDL Earth System Model Version 4.1 (GFDL-ESM 4.1): Overall Coupled Model Description and Simulation Characteristics, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002015, 1005 <https://doi.org/https://doi.org/10.1029/2019MS002015>, e2019MS002015 2019MS002015, 2020.
- EC-Earth, E.-E. C.: EC-Earth-Consortium EC-Earth3-Veg model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.4706>, 2019a.
- EC-Earth, E.-E. C.: EC-Earth-Consortium EC-Earth3 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.4700>, 2019b.
- 1010 EC-Earth, E.-E. C.: EC-Earth-Consortium EC-Earth3-AerChem model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.4701>, 2020.
- Ekman, A. M. L.: Do sophisticated parameterizations of aerosol-cloud interactions in CMIP5 models improve the representation of recent observed temperature trends?, *Journal of Geophysical Research: Atmospheres*, 119, 817–832, <https://doi.org/10.1002/2013JD020511>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2013JD020511>, 2014.

- 1015 Emberson, L.: Effects of ozone on agriculture, forests and grasslands, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 378, 20190327, <https://doi.org/10.1098/rsta.2019.0327>, 2020.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, publisher: Copernicus GmbH, 2016.
- 1020 Fiedler, S., Naik, V., O'Connor, F. M., Smith, C. J., Griffiths, P., Kramer, R. J., Takemura, T., Allen, R. J., Im, U., Kasoar, M., Modak, A., Turnock, S., Voulgarakis, A., Watson-Parris, D., Westervelt, D. M., Wilcox, L. J., Zhao, A., Collins, W. J., Schulz, M., Myhre, G., and Forster, P. M.: Interactions between atmospheric composition and climate change – Progress in understanding and future opportunities from AerChemMIP, PDRMIP, and RFMIP, *Geoscientific Model Development*, 17, 2387–2417, <https://doi.org/10.5194/gmd-17-2387-2024>, 2024.
- 1025 Finney, D. L., Doherty, R. M., Wild, O., Young, P. J., and Butler, A.: Response of lightning NO_x emissions and ozone production to climate change: Insights from the Atmospheric Chemistry and Climate Model Intercomparison Project, *Geophysical Research Letters*, 43, 5492–5500, <https://doi.org/10.1002/2016GL068825>, 2016.
- Fiore, A. M., Milly, G. P., Hancock, S. E., QuiÃ±ones, L., Bowden, J. H., Helstrom, E., Lamarque, J.-F., Schnell, J., West, J. J., and Xu, Y.: Characterizing Changes in Eastern U.S. Pollution Events in a Warming World, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035985, <https://doi.org/https://doi.org/10.1029/2021JD035985>, e2021JD035985 2021JD035985, 2022.
- 1030 Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M., and Zhang, H.: The earth's energy budget, climate feedbacks, and climate sensitivity, in: *Climate change 2021: The physical science basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, UK and New York, NY, USA, <https://doi.org/10.1017/9781009157896.009>, section: 7 Type: Book section, 2021.
- Forster, P. M., Andrews, T., Good, P., Gregory, J. M., Jackson, L. S., and Zelinka, M.: Evaluating adjusted forcing and model spread for historical and future scenarios in the CMIP5 generation of climate models, *Journal of Geophysical Research: Atmospheres*, 118, 1139–1150, <https://doi.org/10.1002/jgrd.50174>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/jgrd.50174>, 2013.
- 1040 Forster, P. M., Richardson, T., Maycock, A. C., Smith, C. J., Samset, B. H., Myhre, G., Andrews, T., Pincus, R., and Schulz, M.: Recommendations for diagnosing effective radiative forcing from climate models for CMIP6, *Journal of Geophysical Research: Atmospheres*, 121, 12,460–12,475, <https://doi.org/10.1002/2016JD025320>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2016JD025320>, 2016.
- Forster, P. M., Smith, C. J., Walsh, T., Lamb, W. F., Lamboll, R., Hauser, M., Ribes, A., Rosen, D., Gillett, N., Palmer, M. D., Rogelj, J., von Schuckmann, K., Seneviratne, S. I., Trewin, B., Zhang, X., Allen, M., Andrew, R., Birt, A., Borger, A., Boyer, T., Broersma, J. A.,
- 1045 Cheng, L., Dentener, F., Friedlingstein, P., Gutiérrez, J. M., Gütschow, J., Hall, B., Ishii, M., Jenkins, S., Lan, X., Lee, J.-Y., Morice, C., Kadow, C., Kennedy, J., Killick, R., Minx, J. C., Naik, V., Peters, G. P., Pirani, A., Pongratz, J., Schleussner, C.-F., Szopa, S., Thorne, P., Rohde, R., Rojas Corradi, M., Schumacher, D., Vose, R., Zickfeld, K., Masson-Delmotte, V., and Zhai, P.: Indicators of Global Climate Change 2022: annual update of large-scale indicators of the state of the climate system and human influence, *Earth System Science Data*, 15, 2295–2327, <https://doi.org/10.5194/essd-15-2295-2023>, 2023.
- 1050 Fyfe, J. C., Kharin, V. V., Santer, B. D., Cole, J. N. S., and Gillett, N. P.: Significant impact of forcing uncertainty in a large ensemble of climate model simulations, *Proceedings of the National Academy of Sciences*, 118, e2016549118, <https://doi.org/10.1073/pnas.2016549118>, 2021.

Gaudel, A., Cooper, O. R., Ancellet, G., Barret, B., Boynard, A., Burrows, J. P., Clerbaux, C., Coheur, P.-F., Cuesta, J., Cuevas, E., Doniki, S., Dufour, G., Ebojie, F., Foret, G., Garcia, O., Granados-Muñoz, M. J., Hannigan, J. W., Hase, F., Hassler, B., Huang, G., Hurtmans, D., Jaffe, D., Jones, N., Kalabokas, P., Kerridge, B., Kulawik, S., Latter, B., Leblanc, T., Le Flochmoën, E., Lin, W., Liu, J., Liu, X., Mahieu, E., McClure-Begley, A., Neu, J. L., Osman, M., Palm, M., Petetin, H., Petropavlovskikh, I., Querel, R., Rahpoe, N., Rozanov, A., Schultz, M. G., Schwab, J., Siddans, R., Smale, D., Steinbacher, M., Tanimoto, H., Tarasick, D. W., Thouret, V., Thompson, A. M., Trickl, T., Weatherhead, E., Wespes, C., Worden, H. M., Vigouroux, C., Xu, X., Zeng, G., and Ziemke, J.: Tropospheric Ozone Assessment Report: Present-day distribution and trends of tropospheric ozone relevant to climate and global atmospheric chemistry model evaluation, *Elementa: Science of the Anthropocene*, 6, 39, <https://doi.org/10.1525/elementa.291>, 2018.

Ghan, S. J.: Technical Note: Estimating aerosol effects on cloud radiative forcing, *Atmospheric Chemistry and Physics*, 13, 9971–9974, <https://doi.org/10.5194/acp-13-9971-2013>, 2013.

Gidden, M. J., Riahi, K., Smith, S. J., Fujimori, S., Luderer, G., Kriegler, E., van Vuuren, D. P., van den Berg, M., Feng, L., Klein, D., Calvin, K., Doelman, J. C., Frank, S., Fricko, O., Harmsen, M., Hasegawa, T., Havlik, P., Hilaire, J., Hoesly, R., Horing, J., Popp, A., Stehfest, E., and Takahashi, K.: Global emissions pathways under different socioeconomic scenarios for use in CMIP6: a dataset of harmonized emissions trajectories through the end of the century, *Geoscientific Model Development*, 12, 1443–1475, <https://doi.org/10.5194/gmd-12-1443-2019>, 2019.

Gillett, N. P., Shiogama, H., Funke, B., Hegerl, G., Knutti, R., Matthes, K., Santer, B. D., Stone, D., and Tebaldi, C.: The Detection and Attribution Model Intercomparison Project (DAMIP v1.0) contribution to CMIP6, *Geoscientific Model Development*, 9, 3685–3697, <https://doi.org/10.5194/gmd-9-3685-2016>, publisher: Copernicus GmbH, 2016.

Gliß, J., Mortier, A., Schulz, M., Andrews, E., Balkanski, Y., Bauer, S. E., Benedictow, A. M. K., Bian, H., Checa-Garcia, R., Chin, M., Ginoux, P., Griesfeller, J. J., Heckel, A., Kipling, Z., Kirkevåg, A., Kokkola, H., Laj, P., Le Sager, P., Lund, M. T., Lund Myhre, C., Matsui, H., Myhre, G., Neubauer, D., van Noije, T., North, P., Olivie, D. J. L., Rémy, S., Sogacheva, L., Takemura, T., Tsigaridis, K., and Tsyro, S. G.: AeroCom phase III multi-model evaluation of the aerosol life cycle and optical properties using ground- and space-based remote sensing as well as surface in situ observations, *Atmospheric Chemistry and Physics*, 21, 87–128, <https://doi.org/10.5194/acp-21-87-2021>, 2021.

Golaz, J.-C., Caldwell, P. M., Van Roekel, L. P., Petersen, M. R., Tang, Q., Wolfe, J. D., Abeshu, G., Anantharaj, V., Asay-Davis, X. S., Bader, D. C., Baldwin, S. A., Bisht, G., Bogenschutz, P. A., Branstetter, M., Brunke, M. A., Brus, S. R., Burrows, S. M., Cameron-Smith, P. J., Donahue, A. S., Deakin, M., Easter, R. C., Evans, K. J., Feng, Y., Flanner, M., Foucar, J. G., Fyke, J. G., Griffin, B. M., Hannay, C., Harrop, B. E., Hoffman, M. J., Hunke, E. C., Jacob, R. L., Jacobsen, D. W., Jeffery, N., Jones, P. W., Keen, N. D., Klein, S. A., Larson, V. E., Leung, L. R., Li, H.-Y., Lin, W., Lipscomb, W. H., Ma, P.-L., Mahajan, S., Maltrud, M. E., Mametjanov, A., McClean, J. L., McCoy, R. B., Neale, R. B., Price, S. F., Qian, Y., Rasch, P. J., Reeves Eyre, J. E. J., Riley, W. J., Ringler, T. D., Roberts, A. F., Roesler, E. L., Salinger, A. G., Shaheen, Z., Shi, X., Singh, B., Tang, J., Taylor, M. A., Thornton, P. E., Turner, A. K., Veneziani, M., Wan, H., Wang, H., Wang, S., Williams, D. N., Wolfram, P. J., Worley, P. H., Xie, S., Yang, Y., Yoon, J.-H., Zelinka, M. D., Zender, C. S., Zeng, X., Zhang, C., Zhang, K., Zhang, Y., Zheng, X., Zhou, T., and Zhu, Q.: The DOE E3SM Coupled Model Version 1: Overview and Evaluation at Standard Resolution, *Journal of Advances in Modeling Earth Systems*, 11, 2089–2129, <https://doi.org/10.1029/2018MS001603>, 2019.

Griffiths, P., Murray, L., Zeng, G., Shin, Y. M., Abraham, N. L., Archibald, A. T., Deushi, M., Emmons, L. K., Galbally, I. E., Hassler, B., Horowitz, L. W., Keeble, J., Liu, J., Moeini, O., Naik, V., O'Connor, F. M., Oshima, N., Tarasick, D., Tilmes, S., Turnock, S. T., Wild, O., Young, P. J., and Zanis, P.: Tropospheric Ozone in CMIP6 Simulations, *Atmospheric Chemistry and Physics*, 21, 4187–4218, <https://doi.org/10.5194/acp-21-4187-2021>, 2021.

- Griffiths, P., Shin, Y., Keeble, J., and Archibald, A.: Tropospheric Ozone Budget in AerChemMIP Experiments, Tech. Rep. EGU23-7364, Copernicus Meetings, <https://doi.org/10.5194/egusphere-egu23-7364>, 2023.
- Guilyardi, E., Balaji, V., Lawrence, B., Callaghan, S., Deluca, C., Denvil, S., Lautenschlager, M., Morgan, M., Murphy, S., and Taylor, K. E.: Documenting Climate Models and Their Simulations, *Bulletin of the American Meteorological Society*, 94, 623–627, <https://doi.org/10.1175/BAMS-D-11-00035.1>, publisher: American Meteorological Society Section: Bulletin of the American Meteorological Society, 2013.
- Guo, H., John, J. G., Blanton, C., McHugh, C., Nikonov, S., Radhakrishnan, A., Rand, K., Zadeh, N. T., Balaji, V., Durachta, J., Dupuis, C., Menzel, R., Robinson, T., Underwood, S., Vahlenkamp, H., Bushuk, M., Dunne, K. A., Dussin, R., Gauthier, P. P., Ginoux, P., Griffies, S. M., Hallberg, R., Harrison, M., Hurlin, W., Lin, P., Malyshev, S., Naik, V., Paulot, F., Paynter, D. J., Ploshay, J., Reichl, B. G., Schwarzkopf, D. M., Seman, C. J., Shao, A., Silvers, L., Wyman, B., Yan, X., Zeng, Y., Adcroft, A., Dunne, J. P., Held, I. M., Krasting, J. P., Horowitz, L. W., Milly, P., Shevliakova, E., Winton, M., Zhao, M., and Zhang, R.: NOAA-GFDL GFDL-CM4 model output historical, <https://doi.org/10.22033/ESGF/CMIP6.8594>, 2018.
- Hajima, T., Abe, M., Arakawa, O., Suzuki, T., Komuro, Y., Ogura, T., Ogochi, K., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Ohgaito, R., Ito, A., Yamazaki, D., Ito, A., Takata, K., Watanabe, S., Kawamiya, M., and Tachiiri, K.: MIROC MIROC-ES2L model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.5602>, 2019.
- Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M.: Development of the MIROC-ES2L Earth system model and the evaluation of biogeochemical processes and feedbacks, *Geoscientific Model Development*, 13, 2197–2244, <https://doi.org/10.5194/gmd-13-2197-2020>, 2020.
- Hassan, T., Allen, R. J., Liu, W., Shim, S., van Noije, T., Le Sager, P., Oshima, N., Deushi, M., Randles, C. A., and O’Connor, F. M.: Air quality improvements are projected to weaken the Atlantic meridional overturning circulation through radiative forcing effects, *Communications Earth and Environment*, 3, 149, <https://doi.org/10.1038/s43247-022-00476-9>, 2022.
- HE, B., YU, Y., BAO, Q., LIN, P., LIU, H., LI, J., WANG, L., LIU, Y., WU, G., CHEN, K., GUO, Y., ZHAO, S., ZHANG, X., SONG, M., and XIE, J.: CAS FGOALS-f3-L model dataset descriptions for CMIP6 DECK experiments, *Atmospheric and Oceanic Science Letters*, 13, 582–588, <https://doi.org/10.1080/16742834.2020.1778419>, 2020.
- Held, I. M., Guo, H., Adcroft, A., Dunne, J. P., Horowitz, L. W., Krasting, J., Shevliakova, E., Winton, M., Zhao, M., Bushuk, M., Wittenberg, A. T., Wyman, B., Xiang, B., Zhang, R., Anderson, W., Balaji, V., Donner, L., Dunne, K., Durachta, J., Gauthier, P. P. G., Ginoux, P., Golaz, J.-C., Griffies, S. M., Hallberg, R., Harris, L., Harrison, M., Hurlin, W., John, J., Lin, P., Lin, S.-J., Malyshev, S., Menzel, R., Milly, P. C. D., Ming, Y., Naik, V., Paynter, D., Paulot, F., Rammaswamy, V., Reichl, B., Robinson, T., Rosati, A., Seman, C., Silvers, L. G., Underwood, S., and Zadeh, N.: Structure and Performance of GFDL’s CM4.0 Climate Model, *Journal of Advances in Modeling Earth Systems*, 11, 3691–3727, <https://doi.org/https://doi.org/10.1029/2019MS001829>, 2019.
- Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitkanen, T., Seibert, J. J., Vu, L., Andres, R. J., Bolt, R. M., Bond, T. C., Dawidowski, L., Kholod, N., Kurokawa, J.-I., Li, M., Liu, L., Lu, Z., Moura, M. C. P., O’Rourke, P. R., and Zhang, Q.: Historical (1750–2014) anthropogenic emissions of reactive gases and aerosols from the Community Emissions Data System (CEDS), *Geoscientific Model Development*, 11, 369–408, <https://doi.org/10.5194/gmd-11-369-2018>, 2018.
- Holland, M. M., Hannay, C., Fasullo, J., Jahn, A., Kay, J. E., Mills, M., Simpson, I. R., Wieder, W., Lawrence, P., Kluzek, E., and Bailey, D.: New model ensemble reveals how forcing uncertainty and model structure alter climate simulated across CMIP generations of the Community Earth System Model, *Geoscientific Model Development*, 17, 1585–1602, <https://doi.org/10.5194/gmd-17-1585-2024>, 2024.

- Huang, W.: THU CIESM model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.8843>, 2019.
- 1130 IPCC: Summary for policymakers, in: Climate change 2021: The physical science basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, UK and New York, NY, USA, <https://doi.org/10.1017/9781009157896.001>, type: Book section, 2021.
- 1135 Jungclaus, J., Bittner, M., Wieners, K.-H., Wachsmann, F., Schupfner, M., Legutke, S., Giorgetta, M., Reick, C., Gayler, V., Haak, H., de Vrese, P., Raddatz, T., Esch, M., Mauritsen, T., von Storch, J.-S., Behrens, J., Brovkin, V., Claussen, M., Crueger, T., Fast, I., Fiedler, S., Hagemann, S., Hohenegger, C., Jahns, T., Kloster, S., Kinne, S., Lasslop, G., Kornblueh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Müller, W., Nabel, J., Notz, D., Peters-von Gehlen, K., Pincus, R., Pohlmann, H., Pongratz, J., Rast, S., Schmidt, H., Schnur, R., Schulzweida, U., Six, K., Stevens, B., Voigt, A., and Roeckner, E.: MPI-M MPI-ESM1.2-HR model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.6594>, 2019.
- 1140 Kalisoras, A., Georgoulas, A. K., Akritidis, D., Allen, R. J., Naik, V., Kuo, C., Szopa, S., Nabat, P., Olivié, D., van Noije, T., Le Sager, P., Neubauer, D., Oshima, N., Mulcahy, J., Horowitz, L. W., and Zanis, P.: Decomposing the Effective Radiative Forcing of anthropogenic aerosols based on CMIP6 Earth System Models, *EGUsphere*, 2023, 1–38, <https://doi.org/10.5194/egusphere-2023-2571>, 2023.
- Karset, I. H. H., Berntsen, T. K., Storelvmo, T., Alterskjær, K., Grini, A., Olivié, D., Kirkevåg, A., Seland, \, Iversen, T., and Schulz, M.: Strong impacts on aerosol indirect effects from historical oxidant changes, *Atmospheric Chemistry and Physics*, 18, 7669–7690, <https://doi.org/10.5194/acp-18-7669-2018>, 2018.
- 1145 Kelley, M., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Del Genio, A. D., Elsaesser, G. S., Faluvegi, G., Kiang, N. Y., Kim, D., Lacis, A. A., Leboissetier, A., LeGrande, A. N., Lo, K. K., Marshall, J., Matthews, E. E., McDermid, S., Mezuman, K., Miller, R. L., Murray, L. T., Oinas, V., Orbe, C., García-Pando, C. P., Perlwitz, J. P., Puma, M. J., Rind, D., Romanou, A., Shindell, D. T., Sun, S., Tausnev, N., Tsigaridis, K., Tselioudis, G., Weng, E., Wu, J., and Yao, M.-S.: GISS-E2.1: Configurations and Climatology, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002025, <https://doi.org/https://doi.org/10.1029/2019MS002025>, e2019MS002025 10.1029/2019MS002025, 2020.
- Kim, Y., Noh, Y., Kim, D., Lee, M.-I., Lee, H. J., Kim, S. Y., and Kim, D.: KIOST KIOST-ESM model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.5296>, 2019.
- 1155 Kok, J. F., Adebisi, A. A., Albani, S., Balkanski, Y., Checa-Garcia, R., Chin, M., Colarco, P. R., Hamilton, D. S., Huang, Y., Ito, A., Klose, M., Leung, D. M., Li, L., Mahowald, N. M., Miller, R. L., Obiso, V., Pérez García-Pando, C., Rocha-Lima, A., Wan, J. S., and Whicker, C. A.: Improved representation of the global dust cycle using observational constraints on dust properties and abundance, *Atmospheric Chemistry and Physics*, 21, 8127–8167, <https://doi.org/10.5194/acp-21-8127-2021>, 2021.
- 1160 Krasting, J. P., John, J. G., Blanton, C., McHugh, C., Nikonov, S., Radhakrishnan, A., Rand, K., Zadeh, N. T., Balaji, V., Durachta, J., Dupuis, C., Menzel, R., Robinson, T., Underwood, S., Vahlenkamp, H., Dunne, K. A., Gauthier, P. P., Ginoux, P., Griffies, S. M., Hallberg, R., Harrison, M., Hurlin, W., Malyshev, S., Naik, V., Paulot, F., Paynter, D. J., Ploshay, J., Reichl, B. G., Schwarzkopf, D. M., Seman, C. J., Silvers, L., Wyman, B., Zeng, Y., Adcroft, A., Dunne, J. P., Dussin, R., Guo, H., He, J., Held, I. M., Horowitz, L. W., Lin, P., Milly, P., Shevliakova, E., Stock, C., Winton, M., Wittenberg, A. T., Xie, Y., and Zhao, M.: NOAA-GFDL GFDL-ESM4 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.8597>, 2018.
- 1165

- Kuhlbrodt, T., Jones, C. G., Sellar, A., Storkey, D., Blockley, E., Stringer, M., Hill, R., Graham, T., Ridley, J., Blaker, A., Calvert, D., Copsey, D., Ellis, R., Hewitt, H., Hyder, P., Ineson, S., Mulcahy, J., Siahaan, A., and Walton, J.: The Low-Resolution Version of HadGEM3 GC3.1: Development and Evaluation for Global Climate, *Journal of Advances in Modeling Earth Systems*, 10, 2865–2888, <https://doi.org/https://doi.org/10.1029/2018MS001370>, 2018.
- 1170 Lamarque, J.-F., Shindell, D. T., Josse, B., Young, P. J., Cionni, I., Eyring, V., Bergmann, D., Cameron-Smith, P., Collins, W. J., Doherty, R., Dalsoren, S., Faluvegi, G., Folberth, G., Ghan, S. J., Horowitz, L. W., Lee, Y. H., MacKenzie, I. A., Nagashima, T., Naik, V., Plummer, D., Righi, M., Rumbold, S. T., Schulz, M., Skeie, R. B., Stevenson, D. S., Strode, S., Sudo, K., Szopa, S., Voulgarakis, A., and Zeng, G.: The Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP): overview and description of models, simulations and climate diagnostics, *Geoscientific Model Development*, 6, 179–206, <https://doi.org/10.5194/gmd-6-179-2013>, publisher: Copernicus GmbH, 2013.
- Lawrence, D. M., Hurtt, G. C., Arneth, A., Brovkin, V., Calvin, K. V., Jones, A. D., Jones, C. D., Lawrence, P. J., de Noblet-Ducoudré, N., Pongratz, J., Seneviratne, S. I., and Shevliakova, E.: The Land Use Model Intercomparison Project (LUMIP) contribution to CMIP6: rationale and experimental design, *Geoscientific Model Development*, 9, 2973–2998, <https://doi.org/10.5194/gmd-9-2973-2016>, 2016.
- Leach, N. J., Jenkins, S., Nicholls, Z., Smith, C. J., Lynch, J., Cain, M., Walsh, T., Wu, B., Tsutsui, J., and Allen, M. R.: FaIRv2.0.0: a generalized impulse response model for climate uncertainty and future scenario exploration, *Geoscientific Model Development*, 14, 3007–3036, <https://doi.org/10.5194/gmd-14-3007-2021>, publisher: Copernicus GmbH, 2021.
- 1180 Lee, J., Kim, J., Sun, M.-A., Kim, B.-H., Moon, H., Sung, H. M., Kim, J., and Byun, Y.-H.: Evaluation of the Korea Meteorological Administration Advanced Community Earth-System model (K-ACE), *Asia-Pacific Journal of Atmospheric Sciences*, 56, 381–395, <https://doi.org/10.1007/s13143-019-00144-7>, 2020a.
- 1185 Lee, J., Gleckler, P. J., Ahn, M.-S., Ordonez, A., Ullrich, P. A., Sperber, K. R., Taylor, K. E., Planton, Y. Y., Guilyardi, E., Durack, P., Bonfils, C., Zelinka, M. D., Chao, L.-W., Dong, B., Doutriaux, C., Zhang, C., Vo, T., Boutte, J., Wehner, M. F., Pendergrass, A. G., Kim, D., Xue, Z., Wittenberg, A. T., and Krasting, J.: Systematic and objective evaluation of Earth system models: PCMDI Metrics Package (PMP) version 3, *Geoscientific Model Development*, 17, 3919–3948, <https://doi.org/10.5194/gmd-17-3919-2024>, 2024.
- Lee, L. A., Carslaw, K. S., Pringle, K. J., Mann, G. W., and Spracklen, D. V.: Emulation of a complex global aerosol model to quantify sensitivity to uncertain parameters, *Atmospheric Chemistry and Physics*, 11, 12 253–12 273, <https://doi.org/10.5194/acp-11-12253-2011>, publisher: Copernicus GmbH, 2011.
- 1190 Lee, W.-L. and Liang, H.-C.: AS-RCEC TaiESM1.0 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.9755>, 2020.
- Lee, W.-L., Wang, Y.-C., Shiu, C.-J., Tsai, I., Tu, C.-Y., Lan, Y.-Y., Chen, J.-P., Pan, H.-L., and Hsu, H.-H.: Taiwan Earth System Model Version 1: description and evaluation of mean state, *Geoscientific Model Development*, 13, 3887–3904, <https://doi.org/10.5194/gmd-13-3887-2020>, 2020b.
- 1195 Li, L.: CAS FGOALS-g3 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.3356>, 2019.
- Li, L., Yu, Y., Tang, Y., Lin, P., Xie, J., Song, M., Dong, L., Zhou, T., Liu, L., Wang, L., Pu, Y., Chen, X., Chen, L., Xie, Z., Liu, H., Zhang, L., Huang, X., Feng, T., Zheng, W., Xia, K., Liu, H., Liu, J., Wang, Y., Wang, L., Jia, B., Xie, F., Wang, B., Zhao, S., Yu, Z., Zhao, B., and Wei, J.: The Flexible Global Ocean-Atmosphere-Land System Model Grid-Point Version 3 (FGOALS-g3): Description and Evaluation, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002 012, <https://doi.org/https://doi.org/10.1029/2019MS002012>, e2019MS002012 2019MS002012, 2020.
- 1200

- Li, Y., Wang, Z., Lei, Y., Che, H., and Zhang, X.: Impacts of reductions in non-methane short-lived climate forcers on future climate extremes and the resulting population exposure risks in Asia, *Atmospheric Chemistry and Physics Discussions*, pp. 1–32, <https://doi.org/10.5194/acp-2022-422>, publisher: Copernicus GmbH, 2022.
- Lin, Y., Huang, X., Liang, Y., Qin, Y., Xu, S., Huang, W., Xu, F., Liu, L., Wang, Y., Peng, Y., Wang, L., Xue, W., Fu, H., Zhang, G. J., Wang, B., Li, R., Zhang, C., Lu, H., Yang, K., Luo, Y., Bai, Y., Song, Z., Wang, M., Zhao, W., Zhang, F., Xu, J., Zhao, X., Lu, C., Chen, Y., Luo, Y., Hu, Y., Tang, Q., Chen, D., Yang, G., and Gong, P.: Community Integrated Earth System Model (CIESM): Description and Evaluation, *Journal of Advances in Modeling Earth Systems*, 12, e2019MS002 036, <https://doi.org/https://doi.org/10.1029/2019MS002036>, e2019MS002036 2019MS002036, 2020.
- Liu, L., Shawki, D., Voulgarakis, A., Kasoar, M., Samset, B. H., Myhre, G., Forster, P. M., Hodnebrog, Ø., Sillmann, J., Aalbergstjø, S. G., Boucher, O., Faluvegi, G., Iversen, T., Kirkevåg, A., Lamarque, J.-F., Olivie, D., Richardson, T., Shindell, D., and Takemura, T.: A PDRMIP Multimodel Study on the Impacts of Regional Aerosol Forcings on Global and Regional Precipitation, *Journal of Climate*, 31, 4429 – 4447, <https://doi.org/10.1175/JCLI-D-17-0439.1>, 2018.
- Liu, Z., Doherty, R. M., Wild, O., O'Connor, F. M., and Turnock, S. T.: Correcting ozone biases in a global chemistry-climate model: implications for future ozone, *Atmospheric Chemistry and Physics Discussions*, pp. 1–20, <https://doi.org/10.5194/acp-2022-196>, publisher: Copernicus GmbH, 2022.
- Lovato, T. and Peano, D.: CMCC CMCC-CM2-SR5 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.3825>, 2020.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenez-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J.-S., Tian, F., Voigt, A., Vrese, P., Wieners, K.-H., Wilkenskeld, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO₂, *Journal of Advances in Modeling Earth Systems*, 11, 998–1038, <https://doi.org/https://doi.org/10.1029/2018MS001400>, 2019.
- Meehl, G. A., Moss, R., Taylor, K. E., Eyring, V., Stouffer, R. J., Bony, S., and Stevens, B.: Climate Model Intercomparisons: Preparing for the Next Phase, *Eos, Transactions American Geophysical Union*, 95, 77–78, <https://doi.org/10.1002/2014EO090001>, <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2014EO090001>, 2014.
- Melton, J. R., Wania, R., Hodson, E. L., Poulter, B., Ringeval, B., Spahni, R., Bohn, T., Avis, C. A., Beerling, D. J., Chen, G., Eliseev, A. V., Denisov, S. N., Hopcroft, P. O., Lettenmaier, D. P., Riley, W. J., Singarayer, J. S., Subin, Z. M., Tian, H., Zürcher, S., Brovkin, V., van Bodegom, P. M., Kleinen, T., Yu, Z. C., and Kaplan, J. O.: Present state of global wetland extent and wetland methane modelling: conclusions from a model inter-comparison project (WETCHIMP), *Biogeosciences*, 10, 753–788, <https://doi.org/10.5194/bg-10-753-2013>, 2013.
- Menary, M. B., Robson, J., Allan, R. P., Booth, B. B. B., Cassou, C., Gastineau, G., Gregory, J., Hodson, D., Jones, C., Mignot, J., Ringer, M., Sutton, R., Wilcox, L., and Zhang, R.: Aerosol-Forced AMOC Changes in CMIP6 Historical Simulations, *Geophysical Research Letters*, 47, e2020GL088 166, <https://doi.org/https://doi.org/10.1029/2020GL088166>, e2020GL088166 2020GL088166, 2020.

- 1240 Miller, R. L., Schmidt, G. A., Nazarenko, L. S., Bauer, S. E., Kelley, M., Ruedy, R., Russell, G. L., Ackerman, A. S., Aleinov, I., Bauer, M., Bleck, R., Canuto, V., Cesana, G., Cheng, Y., Clune, T. L., Cook, B. I., Cruz, C. A., Del Genio, A. D., Elsaesser, G. S., Faluvegi, G., Kiang, N. Y., Kim, D., Lacis, A. A., Leboissetier, A., LeGrande, A. N., Lo, K. K., Marshall, J., Matthews, E. E., McDermid, S., Mezuman, K., Murray, L. T., Oinas, V., Orbe, C., Pérez García-Pando, C., Perlwitz, J. P., Puma, M. J., Rind, D., Romanou, A., Shindell, D. T., Sun, S., Tausnev, N., Tsigaridis, K., Tselioudis, G., Weng, E., Wu, J., and Yao, M.-S.: CMIP6 Historical Simulations (1850–2014) With GISS-E2.1, *Journal of Advances in Modeling Earth Systems*, 13, e2019MS002034, <https://doi.org/https://doi.org/10.1029/2019MS002034>, e2019MS002034 10.1029/2019MS002034, 2021.
- 1245 Monerie, P.-A., Robson, J. I., Dunstone, N. J., and Turner, A. G.: Skilful seasonal predictions of global monsoon summer precipitation with DePreSys3, *Environmental Research Letters*, 16, 104035, <https://doi.org/10.1088/1748-9326/ac2a65>, publisher: IOP Publishing, 2021.
- Morgenstern, O., O'Connor, F. M., Johnson, B. T., Zeng, G., Mulcahy, J. P., Williams, J., Teixeira, J., Michou, M., Nabat, P., Horowitz, L. W., Naik, V., Sentman, L. T., Deushi, M., Bauer, S. E., Tsigaridis, K., Shindell, D. T., and Kinnison, D. E.: Reappraisal of the Climate Impacts of Ozone-Depleting Substances, *Geophysical Research Letters*, 47, e2020GL088295, <https://doi.org/10.1029/2020GL088295>, [eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020GL088295](https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020GL088295), 2020.
- 1250 Mortier, A., Gliß, J., Schulz, M., Aas, W., Andrews, E., Bian, H., Chin, M., Ginoux, P., Hand, J., Holben, B., Zhang, H., Kipling, Z., Kirkevåg, A., Laj, P., Lurton, T., Myhre, G., Neubauer, D., Olivié, D., von Salzen, K., Skeie, R. B., Takemura, T., and Tilmes, S.: Evaluation of climate model aerosol trends with ground-based observations over the last 2 decades – an AeroCom and CMIP6 analysis, *Atmospheric Chemistry and Physics*, 20, 13355–13378, <https://doi.org/10.5194/acp-20-13355-2020>, 2020.
- 1255 Moseid, K. O., Schulz, M., Storelvmo, T., Julsrud, I. R., Olivié, D., Nabat, P., Wild, M., Cole, J. N. S., Takemura, T., Oshima, N., Bauer, S. E., and Gastineau, G.: Bias in CMIP6 models as compared to observed regional dimming and brightening, *Atmospheric Chemistry and Physics*, 20, 16023–16040, <https://doi.org/10.5194/acp-20-16023-2020>, publisher: Copernicus GmbH, 2020.
- 1260 Murray, L. T., Logan, J. A., and Jacob, D. J.: Interannual variability in tropical tropospheric ozone and OH: The role of lightning: IAV IN OZONE AND OH-ROLE OF LIGHTNING, *Journal of Geophysical Research: Atmospheres*, 118, 11468–11480, <https://doi.org/10.1002/jgrd.50857>, 2013.
- Myhre, G., Forster, P. M., Samset, B. H., Hodnebrog, Ø., Sillmann, J., Aalberg, S. G., Andrews, T., Boucher, O., Faluvegi, G., Fläschner, D., Iversen, T., Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Olivié, D., Richardson, T. B., Shindell, D., Shine, K. P., Stjern, C. W., Takemura, T., Voulgarakis, A., and Zwiers, F.: PDRMIP: A Precipitation Driver and Response Model Intercomparison Project—Protocol and Preliminary Results, *Bulletin of the American Meteorological Society*, 98, 1185–1198, <https://doi.org/10.1175/BAMS-D-16-0019.1>, publisher: American Meteorological Society Section: Bulletin of the American Meteorological Society, 2017.
- 1265 NASA/GISS, N. G. I. f. S. S.: NASA-GISS GISS-E2.1G model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.7127>, 2018.
- NASA/GISS, N. G. I. f. S. S.: NASA-GISS GISS-E2-1-G-CC model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.11762>, 2019a.
- NASA/GISS, N. G. I. f. S. S.: NASA-GISS GISS-E2.1H model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.7128>, 2019b.
- 1275 Neubauer, D., Ferrachat, S., Siegenthaler-Le Drian, C., Stoll, J., Folini, D. S., Tegen, I., Wieners, K.-H., Mauritsen, T., Stemmler, I., Barthel, S., Bey, I., Daskalakis, N., Heinold, B., Kokkola, H., Partridge, D., Rast, S., Schmidt, H., Schutgens, N., Stanelle, T., Stier,

- P., Watson-Parris, D., and Lohmann, U.: HAMMOZ-Consortium MPI-ESM1.2-HAM model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.5016>, 2019.
- O'Connor, F. M., Abraham, N. L., Dalvi, M., Folberth, G. A., Griffiths, P. T., Hardacre, C., Johnson, B. T., Kahana, R., Keeble, J., Kim, B.,
1280 Morgenstern, O., Mulcahy, J. P., Richardson, M., Robertson, E., Seo, J., Shim, S., Teixeira, J. C., Turnock, S. T., Williams, J., Wiltshire, A. J., Woodward, S., and Zeng, G.: Assessment of pre-industrial to present-day anthropogenic climate forcing in UKESM1, *Atmos. Chem. Phys.*, 21, 1211–1243, <https://doi.org/10.5194/acp-21-1211-2021>, publisher: Copernicus Publications, 2021.
- O'Connor, F. M., Johnson, B. T., Jamil, O., Andrews, T., Mulcahy, J. P., and Manners, J.: Apportionment of the Pre-Industrial to Present-Day Climate Forcing by Methane Using UKESM1: The Role of the Cloud Radiative Effect, *Journal of Advances in Modeling Earth Systems*,
1285 14, e2022MS002991, <https://doi.org/https://doi.org/10.1029/2022MS002991>, e2022MS002991 2022MS002991, 2022.
- O'Neill, B. C., Tebaldi, C., van Vuuren, D. P., Eyring, V., Friedlingstein, P., Hurtt, G., Knutti, R., Krieger, E., Lamarque, J.-F., Lowe, J., Meehl, G. A., Moss, R., Riahi, K., and Sanderson, B. M.: The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6, *Geoscientific Model Development*, 9, 3461–3482, <https://doi.org/10.5194/gmd-9-3461-2016>, publisher: Copernicus GmbH, 2016.
- Oshima, N., Yukimoto, S., Deushi, M., Koshiro, T., Kawai, H., Tanaka, T. Y., and Yoshida, K.: Global and Arctic effective radiative forcing
1290 of anthropogenic gases and aerosols in MRI-ESM2.0, *Progress in Earth and Planetary Science*, 7, 38, <https://doi.org/10.1186/s40645-020-00348-w>, 2020.
- Pak, G., Noh, Y., Lee, M.-I., Yeh, S.-W., Kim, D., Kim, S.-Y., Lee, J.-L., Lee, H. J., Hyun, S.-H., Lee, K.-Y., Lee, J.-H., Park, Y.-G., Jin, H., Park, H., and Kim, Y. H.: Korea Institute of Ocean Science and Technology Earth System Model and Its Simulation Characteristics, *Ocean Science Journal*, 56, 18–45, <https://doi.org/10.1007/s12601-021-00001-7>, 2021.
- 1295 Park, S. and Shin, J.: SNU SAM0-UNICON model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.7789>, 2019.
- Park, S., Shin, J., Kim, S., Oh, E., and Kim, Y.: Global Climate Simulated by the Seoul National University Atmosphere Model Version 0 with a Unified Convection Scheme (SAM0-UNICON), *Journal of Climate*, 32, 2917 – 2949, <https://doi.org/10.1175/JCLI-D-18-0796.1>, 2019.
- 1300 Pascoe, C., Lawrence, B. N., Guilyardi, E., Juckes, M., and Taylor, K. E.: Documenting numerical experiments in support of the Coupled Model Intercomparison Project Phase 6 (CMIP6), *Geoscientific Model Development*, 13, 2149–2167, <https://doi.org/10.5194/gmd-13-2149-2020>, publisher: Copernicus GmbH, 2020.
- Persad, G., Samset, B. H., Wilcox, L. J., Allen, R. J., Bollasina, M. A., Booth, B. B. B., Bonfils, C., Crocker, T., Joshi, M., Lund, M. T., Marvel, K., Merikanto, J., Nordling, K., Undorf, S., van Vuuren, D. P., Westervelt, D. M., and Zhao, A.: Rapidly evolving aerosol emissions are a
1305 dangerous omission from near-term climate risk assessments, *Environmental Research: Climate*, 2, 032 001, <https://doi.org/10.1088/2752-5295/acd6af>, 2023.
- Pincus, R., Forster, P. M., and Stevens, B.: The Radiative Forcing Model Intercomparison Project (RFMIP): experimental protocol for CMIP6, *Geoscientific Model Development*, 9, 3447–3460, <https://doi.org/10.5194/gmd-9-3447-2016>, publisher: Copernicus GmbH, 2016.
- Prather, M. J., Zhu, X., Tang, Q., Hsu, J., and Neu, J. L.: An atmospheric chemist in search of the tropopause, *Journal of Geophysical
1310 Research*, 116, D04 306, <https://doi.org/10.1029/2010JD014939>, 2011.
- Rao, S., Klimont, Z., Smith, S. J., Van Dingenen, R., Dentener, F., Bouwman, L., Riahi, K., Amann, M., Bodirsky, B. L., van Vuuren, D. P., Aleluia Reis, L., Calvin, K., Drouet, L., Fricko, O., Fujimori, S., Gernaat, D., Havlik, P., Harmsen, M., Hasegawa, T., Heyes, C., Hilaire, J., Luderer, G., Masui, T., Stehfest, E., Streffer, J., van der Sluis, S., and Tavoni, M.: Future air pollution in the Shared Socio-economic Pathways, *Global Environmental Change*, 42, 346–358, <https://doi.org/10.1016/j.gloenvcha.2016.05.012>, 2017.

- 1315 Riahi, K., van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., Bauer, N., Calvin, K., Dellink, R., Fricko, O., Lutz, W., Popp, A., Cuaresma, J. C., Kc, S., Leimbach, M., Jiang, L., Kram, T., Rao, S., Emmerling, J., Ebi, K., Hasegawa, T., Havlik, P., Humpenöder, F., Da Silva, L. A., Smith, S., Stehfest, E., Bosetti, V., Eom, J., Gernaat, D., Masui, T., Rogelj, J., Strefler, J., Drouet, L., Krey, V., Luderer, G., Harmsen, M., Takahashi, K., Baumstark, L., Doelman, J. C., Kainuma, M., Klimont, Z., Marangoni, G., Lotze-Campen, H., Obersteiner, M., Tabeau, A., and Tavoni, M.: The Shared Socioeconomic Pathways and their energy, land use, and greenhouse
- 1320 gas emissions implications: An overview, *Global Environmental Change*, 42, 153–168, <https://doi.org/10.1016/j.gloenvcha.2016.05.009>, 2017.
- Ridley, J., Menary, M., Kuhlbrodt, T., Andrews, M., and Andrews, T.: MOHC HadGEM3-GC31-LL model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.6109>, 2019a.
- Ridley, J., Menary, M., Kuhlbrodt, T., Andrews, M., and Andrews, T.: MOHC HadGEM3-GC31-MM model output prepared for CMIP6
- 1325 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.6112>, 2019b.
- Rong, X.: CAMS CAMS_CSM1.0 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.9754>, 2019.
- Rong, X., Li, J., Chen, H., Xin, Y., Su, J., Hua, L., Zhou, T., Qi, Y., Zhang, Z., Zhang, G., and Li, J.: The CAMS Climate System Model and a Basic Evaluation of Its Climatology and Climate Variability Simulation, *Journal of Meteorological Research*, 32, 839–861, <https://doi.org/10.1007/s13351-018-8058-x>, 2018.
- 1330 Samset, B. H., Myhre, G., Forster, P. M., Hodnebrog, Ø., Andrews, T., Faluvegi, G., Fläschner, D., Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Olivié, D., Richardson, T., Shindell, D., Shine, K. P., Takemura, T., and Voulgarakis, A.: Fast and slow precipitation responses to individual climate forcings: A PDRMIP multimodel study, *Geophysical Research Letters*, 43, 2782–2791, <https://doi.org/https://doi.org/10.1002/2016GL068064>, 2016.
- Schlund, M., Lauer, A., Gentile, P., Sherwood, S. C., and Eyring, V.: Emergent constraints on equilibrium climate sensitivity in CMIP5: do
- 1335 they hold for CMIP6?, *Earth System Dynamics*, 11, 1233–1258, <https://doi.org/10.5194/esd-11-1233-2020>, 2020.
- Schlund, M., Hassler, B., Lauer, A., Andela, B., Jöckel, P., Kazeroni, R., Loosveldt Tomas, S., Medeiros, B., Predoi, V., Sénési, S., Servonnat, J., Stacke, T., Vegas-Regidor, J., Zimmermann, K., and Eyring, V.: Evaluation of native Earth system model output with ESMValTool v2.6.0, *Geoscientific Model Development*, 16, 315–333, <https://doi.org/10.5194/gmd-16-315-2023>, publisher: Copernicus GmbH, 2023.
- Scoccimarro, E., Bellucci, A., and Peano, D.: CMCC CMCC-CM2-HR4 model output prepared for CMIP6 CMIP historical,
- 1340 <https://doi.org/10.22033/ESGF/CMIP6.3823>, 2020.
- Seferian, R.: CNRM-CERFACS CNRM-ESM2-1 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.4068>, 2018.
- Séférian, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C., Berthet, S., Chevallier, M., Sénési, S., Franchisteguy, L., Vial, J., Mallet, M., Joetzjer, E., Geoffroy, O., Guérémy, J.-F., Moine, M.-P., Msadek, R., Ribes, A., Rocher, M.,
- 1345 Roehrig, R., Salas-y Mélia, D., Sanchez, E., Terray, L., Valcke, S., Waldman, R., Aumont, O., Bopp, L., Deshayes, J., Éthé, C., and Madec, G.: Evaluation of CNRM Earth System Model, CNRM-ESM2-1: Role of Earth System Processes in Present-Day and Future Climate, *Journal of Advances in Modeling Earth Systems*, 11, 4182–4227, <https://doi.org/https://doi.org/10.1029/2019MS001791>, 2019.
- Seland, Ø., Bentsen, M., Olivié, D., Toniazzi, T., Gjermundsen, A., Graff, L. S., Debernard, J. B., Gupta, A. K., He, Y.-C., Kirkevåg, A., Schwinger, J., Tjiputra, J., Aas, K. S., Bethke, I., Fan, Y., Griesfeller, J., Grini, A., Guo, C., Ilicak, M., Karset, I. H. H., Landgren,
- 1350 O., Liakka, J., Moseid, K. O., Nummelin, A., Spensberger, C., Tang, H., Zhang, Z., Heinze, C., Iversen, T., and Schulz, M.: Overview of the Norwegian Earth System Model (NorESM2) and key climate response of CMIP6 DECK, historical, and scenario simulations, *Geoscientific Model Development*, 13, 6165–6200, <https://doi.org/10.5194/gmd-13-6165-2020>, 2020.

- Seland, y., Bentsen, M., Olivie, D. J. L., Toniazzo, T., Gjermundsen, A., Graff, L. S., Debernard, J. B., Gupta, A. K., He, Y., Kirkevåg, A., Schwinger, J., Tjiputra, J., Aas, K. S., Bethke, I., Fan, Y., Griesfeller, J., Grini, A., Guo, C., Ilicak, M., Karset, I. H. H., Landgren, O. A., Liakka, J., Moseid, K. O., Nummelin, A., Spensberger, C., Tang, H., Zhang, Z., Heinze, C., Iversen, T., and Schulz, M.: NCC NorESM2-LM model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.8036>, 2019.
- 1355 Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O'Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., de Mora, L., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahhaan, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G., and Zerroukat, M.: UKESM1: Description and Evaluation of the U.K. Earth System Model, *Journal of Advances in Modeling Earth Systems*, 11, 4513–4558, <https://doi.org/https://doi.org/10.1029/2019MS001739>, 2019.
- 1360 Shim, S., Sung, H., Kwon, S., Kim, J., Lee, J., Sun, M., Song, J., Ha, J., Byun, Y., Kim, Y., Turnock, S. T., Stevenson, D. S., Allen, R. J., O'Connor, F. M., Teixeira, J. C., Williams, J., Johnson, B., Keeble, J., Mulcahy, J., and Zeng, G.: Regional Features of Long-Term Exposure to PM_{2.5} Air Quality over Asia under SSP Scenarios Based on CMIP6 Models, *International Journal of Environmental Research and Public Health*, 18, <https://doi.org/10.3390/ijerph18136817>, 2021.
- Shindell, D. T., Lamarque, J.-F., Schulz, M., Flanner, M., Jiao, C., Chin, M., Young, P. J., Lee, Y. H., Rotstajn, L., Mahowald, N., Milly, G., Faluvegi, G., Balkanski, Y., Collins, W. J., Conley, A. J., Dalsoren, S., Easter, R., Ghan, S., Horowitz, L., Liu, X., Myhre, G., Nagashima, T., Naik, V., Rumbold, S. T., Skeie, R., Sudo, K., Szopa, S., Takemura, T., Voulgarakis, A., Yoon, J.-H., and Lo, F.: Radiative forcing in the ACCMIP historical and future climate simulations, *Atmospheric Chemistry and Physics*, 13, 2939–2974, <https://doi.org/10.5194/acp-13-2939-2013>, publisher: Copernicus GmbH, 2013.
- 1370 Shonk, J. K. P., Turner, A. G., Chevuturi, A., Wilcox, L. J., Dittus, A. J., and Hawkins, E.: Uncertainty in aerosol radiative forcing impacts the simulated global monsoon in the 20th century, *Atmospheric Chemistry and Physics*, 20, 14 903–14 915, <https://doi.org/10.5194/acp-20-14903-2020>, 2020.
- 1375 Sidorenko, D., Rackow, T., Jung, T., Semmler, T., Barbi, D., Danilov, S., Dethloff, K., Dorn, W., Fieg, K., Goessling, H. F., Handorf, D., Harig, S., Hiller, W., Juricke, S., Losch, M., Schröter, J., Sein, D. V., and Wang, Q.: Towards multi-resolution global climate modeling with ECHAM6–FESOM. Part I: model formulation and mean climate, *Climate Dynamics*, 44, 757–780, <https://doi.org/10.1007/s00382-014-2290-6>, 2015.
- 1380 Simpson, I. R., Rosenbloom, N., Danabasoglu, G., Deser, C., Yeager, S. G., McCluskey, C. S., Yamaguchi, R., Lamarque, J.-F., Tilmes, S., Mills, M. J., and Rodgers, K. B.: The CESM2 Single-Forcing Large Ensemble and Comparison to CESM1: Implications for Experimental Design, *Journal of Climate*, 36, 5687–5711, <https://doi.org/10.1175/JCLI-D-22-0666.1>, publisher: American Meteorological Society Section: Journal of Climate, 2023.
- 1385 Skeie, R. B., Myhre, G., Hodnebrog, Ø., Cameron-Smith, P. J., Deushi, M., Hegglin, M. I., Horowitz, L. W., Kramer, R. J., Michou, M., Mills, M. J., Olivie, D. J. L., Connor, F. M. O., Paynter, D., Samset, B. H., Sellar, A., Shindell, D., Takemura, T., Tilmes, S., and Wu, T.: Historical total ozone radiative forcing derived from CMIP6 simulations, *npj Climate and Atmospheric Science*, 3, 1–10, <https://doi.org/10.1038/s41612-020-00131-0>, number: 1 Publisher: Nature Publishing Group, 2020.
- Skeie, R. B., Hodnebrog, Ø., and Myhre, G.: Trends in atmospheric methane concentrations since 1990 were driven and modified by anthropogenic emissions, *Communications Earth & Environment*, 4, 317, <https://doi.org/10.1038/s43247-023-00969-1>, 2023.

- 1390 Smith, C., Cummins, D. P., Fredriksen, H.-B., Nicholls, Z., Meinshausen, M., Allen, M., Jenkins, S., Leach, N., Mathison, C., and Partanen, A.-I.: *fair-calibrate v1.4.1: calibration, constraining, and validation of the FaIR simple climate model for reliable future climate projections*, *Geoscientific Model Development*, 17, 8569–8592, <https://doi.org/10.5194/gmd-17-8569-2024>, 2024.
- Smith, C. J., Forster, P. M., Allen, M., Leach, N., Millar, R. J., Passerello, G. A., and Regayre, L. A.: FAIR v1.3: a simple emissions-based impulse response and carbon cycle model, *Geoscientific Model Development*, 11, 2273–2297, <https://doi.org/10.5194/gmd-11-2273-2018>, publisher: Copernicus GmbH, 2018.
- 1395 Smith, C. J., Harris, G. R., Palmer, M. D., Bellouin, N., Collins, W., Myhre, G., Schulz, M., Golaz, J.-C., Ringer, M., Storelvmo, T., and Forster, P. M.: Energy Budget Constraints on the Time History of Aerosol Forcing and Climate Sensitivity, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033622, <https://doi.org/10.1029/2020JD033622>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020JD033622>, 2021.
- 1400 Song, Z., Qiao, F., Bao, Y., Shu, Q., Song, Y., and Yang, X.: FIO-QLNM FIO-ESM2.0 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.9199>, 2019.
- Stephote, H., Wilcox, L. J., and Highwood, E. J.: Is there a robust effect of anthropogenic aerosols on the Southern Annular Mode?, *Journal of Geophysical Research: Atmospheres*, 121, 10,029–10,042, <https://doi.org/10.1002/2015JD024218>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/2015JD024218>, 2016.
- 1405 Stevens, B., Giorgetta, M., Esch, M., Mauritsen, T., Crueger, T., Rast, S., Salzmann, M., Schmidt, H., Bader, J., Block, K., Brokopf, R., Fast, I., Kinne, S., Kornbluh, L., Lohmann, U., Pincus, R., Reichler, T., and Roeckner, E.: Atmospheric component of the MPI-M Earth System Model: ECHAM6, *Journal of Advances in Modeling Earth Systems*, 5, 146–172, <https://doi.org/https://doi.org/10.1002/jame.20015>, 2013.
- Stevenson, D. S., Young, P. J., Naik, V., Lamarque, J.-F., Shindell, D. T., Voulgarakis, A., Skeie, R. B., Dalsoren, S. B., Myhre, G., Bernsten, T. K., Folberth, G. A., Rumbold, S. T., Collins, W. J., MacKenzie, I. A., Doherty, R. M., Zeng, G., van Noije, T. P. C., Strunk, A., Bergmann, D., Cameron-Smith, P., Plummer, D. A., Strode, S. A., Horowitz, L., Lee, Y. H., Szopa, S., Sudo, K., Nagashima, T., Josse, B., Cionni, I., Righi, M., Eyring, V., Conley, A., Bowman, K. W., Wild, O., and Archibald, A.: Tropospheric ozone changes, radiative forcing and attribution to emissions in the Atmospheric Chemistry and Climate Model Intercomparison Project (ACCMIP), *Atmospheric Chemistry and Physics*, 13, 3063–3085, <https://doi.org/10.5194/acp-13-3063-2013>, publisher: Copernicus GmbH, 2013.
- 1410 Stevenson, D. S., Zhao, A., Naik, V., O'Connor, F. M., Tilmes, S., Zeng, G., Murray, L. T., Collins, W. J., Griffiths, P. T., Shim, S., Horowitz, L. W., Sentman, L. T., and Emmons, L.: Trends in global tropospheric hydroxyl radical and methane lifetime since 1850 from AerChem-MIP, *Atmospheric Chemistry and Physics*, 20, 12 905–12 920, <https://doi.org/10.5194/acp-20-12905-2020>, publisher: Copernicus GmbH, 2020.
- 1415 Stohl, A., Aamaas, B., Amann, M., Baker, L. H., Bellouin, N., Bernsten, T. K., Boucher, O., Cherian, R., Collins, W., Daskalakis, N., Dusinga, M., Eckhardt, S., Fuglestad, J. S., Harju, M., Heyes, C., Hodnebrog, Ø., Hao, J., Im, U., Kanakidou, M., Klimont, Z., Kupiainen, K., Law, K. S., Lund, M. T., Maas, R., MacIntosh, C. R., Myhre, G., Myriokefalitakis, S., Olivié, D., Quaas, J., Quennehen, B., Raut, J.-C., Rumbold, S. T., Samset, B. H., Schulz, M., Seland, Ø., Shine, K. P., Skeie, R. B., Wang, S., Yttri, K. E., and Zhu, T.: Evaluating the climate and air quality impacts of short-lived pollutants, *Atmospheric Chemistry and Physics*, 15, 10 529–10 566, <https://doi.org/10.5194/acp-15-10529-2015>, 2015.
- 1420 Stouffer, R. J., Eyring, V., Meehl, G. A., Bony, S., Senior, C., Stevens, B., and Taylor, K. E.: CMIP5 Scientific Gaps and Recommendations for CMIP6, *Bulletin of the American Meteorological Society*, 98, 95–105, <https://doi.org/10.1175/BAMS-D-15-00013.1>, publisher: American Meteorological Society Section: Bulletin of the American Meteorological Society, 2017.
- 1425

- Swart, N. C., Cole, J. N., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Solheim, L., von Salzen, K., Yang, D., Winter, B., and Sigmond, M.: CCCma CanESM5 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.3610>, 2019a.
- 1430 Swart, N. C., Cole, J. N. S., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Hanna, S., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Sigmond, M., Solheim, L., von Salzen, K., Yang, D., and Winter, B.: The Canadian Earth System Model version 5 (CanESM5.0.3), *Geoscientific Model Development*, 12, 4823–4873, <https://doi.org/10.5194/gmd-12-4823-2019>, 2019b.
- 1435 Szopa, S., Naik, V., Adhikary, B., Artaxo, P., Berntsen, T., Collins, W., Fuzzi, S., Gallardo, L., Kiendler-Scharr, A., Klimont, Z., Liao, H., Unger, N., and Zanis, P.: Short-lived climate forcers, in: *Climate change 2021: The physical science basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, UK and New York, NY, USA, <https://doi.org/10.1017/9781009157896.008>, section: 6 Type: Book section, 2021a.
- 1440 Szopa, S., Naik, V., Adhikary, B., Artaxo, P., Berntsen, T., Collins, W., Fuzzi, S., Gallardo, L., Kiendler-Scharr, A., Klimont, Z., Liao, H., Unger, N., and Zanis, P.: Short-lived climate forcers supplementary material, in: *Climate change 2021: The physical science basis. Contribution of working group I to the sixth assessment report of the intergovernmental panel on climate change*, edited by Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press, Cambridge, UK and New York, NY, USA, https://www.ipcc.ch/report/ar6/wg1/downloads/report/IPCC_AR6_WGI_Chapter06_SM.pdf, section: 6 Type: Book section, 2021b.
- 1445 Tang, Y., Rumbold, S., Ellis, R., Kelley, D., Mulcahy, J., Sellar, A., Walton, J., and Jones, C.: MOHC UKESM1.0-LL model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.6113>, 2019.
- Tatebe, H. and Watanabe, M.: MIROC MIROC6 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.5603>, 2018.
- 1450 Tatebe, H., Ogura, T., Nitta, T., Komuro, Y., Ogochi, K., Takemura, T., Sudo, K., Sekiguchi, M., Abe, M., Saito, F., Chikira, M., Watanabe, S., Mori, M., Hirota, N., Kawatani, Y., Mochizuki, T., Yoshimura, K., Takata, K., O'ishi, R., Yamazaki, D., Suzuki, T., Kurogi, M., Kataoka, T., Watanabe, M., and Kimoto, M.: Description and basic evaluation of simulated mean state, internal variability, and climate sensitivity in MIROC6, *Geoscientific Model Development*, 12, 2727–2765, <https://doi.org/10.5194/gmd-12-2727-2019>, 2019.
- 1455 Thornhill, G., Collins, W., Olivié, D., Skeie, R. B., Archibald, A., Bauer, S., Checa-Garcia, R., Fiedler, S., Folberth, G., Gjermundsen, A., Horowitz, L., Lamarque, J.-F., Michou, M., Mulcahy, J., Nabat, P., Naik, V., O'Connor, F. M., Paulot, F., Schulz, M., Scott, C. E., Séférian, R., Smith, C., Takemura, T., Tilmes, S., Tsigaridis, K., and Weber, J.: Climate-driven chemistry and aerosol feedbacks in CMIP6 Earth system models, *Atmospheric Chemistry and Physics*, 21, 1105–1126, <https://doi.org/10.5194/acp-21-1105-2021>, publisher: Copernicus GmbH, 2021a.
- 1460 Thornhill, G. D., Collins, W. J., Kramer, R. J., Olivié, D., Skeie, R. B., O'Connor, F. M., Abraham, N. L., Checa-Garcia, R., Bauer, S. E., Deushi, M., Emmons, L. K., Forster, P. M., Horowitz, L. W., Johnson, B., Keeble, J., Lamarque, J.-F., Michou, M., Mills, M. J., Mulcahy, J. P., Myhre, G., Nabat, P., Naik, V., Oshima, N., Schulz, M., Smith, C. J., Takemura, T., Tilmes, S., Wu, T., Zeng, G., and Zhang, J.: Effective radiative forcing from emissions of reactive gases and aerosols – a multi-model comparison, *Atmospheric Chemistry and Physics*, 21, 853–874, <https://doi.org/10.5194/acp-21-853-2021>, publisher: Copernicus GmbH, 2021b.

- 1465 Turnock, S. T., Allen, R. J., Andrews, M., Bauer, S. E., Deushi, M., Emmons, L., Good, P., Horowitz, L., John, J. G., Michou, M., Nabat, P., Naik, V., Neubauer, D., O'Connor, F. M., Olivié, D., Oshima, N., Schulz, M., Sellar, A., Shim, S., Takemura, T., Tilmes, S., Tsigaridis, K., Wu, T., and Zhang, J.: Historical and future changes in air pollutants from CMIP6 models, *Atmospheric Chemistry and Physics*, 20, 14 547–14 579, <https://doi.org/10.5194/acp-20-14547-2020>, 2020.
- Turnock, S. T., Allen, R., Archibald, A. T., Dalvi, M., Folberth, G., Griffiths, P. T., Keeble, J., Robertson, E., and
1470 O'Connor, F. M.: The Future Climate and Air Quality Response From Different Near-Term Climate Forcer, *Climate, and Land-Use Scenarios Using UKESM1*, *Earth's Future*, 10, e2022EF002 687, <https://doi.org/10.1029/2022EF002687>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2022EF002687>, 2022.
- Turnock, S. T., Reddington, C. L., West, J. J., and O'Connor, F. M.: The Air Pollution Human Health Burden in Different Future Scenarios That Involve the Mitigation of Near-Term Climate Forcers, *Climate and Land-Use*, 7, e2023GH000 812,
1475 <https://doi.org/10.1029/2023GH000812>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2023GH000812>, 2023.
- van Noije, T., Bergman, T., Le Sager, P., O'Donnell, D., Makkonen, R., Gonçalves-Ageitos, M., Döscher, R., Fladrich, U., von Hardenberg, J., Keskinen, J.-P., Korhonen, H., Laakso, A., Myriokefalitakis, S., Ollinaho, P., Pérez García-Pando, C., Reerink, T., Schrödner, R., Wyser, K., and Yang, S.: EC-Earth3-AerChem: a global climate model with interactive aerosols and atmospheric chemistry participating in CMIP6, *Geoscientific Model Development*, 14, 5637–5668, <https://doi.org/10.5194/gmd-14-5637-2021>, 2021.
- 1480 Voldoire, A.: CMIP6 simulations of the CNRM-CERFACS based on CNRM-CM6-1 model for CMIP experiment historical, <https://doi.org/10.22033/ESGF/CMIP6.4066>, 2018.
- Voldoire, A., Saint-Martin, D., Sénési, S., Decharme, B., Alias, A., Chevallier, M., Colin, J., Guérémy, J.-F., Michou, M., Moine, M.-P., Nabat, P., Roehrig, R., Salas y Mélia, D., Séférian, R., Valcke, S., Beau, I., Belamari, S., Berthet, S., Cassou, C., Cattiaux, J., Deshayes, J., Douville, H., Ethé, C., Franchistéguy, L., Geoffroy, O., Lévy, C., Madec, G., Meurdesoif, Y., Msadek, R., Ribes, A., Sanchez-Gomez, E., Terray, L., and Waldman, R.: Evaluation of CMIP6 DECK Experiments With CNRM-CM6-1, *Journal of Advances in Modeling Earth Systems*, 11, 2177–2213, <https://doi.org/https://doi.org/10.1029/2019MS001683>, 2019.
- 1485 Volodin, E., Mortikov, E., Gritsun, A., Lykossov, V., Galin, V., Diansky, N., Gusev, A., Kostrykin, S., Iakovlev, N., Shestakova, A., and Emelina, S.: INM INM-CM4-8 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.5069>, 2019a.
- Volodin, E., Mortikov, E., Gritsun, A., Lykossov, V., Galin, V., Diansky, N., Gusev, A., Kostrykin, S., Iakovlev, N., Shestakova, A., and
1490 Emelina, S.: INM INM-CM5-0 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.5070>, 2019b.
- Volodin, E. M. and Kostrykin, S. V.: The aerosol module in the INM RAS climate model, *Russian Meteorology and Hydrology*, 41, 519–528, <https://doi.org/10.3103/S106837391608001X>, 2016.
- Volodin, E. M., Dianskii, N. A., and Gusev, A. V.: Simulating present-day climate with the INMCM4.0 coupled model of the atmospheric and oceanic general circulations, *Izvestiya, Atmospheric and Oceanic Physics*, 46, 414–431, <https://doi.org/10.1134/S000143381004002X>,
1495 2010.
- Volodin, E. M., Mortikov, E. V., Kostrykin, S. V., Galin, V. Y., Lykossov, V. N., Gritsun, A. S., Diansky, N. A., Gusev, A. V., and Iakovlev, N. G.: Simulation of the present-day climate with the climate model INMCM5, *Climate Dynamics*, 49, 3715–3734, <https://doi.org/10.1007/s00382-017-3539-7>, 2017.
- Watson-Parris, D., Williams, A., Deaconu, L., and Stier, P.: Model calibration using ESEm v1.1.0 – an open, scalable Earth system emulator, *Geoscientific Model Development*, 14, 7659–7672, <https://doi.org/10.5194/gmd-14-7659-2021>, 2021.
- 1500 Watson-Parris, D., Rao, Y., Olivié, D., Seland, Ø., Nowack, P., Camps-Valls, G., Stier, P., Bouabid, S., Dewey, M., Fons, E., Gonzalez, J., Harder, P., Jeggle, K., Lenhardt, J., Manshausen, P., Novitasari, M., Ricard, L., and Roesch, C.: ClimateBench

- v1.0: A Benchmark for Data-Driven Climate Projections, *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002954, <https://doi.org/10.1029/2021MS002954>, eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021MS002954>, 2022.
- 1505 Wieners, K.-H., Giorgetta, M., Jungclaus, J., Reick, C., Esch, M., Bittner, M., Legutke, S., Schupfner, M., Wachsmann, F., Gayler, V., Haak, H., de Vrese, P., Raddatz, T., Mauritsen, T., von Storch, J.-S., Behrens, J., Brovkin, V., Claussen, M., Crueger, T., Fast, I., Fiedler, S., Hagemann, S., Hohenegger, C., Jahns, T., Kloster, S., Kinne, S., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Müller, W., Nabel, J., Notz, D., Peters-von Gehlen, K., Pincus, R., Pohlmann, H., Pongratz, J., Rast, S., Schmidt, H., Schnur, R., Schulzweida, U., Six, K., Stevens, B., Voigt, A., and Roeckner, E.: MPI-M MPI-ESM1.2-LR model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.6595>, 2019.
- 1510 Wilcox, L.: A summary of summary of aerosol and chemistry representations in CMIP6, <https://doi.org/10.17605/OSF.IO/8FWJ3>.
- Wilcox, L. J., Highwood, E. J., and Dunstone, N. J.: The influence of anthropogenic aerosol on multi-decadal variations of historical global climate, *Environmental Research Letters*, 8, 024033, <https://doi.org/10.1088/1748-9326/8/2/024033>, publisher: IOP Publishing, 2013.
- Wilcox, L. J., Liu, Z., Samset, B. H., Hawkins, E., Lund, M. T., Nordling, K., Undorf, S., Bollasina, M., Ekman, A. M. L., Krishnan, S., Merikanto, J., and Turner, A. G.: Accelerated increases in global and Asian summer monsoon precipitation from future aerosol reductions, *Atmospheric Chemistry and Physics*, 20, 11955–11977, <https://doi.org/10.5194/acp-20-11955-2020>, 2020.
- 1515 Wilcox, L. J., Allen, R. J., Samset, B. H., Bollasina, M. A., Griffiths, P. T., Keeble, J., Lund, M. T., Makkonen, R., Merikanto, J., O'Donnell, D., Paynter, D. J., Persad, G. G., Rumbold, S. T., Takemura, T., Tsigaridis, K., Undorf, S., and Westervelt, D. M.: The Regional Aerosol Model Intercomparison Project (RAMIP), *Geoscientific Model Development*, 16, 4451–4479, <https://doi.org/10.5194/gmd-16-4451-2023>, 2023.
- 1520 Wu, T., Chu, M., Dong, M., Fang, Y., Jie, W., Li, J., Li, W., Liu, Q., Shi, X., Xin, X., Yan, J., Zhang, F., Zhang, J., Zhang, L., and Zhang, Y.: BCC BCC-CSM2MR model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.2948>, 2018.
- Wu, T., Lu, Y., Fang, Y., Xin, X., Li, L., Li, W., Jie, W., Zhang, J., Liu, Y., Zhang, L., Zhang, F., Zhang, Y., Wu, F., Li, J., Chu, M., Wang, Z., Shi, X., Liu, X., Wei, M., Huang, A., Zhang, Y., and Liu, X.: The Beijing Climate Center Climate System Model (BCC-CSM): the main progress from CMIP5 to CMIP6, *Geoscientific Model Development*, 12, 1573–1600, <https://doi.org/10.5194/gmd-12-1573-2019>, 2019.
- 1525 Wu, T., Zhang, F., Zhang, J., Jie, W., Zhang, Y., Wu, F., Li, L., Yan, J., Liu, X., Lu, X., Tan, H., Zhang, L., Wang, J., and Hu, A.: Beijing Climate Center Earth System Model version 1 (BCC-ESM1): model description and evaluation of aerosol simulations, *Geoscientific Model Development*, 13, 977–1005, <https://doi.org/10.5194/gmd-13-977-2020>, 2020.
- YU, Y.: CAS FGOALS-f3-L model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.3355>, 2019.
- 1530 YUKIMOTO, S., KAWAI, H., KOSHIRO, T., OSHIMA, N., YOSHIDA, K., URAKAWA, S., TSUJINO, H., DEUSHI, M., TANAKA, T., HOSAKA, M., YABU, S., YOSHIMURA, H., SHINDO, E., MIZUTA, R., OBATA, A., ADACHI, Y., and ISHII, M.: The Meteorological Research Institute Earth System Model Version 2.0, MRI-ESM2.0: Description and Basic Evaluation of the Physical Component, *Journal of the Meteorological Society of Japan. Ser. II*, 2019.
- Yukimoto, S., Koshiro, T., Kawai, H., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T., Hosaka, M., Yoshimura, H., Shindo, E., Mizuta, R., Ishii, M., Obata, A., and Adachi, Y.: MRI MRI-ESM2.0 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.6842>, 2019.
- 1535 Zanis, P., Akritidis, D., Georgoulas, A. K., Allen, R. J., Bauer, S. E., Boucher, O., Cole, J., Johnson, B., Deushi, M., Michou, M., Mulcahy, J., Nabat, P., Olivieri, D., Oshima, N., Sima, A., Schulz, M., Takemura, T., and Tsigaridis, K.: Fast responses on pre-industrial climate from present-day aerosols in a CMIP6 multi-model study, *Atmospheric Chemistry and Physics*, 20, 8381–8404, <https://doi.org/10.5194/acp-20-8381-2020>, 2020.
- 1540

- Zanis, P., Akritidis, D., Turnock, S., Naik, V., Szopa, S., Georgoulas, A. K., Bauer, S. E., Deushi, M., Horowitz, L. W., Keeble, J., Sager, P. L., O'Connor, F. M., Oshima, N., Tsigaridis, K., and Noije, T. v.: Climate change penalty and benefit on surface ozone: a global perspective based on CMIP6 earth system models, *Environmental Research Letters*, 17, 024 014, <https://doi.org/10.1088/1748-9326/ac4a34>, publisher: IOP Publishing, 2022.
- 1545 Zeng, G., Morgenstern, O., Williams, J. H. T., O'Connor, F. M., Griffiths, P. T., Keeble, J., Deushi, M., Horowitz, L. W., Naik, V., Emmons, L. K., Abraham, N. L., Archibald, A. T., Bauer, S. E., Hassler, B., Michou, M., Mills, M. J., Murray, L. T., Oshima, N., Sentman, L. T., Tilmes, S., Tsigaridis, K., and Young, P. J.: Attribution of Stratospheric and Tropospheric Ozone Changes Between 1850 and 2014 in CMIP6 Models, *Journal of Geophysical Research: Atmospheres*, 127, e2022JD036 452, <https://doi.org/10.1029/2022JD036452>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2022JD036452>, 2022.
- 1550 Zhang, H., Zhang, M., Jin, J., Fei, K., Ji, D., Wu, C., Zhu, J., He, J., Chai, Z., Xie, J., Dong, X., Zhang, D., Bi, X., Cao, H., Chen, H., Chen, K., Chen, X., Gao, X., Hao, H., Jiang, J., Kong, X., Li, S., Li, Y., Lin, P., Lin, Z., Liu, H., Liu, X., Shi, Y., Song, M., Wang, H., Wang, T., Wang, X., Wang, Z., Wei, Y., Wu, B., Xie, Z., Xu, Y., Yu, Y., Yuan, L., Zeng, Q., Zeng, X., Zhao, S., Zhou, G., and Zhu, J.: Description and Climate Simulation Performance of CAS-ESM Version 2, *Journal of Advances in Modeling Earth Systems*, 12, e2020MS002 210, <https://doi.org/https://doi.org/10.1029/2020MS002210>, e2020MS002210 2020MS002210, 2020.
- 1555 Zhang, J., Wu, T., Shi, X., Zhang, F., Li, J., Chu, M., Liu, Q., Yan, J., Ma, Q., and Wei, M.: BCC BCC-ESM1 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.2949>, 2018.
- Zhang, J., Furtado, K., Turnock, S. T., Mulcahy, J. P., Wilcox, L. J., Booth, B. B., Sexton, D., Wu, T., Zhang, F., and Liu, Q.: The role of anthropogenic aerosols in the anomalous cooling from 1960 to 1990 in the CMIP6 Earth system models, *Atmospheric Chemistry and Physics*, 21, 18 609–18 627, <https://doi.org/10.5194/acp-21-18609-2021>, publisher: Copernicus GmbH, 2021a.
- 1560 Zhang, Y., Ciais, P., Boucher, O., Maignan, F., Bastos, A., Goll, D., Lurton, T., Viovy, N., Bellouin, N., and Li, L.: Disentangling the Impacts of Anthropogenic Aerosols on Terrestrial Carbon Cycle During 1850–2014, *Earth's Future*, 9, e2021EF002 035, <https://doi.org/10.1029/2021EF002035>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021EF002035>, 2021b.
- Zhao, A., Ryder, C. L., and Wilcox, L. J.: How well do the CMIP6 models simulate dust aerosols?, *Atmospheric Chemistry and Physics*, 22, 2095–2119, <https://doi.org/10.5194/acp-22-2095-2022>, 2022.
- 1565 Ziehn, T., Chamberlain, M., Lenton, A., Law, R., Bodman, R., Dix, M., Wang, Y., Dobrohotoff, P., Srbinovsky, J., Stevens, L., Vohralik, P., Mackallah, C., Sullivan, A., O'Farrell, S., and Druken, K.: CSIRO ACCESS-ESM1.5 model output prepared for CMIP6 CMIP historical, <https://doi.org/10.22033/ESGF/CMIP6.4272>, 2019.
- Ziehn, T., Chamberlain, M. A., Law, R. M., Lenton, A., Bodman, R. W., Dix, M., Stevens, L., Wang, Y.-P., and Srbinovsky, J.: The Australian Earth System Model: ACCESS-ESM1.5, *Journal of Southern Hemisphere Earth Systems Science*, 70, 193–214, <https://doi.org/10.1071/ES19035>, 2020.
- 1570