

# Riverine nutrient impact on global ocean nitrogen cycle feedbacks and marine primary production in an Earth System Model

Miriam Tivig<sup>1,2</sup>, David P. Keller<sup>1,3</sup>, and Andreas Oschlies<sup>1</sup>

<sup>1</sup>GEOMAR Helmholtz-Zentrum für Ozeanforschung Kiel, Wischhofstr. 1-3, D-24148 Kiel, Germany

<sup>2</sup>Deutscher Wetterdienst, Abteilung Klima und Umwelt, Michendorfer Chaussee 23, D-14473 Potsdam, Germany.

<sup>3</sup>Carbon to Sea Initiative

**Correspondence:** Miriam Tivig (mtivig@geomar.de)

## Abstract.

Riverine nutrient export is an important process in marine coastal biogeochemistry and also impacts global marine biology. The nitrogen cycle is a key player here. Internal feedbacks regulate not only nitrogen distribution, but also primary production and thereby oxygen concentrations. Phosphorus is another essential nutrient and interacts with the nitrogen cycle via different feedback mechanisms. After a previous study of the N-marine nitrogen cycle response to riverine nitrogen supply, we here ~~additionally~~ include phosphorus from river export with different phosphorus burial scenarios and study the impact of phosphorus alone and in combination with nitrogen in a global 3-D ocean biogeochemistry model. Again, we analyse the effects on near coastal and open ocean biogeochemistry. We find that the addition of bio-available riverine phosphorus alone or together with nitrogen affects marine biology on millennial timescales more than riverine nitrogen alone. Biogeochemical feedbacks in the marine nitrogen cycle are strongly influenced by ~~the~~ additional phosphorus. Where bio-available phosphorus is increased by river input, nitrogen ~~concentrations increase~~ concentration increases as well, except for regions with high denitrification rates. High phosphorus burial rates ~~on the other hand~~ decrease biological production significantly. Globally, riverine phosphorus leads to elevated primary production rates in the coastal and open oceans.

## Plain Language Summary

Coastal oceans are the most productive parts of the global ocean and, arguably, most sensitive to environmental change, but poorly resolved in most global models. In reality, rivers influence the coastal oceans, because they transport nutrients from land to the sea. This nutrient supply is often not included in global models, even though it may impact not only the coastal biology, but also the marine biology in regions farther away. We here include dissolved inorganic nitrogen and phosphorus from river export in a global ocean model and study the effects on the near coastal and the open ocean. We find that the addition of riverine phosphorus affects marine biology on millennial timescales more than riverine nitrogen alone. Where phosphorus is increased by river input, nitrogen concentrations increase as well, except for regions with high rates of bacterial consumption of nitrate (denitrification). High phosphorus burial rates on the other hand, decrease biological production significantly. ~~In ocean regions with low oxygen concentration, additional nitrogen supply can paradoxically lead to a loss in nitrogen and a dampening of~~

~~primary production. This vicious cycle is locally enhanced by increased phosphorus supply. Nevertheless, riverine phosphorus~~  
25 ~~globally~~ Globally, riverine phosphorus leads to higher primary production rates in the coastal and open oceans.

*Copyright statement.* TEXT

## 1 Introduction

Nitrogen and phosphorus are both considered as limiting nutrients in the global ocean, ~~nitrogen as~~. 'Limiting' refers here to the  
concept of Liebig's law of growth rate determined by the availability of the most limiting substrate (Allaby, 2010). It states that  
30 growth is dictated not by total resources available, but by the scarcest resource (limiting factor). Nitrogen can be considered  
as the 'proximate limiting' and phosphorus as the 'ultimate limiting' nutrient, according to the definition by Tyrrell (1999).  
Changes in the availability of oceanic fixed-nitrogen (N) are known to have driven marine productivity changes, thereby reg-  
ulating the strength of the biological carbon pump ~~and thereby the partitioning of carbon dioxide (CO<sub>2</sub>) between~~. By this, the  
availability of N also influences the carbon cycle in atmosphere and ocean (Falkowski, 1997). Phosphorus (P) is limiting marine  
35 productivity on geological and global scales and plays an important role in regulating oceanic oxygen inventories (Monteiro  
et al., 2012; Palastanga et al., 2011).

Oceanic fixed nitrogen concentrations are mainly controlled by the balance between nitrogen fixation and denitrification, but  
atmospheric deposition and riverine input also contribute to the global N budget (Somes et al., 2013; Deutsch et al., 2007;  
Gruber, 2004; Ruttenger, 2003). Although several studies question the stability of the marine N inventory (Zehr and Capone,  
40 2020; Codispoti et al., 2001; Gruber and Sarmiento, 1997; Codispoti, 1995), the pre-industrial global nitrogen cycle is often  
assumed to reflect a steady state (Deutsch et al., 2007; Altabet, 2006; Gruber, 2004; Tyrrell, 1999; Redfield et al., 1963).

This balance is assured by negative N-cycle feedbacks that stabilize marine N concentration: Where fixed N is sparse, dia-  
zotrophs can fix atmospheric N<sub>2</sub> instead of depending on dissolved inorganic nitrogen (DIN), such as ammonia or nitrate. But  
as the process of atmospheric N fixation is slower and requires more energy, mostly to keep oxygen away from the oxygen-  
45 sensitive enzyme nitrogenase, diazotrophs are rapidly outcompeted by other phytoplankton who do not have this energetic cost,  
if sufficient DIN is available. Growth of diazotrophs is also limited by the availability of phosphate, light and iron.

On the other hand, the global N budget is regulated by loss of fixed N, predominantly via denitrification. This process describes  
anaerobic respiration of organic matter via bacterial reduction of nitrate to N<sub>2</sub>, and occurs in the water column as well as in  
sediments if the oxygen concentration is low (Gruber, 2004; Deutsch et al., 2001). Denitrification limits itself, as the consump-  
50 tion of nitrate leads to a reduction in the production of organic matter and hence to less oxygen consumption.

Generally the processes of N<sub>2</sub>-fixation and denitrification take place in different regions of the world ocean. Still, feedbacks  
link these processes globally and are generally assumed to restore the balance in the global marine N budget. In some regions,  
however, N<sub>2</sub>-fixation and denitrification occur in geographical proximity and may produce a "vicious cycle" with a runaway  
local loss of fixed N (Landolfi et al., 2013). Estimates of the mean residence time of fixed nitrogen in the global ocean amount

55 to a few thousand years (Gruber, 2004). Due to the complexity of internal feedbacks and the dynamic role of nitrogen in the marine biological production it is difficult to assess, how sensitive the global N concentration is to perturbations of marine biogeochemistry.

~~Of particular interest is the interaction between fixed N and phosphorus (P), as P has an impact on the contributors of~~ In a previous study (Tivig et al., 2021), we used the Earth system climate model of intermediate complexity of the University of Victoria (UVic) version 2.9 (Eby et al., 2009; Weaver et al., 2001) to study N cycle feedbacks ~~-P has been termed the 'ultimate' limiting nutrient (compared to the 'proximate' limiting N, citepTyrrell99), and the~~ in the modelled ocean in response to the addition of riverine dissolved inorganic nitrogen (Tivig et al., 2021). We found that, although in hot-spots near the river mouths marine primary production increased due to the additional N, globally biogeochemical feedbacks stabilized DIN concentrations and primary production, and could even lead to a local decline in DIN and productivity in proximity to low oxygen regions. In those idealized simulations, N was the only nutrient supplied via rivers.

With this, we confirmed an early study by Tyrrell (1999), who used a box model and found, that "an increase in the river delivery of nitrate has no long-term effect on productivity". But Tyrrell (1999) also concluded, that "an increase in the river delivery of phosphate, on the other hand, causes a sustained and proportionate increase in productivity". The interaction between fixed N and phosphorus (P) is of particular interest for marine primary production. The availability of P is one of the limiting factors for N<sub>2</sub> fixation (Wang et al., 2019; Landolfi et al., 2015).

In Tivig et al. (2021) we also found, that locally, marine productivity has been increased by riverine N supply. Riverine supply of P, is predicted to regulate total ocean productivity globally (Tyrrell, 1999). Nevertheless, locally, in coastal oceans and in regions with low oxygen concentrations, marine biogeochemistry can differ from global averages.

The total P inventory in the global ocean is mainly controlled by riverine input and burial at the seafloor (Wallmann, 2010; 75 Ruttenberg, 2003; Baturin, 2003; Delaney, 1998). The residence time of P in the global ocean is approximately one order of magnitude longer than that of fixed N (Delaney, 1998). (Bio-)chemical weathering on continents represents the main source of riverine bio-available P (Filipelli, 2008; Föllmi, 1995). Changes in marine P are mostly a consequence of changes in terrestrial weathering and therefore have generally occurred on geological timescales.

Nevertheless, climate change and direct anthropogenic interventions in the P cycle have started to alter P fluxes during the last 80 century. Due to the wide-spread use of fertilizers, to deforestation and sewage sources, the riverine load of phosphorus has increased globally (Seitzinger et al., 2010; Filipelli, 2008). In addition, river flows have also been dramatically altered by land use change and the damming of rivers (Cappellen and Maavara, 2016). In order to predict future changes in marine biogeochemistry, it is therefore relevant to understand, how P fluxes impact the marine N cycle, the N cycle feedbacks and **primary production** marine productivity globally and regionally.

85 As global observations and measurements of ocean nutrients and fluxes are difficult and observations still relatively sparse, models are often used to investigate large-scale marine biogeochemistry over long time scales. However, only recently have there been more global modelling studies with riverine nutrient input. In one of the first studies, Giraud et al. (2008) analysed coastal fluxes of P, silicate, and dissolved iron in a global ocean model and found that including nutrients in the coastal ocean impacts biological activity locally but also in the open ocean. They also found, that excess nutrients in the coastal ocean can

90 impact the open ocean biogeochemistry depending on which nutrient is advected from the coastal region. Nutrient availability and its consumption in the coastal domain control this advection. Additional P does affect coastal oceans especially if they are P-limited and if they are not limited by other nutrients like iron. In this case, Giraud et al. (2008) found that increased primary production in the coastal oceans can lead to a depletion of nutrients in the open ocean, reducing biological activity there. This "seesaw effect" was detected by Giraud et al. (2008) on local and global scale. If P is not consumed in the coastal oceans, 95 it may be advected offshore, eventually increasing primary production there. N was not simulated explicitly in that study but coupled to P via Redfield ~~ration. In addition, the ratio. The~~ simulation performed by (Giraud et al., 2008) was only run for 10 years, hence not long enough for N-cycle feedbacks to materialise.

In a more recent study (Lacroix et al., 2020) implemented estimated riverine nutrient loads of P, N, iron, carbon (C) and silica in a global ocean model and compared the results with those by a reference simulation, where the same nutrients were added 100 directly and homogeneously to the open ocean surface. They found that even if the ocean circulation remains the main driver for biogeochemical distributions in the open ocean, it appeared necessary to include riverine inputs for the representation of heterogeneous features in the coastal ocean. They identified the catchments of the tropical Atlantic, the Arctic Ocean, Southeast Asia and Indo-Pacific islands as regions of dominant contributions of riverine supplies to the ocean, leading to a strong primary production increase in the tropical west Atlantic, Bay of Bengal and the East China Sea. Nevertheless the focus here 105 was mainly on C export, and N feedbacks have not been considered by Lacroix et al. (2020).

~~In a previous study, we used the Earth system climate model of intermediate complexity of the University of Victoria (UVic) version 2.9 (Eby et al., 2009; Weaver et al., 2001) to study N cycle feedbacks in the modelled ocean in response to the addition of riverine dissolved inorganic nitrogen (Tivig et al., 2021). We found that although marine primary production increased due to the additional N in hot-spots near the river mouths, globally biogeochemical feedbacks stabilized DIN concentrations and primary production, but could even lead to a local decline in DIN and productivity in proximity to low oxygen regions. In those idealized simulations, N was the only nutrient supplied via rivers.~~ As P concentrations have been increasing in many water bodies over the world, rivers transport more P to the coastal oceans. Beusen and Bouwman (2022) showed that human-dominated river supply of N and P has not only increased in the past, but will do so in the future, due to legacies of past nutrient management, even if efforts are made to reduce these nutrient loads. Regarding P, not only is the addition of nutrient *per se* 115 relevant for ocean marine biogeochemistry, but the stoichiometric ratio of N and P is also essential ~~for marine biogeochemistry~~ (Garnier et al., 2010; Redfield et al., 1963; Beusen and Bouwman, 2022). In the present study we have extended our experiment from Tivig et al. (2021) and include riverine supply of ~~both N and P in order to~~ P in addition to N. We again aim to study the feedbacks ~~of the N cycle in the N-cycle but this time face~~ to the combined input of both limiting nutrients.

Specifically, we address the following questions:

- 120
- How does riverine N and P input together, rather than riverine N input alone, affect the representation of ocean biogeochemistry including marine primary production in our model?
  - How does the addition of riverine P input affect specifically the N cycle and N-cycle feedbacks?
  - What effect has the inclusion of riverine P fluxes specifically on marine oxygen concentrations?




To address these questions, an Earth system model of intermediate complexity (EMIC, Claussen et al. (2002)) that resolves the relevant biogeochemical feedbacks is employed. It allows the integration of a large number of processes at reduced computational costs due to a coarser resolution and simplified assumptions, in case of UVic with respect to atmospheric dynamics. This form of model makes it possible to run millennial-scale simulations with different assumptions, allowing the analysis of processes and feedbacks operating in the climate system on such timescales (Weaver et al., 2001). Compared to the study of Tyrrell (1999), where a simple box model has been used to study the relative influences of nitrogen and phosphorus on oceanic primary production, the current study is based on a global 3D Earth system model, where the global and regional distribution of N and P can be analysed in more detail. This is the prerequisite for the inclusion of river exports to the ocean, with different amounts of nutrients depending on the individual river. Hence, regional feedbacks in the N cycle and their effects on marine primary production can be assessed as well as localised biogeochemical reactions for example near oxygen minimum zones.

## 2 Methods

### 2.1 Earth system model UVic

The University of Victoria Earth System Climate model (UVic ESCM) version 2.9 (Eby et al., 2009; Weaver et al., 2001) consists of a three-dimensional ( $1.8^\circ \times 3.6^\circ$ , 19 levels) general circulation model of the ocean, a two-dimensional, single-layer energy-moisture balance atmospheric model, a dynamic-thermodynamic sea ice model, and a terrestrial vegetation model. The atmospheric energy-moisture balance model (Fanning and Weaver, 1996) dynamically calculates heat and water fluxes between the atmosphere and the ocean, land and sea ice, and is forced by monthly climatological winds prescribed from NCEP/NCAR. The 19 vertical levels of the oceanic component, Modular Ocean Model 2 (MOM2), are 50 m thick near the surface and up to 500 m in the deep ocean. The oceanic physical settings are the same as in Keller et al. (2012). The ocean model includes a marine ecosystem module based on Keller et al. (2012) with updates as noted in Partanen et al. (2016). The ocean ecosystem and biogeochemical model is an improved NPZD (nutrient, phytoplankton, zooplankton, detritus) ecosystem model based on ~~(Schmittner et al., 2008)~~ Schmittner et al. (2008) and includes seven prognostic variables: two phytoplankton classes (nitrogen fixing diazotrophs  $P_D$  and other phytoplankton  $P_O$ ), zooplankton (Z), sinking particulate detritus (D), nitrate ( $\text{NO}_3$ ), phosphate ( $\text{PO}_4$ ) and oxygen ( $\text{O}_2$ ) (Fig. 1).  $\text{NO}_3$  and  $\text{PO}_4$  are linked through exchanges with the biological variables by constant (Redfield) stoichiometry (Schmittner et al., 2008). Since diazotrophs can fix nitrogen gas dissolved in seawater, they are not limited by  $\text{NO}_3$  ~~nor by a maximum  $\text{NO}_3$  concentration~~, while the growth of other phytoplankton is limited by  $\text{NO}_3$  and  $\text{PO}_4$ . All phytoplankton are additionally limited by iron, light and temperature. For the current study of nitrogen cycle feedbacks it is a clear advantage, that UVic explicitly calculates diazotrophs and  $\text{N}_2$ -fixation. Keller et al. (2012) found that patterns and global amounts of modelled  $\text{N}_2$  fixation were mostly consistent with the relatively sparse available observations (Sohm et al., 2011). The main differences are discussed within the framework of our results. See Keller et al. (2012) for a full description and evaluation of simulated marine biogeochemistry.

As in our previous study (Tivig et al., 2021) we use empirical transfer functions derived from benthic flux measurements



**Figure 1.** Ecosystem model schematics for the NPZD model with the prognostic variables (in square boxes) and the fluxes of matter between them, indicated by arrows. See details in the text. Figure updated from Keller et al. (2012) and Tivig et al. (2021).

to calculate benthic denitrification following Bohlen et al. (2012), combined with a subgrid bathymetry scheme for shallow continental shelves and other topographical features that are too fine to be resolved on the coarse UVic grid (see details in Somes et al. (2010b)).

## 160 2.2 Including riverine Nitrogen and Phosphorus

### 2.2.1 Global Nutrient Export from WaterSheds 2 (NEWS2)

The basic UVic model and ecosystem module do not account for riverine nutrient input. The only source of N in the ocean model consists in  $N_2$ -fixation. In Tivig et al. (2021) we included riverine N as calculated by a global, spatially explicit model of nutrient exports by rivers, NEWS2 (Mayorga et al., 2010). This second version of a system of sub-models estimates the present-day annual export yield at the river mouth for each of 6081 river catchment areas and for dissolved and particulate forms of organic and inorganic N and P, as well as dissolved organic and particulate carbon. In our model study, parameterization of riverine N fluxes is identical to Tivig et al. (2021). Since nitrate is the only nitrogen nutrient explicitly resolved in the

UVic version used, all bioavailable N has been included in the nitrate compartment of the model. See Mayorga et al. (2010) for more details on the model configuration and Dumont et al. (2005) for further details on the validation of NEWS dissolved inorganic nitrogen (DIN).  
170


### 2.2.2 Riverine reactive Phosphorus

Earlier applications of the UVic ESCM assumed a fixed marine P inventory (Keller et al., 2012; Oschlies et al., 2019; Schmittner et al., 200). In addition to DIN, we here include also P from river discharge. We ~~here~~ focus on the total amount of reactive P, i.e. that P that exchanges with the dissolved oceanic P-reservoir and thus is available for biological uptake (Filipelli, 2008; Ruttenberg, 2003).  
175 Estimates of globally integrated pre-industrial riverine supply of bioavailable P range from 0.1 Tmol P yr<sup>-1</sup> to 0.3 Tmol P yr<sup>-1</sup> (Kemena et al., 2019; Ruttenberg, 2003; Filipelli, 2008; Compton et al., 2000; Colman and Holland, 2000). Taking into account only dissolved inorganic P from rivers would underestimate the amount of bioavailable P from river discharge, as most studies estimate DIP export from rivers to be significantly lower, between 0.01 and 0.05 Tmol P yr<sup>-1</sup> (Mayorga et al., 2010; Filipelli,  
180 2008; Harrison et al., 2005). Following ~~other studies~~ the results of Colman and Holland (2000); Ruttenberg (2003), we decided to include dissolved organic and inorganic P (DOP, DIP), as well as 45 % of total particulate P (TPP). This represents the upper range of the fraction of riverine TPP flux estimated as reactive-P (Ruttenberg, 2003; Colman and Holland, 2000). The numbers for DIP, DOP and PP export at the river mouths have been taken from the NEWS2 data set. DIP and DOP were taken as such, PP was multiplied by 0.45 to obtain the desired fraction of total particulate P. Consistent with Tivig et al. (2021), ~~for the current~~  
185 ~~study the~~ the nutrients from NEWS2 were interpolated onto the coarser UVic grid. We assumed a periodic seasonal cycle in runoff and that concentrations in the discharged river water are constant throughout the seasonal cycle. The annual P load is thus distributed over the months using the fractions of monthly freshwater discharge as respective weights. The global amount of P we added to the UVic ocean is 0.17 Tmol P yr<sup>-1</sup> (5.4 Tg P yr<sup>-1</sup>), which lays in the range estimated by previous studies (Kemena et al., 2019; Ruttenberg, 2003; Benitez-Nelson, 2000). Since phosphate is the only phosphorus nutrient explicitly  
190 resolved in the UVic version used, we decided like for N to put all bioavailable P into the phosphate compartment of the model.

### 2.2.3 Burial of Nitrogen and Phosphorus ~~burial~~

In the global ocean, N input via N<sub>2</sub> fixation and riverine is balanced by denitrification (here implicitly including anammox). In particular denitrification in the benthic zone is considered as the major sink for fixed N (Voss et al., 2013; Galloway et al., 2004). As in Tivig et al. (2021), we include empirical transfer functions derived from benthic flux measurements (Bohlen et al., 2012)  
195 to simulate benthic denitrification. These functions are based on dynamic vertically integrated sediment models and estimate denitrification from the rain rate of particulate organic carbon to the seafloor, bottom water O<sub>2</sub> and NO<sub>3</sub> concentrations. We use a subgrid bathymetry scheme for shallow continental shelves to better resolve particulate organic matter sinking and remineralization at the seafloor (Somes and Oschlies, 2015; Somes et al., 2013).

UVic does not contain a prognostic and vertically resolved sediment model. Therefore, the input of reactive P to the ocean has  
200 to be counterbalanced by a ~~parameterised~~ parameterized sink. For this purpose ~~we included~~, we include burial functions based



**Figure 2.** [Export yield of total reactive phosphorus for each discharge point in  \$\text{mmol P m}^{-2} \text{ month}^{-1}\$  from NEWS2 data set interpolated on the UVic grid for January \(a\), April \(b\), July \(c\) and October \(d\).](#)

on Flögel et al. (2011) and Wallmann (2010), which have been tested in previous studies with UVic by Kemena et al. (2019) and Niemeyer et al. (2017). With these functions, the burial of P in the sediment ( $BUR_p$ ) is calculated in every ocean grid box column from the difference between the simulated detritus P rain rate to the sediment ( $RR_p$ ) and the benthic release flux of phosphate from the sediment ( $BEN_p$ ):

$$205 \quad BUR_p = RR_p - BEN_p, \quad (1)$$

Following Flögel et al. (2011), the burial of P is evaluated separately for each grid box column on the shelf and in the deep-sea. Deep-sea is defined as ocean grid box columns where the ocean is deeper than 1000 m.

Benthic P release ( $BEN_p$ ) is calculated locally with:

$$BEN_p = \frac{BEN_c}{r_{c/p}}, \quad (2)$$

210  $BEN_c$  represents the benthic fluxes of carbon and was computed from the difference of the carbon rain-rate to the sediment ( $RR_c$ ) and a virtual burial flux of organic carbon ( $BUR_c$ ):

$$BEN_c = RR_c - BUR_c, \quad (3)$$

Depending on the ocean depth of the considered grid box,  $BUR_c$  is computed from the modelled detritus export in terms of carbon ([Kemena et al., 2019](#)), on the shelf and continental margin (Kemena et al., 2019):

215 
$$BUR_c = 0.14RR_c^{1.11}, \tag{4}$$

and in the deep sea:

$$BUR_c = 0.14RR_c^{1.05}, \tag{5}$$

$RR_c$  is calculated by the model in  $\text{mmol C m}^{-2} \text{ yr}^{-1}$ . The  $r_{c/p}$  ratio depends on bottom water oxygen concentration and is calculated following Kemena et al. (2019) and Wallmann (2010):

220 
$$r_{c/p} = Y_F - A * \exp(-O_2/r) \tag{6}$$

with  $O_2$  in  $\text{mmol m}^{-3}$  and the coefficients  $Y_F = 123 \pm 24$ ,  $A = -112 \pm 24$  and  $r = 32 \pm 19 \text{ mmol m}^{-3}$ . Variations in the coefficients determine the strength of the burial. Because there are large uncertainties in these numbers, different experiments have been performed to evaluate the ~~the~~-model response to variations in burial.

### 2.3 Experimental design

225 To analyse the effect of riverine nutrient supply in the UVic model, six experiments were performed (Table 1). In each experiment the model was run for 10,000 years, starting from an already-spun-up steady state with the standard model version without riverine nutrients, with pre-industrial conditions for insolation and a fixed atmospheric  $\text{CO}_2$  concentration of 283 ppm (Keller et al., 2012), i.e. a stable climate. ~~All simulations included~~ Note that the spin-up did not include benthic denitrification and subgrid bathymetry. These two features have been included at the start of each simulation, including the control simulation

230 (CTR). Hence, at the start of each simulations and for approximately 2000 years, there is a drift in N concentration.

Like in our previous study Tivig et al. (2021), a control simulation (CTR) was performed without riverine nutrients. The simulation NEWS-N is identical to the main experiment analysed in Tivig et al. (2021), where only DIN from river discharge was added to the coastal ocean. In a follow-up experiment, rivers exported only P to the modelled ocean (NEWS-P), but without any burial to balance the P budget. NEWS-N+P tested the impacts on marine biology and biochemistry with regard to additional N

235 and P (without P burial). Finally, two P burial variation experiments have been performed, where riverine N and P supply from the NEWS2 model was applied. In a low burial configuration (N+P-BURLOW) we used the coefficients tested by Kemena et al. (2019) in their simulation low-bur. The second burial experiment (N+P-BURHIGH) includes the original burial functions of Flögel et al. (2011) with the coefficients for  $Y_F$ ,  $A$  and  $r$  described there as well as in the reference burial experiment in Kemena et al. (2019) (Table 1).

240 Note that previous studies using these burial functions have balanced the modelled P budget by an equivalent weathering flux providing P to the ocean via river discharge in the same amount as P is buried in the marine sediments ([Kemena et al., 2019](#); [Niemeyer et al., 2010](#)). In the current study there is no direct link between the two fluxes: riverine P is calculated from the NEWS2 model (Mayorga et al., 2010), while P burial is calculated independently, using only detritus export and bottom-water oxygen concentration

**Table 1.** Overview of simulations and riverine nutrient fluxes:


Simulation	N-flux in $\text{Tg-Tmol N yr}^{-1}$	P-flux in $\text{Tg-Tmol P yr}^{-1}$	Description
CTR	0	0	UVic simulation without NEWS (Control)
NEWS-N	<del>22.8</del> -1.6	0	UVic simulation with DIN from NEWS
NEWS-P	0	<del>5.4</del> -0.17	UVic simulation with P from NEWS, no burial
NEWS-N+P	<del>22.8</del> -1.6	<del>5.4</del> -0.17	UVic simulation with N and P from NEWS, no burial
N+P-BURLOW	<del>22.8</del> -1.6	<del>5.4</del> -0.17	UVic simulation with N and P <u>from NEWS</u> , low burial configuration: $Y_F = 100.5$ ; $A = 90$ ; $r = 38$ . (Kemena et al., 2019)
N+P-BURHIGH	<del>22.8</del> -1.6	<del>5.4</del> -0.17	UVic simulation with N and P <u>from NEWS</u> , high burial configuration: $Y_F = 123$ ; $A = 112$ ; $r = 32$ (Flögel et al., 2011)

(see Table 1 for the overview of the fluxes). In all simulations where riverine P is included, the global P budget is therefore not exactly balanced ~~-The~~ (Figure 3). While P concentration only slowly increases in the scenario with a low burial rate, the increase is stronger in the two scenarios without P burial (NEWS-P and NEWS-N+P). In the scenario with high burial rates, P concentrations decrease and the total P flux is negative. In the timeseries (Figure 3) it is also visible, that this flux is slowly increasing over the simulation and hence P inventory is following an asymptotic evolution. These different simulations permit us to ~~analyze-analyse~~ the sensitivity of the N cycle feedbacks under different conditions of P supply and do not pretend to reproduce the exact reality. Furthermore, the phosphorus balance of the (pre-anthropogenic) ocean is still poorly defined and the input and output fluxes are only rudimentarily constrained (Wallmann, 2010; Föllmi, 1995).

### 3 Results

#### 3.1 Phosphorus dynamics

In the four simulations with riverine P input, globally ~~5.4 Tg-0.17 Tmol~~  $\text{P yr}^{-1}$  (~~0.17 Tmol-5.4 Tg~~  $\text{P yr}^{-1}$ ) are added to the coastal oceans as reactive P, ~~which~~ (Figure 2). This is close to the amount estimated by Föllmi (1995), and within the range of other literature values for global fluvial fluxes of bioavailable P (Kemena et al., 2019; Ruttenberg, 2003; Benitez-Nelson, 2000; Compton et al., 2000). In NEWS-P and NEWS-N+P no sink of P was implemented, and these simulations are characterized by continually increasing P inventory (Figure 3). Benthic burial fluxes of P amount to  $0.16 \text{ Tmol yr}^{-1}$  in N+P-BURLOW and  $0.24 \text{ Tmol yr}^{-1}$  in N+P-BURHIGH, leading to an imbalance of  $+0.01 \text{ Tmol yr}^{-1}$  and  $-0.07 \text{ Tmol yr}^{-1}$  respectively. These results range within the estimates for burial rates from observations, which globally vary around  $0.2 \text{ Tmol P yr}^{-1}$ :  $0.11\text{--}0.34 \text{ Tmol P yr}^{-1}$  in Benitez-Nelson (2000),  $0.17\text{--}0.24 \text{ Tmol P yr}^{-1}$  for Ruttenberg (2003) and  $0.21 \text{ Tmol P yr}^{-1}$  in Filipelli (2008).



**Figure 3.** Top: Timeseries of global phosphorus in all simulations over the 10 000 simulation years in Pg P. Bottom: Timeseries of total phosphorus fluxes (riverine input and burial flux) in Tmol P per year for the simulations with low and high burial. Simulation descriptions can be found in Table 1.

The highest simulated P burial fluxes can be found in coastal regions, especially in the western Pacific and Atlantic ocean (Figure A2). Simulated burial hot spots are situated in proximity to the coast of China, East Russia and Alaska, next to the



265 eastern coast of North- and South-America and in the North Sea. There are no global observational ~~data-sets~~ data sets of P burial. Nevertheless, the distribution of the benthic fluxes is similar to other model studies that have used a similar algorithm (e.g. Bohlen et al., 2012).


#### ~~Burial flux of P in the N+P-BURHIGH simulation in g P yr<sup>-1</sup>.~~

The inclusion of riverine P has a significant impact on global P concentrations. In simulation CTR without riverine nutrient  
270 supply, surface phosphate concentrations range from 0 mmol P m<sup>-3</sup> in most of the western tropical and subtropical ocean basins to more than 2 mmol P m<sup>-3</sup> in the Southern Ocean (Figure 4). The addition of riverine N (in NEWS-N) leads only to small changes in the surface P concentrations, in particular P declines reflecting enhanced biological processes (P uptake) in the coastal oceans (~~Figure 4b~~ not explicitly shown here). The regions with a decrease in P can be found in the shelf and coastal oceans, corresponding to the regions, where N concentrations ~~are increased~~ increase in Tivig et al. (2021). Nevertheless, the  
275 magnitude of P changes is small compared to the spatial variance of surface phosphate concentrations of 0.64 mmol P m<sup>-3</sup> in CTR and 0.85 mmol P m<sup>-3</sup> after 10,000 years of riverine N supply in NEWS-N (0.65 mmol P m<sup>-3</sup> at the start of the simulation). For simulations with inclusion of riverine P without benthic burial of P, surface P concentrations after 10,000 years of ~~rivering~~ riverine P supply are much higher than in the control simulation (Figure 4 ~~c-~~ and ~~d.~~), independent of the additional riverine N supply. Additional P is distributed over the global oceans except for the tropical Atlantic and the north-  
280 western tropical Pacific Ocean. These regions correspond to the northern subtropical gyres and are known to be oligotrophic and phosphate limited (Mather et al., 2008; Martiny et al., 2019). The other extreme is the simulation N+P-BURHIGH, where the modelled ocean is loosing P in all those regions, where surface P concentrations are different from 0 in CTR (Figure 4 f.). Only moderate changes and a spatial variance of 0.69 mmol P m<sup>-3</sup> are the result of the simulation with riverine N and P and low burial functions (N+P-BURLOW, Figure 4 e.).

285 After 10,000 years of ~~simulations with the different scenarios, surface phosphate of simulation, of all experiments including riverine phosphate, the global average P concentration in~~ N+P-BURLOW ~~appears most similar is nearest~~ to observed present-day oceanic conditions (compare observations with data from the World Ocean Atlas (Garcia et al., 2019b) in Figure A4). Nevertheless, each scenario provides a different insight in marine biogeochemical feedbacks.

### 3.2 P influence on oceanic N inventory and N-cycle feedbacks

290 The addition of riverine P to the modelled ocean has a considerable impact on the distribution of simulated NO<sub>3</sub>. At the end of CTR, NEWS-N and the simulation N+P-BURLOW, the N sink by denitrification and the N sources by N<sub>2</sub>-fixation and riverine input are balanced (Figure ~~??3~~ Figure 5). This is not the case for NEWS-P and NEWS-N+P, where the oceanic N inventory slightly increases even at the end of the simulations, and very different from N+P-BURHIGH, where N concentrations decrease continuously. This evolution is also different from P inventory, with its asymptotic shape (Figure 3). In NEWS-P, where only  
295 P is added via river runoff, global oceanic N is only slightly lower than in NEWS-N+P and considerably higher (around ~~200 Pg~~ 6 Pmol N) than in NEWS-N. This is mainly the result of the increase in N-fixation triggered by the additional P flux (Figure 5). Comparing NEWS-N, NEWS-P and N+P-BURLOW shows, that while global benthic denitrification stays nearly constant during the simulations, water column denitrification and N<sub>2</sub>-fixation develop differently depending on the



**Figure 4.** Global distribution of surface-PO<sub>4</sub> concentrations in  $\text{mmol P m}^{-3}$  averaged over the upper 180 m at the end of the respective simulations over 10,000 years. (a) Global distribution of PO<sub>4</sub> concentrations in CTR ;-(b) NEWS-N ;-(c) Difference in PO<sub>4</sub> concentrations between the simulations NEWS-N, NEWS-P ;-(d) NEWS-N NEWS N+P ;-(e) N+P-BURLOW ;-(f) N+P-BURHIGH compared to CTR. Note the different scale in (bf). The magenta-orange colour indicates regions, where the difference in PO<sub>4</sub> concentrations between the respective simulation and CTR is higher than 1.6 exceeds 3.4  $\text{mmol P m}^{-3}$ .

experimental design (Figure 5): additional P leads to an increase in global denitrification and N<sub>2</sub>-fixation. Including low burial functions stabilizes both fluxes over time, but at higher levels than in NEWS-N. In N+P-BURHIGH N-fixation and water column denitrification decrease significantly over the simulations, but seem to stabilize at the end of the 10,000 years. But the

flux of benthic denitrification decreases slower and has not stabilized at the end of the experiment, causing the continuous N decrease in this simulation. To see a stabilized budget here, the simulations should have been much longer.

305 ~~Surface~~ In the upper oceans, NO<sub>3</sub> concentrations increase ~~almost everywhere~~ with supplemental P, except near the main oxygen minimum zones (OMZ) of the Gulf of Guinea, the northern Indian Ocean and the tropical eastern Pacific. Without P burial, N increases ~~globally and~~ especially in the tropical Pacific, the North Atlantic and the higher latitudes, but decreases ~~near the main oxygen minimum zones (OMZ) e.g. in the tropical eastern Pacific because of enhanced denitrification in a more productive ocean.~~ Including P burial leads to smaller N increase, and even a high N decrease ~~as~~ in the simulation with high P burial (Figure 6).

310 ~~At 300-m depth, NO<sub>3</sub> shows generally higher concentrations compared to the surface, except for the experiments with P burial and in the OMZs of the Gulf of Guinea, the northern Indian Ocean and the tropical eastern Pacific (Figure ??).~~ Including riverine P has more impact than including riverine N supply alone. If more P is buried, N concentrations decrease globally. The above pattern of N show increasing concentrations almost everywhere in experiment NEWS-P and NEWS-N+P compared to CTR, but is almost reversed for experiment N+P-BURHIGH, with N concentrations ~~at 300m depth~~ increasing only near the 315 OMZ in the Bay of Bengal and off the Pacific coast of Central-America (Figure ~~??~~6).

These changes in NO<sub>3</sub> concentrations are the result of P fluxes impacting the two main processes of the N feedback-cycle, denitrification (Figure 9) and N<sub>2</sub>-fixation (Figure 8). It is widely assumed, that the surface NO<sub>3</sub> to PO<sub>4</sub> ratio is a dominant controlling factor of these feedback mechanisms (Gruber, 2008). Diazotrophs are limited by P, especially in regions, where 320 light, temperature and iron availability are not limiting. The regions where N<sub>2</sub>-fixation is enhanced in the simulations with riverine P (NEWS-P, NEWS-N+P and N+P-BURLOW) corresponds well with regions of general P limitation (compare with (Kemena et al., 2019), Figure 8). N<sub>2</sub>-fixation is stimulated here by the addition of P to the ocean and is generally sensitive to changes in P supply: In the experiment with high burial rates, these same regions are characterized by decreasing N<sub>2</sub>-fixation rates (Figure 8, (f)).

325 Denitrification is the other feedback mechanism controlling the global ocean N budget. In the simulations where N concentration increases due to higher N<sub>2</sub>-fixation rates, denitrification is enhanced as well. In the simulation with high P burial rates on the other hand, denitrification rates are much lower than in the control simulations in the Gulf of Guinea, the Gulf Bay of Bengal and the eastern tropical Pacific (Figure 9, (f)), leading also to higher N concentrations in these regions ~~at 300-m depth (Figure ?? (Figure 6)).~~

330 Both processes, N<sub>2</sub>-fixation and denitrification, are increased in the simulation with additional riverine P. ~~Except for the Gulf of Bengal, Where~~ they take place in different separated regions of the tropical and subtropical oceans ~~and, these processes~~ lead to an increase in DIN-N in the global surface ocean (~~N<sub>2</sub>~~N<sub>2</sub>-fixation) and a loss in N in the water columns near the main OMZ of the ocean (denitrification). This is not the case for example in the Bay of Bengal.

In our previous study with ~~riverine N only~~ the addition of only riverine N, the "vicious cycle" defined by Landolfi et al. (2013) 335 was suspected to be the reason for a significant decrease in N concentration in the Gulf Bay of Bengal, even though rivers exported additional N to the sea (~~Figure ?? (d), Tivig et al. (2021))~~ (Tivig et al., 2021). Adding riverine P ~~might enhance~~ enhances

this process here, so that the ocean is losing N not only in the Bay of Bengal, but also in the upper tropical Atlantic ocean basin (Figure 7d,g,j,m) near the Gulf of Guinea and in the upper tropical Pacific ocean (Figure 7e,h,k,n).

340 The main driver for this vigorous vicious cycle ~~is the in these simulations is~~ P supply from the rivers and the interplay of different feedback loops in ~~these regions of the ocean the marine biogeochemistry~~ (Figure 10). Considering N on the one side, additional N from rivers or atmospheric fixation enhances marine biological production, which contributes to higher export rates, leading to remineralization and oxygen consumption. In regions, where oxygen is scarce, denitrification will take place and consume N. N<sub>2</sub>-fixation is limited by P. Therefore, adding more P for example from rivers can enhance this process and  
345 by this the vicious cycle. On the other hand, increasing P can also lead to lower oxygen concentrations and therefore more denitrification.

Note that iron availability also plays an important role. ~~The~~ It is a limiting factor of diazotroph growth and hence N<sub>2</sub>-fixation (Landolfi et al., 2013). Iron availability can lead to a shift in the growth of N fixers to regions with higher iron availability, and therefore also decouple denitrification and N<sub>2</sub>-fixation, breaking by this the vicious cycle (Landolfi et al., 2013; Moore et al., 2009)  
350 . Our UVic model represents iron including a static concentration mask. Therefore no interactive response to perturbations of ocean biogeochemistry are possible and iron availability and limitation will not change during the simulations (Nickelsen et al., 2014). ~~N<sub>2</sub>-fixation is influenced by P only.~~

### 3.3 Phosphorus cycling and oxygen minimum zones

Low ventilation of the water column or high rates of remineralization of organic matter can cause oxygen depleted waters.  
355 Niemeier et al. (2017) found that the increase in the marine phosphorus inventory under assumed business-as-usual global warming conditions could lead to a 4- to 5-fold expansion of the suboxic water volume over millennial timescales. Several studies also suggest, that O<sub>2</sub> depletion in coastal regions caused by eutrophication may enhance P release from sediments, thereby providing additional P (Flögel et al., 2011; Wallmann, 2010; Ingall and Jahnke, 1994). These two processes together form a positive feedback loop, enhancing oxygen depletion and expansion of the oxygen minimum zones (Oschlies et al., 2018)  
360 (Figure 10). The processes relevant for this feedback are all included in the ~~experiments with burial~~ burial experiments. ~~The three~~ Three regions, the Gulf of Guinea, the Bay of Bengal and the eastern tropical Pacific ocean are ~~all~~ characterised by low to very low oxygen concentrations (Figure ~~??~~ 11a).

The black contour lines in Figure 11 indicates the extent of the oxygen minimum zones at ~~177~~ 300 m depth, averaged over the last 100 years of the simulations. The oxygen minimum zone is defined here by O<sub>2</sub> concentrations lower than 70 mmol m<sup>-3</sup>.  
365 The main regions with low oxygen concentrations are known to be situated in subsurface waters of the Arabian Sea and in the areas of the eastern boundary upwelling regions in the tropical oceans off California, Peru and Namibia (Oschlies et al., 2018, e.g.). The model results (Figure 11(a)) show, that UVic misplaces the oxygen minimum zone in the Indian Ocean from the Arabian Sea to the the Bay of Bengal. This is similar to other biogeochemical ocean models (Séférián et al., 2020). In reality, the Bay of Bengal is a region with strong seasonality driven by the Asian monsoon system (Löscher et al., 2020). Therefore  
370 highly variable oxygen concentrations inhibit denitrification while high water column denitrification has been observed in the

Arabian Sea (Johnson et al., 2019; Bange et al., 2005).

In the simulations with increased P from rivers oxygen concentrations decrease significantly in all tropical oceans as well as in the eastern boundary regions. P burial damps the oxygen depletion and can even lead to an increase in oxygen concentrations especially in the tropical oceans and in the OMZs (Figure 11). Changes in oxygen concentrations are not limited to oxygen  
375 minimum zones and regions do not exactly correlate with regions of changed denitrification rates (Figure 9).

An observational estimate of today's suboxic water area and volume equals  $30.4 \pm 3$  millions of  $\text{km}^2$  and  $102 \pm 15$  millions of  $\text{km}^3$  respectively, for oxygen concentrations less than  $20 \text{ mmol m}^{-3}$  (Paulmier and Ruiz-Pino, 2009). In CTR, OMZ defined as regions with oxygen concentration less than  $20 \text{ mmol m}^{-3}$  (less than  $70 \text{ mmol m}^{-3}$ ) extend to 13.1 millions of  $\text{km}^3$  (52 millions of  $\text{km}^3$ ). The addition of riverine N ~~only leads alone leads only~~ to small changes in oxygen concentrations with a volume of  
380 OMZ of 13.2 (54) millions of  $\text{km}^3$ ). Adding P leads to a strong increase in OMZ area with up to a 68 (192) millions of  $\text{km}^3$  expansion in NEWS-N+P. With decreased P concentrations, the  $\text{O}_2$  concentration increases globally and the global volume of suboxic waters is reduced to less than 1 (5) millions of  $\text{km}^3$ . Water column denitrification is mainly controlled by the oxygen concentration (Gruber, 2008). While adding riverine N does not impact the ocean denitrification significantly, the addition of P leads to an increase in denitrification in the three main regions with oxygen deficit waters (Figure 9|12). Consequently this leads  
385 also to reduced N concentrations in these same regions. ~~On the other hand~~ However, high P burial rates ~~lead globally worldwide and hence lower P concentrations lead~~ to significantly higher  $\text{O}_2$  concentrations, decreased rates of denitrification and ~~hence therefore~~ higher N concentrations. This is especially the case in regions, where the N cycle feedbacks have limited higher N concentrations before (Figure ??-13 (f)). Including N from rivers only slightly impacts the  $\text{O}_2$  distribution, but can ~~lead to start~~ a negative feedback loop near oxygen minimum zones. In our first experiment this phenomenon also impacted primary  
390 production in these regions, damping a global increase in marine production compared to the control simulation. Adding P from riverine export to the modelled ocean ~~leads to much more changes in N and in the~~ has a higher impact on N and oxygen concentration. As  $\text{O}_2$  plays an important role in the N feedback cycles, the impact is more significant and not only reduced to the OMZ (refer also to Figure 10). ~~Feedbacks in the marine N and P cycle. Yellow colored boxes refer to processes concerning the marine N inventory. Blue refers to the marine P inventory. The black arrows symbolise positive relations or feedbacks. The dashed arrows symbolise negative relations or feedbacks.~~  
395

### 3.4 Primary Production

In Tivig et al. (2021) we showed, that including riverine N had only limited impact on marine productivity, due at least partially to feedback reactions in the marine N cycle. Additional P changes this result significantly (Figure 14): Comparing the simulation with riverine N only to the simulation with riverine P only shows that, at least in our model and on a millennial  
400 timescale, P is more limiting for primary production than N alone. Primary production amounts to  $67 \text{ Pg C yr}^{-1}$  globally in NEWS-P compared to only  $55 \text{ Pg C yr}^{-1}$  in NEWS-N. Even in the low burial simulation, marine biology is more productive than in the simulation without riverine P addition ( $59 \text{ Tg C yr}^{-1}$ ). In N+P-BURHIGH on the other hand, where the ocean is deprived from P at the end of the simulation, production rates decrease in the global tropical oceans to reach a global integral  $33 \text{ Pg C yr}^{-1}$ . Only near the river mouths, where burial has not been effective yet, NPP rates are higher than in the control

405 simulation. To summarize, NPP rates are sensitive to P addition in our model. In the simulations with N and P addition from rivers, the feedbacks are still active, but NPP rates are nevertheless higher than in our simulation with riverine N alone.

#### 4 **Conclusions**Discussion and limitations

In this ~~follow-up study~~ study we compare simulations with different biogeochemical settings for N and P in the same ocean circulation. We find that the addition of riverine phosphorus to the coastal oceans in our model, has more impact on the nitrogen cycle and its internal feedbacks than the addition of N alone (Figure 10). First it impacts the N budget directly by enhancing N-fixation, especially in P limited regions. Secondly, it fuels marine primary production, which leads to an increase in detritus export and hence remineralization, which in turn reduces oxygen concentration. Lower oxygen concentration increases denitrification and therefore balances the nitrogen budget globally. We have also shown that there are regional patterns and that locally, especially in proximity of regions with low oxygen concentrations, the addition of riverine P can lead to N depletion as a consequence of a feedback loop.

415 Nevertheless, our experimental settings has some limitations that have to be accounted for. First, sedimentary N-loss is simulated according to an empirical transfer function based on organic carbon sinking flux to the sediments and bottom-water dissolved oxygen and nitrate. We apply the empirical function from Bohlen et al. (2012). Since our coarse resolution model does not fully resolve narrow continental shelves and coastal dynamics, a subgrid-scale bathymetry parameterization is included (Somes and Oschlies, 2015). Still, UVic underestimates sedimentary N-loss in continental shelves (see also Somes et al. (2017)). The coarse resolution of coastal regions in UVic, with no explicit simulation of the coastal processes, also implies that nutrient retention on the coastal shelf is not included. Hence, the buffer effects of the coasts is not taken into account. Riverine nutrient export to the open ocean might therefore be overestimated in our experiments. These processes could be parametrized like in Sharples et al. (2017); Izett and Fennel (2018) in a future experiment. Finally, submarine groundwater discharge to the oceans is not simulated, but might be an essential component of biogeochemical budgets (Santos et al., 2021; Slomp and Capellen, 2004)

~  
The current study does not incorporate all the feedbacks that may potentially impact the N cycle (e.g., stoichiometric variability of phytoplankton, anthropogenically induced changes in ocean acidification and iron deposition). Also, the spatial patterns for NPP and N<sub>2</sub> fixation differ from observations. In the Indian Ocean, the model simulates too much N<sub>2</sub> fixation in the Bay of Bengal and not enough in the Arabian Sea, a common problem of coarser resolved models, as already stated in this manuscript (Séférian et al., 2020; Moore and Doney, 2007). Of the three major quasi-permanent sites of water column denitrification (Arabian Sea, the Eastern tropical North Pacific and the Eastern tropical South Pacific off the coast of Chile (Codispoti, 2007)), the model is only able to reasonably simulate the eastern tropical North Pacific one. In the Eastern tropical South Pacific, modelled denitrification is too close to the equator and does not occur off the coast of Chile where it should (Keller et al., 2012).

435 Nevertheless, the UVic model has been tested and used for several previous studies of marine biogeochemistry. (Keller et al., 2012) has found, that surface nitrate and phosphate concentrations are similar to observational data (Garcia et al., 2019a). The model



rates of  $N_2$  fixation are within the wide range ( $\sim 7$ - $15 \text{ Tmol N yr}^{-1}$ ) of global  $N_2$  fixation estimates (Codispoti, 2007; Deutsch et al., 2007; C  
440 Despite the model limitations mentioned before, the University of Victoria Earth System Climate Model of intermediate complexity is a useful tool to investigate long-term changes in marine biogeochemistry and has been used and evaluated in many studies before (e.g. Mengis et al., 2020; Somes et al., 2017; Keller et al., 2012).

For the riverine nutrient input, we used the results of the NEWS2 model. Mayorga et al. (2010) and Dumont et al. (2005) evaluated the individual models for the river export and found, that despite uncertainties associated with model inputs, nutrient yield and export were close to observations and the global export estimates were similar to previous publications.

445 Although the riverine nutrient export from NEWS2 includes the anthropogenic component, based on the year 2000, we did not focus on the question in our experiment and the millennial-scale simulations do not provide much insight into the short time variability. Nevertheless, the question of the influence of human activities on the marine N cycle is most central. Human activities have markedly altered the earth's cycles of the nutrients nitrogen and phosphorus. Beusen and Bouwman (2022) have shown that rivers, which have more than 50 % anthropogenic sources and at the same time elevated N:P ratios ( $> 25$ ) contributed

450 36 % to the total global N export to coastal waters. They also found, that from 1970 to ~~Tivig et al. (2021), a new component was added~~ 2015, global N and P delivery to surface waters has increased from 44 to 71  $\text{Tg N yr}^{-1}$  for N (3.1 to 5.1  $\text{Tmol N yr}^{-1}$ ) and 7.1 to 9.7  $\text{Tg P yr}^{-1}$  (0.23 to 0.31  $\text{Tmol P yr}^{-1}$ ) for P, although the export has been decreased in industrialized countries. Changes do not only affect the absolute amount of the nutrients, but also the N:P ratios. Understanding the interplay between these to nutrients is therefore important, to predict changes in coastal and global oceans (Beusen and Bouwman, 2022).

## 455 5 Conclusions

In a previous study, Tivig et al. (2021) added a new component to the global Earth system model UVic, ~~that had been employed before~~ to simulate the effects of N supply from river discharge. They found that, internal feedbacks in the nitrogen cycle mostly compensated for the imposed yearly addition of 1.6 to 3.3  $\text{Tmol}$  of riverine nitrogen and limited the impact on global marine productivity to  $< 2 \%$ . In the current study, riverine phosphorus has been included separately and in addition to N. After 10,000

460 years, only a simulation with low P burial rates reached a steady state. ~~Nevertheless, all simulations showed that including riverine P in the model has more impact on the modelled marine biogeochemistry and biology than the inclusion of N alone. Therefore, we can answer the questions raised at the start:~~

After 10,000 years of simulations, the global amount of P in the ocean has increased significantly in the simulations where riverine P is not balanced by burial sinks and continues to increase ~~by 5.4  $\text{Tg P yr}^{-1}$~~ . In the simulation with low burial rates,

465 global P ~~increase only by 0.5  $\text{Tg P yr}^{-1}$~~  increases only slightly, while the simulation with high burial rates leads to a continual loss in global P ~~by 2.1  $\text{Tg P yr}^{-1}$~~ . The simulations showed that, on millennial timescales, including riverine P in the model has more impact on the modelled marine biogeochemistry and biology than the inclusion of N alone. Therefore, we can answer the questions raised at the start:

- Additional P from river runoff affects marine biology not only near the river mouths but also in regions far off the coasts
- 470 and in the deep oceans.




- Because the additional P (as well as the loss of P in N+P-BURHIGH) affects the two processes denitrification and N<sub>2</sub>-fixation, the N-cycle is also affected and N concentration increases or decreases locally and globally. In simulations where P is added without burial, N concentrations also increase globally, except in regions where denitrification is dominant, in proximity to regions with low to very low oxygen concentrations. Balancing the addition of P by P burial leads to a decrease in the global marine N concentration. The global amount of N at the end of the 10,000 years is lower in the burial simulations than in the simulations without P compensation. This can be attributed to the internal feedbacks: additional P increases N<sub>2</sub>-fixation but also denitrification. Knowing that denitrification consumes 7 mol of NO<sub>3</sub> for every mole of organic N provided by N<sub>2</sub>-fixation, in regions with low oxygen concentration, a 'vicious cycle' leads to a runaway loss of N. In the simulations where P is not limiting, the N loss can be globally compensated by N<sub>2</sub>-fixation in other regions. Where P is reduced because of sedimental burial, even if only low, N<sub>2</sub>-fixation is not effective enough to compensate for the loss in N due to denitrification.
- Adding P in the coastal oceans has significant impact on marine oxygen concentrations. Simulated oxygen concentrations decrease globally but more in the tropical and Nordic oceans, if P is added without an additional sink. In the simulation where P is lost due to sediment burial, however, O<sub>2</sub> concentrations increase in the whole oceans. In Tivig et al. (2021) we found, that O<sub>2</sub> concentrations is only slightly impacted by the addition of riverine N alone.

Finally, our study showed that additional P from riverine input strongly influences marine productivity not only in the coastal oceans, but also in the open oceans worldwide. Our model simulations suggest that, on millennial timescales, the impact of riverine P on ocean biogeochemistry is more important than the one of riverine N. While this result is linked to our model configuration, ~~it is nevertheless relevant for the real ocean~~, with all its limitations described before, it confirms the conclusion of e.g. Tyrrell (1999). In our experiments we also showed that internal feedbacks in the N and P cycle play a crucial role for marine biogeochemistry and can highly impact regional biology, especially in the coastal oceans.

The main sources of P as the “ultimately limiting nutrient” are rivers transporting P provided by weathering and human activities (Giraud et al., 2008; Föllmi, 1995). While weathering processes are active on millennial timescales and therefore only slowly change the amount of P advected by the rivers, human activities have influenced these nutrient transports more rapidly during the last centuries (Beusen et al., 2016). Additionally, changes in atmospheric carbon dioxide concentrations and resulting climate changes ~~has~~ have already and will in the future impact riverine export as well as ocean biogeochemistry (Gao et al., 2023). Although the uncertainties in the real and the modelled nutrient fluxes are still large, our simulation suggests that global ocean biogeochemistry can be affected by the supply of nutrients from rivers and that the global representation of biological activity may be improved by considering riverine export and coastal processes. For this purpose, a better spatial resolution of the coastal oceans as well as more realistic representation of coastal N and P cycling processes could be helpful, also in order to analyse the variability on shorter timescales.


*Code and data availability.* The data and material that support the findings of this study are available through GEOMAR at <https://hdl.handle.net/20.500.12085/85adfd5-bc86-440c-a205-496749a9025f> (Tivig et al., 2024). More information on the original NEWS2 data set is available from the Global NEWS group at the web site <http://icr.ioc-unesco.org/index.php>. Please email Emilio Mayorga at [mayorga@marine.rutgers.edu](mailto:mayorga@marine.rutgers.edu) to obtain this data set.

## **Appendix A: [Supplemental Figures](#)**




**Figure 5.** Timeseries of global nitrogen ~~in all simulations~~ fluxes from NEWS-N (a), NEWS-P (b), N+P-BURLOW (c) and N+P-BURHIGH (d) over the 10,000 simulation years in Pg-Tmol N per year. ~~Simulation description can be found in Table 1~~ Nitrogen fixation (blue solid line) and riverine N input (blue dotted line) are balanced by water column denitrification and benthic denitrification (red solid and red dotted line, respectively). The global sum of all N fluxes is shown as bold black line. Fluxes are given in Table 1 absolute values. Note the different scales of y-axis. The dotted line in grey is the zero-line of the global sum of N fluxes.

Timeseries of global nitrogen fluxes from NEWS-N (a), NEWS-P (b) and N+P-BURLOW (c) over the 10,000 simulation years in Tg N per year. Nitrogen fixation (blue solid line) and riverine N input (blue dotted line) are balanced by water column denitrification and benthic




**Figure 6.** Global distribution of  $\text{NO}_3$  concentration at-averaged over the surface-upper 180 m in  $\text{mmol N m}^{-3}$ . The panel (a) shows  $\text{NO}_3$  concentration in CTR. Panels (b)-(f) show N concentration in the simulations NEWS-N, NEWS-P, NEWS-N+P, N+P-BURLOW and N+P-BURHIGH as difference to CTR.

Global distribution of  $\text{NO}_3$  concentration at 300 m depth in  $\text{mmol N m}^{-3}$ . The panel (a) shows  $\text{NO}_3$  concentration in CTR. Panels (b)–(f) show N concentration in the simulations NEWS-N, NEWS-P, NEWS-N+P, N+P-BURLOW and N+P-BURHIGH as difference to CTR.




**Figure 7.** Difference in the zonal mean concentrations of  $\text{NO}_3$  in the main ocean basins (Atlantic: left column, Pacific: middle, and Indian Ocean: right) in  $\text{mmol N m}^{-3}$  as difference between the simulations NEWS-N, NEWS-P, NEWS-N+P, N+P-BURLOW and N+P-BURHIGH and CTR.




**Figure 8.** Vertical integration of N<sub>2</sub> fixation in mmol N m<sup>-2</sup> yr<sup>-1</sup>. (a) N<sub>2</sub> fixation in CTR. (b)-(f) difference in N<sub>2</sub> fixation between the simulations NEWS-N, NEWS-P, NEWS-N+P, N+P-BURLOW and N+P-BURHIGH and CTR. The cyan colour indicates regions with very low N<sub>2</sub> fixation rates compared to CTR. The orange colour indicates regions with very high N<sub>2</sub> fixation rates compared to CTR.

### Denitrification




**Figure 9.** Vertical integration of denitrification in  $\text{mmol N m}^{-2} \text{ yr}^{-1}$ . (a) Denitrifications rates in CTR. (b)-(f) difference in denitrification rates between the simulations NEWS-N, NEWS-P, NEWS-N+P, N+P-BURLOW and N+P-BURHIGH and CTR. The light blue-cyan colour indicates regions with very low denitrification rates compared to CTR. The light red-orange colour indicates regions with very high denitrification rates compared to CTR.







**Figure 10.** Feedbacks in the marine N and P cycle, due to riverine nutrient supply to the global oceans. Yellow colored boxes refer to processes concerning the marine N inventory. Blue refers to the marine P inventory. Green boxes show riverine nutrient input. The black arrows symbolise positive effects or feedbacks. The dashed arrows symbolise negative effects or feedbacks. Red bordered arrows symbolise processes involved in the vicious cycle (Landolfi et al., 2013).



**Figure 11.** Oxygen concentration at 302-300 m depth in  $\text{mmol O}_2 / \text{m}^3$ . (a) distribution of oxygen concentration in the control simulation, difference in oxygen concentration between the control-simulation and simulations NEWS (b), NEWS-P-NEWS-P (c), NEWS-N-NEWS-N+P (d), PP N+P-Burlow (e), PP N+P-Burhigh (f), respectively, and CTR. The black bold line shows the limits of the oxygen minimum zone at 302 m depth in the control simulation.



**Figure 12.** Difference between N+P-Burlew and CTR simulation for the average of the upper 300 m for  $PO_4$  concentrations in  $mmol P m^{-3}$  (a),  $NO_3$  concentrations in  $mmol N m^{-3}$  (b) and  $O_2$  concentrations in  $mmol O_2 m^{-3}$  (d) and for the vertical integration of denitrification in  $mmol N m^{-2} yr^{-1}$  (c) at the end of the respective simulations over 10,000 years. The red circle shows the region of the Bay of Bengal.



**Figure 13.** Difference between N+P-Burhigh and CTR simulation for the average of the upper 300 m for PO<sub>4</sub> concentrations in mmol P m<sup>-3</sup> (a), NO<sub>3</sub> concentrations in mmol N m<sup>-3</sup> (b) and O<sub>2</sub> concentrations in mmol O<sub>2</sub> m<sup>-3</sup> (d) and for the vertical integration of denitrification in mmol N m<sup>-2</sup> yr<sup>-1</sup> (c) at the end of the respective simulations over 10,000 years. The red circle shows the region of the Bay of Bengal.






Figure 14. Vertical integration of primary production in  $\text{g C m}^{-2} \text{ yr}^{-1}$ .




**Figure A1.** DIN export yield for each discharge point in mmol N m<sup>-2</sup> month<sup>-1</sup> from NEWS2 data set interpolated on the UVic grid for January (a), April (b), July (c) and October (d).




**Figure A2.** Burial flux of P in the N+P-BURHIGH simulation in  $\text{mmol P yr}^{-1}$ .





**Figure A3.** Global profile of  $PO_4$  in  $\text{mmol m}^{-3}$  in the simulations NEWS, CTR, NEWS-P, NEWS-N+P, NEWS-Burlow, News-Burhigh and from the World Ocean Atlas (WOA). (a) global average of  $PO_4$  (b), Global profiles of misfit between each simulation and the WOA dataset (c), global average profiles of  $PO_4$  in the Atlantic Ocean (d), global average profiles of  $PO_4$  in the Pacific Ocean (e), global average profiles of  $PO_4$  in the Indian Ocean (f).



**Figure A4.** Global profile of  $NO_3$  in  $mmol\ m^{-3}$  in the simulations NEWS, CTR, NEWS-P, NEWS-N+P, NEWS-Burlow, News-Burhigh and from the World Ocean Atlas (WOA). (a) global average of  $NO_3$  (b), Global profiles of misfit between each simulation and the WOA dataset (c), global average profiles of  $NO_3$  in the Atlantic Ocean (d), global average profiles of  $NO_3$  in the Pacific Ocean (e), global average profiles of  $NO_3$  in the Indian Ocean (f).

*Author contributions.* MT developed the research concept in discussion with AO and DPK. DPK provided the initial model code, which was further developed, run, and analysed by MT. MT analysed the model output and visualised the results. MT wrote the manuscript with contributions from all co-authors.

510 *Competing interests.* The authors declare that they have no conflict of interest.

*Acknowledgements.* We gratefully acknowledge E. Mayorga, S. Seitzinger and their co-authors for making their database of Global Nutrient Export from WaterSheds 2 (NEWS2) available for our study. AO and DPK acknowledge funding from the European Union's Horizon 2020 Research and Innovation Program under grant 820989 (project COMFORT, "Our common future ocean in the Earth system — quantifying coupled cycles of carbon, oxygen, and nutrients for determining and achieving safe operating spaces with respect to tipping points") and OceanNETs (grant no. #869357). The work reflects only the author's view; the European Commission and their executive agency are not responsible for any use that may be made of the information the work contains. This work was also supported by the German Research Foundation (DFG) as part of the research project SFB 754 "Climate-Biogeochemistry Interactions in the Tropical Ocean". We would also thank the GEOMAR's Biogeochemical Modelling group for many fruitful discussions.

515

## 520 **References**

- Allaby, M.: Liebig's law of the minimum, <https://doi.org/10.1093/acref/9780199567669.013.3219>, 2010.
- Altabet, M. A.: Constraints on oceanic N balance/imbalance from sedimentary 15N records, *Biogeosciences*, 4, 75–86, <https://doi.org/10.5194/bg-4-75-2007>, 2006.
- Bange, H., Naqvi, S., and Codispoti, L.: The nitrogen cycle in the Arabian Sea, *Progress in Oceanography*, 65(2-4), 145–158,   
525 <https://doi.org/10.1016/j.pocean.2005.03.002>, 2005.
- Baturin, G. N.: Phosphorus Cycle in the Ocean, *Lithology and Mineral Resources*, 38, 101–119, 2003.
- Benitez-Nelson, C. R.: The biogeochemical cycling of phosphorus in marine systems, *Earth-Science Reviews*, 51, 109–135,   
[https://doi.org/10.1016/S0012-8252\(00\)00018-0](https://doi.org/10.1016/S0012-8252(00)00018-0), 2000.
- Beusen, A. H. W. and Bouwman, A. F.: Future projections of river nutrient export to the global coastal ocean show persisting nitrogen and   
530 phosphorus distortion, *Frontiers:Water*, 4, 893 585, <https://doi.org/10.3389/frwa.2022.893585>, 2022.
- Beusen, A. H. W., Bouwman, A. F., Beek, L. P. H. V., Mogollon, J. M., and Middelburg, J. J.: Global riverine N and P transport to ocean increased during the 20th century despite increased retention along the aquatic continuum, *Biogeosciences*, 13, 2441–2451,   
<https://doi.org/10.5194/bg-13-2441-2016>, 2016.
- Bohlen, L., Dale, A. W., and Wallmann, K.: Simple transfer functions for calculating benthic fixed nitrogen losses and C:N:P regeneration   
535 ratios in global biogeochemical models, *Global Biogeochemical Cycles*, 26, GB3029, <https://doi.org/10.1029/2011GB004198>, 2012.
- Cappellen, P. V. and Maavara, T.: Rivers in the Anthropocene: Global scale modifications of riverine nutrient fluxes by damming, *Ecology & Hydrobiology*, 16, 106–111, 2016.
- Claussen, M., Mysak, L. A., Weaver, A. J., Crucifix, M., Fichet, T., Loutre, M.-F., Weber, S., Alcamo, J., Alexeev, V., Berger, A., Calov, R., Ganopolski, A., Goosse, H., Lohmann, G., Lunkeit, F., Mokhov, I., Petoukhov, V., Stone, P., and Wang, Z.: Earth System   
540 Models of Intermediate Complexity: Closing the Gap in the Spectrum of Climate System Models, *Climate Dynamics*, 18, 579–586, <https://doi.org/10.1007/s00382-001-0200-1>, 2002.
- Codispoti, L. A.: Is the ocean losing nitrate?, *Nature*, 376, 1995.
- Codispoti, L. A.: An oceanic fixed nitrogen sink exceeding 400 Tg Na-1 vs the concept of homeostasis in the fixed-nitrogen inventory, *Biogeosciences*, 4, 233–253, 2007.
- 545 Codispoti, L. A., Brandes, J. A., Christensen, J. P., Devol, A. H., Naqvi, S. A., Paerl, H. W., and Yoshinari, T.: The oceanic fixed nitrogen and nitrous oxide budgets: Moving targets as we enter the anthropocene?, *Scientia Marina*, 65, 85–105, 2001.
- Colman, A. S. and Holland, H. D.: The Global Diagenetic Flux of Phosphorus from Marine Sediments to the Oceans: Redox Sensitivity and the Control of Atmospheric Oxygen Levels., in: *Marine Authigenesis: From Global to Microbial*, vol. 66, p. 53–75, SEPM Society for Sedimentary Geology, <https://doi.org/10.2110/pec.00.66.0053>, 2000.
- 550 Compton, J., Mallinson, D., Glenn, C. R., Filippelli, G., Föllmi, K., Shields, G., and Zanin, Y.: Variations in the global phosphorus cycle, in: *Marine Authigenesis: From Global to Microbial*, vol. 66, pp. 21–33., SEPM Society for Sedimentary Geology, <https://doi.org/10.2110/pec.00.66.0021>, 2000.
- Delaney, M. L.: Phosphorus accumulation in marine sediments and the oceanic phosphorus cycle, *Global Biogeochemical Cycles*, 12 No.4, 563–572, 1998.
- 555 Deutsch, C., Gruber, N., Key, R. M., and Sarmiento, J. L.: Denitrification and N<sub>2</sub> fixation in the Pacific Ocean, *Global Biogeochemical Cycles*, 15, 483–506, 2001.

- Deutsch, C., Sarmiento, J. L., Sigman, D. M., Gruber, N., and Dunne, J. P.: Spatial coupling of nitrogen inputs and losses in the ocean, *Nature*, 445, 163–167, 2007.
- 560 Dumont, E., Harrison, J. A., Kroeze, C., Bakker, E. J., and Seitzinger, S. P.: Impacts of Atmospheric Anthropogenic Nitrogen on the Open Ocean, *Global Biogeochemical Cycles*, 19, GB4S02, <https://doi.org/10.1029/2005GB002488>, 2005.
- Eby, M., Zickfeld, K., Montenegro, A., Archer, D., Meissner, K. J., and Weaver, A. J.: Lifetime of Anthropogenic Climate Change: Millennial Time Scales of Potential CO<sub>2</sub> and Surface Temperature Perturbations, *Journal of Climate*, 22, 2501–2511, <https://doi.org/10.1175/2008JCLI2554.1>, 2009.
- Falkowski, P. G.: Evolution of the nitrogen cycle and its influence on the biological sequestration of CO<sub>2</sub> in the ocean, *Nature*, 387, 272–275, <https://doi.org/10.1038/387272a0>, 1997.
- 565 Fanning, A. F. and Weaver, A. J.: An atmospheric energy-moisture balance model: Climatology, interpentadal climate change, and coupling to an ocean general circulation model, *Journal of Geophysical Research*, 101, 15 111–15 128, <https://doi.org/10.1029/96JD01017>, 1996.
- Filipelli, G. M.: The Global Phosphorus Cycle: Past, Present, and Future, *Elements*, 4, 89–95, <https://doi.org/10.2113/GSELEMENTS.4.2.89>, 2008.
- 570 Flögel, S., Wallmann, K., Poulson, C. J., Zhou, J., Oschlies, A., Voigt, S., and Kuhnt, W.: Simulating the biogeochemical effects of volcanic CO<sub>2</sub> degassing on the oxygen-state of the deep ocean during the Cenomanian/Turonian Anoxic Event (OAE2), *Earth Planet Scientific Letters*, 305, 371–384, <https://doi.org/10.1016/j.epsl.2011.03.0181>, 2011.
- Föllmi, K. B.: The Phosphorus Cycle, phosphogenesis and marine phosphate-rich deposits, *Earth-Science Reviews*, 40, 55–124, [https://doi.org/10.1016/0012-8252\(95\)00049-6](https://doi.org/10.1016/0012-8252(95)00049-6), 1995.
- 575 Galloway, J. N., Dentener, F., Capone, D., Boyer, E., Howarth, R., Seitzinger, S., Asner, G., Cleveland, C., Green, P., Holland, E., Karl, D., Michaels, A., Porter, J., Townsend, A., and Vörösmarty, C.: Nitrogen cycles: past, present, and future, *Biogeochemistry*, 70, 153–226, <https://doi.org/10.1007/s10533-004-0370-0>, 2004.
- Gao, S., Schwinger, J., Tjiputra, J., Bethke, I., Hartmann, J., Mayorga, E., and Heinze, C.: Riverine impact on future projections of marine primary production and carbon uptake, *Biogeochemistry*, 20, 93–119, <https://doi.org/10.5194/bg-20-93-2023>, 2023.
- 580 Garcia, H. E., Weathers, K., Paver, C. R., Smolyar, I., Boyer, T. P., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Baranova, O. K., Seidov, D., and Reagan, J. R.: Volume 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Oxygen Saturation, in: *World Ocean Atlas 2018*, edited by Editor, A. M. T., p. 38pp, NOAA Atlas NESDIS 83, <https://www.nodc.noaa.gov/OC5/woa18/pubwoa18.html>, 2019a.
- Garcia, H. E., Weathers, K., Paver, C. R., Smolyar, I., Boyer, T. P., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Baranova, O. K., Seidov, D., and Reagan, J. R.: Volume 4: Dissolved Inorganic Nutrients (phosphate, nitrate and nitrate+nitrite, silicate), in: *World Ocean Atlas 2018*, edited by Editor, A. M. T., p. 35pp, NOAA Atlas NESDIS 84, [https://www.ncei.noaa.gov/data/oceans/woa/WOA18/DOC/woa18\\_vol4.pdf](https://www.ncei.noaa.gov/data/oceans/woa/WOA18/DOC/woa18_vol4.pdf), 2019b.
- 585 Garnier, J., Beusen, A., Thieu, V., Billen, G., and Bouwman, L.: N:P:Si nutrient export ratios and ecological consequences in coastal seas evaluated by the ICEP approach, *Global Biogeochemical Cycles*, 24, GB0A05, <https://doi.org/10.1029/2009GB003583>, 2010.
- Giraud, X., Quéré, C. L., and da Cunha, L. C.: Importance of coastal nutrient supply for global ocean biogeochemistry, *Global Biogeochemical Cycles*, 22, GB2025, <https://doi.org/10.1029/2006GB002717>, 2008.
- 590 Gruber, N.: The Dynamics of the Marine Nitrogen Cycle and its Influence on Atmospheric CO<sub>2</sub> Variations, in: *The Ocean Carbon Cycle and Climate*, edited by Follows, M. and Oguz, T., pp. 97–148, Springer Netherlands, Dordrecht, [https://doi.org/10.1007/978-1-4020-2087-2\\_4](https://doi.org/10.1007/978-1-4020-2087-2_4), 2004.

- 595 Gruber, N.: The marine nitrogen cycle: Overview and challenges, in: Nitrogen in the Marine Environment, edited by et al., D. G. C., chap. 1, pp. 1–50, Academic Press, San Diego, California, 2 edn., 2008.
- Gruber, N. and Sarmiento, J. L.: Global patterns of marine nitrogen fixation and denitrification, *Global Biogeochemical Cycles*, 11, 235–266, <https://doi.org/10.1029/97GB00077>, 1997.
- 600 Harrison, J. A., Seitzinger, S. P., Bouwman, A. F., Caraco, N. F., Beusen, A. H. W., and Vörösmarty, C. J.: Dissolved inorganic phosphorus export to the coastal zone: Results from a spatially explicit, global model, *Global Biogeochemical Cycles*, 19, GB4S03, <https://doi.org/10.1029/2004GB002357>, 2005.
- Ingall, E. and Jahnke, R.: Evidence for enhanced phosphorus regeneration from marine sediments overlain by oxygen depleted waters, *Geochimica et Cosmochimica Acta*, 58, 2571–2575, 1994.
- Izett, J. G. and Fennel, K.: Estimating the Cross-Shelf Export of Riverine Materials: Part 1. General Relationships From an Idealized Numerical Model, *Global Biogeochemical Cycles*, 32, 160–175, <https://doi.org/10.1002/2017GB005667>, 2018.
- 605 Johnson, K. S., Riser, S. C., and Ravichandran, M.: Oxygen Variability Controls Denitrification in the Bay of Bengal Oxygen Minimum Zone, *Geophysical Research Letters*, 46, 804–811, <https://doi.org/10.1029/2018GL079881>, 2019.
- Karl, D., Micallef, A., Bergman, B., Capone, D., Carpenter, E., Letelier, R., Lipschultz, F., Paerl, H., Sigman, D., and Stal, L.: Dinitrogen fixation in the world’s oceans, *Biogeochemistry*, 57/58, 47–98, 2002.
- Keller, D. P., Oschlies, A., and Eby, M.: A new marine ecosystem model for the University of Victoria Earth System Climate Model, *Geoscientific Model Development*, 5, 1195–1220, <https://doi.org/10.5194/gmd-5-1195-2012>, 2012.
- 610 Kemena, T. P., Landolfi, A., Oschlies, A., Wallmann, K., and Dale, A. W.: Ocean phosphorus inventory: large uncertainties in future projections on millennial timescales and their consequences for ocean deoxygenation, *Earth System Dynamics*, 10, 539–553, <https://doi.org/10.5194/esd-10-539-2019>, 2019.
- Lacroix, F., Ilyina, T., and Hartmann, J.: Oceanic CO<sub>2</sub> outgassing and biological production hotspots induced by pre-industrial river loads of nutrients and carbon in a global modeling approach, *Biogeosciences*, 17, 55–88, <https://doi.org/10.5194/bg-17-55-2020>, 2020.
- 615 Landolfi, A., Dietze, H., Koeve, W., and Oschlies, A.: Overlooked runaway feedback in the marine nitrogen cycle: the vicious cycle, *Biogeosciences*, 10, 1351–1363, <https://doi.org/10.5194/bg-10-1351-2013>, 2013.
- Landolfi, A., Koeve, W., Dietze, H., Kähler, P., and Oschlies, A.: A new perspective on environmental controls of marine nitrogen fixation, *Geophysical Research Letters*, 42, 4482–4489, <https://doi.org/10.1002/2015GL063756>, 2015.
- 620 Löscher, C. R., Mohr, W., Bange, H. W., and Canfield, D. E.: No nitrogen fixation in the Bay of Bengal?, *Biogeosciences*, 17, 851–864, <https://doi.org/10.5194/bg-17-851-2020>, 2020.
- Martiny, A. C., Lomas, M. W., Fu, W., Boyd, P. W., Chen, Y.-I. L., Cutter, G., Ellwood, M. J., Furuya, K., Hashihama, F., Kanda, J., Karl, D. M., Kodama, T., Li, Q. P., Ma, J., Moutin, T., Woodward, E. M. S., and Moore, J. K.: Biogeochemical controls of surface ocean phosphate, *Science Advances*, 5, eaax0341, <https://doi.org/10.1126/sciadv.aax0341>, 2019.
- 625 Mather, R. L., Reynolds, S. E., Wolff, G. A., Williams, R. G., Torres-Valdes, S., Woodward, E. M. S., Landolfi, A., Pan, X., Sanders, R., and Achterberg, E. P.: Phosphorus cycling in the North and South Atlantic Ocean subtropical gyres, *Nature Geoscience*, 1, 439–443, <https://doi.org/10.1038/ngeo232>, 2008.
- Mayorga, E., Seitzinger, S. P., Harrison, J. A., Dumont, E., Beusen, A. H. W., Bouwman, A., Fekete, B. M., Kroeze, C., and van Drecht, G.: Global Nutrient Export from WaterSheds 2 (NEWS 2): Model development and implementation, *Environmental Modelling and Software*, 630 25, 837–853, <https://doi.org/10.1016/j.envsoft.2010.01.007>, 2010.

- Mengis, N., Keller, D. P., MacDougall, A. H., Eby, M., Wright, N., Meissner, K. J., Oeschlies, A., Schmittner, A., MacIsaac, A. J., Matthews, H. D., and Zickfeld, K.: Evaluation of the University of Victoria Earth System Climate Model version 2.10 (UVic ESCM 2.10), *Geoscientific Model Development*, 13, 4183–4204, <https://doi.org/10.5194/gmd-13-4183-2020>, 2020.
- 635 Monteiro, F. M., Pancost, R. D., Ridgwell, A., and Donnadieu, Y.: Nutrients as the dominant control on the spread of anoxia and euxinia across the Cenomanian-Turonian oceanic anoxic event (OAE2): Model-data comparison, *Paleoceanography*, 27, PA4209, <https://doi.org/10.1029/2012PA002351>, 2012.
- Moore, C. M., Mills, M. M., Achterberg, E. P., Geider, R. J., LaRoche, J., Lucas, M. I., McDonagh, E. L., X. Pan, A. J. P., Rijkenberg, M. J. A., Suggett, D. J., Ussher, S. J., and S.Woodward, E. M.: Large-scale distribution of Atlantic nitrogen fixation controlled by iron availability, *Nature Geoscience Letters*, 2, 867–871, <https://doi.org/10.1038/NGEO667>, 2009.
- 640 Moore, J. K. and Doney, S. C.: Iron availability limits the ocean nitrogen inventory stabilizing feedbacks between marine denitrification and nitrogen fixation, *Global Biogeochemical Cycles*, 21, GB2001, <https://doi.org/10.1029/2006GB002762>, 2007.
- Nickelsen, L., Keller, D., and Oeschlies, A.: A dynamic marine iron cycle module coupled to the University of Victoria Earth System Model: the Kiel Marine Biogeochemical Model 2 (KMBM2) for UVic 2.9, *Geoscientific Model Development Discussions*, 7, 8505–8563, <https://doi.org/10.5194/gmdd-7-8505-2014>, 2014.
- 645 Niemeier, D., Kemena, T. P., Meissner, K. J., and Oeschlies, A.: A model study of warming-induced phosphorus–oxygen feedbacks in open-ocean oxygen minimum zones on millennial timescales, *Earth System Dynamics*, 8, 357–367, <https://doi.org/10.5194/esd-8-357-2017>, 2017.
- Oeschlies, A., P.Brandt, Stramma, L., and Schmidtko, S.: Drivers and mechanisms of ocean deoxygenation, *Nature Geoscience*, 11, 467–473, <https://doi.org/10.1038/s41561-018-0152-2>, 2018.
- 650 Oeschlies, A., Koeve, W., Landolfi, A., and Kähler, P.: Loss of fixed nitrogen causes net oxygen gain in a warmer future ocean, *Nature Communications*, <https://doi.org/10.1038/s41467-019-10813-w>, 2019.
- Palastanga, V., Slomp, C. P., and Heinze, C.: Long-term controls on ocean phosphorus and oxygen in a global biogeochemical model, *Global Biogeochemical Cycles*, 25, GB3024, <https://doi.org/10.1029/2010GB003827>, 2011.
- Partanen, A.-I., Keller, D. P., Korhonen, H., and Matthews, H. D.: Impacts of sea spray geoengineering on ocean biogeochemistry, *Geophysical Research Letters*, 43, 7600–7608, <https://doi.org/10.1002/2016GL070111>, 2016.
- 655 Paulmier, A. and Ruiz-Pino, D.: Oxygen minimum zones (OMZs) in the modern ocean, *Global Biogeochemical Cycles*, 25, GB3024, <https://doi.org/10.1016/j.pocean.2008.08.001>, 2009.
- Redfield, A. C., Ketchum, B. H., and Richards, F. A.: The Influence of Organisms on the Composition of Sea-Water, in: *The Sea Vol. 2*, edited by Hill, M. N., pp. 26–27, Interscience, New York, 1963.
- 660 Ruttenger, K. C.: The Global Phosphorus Cycle, in: *Treatise on Geochemistry*, edited by Schlesinger, W., Elsevier, <https://doi.org/https://doi.org/10.1016/B0-08-043751-6/08153-6>, 2003.
- Santos, I. R., Chen, X., Lecher, A. L., Sawyer, A. H., Moosdorf, N., Rodellas, V., Tamborski, J., Cho, H.-M., Dimova, N., Sugimoto, R., Bonaglia, S., Li, H., Hajati, M.-C., and Li, L.: Submarine groundwater discharge impacts on coastal nutrient biogeochemistry, vol. 2, p. 307–323, <https://doi.org/10.1038/s43017-021-00152-0>, 2021.
- 665 Schmittner, A., Oeschlies, A., Giraud, X., Eby, M., and Simmons, H. L.: A global model of the marine ecosystem for long-term simulations: Sensitivity to ocean mixing buoyancy forcing, particle sinking, and dissolved organic matter cycling, *Global Biogeochemical Cycles*, 19, GB3004, <https://doi.org/10.1029/2004GB002283>, 2005.

- Schmittner, A., Oschlies, A., Matthews, H. D., and Galbraith, E. D.: Future changes in climate, ocean circulation, ecosystems, and biogeochemical cycling simulated for a business-as-usual CO<sub>2</sub> emission scenario until year 4000 AD, *Global Biogeochemical Cycles*, 22, GB1013, <https://doi.org/10.1029/2007GB002953>, 2008.
- Seitzinger, S. P., Mayorga, E., Bouwman, A. F., Kroeze, C., Beusen, A. H. W., Billen, G., Drecht, G. V., Dumont, E., Fekete, B. M., Garnier, J., and Harrison, J. A.: Global river nutrient export: A scenario analysis of past and future trends, *Global Biogeochemical Cycles*, 24, GB0A08, <https://doi.org/10.1029/2009GB003587>, 2010.
- Sharples, J., Middelburg, J. J., Fennel, K., and Jickells, T. D.: What proportion of riverine nutrients reaches the open ocean?, *Global Biogeochemical Cycles*, 31, 39–58, <https://doi.org/10.1002/2016GB005483>, 2017.
- Slomp, C. P. and Capellen, P. V.: Nutrient inputs to the coastal ocean through submarine groundwater discharge: controls and potential impact, *Journal of Hydrology*, 295, 64–86, <https://doi.org/10.1016/j.jhydrol.2004.02.018>, 2004.
- Sohm, J. A., Webb, E. A., and Capone, D. G.: Emerging patterns of marine nitrogen fixation, *Nature Reviews Microbiology*, 9, 499–508, <https://doi.org/10.1038/nrmicro2594>, 2011.
- Somes, C. J. and Oschlies, A.: On the influence of “non-Redfield” dissolved organic nutrient dynamics on the spatial distribution of N<sub>2</sub> fixation and the size of the marine fixed nitrogen inventory, *Global Biogeochemical Cycles*, 29, 973–993, <https://doi.org/10.1002/2014GB005050>, 2015.
- Somes, C. J., Schmittner, A., Galbraith, E. D., Lehmann, M. F., Altabet, M. A., Montoya, J. P., Letelier, R. M., Mix, A. C., Bourbonnais, A., and Eby, M.: Simulating the global distribution of nitrogen isotopes in the ocean, *Global Biogeochemical Cycles*, 24, GB4019, <https://doi.org/10.1029/2009GB003767>, 2010b.
- Somes, C. J., Oschlies, A., and Schmittner, A.: Isotopic constraints on the pre-industrial oceanic nitrogen budget, *Biogeosciences*, 10, 5889–5910, <https://doi.org/10.5194/bg-10-5889-2013>, 2013.
- Somes, C. J., Schmittner, A., Muglia, J., and Oschlies, A.: SA Three-Dimensional Model of the Marine Nitrogen Cycle during the Last Glacial Maximum Constrained by Sedimentary Isotopes, *Frontiers in Marine Science*, 4:108, <https://doi.org/10.3389/fmars.2017.00108>, 2017.
- Séférián, R., Berthet, S., Yool, A., Palmiéri, J., Bopp, L., Tagliabue, A., Kwiatkowski, L., Aumont, O., Christian, J., Dunne, J., Gehlen, M., Ilyina, T., John, J. G., Li, H., Long, M. C., Luo, J. Y., Nakano, H., Romanou, A., Schwinger, J., Stock, C., Santana-Falcón, Y., Takano, Y., Tjiputra, J., Tsujino, H., Watanabe, M., Wu, T., Wu, F., and Yamamoto, A.: Tracking Improvement in Simulated Marine Biogeochemistry Between CMIP5 and CMIP6, *Current Climate Change Reports*, 6, 95–119, <https://doi.org/10.1007/s40641-020-00160-0>, 2020.
- Tivig, M., Keller, D. P., and Oschlies, A.: Riverine nitrogen supply to the global ocean and its limited impact on global primary production: a feedback study using an Earth system model, *Biogeosciences*, 18, 5327–5350, <https://doi.org/10.5194/bg-18-5327-2021>, 2021.
- Tivig, M., Keller, D. P., and Oschlies, A.: UVic simulation with riverine nutrient export from NEWS2 dataset., <https://doi.org/hdl:20.500.12085/85adfed5-bc86-440c-a205-496749a9025f>, 2024.
- Tyrrell, T.: The relative influences of nitrogen and phosphorus on oceanic primary production, *Nature*, 400, 525–531, 1999.
- Voss, M., Bange, H. W., Dippner, J. W., Middelburg, J. J., Montoya, J. P., and Ward, B.: The marine nitrogen cycle: recent discoveries, uncertainties and the potential relevance of climate change, *Philosophical Transactions of the Royal Society B*, 368, 20130 121, <https://doi.org/10.1098/rstb.2013.0121>, 2013.
- Wallmann, K.: Phosphorus imbalance in the global ocean?, *Global Biogeochemical Cycles*, 24, GB4030, <https://doi.org/10.1029/2009GB003643>, 2010.



- 705 Wang, W.-L., Moore, J. K., Martiny, A. C., and Primeau, F. W.: Convergent estimates of marine nitrogen fixation, *Nature*, 566, 205–211, <https://doi.org/10.1038/s41586-019-0911-2>, 2019.
- Weaver, A. J., Eby, M., Wiebe, E. C., Bitz, C., Duffy, P. B., Ewen, T. L., Fanning, A. F., Holland, M. M., MacFadyen, A., Matthews, H. D., Meissner, K. J., Saenko, O., Schmittner, A., Wang, H., and Yoshimori, M.: The UVic Earth System Climate Model: Model Description, Climatology, and Applications to Past, Present and Future Climates, *Atmosphere-ocean*, 39(4), 361–428, <https://doi.org/10.1080/07055900.2001.9649686>, 2001.
- 710 Zehr, J. P. and Capone, G.: Changing perspectives in marine nitrogen fixation, *Science*, 368, <https://doi.org/10.1126/science.aay9514>, 2020.