

# Invisible aerosol layers: improved lidar detection capabilities by means of laser-induced aerosol fluorescence

Benedikt Gast<sup>1</sup>, Cristofer Jimenez<sup>1</sup>, Albert Ansmann<sup>1</sup>, Moritz Haarig<sup>1</sup>, Ronny Engelmann<sup>1</sup>, Felix Fritzsch<sup>1</sup>, Athena A. Floutsi<sup>1</sup>, Hannes Griesche<sup>1</sup>, Kevin Ohneiser<sup>1</sup>, Julian Hofer<sup>1</sup>, Martin Radenz<sup>1</sup>, Holger Baars<sup>1</sup>, Patric Seifert<sup>1</sup>, and Ulla Wandinger<sup>1</sup>

<sup>1</sup>Leibniz Institute for Tropospheric Research, Leipzig, Germany

**Correspondence:** Benedikt Gast (bgast@tropos.de)

**Abstract.** One of the most powerful instruments for studying aerosol particles and their interactions with the environment is atmospheric lidar. In recent years, fluorescence lidar has emerged as a useful tool for identifying aerosol particles due to its link with biological content. Since 2022, this technique has been implemented in Leipzig, Germany. This paper describes the experimental setup and data analysis, with a special emphasis on the characterization of the new fluorescence channel centered at 466 nm. The new capabilities of the fluorescence lidar are examined and corroborated through several case studies. Most of the measurement cases considered are from the spring and summer of 2023, when large amounts of biomass-burning aerosol from the huge forest fires in Canada were transported to Europe. The fluorescence of the observed aerosol layers is characterized. For wildfire smoke, the fluorescence capacity was typically in the range of  $2\text{--}7 \times 10^{-4}$ , which aligns well with the values reported in the literature, with slightly larger values. The key aspects of this study are the capabilities of the fluorescence lidar technique, which can potentially improve not only the typing but even the detection of aerosol particles. In several measurement cases with an apparently low aerosol load, the fluorescence channel clearly revealed the presence of aerosol layers that were not detectable with the traditional elastic-backscatter channels. This capability is discussed in detail and linked to the fact that fluorescence backscattering is related to aerosol particles only. A second potential of the fluorescence technique is the distinction between non-activated aerosol particles and hydrometeors, given water's inability to exhibit fluorescence. A smoke-cirrus case study suggests an influence of the aerosol layer on cloud formation, as it seems to affect the elastic backscatter coefficient within the cloud passing time. These mentioned applications promise huge advancements towards a more detailed view of the aerosol-cloud interaction problem.

## 1 Introduction

A crucial player in the atmospheric system are aerosol particles, given their role in various processes that ultimately shape the Earth's energy and hydrological budgets. Firstly, aerosol particles scatter and absorb radiation, affecting the energy balance on a global scale. By serving as cloud condensation nuclei (CCN) or ice nucleating particles (INPs), these particles can impact the microphysical properties of clouds (Liu et al., 2014), making them more or less reflective depending on the aerosol situation. In the case of liquid-water clouds, this effect has been largely studied (Twomey, 1974, 1977; Twomey et al., 1984; Quaas

et al., 2020), for ice clouds it is a rather new topic (Patnaude and Diao, 2020; Maciel et al., 2023). Because the microphysical properties of a cloud play a major role in its development and the formation of precipitation, aerosol conditions can further affect the extension and lifetime of cloud events and therefore the global albedo (Albrecht, 1989; Stevens and Feingold, 2009). Highly absorbing aerosol particles might even affect clouds via the so-called semi-direct effect, which can manifest, e.g., in the evaporation of cloud droplets due to an aerosol-heated environment (Hansen et al., 1997). Aerosol effects on the ice phase of cloud formation only complicate the picture. Multiple efforts have been made to analyze the role of aerosol particles as INPs 30 in mixed-phase clouds via heterogeneous freezing and the global effect (Lohmann, 2017). As for cirrus clouds, recent studies suggest that heterogeneous freezing in cirrus clouds, particularly via smoke particles, needs to be explored in more detail (Ans-40 man et al., 2021; Veselovskii et al., 2022a; Mamouri et al., 2023; Ansmann et al., 2024a, b). To improve our understanding of these complex aerosol-cloud interaction processes, reliable detection and characterization of atmospheric aerosol particles are essential.

35 Multi-wavelength polarization lidars are powerful tools to detect and characterize aerosol particles. After decades of study, several classification schemes are available in the literature (Floutsi et al., 2023; Groß et al., 2013; Burton et al., 2012), mostly relying on intensive (i.e., concentration-independent) optical properties such as the lidar ratio, particle depolarization ratio and Ångström exponent. However, although clear signatures can be expected for some particle types (e.g., low particle depolarization and low lidar ratios for marine aerosol particles), some limitations remain. Distinguishing between stratospheric smoke 40 and volcanic sulfates or separating between tropospheric smoke and urban pollution remain difficult tasks, as their typical ranges of values for particle depolarization and lidar ratios partially overlap. Additional information, such as the fluorescence of atmospheric aerosol particles, may be required to address these typing difficulties (Veselovskii et al., 2022b).

45 Laser-induced fluorescence is a well-established technique, serving as the basis of several remote-sensing applications. Fluorescence lidars have been around for a while, but their application has mostly focused on water composition (Palmer et al., 2013; Cadondon et al., 2020) and vegetation (Edner et al., 1994). In the domain of atmospheric research, efforts have mostly gone towards the detection of single molecules (McIlrath, 1980; Wang et al., 2021). To investigate the fluorescence of atmospheric aerosol particles, the experiments have been mostly performed through in situ probing (Pinnick et al., 2004; Pan, 2015; Zhang et al., 2019; Kawana et al., 2021). As an example, Pan (2015) analyzed the fluorescence of aerosol particles by measurements 50 with an ultraviolet aerodynamic particle sizer (UV-APS) and reported a spectral range for atmospheric fluorescence of 400 to 650 nm, when excited at 355 nm. A first hint towards the observation of atmospheric aerosol fluorescence with lidar came in 2005, when Immler et al. (2005) observed an inelastic backscatter signal in the water vapor Raman channel (i.e., at 407 nm) that was not produced by Raman scattering. They attributed it to the laser-induced fluorescence emission from biomass-burning (BB) aerosol particles and already linked the aerosol fluorescence to organic compounds. In the following, the first atmospheric fluorescence lidars were set up. Sugimoto et al. (2012) constructed a lidar spectrometer and studied the fluorescence 55 of Asian dust and pollution aerosols in the lower troposphere. The first advanced multi-channel atmospheric lidar system with fluorescence technology was implemented at Lindenberg observatory of the German Meteorological Service (DWD). Initially, Reichardt (2014) observed atmospheric aerosol fluorescence with a lidar spectrometer that was dedicated to measurements of atmospheric water content. He established the fluorescence capacity as a new intensive aerosol property/parameter, which is

defined as the ratio of the fluorescence backscatter coefficient to an elastic particle backscatter coefficient. Later, Reichardt et al. (2018) implemented a second spectrometer to measure the laser-induced fluorescence of aerosol particles in the middle and upper troposphere. They characterized the fluorescence of mineral dust and BB aerosol and underlined the capabilities of fluorescence measurements to study aerosol-cloud coexistence by enabling the observation of aerosol particles even inside clouds. Saito et al. (2018) studied the spectral fluorescence of atmospheric pollen with lidar and reported a spectral range of 400 to 600 nm for the fluorescence emission, when excited at 355 nm.

Parallel to the developments in spectrometric fluorescence measurements, a further, more easily accessible approach to measure aerosol fluorescence with a single broadband/discrete lidar channel was pursued. Rao et al. (2018) and Li et al. (2019) used Nd:YAG lasers at 266 nm and 355 nm, respectively, and studied the backscattered light in one elastic-backscatter (at the excitation wavelength) and one fluorescence channel. Both instruments were dedicated to aerosol fluorescence measurements and looked only at the boundary layer. Veselovskii et al. (2020, 2021) extended the concept by adding a single broadband fluorescence channel into an existing multi-wavelength lidar system at Lille, France. They also described a retrieval scheme for the computation of the fluorescence backscatter coefficient out of the signal ratio between the fluorescence and the nitrogen channels. Their observations also confirmed the potential of the fluorescence lidar technique to study aerosol particles inside clouds (Veselovskii et al., 2022a). Veselovskii et al. (2022b) showed that fluorescence measurements can improve the aerosol classification with lidar. They proposed, for the first time, a simple classification scheme that combines the linear depolarization ratio with the fluorescence capacity. With this approach, they were able to discriminate between smoke, mineral dust, pollen and urban aerosol, as pollen and smoke showed significantly higher values of fluorescence capacity than urban aerosol and mineral dust. Reichardt et al. (2023) described a procedure for absolute calibration of spectrometric fluorescence measurements and proposed a method to correct for the systematic fluorescence error in water vapor measurements with Raman lidar, which is significant for dry and strongly fluorescent aerosol layers. They also emphasized that the spectrum's shape is closely related to the aerosol type. Veselovskii et al. (2023) presented an approach to measure rough fluorescence spectra with a lidar with five discrete broadband fluorescence channels at Moscow, Russia. They reported advancements in aerosol typing with this approach compared to a single fluorescence channel. Smoke and urban aerosol particles could be discriminated even at high relative humidity and in the presence of hygroscopic growth.

In this work, we explore the observational capabilities of an atmospheric fluorescence lidar utilizing measurements performed in Leipzig, Germany, with an upgraded system since 2022. A detailed description of the new experimental setup is provided in Sec. 2. The analysis of several measurement case studies is presented in Sec. 3. Our findings corroborate the results obtained by previous studies on the capabilities of fluorescence lidars and deepen the discussion in the field of aerosol studies utilizing fluorescence lidar observations. We discuss a unique new capability that is special to this measurement approach. Because it is sensitive to particles only, a fluorescence channel can potentially improve not only the typing but even the detection of aerosol particles. Sec. 3.2.3 provides an in-depth analysis of the reasons for this increased sensitivity of the fluorescence channel to aerosol particles. An exceptional smoke-cirrus interaction case presented in Sec. 3.2.4 highlights the importance of the ability to detect thin aerosol layers in the upper troposphere and lower stratosphere (UTLS) region for the investigation of cirrus cloud

formation. Furthermore, it corroborates and expands the initial work on the detection of fluorescence signals inside ice clouds.

The paper concludes in Sec. 4.

## 95 2 Experimental Setup

### 2.1 Implementation of a fluorescence channel in the MARTHA lidar system

The Multi-wavelength Atmospheric Raman lidar for Temperature, Humidity and Aerosol profiling (MARTHA) is a lab-based

lidar system at the Leibniz Institute for Tropospheric Research (TROPOS) in Leipzig. It emits electromagnetic radiation at


100 three wavelengths (355 nm, 532 nm and 1064 nm) with an overall pulse energy of about 1.2 J at a repetition rate of 30 Hz and collects the backscattered radiation with a large main mirror, which measures 80 cm in diameter. A detailed description of the MARTHA system is given in Mattis et al. (2002), Schmidt et al. (2013) and Jimenez et al. (2019).

To measure the laser-induced fluorescence of atmospheric aerosol, the MARTHA lidar system was upgraded by adding a discrete fluorescence channel into the receiving unit in 2022. To facilitate comparability, the spectral range of the channel was set

in the same wavelength range as in Veselovskii et al. (2020). A 44 nm-wide interference filter from Alluxa centered at 466 nm

105 is used to select a part of the fluorescence spectrum of fluorescing aerosol particles. Because of the similar features, a first comparison of the results obtained in Lille, France, and Leipzig, Germany, is possible.

The backscattered fluorescence intensity depends on the aerosol situation, but in general, it is much weaker than elastic



**Figure 1.** Setup of the far-range receiving unit of the MARTHA system after implementing the fluorescence channel (graphic adapted from Schmidt (2014)). DM: dielectric mirror; BS: beam splitter.

backscatter signals. Therefore, the signal-to-noise ratio in the new channel must be as high as possible. The features that made the MARTHA system suitable to detect fluorescence are its large telescope area and the high-power laser. The second and third harmonic generation setups allowed an increase in the laser energy at 355 nm, sending 6 ns pulses with an energy of 110 about 350 mJ. The new setup of the MARTHA far-range (FR) receiver, including the fluorescence channel, is displayed in Fig. 1. The new detection channel was placed in the branch with the lower wavelengths. Therefore, the first long-pass beam splitter (BS1) was replaced to ensure the complete reflection of the intended fluorescence spectral band. A second beam splitter (BS2) was added. It transmits the shorter wavelengths to the elastic-backscatter and Raman channels related to the UV laser emission 115 at 355 nm and reflects the longer wavelengths towards the fluorescence channel. As the fluorescence signal can be 4–5 orders of magnitude weaker than elastic backscattering (Veselovskii et al., 2020), sufficient suppression of the elastic returns in the new channel was essential to measure fluorescence. The two new beam splitters received customized coatings from Laseroptik GmbH to guarantee high suppression of the elastic-backscatter (and Raman) lines with minimal loss of fluorescence return. Two interference filters were placed in tandem to suppress the elastic components further.

## 2.2 Analytical scheme of the fluorescence backscatter coefficient

120 The lidar system was operated manually and only when no rain was expected. Complete night measurements have been collected since 2022 and were analyzed with a focus on the fluorescing properties of the observed aerosol layers. A second important step is the derivation of new products. The procedure is described as follows: The aerosol fluorescence backscatter coefficient was obtained by forming the ratio of the fluorescence ( $P_F$ ) to the nitrogen Raman ( $P_R$ ) signal, similar to Veselovskii et al. (2020). Both signals can be described in terms of the lidar equation:

$$125 \quad P_F = \beta_F T_L T_F C_F \quad (1)$$

$$P_R = \beta_R T_L T_R C_R. \quad (2)$$

$T_L$  is the atmospheric transmission at the emitted laser wavelength.  $T_R$  and  $T_F$  denote the atmospheric transmission at the Raman and fluorescence wavelength ranges, respectively, and  $C_R$  and  $C_F$  represent the corresponding lidar calibration constants. By dividing Eq. (1) by Eq. (2), the following expression for the aerosol fluorescence backscatter coefficient  $\beta_F$  can 130 be derived:

$$\beta_F = \frac{P_F}{P_R} \frac{T_R}{T_F} \frac{C_R}{C_F} \beta_R. \quad (3)$$

The Raman backscattering  $\beta_R$  is computed using the following expression in terms of the Rayleigh molecular backscatter coefficient ( $\beta_{\text{mol}}$ ):

$$\beta_R = D_R N_{N_2} = 0.78 D_R N_{\text{mol}} = 0.78 \frac{D_R}{D_{\text{mol}}} \beta_{\text{mol}}, \quad (4)$$

135 with  $N_{N_2}$  and  $N_{\text{mol}}$  the number density of nitrogen and air molecules, respectively.  $D_R/D_{\text{mol}}$  accounts for the Raman to Rayleigh backscatter differential-cross-section ratio. These cross sections were determined theoretically using Eqs. (20) and (14) in Adam (2009), resulting in theoretical values of  $D_R^* = 2.7344 \times 10^{-34} \text{ m}^2 \text{ sr}^{-1}$  and  $D_{\text{mol}} = 3.10875 \times 10^{-31} \text{ m}^2 \text{ sr}^{-1}$ .

### 2.3 Technical considerations for the calibration of the fluorescence channel

To derive the particle fluorescence backscatter coefficient from Eq. (3), the traditional method, using a particle-free reference height, cannot be applied, due to the unknown fluorescence response of the background aerosol. Instead, a characterization of the channel's system efficiencies is needed. The contribution of each component in the respective detection path was carefully determined to infer the overall efficiencies and build the lidar-constant ratio  $C_R/C_F$ .

The first point to consider is the bandwidth of the interference filters. For the 387 nm nitrogen Raman channel, with a bandwidth of 2.7 nm, only 95 % of the theoretical Raman cross section can reach the detector, reducing the actual cross section at the detector to  $D_R = 0.95 \times D_R^* = 2.59768 \times 10^{-34} \text{ m}^2 \text{ sr}^{-1}$ . This value is then used in Eq. (4) together with the molecular backscatter computed based on the temperature and pressure profiles provided by the Global Data Assimilation System (GDAS) (Rodell et al., 2004) to derive the Raman backscatter coefficient. As the laser power, pulse length, and telescope area are the same for both detection channels, the lidar-constant ratio  $C_R/C_F$  simplifies to the ratio of the channel efficiencies. This ratio comprises the transmittances or reflectances of the optical elements (such as beam splitters, mirrors, interference filters and neutral-density filters) and the detection efficiencies of the detectors. The transmittances and reflectances of the optical components are collected in Tab. 1. As the neutral-density (ND) filters are eventually changed depending on the atmospheric and system conditions, only one exemplary set of ND filters, which is representative of most of the cases studied in this manuscript, was chosen for Tab. 1. When determining the ND filter transmission, the spectral dependence provided by the manufacturer (Thorlabs) was considered. The detection efficiencies of the photomultiplier tubes (PMTs) are split into electrical gain and detector efficiency. The ratio of the electrical gains was obtained by swapping the detectors in the nitrogen Raman and fluorescence channels and building the ratio of the mean signals measured by both detectors for each channel. This test yielded a PMT gain ratio ( $\eta_{\text{gain},R}/\eta_{\text{gain},F}$ ) of 1.4155, indicating a higher gain of the nitrogen Raman channel's PMT. As for the detector surface, the so-called quantum efficiency accounts for the amount of photoelectrons generated by the cathode divided by the number of incident photons. This efficiency depends on the photon wavelength (Wright, 2017). The spectrally resolved quantum-efficiency data provided by Hamamatsu were considered to assess the PMT type used in the MARTHA system. The maximum efficiency of the detectors is about 35 %, and the values at the wavelength ranges of the two lidar channels were determined by interpolation from the provided data points and averaging over the filter width of the interference filter in the fluorescence channel. This resulted in values of  $\eta_{\text{qe},R} = 34.66\%$  and  $\eta_{\text{qe},F} = 25.13\%$  for the quantum efficiencies of the used PMT type in the nitrogen Raman (386–388 nm) and fluorescence channel (444–488 nm), respectively, and we finally obtained a ratio of  $\eta_{\text{qe},R}/\eta_{\text{qe},F} = 1.379$ .

After these considerations, the ratio of the lidar calibration constants can be calculated from the efficiency ratios of the optical elements, detector gain and spectral response as follows:

$$\frac{C_R}{C_F} = \frac{\mathcal{R}_2 \mathcal{T}_1 \mathcal{T}_2 \mathcal{T}_3 \mathcal{T}_{\text{ND}}}{\mathcal{R}_1 \mathcal{R}_3 \mathcal{T}_4^2} \frac{\eta_{\text{gain},R}}{\eta_{\text{gain},F}} \frac{\eta_{\text{qe},R}}{\eta_{\text{qe},F}} \quad (5)$$

For the set of ND filters considered in Tab. 1 (OD 1.3 in the nitrogen Raman and no ND filters in the fluorescence channel) it results in a value of  $C_R/C_F \approx 0.0303$ .

The remaining unknown in Eq. (3) is the ratio  $\mathcal{T}_R/\mathcal{T}_F$  of the atmospheric transmissions (ground to target) at the Raman and

**Table 1.** Transmittances  $\mathcal{T}$  or reflectances  $\mathcal{R}$  of all optical elements in the 387 nm and 466 nm FR channels. PMT quantum efficiencies, PMT gain ratio and overall ratio of the lidar calibration constants in both channels.

FR channel	Nitrogen Raman (387 nm)	Fluorescence (466 nm)
Common beam splitter (BS2)	$\mathcal{T}_1 = 97.1\%$	$\mathcal{R}_1 = 98\%$
Further (unique) optical elements	beam splitter 407 nm: $\mathcal{T}_2 = 94.5\%$ beam splitter 387 nm: $\mathcal{R}_2 = 95\%$	dielectric mirror (DM) $\mathcal{R}_3 = 99.75\%$
Interference filters	$\mathcal{T}_3 = 70\%$	$\mathcal{T}_4 = 92.5\%$ ( $\mathcal{T}_4^2$ because of 2 filters)
Neutral-density filters (example)	OD = 1.3 $\mathcal{T}_{\text{ND}} \approx 0.0213$	no neutral-density filters $\mathcal{T}_{\text{ND}} = 1$
Product	$\prod_{i=1}^N \mathcal{T}_i \prod_{j=1}^M \mathcal{R}_j$	$\mathcal{R}_2 \mathcal{T}_1 \mathcal{T}_2 \mathcal{T}_3 \mathcal{T}_{\text{ND}} = 0.01299$ $\mathcal{R}_1 \mathcal{R}_3 \mathcal{T}_4^2 = 0.8364$
PMT quantum efficiency	34.66 %	25.13 %
PMT gain ratio	$\eta_{\text{gain,R}} / \eta_{\text{gain,F}} = 1.4155$	
Overall ratio of the lidar calibration constants	$C_{\text{R}} / C_{\text{F}} \approx 0.0303$	

fluorescence wavelengths, respectively. The molecular part ( $\mathbf{T}_{\text{R}} / \mathbf{T}_{\text{F}}|_{\text{mol}}$ ) is calculated straightforwardly from the extinction and backscatter coefficients; the aerosol contribution to the transmission ratio ( $\mathbf{T}_{\text{R}} / \mathbf{T}_{\text{F}}|_{\text{par}}$ ) requires previous knowledge about the aerosol backscatter coefficient. For the profile-based analysis, the aerosol optical properties are determined with the traditional Raman technique (Ansmann et al., 1990, 1992). The particle backscatter coefficient at high temporal resolution was obtained via a constant-based approach, in which a previous profile-based retrieval is needed to calculate the lidar constants, which are then used to compute high-resolution products out of the elastic-backscatter and Raman signals (Baars et al., 2017). In general, the particle atmospheric transmission differs little at the two wavelengths, making the effect on the fluorescence backscatter coefficient small, partially because only about 80 nm separate the central wavelengths of the Raman and fluorescence channels. For the cases with low and medium aerosol loads (see Sec. 3.2), the error in case of non-consideration of the differential transmission was in the range of 2–6 %. In case of an unusually high aerosol optical depth, like on the 4 July 2023 (see Sec. 3.1), the error was in the order of 10 %. Thus, the differential particle transmission at the two wavelengths was considered to guarantee the quality of the fluorescence backscatter coefficient, even above strongly backscattering aerosol layers. But still, the assumption of an appropriate Ångström exponent is necessary, which imposes an uncertainty of  $\pm 1\text{--}7\%$  on the

185 determined  $T_R/T_F|_{\text{par}}$ , depending on the optical thickness of the present aerosol layers.

The data set acquired in Leipzig was then analyzed in a semi-automatic manner, setting the calibration constants and the reference height (particle-free) manually for each case. The fluorescence capacity  $G_F$ ,

$$G_F = \frac{\beta_F}{\beta_{532}}, \quad (6)$$

190 was calculated as the ratio of the fluorescence backscatter coefficient ( $\beta_F$ ) to the elastic particle backscatter coefficient at 532 nm ( $\beta_{532}$ ). To encourage comparability with different setups (fluorescence lidars which do not deploy the second-harmonic generation wavelength and/or use a spectrometric approach or broadband fluorescence channels with different interference filter bandwidth), the general and season-mean values are also provided as spectral fluorescence capacity with respect to the third harmonic generation wavelength 355 nm ( $G_F^{355}$ ),

$$G_F^{355} = \frac{\beta_F}{\beta_{355} d_{\text{IF}}}, \quad (7)$$


195 where  $d_{\text{IF}} = 44 \text{ nm}$  is the bandwidth of the interference filter in the fluorescence channel. Furthermore, data from the collocated portable Raman lidar Polly<sup>XT</sup> (Engelmann et al., 2016) at TROPOS were used to provide quality-assured depolarization profiles.

### 3 Observational results

200 Due to the broad bandwidth of the fluorescence channel and the low intensity of the fluorescence signal, measurements were only possible during the night. At daytime, scattered solar radiation would cause too much noise in the fluorescence channel. As the MARTHA system is operated manually, the number of measurements remains limited. Since August 2022, about 50 measurements have been performed, providing more than 250 hours of atmospheric fluorescence observations. Typical atmospheric values of the fluorescence backscatter coefficient and fluorescence capacity, that were obtained at Leipzig during the time period from August 2022 to October 2023, are presented in the next paragraph.

205 In general,  $\beta_F$  ranged between  $1 \times 10^{-5} \text{ Mm}^{-1} \text{ sr}^{-1}$  for background aerosol and more than  $1 \times 10^{-3} \text{ Mm}^{-1} \text{ sr}^{-1}$  for optically extraordinarily thick wildfire smoke layers. Correspondingly,  $G_F$  ( $G_F^{355}$ ) varied from  $\sim 10^{-6}$  ( $\sim 10^{-8} \text{ nm}^{-1}$ ) for clouds and  $1 \times 10^{-5}$  ( $1 \times 10^{-7} \text{ nm}^{-1}$ ) for background aerosol to  $1.3 \times 10^{-3}$  ( $1.3 \times 10^{-5} \text{ nm}^{-1}$ ), whereas most of the measurement points were in the range of  $5 \times 10^{-5}$  to  $7 \times 10^{-4}$  ( $6 \times 10^{-7} \text{ nm}^{-1}$  to  $9 \times 10^{-6} \text{ nm}^{-1}$ ). I.e., the fluorescence backscatter coefficient was about four orders of magnitude lower than the elastic ones, which agrees with the findings by Veselovskii et al. (2020).

210 In the following, four interesting case studies are presented in several subsections. In Sec. 3.1, the fluorescence properties of wildfire smoke are discussed by analyzing an optically and geometrically thick smoke layer on 4 July 2023. In Sec. 3.2, we first demonstrate the ability of the fluorescence lidar technique to detect optically thin aerosol layers by presenting two case studies (Sec. 3.2.1 and Sec. 3.2.2). Subsequently, we discuss the reasons for the increased sensitivity of the fluorescence channel to aerosol particles in Sec. 3.2.3. Finally, we underline the importance of this new capability by presenting a striking smoke-cirrus interaction case study in Sec. 3.2.4.



**Figure 2.** Height-time distributions of (a) particle backscatter coefficient at 532 nm and (b) fluorescence backscatter coefficient ( $\beta_F$ ) measured with the MARTHA system and (c) particle depolarization ratio at 532 nm from Polly<sup>XT</sup> on 4–5 July 2023. Vertical profiles of (d) the elastic backscatter coefficients and (e)  $\beta_F$  together with the fluorescence capacity ( $G_F$ ) from 21:00 to 22:00 UTC on 4 July 2023.

### 3.1 Fluorescence of wildfire smoke – 4 July 2023

In the spring and summer of 2023, huge wildfires raged across Canada, with unusual intensity in the provinces of Alberta and British Columbia. With the prevailing westerly winds, large amounts of biomass-burning aerosol were transported towards 220 Europe. As a result, we frequently observed wildfire smoke layers over Leipzig from mid-May to mid-July 2023.

As a first example, the fluorescence of an optically thick plume of wildfire smoke on 4–5 July 2023 shall be characterized. Figure 2 displays the height-time distributions of the particle backscatter coefficient at 532 nm, the fluorescence backscatter coefficient and the particle depolarization ratio at 532 nm for this night. Figure 2(a) shows a highly polluted troposphere, with an overall aerosol optical depth (AOD) of around 0.8 at 532 nm. This agrees well with data from the Aerosol Robotic Network (AERONET), where AOD values of around 0.75–0.8 were retrieved at 500 nm at 17:00 UTC on 4 July 2023. An optically thick aerosol layer extended from 3.4 to 5.8 km height. To determine its optical properties, a 1-hour time period was considered 225 for temporal averaging. Figure 2(d) shows the vertical profiles of the fluorescence and elastic backscatter coefficients, together with the fluorescence capacity averaged over the time period from 21:00 to 22:00 UTC. At the optically thickest part,  $\beta_{532}$  reached values of up to  $5 \text{ Mm}^{-1} \text{ sr}^{-1}$ . The 532 nm AOD of the whole layer amounted to around 0.48.

230 The optical properties of this aerosol layer, ranging from 3.4 to 5.8 km height, were then used to determine the aerosol type. The fact that the lidar ratio at 532 nm (60 sr) was significantly larger than the one at 355 nm (38 sr) and the high backscatter-

related Ångström exponent (1.66) are characteristic for aged BB aerosol (Müller et al., 2005; Ansmann et al., 2009; Ohneiser et al., 2021, 2022; Hu et al., 2022; Janicka et al., 2023). Furthermore, these retrieved lidar ratio values are in the same range as reported for aged wildfire smoke in previous studies (e.g., Murayama et al., 2004; Ansmann et al., 2009; Haarig et al., 2018; 235 Hu et al., 2019). The low particle depolarization ratio ( $\delta_{532} \leq 0.07$ , cf. Fig. 2(c)) in the layer from 3.4 to 5.8 km height points to a spherical shape of the particles, which is also typical for aged wildfire smoke in the middle free troposphere (Haarig et al., 2018). Thus, it can be concluded that this tropospheric aerosol layer consisted of aged BB aerosol particles.

Figs. 2(b) and (d) show a very high fluorescence backscatter coefficient ( $\beta_F \approx 2.75 \times 10^{-3} \text{ Mm}^{-1} \text{ sr}^{-1}$ ) for this smoke layer and a corresponding fluorescence capacity of  $G_F \approx 7.8 \times 10^{-4}$ . In other words, smoke shows very high values of fluorescence 240 capacity compared to other particle types and can thus be clearly identified through this new quantity. These values witnessed in our observations agree with the findings by Veselovskii et al. (2020).


Considering the entire 2023 wildfire season, the fluorescence capacity  $G_F$  (spectral fluorescence capacity  $G_F^{355}$ ) of smoke varied from  $1 \times 10^{-4}$  to  $13 \times 10^{-4}$  ( $1.5 \times 10^{-6} \text{ nm}^{-1}$  to  $13 \times 10^{-6} \text{ nm}^{-1}$ ). Thereby, values of  $2-7 \times 10^{-4}$  ( $2-9 \times 10^{-6} \text{ nm}^{-1}$ ) were observed most frequently, which agrees with the results of Hu et al. (2022) and Veselovskii et al. (2022a), who reported 245 values of  $G_F$  in the range of  $1-4.5 \times 10^{-4}$  for their observations at Lille, France. The observed values of  $G_F^{355}$  are also in a range similar to the spectral fluorescence capacities of BB aerosol that were reported by Reichardt et al. (2018), although for a broader wavelength range (455–530 nm). The particle depolarization ratio at 532 nm was low (below 0.07) for most (95 %) of the investigated smoke layers.

### 3.2 Detection of optically thin aerosol layers with the fluorescence channel

250 Besides its relevance for aerosol type identification, our results suggest an additional capability of a fluorescence lidar: to detect optically thin aerosol layers. In several measurements with the new fluorescence channel, an enhanced fluorescence signal revealed the presence of aerosol layers that went unnoticed when employing only the elastic-backscatter detection channels. Three exemplary measurement cases are discussed in the following sections.

#### 3.2.1 Hidden smoke layers – 21 September 2022


255 Figure 3 shows the height-time distributions of the range-corrected lidar signal at 1064 nm (a), the fluorescence backscatter coefficient (b) and the fluorescence capacity (c). According to the elastic backscatter signal in Fig. 3(a), the upper troposphere appears to be rather aerosol-free. Only the polluted boundary layer and some thin layers up to 4 km height indicated aerosol presence, and a thin cloud was visible at around 4 km height from 21:00 to 22:00 UTC. However, an enhanced fluorescence backscatter coefficient in Fig. 3(b) reveals several other fluorescing aerosol structures throughout the middle and upper troposphere (at around 5, 6.5, 9 and 9.75 km height). This already illustrates that with measurements of aerosol fluorescence, thin aerosol layers can be identified more easily from lidar quicklooks and therefore chosen for detailed analysis. Looking at the vertical profiles, this measurement case appears even more impressive. Figure 3(d) displays the time-averaged vertical profiles of the fluorescence and elastic backscatter coefficients together with the fluorescence capacity. The profiles were averaged over 260 the 2-hour time period from 19:04 to 21:04 UTC to exclude the cloud, which was present at around 4 km height from that



**Figure 3.** Height-time distributions of (a) range-corrected lidar signal at 1064 nm, (b) fluorescence backscatter coefficient ( $\beta_F$ ) and (c) fluorescence capacity measured with the MARTHA system on 21 September 2022. Vertical profiles of (d) the elastic backscatter coefficients and (e)  $\beta_F$  together with the fluorescence capacity ( $G_F$ ) from 19:04 to 21:04 UTC on 21 September 2022.

265 point onwards. The lowest (3.3 km) and most fluorescent ( $\beta_F \approx 2.5 \times 10^{-5} \text{ Mm}^{-1} \text{ sr}^{-1}$ ) layer above the boundary layer still shows clearly enhanced elastic backscatter coefficients at all three wavelengths. In the mid-level layers at around 5 and 6.5 km height, the 532 nm and 1064 nm backscatter coefficients are only slightly enhanced compared to the background. For the layer at 6.5 km height, their corresponding maxima are at higher altitudes than the distinct maximum in the fluorescence backscatter coefficient. The 355 nm backscatter coefficient even fails to resolve the aerosol layer at 6.5 km. However, all elastic-backscatter 270 detection channels reach their limits with the two high layers at 9 and 9.75 km altitude. While the 355 nm backscatter coefficient is completely noisy in this altitude range,  $\beta_{532}$  and  $\beta_{1064}$  do show maxima in the altitude range of the increased  $\beta_F$ . But these maxima are difficult to distinguish from the background, which is likewise already quite noisy. Thus, it is unlikely that these two higher layers would have been detected as aerosol layers without the additional fluorescence information, especially because the particle depolarization ratio (not shown) is also quite low, around 2%.

275 The overall AOD of this measurement case was around 0.13 at 532 nm, which is consistent with AERONET data showing 0.1 at 500 nm, whereas the majority of the aerosol was found in the boundary layer ( $\text{AOD} \approx 0.1$ ). The smoke layers above the boundary layer only added up to an AOD of around 0.03. The two thinnest layers at around 9 and 9.75 km height even had an AOD of only 0.002 each at 532 nm.



**Figure 4.** Height-time distributions of (a) range-corrected lidar signal at 532 nm and (b) fluorescence backscatter coefficient ( $\beta_F$ ) measured with the MARTHA system and (c) particle depolarization ratio at 532 nm from Polly<sup>XT</sup> on 15–16 May 2023. (d) Vertical profiles of  $\beta_F$  and the elastic backscatter coefficients together with the fluorescence capacity ( $G_F$ ) from 01:15 to 02:15 UTC on 16 May 2023.

### 3.2.2 A thin smoke layer in the UTLS – 15 May 2023

280 Another example of such "unnoticeable" layers is the night of 15–16 May 2023. Figure 4(a–c) displays the height-time distributions of the range-corrected lidar signal at 532 nm, the fluorescence backscatter coefficient and the particle depolarization ratio at 532 nm. The vertical profiles of the backscatter coefficients together with the fluorescence capacity for the period of 01:15 – 02:15 UTC are shown in Fig. 4(d). This measurement case is characterized by pronounced fluorescent aerosol layers (532 nm AOD  $\approx 0.05$ ), ranging from 4 to 6.7 km height. The high fluorescence capacity and lidar ratio values allow us to identify the 285 aerosol particles present as wildfire smoke. For further discussion, we consider a height-constant and rather homogeneous layer, ranging from 4.6 to 6.1 km (layer 1). Layer 1 shows a mean fluorescence capacity of around  $5.6 \times 10^{-4}$  and lidar ratios of around 40sr at 355 nm and 70sr at 532 nm. The particle depolarization ratio at 532 nm is low (around 1.7 %), indicating a well-advanced aging process of the smoke particles. At around 11 km height, the range-corrected signal at 532 nm in Fig. 4(a) shows another highly fluorescent smoke layer ( $G_F \approx 6.5 \times 10^{-4}$ ), which we will refer to as layer 2 in the following. This

290 higher value of the fluorescence capacity indicates a more efficient fluorescence emission in layer 2 than in layer 1. The reason for this remains unclear. On the one hand, this could be purer smoke, while layer 1 could also contain a small proportion of another less fluorescent aerosol type. On the other hand, the BB aerosol in both layers could differ in chemical composition and optical properties due to different fire sources and transport mechanisms. Backward trajectory analyses point generally to the same source region (the northern part of the North American continent) for both altitudes. However, this does not exclude 295 the possibility of slightly different fire sources. The optical properties of the smoke particles differ slightly between the two layers. The lidar ratios in layer 2 (55sr at 355 nm and 75sr at 532 nm) are slightly higher than the lidar ratios in layer 1. Furthermore, in layer 2, the particle depolarization ratio was slightly enhanced ( $\delta_{532} \approx 6.5\%$ ) compared to layer 1, indicating 300 a more irregular shape of the smoke particles in layer 2 (although in general terms this is still almost spherical). The particle size seems to play a minor role for depolarization in this case, as the backscatter-related Angström exponent for the 355 and 532 nm wavelength pair (1.75 for layer 1 and 1.55 for layer 2) indicates only slightly larger particles in layer 2.

All in all, these differences in aerosol optical properties suggest different fire sources and/or transport mechanisms for both smoke layers. The observed difference in fluorescence capacity is therefore expectable, but its cause remains unclear.


Furthermore, Fig. 4(b) reveals another aerosol layer (layer 3) with enhanced fluorescence at around 11.7 km, directly above 305 layer 2. The complete structure of layer 3 remains unnoticed in the range-corrected signal in Fig. 4(a). This impression is confirmed by the vertical profiles in Fig. 4(d). The thin layer 3 cannot be distinguished from the background noise in  $\beta_{355}$ .  $\beta_{532}$  and  $\beta_{1064}$  exhibit a slight increase, although this increase is only very weakly pronounced at 1064 nm. So, only the 532 nm backscatter coefficient shows a clear peak for this layer 3. At a closer look, also the time-height distribution of the particle depolarization ratio at 532 nm in Fig. 4(c) indicates layer 3 by slightly increased values at this altitude. But again, it would have been hard to recognize this layer from the elastic backscattering products alone, without having a clearer picture of the aerosol 310 situation from the fluorescence channel. This underlines the potential of the fluorescence lidar technique beyond aerosol characterization. Fluorescence backscatter can be used for the detection of aerosol layers in scenarios where concentrations are below the lower detection limit of the elastic-backscatter channels.

### 3.2.3 On the capabilities of a dedicated (aerosol) fluorescence channel

The measurement cases presented demonstrate the advantages of adding fluorescence observations to the analysis. The following 315 paragraph discusses the enhanced capabilities of a fluorescence backscatter channel compared to the classical elastic backscatter channel.

The three elastic-backscatter channels rely on the principle of elastic backscattering of the emitted laser radiation, which occurs at both air molecules and aerosol particles. Because of the strong decrease of air and aerosol density, the scattering intensity 320 strongly decreases with height. In addition, the backscatter detected by the lidar further decreases with height due to the solid angle of the telescope and atmospheric extinction. As a result, the lidar-signal detection needs to cover a wide dynamic range. Elastic signals are usually attenuated in the detection unit to keep them in a manageable range for a single channel. This maximizes the vertical coverage, but at the expense of sensitivity to aerosol particle changes.

A fluorescence channel is dedicated to aerosol particles only. Thus, it can help to increase the sensitivity to aerosol particles by



**Figure 5.** Vertical profiles of (a) particle (magenta), molecular (blue) and total (green) backscatter coefficients at 532 nm and fluorescence backscatter coefficient (black) and (b) background-corrected signals at 532 nm (green) and 466 nm (black) on 16 May 2023 for the time period from 01:15 to 02:15 UTC.

eliminating the molecular component. Furthermore, the fluorescence return does not only scale to the number of particles but also to its cross-section, which is directly related to the fluorescence capacity of the aerosol particles. I.e., a smoke layer will contrast more to the background than a dust layer because of the higher ability of smoke particles to fluoresce. This feature enhances the capabilities of such a channel to detect smoke particles in the atmosphere.

Figure 5(a) illustrates this context by showing the molecular, particle and total (molecular + particle) backscatter coefficient at 532 nm for a measurement on 16 May 2023. The contrast between the smoke layer (4–7 km) and the background aerosol (8–10 km) in the fluorescence backscatter is significantly more pronounced than in the elastic particle backscatter coefficient. The enhanced detection ability in the case of fluorescing aerosol particles becomes evident when comparing the observed lidar signals (background corrected), as depicted in Fig. 5(b) for this measurement case. Especially the aerosol layers at around 11 km stand out much clearer from the background in the fluorescence signal than in the elastic signal.

### 3.2.4 Atmospheric implication: Smoke-cirrus interaction – 29 May 2023

Now, after discussing the possibility of detecting such thin aerosol layers, the question of their relevance in atmospheric research arises. Because of their low optical thicknesses (of typically  $\leq 0.01$ ), such aerosol layers might not have a relevant

radiative effect, but they may impact cloud formation, e.g., by serving as INPs. In both cases presented above (Sec. 3.2.1 and 340 3.2.2), the measurements of the fluorescence backscatter coefficient revealed thin wildfire smoke layers at rather high altitudes around the tropopause. This altitude range, also referred to as UTLS region, is a common site for the formation of cirrus clouds. However, the ability and relevance of smoke particles to act as INPs is an open question in the literature. Although a few available 345 observations showed enhanced immersion-mode INP concentrations inside of BB aerosol plumes (Barry et al., 2021; McCluskey et al., 2014), wildfire smoke is considered to be a rather inefficient INP at temperatures above -30 °C compared to other aerosol types such as dust (e.g., Barry et al., 2021; Knopf et al., 2018). Thus, BB aerosol is, in general, not considered a relevant INP source in mixed-phase cloud processes. Likewise, in situ assessments have suggested that BB aerosol particles 350 rarely freeze to form cirrus clouds (Froyd et al., 2009, 2010). However, the authors could not exclude the INP ability of BB particles due to temperature limitations in their experimental setup. Recent lidar-based studies discussed the potential of smoke particles to promote freezing via deposition and provided evidence of BB aerosol acting as the main INP source in cirrus clouds observed at Limassol, Cyprus (Mamouri et al., 2023) and in the Arctic (Ansmann et al., 2024a, b). Simulations considering gravity waves further explain how heterogeneous freezing overtakes the main role, consuming quickly the water vapor and reducing supersaturation, hampering in this way homogeneous freezing (Ansmann et al., 2024a). Thus, investigations of possible 355 smoke-cirrus interactions in the UTLS region are an important topic for future studies.


Also, several of our measurement cases during the 2023 wildfire season showed cirrus clouds directly below thin smoke layers. One example (29-30 May 2023) is displayed in Fig. 6 and will be discussed in the following.

The range-corrected lidar signal at 532 nm in Fig. 6(a) shows cirrus clouds, that extended from 7 to 11.5 km at the beginning 360 of the measurement. Above, enhanced values of the fluorescence backscatter coefficient (see Fig. 6(b)) reveal the presence of a smoke layer at 10.5 to 12 km height, that was not visible in the elastic-backscatter lidar signal over large parts of the observation period. Only at the end of this measurement (around 02:00 UTC), when the clouds became thinner and more scattered, the aerosol layer could be anticipated vaguely from weak signatures in the range-corrected signal (cf. Fig. 6(a)).

The time-height plot of the fluorescence capacity in Fig. 6(c) reveals another feature of fluorescence backscattering that is 365 exclusive to aerosol particles. Pure water does not fluoresce, and, as our measurements from 2022–2023 showed, hydrometeors such as cloud droplets and ice crystals, exhibit the lowest values of fluorescence capacity. In combination, elastic-backscatter and fluorescence channels can unambiguously differentiate aerosol particles and hydrometeors that coexist within the same air volume. This feature has also been pointed out in previous studies (Reichardt et al., 2018; Veselovskii et al., 2022a) and opens a new door into aerosol and cloud detection.

370 Due to this characteristic, the fluorescence capacity clearly shows the positions of the aerosol and cloud layers relative to each other in one plot. Remarkably, the upper boundary of the cirrus clouds coincides with the lower boundary of the fluorescing smoke layer for large parts of the observation period. The elastic-backscatter signal in Fig. 6(a) clearly shows pronounced virga structures (i.e., stripes of falling ice crystals). Such an arrangement has already been reported in the literature for smoke layers and cirrus clouds observed over the eastern Mediterranean and in the Arctic (Mamouri et al., 2023; Ansmann et al., 2024b).

375 At the beginning of the measurement in the night of 29 May, parts of the cirrus clouds were even embedded in the smoke layer. Furthermore, the smoke layer slowly rose in altitude towards the end of the measurement. At the same time, the cloud top rose




**Figure 6.** Height-time distributions of (a) range-corrected lidar signal at 532 nm, (b) fluorescence backscatter coefficient ( $\beta_F$ ) and (c) fluorescence capacity measured with the MARTHA system on 29 - 30 May 2023. (d) Vertical profiles of  $\beta_F$  and the elastic backscatter coefficient at 532 nm together with the fluorescence capacity ( $G_F$ ) from 21:00 to 22:00 UTC on 29 May 2023. (e) Vertical profiles of temperature and relative humidity over ice from a radiosounding at Lindenberg at 18 UTC on 29 May 2023.

first, and later, the clouds even became scattered and the ice nucleation and cloud formation seemed to stop. All these facts indicate that the smoke particles may have triggered the cloud formation by serving as INPs.

This hypothesis is further supported by Figs. 6(d-e), which display the vertical profiles of the elastic and fluorescence backscatter coefficients, together with the fluorescence capacity from 21:00 to 22:00 UTC and the temperature and relative humidity over ice from a radiosonde launched at the nearest station Lindenberg (150 km away) at 18 UTC on 29 May 2023. From the high elastic backscatter coefficients, two nucleation sections can be identified. There is a lower part with ice crystals falling from about 10.5 km (N1 in Fig. 6(d)), and an upper part where ice crystals start falling from 11.75 km (N2). These falling ranges somehow coincide with the aerosol layers observed with the fluorescence channel. Cloud-top temperatures were also estimated from the radiosonde at Lindenberg and ranged from -58 to -51 °C, a temperature range in which deposition ice nucleation is particularly efficient (Ansmann et al., 2024b). The arrangement of the cloud and the aerosol layer in this case indicates that the ice nucleation happened at the cloud top, from where the freshly formed ice crystals were falling down, thus producing the aforementioned falling stripes.

At around 12 km height, the high fluorescence capacity (up to  $G_F = 7.5 \times 10^{-4}$  at the maximum) indicates the smoke layer. However, the fluorescence backscatter coefficient shows enhanced values over a wider altitude range, even down to 9 km alti-



**Figure 7.** Height-time distributions of the particle backscatter coefficient at 532 nm in the night of 29 - 30 May 2023. Height-time bins with a high fluorescence backscatter coefficient ( $> 2.5 \times 10^{-5} \text{ Mm}^{-1} \text{ sr}^{-1}$ ) are colored in gray. The top of the ice cloud layer was marked in black.

tude, which supports the hypothesis that wildfire smoke particles triggered the ice cloud formation. An interesting feature in

Fig. 6(d) is the reduction of the fluorescence backscatter at the cloud top of the upper cirrus part ( $\beta_F \approx 2.9 \times 10^{-5} \text{ Mm}^{-1} \text{ sr}^{-1}$ ) compared to the higher values above this upper cloud layer ( $\beta_F \approx 5 \times 10^{-5} \text{ Mm}^{-1} \text{ sr}^{-1}$ ) and at the top of the lower part of the cirrus cloud at around 10.9 km height ( $\beta_F \approx 6.4 \times 10^{-5} \text{ Mm}^{-1} \text{ sr}^{-1}$ ). A possible reason for this reduction could be fluorescence quenching (Lakowicz, 2006) by the ice crystals within the upper cloud layer. Water is known to act as a fluorescence quencher for organic fluorophores (e.g., Stryer, 1966; Dobretsov et al., 2014). However, a more plausible explanation would be that some smoke particles acted as INPs and subsequent falling of the ice crystals (indicated by the virga) reduced the number of smoke particles over time. This hypothesis is supported by the fact that the altitude of the reduced  $\beta_F$  coincides with the upper nucleation zone N2.

390 Another reduction of the fluorescence backscatter coefficient was observed inside the ice virga. In the middle part of the cloud, between 8 and 10 km,  $\beta_F$  ranged at very low values of about  $1 \times 10^{-6}$  to  $1 \times 10^{-5} \text{ Mm}^{-1} \text{ sr}^{-1}$ . Aerosol scavenging arises as a possible explanation. I.e., the falling ice crystals collected most of the aerosol particles (impaction), reducing the aerosol load in the cloud layer. In this case, one would expect an accumulation of smoke particles at or directly below the cloud base. Indeed, near the cloud base at around 7 km height, the fluorescence backscatter increases again up to  $1.4 \times 10^{-5} \text{ Mm}^{-1} \text{ sr}^{-1}$ . A

400 similar situation with a smoke layer directly below the cloud base is visible around 8 km from 01:15 to 02:00 UTC in Fig. 6(c), further supporting the scavenging hypothesis. However, both reductions discussed here could also be due to different aerosol loads and characteristics at the different altitudes. The situation is, in any case, complex, and further investigations of similar cases are needed to characterize aerosol particles inside clouds by fluorescence observations.

405 In summary, our measurement results suggest two possible interaction pathways between the observed smoke layer and the cirrus clouds: fluorescence quenching and heterogeneous ice nucleation combined with aerosol scavenging. For further illustration, Fig. 7 shows the elastic and fluorescence backscatter coefficients together in one plot. The height-time bins with pronounced aerosol fluorescence (in gray color) along with the elastic backscattering at 532 nm clearly show a major aerosol-cloud interplay. Just before 01:00 UTC, an interesting situation arose. The smoke particles were deeply embedded in the cloud, 410 exhibiting two layers: one around 10 km and one between 8 and 9 km, accompanied by a significant increase in the elastic backscatter coefficient.

A further potential application of fluorescence lidar is to provide INP information in such cases with low but relevant aerosol presence in the cloud surroundings, especially at the cloud top. A conversion from the unambiguous fluorescence backscatter 415 coefficient to an INP number concentration ( $N_{\text{INP}}$ ) is desirable. An approach applying conversion factors, which link the fluorescence backscatter coefficient with the previously inverted microphysical properties of the fluorescing aerosol particles from multi-wavelength lidar data, was suggested by Veselovskii et al. (2022a). In the case of a low aerosol load or inside a cloud layer, the resulting mean conversion factors, together with the fluorescence backscatter coefficient, can then be used to derive the aerosol surface area concentration, which is needed as input to the INP parameterization (Veselovskii et al., 2022a). An 420 alternative approach would be to determine  $N_{\text{INP}}$  directly from ice crystal number information provided by lidar-radar synergy and find a conversion factor between  $\beta_F$  and  $N_{\text{INP}}$ . Such a factor  $\sigma_F$  would be in the form of  $\sigma_F = N_{\text{INP}}/\beta_F$  and could be used for cirrus cloud scenes with comparable temperature and humidity. Preliminary assessments of INP concentrations via the POLIPHON method (Ansmann et al., 2012; Mamouri and Ansmann, 2014) and ice crystal number concentrations from lidar-radar synergy (Bühl et al., 2019) suggest a conversion factor in the range of  $3\text{--}8 \times 10^4 \text{ Mm sr L}^{-1}$ .  
425 A reliable conversion to link the fluorescence backscatter coefficient to ice nuclei concentrations would be beneficial to investigate aerosol-cloud interactions, especially in those situations with low aerosol amounts. Further aerosol-cloud cases will be investigated in the future to evaluate this potential application of fluorescence backscatter information specifically.

## 4 Conclusions

In this study, we presented the newly implemented fluorescence channel in the lidar system MARTHA, located at TROPOS, 430 Leipzig, Germany. Some of the first measurements performed with the upgraded system were during the summer wildfire season of 2023. The fluorescence capacity of wildfire smoke mainly ranged between  $2 \times 10^{-4}$  and  $7 \times 10^{-4}$ , thus confirming previously reported values in the literature (Hu et al., 2022; Veselovskii et al., 2022a).

Special care was put into the characterization of the fluorescence lidar, where each component along the optical path was considered to determine the system efficiency constants needed to derive the new fluorescence parameters. The detection of 435 optically thin aerosol layers that are only recognizable in the fluorescence signal can significantly improve the detection capabilities of a lidar, which could be critical for low-particle-concentration situations. The enhanced sensitivity results from the fact that laser-induced fluorescence emission originates exclusively from aerosol particles, while air molecules and hydrome-

teors are excluded from this scattering process. Furthermore, as our observations showed, the new dedicated "particle" channel enables an unambiguous differentiation between coexisting unactivated aerosol particles and hydrometeors within clouds.

440

Because of their strong fluorescence and rather low depolarization, the aerosol layers presented in the case studies could be identified as biomass-burning aerosol. The measurements show that such optically thin smoke layers are not so rare in the UTLS region. This suggests that the atmosphere over Europe might be more polluted than previously thought, especially during the summer wildfire season. Those thin layers might not have a strong direct radiative impact, but at these altitudes, smoke 445 particles could become an additional INP source in an otherwise relatively clean atmosphere. Investigating such aerosol layers with a fluorescence lidar, combined with advanced remote-sensing techniques to assess cloud microphysics, could provide more clarity about the relevance of heterogeneous freezing of smoke particles in cirrus cloud formation compared to homogeneous nucleation onto small droplets from background particles. Several observations of cirrus clouds directly below thin 450 biomass-burning aerosol layers suggest that these might be the primary INP source, indicating that heterogeneous freezing is the dominant process. To thoroughly explore this potential aerosol-cloud effect, a larger data set would be beneficial and might provide stronger evidence and more detailed insights into this hypothesis.

Further instrumental upgrades are currently ongoing in the MARTHA system. A new powerful laser, together with a 32-channel spectrometer, will extend the observational sharpness and aim to provide state-of-the-art information about aerosol 455 and clouds from the ground up to the stratosphere.

*Data availability.* Lidar data and products are available upon request at [info@tropos.de](mailto:info@tropos.de) or [polly@tropos.de](mailto:polly@tropos.de). The backward trajectory analysis is based on air mass transport computation with the NOAA (National Oceanic and Atmospheric Administration) HYSPLIT (HYbrid Single-Particle Lagrangian Integrated Trajectory) model ([http://ready.arl.noaa.gov/HYSPLIT\\_traj.php](http://ready.arl.noaa.gov/HYSPLIT_traj.php)). AERONET photometer observations of Leipzig are available in the AERONET database (<http://aeronet.gsfc.nasa.gov/>). GDAS1 (Global Data Assimilation System 1) re-analysis 460 products from the National Weather Service's National Centers for Environmental Prediction are available at <https://www.ready.noaa.gov/gdas1.php>.

*Author contributions.* BG and CJ conceptualized and organized the study. BG wrote the manuscript with the help of CJ. BG took care of the fluorescence products supported by CJ, AA and HB. BG, CJ, RE, AA, UW and MH worked on the experimental setup of the new detection channel. MH, FF, AAF, HG, JH, KO, CJ and BG performed the lidar measurements. MR contributed to the monitoring of the daily aerosol-465 cloud situation. PS, RE, and KO took care of the Polly<sup>XT</sup> lidar system. All co-authors contributed to the several discussions about the new technique and proofreading.

*Competing interests.* The contact author has declared that none of the authors has any competing interests.

*Acknowledgements.* For the fruitful discussions and openness about the new fluorescence technique, we would like to thank Igor Veselovskii, Qiaoyun Hu and Jens Reichardt. We acknowledge the technical team from TROPOS for the experimental support. This research has been 470 supported by the German Federal Ministry of Education and Research (BMBF) under the FONA Strategy "Research for Sustainability" (grant no. 01LK2001A). The contribution of BG was supported from tax revenues on the basis of the budget adopted by the Saxon State Parliament. This work was carried out as part of the Leibniz ScienceCampus 'Smoke and bioaerosols in a changing climate' (BioSmoke) and supported by the Leibniz Association (project number W86/2023).

## References

475 Adam, M.: Notes on temperature-dependent lidar equations, *Journal of Atmospheric and Oceanic Technology*, 26, 1021–1039, <https://doi.org/10.1175/2008JTECHA1206.1>, 2009.

Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245, 1227–1230, <https://doi.org/10.1126/science.245.4923.1227>, 1989.

Ansmann, A., Riebesell, M., and Weitkamp, C.: Measurement of atmospheric aerosol extinction profiles with a Raman lidar, *Optics Letters*, 15, 746–748, <https://doi.org/10.1364/OL.15.000746>, 1990.

Ansmann, A., Wandinger, U., Riebesell, M., Weitkamp, C., and Michaelis, W.: Independent measurement of extinction and backscatter profiles in cirrus clouds by using a combined Raman elastic-backscatter lidar, *Applied Optics*, 31, 7113–7131, <https://doi.org/10.1364/AO.31.007113>, 1992.

Ansmann, A., Baars, H., Tesche, M., Müller, D., Althausen, D., Engelmann, R., Pauliquevis, T., and Artaxo, P.: Dust and smoke transport from Africa to South America: Lidar profiling over Cape Verde and the Amazon rainforest, *Geophysical Research Letters*, 36, [https://doi.org/https://doi.org/10.1029/2009GL037923](https://doi.org/10.1029/2009GL037923), 2009.

Ansmann, A., Seifert, P., Tesche, M., and Wandinger, U.: Profiling of fine and coarse particle mass: case studies of Saharan dust and Eyjafjallajökull/Grímsvötn volcanic plumes, *Atmospheric Chemistry and Physics*, 12, 9399–9415, <https://doi.org/10.5194/acp-12-9399-2012>, 2012.

490 Ansmann, A., Ohneiser, K., Mamouri, R.-E., Knopf, D. A., Veselovskii, I., Baars, H., Engelmann, R., Foth, A., Jimenez, C., Seifert, P., and Barja, B.: Tropospheric and stratospheric wildfire smoke profiling with lidar: mass, surface area, CCN, and INP retrieval, *Atmospheric Chemistry and Physics*, 21, 9779–9807, <https://doi.org/10.5194/acp-21-9779-2021>, 2021.

Ansmann, A., Jimenez, C., Knopf, D. A., Roschke, J., Bühl, J., Ohneiser, K., and Engelmann, R.: Impact of wildfire smoke on Arctic cirrus formation, part 2: simulation of MOSAiC 2019–2020 cases, *EGUphere*, 2024, 1–29, <https://doi.org/10.5194/egusphere-2024-2009>, 495 2024a.

Ansmann, A., Jimenez, C., Roschke, J., Bühl, J., Ohneiser, K., Engelmann, R., Radenz, M., Griesche, H., Hofer, J., Althausen, D., Knopf, D. A., Dahlke, S., Gaudek, T., Seifert, P., and Wandinger, U.: Impact of wildfire smoke on Arctic cirrus formation, part 1: analysis of MOSAiC 2019–2020 observations, *EGUphere*, 2024, 1–29, <https://doi.org/10.5194/egusphere-2024-2008>, 2024b.

Baars, H., Seifert, P., Engelmann, R., and Wandinger, U.: Target categorization of aerosol and clouds by continuous multiwavelength-500 polarization lidar measurements, *Atmospheric Measurement Techniques*, 10, 3175–3201, <https://doi.org/10.5194/amt-10-3175-2017>, 2017.

Barry, K. R., Hill, T. C. J., Levin, E. J. T., Twohy, C. H., Moore, K. A., Weller, Z. D., Toohey, D. W., Reeves, M., Campos, T., Geiss, R., Schill, G. P., Fischer, E. V., Kreidenweis, S. M., and DeMott, P. J.: Observations of Ice Nucleating Particles in the Free Troposphere From Western US Wildfires, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033752, 505 <https://doi.org/https://doi.org/10.1029/2020JD033752>, 2020JD033752, 2021.

Bühl, J., Seifert, P., Radenz, M., Baars, H., and Ansmann, A.: Ice crystal number concentration from lidar, cloud radar and radar wind profiler measurements, *Atmospheric Measurement Techniques*, 12, 6601–6617, <https://doi.org/10.5194/amt-12-6601-2019>, 2019.

Burton, S. P., Ferrare, R. A., Hostetler, C. A., Hair, J. W., Rogers, R. R., Obland, M. D., Butler, C. F., Cook, A. L., Harper, D. B., and Froyd, K. D.: Aerosol classification using airborne High Spectral Resolution Lidar measurements – methodology and examples, *Atmospheric 510 Measurement Techniques*, 5, 73–98, <https://doi.org/10.5194/amt-5-73-2012>, 2012.

Cadondon, J. G., Napal, J. P. D., Abe, K., Lara, R. D., Vallar, E. A., Orbecido, A. H., Belo, L. P., and Galvez, M. C. D.: Characterization of water quality and fluorescence measurements of dissolved organic matter in Cabuyao river and its tributaries using excitation-emission matrix spectroscopy, *Journal of Physics: Conference Series*, 1593, 012 033, <https://doi.org/10.1088/1742-6596/1593/1/012033>, 2020.

Dobretsov, G. E., Syrejschikova, T. I., and Smolina, N. V.: On mechanisms of fluorescence quenching by water, *Biophysics*, 59, 183–188, 515 <https://doi.org/10.1134/S0006350914020079>, 2014.

Edner, H., Johansson, J., Svanberg, S., and Wallinder, E.: Fluorescence lidar multicolor imaging of vegetation, *Appl. Opt.*, 33, 2471–2479, <https://doi.org/10.1364/AO.33.002471>, 1994.

Engelmann, R., Kanitz, T., Baars, H., Heese, B., Althausen, D., Skupin, A., Wandinger, U., Komppula, M., Stachlewska, I. S., Amiridis, V., Marinou, E., Mattis, I., Linné, H., and Ansmann, A.: The automated multiwavelength Raman polarization and water-vapor lidar Polly<sup>XT</sup>: 520 the neXT generation, *Atmospheric Measurement Techniques*, 9, 1767–1784, <https://doi.org/10.5194/amt-9-1767-2016>, 2016.

Floutsi, A. A., Baars, H., Engelmann, R., Althausen, D., Ansmann, A., Bohlmann, S., Heese, B., Hofer, J., Kanitz, T., Haarig, M., Ohneiser, K., Radenz, M., Seifert, P., Skupin, A., Yin, Z., Abdullaev, S. F., Komppula, M., Filioglou, M., Giannakaki, E., Stachlewska, I. S., Janicka, L., Bortoli, D., Marinou, E., Amiridis, V., Gialitaki, A., Mamouri, R.-E., Barja, B., and Wandinger, U.: DeLiAn – a growing collection of depolarization ratio, lidar ratio and Ångström exponent for different aerosol types and mixtures from ground-based lidar observations, 525 *Atmospheric Measurement Techniques*, 16, 2353–2379, <https://doi.org/10.5194/amt-16-2353-2023>, 2023.

Froyd, K. D., Murphy, D. M., Sanford, T. J., Thomson, D. S., Wilson, J. C., Pfister, L., and Lait, L.: Aerosol composition of the tropical upper troposphere, *Atmospheric Chemistry and Physics*, 9, 4363–4385, <https://doi.org/10.5194/acp-9-4363-2009>, 2009.

Froyd, K. D., Murphy, D. M., Lawson, P., Baumgardner, D., and Herman, R. L.: Aerosols that form subvisible cirrus at the tropical tropopause, *Atmospheric Chemistry and Physics*, 10, 209–218, <https://doi.org/10.5194/acp-10-209-2010>, 2010.

Groß, S., Esselborn, M., Weinzierl, B., Wirth, M., Fix, A., and Petzold, A.: Aerosol classification by airborne high spectral resolution lidar 530 observations, *Atmospheric Chemistry and Physics*, 13, 2487–2505, <https://doi.org/10.5194/acp-13-2487-2013>, 2013.

Haarig, M., Ansmann, A., Baars, H., Jimenez, C., Veselovskii, I., Engelmann, R., and Althausen, D.: Depolarization and lidar ratios at 355, 532, and 1064 nm and microphysical properties of aged tropospheric and stratospheric Canadian wildfire smoke, *Atmospheric Chemistry and Physics*, 18, 11 847–11 861, <https://doi.org/10.5194/acp-18-11847-2018>, 2018.

Hansen, J., Sato, M., and Ruedy, R.: Radiative forcing and climate response, *Journal of Geophysical Research: Atmospheres*, 102, 6831–535 6864, <https://doi.org/10.1029/96JD03436>, 1997.

Hu, Q., Goloub, P., Veselovskii, I., Bravo-Aranda, J.-A., Popovici, I. E., Podvin, T., Haeffelin, M., Lopatin, A., Dubovik, O., Pietras, C., Huang, X., Torres, B., and Chen, C.: Long-range-transported Canadian smoke plumes in the lower stratosphere over northern France, *Atmospheric Chemistry and Physics*, 19, 1173–1193, <https://doi.org/10.5194/acp-19-1173-2019>, 2019.

Hu, Q., Goloub, P., Veselovskii, I., and Podvin, T.: The characterization of long-range transported North American biomass burning plumes: 540 what can a multi-wavelength Mie–Raman-polarization-fluorescence lidar provide?, *Atmospheric Chemistry and Physics*, 22, 5399–5414, <https://doi.org/10.5194/acp-22-5399-2022>, 2022.

Immler, F., Engelbart, D., and Schrems, O.: Fluorescence from atmospheric aerosol detected by a lidar indicates biogenic particles in the lowermost stratosphere, *Atmospheric Chemistry and Physics*, 5, 345–355, <https://doi.org/10.5194/acp-5-345-2005>, 2005.

Janicka, L., Davuliene, L., Bycenkiene, S., and Stachlewska, I. S.: Long term observations of biomass burning aerosol over Warsaw by means 545 of multiwavelength lidar, *Opt. Express*, 31, 33 150–33 174, <https://doi.org/10.1364/OE.496794>, 2023.

Jimenez, C., Ansmann, A., Engelmann, R., Haarig, M., Schmidt, J., and Wandinger, U.: Polarization lidar: an extended three-signal calibration approach, *Atmospheric Measurement Techniques*, 12, 1077–1093, <https://doi.org/10.5194/amt-12-1077-2019>, 2019.

Kawana, K., Matsumoto, K., Taketani, F., Miyakawa, T., and Kanaya, Y.: Fluorescent biological aerosol particles over the central Pacific Ocean: covariation with ocean surface biological activity indicators, *Atmospheric Chemistry and Physics*, 21, 15 969–15 983, 550 <https://doi.org/10.5194/acp-21-15969-2021>, 2021.

Knopf, D. A., Alpert, P. A., and Wang, B.: The Role of Organic Aerosol in Atmospheric Ice Nucleation: A Review, *ACS Earth and Space Chemistry*, 2, 168–202, <https://doi.org/10.1021/acsearthspacechem.7b00120>, 2018.

Lakowicz, J. R.: Introduction to Fluorescence, in: *Topics in Fluorescence Spectroscopy: Principles*, edited by Lakowicz, J. R., chap. 1, pp. 555 1–26, Springer Science & Business Media, Boston, third edn., 2006.

Li, B., Chen, S., Zhang, Y., Chen, H., and Guo, P.: Fluorescent aerosol observation in the lower atmosphere with an integrated fluorescence-Mie lidar, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 227, 211–218, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2019.02.019>, 2019.

Liu, Y., Jia, R., Dai, T., Xie, Y., and Shi, G.: A review of aerosol optical properties and radiative effects, *Journal of Meteorological Research*, 560 28, 1003–1028, <https://doi.org/10.1007/s13351-014-4045-z>, 2014.

Lohmann, U.: Anthropogenic Aerosol Influences on Mixed-Phase Clouds, *Current Climate Change Reports*, 13, 32–44, <https://doi.org/10.1007/s40641-017-0059-9>, 2017.

Maciel, F. V., Diao, M., and Patnaude, R.: Examination of aerosol indirect effects during cirrus cloud evolution, *Atmospheric Chemistry and Physics*, 23, 1103–1129, <https://doi.org/10.5194/acp-23-1103-2023>, 2023.

Mamouri, R. E. and Ansmann, A.: Fine and coarse dust separation with polarization lidar, *Atmospheric Measurement Techniques*, 7, 3717–565 3735, <https://doi.org/10.5194/amt-7-3717-2014>, 2014.

Mamouri, R.-E., Ansmann, A., Ohneiser, K., Knopf, D. A., Nisantzi, A., Bühl, J., Engelmann, R., Skupin, A., Seifert, P., Baars, H., Ene, D., Wandinger, U., and Hadjimitsis, D.: Wildfire smoke triggers cirrus formation: lidar observations over the eastern Mediterranean, *Atmospheric Chemistry and Physics*, 23, 14 097–14 114, <https://doi.org/10.5194/acp-23-14097-2023>, 2023.

Mattis, I., Ansmann, A., Althausen, D., Jaenisch, V., Wandinger, U., Müller, D., Arshinov, Y. F., Bobrovnikov, S. M., and Serikov, I. B.: 570 Relative-humidity profiling in the troposphere with a Raman lidar, *Appl. Opt.*, 41, 6451–6462, <https://doi.org/10.1364/AO.41.006451>, 2002.

McCluskey, C. S., DeMott, P. J., Prenni, A. J., Levin, E. J. T., McMeeking, G. R., Sullivan, A. P., Hill, T. C. J., Nakao, S., Carrico, C. M., and Kreidenweis, S. M.: Characteristics of atmospheric ice nucleating particles associated with biomass burning in the US: Prescribed burns 575 and wildfires, *Journal of Geophysical Research: Atmospheres*, 119, 10 458–10 470, <https://doi.org/https://doi.org/10.1002/2014JD021980>, 2014.

McIlrath, T. J.: Fluorescence Lidar, *Optical Engineering*, 19, 194 494, <https://doi.org/10.1117/12.7972549>, 1980.

Murayama, T., Müller, D., Wada, K., Shimizu, A., Sekiguchi, M., and Tsukamoto, T.: Characterization of Asian dust and 580 Siberian smoke with multi-wavelength Raman lidar over Tokyo, Japan in spring 2003, *Geophysical Research Letters*, 31, <https://doi.org/https://doi.org/10.1029/2004GL021105>, 2004.

Müller, D., Mattis, I., Wandinger, U., Ansmann, A., Althausen, D., and Stohl, A.: Raman lidar observations of aged Siberian and Canadian forest fire smoke in the free troposphere over Germany in 2003: Microphysical particle characterization, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/10.1029/2004jd005756>, 2005.

Ohneiser, K., Ansmann, A., Chudnovsky, A., Engelmann, R., Ritter, C., Veselovskii, I., Baars, H., Gebauer, H., Griesche, H., Radenz, M., 585 Hofer, J., Althausen, D., Dahlke, S., and Maturilli, M.: The unexpected smoke layer in the High Arctic winter stratosphere during MOSAiC 2019–2020, *Atmospheric Chemistry and Physics*, 21, 15 783–15 808, <https://doi.org/10.5194/acp-21-15783-2021>, 2021.

590 Ohneiser, K., Ansmann, A., Kaifler, B., Chudnovsky, A., Barja, B., Knopf, D. A., Kaifler, N., Baars, H., Seifert, P., Villanueva, D., Jimenez, C., Radenz, M., Engelmann, R., Veselovskii, I., and Zamorano, F.: Australian wildfire smoke in the stratosphere: the decay phase in 2020/2021 and impact on ozone depletion, *Atmospheric Chemistry and Physics*, 22, 7417–7442, <https://doi.org/10.5194/acp-22-7417-2022>, 2022.

Palmer, S. C., Pelevin, V. V., Goncharenko, I., Kovács, A. W., Zlinszky, A., Présing, M., Horváth, H., Nicolás-Perea, V., Balzter, H., and Tóth, V. R.: Ultraviolet Fluorescence LiDAR (UFL) as a Measurement Tool for Water Quality Parameters in Turbid Lake Conditions, *Remote Sensing*, 5, 4405–4422, <https://doi.org/10.3390/rs5094405>, 2013.

Pan, Y.-L.: Detection and characterization of biological and other organic-carbon aerosol particles in atmosphere using fluorescence, *Journal 595 of Quantitative Spectroscopy and Radiative Transfer*, 150, 12–35, <https://doi.org/https://doi.org/10.1016/j.jqsrt.2014.06.007>, 2015.

Patnaude, R. and Diao, M.: Aerosol Indirect Effects on Cirrus Clouds Based on Global Aircraft Observations, *Geophysical Research Letters*, 47, e2019GL086550, <https://doi.org/https://doi.org/10.1029/2019GL086550>, e2019GL086550 2019GL086550, 2020.

Pinnick, R. G., Hill, S. C., Pan, Y.-L., and Chang, R. K.: Fluorescence spectra of atmospheric aerosol at Adelphi, Maryland, USA: measurement and classification of single particles containing organic carbon, *Atmospheric Environment*, 38, 1657–1672, 600 <https://doi.org/10.1016/j.atmosenv.2003.11.017>, 2004.

Quaas, J., Arola, A., Cairns, B., Christensen, M., Deneke, H., Ekman, A. M. L., Feingold, G., Fridlind, A., Gryspeerdt, E., Hasekamp, O., Li, Z., Lippinen, A., Ma, P.-L., Mülmenstädt, J., Nenes, A., Penner, J. E., Rosenfeld, D., Schrödner, R., Sinclair, K., Sourdeval, O., Stier, P., Tesche, M., van Diedenhoven, B., and Wendisch, M.: Constraining the Twomey effect from satellite observations: issues and perspectives, *Atmospheric Chemistry and Physics*, 20, 15 079–15 099, <https://doi.org/10.5194/acp-20-15079-2020>, 2020.

605 Rao, Z., He, T., Hua, D., Wang, Y., Wang, X., Chen, Y., and Le, J.: Preliminary measurements of fluorescent aerosol number concentrations using a laser-induced fluorescence lidar, *Appl. Opt.*, 57, 7211–7215, <https://doi.org/10.1364/AO.57.007211>, 2018.

Reichardt, J.: Cloud and Aerosol Spectroscopy with Raman Lidar, *Journal of Atmospheric and Oceanic Technology*, 31, 1946 – 1963, <https://doi.org/10.1175/JTECH-D-13-00188.1>, 2014.

610 Reichardt, J., Leinweber, R., and Schwebe, A.: Fluorescing aerosols and clouds: investigations of co-existence, *EPJ Web Conferences*, 176, 05 010, <https://doi.org/10.1051/epjconf/201817605010>, The 28th International Laser Radar Conference (ILRC 28), 2018.

Reichardt, J., Behrendt, O., and Lauermann, F.: Spectrometric fluorescence and Raman lidar: absolute calibration of aerosol fluorescence spectra and fluorescence correction of humidity measurements, *Atmospheric Measurement Techniques*, 16, 1–13, <https://doi.org/10.5194/amt-16-1-2023>, 2023.

615 Rodell, M., Houser, P. R., Jambor, U., Gottschalck, J., Mitchell, K., Meng, C.-J., Arsenault, K., Cosgrove, B., Radakovich, J., Bosilovich, M., Entin, J. K., Walker, J. P., Lohmann, D., and Toll, D.: The Global Land Data Assimilation System, *Bulletin of the American Meteorological Society*, 85, 381 – 394, <https://doi.org/10.1175/BAMS-85-3-381>, 2004.

Saito, Y., Ichihara, K., Morishita, K., Uchiyama, K., Kobayashi, F., and Tomida, T.: Remote Detection of the Fluorescence Spectrum of Natural Pollens Floating in the Atmosphere Using a Laser-Induced-Fluorescence Spectrum (LIFS) Lidar, *Remote Sensing*, 10, 1533, <https://doi.org/10.3390/rs10101533>, 2018.

620 Schmidt, J.: Dual-field-of-view Raman lidar measurements of cloud microphysical properties: Investigation of aerosol-cloud interactions, Ph.D. thesis, Dissertation, Universität Leipzig, 2014, 2014.

Schmidt, J., Wandinger, U., and Malinka, A.: Dual-field-of-view Raman lidar measurements for the retrieval of cloud microphysical properties, *Appl. Opt.*, 52, 2235–2247, <https://doi.org/10.1364/AO.52.002235>, 2013.

Stevens, B. and Feingold, G.: Untangling aerosol effects on clouds and precipitation in a buffered system, *Nature*, 461, 607–613, 625 <https://doi.org/10.1038/nature08281>, 2009.

Stryer, L.: Excited-state proton-transfer reactions. A deuterium isotope effect on fluorescence, *Journal of the American Chemical Society*, 88, 5708–5712, 1966.

Sugimoto, N., Huang, Z., Nishizawa, T., Matsui, I., and Tatarov, B.: Fluorescence from atmospheric aerosols observed with a multi-channel lidar spectrometer, *Opt. Express*, 20, 20 800–20 807, <https://doi.org/10.1364/OE.20.020800>, 2012.

630 Twomey, S.: Pollution and the planetary albedo, *Atmospheric Environment* (1967), 8, 1251–1256, [https://doi.org/https://doi.org/10.1016/0004-6981\(74\)90004-3](https://doi.org/https://doi.org/10.1016/0004-6981(74)90004-3), 1974.

Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *Journal of Atmospheric Sciences*, 34, 1149 – 1152, [https://doi.org/10.1175/1520-0469\(1977\)034<1149:TIOPOT>2.0.CO;2](https://doi.org/10.1175/1520-0469(1977)034<1149:TIOPOT>2.0.CO;2), 1977.

635 Twomey, S. A., Piepgrass, M., and Wolfe, T. L.: An assessment of the impact of pollution on global cloud albedo, *Tellus B*, 36B, 356–366, <https://doi.org/https://doi.org/10.1111/j.1600-0889.1984.tb00254.x>, 1984.

Veselovskii, I., Hu, Q., Goloub, P., Podvin, T., Korenskiy, M., Pujol, O., Dubovik, O., and Lopatin, A.: Combined use of Mie–Raman and fluorescence lidar observations for improving aerosol characterization: feasibility experiment, *Atmospheric Measurement Techniques*, 13, 6691–6701, <https://doi.org/10.5194/amt-13-6691-2020>, 2020.

640 Veselovskii, I., Hu, Q., Goloub, P., Podvin, T., Choël, M., Visez, N., and Korenskiy, M.: Mie–Raman–fluorescence lidar observations of aerosols during pollen season in the north of France, *Atmospheric Measurement Techniques*, 14, 4773–4786, <https://doi.org/10.5194/amt-14-4773-2021>, 2021.

Veselovskii, I., Hu, Q., Ansmann, A., Goloub, P., Podvin, T., and Korenskiy, M.: Fluorescence lidar observations of wildfire smoke inside cirrus: a contribution to smoke–cirrus interaction research, *Atmospheric Chemistry and Physics*, 22, 5209–5221, <https://doi.org/10.5194/acp-22-5209-2022>, 2022a.

645 Veselovskii, I., Hu, Q., Goloub, P., Podvin, T., Barchunov, B., and Korenskii, M.: Combining Mie–Raman and fluorescence observations: a step forward in aerosol classification with lidar technology, *Atmospheric Measurement Techniques*, 15, 4881–4900, <https://doi.org/10.5194/amt-15-4881-2022>, 2022b.

Veselovskii, I., Kasianik, N., Korenskii, M., Hu, Q., Goloub, P., Podvin, T., and Liu, D.: Multiwavelength fluorescence lidar observations of smoke plumes, *Atmospheric Measurement Techniques*, 16, 2055–2065, <https://doi.org/10.5194/amt-16-2055-2023>, 2023.

650 Wang, Y., Hu, R., Xie, P., Chen, H., Wang, F., Liu, X., Liu, J., and Liu, W.: Measurement of tropospheric HO<sub>2</sub> radical using fluorescence assay by gas expansion with low interferences, *Journal of Environmental Sciences*, 99, 40–50, <https://doi.org/https://doi.org/10.1016/j.jes.2020.06.010>, 2021.

Wright, A. G.: Why photomultipliers?, in: *The Photomultiplier Handbook*, Oxford University Press, ISBN 9780199565092, <https://doi.org/10.1093/oso/9780199565092.003.0001>, 2017.

655 Zhang, M., Klimach, T., Ma, N., Könemann, T., Pöhlker, C., Wang, Z., Kuhn, U., Scheck, N., Pöschl, U., Su, H., and Cheng, Y.: Size-Resolved Single-Particle Fluorescence Spectrometer for Real-Time Analysis of Bioaerosols: Laboratory Evaluation and Atmospheric Measurements, *Environmental Science & Technology*, 53, 13 257–13 264, <https://doi.org/10.1021/acs.est.9b01862>, pMID: 31589819, 2019.