

Review Article

AntArchitecture:

Building an age–depth model from Antarctica’s radiostratigraphy to explore ice-sheet evolution

Robert G. Bingham^{1*}, Julien A. Bodart^{2,3*}, Marie G. P. Cavitte^{4*}, Ailsa Chung^{5*}, Rebecca J. Sanderson^{6*}, Johannes C. R. Sutter^{2,3*}, Olaf Eisen^{7,8}, Nanna B. Karlsson⁹, Joseph A. MacGregor¹⁰, Neil Ross⁶, Duncan A. Young¹¹, David W. Ashmore¹², Andreas Born^{13,14}, Winnie Chu¹⁵, Xiangbin Cui¹⁶, Reinhard Drews¹⁷, Steven Franke^{7,17}, Vikram Goel¹⁸, John W. Goodge^{19,20}, A. Clara J. Henry²¹, Antoine Hermant^{2,3}, Benjamin H. Hills²², Nicholas Holschuh²³, Michelle R. Koutnik²⁴, Gwendolyn J.-M. C. Leysinger Vieli²⁵, Emma J. MacKie²⁶, Elisa Mantelli^{7,27}, Carlos Martín²⁸, Felix S. L. Ng²⁹, Falk M. Oraschewski⁹, Felipe Napoleoni¹, Frédéric Parrenin⁵, Sergey V. Popov^{30,31}, Therese Rieckh^{13,14}, Rebecca Schlegel¹⁷, Dustin M. Schroeder^{32,33}, Martin J. Siegert³⁴, Xueyuan Tang^{16,35}, Thomas O. Teisberg³³, Kate Winter³⁶, Shuai Yan¹¹, Harry Davis¹, Christine F. Dow^{37,38}, Tyler J. Fudge²⁴, Tom A. Jordan²⁸, Bernd Kulesa³⁹, Kenichi Matsuoka⁴⁰, Clara J. Nyqvist¹, Maryam Rahnemoonfar^{41,42}, Matthew R. Siegfried²², Shivangini Singh^{11,43}, Vjeran Višnjević^{2,3}, Rodrigo Zamora⁴⁴, Alexandra Zuhr¹⁷

* These authors contributed equally to this work.

Correspondence to: Robert G. Bingham (r.bingham@ed.ac.uk)

Affiliations

¹ School of GeoSciences, University of Edinburgh, UK

² Climate and Environmental Physics, Physics Institute, University of Bern, Switzerland

³ Oeschger Centre for Climate Change Research, University of Bern, Switzerland

⁴ Department of Water and Climate, Vrije Universiteit Brussel, Belgium

⁵ Univ. Grenoble Alpes, CNRS, INRAE, IRD, Grenoble INP, IGE, 38000 Grenoble, France

⁶ School of Geography, Politics and Sociology, Newcastle University, UK

⁷ Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

⁸ Department of Geosciences, University of Bremen, Germany

⁹ Geological Survey of Denmark and Greenland (GEUS), Copenhagen, Denmark

¹⁰ Cryospheric Sciences Laboratory, NASA Goddard Space Flight Centre, Greenbelt, Maryland, USA

- 30 ¹¹ University of Texas Institute for Geophysics, Jackson School of Geosciences, University of Texas at Austin,
31 Texas, USA
- 32 ¹² Met Office, FitzRoy Road, Exeter, UK
- 33 ¹³ Department of Earth Science, University of Bergen, Norway
- 34 ¹⁴ Bjerknes Centre for Climate Research, University of Bergen, Norway
- 35 ¹⁵ School of Earth and Atmospheric Science, Georgia Institute of Technology, Atlanta, Georgia, USA
- 36 ¹⁶ Polar Research Institute of China, Shanghai, China
- 37 ¹⁷ Department of Geosciences, University of Tübingen, Germany
- 38 ¹⁸ National Centre for Polar and Ocean Research (NCPOR), Ministry of Earth Sciences, Vasco da Gama, Goa,
39 India
- 40 ¹⁹ Department of Earth and Environmental Sciences, University of Minnesota Duluth, Duluth, Minnesota, USA
- 41 ²⁰ Planetary Science Institute, Tucson, Arizona, USA
- 42 ²¹ Department of Mathematics, Stockholm University, Sweden
- 43 ²² Department of Geophysics, Colorado School of Mines, Golden, Colorado, USA
- 44 ²³ Department of Geology, Amherst College, Massachusetts, USA
- 45 ²⁴ Department of Earth and Space Sciences, University of Washington, Seattle, Washington, USA
- 46 ²⁵ Department of Geography, University of Zurich, Switzerland
- 47 ²⁶ Department of Geological Sciences, University of Florida, Gainesville, Florida, USA
- 48 ²⁷ Department of Earth and Environmental Sciences, Ludwig-Maximilians-Universität, Munich, Germany
- 49 ²⁸ British Antarctic Survey, Cambridge, UK
- 50 ²⁹ Department of Geography, University of Sheffield, Sheffield, UK
- 51 ³⁰ St Petersburg State University, Russia
- 52 ³¹ Polar Marine Geosurvey Expedition, St. Petersburg, Russia
- 53 ³² Department of Geophysics, Stanford University, California, USA
- 54 ³³ Department of Electrical Engineering, Stanford University, California, USA
- 55 ³⁴ Tremough House, Penryn Campus, University of Exeter, Cornwall, UK
- 56 ³⁵ School of Oceanography, Shanghai Jiao Tong University, China
- 57 ³⁶ Department of Geography and Environmental Sciences, Northumbria University, Newcastle, UK

- 58 ³⁷ Department of Geography and Environmental Management, University of Waterloo, Ontario, Canada
- 59 ³⁸ Department of Applied Mathematics, University of Waterloo, Ontario, Canada
- 60 ³⁹ School of Biosciences, Geography and Physics, Swansea University, UK
- 61 ⁴⁰ Norwegian Polar Institute, Tromsø, Norway
- 62 ⁴¹ Department of Computer Science and Engineering, Lehigh University, Pennsylvania, USA
- 63 ⁴² Department of Civil and Environmental Engineering, Lehigh University, Pennsylvania, USA
- 64 ⁴³ Department of Earth and Planetary Sciences, University of Texas at Austin, Austin, Texas, USA
- 65 ⁴⁴ Centro de Estudios Científicos, Valdivia, Chile

Abstract. Radio-echo sounding (RES) has revealed an internal architecture within both the West and East Antarctic ice sheets that records their depositional, deformational and melting histories. Crucially, RES-imaged internal-reflecting horizons, tied to ice-core age-depth profiles, can be treated as isochrones that record the age-depth structure across the Antarctic ice sheets. These enable the reconstruction of past climate and ice-dynamical processes on large scales, which are complementary to but more spatially extensive than commonly used proxy records (e.g., former ice limits constrained by cosmogenic dating, or offshore sediment sequences) around Antarctica. We review progress towards building a pan-Antarctic age-depth model from these data by first introducing the relevant RES datasets that have been acquired across Antarctica over the last six decades (focussing specifically on those that detected internal-reflecting horizons), and outlining the processing steps typically undertaken to visualise, trace and date (by intersection with ice cores, or modelling) the RES-imaged isochrones. We summarise the scientific applications to which Antarctica's internal architecture has been applied to date and present a pathway to expanding Antarctic radiostratigraphy across the continent to provide a benchmark for a wider range of investigations: (1) Identification of optimal sites for retrieving new ice-core palaeoclimate records targeting different periods; (2) Reconstruction of surface mass balance on millennial or historical timescales; (3) Estimates of basal melting and geothermal heat flux from radiostratigraphy and comprehensively mapping basal-ice units, to complement inferences from other geophysical and geological methods; (4) Advancing knowledge of volcanic activity and fallout across Antarctica; (5) The refinement of numerical models that leverage radiostratigraphy to tune time-varying accumulation, basal melting and ice flow, firstly to reconstruct past behaviour, and then to reduce uncertainties in projecting future ice-sheet behaviour.

86 1 Introduction

87 Throughout the Quaternary (2.58 Ma to present), Antarctica’s ice cover has waxed and waned, inducing
88 concomitant rises and falls in global sea level on the order of several tens of metres (e.g., Drewry, 1983;
89 Pollard and DeConto, 2009; Dutton et al., 2015). It is critical to understand the rates and drivers of these past
90 oscillations in order to contextualise current observations of persistent and accelerating losses from the
91 contemporary Antarctic ice sheets (e.g., Fox-Kemper et al., 2021; Otosaka et al., 2023) and thereby project
92 as accurately as possible the rates at which future global sea-level rise fuelled by ice melt will occur (e.g.,
93 Scambos et al., 2017; Oppenheimer et al., 2019). The evidence for past Antarctic ice-sheet fluctuations has
94 been derived predominantly from sampling sediments deposited offshore around the continent (Escutia et
95 al., 2009; Naish et al., 2009; Cook et al., 2013; Bentley et al., 2014; Gulick et al., 2017; Hillenbrand et al., 2017),
96 dating the exposure history of onshore bedrock and moraine boulders (Brook and Kurz, 1993; Stone et al.,
97 2003; Johnson et al., 2008; Mackintosh et al., 2014; Hein et al., 2016; Hillebrand et al., 2021), and by analysing
98 the ice itself recovered from ice-core sites (e.g., EPICA Community Members, 2004; Jouzel et al., 2007; Higgins
99 et al., 2015; WAIS Divide Project Members, 2015; Dome Fuji Ice Core Project Members, 2017; Yan et al., 2021)
100 (see Brook and Buizert, 2018 for an overview). Together, these form the palaeoclimate records that underpin
101 numerical-modelling reconstructions of past and present ice-sheet extents and inform projections of how
102 these may evolve into the future and affect sea-level change (e.g., Gasson et al., 2016; Golledge et al., 2019;
103 DeConto et al., 2021; Pittard et al., 2022). Recovery of further sediment and ice cores around Antarctica to
104 refine these records and projections remains a scientific imperative – and yet these records are intrinsically
105 spatially limited, are often restricted on the timescales of observation, and for the most part are indirect with
106 respect to ice conditions. Radio-echo sounding across Antarctica complements these records by providing
107 *spatially continuous* data that record past and present ice conditions and, by extension, past and present
108 climate conditions, across the ice sheets.

109 Radio-echo sounding (RES) describes the investigation of the subsurface of ice sheets using electromagnetic
110 waves, and has been conducted from both airborne and ground-based platforms across the Antarctic ice
111 sheets for over 60 years (see reviews by Dowdeswell and Evans, 2004; Bingham and Siegert, 2007; Allen,
112 2008; Schroeder et al., 2020). Primarily deployed for mapping the ice-sheet bed and thereby measuring ice
113 thickness and thus ice volume, the majority of RES surveys have also imaged numerous englacial features,
114 predominantly internal-reflection horizons (a.k.a. internal or englacial layers), crevasses and rheologically-
115 distinct “basal units” of ice that occur between the more obvious reflections of the ice surface and bed (Fig.
116 1). For this review, we collectively term all of the Antarctic ice sheets’ RES-imaged englacial features its
117 *internal architecture*. We will demonstrate that although great progress has already been made in using some
118 of this resource to elucidate ice and climate history, Antarctica’s internal architecture has yet to be exploited
119 to its full potential in refining our understanding of past, present and future ice-sheet behaviour.




Figure 1. Schematic illustration of Antarctica’s internal architecture and the key processes governing its structure. Internal-reflection horizons - the ice sheet’s “radiostratigraphy” - are represented by grey lines between the surface and bed.

In Greenland, a comprehensive archive of internal architecture has already been assembled (see MacGregor et al., 2015a; 2025), facilitating the ice-sheet-wide reconstruction of past accumulation and dynamics, to improve past and future sea-level estimates (MacGregor et al., 2016; Born and Robinson, 2021). However, several major issues have confounded progress in capturing and applying internal architecture across Antarctica, including:

- 1) The Antarctic ice sheets together cover eight times the area of the Greenland Ice Sheet.
- 2) RES data have been collected, processed and archived by multiple international groups across the Antarctic ice sheets, and hence are not available in a standardised form across Antarctica.
- 3) A comprehensive suite of strategies for using internal architecture in numerical ice-sheet models has not been developed.
- 4) Much internal architecture in RES data is highly challenging to identify and map with automated methods.

To address these challenges and work collectively towards consistently capturing and utilising Antarctica’s internal architecture, an international community called *AntArchitecture* was formed in 2018. This community, coordinated via the *Scientific Committee for Antarctic Research* (SCAR), aspires to the ultimate scientific aim of using Antarctica’s internal architecture to deconvolve its ice sheets’ histories and thereby facilitate improved projections of their future behaviour in the face of global climate warming. A first step in

137 this process, and one of the aims of this review, collectively written by the *AntArchitecture* community, is to
138 compile the international community's understanding of the present state of the field in terms of available
139 RES data across the Antarctic ice sheets and their potential applications. Additionally, we seek here to relay
140 community aspirations to address the aforementioned challenges and position Antarctica's internal
141 architecture as a valuable resource for improving our understanding of its ice/climate interactions.

142 We begin with a brief overview of what gives rise to internal architecture in ice, especially the internal-
143 reflection horizons (hereafter IRHs) that are measured by RES (Sect. 2). We then describe how RES data have
144 been, and can be, processed to optimise the extraction of internal architecture and its visualisation, and
145 discuss the common methods currently used to characterise and date IRHs (Section 3). In Sect. 4, we
146 summarise the key RES datasets acquired across Antarctica that image internal architecture, to contextualise
147 in a single place the type and quality of information recorded by each institute and survey in the past six
148 decades, and present an inventory of which existing RES data have so far had several IRHs traced through
149 them. In Sect. 5, we review how internal architecture has been used to reconcile ice-core records, calculate
150 changes to past surface mass balance, explore basal melting in association with subglacial lakes and areas of
151 enhanced geothermal heat flux, and investigate ice-sheet dynamics and other glaciological questions; and
152 outline how the internal architecture has begun to be used in numerical-modelling applications to date. In
153 Sect. 6, we outline a recommended pathway to building a pan-Antarctic database of Antarctica's internal
154 architecture, and discuss key science activities that can be facilitated by its delivery.

155 **2 Internal architecture in ice sheets**

156 The most common way in which internal architecture is viewed and assessed is as radargrams, which are
157 two-dimensional profiles of echo power arrayed in the along-track direction (e.g., Fig. 2). Antarctic
158 radargrams commonly display clear *radiostratigraphy*, the collective term for the multiple sub-parallel and
159 closely-spaced IRHs that are seen in radargrams and often, although do not always, broadly follow the shape
160 of the ice-bed interface (e.g., Fig. 2). IRHs occur as radio-waves propagate down through the ice column and
161 reflect off any boundary where there is a contrast in the dielectric properties within the ice. The propagation
162 of radio-waves through snow, firn and ice is controlled by the relative permittivities of these materials, which
163 are functions of density, electrical conductivity, and/or the development of ice-fabric anisotropy where ice
164 crystals align into a preferential orientation as a result of large englacial stress. Where contrasts in any of
165 these properties are sufficiently strong and sharp, the incident energy will partition and a small fraction of it
166 will be reflected back to the RES receiver at or above the ice surface.

167 In the upper and middle part of the ice column, radiostratigraphy typically arises from (a) density variations,
168 as snow compacts into ice (as explained in pioneering work by Robin et al. (1969) and Clough (1977)) and (b)
169 variations in electrical conductivity, as volcanic aerosols present in the air during snow deposition are




Figure 2. Radargrams from Institute Ice Stream, West Antarctica, obtained by the British Antarctic Survey PASIN RES system in (a) pulse (shallow-sounding) and (b) chirp (deep-sounding) radar modes (Frémand et al., 2022), vertically differentiated to accentuate fine detail. Symbols highlight three IRHs found widely across West Antarctica in airborne radar data. The bed reflection (black-white interface) is partially visible in (a) and clearly visible in (b), marked by the black arrow. Figure modified from Ashmore et al. (2020).

incorporated into the firn (Hammer, 1980; Millar, 1981; Millar, 1982). These density- and electrical-conductivity-derived IRHs are related to snow and ice layers of a specific age buried under subsequent snow accumulation, and thus may be considered isochronous (Hempel et al., 2000; Eisen et al., 2006). Such RES-imaged isochrones may often represent composites of multiple real horizons in the ice, and their thickness is dependent on RES-system resolution (Harrison, 1973; Winter et al., 2017). They are often traceable for considerable distances on RES profiles: some IRHs in the Antarctic and Greenland ice sheets are continuous for hundreds or even thousands of kilometres (e.g., MacGregor et al., 2015a; Winter et al., 2019a; Ashmore et al., 2020). For the focus of this review, isochronous reflections arising from density and electrical conductivity are of significant interest, and IRHs that can be dated at ice cores and traced continuously over long distances to form a “dated radiostratigraphy” are particularly valuable (as explored in-depth in Sect. 4 and 5). There are, however, some cases, especially in the lower part of the ice column, where diachronous IRHs (i.e. IRHs that cannot be treated as single time markers) may be visualised in radargrams. The most common such examples are IRHs that are thought to manifest sudden changes in ice-crystal-orientation fabric that cause anisotropic radio-wave propagation, or cold-warm ice transitions where the pore space on the warm side is filled with meltwater instead of air (Harrison, 1973; Fujita et al., 1999; Eisen et al., 2007). Over ice shelves, pervasive IRHs can mark the boundary between atmospherically-derived (meteoric) and subglacially/submarine-accreted (marine) ice (Holland et al., 2009; Das et al., 2020).

188 The specular behaviour of IRHs also positions them as ideal targets for repeated observations of vertical
189 velocity over time, directly tracking the deformation of the ice sheet, via static phase-sensitive repeat
190 measurements at a point (autonomous phase-sensitive radio-echo sounder, or ApRES; Nicholls et al., 2015)
191 or from airborne re-flights of transects with coherent RES systems (Castelletti et al., 2021). Although these
192 methods have been practised in recent field campaigns (e.g., Hills et al., 2022; Chung et al., 2023; Fudge et
193 al., 2023), we do not discuss this aspect of radiostratigraphy further in this review, beyond noting that
194 establishing the distribution of appropriate IRHs could be a valuable component in expedition planning. A
195 review of static techniques is found in Kingslake et al. (2014), while repeat-pass airborne interferometry of
196 IRH is an active field of research (Castelletti et al., 2021).

197 While the imaging and analysis of radiostratigraphy and its application to assessing ice-sheet evolution form
198 the main focus of this paper, other significant features of internal architecture also convey information that
199 can be used to help understand current and past ice-sheet processes (as depicted in Fig. 1). These include
200 basal units which exhibit different dielectric properties to the surrounding ice and may result from ice-folding
201 due to contrasts in material properties, to accretion, melting due to high rates of geothermal heat flux or
202 overburden pressure from the ice above, or freeze-on processes taking place at the base of the ice sheet (Bell
203 et al., 2011; Bell et al., 2014; Bons et al., 2016; Leysinger Vieli et al., 2018; Wrona et al., 2018; Ross et al.,
204 2020; Franke et al., 2023). Additionally, buried near-surface and basal crevassing imaged by RES systems may
205 be indicative of past grounding-line evolution or ice-stream stagnation events (Retzlaff et al., 1993; Matsuoka
206 et al., 2009; Catania et al., 2010; Kingslake et al., 2018; Wearing and Kingslake, 2019). We elaborate further
207 on these other significant features of internal architecture in Sect. 5.5.

208 **3 Extracting and dating internal architecture from RES data**

209 The information available from radargrams (e.g. Fig. 2), and the degree to which the internal architecture can
210 be used for different applications, depend firstly on the settings of the RES system acquiring the data and
211 secondly on choices made in processing the data. Below we summarise the typical processing workflow for
212 radargram generation and highlight key decisions that influence interpretation of the resulting
213 radiostratigraphy. Figure 3 presents a conceptual support to this discussion. We then discuss the different
214 methods used to trace radiostratigraphy through radargrams, and to assign dates to key IRHs.

215 **3.1 Data processing for optimising IRH tracing**

216 RES data acquisition (Fig. 3a) can be categorised broadly based on two criteria: (a) Phase control of the
217 transmitter or phase sampling by the receiver (i.e., coherent vs. incoherent); and (b) the nature of the
218 transmitted wave (pulsed versus chirped; Gogineni et al. (1998); Peters et al. (2005)). Processing (depicted in
219 Fig. 3b) is similar for all systems, so here we highlight differences that affect radargram quality. Direct




Figure 3. Flowchart illustrating key steps for the processing of RES data from chirp and pulse systems for subsequent radiostratigraphic analysis. (a) Basic configurations and parameters defined on data acquisition. (b) Fundamental and (c) additional steps commonly taken when processing data to visualise IRHs. (d) Depiction of some common ways of tracing or otherwise quantifying IRH geometry.

220
 221 measurements of the dielectric properties of ice cores show that ice conductivity varies on much smaller
 222 length scales than can be imaged by RES (Harrison, 1973; Eisen et al., 2003). Therefore, each RES system
 223 represents subsurface reflectors differently, and data acquired from the same area but by different RES
 224 systems may show different IRHs on intersecting radargrams due to the differences in RES imaging
 225 capabilities (see Fig. 4, after Winter et al. (2017), for an example of a comparison between different RES
 226 systems). For pulsed systems, processing cannot improve the vertical resolution, which is controlled by the
 227 bandwidth and the rate of sampling of the received waveform. For chirped systems, the waveform must be




Figure 4. RES profiles of a few km length for five RES systems, that have profiled across or near Dome C (location in Fig. 6d). The vertical red line in each profile marks the position of the trace closest to Dome C. The surface reflections are shifted to time zero and the length of the RES profiles is indicated on the horizontal axes. For the bottom UTIG and CReSIS panels a 2-D-focused processing is applied. The RES data were acquired with: 1. AWI 150 MHz Aero-EMR; 2. UTIG 60 MHz HiCARS; 3. CReSIS 194 MHz MCoRDS; 4. Italian National Institute of Geophysics and Volcanology (INGV) 150 MHz RES system; and 5. BAS 150 MHz PASIN; for full details and original figure from which this is modified, see Winter et al. (2017).

229 fully sampled first and then match-filtered, integrating the received power while also finely resolving
230 radiostratigraphy targets based on the chirp's bandwidth (Hélière et al., 2007; Peters et al., 2007). This “pulse
231 compression” is the first step in producing a radargram from a chirped system.

232 Following initial data acquisition, RES data are typically processed using geophysical techniques of varying
233 sophistication (Fig. 3b). For example, incoherent noise is typically reduced by various forms of horizontal
234 averaging, and bandpass-filtering can remove irrelevant components of the measured signal. Finally, if
235 possible the data should be focused or migrated to reposition the received signal energy as precisely as
236 possible to their true subsurface locations. This can be done via several methods: (a) Incoherent echo
237 summation, often termed *migration* as in reflection seismology (Yilmaz, 2001); (b) SAR-focusing for point
238 scatterers, common in satellite applications (Ulaby and Lang, 2015); or (c) applying algorithms designed
239 specifically for RES of specular reflections (Heister and Scheiber, 2018; Castelletti et al., 2019; Xu et al., 2022).
240 SAR-focusing has a proven ability to reduce image artefacts and improve along-track resolution, especially in
241 areas with steeply-sloping radiostratigraphy (Holschuh et al., 2014; Castelletti et al., 2019). Multiple SAR-
242 processing techniques currently exist for coherent RES systems, including: (a) unfocused SAR (short apertures
243 without phase correction and equivalent in name to Doppler filtering or coherent echo summation; Hélière
244 et al. (2007)); or (b) more advanced focused SAR, using either 1-D correlations resulting in intermediate
245 apertures, or 2-D correlations resulting in longer apertures (Peters et al., 2005; Peters et al., 2007). The latter
246 is the processing of choice for modern coherent systems for the detection of IRHs in areas with steeply
247 dipping reflections. Unfocused and 1-D SAR approaches will emphasise flat specular reflectors and reduce
248 clutter, at a cost of dipping specular horizons. Large SAR apertures are critical for tracking steeply dipping
249 IRHs, but present greater computational costs and an overall reduction of signal to noise ratio. Cross-track
250 antenna arrays can allow for determination of cross-track IRH slopes.

251 A series of additional corrections and image-processing steps can also be taken to optimise RES data for
252 tracing radiostratigraphy (Fig. 3c). For radar data acquired by airborne platforms, the aircraft-to-ice surface
253 space on the radargram must be removed to obtain true depths below the ice surface; this is often conducted
254 by shifting the vertical axis of the radargram to time zero for each RES trace and flattening the surface based
255 on the location of the surface reflection on the radargram. This can be done using data from the altimeter
256 and/or LIDAR onboard the aircraft, high-resolution surface DEMs, or using the picked surface reflection from
257 the radargram itself (e.g., MacGregor et al., 2015a). Localised density corrections, based on ground-truthing
258 measurements in the upper section of ice cores or other geophysical measurements (e.g., radar data acquired
259 by airborne platforms; Eisen et al., 2002), may also be applied to convert the two-way-travel time from the
260 RES data to ice-equivalent depths. Alternatively, for depth-correcting RES below the pore close-off depth, a
261 spatially uniform firn-depth correction that is typically of the order of several metres may be used to obtain
262 ice-equivalent depths (e.g., Ashmore et al., 2020), although this assumption may only be valid in dry and
263 stable parts of the ice sheet and not in highly dynamic regions (Dowdeswell and Evans, 2004). Others have

also vertically rescaled (or flattened) RES data to facilitate the tracing of continuous reflections by semi-automatic pickers (e.g., Fahnestock et al., 2001a; Sect. 3.2; MacGregor et al., 2015a; 2025). Finally, specific image-processing filters can also be applied to enhance the gain and reduce incoherent noise, which can facilitate IRH tracing on RES data (Ashmore et al., 2020; Bodart et al., 2021; Wang et al., 2023; Franke et al., 2025).

Importantly for users interested in tracing IRHs, and especially the deepest IRHs, most RES data over Antarctica, including those available from open-access repositories, are not optimised for detecting radiostratigraphy. Typically the data have been acquired and processed to optimise retrieval of the bed echo, and some datasets require considerable reprocessing from the raw data to improve the clarity of the radiostratigraphy between the ice surface and the bed (Castelletti et al., 2019). In particular, for thick or unusually heterogenous ice, the best strategy is often to experiment with filtering data differently at different depths until the IRHs at selected depths are most clearly visualised.

3.2 Tracing radiostratigraphy

The primary method for extracting internal architecture from radargrams (e.g. Fig. 3d) has been to trace or “pick” IRHs, typically using semi-automated techniques (e.g., Cavitte et al., 2016; Koch et al., 2023). Where radargram quality is high, IRHs are easily traced and continuous, and fully automated methods may also perform well (e.g., Panton, 2014; Xiong et al., 2018; Delf et al., 2020). Machine-learning methods are in their infancy but show promise for more rapidly tracing radiostratigraphy, as demonstrated recently by Moqadam et al. (2025) on deep IRHs. However, so far most successful applications have been limited to near-surface IRHs in the upper few tens of metres of the ice column (e.g., Dong et al., 2021; Rahnemoonfar et al., 2021; Yari et al., 2021), primarily due to the lack of vertical disturbances and low noise in surface-conformable IRHs. Thus, for most radargrams and deep-ice applications, semi-automated tracing of IRHs is still required. This relies on algorithms that typically follow the local maxima in return power between adjacent traces within a predetermined vertical window, using either open-source or commercial and bespoke software from the seismic industry (e.g., Winter et al., 2019a; Ashmore et al., 2020; Chung et al., 2023; Sanderson et al., 2024). A comprehensive overview of IRH-tracing methods is provided by Moqadam et al. (2025).

The process of tracing IRHs can be categorised into two approaches: (a) tracing as many IRHs as possible regardless of their amplitudes or continuity (MacGregor et al., 2015a; 2025); or (b; more commonly) by identifying IRHs that have a high echo-power, appear distinguishably brighter than adjacent IRHs on radargrams and are continuous for long distances (>100 km), using crossovers between intersecting RES profiles to ensure reliability in the tracing process (e.g., Cavitte et al., 2016; Winter et al., 2019a; Ashmore et al., 2020; Bodart et al., 2021; Wang et al., 2023).

296 Importantly, the thickness of a given IRH in a radargram is dependent on the range resolution of the RES
297 system used to image it, such that RES systems with high pulse-width, and thus finer vertical resolution, may
298 detect several thinner IRHs that would otherwise appear as a single, broader reflection in coarser-resolution
299 systems (see Fig. 4 and Harrison, 1973; Millar, 1982; Karlsson et al., 2014; Winter et al., 2017; Bodart et al.,
300 2021; Cavitte et al., 2021; Franke et al., 2025). This must be accounted for when comparing the position and
301 aspect of IRHs traced in data from RES systems operating with different frequencies and system
302 characteristics (Winter et al., 2017, Franke et al., 2025).

303 **3.3 Complementary approaches to tracing IRHs for characterising radiostratigraphy**

304 Even having applied all possible data processing strategies described above, radiostratigraphy may remain
305 challenging or impossible to trace over some regions due to the innate physical properties of ice in such areas.
306 For example, IRHs may become warped/buckled or disrupted by differential ice flow or flow over steep
307 topography (e.g., Siegert et al., 2003b; Ross et al., 2011; Bingham et al., 2015; Franke et al., 2023; Jansen et
308 al., 2024), while unconformities can be introduced by significant wind scouring of the ice surface (e.g., Welch
309 and Jacobel, 2005; Luo et al., 2022). This variability in itself provides important information about past and
310 present ice behaviour (as we explore further in Sect. 5), and hence warrants alternate methods to
311 characterise the radiostratigraphy where IRHs cannot readily be traced.

312 One method for assessing the general variability of radiostratigraphy across large regions of ice sheets is the
313 Internal Layering Continuity Index (ILCI) developed by Karlsson et al. (2012). This tool maps the variability in
314 vertical signal strength for individual RES traces, acting as a relative measure of the number of dielectric
315 contrasts compared to signal-to-noise ratio. High ILCI values typically indicate regions of an ice sheet
316 characterised by multiple, traceable IRHs, while low ILCI values tend to indicate regions of ice sheet with
317 disrupted or discontinuous IRHs or regions with very few or no IRHs detected by the RES system. Although
318 the method is not easily transferable between different RES systems due to acquisition and processing
319 differences, ILCI has been extensively applied to several regions both in Antarctica (Fig. 5) and Greenland as
320 a mechanism for identifying rapidly the specific sub-regions in which IRHs are likely to be traceable (e.g., Sime
321 et al., 2014; Bingham et al., 2015; Karlsson et al., 2018; Frémand et al., 2022; Tang et al., 2022; Sanderson et
322 al., 2023).

323 Alternative methods have focussed on the extraction of IRH slopes. This avenue acknowledges the challenges
324 of tracing and dating radiostratigraphy in areas of fast or complex ice flow, or where the acquisition or
325 processing methods that have been used were not tailored to the recovery of radiostratigraphy. For
326 discontinuous radiostratigraphy, local slope information is valuable, because radiostratigraphic slope is
327 closely related to particle trajectories within the ice sheet (Hindmarsh et al., 2006; Parrenin and Hindmarsh,
328 2007; Ng and King, 2011; Holschuh et al., 2017). Several methods have therefore been developed to extract
329 slope information, such as incoherent averaging methods (Sime et al., 2011; Holschuh et al., 2017; Delf et al.,




Figure 5. Radiostratigraphic continuity (ILCI) calculated over 10 airborne RES datasets acquired by BAS. Continuous and readily traceable IRHs are indicated in the slow-flowing regions of the ice sheet (high ILCI; bright yellow) whereas disrupted or absent IRHs are likely in the faster-flowing sections of ice streams or where subglacial topography is highly variable (low ILCI; dark purple). The background maps show ice-flow velocities from MEaSUREs (Rignot et al., 2017) and a hillshade of the bedrock from BedMachine (Morlighem, 2020). Figure modified from Frémand et al. (2022).

2020) and methods that use along-track phase information during SAR processing to estimate IRH slope (MacGregor et al., 2015a; Castelletti et al., 2019; Oraschewski et al., 2024).

3.4 Dating internal-reflection horizons (isochrones)

As introduced in Sect. 2, most RES-imaged IRHs have been shown to be isochronous, and the majority of those we treat in this review (i.e. that are imaged in between approximately the first and last few hundreds of metres of the ice column) arise due to the RES systems imaging variations in the electrical conductivity (i.e. acidic content) of the ice with depth. Hereafter in this paper, reiterating that most IRHs are isochrones, we will use the term isochrones to refer to IRHs, and will only re-use the term IRH where it may be ambiguous concerning whether IRHs are isochronous.

340 Ages can be assigned to isochrones at intersections with deep ice cores where age-depth models have already
341 been derived from chemistry analyses (e.g., McConnell et al., 2017; Cole-Dai et al., 2021; Bouchet et al., 2023),
342 but also using modelling techniques where this is not possible. Before any age can be assigned, the age
343 uncertainty that arises from the RES system itself must first be assessed. Uncertainty in reflector depth arises
344 from several sources: (a) proximity of the RES profile to the ice-core site, otherwise a specific reflector
345 geometry (typically flat) must be assumed between the point of closest approach and the ice-core site
346 (MacGregor et al., 2015a); (b) the radio-wave speed, which varies based on permittivity variations as a
347 function of englacial density and anisotropy (e.g., Kovacs et al., 1995; Fujita et al., 2000); (c) the range
348 resolution of the RES system and the signal-to-noise ratio of each traced reflection at (or near) the ice-core
349 site, which enable an estimate of the depth precision to which each traced reflection can be known (e.g.,
350 Cavitte et al., 2016); and (d) the picking accuracy of both the ice surface and the isochrones themselves,
351 which can add several metres of uncertainty. This latter point may include the uncertainty arising from the
352 source of the surface product (i.e. either from cm-resolution onboard altimeter/LIDAR), or directly from the
353 RES data which have much lower resolution of the order of several metres); and whether the picking
354 algorithm is tailored to extract the onset of the reflection, the half-amplitude, or the peak value.

355 The ideal scenario for assigning ages to isochrones is that a RES profile intersects or passes sufficiently close
356 (~500 m vicinity) to the location of an ice-core site for the ice core's depth-age scale (from chemical profiling
357 or layer counting) to be useable for directly assigning ages to the RES-imaged isochrones. In such cases, direct
358 dating at the ice core can be done in two ways: (1) direct age-matching by comparing the isochrone depth
359 from the RES to the age-depth scale of the ice core (e.g. MacGregor et al., 2015a; Cavitte et al., 2016; Bodart
360 et al., 2021), or (2) by matching observed RES isochrones to simulated RES isochrones based on the measured
361 dielectric profiling at the ice core (e.g. Eisen et al., 2003; Winter et al., 2017; Franke et al., 2025). The former
362 can be applied to any isochrones but results in larger uncertainties due to the conversion of the two-way-
363 travel time from the RES data to ice-equivalent depths, whereas the latter is more accurate as the
364 conductivity peaks represent the true physical origin of the isochrones. However, the condition that a
365 reflection must rise above the background noise of the RES system limits its application to specific isochrones
366 (Franke et al., 2025). For either case, the isochrone-depth uncertainty can then be combined with the ice-
367 core age uncertainty to assign a total age uncertainty to the mapped reflections; in these cases, uncertainty
368 is generally dominated by the ice-core-derived age uncertainty in the upper third of the ice column, while
369 the RES-derived depth uncertainty increasingly dominates at larger depth (e.g., MacGregor et al., 2015a;
370 Cavitte et al., 2016; Muldoon et al., 2018; Winter et al., 2019a; Wang et al., 2023). More recently, some
371 isochrones have been dated not by their direct intersection with an ice core, but rather by intersecting other
372 RES datasets that in turn have already been dated by their intersection with a distant ice core. As a result,
373 the age-depth profile is transferred to the new dataset at the crossover(s) between the intersecting RES
374 datasets (e.g., Ashmore et al., 2020; Bodart et al., 2021). In these cases, the relative uncertainties of the

different RES systems at the intersections between RES datasets additionally need be factored into the final age estimation, and the final age estimates are commonly checked using the modelling techniques introduced next (e.g., Bodart et al., 2021; Sanderson et al., 2024).

Where isochrones cannot be directly correlated to an ice-core age-depth relationship due to a lack of nearby ice cores, any intersections with previously dated isochrones, or missing sections in the record (e.g., due to disrupted englacial stratigraphy), age-depth modelling is required to assign ages to isochrones. This is typically done using 1-D models in stable parts of the ice sheet such as at ice divides (e.g., Nye, 1957; Dansgaard and Johnsen, 1969; Ashmore et al., 2020; Bodart et al., 2021; Sanderson et al., 2024); or using more complex multidimensional (2D/3D) models in areas with challenging ice-flow or bed conditions (e.g., Waddington et al., 2007; MacGregor et al., 2015a; Parrenin et al., 2017; Lilien et al., 2021).

4 RES data availability and quality for characterising Antarctica’s internal architecture

RES data have been acquired across Antarctica spanning six decades. An impression of the history can be gained from the periodic release of maps of subglacial topography, the first by Drewry (1975) and Drewry (1983; Antarctica Glaciological and Geophysical Folio Sheet 9), and then through the Bedmap series, now in its third iteration (Frémand et al., 2023, their Fig. 1; Pritchard et al., 2025). However, those maps and associated papers focus only on where the RES data were used to pick an echo at the bed, and do not provide information on whether the constituent surveys also imaged internal architecture. Here, therefore, we will outline the sequential development of airborne RES systems and surveys across Antarctica focussing on their attributes and availability for imaging and analysing internal architecture (Section 4.1). We will then briefly introduce some wide-ranging *ground-based* RES surveys that complement the overall database available for interrogating internal architecture (Section 4.2), and conclude with a “progress report” of existing traced radiostratigraphy across Antarctica (Section 4.3).

4.1 Evolution and availability of airborne RES for internal architecture

Airborne RES surveying across Antarctica has undergone three key technology-led developments as follows. Firstly, there has been a transition from initially recording RES data onto analogue tape recorders or film to inputting data directly into more sophisticated digital recording systems. Secondly, there has been the introduction of Global Navigation Satellite System (GNSS) systems to onboard flight navigation. Thirdly, there has been a progression from all datasets being acquired incoherently, wherein RES data receipt and recording were not phase-matched, to RES acquisition with coherent RES systems, for which the phase of the transmitted signal is preserved in a reference signal with which the received signal is compared, greatly increasing the clarity and quality of the imaging, which is especially pertinent for the analysis of internal architecture. The following subsections and Fig. 6 describe what may accordingly be termed three main eras of airborne RES surveying over the Antarctic ice sheets.




Figure 6. RES coverage and suitability for tracing internal architecture across Antarctica. (a) Airborne RES profiles acquired with analogue RES systems between 1966 and 2003. SPRI/NSF/DTU = Scott Polar Research Institute/National Science Foundation/Technical University of Denmark; BGR = Bundesanstalt für Geowissenschaften und Rohstoffe (Germany's Federal Institute for Geosciences and Natural Resources). (b) Airborne RES profiles acquired digitally but incoherently between 1991 and 2019. INGV = Istituto Nazionale di Geofisica e Vulcanologia (Italy's National Institute of Geophysics and Volcanology). (c) Airborne RES profiles acquired digitally and coherently since 2007. Univ. Kansas / CREsis / OIB = data acquired using systems designed by University of Kansas, often under the auspices of the USA's Centre for Remote Sensing and Integrated Systems and/or NASA's Operation IceBridge. (d) Map of all digitally-acquired airborne RES profiles (two shades of grey) and long-range ground-based RES profiles (coloured tracks) that collectively represent all data presently available for analysing internal architecture through Antarctica. The coverage of digitally-recorded incoherent RES (light grey) includes the originally analogue but now digitised SPRI/NSF/TUD data collected between 1971 and 1974. RES profile locations depicted in this map from Frémand et al. (2023, Tables S1 and S2). US/ITASE/St. Olaf = data acquired by radar system and operators from St. Olaf College, USA, often under the auspices of the International Trans-Antarctic Expedition (Mayewski et al., 2005). Also annotated in panel (d) are key locations mentioned in this paper, with yellow triangles marking key deep-ice-core sites. The yellow dashed line marks the nominal divide between the West Antarctic and

409 **4.1.1 Analogue data recording**

410 Before the 2000s, the majority of airborne RES data across Antarctica were recorded onto analogue tape
411 recorders or film, and most of these data, including all data acquired prior to the 1990s, were acquired
412 without GNSS being available for accurate navigation. Due to the analogue recording, all such systems
413 recorded data incoherently. The coverage of these analogue surveys is shown in Fig. 6a.

414 Continent-spanning RES surveying was pioneered across West Antarctica and around half of East Antarctica
415 by the UK-based Scott Polar Research Institute (SPRI), with key logistical support from the USA's National
416 Science Foundation and, from 1971, using antennas designed and installed by engineers from the Technical
417 University of Denmark (DTU), which fundamentally improved the reflection of IRHs (Swithinbank, 1969; Evans
418 and Smith, 1970; Gudmandsen et al., 1975; Drewry, 2023). Also through the late 1960s and 1970s, and
419 continuing through the 1980s, the Soviet Antarctic Expedition conducted airborne surveying across parts of
420 East Antarctica (Popov, 2020) (Fig. 6a). Both the SPRI-NSF-TUD and Soviet campaigns used a 60 MHz centre-
421 frequency RES system, inspiring the British Antarctic Survey (BAS) to adopt the same centre frequency as it
422 commenced progressive regional RES surveys through the 1980s (Fig. 6a).

423 The 1990s saw a step-change improvement in flight navigation and concomitant RES-dataset positioning with
424 the introduction of GNSS. The 1990s was also a period of transition, with some RES operators developing the
425 capacity to acquire data digitally and coherently (as expanded in Sections 4.1.2 and 4.1.3), but other major
426 data providers, such as the British Antarctic Survey and Russian (formerly Soviet) groups continuing to record
427 data only in analogue format, hence also incoherently. These analogue surveys, supplemented by some
428 regional surveys across East Antarctica over the same era by groups from Australia (e.g., Morgan et al., 1982)
429 and Germany (e.g., Thyssen and Grosfeld (1988); Damaske and McLean (2005)), are depicted in Fig. 6a.

430 **4.1.2 Digital data recording, incoherent RES**

431 Digital recording systems were first implemented in Antarctic airborne RES surveying in the early 1990s by
432 the USA-based University of Texas Institute of Geophysics (UTIG). UTIG's surveys through the 1990s,
433 principally of West Antarctica, used adapted versions of the 60 MHz SPRI-NSF-DTU incoherent RES system
434 (Blankenship et al., 2001; Carter et al., 2007; Young et al., 2016). In the mid-1990s, national operators from
435 Germany and Italy also commenced digital acquisition of incoherent RES surveys across parts of East
436 Antarctica, both using RES systems operating around a higher centre frequency of 150 MHz (Steinhage et al.,
437 2001; Eisen et al., 2007; Tabacco et al., 2008; Zirizzoti et al., 2008). The full coverage of airborne RES data
438 acquired digitally with incoherent RES systems is depicted in Fig. 6b. It includes data acquired broadly from
439 the early 1990s to the mid 2000s, although some operators continued to acquire data incoherently into the
440 mid 2010s.

441 **4.1.3 Digital data recording, coherent RES**

442 Fully coherent RES systems were first operated in Antarctica by the USA-based University of Kansas on a joint
443 USA (NASA; National Aeronautics and Space Administration) / Chile (CECs; Centro de Estudios Científicos)
444 mission to survey fast-changing regions of West Antarctica (Rignot et al., 2004). In 2005, Kansas became host
445 to the USA's Center for Remote Sensing and Integrated Systems (previously Center for Remote Sensing of Ice
446 Sheets; CReSIS), an NSF-designated national Science and Technology Centre with a focus on ice-sheet
447 sounding; and began to operate an upgraded series of deep-looking RES systems with centre frequencies of
448 ~190-194 MHz named Multichannel Coherent Radar Depth Sounders (MCoRDS), which were deployed widely
449 across Antarctica between 2009 and 2019 as part of NASA's Operation IceBridge programme (Fig. 6c;
450 Rodriguez-Morales et al., 2013; MacGregor et al., 2021).

451 From 2004 onwards, both BAS and UTIG transitioned their RES systems into coherent data acquisition, with
452 this year also marking BAS' first implementation of digital data recording. BAS' Polarimetric Radar Airborne
453 Science Instrument (PASIN) has a centre frequency of 150 MHz, and transmits two waveforms, a narrow pulse
454 (0.1 μ s) for detecting shallow radiostratigraphy in the upper 2 km of the ice column, and a deep-sounding
455 chirp (4 μ s) for detecting deeper radiostratigraphy and the bed (Corr et al., 2007; Hélière et al., 2007; see Fig.
456 2 for examples of each). PASIN was upgraded in the mid-2010s to enable the acquisition of swaths (i.e. wide
457 strips) of RES data to map the ice-sheet bed (Arenas-Pingarrón et al., 2023). UTIG integrated a coherent 60
458 MHz centre-frequency RES system (Moussessian et al., 2000) with radio-frequency hardware from DTU to
459 allow high-power coherent recording, which enabled synthetic-aperture-radar (SAR) processing of acquired
460 data (Peters et al., 2005; Peters et al., 2007; refer back to Sect. 3.1 for description of SAR processing). This
461 initial High-Capability Radar Sounder (HiCARS) system was translated to commercially available components
462 (HiCARSII) which were incorporated from the mid-2010s into the subsequent Multifrequency Airborne Radar-
463 sounder for Full-phase Assessment (MARFA), capable of cross-track interferometry for clutter discrimination
464 (Castelletti et al., 2017; Scanlan et al., 2020).

465 Other airborne RES operators transitioned into coherent RES across Antarctica during the mid-2010s.
466 Alongside continuing surveying by BAS, CReSIS and UTIG, large volumes of coherent RES data have been
467 acquired across Antarctica over the last decade by the Russian Antarctic Expedition (Popov, 2020; Popov,
468 2022), Germany's AWI using an improved version of CReSIS' MCoRDS system (Humbert et al., 2018; Karlsson
469 et al., 2018; Winter et al., 2019a; Wang et al., 2023; Franke et al., 2025), as well as the Polar Research Institute
470 of China (PRIC; Cui et al., 2020ba; 2020b; 2020c) and the Korean Polar Research Institute (KOPRI; Lindzey et
471 al., 2020; Lee et al., 2021) (Fig. 6c).

472 **4.2 Ground-based RES datasets**

473 Since the 1960s, groups from at least twelve institutions have acquired ground-based RES datasets focussed
474 on sounding Antarctica's subglacial bed and have also typically imaged internal architecture in the process.
475 Typically, ground-based surveys have been confined to smaller regions or shorter profiles than covered by
476 the airborne RES surveys, befitting the more common application of ground-based RES to detailed site
477 surveys in preparation for retrieving ice cores, or for accessing the ice bed or subglacial lakes (e.g., Frezzotti
478 et al., 2004; Laird et al., 2010; Christianson et al., 2012; Ross et al., 2020). From these surveys, several local
479 radiostratigraphies have been published (e.g., Eisen et al., 2005; Jacobel and Welch, 2005; Koutnik et al.,
480 2016; Cavitte et al., 2023; Chung et al., 2023). These detailed studies provide invaluable seeding points for
481 extending radiostratigraphies much more widely across the ice sheets (e.g., Winter et al., 2019a) and for
482 understanding better ice-sheet history and glaciological processes.

483 Supplementing the more local surveys, some ground-based profiles have been acquired over traverses of
484 multiple 100s of km over the Antarctic ice sheets, and these traverses, marked on Fig. 6d, merit special
485 attention as potential resources for analysing pan-continental radiostratigraphy. A particularly extensive
486 programme of ground-based surveys has been conducted since 1969 by the Japanese Antarctic Research
487 Expedition (JARE) connecting coastal East Antarctica in Dronning Maud and Enderby Land to Dome F, with
488 data from some of these traverses conducted in the 1990s underpinning seminal work on the origins of IRHs
489 (Fujita et al., 1999; Matsuoka et al., 2003). Today, data from JARE represent some of the most spatially
490 extensive of Antarctica's ground-based RES datasets and a rich repository of internal architecture (Fujita et
491 al., 2011; Van Liefferinge et al., 2021; Tsutaki et al., 2022). Further long ground-based RES traverses were
492 acquired by several national and international teams in the 2000s under the auspices of the International
493 Trans-Antarctic Scientific Expedition (ITASE). RES profiles containing particularly rich internal architecture
494 were acquired by the USA-NSF's ITASE traverses across both West (Welch and Jacobel, 2003; Jacobel and
495 Welch, 2005) and East Antarctica (Welch et al., 2009), with findings from Arcone et al. (2012a) suggesting
496 that in some parts of East Antarctica the radiostratigraphy is unconformable and may present significant
497 challenges to tracking radiostratigraphy.

498 **4.3 Progress report**

499 **4.3.1 Summary of available data**

500 Fig. 6d shows the coverage of RES profiles acquired across Antarctica that are, in principle, available for
501 developing a pan-Antarctic stratigraphy. This includes all RES data acquired digitally, whether coherent or
502 incoherent. Generally, RES data that are acquired coherently are the highest quality for tracing unbroken
503 IRHs over long distances, but incoherent data can also be utilised for IRH tracing. Fig. 6d therefore also only
504 includes data that were acquired by GNSS navigation, with one exception: the 1970s SPRI-NSF-TUD data are

505 included because they imaged IRHs across Antarctica with a clarity that rivals many modern RES surveys and
506 form potentially vital links across otherwise unsurveyed regions, and consequently have been “revived” by a
507 dedicated fine-resolution digitisation and distribution programme (Schroeder et al., 2019; Schroeder et al.,
508 2022). Navigational uncertainties inherent to pre-GNSS navigation remain with these data (of the order of
509 several kms), but there is a prospect of using crossovers with more modern datasets to reconstruct the
510 navigation with improved accuracy (Teisberg and Schroeder, 2023).

511 **4.3.2 Existing dated radiostratigraphy across Antarctica**

512 Prior to the inception of *AntArchitecture* in 2018, several studies had produced radiostratigraphies spanning
513 the last 17.5 ka across West Antarctica and 352 ka for East Antarctica (e.g., Hodgkins et al., 2000; Siegert and
514 Hodgkins, 2000; Siegert, 2003; Siegert and Payne, 2004; Jacobel and Welch, 2005; Leysinger Vieli et al., 2011;
515 Steinhage et al., 2013; Karlsson et al., 2014; Wang et al., 2016). However, the spatial extents of these
516 radiostratigraphies were relatively limited. Through *AntArchitecture*, a more coordinated and focused
517 approach to characterising Antarctic radiostratigraphy has been conducted, as depicted in Fig. 7 and detailed
518 in Table 1. This programme has facilitated the recovery and characterisation of several isochrones with ages
519 up to 25 ka across much of the Amundsen and Weddell Sea sectors of West Antarctica (Muldoon et al., 2018;
520 Ashmore et al., 2020; Bodart et al., 2021; Bodart et al., 2023). Over East Antarctica, a much older record has
521 been extracted, owing to the more stable and slow-flowing ice conditions in the area, including isochrones
522 dating back to the last 705 ka (Cavitte et al., 2016; Winter et al., 2019a; Beem et al., 2021; Cavitte et al., 2021;
523 Chung et al., 2023; Wang et al., 2023; Sanderson et al., 2024; Franke et al., 2025).

524 A notable finding is the presence of widespread and ubiquitous isochrones that have been imaged by
525 different RES systems and are found in several ice-core records. Across West Antarctica, the most prevalent
526 isochrone, dated precisely and independently at Byrd and WAIS Divide ice cores to ~4.7 ka, has been
527 identified by several studies (Jacobel and Welch, 2005; Karlsson et al., 2014; Holschuh et al., 2018; Muldoon
528 et al., 2018; Ashmore et al., 2020; Bodart et al., 2021; Bodart et al., 2023). There is evidence that this same
529 isochrone may also be found widely across East Antarctica, based on sulphate concentrations in ice cores and
530 findings from individual RES surveys across the region (Steinhage et al., 2013; Winski et al., 2019; Beem et al.,
531 2021; Cole-Dai et al., 2021; Sigl et al., 2022). Additionally, across much of the West Antarctic Ice Sheet an
532 isochrone dated at 17.5 ka has been observed in both ground-based and airborne RES data (Jacobel and
533 Welch, 2005; Muldoon et al., 2018; Bodart et al., 2021). This 17.5 ka RES isochrone has been identified and
534 linked to an eruption from West Antarctica’s Mount Takahe in both the Byrd (Hammer et al., 1997) and WAIS
535 Divide (McConnell et al., 2017) ice cores. Over East Antarctica, packages of closely spaced isochrones of ages
536 ~38 ka, ~73 ka, ~128 ka, ~160 ka, and ~170 ka have been traced from ice cores (Leysinger Vieli et al., 2011;
537 Winter et al., 2019a; Cavitte et al., 2021; Wang et al., 2023; Sanderson et al., 2024; Franke et al., 2025);
538 notably, the ~73 ka isochrone has been linked by ice-core profiling to the Toba Eruption in Indonesia




Figure 7. Existing open-access dated stratigraphies across Antarctica obtained from the Digital Object Identifiers (DOIs) provided in Table 1, with RES profiles for Bedmap-2 and Bedmap-3 products shown in the background (grey; Frémand et al., 2023). Existing deep ice cores (defined here as ice cores that have been drilled to near the ice-bed interface and that provide a multi-millennial record) are shown as red triangles. (a) Maximum number of isochrones traced through each dataset (from 1 to >8); see Table 1 for full details (b-c) Age of the deepest (oldest) isochrone across each dataset for the WAIS (b) and EAIS (c) regions respectively. Note that the scale used for (c) is logarithmic.

539
 540 (Svensson et al., 2013). Together, such distinct isochrones, imaged by and from multiple RES systems and
 541 platforms, provide important regional or continental-wide time markers, equivalent to Greenland’s highly
 542 recognisable “three sisters” (Fahnestock et al., 2001a; MacGregor et al., 2015a; 2025) for inferring past
 543 changes at specific time intervals.

544 Despite the advances discussed here, the established radiostratigraphy across the Antarctic ice sheets
 545 currently represents only a small subset of the total available RES data (Fig. 7, and refer back to Sect. 4.3 and
 546 Fig. 6). The establishment of the *AntArchitecture* community, and its commitment to establish protocols for
 547 sharing and processing internal architecture across the multiple datasets, is expected to facilitate further
 548 isochrone tracing, which will in turn contribute to the development of the first three-dimensional age-depth
 549 model of the ice sheet.

Table 1. Inventory of expansive radiostratigraphic datasets for the Antarctic ice sheets, ordered by region and cumulative distance of dataset (cumulative distance here corresponds to the total km of isochrone profiles summed for each isochrone where the data are openly available (via use of a DOI), or approximated value where datasets are not publicly available). Openly available datasets are shown in Figure 7; locations of ice cores are marked on Fig. 6d. Data-provider acronyms provided at foot of table; in most cases we also list here a specific project acronym for each survey which can be cross-referenced through the reference and/or dataset listed in each row.

(For EGU sphere formatting, this 10-column table is presented across two pages.)

Region	Survey dates	Data provider (acronyms expanded at foot of table)	Survey name / acronym	Ice-core intersection(s)	No. of traced isochrones
EAIS	1996 – 2023	AWI / CRISIS	EPICA / DML	Kohnen / Dome F	9
EAIS	2016 - 2017	AWI	Beyond EPICA Dome Fuji	Dome F	7
EAIS	1998 - 2008	AWI / CRISIS	DoCo / EPICA / AGAP	Kohnen / Dome F / Vostok / Dome C	5
EAIS	2007 - 2016	BAS	AGAP / PolarGap	South Pole	3
EAIS	1974 – 1979	SPRI-NSF-DTU	-	Vostok / Dome C	12
EAIS	2016 – 2018	BAS	Beyond EPICA Little Dome C	Dome C	20
EAIS	2008 - 2018	UTIG	ICECAP	Dome C	26
EAIS	1974 – 1979	SPRI/NSF/DTU	-	Vostok	15
EAIS	1974 – 1979	SPRI/NSF/DTU	-	Vostok / Dome C / Dome A / South Pole	15
EAIS	2016 – 2017	PRIC	South Pole Corridor	South Pole	8
EAIS	2002 – 2003	AWI	-	Kohnen / Dome F	8
EAIS	2019 – 2020	UA / AWI	Beyond EPICA Little Dome C	Dome C	19
EAIS	2004 – 2005	PRIC	Dome A (CHINARE-21)	Vostok	6
WAIS	2004 – 2018	BAS / CRISIS	BBAS / OIB	WAIS Divide	4
WAIS	1991 – 2014	UTIG	CASERTZ / SOAR / AGASEA / GIMBLE	Byrd / WAIS Divide	1
WAIS	2010 – 2011	BAS	IMAFI	-	3
WAIS	1977 – 1978	SPRI-NSF-DTU	-	Byrd	5
WAIS	2000 – 2001	NSF	ITASE	Byrd	1

Table 1 continued: Columns 6-10.

Isochrone age range (ka)	Cumulative distance of traced IRHs (km)	Dataset Reference	Dataset DOI
4.8 – 91.0	203 500	Franke et al. (2025)	10.1594/PANGAEA.973266
31.4 – 232.7	110 000	Wang et al. (2023)	10.1594/PANGAEA.958462
38.0 – 161.0	40 000	Winter et al. (2019a)	10.1594/PANGAEA.895528
38.0 – 162.0	30 400	Sanderson et al. (2024)	10.5285/cfab639-991a-422f-9caa-7793c195d316
17.5 – 352.4	30 245	Leysinger Vieli et al. (2011)	10.1029/2010JF001785
10.5 – 414.6	24 820	Chung et al. (2023)	10.1594/PANGAEA.963470
10.0 – 705.0	15 500	Cavitte et al. (2021)	10.15784/601411
17.0 – 211.0	15 000	Leysinger Vieli et al. (2004)	-
45.9 – 169.7	13 300	Siegert (2003)	-
4.7 – 93.9	12 100	Beem et al. (2021)	10.15784/601437
4.7 – 72.4	9 700	Steinhage et al. (2013)	-
73.7 - 476.4	3 000	Chung et al. (2023)	10.1594/PANGAEA.957176
34.3 – 161.4	1 300	Wang et al. (2016)	-
2.3 – 16.5	30 700	Bodart et al. (2021)	10.5285/f2de31af-9f83-44f8-9584-f0190a2cc3eb
4.7	19 000	Muldoon et al. (2018)	10.15784/601673
1.9 – 8.1	13 700	Ashmore et al. (2020)	10.5281/zenodo.4945301
0.8 – 16.0	2 400	Siegert and Payne (2004)	10.1002/esp.1238
17.5	1 000	Jacobel and Welch (2005)	10.7265/N5R20Z9T

Data providers:

AWI = Alfred-Wegener Institute, Germany

BAS = British Antarctic Survey, UK

CRISIS = Centre for Remote Sensing and Integrated Systems, USA

NSF = National Science Foundation, USA

PRIC = Polar Research Institute of China

SPRI/NSF/DTU = Scott Polar Research Institute/National Science Foundation/Technical University of Denmark

UA = University of Alabama, USA

UTIG = University of Texas Institute of Geophysics, USA

551

552

5 Applications of internal architecture to wider Antarctic science

553

554

555

556

557

558

559

Here, we now review to what scientific purposes internal architecture has already been exploited. Sect. 5.1 to 5.4, supported by Fig. 8, exemplify four primary applications of RES-imaged isochrones, Sect. 5.5 explores the scientific applications of other forms of internal architecture, and Sect. 5.6 discusses how radiostratigraphic data have been incorporated into numerical modelling, and their use in calibrating ice-sheet models of varying complexity. This section contextualises the following Sect. 6 which then suggests priorities for future research that will be enabled as Antarctica's internal architecture, and particularly its radiostratigraphy, continue to be explored and made available.

560 **5.1 Radiostratigraphy and ice cores**

561 Ice cores from Antarctica provide fundamental palaeoclimate records (e.g., EPICA Community Members,
562 2004; WAIS Divide Project Members, 2015). The layering found in ice cores is also visible in radiostratigraphy,
563 as a function of the RES-system resolution (Section 2), and we have already introduced the concept that RES
564 records tied to existing ice cores provide a basis for extending these “point-source” age-depth chronologies
565 into 3-D age-depth fields that extend widely across the Antarctic ice sheets (cf. Sect. 3.4 and 4.3.2).
566 Conversely, RES-imaged radiostratigraphy can be used to guide the best locations for recovering future ice
567 cores. Accumulation rate, ice dynamics and age-depth relationships extracted from isochrones have
568 previously informed the appropriateness of coring sites (e.g., Neumann et al., 2008; Parrenin et al., 2017;
569 Beem et al., 2021; Wang et al., 2023) and have been essential for pre-site survey of potential future ice coring,
570 e.g. for the *Oldest Ice* endeavour of the International Partnerships for Ice Core Sciences (IPICS; e.g., Fischer
571 et al., 2013; Van Liefferinge and Pattyn, 2013; Karlsson et al., 2018; Lilien et al., 2021; Chung et al., 2023).

572 Radiostratigraphy has also provided opportunities for synchronising and reducing uncertainties in ice-core
573 chronologies by facilitating the direct tracing of isochrones between two or more ice cores in order to
574 correlate ice-core chronologies (as achieved for the Greenland Ice Sheet by MacGregor et al., 2015a; see Fig.
575 8b). In Antarctica, previous studies that have used isochrones to correlate chronologies between ice cores
576 include Siegert et al. (1998), Steinhage et al. (2013), Cavitte et al. (2016), Le Meur et al. (2018) and Winter et
577 al. (2019a) for East Antarctica, and Muldoon et al. (2018) for West Antarctica. These studies have provided
578 confidence that ice cores obtained from locations separate by 100s of km capture analogous variations in
579 palaeoclimate at regional scales, and that the signals recorded by RES correspond to genuine physical
580 variations in the ice (typically variations in electrical conductivity, often related to fallout from past volcanic
581 eruptions; as noted in Sect. 4.5).

582 The key challenge in synchronising ice-core records between distant sites using RES has been in resolving the
583 radiostratigraphically- and ice-core-derived chronologies between each ice-core site, given the order-of-
584 magnitude difference in resolution of chronologies recoverable from RES (on the order of metres) versus ice-
585 core records (on the order of centimetres). This has typically been dealt with using forward modelling based
586 on electrical-conductivity measurements or dielectric profiling of the ice cores to provide a transfer function
587 (e.g., Miners et al., 1997; Hempel et al., 2000; Eisen et al., 2003; Eisen et al., 2006; Winter et al., 2017;
588 Mojtabavi et al., 2022), or by adopting Bayesian frameworks which provide a probability distribution of the
589 age of the isochrones (Muldoon et al., 2018). Thus, while the age-depth fields compiled from isochrones will
590 never match the precision and accuracy of ice-core age-depth relationships (MacGregor et al., 2015a; Winter
591 et al., 2017), they provide the spatial context that ‘point-source’ ice cores cannot. Through isochrone-
592 constraint modelling (see Sect. 5.6), the age of the ice and its spatial distribution can be more effectively
593 constrained in regions distant from the current drilling sites (Born and Robinson, 2021; Sutter et al., 2021).




Figure 8. Schematic illustration of radiostratigraphic observations within an ice sheet and their scientific applications; (a), in the centre, depicts typical ice-sheet locations for applications shown in subsequent panels. (b) Connecting and validating ice cores in Greenland (after MacGregor et al., 2015a). (c) Imaging intersections of IRHs with ice surface in region of surface wind scouring (after Winter et al., 2016). (d) Using isochrones to calculate basal melting across Subglacial Lake Vostok (after Siegert et al., 2001). (e) Using isochrone drawdown to locate region of elevated geothermal heat flux near South Pole (after Jordan et al., 2018). (f) Application of “Internal Layering Continuity Index” (ILCI) to quantify disruption (folding/warping) to otherwise continuous isochrones (after Bingham et al., 2015). (g) Using intersecting RES profiles to explore ice anisotropy (after Gerber et al., 2023). (h) Raymond Arch imaged in near-surface (top panel) and deep RES across Derwael Ice Rise, Dronning Maud Land (after Drews et al., 2015). (i) Basal-ice units and suggested accreted basal ice in East Antarctica (after Bell et al., 2011). (j) Basal crevasses imaged in West Antarctica and used to date regrounding of previously floating ice (after Kingslake et al., 2018). (k) Prominent tephra horizon imaged by RES across Pine Island Glacier, West Antarctica (after Corr and Vaughan, 2008).

595 In marginal locations of the ice sheets, or around nunataks, where persistent pronounced surface scouring is
596 co-located with upward ice flow over subglacial topography – i.e., in regions of so-called “blue ice” – very old
597 ice may outcrop obliquely to the ice surface and hence allow the recovery of a “horizontal ice core” along
598 the ice surface (Spaulding et al., 2013). Dated isochrones have been used to trace the age-depth model
599 recovered from horizontal ice cores back into the ice sheet (Reeh et al., 2002; Siegert et al., 2003a; Winter et
600 al., 2016; Fogwill et al., 2017; Baggenstos et al., 2018; see Fig. 8c). However, shearing and folding can disrupt
601 the stratigraphic order of the outcropping IRHs, rendering the interpretation of their radiostratigraphy more
602 complex than for most vertical ice cores.

603 **5.2 Surface mass balance**

604 Successive snowfall events create a record of progressively buried isochrones which can be observed in
605 radargrams. In slow-flowing ice and especially around ice divides, the depth of isochrones is largely controlled
606 by surface mass balance and therefore dated radiostratigraphy has made it possible to reconstruct past
607 surface mass balance over millennial timescales across spatially extensive regions (e.g., Nereson et al., 2000;
608 Siegert, 2003; Siegert and Payne, 2004; Eisen et al., 2005; Waddington et al., 2007; Neumann et al., 2008;
609 MacGregor et al., 2009; Leysinger Vieli et al., 2011; Karlsson et al., 2014; Koutnik et al., 2016; Cavitte et al.,
610 2018; Bodart et al., 2023). Such records have fundamentally informed us about how mass balance has
611 changed with time over past millenia, for example showing that accumulation rates changed significantly
612 over central (Siegert and Payne, 2004; Neumann et al., 2008; Koutnik et al., 2016; Bodart et al., 2023) and
613 coastal (Karlsson et al., 2014) West Antarctica throughout the Holocene. Typically, vertical strain rates must
614 be corrected for the whole ice column, particularly in regions of (present or past) fast flow, or there is a need
615 to account for basal processes such as enhanced basal melting (e.g., Leysinger Vieli et al., 2011; Chung et al.,
616 2023), because in such cases the isochrone depths will be dynamically modified and therefore will not
617 represent the surface mass balance at the time of deposition (e.g., Koutnik et al., 2016). Where the
618 radiostratigraphy has not been impacted significantly by strain, the shallow-layer approximation can be
619 applied, which allows us to ignore these strain-rate corrections (Waddington et al., 2007). If horizontal
620 advection influences the stratigraphy 2D, 2.5D or 3-D modelling is required (see Sect. 5.6).

621 Regions of unconformable radiostratigraphy occurring throughout the ice column in parts of Antarctica have
622 partly limited the extent to which some surface mass balance records could be more widely extrapolated
623 (Arcone et al., 2012b; Cavitte et al., 2016). RES surveys of the upper ~100 m of the ice column in the affected
624 regions typically reveal widespread conformal, annual horizons modified by local variations in accumulation
625 or ice flow (Eisen, 2008), and the majority of them have been ascribed to wind scouring out surface deposits
626 and forming “megadunes” (Das et al., 2013; Traversa et al., 2023) that then become progressively buried as
627 sets of unconformable IRHs. Studies have identified such unconformities in several locations in East

628 Antarctica (Welch and Jacobel, 2005; Traversa et al., 2023) and West Antarctica (Woodward and King, 2009;
629 Holschuh et al., 2018).

630 **5.3 Basal melting and geothermal heat flux**

631 The presence of a subglacial water body or enhanced geothermal heat flux draws isochrones down towards
632 the ice base. Exploiting this principle, isochrones have been used to calculate melting at the base of the ice.
633 Mismatches between surface-accumulation-driven modelled isochrones and traced isochrones have been
634 used to infer regions of enhanced basal melting in Greenland (Dahl-Jensen et al., 1997; Fahnestock et al.,
635 2001b) and Antarctica (Carter et al., 2009) on the principle that removal of ice at the base by basal melting
636 thins annual layers above. However, for locating areas of enhanced geothermal heat flux (or subglacial lakes,
637 which may sometimes owe their existence to enhanced geothermal heat flux) researchers now typically rely
638 more on analysing the reflectivity or specularity of the ice-bed echo in RES data (e.g., Young et al., 2016; Chu
639 et al., 2021), and only use isochrones to guide derivations of basal melting where such more direct data are
640 lacking.

641 Isochrones have been analysed in more detail over parts of Antarctica to constrain basal melting in more
642 localised settings. For example, Siegert et al. (2000) used deviations in the dip of deep isochrones away from
643 parallelism with the ice-bed/subglacial-lake surface over Subglacial Lake Vostok to calculate basal melting
644 and water exchange between the lake and the overlying ice sheet (Fig. 8d). Jordan et al. (2018) identified
645 isochrones dipping towards the bed ~200 km from the South Pole (Fig. 8e), and used these to model how
646 much basal melt would be required to draw the isochrones down towards the bed. By assuming that minimal
647 frictional melting would be generated by the slow ice flow in this region, they showed that the most likely
648 cause of the isochrones being drawn down towards the bed must be enhanced geothermal heat flux in this
649 region. Ross and Siegert (2020) undertook a detailed survey of isochrone geometry over Subglacial Lake
650 Ellsworth, West Antarctica, and showed that the isochrones were preferentially drawn down over the NW
651 shoreline of the lake, rather than the lake itself. This conclusion was in agreement with the pattern of basal
652 mass balance derived from previous numerical modelling of water circulation in the lake and indicated very
653 high basal melting of ~16 cm a⁻¹ on its northern shoreline.

654 **5.4 Ice-flow dynamics**

655 Moving ice causes IRHs that were originally deposited flat at the surface to deform through folding, tilting
656 and disruption. Therefore, deformed isochrones may be analysed to interpret past ice-flow dynamics.
657 Present-day (last ~35 years) information on ice-flow dynamics is derived from satellite monitoring of ice-
658 surface flow (Rignot et al., 2017), but to understand fully where and how ice-flow dynamics have changed
659 over the past several thousand years, and hence may be likely to do so again, researchers have interrogated
660 how changes to ice-flow dynamics have been imprinted into the RES-imaged internal architecture. The most

661 common methodology has been to explore and classify where the radiostratigraphy diverges from relatively
662 flat isochrones to profiles that show folding (a.k.a. buckling, warping or disruption) of the isochrones (Fig. 8f).
663 Wherever there is folding of isochrones, it is an indication that the ice has experienced considerable strain,
664 often as a result of flowing around or over significant bedrock obstacles (Robin and Millar, 1982; Hindmarsh
665 et al., 2006; Tang et al., 2022) or becoming variously stretched and compressed as it flows through an ice-
666 stream onset region or through ice-stream shear margins (Jacobel et al., 1993; Bell et al., 1998; Ng and
667 Conway, 2004; King, 2011). Overall, isochrone folding can indicate convergent ice flow, anisotropic rheology,
668 basal freeze-on, basal sliding, non-negligible transverse velocity gradients, or the abutting of units of
669 contrasting rheology. Importantly, the signature recorded by these processes is often advected downstream,
670 so that where it is observed does not necessarily indicate where the folding took place (Weertman, 1976;
671 Jacobel et al., 1993; Leysinger Vieli et al., 2004; NEEM Community Members, 2013; Wolovick et al., 2014;
672 Bons et al., 2016; Leysinger Vieli et al., 2018; Ross et al., 2020; Franke et al., 2021; Jennings and Hambrey,
673 2021; Jansen et al., 2024). In certain cases, relict folds that do not correspond to the current ice-flow direction
674 indicate a past change in ice-flow direction (Conway et al., 2002; Siegert et al., 2004; Rippin et al., 2006;
675 Franke et al., 2022).

676 While, therefore, there are multiple origins for isochrone folding, their geographical association with fast ice
677 flow has led to their presence being used as a broad diagnostic of the long-term stability (or otherwise) of ice
678 flow around Antarctica (e.g., Rippin et al., 2003; Siegert et al., 2003b; Bingham et al., 2007; Karlsson et al.,
679 2009; Ross et al., 2011; Bingham et al., 2015; Winter et al., 2015; Sanderson et al., 2023). In areas where
680 isochrones are strongly disrupted by (past or present) enhanced flow, extracting ILCI or isochrone-slope
681 products from the radiostratigraphy (as introduced in Sect. 3.3) has helped to complement reconstructions
682 of past or present ice-flow dynamics (e.g., Karlsson et al., 2012; Bingham et al., 2015; Holschuh et al., 2017;
683 Ashmore et al., 2020; Luo et al., 2020; Sanderson et al., 2023). In some cases, sequences of folded isochrones
684 have been observed beneath sequences of conformable isochrones, indicative of a past sudden change from
685 fast to slow ice flow (e.g., Conway et al., 2002; Siegert et al., 2013; Kingslake et al., 2016). To obtain more
686 complex information on past ice-dynamic changes falls into the realm of applying numerical modelling, which
687 is taken up in Sect. 5.6.

688 An important outcome of most ice flow is that the ice crystals themselves develop a preferred orientation,
689 typically termed anisotropic crystal-orientation fabric, which may then influence the direction-dependent
690 propagation speed of radio waves through ice (Gow and Williamson, 1976; Robin and Millar, 1982; Fujita et
691 al., 1999; Matsuoka et al., 2003; Eisen et al., 2007; Drews et al., 2012; Jordan et al., 2020; Jordan et al., 2022).
692 Studies have reconstructed and constrained the mechanical anisotropy of ice and histories of ice deformation
693 by calculating the travel-time difference for IRHs across intersecting RES profiles where the radio waves have
694 been polarised in different directions (e.g., Fig. 8g; Ershadi et al., 2022; Jordan et al., 2022; Gerber et al., 2023;
695 Zeising et al., 2023). A special case of isochrone folding due to changes in ice-crystal fabric occurs at ice

divides, where upward-pointing folds termed Raymond Arches (Fig. 8h) form due to the interplay of the strain-rate dependence of ice viscosity, which leads to stiffer ice beneath the divide, slowing isochrone thinning down relative to the flanks (Raymond, 1983; Vaughan et al., 1999; Martín et al., 2009; Hindmarsh et al., 2011; Matsuoka et al., 2015). The special geometry of these isochrone arches has been used to infer local ice-flow history including the onset of divide flow (Conway et al., 1999; Kingslake et al., 2016), divide migration (Nereson et al., 1998; Martín et al., 2009; Schannwell et al., 2019) and ice-thickness changes (Drews et al., 2015). With stable ice-divide positions over extended periods of time, these arches can evolve further into double-peaked Raymond Arches, as observed (Drews et al., 2013) and simulated by incorporating anisotropy into the ice-flow models (Pettit et al., 2007; Martín and Gudmundsson, 2012; Martín et al., 2014). In terms of efforts to trace isochrones widely across the Antarctic ice sheets, Raymond Arches have the greatest relevance in how they affect site selection for deep ice cores that are ideally used to assign ages to Antarctic-wide isochrones (as introduced in Sect. 3.4). The relative thinness of isochrones at the apex of Raymond Arches implies that better resolution age-depth records reaching further back in time would be obtained around the flanks, rather than on the apexes, of ice divides where arches are present.

5.5 Applications of internal architecture complementary to radiostratigraphy

Ice located near to the bed of an ice sheet is typically expected to have undergone strong deformation due to shear, or to originate from processes other than earlier surface accumulation. The basal ice of Antarctica and Greenland is typically characterised by an echo-free or low-backscatter zone lacking coherent layered reflections, termed an *echo-free zone* (EFZ) in early observations (Drewry and Meldrum, 1978; Robin and Millar, 1982; Fujita et al., 1999). With modern RES systems, this zone now appears as a basal unit in which IRHs are often warped, folded and pinched out, and consequently lack coherent reflections (Drews et al., 2009), but even without traceable radiostratigraphy this architecture contains useful information about ice properties and origins. With the progressive enhancement of RES-system range resolution, a variety of reflection sub-units distinctly standing out from the otherwise low-backscatter zone have been identified (e.g., Fig 8i; Bell et al., 2011; Bell et al., 2014; Wrona et al., 2018; Ross et al., 2020; Lilien et al., 2021; Franke et al., 2024b). Some of these features manifest as zones with nearly continuous high backscatter spanning several hundred metres in thickness. Some features drape over mountainous subglacial regions (e.g., in Antarctica's Gamburtsev Mountains and Jutulstraumen drainage basin; Bell et al., 2011; Wrona et al., 2018; Franke et al., 2024b), while others build plume-like structures within the cores of englacial folds (e.g., in northern Greenland and Antarctica's Institute Ice Stream; Bell et al., 2014; Ross et al., 2020). These basal units are likely of different origins and exhibit different dielectric properties compared to their low-backscatter surroundings, offering insights into potential formation mechanisms. Current hypotheses include strong deformation on the micro-scale by ice dynamics (Drews et al., 2009), freeze-on of subglacial water at the ice base (Bell et al., 2011; Creyts et al., 2014; Leysinger Vieli et al., 2018), and the incorporation of point reflectors (e.g., basal sediment; Winter et al., 2019b; Franke et al., 2024b), as well as ice flowing over regions

731 with changes in basal friction (Wolovick et al., 2014; Wolovick and Creyts, 2016) or convergent flow (Bons et
732 al., 2016; Ross et al., 2020). The presence of these basal units can influence the rheological properties and
733 fabric structure of the ice column, as well as impact the continuity of climatic records, highlighting their
734 significance for ice-core drilling projects and ice-flow-modelling endeavours (Bell et al., 2014; MacGregor et
735 al., 2015a; Panton and Karlsson, 2015).

736 Buried surface crevasses imaged in RES data have been used as key evidence for timing the shutdown of
737 Kamb Ice Stream (Retzlaff et al., 1993; Jacobel et al., 2000; Smith et al., 2002; Catania et al., 2006) and the
738 reorganisation of flow through Whillans Ice Stream (Conway et al., 2002). The locations and geometry of
739 basal crevasses formed near the grounding line (Fig. 8j) have also been used to identify previously floating
740 ice, and time the formation of ice rises and ice-flow reorganisation during the Holocene in Antarctica's
741 Weddell Sea Sector (Kingslake et al., 2018; Wearing and Kingslake, 2019).

742 Finally, some particularly bright isochrones have been used to constrain the timing of past volcanic eruptions
743 and constrain the ranges of their tephra fallout. Most such reflectors are relatively bright through chemical
744 signatures alone (e.g., Welch and Jacobel, 2003), but a particularly prominent isochrone, ~30 dB stronger
745 than other typical isochrone-reflection strengths, and thus interpreted as containing physical tephra
746 fragments in addition to chemical residues, was mapped and interpreted by Corr and Vaughan (2008) to
747 demonstrate a volcanic eruption occurred ~2000 years ago in West Antarctica and covered much of the Pine
748 Island Glacier basin (Fig 8k).

749 **5.6 Using isochrones in ice-sheet models**

750 Ice-flow models of different complexities comprise the foremost tools for projecting future ice-sheet and
751 glacier evolution (e.g., Gagliardini et al., 2013; Cornford et al., 2015; DeConto and Pollard, 2016; Seroussi et
752 al., 2020; 2024). Incorporating radiostratigraphic data into ice-sheet models provides a means for validation,
753 improves their calibration and might be essential for making more robust projections by models seeking to
754 constrain ice-sheet evolution over the past few centuries to the late Quaternary (Hindmarsh et al., 2009;
755 Leysinger Vieli et al., 2011; Holschuh et al., 2017; Born and Robinson, 2021; Sutter et al., 2021). Palaeo-proxy
756 records such as exposure-age dating (Brook and Kurz, 1993; Mackintosh et al., 2014; Hillebrand et al., 2021),
757 grounding-line reconstructions (Bentley et al., 2014; Wearing and Kingslake, 2019) or estimates of past sea-
758 level highstands (Dutton et al., 2015) provide invaluable snapshots of ice-sheet variability on local, regional
759 and continental scales (Lecavalier et al., 2023, present a state-of-the-art database), but their interpretation
760 remains challenging in terms of attribution of ice volume, and changes to the grounding zone and ice
761 elevation. Dated radiostratigraphy, on the other hand, contains detailed information on the evolution of ice
762 flow on the relevant timescales (as compiled for today in Sect. 4.5) and thus provides a much-refined
763 calibration target bridging gaps in between snapshot proxy data. Although the theoretical link between ice
764 flow and isochrone geometry has been established for steady tube flow of an ice sheet (Parrenin and

765 Hindmarsh, 2007), the general 3D and transient case remains far more challenging. In this section, we
766 overview recent developments in ice-sheet modelling that incorporate or exploit isochronal data from RES
767 surveys.

768 **5.6.1 Modelling past climate and ice-dynamic changes**

769 Radiostratigraphy is an ideal tuning target for ice-sheet models on continental, regional (catchment) and local
770 scales, because it inherently records the history of the ice flow as well as its response to changing climate
771 conditions in its geometry. As opposed to traditionally-employed tuning targets such as surface flow, ice-
772 sheet geometry or ice volume, which only represent snapshots of ice-sheet evolution, radiostratigraphy
773 provides a 3-D structure which has been formed by the transient palaeo-evolution of the ice sheet. Modelling
774 isochronal geometry and age is technically relatively straightforward, with the main challenge being
775 pervasive uncertainties in boundary conditions (e.g. climate forcing and geothermal heat flux) and the
776 intrinsic uncertainties of ice-sheet models due to their parameterisations of physical processes (Sutter et al.,
777 2021). Isochrones in RES data, age-depth profiles in ice cores and the isotopic content of ice sheets have
778 been modelled either by employing Lagrangian (Sutter et al., 2021) or semi-Lagrangian (Tarasov and Peltier,
779 2003; Clarke et al., 2005; Goelles et al., 2014) advection or isochronal models (Born, 2017; Rieckh et al., 2024).
780 Models that simulate stratigraphy can thus be used to explore the effects of palaeoclimate evolution on ice-
781 dynamic changes, such as marine ice-sheet instabilities or the evolution of ice-sheet drainage systems.

782 Continental-scale ice-sheet models employing approximations of the full-Stokes equations have allowed the
783 computation of ice flow on time scales of centuries to millions of years, albeit at the cost of resolution, which
784 is usually ~5–40 km (Pollard and DeConto, 2009; Golledge et al., 2015; Sutter et al., 2019; Albrecht et al.,
785 2020; Seroussi et al., 2024). While these relatively coarse grid sizes (compared to applications of full-Stokes
786 models; e.g. Zhao et al., 2018) preclude a meaningful interpretation of small-scale processes that influence
787 radiostratigraphy (e.g. local freezing, melting, bedrock features etc.), large-scale models have the advantage
788 that they incorporate the whole thermomechanically-coupled ice-sheet system and its response to changing
789 climate conditions. Consequently, large-scale models are also the main tools for projections of sea-level
790 contributions from the Antarctic and Greenland ice sheets (e.g., Goelzer et al., 2020; Seroussi et al., 2020;
791 2024).

792 The analysis of isochrones to inform on past ice flow need not be limited to the grounded parts of an ice
793 sheet and has been extended to ice shelves (Višnjević et al., 2022; Moss et al., 2023), ice rises (Goel et al.,
794 2018; Goel et al., 2024), and the ice-rise/ice-shelf system (Henry et al., 2024). In these studies, isochrones
795 have served as valuable resources for reconstructing both the surface and/or basal mass balance of ice
796 shelves and ice rises using forward and inverse modelling along the flowline (in 2D), and for investigating
797 rheological properties of ice rise/ice shelf systems in 3D (Henry et al., 2024). Extending this approach to
798 include the past ice-shelf evolution and linking the isochronal structure to its grounded counterparts remains

799 challenging due to the lack of tie points to dated isochrones and a lack of observable isochronal structure
800 across the grounding line.

801 **5.6.2 Model integration of isochronal data**

802 A range of models has been used to calculate the age-depth relationship in ice over both large and small
803 portions of Antarctica and compare this with existing radiostratigraphies; an exercise that can offer valuable
804 insights into ice-sheet processes and how these are represented in ice-sheet models (Fig. 9). When
805 integrating isochronal data in models, multiple factors play a role in the choice of model set up, such as the
806 size of the area of interest (e.g. regional or continental) and the type of flow regime present (e.g. dome,
807 vertical shearing, extension). Various types of flow regime are found in Antarctica, ranging from vertical
808 compression at domes moving to vertical shear and finally to longitudinal extension in ice streams and ice
809 shelves. Consequently, it is important to use a model with the most suitable dimensionality (1D, 2D or 3D)
810 for the specific glaciological conditions in the area being studied. 2.5D models, i.e. 2D models that take into
811 account some aspects of a third dimension, provide another option (Chung et al., 2024).

812 1D models typically assume negligible horizontal flow, making simplifying assumptions such as a steady-state
813 velocity field and the local layer approximation (Waddington et al., 2007, provide guidelines on its
814 applicability) and have predominantly been used at domes such as Dome C (Parrenin et al., 2017; Lilien et al.,
815 2021; Chung et al., 2023) and Dome F (Obase et al., 2023; Wang et al., 2023), where vertical compression
816 dominates. Dated isochrones have been used in multiple studies to constrain 1D age-depth models of
817 different complexity to determine millennial-scale accumulation rates in Antarctica (e.g., Leysinger Vieli et
818 al., 2004; Siegert and Payne, 2004; MacGregor et al., 2009; Karlsson et al., 2014; Koutnik et al., 2016; Cavitte
819 et al., 2018; Zhao et al., 2018; Ashmore et al., 2020; Bodart et al., 2023; Sanderson et al., 2024) and retrieve
820 horizontal flow velocity from 2D isochrone architecture (Eisen, 2008). While most such studies have been
821 restricted to using steady-state due to temporal limitations in available data, some models have allowed for
822 temporal changes in boundary conditions (Callens et al., 2016; Parrenin et al., 2017; Chung et al., 2023).

823 3D modelling of ice-rise stratigraphy (Henry et al., 2024) has provided a step towards constraining long-term
824 simulations in coastal areas. The influence of model physics on this stratigraphy was first investigated in 2D
825 idealised studies of Raymond arches (Pettit and Waddington, 2003; Pettit et al., 2007; Martín and
826 Gudmundsson, 2012), with Hindmarsh et al. (2011) extending this work in 3D idealised simulations.
827 Modelling studies have examined the influence of Glen's flow law exponent on Raymond-arch amplitude
828 (Pettit and Waddington, 2003; Martín et al., 2006; Martín and Gudmundsson, 2012). This methodology has
829 been extended to 2D simulations of real-world ice rises and domes in coastal Antarctica with the comparison
830 of modelled and observed Raymond arches at ice divides (Martín et al., 2009; Hindmarsh et al., 2011; Pettit
831 et al., 2011; Martín et al., 2014; Drews et al., 2015; Goel et al., 2018; Goel et al., 2024).




Figure 9. 3D visualisation of selected traced and dated isochrones in East And West Antarctica, and locations where different modelling applications have been conducted. (a) 2.5 ka (black lines) and 16.5 ka (grey lines) isochrones across the Pine Island/Thwaites Glacier catchment area (Bodart et al., 2021). (b) 2.5, 4.7 and 6.9 ka isochrones spanning Institute Ice Stream (Ashmore et al., 2020). (c) Map of Antarctic traced and dated isochrone transects (black lines) and areas where at least one modelling study is available (grey boxes); red boxes denote areas of the 3D visualisations. (d) Traced and dated (38, 48, 90, 160 ka) isochronal structure around Dome C from Winter et al. (2019a) and (e) Cavitte et al. (2021).

Isochrones have also been used to estimate ice temperature on catchment- to continent-wide scales. Because the electrical conductivity of ice varies exponentially with temperature, resulting in higher dielectric attenuation in warmer ice (MacGregor et al., 2007), temperature variability across the ice sheets leaves a signature in the returned power of measured radio waves. To date, studies have concentrated on using thermomechanical ice-sheet models to improve interpretation of RES data by using modelled temperature fields to remove attenuation effects and strengthen interpretations of bed properties based on basal reflectivity (Matsuoka et al., 2012; MacGregor et al., 2015b; Chu et al., 2021; Dawson et al., 2022). This approach assumes that thermomechanical models can estimate the ice temperature field to high confidence. Additionally, 1D age-depth models that incorporate a thermomechanical component (Parrenin et al., 2017; Passalacqua et al., 2017; Obase et al., 2023) have been used to infer basal melt rates in Antarctica close to domes. Temperature modelling, however, can be challenging in fast-flowing areas where heat production by viscous dissipation is substantial, such as along shear margins or ice streams. As efforts to reduce ambiguity in the direct inference of temperature from RES reflection strength develop, it will become possible to assimilate RES measurements of temperature to improve model performance, as has been done with other direct and indirect observations of subsurface temperature (Pattyn, 2010; Van Liefferinge and Pattyn, 2013). While a combined evaluation of model temperature and velocity data from RES data has been performed qualitatively (Holschuh et al., 2019), there is a growing desire to incorporate *both* radiometric and structural information in a formal modelling framework.

6 Future directions

In this review, we have considered how the internal architecture of the Antarctic ice sheets, and in particular their radiostratigraphy, is increasingly being exploited to elucidate ice and climate history. The ultimate aim of these endeavours is to constrain in ever finer detail the rates, locations and underlying processes of past ice-sheet changes in response to climate forcing. This is crucial to inform and reduce uncertainties in models projecting future ice-sheet changes and concomitant global sea-level rise. Yet, despite the progress reported above, Antarctica's internal architecture remains an underutilised resource for this purpose. In this final section, we set out recommendations for future research activities to be underpinned by an expanded and accessible database of Antarctica's internal architecture. Firstly (Sect. 6.1), we present a pathway towards expanding the volume of radiostratigraphy across Antarctica towards the goal of building a 3-D age-depth model of the ice; secondly (Sect. 6.2), we set out a number of future science challenges that a comprehensive database of Antarctica's englacial architecture can help to address; and finally (Sect. 6.3), we make some recommendations for community actions to facilitate the delivery of these goals.

864 **6.1 Pathway to expanding Antarctic radiostratigraphy**

865 We have identified throughout this review a clear need to expand significantly the traced radiostratigraphy
866 across the Antarctic ice sheets, covering both more area and a greater depth range through the ice. To
867 achieve this requires the following steps:

868 ***6.1.1 Numerical modelling to guide where radiostratigraphic constraints are most needed***

869 We recommend that future targets for tracing radiostratigraphy across different regions of Antarctica, from
870 existing RES data or guiding new RES surveys, are informed directly by the needs of the ice-sheet modelling
871 community to benchmark and constrain their models. Modelling can guide location-based suggestions (e.g.
872 to recover more radiostratigraphy away from ice divides and into more dynamic regions where simple model
873 heuristics may misrepresent englacial conditions), or require targeting of particular time periods (e.g.
874 targeting older isochrones that could advance understanding of glacial-interglacial transitions, amongst
875 others).

876 ***6.1.2 Systematic assessment of the potential of existing data for tracing radiostratigraphy***

877 For this review, we have compiled the spatial coverage of existing published RES data across Antarctica that
878 have high-quality (GNSS) navigation and were acquired digitally, and often coherently (Fig. 6d). In principle,
879 this demonstrates the present coverage of RES data from which radiostratigraphy could be extracted and
880 mapped, and indicates that RES datasets range and interconnect widely across both the East and West
881 Antarctic ice sheets. While this presents a positive message of the potential for pan-Antarctic tracing of
882 radiostratigraphy, whether and how much radiostratigraphy *can* be extracted so widely across the ice sheets
883 from all of these profiles remains unknown. Not all of the RES tracks necessarily contain traceable
884 radiostratigraphy, for reasons that range from inherent RES-system limitations upon data acquisition,
885 decisions made in the processing of the data that are available (see Sect. 3), to the presence of physical
886 phenomena in the ice that disrupt radiostratigraphy or steeply sloping basal topography that makes
887 isochrones too steep to be traced (Sect. 5).

888 A community effort is therefore required to investigate the full potential for mapping radiostratigraphy
889 through these existing datasets. A useful first step, which was beyond the scope of this paper, would be to
890 apply the ILCI to all of the modern datasets presented in Fig. 6d to assess their viability for tracing isochrones
891 across different regions, i.e., to produce a more comprehensive version of Fig. 5 expanded to all the datasets
892 discussed in Sect. 4.

893 **6.1.3 Reprocessing of existing datasets to accentuate internal architecture**

894 While the visibility of internal architecture is partly determined by the initial acquisition parameters and
895 varies across Antarctica, the information visible in RES data is also influenced significantly by the processing
896 applied to the data *after* they have been acquired (Sect. 3.1). Where the raw data exist, the data can be
897 reprocessed, which may significantly enhance the value of some existing datasets for tracing their
898 radiostratigraphy. For much of Antarctica's RES data, the only processing that has been applied was
899 implemented to emphasise and pick the bed echo. In some cases, the same processing accentuated
900 radiostratigraphy in parallel but, in others, it has suppressed the imaging of isochrones or induced artefacts
901 in the radargrams that have hampered or precluded any tracing of radiostratigraphy. Therefore, where
902 existing data lack distinct isochrones in locations identified by numerical modelling as optimal candidates for
903 radiostratigraphy, we recommend, where feasible, firstly reprocessing the raw data to enhance internal
904 architecture. Such an initiative is currently being trialled as part of the Open Polar Radar project using AWI,
905 BAS and USA-acquired RES data across Antarctica (Paden et al., 2021).

906 **6.1.4 New data acquisition**

907 Importantly, new RES data for radiostratigraphic constraints need only be acquired where the processes
908 described above have highlighted that existing data cannot provide the radiostratigraphic constraints
909 required by modelling applications. Such areas will fall into three categories:

- 910 (a) Regions that are still unsurveyed or undersurveyed. Clear examples of this situation, from Fig. 6d,
911 comprise data gaps > 100 km wide in East Antarctica in Enderby Land; between South Pole and Vostok;
912 and between Wilkes and Kemp lands; and we also note that the Filchner-Ronne Ice Shelf does not have
913 dense survey cover.
- 914 (b) Regions where RES surveys have occurred but where the existing data – even after reprocessing – do not
915 contain any internal architecture. These regions typically comprise those last surveyed by RES several
916 decades ago with less sophisticated RES systems. From Fig. 6d, we identify the Siple Coast region of West
917 Antarctica as one such data gap. Although this region was intensively studied and surveyed during the
918 1980s and 1990s, its last major RES surveys predate widespread use of coherent RES systems.
- 919 (c) Regions where RES surveys have occurred but where the existing data – even after reprocessing – contain
920 some internal architecture, but which does not meet modelling needs. Likely scenarios here are that age-
921 depth information is needed at finer resolution than is retrievable in the existing data, or there is a
922 requirement to recover radiostratigraphy deeper into the ice than has been imaged by the existing survey.
923 This situation is common amongst existing datasets that were acquired for projects focussed on other
924 scientific priorities. For example, where some airborne RES datasets have been acquired in combination

with potential-field data (gravity and magnetics), the requirement to fly the aircraft at a stable elevation has sometimes led to poor-quality radiostratigraphy where the range from aircraft to ice surface was too large.

These cases should fundamentally guide the locations, nature and platforms of any new RES data acquisition for internal architecture. As reviewed in Sect. 4, modern airborne RES systems and processing algorithms are adept at detecting multiple isochrones over large regions. In some cases, such as through regions of complex topography, complex flow dynamics or a requirement for very fine resolution of isochrones over regional scales, ground-based RES systems that can typically sound more IRHs and deeper into the underlying ice may still represent the optimal tool and justify the resources required to emplace deep-field parties. However, uncrewed aerial vehicles capable of carrying RES systems (Arnold et al., 2020; Teisberg et al., 2022), when routinely operationalised, may offer a cheaper and safer solution over remote and challenging terrains.

6.1.5 Advances in deep learning to expedite the extraction of internal architecture from RES data

As reviewed in Sect. 3, all of the present radiostratigraphy mapped across Antarctica (Fig. 7) has been generated in the absence of a fully automated isochrone-picking algorithm. Although substantial progress has been made, the need for frequent manual intervention has slowed the generation of pan-Antarctic radiostratigraphy. The greatest promise for a step-change in our ability to trace radiostratigraphy significantly faster lies in the application of deep-learning methods to the challenge. As we discussed in Sect. 3.2, deep-learning applications for isochrone tracing are in their infancy, but have already shown great promise for the fast extraction of both near-surface and (more recently) deeper isochrones. While surface-conformable isochrones are relatively more straightforward to trace by machine-learning models, tracing isochrones deeper in the ice column is challenged by IRH fading, unconformities, and/or merging and splitting of isochrones as ice flows over or around large bedrock obstacles. The significant volume of traced radiostratigraphic data now assembled to date across Antarctica (Fig. 7) may now contribute training data to facilitate the advance and wider application of deep learning to tracing Antarctica's deeper isochrones. However, for recently-established machine-learning pipelines (e.g., Moqadam et al., 2025) to use Antarctic radiostratigraphies most effectively, ideally the number of isochrones traced through different regions would need to be increased significantly. A further limitation to this goal is that machine-learning models are highly data-dependent, such that it is still challenging to analyse in parallel datasets derived from different RES systems and/or derived through diverse processing flows. Despite these challenges, the fast progress made towards successful implementation of such applications and the growing availability of traced radiostratigraphy and underlying RES data now being available, are collectively expected to facilitate a step-change in the growing coverage of Antarctic radiostratigraphy in the years to come.

957 6.2 Scientific challenges to be addressed using internal architecture

958 6.2.1 Identification of optimal areas for retrieving new palaeoclimate records

959 As outlined in Sect. 5.1, Antarctica's deep ice cores have provided invaluable palaeoclimate records from
960 both West and East Antarctica and yet there remain two outstanding directives in the quest for augmenting
961 these existing datasets. One, presently the primary focus of the SCAR IPICS *Oldest Ice* programme, is to
962 identify where a potential climate record extending further back in time than Antarctica's current record
963 (back to ~800,000 k.a. from Dome C; Bouchet et al., 2023) can be sampled. This would address the substantial
964 unknown of whether Antarctica's ice holds a direct continuous record of the mid-Pleistocene transition
965 switch from 41-kyr to 100-kyr glacial-interglacial cycles that is inferred to have occurred between ~1.25-0.8
966 M k.a. from marine-sediment oxygen-isotope records (Hays et al., 1976; Clark et al., 2006; Legrain et al.,
967 2023). A second requirement is to locate sites in the Antarctic ice sheets that preserve higher-resolution
968 palaeoclimate records of epochs than are currently represented in the already-sampled sites. In particular,
969 regions with relatively high present or past accumulation rates can potentially preserve high-resolution
970 climate records of the last millenia. We contend that the development of a pan-continental radiostratigraphy
971 could form a crucial tool for identifying most future ice-core locations around Antarctica.

972 We further recommend that attention is placed on tracing radiostratigraphy around Antarctica's blue-ice
973 zones which, as discussed in Sect. 5.1, have and can represent sites for retrieving ice older than 800 k.a.
974 Targeted studies on their radiostratigraphy could improve understanding of how ice deforms to produce the
975 sampled structures, and hence better contextualise how the ice outcropping in such regions is related to ice
976 buried at depth in interior Antarctica.

977 These initiatives may be complemented by the strategic deployment of rapid-access drilling techniques that
978 could be deployed, alongside intersections with ice cores (discussed in Sect. 5.1), to date and validate the
979 radiostratigraphy. Rapid-access drilling (e.g., Goodge and Severinghaus, 2016; Rix et al., 2019; Goodge et al.,
980 2021; Schwander et al., 2023) can provide borehole access into the ice for deploying sensors to record
981 physical characteristics that correlate with RES isochrones (IceCube Collaboration, 2013; Goodge et al., 2021;
982 Schwander et al., 2023). Additionally, rapid-access drilling allows direct sampling of ice that can be used for
983 radiometric-age dating that can validate the radiostratigraphy (e.g., Bender et al., 2008; Rowell et al., 2023).
984 A dedicated programme of rapid-access ice drilling coordinated with *AntArchitecture* could therefore both
985 help to validate radiostratigraphic age-depth models, and provide a relatively quick and cost-effective
986 methodology for targeting potential future sites for both vertical and horizontal ice coring.

987 **6.2.2 Reconstruction of surface mass balance – millennial timescales**

988 In Sect. 5.2, we discussed that tracing deep (>200 m below the ice surface) isochrones across the Antarctic
989 ice sheets enables reconstruction of changes in surface mass balance over the past several millenia. While
990 the few existing studies have mostly focussed at or near ice divides, where horizontal flow and its associated
991 complexities can mostly be neglected, an expanded pan-continental radiostratigraphy that more
992 comprehensively spans and connects all of Antarctica’s central divide regions will enable these simple
993 applications to be expanded, and can provide a spatially widespread record of how surface mass balance has
994 varied regionally at millennial timescales. Such a record would help us to understand the pervasiveness of
995 synoptic snow-accumulation patterns (e.g., Le Meur et al., 2018; Pauling et al., 2023), and could inform
996 scenarios of future plausible surface-mass-balance variability to be incorporated into model projections (see
997 Lenaerts et al., 2019, for a review). In turn, such refined surface-mass-balance reconstructions would greatly
998 improve the climate forcings employed by palaeo-ice-sheet-modelling studies and increase confidence in
999 their conclusions.

1000 **6.2.3 Reconstruction of surface mass balance – historical timescales**

1001 To reduce uncertainties in near-term (i.e., ~next 200 years) projections of Antarctica’s future evolution, and
1002 thereby improve global sea-level projections, there is a critical need to constrain further the regional climate
1003 models (e.g., Pratap et al., 2022) that are fundamental to forcing ice-sheet models. Important validation for
1004 these models comes from the historical record provided primarily by ice cores, but also by near-surface
1005 radiostratigraphy sounded in the upper few 100 m of the ice sheet. Neither this review, nor the
1006 *AntArchitecture* community to date, has focussed on near-surface IRHs. However, the majority of RES surveys
1007 depicted in Fig. 6 also detected near-surface radiostratigraphy, and many additional surveys have been
1008 undertaken over the past decades across Antarctica using a range of airborne and ground-based platforms
1009 that focussed on detecting shallow isochrones, often for local, but sometimes also for more regional,
1010 scientific applications (e.g., Medley et al., 2013; Medley et al., 2014; Konrad et al., 2019; Kowalewski et al.,
1011 2021; Cavitte et al., 2022). We therefore propose that an important future activity should be the
1012 development of a “near-surface” pan-Antarctic radiostratigraphy complementary to the deeper version that
1013 has primarily formed the focus of this review. In parallel with the techniques and philosophy we have
1014 discussed for dating deep isochrones across Antarctica, near-surface radiostratigraphy can be dated from
1015 intersections with near-surface ice-core records; and the product could be progressively refined by using it
1016 to identify where future near-surface ice cores should be drilled to provide finer dating control. The overall
1017 task of tracing near-surface isochrones across Antarctica should benefit from the application of machine
1018 learning to isochrone tracing, as already exemplified by several studies (e.g., Dong et al., 2021; Rahnemoonfar
1019 et al., 2021; Yari et al., 2021).

1020 **6.2.4 Estimate geothermal heat flux from radiostratigraphy**

1021 The studies mentioned in Sect. 5.3 speak to the significant potential for Antarctica's radiostratigraphy to be
1022 used as a resource for constraining variations to the continent's geothermal heat flux, which remains
1023 enigmatic (Burton-Johnson et al., 2020). As exemplified by Fahnestock et al. (2001b) across the Greenland
1024 Ice Sheet, and more locally in Antarctica by Jordan et al. (2018), it is possible to quantify basal melt with
1025 isochrones by calculating how much melting is required to draw isochrones down towards the base. However,
1026 the relationship between isochrone geometry and basal melting is complex, multi-dimensional and partly
1027 controversial (Leysinger Vieli et al., 2007; Carter et al., 2009; Bons et al., 2021; Wolovick et al., 2021b;
1028 Wolovick et al., 2021a). For a continental-scale application of this technique, a more detailed pan-Antarctic
1029 radiostratigraphy is needed. The optimal data product to invert for geothermal heat flux would be the most
1030 widespread tracings of the deepest undisrupted isochrones across the ice sheets, which is challenging
1031 because deeper isochrones are harder to image and significant drawdown of isochrones where basal melting
1032 is high can prohibit widespread tracing (e.g., Ross and Siegert, 2020). Nevertheless, there is significant
1033 potential to use deep isochrone geometry as further calibration for numerical models seeking to invert
1034 geothermal heat flux (Pattyn, 2010; Van Liefferinge and Pattyn, 2013; Burton-Johnson et al., 2020).

1035 **6.2.5 Comprehensive mapping of basal-ice units and deep-isochrone geometry**

1036 In Sect. 5.5, we noted that in some regions of the Antarctic ice sheets, RES data indicate that the deeper ice
1037 has distinctive physical characteristics compared with the ice above, i.e., where this deeper ice obscures or
1038 precludes imaging of IRHs, and where distinct basal-ice units exist around which the overlying IRHs have
1039 become folded or warped. An improved understanding of the distribution of these features across Antarctica
1040 is important for several reasons. Firstly, it would identify where deep-ice palaeoclimate records would be
1041 compromised by ice deformation or basal melting, thus critically informing ice-core site identification.
1042 Secondly, it would act as an observationally-informed broad-scale indicator of which areas of the ice sheet
1043 are prone to basal melting and hence inform mapping of geothermal heat flux. Thirdly, it would provide
1044 information towards a better understanding of how the rheology of Antarctica's ice varies, what are the
1045 causes of this variation, and how these effects impact on Antarctica's ice dynamics. Some of these issues
1046 would be informed by some specific rapid-access drilling into basal-ice units, and a comprehensive mapping
1047 exercise of basal-unit distribution would inform which targets might be most easily accessed. In addition to
1048 mapping basal units themselves, a complementary activity could be to map the degree to which deep-ice
1049 radiostratigraphy follows or diverges from the ice-bed interface across Antarctica. This exercise would inform
1050 modelling aimed to deconvolve how much isochrone geometry is affected by basal topography versus ice
1051 dynamics versus basal melt. This, in turn, will better inform projections of the ice sheets' future with
1052 radiostratigraphic constraints.

1053 **6.2.6 Advance knowledge of volcanic activity and fallout across Antarctica**

1054 Given that most isochrones traced across the Antarctic ice sheets manifest changes to acidity, and that some
1055 of the brightest have been linked to precipitated fallout from volcanic eruptions within and beyond Antarctica,
1056 there is significant potential to use isochrones across Antarctica more comprehensively to trace the spatial
1057 distribution of volcanic fallout from the numerous past eruptions that have been identified by chemical
1058 analyses of Antarctica's ice cores (Narcisi and Petit, 2021). Despite many tephra and cryptotephra
1059 (microscopic layers of volcanic ash) having been detected in Antarctica's ice cores, few have explicitly been
1060 traced widely beyond the ice cores using radiostratigraphy, and most isochrones that have been linked to
1061 past volcanic events have been used as time markers for other purposes, e.g. calculating past accumulation,
1062 rather than having been traced to focus on the origins and properties of the volcanic events themselves (e.g.,
1063 Jacobel and Welch, 2005; Bodart et al., 2023). There is therefore significant potential, already with existing
1064 data, to use Antarctica's radiostratigraphy to trace the geographical distribution of volcanic fallout from
1065 numerous eruptions that have been detected in ice-core records, and this information may be used to help
1066 trace further the origins and nature of past eruptions beyond that which can be gleaned solely from the ice-
1067 core chemistry. This objective would complement the ongoing activities and recent recommendations for
1068 future research on volcanism presented by the SCAR AntVolc group (Geyer et al., 2023).

1069 **6.2.7 Development of a new model benchmark for the Antarctic ice sheets**

1070 As reviewed in Sect. 5.6, the vast majority of ice-sheet models presently employed for ice-sheet
1071 reconstruction and future projections are initialised with present-day snapshots of the ice-sheet state (e.g.,
1072 surface velocity, ice thickness). An Antarctic-wide radiostratigraphy would provide a much better
1073 initialisation and tuning target for ice-sheet models, as it inherently records both ice-flow history and the ice
1074 sheet's response to changing external forcings (e.g., atmospheric and ocean conditions) – all within a tangible
1075 set of physical horizons that can be reproduced by existing models. The development of an Antarctic-wide
1076 radiostratigraphically-calibrated model benchmark is therefore a primary scientific objective for SCAR's
1077 *AntArchitecture* community.

1078 **6.3 Community actions**

1079 The greatest challenge for attaining the outcomes described above is how to foster and maintain engagement
1080 between scientists working across numerous different disciplines and operating at institutions spread across
1081 Earth. Even within the scientific community who self-describe as RES, radar, or even radioglaciology
1082 specialists, this challenge is innate. As we have reviewed, the history and ongoing practices of Antarctic RES
1083 surveying encompass multiple agencies whose foci are typically on medium-term projects of a few years'
1084 duration. The intent of this review was to communicate to a wider audience (both within and beyond the
1085 radioglaciology community) the baseline availability and potential of the present archive of existing RES data

1086 spanning both East and West Antarctica's ice sheets, and to showcase their value for tackling major science
1087 questions concerning Antarctica's ice and climate history and future.

1088 A major challenge to greater progress in the study of Antarctica's internal architecture has been the lack of
1089 a common framework for archiving RES data and metadata between different operators and potential users.
1090 The establishment of the FAIR (Findable, Accessible, Interoperable, and Reusable; Wilkinson et al., 2016)
1091 data-exchange guidelines has provided a clear framework making possible the release of RES data in open-
1092 access repositories, facilitating open-access releases of some of the datasets discussed in Sect. 4.3.2. These
1093 releases have been accompanied by interactive data portals and FAIR-compliant data standards, including
1094 rich metadata relating to the acquisition, processing and quality of the data, and provide examples for
1095 releasing further data in the future. We recommend that the next significant community data focus should
1096 be on developing common protocols for processing RES data, formatting and sharing raw data files, and in
1097 some cases reprocessing existing data to facilitate much greater interoperability of the data moving into the
1098 future. This recommendation falls into the remit of the Open Polar Radar project currently being trialled with
1099 AWI, BAS and USA-acquired RES data (Paden et al., 2021) but, specifically with regards to publishing and
1100 sharing future radiostratigraphy datasets, there remains a need to set a common standard. We suggest a
1101 standardised structure here in Appendix 1.

1102 A core principle moving forwards with our science must also be on improving sustainability, given the
1103 significant resource and carbon impact of using aircraft and establishing deep-field camps in Antarctica.
1104 When proposing new Antarctic RES acquisition, we suggest that it first be demonstrated that it is needed,
1105 following the procedures laid out in Sect. 6.1. Although crewed airborne and ground-based RES platforms
1106 currently presently continue to provide the most reliable options, where new data are clearly needed n
1107 pathways for improving the sustainability of data collection are opening up with the development of
1108 uncrewed aerial vehicles capable of hosting RES systems (Arnold et al., 2020; Teisberg et al., 2022).

1109 **7. Conclusions**

1110 In this review we have highlighted the vast scientific potential that is contained in Radio-Echo Sounding (RES)
1111 data that have been acquired across the Antarctic ice sheets. The majority of these data have been analysed
1112 only to measure ice thickness, using only the bed echoes which are just one component of the complex data
1113 that RES surveys routinely acquire. However, Antarctic RES surveys, conducted for the last six decades, have
1114 also generated vast archives of "internal architecture" (typically 3-D fields of RES-imaged isochrones) that
1115 record the depositional, deformational and melting histories of ice around Antarctica. Until recently, this vast
1116 archive has been relatively little utilised, for reasons ranging from the challenges of working with datasets
1117 acquired with differing RES systems by multiple operators from different countries, to limitations with
1118 processing big datasets and limited capacity, to trace many 10s to 100s of RES-imaged isochrones through

1119 hundreds of thousands of km of RES profiles. We have detailed how RES data are processed and can be
1120 optimised to make them scientifically useful for a wide range of scientific applications exploring the past and
1121 future evolution of the Antarctic ice sheets (Sect. 3); inventoried where RES data are available to analyse
1122 (Sect. 4.1, 4.2 and Fig. 6) and detailed where this process has begun (Sect. 4.3, Fig. 7 and Table 1); and
1123 reviewed how internal architecture has been applied so far to make progress in linking between and verifying
1124 ice-core chronologies, reconstruct surface mass balance, basal melting and ice-flow dynamics, and been
1125 integrated into numerical modelling of ice-sheet evolution (Sect. 5). We have presented a vision for future
1126 research in Antarctic science that can be underpinned by RES-imaged internal architecture of the ice (Sect.
1127 6), which can inform (1) Identification of optimal sites for retrieving new ice-core palaeoclimate records
1128 targeting different periods; (2) Reconstruction of surface mass balance on millennial or historical timescales;
1129 (3) Estimates of basal melting and geothermal heat flux from radiostratigraphy and comprehensively
1130 mapping basal-ice units, to complement inferences from other geophysical and geological methods; (4)
1131 Advancing knowledge of volcanic activity and fallout across Antarctica; and (5) The refinement of numerical
1132 models that leverage radiostratigraphy to tune time-varying accumulation, basal melting and ice flow, firstly
1133 to reconstruct past behaviour, and then to reduce uncertainties in projecting future ice-sheet behaviour.

1134 To address our scientific goals, we call for continued efforts to build and enhance the inclusion and diversity
1135 of researchers involved in acquiring and analysing RES datasets towards understanding better Antarctica's
1136 past and future. This paper has benefitted immeasurably from including perspectives from authors spread
1137 across the world, navigating different stages of their careers, and identifying as different genders, ethnicities,
1138 nationalities and religions; and from including the expertise of field- and data-focussed scientists in the same
1139 space as the expertise of practitioners whose focus is on applying the data and integrating them into
1140 numerical models. We conclude by reiterating our core scientific ambitions for *AntArchitecture* above: to
1141 build a pan-Antarctic database of isochrones that are accessible, sustainable over the long term, and useful
1142 for multiple scientific applications across multiple users, for example ice-sheet modellers and the substantial
1143 ice-core community. Alongside this, and of equal importance, the community that is active both in acquiring
1144 and analysing Antarctica's internal architecture must continue to diversify.

1145 **Author contributions**

1146 The paper was jointly written by RGB, JAB, MGPC, AC, RJS and JCRS (the lead-writing team). All co-authors
1147 contributed ideas, perspectives and edits. The review was conceptualised by RGB, OE, NBK, JAM, NR and DAY
1148 as a deliverable for the SCAR *AntArchitecture* 2018-2022 Action Group. RGB coordinated the writing process.
1149 DWA, RGB, JAB, AB, MGPC, WC, OE, NH, NBK, MRK, GJMCLV, JAM, EJM, EM, CM, FP, NR, JCRS, KW & DAY
1150 made significant contributions to first draft compiled during the covid-19 pandemic in 2020-21, forming the
1151 framework for the current version handled by the lead-writing team since 2023. Original figures were drawn
1152 by KW, NBK & JAB (Fig. 1), DWA (Fig. 2), SF (Fig. 3), JAB (Fig. 5 & 7), RGB (Fig. 6), , MGCP & RJS (Fig. 8), and
1153 JCRS (Fig. 9), and Table 1 was assembled by JAB. Prior to submission, DWA, AB, RD, JWG, MRK, CM, FN, SVP,
1154 DMS, TOT, XC & XT provided substantive edits; SF, VG, ACJC, AH, BHH, FMO, TR & SY led detailed reviews of
1155 each section of the manuscript which shaped further edits; and OE, NBK, GJMCLV, JAM, FSLN, NR, RS, MJS &
1156 DAY contributed final checks and perspectives informing the final version of the paper.

1157 **Competing interests**

1158 Nanna Karlsson is Co-Editor-in-Chief, Olaf Eisen is Advisory Editor, and Reinhard Drews, Joseph MacGregor,
1159 Elisa Mantelli, Carlos Martín and Johannes Sutter are Editors of *The Cryosphere*.

1160 **Disclaimer**

1161 The views and opinions expressed here are those of the author(s) only and do not necessarily reflect those
1162 of the European Union or the European Research Council Executive Agency. Neither the European Union nor
1163 the granting authority can be held responsible for them.

1164 **Acknowledgements**

1165 This research is a contribution to the Scientific Committee for Antarctic Research's *AntArchitecture* Action
1166 Group, and we thank members of SCAR's Physical Sciences and Geosciences Divisions for ongoing support of
1167 the group since 2018. All of the UK-based authors acknowledge research funding support from the UK Natural
1168 Environment Research Council, including Doctoral Training Scholarships to CJN (Edinburgh E4), RJS (One
1169 Planet) and HD (SENSE). JCRS, JAB, AH, and VV acknowledge funding from the Swiss National Science
1170 Foundation (grant no. 211542). MGPC is a postdoctoral researcher of the FRS-FNRS. AC, OE, EM and FP
1171 acknowledge funding from the European Union: AC, OE and FP via the Horizon 2020 Marie Skłodowska-Curie
1172 grant agreement no. 955750 (DEEPICE), EM from European Research Council Starting Grant 101076793. JAM,
1173 DAY, TOT, SY and SS acknowledge support from the US National Aeronautical and Space Administration; TOT
1174 was supported by a NASA FINESST Grant (80NSSC23K0271) and the TomKat Center for Sustainable Energy.
1175 DAY, BHH, NH, MK, EJM, DMS, SY, MR and SS acknowledge funding from the US National Science Foundation:

1176 for DAY, NH, MK, SY and SS through the Center for Oldest Ice Exploration, an NSF Science and Technology
1177 Center (NSF 2019719); for DAY, SY and SS additionally from Earthcube (NSF 2127606); for BHH from an Office
1178 of Polar Programs Postdoctoral Research Fellowship (NSF 2317927); for EJM from Geosciences Open System
1179 Ecosystem Award NSF 2324092; for DMS from Office of Polar Programs Award NSF 1745137; and for MR
1180 from BIGDATA (IIS-1838230, 2308649) and NSF Leadership Class Computing (OAC-2139536) awards. DAY, SY
1181 and SS were also supported by the G. Unger Vetlesen Foundation. This paper is University of Texas Institute
1182 for Geophysics contribution #####. AB and TR acknowledge funding from the Norwegian Research Council
1183 Grant 314614 (Simulating Ice Cores and Englacial Tracers in the Greenland Ice Sheet). RD was supported by
1184 an Emmy Noether Grant from the Deutsche Forschungsgemeinschaft (DR 822/3-1). SF was funded by the
1185 Walter Benjamin Programme of the German Research Foundation (DFG; project number 506043073). ACJH
1186 is supported by the Wallenberg Foundation (KAW 2021.0275). FMO acknowledges support from the German
1187 Academic Scholarship Foundation. XC (Grant 42376253) and XT (Grant 42276257) were supported by
1188 National Natural Science Foundation of China. CFD was funded by the Natural Sciences and Engineering
1189 Research Council of Canada (NSERC; RGPIN-03761-2017) and the Canada Research Chairs Program (950-
1190 231237).

1191 **References**

- 1192 Albrecht, T., R. Winkelmann & A. Levermann (2020) Glacial-cycle simulations of the Antarctic Ice Sheet with
 1193 the Parallel Ice Sheet Model (PISM) – Part 2: Parameter ensemble analysis. *The Cryosphere*, 14, 633-
 1194 656.
- 1195 Allen, C. (2008) A brief history of radio-echo sounding of ice. *Earthzine Monthly Newsletter*.
 1196 <https://earthzine.org/a-brief-history-of-radio-echo-sounding-of-ice-2/> (last accessed 21 May 2024).
- 1197 Arcone, S. A., R. Jacobel & G. Hamilton (2012a) Unconformable stratigraphy in East Antarctica: Part I. Large
 1198 firm cosets, recrystallized growth, and model evidence for intensified accumulation. *Journal of*
 1199 *Glaciology*, 58, 240-252.
- 1200 Arcone, S. A., R. Jacobel & G. Hamilton (2012b) Unconformable stratigraphy in East Antarctica: Part II.
 1201 Englacial cosets and recrystallized layers. *Journal of Glaciology*, 58, 253-264.
- 1202 Arenas-Pingarrón, Á., H. F. J. Corr, C. Robinson, T. A. Jordan & P. V. Brennan (2023) Polarimetric Airborne
 1203 Scientific Instrument, Mark 2: An ice-sounding airborne synthetic-aperture radar for subglacial
 1204 3D imagery. *IET Radar, Sonar & Navigation*, 17, 1391-1404.
- 1205 Arnold, E., C. Leuschen, F. Rodriguez-Morales, J. Li, J. Paden, R. Hale, et al. (2020) CReSIS airborne radars
 1206 and platforms for ice and snow sounding. *Annals of Glaciology*, 61, 58-67.
- 1207 Ashmore, D. W., R. G. Bingham, N. Ross, M. J. Siegert, T. A. Jordan & D. W. F. Mair (2020) Englacial
 1208 architecture and age-depth constraints across the West Antarctic Ice Sheet. *Geophysical Research*
 1209 *Letters*, 47, e2019GL086663.
- 1210 Baggenstos, D., J. P. Severinghaus, R. Mulvaney, J. R. McConnell, M. Sigl, O. J. Maselli, et al. (2018) A
 1211 horizontal ice core from Taylor Glacier, its implications for Antarctic climate history, and an improved
 1212 Taylor Dome Ice Core time scale. *Paleoceanography and Paleoclimatology*, 33, 778-794.
- 1213 Beem, L. H., D. A. Young, J. S. Greenbaum, D. D. Blankenship, M. G. P. Cavitte, J. Guo, et al. (2021)
 1214 Aerogeophysical characterization of Titan Dome, East Antarctica, and potential as an ice-core target.
 1215 *The Cryosphere*, 15, 1719-1730.
- 1216 Bell, R. E., D. D. Blankenship, C. A. Finn, D. L. Morse, T. A. Scambos, J. M. Brozena, et al. (1998) Influence
 1217 of subglacial geology on the onset of a West Antarctic ice stream from aerogeophysical observations.
 1218 *Nature*, 394, 58-62.
- 1219 Bell, R. E., F. Ferraccioli, T. T. Creyts, D. Braaten, H. Corr, I. Das, et al. (2011) Widespread persistent
 1220 thickening of the East Antarctic Ice Sheet by freezing from the base. *Science*, 331, 1592-1595.
- 1221 Bell, R. E., K. Tinto, I. Das, M. Wolovick, W. Chu, T. T. Creyts, et al. (2014) Deformation, warming and softening
 1222 of Greenland's ice by refreezing meltwater. *Nature Geoscience*, 7, 497-502.
- 1223 Bender, M. L., B. Barnett, G. Dreyfus, J. Jouzel & D. Porcelli (2008) The contemporary degassing rate of ⁴⁰Ar
 1224 from the solid Earth. *Proceedings of the National Academy of Sciences*, 105, 8232-8237.
- 1225 Bentley, M. J., C. Ó Cofaigh, J. B. Anderson, H. Conway, B. Davies, A. G. C. Graham, et al. (2014) A
 1226 community-based geological reconstruction of Antarctic Ice Sheet deglaciation since the Last Glacial
 1227 Maximum. *Quaternary Science Reviews*, 100, 1-9.
- 1228 Bingham, R. G., D. M. Rippin, N. B. Karlsson, H. F. J. Corr, F. Ferraccioli, T. A. Jordan, et al. (2015) Ice-flow
 1229 structure and ice-dynamic changes in the Weddell Sea sector of West Antarctica from radar-imaged
 1230 internal layering. *Journal of Geophysical Research: Earth Surface*, 120, 655-670.
- 1231 Bingham, R. G. & M. J. Siegert (2007) Radio-echo sounding over polar ice masses. *Journal of Environmental*
 1232 *and Engineering Geophysics*, 12, 47-62.
- 1233 Bingham, R. G., M. J. Siegert, D. A. Young & D. D. Blankenship (2007) Organized flow from the South Pole to
 1234 the Filchner-Ronne Ice Shelf: An assessment of balance velocities in interior East Antarctica using
 1235 radio-echo sounding data. *Journal of Geophysical Research: Earth Surface*, 112, F03S26.
- 1236
- 1237 Blankenship, D. D., D. L. Morse, C. A. Finn, R. E. Bell, M. E. Peters, S. D. Kempf, et al. (2001) Geologic
 1238 controls on the initiation of rapid basal motion for West Antarctic ice streams: A geophysical
 1239 perspective including new airborne radar sounding and laser altimetry results. In *The West Antarctic*
 1240 *Ice Sheet: Behavior and Environment*, 105-121.
- 1241 Blankenship, D. D., S. D. Kempf, D. A. Young, T. G. Richter, D. M. Schroeder, J. S. Greenbaum, et al. (2017a)
 1242 IceBridge HiCARS 1 L1B Time-Tagged Echo Strength Profiles, Version 1. [Indicate subset used].
 1243 Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center.
- 1244
- 1245 Bodart, J. A., R. G. Bingham, D. W. Ashmore, N. B. Karlsson, A. S. Hein & D. G. Vaughan (2021) Age-depth
 1246 stratigraphy of Pine Island Glacier inferred from airborne radar and ice-core chronology. *Journal of*
 1247 *Geophysical Research: Earth Surface*, 126, e2020JF005927.
- 1248 Bodart, J. A., R. G. Bingham, D. A. Young, J. A. MacGregor, D. W. Ashmore, E. Quartini, et al. (2023) High
 1249 mid-Holocene accumulation rates over West Antarctica inferred from a pervasive ice-penetrating radar
 1250 reflector. *The Cryosphere*, 17, 1497-1512.

1251 Bons, P. D., T. de Riese, S. Franke, M. G. Llorens, T. Sachau, N. Stoll, et al. (2021) Comment on “Exceptionally
 1252 high heat flux needed to sustain the Northeast Greenland Ice Stream” by Smith-Johnsen et al. (2020).
 1253 *The Cryosphere*, 15, 2251-2254.

1254 Bons, P. D., D. Jansen, F. Mundel, C. C. Bauer, T. Binder, O. Eisen, et al. (2016) Converging flow and
 1255 anisotropy cause large-scale folding in Greenland's ice sheet. *Nature Communications*, 7, 11427.

1256 Born, A. (2017) Tracer transport in an isochronal ice-sheet model. *Journal of Glaciology*, 63, 22-38.

1257 Born, A. & A. Robinson (2021) Modeling the Greenland englacial stratigraphy. *The Cryosphere*, 15, 4539-4556.

1258 Bouchet, M., A. Landais, A. Grisart, F. Parrenin, F. Prié, R. Jacob, et al. (2023) The Antarctic Ice Core
 1259 Chronology 2023 (AICC2023) chronological framework and associated timescale for the European
 1260 Project for Ice Coring in Antarctica (EPICA) Dome C ice core. *Climate of the Past*, 19, 2257-2286.

1261 Brook, E. J. & C. Buizert (2018) Antarctic and global climate history viewed from ice cores. *Nature*, 558, 200-
 1262 208.

1263 Brook, E. J. & M. D. Kurz (1993) Surface-exposure chronology using in situ cosmogenic ^3He in Antarctic quartz
 1264 sandstone boulders. *Quaternary Research*, 39, 1-10.

1265 Burton-Johnson, A., R. Dziadek & C. Martín (2020) Geothermal heat flow in Antarctica: Current and future
 1266 directions. *The Cryosphere*, 14, 3843-3873.

1267 Callens, D., R. Drews, E. Witrant, M. Philippe & F. Pattyn (2016) Temporally stable surface mass balance
 1268 asymmetry across an ice rise derived from radar internal-reflection horizons through inverse modeling.
 1269 *Journal of Glaciology*, 62, 525-534.

1270 Carter, S. P., D. D. Blankenship, M. E. Peters, D. A. Young, J. W. Holt & D. L. Morse (2007) Radar-based
 1271 subglacial-lake classification in Antarctica. *Geochemistry, Geophysics, Geosystems*, 8, Q03016.

1272 Carter, S. P., D. D. Blankenship, D. A. Young & J. W. Holt (2009) Using radar-sounding data to identify the
 1273 distribution and sources of subglacial water: Application to Dome C, East Antarctica. *Journal of*
 1274 *Glaciology*, 55, 1025-1040.

1275 Castelletti, D., D. M. Schroeder, S. Hensley, C. Grima, G. Ng, D. Young, et al. (2017) An interferometric
 1276 approach to cross-track clutter detection in two-channel VHF radar sounders. *IEEE Transactions on*
 1277 *Geoscience and Remote Sensing*, 55, 6128-6140.

1278 Castelletti, D., D. M. Schroeder, T. M. Jordan & D. Young (2021) Permanent scatterers in repeat-pass airborne
 1279 VHF radar sounder for layer-velocity estimation. *IEEE Geoscience and Remote Sensing Letters*, 18,
 1280 1766-1770.

1281 Castelletti, D., D. M. Schroeder, E. Mantelli & A. Hilger (2019) Layer-optimized SAR processing and slope
 1282 estimation in radar-sounder data. *Journal of Glaciology*, 65, 983-988.

1283 Catania, G., C. Hulbe & H. Conway (2010) Grounding-line basal melt rates determined using radar-derived
 1284 internal stratigraphy. *Journal of Glaciology*, 56, 545-554.

1285 Catania, G. A., T. A. Scambos, H. Conway & C. F. Raymond (2006) Sequential stagnation of Kamb Ice Stream,
 1286 West Antarctica. *Geophysical Research Letters*, 33, L14502.

1287 Cavitte, M. G. P., D. D. Blankenship, D. A. Young, D. M. Schroeder, F. Parrenin, E. Le Meur, et al. (2016)
 1288 Deep radiostratigraphy of the East Antarctic plateau: Connecting the Dome C and Vostok ice-core
 1289 sites. *Journal of Glaciology*, 62, 323-334.

1290 Cavitte, M. G. P., H. Goosse, K. Matsuoka, S. Wauthy, V. Goel, R. Dey, et al. (2023) Investigating the spatial
 1291 representativeness of East Antarctic ice cores: A comparison of ice core and radar-derived surface
 1292 mass balance over coastal ice rises and Dome Fuji. *The Cryosphere*, 17, 4779-4795.

1293 Cavitte, M. G. P., H. Goosse, S. Wauthy, T. Kausch, J.-L. Tison, B. Van Liefferinge, et al. (2022) From ice core
 1294 to ground-penetrating radar: Representativeness of SMB at three ice rises along the Princess Ragnhild
 1295 Coast, East Antarctica. *Journal of Glaciology*, 68, 1221-1233.

1296 Cavitte, M. G. P., F. Parrenin, C. Ritz, D. A. Young, B. Van Liefferinge, D. D. Blankenship, et al. (2018)
 1297 Accumulation patterns around Dome C, East Antarctica, in the last 73 kyr. *The Cryosphere*, 12, 1401-
 1298 1414.

1299 Cavitte, M. G. P., D. A. Young, R. Mulvaney, C. Ritz, J. S. Greenbaum, G. Ng, et al. (2021) A detailed
 1300 radiostratigraphic data set for the central East Antarctic Plateau spanning from the Holocene to the
 1301 mid-Pleistocene. *Earth System Science Data*, 13, 4759-4777.

1302 Christianson, K., R. W. Jacobel, H. J. Horgan, S. Anandakrishnan & R. B. Alley (2012) Subglacial Lake
 1303 Whillans: Ice-penetrating radar and GPS observations of a shallow active reservoir beneath a West
 1304 Antarctic ice stream. *Earth and Planetary Science Letters*, 331-332, 237-245.

1305 Chu, W., A. M. Hilger, R. Culberg, D. M. Schroeder, T. M. Jordan, H. Seroussi, et al. (2021) Multisystem
 1306 synthesis of radar sounding observations of the Amundsen Sea Sector from the 2004–2005 field
 1307 season. *Journal of Geophysical Research: Earth Surface*, 126, e2021JF006296.

1308 Chung, A., F. Parrenin, R. Mulvaney, L. Vittuari, M. Frezzotti, A. Zanutta, et al. (2024) Age, thinning and spatial
 1309 origin of the Beyond EPICA ice from a 2.5D ice flow model. *EGU sphere*, 2024, 1-21.

1310 Chung, A., F. Parrenin, D. Steinhage, R. Mulvaney, C. Martín, M. G. P. Cavitte, et al. (2023) Stagnant ice and
 1311 age modelling in the Dome C region, Antarctica. *The Cryosphere*, 17, 3461-3483.

1312 Clark, P. U., D. Archer, D. Pollard, J. D. Blum, J. A. Rial, V. Brovkin, et al. (2006) The middle Pleistocene
1313 transition: Characteristics, mechanisms, and implications for long-term changes in atmospheric pCO₂.
1314 *Quaternary Science Reviews*, 25, 3150-3184.

1315 Clarke, G. K. C., N. Lhomme & S. J. Marshall (2005) Tracer transport in the Greenland Ice Sheet: Three-
1316 dimensional isotopic stratigraphy. *Quaternary Science Reviews*, 24, 155-171.

1317 Clough, J. W. (1977) Radio-echo sounding: Reflections from internal layers in ice sheets. *Journal of Glaciology*,
1318 18, 3-14.

1319 Cole-Dai, J., D. G. Ferris, J. A. Kennedy, M. Sigl, J. R. McConnell, T. J. Fudge, et al. (2021) Comprehensive
1320 record of volcanic eruptions in the Holocene (11,000 years) from the WAIS Divide, Antarctica, ice core.
1321 *Journal of Geophysical Research: Atmospheres*, 126, e2020JD032855.

1322 Conway, H., G. Catania, C. F. Raymond, A. M. Gades, T. A. Scambos & H. Engelhardt (2002) Switch of flow
1323 direction in an Antarctic ice stream. *Nature*, 419, 465-467.

1324 Conway, H., B. L. Hall, G. H. Denton, A. M. Gades & E. D. Waddington (1999) Past and future grounding-line
1325 retreat of the West Antarctic Ice Sheet. *Science*, 286, 280-283.

1326 Cook, C. P., T. van de Flierdt, T. Williams, S. R. Hemming, M. Iwai, M. Kobayashi, et al. (2013) Dynamic
1327 behaviour of the East Antarctic Ice Sheet during Pliocene warmth. *Nature Geoscience*, 6, 765-769.

1328 Cornford, S. L., D. F. Martin, A. J. Payne, E. G. Ng, A. M. Le Brocq, R. M. Gladstone, et al. (2015) Century-
1329 scale simulations of the response of the West Antarctic Ice Sheet to a warming climate. *The Cryosphere*, 9, 1579-1600.

1331 Corr, H., F. Ferraccioli, N. Frearson, T. Jordan, C. Robinson, E. Armadillo, et al. (2007) Airborne radio-echo
1332 sounding of the Wilkes Subglacial Basin, the Transantarctic Mountains, and the Dome C region. *Terra
1333 Antarctica Reports*, 13, 55-63.

1334 Corr, H. F. J. & D. G. Vaughan (2008) A recent volcanic eruption beneath the West Antarctic Ice Sheet. *Nature
1335 Geoscience*, 1, 122-125.

1336 .

1337 Creyts, T. T., F. Ferraccioli, R. E. Bell, M. Wolovick, H. Corr, K. C. Rose, et al. (2014) Freezing of ridges and
1338 water networks preserves the Gamburtsev Subglacial Mountains for millions of years. *Geophysical
1339 Research Letters*, 41, 8114-8122.

1340 Cui, X., J. S. Greenbaum, S. Lang, X. Zhao, L. Li, J. Guo, et al. (2020a) The scientific operations of Snow
1341 Eagle 601 in Antarctica in the past five austral seasons. *Remote Sensing*, 12, 2994.

1342 Cui, X., H. Jeofry, J. S. Greenbaum, J. Guo, L. Li, L. E. Lindzey, et al. (2020b) Bed topography of Princess
1343 Elizabeth Land in East Antarctica. *Earth System Science Data*, 12, 2765-2774.

1344 Cui, X. B., W. J. Du, H. Xie & B. Sun (2020c) The ice flux to the Lambert Glacier and Amery Ice Shelf along
1345 the Chinese inland traverse and implications for mass balance of the drainage basins, East Antarctica.
1346 *Polar Research*, 39, 3582.

1347 Dahl-Jensen, D., N. S. Gundestrup, K. Keller, S. J. Johnsen, S. P. Gogineni, C. T. Allen, et al. (1997) A search
1348 in north Greenland for a new ice-core drill site. *Journal of Glaciology*, 43, 300-306.

1349 Damaske, D. & M. McLean (2005) An aerogeophysical survey south of the Prince Charles Mountains, East
1350 Antarctica. *Terra Antarctica*, 12, 87-98.

1351 Dansgaard, W. & S. J. Johnsen (1969) A flow model and a time scale for the ice core from Camp Century,
1352 Greenland. *Journal of Glaciology*, 8, 215 - 223.

1353 Das, I., R. E. Bell, T. A. Scambos, M. Wolovick, T. T. Creyts, M. Studinger, et al. (2013) Influence of persistent
1354 wind scour on the surface mass balance of Antarctica. *Nature Geoscience*, 6, 367-371.

1355 Das, I., L. Padman, R. E. Bell, H. A. Fricker, K. J. Tinto, C. Hulbe, et al. (2020) Multidecadal basal-melt rates
1356 and structure of the Ross Ice Shelf, Antarctica, using airborne ice-penetrating radar. *Journal of
1357 Geophysical Research: Earth Surface*, 125, e2019JF005241.

1358 Dawson, E. J., D. M. Schroeder, W. Chu, E. Mantelli & H. Seroussi (2022) Ice mass-loss sensitivity to the
1359 Antarctic Ice Sheet basal thermal state. *Nature Communications*, 13, 4957.

1360 DeConto, R. M. & D. Pollard (2016) Contribution of Antarctica to past and future sea-level rise. *Nature*, 531,
1361 591-597.

1362 DeConto, R. M., D. Pollard, R. B. Alley, I. Velicogna, E. Gasson, N. Gomez, et al. (2021) The Paris Climate
1363 Agreement and future sea-level rise from Antarctica. *Nature*, 593, 83-89.

1364 Delf, R., D. M. Schroeder, A. Curtis, A. Giannopoulos & R. G. Bingham (2020) A comparison of automated
1365 approaches to extracting englacial-layer geometry from radar data across ice sheets. *Annals of
1366 Glaciology*, 61, 234-241.

1367 Dome Fuji Ice Core Project Members (2017) State dependence of climatic instability over the past 720,000
1368 years from Antarctic ice cores and climate modeling. *Science Advances*, 3, e1600446.

1369 Dong, S., X. Tang, J. Guo, L. Fu, X. Chen & B. Sun (2021) EisNet: Extracting bedrock and internal layers from
1370 radiostratigraphy of ice sheets with machine learning. *IEEE Transactions on Geoscience and Remote
1371 Sensing*, 60, 4303212.

1372 Dowdeswell, J. A. & S. Evans (2004) Investigations of the form and flow of ice sheets and glaciers using radio-
1373 echo sounding. *Reports on Progress in Physics*, 67, 1821-1861.

1374 Drewry, D. J. (1975) Terrain units in eastern Antarctica. *Nature*, 256, 194-195.

- 1375 Drewry, D. J. (Ed.) (1983) *Antarctica: Glaciological and Geophysical Folio*. Cambridge: Cambridge University
1376 Press.
- 1377 Drewry, D. J. (2023) *The Land Beneath the Ice: The Pioneering Years of Radar Exploration in Antarctica*.
1378 Princeton: Princeton University Press.
- 1379 Drewry, D. J. & D. T. Meldrum (1978) Antarctic airborne radio-echo sounding, 1977–78. *Polar Record*, 19,
1380 267–273.
- 1381 Drews, R., O. Eisen, D. Steinhage, I. Weikusat, S. Kipfstuhl & F. Wilhelms (2012) Potential mechanisms for
1382 anisotropy in ice-penetrating radar data. *Journal of Glaciology*, 58, 613–624.
- 1383 Drews, R., O. Eisen, I. Weikusat, S. Kipfstuhl, A. Lambrecht, D. Steinhage, et al. (2009) Layer disturbances
1384 and the radio-echo free zone in ice sheets. *The Cryosphere*, 3, 195–203.
- 1385 Drews, R., C. Martín, D. Steinhage & O. Eisen (2013) Characterizing the glaciological conditions at
1386 Halvfarryggen Ice Dome, Dronning Maud Land, Antarctica. *Journal of Glaciology*, 59, 9–20.
- 1387 Drews, R., K. Matsuoka, C. Martín, D. Callens, N. Bergeot & F. Pattyn (2015) Evolution of Derwael Ice Rise in
1388 Dronning Maud Land, Antarctica, over the last millennia. *Journal of Geophysical Research: Earth
1389 Surface*, 120, 564–579.
- 1390 Dutton, A., A. E. Carlson, A. J. Long, G. A. Milne, P. U. Clark, R. DeConto, et al. (2015) Sea-level rise due to
1391 polar ice-sheet mass loss during past warm periods. *Science*, 349, aaa4019.
- 1392 Eisen, O. (2008) Inference of velocity pattern from isochronous layers in firn, using an inverse method. *Journal
1393 of Glaciology*, 54, 613–630.
- 1394 Eisen, O., I. Hamann, S. Kipfstuhl, D. Steinhage & F. Wilhelms (2007) Direct evidence for continuous radar
1395 reflector originating from changes in crystal-orientation fabric. *The Cryosphere*, 1, 1–10.
- 1396 Eisen, O., U. Nixdorf, F. Wilhelms & H. Miller (2002) Electromagnetic wave speed in polar ice: Validation of
1397 the common-midpoint technique with high-resolution dielectric-profiling and γ -density measurements.
1398 *Annals of Glaciology*, 34, 150–156.
- 1399 Eisen, O., W. Rack, U. Nixdorf & F. Wilhelms (2005) Characteristics of accumulation around the EPICA deep-
1400 drilling site in Dronning Maud Land, Antarctica. *Annals of Glaciology*, 41, 41–46.
- 1401 Eisen, O., F. Wilhelms, U. Nixdorf & H. Miller (2003) Revealing the nature of radar reflections in ice: DEP-
1402 based FDTD forward modeling. *Geophysical Research Letters*, 30, 1218.
- 1403 Eisen, O., F. Wilhelms, D. Steinhage & J. Schwander (2006) Improved method to determine radio-echo
1404 sounding reflector depths from ice-core profiles of permittivity and conductivity. *Journal of Glaciology*,
1405 52, 299–310.
- 1406 EPICA Community Members (2004) Eight glacial cycles from an Antarctic ice core. *Nature*, 429, 623–628.
- 1407 Ershadi, M. R., R. Drews, C. Martín, O. Eisen, C. Ritz, H. Corr, et al. (2022) Polarimetric radar reveals the
1408 spatial distribution of ice fabric at domes and divides in East Antarctica. *The Cryosphere*, 16, 1719–
1409 1739.
- 1410 Escutia, C., M. A. Bárcena, R. G. Lucchi, O. Romero, A. M. Ballegeer, J. J. Gonzalez, et al. (2009) Circum-
1411 Antarctic warming events between 4 and 3.5 Ma recorded in marine sediments from the Prydz Bay
1412 (ODP Leg 188) and the Antarctic Peninsula (ODP Leg 178) margins. *Global and Planetary Change*,
1413 69, 170–184.
- 1414 Evans, S. & B. M. E. Smith (1970) Radio-echo exploration of the Antarctic Ice Sheet, 1969–70. *Polar Record*,
1415 15, 336–338.
- 1416 Fahnestock, M., W. Abdalati, S. Luo & S. Gogineni (2001a) Internal-layer tracing and age-depth-accumulation
1417 relationships for the northern Greenland Ice Sheet. *Journal of Geophysical Research: Atmospheres*,
1418 106, 33789–33797.
- 1419 Fahnestock, M. A., W. Abdalati, I. R. Joughin, J. M. Brozena & P. Gogineni (2001b) High geothermal heat flow,
1420 basal melt, and the origin of rapid ice flow in central Greenland. *Science*, 294, 2338 – 2342.
- 1421 Fischer, H., J. Severinghaus, E. Brook, E. Wolff, M. Albert, O. Alemany, et al. (2013) Where to find 1.5 million
1422 yr old ice for the IPICS "Oldest-Ice" ice core. *Climate of the Past*, 9, 2489–2505.
- 1423 Fogwill, C. J., C. S. M. Turney, N. R. Golledge, D. M. Etheridge, M. Rubino, D. P. Thornton, et al. (2017)
1424 Antarctic ice-sheet discharge driven by atmosphere-ocean feedbacks at the Last Glacial Termination.
1425 *Scientific Reports*, 7, 39979.
- 1426 Fox-Kemper, B., H. T. Hewitt, C. Xiao, G. Aðalgeirsdóttir, S. S. Drijfhout, T. L. Edwards, et al. 2021. Ocean,
1427 cryosphere and sea-level change. In *Climate Change 2021: The Physical Science Basis. Contribution
1428 of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*,
1429 eds. V. Masson-Delmotte, P. Zhai, A. Pirani, S. L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L.
1430 Goldfarb, M. I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J. B. R. Matthews, T. K. Maycock, T. Waterfield,
1431 O. Yelekçi, R. Yu & B. Zhou, 1211–1362. Cambridge: Cambridge University Press.
- 1432 Franke, S., H. Eisermann, W. Jokat, G. Eagles, J. Asseng, H. Miller, et al. (2021) Preserved landscapes
1433 underneath the Antarctic Ice Sheet reveal the geomorphological history of Jutulstraumen Basin. *Earth
1434 Surface Processes and Landforms*, 46, 2728–2745.
- 1435 Franke, S., T. Gerber, C. Warren, D. Jansen, O. Eisen & D. Dahl-Jensen (2023) Investigating the radar
1436 response of englacial-debris-entrained basal-ice units in East Antarctica using electromagnetic
1437 forward modeling. *IEEE Transactions on Geoscience and Remote Sensing*, 61, 4301516.

- 1438 Franke, S., D. Jansen, T. Binder, J. D. Paden, N. Dörr, T. A. Gerber, et al. (2022) Airborne ultra-wideband
1439 radar sounding over the shear margins and along flow lines at the onset region of the Northeast
1440 Greenland Ice Stream. *Earth System Science Data*, 14, 763-779.
- 1441 Franke, S., D. Steinhage, V. Helm, A. M. Zuhr, J. A. Bodart, O. Eisen & P. Bons (2025) Age-depth distribution
1442 in western Dronning Maud Land, East Antarctica, and Antarctic-wide comparisons of internal reflection
1443 horizons. *The Cryosphere*, 19, 1153-1180.
- 1444 Franke, S., M. Wolovick, R. Drews, D. Jansen, K. Matsuoka & P. D. Bons (2024b) Sediment freeze-on and
1445 transport near the onset of a fast-flowing glacier in East Antarctica. *Geophysical Research Letters*, 51,
1446 e2023GL107164.
- 1447 Frémand, A. C., J. A. Bodart, T. A. Jordan, F. Ferraccioli, C. Robinson, H. F. J. Corr, et al. (2022) British
1448 Antarctic Survey's aerogeophysical data: Releasing 25 years of airborne gravity, magnetic, and radar
1449 datasets over Antarctica. *Earth System Science Data*, 14, 3379-3410.
- 1450 Frémand, A. C., P. Fretwell, J. A. Bodart, H. D. Pritchard, A. Aitken, J. L. Bamber, et al. (2023) Antarctic
1451 Bedmap data: Findable, Accessible, Interoperable, and Reusable (FAIR) sharing of 60 years of ice
1452 bed, surface, and thickness data. *Earth System Science Data*, 15, 2695-2710.
- 1453 Frezzotti, M., G. Bitelli, P. De Michelis, A. Deponti, A. Forieri, S. Gandolfi, et al. (2004) Geophysical survey at
1454 Talos Dome, East Antarctica: The search for a new deep-drilling site. *Annals of Glaciology*, 39, 423-
1455 432.
- 1456 Fudge, T. J., B. H. Hills, A. N. Horlings, N. Holschuh, J. E. Christian, L. Davidge, et al. (2023) A site for deep
1457 ice coring at West Hercules Dome: Results from ground-based geophysics and modeling. *Journal of*
1458 *Glaciology*, 69, 538-550.
- 1459 Fujita, S., P. Holmlund, I. Andersson, I. Brown, H. Enomoto, Y. Fujii, et al. (2011) Spatial and temporal
1460 variability of snow accumulation rate on the East Antarctic ice divide between Dome Fuji and EPICA
1461 DML. *The Cryosphere*, 5, 1057-1081.
- 1462 Fujita, S., H. Maeno, S. Uratsuka, T. Furukawa, S. Mae, Y. Fujii, et al. (1999) Nature of radio-echo layering in
1463 the Antarctic Ice Sheet detected by a two-frequency experiment. *Journal of Geophysical Research:*
1464 *Solid Earth*, 104, 13013-13024.
- 1465 Fujita, S., T. Matsuoka, T. Ishida, K. Matsuoka & S. Mae (2000) A summary of the complex dielectric
1466 permittivity of ice in the megahertz range and its applications for radar sounding of polar ice sheets. In
1467 *International Symposium on Physics of Ice Core Records*, 185 - 212. Shikotsukohan, Hokkaido, Japan.
- 1468 Gagliardini, O., T. Zwinger, F. Gillet-Chaulet, G. Durand, L. Favier, B. de Fleurian, et al. (2013) Capabilities
1469 and performance of Elmer/Ice, a new-generation ice-sheet model. *Geoscientific Model Development*,
1470 6, 1299-1318.
- 1471 Gasson, E., R. DeConto, D. Pollard & R. H. Levy (2016) Dynamic Antarctic ice sheet during the early to mid-
1472 Miocene. *Proceedings of the National Academy of Sciences*, 113, 3459-3464.
- 1473 Gerber, T. A., D. A. Lilien, N. M. Rathmann, S. Franke, T. J. Young, F. Valero-Delgado, et al. (2023) Crystal-
1474 orientation-fabric anisotropy causes directional hardening of the Northeast Greenland Ice Stream.
1475 *Nature Communications*, 14, 2653.
- 1476 Geyer, A., A. Di Roberto, J. L. Smellie, M. Van Wyk de Vries, K. S. Panter, A. P. Martin, et al. (2023) Volcanism
1477 in Antarctica: An assessment of the present state of research and future directions. *Journal of*
1478 *Volcanology and Geothermal Research*, 444, 107941.
- 1479 Goel, V., C. Martín & K. Matsuoka (2018) Ice-rise stratigraphy reveals changes in surface mass balance over
1480 the last millennia in Dronning Maud Land. *Journal of Glaciology*, 64, 932-942.
- 1481 Goel, V., C. Martín & K. Matsuoka (2024) Evolution of ice rises in the Fimbul Ice Shelf, Dronning Maud Land,
1482 over the last millennium. *Antarctic Science*, 36, 110-124.
- 1483 Goelles, T., K. Grosfeld & G. Lohmann (2014) Semi-Lagrangian transport of oxygen isotopes in polythermal
1484 ice sheets: implementation and first results. *Geoscientific Model Development*, 7, 1395-1408.
- 1485 Goelzer, H., S. Nowicki, A. Payne, E. Larour, H. Seroussi, W. H. Lipscomb, et al. (2020) The future sea-level
1486 contribution of the Greenland Ice Sheet: A multi-model ensemble study of ISMIP6. *The Cryosphere*,
1487 14, 3071-3096.
- 1488 Gogineni, P., T. Chuah, C. Allen, K. Jezek & R. K. Moore (1998) An improved coherent radar depth sounder.
1489 *Journal of Glaciology*, 44, 659-669.
- 1490 Golledge, N. R., E. D. Keller, N. Gomez, K. A. Naughten, J. Bernales, L. D. Trusel, et al. (2019) Global
1491 environmental consequences of twenty-first-century ice-sheet melt. *Nature*, 566, 65-72.
- 1492 Golledge, N. R., D. E. Kowalewski, T. R. Naish, R. H. Levy, C. J. Fogwill & E. G. W. Gasson (2015) The multi-
1493 millennial Antarctic commitment to future sea-level rise. *Nature*, 526, 421-425.
- 1494 Goodge, J. W. & J. P. Severinghaus (2016) Rapid Access Ice Drill: A new tool for exploration of the deep
1495 Antarctic ice sheets and subglacial geology. *Journal of Glaciology*, 62, 1049-1064.
- 1496 Goodge, J. W., J. P. Severinghaus, J. Johnson, D. Tosi & R. Bay (2021) Deep ice drilling, bedrock coring and
1497 dust logging with the Rapid Access Ice Drill (RAID) at Minna Bluff, Antarctica. *Annals of Glaciology*,
1498 62, 324-339.

1499 Gow, A. J. & T. Williamson (1976) Rheological implications of the internal structure and crystal fabrics of the
1500 West Antarctic Ice Sheet as revealed by deep core drilling at Byrd Station. *Geological Society of*
1501 *America Bulletin*, 87, 1665-1677.

1502 Gudmandsen, P., E. Nilsson, M. Pallisgaard, N. Skou & F. Søndergaard (1975) New equipment for radio-echo
1503 sounding. *Antarctic Journal of the United States*, 10, 234-236.

1504 Gulick, S. P. S., A. E. Shevenell, A. Montelli, R. Fernandez, C. Smith, S. Warny, et al. (2017) Initiation and
1505 long-term instability of the East Antarctic Ice Sheet. *Nature*, 552, 225-229.

1506 Hammer, C. U. (1980) Acidity of polar ice cores in relation to absolute dating, past volcanism, and radio-
1507 echoes. *Journal of Glaciology*, 25, 359-372.

1508 Hammer, C. U., H. B. Clausen & C. C. Langway (1997) 50,000 years of recorded global volcanism. *Climatic*
1509 *Change*, 35, 1-15.

1510 Harrison, C. H. (1973) Radio-echo sounding of horizontal layers in ice. *Journal of Glaciology*, 12, 383-397.

1511 Hays, J. D., J. Imbrie & N. J. Shackleton (1976) Variations in the Earth's orbit: Pacemaker of the ice ages.
1512 *Science*, 194, 1121-1132.

1513 Hein, A., S. M. Marrero, J. Woodward, S. A. Dunning, K. Winter, M. J. Westoby, et al. (2016) Mid-Holocene
1514 pulse of thinning in the Weddell Sea sector of the West Antarctic ice sheet. *Nature Communications*,
1515 7, 12511.

1516 Heister, A. & R. Scheiber (2018) Coherent large-beamwidth processing of radio-echo sounding data. *The*
1517 *Cryosphere*, 12, 2969-2979.

1518 Hélière, F., C.-C. Lin, H. F. J. Corr & D. G. Vaughan (2007) Radio-echo sounding of Pine Island Glacier, West
1519 Antarctica: Aperture-synthesis processing and analysis of feasibility from space. *IEEE Transactions*
1520 *on Geoscience and Remote Sensing*, 45, 2573-2582.

1521 Hempel, L., F. Thyssen, N. Gundestrup, H. B. Clausen & H. Miller (2000) A comparison of radio-echo sounding
1522 data and electrical conductivity of the GRIP Ice Core. *Journal of Glaciology*, 46, 369-374.

1523 Henry, A. C. J., C. Schannwell, V. Višnjević, J. Millstein, P. D. Bons, O. Eisen, et al. (2024) Predicting the
1524 three-dimensional age-depth field of an ice rise. *Authorea*, DOI:
1525 10.22541/essoar.169230234.44865946/v1.

1526 Higgins, J. A., A. V. Kurbatov, N. E. Spaulding, E. Brook, D. S. Introne, L. M. Chimiak, et al. (2015) Atmospheric
1527 composition 1 million years ago from blue ice in the Allan Hills, Antarctica. *Proceedings of the National*
1528 *Academy of Sciences*, 112, 6887-6891.

1529 Hillebrand, T. R., J. O. Stone, M. Koutnik, C. King, H. Conway, B. Hall, et al. (2021) Holocene thinning of
1530 Darwin and Hatherton glaciers, Antarctica, and implications for grounding-line retreat in the Ross Sea.
1531 *The Cryosphere*, 15, 3329-3354.

1532 Hillenbrand, C.-D., J. A. Smith, D. A. Hodell, M. Greaves, C. R. Poole, S. Kender, et al. (2017) West Antarctic
1533 Ice Sheet retreat driven by Holocene warm water incursions. *Nature*, 547, 43-48.

1534 Hills, B. H., K. Christianson, A. O. Hoffman, T. J. Fudge, N. Holschuh, E. C. Kahle, et al. (2022) Geophysics
1535 and thermodynamics at South Pole Lake indicate stability and a regionally thawed bed. *Geophysical*
1536 *Research Letters*, 49, e2021GL096218.

1537 Hindmarsh, R. C. A., E. C. King, R. Mulvaney, H. F. J. Corr, G. Hiess & F. Gillet-Chaulet (2011) Flow at ice-
1538 divide triple junctions: 2. Three-dimensional views of isochrone architecture from ice-penetrating radar
1539 surveys. *Journal of Geophysical Research: Earth Surface*, 116, F02024.

1540 Hindmarsh, R. C. A., G. J.-M. C. Leysinger Vieli & F. Parrenin (2009) A large-scale numerical model for
1541 computing isochrone geometry. *Annals of Glaciology*, 50, 130-140.

1542 Hindmarsh, R. C. A., G. J.-M. C. Leysinger Vieli, M. J. Raymond & G. H. Gudmundsson (2006) Draping or
1543 overriding: The effect of horizontal stress gradients on internal layer architecture in ice sheets. *Journal*
1544 *of Geophysical Research: Earth Surface*, 111, F02018.

1545 Hodgkins, R., M. J. Siegert & J. A. Dowdeswell (2000) Geophysical investigations of ice-sheet internal layering
1546 and deformation in the Dome C region of central East Antarctica. *Journal of Glaciology*, 46, 161-166.

1547 Holland, P. R., H. F. J. Corr, D. G. Vaughan, A. Jenkins & P. Skvarca (2009) Marine ice in Larsen Ice Shelf.
1548 *Geophysical Research Letters*, 36, L11604.

1549 Holschuh, N., K. Christianson & S. Anandakrishnan (2014) Power loss in dipping internal reflectors, imaged
1550 using ice-penetrating radar. *Annals of Glaciology*, 55, 49-56.

1551 Holschuh, N., K. Christianson, H. Conway, R. Jacobel & B. Welch (2018) Persistent tracers of historic ice flow
1552 in glacial stratigraphy near Kamb Ice Stream, West Antarctica. *The Cryosphere*, 12, 2821 - 2829.

1553 Holschuh, N., D. A. Lilien & K. Christianson (2019) Thermal weakening, convergent flow, and vertical heat
1554 transport in the Northeast Greenland Ice Stream shear margins. *Geophysical Research Letters*, 46,
1555 8184-8193.

1556 Holschuh, N., B. R. Parizek, R. B. Alley & S. Anandakrishnan (2017) Decoding ice-sheet behavior using
1557 englacial layer slopes. *Geophysical Research Letters*, 44, 5561-5570.

1558 Humbert, A., D. Steinhage, V. Helm, S. Beyer & T. Kleiner (2018) Missing evidence of widespread subglacial
1559 lakes at Recovery Glacier, Antarctica. *Journal of Geophysical Research: Earth Surface*, 123, 2802-
1560 2826.

IceCube Collaboration (2013) South Pole glacial climate reconstruction from multi-borehole laser particulate stratigraphy. *Journal of Glaciology*, 59, 1117-1128.

Jacobel, R. W., A. M. Gades, D. L. Gottschling, S. M. Hodge & D. L. Wright (1993) Interpretation of radar-detected internal layer folding in West Antarctic ice streams. *Journal of Glaciology*, 39, 528-537.

Jacobel, R. W., T. A. Scambos, N. A. Nereson & C. F. Raymond (2000) Changes in the margin of Ice Stream C, Antarctica. *Journal of Glaciology*, 46, 102-110.

Jacobel, R. W. & B. C. Welch (2005) A time marker at 17.5 kyr BP detected throughout West Antarctica. *Annals of Glaciology*, 41, 47-51.

Jansen, D., S. Franke, C. C. Bauer, T. Binder, D. Dahl-Jensen, J. Eichler, et al. (2024) Shear margins in upper half of Northeast Greenland Ice Stream were established two millennia ago. *Nature Communications*, 15, 1193.

Jennings, S. J. A. & M. J. Hambrey (2021) Structures and deformation in glaciers and ice sheets. *Reviews of Geophysics*, 59, e2021RG000743.

Johnson, J. S., M. J. Bentley & K. Gohl (2008) First exposure ages from the Amundsen Sea Embayment, West Antarctica: The Late Quaternary context for recent thinning of Pine Island, Smith, and Pope Glaciers. *Geology*, 36, 223-226.

Jordan, T. A., C. Martín, F. Ferraccioli, K. Matsuoka, H. Corr, R. Forsberg, et al. (2018) Anomalously high geothermal flux near the South Pole. *Scientific Reports*, 8, 16785.

Jordan, T. M., C. Martín, A. M. Brisbourne, D. M. Schroeder & A. M. Smith (2022) Radar characterization of ice-crystal-orientation fabric and anisotropic viscosity within an Antarctic ice stream. *Journal of Geophysical Research: Earth Surface*, 127, e2022JF006673.

Jordan, T. M., D. M. Schroeder, C. W. Elsworth & M. R. Siegfried (2020) Estimation of ice fabric within Whillans Ice Stream using polarimetric phase-sensitive radar sounding. *Annals of Glaciology*, 61, 74-83.

Jouzel, J., V. Masson-Delmotte, O. Cattani, G. Dreyfus, S. Falourd, G. Hoffmann, et al. (2007) Orbital and millennial Antarctic climate variability over the past 800,000 years. *Science*, 317, 793-796.

Karlsson, N. B., T. Binder, G. Eagles, V. Helm, F. Pattyn, B. Van Liefferinge, et al. (2018) Glaciological characteristics in the Dome Fuji region and new assessment for "Oldest Ice". *The Cryosphere*, 12, 2413-2424.

Karlsson, N. B., R. G. Bingham, D. M. Rippin, R. C. A. Hindmarsh, H. F. J. Corr & D. G. Vaughan (2014) Constraining past accumulation in the central Pine Island Glacier basin, West Antarctica, using radio-echo sounding. *Journal of Glaciology*, 60, 553-562.

Karlsson, N. B., D. M. Rippin, R. G. Bingham & D. G. Vaughan (2012) A 'continuity-index' for assessing ice-sheet dynamics from radar-sounded internal layers. *Earth and Planetary Science Letters*, 335-336, 88-94.

Karlsson, N. B., D. M. Rippin, D. G. Vaughan & H. F. J. Corr (2009) The internal layering of Pine Island Glacier, West Antarctica, from airborne radar-sounding data. *Annals of Glaciology*, 50, 141-146.

King, E. C. (2011) Ice stream or not? Radio-echo sounding of Carlson Inlet, West Antarctica. *The Cryosphere*, 5, 907-916.

Kingslake, J., R. C. A. Hindmarsh, G. Aðalgeirsdóttir, H. Conway, H. F. J. Corr, F. Gillet-Chaulet, et al. (2014) Full-depth englacial vertical ice-sheet velocities measured using phase-sensitive radar. *Journal of Geophysical Research: Earth Surface*, 119, 2604-2618.

Kingslake, J., C. Martín, R. J. Arthern, H. F. J. Corr & E. C. King (2016) Ice-flow reorganization in West Antarctica 2.5 kyr ago dated using radar-derived englacial flow velocities. *Geophysical Research Letters*, 43, 9103-9112.

Kingslake, J., R. P. Scherer, T. Albrecht, J. Coenen, R. D. Powell, R. Reese, et al. (2018) Extensive retreat and re-advance of the West Antarctic Ice Sheet during the Holocene. *Nature*, 558, 430-434.

Koch, I., R. Drews, O. Eisen, S. Franke, D. Jansen, K. Matsuoka, et al. (2023) Radar internal reflection horizons from multisystem data reflect ice dynamic and surface accumulation history along the Princess Ragnhild Coast, Dronning Maud Land, East Antarctica. *Journal of Glaciology*, 70, e18.

Konrad, H., A. E. Hogg, R. Mulvaney, R. Arthern, R. J. Tuckwell, B. Medley, et al. (2019) Observations of surface mass balance on Pine Island Glacier, West Antarctica, and the effect of strain history in fast-flowing sections. *Journal of Glaciology*, 65, 595-604.

Koutnik, M. R., T. J. Fudge, H. Conway, E. D. Waddington, T. A. Neumann, K. M. Cuffey, et al. (2016) Holocene accumulation and ice flow near the West Antarctic Ice Sheet Divide ice-core site. *Journal of Geophysical Research: Earth Surface*, 121, 907-924.

Kovacs, A., A. J. Gow & R. M. Morey (1995) The in-situ dielectric constant of polar firn revisited. *Cold Regions Science and Technology*, 23, 245-256.

Kowalewski, S., V. Helm, E. M. Morris & O. Eisen (2021) The regional-scale surface mass balance of Pine Island Glacier, West Antarctica, over the period 2005–2014, derived from airborne radar soundings and neutron probe measurements. *The Cryosphere*, 15, 1285-1305.

- Laird, C. M., W. A. Blake, K. Matsuoka, H. Conway, C. T. Allen, C. J. Leuschen, et al. (2010) Deep ice stratigraphy and basal conditions in central West Antarctica revealed by coherent radar. *IEEE Geoscience and Remote Sensing Letters*, 7, 246-250.
- Le Meur, E., O. Magand, L. Arnaud, M. Fily, M. Frezzotti, M. Cavitte, et al. (2018) Spatial and temporal distributions of surface mass balance between Concordia and Vostok stations, Antarctica, from combined radar and ice-core data: First results and detailed error analysis. *The Cryosphere*, 12, 1831-1850.
- Lecavalier, B. S., L. Tarasov, G. Balco, P. Spector, C. D. Hillenbrand, C. Buizert, et al. (2023) Antarctic Ice Sheet paleo-constraint database. *Earth System Science Data*, 15, 3573-3596.
- Lee, H., H. Seo, H. Han, H. Ju & J. Lee (2021) Velocity anomaly of Campbell Glacier, East Antarctica, observed by double-differential interferometric SAR and ice-penetrating radar. *Remote Sensing*, 13, 2691.
- Legrain, E., F. Parrenin & E. Capron (2023) A gradual change is more likely to have caused the Mid-Pleistocene Transition than an abrupt event. *Communications Earth & Environment*, 4, 90.
- Lenaerts, J. T. M., B. Medley, M. R. van den Broeke & B. Wouters (2019) Observing and modeling ice-sheet surface mass balance. *Reviews of Geophysics*, 57, 376-420.
- Leysinger Vieli, G. J.-M. C., R. C. A. Hindmarsh & M. J. Siegert (2007) Three-dimensional flow influences on radar layer stratigraphy. *Annals of Glaciology*, 46, 22-28.
- Leysinger Vieli, G. J.-M. C., R. C. A. Hindmarsh, M. J. Siegert & B. Sun (2011) Time-dependence of the spatial pattern of accumulation rate in East Antarctica deduced from isochronic radar layers using a 3-D numerical ice flow model. *Journal of Geophysical Research: Earth Surface*, 116, F02018.
- Leysinger Vieli, G. J.-M. C., C. Martín, R. C. A. Hindmarsh & M. P. Lüthi (2018) Basal freeze-on generates complex ice-sheet stratigraphy. *Nature Communications*, 9, 4669.
- Leysinger Vieli, G. J.-M. C., M. J. Siegert & A. J. Payne (2004) Reconstructing ice-sheet accumulation rates at Ridge B, East Antarctica. *Annals of Glaciology*, 39, 326-330.
- Lilien, D. A., D. Steinhage, D. Taylor, F. Parrenin, C. Ritz, R. Mulvaney, et al. (2021) New radar constraints support presence of ice older than 1.5 Myr at Little Dome C. *The Cryosphere*, 15, 1881-1888.
- Lindzey, L. E., L. H. Beem, D. A. Young, E. Quartini, D. D. Blankenship, C. K. Lee, et al. (2020) Aero-geophysical characterization of an active subglacial lake system in the David Glacier catchment, Antarctica. *The Cryosphere*, 14, 2217-2233.
- Luo, K., S. Liu, J. Guo, T. Wang, L. Li, X. Cui, et al. (2020) Radar-derived internal structure and basal roughness characterization along a traverse from Zhongshan Station to Dome A, East Antarctica. *Remote Sensing*, 12, 1079.
- Luo, K., X. Tang, S. Liu, J. Guo, J. S. Greenbaum, L. Li, et al. (2022) Deep radiostratigraphy constraints support the presence of persistent wind scouring behavior for more than 100 ka in the East Antarctic Ice Sheet. *IEEE Transactions on Geoscience and Remote Sensing*, 60, 4306213.
- MacGregor, J. A., L. N. Boisvert, B. Medley, A. A. Petty, J. P. Harbeck, R. E. Bell, et al. (2021) The scientific legacy of NASA's Operation IceBridge. *Reviews of Geophysics*, 59, e2020RG000712.
- MacGregor, J. A., W. T. Colgan, M. A. Fahnestock, M. Morlighem, G. A. Catania, J. D. Paden, et al. (2016) Holocene deceleration of the Greenland Ice Sheet. *Science*, 351, 590-593.
- MacGregor, J. A., M. A. Fahnestock, G. A. Catania, J. D. Paden, S. P. Gogineni, S. K. Young, et al. (2015a) Radiostratigraphy and age structure of the Greenland Ice Sheet. *Journal of Geophysical Research: Earth Surface*, 120, 212-241.
- MacGregor, J. A., M. A. Fahnestock, J. D. Paden, J. Li, J. P. Harbeck & A. Aschwanden (2025) A revised and expanded deep radiostratigraphy of the Greenland Ice Sheet from airborne radar sounding surveys between 1993–2019. *Earth System Science Data Discussions*, <https://doi.org/10.5194/essd-2024-578>, in review, 2025.
- MacGregor, J. A., J. L. Li, J. D. Paden, G. A. Catania, G. D. Clow, M. A. Fahnestock, et al. (2015b) Radar attenuation and temperature within the Greenland Ice Sheet. *Journal of Geophysical Research: Earth Surface*, 120, 983-1008.
- MacGregor, J. A., K. Matsuoka, M. R. Koutnik, E. D. Waddington, M. Studinger & D. P. Winebrenner (2009) Millennially averaged accumulation rates for the Vostok Subglacial Lake region inferred from deep internal layers. *Annals of Glaciology*, 50, 25-34.
- MacGregor, J. A., D. P. Winebrenner, H. Conway, K. Matsuoka, P. A. Mayewski & G. D. Clow (2007) Modeling englacial radar attenuation at Siple Dome, West Antarctica, using ice chemistry and temperature data. *Journal of Geophysical Research: Earth Surface*, 112, F03008.
- Mackintosh, A. N., E. Verleyen, P. E. O'Brien, D. A. White, R. S. Jones, R. McKay, et al. (2014) Retreat history of the East Antarctic Ice Sheet since the Last Glacial Maximum. *Quaternary Science Reviews*, 100, 10-30.
- Martín, C. & G. H. Gudmundsson (2012) Effects of nonlinear rheology, temperature and anisotropy on the relationship between age and depth at ice divides. *The Cryosphere*, 6, 1221-1229.
- Martín, C., G. H. Gudmundsson & E. C. King (2014) Modelling of Kealey Ice Rise, Antarctica, reveals stable ice-flow conditions in East Ellsworth Land over millennia. *Journal of Glaciology*, 60, 139-146.

- 1683 Martín, C., R. C. A. Hindmarsh & F. J. Navarro (2006) Dating ice flow change near the flow divide at Roosevelt
1684 Island, Antarctica, by using a thermomechanical model to predict radar stratigraphy. *Journal of*
1685 *Geophysical Research: Earth Surface*, 111, F01011.
- 1686 Martín, C., R. C. A. Hindmarsh & F. J. Navarro (2009) On the effects of divide migration, along-ridge flow, and
1687 basal sliding on isochrones near an ice divide. *Journal of Geophysical Research: Earth Surface*, 114,
1688 F02006.
- 1689 Matsuoka, K., T. Furukawa, S. Fujita, H. Maeno, S. Uratsuka, R. Naruse, et al. (2003) Crystal-orientation
1690 fabrics within the Antarctic Ice Sheet revealed by a multipolarization plane and dual-frequency radar
1691 survey. *Journal of Geophysical Research: Solid Earth*, 108, 2499.
- 1692 Matsuoka, K., A. Gades, H. Conway, G. Catania & C. F. Raymond (2009) Radar signatures beneath a surface
1693 topographic lineation near the outlet of Kamb Ice Stream and Engelhardt Ice Ridge, West Antarctica.
1694 *Annals of Glaciology*, 50, 98-104.
- 1695 Matsuoka, K., R. C. A. Hindmarsh, G. Moholdt, M. J. Bentley, H. D. Pritchard, J. Brown, et al. (2015) Antarctic
1696 ice rises and rumpled: Their properties and significance for ice-sheet dynamics and evolution. *Earth-*
1697 *Science Reviews*, 150, 724-745.
- 1698 Matsuoka, K., J. A. MacGregor & F. Pattyn (2012) Predicting radar attenuation within the Antarctic Ice Sheet.
1699 *Earth and Planetary Science Letters*, 359-360, 173-183.
- 1700 Mayewski, P. A., M. Frezzotti, N. Bertler, T. van Ommen, G. Hamilton, T. H. Jacka, et al. (2005) The
1701 International Trans-Antarctic Scientific Expedition (ITASE): an overview. *Annals of Glaciology*, 41,
1702 180-185.
- 1703 McConnell, J. R., A. Burke, N. W. Dunbar, P. Köhler, J. L. Thomas, M. M. Arienzo, et al. (2017) Synchronous
1704 volcanic eruptions and abrupt climate change ~17.7 ka plausibly linked by stratospheric ozone
1705 depletion. *Proceedings of the National Academy of Sciences*, 114, 10035-10040.
- 1706 Medley, B., I. Joughin, S. B. Das, E. J. Steig, H. Conway, S. Gogineni, et al. (2013) Airborne-radar and ice-
1707 core observations of annual snow accumulation over Thwaites Glacier, West Antarctica, confirm the
1708 spatiotemporal variability of global and regional atmospheric models. *Geophysical Research Letters*,
1709 40, 3649-3654.
- 1710 Medley, B., I. Joughin, B. E. Smith, S. B. Das, E. J. Steig, H. Conway, et al. (2014) Constraining the recent
1711 mass balance of Pine Island and Thwaites glaciers, West Antarctica, with airborne observations of
1712 snow accumulation. *The Cryosphere*, 8, 1375-1392.
- 1713 Millar, D. H. M. (1981) Radio-echo layering in polar ice sheets and past volcanic activity. *Nature*, 292, 441-443.
- 1714 Millar, D. H. M. (1982) Acidity levels in ice sheets from radio echo-sounding. *Annals of Glaciology*, 3, 199-203.
- 1715 Miners, W. D., A. Hildebrand, S. Gerland, N. Blindow, D. Steinhage & E. W. Wolff (1997) Forward modeling of
1716 the internal layers in radio-echo sounding using electrical and density measurements from ice cores.
1717 *Journal of Physical Chemistry B*, 101, 6201-6204.
- 1718 Mojtavavi, S., O. Eisen, S. Franke, D. Jansen, D. Steinhage, J. Paden, et al. (2022) Origin of englacial
1719 stratigraphy at three deep ice core sites of the Greenland Ice Sheet by synthetic radar modelling.
1720 *Journal of Glaciology*, 68, 799-811.
- 1721 Moqadam, H., D. Steinhage, A. Wilhelm & O. Eisen (2025) Going deeper with deep learning: Automatically
1722 tracing internal reflection horizons in ice sheets. In review for *Machine Learning and Computation*,
1723 <https://doi.org/10.22541/essoar.172987463.39597493/v1>.
- 1724 Morgan, V., T. Jacka, G. Akerman & A. Clarke (1982) Outlet glacier and mass-budget studies in Enderby,
1725 Kemp, and Mac. Robertson Lands, Antarctica. *Annals of Glaciology*, 3, 204-210.
- 1726 Morlighem, M. (2020) MEaSUREs BedMachine Antarctica, Version 2. NASA National Snow and Ice Data
1727 Center Distributed Active Archive Center.
- 1728 Moss, G., V. Višnjević, O. Eisen, F. M. Oraschewski, C. Schröder, J. H. Macke, et al. (2023) Simulation-based
1729 inference of surface accumulation and basal-melt rates of an Antarctic ice shelf from isochronal layers.
1730 *ArXiv*, abs/2312.02997.
- 1731 Moussessian, A., R. L. Jordan, E. Rodriguez, A. Safaeinili, T. L. Akins, W. N. Edelstein, et al. (2000) A new
1732 coherent radar for ice sounding in Greenland. In *International Geoscience and Remote Sensing*
1733 *Symposium (IGARRS) Proceedings*, 484-486 vol.2.
- 1734 Muldoon, G. R., C. S. Jackson, D. A. Young & D. D. Blankenship (2018) Bayesian estimation of englacial radar
1735 chronology in Central West Antarctica. *Dynamics and Statistics of the Climate System*, 3, dzy004.
- 1736 Naish, T., R. Powell, R. Levy, G. Wilson, R. Scherer, F. Talarico, et al. (2009) Obliquity-paced Pliocene West
1737 Antarctic ice sheet oscillations. *Nature*, 458, 322-328.
- 1738 Narcisi, B. & J. R. Petit (2021) Englacial tephtras of East Antarctica. *Geological Society, London, Memoirs*, 55,
1739 649-664.
- 1740 NEEM Community Members (2013) Eemian interglacial reconstructed from a Greenland folded ice core.
1741 *Nature*, 493, 489-494.
- 1742 Nereson, N. A., C. F. Raymond, R. W. Jacobel & E. D. Waddington (2000) The accumulation pattern across
1743 Siple Dome, West Antarctica, inferred from radar-detected internal layers. *Journal of Glaciology*, 46,
1744 75-87.

- 1745 Nereson, N. A., C. F. Raymond, E. D. Waddington & R. W. Jacobel (1998) Migration of the Siple Dome ice
1746 divide, West Antarctica. *Journal of Glaciology*, 44, 643-652.
- 1747 Neumann, T. A., H. Conway, S. F. Price, E. D. Waddington, G. A. Catania & D. L. Morse (2008) Holocene
1748 accumulation and ice-sheet dynamics in central West Antarctica. *Journal of Geophysical Research:*
1749 *Earth Surface*, 113, F02018.
- 1750 Ng, F. & H. Conway (2004) Fast-flow signature in the stagnated Kamb Ice Stream, West Antarctica. *Geology*,
1751 32, 481-484.
- 1752 Ng, F. & E. C. King (2011) Kinematic waves in polar firn stratigraphy. *Journal of Glaciology*, 57, 1119-1134.
- 1753 Nicholls, K. W., H. F. J. Corr, C. L. Stewart, L. B. Lok, P. V. Brennan & D. G. Vaughan (2015) A ground-based
1754 radar for measuring vertical strain rates and time-varying basal melt rates in ice sheets and shelves.
1755 *Journal of Glaciology*, 61, 1079-1087.
- 1756 Nye, J. (1957) The distribution of stress and velocity in glaciers and ice sheets. *Proceedings of the Royal*
1757 *Society A: Mathematical, Physical and Engineering Sciences*, 239, 113-133.
- 1758 Obase, T., A. Abe-Ouchi, F. Saito, S. Tsutaki, S. Fujita, K. Kawamura, et al. (2023) A one-dimensional
1759 temperature and age modeling study for selecting the drill site of the oldest ice core near Dome Fuji,
1760 Antarctica. *The Cryosphere*, 17, 2543-2562.
- 1761 Oppenheimer, M., B. C. Glavovic, J. Hinkel, R. van de Wal, A. K. Magnan, A. Abd-Elgawad, et al. 2019. Sea-
1762 level rise and implications for low-lying islands, coasts and communities. In *IPCC Special Report on*
1763 *the Ocean and Cryosphere in a Changing Climate*, eds. H.-O. Pörtner, D. C. Roberts, V. Masson-
1764 Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegria, M. Nicolai, A. Okem, J.
1765 Petzold, B. Rama & N. M. Weyer, 321-445. Cambridge: Cambridge University Press.
- 1766 Oraschewski, F. M., I. Koch, M. R. Ershadi, J. Hawkins, O. Eisen & R. Drews (2023) Layer-optimized SAR
1767 processing with a mobile phase-sensitive radar for detecting the deep englacial stratigraphy of Colle
1768 Gnifetti, Switzerland/Italy. *The Cryosphere*, 18, 3875-3889.
- 1769 Otosaka, I. N., A. Shepherd, E. R. Ivins, N. J. Schlegel, C. Amory, M. R. van den Broeke, et al. (2023) Mass
1770 balance of the Greenland and Antarctic ice sheets from 1992 to 2020. *Earth System Science Data*, 15,
1771 1597-1616.
- 1772 Paden, J., K. Tinto, D. Young, K. Christianson, D. Schroeder, M. Rahnmooonfar, et al. (2021) Open Polar
1773 Radar software and services to standardize radar echograms and integrate into a geospatial
1774 database. . In *AGU Fall Meeting 2021*, 2021AGUFM.C51A.08P.
- 1775 Panton, C. (2014) Automated mapping of local layer slope and tracing of internal layers in radio echograms.
1776 *Annals of Glaciology*, 55, 71-77.
- 1777 Panton, C. & N. B. Karlsson (2015) Automated mapping of near-bed radio-echo layer disruptions in the
1778 Greenland Ice Sheet. *Earth and Planetary Science Letters*, 432, 323-331.
- 1779 Parrenin, F., M. G. P. Cavitte, D. D. Blankenship, J. Chappellaz, H. Fischer, O. Gagliardini, et al. (2017) Is
1780 there 1.5-million-year-old ice near Dome C, Antarctica? *The Cryosphere*, 11, 2427-2437.
- 1781 Parrenin, F. & R. Hindmarsh (2007) Influence of a non-uniform velocity field on isochrone geometry along a
1782 steady flowline of an ice sheet. *Journal of Glaciology*, 53, 612-622.
- 1783 Passalacqua, O., C. Ritz, F. Parrenin, S. Urbini & M. Frezzotti (2017) Geothermal flux and basal melt rate in
1784 the Dome C region inferred from radar reflectivity and heat modelling. *The Cryosphere*, 11, 2231-2246.
- 1785 Pattyn, F. (2010) Antarctic subglacial conditions inferred from a hybrid ice-sheet/ice-stream model. *Earth and*
1786 *Planetary Science Letters*, 295, 451-461.
- 1787 Pauling, A. G., C. M. Bitz & E. J. Steig (2023) Linearity of the climate-system response to raising and lowering
1788 West Antarctic and coastal Antarctic topography. *Journal of Climate*, 36, 6195-6212.
- 1789 Peters, M. E., D. D. Blankenship, S. P. Carter, S. D. Kempf, D. A. Young & J. W. Holt (2007) Along-track
1790 focusing of airborne radar sounding data from West Antarctica for improving basal reflection analysis
1791 and layer detection. *IEEE Transactions on Geoscience and Remote Sensing*, 45, 2725-2736.
- 1792 Peters, M. E., D. D. Blankenship & D. L. Morse (2005) Analysis techniques for coherent airborne radar
1793 sounding: Application to West Antarctic ice streams. *Journal of Geophysical Research: Solid Earth*,
1794 110, B06303.
- 1795 Pettit, E. C., T. Thorsteinsson, H. P. Jacobson & E. D. Waddington (2007) The role of crystal fabric in flow near
1796 an ice divide. *Journal of Glaciology*, 53, 277-288.
- 1797 Pettit, E. C. & E. D. Waddington (2003) Ice flow at low deviatoric stress. *Journal of Glaciology*, 49, 359-369.
- 1798 Pettit, E. C., E. D. Waddington, W. D. Harrison, T. Thorsteinsson, D. Elsberg, J. Morack, et al. (2011) The
1799 crossover stress, anisotropy and the ice flow law at Siple Dome, West Antarctica. *Journal of Glaciology*,
1800 57, 39-52.
- 1801 Pittard, M. L., P. L. Whitehouse, M. J. Bentley & D. Small (2022) An ensemble of Antarctic deglacial simulations
1802 constrained by geological observations. *Quaternary Science Reviews*, 298, 107800.
- 1803 Pollard, D. & R. M. DeConto (2009) Modelling West Antarctic Ice Sheet growth and collapse through the past
1804 five million years. *Nature*, 458, 329-332.
- 1805 Popov, S. (2020) Fifty-five years of Russian radio-echo sounding investigations in Antarctica. *Annals of*
1806 *Glaciology*, 61, 14-24.

- 1807 Popov, S. (2022) Ice cover, subglacial landscape, and estimation of bottom melting of Mac. Robertson,
1808 Princess Elizabeth, Wilhelm II, and western Queen Mary Lands, East Antarctica. *Remote Sensing*, 14,
1809 241.
- 1810 Pratap, B., R. Dey, K. Matsuoka, G. Moholdt, K. Lindbäck, V. Goel, et al. (2022) Three-decade spatial patterns
1811 in surface mass balance of the Nivlisen Ice Shelf, central Dronning Maud Land, East Antarctica.
1812 *Journal of Glaciology*, 68, 174-186.
- 1813 Pritchard, H. D., P. T. Fretwell, A. C. Frémand, J. A. Bodart, J. D. Kirkham, et al. (2025) Bedmap3 updated ice
1814 bed, surface and thickness datasets for Antarctica. *Scientific Data*, 12, 414.
- 1815 Rahnemoonfar, M., M. Yari, J. Paden, L. Koenig & O. Ibikunle (2021) Deep multi-scale learning for automatic
1816 tracking of internal layers of ice in radar data. *Journal of Glaciology*, 67, 39-48.
- 1817 Raymond, C. F. (1983) Deformation in the vicinity of ice divides. *Journal of Glaciology*, 29, 357-373.
- 1818 Reeh, N., H. Oerter & H. H. Thomsen (2002) Comparison between Greenland ice-margin and ice-core oxygen-
1819 18 records. *Annals of Glaciology*, 35, 136-144.
- 1820 Retzlaff, R., N. Lord & C. R. Bentley (1993) Airborne-radar studies: Ice Streams A, B and C, West Antarctica.
1821 *Journal of Glaciology*, 39, 495-506.
- 1822 Rieckh, T., A. Born, A. Robinson, R. Law & G. Gülle (2024) Introducing ELSA v2.0: An isochronal model for
1823 ice-sheet layer tracing. *Geoscientific Model Development*, 17, 6987-7000.
- 1824 Rignot, E., J. Mouginot & B. Scheuchl (2017) MEaSUREs InSAR-Based Antarctica Ice Velocity Map, Version
1825 2. NASA National Snow and Ice Data Center Distributed Active Archive Center.
- 1826 Rignot, E., R. H. Thomas, P. Kanagaratnam, G. Casassa, E. Frederick, S. Gogineni, et al. (2004) Improved
1827 estimation of the mass balance of glaciers draining into the Amundsen Sea sector of West Antarctica
1828 from the CECS/NASA 2002 campaign. *Annals of Glaciology*, 39, 231-237.
- 1829 Rippin, D. M., M. J. Siegert & J. L. Bamber (2003) The englacial stratigraphy of Wilkes Land, East Antarctica,
1830 as revealed by internal radio-echo sounding layering, and its relationship with balance velocities.
1831 *Annals of Glaciology*, 36, 189-196.
- 1832 Rippin, D. M., M. J. Siegert, J. L. Bamber, D. G. Vaughan & H. F. J. Corr (2006) Switch-off of a major enhanced
1833 ice-flow unit in East Antarctica. *Geophysical Research Letters*, 33, L15501.
- 1834 Rivera, A., J. Uribe, R. Zamora & J. Oberreuter (2015) Subglacial Lake CECs: Discovery and in situ survey of
1835 a privileged research site in West Antarctica. *Geophysical Research Letters*, 42, 3944-3953.
- 1836 Rix, J., R. Mulvaney, J. Hong & D. A. N. Ashurst (2019) Development of the British Antarctic Survey Rapid-
1837 Access Isotope Drill. *Journal of Glaciology*, 65, 288-298.
- 1838 Robin, G. d. Q., S. Evans & J. T. Bailey (1969) Interpretation of radio echo sounding in polar ice sheets.
1839 *Philosophical Transactions of the Royal Society of London Series A - Mathematical and Physical
1840 Sciences*, 265, 437-505.
- 1841 Robin, G. d. Q. & D. H. M. Millar (1982) Flow of ice sheets in the vicinity of subglacial peaks. *Annals of
1842 Glaciology*, 3, 290-294.
- 1843 Rodriguez-Morales, F., S. Gogineni, C. J. Leuschen, J. D. Paden, J. Li, C. C. Lewis, et al. (2013) Advanced
1844 multifrequency radar instrumentation for polar research. *IEEE Transactions on Geoscience and
1845 Remote Sensing*, 52, 2824-2842.
- 1846 Ross, N., H. Corr & M. Siegert (2020) Large-scale englacial folding and deep-ice stratigraphy within the West
1847 Antarctic Ice Sheet. *The Cryosphere*, 14, 2103-2114.
- 1848 Ross, N. & M. Siegert (2020) Basal melting over Subglacial Lake Ellsworth and its catchment: Insights from
1849 englacial layering. *Annals of Glaciology*, 61, 198-205.
- 1850 Ross, N., M. J. Siegert, J. Woodward, A. M. Smith, H. F. J. Corr, M. J. Bentley, et al. (2011) Holocene stability
1851 of the Amundsen-Weddell ice divide, West Antarctica. *Geology*, 39, 935-938.
- 1852 Rowell, I. F., R. Mulvaney, J. Rix, D. R. Tetzner & E. W. Wolff (2023) Viability of chemical and water isotope
1853 ratio measurements of RAID ice chippings from Antarctica. *Journal of Glaciology*, 69, 623-638.
- 1854 Sanderson, R. J., N. Ross, K. Winter, R. G. Bingham, S. L. Callard, T. A. Jordan, et al. (2024) Dated radar-
1855 stratigraphy between Dome A and South Pole, East Antarctica: Old-ice potential and ice-sheet history.
1856 *Journal of Glaciology*.
- 1857 Sanderson, R. J., K. Winter, S. L. Callard, F. Napoleoni, N. Ross, T. A. Jordan, et al. (2023) Englacial
1858 architecture of Lambert Glacier, East Antarctica. *The Cryosphere*, 17, 4853-4871.
- 1859 Scambos, T. A., R. E. Bell, R. B. Alley, S. Anandakrishnan, D. H. Bromwich, K. Brunt, et al. (2017) How much,
1860 how fast?: A science review and outlook for research on the instability of Antarctica's Thwaites Glacier
1861 in the 21st century. *Global and Planetary Change*, 153, 16-34.
- 1862 Scanlan, K. M., A. Rutishauser, D. A. Young & D. D. Blankenship (2020) Interferometric discrimination of cross-
1863 track bed clutter in ice-penetrating radar sounding data. *Annals of Glaciology*, 61, 68-73.
- 1864 Schannwell, C., R. Drews, T. A. Ehlers, O. Eisen, C. Mayer & F. Gillet-Chaulet (2019) Kinematic response of
1865 ice-rise divides to changes in ocean and atmosphere forcing. *The Cryosphere*, 13, 2673-2691.
- 1866 Schroeder, D. M., R. G. Bingham, D. D. Blankenship, K. Christianson, O. Eisen, G. E. Flowers, et al. (2020)
1867 Five decades of radioglaciology. *Annals of Glaciology*, 61, 1-13.
- 1868 Schroeder, D. M., A. L. Broome, A. Conger, A. Lynch, E. J. Mackie & A. Tarzona (2022) Radiometric analysis
1869 of digitized Z-scope records in archival radar sounding film. *Journal of Glaciology*, 68, 733-740.

- Schroeder, D. M., J. A. Dowdeswell, M. J. Siegert, R. G. Bingham, W. N. Chu, E. J. MacKie, et al. (2019) Multidecadal observations of the Antarctic Ice Sheet from restored analog radar records. *Proceedings of the National Academy of Sciences*, 116, 18867-18873.
- Schwander, J., T. F. Stocker, R. Walther & S. Marending (2023) Progress of the RADIX (Rapid Access Drilling and Ice eXtraction) fast-access drilling system. *The Cryosphere*, 17, 1151-1164.
- Seroussi, H., S. Nowicki, A. J. Payne, H. Goelzer, W. H. Lipscomb, A. Abe-Ouchi, et al. (2020) ISMIP6 Antarctica: a multi-model ensemble of the Antarctic Ice Sheet evolution over the 21st century. *The Cryosphere*, 14, 3033-3070.
- Seroussi, H., T. Pelle, W. H. Lipscomb, et al. (2024) Evolution of the Antarctic Ice Sheet over the next three centuries from an ISMIP6 model ensemble. *Earth's Future*, 12, e2024EF004561.
- Siegert, M., N. Ross, H. Corr, J. Kingslake & R. Hindmarsh (2013) Late Holocene ice-flow reconfiguration in the Weddell Sea sector of West Antarctica. *Quaternary Science Reviews*, 78, 98-107.
- Siegert, M. J. (2003) Glacial–interglacial variations in central East Antarctic ice accumulation rates. *Quaternary Science Reviews*, 22, 741-750.
- Siegert, M. J., J. C. Ellis-Evans, M. Tranter, C. Mayer, J.-R. Petit, A. Salamatin, et al. (2001) Physical, chemical and biological processes in Lake Vostok and other Antarctic subglacial lakes. *Nature*, 414, 603-609.
- Siegert, M. J., R. C. A. Hindmarsh & G. S. Hamilton (2003a) Evidence for a large surface ablation zone in central East Antarctica during the last Ice Age. *Quaternary Research*, 59, 114-121.
- Siegert, M. J. & R. Hodgkins (2000) A stratigraphic link across 1100 km of the Antarctic Ice Sheet between the Vostok ice-core site and Titan Dome (near South Pole). *Geophysical Research Letters*, 27, 2133-2136.
- Siegert, M. J., R. Hodgkins & J. A. Dowdeswell (1998) A chronology for the Dome C deep ice-core site through radio-echo layer correlation with the Vostok Ice Core, Antarctica. *Geophysical Research Letters*, 25, 1019-1022.
- Siegert, M. J., R. Kwok, C. Mayer & B. Hubbard (2000) Water exchange between the subglacial Lake Vostok and the overlying ice sheet. *Nature*, 403, 643-646.
- Siegert, M. J. & A. J. Payne (2004) Past rates of accumulation in central West Antarctica. *Geophysical Research Letters*, 31, L12403.
- Siegert, M. J., A. J. Payne & I. Joughin (2003b) Spatial stability of Ice Stream D and its tributaries, West Antarctica, revealed by radio-echo sounding and interferometry. *Annals of Glaciology*, 37, 377-382.
- Siegert, M. J., B. Welch, D. Morse, A. Vieli, D. D. Blankenship, I. Joughin, et al. (2004) Ice-flow direction change in interior West Antarctica. *Science*, 305, 1948-1951.
- Sigl, M., M. Toohey, J. R. McConnell, J. Cole-Dai & M. Severi (2022) Volcanic stratospheric sulfur injections and aerosol optical depth during the Holocene (past 11,500 years) from a bipolar ice-core array. *Earth System Science Data*, 14, 3167-3196.
- Sime, L. C., R. C. A. Hindmarsh & H. Corr (2011) Automated processing to derive dip angles of englacial radar reflectors in ice sheets. *Journal of Glaciology*, 57, 260-266.
- Sime, L. C., N. B. Karlsson, J. D. Paden & S. P. Gogineni (2014) Isochronous information in a Greenland ice sheet radio echo sounding data set. *Geophysical Research Letters*, 41, 1593-1599.
- Smith, B. E., N. E. Lord & C. R. Bentley (2002) Crevasse ages on the northern margin of Ice Stream C, West Antarctica. *Annals of Glaciology*, 34, 209-216.
- Spaulding, N. E., J. A. Higgins, A. V. Kurbatov, M. L. Bender, S. A. Arcone, S. Campbell, et al. (2013) Climate archives from 90 to 250 ka in horizontal and vertical ice cores from the Allan Hills Blue Ice Area, Antarctica. *Quaternary Research*, 80, 562-574.
- Steinhage, D., S. Kipfstuhl, U. Nixdorf & H. Miller (2013) Internal structure of the ice sheet between Kohnen station and Dome Fuji, Antarctica, revealed by airborne radio-echo sounding. *Annals of Glaciology*, 54, 163-167.
- Steinhage, D., U. Nixdorf, U. Meyer & H. Miller (2001) Subglacial topography and internal structure of central and western Dronning Maud Land, Antarctica, determined from airborne radio-echo sounding. *Journal of Applied Geophysics*, 47, 183-189.
- Stone, J.O., G. A. Balco, D. E. Sugden, M. W. Caffee, L. C. Sass III, S. G. Cowdery & C. Siddoway (2003) Holocene deglaciation of Marie Byrd Land, West Antarctica, *Science*, 299, 99-102.
- Sutter, J., H. Fischer & O. Eisen (2021) Investigating the internal structure of the Antarctic Ice Sheet: The utility of isochrones for spatiotemporal ice-sheet model calibration. *The Cryosphere*, 15, 3839-3860.
- Sutter, J., H. Fischer, K. Grosfeld, N. B. Karlsson, T. Kleiner, B. Van Liefferinge, et al. (2019) Modelling the Antarctic Ice Sheet across the mid-Pleistocene transition: Implications for Oldest Ice. *The Cryosphere*, 13, 2023-2041.
- Svensson, A., M. Bigler, T. Blunier, H. B. Clausen, D. Dahl-Jensen, H. Fischer, et al. (2013) Direct linking of Greenland and Antarctic ice cores at the Toba eruption (74 ka BP). *Climate of the Past*, 9, 749-766.
- Swithinbank, C. (1969) Airborne radio-echo sounding by the British Antarctic Survey. *The Geographical Journal*, 135, 551-553.
- Tabacco, I., C. Bianchi, J. Baskaradas, L. Cafarella, S. Umberto, A. Zirizzotti, et al. (2008) Italian RES investigation in Antarctica: The new radar system. *Terra Antarctica Reports*, 14, 213 - 216.

- 1932 Tang, X., K. Luo, S. Dong, Z. Zhang & B. Sun (2022) Quantifying basal roughness and internal-layer continuity
1933 index of ice sheets by an integrated means with radar data and deep learning. *Remote Sensing*, 14,
1934 4507.
- 1935 Tarasov, L. & W. R. Peltier (2003) Greenland glacial history, borehole constraints, and Eemian extent. *Journal*
1936 *of Geophysical Research: Solid Earth*, 108, 2143.
- 1937 Teisberg, T. O. & D. M. Schroeder (2023) Digital tools for analog data: Reconstructing the first ice-penetrating
1938 radar surveys of Antarctica and Greenland. In *International Geoscience and Remote Sensing*
1939 *Symposium (IGARRS) Proceedings*, 44-47.
- 1940 Teisberg, T. O., D. M. Schroeder, A. L. Broome, F. Lurie & D. Woo (2022) Development of a UAV-borne pulsed
1941 ice-penetrating radar system. In *IGARSS 2022 - 2022 IEEE International Geoscience and Remote*
1942 *Sensing Symposium*, 7405-7408.
- 1943 Thyssen, F. & K. Grosfeld (1988) Ekström Ice Shelf, Antarctica. *Annals of Glaciology*, 11, 180-183.
- 1944 Tinto, K. J., L. Padman, C. S. Siddoway, S. R. Springer, H. A. Fricker, I. Das, et al. (2019) Ross Ice Shelf
1945 response to climate driven by the tectonic imprint on seafloor bathymetry. *Nature Geoscience*, 12,
1946 441-449.
- 1947 Traversa, G., D. Fugazza & M. Frezzotti (2023) Megadunes in Antarctica: Migration and characterization from
1948 remote and in situ observations. *The Cryosphere*, 17, 427-444.
- 1949 Tsutaki, S., S. Fujita, K. Kawamura, A. Abe-Ouchi, K. Fukui, H. Motoyama, et al. (2022) High-resolution
1950 subglacial topography around Dome Fuji, Antarctica, based on ground-based radar surveys over 30
1951 years. *The Cryosphere*, 16, 2967-2983.
- 1952 Ulaby, F. & D. B. Lang. 2015. A Strategy for Active Remote Sensing Amid Increased Demand for Radio
1953 Spectrum. Washington DC: The National Academies Press.
- 1954 Van Liefferinge, B. & F. Pattyn (2013) Using ice-flow models to evaluate potential sites of million year-old ice
1955 in Antarctica. *Climate of the Past*, 9, 2335-2345.
- 1956 Van Liefferinge, B., D. Taylor, S. Tsutaki, S. Fujita, P. Gogineni, K. Kawamura, et al. (2021) Surface mass
1957 balance controlled by local surface slope in inland Antarctica: Implications for ice-sheet mass balance
1958 and Oldest Ice delineation in Dome Fuji. *Geophysical Research Letters*, 48, e2021GL094966.
- 1959 Vaughan, D. G., H. F. J. Corr, C. S. M. Doake & E. D. Waddington (1999) Distortion of isochronous layers in
1960 ice revealed by ground-penetrating radar. *Nature*, 398, 323-326.
- 1961 Višnjević, V., R. Drews, C. Schannwell, I. Koch, S. Franke, D. Jansen, et al. (2022) Predicting the steady-state
1962 isochronal stratigraphy of ice shelves using observations and modeling. *The Cryosphere*, 16, 4763-
1963 4777.
- 1964 Waddington, E. D., T. A. Neumann, M. R. Koutnik, H.-P. Marshall & D. L. Morse (2007) Inference of
1965 accumulation-rate patterns from deep layers in glaciers and ice sheets. *Journal of Glaciology*, 53, 694-
1966 712.
- 1967 WAIS Divide Project Members (2015) Precise interpolar phasing of abrupt climate change during the last ice
1968 age. *Nature*, 520, 661-665.
- 1969 Wang, T. T., B. Sun, X. Y. Tang, X. P. Pang, X. B. Cui, J. X. Guo, et al. (2016) Spatio-temporal variability of
1970 past accumulation rates inferred from isochronous layers at Dome A, East Antarctica. *Annals of*
1971 *Glaciology*, 57, 87-93.
- 1972 Wang, Z., A. Chung, D. Steinhage, F. Parrenin, J. Freitag & O. Eisen (2023) Mapping age and basal conditions
1973 of ice in the Dome Fuji region, Antarctica, by combining radar internal-layer stratigraphy and flow
1974 modeling. *The Cryosphere*, 17, 4297-4314.
- 1975 Wearing, M. G. & J. Kingslake (2019) Holocene formation of Henry Ice Rise, West Antarctica, inferred from
1976 ice-penetrating radar. *Journal of Geophysical Research: Earth Surface*, 124, 2224-2240.
- 1977 Weertman, J. (1976) Sliding-no sliding zone effect and age determination of ice cores. *Quaternary Research*,
1978 6, 203-207.
- 1979 Welch, B. C. & R. W. Jacobel (2003) Analysis of deep-penetrating radar surveys of West Antarctica, US-ITASE
1980 2001. *Geophysical Research Letters*, 30, 1444.
- 1981 Welch, B. C. & R. W. Jacobel (2005) Bedrock topography and wind erosion sites in East Antarctica:
1982 observations from the 2002 US-ITASE traverse. *Annals of Glaciology*, 41, 92-96.
- 1983 Welch, B. C., R. W. Jacobel & S. A. Arcone (2009) First results from radar profiles collected along the US-
1984 ITASE traverse from Taylor Dome to South Pole (2006–2008). *Annals of Glaciology*, 50, 35-41.
- 1985 Wilkinson, M. D., M. Dumontier, I. J. Aalbersberg, G. Appleton, M. Axton, A. Baak, et al. (2016) The FAIR
1986 guiding principles for scientific data management and stewardship. *Scientific Data*, 3, 160018.
- 1987 Winski, D. A., T. J. Fudge, D. G. Ferris, E. C. Osterberg, J. M. Fegyveresi, J. Cole-Dai, et al. (2019) The SP19
1988 chronology for the South Pole Ice Core – Part 1: volcanic matching and annual layer counting. *Climate*
1989 *of the Past*, 15, 1793-1808.
- 1990 Winter, A., D. Steinhage, E. J. Arnold, D. D. Blankenship, M. G. P. Cavitte, H. F. J. Corr, et al. (2017)
1991 Comparison of measurements from different radio-echo sounding systems and synchronization with
1992 the ice core at Dome C, Antarctica. *The Cryosphere*, 11, 653-668.
- 1993 Winter, A., D. Steinhage, T. T. Creyts, T. Kleiner & O. Eisen (2019a) Age stratigraphy in the East Antarctic Ice
1994 Sheet inferred from radio-echo sounding horizons. *Earth System Science Data*, 11, 1069-1081.

1995 Winter, K., J. Woodward, S. A. Dunning, C. S. M. Turney, C. J. Fogwill, A. S. Hein, et al. (2016) Assessing the
1996 continuity of the blue-ice climate record at Patriot Hills, Horseshoe Valley, West Antarctica.
1997 *Geophysical Research Letters*, 43, 2019-2026.

1998 Winter, K., J. Woodward, N. Ross, S. A. Dunning, R. G. Bingham, H. F. J. Corr, et al. (2015) Airborne-radar
1999 evidence for tributary flow switching in Institute Ice Stream, West Antarctica: Implications for ice-sheet
2000 configuration and dynamics. *Journal of Geophysical Research: Earth Surface*, 120, 1611-1625.

2001 Winter, K., J. Woodward, N. Ross, S. A. Dunning, A. S. Hein, M. J. Westoby, et al. (2019b) Radar-detected
2002 englacial debris in the West Antarctic Ice Sheet. *Geophysical Research Letters*, 46, 10454-10462.

2003 Wolovick, M. J. & T. T. Creyts (2016) Overturned folds in ice sheets: Insights from a kinematic model of
2004 traveling sticky patches and comparisons with observations. *Journal of Geophysical Research: Earth
2005 Surface*, 121, 1065-1083.

2006 Wolovick, M. J., T. T. Creyts, W. R. Buck & R. E. Bell (2014) Traveling slippery patches produce thickness-
2007 scale folds in ice sheets. *Geophysical Research Letters*, 41, 8895-8901.

2008 Wolovick, M. J., J. C. Moore & L. Zhao (2021a) Joint inversion for surface accumulation rate and geothermal
2009 heat flow from ice-penetrating radar observations at Dome A, East Antarctica. Part I: Model description,
2010 data constraints, and inversion results. *Journal of Geophysical Research: Earth Surface*, 126,
2011 e2020JF005937.

2012 Wolovick, M. J., J. C. Moore & L. Zhao (2021b) Joint inversion for surface accumulation rate and geothermal
2013 heat flow from ice-penetrating radar observations at Dome A, East Antarctica. Part II: Ice-sheet state
2014 and geophysical analysis. *Journal of Geophysical Research: Earth Surface*, 126, e2020JF005936.

2015 Woodward, J. & E. C. King (2009) Radar surveys of the Rutford Ice Stream onset zone, West Antarctica:
2016 Indications of flow (in)stability? *Annals of Glaciology*, 50, 57-62.

2017 Wrona, T., M. J. Wolovick, F. Ferraccioli, H. Corr, T. Jordan & M. J. Siegert (2018) Position and variability of
2018 complex structures in the central East Antarctic Ice Sheet. *Geological Society Special Publications*,
2019 461, 113-129.

2020 Xiong, S., J.-P. Muller & R. C. Carretero (2018) A new method for automatically tracing englacial layers from
2021 MCoRDS data in NW Greenland. *Remote Sensing*, 10, 43.

2022 Xu, B., S. Lang, X. Cui, L. Li, X. Liu, J. Guo, et al. (2022) Focused synthetic-aperture-radar processing of ice-
2023 sounding data collected over East Antarctic Ice Sheet via spatial-correlation-based algorithm using
2024 fast back projection. *IEEE Transactions on Geoscience and Remote Sensing*, 60, 5233009.

2025 Yan, Y., E. J. Brook, A. V. Kurbatov, J. P. Severinghaus & J. A. Higgins (2021) Ice-core evidence for
2026 atmospheric oxygen decline since the Mid-Pleistocene Transition. *Science Advances*, 7, eabj9341.

2027 Yari, M., O. Ibikunle, D. Varshney, T. Chowdhury, A. Sarkar, J. Paden, et al. (2021) Airborne snow-radar data
2028 simulation with deep learning and physics-driven methods. *IEEE Journal of Selected Topics in Applied
2029 Earth Observations and Remote Sensing*, 14, 12035-12047.

2030 Yilmaz, Ö. 2001. *Seismic Data Analysis: Processing, Inversion, and Interpretation of Seismic Data*. Tulsa, OK:
2031 Society of Exploration Geophysicists.

2032 Young, D. A., D. M. Schroeder, D. D. Blankenship, S. D. Kempf & E. Quartini (2016) The distribution of basal
2033 water between Antarctic subglacial lakes from radar sounding. *Philosophical Transactions of the Royal
2034 Society A: Mathematical, Physical and Engineering Sciences*, 374, 20140297.

2035 Zeising, O., T. A. Gerber, O. Eisen, M. R. Ershadi, N. Stoll, I. Weikusat, et al. (2023) Improved estimation of
2036 the bulk ice-crystal-fabric asymmetry from polarimetric phase co-registration. *The Cryosphere*, 17,
2037 1097-1105.

2038 Zhao, L. Y., J. C. Moore, B. Sun, X. Y. Tang & X. R. Guo (2018) Where is the 1-million-year-old ice at Dome
2039 A? *The Cryosphere*, 12, 1651-1663.

2040 Zirizzotti, A., J. A. Baskaradas, C. Bianchi, U. Sciacca, I. E. Tabacco & E. Zuccheretti (2008) Glacio RADAR
2041 systems and results. Proceedings of 2008 IEEE National Radar Conference, doi:
2042 10.1109/RADAR.2008.4720993.

2043

2044 **Appendix: Suggested standardised structure for the publication of traced IRHs across Antarctica**

2045 For publishing future radiostratigraphy datasets, we recommend scientists to follow the structure and
2046 naming convention specified in Table A1 for the first ten columns, after which additional columns may be
2047 added at the discretion of the scientists.

2048 In the metadata, we recommend that authors also provide at least the following information:

2049 (a) Name(s), version(s) and frequency of RES system(s) used.

2050 (b) Value for speed of radar wave in ice used to convert IRH depths to metres below the ice surface.

2051 (c) Value for any firn correction applied.

2052 (d) The coordinate system(s) used following the World Geodetic System 1984 datum and appropriate
2053 projection (i.e., EPSG:3031 for Antarctica).

2054 (e) If applicable, the type of radar product (e.g. waveform) on which the IRHs were traced.

2055 (f) The uncertainties associated with either the IRH age or depth based on RES system resolution and IRH
2056 picking, amongst others. Ideally, if the metadata vary throughout the dataset, then such information should
2057 be attached to each data point as additional columns to those shown in Table A1.

2058 (g) The source of age control (i.e., ice-core age scale, model).

2059 Additional information may also be added to the metadata, such as the type of processing used to extract
2060 the IRHs (if different from the processing used to trace the bed); the distance in the along-track direction
2061 along the RES transect for each data point; a flag number indicating whether the ice thickness, surface and
2062 bed elevations come directly from the along-track radar or from an interpolated gridded product, if
2063 applicable; the spatial resolution (or spacing distance between each data point); the dating method (s) used
2064 to provide an age for each IRH; and the type of software and tools used to pick the IRHs. Missing values in
2065 the float data should be set to NaN and specified in the metadata. We also recommend the use of open-
2066 access and FAIR data formats for storing the data, such as CSV or tabular data file (or netcdf if CSV or tabular
2067 data file is not suitable) where metadata can be easily embedded together with the data. Finally, we
2068 recommend scientists to publish their data in open-access repositories alongside the paper publication, with
2069 a DOI that can be linked back to the original paper. Together, these suggested protocols will ensure the
2070 longevity of the data products for future applications and enable faster retrieval thereof, particularly with
2071 regards to the large data volumes expected from automatic IRH tracking algorithms in the future.

2072 **Table A1.** Suggested standardised structure for the publication of IRH datasets associated with the
2073 AntArchitecture community effort following FAIR data standards.

2074 (For EGUsphere formatting, this 12-column table is presented across two rows.)

2075 Table rows 1-6:

Line ID or transect name	Trace timestamp (GPS time)	Longitude (decimal degrees)	Latitude (decimal degrees)	X coordinate (EPSG:3031; metres)	Y coordinate (EPSG:3031; metres)
--------------------------------	----------------------------------	-----------------------------------	----------------------------------	--	--

2076

2077 Table rows 7-12

IRH name	IRH (two- way travel- time through ice only)	IRH depth below ice surface (metres)	Ice thickness (metres)	Surface elevation (WGS84 ellipsoid; metres)	Bed elevation (WGS84 ellipsoid; metres)
-------------	--	---	------------------------------	---	---

2078