



Source contribution to ozone pollution during June 2021 in Arizona: Insights from WRF-Chem tagged O₃ and CO

Yafang Guo^{1,2}, Mohammad Amin Mirrezaei¹, Armin Sorooshian^{1,3}, Avelino F. Arellano^{1,3}

¹ *Department of Hydrology and Atmospheric Sciences, The University of Arizona, Tucson, AZ, USA*

² *Sonoma Technology, 1450 N. McDowell Blvd., Suite 200, Petaluma, CA, USA*

³ *Department of Chemical and Environmental Engineering, The University of Arizona, Tucson, AZ, USA*

Corresponding to: Avelino F. Arellano (afarellano@arizona.edu)

Abstract. This study reports the contribution of fire emissions on ozone (O₃) pollution in Arizona compared to local and regional anthropogenic emissions. Using the WRF-Chem modeling system with different O₃ and CO tags, we quantified the contributions of these emissions to O₃ levels during June 2021, a period when the region was experiencing both drought conditions and extreme heat. Our findings indicate that background O₃ levels accounted for about 50% of the total O₃, with local anthropogenic emissions contributing between 24% and 40%. During the peak smoky time period, fire-contributed O₃ was significant across the Phoenix metropolitan area, ranging from 5 to 23 ppb or 5% to 21 % of total O₃ levels, with an average of 15 ppb or 15%. We verify these O₃ fire tags by conducting a model sensitivity test that excluded fire emissions, which showed strong agreement on the spatiotemporal pattern of O₃ due to fire emissions, although the magnitude of the contribution is underestimated by a factor of 1.4. This further demonstrates that wildfires exacerbate O₃ exceedances over urban areas. Our analysis also showed that the O₃ levels in Yuma are significantly influenced by transboundary pollution from California and Mexico, whereas Phoenix's O₃ levels are mainly driven by local anthropogenic emissions, with much smaller contributions from external sources during the study period. Consistent with previous reports, our findings highlight the role of wildfires and regional emissions in confounding the assessment of local O₃ pollution in urban environments, especially during dry and extremely hot summer in semi-arid/arid regions.



1. Introduction

30 Ozone (O_3) pollution remains a pressing environmental and public health concern, especially in regions prone to wildfire activity (Jaffe et al., 2020; Jaffe and Wigder, 2012; Abatzoglou and Williams, 2016). Elevated O_3 levels can lead to a range of respiratory issues, cardiovascular problems, and other health complications, underscoring the importance of identifying and mitigating the sources of O_3 exceedances (Turner et al., 2016; Adhikari and Yin, 2020; Huangfu
35 and Atkinson, 2020). Arizona, particularly the south/southeast region, which is part of the Sonoran Desert and characterized by unique meteorological conditions and susceptibility to wildfires, presents a particularly challenging context for O_3 management (Betito et al., 2024; Guo et al., 2024; Miech et al., 2024; Sorooshian et al., 2024). During wildfire seasons, the complex interplay between local emissions, wildfire smoke, and meteorological factors contributes to significant O_3
40 exceedances, posing risks to both human health and ecological systems (Jaffe and Wigder, 2012; Jaffe et al., 2013; Selimovic et al., 2020).

Wildfires can significantly impact urban O_3 levels by releasing large quantities of pollutants that act as precursors to O_3 formation, including volatile organic compounds (VOCs) and nitrogen oxides (NO_x) (Andreae, 2019; Akagi et al., 2012). The pollutants can be transported over long
45 distances by wind, affecting urban air quality far from the fire source and combining with local emissions to exacerbate O_3 pollution (Xu et al., 2021; Jin et al., 2023; Jaffe and Wigder, 2012; Ninneman and Jaffe, 2021). In addition to O_3 , wildfire smoke leads to an increase in other atmospheric oxidants, such as hydroxyl radicals (OH) and hydroperoxyl radicals (HO_2), while reducing NO_2 photolysis rates due to the shading effect of smoke plumes (Buyssse et al., 2019).
50 This shading effect reduces the amount of sunlight available for the photolysis of NO_2 , which is a crucial step in O_3 formation. During wildfire events, stagnant air conditions often prevail in urban regions, preventing the dispersion of pollutants and allowing them to accumulate. For example, temperature inversions, which are more common during these events, trap pollutants near the ground, leading to higher concentrations of O_3 and other harmful substances (Alonso-Blanco et al., 2018; Burke et al., 2023; Jaffe et al., 2020; Xu et al., 2021).
55

Case studies, such as during various episodes of California wildfires, have demonstrated significant increases in urban O_3 levels, affecting cities far from the fire areas (Xu et al., 2021; Jin



et al., 2023; McClure and Jaffe, 2018). Understanding the specific contributions of different emission sources to O₃ pollution during wildfire events is crucial for developing effective air quality management strategies. Regional and local O₃ levels are influenced by local production, in addition to regional and long-range transport of O₃ and its precursors. Anthropogenic activities, such as vehicle combustion, industrial combustion, and power plants, as well as the biogenic emissions, are the most significant sources of NO_x and VOCs. When wildfires contribute additional NO_x and VOCs to the atmosphere, the overall levels of ground-level O₃ can rise dramatically as these plumes penetrate into urban areas, further degrading air quality (Pfister et al., 2006).

Source attribution techniques offer an alternative perspective to quantify the primary contributors to enhanced O₃ levels during smoky periods by identifying the contributions of specific sources and regions, such as anthropogenic, fire, and biogenic emissions, regional and international transport and stratospheric transport. In general, there are two main modeling approaches for O₃ source attribution or source apportionment: 1) model sensitivity experiments; 2) species tagging methods (Thunis et al., 2019; Kwok et al. 2015). The latter modeling approach tracks O₃ formation by tagging precursors from particular source types and areas throughout the model simulation, providing a direct attribution of modeled O₃ levels to these sources. The tagging technique entails modifying the model's source code to incorporate tracers into the chemistry mechanism. Models of atmospheric chemistry and transport that have implemented a tagging technique to perform O₃ source attribution include among others the Community Multiscale Air Quality (CMAQ) model with a new version of Integrated Source Apportionment Method (ISAM) (De La Paz et al., 2024; Shu et al., 2023), a submodel called TAGGING in the EMAC (European Centre for Medium-Range Weather Forecasts – Hamburg (ECHAM)/Modular Earth Submodel System (MESSy) (Grewe et al., 2017), the global Model for Ozone and Related chemical Tracers (MOZART- 4) (Emmons et al., 2012), the Nested Air Quality Prediction Model System (NAQPMS) (Zhang et al., 2020), CAM4-chem (Community Atmosphere Model version 4 with chemistry) within the Community Earth System Model (CESM) (Butler et al., 2018; Butler et al., 2020; Li et al., 2023; Nalam et al., 2024), the University of California Davis/Caltech air quality model (Zhao et al. 2022) and the Weather Research and Forecasting model coupled with Chemistry (WRF-Chem) (Pfister et al., 2013; Gao et al. 2016; Lupaşcu and Butler, 2019; Lupaşcu et al., 2022).



Typical model sensitivity analysis to determine the impact of a particular process (like fire) on target variables (like O₃) are usually conducted in practice as a suite of process-denial experiments and/or a series of model simulations with brute-force incremental changes on particular parameters or input datasets, along with developing model forward (decoupled direct method) and backward (adjoint) algorithms for sensitivity to emission calculations. Here, differences in simulated O₃ with and without fire emissions is interpreted to be the contribution of fire to modeled O₃ abundance. Unlike the tagging method, it utilizes the current model as is, without needing modifications. Models (and algorithms) that are used to predict how O₃ responds to changes in specific sources of emissions include among others those using WRF-SMOKE-CAMx (SMOKE: Sparse Matrix Operator Kernel Emissions model; CAMx: Comprehensive Air quality Model with extensions) (Zhang et al., 2017, Goldberg et al. 2016), WRF-Chem (Li et al., 2015), High-Order Decoupled Direct Method in Three Dimensions (HDDM-3D) (e.g., Cohan et al., 2005), CMAQ (Yeganeh et al., 2024; Collet et al., 2014, Hakami et al., 2007), STEM (Hakami et al., 2006), and climate chemistry model (E39C) (Grewe et al., 2012; Grewe et al., 2010).

These recent model and algorithm developments have shown the importance of integrating sophisticated modeling approaches and comprehensive data analysis to help better inform policies aimed at reducing O₃ pollution and its associated health impacts. Building on the work of Lupaşcu and Butler (2019) and Emmons et al. (2012), this study employs the tagging technique within the WRF-Chem modeling system to investigate the sources contributing to O₃ exceedances during a recent Arizona wildfire season, particularly examining the impacts of fires on O₃ levels. By utilizing this advanced modeling framework, our objective is to untangle the contributions of wildfire emissions, local anthropogenic activities, and regional transport to urban O₃ pollution. We verify this tagging approach by also conducting sensitivity experiments using the WRF-Chem model configuration by excluding fire emissions (i.e., zero-out emission scenario). Our focus is on June 2021 in Arizona, a period marked by multiple active wildfires and high O₃ concentrations. Specifically, we analyze two selected cases when Phoenix, the largest city in Arizona, was significantly impacted by wildfire smoke.

Section 2 presents the observational datasets from ground-based Environmental Protection Agency's Air Quality System (EPA AQS) sites and satellites, alongside the model setup. Section 3 begins with a detailed introduction of the selected cases, followed by an analysis of



comprehensive O₃ source apportionment. The discussion and conclusions are presented in Section 4.

2. Data and Methods

120 2. Study region and period

As mentioned, this is a case study focusing on 2021 dry summer (June) in Arizona. This period was notably severe, exacerbated by a combination of prolonged drought and an intense heatwave. The state experienced one of its hottest Junes on record, with temperatures frequently exceeding 115°F (46°C), with no significant precipitation recorded. This extreme heat, combined with
125 exceptionally dry conditions, were pivotal in the ignition and spread of multiple wildfires, leading to numerous large wildfires. In total, dozens of fires were reported across Arizona and New Mexico during this period, many sparked by lightning strikes on desert landscapes.

The Telegraph Fire, one of the largest wildfires in Arizona's history, began on 4 June 2021, near Superior, Arizona. By the time it was fully contained on 3 July 2021, the fire had burned over
130 180,000 acres. The Rafael Fire started on 18 June 2021 to the southwest of Flagstaff, which prompted widespread evacuations and road closures. The Rafael Fire had burned over 38 square miles by late June.

Figure 1 provides a detailed overview of the study region, encompassing the Phoenix metropolitan area in Arizona. Panel (a) displays a topographic map highlighting the Phoenix metropolitan area, with the red circle marking the AQS JLG Supersite, which serves as a central monitoring location.
135 The O₃ nonattainment area is delineated by red curves. This area is of particular interest due to its regulatory significance and the need for targeted air quality management strategies.

Additionally, the map indicates the locations of the Rafael and Telegraph fires, marked with stars. The Telegraph Fire is located to the southeast of Phoenix and Rafael Fire is located to the north of
140 Phoenix. Panel (b) of the figure illustrates the distribution of AQS monitoring sites within the nonattainment area. Each site is numbered and geolocated, offering an observational network for tracking O₃ and other air pollutants. Lists of the site locations and associated names are provided in Supplement Table S1. The spatial arrangement of monitoring sites facilitates a spatiotemporal analysis and assessment of pollutant level enhancements from wildfire smoke and other sources
145 of emissions.




Figure 1. (a) Topographic map of the Phoenix metropolitan area in Arizona, with the red circle representing USEPA/AQS JLG Supersite. The O₃ nonattainment area is outlined by a red border. The locations of the two largest wildfires during June 2021, Rafael and Telegraph, are marked with stars. (b) EPA AQS sites within the nonattainment area are shown. The sites are numbered and positioned according to their geographical locations with JLG noted as AQS site 23.

2.2 WRF-Chem setup

The Weather Research Forecasting with Chemistry (WRF-Chem) (Grell et al., 2005) model v4.4 is utilized here to simulate wildfire activities and study tropospheric O₃ pollution. Meteorological
150 initial and lateral boundary conditions are supplied every six hours by the Global Forecast System (GFS) with a horizontal grid spacing of 1° and 12-km NAM (North American Mesoscale Forecast System), while chemical initial and boundary conditions are provided by the Whole Atmosphere Community Climate Model (WACCM) for chemistry (Marsh et al., 2013; Tilmes et al., 2015). Biogenic emissions are calculated online using the Model of Emissions of Gases and Aerosols
155 from Nature (MEGAN, version 2.1) (Guenther, 2007; Guenther et al., 2006), based on the simulated meteorological conditions during the WRF-Chem runs. The anthropogenic emissions used in this study are obtained from 2017 National Emissions Inventories (NEI2017) data provided by the US EPA (<https://www.epa.gov/air-emissions-inventories/2017-national-emissions-inventory-nei-data>) with a 4 km grid spacing covering the U.S. and surrounding land areas.
160 Biomass burning emissions are calculated using the Fire Inventory from NCAR (FINNv2.5) (Wiedinmyer et al., 2023) and the online plume-rise model (Freitas et al., 2007). FINNv2.5 is



based on fire counts derived from Moderate Resolution Imaging Spectroradiometer (MODIS) and Visible Infrared Imaging Radiometer Suite (VIIRS) active fire detection (Wiedinmyer et al., 2023). A summary of the model configuration, parameterization, and a comprehensive model evaluation
165 against multiple observational and reanalysis datasets are provided in Guo et al. (2024). The WRF-Chem simulation period focuses on June 2021, targeting multiple wildfire activities near Phoenix.

2.3 O₃ tags and experiment design

To better understand the impacts of wildfire emissions on urban environmental settings, a species tagging technique was employed within the WRF-Chem model following recent demonstrations
170 (Emmons et al., 2012; Gao et al. 2016; Lupaşcu and Butler, 2019; Butler et al., 2018; Butler et al., 2020). Emmons et al. (2012) first introduced a method for tracking the sources of O₃ in the troposphere using a tagging approach within various chemical transport models, specifically MOZART-4. This tagging mechanism allows for a detailed attribution of O₃ to its precursor emissions sources, providing insights into how different sources contribute to overall O₃ levels.
175 Later on, Butler et al. (2018) applied the tagging mechanism for tracking the sources of tropospheric O₃ within the Community Earth System Model (CESM) version 1.2.2 and presented an updated version with comparison to Emmons et al. (2012). Lupaşcu and Butler (2019) then implemented the tagging mechanism within the WRF-Chem model to explore the origins of surface O₃ across Europe by distinguishing the contributions of different NO_x emission sources to
180 O₃ concentrations in various European regions.

Following Lupaşcu and Butler (2019), here we apply the tagging technique in the WRF-Chem model to quantify the contributions of different NO_x sources by not just tagging different regions but also different types of emissions. Our tags include four main categories: 1) regions that are local and adjacent, such as Arizona, California, and Mexico; 2) emission types, including
185 anthropogenic sources and fires; 3) tracers, including NO, NO₂, CO, and reaction products like O₃, O, and the corresponding NO_y reservoir species; and 4) background O₃ from initial and boundary O₃ levels. Note that these tracers undergo the same processes (advection, mixing, convection, chemical loss, deposition) within the continuity equation associated for each species in the model but they do not interact and affect changes in the modeled chemical system.

190 To implement the tagging technique in WRF-Chem, several steps must be completed before running the model. First, a tagged gas-phase chemical mechanism is created to incorporate tagged



tracers and reactions representing these tagged species, as well as the production and loss of O₃ to account for the tagged NO_x emissions. Here, we assume that O₃ peaks in urban areas of Arizona during this study period (June) is under a NO_x-limited chemical regime based on our previous studies (e.g., Guo et al., 2024; Greenslade et al., 2024). This new tagged mechanism is modified from the source code of the original MOZART mechanism within the WRF-Chem model. Next, both the anthropogenic and fire emission input files are modified to include tags related to different regions. For each regional tag, such as Arizona, NO_x and CO concentrations from outside Arizona are set to zero. Finally, the tags are initialized, and their boundary conditions are determined by the WACCM model output. The advantage of using WACCM is that it provides tagged CO tracers, including global biomass burning, North American anthropogenic emissions, and continental transport from regions such as East Asia, Europe, and Africa.

Since meteorological conditions, particularly wind speed and direction, have a significant impact on wildfire activities and plume coverage, we also apply the higher resolution 12-km NAM (North American Mesoscale Forecast System) dataset as the initial and boundary conditions. Evaluations are conducted for each selected case, comparing them against two boundary conditions. The simulations featuring winds and smoke plumes that best match satellite observations are selected. To help verify the contribution of wildfire emissions to O₃ levels, another set of simulations is performed by removing fire emissions. This serves as a sensitivity test for evaluating the model results with tags.

2.4 EPA AQS surface observations

To evaluate the accuracy of our model simulations, we use the surface observations from the Environmental Protection Agency's Air Quality System (EPA AQS). The AQS provides comprehensive air quality data from monitoring stations across the U.S., offering measurements of various pollutants, including O₃, particulate matter (PM_{2.5} and PM₁₀), and other criteria pollutants. The hourly and daily surface in situ observations of O₃ (including MDA8), CO, NO₂, and meteorological fields such as temperature, relative humidity, and winds from the EPA AQS monitoring network are used in this study (Demerjian, 2000). A total of 23 sites within the nonattainment area were selected based on their availability of O₃ measurements during the study periods, as shown in Figure 1b. The dataset undergoes quality control procedures to filter out any



erroneous or incomplete records, ensuring that only high-quality observations are used in our evaluation.

2.5 HMS smoke products

The Hazard Mapping System (HMS) smoke products provide detailed daily maps showing the geographic extent and concentration of smoke plumes across the U.S. and surrounding regions. The system integrates various satellite data sources, including the MODIS and VIIRS sensors, GOES (Geostationary Operational Environmental Satellite) imagery, to detect fire locations and estimate smoke coverage. The HMS smoke analysis has been a useful tool in monitoring wildfire impacts, supporting meteorological forecasting, and informing public safety measures related to air quality (e.g., Brey et al., 2018; Rolph et al., 2009).

The smoke products typically include three types of shapefiles: light, medium, and heavy (NOAA, 2023). Each category includes one or more shapefiles representing the smoke coverage estimated from satellite observations or images. These smoke products are used in this study to identify and select cases when Phoenix is defined as experiencing heavy smoke days.

2.6 TROPOMI satellite retrievals

The TROPospheric Monitoring Instrument (TROPOMI) is a state-of-the-art satellite sensor onboard the European Space Agency's (ESA) Sentinel-5 Precursor (S5P) satellite, launched in October 2017. TROPOMI actively measures tropospheric columnar atmospheric constituents including O₃, CO, NO₂, formaldehyde (HCHO). The TROPOMI dataset over Arizona has a spatial resolution of approximately 5.5 × 3.5 km² at nadir and provides daily data with an early afternoon (~12-14 PM) overpass time (Ludewig et al., 2020; Van Geffen et al., 2020). The data utilized in this research underwent a quality control process, where a quality assurance value (qa_value) greater than 0.50 was applied for HCHO and CO and a qa_value greater than 0.75 was applied for NO₂. The quality-controlled datasets were then gridded to a resolution of 0.07° × 0.07° for spatial analysis. For days with a lack of good quality data over the study domain, the data were further gridded to a coarser spacing of 0.2° × 0.2° to better capture the general spatial pattern of NO₂ and HCHO tropospheric columns.



3. Results and discussions

250 3.1 Case selection

Throughout June 2021, Phoenix experienced an intensified heatwave and drought conditions conducive for wildfire activity. Figure 2 presents the observed daily variations of pollutant levels for June 2021 from the EPA AQS at the Phoenix JLG Supersite compared to results from the TROPOMI satellite. Data on temperature (T) shows the heatwave began on June 12, with the daily
255 maximum temperature reaching 43°C and remaining at least that high until June 20. The Maximum Daily 8-Hour Average Ozone (MDA8 O₃) levels were around 50-70 ppb until June 10, when O₃ began to increase, exceeding the NAAQS standard (70 ppb) along with elevated surface concentrations of CO and NO₂, and T. A notably high MDA8 O₃ level (100 ppb) was observed on June 15. MDA8 O₃ then decreased to below NAAQS levels on June 17 up until June 27. Surface
260 CO levels generally followed the O₃ variation, ranging between 400-700 ppb. However, a noticeable peak in CO, exceeding 1000 ppb, was observed on June 11. NO₂ levels were compared between the EPA AQS and TROPOMI satellite data. On June 11, both surface and tropospheric column NO₂ levels exhibited a significant peak, indicating that emissions were mostly within the planetary boundary layer. Conversely, the discrepancies between NO₂ surface and column levels
265 beginning June 13 suggest different sources for surface and tropospheric NO₂ emissions, particularly on June 15, when MDA8 O₃ exceeded 100 ppb; surface NO₂ was relatively low, while column NO₂ was high. The peak period beginning June 14 of HCHO concentration was also captured by both AQS and TROPOMI observations. In Guo et al. (2024), they showed that during the period with elevated temperature (June 12-20), relative humidity is low but normal for June in
270 Arizona. The high temperature resulted in an increase of isoprene and HCHO simultaneously.




Figure 2. Observational daily variations of surface (a) MDA8 O₃, (b) CO, (c) NO₂, (d) HCHO, and (e) temperature (T) from EPA AQS at Phoenix JLG supersite (blue), as well as column (c) NO₂ and (d) HCHO concentrations from TROPOMI satellite (red) in June 2021. The NAAQS 2015 standard is denoted as black dash line. Note that the daily AQS CO, NO₂, and T values shown represent the daily maximum while the HCHO value is the daily mean.

As mentioned earlier, the Telegraph Fire began on June 3 and lasted for one month, while the Rafael Fire started on June 18. Guo et al. (2024) showed that in the month of June, the prevailing wind over Phoenix was mostly southwesterly, limiting the impacts of these wildfires on Phoenix to certain days when winds shifted direction and brought smoke plumes to the city. After reviewing the HMS smoke data for June 2021, we identified two smoky periods that might have potentially influenced surface O₃ concentrations over Phoenix.

The first selected case is on 15 June 2021. On this day, multiple sites within the nonattainment area observed O₃ exceedances (>70 ppb). An excessive heat warning was issued and remained in effect through the end of the week, with temperatures 10 to 15 degrees above average (Gard and Garrett, 2021). The wind shifted from southwesterly to northeasterly, bringing the Telegraph Fire plumes to Phoenix. The second case is on 26 June 2021, when smoke from the Rafael Fire spread to the north of Phoenix with a change of wind direction from southwesterly to northerly.



Shown in Figure 3 is the heavy smoke coverage from the HMS smoke products over the Phoenix
285 area during two selected cases in June 2021, highlighting the impact of the Rafael and Telegraph
fires. In Case I, on 14 June 2021, at 16:00 LT, smoke from both the active Telegraph (southeast of
Phoenix) Fire spread primarily to the northeast. By 15 June 2021, at 16:20 LT, the smoke coverage
had expanded significantly, with a dense plume covering central Arizona, including Phoenix. On
16 June 2021, at 11:00 LT, the dense and widespread smoke continued to affect the periphery of
290 Phoenix. In Case II, on 25 June 2021, at 07:00 LT, smoke primarily from the Rafael Fire extends
to the east, far away from Phoenix. By 26 June 2021, at 17:00 LT, the smoke plume from Rafael
Fire changed direction to the south and covered the north of Phoenix. Active wildfires contributing
to the smoke over Phoenix are marked with yellow stars, indicating the origin and spread direction
of the smoke plumes. These two cases are selected for further modeling studies to help understand
295 how near-range wildfires affect the Phoenix metropolitan area.




Figure 3. (a-c) Heavy smoke coverage from Hazard Mapping System (HMS) smoke products for Case I, and (d-f) for Case II over Phoenix area. The active wildfire that was accountable for the smoke over Phoenix is marked as yellow star.

In Figure S1 we present a series of screenshots from the MODIS Terra Corrected Reflectance map, overlaid with MODIS fires and thermal anomaly products, to depict wildfire activities in Arizona for the above two cases. Similar to Figure 3, the top panel illustrates Case I, focusing on the



300 Telegraph Fire from June 14 to 16. The bottom panel captures Case II, highlighting the Rafael Fire from June 25 to 27. Both cases show visible smoke plumes and thermal anomalies (orange color) indicating active fire regions, with the fire spreading and producing significant amounts of smoke passing Phoenix.

In addition to HMS smoke products, we show in Figure 4 the daily TROPOMI tropospheric
305 columns of HCHO, NO₂ and CO during smoke periods for Case I over the Phoenix area. In Case I, the HCHO levels initially show low levels, with scattered low concentrations on June 13 except the east of Telegraph Fire. By June 14, there is a significant increase in HCHO, especially northeast of Phoenix, correlating with the smoke plume from the Telegraph Fire, as seen in Figure 3a. On June 15, the elevated HCHO levels were more dispersed, affecting mainly the south of Phoenix.
310 By June 16, HCHO tropospheric column decreased to the normal levels over Phoenix. For NO₂, June 13 shows low levels with a typical urban anthropogenic emissions spatial profile. June 14 exhibits a significant increase in NO₂, particularly northeast of Phoenix, similar to the HCHO distribution. On June 15, NO₂ levels are high over a wider area, including Phoenix and the path of plumes (Figure 3b). By June 16, NO₂ levels decrease but remain elevated. A similar pattern has
315 been observed in CO, where its high concentrations are closely correlated with HCHO, NO₂, and smoke coverage, as shown in Figure 3. The TROPOMI results for Case II are presented in Figure S2.




Figure 4. TROPOMI tropospheric columnar HCHO (top), NO₂ (middle), and CO (bottom) during the smoke periods for Case I over Phoenix area. The black polygon lines represent the EPA designated Phoenix-Mesa nonattainment area. The grid resolution is 0.2°. Grids without available data are marked as white space.

Similar to the TROPOMI satellite observations, we also examined the model simulation results. Figure 5 presents the WRF-Chem simulated tropospheric columnar values of HCHO, NO₂, and CO at 14:00 local time (same time as the TROPOMI observations) during the smoke periods for Case I. Comparing Figures 4 and 5, on June 13 as a pre-smoke day, the HCHO levels in the city region are comparable, with values primarily below 6×10^{15} molecules/cm², while the Telegraph Fire burning area reached over 10×10^{15} molecules/cm². This day represents a typical distribution of urban pollutants over Phoenix. By June 14, levels of HCHO, NO₂, and CO increased significantly, particularly in the southeastern part of Phoenix, although the magnitude and spatial patterns appear to differ from the satellite observations in Figure 4. On June 15, the tropospheric columnar values decreased but the wildfire signal remained significant until June 16, when the spatial pattern returned to typical conditions. Additional WRF-Chem simulated results for Case II are available in Figure S3-4.

In summary, observations from HMS, TROPOMI, and WRF-Chem models indicate that the Telegraph Fire had a significant impact on Phoenix air quality during June 14-15. While the rise of plumes during wildfires greatly influences the columnar concentrations of pollutants by transporting smoke and emissions higher into the atmosphere, the mixing levels within the surface



335 or planetary boundary layer are more important to the overall air quality and pollutant distribution
as the more immediate impact on public health is expected at ground-level.




Figure 5. WRF-Chem simulated HCHO, NO₂, and CO tropospheric columns at local time 14:00 during the smoke periods for Case I over Phoenix area.

We show in Figure 6 the WRF-Chem simulated surface concentrations of HCHO, NO₂, and CO. Since each case involves two sets of simulations using GFS and NAM meteorological boundary
340 conditions, the selection of the model results is based on an initial evaluation against AQS and satellite observations. For Case I, the results from the GFS simulations demonstrate better agreement, while for Case II, the NAM simulations show better alignment of smoke. Comparing these with the columnar levels of NO₂ and CO in Figure 5, it is evident that the extent of the smoky day on June 14, as observed from HMS and TROPOMI, is not reflected at the surface level,
345 whereas the smoky day on June 15 is apparent in both surface and columnar concentrations. Additionally, for HCHO, increases are also observed at the surface on June 14. This discrepancy indicates that on June 14, the wildfire smoke was primarily affecting atmospheric layers aloft without significantly impacting the ground level, while on June 15, the smoke was more distributed in the lowermost troposphere, increasing the surface pollution concentrations. Model results of tropospheric column and surface HCHO, NO₂, and CO for Case II are provided for reference
350 in the Supplement as Figures S1-S2, respectively.




Figure 6. Same as Figure 5, but for surface concentrations.

3.2 Source attribution with tags

An extensive evaluation of the same configuration of the WRF-Chem model using the MOZART
355 chemical mechanism, except for the tags, has been presented previously by Guo et al. (2024).
Briefly, our evaluation showed a Pearson correlation coefficient (R) of 0.81 for modeled and
observed O_3 over Phoenix with a mean bias (MB) of -2.9 ppb and 1.0 ppb for hourly and MDA8
 O_3 , respectively. For CO and NO_2 , the normalized bias is 7.1% and 5.3%, respectively. The model
simulations also show that surface formaldehyde-to-nitrogen dioxide ratio (FNR), which is an
360 indicator of chemical regime affecting O_3 production varies from a VOC-limited regime in the
most populated areas to a transition between VOC-limited and NO_x -limited regimes throughout
the metro area. Here in this study, our discussion of the model results is focused on the month of
June 2021, a period marked by active wildfires over Arizona against a backdrop of not only an O_3
chemical regime that is in transition to NO_x -limited but also of drought and heat wave conditions.
365 We first provide an analysis of the contribution of different source regions and emission types to
the monthly CO and MDA8 O_3 concentrations to understand the overall pollution sources in the
State of Arizona. Then, we focus on the analysis of O_3 during smoky days by examining the two
selected cases described in Section 3.1.



370 **3.2.1 Monthly CO and O₃ extremes for June 2021**




Figure 7. WRF-Chem simulated 90th percentile of surface CO concentrations during June 2021 for total CO (a), and the contributions from different CO sources (b-h). Each panel represents different aspects of CO: (a) total CO, (b) background CO, (c) anthropogenic CO sources, (d) CO from California anthropogenic sources, (e) CO from Arizona anthropogenic sources, (f) CO from Mexico anthropogenic sources, (g) CO from Arizona wildfires, and (h) CO from Mexico wildfires. Key locations such as Phoenix (PHX), Tucson (TUS), and Yuma are marked as stars on the maps. Telegraph and Rafael fires are denoted as unfilled circles.

Shown in Figure 7 is an overview of CO concentrations in Arizona during June 2021, highlighting the impact of various sources on CO distribution. Each panel represents the 90th percentile for the entire month of different CO and its sources: (a) total CO, (b) background CO levels, (c) anthropogenic CO sources, (d) CO from California anthropogenic sources, (e) CO from Arizona anthropogenic sources, (f) CO from Mexico anthropogenic sources, (g) CO from Arizona wildfires, and (h) CO from Mexico wildfires. Comparing the total CO concentrations (Figure 7a) with anthropogenic CO (Figure 7c), we can see a clear signature of anthropogenic activities in



380 cities such as Phoenix (PHX), Tucson (TUS), and Las Vegas, located in the upper left corner of the map.

The “background” CO levels (Figure 7b) are generally constant, ranging between 50 and 70 ppb across the region, which is closely related to international or long-range transport as well as global secondary CO formation. When examining anthropogenic sources, contributions are tagged separately for California (Figure 7d), Arizona (Figure 7e), and Mexico (Figure 7f). The dominant 385 contributions are seen around Arizona's urban areas, particularly Phoenix and Tucson, highlighting the impact of local urban emissions. CO from Mexico also influences southwestern boundaries with Arizona, particularly the city of Yuma, with an estimate of 30 ppb. Contributions from California (Figure 7d) are limited to the state boundaries, with only minor impacts to surface CO (~5 ppb) during this period.

390 Besides the anthropogenic sources, CO is emitted directly from wildfires, as a result of incomplete combustion of organic materials such as trees, plants, and other vegetation. During a wildfire, the high temperature during its flaming phase causes rapid oxidation of carbon-containing materials, but not all the carbon is fully oxidized to carbon dioxide (CO₂) during its (lower temperature) smoldering phase. Emissions from wildfires can have far-reaching impacts, as CO is a relatively 395 long-lived gas in the atmosphere, with a typical lifetime of several weeks to a few months. This is facilitated as well by associated plume rise especially during the fire’s flaming phase. As shown in Figure 7g, wildfires in Arizona notably elevate CO levels, especially in areas downwind of active fires with six major wildfire activities identified.




Figure 8. WRF-Chem simulated 90th percentile of O₃ concentrations during June 2021 for (a) MDA8 O₃, and contributions from different sources as (b) background O₃, (c) O₃ from anthropogenic sources, (d) O₃ from California anthropogenic sources, (e) O₃ from Arizona anthropogenic sources, (f) O₃ from Mexico anthropogenic sources, (g) O₃ from Arizona wildfires, and (h) O₃ from Mexico wildfires. Key locations such as Phoenix (PHX), Tucson (TUS), and Yuma are marked as stars on the maps. Telegraph and Rafael fires are denoted as unfilled circles.

400 Additionally, we examined MDA8 O₃ using the tags, as presented in Figure 8. We see that the MDA8 O₃ concentrations are predominantly high across the region, with the highest levels observed around Phoenix. Figure 8b indicates that the background O₃ levels are uniformly high, approximately 50 ppb, suggesting that even in the absence of local sources, O₃ concentrations remain elevated due to regional and global influences on a monthly basis.

405 We show in Figures 8c to 8f a regional decomposition of the anthropogenic contributions to O₃ levels. Figure 8c represents all anthropogenic sources, revealing significant contributions, especially around urban centers like Phoenix and Tucson. Figure 8d shows the small impact of California's anthropogenic emissions on Arizona's O₃ levels during this period only reaching ~3 ppb in Yuma. In contrast, Arizona's anthropogenic contributions to Arizona's O₃ levels (Figure 8e)



410 are substantial (as expected), ranging from 25 to 30 ppb within the nonattainment area. Mexico's anthropogenic contributions (Figure 8f) have a larger impact than CO (Figure 7f) in terms of spatial coverage, affecting most of the southern Arizona regions and even reaching Phoenix at 3 ppb. The magnitude is also higher, reaching 10 ppb for Yuma.

Similar to CO, Figures 8g and 8h focus on O₃ contributions from wildfires in Arizona and Mexico, 415 respectively. However, while CO is directly emitted from wildfires, O₃ is chemically formed from precursors such as VOCs and NO_x transported with the smoke. Consequently, the patterns of O₃ differ from those of CO. O₃ can have a larger impact due to the transport of these precursors, leading to significant O₃ formation even far from the wildfire sources. Figure 8g shows that wildfires in Arizona contribute notably to O₃ levels, particularly in areas close to and downwind 420 of the fires. O₃ concentrations range from 1 to 10 ppb, with the highest levels observed near the wildfire locations. The influence of these wildfires extends towards the east and southeast, consistent with the prevailing winds being eastward, and indicating the transport of O₃ precursors and subsequent formation of O₃ in these areas.

Figure 8h highlights the influence of wildfires in Mexico on O₃ levels in Arizona, particularly 425 affecting the southern and southwestern parts of the state. The contributions from Mexico wildfires are less than 3 ppb. The transport of smoke and O₃ precursors from Mexico affects a broader area than CO, reaching as far as Phoenix and diminishing farther north. This underscores the effect of cross-border wildfire emissions on O₃ levels and air quality in southern Arizona, particularly in border regions like Yuma.

430 We can see in Figures 7 and 8 that Yuma, which is located at the boundaries of Mexico and California, are influenced by local, regional, and transboundary CO and O₃. Here, we present the modeled and observed hourly O₃ concentrations at local time from Yuma monitoring site (AQS site number: 04-027-8011) for the period between June 14 and June 19, highlighting the contributions from various sources. Two episodes of hourly surface O₃ exceeding 70 ppb are 435 observed on June 15 and June 17, which the WRF-Chem model generally captures, although some discrepancies exist.

The shaded areas reveal the contributions from different sources: background O₃, local and regional anthropogenic emissions, and wildfire emissions from Arizona and Mexico. Figure 9 shows that O₃ levels in Yuma are clearly impacted by emissions from Mexico. For instance, on



440 June 17, the contributions from sources outside Arizona exceeded those from local emissions, suggesting the influence of transboundary pollution on Yuma's air quality.




Figure 9. Hourly O₃ concentrations (in ppb) at the Yuma monitoring site (site number: 04-027-8011) between 14-19 June 2021, at local time. The dashed black line represents observed O₃ levels from the AQS, while the solid blue line shows WRF-Chem simulated O₃ concentrations. Shaded areas indicate contributions from various sources: background O₃ (gray), Arizona wildfires (orange), anthropogenic emissions from Mexico (green), anthropogenic emissions from Arizona (red), anthropogenic emissions from California (purple), and Mexico wildfires (yellow).

Figures 7-9 demonstrate the complex interplay of local, regional, and transboundary sources in determining CO and O₃ levels. By examining the contributions of local anthropogenic emissions, wildfire emissions, and regional influences from neighboring states and countries, as well as background levels, these figures provide new perspectives of air quality in the region.

3.2.2 Smoky day O₃ analysis

Fire Contributions to Phoenix O₃. The detailed analysis presented in Section 3.2.1 provides an overview of the key sources of pollution during a fire season in June. In this section, we examine the impact of wildfire smoke plumes on urban areas by examining two specific smoky days (two cases) with a focus on the Phoenix metropolitan area, where the cases are described in Section 3.1. To assess the effects of fire emissions on O₃ concentrations, we conducted an additional set of



WRF-Chem simulations without fire emissions for the same period. The simulations without fire emissions serve as a model sensitivity test to evaluate the impact of wildfires.

455 Figure 10 illustrates the impact of fire emissions on the MDA8 O₃ concentrations for the two cases. The top panels represent Case I for June 15 (a) without fire emissions and (b) with fire emissions. Similarly, the bottom panels depict Case II for June 26 (c) without fire emissions and (d) with fire emissions. The comparison between the left and right panels highlights the significant contribution of wildfire emissions to O₃ levels in the Phoenix metropolitan area.

460 In Case I (June 15), the presence of fire emissions (Figure 10b) leads to a substantial increase in MDA8 O₃ concentrations, exceeding 110 ppb in areas directly affected by the wildfire plumes. This is in stark contrast to the scenario without fire emissions (panel a), where O₃ levels remain below 90 ppb. The path of the elevated MDA8 O₃ in Figure 10(b) aligns with the HMS smoke coverage depicted in Figure 3.

465 For Case II (June 26), a similar pattern is observed, albeit with a much weaker intensity. The inclusion of fire emissions (Figure 10d) also results in elevated MDA8 O₃, with peak values reaching around 90 ppb, while without fire emissions (Figure 10c), O₃ levels are significantly lower, generally below 70 ppb. The spatial distribution of MDA8 O₃ also aligns with the mean transport pathway of the wildfire plumes.

470 The AQS observations, indicated by the colored circles, are generally consistent with the model results when fire emissions are included, demonstrating the model's ability in capturing the impact of wildfire emissions on ground-level O₃ concentrations. The mean bias between model without fire emissions and observations is -7.9 ppb for case I and 9.7 ppb for case II. When fire emissions are included, the mean bias is reduced to -1.8 ppb and 2.9 ppb for the two cases, respectively.

475 Overall, the sensitivity simulation suggests that wildfires exacerbate O₃ pollution, especially when fire smoke passes through urban areas when photolysis is high. Additionally, it enables us to verify the O₃ fire tags. Ideally, the difference in O₃ concentrations when fire emissions are excluded should match the O₃ fire tags. However, studies have shown that this is not always the case mainly due to non-linearity of O₃ chemistry to precursor emissions as well as the spatiotemporal
480 heterogeneity of O₃ chemical regime. The differences between attributing source contributions through sensitivity or tagging approaches have been noted by several studies (e.g., Grewe et al., 2010; Kwok et al., 2015). These studies reported that the sensitivity method could potentially



induce large errors (factor of 2), which depend on the degree of linearity of the chemical system. To verify our tagging approach, we show in Figure 11 the WRF-Chem simulated daytime (7:00-
485 19:00 LT) average of O_3 concentrations for two different cases: Case I on 15 June 2021 (top panels) and Case II on 26 June 2021 (bottom panels). The left panels display the differences in O_3 levels between scenarios with and without fire emissions. The right panels show the daytime average O_3 concentrations attributed to fire emissions (fire tag). The spatial variations observed in the two methods are evidently similar across both cases. However, the values differ by a factor of 1.4, as
490 indicated by the color bar scales, which aligns with previous expectations.




Figure 10. WRF-Chem simulated MDA8 O_3 concentrations for Case I (15 June 2021, top panels) and Case II (26 June 2021, bottom panels) under two conditions: without fire emissions (left panels) and with fire emissions (right panels). AQS observations are represented by colored circles, excluding sites with missing or low-quality data. Stars indicate the locations of the wildfires (top: Telegraph; bottom: Rafael). The red outline represents the designated nonattainment area.




Figure 11. Daytime (7:00-19:00 LT) average O₃ concentrations simulated by WRF-Chem for Case I (15 June 2021, top panels) and Case II (26 June 2021, bottom panels). The left panels show the difference between scenarios with and without fire emissions, while the right panels depict the daytime average O₃ fire tag. Stars mark the wildfire locations (Telegraph Fire at the top and Rafael Fire at the bottom). The red outline denotes the designated nonattainment area. Note that the color bar scales for left and right panels are different. The black arrows indicate the path of smoke plumes.

In addition to examining the spatial variations of O₃ concentrations, we also present the temporal variations of surface hourly O₃ within the Phoenix area in Figure 12, which includes a detailed
495 look at each individual AQS site and the contribution of each O₃ tag to the overall O₃ levels. First, a site located under the plume path with significant O₃ elevation from smoke is selected for each case. Next, a timestamp is chosen when the O₃ fire tag is at its peak to review and compare observations from all AQS sites. The top panels of Figure 14 show the hourly O₃ concentrations



at AQS sites 7024 and 1010 for Case I and Case II, respectively. The locations and site numbers
500 are detailed in Figure 1 and Table S1. For Case I between June 15 and 17, the peak hourly O₃
concentration reached approximately 115 ppb on June 15 at 17:00 local time, aligning with AQS
measurements. The contribution from Arizona fire emissions is evident, as indicated by the orange
segments in the stacked area chart (Figure 14a). Background O₃ levels (gray shading) constitute
the largest portion of the total O₃, accounting for approximately 50%. Local anthropogenic
505 emissions are the next significant contributor, varying between 24% and 40%, depending on the
urban setting of the site. A closer examination of other sites during the O₃ peak hour on June 15
reveals that fire contributed O₃ is significant across the area, with values around 15 ppb or 15%
(Figure 14b). This indicates that the wildfire events during this period had a substantial impact on
elevating O₃ levels.

510 For Case II, O₃ levels are much lower, peaking at about 80 ppb on June 26 at 11:00. Compared to
Case I (Figure 14a), the impact of fires on O₃ levels is less pronounced. After June 26, O₃ levels
returned to non-smoky day patterns, with most contributions from local anthropogenic emissions.
Figure 14d further illustrates the distribution of O₃ sources across multiple sites at 14:00 on June
26, showing fire contributions of 5-10 ppb or approximately 10%. The background O₃ levels
515 remain consistent with Case I. The differences between these two cases may be attributed to
varying meteorological conditions, fire intensity, and/or the spatial distribution of emissions during
the two periods. During Case I, Arizona experienced excessive heat and record high temperatures
(Figure 2), and the Telegraph Fire had a larger and longer smoke impact than the Rafael Fire in
Case II. Unlike Yuma, as shown in Figure 9, O₃ levels in Phoenix are primarily influenced by local
520 emissions, with much smaller contributions from California or Mexico, even with significant
contributions from wildfire smoke.

An additional figure comparing the effects of anthropogenic and fire-related emissions on O₃ levels
for Case I is provided in Figure S5. This figure shows a pronounced diurnal cycle, with O₃ levels
increasing from early morning, peaking around noon to early afternoon (12 to 1pm), and then
525 declining towards the evening. Our results show significant differences between these two
emission sources across three urban settings: suburban, urban, and rural. In the early morning and
early afternoon, O₃ levels are predominantly influenced by anthropogenic emissions at most AQS
sites. However, in the late afternoon, when a fire smoke plume passed through the Phoenix urban



530 area, the contribution of fire related O₃ increases significantly and, in some rural sites, even surpasses local anthropogenic production.




Figure 12. Contribution of each tagged O₃ source to the hourly O₃ concentrations (ppb) for Case I (a-b) and Case II (c-d). The top panels (a, c) show the hourly variations in O₃ concentrations at a single AQS site (#7024 for Case I and #1010 for Case II) from 15-18 June 2021 and 26-28 June 2021, respectively. The bottom panels (b, d) display the contributions of different O₃ sources at multiple sites at the time stamps indicated by the blue vertical lines in (a) and (c). O₃ sources include background O₃ (BG O₃), Arizona anthropogenic (AZ anthro), California anthropogenic (CA anthro), rest of the anthropogenic (Rest anthro), Arizona fire (AZ fire), and rest of the fire (Rest fire).

We also present in Figure 13 the WRF-Chem simulated surface CO concentrations on 15 June 2021 (Case I). By comparing the difference in CO concentrations with fire emissions (Figure 13b) and without fire emissions (Figure 13a) to the CO fire tag (Figure 13d), we observe a similar spatial pattern to that of O₃ in Figure 12. However, the CO fire tag indicates a more extensive area of low CO concentration coverage compared to the sensitivity method.

535




Figure 13. Daytime (7:00-19:00 LT) average surface CO concentrations simulated by WRF-Chem during Case I (15 June 2021) for (a) without fire emissions, (b) with fire emissions, (c) difference between (b) and (a), and (d) CO fire tags. Stars mark the wildfire locations (Telegraph Fire at the top and Rafael Fire at the bottom). The red outline denotes the designated nonattainment area.

Impact of Fire on Chemistry and Meteorology. We show in Figure 14 the temporal variations in the photolysis rate of NO_2 (J_{NO_2}), NO_x ($\text{NO}+\text{NO}_2$), HO_x ($\text{OH}+\text{HO}_2$), and O_3 concentrations in metro Phoenix (Site 7024) at local time 16:00 over a seven-day period in June 2021, covering Case I under two conditions: with and without fire emissions. This site (see Figure 1) is situated along the plume coverage downwind of the fire. We also included key meteorological variables (net and outgoing longwave radiation, winds, surface temperature and PBL height) and concentration of black carbon aerosols (which is a light absorbing particle) to elucidate the direct radiative impact of the fires. In Figure 14(a), the photolysis rates of NO_2 (J_{NO_2}) are consistently only slightly higher without fire emissions while NO_x concentrations vary across the week (lower in June 14 but



slightly higher in June 15 with fire). HO_x levels vary similarly with NO_x during this fire event, possibly associated with VOCs from fires. This results to O₃ in June 15 at 4pm that is significantly higher in the simulation with fire compared to simulation without fire. The net and outgoing
550 longwave radiation, along with black carbon concentration are also higher with fire. Note that there is a significant wind shift from northward to southward (along with lower wind speed) in June 15 when fire is included (Figure 14d), along resulting in the displacement of O₃ and CO hotspot observed in Figure 10 and Figure 13, respectively. This is consistent with the observed exceedance of O₃ levels on the same day.




Figure 14. WRF-Chem simulated time series (11-17 June 2021) of daily photolysis rate of NO_2 (J_{NO_2}), concentrations of NO_x ($\text{NO}+\text{NO}_2$; ppb), HO_x ($\text{OH}+\text{HO}_2$; ppb), and O_3 (ppb), meteorological conditions such as net and outgoing longwave (OLR) radiation (watts/m^2), 10-meter zonal and meridional and zonal wind and wind speed (10m U and V, Wind Speed (m/s), 2-meter air temperature (K), concentration of black carbon (BC) aerosols ($\mu\text{g}/\text{m}^3$), and planetary boundary layer height (PBLH). All these are sampled at 16:00 local time in Phoenix (Site 19 - #7024). The black markers represent the values with fire emissions, while the red markers indicate values without fire emissions.



To further investigate the fire impact, we present in Figure 15 a cross-sectional view of the smoke plume as it travels towards Phoenix during Case I, highlighting the concentrations of multiple atmospheric pollutants, including O₃, CO, NO_x, HCHO (formaldehyde), PM_{2.5} (particulate matter), and PAN (peroxyacetyl nitrate). Near the fire location, concentrations of CO, NO_x, HCHO, and PM_{2.5}, which are primary pollutants directly emitted from the fire, are high, whereas O₃ concentrations are lower. As the smoke moves closer to the urban region, NO_x levels in the boundary layer increase significantly, along with O₃ levels reaching 100 ppb above the ground. Levels of NO_x from fires diminishes at a faster rate than HCHO and PM_{2.5} levels along the trajectory. It is clear from the figure that pollutants from the fires are transported towards the valley.

This is particularly true for PAN which shows an enhancement above the valley along with CO and PM_{2.5}. These enhancements aloft are not present in the cross-section of WRF-Chem simulation without fire emissions (see Supplemental Figure S7). Previous studies have indicated that the rapid conversion of NO_x to PAN can limit O₃ production near fires, especially at low temperatures, but the decomposition of PAN can lead to additional O₃ production further downwind of the fires especially in the presence of higher amounts of VOCs (Alvarado et al., 2010; Jaffe et al., 2013). The concentrations of O₃, CO, NO_x, and PAN from fire tags presented in Figure S6 (alongside Figure S7) further demonstrate that fire smoke exacerbates urban O₃ levels, while the exceedance is predominantly from local production.

In summary, this cross-sectional analysis illustrates the complex vertical and horizontal distribution of various pollutants and their transformations within a smoke plume traveling towards Phoenix. The interaction between primary emissions from fires, secondary pollutants formed during transport, and the presence of local anthropogenic emissions in the urban environment highlights the multifaceted nature of urban air quality impacts during wildfire events.




Figure 15. Cross-sectional analysis of a smoke plume traveling towards Phoenix during 15 June 2021, showing vertical and horizontal distribution of various pollutants. The subplots represent concentrations of (a) O₃, (b) CO, (c) NO_x, (d) HCHO, (e) PM_{2.5}, and (f) PAN (Peroxyacetyl Nitrate) across different altitudes and distances from the fire where TEL means Telegraph. The plume path is denoted in Figure 11. The gray shading represents the topography heights.

Fire-induced Changes in Chemical Regime. We show in Figure 16 the associated impact of fires on the chemical regimes of O₃ formation over Phoenix at local time 14:00. Here, two key indicators are chosen to illustrate this impact: the HCHO/NO₂ ratio, also known as the Formaldehyde to NO₂ Ratio (FNR), and the H₂O₂/HNO₃ ratio (Sillman, 1995; Tonnesen et al., 2000). The HCHO/NO₂ ratio (FNR) has been used in previous studies as an indicator for determining the sensitivity of O₃ formation to either VOCs or NO_x (Martin et al., 2004; Jin et al., 2020; Mirrezaei et al., 2024). A higher FNR indicates a NO_x-limited regime, where O₃ formation is more sensitive to changes in

585

590



responsive to changes in VOCs. In the context of wildfire smoke, the influx of VOCs from the fires can shift the chemical regime from NO_x -limited to VOC-limited, altering the dynamics of O_3 production in the urban area. This shift can lead to unexpected increases in O_3 levels as the balance of precursors is altered by the incoming smoke plume. The $\text{H}_2\text{O}_2/\text{HNO}_3$ ratio serves as another
595 indicator of the oxidizing capacity of the atmosphere and the relative contributions of different chemical pathways to O_3 formation. H_2O_2 (hydrogen peroxide) is a product of VOC oxidation, while HNO_3 (nitric acid) results from the oxidation of NO_x . A higher ratio suggests an environment with abundant VOC oxidation, often associated with high levels of O_3 production (Sillman, 1995). Conversely, a lower ratio indicates a dominance of NO_x oxidation pathways, which can suppress
600 O_3 formation under certain conditions. The presence of wildfire emissions can increase the levels of both VOCs and NO_x , thereby influencing this ratio and providing insights into the changing oxidative environment over Phoenix. The relative change in VOCs and NO_x will affect O_3 sensitivity, depending on which of these pollutants has a larger percentage change. Miech et al. (2024) found that at Phoenix JLG supersite, when the sensitivity is under VOC-limited, FNR is
605 higher than normal suggesting elevated VOCs relative to NO_2 under a smoke event and shifting the sensitivity towards a transitional or NO_x -limited. This is also seen in Figure 15 where levels of CO and HCHO are relatively elevated than NO_x along the fire plume trajectory.

In Figure 16, the analysis of these two surface ratios reveals how wildfire smoke alters the chemical regime over Phoenix at local time 14:00 when O_3 production is expected to peak. Without the
610 smoke plume, the majority of Phoenix urban area in the early afternoon, when the photolysis is highest, is under a transitional/ NO_x -limited regime (Figure 16a). With the presence of smoke, additional NO_x and VOCs are brought to the region and the regime shifts towards more NO_x -limited in the central urban region, as seen by the increase (orange contours) in the FNR (Figure 16c), consistent with Miech et al. (2024). In contrast, FNR decreases across the broader extent of
615 the fire, (blue contours), most likely with the introduction of NO_x from PAN decomposition further downwind. Accordingly, the $\text{H}_2\text{O}_2/\text{HNO}_3$ ratio provides a similar picture of the increasing ratios in the metropolitan area due to the influx of emissions from the fires.




Figure 16. Sensitivity analysis of different ratios in Case I at local time 14:00 under conditions with and without fire. The top row (a-c) is the surface HCHO/NO₂ ratio, while the bottom row (d-f) depicts the surface H₂O₂/HNO₃ ratio. Ratios larger than 5 are shown as white space. (a) and (d) show the respective ratios without fire, and (b) and (e) display the ratios with fire. (c) and (f) represent the differences in these ratios between the scenarios with and without fire. The color scales indicate the magnitude of these ratios and differences, with red and blue colors denoting higher and lower values, respectively.

4. Conclusion

620 This study focuses on the contribution of wildfire emissions on surface O₃ levels in the Phoenix metropolitan area during June 2021, a period characterized by extreme heat and drought conditions. We apply model sensitivity and species tagging approaches in the WRF-Chem model to quantify its contribution relative to other sources. Two specific cases (Telegraph and Rafael fires) are chosen for analyzing the wildfire impact on urban regions. During these cases, the MDA8 O₃ is
625 observed to exceed the NAAQS standard (70 ppb), especially for case I (Telegraph fire), with concentrations reaching 110 ppb.

Overall, the O₃ levels in Arizona are influenced by a combination of background levels, local anthropogenic emissions, and wildfire contributions. The highest O₃ levels were observed around urban centers, with wildfires significantly contributing to elevated O₃, especially near the fire sites
630 and downwind areas. The spatial and temporal distributions of CO and O₃, as well as the contributions from different tags in Arizona during the wildfire season, reveal significant contributions of both anthropogenic and wildfire emissions to CO levels across the state, with local



urban emissions still playing a dominant role in areas like Phoenix and Tucson. However, wildfire emissions were particularly impactful in regions downwind of the fires.

635 In fact, our simulations show that wildfire emissions notably increased the MDA8 O₃ levels during two fire plume case studies that we examined. The Telegraph Fire, in particular, contributed to significant O₃ levels on June 15. The results demonstrated that background O₃ levels account the bulk of total O₃ (around 50%), with local anthropogenic emissions contributing significantly (24% to 40%) depending on the urban setting. During peak O₃ hours, fire-contributed O₃ was significant
640 across multiple sites, ranging from 5 to 23 ppb or 5% to 21 % of total O₃ levels, with an average of 15 ppb or 15%. Without smoke, the Phoenix urban area is primarily under a transition to NO_x-limited regime in the early afternoon when O₃ photolysis rate is highest. With smoke present, the central urban region becomes more NO_x-limited due to the addition of VOCs transported from the fires relative to NO_x which is already high from local anthropogenic emissions. In contrast, the
645 suburban and rural areas downwind of the fires generally experience a decrease in the ratio shifting towards a more VOC-limited regime, which is likely due to the addition of NO_x from fires as a result of thermal decomposition of PAN from the fires transported to these areas.

By closely investigating these tags, we also find differences between Phoenix and Yuma. Unlike Yuma, where O₃ levels are significantly influenced by transboundary emissions from California
650 and Mexico, Phoenix's O₃ levels are primarily driven by local emissions, with much smaller contributions from these external sources during the study period. Specifically for a smoky day, during the diurnal cycle of O₃ levels, anthropogenic emission contributed local O₃ production dominate in the early morning and early afternoon, while fire-related O₃ contributions increase significantly in the late afternoon when a smoke plume passes through. This pattern is observed
655 across suburban, urban, and rural settings, with fire related O₃ sometimes surpassing local anthropogenic production in rural areas.

As has been suggested by previous studies, the substantial enhancements in O₃ concentrations due to wildfire emissions highlight the necessity of accounting for wildfire impacts in formulating effective air quality management strategies. Such strategies should consider the influence of fire
660 emissions on urban O₃ and more notably their subsequent interactions with local emissions, chemistry, and meteorology (e.g., He et al., 2024) to help provide additional perspectives on current O₃ pollution assessments. This is especially the case over urban areas in semi-arid/arid



environments like southwest United States, where confounding (and compounding) factors arising from meteorological extremes and dynamical challenges due to complex topography are present; 665 notwithstanding the effect of climate change to increasing global aridity and fire frequency and intensity. The adoption of source attribution approaches (including the use of data-driven techniques) within coupled weather-air quality models, accompanied by more comprehensive evaluation, can be a useful complementary activity to consider in helping support current approaches in understanding O₃.

670

Code and Data Availability Statement.

The WRF-Chem model is publicly available at <https://github.com/wrf-model/WRF> (last access: 25 June 2022). The model outputs can be provided upon request to the corresponding author. EPA AQS and PAMS hourly and daily datasets are available at 675 https://aq5.epa.gov/aqsweb/airdata/download_files.html.

Author contributions.

YG and AA designed the research. YG performed the model runs and subsequent analysis. YG wrote the paper with contributions from AA and AS. AM helped with the satellite data analysis.

Competing Interests.

680 Some authors are members of the editorial board of journal Atmospheric Chemistry and Physics.

Acknowledgments.

We especially acknowledge Dr. Gabriele Pfister and Dr. Louisa Emmons at NCAR/ACOM for kindly providing the resources for implementing tagging in WRF-Chem, for which this study is being built upon. We also thank Matthew Pace, Rene Nsanzineza, and Michael Graves at the 685 Arizona Department of Environmental Quality (ADEQ) for the help with observations for the state of Arizona.

Financial Support.

This work is supported by the Arizona Board of Regents (ABOR) Grant from the Technology and Research Initiative Fund (TRIF).

690



References:

- Abatzoglou, J. T. and Williams, A. P.: Impact of anthropogenic climate change on wildfire across western US forests, *Proceedings of the National Academy of Sciences*, 113, 11770-11775, doi:10.1073/pnas.1607171113, 2016.
- 695 Adhikari, A. and Yin, J.: Short-Term Effects of Ambient Ozone, PM_{2.5}, and Meteorological Factors on COVID-19 Confirmed Cases and Deaths in Queens, New York, *International Journal of Environmental Research and Public Health*, 17, 4047, doi:10.3390/ijerph17114047, 2020.
- Akagi, S. K., Craven, J. S., Taylor, J. W., McMeeking, G. R., Yokelson, R. J., Burling, I. R., Urbanski, S. P., Wold, C. E., Seinfeld, J. H., Coe, H., Alvarado, M. J., and Weise, D. R.:
700 Evolution of trace gases and particles emitted by a chaparral fire in California, *Atmos. Chem. Phys.*, 12, 1397-1421, 10.5194/acp-12-1397-2012, 2012.
- Alonso-Blanco, E., Castro, A., Calvo, A. I., Pont, V., Mallet, M., and Fraile, R.: Wildfire smoke plumes transport under a subsidence inversion: Climate and health implications in a distant urban area, *Science of The Total Environment*, 619-620, 988-1002,
705 <https://doi.org/10.1016/j.scitotenv.2017.11.142>, 2018.
- Andreae, M. O.: Emission of trace gases and aerosols from biomass burning – an updated assessment, *Atmos. Chem. Phys.*, 19, 8523-8546, 10.5194/acp-19-8523-2019, 2019.
- Betito, G., Arellano, A., and Sorooshian, A.: Influence of Transboundary Pollution on the Variability of Surface Ozone Concentrations in the Desert Southwest of the U.S.: Case Study for
710 Arizona, *Atmosphere*, 15, 401, 10.3390/atmos15040401, 2024.
- Brey, S. J., Ruminski, M., Atwood, S. A., and Fischer, E. V.: Connecting smoke plumes to sources using Hazard Mapping System (HMS) smoke and fire location data over North America, *Atmos. Chem. Phys.*, 18, 1745–1761, <https://doi.org/10.5194/acp-18-1745-2018>, 2018.
- Burke, M., Childs, M. L., de la Cuesta, B., Qiu, M., Li, J., Gould, C. F., Heft-Neal, S., and Wara, M.: The contribution of wildfire to PM(2.5) trends in the USA, *Nature*, 622, 761-766,
715 10.1038/s41586-023-06522-6, 2023.
- Butler, T., Lupascu, A., and Nalam, A.: Attribution of ground-level ozone to anthropogenic and natural sources of nitrogen oxides and reactive carbon in a global chemical transport model, *Atmos. Chem. Phys.*, 20, 10707-10731, 10.5194/acp-20-10707-2020, 2020.
- 720 Butler, T., Lupascu, A., Coates, J., and Zhu, S.: TOAST 1.0: Tropospheric Ozone Attribution of Sources with Tagging for CESM 1.2.2, *Geosci. Model Dev.*, 11, 2825-2840, 10.5194/gmd-11-2825-2018, 2018.
- Buysse, C. E., Kaulfus, A., Nair, U., and Jaffe, D. A.: Relationships between Particulate Matter, Ozone, and Nitrogen Oxides during Urban Smoke Events in the Western US, *Environmental Science & Technology*, 53, 12519-12528, 10.1021/acs.est.9b05241, 2019.
- 725 Cohan, D. S., Hakami, A., Hu, Y., and Russell, A. G.: Nonlinear Response of Ozone to Emissions: Source Apportionment and Sensitivity Analysis, *Environmental Science & Technology*, 39, 6739-6748, 10.1021/es048664m, 2005.
- Collet, S., Minoura, H., Kidokoro, T., Sonoda, Y., Kinugasa, Y., Karamchandani, P., Johnson, J., Shah, T., Jung, J., and DenBleyker, A.: Future year ozone source attribution modeling studies for
730



- the eastern and western United States, *Journal of the Air & Waste Management Association*, 64, 1174-1185, 10.1080/10962247.2014.936629, 2014.
- De la Paz, D., Borge, R., de Andrés, J. M., Tovar, L., Sarwar, G., and Napelenok, S. L.:
735 Summertime tropospheric ozone source apportionment study in the Madrid region (Spain),
Atmos. Chem. Phys., 24, 4949-4972, 10.5194/acp-24-4949-2024, 2024.
- Demerjian, K. L.: A review of national monitoring networks in North America, *Atmospheric Environment*, 34, 1861-1884, [https://doi.org/10.1016/S1352-2310\(99\)00452-5](https://doi.org/10.1016/S1352-2310(99)00452-5), 2000.
- Emmons, L. K., Hess, P. G., Lamarque, J. F., and Pfister, G. G.: Tagged ozone mechanism for
740 MOZART-4, CAM-chem and other chemical transport models, *Geosci. Model Dev.*, 5, 1531-
1542, 10.5194/gmd-5-1531-2012, 2012.
- Freitas, S. R., Longo, K. M., Chatfield, R., Latham, D., Silva Dias, M. A. F., Andreae, M. O.,
Prins, E., Santos, J. C., Gielow, R., and Carvalho Jr, J. A.: Including the sub-grid scale plume
rise of vegetation fires in low resolution atmospheric transport models, *Atmos. Chem. Phys.*, 7,
3385-3398, 10.5194/acp-7-3385-2007, 2007.
- 745 Gard, H. and Garrett, M.: Wildfire smoke stopped Phoenix from breaking record high of 115
degrees, 2021.
- Goldberg, D. L., Vinciguerra, T. P., Anderson, D. C., Hemberck, L., Canty, T. P., Ehrman, S. H.,
... & Dickerson, R. R.: CAMx ozone source attribution in the eastern United States using
guidance from observations during DISCOVER-AQ Maryland. *Geophysical research letters*,
750 43(5), 2249-2258, doi:10.1002/2015GL067332, 2016.
- Grell, G. A., Peckham, S. E., Schmitz, R., McKeen, S. A., Frost, G., Skamarock, W. C., and
Eder, B.: Fully coupled “online” chemistry within the WRF model, *Atmospheric Environment*,
39, 6957-6975, <https://doi.org/10.1016/j.atmosenv.2005.04.027>, 2005.
- Grewe, V., Tsati, E., and Hoor, P.: On the attribution of contributions of atmospheric trace gases
755 to emissions in atmospheric model applications, *Geosci. Model Dev.*, 3, 487-499, 10.5194/gmd-
3-487-2010, 2010.
- Grewe, V., Dahlmann, K., Matthes, S., and Steinbrecht, W.: Attributing ozone to NO_x
emissions: Implications for climate mitigation measures, *Atmospheric Environment*, 59, 102-
107, <https://doi.org/10.1016/j.atmosenv.2012.05.002>, 2012.
- 760 Grewe, V., Tsati, E., Mertens, M., Frömming, C., and Jöckel, P.: Contribution of emissions to
concentrations: the TAGGING 1.0 submodel based on the Modular Earth Submodel System
(MESSy 2.52), *Geosci. Model Dev.*, 10, 2615-2633, 10.5194/gmd-10-2615-2017, 2017.
- Guenther, A.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of
Emissions of Gases and Aerosols from Nature), *Atmospheric Chemistry and Physics*, 7, 4327-
765 4327, 10.5194/acp-6-3181-2006, 2007.
- Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of
global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols
from Nature), *Atmos. Chem. Phys.*, 6, 3181-3210, 10.5194/acp-6-3181-2006, 2006.
- 770 Gao, J., Zhu, B., Xiao, H., Kang, H., Hou, X., & Shao, P., A case study of surface ozone source
apportionment during a high concentration episode, under frequent shifting wind conditions over



- the Yangtze River Delta, China. *Science of the Total Environment*, 544, 853-863, 10.1016/j.scitotenv.2015.12.039, 2016
- 775 Guo, Y., Roychoudhury, C., Mirrezaei, M. A., Kumar, R., Sorooshian, A., and Arellano, A. F.: Investigating ground-level ozone pollution in semi-arid and arid regions of Arizona using WRF-Chem v4.4 modeling, *Geosci. Model Dev.*, 17, 4331-4353, 10.5194/gmd-17-4331-2024, 2024.
- Hakami, A., Seinfeld, J. H., Chai, T., Tang, Y., Carmichael, G. R., & Sandu, A., Adjoint sensitivity analysis of ozone nonattainment over the continental United States. *Environmental science & technology*, 40(12), 3855-3864, 10.1021/es052135g, 2006.
- 780 Hakami, A., Henze, D. K., Seinfeld, J. H., Singh, K., Sandu, A., Kim, S., Li, Q., The adjoint of CMAQ. *Environmental science & technology*, 41(22), 7807-7817, 10.1021/es070944p, 2007.
- He C., Kumar R., Tang, W., Pfister, G., Xu, Y., Qian, Y., Brasseur, G., Air Pollution Interactions with Weather and Climate Extremes: Current Knowledge, Gaps, and Future Directions. *Current Pollution Reports*, 22, 1-3, 10.1007/s40726-024-00296-9, 2024.
- 785 Huangfu, P. and Atkinson, R.: Long-term exposure to NO₂ and O₃ and all-cause and respiratory mortality: A systematic review and meta-analysis, *Environment International*, 144, 105998, <https://doi.org/10.1016/j.envint.2020.105998>, 2020.
- Jaffe, D. A. and Wigder, N. L.: Ozone production from wildfires: A critical review, *Atmospheric Environment*, 51, 1-10, <https://doi.org/10.1016/j.atmosenv.2011.11.063>, 2012.
- 790 Jaffe, D. A., Wigder, N., Downey, N., Pfister, G., Boynard, A., and Reid, S. B.: Impact of Wildfires on Ozone Exceptional Events in the Western U.S, *Environmental Science & Technology*, 47, 11065-11072, 10.1021/es402164f, 2013.
- Jaffe, D. A., O'Neill, S. M., Larkin, N. K., Holder, A. L., Peterson, D. L., Halofsky, J. E., and Rappold, A. G.: Wildfire and prescribed burning impacts on air quality in the United States, *Journal of the Air & Waste Management Association*, 70, 583-615, 10.1080/10962247.2020.1749731, 2020.
- 795 Jin, X., Fiore, A., Boersma, K. F., Smedt, I. D., & Valin, L., Inferring changes in summertime surface Ozone-NO_x-VOC chemistry over US urban areas from two decades of satellite and ground-based observations, *Environmental science & technology*, 54(11), 6518-6529, doi: 10.1021/acs.est.9b07785, 2020
- 800 Jin, X., Fiore, A. M., and Cohen, R. C.: Space-Based Observations of Ozone Precursors within California Wildfire Plumes and the Impacts on Ozone-NO_x-VOC Chemistry, *Environmental Science & Technology*, 57, 14648-14660, 10.1021/acs.est.3c04411, 2023.
- Kwok, R. H. F., Baker, K. R., Napelenok, S. L., & Tonnesen, G. S., Photochemical grid model implementation and application of VOC, NO_x, and O₃ source apportionment, *Geoscientific Model Development*, 8(1), 99-114, 10.5194/gmd-8-99-2015, 2015.
- 805 Li, J., Georgescu, M., Hyde, P., Mahalov, A., and Moustaoui, M.: Regional-scale transport of air pollutants: impacts of Southern California emissions on Phoenix ground-level ozone concentrations, *Atmos. Chem. Phys.*, 15, 9345-9360, 10.5194/acp-15-9345-2015, 2015.
- 810 Li, P., Yang, Y., Wang, H., Li, S., Li, K., Wang, P., Li, B., and Liao, H.: Source attribution of near-surface ozone trends in the United States during 1995-2019, *Atmos. Chem. Phys.*, 23, 5403-5417, 10.5194/acp-23-5403-2023, 2023.



- 815 Ludewig, A., Kleipool, Q., Bartstra, R., Landzaat, R., Leloux, J., Loots, E., Meijering, P., van der Plas, E., Rozemeijer, N., Vonk, F., and Veeffkind, P.: In-flight calibration results of the TROPOMI payload on board the Sentinel-5 Precursor satellite, *Atmos. Meas. Tech.*, 13, 3561-3580, 10.5194/amt-13-3561-2020, 2020.
- Lupaşcu, A. and Butler, T.: Source attribution of European surface O₃ using a tagged O₃ mechanism, *Atmos. Chem. Phys.*, 19, 14535-14558, 10.5194/acp-19-14535-2019, 2019.
- 820 Lupaşcu, A., Otero, N., Minkos, A., and Butler, T.: Attribution of surface ozone to NO_x and volatile organic compound sources during two different high ozone events, *Atmos. Chem. Phys.*, 22, 11675-11699, 10.5194/acp-22-11675-2022, 2022.
- Marsh, D. R., Mills, M. J., Kinnison, D. E., Lamarque, J.-F., Calvo, N., and Polvani, L. M.: Climate Change from 1850 to 2005 Simulated in CESM1(WACCM), *Journal of Climate*, 26, 7372-7391, <https://doi.org/10.1175/JCLI-D-12-00558.1>, 2013.
- 825 Martin, R. V., Fiore, A. M., & Van Donkelaar, A., Space-based diagnosis of surface ozone sensitivity to anthropogenic emissions, *Geophysical Research Letters*, 31(6), doi:10.1029/2004GL019416, 2004.
- McClure, C. D. and Jaffe, D. A.: Investigation of high ozone events due to wildfire smoke in an urban area, *Atmospheric Environment*, 194, 146-157, <https://doi.org/10.1016/j.atmosenv.2018.09.021>, 2018.
- 830 Miech, J. A., Herckes, P., Fraser, M. P., Arellano, A. F., Mirrezaei, M. A., and Guo, Y.: Evaluating Phoenix Metropolitan Area Ozone Behavior Using Ground-Based Sampling, Modeling, and Satellite Retrievals, *Atmosphere*, 15, 555, doi:10.3390/atmos15050555, 2024.
- 835 Mirrezaei, M. A., Arellano, A., Guo, Y., Roychoudhury, C., & Sorooshian, A.. Ozone production over arid regions: insights into meteorological and chemical drivers. *Environmental Research Communications*, 6(5), 051009, doi:10.1088/2515-7620/ad484c, 2024.
- Nalam, A., Lupascu, A., Ansari, T., and Butler, T.: Regional and sectoral contributions of NO_x and reactive carbon emission sources to global trends in tropospheric ozone during the 2000-2018 period, *EGUsphere*, 2024, 1-39, 10.5194/egusphere-2024-432, 2024.
- 840 Ninneman, M. and Jaffe, D. A.: The impact of wildfire smoke on ozone production in an urban area: Insights from field observations and photochemical box modeling, *Atmospheric Environment*, 267, 118764, <https://doi.org/10.1016/j.atmosenv.2021.118764>, 2021.
- National Oceanic and Atmospheric Administration (NOAA) (2023). Hazard Mapping System Fire and Smoke Product. Retrieved from <https://www.ospo.noaa.gov/Products/land/hms.html>. Accessed on April 8, 2024.
- 845 Pfister, G. G., Walters, S., Emmons, L. K., Edwards, D. P., and Avise, J.: Quantifying the contribution of inflow on surface ozone over California during summer 2008, *Journal of Geophysical Research: Atmospheres*, 118, 12,282-212,299, <https://doi.org/10.1002/2013JD020336>, 2013.
- 850 Pfister, G. G., Emmons, L. K., Hess, P. G., Honrath, R., Lamarque, J.-F., Val Martin, M., Owen, R. C., Avery, M. A., Browell, E. V., Holloway, J. S., Nedelec, P., Purvis, R., Ryerson, T. B., Sachse, G. W., and Schlager, H.: Ozone production from the 2004 North American boreal fires,



- Journal of Geophysical Research: Atmospheres, 111, <https://doi.org/10.1029/2006JD007695>, 2006.
- 855 Rolph, G. D., Draxler, R. R., Stein, A. F., Taylor, A., Ruminski, M. G., Kondragunta, S., Zeng, J., Huang, H., Manikin, G., McQueen, J. T., & Davidson, P. M., Description and Verification of the NOAA Smoke Forecasting System: The 2007 Fire Season, *Weather and Forecasting*, 24(2), 361-378. <https://doi.org/10.1175/2008WAF2222165.1>, 2009.
- 860 Ruminski, M., Kondragunta, S., Monitoring fire and smoke emissions with the hazard mapping system, *Proc. SPIE 6412, Disaster Forewarning Diagnostic Methods and Management*, 64120B, <https://doi.org/10.1117/12.694183>, 2006.
- Selimovic, V., Yokelson, R. J., McMeeking, G. R., and Coefield, S.: Aerosol Mass and Optical Properties, Smoke Influence on O₃, and High NO₃ Production Rates in a Western U.S. City Impacted by Wildfires, *Journal of Geophysical Research: Atmospheres*, 125, e2020JD032791, <https://doi.org/10.1029/2020JD032791>, 2020.
- 865 Shu, Q., Napelenok, S. L., Hutzell, W. T., Baker, K. R., Henderson, B. H., Murphy, B. N., and Hogrefe, C.: Comparison of ozone formation attribution techniques in the northeastern United States, *Geosci. Model Dev.*, 16, 2303-2322, 10.5194/gmd-16-2303-2023, 2023.
- 870 Sillman, S.: The use of NO_y, H₂O₂ and HNO₃ as indicators for ozone-NO_x-hydrocarbon sensitivity in urban locations, *Journal of Geophysical Research: Atmospheres*, 100, 14175-14188, 10.1029/94JD02953, 1995.
- Sorooshian, A., Arellano, A. F., Fraser, M. P., Herckes, P., Betito, G., Betterton, E. A., Braun, R. A., Guo, Y., Mirrezaei, M. A., and Roychoudhury, C.: Ozone in the Desert Southwest of the United States: A Synthesis of Past Work and Steps Ahead, *ACS ES&T Air*, 1, 62-79, 10.1021/acsestair.3c00033, 2024.
- 875 Thunis, P., Clappier, A., Tarrason, L., Cuvelier, C., Monteiro, A., Pisoni, E., Wesseling, J., Belis, C. A., Pirovano, G., Janssen, S., Guerreiro, C., and Peduzzi, E.: Source apportionment to support air quality planning: Strengths and weaknesses of existing approaches, *Environment International*, 130, 104825, <https://doi.org/10.1016/j.envint.2019.05.019>, 2019.
- 880 Tilmes, S., Lamarque, J. F., Emmons, L. K., Kinnison, D. E., Ma, P. L., Liu, X., Ghan, S., Bardeen, C., Arnold, S., Deeter, M., Vitt, F., Ryerson, T., Elkins, J. W., Moore, F., Spackman, J. R., and Val Martin, M.: Description and evaluation of tropospheric chemistry and aerosols in the Community Earth System Model (CESM1.2), *Geosci. Model Dev.*, 8, 1395-1426, 10.5194/gmd-8-1395-2015, 2015.
- 885 Tonnesen, G.S., Dennis, R.L., Analysis of radical propagation efficiency to assess ozone sensitivity to hydrocarbons and NO_x: 2. Long-lived species as indicators of ozone concentration sensitivity. *J. Geophys. Res. Atmos.* 105, 9227-9241, <https://doi.org/10.1029/1999JD900372>, 2000.
- 890 Turner, M. C., Jerrett, M., III, C. A. P., Krewski, D., Gapstur, S. M., Diver, W. R., Beckerman, B. S., Marshall, J. D., Su, J., Crouse, D. L., and Burnett, R. T.: Long-Term Ozone Exposure and Mortality in a Large Prospective Study, *American Journal of Respiratory and Critical Care Medicine*, 193, 1134-1142, 10.1164/rccm.201508-1633OC, 2016.



- van Geffen, J., Boersma, K. F., Eskes, H., Sneep, M., ter Linden, M., Zara, M., and Veeffkind, J. P.: S5P TROPOMI NO₂ slant column retrieval: method, stability, uncertainties and comparisons with OMI, *Atmos. Meas. Tech.*, 13, 1315-1335, 10.5194/amt-13-1315-2020, 2020.
- 895 Wiedinmyer, C., Kimura, Y., McDonald-Buller, E. C., Emmons, L. K., Buchholz, R. R., Tang, W., Seto, K., Joseph, M. B., Barsanti, K. C., Carlton, A. G., and Yokelson, R.: The Fire Inventory from NCAR version 2.5: an updated global fire emissions model for climate and chemistry applications, *Geosci. Model Dev.*, 16, 3873-3891, 10.5194/gmd-16-3873-2023, 2023.
- 900 Xu, L., Crounse, J. D., Vasquez, K. T., Allen, H., Wennberg, P. O., Bourgeois, I., Brown, S. S., Campuzano-Jost, P., Coggon, M. M., Crawford, J. H., DiGangi, J. P., Diskin, G. S., Fried, A., Gargulinski, E. M., Gilman, J. B., Gkatzelis, G. I., Guo, H., Hair, J. W., Hall, S. R., Halliday, H. A., Hanisco, T. F., Hannun, R. A., Holmes, C. D., Huey, L. G., Jimenez, J. L., Lamplugh, A., Lee, Y. R., Liao, J., Lindaas, J., Neuman, J. A., Nowak, J. B., Peischl, J., Peterson, D. A., Piel, F., Richter, D., Rickly, P. S., Robinson, M. A., Rollins, A. W., Ryerson, T. B., Sekimoto, K., Selimovic, V., Shingler, T., Soja, A. J., St. Clair, J. M., Tanner, D. J., Ullmann, K., Veres, P. R., Walega, J., Warneke, C., Washenfelder, R. A., Weibring, P., Wisthaler, A., Wolfe, G. M., Womack, C. C., and Yokelson, R. J.: Ozone chemistry in western U.S. wildfire plumes, *Science Advances*, 7, eabl3648, doi:10.1126/sciadv.abl3648, 2021.
- 910 Yeganeh, A. K., Momeni, M., Choi, Y., Park, J., and Jung, J.: A case study of surface ozone source contributions in the Seoul metropolitan area using the adjoint of CMAQ, *Journal of the Air & Waste Management Association*, null-null, 10.1080/10962247.2024.2361021, 2024.
- Zhang, R., Cohan, A., Pour Biazar, A., and Cohan, D. S.: Source apportionment of biogenic contributions to ozone formation over the United States, *Atmospheric Environment*, 164, 8-19, <https://doi.org/10.1016/j.atmosenv.2017.05.044>, 2017.
- 915 Zhang, Y., Zhao, Y., Li, J., Wu, Q., Wang, H., Du, H., Yang, W., Wang, Z., and Zhu, L.: Modeling Ozone Source Apportionment and Performing Sensitivity Analysis in Summer on the North China Plain, *Atmosphere*, 11, 992, doi:10.3390/atmos11090992, 2020.
- 920 Zhao, Y., Li, Y., Kumar, A., Ying, Q., Vandenberghe, F., & Kleeman, M. J., Separately resolving NO_x and VOC contributions to ozone formation. *Atmospheric Environment*, 285, 119224, 10.1016/j.atmosenv.2022.119224, 2022.