

A thicker-than-present East Antarctic Ice Sheet plateau during the Last Glacial Maximum

Cari Rand¹, Richard S. Jones¹, Andrew N. Mackintosh¹, Brent Goehring², Kat Lilly³

¹Securing Antarctica's Environmental Future, School of Earth, Atmosphere and Environment, Monash University, Wellington Road, Clayton, Melbourne, Victoria 3800, Australia

²Los Alamos National Laboratory, Bikini Atoll Road, Los Alamos, New Mexico 87545, USA

³RSC, PO Box 5647, Dunedin, New Zealand

Correspondence to: Cari Rand (cari.rand@monash.edu)

Abstract. In this study, we present a surface-exposure chronology of past ice-thickness change derived from *in-situ* cosmogenic-¹⁴C dating at a site in the Grove Mountains, located on the edge of the East Antarctic plateau, 380 km inland from the coastline in the Lambert Glacier-Amery Ice Shelf sector. At the Last Glacial Maximum (LGM), observations and models suggest that increased ice-sheet volume was accommodated by thicker ice near the coast and grounding-line advance towards the continental-shelf edge. In contrast, the ice-sheet interior maintained a relatively stable thickness until present, with ice-core evidence even suggesting thinner ice relative to today. However, the magnitude of these thickness changes and the transition point from thicker-than-present to thinner-than-present LGM ice are poorly constrained. Here, we reconstruct changes in the thickness of the East Antarctic Ice Sheet since the LGM at a nunatak in the Grove Mountains using *in-situ* ¹⁴C, which circumvents the common issue of long-lived nuclide inheritance that leads to inaccurate records of LGM ice thickness. Samples between 1,912 m above sea level (a.s.l.) and the modern ice margin (~1,825 m a.s.l.) yield ¹⁴C ages of 0.18–5.26 ka. Samples at and above 1,912 m a.s.l. have saturated ¹⁴C concentrations, implying exposure of the nunatak summit through the LGM. We therefore place the LGM ice surface in the Grove Mountains ~70 m higher than at present. The unsaturated samples below 1,912 m a.s.l. indicate that gradual thinning began ~16 ka, with some (25–45%) post-LGM thinning recorded ~16–11 ka and most (55–75%) recorded during the Holocene. Ice-sheet models that do not account for this thickness change would inaccurately characterize the LGM geometry of the EAIS and underestimate its contributions to deglacial sea-level rise.

1 Introduction

The East Antarctic Ice Sheet (EAIS) is the largest contiguous mass of ice on Earth (Rignot *et al.*, 2019). Loss of ice to melting and calving is predicted to be offset by increases in snow accumulation over the coming century, but beyond 2100 CE, the ice sheet is expected to lose mass and contribute to sea-level rise (Stokes *et al.*, 2022). Characterizing past changes of the EAIS is necessary for several reasons:

1. Satellite observations of Antarctic glaciers extend back only to the 1960s, so other records of past ice-sheet states are needed in order to reliably distinguish long-term trends from natural variability (Hanna *et al.*, 2020; Jones *et al.*, 2022);
2. Geodetic data used to estimate modern ice-mass changes must be corrected for glacial isostatic adjustment (e.g., Coulon *et al.*, 2021), the magnitude of which is dependent on the past configuration of the ice sheet;

3. Determining the magnitude and timing of ice loss can identify or exclude potential sources of meltwater input to oceans during past periods of rapid sea-level rise (e.g., Lin *et al.*, 2021); and
4. Numerical models informed by records of past ice-sheet change can be used to estimate the future contributions to sea-level rise (e.g., DeConto *et al.*, 2021).

However, reconstructing the geometry of the EAIS is challenging. Evidence of past ice thickness comes from radar, ice-core and geological data, which are sparse owing to the remoteness of East Antarctica, the large area of the ice sheet, and the sparsity of ice-free areas. Furthermore, different records of LGM ice thickness are often in disagreement with one another.

During the Last Glacial Maximum (LGM), at approximately 20 ka, available evidence points towards a more extensive but shallower-gradient ice sheet (Mackintosh *et al.*, 2014). Dated acid-insoluble organic matter in sediments from the East Antarctic coast indicate that the EAIS advanced to the edge of the continental shelf in most locations during the LGM (Bentley *et al.*, 2014), with constraints from cosmogenic ^{10}Be and ^{26}Al indicating the presence of ice near the coast that was thicker than it is now (e.g., Mackintosh *et al.*, 2007; White *et al.*, 2011a; Yamane *et al.*, 2011). Meanwhile, snow-accumulation rates interpolated between ice domes suggest a thinner ice sheet across the East Antarctic plateau (Buizert *et al.*, 2021) at the LGM. A “hinge zone” thus likely existed between thicker ice at the coast and thinner ice in the interior during the LGM relative to today (Bockheim *et al.*, 1989; Andersen *et al.*, 2023), but the location of this transition point across East Antarctica is unclear. Cosmogenic ^{10}Be and ^{26}Al ages from ice-free areas on the edge of the East Antarctic plateau such as the Grove Mountains or southern Prince Charles Mountains (PCMs) are older than the LGM (e.g., Lilly *et al.*, 2010; White *et al.*, 2011a), implying no change since or slightly thinner ice in these locations at the LGM (Fig. 1).

Existing cosmogenic-nuclide data from regions of cold-based non-erosive ice, however, may not provide reliable constraints on LGM ice thickness. Many samples have apparently pre-LGM and inconsistent ^{10}Be ($t_{0.5}$) and ^{26}Al ($t_{0.5}$) exposure ages, indicating nuclide inventories inherited from previous periods of exposure (Balco *et al.*, 2014). Due to the short half-life of *in situ* ^{14}C (5.7 kyr; Nichols, 2022), its concentration decays quickly when shielded (e.g., when covered by ice; Goehring, *et al.*, 2019a); this makes *in situ* ^{14}C a useful tool for investigating post-LGM glacial history (Nichols, 2022).




Figure 1: Constraints on central East Antarctic ice thickness at the Last Glacial Maximum (LGM). (a) Inferred LGM-to-present ice-thickness differences near Lambert Glacier. Dashed black line shows the 1900 m a.s.l. contour (Liu *et al.* 2015), the elevation of our sampled nunatak. This line represents the most interior geological evidence and reflects a potential hinge zone between coastal and interior LGM-ice-thickness change. **If coastal ice thins and interior ice thickens after the LGM, the modern ice-surface profile would intersect the LGM surface profile somewhere in the middle; this intersection is the “hinge zone”, at which ice there has been no net change in ice thickness since the LGM.** LGM-thickness reconstructions in White *et al.* (2011a) and Lilly *et al.* (2010) placed the “hinge zone” in areas equivalent to a present-day ice-surface elevation of ~1900 m a.s.l. Elements of this map were provided by the Quantarctica 3 GIS package provided by the Norwegian Polar Institute (Matsuoka *et al.*, 2018), including ice-free areas (Burton-Johnson *et al.*; 2016), the current Antarctic-ice-sheet grounding line (Bindshadler *et al.*, 2011), and the inferred East Antarctic grounding line 20 ka (Bentley *et al.*, 2014). Pink-shaded ice indicates the extent of the catchment of Lambert Glacier (Zwally *et al.*, 2012). LGM thickness data for this figure come from Buizert *et al.* (2021; Dome Fuji and EPICA Dome C), Lilly *et al.* (2010; Grove Mountains), Mackintosh *et al.* (2007; Framnes Mountains), Mackintosh *et al.* (2014) and references therein (Bunger Hills, Law Dome, Vestfold Hills, Vostok Station, and Windmill Islands), and White *et al.* (2011a; Prince Charles Mountains). (b) Satellite view of the study area with the sampled nunatak (Nunatak 1921). Bedrock and erratic samples were

collected in a transect extending from the modern ice surface to nunatak summit. Ice at this site flows slowly (blue arrow; Rignot *et al.*, 2011) northwest, towards the Amery Ice Shelf, though flow speeds are low and directions are strongly influenced by topography in the vicinity of nunataks. (c) Diagram illustrating the concept of a “hinge zone” in ice-thickness change. Image shows hypothetical vertically exaggerated cross-sections of the East Antarctic Ice Sheet at the LGM (dotted line) and present day (dashed line). The Grove Mountains are a key site for testing the location of this “hinge zone” as they lie close to the elevation of the “hinge zone” identified in previous studies of the region (e.g., White *et al.*, 2011a).

In this study, we aim to constrain the position of the “hinge zone” in this region by measuring *in-situ* ^{14}C in bedrock and erratic samples previously measured for ^{10}Be and ^{26}Al from the Grove Mountains, a key site in the ice-sheet interior. Rocks exposed since before the LGM should have concentrations of ^{14}C at secular equilibrium between production and decay (saturation), a state that requires ca. 5 half-lives of continuous exposure (Dunai, 2010). Conversely, rocks with less-than-saturated concentrations of ^{14}C from a site in East Antarctica imply that those samples were likely covered for some duration post-LGM by a thicker-than-present EAIS that subsequently thinned. The concentration of a cosmogenic nuclide in a sample will remain at secular equilibrium indefinitely unless disturbed by cover, erosion, or transport; thus, only a minimum age can be assigned to saturated samples. Measuring samples from an elevation transect with *in situ* ^{14}C thus allows us to reevaluate the ice-thickness history at the site: the ice must have been thick enough to cover at least the highest-elevation unsaturated sample, and had to have been that thick within the time it would have taken for the ^{14}C concentration of that sample to reach saturation again.

1.1 Study Area

The Grove Mountains are well located to assess how far inland the EAIS was thicker at the LGM than it is at present and whether previously measured concentrations of ^{10}Be and ^{26}Al from this site likely reflect a component of nuclides inherited from a previous period of exposure. These isolated nunataks are located ~200 km upstream of the main trunk of Lambert Glacier and ~400 km inland/south of the Antarctic coast (Fig. 1) and are the most interior ice-free area in this region. The summits of the nunataks rise 100-200 m above the modern ice surface (~1800 m a.s.l.), providing the potential to record past EAIS-thickness changes. Ice flows slowly (<5 m yr⁻¹) to the west-northwest between these nunataks (Rignot *et al.*, 2011).

At Nunatak 1921 (named for its summit elevation; local ice-surface elevation \approx 1825 m a.s.l.) evidence of past ice cover is apparent from the occurrence of felsic cobbles atop very weathered orthogneiss bedrock (Lilly, 2008). Given the sparsity of outcrops and non-channelized nature of ice flow in the interior EAIS, we are not able to identify the provenance of these cobbles beyond stating that they are not derived from Nunatak 1921 (i.e. they are erratics).

2 Methods

Here we reanalyze samples first presented in Lilly *et al.* (2010), which were collected from the Grove Mountains for ^{10}Be and ^{26}Al analysis as part of a study of the long-term glacial history of the region. Measurements of ^{10}Be and ^{26}Al were carried out in 2004 at the ANTARES Accelerator Mass Spectrometry facility. Nuclide

concentrations below saturation were recorded for all samples, indicating 40-700 kyr of exposure since the bedrock was last reset. For full details, see Lilly (2008) and Lilly et al. (2010).

The samples were collected in an elevation transect from the present-day ice surface on the upstream face of Nunatak 1921 in 2003/4 and 2004/5 (Lilly *et al.*, 2010; **Table 1**). Pairs of bedrock and erratic samples showed no evidence of post-depositional movement, cover by sediments, or subaerial erosion. Samples were preferentially collected from ridgetops to minimize the chances of shielding by snow. As neither plucking scars nor glacial striae were observed at the site (Lilly *et al.*, 2010), indicating low or negligible rates of subglacial erosion, and because of the scatter observed in the ^{10}Be and ^{26}Al data from this site, we anticipate that the existing nuclide concentrations do not accurately record LGM ice thickness.

To provide a test of LGM ice thickness, we carried out *in situ* ^{14}C analysis on the ten of these samples from which available material remained. These samples form a transect covering 96 m of elevation (1825-1921 m a.s.l.). Seven of the samples were erratic cobbles. The remaining three (marked in **Table 1** with “(BR)” appended to the Sample ID) were bedrock samples.

Table 1: Sample locations. Seven samples were erratic cobbles. Three samples were bedrock (BR). A density of 2.7 g cm^{-3} is assumed for all samples.

Sample ID	Elevation (m a.s.l.)	Elevation above modern ice surface (m)	Latitude (degrees S)	Longitude (degrees E)	Thickness (cm)	Topographic shielding	Lithology
GR01	1832	7	72.9115	74.9096	2	0.985	Felsic metamorphic
GR03	1854	29	72.9115	74.9079	2	0.992	Quartzite
GR04	1870	45	72.9110	74.9067	2	1.000	Quartzite
GR06	1894	69	72.9099	74.9044	2	0.998	Fine-grained felsic
GR07	1921	96	72.9088	74.9045	2	1.000	Quartzite
GR12(BR)	1825	0	72.9112	74.9097	2	0.985	Orthogneiss
GR13	1839	14	72.9115	74.9094	2	0.993	Unrecorded*
GR15(BR)	1847	22	72.9115	74.9088	3	0.993	Orthogneiss
GR18	1873	48	72.9108	74.9061	4	1.000	Vein quartz
GR21(BR)	1912	87	72.9088	74.9045	3	0.999	Orthogneiss

*Sample GR13 was crushed prior to the beginning of this study, and its lithology was not recorded prior to crushing.

Quartz was isolated through physical and chemical processing at the Tulane University Cosmogenic Nuclides Laboratory (TUCNL; Goehring *et al.*, 2019). Whole samples were crushed and milled, then all samples were sieved to select their 125-500-micron fractions. Sieved samples were then rinsed with tap water to remove clay-sized grains. A roller-type magnetic separator was then used to remove magnetic minerals. Froth flotation was used to separate quartz and feldspar grains, followed by etching for at least two days in 5% HF/HNO₃ on a shaker table and at least two days in a sonicator in 1% HF/HNO₃ in order to remove adsorbed carbon species (Nichols and Goehring, 2019).

Following the isolation and purification, 0.6-5 g aliquots were separated from the cleaned quartz for ^{14}C extraction. Before extraction, each aliquot was sonicated in 50% HNO₃ for 0.5 hr, then rinsed with Type I water and dried

overnight in a vacuum oven. The dried quartz was then loaded into a Pt combustion boat containing LiBO₂ flux which had been degassed, fused, and cooled. This boat containing flux and sample was then step heated in O₂ for 0.5 hr at 500 °C for cleaning. The quartz is then heated for a further 3 hr at 1,100 °C in the Tulane University Carbon Extraction and Graphitization System to extract gases for measurement. Carbon species released were oxidized to CO₂ over 0.64-0.86 mm graded, crushed quartz chips at 850 °C, then cryogenically purified, collected, and diluted to ~110 µg with ¹⁴C-free CO₂ (Goehring *et al.*, 2019). An aliquot of this gas was separated for δ¹³C analysis and the remainder graphitized via Fe-catalyzed H₂ reduction. For further details, see Goehring *et al.* (2019) and Nichols and Goehring (2019). Carbon-isotope ratios were then measured at the National Ocean Sciences Accelerator Mass Spectrometry facility at the Woods Hole Oceanographic Institution (see Table A2), and data reduction followed Hippe and Lifton (2014).

A blank value of 43,661 ± 11,279 atoms was subtracted from the total measured atoms from each sample; this value represents the mean and standard deviation of process blanks run at the TUCNL (Goehring *et al.*, 2019) through the timespan within which samples for this study were measured (July 10, 2021-August 27, 2021). This blank-corrected measurement was divided by the sample mass to determine the ¹⁴C concentration of each sample. Exposure ages were calculated using the “LSDn” nuclide-specific production rate scaling scheme of Lifton *et al.* (2014). The production rate of *in-situ* ¹⁴C was calibrated using the CRONUS-A interlaboratory comparison material (Goehring *et al.*, 2019). The CRONUS-A material is assumed to be saturated with *in-situ* ¹⁴C based on geological observations indicating that its collection site has not been covered in the last 11.3 Myr (Goehring *et al.*, 2019; Nichols *et al.*, 2019). Repeated measurements of CRONUS-A material at the TUCNL show ~6% variation in ¹⁴C concentrations (Goehring, *et al.*, 2019a); thus, we use a minimum uncertainty equal to 6% of the calculated ¹⁴C concentration of our samples for exposure-age calculation (Table 2).

Table 2: Sample ¹⁴C concentrations and exposure ages. All measurements of ¹⁴C atoms per sample corrected by subtracting a 0.44 ± 0.11 × 10⁵ atom blank prior to concentration calculation. Where the 1 σ [¹⁴C] uncertainty and 6% [¹⁴C] differ, the larger uncertainty value is reported. “Internal” ¹⁴C-age uncertainties include only instrumental uncertainty. “External” ¹⁴C-age uncertainties include both instrumental and production-rate uncertainties. All ¹⁴C-age uncertainties are here presented at 1 σ. For further sample details, see Table S1. For details of the dataset used to calibrate production rates at our site, see Table S3.

Sample number	[¹⁴ C] (10 ⁵ atoms g ⁻¹)	¹⁴ C Age (ka)	Internal ¹⁴ C-age uncertainty (ka)	External ¹⁴ C-age uncertainty (ka)
GR01	1.10 ± 0.20	1.480	0.093	0.190
GR03	2.82 ± 0.17	4.227	0.100	0.573
GR04	5.43 ± 0.33	11.895	0.365	2.742
GR06	6.23 ± 0.37	16.289	0.694	5.262
GR07	8.18 ± 0.49	Saturated	N/A	N/A
GR12(BR)	1.20 ± 0.07	1.417	0.034	0.162
GR13	1.62 ± 0.10	2.142	0.038	0.253
GR15(BR)	0.18 ± 0.01	0.183	0.013	0.023
GR18	5.50 ± 0.33	12.466	0.399	2.999
GR21(BR)	7.81 ± 0.47	Saturated	N/A	N/A

3 Results

Our samples have ^{14}C concentrations between $1.8 \pm 2.4 \times 10^3$ atoms g^{-1} (GR15) and $8.18 \pm 49.1 \times 10^3$ atoms g^{-1} (GR07, Table 2). The sample with the lowest concentration has an exposure age of 0.18 ± 0.02 ka, and the samples with the highest concentrations are saturated (Table 2). These exposure ages are 38 ± 9.6 (GR01) to 295 ± 27.1 kyr (GR03) less than ^{10}Be and ^{26}Al exposure ages from each sample (for full sample-measurement details, see Tables S1-6). Samples form a thinning transect with concentrations and ages mostly increasing monotonically with elevation (Fig. 2). There are however two exceptions, both low-elevation bedrock samples (GR15 and GR12). We suspect that the site of GR15 may have been covered by snow or other sediment, though we have not acquired any field evidence to this effect. GR12 may instead not be an outlier, as its concentration lies within the uncertainty window of that of GR01.




Figure 2: Sample nuclide concentrations and exposure ages. The current ice surface at this site is approximately coincident with the elevation of sample GR12. (a) Locations of samples noted on a photograph of the south face of Nunatak 1921. IDs of samples from which ^{14}C was measured in this study are highlighted, and sample locations marked with red circles. Black circles represent the locations of samples from which ^{14}C was not measured, as no sample material was available. This image modified from Lilly *et al.* (2010). (b) ^{14}C concentrations plotted against elevation. Isochrons (dotted lines) show corresponding exposure ages at each elevation. Tilted vertical gray band to right represents the saturation error envelope as calculated using the online exposure-age calculator formerly known as the CRONUS-Earth online exposure-age calculator using the CRONUS-A measurements listed in Table S3. Samples GR07 and GR21 are saturated, indicating >25 kyr of exposure and implying the summit of the nunatak was exposed during the LGM. Horizontal red band indicates the range of possible LGM ice-surface elevations limited by the elevations of GR06, the highest-elevation unsaturated sample, and GR21, the lowest-elevation saturated sample. We consider samples GR12 and GR15 outliers, as the trend of

decreasing elevation with decreasing age recorded by all of the erratic samples places these two samples out of order. (c) Sample exposure ages plotted against elevation, calculated from concentrations of ^{14}C (this work) and ^{26}Al and ^{10}Be (Lilly *et al.*, 2010). Note the younger exposure ages calculated from ^{14}C . (d) As plot (c), but only showing ^{14}C exposure ages for the last 22 ka. Light blue bar indicates the timing of meltwater pulse 1a (Deschamps *et al.*, 2012). Samples GR07 and GR21 are saturated with ^{14}C and thus omitted from this plot. See the Supplementary tables for all sample information, nuclide concentrations and calculated exposure ages.

If the ice-sheet thickness was similar to or thinner than at present in the vicinity of the Grove Mountains at the LGM, all of our samples would be saturated with ^{14}C . However, our samples show a clear trend of increasing ^{14}C concentrations with elevation (Fig. 2). Only two samples (GR07 and GR21, Table 2) show clear evidence of saturation, both near the summit of the nunatak. These results thus show that ice was thicker at the LGM than at present in the Grove Mountains but not sufficiently thick as to override the summit for a significant duration, though saturation precludes neither a short period (<3 kyr) of cover nor cover by thin (<10 m) ice (Fig. 3).




Figure 3: Burial-history contour plot for a sample at 1,912 m above sea level (a.s.l.) in the Grove Mountains. Modeled glacial histories start at 50 ka with one episode of burial under >10 m of ice. No ^{14}C is produced while buried. ^{14}C concentrations in the sample are saturated at model start. The black-shaded part of the graph shows impossible histories (i.e., histories that require future burial). The grey-shaded part of the graph shows histories that would result in sample GR21 having a concentration below saturation for ^{14}C . The unshaded portion of the graph shows the uncertainty window of a saturated sample at this latitude and elevation (7.3×10^5 atoms g^{-1} ; 72.9088°S ; 1,912 m a.s.l.). Only the lower-concentration end of the saturation window is consistent with any significant degree of burial under enough ice to effectively stop production (~ 10 m); thus, only samples that were buried a long time ago or for a very short duration could show

concentrations approaching saturation. Sample GR21 plots off the bottom-left corner of this figure; its mean ^{14}C concentration (7.81×10^5 atoms g^{-1} , see Table 1) is thus inconsistent with any episode of burial longer than 3 kyr in the last 30 kyr, indicating that, if there was any significant duration of cover experienced by these samples, it occurred predominantly prior to the LGM. Permittable episodes of cover become shorter and occur earlier if samples are not assumed to be saturated at 50 ka.

The LGM ice surface must have been between the lowest of our saturated and highest of our unsaturated samples, corresponding to an elevation between 1894 and 1912 m a.s.l. This equates to ice 63-87 m thicker at the LGM than at present, with subsequent thinning.

Additionally, exposure ages calculated from ^{14}C concentrations allow us to infer a simple thinning history at Nunatak 1921. The highest-elevation unsaturated sample (GR06) provides a minimum post-LGM age for the onset of thinning at the site of 16.3 ± 5.3 ka (Fig. 2, Table 2). Up to 18 m (21-29%) of thinning could have occurred before and up to 21 m (24-33%) during meltwater pulse 1a (MWP-1a; Fig. 2d), ~13.5-14.7 ka, assuming the mean exposure ages of GR04 and GR18 and a linear thinning history. The potential for glacial overshoot, whereby the glacier thins beyond its new equilibrium thickness and subsequently rethickens, however, makes these minimum estimates. Note, too, the large uncertainties on the pre-Holocene ages presented here relative to the duration of MWP-1a; more-stringent age control is necessary to state confidently the relative timing of ice loss here. Most post-LGM thinning (55-70%) is recorded during the Holocene (the last 11.7 ka; Walker *et al.*, 2009), as opposed to the earlier stage of deglaciation, when most Antarctic ice loss is modelled (Pittard *et al.* 2020). Based on our lowest-elevation sample (GR12), which was collected less than 1 m above the current ice surface (~1825 m a.s.l.), the present-day ice thickness was reached at 1.4 ± 0.2 ka (Tables 1 & 2).

4 Discussion

New exposure ages calculated from *in situ* ^{14}C concentrations allow us to revise the history of the EAIS at this site. The combination of saturated and unsaturated samples on Nunatak 1921 shows that its summit was exposed during the LGM, yet the ice sheet was modestly thicker (up to 87 m) here at the LGM than at present, contrary to interpretations of previous ^{10}Be and ^{26}Al data at this site and reconstructions of the interior EAIS at the LGM (e.g. Lilly *et al.*, 2010; Buizert *et al.*, 2021).

Our samples were saturated with neither ^{10}Be nor ^{26}Al , but show evidence of long, complex exposure histories (Lilly *et al.*, 2010; Table S5). The high contribution of inherited nuclides from pre-LGM exposure prevents an accurate test of the LGM ice thickness and reconstruction of the post-LGM thinning history. Exposure long enough to build these ^{10}Be and ^{26}Al concentrations up would also have left these samples saturated with ^{14}C . Our ^{14}C data indicate that ice cover > ca. 10 m thick occurred at this site up to at least 70 m above the present ice surface, and the period of cover was long enough to allow ^{14}C concentrations in our samples to decay to near-background levels, given the low concentrations of the lowest-elevation samples (Table 2). The summit of the nunatak was either not covered or only covered briefly (≤ 1 kyr) or by thin (≤ 10 m) ice enough for the two summit samples to become re-saturated with ^{14}C following re-exposure (Fig. 3). Following the LGM, the nunataks were progressively re-exposed through the Late Holocene.

Direct constraints from cosmogenic ^{10}Be and ^{26}Al show evidence of the ice being thicker near the Antarctic coast at the LGM than at present (e.g., Mackintosh *et al.*, 2007; White *et al.*, 2011a), but exposure ages derived from the same nuclides from interior sites such as the Grove Mountains pre-date the LGM (Lilly *et al.*, 2010). While we cannot rule out the thicker-than-present ice at the Grove Mountains being an entirely localized phenomenon, we suggest based on the application of ^{14}C in this study and other Antarctic studies (e.g., White *et al.*, 2011b; Fogwill *et al.*, 2014; Nichols *et al.*, 2019; Hillebrand *et al.*, 2021) that at least some previous reconstructions of LGM ice thickness based on longer-lived nuclides (e.g. ^{10}Be and ^{26}Al) which show either thickening or no thinning since the LGM away from the coast and fastest-flowing parts of East Antarctica may be inaccurate. The potential for low levels of Al and Be inheritance in cold, arid regions – such as on the edge of the ice-sheet plateau but possibly also in coastal areas of Antarctica - highlights the usefulness of ^{14}C as a tool for improving ice histories derived from long-lived nuclides.

Our new chronology indicates that the zone of thicker-than-present LGM ice extended further inland than was previously thought (Mackintosh *et al.*, 2014). Cosmogenic dating and geomorphic evidence from elsewhere in the Lambert Glacier catchment support a low-angle ice stream surface at the LGM, with ice 160 m thicker at the most upstream site in the PCMs (Mt. Ruker), and at least 250 m and up to 800 m thicker at sites closer to the coast (White *et al.*, 2011a; **Fig. 1**). The “hinge zone” between interior and coastal change, where the LGM ice thickness was the same as today, was proposed to be at ~1900-2000 m a.s.l. based on the available evidence at the PCMs and Grove Mountains (Mackintosh *et al.*, 2014). A thicker-than-present EAIS at the Grove Mountains during the LGM therefore indicates that this “hinge zone” lies further inland, increasing the amount of LGM ice volume across much of the ice sheet (**Fig. 4**).




Figure 4: Implications of new LGM ice thickness constraints on the East Antarctic “hinge zone”. Modified from Fig. 1c, the diagram shows hypothetical vertically exaggerated cross-sections of the East Antarctic Ice Sheet at the present day (dashed line), and at the LGM based on previous evidence (dotted line) and accounting for our data (dot-dashed line). Our results indicate that ice at the Grove Mountains (near the approximate elevation previously considered the “hinge zone”) was ~70 m thicker than it is today. Assuming that LGM ice-thickness estimates near the coast are accurate, this necessitates a steeper coastal

ice-surface slope to accommodate the increased thickness at the Grove Mountains (and a shallower East Antarctic plateau ice-surface slope if the LGM ice-thickness estimates in the interior are accurate), moving the “hinge zone” further into the interior. The exact gradients of these slopes and location of the “hinge zone” control the volume of ice lost from the East Antarctic Ice Sheet since the LGM. **Note that the distances and slopes displayed in this figure are not to scale, and only schematically illustrate the relative elevations and changes in slope. Ice-surface profiles are here depicted as straight lines to aid visibility. The true profiles would curve, as in Fig. 1c.**

Ice in East Antarctica being thicker at the LGM than at present only within a few hundred kilometers of the coastline would be consistent with reconstructions of MWP-1a that call for only a limited input of meltwater from Antarctica (e.g., Yeung *et al.*, 2019). Our work shows that EAIS thickening extended further inland than indicated by ^{10}Be and ^{26}Al ages (e.g., Lilly *et al.*, 2010), providing a modest additional ice volume for MWP-1a, and that thinning started before and possibly occurred during the period of MWP-1a. We cannot accurately quantify how much EAIS volume was lost during this period, due to the uncertainties of our calculated exposure ages. Our data indicate that likely less than half of the post-LGM ice loss occurred before or during MWP-1a in this region, consistent with studies identifying Antarctica as likely being a minor contributor and the majority of the Antarctic contribution to have been sourced from West Antarctica (e.g., Lin *et al.*, 2021).

An implication of this interior portion of the EAIS being thicker than **previously suggested** at the LGM is that the ice subsequently thinned, allowing us to evaluate deglacial leads and lags between the coast and interior. The earliest deglaciation constraints in this region come from ice-sheet thinning in the PCMs at 18 ka (White *et al.*, 2011a; Bentley *et al.*, 2014), which was possibly coincident with grounding-line retreat on the continental shelf in Prydz Bay (Mackintosh *et al.*, 2014). Ice-shelf retreat began by ~16 ka and ~14 ka in west-central and eastern Prydz Bay, respectively, with the Rauer Group and Vestfold Hills ice-free by ~11 ka (White *et al.*, 2022). Our record **suggests that** ice in the Grove Mountains **began thinning** ~16 ka, ~2 kyr earlier **than** in the PCMs, though the timing at the Grove Mountains is broadly consistent with available evidence of deglaciation at the coast. The modern ice-surface elevation was reached by 9-12 ka at the PCMs (White *et al.*, 2011a) but 1.4 ka in the Grove Mountains, ~8-11 ka later. **Cosmogenic-exposure ages reported here from other studies were recalculated using the online exposure age calculator formerly known as the CRONUS-Earth online exposure age calculator [hess.ess.washington.edu], as with our ^{14}C ages, using the primary ^{10}Be calibration dataset of Borchers *et al.*, 2016.** Part of this discrepancy could be due to inheritance in the samples from the PCMs. Further work to measure *in-situ* ^{14}C in samples from the PCMs would enable an evaluation of the degree of lead and lag between sites.

Deglaciation thus possibly started and likely finished earlier downstream, and the magnitude of thinning was greater at the Antarctic coastline than in its interior. Ice-sheet modeling indicates that responses to sea-level rise, decreased accumulation, and changes in temperature should manifest first at the margins of the ice sheet, causing thinning to propagate into the interior of the ice sheet (Alley and Whillans, 1984; Spector *et al.*, 2019). Such propagation is likely slowed and attenuated by distance and travel over bedrock highs (Johnson *et al.*, 2021), such as the Grove Mountains. Modern observations confirm that such dynamic thinning occurs over decadal timescales (e.g., Felikson *et al.*, 2017), but our data indicate that such processes may continue over centuries to millennia.

If the Grove Mountains are representative of the behavior of similar locations in interior East Antarctica, more of the EAIS may have been thicker-than-present at the LGM and subsequently **ly** thinned **more** than was previously

thought. Ice-sheet models may thus currently underestimate LGM ice volume and rates and magnitudes of deglacial ice loss.

5 Conclusions

Our new *in situ* ^{14}C results provide improved constraints on past East Antarctic Ice Sheet thickness at a site ~400 km inland from the present-day coast. These data show that the ice sheet at the Grove Mountains was thicker than at present at the LGM, but the summits of these nunataks were exposed. Ice-sheet thinning began here ~15 ka and continued through the Holocene, likely in response to changes near the grounding line that propagated upstream. This work demonstrates that the “hinge zone” separating the interior ice (which was thinner at the LGM than it is today) from the ice nearer the coast (which was thicker at the LGM than it is today) was further inland than was previously thought. The additional ice volume implied by these findings therefore needs to be accounted for in numerical ice sheet and glacial isostatic adjustment reconstructions of the last deglaciation.

Code availability

Data availability

All data described in the paper are included in Tables S1-6.

Interactive computing environment

Sample availability

Video supplement

Author contributions

CR processed samples for ^{14}C analysis, wrote the paper, and prepared all figures. CR, RJ, and AM conceived the project. All authors read and commented on the manuscript. BG provided code for exposure-age calculation and plotting. KL undertook fieldwork in the Grove Mountains and collected all the field observations and samples presented here.

Competing interests

The authors declare that they have no conflict of interest.

Disclaimer

Acknowledgements

This work was supported by Australian Research Council grants DE210101923, awarded to RSJ, and ARC Special Research Initiative ‘Securing Antarctica’s Environmental Future’ (SR200100005). CR would also like to acknowledge support from the Monash Graduate and International Tuition Scholarships.

References

- Alley, R. B. and Whillans, I. M.: Response of the East Antarctica ice sheet to sea-level rise, *Journal of Geophysical Research: Oceans*, 89, 6487-6493, 1984.
- Andersen, J. L., Newall, J. C., Fredin, O., Glasser, N. F., Lifton, N. A., Stuart, F. M., Fabel, D., Caffee, M., Pedersen, V. K., and Koester, A. J.: A topographic hinge-zone divides coastal and inland ice dynamic regimes in East Antarctica, *Communications Earth & Environment*, 4, 9, 2023.
- Balco, G., Stone, J. O., Sliwinski, M. G., and Todd, C.: Features of the glacial history of the Transantarctic Mountains inferred from cosmogenic ^{26}Al , ^{10}Be and ^{21}Ne concentrations in bedrock surfaces, *Antarctic Science*, 26, 708-723, 2014.
- Bentley, M. J., Cofaigh, C. O., Anderson, J. B., Conway, H., Davies, B., Graham, A. G., Hillenbrand, C.-D., Hodgson, D. A., Jamieson, S. S., and Larter, R. D.: A community-based geological reconstruction of Antarctic Ice Sheet deglaciation since the Last Glacial Maximum, *Quaternary Science Reviews*, 100, 1-9, 2014.
- Bentley, M. J., Fogwill, C. J., Kubik, P. W., and Sugden, D. E.: Geomorphological evidence and cosmogenic $^{10}\text{Be}/^{26}\text{Al}$ exposure ages for the Last Glacial Maximum and deglaciation of the Antarctic Peninsula Ice Sheet, *Geological Society of America Bulletin*, 118, 1149-1159, 2006.
- Bindschadler, R., H. Choi, and ASAID Collaborators. 2011. High-resolution Image-derived Grounding and Hydrostatic Lines for the Antarctic Ice Sheet. Boulder, Colorado, USA: National Snow and Ice Data Center. Digital media.
- Bockheim, J. G., Wilson, S. C., Denton, G. H., Andersen, B. G., and Stuiver, M.: Late quaternary ice-surface fluctuations of Hatherton Glacier, Transantarctic Mountains, *Quaternary Research*, 31, 229-254, 1989.
- Borchers, B., Marrero, S., Balco, G., Caffee, M., Goehring, B., Lifton, N., Nishiizumi, K., Phillips, F., Schaefer, J., and Stone, J.: Geological calibration of spallation production rates in the CRONUS-Earth project, *Quaternary Geochronology*, 31, 188-198, 2016.
- Buizert, C., Fudge, T., Roberts, W. H., Steig, E. J., Sherriff-Tadano, S., Ritz, C., Lefebvre, E., Edwards, J., Kawamura, K., and Oyabu, I.: Antarctic surface temperature and elevation during the Last Glacial Maximum, *Science*, 372, 1097-1101, 2021.
- Burton-Johnson, A., Black, M., Fretwell, P. T., and Kaluza-Gilbert, J.: An automated methodology for differentiating rock from snow, clouds and sea in Antarctica from Landsat 8 imagery: a new rock outcrop map and area estimation for the entire Antarctic continent, *The Cryosphere*, 10, 1665-1677, 2016.
- Coulon, V., Bulthuis, K., Whitehouse, P. L., Sun, S., Haubner, K., Zipf, L., and Pattyn, F.: Contrasting response of West and East Antarctic ice sheets to glacial isostatic adjustment, *Journal of Geophysical Research: Earth Surface*, 126, e2020JF006003, 2021.
- DeConto, R. M., Pollard, D., Alley, R. B., Velicogna, I., Gasson, E., Gomez, N., Sadai, S., Condron, A., Gilford, D. M., and Ashe, E. L.: The Paris Climate Agreement and future sea-level rise from Antarctica, *Nature*, 593, 83-89, 2021.

Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A. L., Henderson, G. M., Okuno, J. i., and Yokoyama, Y.: Ice-sheet collapse and sea-level rise at the Bølling warming 14,600 years ago, *Nature*, 483, 559-564, 2012.

Dunai, T. J.: *Cosmogenic nuclides: principles, concepts and applications in the earth surface sciences*, Cambridge University Press, 2010.

Felikson, D., Bartholomäus, T. C., Catania, G. A., Korsgaard, N. J., Kjær, K. H., Morlighem, M., Noël, B., Van Den Broeke, M., Stearns, L. A., and Shroyer, E. L.: Inland thinning on the Greenland ice sheet controlled by outlet glacier geometry, *Nature Geoscience*, 10, 366-369, 2017.

Fogwill, C., Turney, C., Golledge, N., Rood, D., Hippe, K., Wacker, L., Wieler, R., Rainsley, E., and Jones, R.: Drivers of abrupt Holocene shifts in West Antarctic ice stream direction determined from combined ice sheet modelling and geologic signatures, *Antarctic Science*, 26, 674-686, 2014.

Goehring, B. M., Balco, G., Todd, C., Moening-Swanson, I., and Nichols, K.: Late-glacial grounding line retreat in the northern Ross Sea, *Antarctica, Geology*, 47, 291-294, 2019a.

Goehring, B. M., Wilson, J., and Nichols, K.: A fully automated system for the extraction of in situ cosmogenic carbon-14 in the Tulane University cosmogenic nuclide laboratory, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 455, 284-292, 2019b.

Hanna, E., Pattyn, F., Navarro, F., Favier, V., Goelzer, H., van den Broeke, M. R., Vizcaino, M., Whitehouse, P. L., Ritz, C., and Bulthuis, K.: Mass balance of the ice sheets and glaciers—Progress since AR5 and challenges, *Earth-Science Reviews*, 201, 102976, 2020.

Hillebrand, T. R., Stone, J. O., Koutnik, M., King, C., Conway, H., Hall, B., Nichols, K., Goehring, B., and Gillespie, M. K.: Holocene thinning of Darwin and Hatherton glaciers, Antarctica, and implications for grounding-line retreat in the Ross Sea, *The Cryosphere*, 15, 3329-3354, 2021.

Hippe, K. and Lifton, N. A.: Calculating isotope ratios and nuclide concentrations for in situ cosmogenic ¹⁴C analyses, *Radiocarbon*, 56, 1167-1174, 2014.

Johnson, J. S., Pollard, D., Whitehouse, P. L., Roberts, S. J., Rood, D. H., and Schaefer, J. M.: Comparing glacial-geological evidence and model simulations of ice sheet change since the last glacial period in the Amundsen Sea sector of Antarctica, *Journal of Geophysical Research: Earth Surface*, 126, e2020JF005827, 2021.

Jones, R. S., Johnson, J. S., Lin, Y., Mackintosh, A. N., Sefton, J. P., Smith, J. A., Thomas, E. R., and Whitehouse, P. L.: Stability of the Antarctic Ice Sheet during the pre-industrial Holocene, *Nature Reviews Earth & Environment*, 3, 500-515, 2022.

Korschinek, G., Bergmaier, A., Faestermann, T., Gerstmann, U., Knie, K., Rugel, G., Wallner, A., Dillmann, I., Dollinger, G., and Von Gostomski, C. L.: A new value for the half-life of ¹⁰Be by heavy-ion elastic recoil detection and liquid scintillation counting, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 268, 187-191, 2010.

Lifton, N., Sato, T., and Dunai, T. J.: Scaling in situ cosmogenic nuclide production rates using analytical approximations to atmospheric cosmic-ray fluxes, *Earth and Planetary Science Letters*, 386, 149-160, 2014.

Lilly, K.: Three million years of East Antarctic ice sheet history from in situ cosmogenic nuclides in the Lambert-Amery Basin, 2008. 2008.

Lilly, K., Fink, D., Fabel, D., and Lambeck, K.: Pleistocene dynamics of the interior East Antarctic ice sheet, *Geology*, 38, 703-706, 2010.

Lin, Y., Hibbert, F. D., Whitehouse, P. L., Woodroffe, S. A., Purcell, A., Shennan, I., and Bradley, S. L.: A reconciled solution of Meltwater Pulse 1A sources using sea-level fingerprinting, *Nature communications*, 12, 2015, 2021.

Liu, H., K. C. Jezek, B. Li, and Z. Zhao. 2015. Radarsat Antarctic Mapping Project Digital Elevation Model, Version 2. [Indicate subset used]. Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center. doi: <http://dx.doi.org/10.5067/8JKNEW6BFRVD>. [Accessed 1/08/2023]

Mackintosh, A., White, D., Fink, D., Gore, D. B., Pickard, J., and Fanning, P. C.: Exposure ages from mountain dipsticks in Mac. Robertson Land, East Antarctica, indicate little change in ice-sheet thickness since the Last Glacial Maximum, *Geology*, 35, 551-554, 2007.

Mackintosh, A. N., Verleyen, E., O'Brien, P. E., White, D. A., Jones, R. S., McKay, R., Dunbar, R., Gore, D. B., Fink, D., and Post, A. L.: Retreat history of the East Antarctic Ice Sheet since the last glacial maximum, *Quaternary Science Reviews*, 100, 10-30, 2014.

Matsuoka, K., Skoglund, A., Roth, G., de Pomereu, J., Griffiths, H., Headland, R., ... Melvær, Y. (2018). Quantarctica [Data set]. Norwegian Polar Institute. <https://doi.org/10.21334/npolar.2018.8516e961>

Miles, B. W., Stokes, C. R., Jenkins, A., Jordan, J. R., Jamieson, S. S., and Gudmundsson, G. H.: Slowdown of Shirase Glacier, East Antarctica, caused by strengthening alongshore winds, *The Cryosphere*, 17, 445-456, 2023.

Nichols, K. A. and Goehring, B. M.: Isolation of quartz for cosmogenic in situ ^{14}C analysis, *Geochronology*, 1, 43-52, 2019.

Nichols, K. A., Goehring, B. M., Balco, G., Johnson, J. S., Hein, A. S., and Todd, C.: New last glacial maximum ice thickness constraints for the Weddell Sea Embayment, Antarctica, *The Cryosphere*, 13, 2935-2951, 2019.

Nichols, K. A.: A decade of in situ cosmogenic ^{14}C in Antarctica, *Annals of Glaciology*, 2022. 1-6, 2022.

Nishiizumi, K.: Preparation of new ^{10}Be and ^{26}Al AMS standard reference materials, *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 530, 43-47, 2022.

Pittard, M. L., Whitehouse, P. L., Bentley, M. J., and Small, D.: An ensemble of Antarctic deglacial simulations constrained by geological observations, *Quaternary Science Reviews*, 298, 107800, 2022.

Rignot, E., Mouginot, J., and Scheuchl, B.: Ice flow of the Antarctic ice sheet, *Science*, 333, 1427-1430, 2011.

Rignot, E., Mouginot, J., Scheuchl, B., Van Den Broeke, M., Van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proceedings of the National Academy of Sciences*, 116, 1095-1103, 2019.

Spector, P., Stone, J., and Goehring, B.: Thickness of the divide and flank of the West Antarctic Ice Sheet through the last deglaciation, *The Cryosphere*, 13, 3061-3075, 2019.

Stokes, C. R., Abram, N. J., Bentley, M. J., Edwards, T. L., England, M. H., Foppert, A., Jamieson, S. S., Jones, R. S., King, M. A., and Lenaerts, J.: Response of the East Antarctic Ice Sheet to past and future climate change, *Nature*, 608, 275-286, 2022.

Walker, M., Johnsen, S., Rasmussen, S. O., Popp, T., Steffensen, J. P., Gibbard, P., Hoek, W., Lowe, J., Andrews, J., and Björck, S.: Formal definition and dating of the GSSP (Global Stratotype Section and Point) for the base of the Holocene using the Greenland NGRIP ice core, and selected auxiliary records, *Journal of Quaternary Science: Published for the Quaternary Research Association*, 24, 3-17, 2009.

471 White, D., Fülöp, R.-H., Bishop, P., Mackintosh, A., and Cook, G.: Can in-situ cosmogenic ^{14}C be used to assess
 472 the influence of clast recycling on exposure dating of ice retreat in Antarctica?, *Quaternary Geochronology*, 6,
 473 289-294, 2011.

474 White, D. A., Fink, D., and Gore, D. B.: Cosmogenic nuclide evidence for enhanced sensitivity of an East
 475 Antarctic ice stream to change during the last deglaciation, *Geology*, 39, 23-26, 2011.

476 White, D. A., Fink, D., Lilly, K., O'Brien, P., Dorschel, B., Berg, S., Bennike, O., Gore, D. B., Fabel, D., and
 477 Blaxell, M.: Rapid ice sheet response to deglacial and Holocene paleoenvironmental changes in eastern Prydz
 478 Bay, East Antarctica, *Quaternary Science Reviews*, 280, 107401, 2022.

479 Yamane, M., Yokoyama, Y., Miura, H., Maemoku, H., Iwasaki, S., and Matsuzaki, H.: The last deglacial history
 480 of Lützow-Holm Bay, East Antarctica, *Journal of Quaternary Science*, 26, 3-6, 2011.

481 Yeung, N., Menviel, L., Meissner, K., and Sikes, E.: Assessing the Spatial Origin of Meltwater Pulse 1A Using
 482 Oxygen-Isotope Fingerprinting, *Paleoceanography and Paleoclimatology*, 34, 2031-2046, 2019.

483 Zwally, H. Jay, Mario B. Giovinetto, Matthew A. Beckley, and Jack L. Saba, 2012, Antarctic and Greenland
 484 Drainage Systems, GSFC Cryospheric Sciences Laboratory, at
 485 http://icesat4.gsfc.nasa.gov/cryo_data/ant_grn_drainage_systems.php