



## “Blooming” of litter-mixing effects: The role of flower and leaf litter interactions on decomposition in terrestrial and aquatic ecosystems

5 Mery Ingrid Guimarães de Alencar<sup>1,2</sup>, Rafael D. Guariento<sup>3</sup>, Bertrand Guenet<sup>2</sup>, Luciana S. Carneiro<sup>1</sup>,  
Eduardo L. Voigt<sup>4</sup> & Adriano Caliman<sup>1</sup>

<sup>1</sup> Departamento de Ecologia, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Natal, 59078-900, Brazil.

<sup>2</sup> Laboratoire de Géologie, Ecole normale supérieure, CNRS, IPSL, Université PSL, Paris, 75005, France

10 <sup>3</sup> Universidade Federal do Mato Grosso do Sul, CCBS, Campo Grande, 79070-900, Brazil.

<sup>4</sup> Departamento de Biologia Celular e Genética, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Natal, 59078-900, Brazil.

Correspondence to: Mery Ingrid Guimarães de Alencar (alencarmery@gmail.com)

15 **Abstract.** The diversity effect on decomposition, through the litter-mixing effects, plays a central role in determining the nutrient and carbon dynamics in ecosystems. However, the litter-mixing effects are centered on a leaf litter perspective. Important aspects related to intraspecific interaction and biomass concentration are rarely evaluated, even though they could be essential to determine the litter decomposition dynamics. Here, we introduced a new perspective to evaluate whether and how the interaction between flower and leaf litter affects the occurrence, direction, and magnitude of litter-mixing effects in terrestrial and aquatic ecosystems.

20 We performed laboratory experiments using flower and leaf litter from the yellow trumpet tree *Tabebuia aurea* (Silva Manso) Benth. & Hook. f. ex. S. Moore as a model. To obtain realistic results and a mechanistic understanding, we used different scenarios of flower:leaf litter biomass proportion and measured 13 functional traits, respectively. Litter-mixing effects were consistent in both aquatic and terrestrial environments, with faster decomposition of both litter types in mixtures compared to their monocultures (synergistic effects). Litter-mixing effects were stronger in the terrestrial environment and higher flower:leaf litter biomass proportion. Our results indicate that synergistic results are mainly associated with complementary effects. Flower litter had a higher concentration of nutrients important to decomposition, such as N, P, K, and water-holding capacity, while leaf litter had a higher concentration of Ca, Mg, and Na. To our knowledge, this is the first study to demonstrate the importance of litter-mixing effects between flower and leaf litter. This result sheds light on the secondary consequences of flower litter on decomposition, suggesting that species with high reproductive investment in flower biomass may play an important role in the nutrient and carbon recycling

25 of diverse plant communities, exerting a pivotal role in biogeochemical dynamics.

30

### 1 Introduction

Decomposition is an important ecosystem process because of its role in the energy and matter flows within and across terrestrial and aquatic ecosystems, which affects the cycling of nutrients and carbon (C) in the biosphere (Waring, 2013). Up to 90% of the primary production accumulates as organic matter (OM) in the soil (Cebrian, 1999), and a considerable proportion of this stock is transported to rivers, lakes, and oceans, contributing to the stocks of OM in aquatic environments, along with autochthonous OM (Tranvik et al., 2009; Aufdenkampe et al., 2011). Decomposition is controlled by abiotic factors such as temperature and humidity, as well as biotic factors such as the abundance and composition of decomposers, and litter quality, with

35



40 the relative importance of each factor varying between biomes and ecosystems (Makkonen et al., 2012; García-Palacios et al.,  
2016; Djukic et al., 2018). While global OM stocks are relatively well known (Schimel et al., 2001; Hengl et al., 2017), the  
mechanisms governing OM dynamics in ecosystems are far less understood (Tian et al., 2015). Given the vast size of OM stocks  
in ecosystems, even minor changes in OM content and dynamics can have major impacts on global C and nutrient budgets (Dignac  
et al., 2017; Minasny et al., 2017). Therefore, understanding the peculiarities of the mechanisms that regulate decomposition  
45 dynamics is crucial for comprehending the flows and stocks of C and nutrients in terrestrial and aquatic ecosystems (Basile-Doelsch  
et al., 2015; Davidson and Janssens, 2006).

Litter quality, which refers to the edibility of litter as a food resource for decomposers, has effects on decomposition that  
are comparable to, or even stronger than, those of abiotic factors across terrestrial and aquatic ecosystems (Makkonen et al., 2012;  
García-Palacios et al., 2016). The chemical compounds and physical structures of plant litter in ecosystems are highly diverse and  
heterogeneous, leading to distinct litter quality. The diversity of plant life forms, organs, and traits corresponds to the myriad of  
50 plant tissues that contain different pools of compounds (Jackson et al., 2013). After senescence, these tissues form litter pools with  
a wide range of resistance to biological (mostly microbial) degradation (Kuzyakov and Blagodatskaya, 2015; Jones et al., 2023).  
Therefore, the variety of functional differences in plant litter can affect decomposition both intrinsically and through mechanisms  
mediated by complex interactive effects among contrasting litter types, resulting in litter-mixing effects (LMEs) on decomposition  
(Schindler and Gessner, 2009; Liu et al., 2020; Hättenschwiler and Jørgensen, 2010).

55 Most biogeochemical models predicting decomposition dynamics in natural ecosystems ignore interactions among plant  
litters of different qualities and assume that the sum of the individual litters can predict decomposition in a mixture (Manzoni and  
Porporato, 2009). However, LMEs on decomposition could either increase (i.e. synergistic) or decrease (i.e. antagonistic) the  
decomposition rate of individual litter compared to their monocultures, and the magnitude of such effects can be highly variable  
depending on litter traits, decomposer community, and environmental contexts (Gartner and Cardon, 2004; Liu et al., 2020; Porre  
60 et al., 2020). The synergistic effects are mainly attributed to mechanisms such as nutrient transfer, niche partitioning among  
decomposers, and improvement of microclimatic conditions (Gessner et al., 2010; Boyero et al., 2011; Hättenschwiler et al., 2005;  
Makkonen et al., 2013). Mechanisms responsible for antagonistic effects include microbial nutrient immobilization, inhibitory  
decomposition by secondary compounds, and decomposer preferential feeding (Kuzyakov, 2002; Hättenschwiler et al., 2005).  
However, these mechanisms have been predominantly demonstrated in studies that manipulate inter-specific leaf litter diversity  
65 and in experiments where species contribute equally to the litter biomass in the mixture (but see Madritch and Hunter 2004;  
Crutsinger et al. 2009; Zhang et al. 2022). Some efforts have been directed towards understanding the intra-specific LMEs from  
different plant organs or varying proportions of litter types in litter mixtures on ecosystem functions (Dearden et al., 2006; de Paz  
et al., 2018; Hou and Lü, 2021; Zhao et al., 2019; Zhang et al., 2022). This is an important research avenue because it reinforces  
the idea that LMEs on decomposition depend more on functional dissimilarity than on the taxonomic richness of litter, and it may  
70 also indicate that within-species LMEs may occur and be particularly relevant for ecosystem functioning in low-diversity plant  
communities.

Studies have demonstrated that the mixture of various sources of dissolved and particulate labile organic matter (LOM) and  
refractory organic matter (ROM) can exert opposing effects on each other's decomposition, both in terrestrial and aquatic  
ecosystems (Guenet et al., 2010). Generally, LOM is expected to accelerate the decomposition of ROM (Guenet et al., 2010) while  
75 ROM is expected to inhibit the decomposition of LOM (Liu et al. 2020; but see Swan and Palmer 2006; Cuchietti et al. 2014).  
However, it is currently unclear how variations in the relative proportions of LOM and ROM in litter mixtures affect the magnitude  
and direction of LMEs, considering the degradation rate of each litter type and the whole litter mixture. This is important because  
the LOM:ROM biomass ratio in the detritus pool varies spatially and temporally within and among ecosystems due to a variety of



causes (McClain et al., 2003). The LOM:ROM biomass ratio in the detritus pool is critical for microbial degradation rates, as the limited number of metabolic pathways available to microbial decomposers have specific energy requirements (German et al., 2011). Therefore, the LOM:ROM proportion in litter mixtures could be pivotal in determining the occurrence, magnitude, and direction of LMEs on decomposition (Smith and Bradford, 2003; Schindler and Gessner, 2009), but this is not well understood (Sayer et al., 2007; Gripp et al., 2018).

In this study, we explored a novel potential after-life role of flower litter in mediating the LMEs on decomposition. For this, we utilized flower and leaf litter (hereafter litter types) from the trumpet tree *Tabebuia aurea* (Silva Manso) Benth. & Hook. f. ex. S. Moore, as sources of LOM and ROM, respectively. To better understand the possible mechanisms underlying LMEs on decomposition, we evaluated the occurrence and magnitude of LMEs on each litter type individually and on whole litter mixture. In doing so, we sought to determine whether the LMEs of flower and leaf litter mixing were reciprocal or unilateral, and whether the magnitude and direction (i.e. synergistic or antagonistic) of such effects were symmetric or asymmetric. Previous studies have shown that the magnitude of LMEs on decomposition is affected by dissimilarity in litter quality (Schindler and Gessner, 2009; Finerty et al., 2016). As a result, much of the research on LMEs has focused on mixing leaf litter from different species (Porre et al., 2020; Hättenschwiler et al., 2005), whereas the interaction among mixed litters from different plant organs has not been well explored. It is important to note that unequal investment among plant organs, which can occur due to differences in organ form and function, may result in variations in the chemical composition of tissues across different plant organs, which has legacy consequences on litter decomposition (Freschet et al., 2013; Jackson et al., 2013). Leaves are specialized organs for photosynthesis and have a longer lifespan than flowers (Roddy et al., 2019). Therefore, accordingly to the growth-rate hypothesis leaves are expected to have higher concentrations of structural compounds such as lignin and secondary metabolites than flowers (Stamp, 2003). Conversely, flowers are fast-growing ephemeral organs specialized in reproduction (Ashman and Schoen, 1994). As a result, flowers, on average, are expected to receive less investment in the production of constitutive defenses against herbivory and structural tissues, but a greater investment in nutrients for growth and in labile C-compounds such as sugars for nectar production to attract pollinators (Mccall and Irwin, 2007; Boaventura et al., 2022). Finally, the differences in litter quality between leaves and flowers could also become more pronounced during senescence, as leaves have been shown to have the highest rate of nutrient resorption among plant organs (Freschet et al., 2010). Thus, since litter characteristics in general reflect environmental pressures that shape the form and function of plant organs when alive, we rationalized that flower litter of *T. aurea* is more labile than its leaf litter, and that this dissimilarity may cause LMEs in their decomposition when both litters are mixed.

We tested the following hypotheses: (i) flower litter quality will be higher (i.e. LOM) than leaf litter quality (i.e. ROM) and consequently flower litter will decompose faster than leaf litter; (ii) the interaction between flower and leaf litter during decomposition will result in LMEs; (iii) litter-mixing effects on each litter type will be mostly asymmetric, with more frequent and/or stronger positive effects of flower litter on the decomposition of leaf litter; and (iv) litter-mixing effects, on each litter type and mixture, will depend on the relative proportion of each litter type in the mixture. Since the species from the *Tabebuia* genus as well as other trumpet trees can colonize floodplains, riparian areas, and seasonally dry forests in the tropics (Ribeiro and Brown 2006), the litter of these species can be decomposed under aquatic or terrestrial ecosystem contexts. Litter-mixing effects on decomposition have been traditionally investigated in aquatic (Boyero et al., 2021) and terrestrial (Makkonen et al., 2012) ecosystems separately (but see Handa et al. 2014; García-Palacios et al. 2016) using different species and methodologies. This hinders testing the generality of the results of these studies for different types of ecosystems. Therefore, considering that *T. aurea* can contribute litter to both terrestrial and aquatic ecosystems, we tested our hypotheses throughout terrestrial and aquatic experiments.



## 2 Methods

### 120 2.1 Study site and species

The experiment was conducted in the laboratory at the Universidade Federal do Rio Grande do Norte, Brazil. The flower and leaf litter of *T. aurea* was sampled in a forest fragment (more details below). The geographic distribution of *T. aurea* in South America extends to most Brazilian biomes, such as the Amazon, Atlantic Forest, Cerrado, and Pantanal (Lorenzi, 1992), and its environmental distribution ranges from dry forests to riparian forests and floodplains (Batalha and Mantovani, 2001; Lorenzi, 125 1992). Thus, the widespread geographic and environmental distribution of *T. aurea* allows its litter to contribute to the flow of matter and energy in both the aquatic and terrestrial ecosystems (Fig. S1; see Supplement A for more details to species in the section *Species used*). Synchronous and massive flowering, which is a common characteristic of Bignoniaceae species, is preceded by the loss of leaves (Barros, 2001). This phenological pattern creates a potentially important scenario for testing the LMEs on decomposition, as a layer of leaf litter is deposited in the soil, which is then covered by a layer of flower litter a few days later (Fig. 130 S1).

### 2.2 Litter sampling and experimental design and setup

We sampled flower and leaf litter under the canopy of *T. aurea* individuals immediately after abscission. Soon after litter sampling, the litter types were separately dried at 60 °C for 72 h until a constant weight was achieved. The litter was conditioned 135 in a dry and dark place to avoid changes in its chemical composition.

The experimental design followed an additive rather than a substitutive design, which is commonly used in experiments designed to test the effects of species diversity and litter mixing on ecosystem functioning (Jolliffe, 2000) (Table S1; see Supplement A for more detailed description about the additive experimental design in *Experimental design and setup*).

We performed controlled laboratory experiments to simulate terrestrial and aquatic environments. The duration of the 140 terrestrial and aquatic experiments was standardized by the time required for approximately 50% of the more labile litter (i.e. flowers) to be decomposed in each environment. The decomposition process is faster in aquatic environments (Lidman et al., 2017); therefore, the aquatic experiment lasted for 3 months, while the terrestrial experiment lasted for 7 months.

To ensure aerobic conditions in both environments, distinct microcosms were used. In the terrestrial experiment, plastic containers (5 cm in diameter and 10 cm in height) were used as microcosms. Each terrestrial microcosm was filled with a soil layer 145 of approximately 5 cm height, collected under the canopy of *T. aurea* individuals in the same area in which litter was collected. The soil was sieved (2-mm mesh size) to remove large litter particles, homogenized, and added to the microcosms. This procedure maintained a substantial part of the soil microflora and micro- and meso-fauna (Swift et al., 1979) while reducing environmental heterogeneity among experimental microcosms. The amounts of litter types were added according to the information in Table S1.

Because *T. aurea* displays a very marked phenological pattern of leaf and flower abscission (where leaves senesce and fall 150 completely one to two weeks before flowering and the consequent flower fall), and because studies have demonstrated that litter spatial position can alter litter decomposition (Berenstecher et al., 2021), we arranged the flower and leaf litter in the microcosms resembling their natural position in the litter layer. First, the leaf litter was added above the soil within the microcosms, followed by the flower litter overtopping it. All microcosms were then randomly arranged in plastic trays and covered with a 1 mm mesh opening screen to prevent the entry of foreign materials. The experiment was conducted in a laboratory room at a constant 155 temperature of approximately 25 °C and a 12:12 h light:dark period. To avoid moisture limitation of litter decomposition in the terrestrial experiment, each microcosm was individually irrigated every three days with approximately 50 ml of tap water using a hand-held sprinkler. The amount of water was based on an estimation of the accumulated average precipitation at the study site during the experiment (January to July; Santos e Silva et al. 2012)



160 In the aquatic experiment, the microcosms were composed of 1 L glass bottles. The amount of each litter type added to the  
respective monocultures and mixtures is shown in Table S1. Tap water was used to fill the aquatic experimental microcosms. The  
microcosms of the mixtures were filled with 1 L of water. However, as the final litter biomass added to the microcosms differed  
between mixtures and their respective monocultures, as well as throughout the monocultures, the volume of water in each  
microcosm was varied to maintain a final litter concentration of 3 g/L across all treatments.

165 As in the terrestrial experiment, litter types were added intact to the microcosms. To prevent litter from floating and/or  
sticking to the inner wall of the microcosms, we packed litter in synthetic bags with a 1 mm mesh size, each containing a small  
metal weight, to keep the litter near the bottom of the microcosms. We ensured aerobic conditions in each microcosm and promoted  
water circulation and constant homogenization of abiotic conditions by providing aeration to each individual microcosm. The  
microcosms were randomly distributed along shelves in a darkened room at a constant temperature of approximately 25 °C to  
avoid primary production.

170

### 2.3 Measurements of the litter mass remaining

At the end of the experiments, litter was carefully removed from the microcosms. For mixtures, the remaining flower and  
leaf litter were visually identified and separated and subsequently dried and weighed to estimate litter mass loss. The procedure  
varied between terrestrial and aquatic experiments. In the terrestrial experiment, we separated the remaining flower and leaf litter  
175 from each other (in mixtures) and from the soil particles and placed them in aluminum trays for subsequent weighing. In the aquatic  
experiment, flower litter fragmentation prohibited the same procedure. Instead, we filtered the litter from each microcosm, for both  
monocultures and mixtures, using a previously weighted paper filter. For monocultures, we removed the flower and leaf litter from  
the bags and placed them on a paper filter at the end of the experiment. We then poured the entire water volume from each  
microcosm containing the leaked particulate material into the corresponding filter. We followed the same procedure for the  
180 mixtures; however, leaf litter fragments, which were tougher than flower litter, were easily identified and collected from the filter  
surface. Leaf litter fragments were placed in previously weighed paper filters. The identified litter on the filter paper was then  
packed in aluminum trays and dried at 60 °C for 72 h. Subsequently, we repeated the weighing procedure and measured the  
remaining mass of each litter type in its respective microcosm, allowing us to estimate the decomposition rate of each litter type  
individually, even in mixtures.

185 We estimated the decomposition rate in both experiments as the percent of litter mass remaining (LMR %) calculated as  
the percentage of the dry mass of each type of litter (decomposing alone or in the mixture) at the end of the experiment concerning  
its respective initial dry mass accordingly to the following Eq. (1):

$$\text{LMR}_i (\%) = (\text{Fdwi}/\text{Idwi}) \times 100, \quad (1)$$

where  $\text{Fdwi}$  and  $\text{Idwi}$  are the final and initial dry weights of litter  $i$  (flowers or leaves), respectively.

190 To estimate the total percentage of litter mass remaining for the two litter types combined in the observed mixture, we used  
Eq. (2):

$$\text{LMR}_{obs}(\%) = \left( \frac{\text{Fdwf} + \text{Fdwf}}{\text{Idwf} + \text{Idwl}} \right) \times 100, \quad (2)$$

where  $\text{Fdwf}$  and  $\text{Fdwf}$  are the final dry weights of flower and leaf litter, respectively, and  $\text{Idwf}$  and  $\text{Idwl}$  are the initial dry weights  
of flower and leaf litter at the beginning of the experiment, respectively.

195

### 2.4 Estimative of flower and leaf litter functional traits

We measured a set of traits to describe the functional differences between the litter types of *T. aurea*. These analyses focused  
on estimating the initial values for litter chemical and physical traits that commonly have an impact on litter decomposition in



200 terrestrial and aquatic ecosystems. For chemical analyses, at least 3 g of each litter type was ground to a fine powder using a mortar  
and pestle. We then estimated the total C concentration using the high-temperature combustion method and infrared CO<sub>2</sub> detection  
with a Shimadzu TOC-5000 total carbon analyzer. Total nitrogen content (N) was estimated by acid digestion using Kjeldahl  
distillation (Allen et al. 1974). The total phosphorus (P) was estimated through strong acid digestion and reaction with molybdate  
(Fassbender, 1973). Potassium (K), calcium (Ca), manganese (Mg), and sodium (Na) content were estimated according to  
analytical protocols (Sarruge and Haag, 1974; Robertson et al., 1999). Structural compounds, such as lignin (Lig) and cellulose  
205 (Cel), were estimated by the sequential method of neutral detergent and second acid detergent digestion (Goering and Van Soest,  
1970). Phenolic compounds (Phe) were estimated by the Folin assay (Graça et al., 2005). We used the Antrona method (Morris,  
1948; Van Handel, 1968) to determine the non-reducing soluble sugars (S-carb).

To evaluate physical traits, we assessed the water-holding capacity (WHC) and leaching of flower and leaf litter, which are  
considered crucial factors in determining litter decomposition (Makkonen et al., 2013). To evaluate WHC, we used dried flower  
210 and leaf litter and moistened the replicates with 50 ml of water (the same volume used to irrigate the terrestrial experiment) two  
hours before the measurements, based on Makkonen et al. (2013). For the leaching measurement, we stimulated the loss of  
hydrolysable water compounds, which is the main form of mass loss in the initial stage of decomposition, based on Pérez-  
Harguindeguy et al. (2013). Our physical traits were measured based on the flower:leaf biomass proportions used in the experiment.  
but due to experimental limitation, we used six treatments encompassing the whole gradient of variation since only flower litter  
215 (1:0), decreasing the flower biomass in 0.8, 0.6, 0.4, and 0.2, until only leaf litter (0:1). Each proportion was replicated three 3  
times. For both structural traits, the material was dried in an oven at 60 °C for 72 h before and after the measurements.

## 2.5 Data analysis

To test the functional differences between the litter types of *T. aurea*, we compared the average concentrations of functional  
220 traits (C, Cel, Lig, Phe, S-Carb, N, P, K Na, Mg, Ca, WHC, leaching, and some stoichiometric ratios (Lig:S-Carb, Lig:N, Lig:P)  
between both litter types using an unpaired t-test.

To compare the LMEs on the decomposition of the whole mixture, that is, the aggregate response of the two or more litter  
types to litter mixing (LMR<sub>obs</sub>), the observed values of the aggregate decomposition of both litter types in the mixture should be  
compared with those expected from the sum of the weighted decomposition of each component litter type in monoculture (Loreau,  
225 1998). The expected LMR for the combined responses of both litter types to litter mixing was calculated using Eq. (3):

$$\text{LMR}_{\text{exp}}(\%) = \left( (\text{LMR}_f) \times (p_i) \right) + \left( (\text{LMR}_l) \times (p_i) \right), \quad (3)$$

where LMR<sub>f</sub> is the percentage of flower litter mass remaining in the monoculture, LMR<sub>l</sub> is the percentage of leaf litter mass  
remaining in the monoculture, and  $p_i$  is the proportion of the biomass of litter  $i$  in the mixture.

We conducted a set of statistical analyses to test our hypotheses. Initially, we conducted unpaired t-tests between the litter  
230 types to evaluate if the decomposition rate of flower and leaf litter differed. Specifically, we compared the LMR of each litter type  
in monocultures in both terrestrial and aquatic experiments. Then, we conducted regression analyses to assess the effect of flower  
litter biomass on the decomposition of each litter type, alone and in mixtures. Specifically, we assessed the LMR of each litter type  
in monocultures and mixtures as a function of the variation in flower litter biomass proportion. Next, we employed the test of  
heterogeneity of slopes to determine if litter mixture affected the biomass-decomposition relationship of each litter type separately  
235 as well as to the mixture. For this analysis we considered the proportion of flower in the litter mixture as the predictor and the  
LMR of each litter type alone or combined as the response variable. Statistically significant effects (i.e. when the slopes of the  
regressions are different from each other) would indicate that the biomass-decomposition relationship differed between litter  
decomposing alone and in mixture in response to flower litter biomass proportion. In cases where the slopes of the regressions did



not differ significantly from each other, we used unpaired t-tests to compare the grand mean decomposition (i.e. irrespective of  
240 litter biomass) of each litter type alone as well as for the whole mixture. The aforementioned analytical procedures were performed  
separately for terrestrial and aquatic experiments because of experimental design differences between them (see details in *Litter  
sampling and experimental design and setup*).

To understand the magnitude of LMEs on the decomposition of individual litter types and the whole mixture in response to  
flower litter biomass proportion, we calculated the relative mixture effect (RME) in each litter type for each ecosystem (Barantal  
245 et al., 2011) using Eq. (4):

$$\text{RME (\%)} = \left( \frac{\text{LMR}_{exp} - \text{LMR}_{obs}}{\text{LMR}_{exp}} \right) \times 100, \quad (4)$$


where RME is the relative mixture effect (%) for the whole combined litter or each litter type in the mixture; for the whole mixture,  
LMR<sub>exp</sub> is the expected litter mass remaining calculated by averaging the LMR values of both litter types in monoculture, and  
LMR<sub>obs</sub> is the observed litter mass remaining of the whole mixture calculated by averaging the observed LMR values of both litter  
250 types in the mixture. For each litter type, LMR<sub>exp</sub> is the flower or leaf litter mass remaining in the monoculture and LMR<sub>obs</sub> is the  
flower or leaf litter mass remaining in litter *i* in the mixture. For RMEs, positive and negative values indicated that litter decomposes  
faster and slower in mixtures than in its respective monoculture. To assess whether the RME for each type of litter and for the  
mixture is a function of the biomass of the respective litter type and the proportion of the mixture between flower and leaf litter,  
we utilized linear regressions for both aquatic and terrestrial experiments.

255 Before linear regression analysis, the data were tested for assumptions of linearity with Run's Test. The homogeneity of the  
residuals as assumptions for linear regressions and unpaired t-tests were tested using the Bartlett Test. All statistical analyses were  
performed using the GraphPad Prism software (version 6.0). A level of significance of  $\alpha = 0.05$  was considered for all analyses.

### 3 Results

#### 260 3.1 Flower and leaf litter chemical and structural composition

Overall, flower litter had a more labile chemical composition and physical traits than leaf litter. Except for the C content,  
all analyzed chemical constituents displayed significantly different concentrations between the litter types (Fig. 1a). Specifically,  
flower litter had significantly higher concentrations of S-carb, N, P, and K than leaf litter, whereas leaf litter had significantly  
higher concentrations of Cel, Lig, Phe, Ca, Mn, and Na than did flower litter (Fig. 1a). Additionally, the Lig:S-Carb ratio, which  
265 indicates the relative proportion of recalcitrant and labile C, was significantly lower in the flower litter than in the leaf litter. The  
same pattern was observed for Lig:N and Lig:P ratios (Fig. 1b). Finally, the physical traits followed the same pattern, with flower  
litter exhibiting higher values of WHC and leaching than leaf litter (Fig. 1c).



270 **Figure 1:** Average ( $n=3 \pm 1SD$ ) functional trait values for a) concentration of chemical constituents, b) stoichiometry, and c) physical traits for flower and leaf litter of *Tabebuia aurea*. Abbreviations are as follow: Cel – cellulose, Lig – lignin, Phe – phenolic compounds, S-carb – soluble carbohydrates, WHC – water holding capacity. Asterisks above the bars indicate significant statistical differences between the flower and leaf litter for the respective functional traits (unpaired t-test; \*\* $p<0.001$ ; \*\*\* $p<0.0001$ ).





### 275 3.2 Differences in the decomposition rate between the litter types in monoculture


Considering only the values of leaf and flower litter decomposing in monocultures, the leaf litter decomposed significantly slower than flower litter in the terrestrial experiment, with the average leaf LMR (84.4%) was significantly higher compared to the average flower LMR (41.8%) (Fig. 2ab;  $t=72.4$ ;  $p<0.0001$ ; unpaired t-test). This pattern was consistent to the aquatic experiment, the values of leaf and flower litter decomposing in monoculture were significantly different from each other. The leaf litter decomposed significantly slower than flower litter, with average leaf LMR (72.7%) significantly higher than the average flower LMR (50.3%) (Fig. 3ab;  $t=13.3$ ;  $p<0.0001$ ; unpaired t-test).

### 285 3.3 Litter-mixing effects of mixed flower and leaf litter on decomposition in the terrestrial experiment

Leaf litter decomposition rates, expressed as LMR, did not significantly vary across the gradient of leaf litter biomass in the monoculture (Fig. 2a). However, leaf litter decomposition was significantly altered when mixed with flower litter. In general, an increase in the amount of flower litter had significant, positive (i.e. lower LMR values) and linear effects on leaf litter decomposition rates (Fig. 2a;  $F_{1, 14} = 215.9$ ;  $p<0.0001$ ). However, interestingly, in the two mixtures with the lowest flower:leaf litter biomass proportion, the decomposition of leaf litter were lower than those observed in their respective monocultures (i.e. higher LMR values; Fig. 2a).

290 Similar to leaf litter, flower litter decomposition rates did not vary significantly in response to its biomass variation in the monoculture (Fig. 2b). However, as observed for leaf litter decomposition, flower litter decomposition was significantly altered when mixed with leaf litter. In this scenario, the increasing amounts of leaf litter in mixtures had significant but negative (i.e., higher LMR values), linear, and LMEs on flower litter decomposition rates (Fig. 2b;  $F_{1, 14} = 65.9$ ;  $p<0.0001$ ).

295 Finally, we observed similar significant effects on the variation in LMR for both litter types combined in response to the proportion of flower litter mass in the mixture (Fig. 2c). Decreasing the amount of flower litter in the mixture significantly increased both the expected and observed values of LMR for the litter mixture as a whole, although the slopes of both relationships did not differ significantly (Fig. 2c;  $F_{1, 14} = 3.5$ ;  $p=0.08$ ). Litter-mixing effects on the whole mixture was significant (Fig. 2c – right panel;  $t=16.5$ ;  $p<0.0001$  unpaired t-test), as the average observed LMR for the whole mixture (37%) was significantly lower than its average expected value (63%) calculated from the decomposition of both litter types alone. These results indicated that, on average, litter mixing had a stimulating effect of 26% on the decomposition of the whole mixture treatment.



**Figure 2:** Patterns of flower and leaf litter mass remaining in the terrestrial experiment for single a) leaf, b) flower, and c) whole mixture (leaf +flower) decomposing alone (i.e. monocultures) or mixed (i.e. mixtures). Litter mass remaining for each litter type alone and in combination were fitted as linear functions of flower:leaf litter proportion. Values in left panels are mean ( $n=10 \pm 95\%$  CI). Slope values in bold indicate significant statistical differences regarding the interactive effects between the explanatory variable and litter mass remaining ( $p<0.05$ ; F-test for Homogeneity of slopes analysis). Values in right panel of Fig. 2c depict the grand mean ( $n=90 \pm 95\%$  CI). Different letters above the grand mean values indicate significant statistical difference ( $p<0.05$ ; unpaired t-test).




### 310 3.4 Litter-mixing effects of mixed flower and leaf litter on decomposition in the aquatic experiment

Leaf litter biomass did not significantly affect the variation in leaf LMR, either alone or in combination with flower litter (Fig. 3a;  $F_{1, 14} = 0.1$ ;  $p=0.76$ ). However, when mixed with flower litter, the average leaf LMR was significantly lower (69.5%) than that of its monoculture (72.7%), indicating that leaf litter decomposed 4.5% faster on average in the presence of flower litter (Fig. 3a, right panel;  $t=2.1$ ;  $p=0.04$ , unpaired t-test).

315 Similar patterns were observed for flower litter decomposition. Variations in flower litter biomass had no effect on flower LMR, either alone or in combination with leaf litter (Fig. 3b;  $F_{1, 14} = 0.3$ ;  $p=0.62$ ). However, when mixed with leaf litter, the average flower LMR was significantly lower (47.1%) than that of its monoculture (50.3%), indicating that flower litter decomposed on average 6.3% faster in the presence of leaf litter (Fig. 3b, right panel;  $t=2.2$ ;  $p=0.04$ , unpaired t-test).

320 Finally, the expected and observed values for the LMR of the whole mixture increased significantly as a function of the decreasing proportion of flower litter mass in the mixture (Fig. 3c). However, similar to the observations in the terrestrial experiment, the slopes of both relationships were not significantly different from each other (Fig. 3c;  $F_{1, 14} = 1.3$ ;  $p=0.27$ ). Nevertheless, the LMEs on the decomposition of the whole mixture were significant (Fig. 3c, right panel;  $t=2.3$ ;  $p=0.03$ , unpaired t-test) on average. The observed LMR for the litter mixture (57.5%) was significantly lower than its expected value (61.5%),  
325 6.5% on the decomposition of the whole litter mixture.



**Figure 3:** Patterns of flower and leaf litter mass remaining in the aquatic experiment for single a) leaf, b) flower, and c) whole mixture (leaf +flower) decomposing alone (i.e. monocultures) or mixed (i.e. mixtures). Litter mass remaining for each litter type alone and in combination were fitted as linear functions of flower:leaf litter proportion. Values in left panels are mean ( $n=3 \pm 95\%$  CI) for monocultures and ( $n=6 \pm 95\%$  CI) for mixtures. Slope values in bold indicate significant statistical differences regarding the interactive effects between the explanatory variable and litter mass remaining ( $p<0.05$ ; F-test for Homogeneity of slopes analysis). Values in right panels depict the grand mean ( $n=27 \pm 95\%$  CI) for monocultures and ( $n=54 \pm 95\%$  CI) for mixtures. Different letters above the grand mean values indicate significant statistical difference ( $p<0.05$ ; unpaired t-test).

330

### 3.5 The magnitude of RME for terrestrial and aquatic experiments


335



Variations of RME values for the two litter types and for the whole litter mixture in response to flower:leaf litter biomass proportion showed distinct patterns for terrestrial and aquatic experiments (Fig. 4). In general, RME values were higher for the terrestrial experiment compared to the aquatic experiment considering each litter type and the whole mixture. In the terrestrial experiment, RME values for flower, leaf, and both litter types combined, varied significantly and negatively as a function of flower:leaf litter proportion (Fig 4a). However, the variation of RME values for flower and leaf litter as a function of flower:leaf biomass proportion were parallel and not statistically different from each other ( $F_{1, 14} = 0.04$ ;  $p=0.85$ ) but they were both statistically different (steeper) from the variation of RME values for the whole mixture ( $F_{2, 21} = 21.2$ ;  $p<0.0001$ ). On average, the decomposition of flower litter increased 49% in the presence of leaf litter and the decomposition of leaf litter increased, on average, 31.2% in the presence of flower litter, but such a difference was not statistically significant (Fig. 4a right panel;  $t=1.2$ ;  $p=0.25$  unpaired t-test).

Contrary to what was observed in the terrestrial experiment, we did not observe significant effects of flower:leaf litter biomass proportion on the variation of RME values for leaf and flower litter as well as for whole mixture (Fig. 4b). Furthermore, average values of RMEs for flower (7%) and leaf litter (6%) were not significantly different from each other (Fig. 4a right panel;  $t=0.34$ ;  $p=0.74$  unpaired t-test).

350



**Figure 4:** Response of the relative mixture effect (RME) of flower, leaf and whole mixture to the gradient of flower:leaf litter proportion in a) terrestrial and b) aquatic ecosystems. Regression lines sided by different letters indicate their slopes are statistically different from each other ( $p<0.05$ ; F-test for Homogeneity of slopes analysis). Values in right panel of Fig. 4a depict the grand mean. Symbols represent mean  $\pm 1SD$ . Errors bars were shorter than the size of the symbols.

355

#### 4 Discussion



Our study is the first to assess the LMEs of mixing flower and leaf litter on decomposition across terrestrial and aquatic ecosystems. Our findings suggest that flowers have a lasting interactive effect on litter decomposition beyond their role in reproduction, providing evidence of LMEs resulting from the mixture of litter from different plant organs, even at the intra-specific level. First, our findings reveal that *T. aurea*'s flower and leaf litter have distinct functional traits and decomposition rates. Consistent with our first hypothesis, *T. aurea*'s flower litter exhibits chemical and physical functional trait values indicative of a more labile detritus compared to *T. aurea*'s leaf litter. Such functional divergences between flower and leaf litter quality were confirmed by the higher decomposition rates of flower litter compared to leaf litter in both terrestrial and aquatic environments. Secondly, our results strongly supported our second prediction since the LMEs of flower and leaf mixing were consistent in terrestrial and aquatic environments. Third, we hypothesized that the LMEs of flower litter on leaf decomposition would be stronger and more positive than those from leaf litter on flower litter. However, flower and leaf litter mixing had reciprocal effects on the decomposition of each other, with symmetric LMEs (both in magnitude and direction) on the decomposition of both litter types and in both terrestrial and aquatic ecosystems. Finally, our fourth prediction that the LMEs resulting from the mixture of flower and leaf litter would vary in magnitude and direction depending on the proportion of flower and leaf biomass in the litter mixture, was supported only in the terrestrial environment, where the decomposition rate of leaves, flowers, and the whole mixture, was faster with increasing proportion of flower litter in the litter mixture. This result has two important ramifications. A more specific one suggests that the unbalance distribution of nutrients in flower and leaf tissues has consequences on litter decomposability after the senescence process, with further effects to the interaction between flower and leaf litter decomposition (Schmitt and Perfecto, 2020). This result expands what have been reported by recent studies showing the effects of litter mixing from different plant organs for the occurrence of LMEs on decomposition (de Paz et al., 2018; Lin et al., 2019; Hou and Lü, 2021). Secondly, a more general implication of our results indicates that since the increasing proportion of flower litter in the mixture increased the decomposition of litter mixture as a whole, the LOM:ROM biomass proportion in the detritus pool can be a crucial factor mediating the mechanisms controlling the decomposition process in ecosystems (Guenet et al., 2010). Finally, our results also reinforce the notion of what has been observed in studies that seek to synthesize the effects of litter functional diversity on decomposition, which advocate that LMEs are more consistent in terrestrial than in aquatic ecosystems (Gessner et al., 2010).

Leaves and flowers are plant organs with distinct functions and forms, resulting in differences in their chemical and physical characteristics. Recent research has emphasized the significance of indirect effects of ecological and evolutionary mechanisms in shaping litter decomposition through legacy effects from functional traits of living plant tissues that persist after tissue death and impact litter decomposition through after-life effects (Freschet et al., 2012; Cornelissen et al., 2023). The longer lifespan and persistence of leaves require the plant to invest more in structural tissues to provide greater physical resistance for these organs, as well as in secondary compounds that act as constitutive defenses against herbivory and photo-damage by UV radiation (Stamp, 2003). Additionally, leaf senescence is generally slow, and many species have efficient nutrient resorption prior to leaf abscission, especially deciduous species living in infertile soils (Brant and Chen, 2015) like the one used in our study. On the other hand, flowers are ephemeral reproductive organs with high concentrations of nutrients and soluble compounds, as well as labile C compounds that make up nectar to attract pollinators (Freeman et al., 1991; Galetto and Bernardello, 2004). Compared to leaves, flowers are generally fast-growing and short-lived organs (Ashman and Schoen, 1994), which results in higher nutrient investment, lower nutrient resorption and lesser herbivore attack than leaves (Mccall and Irwin, 2007). As a result, flower litter may contain higher concentrations of nutrients and labile C compounds, while having lower amounts of structural tissues and deterrent secondary compounds (Stamp, 2003). These conjectures were most supported by our data and confirmed our hypothesis that flower litter is a more LOM than leaf litter and decomposes faster than leaf litter in both terrestrial and aquatic ecosystems. Flower litter has higher concentrations of N, P, K, and labile C, as well as higher leaching capacity and WHC, while leaf litter



were richer in micronutrients such as Mg, Ca, and Na. These results also supported the assumptions we have used to rationalize our second hypothesis, which predicted that due to functional differences in litter quality, the mixture of flower and leaf litter would cause LMEs on the decomposition of both litter types. Studies in both terrestrial and aquatic ecosystems have shown that the litter functional dissimilarity rather than litter species number is the most important factor causing LMEs on decomposition (Epps et al., 2007; Lecerf et al., 2011; Violle et al., 2017). Our study supports this paradigm in demonstrating that LMEs on decomposition can also occur intraspecifically via the interaction of flower and leaf litter, and call attention to the importance of LMEs on decomposition even in low-diverse systems through the interactions of litter from different plant organs.

405           Meanwhile, according to our third hypothesis, we expected the LMEs between flower and leaf litter to be asymmetrical, with the leaf litter decomposing more quickly in the presence of flower litter than vice versa. Our conjecture relied on the results of past studies that show that litter with contrasting qualities affects and/or responds to litter mixing in distinct ways. In general, studies have shown that LOM stimulates the decomposition of ROM mainly due to nutrient transfer and/or priming effects (Guenet et al., 2010; Liu et al., 2020), while ROM may inhibit the decomposition of LOM due to the presence of deterrent secondary metabolites (Hättenschwiler et al., 2005). However, our findings only partially supported our third prediction. While we observed that, on average, flower litter (LOM) accelerated the decomposition of leaf litter (ROM), we also found that leaf litter reciprocally accelerated the decomposition of flower litter, and the magnitudes of these effects were statistically indistinguishable from each other in both terrestrial and aquatic experiments. These results are due to complementary effects. The two litter types have highly contrasting chemical and physical characteristics. Flower litter has a higher concentration of labile C and nutrients (N, P, K), and WHC and leaching potential than leaf litter. This pattern points out to the possibility of flower litter to accelerate the decomposition of leaf litter through mechanisms such as nutrient transfer and/or improved microenvironment conditions (Hättenschwiler and Jørgensen, 2010; Makkonen et al., 2013). On the other hand, leaf litter has higher concentrations of micronutrients such as Mg, Ca, Na, which could, via micronutrient transfer, compensate for possible limitations of flower litter decomposition by these elements. In fact, studies have already shown that litter decomposition is co-limited by macro and micronutrients in tropical forests (Kaspari et al., 2008; García-Palacios et al., 2016). Additionally, specifically for the terrestrial experiment, the higher toughness of leaf litter may have influenced microenvironmental conditions of the litter layer inside microcosms (Makkonen et al., 2013), preventing the compaction of the flower litter layer and avoiding anaerobic conditions, which could negatively affect the decomposition of flower litter in monocultures.

425           However, for a better understanding of the aforementioned mechanisms underlying the LMEs of flower and leaf litter mixing on decomposition, it is essential to consider how these effects, respective to each litter type, varied in response to the flower-to-leaf litter proportion. In the terrestrial experiment, we observed that RME of leaf litter increased in response to an increase in flower litter biomass in the litter mixtures, and negative RME values for leaf litter decomposition were observed only in the two mixtures with the lowest flower litter biomass. As discussed above, the variation in positive values of RME in response to the increasing biomass of flower litter in mixtures points out to mechanisms that are generally attributed to the enhancing effects of LOM on the decomposition of ROM in litter mixtures, such as nutrient transfer or mining and priming effects (Guenet et al., 2010; Chen et al., 2014). The resource concentration hypothesis posits that resource quantity drives resource use efficiency (Charnov, 1976; Hambäck and Englund, 2005). The optimal foraging efficiency of microbial decomposers depends on mechanisms that maximize the balance between enzyme production and energy gain. We conjectured that the increase in flower litter biomass could have optimized the enzyme production and energy for the maintenance of metabolic processes, known as the substrate induction hypothesis (Allison et al., 2014; Schimel and Weintraub, 2003), which might have enhanced the leaf litter decomposition. Another possibility is the occurrence of priming effects mechanisms, such as co-metabolism, which posits that the decomposition of ROM



440 may be enhanced by LOM targeting enzymes capable of degrading the ROM, and/or that LOM decomposition may supply energy for microorganisms to produce extracellular enzymes capable of degrading ROM (Guenet et al., 2010). Such LMEs, resource-mediated mechanisms of flower litter on leaf litter decomposition in the terrestrial experiment may have acted in combination with the improvement of microenvironmental conditions promoted by the higher WHC of flower litter. However, such synergistic effects of flower litter on leaf litter turned into antagonistic effects when the biomass of flower litter was too low in mixtures, a pattern possible explained by preferential feeding of decomposers on flower litter and the limitation of substrate induction mechanisms when LOM is scarce in litter mixture (Cheng, 2009; Wang et al., 2015).

445 However, what could explain the unexpected variation of flower litter decomposition in the terrestrial experiment as the proportion of flower-to-leaf litter varied in litter mixtures? Although the decomposition of flower litter was enhanced in the presence of leaf litter irrespective of the flower-to-leaf litter proportion, these effects consistently weakened as the amount of leaf litter increased and the amount of flower litter decreased in the litter mixture. We conjectured that such results might have occurred due to the combination of two potential mechanisms. First, the presence of even a small amount of leaf litter could have an enhancing effect on the decomposition of flower litter if it is enough to meet the microbial decomposer community's demand for a specific limiting nutrient in the mixture. This, for example, might have occurred in litter mixtures due to the higher concentrations of micronutrients such as Ca, Mg, and Na, in leaf litter compared to flower litter. These micronutrients are considered important for litter decomposition in tropical forests (Kaspari et al., 2009), and their demand for decomposers is comparatively lower than macronutrients, such as N and P (Tyler, 2005). Therefore, even the low proportion of leaf litter might have been sufficient to meet the micronutrient demand of decomposers for decomposing flower litter. Secondly, the interaction between different types of litters can affect their decomposition through two non-mutually exclusive ways: through the effect a given litter can have on the other and/or through the response a given litter can exhibit to the interaction with another litter in the mixture. For example, labile litter can both expedite the breakdown of another litter (as discussed earlier), but also it may be more reactive to stimulation from another litter. This is because labile litter typically offers fewer resources that limit decomposers. Consequently, when a stimulus results from interactions with another litter, it's more likely to boost the decomposition of labile litter compared to refractory litter. We believe that this mechanism may have been relevant in determining the observed results in the terrestrial experiment, as the synergistic effects of the mixture of litter on flower litter decomposition increased with the rise in flower litter biomass in the mixture, while decreased with the increasing in the amount of leaf litter in the mixture.

465 The patterns resulting from the mixture of flower and leaf litter and the variation in the relative biomass of these litter types in the mixture were much less pronounced in the aquatic experiment. Although litter mixing also resulted in synergistic effects in aquatic experiment, such effects were weaker compared to those observed in the terrestrial experiment and did not consistently vary with the variation in the flower-to-leaf litter biomass proportion. Overall, these results support commonly reported observations in the literature regarding the effects of detritus mixture on decomposition, showing that LMEs are generally weaker or absent in aquatic ecosystems compared to terrestrial ones (Gessner et al., 2010). These results may be, in their entirety or in part, attributed to the fact that mechanisms potentially relevant for triggering LMEs in terrestrial ecosystem, such as moisture exchange between different types of litters and LMEs on physical properties of litter layer, lose relevance in the aquatic environment (Schmidt et al., 2011; Bengtsson et al., 2018). However, we may also have underestimated the LMEs in the aquatic experiment since, in this environment, a considerable portion of organic matter is leached from the litter and degraded in the water column in its dissolved form, which was not quantified in our experiment. In fact, our results showed that leaching is responsible to provoke 25% and 9% of mass loss on flower and leaf litter, respectively (Fig 1c). Several studies have demonstrated that litter-mixing interactions between dissolved organic matter from litter of different qualities also occur and accelerate decomposition in the water column (Farjalla et al., 2009; Fonte et al., 2013). Hence, forthcoming studies should explore the impact of flower and






leaf litter mixture on both particulate and dissolved organic matter decomposition to achieve a more comprehensive understanding of the LMEs of flower and leaf litter on decomposition.

480 Although the results of our experiment have demonstrated consistent patterns of flower and leaf litter mixture in the occurrence, magnitude, and direction of LMEs in decomposition, it is important to consider some caveats of our experiment. Firstly, we were unable to measure how the effect of the mixture and the variation in the proportion of flower and leaf litter affected the microbial community, which was the primary group of decomposers mediating our results. Therefore, our interpretations regarding how the mixture of flower and leaf litter affected the mechanisms arising from the interaction between litter and microbial decomposers were made considering the plausibility with which such mechanisms are presented in the literature on the subject.  
485 Secondly, although we made an effort to maintain environmental conditions similar to those observed in nature, laboratory conditions will always suppress features of the environment that may be relevant to the study at hand. Soil macrofauna was suppressed in our experiment, which could alter the realism of our results. However, the effect of the presence of soil invertebrates has been repeatedly shown to be an important factor in determining the occurrence of LMEs in decomposition. Even with the absence of soil macrofauna, our results showed consistent LMEs, and these were mediated by the relative biomass of flower and  
490 leaf litter in the terrestrial experiment. This suggests that the presence of soil macrofauna could reinforce or further amplify the observed patterns but probably would not alter them.

Massive flowering is a phenology pattern found in a range of species in different ecosystems, beyond the Bignoniaceae family (Conceição et al., 2013; Whigham, 2013; Zheng et al., 2020). This phenology pattern has the potential to be a significant source of LOM and could have secondary consequences on litter decomposition. The litter-mixing interactions between flower  
495 and leaf litter could be widespread in natural ecosystems, caused by differences in quality between flowers and leaves litter. These differences may be primarily attributed to variations in the form and function of these organs, creating a legacy effect for decomposition (Freschet et al., 2013; Cornelissen et al., 2023). In particular, our results indicate that species that present massive flowering phenology may represent key roles mediating the occurrence of temporal and/or spatial biogeochemical hotspots (Kuzakov, 2010), both through direct effects, where the flowers themselves generally represent a more labile and rapidly  
500 decomposing litter, thus being recycled more quickly and efficiently, and through indirect effects, where flower litter can interact with leaf litter complementarily stimulating the decomposition of both litters at the litter layer around the flowering tree. However, the results observed in the terrestrial experiment, which highlight that the magnitude of LMEs depends on the flower-to-leaf litter biomass proportion in the litter mixture, may represent the occurrence of a phenomenon analogous to the Janzen-Connell Hypothesis (Janzen, 1970; Connell, 1971). This hypothesis predicts that patterns of seed predation (Janzen, 1970) and herbivory  
505 on seedlings (Connell, 1971) are more intense near the parent tree because of resource concentration effects. In forests, the interaction between flower and leaf litter could occur beyond the taxonomic level, if differences of litter quality between flower and leaf litter were widespread. For example, in dense forests the presence of a few scattered trees presenting massive flowering can generate LMEs on leaf litter at either intra- or inter-specific levels (Fig. 5a). Analogously to what is expected for seed and seedlings distribution according to the Janzen-Connell Hypothesis, flower litter biomass should be more concentrated below the  
510 flowering tree and gradually decrease farther from it. On the other hand, leaf litter biomass would be more homogeneously distributed in the litter layer due to the contribution from the neighboring trees (Fig. 5b). This differential input of flower and leaf litterfall to the litter layer could generate a pattern of variation both in the concentration of flower litter and in the proportion of flower-to-leaf litter in the litter layer in relation to the blooming tree. Therefore, the rate of nutrient recycling due to decomposition should be higher near the flowering tree due to the decomposition of the high flower litter biomass itself and because, according  
515 to our results, synergistic effects of flower and leaf litter on the decomposition of both litter types are stronger in high flower-to-



leaf litter biomass proportions (Fig 5b). These potential effects would be more important in terrestrial ecosystems, both because the LMEs of litter mixing were responsive to variation in the flower-to-leaf litter biomass proportion only in the terrestrial experiment and because the arrangement of higher flower litter concentration around the flowering tree is more likely to be found in terrestrial ecosystems. However, the conjecture presented in this conceptual model must be properly tested to validate its expected results since our experiment, although allowing us to speculate on potential hypotheses and patterns, does not enable us to test or confirm them.



**Figure 5:** Conceptual model expanding the importance of flower and leaf litter spatial distribution and potential interactions in relation to the distance of the flowering tree in natural forests. a) Scattered distribution of massive flowering trees in a natural forest; Photo by S. Joseph Wright, Smithsonian Tropical Research Institute. b) We adapted a theoretical scheme based on the Janzen-Connell Hypothesis, which assumes that predation on seed and herbivory on seedlings decreases along the distance from the parent tree as seeds and seedlings become rarer on the forest floor. In our case, we assume that nutrient recycling, measured as litter decomposition rate, is a function of the absolute and relative biomass of flower and leaf litter in the litter layer. Absolute and relative biomass of flower litter decreases along the distance from the flowering tree. This is because the dispersion of flower litter to the litter layer is stronger near the flowering tree, becoming increasingly weaker with distance from the flowering tree, and the quantity of leaf litter from all neighboring tree species in the litter layer is independent of the distance from the flowering tree. Near the flowering tree, recycling through decomposition is expected to be higher due to the large amount of flower litter, which decomposes quickly because of its high quality, but also because the litter-mixing effects of the interaction between flower and leaf litter are stronger in the litter mixtures with a high proportion of flower-to-leaf litter. The results used to conjecture the predictions of this conceptual model are presented in Fig.4a.



## 5 Conclusions

Our findings highlight the importance litter from plant reproductive organs for LMEs in ecosystems, which could substantially contribute to changes in nutrient and carbon dynamics. Although, recent studies have evidenced about the role of reproductive organs to increase the decomposition of organic matter in the natural environment in both terrestrial (de Paz et al., 2018; Schmitt and Perfecto, 2020) and aquatic (Rezende et al., 2017) ecosystems, it is necessary to evaluate which mechanisms are driving the effects in each environment. We found evidence of the central role of complementarity in the occurrence, magnitude, and direction of these effects. In the terrestrial environment, the complementary effect was related to proportion of litter types in the litter mixture, indicating the potential importance of interplay between resource quality and quantity in determining niche partitioning among microbial decomposers. Future studies may investigate the generalities of flower and leaf litter on LMEs at intra- and inter-specific levels, as well as the potential role of flower litter in regulating LMEs on litter decomposition, across a large variety of plant species, an aspect that has been completely neglected in the literature about the effects of litter mixing on decomposition.

540  
545  
\$50

### Code availability

Code is available upon request from the corresponding author.

### Data availability

555 Data is available upon request from the corresponding author.

### Authors contribution

MIGA was responsible for running the experiments and writing the original draft. RDG, LSC, and AC were responsible for conceptualizing the experiments. BG and AC were responsible for the funding acquisition. MIGA and AC were responsible for the formal analyses. BG, MIGA, and AC were responsible for data interpretation. ELV was responsible for providing material and reagents for the chemical analyses. All co-authors contributed by reviewing and editing the submitted version.

### Competing interests

565 At least one of the (co-)authors is a member of the editorial board of Biogeosciences.

## References

- Allen, S. E., Grimshaw, H. M., Parkinson, J. A., and Quarmby, C.: Chemical analysis of ecological material., Blackwell, Oxford, 1974.
- Allison, S. D., Chacon, S. S., and German, D. P.: Substrate concentration constraints on microbial decomposition, *Soil Biol. Biochem.*, 79, 43–49, <https://doi.org/10.1016/j.soilbio.2014.08.021>, 2014.
- 570 Ashman, T.-L. and Schoen, D. J.: How long should flowers live?, *Nature*, 371, 788–791, <https://doi.org/10.1038/371788a0>, 1994.
- Aufdenkampe, A. K., Mayorga, E., Raymond, P. A., Melack, J. M., Doney, S. C., Alin, S. R., Aalto, R. E., and Yoo, K.: Riverine coupling of biogeochemical cycles between land, oceans, and atmosphere, *Front. Ecol. Environ.*, 9, 53–60, <https://doi.org/10.1890/100014>, 2011.
- 575 Barantal, S., Roy, J., Fromin, N., Schimann, H., and Hättenschwiler, S.: Long-term presence of tree species but not chemical



- diversity affect litter mixture effects on decomposition in a neotropical rainforest., *Oecologia*, 167, 241–252,  
<https://doi.org/10.1007/s00442-011-1966-4>, 2011.
- Barros, M. G.: Pollination ecology of *Tabebuia aurea* (Manso) Benth. & Hook. and *T. ochracea* (Cham.) Standl. (Bignoniaceae)  
580 in Central Brazil cerrado vegetation, *Rev. Bras. Botânica*, 24, 255–261, <https://doi.org/10.1590/S0100-84042001000300003>,  
2001.
- Basile-Doelsch, I., Balesdent, J., and Rose, J.: Are interactions between organic compounds and nanoscale weathering minerals  
the key drivers of carbon storage in soils?, *Environ. Sci. Technol.*, 49, 3997–3998, <https://doi.org/10.1021/acs.est.5b00650>, 2015.
- Batalha, M. A. and Mantovani, W.: Floristic composition of the cerrado in the Pé-de-Gigante Reserve (Santa Rita do Passa  
585 Quatro, southeastern Brazil), *Acta Bot. Brasilica*, 15, 289–304, <https://doi.org/10.1590/S0102-33062001000300001>, 2001.
- Bengtsson, M. M., Attermeyer, K., and Catalán, N.: Interactive effects on organic matter processing from soils to the ocean: are  
priming effects relevant in aquatic ecosystems?, *Hydrobiologia*, <https://doi.org/10.1007/s10750-018-3672-2>, 2018.
- Berenstecher, P., Araujo, P. I., and Austin, A. T.: Worlds apart: Location above- or below-ground determines plant litter  
decomposition in a semi-arid Patagonian steppe, *J. Ecol.*, 109, 2885–2896, <https://doi.org/https://doi.org/10.1111/1365->  
590 2745.13688, 2021.
- Boaventura, M. G., Villamil, N., Teixido, A. L., Tito, R., Vasconcelos, H. L., Silveira, F. A. O., and Cornelissen, T.: Revisiting  
florivory: an integrative review and global patterns of a neglected interaction, *New Phytol.*, 233, 132–144,  
<https://doi.org/10.1111/nph.17670>, 2022.
- Boyer, L., Pearson, R. G., Gessner, M. O., Barmuta, L. A., Ferreira, V., Graça, M. A. S., Dudgeon, D., Boulton, A. J., Callisto,  
595 M., Chauvet, E., Helson, J. E., Bruder, A., Albariño, R. J., Yule, C. M., Arunachalam, M., Davies, J. N., Figueroa, R., Flecker,  
A. S., Ramírez, A., Death, R. G., Iwata, T., Mathooko, J. M., Mathuriau, C., Gonçalves, J. F., Moretti, M. S., Jinggut, T.,  
Lamothe, S., M’Erimba, C., Ratnarajah, L., Schindler, M. H., Castela, J., Buria, L. M., Cornejo, A., Villanueva, V. D., and West,  
D. C.: A global experiment suggests climate warming will not accelerate litter decomposition in streams but might reduce carbon  
sequestration, *Ecol. Lett.*, 14, 289–294, <https://doi.org/10.1111/j.1461-0248.2010.01578.x>, 2011.
- 600 Boyero, L., Pérez, J., López-Rojo, N., Tonin, A. M., Correa-Araneda, F., Pearson, R. G., Bosch, J., Albariño, R. J., Anbalagan,  
S., Barmuta, L. A., Beesley, L., Burdon, F. J., Caliman, A., Callisto, M., Campbell, I. C., Cardinale, B. J., Casas, J. J., Chará-  
Serna, A. M., Ciapala, S., Chauvet, E., Colón-Gaud, C., Cornejo, A., Davis, A. M., Degebrodt, M., Dias, E. S., Díaz, M. E.,  
Douglas, M. M., Elozegi, A., Encalada, A. C., de Eyto, E., Figueroa, R., Flecker, A. S., Fleituch, T., Frainer, A., França, J. S.,  
García, E. A., García, G., García, P., Gessner, M. O., Giller, P. S., Gómez, J. E., Gómez, S., Gonçalves, J. F., Graça, M. A. S.,  
605 Hall, R. O., Hamada, N., Hepp, L. U., Hui, C., Imazawa, D., Iwata, T., Junior, E. S. A., Kariuki, S., Landeira-Dabarca, A., Leal,  
M., Lehosmaa, K., M’Erimba, C., Marchant, R., Martins, R. T., Masese, F. O., Camden, M., McKie, B. G., Medeiros, A. O.,  
Middleton, J. A., Muotka, T., Negishi, J. N., Pozo, J., Ramírez, A., Rezende, R. S., Richardson, J. S., Rincón, J., Rubio-Ríos, J.,  
Serrano, C., Shaffer, A. R., Sheldon, F., Swan, C. M., Tenkiano, N. S. D., Tieghe, S. D., Tolod, J. R., Vernasky, M., Watson, A.,  
Yegon, M. J., and Yule, C. M.: Latitude dictates plant diversity effects on instream decomposition, *Sci. Adv.*, 7, eabe7860,  
610 <https://doi.org/10.1126/sciadv.abe7860>, 2021.
- Brant, A. N. and Chen, H. Y. H.: Patterns and Mechanisms of Nutrient Resorption in Plants, *CRC. Crit. Rev. Plant Sci.*, 34, 471–  
486, <https://doi.org/10.1080/07352689.2015.1078611>, 2015.
- Cebrian, J.: Patterns in the Fate of Production in Plant Communities, *Am. Nat.*, 154, 449–468, <https://doi.org/10.1086/303244>,  
1999.
- 615 Charnov, E. L.: Optimal foraging, the marginal value theorem, *Theor. Popul. Biol.*, 9, 129–136, [20](https://doi.org/10.1016/0040-</a></p></div><div data-bbox=)



- 5809(76)90040-X, 1976.
- Chen, R., Senbayram, M., and Blagodatsky, S.: Soil C and N availability determine the priming effect : microbial N mining and stoichiometric decomposition theories, *Glob Chang Biol.*, 20, 2356–2367, <https://doi.org/10.1111/gcb.12475>, 2014.
- Cheng, W.: Rhizosphere priming effect: Its functional relationships with microbial turnover, evapotranspiration, and C-N budgets, *Soil Biol. Biochem.*, 41, 1795–1801, <https://doi.org/10.1016/j.soilbio.2008.04.018>, 2009.
- 620 Conceição, A. A., Alencar, T. G., Souza, J. M., Moura, A. D. C., and Silva, G. A.: Massive post-fire flowering events in a tropical mountain region of Brazil: high episodic supply of floral resources, *Acta Bot. Bra.*, 27, 847-850, <https://doi.org/10.1590/S0102-33062013000400025>, 2013.
- Connell, J.: On the role of the natural enemies in preventing competitive exclusion in some marine animals and in rain forest trees, In: Den Boer, P. J. and Gradwell, G.R. (eds) *Dynamics of Populations*, Centre for Agricultural Publishing and Documentation, Wageningen, The Netherlands, pp 298-312, 1971.
- 625 Cornelissen, J. H. C., Cornwell, W. K., Freschet, G. T., Weedon, J. T., Berg, M. P., and Zanne, A. E.: Coevolutionary legacies for plant decomposition, *Trends Ecol. Evol.*, 38, 44–54, <https://doi.org/10.1016/j.tree.2022.07.008>, 2023.
- Crutsinger, G. M., Sanders, N. J., and Classen, A. T.: Comparing intra- and inter-specific effects on litter decomposition in an old-field ecosystem, *Basic Appl. Ecol.*, 10, 535–543, <https://doi.org/10.1016/j.baae.2008.10.011>, 2009.
- 630 Davidson, E. A. and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change., *Nature*, 440, 165–173, <https://doi.org/10.1038/nature04514>, 2006.
- Dearden, F. M., Dehlin, H., Wardle, D. A., and Nilsson, M.-C.: Changes in the ratio of twig to foliage in litterfall with species composition, and consequences for decomposition across a long term chronosequence, *Oikos*, 115, 453–462, <https://doi.org/10.1111/j.2006.0030-1299.15354.x>, 2006.
- 635 Dignac, M.-F., Derrien, D., Barré, P., Barot, S., Cécillon, L., Chenu, C., Chevallier, T., Freschet, G. T., Garnier, P., Guenet, B., Hedde, M., Klumpp, K., Lashermes, G., Maron, P.-A., Nunan, N., Roumet, C., and Basile-Doelsch, I.: Increasing soil carbon storage: mechanisms, effects of agricultural practices and proxies. A review, *Agron. Sustain. Dev.*, 37, 14, <https://doi.org/10.1007/s13593-017-0421-2>, 2017.
- 640 Djukic, I., Kepfer-Rojas, S., Schmidt, I., Larsen, K., Beier, C., Berg, B., Verheyen, K., Caliman, A., Paquette, A., Gutiérrez, A., Humber, A., Valdecantos, A., Petraglia, A., Alexander, H., Augustaitis, A., Saillard, A., Fernández, A., Sousa, A., Lillebø, A. I., and Tóth, Z.: Early stage litter decomposition across biomes, *Sci. Total Environ.*, 628–629, 1369–1394, <https://doi.org/10.1016/j.scitotenv.2018.01.012>, 2018.
- Epps, K. Y., Comerford, N. B., Reeves James B., I. I. I., Cropper Wendell P., J., and Araujo, Q. R.: Chemical diversity – highlighting a species richness and ecosystem function disconnect, *Oikos*, 116, 1831–1840, <https://doi.org/10.1111/j.0030-1299.2007.15853.x>, 2007.
- 645 Farjalla, V. F., Marinho, C. C., Faria, B. M., Amado, A. M., Esteves, F. de A., Bozelli, R. L., and Giroldo, D.: Synergy of Fresh and Accumulated Organic Matter to Bacterial Growth, *Microb. Ecol.*, 57, 657–666, <https://doi.org/10.1007/s00248-008-9466-8>, 2009.
- 650 Fassbender, H. .: Simultane P-Bestimmung in N-Kjeldahl-Ausfischluß von Bodenproben., *Die Phosphorsäure*, 30, 44–53, 1973.
- Finerty, G. E., de Bello, F., Bílá, K., Berg, M. P., Dias, A. T. C., Pezzatti, G. B., and Moretti, M.: Exotic or not, leaf trait dissimilarity modulates the effect of dominant species on mixed litter decomposition, *J. Ecol.*, 104, 1400–1409, <https://doi.org/10.1111/1365-2745.12602>, 2016.
- Fonte, E. S., Amado, A. M., Meirelles-Pereira, F., Esteves, F. A., Rosado, A. S., and Farjalla, V. F.: The Combination of



- 655 Different Carbon Sources Enhances Bacterial Growth Efficiency in Aquatic Ecosystems, *Microb. Ecol.*, 66, 871–878, <https://doi.org/10.1007/s00248-013-0277-1>, 2013.  
Freeman, C. E., Worthington, R. D., and Jackson, M. S.: Floral Nectar Sugar Compositions of Some South and Southeast Asian Species, *Biotropica*, 23, 568–574, <https://doi.org/10.2307/2388394>, 1991.  
Freschet, G. T., Cornelissen, J. H. C., van Logtestijn, R. S. P., and Aerts, R.: Substantial nutrient resorption from leaves, stems  
660 and roots in a subarctic flora: what is the link with other resource economics traits?, *New Phytol.*, 186, 879–889, <https://doi.org/10.1111/j.1469-8137.2010.03228.x>, 2010.  
Freschet, G. T., Aerts, R., and Cornelissen, J. H. C.: A plant economics spectrum of litter decomposability, *Funct. Ecol.*, 26, 56–65, <https://doi.org/10.1111/j.1365-2435.2011.01913.x>, 2012.  
Freschet, G. T., Cornwell, W. K., Wardle, D. A., Elumeeva, T. G., Liu, W., Jackson, B. G., Onipchenko, V. G., Soudzilovskaia,  
665 N. A., Tao, J., and Cornelissen, J. H. C.: Linking litter decomposition of above- and below-ground organs to plant–soil feedbacks worldwide, *J. Ecol.*, 101, 943–952, <https://doi.org/10.1111/1365-2745.12092>, 2013.  
Galetto, L. and Bernardello, G.: Floral nectaries, nectar production dynamics and chemical composition in six *Ipomoea* species (*Convolvulaceae*) in relation to pollinators, *Ann. Bot.*, 94, 269–280, <https://doi.org/10.1093/aob/mch137>, 2004.  
García-Palacios, P., Mckie, B. G., Handa, I. T., Frainer, A., and Hättenschwiler, S.: The importance of litter traits and  
670 decomposers for litter decomposition: A comparison of aquatic and terrestrial ecosystems within and across biomes, *Funct. Ecol.*, 30, <https://doi.org/10.1111/1365-2435.12589>, 2016.  
Gartner, T. B. and Cardon, Z. G.: Decomposition dynamics in mixed-species leaf litter, *Oikos*, 104, 230–246, <https://doi.org/10.1111/j.0030-1299.2004.12738.x>, 2004.  
German, D. P., Chacon, S. S., and Allison, S. D.: Substrate concentration and enzyme allocation can affect rates of microbial  
675 decomposition, *Ecology*, 92, 1471–1480, <https://doi.org/10.1890/10-2028.1>, 2011.  
Gessner, M. O., Swan, C. M., Dang, C. K., McKie, B. G., Bardgett, R. D., Wall, D. H., and Hättenschwiler, S.: Diversity meets decomposition, *Trends Ecol. Evol.*, 25, 372–380, <https://doi.org/10.1016/j.tree.2010.01.010>, 2010.  
Goering, H.K., and Van Soest, P. J.: Forage fiber analysis (Apparatus, reagents, procedures and some applications), 379 pp., 1970.
- 680 Graça, M., Bärlocher, F., and Gessner, M.: *Methods to Study Litter Decomposition*, 1st edn, Springer, The Netherlands, <https://doi.org/10.1007/1-4020-3466-0>, 2005.  
Gripp, A. R., Esteves, F. de A., Carneiro, L. S., Guariento, R. D., Figueiredo-Barros, M. P., Coq, S., Milcu, A., and Caliman, A.: Weak to no effects of litter biomass and mixing on litter decomposition in a seasonally dry tropical forest, *Pedobiologia (Jena)*, 68, 20–23, <https://doi.org/https://doi.org/10.1016/j.pedobi.2018.02.003>, 2018.
- 685 Guenet, B., Danger, M., Abbadie, L., and Lacroix, G.: Priming effect: Bridging the gap between terrestrial and aquatic ecology, *Ecology*, 91, 2850–2861, <https://doi.org/10.1890/09-1968.1>, 2010.  
Hambäck, P. A. and Englund, G.: Patch area, population density and the scaling of migration rates: the resource concentration hypothesis revisited, *Ecol. Lett.*, 8, 1057–1065, <https://doi.org/10.1111/j.1461-0248.2005.00811.x>, 2005.  
Handa, I. T., Aerts, R., Berendse, F., Berg, M. P., Bruder, A., Butenschoen, O., Chauvet, E., Gessner, M. O., Jabiol, J.,  
690 Makkonen, M., McKie, B. G., Malmqvist, B., Peeters, E. T. H. M., Scheu, S., Schmid, B., van Ruijven, J., Vos, V. C. A., and Hättenschwiler, S.: Consequences of biodiversity loss for litter decomposition across biomes., *Nature*, 509, 218–221, <https://doi.org/10.1038/nature13247>, 2014.  
Van Handel, E.: Direct microdetermination of sucrose., *Anal. Biochem.*, 22, 280–283, 1968.



- Hättenschwiler, S. and Jørgensen, H. B.: Carbon quality rather than stoichiometry controls litter decomposition in a tropical rain forest, *J. Ecol.*, 98, 754–763, <https://doi.org/10.1111/j.1365-2745.2010.01671.x>, 2010.
- Hättenschwiler, S., Tiunov, A. V. and Scheu, S.: Biodiversity and Litter Decomposition in Terrestrial Ecosystems, *Annu. Rev. Ecol. Evol. Syst.*, 36, 191–218, <https://doi.org/10.1146/annurev.ecolsys.36.112904.151932>, 2005.
- Hengl, T., Mendes de Jesus, J., Heuvelink, G. B. M., Ruiperez Gonzalez, M., Kilibarda, M., Blagotić, A., Shangguan, W., Wright, M. N., Geng, X., Bauer-Marschallinger, B., Guevara, M. A., Vargas, R., MacMillan, R. A., Batjes, N. H., Leenaars, J. G. B., Ribeiro, E., Wheeler, I., Mantel, S., and Kempen, B.: SoilGrids250m: Global gridded soil information based on machine learning, *PLoS One*, 12, e0169748, <https://doi.org/10.1371/journal.pone.0169748>, 2017.
- Hou, S.-L. and Lü, X.-T.: Mixing effects of litter decomposition at plant organ and species levels in a temperate grassland, *Plant Soil*, 459, 387–396, <https://doi.org/10.1007/s11104-020-04773-0>, 2021.
- Jackson, B. G., Peltzer, D. A., and Wardle, D. A.: Are functional traits and litter decomposability coordinated across leaves, twigs and wood? A test using temperate rainforest tree species, *Oikos*, 122, 1131–1142, <https://doi.org/10.1111/j.1600-0706.2012.00056.x>, 2013.
- Janzen, D. H. Herbivores and number of tree species in tropical forests. *American Naturalist*, *Am. Nat.*, 104, 501–528, <https://doi.org/10.1086/282687>, 1970.
- Jolliffe, P. A.: The replacement series, *J. Ecol.*, 88, 371–385, <https://doi.org/10.1046/j.1365-2745.2000.00470.x>, 2000.
- Jones, A. R., Dalal, R. C., Gupta, V. V. S. R., Schmidt, S., Allen, D. E., Jacobsen, G. E., Bird, M., Grandy, A. S., and Sanderman, J.: Molecular complexity and diversity of persistent soil organic matter, *Soil Biol. Biochem.*, 184, 109061, <https://doi.org/10.1016/j.soilbio.2023.109061>, 2023.
- Kaspari, M., Garcia, M. N., Harms, K. E., Santana, M., Wright, S. J., and Yavitt, J. B.: Multiple nutrients limit litterfall and decomposition in a tropical forest, *Ecol. Lett.*, 11, 35–43, <https://doi.org/10.1111/j.1461-0248.2007.01124.x>, 2008.
- Kaspari, M., Yanoviak, S. P., Dudley, R., Yuan, M., and Clay, N. A.: Sodium shortage as a constraint on the carbon cycle in an inland tropical rainforest, *Proc. Natl. Acad. Sci.*, 106, 19405–19409, <https://doi.org/10.1073/pnas.0906448106>, 2009.
- Kuzyakov, Y.: Review: Factors affecting rhizosphere priming effects, *J. Plant Nutr. Soil Sci.*, 165, 382–396, [https://doi.org/10.1002/1522-2624\(200208\)165:4<382::AID-JPLN382>3.0.CO;2-#](https://doi.org/10.1002/1522-2624(200208)165:4<382::AID-JPLN382>3.0.CO;2-#), 2002.
- Kuzyakov, Y.: Priming effects: Interactions between living and dead organic matter, *Soil Biol. Biochem.*, 42, 1363–1371, <https://doi.org/10.1016/j.soilbio.2010.04.003>, 2010.
- Kuzyakov, Y. and Blagodatskaya, E.: Microbial hotspots and hot moments in soil: Concept & review, *Soil Biol. Biochem.*, 83, 184–199, <https://doi.org/10.1016/j.soilbio.2015.01.025>, 2015.
- Lecerf, A., Marie, G., Kominoski, J. S., LeRoy, C. J., Bernadet, C., and Swan, C. M.: Incubation time, functional litter diversity, and habitat characteristics predict litter-mixing effects on decomposition, *Ecology*, 92, 160–169, <https://doi.org/10.1890/10-0315.1>, 2011.
- Lidman, J., Jonsson, M., Burrows, R. M., Bundschuh, M., and Sponseller, R. A.: Composition of riparian litter input regulates organic matter decomposition: Implications for headwater stream functioning in a managed forest landscape, *Ecol. Evol.*, 7, 1068–1077, <https://doi.org/10.1002/ece3.2726>, 2017.
- Lin, D., Pang, M., Fanin, N., Wang, H., Qian, S., Zhao, L., Yang, Y., Mi, X., and Ma, K.: Fungi participate in driving home-field advantage of litter decomposition in a subtropical forest, *Plant Soil*, 434, 467–480, <https://doi.org/10.1007/s11104-018-3865-5>, 2019.
- Liu, J., Liu, X., Song, Q., Compson, Z. G., LeRoy, C. J., Luan, F., Wang, H., Hu, Y., and Yang, Q.: Synergistic effects: a



- common theme in mixed-species litter decomposition, *New Phytol.*, 227, 757–765, <https://doi.org/10.1111/nph.16556>, 2020.
- Loreau, M.: Separating Sampling and Other Effects in Biodiversity Experiments, *Oikos*, 82, 600–602, <https://doi.org/10.2307/3546381>, 1998.
- 735 Lorenzi, H.: *Árvores brasileiras: manual de identificação e cultivo de plantas arbóreas nativas do Brasil*, Instituto Plantarum de Estudos da Flora, 1st edn. Instituto Plantarum de Estudos da Flora, Brazil, 1992.
- Madritch, M. D. and Hunter, M. D.: Phenotypic diversity and litter chemistry affect nutrient dynamics during litter decomposition in a two species mix, *Oikos*, 105, 125–131, <https://doi.org/10.1111/j.0030-1299.2004.12760.x>, 2004.
- 740 Makkonen, M., Berg, M. P., Handa, I. T., Hättenschwiler, S., van Ruijven, J., van Bodegom, P. M., and Aerts, R.: Highly consistent effects of plant litter identity and functional traits on decomposition across a latitudinal gradient, *Ecol. Lett.*, 15, 1033–1041, <https://doi.org/10.1111/j.1461-0248.2012.01826.x>, 2012.
- Makkonen, M., Berg, M. P., van Logtestijn, R. S. P., van Hal, J. R., and Aerts, R.: Do physical plant litter traits explain non-additivity in litter mixtures? A test of the improved microenvironmental conditions theory, *Oikos*, 122, 987–997, <https://doi.org/10.1111/j.1600-0706.2012.20750.x>, 2013.
- 745 Manzoni, S. and Porporato, A.: Soil carbon and nitrogen mineralization: Theory and models across scales, *Soil Biol. Biochem.*, 41, 1355–1379, <https://doi.org/10.1016/j.soilbio.2009.02.031>, 2009.
- Mccall, A. and Irwin, R.: Florivory: The intersection of pollination and herbivory, *Ecol. Lett.*, 9, 1351–1365, <https://doi.org/10.1111/j.1461-0248.2006.00975.x>, 2007.
- 750 McClain, M. E., Boyer, E. W., Dent, C. L., Gergel, S. E., Grimm, N. B., Groffman, P. M., Hart, S. C., Harvey, J. W., Johnston, C. A., Mayorga, E., McDowell, W. H., and Pinay, G.: Biogeochemical Hot Spots and Hot Moments at the Interface of Terrestrial and Aquatic Ecosystems, *Ecosystems*, 6, 301–312, <https://doi.org/10.1007/s10021-003-0161-9>, 2003.
- Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.-S., Cheng, K., Das, B. S., Field, D. J., Gimona, A., Hedley, C. B., Hong, S. Y., Mandal, B., Marchant, B. P., Martin, M., McConkey, B. G.,
- 755 Mulder, V. L., O'Rourke, S., Richer-de-Forges, A. C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C.-C., Vågen, T.-G., van Wesemael, B., and Winowiecki, L.: Soil carbon 4 per mille, *Geoderma*, 292, 59–86, <https://doi.org/10.1016/j.geoderma.2017.01.002>, 2017.
- Morris, D. L.: Quantitative determination of carbohydrates with Dreywood's anthrone reagent., *Science*, 107, 111–114, 1948.
- de Paz, M., Gobbi, M. E., and Raffaele, E.: Fallen fruits stimulate decomposition of leaf litter of dominant species in NW
- 760 Patagonia shrublands, *Plant Soil*, 425, 433–440, <https://doi.org/10.1007/s11104-018-3590-0>, 2018.
- Pérez-Harguindeguy, N., Díaz, S., Garnier, E., Lavorel, S., Poorter, H., Jaureguiberry, P., Bret-Harte, M. S., Cornwell, W. K., Craine, J. M., Gurvich, D. E., Urcelay, C., Veneklaas, E. J., Reich, P. B., Poorter, L., Wright, I. J., Ray, P., Enrico, L., Pausas, J. G., de Vos, A. C., Buchmann, N., Funes, G., Quétier, F., Hodgson, J. G., Thompson, K., Morgan, H. D., ter Steege, H., Sack, L., Blonder, B., Poschlod, P., Vaieretti, M. V., Conti, G., Staver, A. C., Aquino, S., and Cornelissen, J. H. C.: New handbook for standardised measurement of plant functional traits worldwide, *Aust. J. Bot.*, 61, 167–234, 2013.
- 765 Porre, R. J., van der Werf, W., De Deyn, G. B., Stomph, T. J., and Hoffland, E.: Is litter decomposition enhanced in species mixtures? A meta-analysis, *Soil Biol. Biochem.*, 145, 107791, <https://doi.org/10.1016/j.soilbio.2020.107791>, 2020.
- Rezende, R., R.S. Correia, P., Goncalves Jr, J., and Santos, A.: Organic matter dynamics in a savanna transition riparian zone: Input of plant reproductive parts increases leaf breakdown process, *J. Limnol.*, 76, 514–523, <https://doi.org/10.4081/jlimnol.2017.1601>, 2017.
- 770 Ribeiro, S. P. and Brown, V. K.: Prevalence of monodominant vigorous tree populations in the tropics: herbivory pressure on





- Tabebuia species in very different habitats, *J. Ecol.*, 94, 932–941, <https://doi.org/10.1111/j.1365-2745.2006.01133.x>, 2006.
- Robertson, G. P., Coleman, D., Bledsoe, C. S., and Sollins, P.: Standard soil methods for long-term ecological research, 1st edn. Oxford University Press, New York, 1999.
- 775 Roddy, A. B., Jiang, G.-F., Cao, K., Simonin, K. A., and Brodersen, C. R.: Hydraulic traits are more diverse in flowers than in leaves, *New Phytol.*, 223, 193–203, <https://doi.org/10.1111/nph.15749>, 2019.
- Santos e Silva, C. M., Lúcio, P. S., and Spyrides, M. H. C.: Distribuição espacial da precipitação sobre o Rio Grande do Norte: estimativas via satélites e medidas por pluviômetros, *Rev. Bras. Meteorol.*, 27, 337–346, <https://doi.org/10.1590/S0102-77862012000300008>, 2012.
- 780 Sarruge, J. . and Haag, H. P.: Análise química das plantas., 1st edn. Livrocetes, Piracicaba, 1974.
- Sayer, E. J., Powers, J. S., and Tanner, E. V. J.: Increased Litterfall in Tropical Forests Boosts the Transfer of Soil CO<sub>2</sub> to the Atmosphere, *PLoS One*, 2, e1299, <https://doi.org/10.1371/journal.pone.0001299>, 2007.
- Schimel, D. S., House, J. I., Hibbard, K. A., Bousquet, P., Ciais, P., Peylin, P., Braswell, B. H., Apps, M. J., Baker, D., Bondeau, A., Canadell, J., Churkina, G., Cramer, W., Denning, A. S., Field, C. B., Friedlingstein, P., Goodale, C., Heimann, M.,
- 785 Houghton, R. A., Melillo, J. M., Moore, B., Murdiyarso, D., Noble, I., Pacala, S. W., Prentice, I. C., Raupach, M. R., Rayner, P. J., Scholes, R. J., Steffen, W. L., and Wirth, C.: Recent patterns and mechanisms of carbon exchange by terrestrial ecosystems, *Nature*, 414, 169–172, <https://doi.org/10.1038/35102500>, 2001.
- Schimel, J. P. and Weintraub, M. N.: The implications of exoenzyme activity on microbial carbon and nitrogen limitation in soil: A theoretical model, *Soil Biol. Biochem.*, 35, 549–563, [https://doi.org/10.1016/S0038-0717\(03\)00015-4](https://doi.org/10.1016/S0038-0717(03)00015-4), 2003.
- 790 Schindler, M. and Gessner, M. O.: Functional leaf traits and biodiversity effects on litter decomposition in a stream, *Ecology*, 90, 1641–1649, <https://doi.org/10.1890/0012-9658-91.6.1869>, 2009.
- Schmidt, M. W. I., Torn, M. S., Abiven, S., Dittmar, T., Guggenberger, G., Janssens, I. A., Kleber, M., Kögel-Knabner, I., Lehmann, J., Manning, D. A. C., Nannipieri, P., Rasse, D. P., Weiner, S., and Trumbore, S. E.: Persistence of soil organic matter as an ecosystem property, *Nature*, 478, 49, <https://doi.org/10.1038/nature10386>, 2011.
- 795 Schmitt, L. and Perfecto, I.: Who gives a flux? Synchronous flowering of *Coffea arabica* accelerates leaf litter decomposition, *Ecosphere*, 11, e03186, <https://doi.org/https://doi.org/10.1002/ecs2.3186>, 2020.
- Smith, V. C. and Bradford, M. A.: Do non-additive effects on decomposition in litter-mix experiments result from differences in resource quality between litters?, *Oikos*, 102, 235–242, <https://doi.org/10.1034/j.1600-0706.2003.12503.x>, 2003.
- Stamp, N.: Out Of The Quagmire Of Plant Defense Hypotheses, *Q. Rev. Biol.*, 78, 23–55, <https://doi.org/10.1086/367580>, 2003.
- 800 Swift, M. J., Heal, O. W., Anderson, J. M., and Anderson, J. M.: Decomposition in terrestrial ecosystems, 5th edn. University of California Press, Berkeley, 1979.
- Tian, H., Lu, C., Yang, J., Banger, K., Huntzinger, D. N., Schwalm, C. R., Michalak, A. M., Cook, R., Ciais, P., Hayes, D., Huang, M., Ito, A., Jain, A. K., Lei, H., Mao, J., Pan, S., Post, W. M., Peng, S., Poulter, B., Ren, W., Ricciuto, D., Schaefer, K., Shi, X., Tao, B., Wang, W., Wei, Y., Yang, Q., Zhang, B., and Zeng, N.: Global patterns and controls of soil organic carbon
- 805 dynamics as simulated by multiple terrestrial biosphere models: Current status and future directions, *Global Biogeochem. Cycles*, 29, 775–792, <https://doi.org/10.1002/2014GB005021>, 2015.
- Tranvik, L. J., Downing, J. A., Cotner, J. B., Loiselle, S. A., Striegl, R. G., Ballatore, T. J., Dillon, P., Finlay, K., Fortino, K., Knoll, L. B., Kortelainen, P. L., Kutser, T., Larsen, S., Laurion, I., Leech, D. M., Leigh McCallister, S., McKnight, D. M., Melack, J. M., Overholt, E., Porter, J. A., Prairie, Y., Renwick, W. H., Roland, F., Sherman, B. S., Schindler, D. W., Sobek, S.,
- 810 Tremblay, A., Vanni, M. J., Verschoor, A. M., Von Wachenfeldt, E., and Weyhenmeyer, G. A.: Lakes and reservoirs as



regulators of carbon cycling and climate, *Limnol. Oceanogr.*, 54, 2298–2314, [https://doi.org/10.4319/lo.2009.54.6\\_part\\_2.2298](https://doi.org/10.4319/lo.2009.54.6_part_2.2298), 2009.

Tyler, G.: Changes in the concentrations of major, minor and rare-earth elements during leaf senescence and decomposition in a *Fagus sylvatica* forest, *For. Ecol. Manage.*, 206, 167–177, <https://doi.org/10.1016/j.foreco.2004.10.065>, 2005.

815 Violle, C., Thuiller, W., Mouquet, N., Munoz, F., Kraft, N. J. B., Cadotte, M. W., Livingstone, S. W., and Mouillot, D.: Functional Rarity: The Ecology of Outliers, *Trends Ecol. Evol.*, 32, 356–367, <https://doi.org/10.1016/j.tree.2017.02.002>, 2017.

Wang, J., Liu, L., Wang, X., and Chen, Y.: The interaction between abiotic photodegradation and microbial decomposition under ultraviolet radiation, *Glob. Chang. Biol.*, 21, 2095–2104, <https://doi.org/10.1111/gcb.12812>, 2015.

820 Waring, B. G.: Exploring relationships between enzyme activities and leaf litter decomposition in a wet tropical forest, *Soil Biol. Biochem.*, 64, 89–95, <https://doi.org/10.1016/j.soilbio.2013.04.010>, 2013.

Whigham, A. E.: Senescent Neotropical flowers (Lecythidaceae) offer a rich nutrient source to ground-foraging arthropods, *J. Torrey Bot. Soc.*, 140, 31–40, <https://doi.org/10.3159/TORREY-D-12-00028.1>, 2013.

Zhang, X., Zhang, Y., Jiang, S., Song, C., Zhang, J., and Mao, R.: Dominant species and evenness level co-regulate litter mixture decomposition in a boreal peatland, *Plant Soil*, 474, 423–436, <https://doi.org/10.1007/s11104-022-05346-z>, 2022.

825 Zhao, W., van Logtestijn, R. S. P., van Hal, J. R., Dong, M., and Cornelissen, J. H. C.: Non-additive effects of leaf and twig mixtures from different tree species on experimental litter-bed flammability, *Plant Soil*, 436, 311–324, <https://doi.org/10.1007/s11104-019-03931-3>, 2019.

Zheng, X., Lin, S., Fu, H., Wan, Y., and Ding, Y.: The Bamboo Flowering Cycle Sheds Light on Flowering Diversity, *Front. Plant Sci.*, 11, <https://www.frontiersin.org/articles/10.3389/fpls.2020.00381>, 2020.

830