

Will ~~Are~~ rivers becoming more intermittent in France? Learning from an extended set of hydrological-climate projections ~~based on the Coupled Model Intercomparison Project phase 5 (CMIP5)~~

Tristan Jaouen¹, Lionel Benoit², Louis Héraut¹, Eric Sauquet¹

¹INRAE Lyon-Grenoble Auvergne-Rhône-Alpes, RiverLy, Villeurbanne, France

²INRAE PACA, Biostatistique et processus SPatiaux, Avignon, France

Correspondence to: Tristan Jaouen (tristan.jaouen@inrae.fr) and Eric Sauquet (eric.sauquet@inrae.fr)

Abstract

This study aims to assess the changes in the intermittency of river flows across France in the context of climate change. Projections of flow intermittence are derived from the results of the Explore2 project, which is the latest national study that proposes a wide range of potential hydrological futures for the 21st century. The multi-model approach developed within the Explore2 project enables to characterize uncertainties in future flow intermittence. Combined with discrete observations of flow states, hydrological projections are post-processed to compute the daily probability of flow intermittency (PFI) on each element of the partition of France in Hydro-EcoRegions hydroecoregions (HER2).

The post-processing consists of-in calibrating logistic regressions between the historical flow states of the Observatoire National des Étiages National Low-Flow Observatory (ONDE) network and the flow data simulated by the hydrological models (HMs) involved in Explore2 projected-run with the SAFRANafran atmospheric reanalysis as inputs. After calibration, these regressions are used to project daily PFIs for the whole-of-theentire 21st century, based on flow simulations from five hydrological modelsHMs driven by up to 17 climate projections under RCP 2.6, 4.5, and 8.5 climate change scenarios.

The results show good agreement among the hydrological modelsHMs regarding the increase in flow intermittency under RCP 4.5 and 8.5. The changes-projected increase in mean daily PFI between July and October-and-as well as the shifts-in of the first and last days when PFI exceeds 20%-,% both suggest a gradual intensification and extension of dry spells throughout the century. The southern regions of France are likely to experience greater increases in runoff intermittency than the northern regions,- and mountainous regions such as the Alps and the Pyrenees are likely to experience changes in their dynamics of intermittence with a reduction of winter intermittence and the apparition or increase of summer intermittence. The uncertainty of these projected changes is larger in northern France due to greater intermodel variability in this region. Uncertainty is greater in northern France, due to the variability of rainfall. Mountainous regions such as the Alps and the Pyrenees are likely to experience changes in the dynamics of snowmelt and groundwater recharge, which could lead to changes in their runoff regimes.

Commenté [T1]: RC3.1

Commenté [T2]: RC3.12

Commenté [T3]: RC3.12

Commenté [T4]: RC3.11

Commenté [T5]: RC3.11

Commenté [T6]: RC3.11

Commenté [T7]: RC2.3

30 1 Introduction

35 Intermittent Rivers and Ephemeral Streams (IRES) are watercourses characterized by the absence of continuous year-round water flow (Sefton et al., 2019). While water management practices can exacerbate flow intermittence by altering runoff patterns, climatic factors - particularly aridity - remain the primary determinants of these systems (Addor et al., 2018; Hammond and Fleming, 2021; Sauquet et al., 2021; Zipper et al., 2021). In recent decades, climate change has intensified these dynamics, significantly impacting the global water cycle: summer precipitations tend to decrease, and rising temperatures have increased evapotranspiration and driven more frequent heatwaves (Douville et al., 2021). These changes have led to more severe low-flow periods and more frequent and widespread dry spells in Western Europe (Delso et al., 2017; Trambly et al., 2020; Vicente-Serrano et al., 2020). Meanwhile, IRES play a critical role as interfaces between terrestrial and aquatic ecosystems. Altered patterns of flow interruption can therefore compromise water quantity and quality in downstream perennial rivers, increase extinction risks for specialized species, and disrupt organic matter, nutrient, and sediment cycles (Bertassello et al., 2022; Gallart et al., 2012; Giezendanner et al., 2021; Finn et al., 2011; Meyer et al., 2007; Sarremejane et al., 2017; Larned et al., 2010). Hence, a better understanding of IRES behaviours and their interaction with perennial streams is needed to assess the impacts of ongoing and future changes on river ecosystems (Döll and Schmied, 2012; Jaeger et al., 2014; Pumo et al., 2016; Leigh and Datry, 2017), and to address water management challenges (Acuña et al., 2014).

45 Globally, IRES account for nearly two-thirds of river networks (Schneider et al., 2017; Messenger et al., 2021), with this proportion reaching 39% in France (Snelder et al., 2013). Their spatial and temporal dynamics can be characterized using conceptual models of network contraction (Shaw et al., 2017; Botter and Durighetto, 2020) or through semi-conceptual and machine learning frameworks that integrate hydroclimatic and physiographic factors (Lapides et al., 2021; Durighetto et al., 2022). The strongest predictors of flow intermittence are aridity indices (Vicente-Serrano et al., 2019 ; Jaeger et al., 2019). However, local variations in permeability, lithology and topography also influence intermittence patterns in specific regions, including West Africa (Yu et al., 2018; Belemougri et al., 2021), northern Spain (González-Ferreras and Barquín, 2017) and northwestern Australia (Bourke et al., 2021). Studying intermittence at a large scale therefore requires considering the diversity of hydrological processes at play, particularly when multiple watersheds exhibit diverse sensitivities to hydroclimatic conditions. This variability highlights the need for robust methodologies to assess intermittence at national to (semi-)continental scales (Addor et al., 2018; Hammond and Fleming, 2021; Sando et al., 2022; Döll et al., 2024).

60 Understanding the drivers of river intermittence is also crucial for analysing historical trends and projecting future changes in IRES behaviour. Climate change has already expanded the spatial and temporal extent of intermittent flows worldwide (Zipper et al., 2021; Jaeger et al., 2019; Sando et al., 2022; Gudmundsson et al., 2021), and the Mediterranean basin is particularly affected (Trambly et al., 2020; De Girolamo et al., 2022). However, interpreting climate projections and emission scenarios at the scale of headwater basins remains rare (Schneider et al., 2013). Indeed, many climate models have a coarse spatial

Commenté [T8]: RC2.1, RC3.2

Commenté [T9]: RC1.6

Commenté [T10]: RC1.6

Commenté [T11]: RC1.6

Commenté [T12]: RC1.6

resolution and high uncertainties emerge from multi-model ensembles (MMEs), which makes accurate predictions of future IRES changes difficult.

To address these challenges, France has implemented the Explore2 project (<https://entrepot.recherche.data.gouv.fr/dataverse/explore2>; Sauquet et al., in prep.) that uses a multi-model, multi-scenario approach to simulate hydrological conditions throughout for the 21st century. It assesses uncertainties at each step of the modelling process. However, the resolution of the climate projections involved in Explore2 is too coarse to accurately capture IRES dynamics. Hence, this study extends the results of Explore2 for purposes of generating flow intermittence projections for French headwater streams. To this end, we follow a statistical approach that estimates the daily Probability of Flow Intermittence (*PFI*) in small streams based on streamflow data simulated by Explore2 for larger perennial rivers. The estimated *PFI* serves as a proxy indicator for the intensity and duration of dry periods. The method is calibrated using field observations carried out regularly at more than 3200 upstream river sites prone to drying.

To achieve this, the present work builds on two key studies: Beaufort et al. (2018), which demonstrated consistent performance of *PFI* estimation across the heterogeneous climate of France using daily discharge, and Sauquet et al. (2021), which projected *PFI* under a limited set of climate projections with coarse spatial resolution. In contrast, this study incorporates a finer spatial resolution, a broader range of climate scenarios, and a wider variety of hydrological models thanks to the integration of Explore2 projections. It therefore addresses a critical research gap by generating projections of small stream intermittence across France and throughout the 21st century while quantifying the associated uncertainties.

The remainder of the paper is organized into five sections. The data used are outlined in the second section. Section 3 describes the statistical method linking *PFI* to hydrological projections. Section 4 presents the results, which focus on the mean daily *PFI* between July and October (*mPFI₇₋₁₀*) and the median of the first and last days (respectively, *Tf* and *Tl*) when *PFI* exceeds 20%. Finally, Section 5 discusses these results and Section 6 concludes the study.

Intermittent Rivers and Ephemeral Streams (IRES) are watercourses characterized by the absence of continuous year-round water flow (Sefton et al., 2019). Flow intermittence results from various factors such as limited rainfall or freezing conditions, or human-induced alterations of the runoff pattern. Climate is a primary driver of streamflow patterns, though geology and topography can also play significant roles in their development. Climate change is expected to modify the water cycle, with a significant reduction in summer precipitation and an increase in frequency of the heat waves in Western Europe (Vicente-Serrano et al., 2020), which is likely to result in more severe low flow periods and more frequent and widespread dry periods (Delso et al., 2017). Several features of flow intermittence such as its seasonality, duration, and spatial extent are likely to be altered, and a shift from perennial to intermittent flow may also affect many rivers (Van Meerveld et al., 2020). IRES play a central role as an interface between terrestrial and aquatic ecosystems. As a result, altered patterns of flow interruption can compromise water quantity and quality in downstream perennial rivers, endanger their specific biodiversity, and disrupt organic matter, nutrient, and sediment cycles, etc. (Meyer et al., 2007; Finn et al., 2011; Clarke et al., 2008; Gallart et al., 2012; Sarremejane et al.,

2017). A better understanding of the behaviour of IRES and their interaction with perennial streams is therefore necessary to assess the impact of ongoing and future changes on river ecosystems (Döll and Schmied, 2012; Jaeger et al., 2014; Pumo et al., 2016; Leigh and Datry, 2017), and to address water management challenges (Acuña et al., 2014). Note that most past studies addressed the impact of climate change on flow intermittence at the local scale (e.g. in southern Italy, cf. (De Girolamo et al., 2022)) so there is a need to place these results in a wider context and to determine the extent of the changes and their consequences for the main rivers downstream.

To update the knowledge about the impact of climate change on water resources throughout its territory, France has recently implemented the project Explore2 (<https://entrepot.recherche.data.gouv.fr/dataverse/explore2>; Sauquet et al., in prep.). Through a multi-model and multi-scenario approach, Explore2 encompasses a wide range of possible hydrological futures for the whole 21st century, and therefore allows assessing the uncertainties arising at each step of the modelling process. However, the spatial resolution of the climate projections involved in Explore2 is too coarse to properly resolve IRES, in particular because IRES are usually small streams located in headwater catchments. To overcome this limitation and gain insight into IRES behaviour under climate change, the present study aims to extend the results of Explore2 to obtain projections of flow intermittence in headwater streams. We follow a statistical approach using streamflow data from large and perennial rivers to predict the daily Probability of Flow Intermittence (*PFI*). The *PFI* is defined as the proportion of headwater streams under drying conditions at a regional scale and is used as a proxy indicator for the intensity and duration of dry periods. Usually obtained in small watersheds (Sefton et al., 2019), it is calibrated using field campaigns specific to France, carried out regularly at more than 3200 upstream river sites prone to drying.

This approach has been initially developed and validated by Beaufort et al. (2018) using daily discharge and monthly flow intermittence observations over the period 2011–2017. Subsequently, this approach was leveraged by Sauquet et al. (2021) to project the *PFI* in a changing climate, but the spatial resolution of this “proof-of-concept” study was coarse and only a handful of climate scenarios were considered. The present study continues the enhancement on this *PFI* modelling framework, and in particular aims to increase the spatial resolution of future *PFI* projections, to consider a larger number of hydrological and climate projections (thanks to Explore2 datasets), and to carefully quantify the uncertainties associated with the projected changes in river flow intermittence over the 21st century across France.

The paper is organized in five parts. The data used are outlined in the second section. The statistical method linking *PFI* to hydrological projections is described in Section 3, with results presented in Section 4. The results will focus on the mean daily *PFI* between July and October (*mPFI_{July–Oct}*), and the median of the first and last days (respectively, *T_f* and *T_l*) with *PFI* exceeded 20%. Finally Section 5 discusses these results, and Section 6 concludes the study.

2 Data

2.1 Monitoring river flow intermittence

The French Biodiversity Office (OFB, <https://ofb.gouv.fr/>) initiated in 2012 the National Low-Flow Observatory (ONDE) in 2012 to gain a better understanding of low flows and intermittent rivers. ONDE is a stream intermittence monitoring network comprising 3248 observation sites strategically distributed across the French river network with the aim of characterizing the

135 occurrence and the intensity of summer (low-) flows (Nowak and Durozoi, 2012). The network has been designed to focus on
streams with a Strahler ~~orders~~ ranging from 1 to 4, which are prone to natural and/or anthropogenic intermittent flow
conditions. Most sites (85%) are located on “small streams” with a drainage area $\leq 100 \text{ km}^2$, 75% are situated on streams with
a Strahler index of 1 or 2, and 20% of the sites have a drainage area $\leq 10 \text{ km}^2$. (Appendix Fig. G1 and G3).

Commenté [T13]: RC1.2, RC1.3

Commenté [T14]: RC3.12

Commenté [T15]: RC3.5

140 A key feature of the ONDE network is that it involves the flow state of watercourses, which results from the visual inspection
of streams by OFB observers at each location, instead of a sensor-based measurement of flow rates. In the present study, two
binary categories are defined for ONDE observations: (1) "visible flows" and (2) "dry conditions", which gather non-visible
flows (standing water in isolated pools) and totally dry conditions (dry riverbed at or near the ONDE site). We use only data
from regular and structured field campaigns (referred to as “usual campaigns” in the ONDE nomenclature), during which t
145 ONDE sites used in this study are monitored ~~during field campaigns carried out~~ systematically in mainland France around the
25th day of each month from May to September. Here we use ONDE~~We consider~~ data from 2012 to 2022, ~~and use these data~~
for model calibration and evaluation. ~~To ensure that the drying probability computed for each HER2 is not biased, ONDE field~~
~~campaigns are only taken into account if at least 75% of the ONDE sites within a given HER2 are actually monitored during~~
~~the campaign at hand.~~

Commenté [T16]: RC1.4

150 Based on ONDE data, the *PFI* is defined as the proportion of small streams under drying conditions at the regional scale. It is
used as a proxy indicator for the intensity and duration of dry periods. Long-term analysis of the *PFI* characterises the spatial
and temporal trends of drought across regions and provides valuable information about IRES intermittence behaviour.

Commenté [T17]: RC1.2

2.2 Delineating areas with homogeneous hydrological behaviour: the Hydro-EcoRegions

155 Hydro-EcoRegions (HER) are spatially homogeneous areas defined over France based on natural drivers involved in river
ecosystem functioning, such as geology, topography, and climate. There are 85 level-2 HER (HER2) across France, and they
are derived from the ~~sub-~~divisions of the 22 level-1 HER (HER1) (Wasson et al., 2002), which were used in the previous study
about the impact of climate change on *PFI* in France (Sauquet et al., 2021). In contrast, the present study investigates flow
intermittence at the scale of HER2 regions in order to model *PFI* at a higher spatial resolution. The statistical approach
160 developed hereafter requires sufficient observations of flow states for calibration purposes (see Section Sect. 2.4.3.1), which
led us to perform five groupings of HER2 in order to ensure that enough observations are available for model calibration.
When merging HER2 regions we checked that they belonged to the same level-1 HER to ensure that they share similar
environmental characteristics at the large scale. In the end, a partition of France into 75 entities (HER2 or pool of HER2) was
considered for this study (~~median area: 4990 km²~~-Fig. 1).

Commenté [T18]: RC3.12

Commenté [T19]: RC3.13

165 These study involves 75 HER2s with have a median surface area of 4990 km² and , of which more than 20% (17/75) have a
surface area greater than 10000 km². The HER2s were the subject of ONDE field campaigns from May to September between

2012 and 2022, thus resulting in 4125 campaigns. A total of 78 campaigns are excluded from the analysis due to their failure to cover more than 75% of the monitoring stations within a given HER2. Thus, the data exclusion rate from the ONDE network is less than 2%, leaving 4047 field campaigns with usable data.

To ensure that the drying probability computed for each HER2 is not biased, ONDE field campaigns are only taken into account if at least 75% of the ONDE sites within a given HER2 are actually monitored during the campaign at hand.




Figure 1: Delineation of level 2 Hydro-EcoRegions (HER2) across France. The four HER2 that are used for illustration in this study are highlighted in purple (see Sect. 3.7).

2.3 Modelling of future discharge hydrological projections over France: the Explore2 project

The assessment of changes in the PFI over the 21st century are based on the large set of hydrological projections produced by the Explore2 project (Sauquet et al., in prep). The daily discharge projections were obtained at about 4000 simulation points distributed along the French river network, with a constraint on the minimum drainage area imposed by the spatial resolution of climate projections (64 km²) and another constraint of an even coverage across France. Most Explore2 simulation points were selected since data collected at these locations contribute to actual environmental monitoring or water management issues.

Commenté [T20]: RC1.12

185 Explore2 is a multi-model simulation experiment, involving nine different hydrological models (HM), but some of them are restricted to regional applications. On average, each simulation point has discharge projections simulated by four hydrological models HM (Interquartile Q1 and Q3 (IQ): 3-5). Finally, with the objective of predicting flow intermittence at the national scale, we select the five models with the largest simulation domain: CTRIP (Decharme et al., 2019), GRSD (De Lavenne et al., 2016), J2000 (Morel et al., 2023), ORCHIDEE (Huang et al., 2024), and SMASH (Jay-Allemand et al., 2020).

190 The Explore2 project considers only natural flows, although some of the HMs hydrological models involved in the project have the capability of including human-induced influences. The hydrological model evaluation of hydrological models was performed using an extended dataset of near-pristine catchments (Strohmenger et al., 2023). This observation-based simulation for each HM was. The simulations used for model evaluation were carried out for the period 1976-2022 using the French near-surface SAFRAN meteorological reanalysis (Vidal et al. 2010) as input. In addition, these hydrological models HM included

195 in Explore2 have been evaluated on the common period of availability 1976-2019 (Sauquet et al., in prep). After validation, the hydrological models HM are forced by an ensemble dataset derived from 17 pairs of Global and Regional Climate Models (GCM-RCM) corrected by the statistical adjustment method ADAMONT (Verfaillie et al., 2018) for both historical (1976-2004) and future climate scenarios RCP 2.6, 4.5 and 8.5 (2005-2100; Appendix Sect. A). In the end, these bias-corrected climate projections were used as inputs for the hydrological models: all 1740 GCM-RCM projections (resp. 9 and 17) are used

200 for under RCP8.5 scenario, whereas 10 of these projections are used under the RCP2.6 and 9 under RCP4.5 scenario (resp. RCP4.5 and RCP8.5). The baseline period H0 is set to 1976-2005 and used thereafter for analysing changes in IRES behaviour. Overall, the hydrological projections used thereafter to predict PFI result from combinations RCP-GCM-RCM-Hydrological Model HM.

Commenté [T21]: RC3.11

Commenté [T22]: RC2.4

Commenté [T23]: RC3.11

Commenté [T24]: RC3.11

Commenté [T25]: RC3.11

Commenté [T26]: RC3.11

Commenté [T27]: RC2.5

Commenté [T28]: RC2.8

Commenté [T29]: RC3.11

3 Modelling framework

205 3.1 Linking intermittence to stream discharge at the HER2 scale

The empirical method suggested by Beaufort et al. (2018; Eq. (4)) is applied here to estimate the Probability of Flow Intermittence (PFI) at the scale of HER2 regions. This method follows a two-step process that links observed flow states at ONDE sites to daily streamflows (either gauged or modelled) using logistic regressions.

First, observations from ONDE sites are used to determine the percentage of sites with “dry conditions” for each HER2 (h) and each campaign date (j). Under current climate conditions, (f) values are given by the percentage of ONDE sites with “dry conditions” for each HER2 (h) and each campaign date (j) (Fig. 2). This percentage of dry ONDE sites is considered as representative of the regional PFI. Flow intermittence evolves over relatively long time scales, with seasonal transitions between flowing and dry states occurring gradually within the year, except in cases of significant rainfall events that can temporarily trigger flow conditions. Given this inertia, the estimation of flow intermittence remains appropriate even if not all sites within a HER2 are monitored on the exact same day, with a shift of only a few days during a given campaign. To further ensure the temporal consistency across campaigns the monitoring is systematically conducted at fixed sites, and data from a

Commenté [T30]: RC3.3, RC3.5

given ONDE campaign and HER2 are only considered if at least 75% of the ONDE sites within the HER2 are monitored during the campaign. This approach minimizes potential biases associated with incomplete observations, and ensures that the percentage of dry ONDE sites referred hereafter as $PFI_h(j)$ can be considered representative of the PFI .

Subsequently, a logistic regression model is calibrated for each HER2 in order to ~~estimate-link the~~ PFI values for day j to ~~streamflow conditions~~ (Eq. (1), Fig. 2):-

$$PFI_h(j) = \frac{e^{\beta_{0-HER2h} + \beta_{1-HER2h} \times F_{Q-HER2h}(j)}}{1 + e^{\beta_{0-HER2h} + \beta_{1-HER2h} \times F_{Q-HER2h}(j)}}, \quad (1)$$

where $\beta_{0-HER2h}$ and $\beta_{1-HER2h}$ are respectively the logistic regression intercept and slope coefficient associated with the predictor $F_{Q-HER2h}$.


The model uses the mean non-exceedance frequencies of discharge $F_{Q-HER2h}(j)$ as ~~explanatory variable~~ predictor for $PFI_h(j)$ (Eq. (2)). $F_{Q-HER2h}$ is regarded as a proxy ~~for~~ characterising the current wet versus dry hydrological conditions at the regional scale. ~~It is derived from the non-exceedance frequency curves of streams, which describe the probability that a given discharge value is not exceeded. More specifically, for each specific date~~ (j) corresponding to an ONDE field campaign and each HER2 (h), the daily empirical non-exceedance frequencies of discharge are spatially averaged over all available n streams whose drainage areas intersect the HER2 of interest (weighted by their drainage area) and temporally averaged over the period $[j - 6; j]$. This seven-day window allows the ~~integration-inclusion~~ of non-simultaneous response times caused by ~~heterogeneous~~ propagation times in the underground and the river networks (the choice for a seven-day window is the result of an optimisation process, see Appendix Sect. B). ~~In the end, $F_{Q-HER2h}$ is defined by:~~

$$F_{Q-HER2h}(j) = \frac{1}{7 \times n} \sum_{k=j-6}^j \sum_{s=1}^n F_Q(k, s) \quad (2)$$

where $F_Q(k, s)$ represents the non-exceedance frequency of each stream s among the n streams whose drainage areas intersect the HER2 h . This frequency is weighted by the drainage area of stream s . The summation is computed over each day k within the $[j - 6; j]$ period around campaign j .

Commenté [T31]: RC2.6

Once calibrated, the logistic regressions are used to convert the hydrological projections from the Explore2 project into *PFI* projections.






Figure 2: Schematic view of the approach adapted to derive the regional Probability of Flow Intermittence (*PFI*) from ONDE sites using and daily gauging flows either gauged or modelled stations within a given HER2. The daily flows simulated by Explore2 at three simulation points generate flow time series (column 1) and their corresponding non-exceedance frequency curves (column 2). A logistic regression is then used to link daily flows to ONDE observations, as illustrated here for the campaign of September 25, 2012. For this campaign, daily discharge values simulated between September 18 and September 25, 2012, are extracted for each simulation point whose drainage area intersects the HER2. These seven flow values are matched to their respective non-exceedance frequency using the non-exceedance frequency curve. The mean non-exceedance frequency across all simulation points is then associated with the proportion of ONDE sites exhibiting “dry conditions” during this campaign (observed *PFI*, column 3). The logistic regression is calibrated using data from all ONDE campaigns. Once calibrated, these models allow for the conversion of daily discharge time series into daily *PFI* time series for the HER2 (column 4).

3.2 Calibrating the *PFI* logistic regression models using observed daily discharge data

Logistic regression models are preliminarily fitted between 2012 and 2022 using observed discharge data. The daily discharge data used in the calculation of $F_{Q,HER2h}$ were compiled using gauging station records from the French reference hydrologic network HydroPortail (Leleu et al., 2014) (www.hydro.eaufrance.fr). The set of gauging stations were selected on the basis of three different criteria:

- Daily discharge data must be available during the ONDE field campaigns, i.e., within the $[j-10;j+10]$ interval centred on 25th of each month from May to September, for the years 2012 to 2022.
- The human disturbance to flow must be minimal at the gauging stations.
- All selected gauging stations must have a drainage area of less than 2000 km². Large gauged basins with a high Strahler order have been discarded since they are unlikely to behave in the same way as headwater small streams.

This selection process results in a final dataset including 1008 stations (Fig. 3). The flow duration non-exceedance frequency curve of each gauging station is computed using its entire available discharge time series, leading to variations in the time

Commenté [T32]: RC1.7, RC2.6

Mis en forme : Police :Italique

Commenté [T33]: RC1.11

Commenté [T34]: RC2.6

Commenté [T35]: RC1.5

265 periods covered by the discharge data across stations. Nevertheless, limiting the analysis to the period 2012-2022 would likely
exclude critical information on hydrological extremes and fail to fully capture the variability of high and low flow events,
which are pivotal for understanding a river's behaviour during extreme conditions.

3.3 Assessing the performance of *PFI* logistic regression models under current climate

270 ~~Logistic regression models are preliminarily fitted between 2012 and 2022 using observed discharge data.~~ Several cross-
validation schemes are then considered to test the robustness of these models. A leave-one-year-out cross-validation is first
carried out by excluding one year at a time from the training dataset, learning from the data of the remaining years and then
making estimations for the excluded year. The robustness under different climate conditions is also assessed through
differential split-sample test (DSST; Dakhloui et al., 2017), which is a k-fold cross-validations performed on three distinct
groups of hydrological years categorized as dry, intermediate, or wet (Appendix Sect. C). The years are discriminated ~~selected~~
according to the aridity index (Barrow, 1992), which is one of the main drivers of flow intermittence at the global scale
275 (Sauquet et al., 2021).

The model performance is assessed using several skill scores: the bias (to detect underestimation or overestimation), the mean
absolute error (MAE, to quantify the magnitude of prediction errors), and the root-mean-square error (RMSE, which provides
a quadratic estimation that assigns relatively higher weight to large errors). Additionally, the Nash-Sutcliffe model efficiency
280 coefficient (NSE) is computed to compare the variance of the estimation error to the variance of the observed time series (Nash
and Sutcliffe, 1970). The Kling-Gupta Efficiency (KGE) complements the NSE metric as KGE is the Euclidean distance
between observed and estimated *PFI*, computed using the coordinates of bias, standard deviation, and correlation (Gupta et
al., 2009). The Leave One Year Out analysis results are obtained by averaging the validation metrics computed for each year.

3.4 ~~Simulating *PFI* in a changing climate~~ Selecting simulation points for *PFI* projections under climate change

285 The modelling framework suggested by (Beaufort et al., 2018; Sauquet et al., 2021) was applied to the 75 HER2s under
regional climate projections to assess the potential impact of climate change on flow intermittence dynamics.

Discharge data derived from the Explore2 database are used as input for the logistic regression models introduced in Sect.
2.1.3.1. However, not all the stations in the HydroPortail database have been simulated by the hydrological models ~~HMs~~. To
290 overcome this co-location problem, the choice ~~was-is~~ made to identify the simulation points closest to the 1008 gauging stations
selected for the diagnosis in Section Sect. 2.1.4.3.2 (Fig. 3). The conditions of the application experiment are thus similar to
the ones of the diagnostic experiment. A constraint ~~was-is~~ applied to ensure that no simulation point ~~was-is~~ assigned multiple
times, unless there ~~was-is~~ no alternative point within a 50 km distance from the gauging station (Appendix Sect. D). This
selection ~~was-is~~ made independently for each hydrological model. Only 483 points ~~awere~~ assigned for the J2000 model because
295 this model simulates streamflow only for the Loire and Rhône river basins.


Mis en forme : Police :Italique

Commenté [T36]: RC3.4

Commenté [T37]: RC3.13

Commenté [T38]: RC3.11

Commenté [T39]: RC3.13






Figure 3: Location of the 3302 ONDE sites, the 1008 gauging stations selected from the HydroPortail database, and the Explore2 simulation points selected for each GRSD and J2000 hydrological model (line 1). The maps on the second and third line respectively show the ratio between the sum of the catchment areas defined by the gauging stations or the simulation points and the surface area of the HER2 they intersect (line 2). These catchment areas may overlap and their sums are computed without considering their spatial distribution. The maps on the third line show the density of gauging stations or simulation points (line 3). The same results are available for the other hydrological models in Appendix Sect. E.

Commenté [T40]: RC1.8, RC1.9, RC2.7

Commenté [T41]: RC3.11

3.5 Constructing synthetic non-exceedance frequency curves for PFI projections

Commenté [T42]: RC3.4

Mis en forme : Titre 2

Applying empirical formulas (i.e., fitted using observed discharges) to simulated discharges may lead to poor estimates of PFI due to bias in the modelling chains. Thus, before being applied to future climate conditions, Eq. (1) needs to be recalibrated with past discharges simulated by each HM (here discharges simulated by the different HMs using SAFRAN reanalysis data available during the period 2012–2022 are used). In our framework, non-exceedance frequency curves must be computed for each simulation point prior to PFI estimation. To avoid problems of extrapolating non-exceedance frequency for-of discharge values outside the range of historical flows simulated with SAFRAN when performing PFI projection under changing climate, we decided to build a synthetic flow time series with a period length of 169 years, by combining that combine the historical available daily discharge data simulated with SAFRAN over the period 1976–2022 with projected discharge data simulated by the different modelling chains RCP-GCM-RCM-Hydrological Model over the period 1976–2100. This concatenation was-is made independently for each modelling chain RCP-GCM-RCM-HM at each and simulation point. As a consequence, while the assessment of the performance of PFI logistic regression models under current climate is conducted using observed discharge data that can be unbalanced (i.e., covering different observation periods) due to heterogeneous data availability across stations (see Sect. 3.3), the calibration used for projecting PFI under future climate scenarios relies on

Commenté [T43]: RC1.5

standardized and balanced synthetic time series which ensures the consistency of the modelling across simulation points and modelling chains.

Finally, each synthetic discharge time series generated by a specific RCP-GCM-RCM-~~Hydrological Model~~~~HM-modelling~~ chain results in a ~~flow-duration~~~~non-exceedance frequency~~ curve which is used thereafter to calibrate ~~the a~~ logistic regression (Eq. (1)). ~~The resulting logistic regressions are evaluated for each HM using the median of the bias, MAE, RMSE, NSE, and KGE skill scores over all RCP-GCM-RCM modelling chains.~~

Commenté [T44]: RC3.11

Commenté [T45]: RC2.6

3.6 Recalibrating logistic regressions and projecting *PFI* under climate scenarios

Before being applied to future climate conditions, the logistic regression model of Eq. (1) needs to be recalibrated with past discharges simulated by each hydrological model (here discharges simulated by the different hydrological models using SAFRAN reanalysis data available during the period 2012-2022 are used). The resulting logistic regressions are then re-evaluated for each hydrological model using the median of the bias, MAE, RMSE, NSE, and KGE skill scores over all RCP-GCM-RCM modelling chains.

Commenté [T46]: RC3.4

Mis en forme : Titre 2

Finally, after re-calibration, these logistic regressions are used to project *PFI* based on discharge data simulated for both the historical period (1976-2004) and the projection period (2005-2100) under various climate scenarios (RCP 2.6, 4.5 and 8.5).

3.7 Assessing changes in *PFI* and associated uncertainties

We compare *PFI* between a reference period ~~H0~~ (1976-2005), and medium-term (~~H1~~, 2041-2070) as well as long-term (~~H2~~, 2070-2099) horizons. Results are presented at the HER2 scale across France and ~~illustrated-discussed~~ in more details for four ~~contrasted~~ HER2s (Fig. 1), ~~deliberately~~ selected to represent contrasting hydro-climatic conditions and behaviours: “Haute Normandie Picardie” (HER2-57, North, oceanic temperate climate), “Plaine de Bourgogne” (HER2-81, Northeast, continental temperate climate), “Massif de l’Oisans” (HER2-12, Alps, mountainous climate), and “Plaine méditerranéenne” (HER2-105, Southeast, Mediterranean climate). In the rest of the article, “dry periods” of HER2s are defined as the period of the year when *PFI* is greater than 20%. For each HER2, the changes in intensity and seasonality of flow intermittence for ~~2041-2070~~~~H1~~ and ~~2070-2099~~~~H2~~ relative to ~~1976-2005~~~~H0~~, are characterized using the mean daily *PFI* between July and October ($mPFI_{7-10}$), and the median of the first and last days (respectively, T_f and T_l) of the year with *PFI* exceeding 20%.

Mis en forme : Police :Italique

Commenté [T47]: RC1.12

The analysis of the uncertainty propagation in the modelling framework is examined using the QUALYPSO method, applied to the national average of $mPFI_{7-10}$ ($mPFI_{7-10}[France]$), weighted by HER2 areas and obtained under RCP 2.6, 4.5 and 8.5 climate change scenarios (Evin et al., 2019; Evin et al., 2021). The QUALYPSO method is used to characterise the changes in $mPFI_{7-10}[France]$ by estimating its ensemble mean over all the projections, as well as the total uncertainty associated with this set of projections, the contribution of the various sources of uncertainty (RCP, GCM, RCM, ~~hydrological models~~~~HM~~, residual and internal climate variability) to the total uncertainty and the main effect of each individual model and its contribution to the total uncertainty. The contribution of ~~one specific model each component to the total uncertainty~~ is estimated as its fixed effect

Commenté [T48]: RC3.11

in a linear regression model, which adjusts for imbalances in the number of simulations across components within modelling chains by the difference between the mean of all modelling chains considering the model at hand and the ensemble mean (Appendix Sect. F).

Commenté [T49]: RC2.9

A multi-model index of agreement (*MIA*) is also computed on $mPFI_{7,10}$ time series to highlight convergence in changes over the projections (Tramblay and Somot, 2018):

$$MIA = \frac{1}{n} \sum_{k=1}^n i_k, \quad (3)$$

where n is the number of projections, $i_k = 1$ for a significant positive trend of the $mPFI_{7,10}$ according to the Mann-Kendall test (with $\alpha = 0.1$), $i_k = -1$ for a significant negative trend and $i_k = 0$ for no significant trend.

4 Results

4.1 Data pre-processing

The study involves 75 HER2s with a median surface area of 4990 km², of which more than 20% (17/75) have a surface area greater than 10000 km². The HER2s were the subject of ONDE field campaigns from May to September between 2012 and 2022, thus resulting in 4125 campaigns. A total of 78 campaigns are excluded from the analysis due to their failure to cover more than 75% of the monitoring stations within a given HER2. Thus, the data exclusion rate from the ONDE network is less than 2%, leaving 4047 field campaigns with usable data.

The 3302 ONDE sites are located in headwaters small streams while gauging stations in France are monitoring medium size catchments (Van Meerveld et al., 2020). In addition, all Explore2 simulation sites have a drainage area larger than 64 km². Unsurprisingly, there is little overlap between the distribution of areas drained by ONDE sites (median: 24 km²; quartiles Q1-Q3: 12-50) and those drained by the two sets of 1008 gauging stations (median: 173 km²; IQR: 85-396) and 1008 simulation points (median: 178 km²; Q1-Q3: 95-400) (Appendix Sect Fig. E1). ONDE sites, gauging stations and simulation points are located at similar elevations (overall median: 168 m; Q1-Q3: 75-308; Appendix Fig. G2). The drainage areas of ONDE sites, gauging stations and simulation points represent respectively a median coverage of 21% (Q1-Q3: 15-35), 54% (Q1-Q3: 31-94) and 55% (Q1-Q3: 32-96) of the HER2 areas. This corresponds to a median number of 6.1 sites (Q1-Q3: 5-8), 3.6 stations (Q1-Q3: 1.9-5.7) and 3.7 simulation points (Q1-Q3: 2-6) per 1000 km² of HER2 (Fig. 3). These statistics highlight the question of representativeness of the ONDE sites despite the even coverage of the ONDE network.

Commenté [T50]: RC3.5

4.2 Model performance

The logistic regressions calibrated using observed flows accurately predict *PFI* under current climate conditions (Fig. 4; Appendix [SeetFig. GH1](#)). The median performance across HER2s is, for example, 0.85 for KGE ([Q1-Q3](#): 0.77-0.89) and 0.79 for NSE ([Q1-Q3](#): 0.64-0.84) when the leave-one-year-out cross-validation is considered. The KGE skill score exceeds 80% in 50 out of 75 HER2 and the bias is very low. The best performances are observed in sedimentary plains and in Aquitaine while lower performances are obtained in mountainous areas such as the Alps, the Massif Central and the Pyrenees. The mean absolute error and RMSE show that the largest absolute deviations between observations and predictions occur in the southeastern part of France. This result is consistent with previous findings (e.g. Fig. 3 in Sauquet et al., 2021): (1) the performance of the logistic regressions is partly correlated with the level of intermittence, and (2) no-flow conditions which mainly occur in winter due to freezing in the Alps are not captured by the ONDE network. Observed values of *mPFI*₇₋₁₀ range between 3 and 37% (median: 14.3%) across France while values of *mPFI*₇₋₁₀ obtained with SAFRAN range between 2 and 37 % (median: 14.4%). The modelling approach is able to reproduce the spatial pattern of flow intermittence (Fig. 4), although the results obtained with the DSST show less satisfactory performance (Appendix [SeetFig. GH2 and Table H1](#)). The two skill scores KGE and NSE are sensitive to wet, intermediate and dry conditions while the other skill scores show no major change. The values of RMSE and MAE skill scores do not increase much when the calibration dataset is stratified based on climate conditions, which indicates that the proposed model is quite robust to climate fluctuations. The bias, however, indicates an underestimation of *PFI* in the southwest and in the north during dry years. This lower performance is partly due to a reduced number of years used for calibration purposes, and to the difficulty to extrapolate the logistic regression equations under unobserved climate conditions.

Mis en forme : Police :Italique
Commenté [T51]: RC3.5

Mis en forme : Police :Italique




Figure 4: Leave-One-Year-Out cross-validation assessing the calibration of the logistic regressions using discharge measurements from gauging stations. From left to right: observed mean *PFI* measured during the calibration period (2012-2022, left), Kling-Gupta Efficiency (KGE, center) and absolute bias of *PFI* predictions (right).

The logistic regressions fitted with discharge data simulated by hydrological models^{HM} also accurately reconstruct *PFI* from May to September. Compared to the results obtained with observed discharge data, the performance is slightly lower due to the inherent imperfections of the SAFRAN reanalysis compared to the observed climate (Fig. 5, 6; Table 1; Appendix Sect. HJ). The inter-HER2 and inter-projections median estimated $mPFI_{7-10}$ across France is 11.3-13.5% depending on hydrological models^{HM}, which is close to the 14.3% value (Q1-Q3IQR: 8.4-20.3) derived from the ONDE network. The KGE is slightly lower but exceeds 80% in more than half of the HER2 using the CTRIP (38/75), SMASH (38/75), and GRSD (43/75) models. The inter-HER2 median of the NSE skill score (calculated for all RCP-GCM-RCM modelling chains and for each of these three hydrological models^{HM}) indicates that between 71% and 74% of the variance in $mPFI_{7-10}$ is explained by the logistic regressions. The other skill scores do not show a loss of performance (KGE between 0.80 and 0.81; MAE between 0.04 and 0.05, and RMSE between 0.06 and 0.07) when comparing the results of the Leave-One-Year-Out cross-validation obtained using observed and simulated discharge data. Lower performance is observed with the ORCHIDEE and J2000 models, as the KGE exceeds 80% in only 7/75 and 5/37 HER2, respectively. Their median KGE (0.60 and 0.69, respectively), NSE (0.49 and 0.62), MAE (0.06), and RMSE (0.08 and 0.09) are lower than the values obtained during calibration using observed flow data, indicating that these two hydrological models seem less sensitive.

Regardless of the hydrological model^{HM} of interest, the maps of skill scores show spatial patterns that are consistent with the results obtained with observed discharge data (Fig. 6; Appendix Sect. HJ). Both NSE and KGE remain high in the sedimentary plains and the southern part of the country, while they are less favorable/favourable in mountainous regions (Alps, Massif Central, and Pyrenees). An overestimation persists in the southwestern part of France, while underestimations of *PFI* values are found in the northwestern, northern, and southeastern parts of France, and are especially pronounced during dry years. The

Commenté [T52]: RC1.10

Mis en forme : Police :Italique

Mis en forme : Police :Italique

Commenté [T53]: RC3.11

Mis en forme : Police :Italique

Commenté [T54]: RC3.11

Commenté [T55]: RC3.11

Commenté [T56]: RC3.11

Commenté [T57]: 3.12

425

430

models also successfully reproduce the inter-annual variability of *PFI*, particularly the alternation of dry (e.g., 2019) and wet (e.g., 2015) years (Fig. 5: ~~Appendix-Set-G~~). Thus, the modelling chains demonstrate their ability to simulate *PFI* over the period 2012–2022 and are therefore deemed reliable to attempt projecting changes in flow intermittence under modified climate conditions. However, note that *PFI* values are slightly underestimated when the logistic regressions are calibrated on the driest years and ~~then applied to wetter years than those used for calibration~~. This suggests that our future projections are likely to be biased in the same way, since the calibration is based on current conditions before applying the regression models to a potentially drier future climate.

Commenté [T58]: RC2.10

Table 1: Evaluation of the logistic regressions calibrated using observed discharge data (results of the cross validation) and using flows simulated by the CTRIP, GRSD, ~~J2000~~, ORCHIDEE and SMASH hydrological models. Statistics of the skill scores are summarized by medians and ~~the first and the third interquartiles ranges (IQ1-Q3, into brackets)~~ across all HER2s.

Commenté [T59]: RC2.4

HM	NSE (unitless)	KGE (unitless)	MAE (unitless)	RMSE (unitless)
Observed-discharge				
data	0.79	0.85	0.10	0.07
Leave-One-Year Out (2012-2022)	(IQ: 0.64-0.84)	(IQ: 0.77-0.89)	(IQ: 0.06-0.13)	(IQ: 0.05-0.09)
CTRIP	0.74 (IQ: 0.61-0.80)	0.80 (IQ: 0.69-0.85)	0.05 (IQ: 0.04-0.07)	0.07 (IQ: 0.05-0.10)
GRSD	0.74 (IQ: 0.64-0.80)	0.81 (IQ: 0.71-0.86)	0.04 (IQ: 0.03-0.06)	0.06 (IQ: 0.04-0.08)
J2000	0.62 (IQ: 0.52-0.71)	0.69 (IQ: 0.63-0.79)	0.06 (IQ: 0.04-0.09)	0.09 (IQ: 0.07-0.13)
ORCHIDEE	0.49 (IQ: 0.39-0.63)	0.60 (IQ: 0.50-0.72)	0.06 (IQ: 0.04-0.09)	0.08 (IQ: 0.06-0.11)
SMASH	0.71 (IQ: 0.65-0.80)	0.80 (IQ: 0.72-0.86)	0.04 (IQ: 0.03-0.06)	0.06 (IQ: 0.05-0.08)


Tableau mis en forme

435

<u>Hydrological model</u>	<u>NSE (unitless)</u>	<u>KGE (unitless)</u>	<u>MAE (unitless)</u>	<u>RMSE (unitless)</u>
<u>Observed discharge</u>				
<u>data</u>	0.79	0.85	0.10	0.07
<u>Leave One Year</u>	(Q1-Q3: 0.64-0.84)	(Q1-Q3: 0.77-0.89)	(Q1-Q3: 0.06-0.13)	(Q1-Q3: 0.05-0.09)
<u>Out (2012-2022)</u>				
<u>CTrip</u>	0.74	0.80	0.05	0.07
	(Q1-Q3: 0.61-0.80)	(Q1-Q3: 0.69-0.85)	(Q1-Q3: 0.04-0.07)	(Q1-Q3: 0.05-0.10)
<u>GRSD</u>	0.74	0.81	0.04	0.06
	(Q1-Q3: 0.64-0.80)	(Q1-Q3: 0.71-0.86)	(Q1-Q3: 0.03-0.06)	(Q1-Q3: 0.04-0.08)
<u>J2000</u>	0.62	0.69	0.06	0.09
	(Q1-Q3: 0.52-0.71)	(Q1-Q3: 0.63-0.79)	(Q1-Q3: 0.04-0.09)	(Q1-Q3: 0.07-0.13)
<u>ORCHIDEE</u>	0.49	0.60	0.06	0.08
	(Q1-Q3: 0.39-0.63)	(Q1-Q3: 0.50-0.72)	(Q1-Q3: 0.04-0.09)	(Q1-Q3: 0.06-0.11)
<u>SMASH</u>	0.71	0.80	0.04	0.06
	(Q1-Q3: 0.65-0.80)	(Q1-Q3: 0.72-0.86)	(Q1-Q3: 0.03-0.06)	(Q1-Q3: 0.05-0.08)


440

445



450 **Figure 5:** Time series of *PFI* for the four HER2s selected for illustrative purposes. Dots and lines are observed *PFI* derived from the
 455 **ONDE** network and estimated *PFI* computed by the logistic regressions using simulated discharge by each hydrological modelHMs,
 respectively.

Commenté [T60]: RC3.11



455 **Figure 6:** Kling-Gupta Efficiency (KGE) assessing the calibration of logistic regression using discharge simulated by each
 hydrological modelHM with SAFRAN-based simulations available during the 2012-2022 period

Commenté [T61]: RC3.11

4.3 Flow intermittence projections

The logistic regressions fitted for each HER2 are applied to derive the *PFI* values from the historical (1976-2004) and future (2005-2100) discharges simulated by the hydrological modelsHM4s forced by RCP-GCM-RCM climate projections over the period 1976-2100.

Commenté [T62]: RC3.11

Figure 7 shows the evolution of *mPFI*₇₋₁₀ and of the start date (*Tf*) and end date (*Tl*) of the dry period for 2041-2070H1 and 2070-2099H2 compared to 1976-2005H0. The results are illustrated in Fig. 7 with the median projection given by the GRSD model (see Appendix SetFig. HJ1-J4 for other models) under RCP2.6, RCP4.5, and RCP8.5 climate scenarios. Table 2 details the same results for the four HER2 under RCP8.5.

Commenté [T63]: RC2.8

Overall, the average *PFI* shows an increase, indicating a gradual intensification of dry periods over the 21st century under the RCP 4.5 and RCP 8.5 scenarios. The inter-HER2 and inter-projections median of *mPFI*₇₋₁₀ calculated at the national scale (*mPFI*₇₋₁₀[France]) for the baseline period 1976-2005H0 ranges from 10 to 12% depending on the hydrological modelHM. However, the median projection of *mPFI*₇₋₁₀ during 1976-2005H0 is higher than 20% in 12 to 20 of the 75 HER2s (depending on the hydrological modelHM), although although it does not exceed 42% in any HER2. In contrast, uUnder RCP 8.5, *mPFI*₇₋₁₀[France] could reach values between 13 and 21% in the mid-term of the century 2041-2070H1 and between 16 and 29% by the end of the century 2070-2099H2. By this time, six HER2s could have inter-projections median of *mPFI*₇₋₁₀ exceeding 50% according to at least two hydrological modelsHM4: four in the south-east of France; one in the southwest; and one in the northwest of France. Under RCP 4.5, changes are more moderate by the end of the century 2070-2099H2, with *mPFI*₇₋₁₀[France] ranging between 14 and 20%. The spatial pattern of the changes is not uniform but looks similar between RCP 8.5 and RCP4.5. A strengthening divide between the south-west and the north-east of the country is projected. Regions already prone to intermittence are expected to experience an increase in this phenomenon under both emission scenarios. The ratio of the ensemble median of *mPFI*₇₋₁₀ at the end of the century compared to the baseline period evolves around 1.4 for RCP 4.5 and 1.9 for RCP 8.5. Mountainous regions see the intensity of summer dry periods increase but remain relatively spared.

Commenté [T64]: RC2.8

Commenté [T65]: RC3.11

Commenté [T66]: RC3.11

Commenté [T67]: RC2.11

Commenté [T68]: RC3.11

Shifts of the start and the end of the dry period are partly correlated with changes in *mPFI*₇₋₁₀ (Fig. 7). It appears that climate change results in both earlier and later dry periods in a fairly symmetrical manner: dry periods are advanced and extended, but with different sensitivities structured along a north-south gradient. Under RCP 8.5 climate conditions, the dry periods are projected to get (on median) longer in southern France at the end of the century. According to at least two hydrological modelsHM, the highest shifts of *Tf* or *Tl* could exceed 5 weeks in several mountainous HER2 (four HER2 in the Alps, three in the Pyrenees, two in the Jura). In contrast in the northern part of France, the dry periods could be advanced and extended by only one or two weeks, as observed in HER2-81 “Plaine de Bourgogne” (Table 2). HER2 28 could be one of the most affected in the north due to its impermeable clay-sandy formations, which differs from the neighbouring sedimentary formations. One

Commenté [T69]: RC3.11

Table 2: Statistics of flow intermittence characteristics for the four illustrative HER2s under RCP 8.5. The minimum (min), median (med) and maximum (max) are the ensemble minimum, median and maximum across all the projections and hydrological models. *Nb Days PFI > 20%: Number of days in dry periods (period of the year when PFI exceeds 20%).*




Figure 7: Ensemble median of changes in $mPFI_{7-10}$ (columns 1 to 3), and ensemble median shift of the start date Tf (columns 4 and 5) and the end date Tl (columns 6 and 7) of the dry period over the two periods $2041-2070H1$ and $2070-2099H2$ under the three RCPs for the GRSD hydrological model, relative to the baseline period $1976-2005H0$. The shift, expressed in week, takes a positive value when the duration of the dry period increases. Grey HER2s had no period with a $PFI > 20\%$ during the reference period. The same results are available for the other hydrological models HM in Appendix Set Fig. HJ1-J4.

Commenté [T70]: RC3.6

Commenté [T71]: RC3.9

Commenté [T72]: RC3.11

Average proportion of zero flows smoothed over 5 days
Hydrological model: GRSD - HER2 12




Figure 8: Seasonal pattern of *PFI* for the periods 1976-2005, 2041-2070 and 2071-2100 under RCP 2.6, 4.5 and 8.5 scenarios for GRSD hydrological models in the Alps (HER2 12). The storylines highlighted in this figure are introduced in Appendix Sect. H.1. The same results are available for HER2s 57, 81 and 105, and for the other hydrological models in Appendix Sect. H.5-21.

4.4 Uncertainty analysis in hydro-climatic projections and their impact on *PFI* projections

Figure 9 illustrates the results of the uncertainty analysis, showing the variability-uncertainty range of $mPFI_{7-10}[France]$ throughout the 21st century. These results confirm that dry conditions may occur more frequently in a changing climate, with 22% projected increase in $mPFI_{7-10}[France]$ by the end of the century. The total variability-uncertainty of $mPFI_{7-10}[France]$ over all projections results from the accumulation of uncertainties related to RCP scenarios, GCMs and RCMs, hydrological models, residual variability, and internal variability. The confidence interval, and therefore the extent of change, increases over the course of the 21st century due to the divergent results of the modelling chain. This finding is consistent with studies using a similar methodology to assess the uncertainty of panels of flow projections (Evin et al., 2019; Aitken et al., 2023).

Commenté [T73]: RC2.8

Commenté [T74]: RC3.11

Commenté [T75]: RC3.7

Commenté [T76]: RC2.9

Commenté [T77]: RC2.9

Commenté [T78]: RC2.12

The contribution of each component to the total uncertainty varies over time. The fraction of total variance due to residual variability is small and remains stable over time. The main part of the variability uncertainty in climate change responses is thus explained by RCP, GCM, RCM and hydrological modelsHM. In addition, the contribution of the internal variability is highly predominant and represents more than 75% of the total uncertainty until the mid-century. This contribution then decreases with time as the total uncertainty increases. It becomes less than 50% of the total uncertainty by the end of the century but remains the greatest contributor to the total uncertainty over the whole simulation period.

Regarding the spatial distribution of the different contributions to the total uncertainty, it appears that climate data models are predominant source of the uncertainty, and that they are distributed in a spatially heterogeneous manner over France. The uncertainty related to the RCP scenarios increases over time and becomes predominant in the south, surpassing uncertainties related to other steps of the modelling chain (GCM, RCM, and hydrological modelsHM). In the mountainous regions, differences between results obtained with RCP4.5 and RCP8.5 are clearer by the end of the century, leading to a contribution of RCP to the total uncertainty reaching 20% at the end of the century. In northern France, the contribution of RCP to the uncertainty is lower compared to hydrological modelsHM, GCM, or RCM contributions. Uncertainty related to hydrological models is also spatially structured, with greater uncertainty in the northwestern part of France.

Commenté [T79]: RC2.9

Commenté [T80]: RC3.11

Commenté [T81]: RC2.9

Commenté [T82]: RC3.11

Commenté [T83]: RC3.11




Figure 9: Decomposition of the effects contributing to the variance of projections for relative changes (unitless) of $mPFI_{7-10}[France]$. In the top-left graph, the grey line and the coloured areas respectively show the climate change response and the contribution of time for each step of the modelling chain to the 90% confidence interval, respectively. For each model uncertainty and internal variability component, the vertical extent of the corresponding area is proportional to the fraction of total uncertainty explained by the component.

4.5 Agreement between changes

Despite uncertainties about the intensity of change, *PFI* projections still show a good degree of convergence in the direction of change when considering all climate projections driving each hydrological model individually, as well as in the overall convergence of *PFI* projections (Fig. 10). *MIA* values are first calculated independently for each *hydrological model*, before all projections are combined for multi-model assessment.

Figure 10 highlights spatial contrasts of the *MIA* multi-model approach: unsurprisingly, the projections of *mPFI*₇₋₁₀ are more uncertain in the northern part of France. The climate change signal on the proportion of dry periods is not homogeneous across France, with significant uncertainties remaining for the northern part.

The agreement of projections is high for the GRSD and SMASH models across France (*MIA* values close to +100% under RCP8.5 in Fig. 10 and +80% under RCP4.5, see Appendix Sect. [HK](#)). In contrast, CTRIP, J2000, and ORCHIDEE suggest contrasted regional impacts of climate change on *PFI* under RCP4.5, with a reduction of the *mPFI*₇₋₁₀ in the northern part of France, indicating a differentiated sensitivity of hydrological models to climate changes. This trend is also observed for ORCHIDEE under RCP8.5, while the other models agree on an increasing drying.

Mis en forme : Police :Italique




Figure 10: Agreement between projections of *mPFI*₇₋₁₀ for each hydrological model and inter-model agreement on the change signal of *mPFI*₇₋₁₀ under the RCP 8.5 scenario.

5 Discussion

5.1 Modelling framework and assumptions

This study presents projections of the *PFI* in ~~headwater small~~ streams by HER2 in France over the 21st century. It extends the previous analysis of Sauquet et al. (2021) on the 2012-2018 period by exploiting an extensive national dataset of stream intermittence observations collected on low-order streams (3302 observation sites), which are the most affected by shifts from perennial to intermittent flows (Reynolds et al., 2015; Dhungel et al., 2016). For the first time, we project the future evolution of the *PFI* using discharge data obtained from 5 hydrological models with conceptual (GRSD, SMASH), surface (CTRIP, ORCHIDEE), or process-oriented distributed (J2000) structures, under a multitude of possible climate scenarios informed by 17 pairs of CMIP5 GCM-RCM models.

The results of this study should be interpreted within the context of its underlying assumptions. Firstly, the study focuses on unregulated streams to characterize the "natural" hydrology (i.e., without considering water abstraction by anthropogenic activities or the impact of hydraulic engineering structures). This assumption was made in the Explore2 project, which provides the input streamflow simulations. Nevertheless, global water model simulations including direct human impacts by Döll and Zhang (2010) ~~or Gudmundsson et al. (2021)~~ concluded that ecologically relevant flow characteristics will be more altered by climate change than by withdrawals and dams. However, we believe that quantitatively estimating the extent to which flow intermittence is due to direct anthropogenic stressors is ~~important-crucial~~. ~~Such an estimation could for~~ ~~improving~~ our projections of flow intermittence ~~as well as for and inform~~ decision-making ~~aimed at in view to~~ ~~regulating~~ and preventing water stress situations for populations.

~~Secondly~~, groundwater levels are not incorporated into the model, although they could potentially enhance the accuracy of the projections. ~~Indeed, similar logistic regression models were used in Beaufort et al. (2018) and they incorporated groundwater levels measured by piezometers. Likewise, in Beaufort et al. (2019), climate, hydrological and morphological data were supplemented with groundwater levels to predict flow intermittence at a local scale. In these studies incorporating subsurface water in the regression models was thought to improve performance, in particular because the contribution of subsurface processes is known to play a decisive role in maintaining baseflow and mitigating flow intermittence. However, information about groundwater levels has not been included in the present study because hydrological projections incorporating groundwater levels have not yet been conducted across entire France (for instance, within the Explore2 project, P~~projections of groundwater levels have ~~only~~ been produced ~~for a restricted set of areas in France).~~ ~~by the Explore2 project but only for a limited area of France.~~

Thirdly, with only five annual discrete observations of streamflow intermittence over eleven years for the calibration of the logistic regression models, our ability to capture the full range of extremes and variability in these regression models remains imperfect. Yet, visual monitoring remains the most common technique for observing non-perennial streams. An alternative approach would be to consider citizen science to augment our database, although concerns about data reliability persist, in particular because past studies have shown participant agreement rates ranging from 46% to 70% (Scheller et al., 2024). Data

Commenté [T84]: RC1.2

Commenté [T85]: RC1.13

Commenté [T86]: RC1.14

scarcity necessitated conducting this analysis at the HER2 scale, which nonetheless represents a significant improvement of the spatial resolution by increasing the number of modelling subdomains from 22 to 75 compared to previous studies (Sauquet et al., 2021). In future work, a downscaling process could enhance the usability of *PFI* projections for local stakeholders in water management.

590 Fourthly, while projections of no-flow events often exhibit significant uncertainties when derived directly from discharge simulated by ~~HMs~~ *hydrological models* (Evin et al., ~~2019~~2024; Aitken et al., 2023), this study illustrates how categorical and discrete (in space and time) field observations can be combined with conventional stream gauge data, to enhance the understanding of *headwater* ~~small~~ stream drying dynamics. In this way, despite the divergences between projections induced by climate data, the agreement within this multi-model approach is relatively strong, indicating a consensus toward increasing *PFI* throughout the 21st century.

5.2 A consistent signal across projections

This study confirms that logistic regressions properly capture the relationships between flows in large watersheds and the *PFI* of *headwater* ~~small~~ streams. These regressions are calibrated and validated using discharge measurements from gauging stations, and subsequently using discharge simulated from SAFRAN climate data. Following this second calibration, projections based on regressions consistently indicate an increase in average *PFI* and a shift in the start and end dates of dry periods under RCP 4.5 and RCP 8.5 climate projections, suggesting a progressive intensification and extension of dry periods over the 21st century.

These results are consistent with previous studies indicating a transition of many streams from perennial to intermittent regimes (Jaeger et al., 2014; Reynolds et al., 2015; Dhungel et al., 2016; Schneider et al., 2013). ~~Rising temperatures attributed to climate change lead to an increased evapotranspiration and aridity, consequently resulting in an increase in dry streams (Tramblay et al., 2020; Zipper et al., 2021).~~ In regions already affected by intermittence, an increase in the intensity and duration of drying periods is likely, a phenomenon also anticipated in other areas around the world (Jaeger et al., 2014; Dai et al., 2011). Increasing intermittence and decreasing flow rates are observed in catchment-scale studies: the fraction of time with zero discharge increases from 0.05% to 4.30% in a Swiss Alpine catchment between 2020–2040 and 2080–2100 (Halloran et al., 2023) while the no-flowing phase could extend by up to 12 days between 1980–2009 and 2030–2059 for a catchment in southern Italy (De Girolamo et al., 2022). This trend is also noticeable on a larger scale, as monitoring of gauging stations in five regions with Mediterranean climates around the world between 1980 and 2019 (Carlson et al., 2024) and over 452 rivers on the European continent between 1970 and 2010 (Tramblay et al., 2020) both show that approximately 30% of them have already experienced drier conditions due to climate change, with modified flow regimes or extended periods of drying.

615 5.3 Uncertainties in northern France

In the northern part of France, discrepancies between GCM-RCM-~~Hydrological Models~~*HM* projections result in higher uncertainties in *PFI* projections and pronounced geographical contrasts on the multi-model-based *MIA* map (Fig. 10). ~~Part of~~

Mis en forme : Police :Italique

Commenté [T87]: RC3.11

Commenté [T88]: RC2.8

Commenté [T89]: RC2.8

Commenté [T90]: RC3.2

Commenté [T91]: RC3.11

Commenté [T92]: RC3.8

620 these uncertainties can be attributed to the underestimation of *PFI* in this region, particularly during dry years (Sect. 4.2, Appendix Fig. H2 and Table H1). However, the primary source of t~~These uncertainties likely stems primarily arise~~from
uncertainties in future rainfall patterns in this region where the majority of Explore2 projections indicate an increase in winter
rainfall and winter mean flows (for 8 out of 9 hydrological models~~HMs~~) and a decrease in summer precipitation (Evin et al.,
2024; Sauquet et al., 2024). The annual precipitation by the end of the century remains uncertain, due to the compensatory
effect between increased winter recharge and an increased evapotranspiration. (Ribes et al., 2022; Douville et al., 2021). As a
result, some hydrological models predict a shortening of the dry period for certain northern HER2 regions, including under
635 RCP 8.5. Similar uncertainties are observed along the east coast of the USA, where precipitation changes could turn
intermittent streams into perennial ones (Dhungel et al., 2016) while alternative climate change scenarios projected that by the
2040s, approximately half of the streams in Washington state would shift from snow-fed to rain-fed, resulting in reduced
annual discharge (Reidy Liermann et al., 2012).

5.4 Transformation of the snowmelt regime in the Alps

630 The mountain ranges could be moderately affected by the increased probability of dry conditions, consistent with our previous
modelling efforts (Sauquet et al., 2021), but our projections anticipate two specific phenomena in these regions. First, the
Pyrenees are more affected than the other mountain ranges, with significant changes impacting the massif and the dependent
basins in southwestern France. Additionally, Alpine HER2s will probably undergo hydrological regime changes. Higher winter
temperatures will lead to reduced snowfall and snow-related intermittence. This snowpack reduction will also reduce
635 groundwater recharge by spring melt. In addition, since soils tend to retain less water in summer due to more intense and brief
rainstorms (Rutkowska et al., 2023), an increased summer intermittence is expected despite increasing summer precipitations.
Using 16 hydrographic variables describing the magnitude, frequency, timing, duration, and rate of change of the flow regime
at 59 primarily selected sites with a Strahler order of 5, Dhungel et al. (2016) also observed the reduction in the snowmelt
regime at 2 Rocky Mountain sites under climate change. Halloran et al. (2023) also conclude that groundwater will play an
640 increasingly important role in ensuring flow in alpine streams and that the shift from perennial to intermittent could occur for
alpine streams over the course of the current century.
Furthermore, the reduction of the snowmelt regime in the Alps indicate that we can still make projections consistent with the
literature beyond the calibration period (May to September). More generally, the hypothesis of temporal transferability of the
models is a strong assumption in data exploitation, as it assumes that climate models, statistical adjustment methods and
645 hydrological models can simulate the behaviour of the systems they represent in a future hydro-climatic context that is very
different from the one in which they were developed (Evin et al., 2024). In this context, it is important to recall the risk that
the projections may be underestimated, as demonstrated by the validation of logistic regressions calibrated on wetter years
compared to dry years.

Commenté [T93]: RC3.11

Commenté [T94]: RC2.13

Commenté [T95]: RC2.13

6 Conclusion

650 ~~This~~ study assesses the changes in the intermittency of river flows across France in the context of climate change. For the first
time, multi-model and multi-scenario hydro-climatic projections are used as a predictor to explore the possible evolution of
the daily probability of flow intermittency at the scale of (level 2) Hydro-EcoRegions. ~~In conclusion, HMs often show~~
~~weaknesses in characterizing low flows and their responses to climate change. This study and other work in progress (e.g. Döll~~
~~et al., 2024) show that the simulation of intermittent rivers on a national or global scale is a complex subject with real scope~~
655 ~~for progress.~~ Leveraging monthly monitoring of headwaters ~~small~~ streams over ten summers, we calibrated logistic regressions
to transform hydrological projections of large watersheds into regional proportions of flow intermittence for the 21st century.
Under both RCP 4.5 and 8.5 scenarios, robust signals indicate an intensification of dry events, marked by increased ~~probability~~
~~of flow intermittence~~ ~~PF~~ and longer dry periods throughout the year. These changes are projected to be more pronounced in
southern France, with greater uncertainty in the northern half of the country. Mountain areas could remain relatively spared
660 from summer dry periods but shifts in hydrological regimes are anticipated. By the end of the century under RCP8.5, dry
stream phenomena along the Atlantic coast could surpass those currently observed in the Mediterranean region by the usual
monitoring campaigns. The evolution of droughts and reduced water availability suggested by these results could lead to
significant ecological impacts, including alterations in the structure and function of freshwater ecosystems (such as changes in
microbial activity and habitat loss), shifts in soil chemistry, increased carbon and solute fluxes, and sediment mobilization
665 (Geris et al., 2015).

Commenté [T96]: RC3.10

Commenté [T97]: 1.2

Appendix A: Explore2 modelling

Global Climate Model	Regional Climate Model	Histo- rical	RCP 2.6	RCP 4.5	RCP 8.5
CNRM-CERFACS-CNRM-CM5	CNRM-ALADIN6	×	×	×	×
CNRM-CERFACS-CNRM-CM5	KNMI-RACMO22E	×	×	×	×
CNRM-CERFACS-CNRM-CM5	MOHC-HadREM3-GA7-05	×	-	-	×
ICHEC-EC-EARTH	KNMI-RACMO22E	×	×	×	×
ICHEC-EC-EARTH	SMHI-RCA4	×	×	×	×
ICHEC-EC-EARTH	MOHC-HadREM3-GA7-05	×	×	-	×
MOHC-HadGEM2-ES	CNRM-ALADIN63	×	-	-	×
MOHC-HadGEM2-ES	CLMcom-CCLM4-8-17	×	-	×	×
MOHC-HadGEM2-ES	ICTP-RegCM4-6	×	×	-	×
MOHC-HadGEM2-ES	MOHC-HadREM3-GA7-05	×	×	-	×
IPSL-IPSL-CM5A-MR	DMI-HIRHAM5	×	-	-	×
IPSL-IPSL-CM5A-MR	SMHI-RCA4	×	-	×	×
IPSL-IPSL-CM5A-MR	IPSL-WRF384P	×	-	×	×
MPI-M-MPI-ESM-LR	CLMcom-CCLM4-8-17	×	×	×	×

MPI-M-MPI-ESM-LR	ICTP-RegCM4-6	×	×	-	×
MPI-M-MPI-ESM-LR	MPI-CSC-REMO2009	×	×	×	×
NCC-NorESM1-M	DMI-HIRHAM5	×	-	×	×
NCC-NorESM1-M	GERICS-REMO2015	×	×	×	×
NCC-NorESM1-M	IPSL-WRF381P	×	-	-	×

Table A1: Global and Regional Climate Models combinations driving the Explore2 Hydrological Models selected for *PFI* simulation in the 21st century

Mis en forme : Police :Italique

Appendix B: Sensitivity analysis of the duration of flow measurement interval used as input for the logistic regression to calculate the *PFI*

Mis en forme : Police :Italique




Figure B1: Kling-Gupta Efficiency (KGE) computed on results obtained by Leave One Year Out validations, testing the sensitivity to the time window of daily discharge used for calibrating the logistic regressions. Boxes represent the quartiles Q1 and Q3, the whiskers extend up to 1.5 times the *IQR-interquartile range* above Q3 and below Q1 and points located beyond the whiskers are displayed individually. The dashed line represents the median KGE of the [-6:j]- window

680 A sensitivity analysis was performed to fix the time window of daily discharge that optimizes the calibration of logistic regressions presented in Sect. 3.1. The calibration and validation were performed using the Leave One Year Out method, detailed in Sect. 3.3. The model performance was assessed using the Kling-Gupta Efficiency (KGE) (Gupta et al., 2009). The KGE values for the different HER2 are summarized using boxplots for each tested window size. The [j-6;j] window was selected because it corresponds to the highest median KGE score, the narrowest interquartile range, and the highest minimum
685 KGE.

Appendix C: Separation of dataset based on dry, intermediate and wet years

The data from ONDE sites used for validation were collected between May and September over 11 years from 2012 to 2022. The robustness of the model was assessed through a validation process involving three sets of dry, intermediate, and wet years. For each test, model calibration was performed using the years excluded from the validation set.

The hydrological years are distributed into three equal groups of hydrological years (dry years, intermediate years and wet years) according to the annual aridity index, calculated at the national scale. The aridity index AI was given by the ratio between the total annual precipitation and potential evapotranspiration from August 1 of the previous year to July 31 of the current year (Barrow, 1992; Figure A2.1). A set of dry years was formed using hydrological years where potential evapotranspiration exceeded annual precipitation ($AI < 1$ for 2017, 2019, and 2022). Two sets were then created: the four years with AI greater than 1.4 were classified as wet years (2012, 2015, 2020 and 2021), and the remaining four years with AI between 1.15 and 1.37 were classified as intermediate years (2013, 2014, 2016, 2018).




Figure C1: Aridity index for the hydrological years 2012 to 2022

Appendix D: Assignment of simulation points

720 After establishing the models' reliability through validation at secure monitoring stations, the next step involved extrapolation, which entailed linking these monitoring sites to the nearest simulation points in Explore 2 to project future hydrological scenarios while preserving the existing data structure. The proximity between the gauging station A and a simulation point in the Explore2 project is measured by the distance:

$$Dist(A, B) = \sqrt{(X(A) - X(B))^2 + (Y(A) - Y(B))^2 + (\alpha \times \Delta S_{rel})^2} \quad (D1)$$

725

Here, $(X(A), Y(A))$ and $(X(B), Y(B))$ are coordinates of A and B (in km), respectively, and $Surf(A)$ and $Surf(B)$ are the drainage areas of A and B, respectively that are used to compute the relative difference between the drainage areas (absolute value).

$$\Delta S_{rel} = 2 \times \frac{|Surf(A) - Surf(B)|}{|Surf(A) + Surf(B)|} \quad (D2)$$

730 The coefficient α is used to balance the importance of geographical distance and the relative difference in surface and was set to 100.

Appendix E: Location of the ONDE sites, the gauging stations and the Explore2 simulation points selected for each hydrological model

Commenté [T98]: RC2.7, RC1.8
Mis en forme : Titre 2




Figure E1: Location of the 3302 ONDE sites, the 1008 gauging stations selected from the HydroPortail database, and the Explore2 simulation points selected for each hydrological model (line 1). Maps on the second and third line respectively show the ratio between the sum of the catchment areas defined by the gauging stations or the simulation points and the surface area of the HER2 they intersect (line 2), and the density of gauging stations or simulation points (line 3).

Mis en forme : Légende

Appendix F: QUALYPSO method

QUALYPSO method is applied to the multi-scenario multi-model ensembles of projections of the national average of $mPFI_{7-10}$ ($mPFI_{7-10}[France]$), weighted by HER2 areas and obtained under RCP 2.6, 4.5 and 8.5 climate change scenarios. According to Evin et al. (2019), we assume that the variable $mPFI_{7-10\ i,j,k,l}^*(t)$ characterizes the change of $mPFI_{7-10}$ between a year t and the control year c (e.g., $c = 1990$ in this study), for a given combination of RCP scenario i , GCM j , RCM k , hydrological model l :

$$mPFI_{7-10\ i,j,k,l}^*(t) = mPFI_{7-10\ i,j,k,l}(t) - mPFI_{7-10\ i,j,k,l}(c)$$

The change of $mPFI_{7-10}$ can be split up into:

$$mPFI_{7-10\ i,j,k,l}^*(t) = \varphi_{i,j,k,l}^*(t) + \eta_{i,j,k,l}^*(t)$$

Where

$\varphi_{i,j,k,l}^*(t)$ is the climate change response of the RCP/GCM/RCM/Hydrological model combination,

$\eta_{i,j,k,l}^*(t)$ is the deviation from the climate change response for this RCP/GCM/RCM/Hydrological model combination, as a result of internal variability, representing natural and stochastic variability in the climate system.

For each time t , the climate change response characterized by $\varphi_{i,j,k,l}^*(t)$ of any RCP/GCM/RCM/Hydrological model combination can be expressed as:

$$\varphi_{i,j,k,l}^*(t) = \mu(t) + \alpha_i(t) + \beta_j(t) + \gamma_k(t) + \theta_l(t) + \varepsilon_{i,j,k,l}(t)$$

Where

$\mu(t)$ is the ensemble mean climate change response,

$\alpha_i(t)$ is the main effect of emission scenario i (i.e., the mean deviation of RCP scenario i from $\mu(t)$),

$\beta_j(t)$ is the main effect of GCM j ,

$\gamma_k(t)$ is the main effect of RCM k ,

$\theta_l(t)$ is the main effect of the hydrological model l ,

$\varepsilon_{i,j,k,l}(t) = \varphi_{i,j,k,l}^*(t) - \mu(t) - \alpha_i(t) - \beta_j(t) - \gamma_k(t) - \theta_l(t)$ correspond to residual terms. For each year t , $\varepsilon_{i,j,k,l}(t)$ are assumed to be independent and identically distributed over all scenarios, GCMs, RCMs, and hydrological models, and to follow normal distributions, with mean 0 and variance $\sigma^2(t)$.

Commenté [T99]: RC2.9

Mis en forme : Justifié, Interligne : 1.5 ligne

Mis en forme

Mis en forme

Mis en forme

Mis en forme

Mis en forme : Police :10 pt

Mis en forme : Police :10 pt

Mis en forme : Police :10 pt

Mis en forme : Police :10 pt

Mis en forme

Mis en forme : Justifié, Interligne : 1.5 ligne

Mis en forme

Mis en forme : Justifié, Interligne : 1.5 ligne

Mis en forme

Mis en forme

Mis en forme

Mis en forme

Mis en forme : Justifié, Interligne : 1.5 ligne

Mis en forme : Police : (Par défaut) +Corps (Times New Roman), Non Italique

Mis en forme : Police :10 pt

Mis en forme

Mis en forme

Mis en forme : Police :10 pt

Mis en forme : Police :10 pt

Mis en forme : Police :10 pt

Mis en forme

Mis en forme

Mis en forme

Mis en forme

Mis en forme

Mis en forme : Police :10 pt

If $\varphi_{i,j,k,l}^*(t)$ and $\eta_{i,j,k,l}^*(t)$ are assumed to be independent, the total variance of the change variable $mPFI_{7-10\ i,j,k,l}^*(t)$ is given by:

$$Var\left(mPFI_{7-10\ i,j,k,l}^*(t)\right) = Var\left(\varphi_{i,j,k,l}^*(t)\right) + Var\left(\eta_{i,j,k,l}^*(t)\right)$$

775 Where

$Var(\eta_{i,j,k,l}^*(t))$ is the uncertainty associated to internal variability of $mPFI_{7-10\ i,j,k,l}^*(t)$.

$Var(\varphi_{i,j,k,l}^*(t))$ is the uncertainty in the climate change response, calculated as the sum of different uncertainty components:

$$Var(\varphi_{i,j,k,l}^*(t)) = Var(\mu(t)) + Var(\alpha_i(t)) + Var(\beta_j(t)) + Var(\gamma_k(t)) + Var(\theta_l(t)) + Var(\varepsilon_{i,j,k,l}(t))$$

780 RCP uncertainty $Var(\alpha_i(t))$ quantifies the variability in $mPFI_{7-10\ i,j,k,l}^*(t)$ values across the RCP scenarios. Specifically, $\alpha_i(t)$ represents the differences between the means of $mPFI_{7-10\ i,j,k,l}^*(t)$ involving each scenario i and the overall mean of $mPFI_{7-10\ i,j,k,l}^*(t)$ across all scenarios. This component illustrates how differences between emission scenarios contribute to the total uncertainty in the projected $mPFI_{7-10\ i,j,k,l}^*(t)$.

Similarly, GCM, RCM and hydrological model contributions to $mPFI_{7-10\ i,j,k,l}^*(t)$ uncertainties are characterized by the variances for GCMs ($Var(\beta_j(t))$), RCMs ($Var(\gamma_k(t))$), and hydrological models ($Var(\theta_l(t))$).

785 Residual variability ($Var(\varepsilon_{i,j,k,l}(t))$) captures interaction effects and unexplained variability not accounted for by the main effects.

Mis en forme

Mis en forme : Justifié, Interligne : 1.5 ligne

Mis en forme

Mis en forme

Mis en forme : Police :10 pt

Mis en forme

Mis en forme : Police :10 pt

Mis en forme

Mis en forme : Police :10 pt

Mis en forme : Justifié, Interligne : 1.5 ligne

Mis en forme

Mis en forme : Police :10 pt

Mis en forme

Mis en forme : Police :10 pt

Mis en forme

Mis en forme

Mis en forme

Mis en forme

Mis en forme

Mis en forme : Police :10 pt

Mis en forme

Mis en forme

Mis en forme

Mis en forme : Justifié, Interligne : 1.5 ligne

Mis en forme : Police :10 pt

Mis en forme

Appendix EG: Metadata




Figure EG1: Frequency distribution of the number of ONDE sites, the 1008 gauging stations selected from the Hydroportail database, and the 1008 simulation points from Explore2 as a function of catchment area

NA: Missing values




Figure EG2: Frequency distribution of the number of ONDE sites, the 1008 gauging stations selected from the Hydroportail database, and the 1008 simulation points from Explore2 as a function of elevation

NA: Missing value




Figure G3: Strahler index of ONDE sites

NA: Missing value

Appendix FH: Model performance using observed discharge data from gauging stations as predictors

Over the observation period (2012-2022), logistic regression models estimate the PFI values at HER2 scale (Figure-Fig. 4). The median observed PFI across all HER2 and all campaigns is 14.3% (Q1-Q3IQR: 8.4-20.3) while the logistic regression models yield a median value of 14.4% (Q1-Q3IQR: 8.5-20.4). With results of the leave-one-year-out cross-validation, the models explain 73% of the PFI variability according to the NSE (Q1-Q3IQR: 64-84%) (Table 1; Figure-Fig. HF2), the KGE exceeds 80% in 50 out of 75 HER2 (Figure-Fig. HF2) and the bias is very low.

The models are also able to describe the inter-annual variability with the alternation of dry and wet years (Figure-Fig. HF1). The median KGE and NSE scores remain above 0.71 during k-fold validations considering dry, intermediate, and wet years for calibration. RMSE and MAE values do not increase much when the calibration dataset is stratified based on climate conditions, which indicates that the proposed model is quite robust to climate variations under current conditions.

Mis en forme : Police :Italique

Mis en forme : Police :Italique

Mis en forme : Police :Italique




Figure FH1: Time series of PFI in HER2 13, 57, 81 and 105. Points and lines respectively represent the PFI derived from the ONDE network and the PFI estimated by the logistic regression models using discharge data from gauging stations during the calibration period (2012-2022).

Mis en forme : Police :Italique

Mis en forme : Police :Italique

Mis en forme : Police :Italique




Figure FH2: Kling-Gupta Efficiency (KGE, line 1), Nash-Sutcliffe model efficiency coefficient (NSE, line 2), Bias (line 3), Mean Absolute Error (MAE, line 4), Root-mean-square error (RMSE, line 5) over the calibration period (2012-2022)

845

Validation years	<i>PFI</i> observed on ONDE network (%)	Predicted <i>PFI</i> (%)	Nash Sutcliffe Efficiency (NSE, unitless)	Kling Gupta Efficiency (KGE, unitless)	Bias (unitless)	Mean Absolute Error (MAE, unitless)	Root Mean Square Error (RMSE, unitless)	Mis en forme : Police :Italique, Anglais (États-Unis)
								Mis en forme : Police :Italique
Leave One Year Out (2012-2022)	14.3	14.4	0.79	0.85	-0.02	0.10	0.07	
	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	
	8;20	9;20	0.6;0.8	0.8;0.9, min-	min-max	0.06;0.13 min-	0.05;0.09	
	min-max	min-max	min-max	max 0.3;1.0	-0.68;0.70	max	min-max	
	3;37	2;37	0.2;0.9			0.02;0.32	0.03;0.15	
Wet years (2013, 2014, 2016, 2018)	8.3	8.7	0.76	0.71	0.17	0.05	0.05	
	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	
	5;15	5;13	0.6;0.9	0.6;0.8	min-max	0.03;0.09 min-	0.03;0.08	
	min-max	min-max	min-max	min-max	-7.09;7.32	max 0.01;0.23	min-max	
	1;29	0;35	-1.7;1.0	-4.0;1.0			0.02;0.15	
Intermediate years (2012, 2015, 2020, 2021)	12.9	13.6	0.76	0.76	0.21	0.07	0.06	
	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	
	7;19	7;19	0.6;0.8	0.7;0.9 min-	min-max	0.04;0.09 min-	0.04;0.09	
	min-max	min-max	min-max	max	-8.85;5.76	max 0;0.25	min-max	
	1;35	2;31	-0.7;1.0	-0.2;1.0			0;0.18	
Dry years (2017, 2019, 2022)	21.5	21.4	0.7	0.75	-0.71	0.09	0.08	
	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	IQQ1-Q3	
	14;30	14;32	0.5;0.8	0.6;0.8 min-	min-max	0.06;0.12	0.06;0.12	
	min-max	min-max	min-max	max	-15.13;12.78	min-max	min-max	
	2;51	4;49	-0.1;1.0	-0.6;0.9		0.01;0.43	0.03;0.24	

Table **FH1**: Validation results of drying probability predictions at the HER2 scale using observed flows from the 1008 gauging stations from the HYDRO database. The results correspond to the inter-HER medians, quartiles, minimum and maximum values. The Leave One Year Out analysis results are obtained by averaging the validation metrics computed for each year.

Drying probability: Percentage of ONDE sites in a dry state computed for each HER2, averaged by month; **FDC**: **Flow Duration Curve**; **Q1-Q3**: **first and third Interquartiles range**; Min: minimum; Max: maximum

Appendix G1: Model performance using discharge data simulated with SAFRAN





Figure G1: Nash–Sutcliffe model efficiency coefficient (NSE, line 1), bias (line 2), Mean Absolute Error (MAE, line 3), Root-mean-square error (RMSE, line 4) assessing the calibration of logistic regression using discharge data simulated with SAFRAN data available during the calibration period (2012–2022)

Mis en forme : Police :Italique

860



865

Figure HJ1: Ensemble median of changes in $mPFI_{7-10}$ (columns 1 to 3), and ensemble median shift of the start date T_s-T_f (columns 4 and 5) and the end date T_e (columns 6 and 7) of the dry period over the two periods 2041-2070H1 and 2070-2099H2 under the three RCPs for the CTRIP hydrological model, relative to the baseline period 1976-2005H0. The shift, expressed in week, takes one a positive value when the duration of the dry period increases. Grey HER2s had no period with a $PFI > 20\%$ during the reference period.

- Mis en forme : Police :Italique
- Mis en forme : Police :Italique, Indice
- Mis en forme : Police :Italique
- Mis en forme : Police :Italique
- Mis en forme : Police :Italique

870

875




Figure HJ2: Ensemble median of changes in $mPFI_{2-10}$ (columns 1 to 3), and ensemble median shift of the start date T_{fs} (columns 4 and 5) and the end date T_{le} (columns 6 and 7) of the dry period over the two periods 2041-2070H1 and 2070-2099H2 under the three RCPs for the J2000 hydrological model, relative to the baseline period 1976-2005H0. The shift, expressed in week, takes one a positive value when the duration of the dry period increases. Grev HER2s had no period with a $PFI > 20\%$ during the reference period.

Mis en forme : Police :Italique

Mis en forme : Police :Italique, Indice

Mis en forme : Police :Italique

Mis en forme : Police :Italique




Figure HJ3: Ensemble median of changes in $mPFI_{7-10}$ (columns 1 to 3), and ensemble median shift of the start date T_s (columns 4 and 5) and the end date T_e (columns 6 and 7) of the dry period over the two periods 2041-2070H1 and 2070-2099H2 under the three RCPs for the ORCHIDEE hydrological model, relative to the baseline period 1976-2005H0. The shift, expressed in week, takes one a positive value when the duration of the dry period increases. Grey HER2s had no period with a $PFI > 20\%$ during the reference period.

Mis en forme : Police :Italique

Mis en forme : Police :Italique, Indice

Mis en forme : Police :Italique

Mis en forme : Police :Italique




Figure HJ4: Ensemble median of changes in $mPFI_{7-10}$ (columns 1 to 3), and ensemble median shift of the start date T_s (columns 4 and 5) and the end date T_e (columns 6 and 7) of the dry period over the two periods 2041-2070 and 2070-2099 under the three RCPs for the SMASH hydrological model, relative to the baseline period 1976-2005. The shift, expressed in week, takes a positive value when the duration of the dry period increases. Grey HER2s had no period with a $PFI > 20\%$ during the reference period.

Mis en forme : Police :Italique

Mis en forme : Police :Italique, Indice

Mis en forme : Police :Italique


Mis en forme : Police :Italique

910 **Narratives**

The following figures present time series of the evolution of *PFI* dynamics at different horizons and under different RCP scenarios for each hydrological model in HER2 12, 57, 81 and 105. Four contrasting scenarios were highlighted among the Explore2 climate projections to illustrate a diversity of potential changes under RCP 8.5. These story lines range from "Strong warming and strong summer (and annual) drying", selected as one of the most extreme climate projections, to "Moderate warming and precipitation change" with less pronounced alterations, with two alternative projections named "Dry all year, reduced winter recharge" and "Hot and humid all seasons". They are also illustrated here to present distinct hydrological nuances.

Mis en forme : Police :Italique

920



925

930

935 **Figure HJ5:** Time series of the evolution of *PFI* dynamics at different horizons under different RCP scenarios for CTRIP hydrological models in HER2 12

Mis en forme : Police :Italique

Average proportion of zero flows smoothed over 5 days Hydrological model: GRSD - HER2 12




Figure HJ6: Time series of the evolution of PFI dynamics at different horizons under different RCP scenarios for GRSD hydrological models in HER2 12

Mis en forme : Police :Italique

Average proportion of zero flows smoothed over 5 days Hydrological model: ORCHIDEE - HER2 12




Figure HJ7: Time series of the evolution of PFI dynamics at different horizons under different RCP scenarios for ORCHIDEE hydrological models in HER2 12

Mis en forme : Police :Italique

Average proportion of zero flows smoothed over 5 days Hydrological model: SMASH - HER2 12




Figure HJ8: Time series of the evolution of PFI dynamics at different horizons under different RCP scenarios for SMASH hydrological models in HER2 12

Mis en forme : Police :Italique

Average proportion of zero flows smoothed over 5 days Hydrological model: CTRIP - HER2 57




Figure HJ9: Time series of the evolution of PFI dynamics at different horizons under different RCP scenarios for CTRIP hydrological models in HER2 57

Mis en forme : Police :Italique

1015

1020

1025


1030

1035

1040

1045

Figure HJ10: Time series of the evolution of PFI dynamics at different horizons under different RCP scenarios for GRSD hydrological models in HER2 57



Mis en forme : Police :Italique

Figure HJ11: Time series of the evolution of PFI dynamics at different horizons under different RCP scenarios for ORCHIDEE hydrological models in HER2 57

Mis en forme : Police :Italique

Average proportion of zero flows smoothed over 5 days Hydrological model: SMASH - HER2 57




Figure HJ12: Time series of the evolution of PFI dynamics at different horizons under different RCP scenarios for SMASH hydrological models in HER2 57

Mis en forme : Police :Italique

Average proportion of zero flows smoothed over 5 days Hydrological model: CTRIP - HER2 81




Figure HJ13: Time series of the evolution of PFI dynamics at different horizons under different RCP scenarios for CTRIP hydrological models in HER2 81

Mis en forme : Police :Italique

Average proportion of zero flows smoothed over 5 days Hydrological model: GRSD - HER2 81




Figure HJ14: Time series of the evolution of *PFI* dynamics at different horizons under different RCP scenarios for GRSD hydrological models in HER2 81

Mis en forme : Police :Italique

Average proportion of zero flows smoothed over 5 days Hydrological model: J2000 - HER2 81




Figure HJ15: Time series of the evolution of *PFI* dynamics at different horizons under different RCP scenarios for J2000 hydrological models in HER2 81

Mis en forme : Police :Italique



1120

1125



1140

1145

1150

1155

1160

1165

1170

1175

1180




Figure HJ18: Time series of the evolution of PFI dynamics at different horizons under different RCP scenarios for CTRIP hydrological models in HER2 105

Mis en forme : Police :Italique




Figure HJ19: Time series of the evolution of PFI dynamics at different horizons under different RCP scenarios for GRSD hydrological models in HER2 105

Mis en forme : Police :Italique

1185

1190

1195

1200

1205

1210

1215




Figure HJ20: Time series of the evolution of *PFI* dynamics at different horizons under different RCP scenarios for ORCHIDEE hydrological models in HER2 105

Mis en forme : Police :Italique




Figure HJ21: Time series of the evolution of *PFI* dynamics at different horizons under different RCP scenarios for SMASH hydrological models in HER2 105

Mis en forme : Police :Italique

Appendix **IK**: Agreement between changes under RCP 4.5




Figure **IK1**: Agreement between projections of $mPFI_{7-10}$ for each hydrological model and inter-model agreement on the change signal of $mPFI_{7-10}$ under the RCP 4.5 scenario.

Code availability

The codes are available at <https://github.com/tjaouen/PostDocINRAE>

Data availability

The Explore2 streamflow projections are available at <https://entrepot.recherche.data.gouv.fr/dataverse/explore2> and are described in detail by Sauquet et al.⁷ (in prep).

ONDE dataset is available at <https://onde.eaufrance.fr/acces-aux-donnees>.

Mis en forme : Police :Italique

Mis en forme : Police :Italique, Indice

Mis en forme : Police :Italique

Mis en forme : Police :Italique, Indice

Commenté [T100]: RC1.15

Author contribution

Dr. Tristan Jaouen, Dr. Lionel Benoit and Dr. Eric Sauquet designed the experiments and Dr. Tristan Jaouen carried them out. Louis Héraut organized the data from the Explore2 project and made it available. Dr. Tristan Jaouen and Dr. Eric Sauquet developed the model code. Dr. Tristan Jaouen performed the simulations. Dr. Tristan Jaouen prepared the manuscript with contributions from all co-authors.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

This research was funded by the Rhône-Méditerranée-Corse Water Agency under the supervision of Dr. Benoît Terrier. Scientific oversight was provided within INRAE by Dr. Éric Sauquet and Dr. Lionel Benoit.
This work builds on an initial exploration led by Dr. Aurélien Beaufort and Dr. Éric Sauquet, along with contributions from all participants in the Explore2 project. The archiving and management of Explore2 data, expertly handled by Louis Héraut, also played a crucial role in this study.
Additionally, this research directly applies the QUALYPSO method, developed and adapted to Explore2 by Dr. Guillaume Evin.
We extend our deep gratitude to all these individuals for their invaluable contributions to this work.

References

Acuña, V., Datry, T., Marshall, J., Barceló, D., Dahm, C. N., Ginebreda, A., McGregor, G., Sabater, S., Tockner, K., and Palmer, M. A.: Why should we care about temporary waterways?, Science, 343, 1080-1081, <https://doi.org/10.1126/science.1246666>, 2014.

Addor, N., Nearing, G., Prieto, C., Newman, A. J., le Vine, N., and Clark, M. P.: A Ranking of Hydrological Signatures Based on Their Predictability in Space, Water Resources Research, 54, 8792–8812, <https://doi.org/10.1029/2018WR022606>, 2018.

Aitken, G., Beevers, L., Parry, S., and Facer-Childs, K.: Partitioning model uncertainty in multi-model ensemble river flow projections, Climatic Change, 176, 153, <https://doi.org/10.1007/s10584-023-03621-1>, 2023.

Mis en forme : Titre 1

Barrow, C. J.: World atlas of desertification (United Nations Environment Programme), edited by N. Middleton and D. S. G. Thomas. Edward Arnold, London, 1992. isbn 0 340 55512 2, £89.50 (hardback), ix + 69 pp., Land Degrad. Dev., 3, 249-249, <https://doi.org/10.1002/ldr.3400030407>, 1992.

1260 Beaufort, A., Lamouroux, N., Pella, H., Detry, T., and Sauquet, E.: Extrapolating regional probability of drying of headwater streams using discrete observations and gauging networks, Hydrol. Earth Syst. Sci., 22, 3033–3051, <https://doi.org/10.5194/hess-22-3033-2018>, 2018.

1265 [Beaufort, A., Carreau, J., and Sauquet, E.: A classification approach to reconstruct local daily drying dynamics at headwater streams, Hydrological Processes, 33, 1896–1912, <https://doi.org/10.1002/hyp.13445>, 2019.](#)

[Belemtougri, A. P., Ducharme, A., Tazen, F., Oudin, L., and Karambiri, H.: Understanding key factors controlling the duration of river flow intermittency: Case of Burkina Faso in West Africa, Journal of Hydrology: Regional Studies, 37, 100908, <https://doi.org/10.1016/j.ejrh.2021.100908>, 2021.](#)

1270 [Bertassello, L. E., Durighetto, N., and Botter, G.: Eco-hydrological modelling of channel network dynamics - Part 2: Application to metapopulation dynamics, Royal Society Open Science, 9, 220945, <https://doi.org/10.1098/rsos.220945>, 2022.](#)

1275 [Botter, G., and Durighetto, N.: The Stream Length Duration Curve: A Tool for Characterizing the Time Variability of the Flowing Stream Length, Water Resources Research, 56, e2020WR027282, <https://doi.org/10.1029/2020WR027282>, 2020.](#)

[Bourke, S. A., Degens, B., Searle, J., de Castro Tayer, T., and Rothery, J.: Geological permeability controls streamflow generation in a remote, ungauged, semi-arid drainage system, Journal of Hydrology: Regional Studies, 38, 100956, <https://doi.org/10.1016/j.ejrh.2021.100956>, 2021.](#)

1280 Carlson, S. M., Ruhf, A., Bogan, M. T., Hazard, C. W., Ayers, J., Grantham, T. E., Batalla, R. J., and Garcia, C.: Losing flow in free-flowing Mediterranean-climate streams, Front Ecol Environ., 22, e2737, <https://doi.org/10.1002/fee.2737>, 2024.

1285 [Clarke, A., Mac Nally, R., Bond, N., and Lake, P. S.: Macroinvertebrate diversity in headwater streams: a review, Freshwater Biology, 53, 1707–1721, <https://doi.org/10.1111/j.1365-2427.2008.02041.x>, 2008.](#)

Dai, Z., Amatya, D. M., Sun, G., Trettin, C. C., Li, C., and Li, H.: Climate variability and its impact on forest hydrology on south Carolina coastal plain, USA, Atmosphere. 2011, 2, 330-357, <https://doi.org/10.3390/atmos2030330>, 2011.

1290 Dakhloui, H., Ruelland, D., Trambay, Y., and Bargaoui, Z.: Evaluating the robustness of conceptual rainfall-runoff models under climate variability in northern Tunisia, *Journal of Hydrology*, 550, 201-217, <https://doi.org/10.1016/j.jhydrol.2017.04.032>, 2017.

De Girolamo, A. M., Barca, E., Leone, M., and Lo Porto, A.: Impact of long-term climate change on flow regime in a Mediterranean basin, *Journal of Hydrology: Regional Studies*, 41, 101061. <https://doi.org/10.1016/j.ejrh.2022.101061>, 2022.

De Lavenne, A., Thirel, G., Andréassian, V., Perrin, C., and Ramos, M.-H.: Spatial variability of the parameters of a semi-distributed hydrological model, *Proc. IAHS*, 373, 87–94, <https://doi.org/10.5194/piahs-373-87-2016>, 2016.

1300 Decharme, B., Delire, C., Minvielle, M., Colin, J., Vergnes, J.-P., Alias, A., et al.: Recent changes in the ISBA-CTRIP land surface system for use in the CNRM-CM6 climate model and in global off-line hydrological applications, *Journal of Advances in Modelling Earth Systems*, 11, 1207–1252, <https://doi.org/10.1029/2018MS001545>, 2019.

Delso, J., Magdaleno, F., and Fernández-Yuste, J. A.: Flow patterns in temporary rivers: a methodological approach applied to southern Iberia, *Hydrological Sciences Journal*, 62, 1551-1563, <https://doi.org/10.1080/02626667.2017.1346375>, 2017.

Dhungel, S., Tarboton, D. G., Jin, J., and Hawkins, C. P.: Potential effects of climate change on ecologically relevant streamflow regimes, *River Res. Applic.*, 32, 1827-1840, <https://doi.org/10.1002/rra.3029>, 2016.

1310 Döll, P., ~~Abbasi, M., Messenger, M. L., Trautmann, T., Lehner, B., Lamouroux, N., Hasan, H. M. M., Schulze, K., Gerdener, H., Börger, L., Shadkam, S., Aekermann, S., Hosseini Moghari, S. M., Müller-Schmied, H., Güntner, A., and Kutsche, J.: Streamflow Intermittence in Europe: Estimating High-Resolution Monthly Time Series by Downscaling of Simulated Runoff and Random Forest Modeling~~
~~Leveraging multi-variable observations to reduce and quantify the output uncertainty of a global hydrological model: evaluation of three ensemble-based approaches for the Mississippi River basin, *Water Resources Research*, 60, e2023WR036900, <https://doi.org/10.1029/2023WR036900>, 2024.~~
1315 *Hydrol. Earth Syst. Sci.*, 28, 2259-2295, <https://doi.org/10.5194/hess-28-2259-2024>, 2024.

Döll, P., and Schmied, H. M.: How is the impact of climate change on river flow regimes related to the impact on mean annual runoff? A global-scale analysis, *Environ. Res. Lett.*, 7, 14037-11, <https://doi.org/10.1088/1748-9326/7/1/014037>, 2012.

1320 Döll, P., and Zhang, J.: Impact of climate change on freshwater ecosystems: a global-scale analysis of ecologically relevant river flow alterations, *Hydrol. Earth Syst. Sci.*, 14, 783–799, <https://doi.org/10.5194/hess-14-783-2010>, 2010.

- 1325 [Douveille, H., Raghavan, K., Renwick, J., Allan, R.P., Arias, P.A., Barlow, M., Cerezo-Mota, R., Cherchi, A., Gan, T.Y., Gergis, J., Jiang, D., Khan, A., Pokam Mba, W., Rosenfeld, D., Tierney, J., and Zolina, O.: Water Cycle Changes. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* \[Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S.L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M.I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J.B.R., Maycock, T.K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B.\], Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1055–1210, doi:10.1017/9781009157896.010, 2021.](#)
- 1330 [Durighetto, N., Bertassello, L. E., and Botter, G.: Eco-hydrological modelling of channel network dynamics - Part 1: Stochastic simulation of active stream expansion and retraction, *Royal Society Open Science*, 9, 220944, <https://doi.org/10.1098/rsos.220944>, 2022.](#)
- 1335 Evin, G., Hingray, B., Blanchet, J., Eckert, N., Morin, S., and Verfaillie, D.: Partitioning uncertainty components of an incomplete ensemble of climate projections using data augmentation, *Journal of Climate*, 32, 2423–2440, <https://doi.org/10.1175/jcli-d-18-0606.1>, 2019.
- 1340 [Evin, G., Hingray, B., Reverdy, A., Ducharne, A., and Sauquet, E.: Ensemble de projections Explore2 : Changements moyens et incertitudes associées, *Recherche Data Gouv*, 90 pp., <https://doi.org/10.57745/KWH320>, 2024.](#)
~~Evin, G., Hingray, B., Ducharne, A., and Sauquet, E.: Ensemble de projections Explore2 : Changements moyens et incertitudes associées, *INRAE—IGE; CNRS—IGE; CNRS—IPSL; INRAE*, <https://hal.science/hal-04609542>, 2024.~~
- 1345 Evin, G., Somot, S., and Hingray, B.: Balanced estimate and uncertainty assessment of European climate change using the large EURO-CORDEX regional climate model ensemble, *Earth System Dynamics*, 12, 1543–1569, <https://doi.org/10.5194/esd-12-1543-2021>, 2021.
- 1350 Finn, D., Bonada, N., Múrria, C., and Hughes, J.: Small but mighty: headwaters are vital to stream network biodiversity at two levels of organization, *Journal of the North American Benthological Society*, 30, 963–980, <https://doi.org/10.1899/11-012.1>, 2011.
- 1355 Gallart, F., Prat, N., García-Roger, E. M., Latron, J., Rieradevall, M., Llorens, P., Barberá, G. G., Brito, D., De Girolamo, A. M., Lo Porto, A., Buffagni, A., Erba, S., Neves, R., Nikolaidis, N. P., Perrin, J. L., Querner, E. P., Quiñero, J. M., Tournoud, M. G., Tzoraki, O., Skoulikidis, N., Gómez, R., Sánchez-Montoya, M. M., and Froebrich, J.: A novel approach to analysing the regimes of temporary streams in relation to their controls on the composition and structure of aquatic biota, *Hydrol. Earth Syst. Sci.*, 16, 3165–3182, <https://doi.org/10.5194/hess-16-3165-2012>, 2012.

- Geris, J., Tetzlaff, D., and Soulsby, C.: Resistance and resilience to droughts: Hydropedological controls on catchment storage and run-off response, *Hydrological Processes*, 29, 4579–4593, <https://doi.org/10.1002/hyp.10480>, 2015.
- Giezendanner, J., Benettin, P., Durigetto, N., Botter, G., and Rinaldo, A.: A Note on the Role of Seasonal Expansions and Contractions of the Flowing Fluvial Network on Metapopulation Persistence, *Water Resources Research*, 57, e2021WR029813, <https://doi.org/10.1029/2021WR029813>, 2021.
- González-Ferreras, A. M., and Barquín, J.: Mapping the temporary and perennial character of whole river networks, *Water Resources Research*, 53, 6709–6724, <https://doi.org/10.1002/2017WR020390>, 2017.
- Gudmundsson, L., Boulange, J., Do, H. X., Gosling, S. N., Grillakis, M. G., Koutroulis, A. G., Leonard, M., Liu, J., Müller Schmied, H., Papadimitriou, L., Pokhrel, Y., Seneviratne, S. I., Satoh, Y., Thiery, W., Westra, S., Zhang, X., and Zhao, F.: Globally observed trends in mean and extreme river flow attributed to climate change, *Science*, 371, 1159–1162, <https://doi.org/10.1126/science.aba3996>, 2021.
- Gupta, H., Kling, H., Yilmaz, K., and Martinez, G.: Decomposition of the mean squared error and NSE performance criteria: Implications for improving hydrological modelling, *Journal of Hydrology*, 377, 80–91, <https://doi.org/10.1016/j.jhydrol.2009.08.003>, 2009.
- Halloran, L., Millwater, J., Hunkeler, D., and Arnoux, M.: Climate change impacts on groundwater discharge-dependent streamflow in an alpine headwater catchment, *Science of The Total Environment*, 902, 166009, <https://doi.org/10.1016/j.scitotenv.2023.166009>, 2023.
- Hammond, J. C., and Fleming, B. J.: Evaluating low flow patterns, drivers and trends in the Delaware River Basin, *Journal of Hydrology*, 598, 126246, <https://doi.org/10.1016/j.jhydrol.2021.126246>, 2021.
- Huang, P., Ducharme, A., Rinchuso, L., Polcher, J., Baratgin, L., Bastrikov, V., and Sauquet, E.: Multi-objective calibration and evaluation of the ORCHIDEE land surface model over France at high resolution, *EGUsphere* [preprint], <https://doi.org/10.5194/egusphere-2024-445>, 2024.
- Jaeger, K. L., Olden, J. D., and Pelland, N. A.: Climate change poised to threaten hydrologic connectivity and endemic fishes in dryland streams, *Proceedings of the National Academy of Sciences of the United States of America*, 111, 13894–13899, <https://doi.org/10.1073/pnas.1320890111>, 2014.

- 1395 [Jaeger, K. L., Sando, R., McShane, R. R., Dunham, J. B., Hockman-Wert, D. P., Kaiser, K. E., Hafen, K., Risley, J. C., and Blasch, K. W.: Probability of Streamflow Permanence Model \(PROSPER\): A spatially continuous model of annual streamflow permanence throughout the Pacific Northwest, Journal of Hydrology X, 2, 100005, <https://doi.org/10.1016/j.hydroa.2018.100005>, 2019.](#)
- 1400 Jay-Allemand, M., Javelle, P., Gejadze, I., Arnaud, P., Malaterre, P.-O., Fine, J.-A., and Organde, D.: On the potential of variational calibration for a fully distributed hydrological model: application on a Mediterranean catchment, Hydrol. Earth Syst. Sci., 24, 5519–5538, <https://doi.org/10.5194/hess-24-5519-2020>, 2020.
- 1405 [Lapides, D. A., Leclerc, C. D., Moidu, H., Dralle, D. N., and Hahm, W. J.: Variability of stream extents controlled by flow regime and network hydraulic scaling, Hydrological Processes, 35, e14079, <https://doi.org/10.1002/hyp.14079>, 2021.](#)
- [Larned, S. T., Datry, T., Arscott, D. B., & Tockner, K.: Emerging concepts in temporary-river ecology. In Freshwater Biology, 55, 717–738, <https://doi.org/10.1111/j.1365-2427.2009.02322.x>, 2010.](#)
- 1410 Leigh, C., and Datry, T.: Drying as a primary hydrological determinant of biodiversity in river systems: A broad-scale analysis, Ecography, 40, 487–499, <https://doi.org/10.1111/ecog.02230>, 2017.
- 1415 [Leleu, I., Tonnelier, I., Puechberty, R., Gouin, P., Viquendi, I., Cobos, L., Foray, A., Baillon, M., and Ndim, P.-O.: Re-founding the national information system designed to manage and give access to hydrometric data, La Houille Blanche, 100, 25–32, <https://doi.org/10.1051/lhb/2014004>, 2014.](#)
- [Messenger, M.L., Lehner, B., Cockburn, C., Lamouroux, N., Pella, H., Snelder, T., Tockner, K., Trautmann, T., Watt, C., Datry, T.: Global prevalence of non-perennial rivers and streams, Nature, 594, 391–397, <https://doi.org/10.1038/s41586-021-03565-5>, 2021.](#)
- 1420 Meyer, J., Strayer, D., Wallace, J., Eggert, S., Helfman, G., and Leonard, N.: The contribution of headwater streams to biodiversity in river networks, JAWRA Journal of the American Water Resources Association, 43, 86–103, <https://doi.org/10.1111/j.1752-1688.2007.00008.x>, 2007.

Morel, M., Pella, H., Branger, F., Sauquet, E., Grenouillet, G., Côte, J., Braud, I., and Lamouroux, N.: Catchment-scale applications of hydraulic habitat models: Climate change effects on fish, *Ecohydrology*, 16, e2513, <https://doi.org/10.1002/eco.2513>, 2023.

Nash, J. E., and Sutcliffe, J. V.: River flow forecasting through conceptual models part I-A discussion of principles, *Journal of Hydrology*, 10, 282-290, [http://dx.doi.org/10.1016/0022-1694\(70\)90255-6](http://dx.doi.org/10.1016/0022-1694(70)90255-6), 1970.

Nowak C., and Durozoi B.: Guide de dimensionnement et de mise en œuvre du suivi national des étiages estivaux, ONEMA, 33 p., 2012.

Pumo, D., Caracciolo, D., Viola, F., and Noto, L. V.: Climate change effects on the hydrological regime of small non-perennial river basins, *Sci Total Environ.*, 542, 76-92, <https://doi.org/10.1016/j.scitotenv.2015.10.109>, 2016.

Reidy Liermann, C., Olden, J., Beechie, T., Kennard, M., Skidmore, P., Konrad, C., and Imaki, H.: Hydrogeomorphic classification of Washington State rivers to support emerging environmental flow management strategies, *River Research and Applications*, 28, 1340-1358, <https://doi.org/10.1002/rra.1541>, 2012.

Reynolds, L. V., Shafroth, P. B., and Poff, L.: Modeled intermittency risk for small streams in the Upper Colorado River Basin under climate change, *Journal of Hydrology*, 523, 768-780, <https://doi.org/10.1016/j.jhydrol.2015.02.025>, 2015.

Ribes, A., Boé, J., Qasmi, S., Dubuisson, B., Douville, H., and Terray, L.: An updated assessment of past and future warming over France based on a regional observational constraint, *Earth Syst. Dynam.*, 13, 1397–1415, <https://doi.org/10.5194/esd-13-1397-2022>, 2022.

Rutkowska, A., Osuch, M., Żelazny, M., Banasik, K., and Klimek, M.: Climatic and anthropogenic drivers of zero-flow events in intermittent rivers in Poland, *Journal of Water and Land Development*, 57, 52-61, <https://doi.org/10.24425/jwld.2023.145335>, 2023.

Sando, R., Jaeger, K. L., Farmer, W. H., Barnhart, T. B., McShane, R. R., Welborn, T. L., Kaiser, K. E., Hafen, K. C., Blasch, K., York, B., and Shallcross, A.: Predictions and drivers of sub-reach-scale annual streamflow permanence for the upper Missouri River basin: 1989–2018, *Journal of Hydrology X*, 17, 100138, <https://doi.org/10.1016/j.hydroa.2022.100138>, 2022.

Sarremejane, R., Cañedo-Argüelles, M., Prat, N., Mykrä, H., Muotka, T., and Bonada, N.: Do metacommunities vary through time? Intermittent rivers as model systems, *Journal of Biogeography*, 44, 2752-2763, <https://doi.org/10.1111/jbi.13077>, 2017.

1460 Sauquet, E., Beaufort, A., Sarremejane, R., and Thirel, G.: Predicting flow intermittence in France under climate change, Hydrological Sciences Journal, 66, 2046-2059, <https://doi.org/10.1080/02626667.2021.1963444>, 2021.

1465 (in prep.) Sauquet, E., Evin, G., Siauue, S., Aissat, R., Arnaud, P., Berel, M., Bonneau, J., Branger, F., Colleoni, F., Ducharne, A., Gailhard, J., Habets, F., Hendrickx, F., Hérault, L., Hingray, B., Huang, P., Jaouen, T., Jeantet, A., Lanini, S., Le Lay, M., Magand, C., Mimeau, L., Monteil, C., Munier, S., Perrin, C., Robelin, O., Rousset, F., Soubeyroux, J.-M., Strohmenger, L., Thirel, G., Tocquer, F., Tramblay, Y., Vergnes, J.-P., Vidal, J.-P.: A large transient multi-scenario multi-model ensemble of future streamflows and groundwater projections in France, Hydrology and Earth System Sciences, Data description paper.

1470 Sauquet, E., Evin, G., Siauue, S., Bornancin-Plantier, A., Jacquin, N., Arnaud, P., Bérel, M., Bernus, S., Bonneau, J., Branger, F., Caballero, Y., Colléoni, F., Collet, L., Corre, L., Drouin, A., Ducharne, A., Fournier, M., Gailhard, J., Habets, F., Hendrickx, F., Hérault, L., Hingray, B., Huang, P., Jaouen, T., Jeantet, A., Lanini, S., Le Lay, M., Loudin, S., Magand, C., Marson, P., Mimeau, L., Monteil, C., Munier, S., Perrin, C., Robin, Y., Rousset, F., Soubeyroux, J.-M., Strohmenger, L., Thirel, G., Tocquer, F., Tramblay, Y., Vergnes, J.-P., Vidal, J.-P., and Vrac, M.: Messages et enseignements du projet Explore2, Recherche Data Gouv, 24 pp., <https://doi.org/10.57745/J3XIPW>, 2024.

1475 Scheller, M., Van Meerveld, I., and Seibert, J.: How well can people observe the flow state of temporary streams?, Frontiers in Environmental Science, 12, <https://doi.org/10.3389/fenvs.2024.1352697>, 2024.

1480 Schneider, C., Laizé, C. L. R., Acreman, M. C., and Flörke, M.: How will climate change modify river flow regimes in Europe?, Hydrol. Earth Syst. Sci., 17, 325–339, <https://doi.org/10.5194/hess-17-325-2013>, 2013.

Schneider, A., Jost, A., Coulon, C., Silvestre, M., Théry, S., and Ducharne, A.: Global-scale river network extraction based on high-resolution topography and constrained by lithology, climate, slope, and observed drainage density, Geophysical Research Letters, 44, 2773–2781, <https://doi.org/10.1002/2016GL071844>, 2017.

1485 Sefton, C. E.M., Parry, S., England, J., and Angell, G.: Visualising and quantifying the variability of hydrological state in intermittent rivers, Fundamental and Applied Limnology [in special issue: The ecohydrology of temporary streams], 193, 21–38, <https://doi.org/10.1127/fal/2019/1149>, 2019.

1490 Shaw, S. B., Bonville, D. B., and Chandler, D. G.: Combining observations of channel network contraction and spatial discharge variation to inform spatial controls on baseflow in Birch Creek, Catskill Mountains, USA, Journal of Hydrology: Regional Studies, 12, 1–12, <https://doi.org/10.1016/j.ejrh.2017.03.003>, 2017.

Commenté [T101]: RC1.1, RC2.2

- 1495 [Snelder, T. H., Datry, T., Lamouroux, N., Larned, S. T., Sauquet, E., Pella, H., and Catalogne, C.: Regionalization of patterns of flow intermittence from gauging station records, *Hydrology and Earth System Sciences*, 17, 2685–2699, <https://doi.org/10.5194/hess-17-2685-2013>, 2013.](#)
- Strohmenger, L., Sauquet, E., Bernard, C., Bonneau, J., Branger, F., Bresson, A., Brigode, P., Buzier, R., Delaigue, O., Devers, A., Evin, G., Fournier, M., Hsu, S.-C., Lanini, S., De Lavenne, A., Lemaitre-Basset, T., Magand, C., Mendoza Guimarães, G., Mentha, M., Munier, S., Perrin, C., Podechard, T., Rouchy, L., Sadki, M., Soutif-Bellenger, M., Tilmant, F., Tramblay, Y., 1500 Véron, A.-L., Vidal, J.-P., and Thirel, G.: On the visual detection of non-natural records in streamflow time series: challenges and impacts, *Hydrol. Earth Syst. Sci.*, 27, 3375–3391, <https://doi.org/10.5194/hess-27-3375-2023>, 2023.
- Tramblay, Y., Rutkowska, A., Sauquet, E., Sefton, C., Laaha, G., Osuch, M., Albuquerque, M. T. D., Alves, M., Banasik, K., Beaufort, A., et al.: Trends in flow intermittence for European rivers, *Hydrological Sciences Journal*, 66, 37–49, 1505 <https://doi.org/10.1080/02626667.2020.1849708>, 2020.
- Tramblay, Y., and Somot, S.: Future evolution of extreme precipitation in the Mediterranean, *Climatic Change*, 151, 289–302, <https://doi.org/10.1007/s10584-018-2300-5>, 2018.
- 1510 Van Meerveld, H. J. I., Sauquet, E., Gallart, F., Sefton, C., Seibert, J., and Bishop, K.: Aqua temporaria incognita, *Hydrological Processes*, 34, 5704–5711, <https://doi.org/10.1002/hyp.13979>, 2020.
- Verfaillie, D., Lafayssse, M., Déqué, M., Eckert, N., Lejeune, Y., and Morin, S.: Multi-component ensembles of future meteorological and natural snow conditions for 1500 m altitude in the Chartreuse mountain range, Northern French Alps, *The Cryosphere*, 12, 1249–1271, <https://doi.org/10.5194/tc-12-1249-2018>, 2018. 1515
- Vicente-Serrano, [MS](#), [SM](#), McVicar, T. R., Miralles, D. G., Yang, Y., and Tomas-Burguera, M.: Unravelling the influence of atmospheric evaporative demand on drought and its response to climate change, *WIREs Clim Change*, 11, e632, <https://doi.org/10.1002/wcc.632>, 2020. 1520
- [Vicente-Serrano, S. M., Peña-Gallardo, M., Hannaford, J., Murphy, C., Lorenzo-Lacruz, J., Dominguez-Castro, F., López-Moreno, J. I., Beguería, S., Noguera, I., Harrigan, S., and Vidal, J. P.: Climate, Irrigation, and Land Cover Change Explain Streamflow Trends in Countries Bordering the Northeast Atlantic, *Geophysical Research Letters*, 46, 10821–10833, <https://doi.org/10.1029/2019GL084084>, 2019.](#) 1525

Vidal, J.-P., Martin, E., Franchistéguy, L., Baillon, M. and Soubeyroux, J.-M.: A 50-year high-resolution atmospheric reanalysis over France with the Safran system, *Int. J. Climatol.*, 30, 1627-1644, <https://doi.org/10.1002/joc.2003>, 2010.

Wasson, J. G., Chandesris, A., Pella, H., and Blanc, L.: Définition des hydro-écorégions françaises métropolitaines. Approche régionale de la typologie des eaux courantes et éléments pour la définition des peuplements de référence d'invertébrés, Irstea, pp.190, hal-02580774, 2002.

Yu, X., and Duffy, C. J.: Watershed hydrology: Scientific advances and environmental assessments, In *Water (Switzerland)*, 10, 288, <https://doi.org/10.3390/w10030288>, 2018.

Zipper, S., Hammond, J., Shanafield, M., Zimmer, M., Datry, T., Jones, C., Kaiser, K., Godsey, S., Burrows, R., and Blaszcak, J., et al.: Pervasive changes in stream intermittency across the United States, *Environmental Research Letters*, 16, 084033, <https://doi.org/10.1088/1748-9326/ac14ec>, 2021.