

Drought propagation in high-latitude catchments: Insights from a 60-Year Analysis Using Standardized Indices

Claudia Teutschbein¹, Thomas Grabs¹, Markus Giese², Andrijana Todorović³, and Roland Barthel²

5 ¹ Uppsala University, Department of Earth Sciences, Program for Air, Water and Landscape Sciences, Villavägen 16, 75236 Uppsala, Sweden

² University of Gothenburg, Department of Earth Sciences, Medicinaregatan 7B, 41262 Gothenburg, Sweden

³ University of Belgrade, Faculty of Civil Engineering, Institute of Hydraulic and Environmental Engineering, Bulevar kralja Aleksandra 73, 11000 Belgrade, Republic of Serbia

10 *Correspondence to:* Claudia Teutschbein (claudia.teutschbein@geo.uu.se)

Abstract. Droughts, traditionally less associated with high-latitude regions, are emerging as significant challenges due to changing climatic conditions. Recent severe droughts in Europe have exposed the vulnerability of northern catchments, where shifts in temperature and precipitation patterns may intensify drought impacts. This study investigates the dynamics of drought propagation in high-latitude regions, focusing on four key aspects: (1) the typical lag time for drought conditions to propagate from initial precipitation deficits to impacts on soil moisture, streamflow, and groundwater systems, (2) the probability of precipitation deficits leading to these droughts, (3) the key catchment characteristics influencing drought propagation, and (4) how drought propagation has evolved under changing climate conditions. By analyzing long-term observational records from 50 Swedish catchments, the study reveals that drought propagation is highly variable and influenced by a complex interplay of catchment characteristics, hydroclimatic conditions, and soil properties. Soil moisture exhibits the shortest propagation times, often responding within a month to precipitation deficits, while groundwater shows the longest and most variable response times, sometimes exceeding several months. The probability of precipitation deficits propagating into soil moisture droughts is highest, followed by streamflow and groundwater, with these probabilities increasing over time. Across all drought types, annual precipitation and streamflow emerge as the most influential factors governing both propagation time and probability. Although most catchments have become wetter year-round due to climate change, southern catchments are increasingly vulnerable to spring droughts (particularly soil moisture drought) driven by increasing evaporative demand. Despite these hydroclimatic shifts, no significant long-term trends in propagation times or probabilities have been observed over the past 60 years. These findings highlight the need for tailored, region-specific water management strategies to address seasonal and regional variations in drought risks, particularly as climate change continues to reshape hydrological regimes.

30 **Non-technical summary:** Droughts are becoming a growing concern in high-latitude regions, like Sweden, due to changing climate patterns. This study looked at how droughts develop and spread through soil, rivers, and groundwater systems by analyzing data from 50 catchments across Sweden. Over a period of 60 years, most areas have gotten wetter throughout the year, but the southern regions have become more prone to spring droughts due to increased evaporation.

The study found that soil moisture responds quickly to a lack of rain, within about a month, while rivers take around two months to show signs of drought, and groundwater takes the longest to respond - up to several months. The chance of a rainfall shortage leading to a drought is highest for soil moisture, followed by rivers and then groundwater. The biggest factors influencing droughts are the amount of rain a region gets and how much water flows through rivers.

35 months to show signs of drought, and groundwater takes the longest to respond - up to several months. The chance of a rainfall shortage leading to a drought is highest for soil moisture, followed by rivers and then groundwater. The biggest factors influencing droughts are the amount of rain a region gets and how much water flows through rivers.

Importantly, despite changes in climate, the study found that the underlying drought dynamics stayed stable over the past 60 years. This means that current water management strategies based on past data are still reliable, but ongoing monitoring will

40 be essential as climate change continues to evolve. The findings highlight the need for region-specific approaches to managing water resources, especially in southern areas where spring droughts are becoming more of a concern.

1 Introduction

Droughts, typically described as episodic, socio-climatologically induced water deficits caused by anomalies in average conditions (Pereira et al., 2006), can occur in any climate zone (Wilhite, 1996; WMO and GWP, 2016). They stand out among natural hazards due to their unique characteristics, progressing gradually with a slow onset and long recovery times, making them difficult to identify precisely (Rajsekhar et al., 2015; Spinoni et al., 2015; UNDRR, 2021). Often referred to as ‘creeping disasters’ (Van Loon, 2015), they are subtle but can persist for extended periods and have considerable impacts across various sectors, including water supply quality and quantity, ecosystems, agriculture, and hydropower production (Blauhut et al., 2022; Teutschbein et al., 2023b; UNDRR, 2021).

Although droughts are typically associated with arid or semi-arid regions, recent events such as the 2018 or the 2022 European droughts (Blauhut et al., 2022; Garrido-Perez et al., 2024; Tripathy and Mishra, 2023) have shown that colder high-latitude regions, including Scandinavia, are also at risk (Bakke et al., 2020; Teutschbein et al., 2022, 2023a). High-latitude catchments, which feature distinct hydrological processes such as snow accumulation or snowmelt, exhibit unique responses to climate change and drought conditions. These regions are particularly sensitive to climate change (IPCC, 2021) and have already seen substantial alterations in streamflow regimes (Arheimer and Lindström, 2015; Blöschl et al., 2017; Teutschbein et al., 2022) and groundwater recharge (Nygren et al., 2020, 2021). As the climate further warms, changes in temperature and changing precipitation patterns are expected to disrupt snow-related processes even more, leading to profound shifts in the hydrological dynamics in these regions (Arheimer and Lindström, 2015; Irannezhad et al., 2015; Skålevåg and Vormoor, 2021; Teutschbein et al., 2015; Wilson et al., 2010). These changes are likely to intensify the development and propagation of droughts (Ahopelto et al., 2023; Spinoni et al., 2018; Teutschbein et al., 2023b), with complex and multifaceted hydrological consequences.

This complexity arises because drought development involves changes in water fluxes that affect various feedback mechanisms within the hydrologic cycle (Van Loon, 2015). Drought propagation is generally understood as a hierarchical




Figure 1: Conceptual framework depicting the onset and hierarchical progression of drought within the water cycle (adopted and modified from Changnon, 1987). The framework begins with meteorological drought and illustrates the cascading effects on snowpack and soil moisture, as well as on streamflow and groundwater.

top-down process (Changnon, 1987), where variations in precipitation (rain/snow) and temperature (as proxy for evaporative demand) cause a meteorological drought (Figure 1). Over time, this can cascade down to other hydrological variables in the water cycle, e.g., to snow pack, soil moisture, streamflow, and groundwater, often occurring in a non-linear manner and with considerable delays (Mukherjee et al., 2018). Deficits in soil moisture that affect soil vegetation and crops are typically framed as agricultural/ecological droughts (Van Loon, 2015), and deficits in streamflow and groundwater as hydrological droughts (Mishra and Singh, 2010). These various types of drought are interlinked through both positive and negative feedback processes (Van Loon, 2015), and the seasonal dynamics of snow and ice add further complexity to the processes. The timing and intensity of snow accumulation and melt can significantly influence water availability, either counteracting or amplifying seasonal precipitation deficits (Staudinger et al., 2014).

While drought propagation has been examined in other parts of the world, particularly in more temperate and arid regions (Bevacqua et al., 2021; Brunner and Chartier-Rescan, 2024; Entekhabi, 2023; 2024a; Heudorfer and Stahl, 2016; Odongo et al., 2023; Sattar et al., 2019), many analyses have either focused on individual events, often framed as 'storylines' (Chan et al., 2022; Gessner et al., 2022), on large-scale comparative studies (Li et al., 2023; Orth and Destouni, 2018), or on the relationship between only two drought types, such as precipitation and soil moisture (Geng et al., 2024a), streamflow (Hellwig et al., 2022; Ma et al., 2021; Zhang et al., 2022a), or groundwater (Bloomfield and Marchant, 2013). In contrast, high-latitude catchments, which are expected to receive more precipitation in a future climate (IPCC, 2021), have not received the same level of systematic and statistical attention, and studies that adopt an integrated approach by examining multiple hydrological components simultaneously remain rare. A key research gap lies in understanding how the heterogeneity in high-latitude catchments, including differences in catchment size, topographic features, land-use patterns, and climatic conditions unique to these regions, influence the propagation of precipitation deficits. This gap hampers our ability to discern common patterns and key factors influencing drought propagation, resulting in an incomplete understanding of how droughts evolve and transition across these landscapes. To address these knowledge gaps, this paper seeks to answer the following research questions in the context of high-latitude catchments:

1. What is the typical lag time for drought conditions to propagate from initial precipitation deficits to subsequent impacts on soil moisture, surface water, and groundwater systems in high-latitude catchments?
2. What is the propagation probability of precipitation deficits translating into droughts in soil moisture, surface water, and groundwater in these regions?
3. Which topographic, hydroclimatic, soil and land-cover characteristics are most strongly linked to drought propagation and progression in high-latitude catchments?
4. How has drought propagation in high-latitude catchments evolved in response to a changing climate?

This study addresses these questions using long-term observational data from 50 Swedish catchments, complemented by ERA5-Land reanalysis data, providing a robust empirical foundation for tracing precipitation deficits through different parts

of the hydrological system. The analysis minimizes uncertainty and captures real-world variability in drought responses, which is particularly valuable for understanding complex hydrological processes across diverse northern landscapes.


110 **2 Methodology**

2.1 Study Sites

The propagation of drought was analyzed using the CAMELS-SE dataset (Teutschbein, 2024a, b), which includes data from 50 high-latitude catchments in Sweden (Figure 2), spanning the years 1961 to 2020. These catchments are distributed across a longitudinal range from 56°N to 68°N and encompass all three major climate zones in Sweden (Figure 2a): the polar tundra climate (ET) in the Scandinavian Mountains of northwestern Sweden, the subarctic boreal climate (Dfc) in central and northern Sweden, and the warm-summer hemiboreal climate zone (Dfb) in southern Sweden (Teutschbein, 2024a; Todorović et al., 2024).

Average elevation across the catchments varies from 12 to 942 m a.s.l. (Figure 2b), with catchment areas ranging from 2 to 8,425 km². Forests dominate the land cover of these catchments, with only very few catchments exhibiting intensive agricultural activities. Glaciers and urbanized areas occupy negligible portions of the catchment area, up to 2% and 3% respectively, while lakes and wetlands are generally scarce, with a median area of 12%. Approximately one-third of the catchments are subject to regulation, though the impact of reservoirs on streamflow is relatively minor (Todorović et al., 2022; Tootoonchi et al., 2023), which is crucial for accurately representing natural drought propagation.

The catchments exhibit different hydroclimatic properties, and can - following the grouping by Teutschbein et al. (2022) - be clustered in five different groups based on their streamflow dynamics (Figure 2c). The selected catchments are predominantly humid, with the wettest areas found in western Sweden, characterized by both high precipitation and streamflow (clusters 1, 3 and 4). Snow-dominated and transitional catchments (clusters 1-3) are more common than those dominated by rainfall (clusters 4-5). A noticeable north-south temperature gradient exists, with catchments in clusters 1 and 2 featuring mean temperatures below zero, catchments in cluster 3 around 1.6°C and in clusters 4 and 5 around 6.5°C (Figure 2c). A similar gradient is observed in the runoff coefficient, which reflects the proportion of precipitation converted into runoff rather than lost through evapotranspiration. The highest runoff coefficient is found in cluster 1 (nearly 97%), followed by clusters 2 (60%) and 3 (57%), with clusters 4 (53%) and 5 (33%) exhibiting lower values. A detailed description of the physiographic and hydroclimatic features of the catchments can be found in Teutschbein (2024a, b).



135 **Figure 2: Illustration of the 50 streamflow stations along with their catchment areas in relation to Sweden's (a) climate zones, as**
 140 **classified by the Köppen-Geiger system (Beck et al., 2018), encompassing the polar tundra (ET), subarctic boreal (Dfc), and warm-**
 145 **summer hemiboreal (Dfb) climates, (b) elevation data sourced from Lantmäteriet, the Swedish Mapping, Cadastral, and Land**
 150 **Registration Authority, and (c) hydroclimatic catchment clusters identified by Teutschbein et al. (2022) with their corresponding**
annual precipitation, mean temperature and streamflow statistics.

140 2.2 Data

Daily precipitation, temperature and streamflow data from 1961 to 2020 were obtained from the freely available CAMELS-SE dataset (Teutschbein, 2024a, 2024b) and aggregated into monthly mean values. The daily temperature and precipitation series in the dataset were originally derived from the Swedish Meteorological and Hydrological Institute (SMHI) national precipitation-temperature grid, which has a spatial resolution of 4 km x 4 km (SMHI, 2005; Johansson, 2000).

145 Due to the lack of long-term soil moisture observations, monthly mean volumetric soil moisture for the uppermost 7 cm of soil over the same period (1961-2020) was obtained from the ERA5-Land reanalysis dataset, which is publicly available at a spatial resolution of 11 km x 11 km through the Copernicus Climate Data Store (Hersbach, et al., 2023). Catchment-specific values for precipitation, and soil moisture were computed by averaging all grid cells whose centers were fully located within the catchment boundaries.

150 Monthly mean groundwater observations were downloaded from the Swedish Geological Survey (SGU), which provides measurements across 1,506 wells across Sweden on its publicly accessible website. For this study, we selected wells based on the following criteria:

- located within the boundaries of any of the 50 study catchments or within a 5 km buffer

- minimum of 20 years of data coverage
- at least 120 data points
- no gaps in the data lasting more than 5 months

A spatial match between groundwater wells meeting these criteria and catchments was found for only 15 out of the 50 catchments, enabling the study of drought propagation throughout the entire hydrological system in these areas. The number of groundwater wells within the 15 catchments varied from 1 to 14. Data gaps up to 5 months were filled using the average of normalized groundwater levels from the remaining wells in the catchment (when more than one well was available), otherwise through linear interpolation.

2.3 Drought Identification through Standardized Drought Indices

Numerous drought indices have been developed over the past decades to simplify the complex task of identifying and quantifying droughts into single, interpretable values (WMO and GWP, 2016). To identify periods of droughts across different components of the water cycle (Figure 1), we selected the following widely-used dimensionless standardized drought indices that express droughts as a deviation from normal hydroclimatic conditions:

- The **standardized precipitation index (SPI)**, originally proposed by McKee et al. (1993), is based exclusively on *precipitation* data. Due to its adaptability across various spatial and temporal scales, it has become a widely used tool in drought comparison studies. The World Meteorological Organization (WMO) and other institutions have recognized the SPI as the primary method for assessing meteorological drought (WMO and GWP, 2016).
- The **standardized soil moisture index (SSMI)**, as described by Sheffield and Wood (2007), is derived from volumetric *soil moisture* values. These values echo the combined effects of a range of hydrological processes, such as plant transpiration, soil evaporation, infiltration, runoff, and the accumulation and melting of snow.
- The **standardized streamflow index (SSFI)** operates on a similar concept as the previous indices, but utilizes *streamflow* data instead (Vicente-Serrano et al., 2011). It is commonly used to quantify hydrological droughts (Vicente-Serrano et al., 2012).
- The **standardized groundwater index (SGI)**, developed by Bloomfield and Marchant (2013), standardizes *groundwater level* time series to characterize groundwater droughts. Given the complexity of groundwater flow systems, the SGI is useful for understanding fluctuations in groundwater levels and storage in response to variations in water input.

Each index was calculated by first fitting a suitable probability distribution to the monthly time series of the respective variable. This allowed the determination of cumulative probabilities, which were subsequently converted into standardized values (so-called z-scores) assuming a normal distribution with a mean of zero and a standard deviation of one. These z-scores represent how many standard deviations a given value deviates from the mean and directly represent the standardized drought index. Positive values reflect conditions that are wetter than average (i.e., no drought), while negative scores correspond to drier-

than-normal conditions, typically classified into extreme drought ≤ -2 , severe ≤ -1.5 , moderate ≤ -1.0 and mild drought ≤ 0 . All indices were calculated for different aggregation periods (i.e., for 1, 3, 6, 12 and 24 months), following the procedures described in Teutschbein et al. (2022). Aggregation over shorter time spans (less than 3 months) highlights short-term variability and is useful for assessing immediate drought impacts. In contrast, longer aggregation periods (more than 6 months) are better suited for identifying prolonged anomalies and the cumulative influence of drought across the water cycle. The computation of all standardized indices is influenced by both the data record length (Wu et al., 2005) and the type of probability distribution applied (Stagge et al., 2015). We followed the methodology outlined by Teutschbein et al. (2022) and tested 16 one-, two and three-parameter distributions probability distributions (Figure 3) for each considered series (4 indices, obtained for 50 gauging stations and 12 months of the year, at 5 different aggregation periods) separately. For a detailed description and corresponding equations of each tested distribution, we refer the reader to Teutschbein et al. (2022). To determine the most suitable distribution for each case, we utilized Kuiper's goodness-of-fit test (Kuiper, 1960).

	Precipitation	Soil moisture	Streamflow	Groundwater	Total
Generalized Extreme Value	8%	39%	38%	17%	26%
Weibull	13%	8%	9%	9%	10%
Logistic	9%	9%	8%	11%	9%
Log-logistic	13%	7%	6%	9%	9%
t Location-scale	8%	8%	10%	8%	9%
Log-normal	9%	13%	6%	6%	8%
Extreme Value	1%	1%	16%	8%	6%
Nakagami	11%	2%	1%	8%	5%
Normal	6%	8%	4%	3%	5%
Gamma	12%	2%	1%	6%	5%
Inverse Gaussian	7%	3%	1%	3%	4%
Birnbaum Saunders	3%	0%	1%	8%	3%
Rayleigh	0%	0%	0%	2%	0%
Generalized Pareto	0%	0%	0%	1%	0%
Exponential	0%	0%	0%	1%	0%
Half-normal	0%	0%	0%	0%	0%

Figure 3: Summary of 16 candidate distributions and the percentage of times each was selected as the best fit for 3-month aggregated data across the four hydrologic variables. The final column shows the overall selection frequency across all fitting instances. Best-fitting distributions are highlighted in varying shades of purple, with darker shades indicating higher selection percentages.

2.4 Detection of Spatiotemporal Patterns

To explore overarching spatiotemporal drought patterns, we examined the standardized drought indices based on both 3-month and 12-month aggregation periods, covering the full observation period from 1961 to 2020 and including all studied catchments. This approach enabled us to detect distinct drought episodes and assess their geographical distribution. Particular
205 attention was given to identifying possible north–south gradients, which were anticipated due to regional differences in evaporation dynamics and snow-related processes such as accumulation and melt.

2.5 Statistical Analysis of Drought Propagation

For all statistical analyses of drought propagation described below, we adopted the 3-month aggregation period, which is regularly used to evaluate drought effects on sectors such as drinking water supply, industrial operations, hydropower
210 generation, and freshwater ecosystems (Bae et al., 2019; Stagge et al., 2015). The choice of a 3-month period strikes a balance between capturing the immediate effects of precipitation deficits and the broader, seasonal impacts.

2.5.1 Propagation Time

Given the inherent complexity of hydrological processes and their interactions, considerable variability in drought responses is expected across different hydrological components. To assess how various components react to precipitation deficits, we
215 calculated the cross-correlation between the SPI and the three other indices (SSMI, SSFI, and SGI), focusing exclusively on periods when SSMI, SSFI, or SGI indicated moderate, severe or extreme drought (i.e. indices < -1), thus excluding mild droughts and non-drought conditions from the analysis. Cross-correlation served as a metric to quantify the similarity between the SPI time series and the lagged time series of the other indices, with lags ranging from 1 to 12 months.

Following the procedure outlined by Bloomfield and Marchant (2013), which was initially applied to groundwater response
220 times, we used the lag time that produced the highest Spearman's rank correlation coefficient θ_s (Spearman, 1904) as an indicator of response times. Lag time is a commonly used metric to describe drought propagation (Odongo et al., 2023; Zhang et al., 2022b), where shorter lag times indicate a faster response of a specific component to SPI changes.

In this study, we determined propagation times from precipitation to soil moisture, streamflow and groundwater separately for each catchment, and the results were grouped and analyzed according to catchment clusters (c.f. Figure 2c) to facilitate the
225 identification of spatial patterns.

2.5.2 Propagation Probability

To investigate the role of precipitation deficits in triggering drought conditions across the hydrological system, we conducted a probabilistic analysis focused on the occurrence of different drought types following a precipitation drought event. After identifying precipitation droughts with SPI < -1 (i.e., excluding mild droughts) as separate events/runs, we evaluated whether

230 corresponding deficits in soil moisture (SSMI), streamflow (SSFI) or groundwater (SGI) could be observed within different drought propagation time frames. This analysis is captured by the conditional probability expressed in Eq. (1).

$$P(index_{lag} < -1 | SPI < -1) = \frac{P(SPI < -1 \cap index_{lag} < -1)}{P(SPI < -1)} \quad (1)$$

235 Here, $index_{lag}$ represents any of the drought indices for soil moisture, streamflow, or groundwater (i.e., SSMI, SSFI or SGI) at a given temporal lag . Therefore, $P(index_{lag} < -1 | SPI < -1)$ is the conditional probability of any of the drought indices indicating drought conditions given that a precipitation drought (indicated by $SPI < -1$) has already occurred.

In this analysis, we tested all possible time lags ranging from 1 to 12 months. This involved starting with the probability that a precipitation drought would cause a soil moisture, streamflow, or groundwater drought within the same month, then
240 extending the analysis to include the same and subsequent months, and so forth, up to 12 months.

We also performed this analysis separately for each season, categorizing months as follows: spring (March–May; MAM), summer (June–August; JJA), autumn (September–November; SON), and winter (December–February; DJF). To assess the variation of clusters within each season, we utilized the coefficient of variation (CV).

This conditional probability framework provides insights into how precipitation deficits propagate through the hydrological
245 system, reflecting the strength and timing of drought signal transmission across different hydrological components (Zhu et al., 2021). It offers a useful approach to highlight how precipitation shortages can trigger subsequent soil moisture, streamflow, or groundwater droughts.

2.5.3 Governing Factors of Drought Propagation

To explore the relationship between propagation time and probability with underlying catchment characteristics and
250 hydroclimatic factors, we employed the non-parametric Spearman's rank correlation coefficient θ_s (Spearman, 1904). This statistical approach allowed us to assess correlations across all pairwise combinations of standardized drought indices and a comprehensive range of potential influencing factors. These factors included geographic and physical attributes such as latitude, catchment area, elevation, and slope, as well as hydrological and climatic variables like the degree of regulation (DOR), regulation volume, annual mean precipitation, annual mean temperature, and annual streamflow. Additionally, we
255 considered the impact of soil types and land cover variations, such as the percentage of forest cover or agricultural land.

The choice of Spearman's rank correlation over the linear Pearson product-moment correlation (Pearson, 1920) was driven by its ability to capture monotonic relationship without assuming any specific form of the relationship, such as linearity or logarithmic behavior. This flexibility is particularly advantageous for drought propagation through the hydrological system, because relationships between variables often do not follow a strict linear pattern (Mukherjee et al., 2018). By using Spearman's
260 correlation, we ensured that our analysis could detect and quantify both linear and non-linear associations, providing a more robust understanding of how different catchment and climatic characteristics influence drought propagation dynamics.

2.6 Droughts and their Propagation in a Changing Climate

The extensive 60-year observational data record encompassed two ‘climate normal periods’ (CNPs), each representing a 30-year climatological baseline as defined by the World Meteorological Organization (WMO Climatological Normals, 2021).
265 This provided a unique opportunity to conduct an assessment of potential long-term shifts in droughts and their propagation in high-latitude catchments. Specifically, we investigated whether there were significant differences in both the drought indices, the propagation time and the propagation probability in each catchment cluster between the CNPs of 1961-1990 and 1991-2020, utilizing the non-parametric Wilcoxon rank sum test (Asadzadeh et al., 2014; Montgomery and Runger, 2010). This analysis focused primarily on precipitation drought propagation into soil moisture and streamflow, as the available
270 groundwater observations were not of sufficient length for such an assessment.

To analyze variations in annual drought conditions, we utilized 12-month aggregated drought indices (SPI-12, SSMI-12, SSFI-12), calculated for September each year. This selection covers the full hydrological year, from the preceding October through to the current September. For seasonal drought conditions, we employed 3-month aggregated drought indices (SPI-3, SSMI-3, SSFI-3) corresponding to the final month of each season: May (spring, MAM), August (summer, JJA), November (autumn,
275 SON), and February (DJF).




Figure 4: Computed standardized indices for all Swedish study catchments from 1961–2020, sorted by latitude on the y-axis (with uneven spacing reflecting variations in catchment density across latitudes) and time on the x-axis. Panels show (a-b) precipitation (SPI), (c-d) soil moisture (SSMI), (e-f) streamflow (SSFI) and (g-h) groundwater (SGI) over two different aggregation periods: 3-month indices (left panels) and less-noisy 12-month indices (right panels). Light blue indicates non-drought conditions, yellow indicates mild drought, and darker red and blue represents more severe drought. White areas in SGI (g-h) denote missing values.

3 Results

3.1 Spatiotemporal Drought Patterns across the Hydrological System

285 The computed standardized indices (SPI, SSMI, SSFI, and SGI) at 3-month aggregation period highlight multiple significant drought periods across the country, with both temporal and spatial variations evident across the different hydrological components (Figure 4, left panels). We also showcase the less noisy 12-month aggregation period to facilitate visual comparisons of different catchments and drought indices (Figure 4, right panels). Notably, two pronounced drought periods, occurring in the late 1960s and mid-1970s, affected the entire country and had major impacts on the hydrological system.

290 These events began with precipitation deficits, as indicated by the SPI (Figure 4a,b), and propagated through to soil moisture (Figure 4c,d), streamflow (Figure 4e,f), and groundwater systems (Figure 4g,h), with each component experiencing drought conditions for several months.

The droughts of 1996 and the 2003 European heatwave are also clearly visible across all hydrological components, demonstrating the widespread effects of these events. Similarly, more recent droughts, particularly those in 2016/17 and 2018,

295 are well represented, affecting soil moisture, streamflow, and groundwater, although the severity and extent vary by region. Beyond these general patterns, the standardized indices also reveal notable spatial variations in how different hydrological components were affected by droughts. For example, the early 1990s drought primarily affected catchments in southern Sweden, where the impact on soil moisture (Figure 4c,d) was substantially stronger compared to other components. Similarly, the drought of 2016/17 and 2018 display distinct spatial patterns: While precipitation and streamflow deficits were observed

300 nationwide, groundwater levels were disproportionately affected in southern Sweden.

3.2 Propagation Time

The cross-correlation analysis uncovered differences in drought propagation times across the five identified clusters (Figure 5). Soil moisture exhibited the shortest propagation times across all clusters, with average response times ranging from 0.1 to 2.7 months, highlighting its sensitivity to short-term climatic fluctuations. Streamflow showed slightly longer propagation

305 times compared to soil moisture, with some regional differences observed. The western clusters 1, 3 and 4 (Figure 5a,c,e) exhibited response times of 0.9 to 2.4 months, whereas the other two clusters - Cluster 2 (Figure 5b) and Cluster 5 (Figure 5d) - had somewhat longer average response times of 3 and 3.5 months, respectively. This suggests that while streamflow responds quickly to precipitation deficits, the speed of this response is influenced by regional hydrological and climatic conditions.

Groundwater displayed the longest and most variable propagation times among the hydrological components analyzed. The

310 response times varied considerably across clusters, with slightly longer response times in western clusters 2 and 5. Note that the northernmost Cluster 1 (Figure 5a) had only one groundwater observation in the region, for which no correlation could be found to precipitation deficits. In the other clusters, average groundwater response times ranged from 2.6 months to 6.7 months. Groundwater also exhibited the highest internal variability within each cluster, with differences of several months between the fastest and slowest responses. Clusters 2 and 5 were particularly notable, with differences of up to 10 months observed between

the shortest and longest groundwater response times. This high variability underscores the complexity of groundwater dynamics and its longer, more delayed response to precipitation deficits compared to other hydrological components. Across all 50 catchments in Sweden (Figure 5f), soil moisture responds to precipitation deficits within an average in one month, streamflow in two months, and groundwater within 4 months.




Figure 5: Propagation times (in months) from precipitation deficits to soil moisture, streamflow and groundwater drought across the five different catchment clusters (a-e) and across all catchments (f) in Sweden.

3.3 Propagation Probability

To quantify how precipitation deficits contribute to the development of more severe droughts in other hydrological components, we analyzed conditional probabilities. The systematic assessment of step-wise time lags (Figure 6) revealed that at the shortest lag of one month, a precipitation deficit resulted in an agricultural/ecological drought in 48% of cases, a streamflow drought in 41%, and a groundwater drought in 38% of cases (Figure 6a). As the lag times are progressively extended month by month, the likelihood of a precipitation drought triggering subsequent droughts in other components increased considerably. Over time, the frequencies of agricultural/ecological, streamflow, and groundwater droughts converged, indicating a more uniform occurrence of drought across the hydrological system.




Figure 6: Probability of a moderate, severe, or extreme precipitation drought ($SPI < -1$) triggering subsequent droughts in soil moisture, streamflow, or groundwater (SSMI, SSFI, or SGI < -1) across different time lags. Panel (a) shows the overall conditional probability for all catchments combined. The boxplots depict the distribution of probabilities across catchments, with the solid black line representing the median probability for a 1-month lag time. The thin grey lines correspond to the median probabilities for step-wise increasing time lags, while the lines for 3 and 12 months are highlighted for clarity. The bottom panels provide a detailed breakdown by cluster for (b) soil moisture (SSMI), (c) streamflow (SSFI), and (d) groundwater (SGI), respectively.

Across the clusters, the conditional probability of a soil moisture drought occurring within one month of a precipitation deficit ranged from 41 to 43% in the northern clusters 1-3, increasing to 51% and 57% in the southern clusters 4-5. As the time lag increased, the probabilities rose, and the differences between northern and southern catchments became less pronounced. This general pattern also applied to streamflow (Figure 6c) and groundwater droughts (Figure 6d). However, these components did not exhibit as strong a north-south gradient. Instead, the propagation probabilities across the different clusters were more balanced, with the exception of cluster 4, located in southwestern Sweden, where the probabilities of precipitation deficits propagating to streamflow or groundwater consistently remained 20-30% higher than in other Swedish regions.





Figure 7: Probability of a moderate, severe, or extreme precipitation drought ($SPI < -1$) in different seasons (spring, summer, autumn and winter) triggering subsequent droughts in (a) soil moisture, (b) streamflow, or (c) groundwater (SSMI, SSFI, or SGI < -1) across different time lags. The boxplots depict the distribution of probabilities across catchments, with the solid black line representing the median probability for a 1-month lag time. The thin grey lines correspond to the medians probabilities for step-wise increasing time lags, while the lines for 3 and 12 months are highlighted for clarity.

The propagation probability did not only vary across clusters, but also across seasons (Figure 7). The biggest difference can be seen in the propagation probability to soil moisture (Figure 7a), where the highest propagation probabilities occurred in summer. This means, a precipitation deficit of $SPI < -1$ in summer had on average a 68% probability to entail a soil moisture drought within one months, while the probability in spring and winter was less than 40%. This pattern was consistent also for longer time lags. For streamflow, the pattern was less pronounced. At a 1-month time lag, the propagation probability was slightly higher in the autumn (45%) than in the other seasons (~40%, Figure 7b), though the differences were not statistically significant. However, when considering longer time-lags, spring consistently showed somewhat higher propagation probabilities (Figure 7b). For groundwater, propagation probabilities remained fairly consistent – ranging from 34% to 39% in spring, summer and winter (Figure 7c). Only autumn stuck out with somewhat lower probabilities (32% at 1-month lag), a pattern that persisted across all time lags, though the differences were modest and statistically insignificant.



365 **Figure 8: Cluster-wise probability of a moderate, severe, or extreme precipitation drought ($SPI < -1$) triggering subsequent droughts by season (rows) and type of subsequent drought (columns). Each subplot also depicts the coefficient of variation (CV) across clusters for time-lag 1.**

A more detailed analysis of propagation probability by cluster revealed distinct north-south patterns during certain seasons and drought types (Figure 8). In spring (MAM), the propagation probability to soil moisture (Figure 8a) was considerably higher
370 for the two southernmost clusters (4 and 5) at 50% and 53%, compared to the northern clusters, where it ranged from 23% to 27%. For streamflow (Figure 8b) and groundwater (Figure 8c), cluster 4 stood out with notably higher propagation probabilities exceeding 60%.

In summer (JJA), the propagation probability for soil moisture increased linearly from 60% to nearly 75% across all time lags as one moves further south (Figure 8d). Conversely, the propagation probability to streamflow (Figure 8e) showed an inverse
375 trend at a 1-month time lag, i.e., decreasing from 47% in the north to 30% in the south, with the exception of cluster 4 (53%). This north-south gradient was less pronounced for longer time lags.

In autumn (SON), the spatial patterns resembled those of summer. However, the overall propagation probabilities to soil moisture were 10% to 20% lower than in summer (Figure 8g), while probabilities for streamflow (Figure 8h) and groundwater (Figure 8i) were slightly higher compared to summer levels.

380 Winter (DJF) exhibited the lowest propagation probabilities to soil moisture (Figure 8j) across all clusters, with the exception of cluster 3, which had the lowest probability in spring. Both winter and spring displayed the highest variations in soil moisture (Figure 8a,j) and streamflow (Figure 8b,k) across clusters, with coefficients of variation (CV) ranging from 0.34 to 0.44. Groundwater showed generally higher variations than both soil moisture and streamflow, with CVs ranging from 0.42 to 0.59, peaking in summer (Figure 8f) and winter (Figure 8l).


385 3.4 Governing Factors of Drought Propagation

Among the three hydrological components, the propagation time to soil moisture generally exhibited the most significant correlations with physical/topographic catchment features (Figure 9a, left). Significant positive correlations (p -value < 0.05) were observed with latitude, elevation, slope, shrubs and grassland, and wetlands ($\theta_s = 0.31$ to 0.43), indicating longer propagation times. Significant negative correlations included annual mean precipitation, mean temperature, silt, urban areas,
390 and agriculture ($\theta_s = -0.34$ to -0.41). For propagation time to streamflow (Figure 9a, center), significant positive correlations were observed with catchment area, till soils, and water features ($\theta_s = 0.29$ to 0.41), while annual precipitation, streamflow and the runoff coefficient exhibited strong negative correlations with drought propagation time ($\theta_s = -0.34$ to -0.59), indicating that higher precipitation and absolute as well as relative streamflow generation are associated with shorter propagation times. Propagation times to groundwater tended towards negative correlations (Figure 9a, right), with significant correlations
395 observed only for annual precipitation ($\theta_s = -0.75$) and areas annual streamflow ($\theta_s = -0.56$). Notably, only one single catchment characteristic exerted a strong influence on drought propagation across all three hydrological components simultaneously, namely annual precipitation, indicating shorter propagation times for catchments with higher annual precipitation.

In terms of propagation probabilities, soil moisture again showed the most extensive correlation (Figure 9b, left). Negative correlations were detected with all catchment features, particularly with latitude, elevation and slope ($\theta_s = -0.57$ to -0.73).

400 Strong negative correlations were also found with till or weathered deposits, open land, shrubs and grassland as well as wetlands ($\theta_s = -0.51$ to -0.61). In contrast, strong positive correlations were observed with annual temperature ($\theta_s = 0.81$) and agriculture ($\theta_s = 0.68$). For propagation probability to streamflow (Figure 9b, center), fewer significant correlations were evident. Negative correlations were found with catchment area, till soils and water bodies ($\theta_s = -0.33$ to -0.50), whereas strong positive correlations were detected for annual precipitation ($\theta_s = 0.73$), streamflow ($\theta_s = 0.59$), and the runoff coefficient ($\theta_s = 0.43$). Propagation probability to groundwater (Figure 9b, right) showed only three significant correlations, all of them positive: annual precipitation ($\theta_s = 0.66$) and streamflow ($\theta_s = 0.59$), as well as clayey and clay till soils ($\theta_s = 0.66$).

405





415 **Figure 10: Shifts in (a-e) annual and (f-j) seasonal drought indices (SPI, SSMI and SSFI) between two climate normal periods, 1961–**
1990 (CNP1) and 1991–2020 (CNP2), for five hydrological clusters. Boxplots in panels (a)-(e) represent the range of index values
(SPI, SSMI and SSFI) across catchments within each cluster, with significant shifts between CNP1 and CNP2 indicated by 'sign.'
420 based on the Wilcoxon rank-sum test ($p < 0.05$). Panels (f)-(j) display the absolute change in indices for different seasons (spring,
summer, autumn, and winter), with darker blue shading representing more pronounced increases in index values (wetter
conditions), and darker purple shading representing stronger decreases (drier conditions). Significant seasonal changes are
highlighted with black text.

3.5 Droughts and their Propagation in a Changing Climate

3.5.1 Changes in Spatiotemporal Drought Patterns

The analysis of 12-month aggregated drought indices revealed a general upward trend from CNP1 to CNP2 (i.e., indices
425 become more positive), indicating a reduction in the severity of droughts overall (Figure 10a-e). In all five clusters, the
Wilcoxon rank-sum test consistently rejected the null hypothesis that the SPI and SSMI values for CNP1 and CNP2 were
drawn from distributions with equal medians. This indicates a statistically significant shift ($p < 0.05$) in precipitation and soil
moisture drought conditions across Sweden between these two periods. For the SSFI, significant differences between CNP1
and CNP2 were only observed in the northernmost clusters 1-3 (Figure 10a-c), implying a notable shift in streamflow drought
430 severity in these areas.

Seasonal analysis of 3-month aggregated drought indices showed a similar pattern, with predominantly positive shifts across
most clusters and seasons (Figure 10f-j), suggesting a reduction in drought severity. However, some exceptions are observed.
In the two northernmost clusters 1 and 2, autumn indices (particularly soil moisture and streamflow) exhibited slight negative
shifts, indicating slightly drier conditions during this season (Figure 10f, g). Similarly, in the southernmost clusters 4 and 5,
435 substantial negative changes were observed in spring soil moisture and streamflow indices (Figure 10i, j), suggesting increased
drought severity during this season in southern Sweden.

3.5.2 Changes in Propagation Time

For propagation time to soil moisture drought (SSMI) (Table 1a), most clusters exhibited no change between CNP1 and CNP2
(i.e., values of 0), except for cluster 2, which showed a modest increase of +0.5 months. For propagation time to streamflow
440 drought (SSFI), cluster 2 also experienced a small increase (+0.5 months), while cluster 3 showed a more noticeable reduction
of -1 month. However, in all cases, the Wilcoxon rank-sum test failed to reject the null hypothesis that the propagation times
to soil moisture and to streamflow during CNP1 and CNP2 were drawn from continuous distributions with equal medians
(Table 1a). This implies that there has been no statistically significant change (all p -values > 0.05) in the drought propagation
times between these two periods.

445

450

Table 1: Drought Propagation Characteristics from precipitation deficits to soil moisture (SSMI) and streamflow droughts (SSFI), including the (a) observed median propagation times and (b) propagation probabilities for drought transitions. The changes in these characteristics are compared between two climate periods, CNP1 (1961-1990) and CNP2 (1991-2020). The p-values reflect the statistical significance of these changes, tested using the Wilcoxon rank-sum test. Significant changes (p-values <0.5) are highlighted in bold.

Propagation Characteristics	SSMI				SSFI			
	CNP1	CNP2	Change	p-value	CNP1	CNP2	Change	p-value
(a) Propagation Time								
All catchments	0	0	0.0	0,299	2	1	- 1.0	0,197
Cluster 1	0	0	0.0	1,000	1	1	0.0	0,734
Cluster 2	0	0.5	+ 0.5	0,657	2	2.5	+ 0.5	0,835
Cluster 3	0	0	0.0	0,056	2	1	- 1.0	0,120
Cluster 4	0	0	0.0	NaN	1	1	0.0	1,000
Cluster 5	0	0	0.0	0,653	2	1.5	- 0.5	0,135
(b) Propagation Probability								
within 1 month	47%	48%	+ 1%	0,197	42%	41%	- 1%	0,796
within 2 months	52%	54%	+ 2%	0,081	53%	52%	- 1%	0,416
within 3 months	55%	59%	+ 5%	0,043	61%	57%	- 4%	0,591
within 4 months	58%	64%	+ 7%	0,070	62%	60%	- 2%	0,828
within 5 months	61%	66%	+ 5%	0,048	64%	63%	- 1%	0,809
within 6 months	64%	71%	+ 7%	0,038	69%	66%	- 3%	0,836
within 7 months	68%	72%	+ 4%	0,091	70%	69%	- 1%	0,728
within 8 months	71%	75%	+ 4%	0,076	74%	70%	- 4%	0,661
within 9 months	73%	75%	+ 2%	0,101	75%	72%	- 4%	0,735
within 10 months	75%	77%	+ 2%	0,236	76%	74%	- 3%	0,715
within 11 months	79%	80%	1%	0,291	78%	74%	- 4%	0,365
within 12 months	81%	82%	1%	0,274	80%	78%	- 2%	0,336

3.5.3 Changes in Propagation Probability

455

Similarly, we investigated as to whether the conditional probability of soil moisture respective streamflow drought following a precipitation drought has changed over time (Table 1b). The probability of propagation to soil moisture (SSMI) generally increased by 1% to 7% in CNP2 compared to CNP1, while the probabilities of propagation to streamflow (SSFI) slightly decrease by -1% to -4%. Despite these observed changes, the Wilcoxon rank sum test again did not reject the hypothesis of equal medians, indicating that no statistically significant shift occurred in drought propagation probabilities over the 60-year observational record (p-values > 0.05, Table 1b).

460

4 Discussion

4.1 Droughts across hydrological components

The computation of standardized drought indices enabled a visual exploration of the complex interactions between precipitation deficits and drought propagation across different hydrological components - soil moisture, streamflow, and groundwater - in high-latitude catchments. A consistent temporal alignment is observed between the standardized indices (SPI, SSMI, SSFI, and SGI), which is visually particularly clear for the longer 12-month aggregation period. This alignment suggests that prolonged precipitation deficits typically result in more severe and widespread drought conditions across all components, affecting soil moisture, surface water, and groundwater.

Spatially, the analysis highlights notable variability, especially across latitudes. Southern regions, particularly during drought events like those in 2016 and 2018, experienced more pronounced impacts on groundwater systems (Bakke et al., 2020). This pattern may be attributed to unique regional hydrological processes, such as the role of snowmelt in the north versus higher evaporative demand, water extraction rates and slower recharge processes in the south. These findings emphasize the importance of considering both spatial and temporal dimensions in drought analysis. They also suggest that groundwater systems in southern areas may be more vulnerable to prolonged dry periods, raising concerns about sustainable water management in these regions (Barthel et al., 2021).

While short-term indices capture the immediate impacts of precipitation deficits (critical for understanding the onset of drought conditions), the long-term indices reveal the persistence and severity of droughts that can strain water resources, ecosystems, and agricultural productivity over extended periods (Stagge et al., 2015; Teutschbein et al., 2023b). Groundwater droughts, in particular, are less frequent but tend to be more severe and prolonged compared to those affecting soil moisture and streamflow (Bloomfield and Marchant, 2013).

4.2 Propagation Time

The analysis of drought propagation times across the five identified clusters revealed notable variations in response times among different hydrological components, specifically soil moisture, streamflow, and groundwater. These variations highlight the complex and region-specific nature of drought dynamics in the study area. Some general observations were that soil moisture consistently had the shortest propagation times, often responding almost immediately (within one month) to precipitation deficits. Therefore, soil moisture in the uppermost soil layer is highly sensitive to changes in precipitation, likely due to its direct exposure to surface conditions (Singh et al., 2021). Streamflow generally exhibits slightly longer propagation times than soil moisture but remains within a relatively short timeframe, averaging about two months across all clusters. This indicates that while streamflow is also responsive to precipitation changes, the routing process through catchments introduces a slight delay compared to soil moisture (Robinson et al., 1995; Singh et al., 2021). In contrast, groundwater consistently displays the longest propagation times, averaging around four months, with significant variation observed across different clusters. This delay in groundwater response is expected, as groundwater systems typically take longer to react to precipitation

deficits due to the slow recharge of aquifers and the movement of water through subsurface layers. Notably, catchments with
495 longer groundwater propagation times may have deeper aquifers, lower recharge rates, or specific hydrogeological conditions
such as confined aquifers or low permeability leading to a delayed response (Gong et al., 2023). Our findings align with
previous research indicating that groundwater responses to precipitation anomalies are more heterogeneous than those of
streamflow (Weider and Boutt, 2010).

Clusters 3, 4, and 5 demonstrate shorter groundwater propagation times compared to clusters 1 and 2, indicating more
500 responsive groundwater systems, possibly due to shallower aquifers or regions where groundwater recharge processes occur
more rapidly (Cochand et al., 2020; Gong et al., 2023).

This variability supports earlier findings that drought propagation into groundwater is highly site-specific (Bloomfield and
Marchant, 2013; Kumar et al., 2016; Nygren et al., 2022; Stoelzle et al., 2014), emphasizing the critical role of hydrogeological
characteristics and subsurface storage processes in shaping groundwater responses. Because of their delayed and prolonged
505 responses, groundwater systems require proactive and sustained adaptive management (Thomann et al., 2022). Early detection
is challenging, and once groundwater droughts develop, they can be slow to recover (Alam et al., 2021; Hellwig et al., 2021).
This highlights the importance of targeted monitoring and preparedness strategies, such as water source diversification and
drought reserves (Langridge and Daniels, 2017), regulatory schemes for groundwater extraction (Vaux, 2011), or strategies
for implementing managed aquifer recharge (Petersen-Perlman et al., 2022), especially in areas where groundwater serves as
510 a critical resource.

In contrast, the quicker responses of soil moisture and streamflow to precipitation deficits across all clusters emphasize the
importance of implementing timely drought response measures for these components as they quickly reflect precipitation
deficits. In agriculture, such measures may include regulated deficit irrigation (Chai et al., 2015), mulching or optimized
seeding dates (Bodner et al., 2015). In water supply systems, options such as adaptive reservoir management or temporary
515 water-saving advisories (Wendt et al., 2021) may be particularly relevant in regions dependent on river water resources.

4.3 Propagation Probability

Our conditional probability analysis provides new insight into how precipitation deficits propagate through different
components of the hydrological system, revealing both temporal and spatial dynamics of drought transmission in Sweden. At
the shortest lag (one month), the probability of a precipitation deficit leading to a soil moisture drought is the highest,
520 emphasizing that agricultural and ecological systems are often the first to be impacted by water shortages. In contrast, the
probabilities of precipitation deficits propagating to streamflow and then to groundwater are lower (Geng et al., 2024b; Meresa
et al., 2023; Wang et al., 2022).

As the lag time systematically extends, the conditional probabilities for all three components (soil moisture, streamflow,
groundwater) begin to converge, indicating that prolonged precipitation deficits increase the likelihood of all components
525 experiencing drought. This suggests that while some systems (like soil moisture) respond quickly, over time, the effects of
prolonged precipitation deficits will eventually impact all components, necessitating a comprehensive approach to drought

risk management, which considers the entire cycle of disaster management from prediction and prevention to practical measures reducing impacts of droughts and supporting recovery in a sustainability context (AghaKouchak et al., 2015; Grobicki et al., 2015).

530 There is clear spatial variability in how droughts propagate across different clusters, with distinct north-south gradients observed in the propagation probabilities of soil moisture droughts. The fact that only 41%-43% of all precipitation deficits ($SPI < -1$) in the northern clusters 1-3 lead to soil moisture deficits within one month suggests that other factors also play a crucial role for the development of agricultural/ecological drought. Snow might be a key player, as is indicated by the probability getting higher when moving towards southern catchments with less snow and more rainfall. We can also speculate

535 that this regional pattern is formed by other factors such as high soil water storage capacities (Geris et al., 2015) or different drought-generating mechanisms in these snow-dominated regions (Van Loon and Van Lanen, 2012). Conversely, clusters 4 and 5 exhibit higher probabilities (up to 57% within one month), indicating a stronger link between precipitation deficits and soil moisture droughts. These regions may have soil types or land cover that are more susceptible to rapid drying during periods of low precipitation, or may be influenced by regional climatic such as higher evaporative demand (Teutschbein, 2024a).

540 However, this gradient is less pronounced for streamflow and groundwater, where probabilities are more balanced across clusters, except for cluster 1 in northwestern and cluster 4 in southwestern Sweden, which consistently show higher conditional drought probabilities. In fact, catchments in clusters 1 and 4 receive considerably higher amounts of annual precipitation (978 mm and 944 mm, respectively, compared to the Swedish average of 784 mm). Especially catchments in cluster 4 also experiences greater evaporative demand (particularly during summer) due to their southern location, while also receiving the

545 least amount of snow (10-20% of annual precipitation compared to 30% in central and 40% in northern Sweden). The delayed but persistent response of groundwater (propagation probabilities of 52-70% within 3 months in clusters 2-5) underscores once again the need for long-term monitoring and proactive management strategies, especially in regions with high groundwater dependence (Saito et al., 2021; Thomann et al., 2022). The variability in drought propagation probabilities and response times across different clusters and hydrological components suggests that a one-size-fits-all approach to drought

550 management is unlikely to be effective across diverse regions and sectors (Stenfors et al., 2024b). Instead, a more tailored approach is required, involving context-specific analysis that considers local climatic, topographic, and hydrological conditions (Kchouk et al., 2022), while also recognizing the dependency of water supply on different resources, such as surface or groundwater (Stenfors et al., 2024a).

It is noteworthy that groundwater droughts featured the highest variability within clusters, which highlights the complexity of

555 subsurface hydrological processes and may reflect differences in regional hydrological conditions, groundwater storage capacity, or data quality (Kumar et al., 2016; Stoelzle et al., 2014). Thus, more research efforts are needed to study the propagation into groundwater (Barthel et al., 2021) and to link it to aquifer properties and recharge rates.

Our analysis also revealed significant seasonal variations in the probability of precipitation deficits leading to droughts in different hydrological components. Generally, summer stands out as the season with the highest propagation probabilities

560 across all time lags, particularly for soil moisture droughts. This can be attributed to several factors: high evapotranspiration

rates, reduced soil moisture recharge, and greater demand for water by vegetation during this period (Andersson, 1989; Cienciala et al., 1999), which make the soil system even more sensitive to precipitation deficits. In contrast, spring and winter show lower propagation probabilities to soil moisture droughts, likely due to lower evapotranspiration demands and a greater influence of snow and snowmelt, which seems to buffer the effects of precipitation deficits and dampen soil moisture droughts (Potopová et al., 2016). These patterns are less pronounced for streamflow and groundwater droughts, which show relatively stable propagation probabilities across all seasons.

North-south differences emerged also in seasonal propagation probabilities, particularly for soil moisture and streamflow. While the overall inter-cluster patterns remain consistent across all four seasons, the absolute probabilities vary. For instance, the conditional probability of soil moisture droughts is generally higher in southern catchments and lower in northern catchments throughout the year. In spring, snowmelt plays a key role in mitigating precipitation deficits, resulting in significantly lower propagation probabilities in the snow-dominated northern clusters (1-3) of around 20-30% - compared to 50% in the southern clusters. During summer and autumn, total precipitation and evaporative demand become the dominant factors (Koster et al., 2019; Wang et al., 2022), driving up the probability of soil moisture droughts, especially in the south. In winter, snow accumulation acts as a buffer, further reducing the propagation probabilities for soil moisture droughts, particularly in regions with substantial snowfall (Potopová et al., 2016). Conversely, streamflow drought propagation probability during summer decreases as one moves south, in line with the southwards-decreasing runoff coefficients (i.e., proportionally less precipitation turns into runoff). These coefficients range from 97% in northern cluster 1 to 37% in southern cluster 5, which is consistent with the results of the correlation analysis, in which the runoff coefficient emerged as an explanatory factor for streamflow drought propagation probability.

It should be noted that inter-cluster variation is largest during spring and winter for both soil moisture and streamflow propagation probabilities. This indicates that northern and southern catchments behave more distinctly in these seasons, with northern clusters showing stronger buffering effects due to snow accumulation and snowmelt. In contrast, summer and autumn see a more uniform response across regions, which emphasizes the proportionally larger influence of snow processes in the north during spring and winter, compared to the evaporative demand in the summer and autumn. However, groundwater propagation exhibits consistently high inter-cluster variability across all seasons, suggesting that additional regional factors, particularly in cluster 4, may outweigh the effects of snow and evaporative demand. This could include regional geological differences, such as aquifer characteristics and groundwater storage capacities, which play a critical role in determining the speed and extent of drought propagation in groundwater systems.

These results suggest that drought propagation and resulting impacts are highly dependent on both seasonal timing and geographical location, reflecting the complex interactions between climate conditions, hydrological processes, and land characteristics, which can either accelerate (e.g., high evaporative demand) or delay drought propagation (e.g., snow-melt in snow-dominated catchments) (Koster et al., 2019; Potopová et al., 2016). In practical terms, this analysis highlights the need for seasonally adaptive water management strategies, especially in the southern clusters where the propagation probability to soil moisture is elevated. It also points to the importance of long-term monitoring, particularly for groundwater, where slow

595 but persistent drought impacts may be underestimated in the short term (Barthel et al., 2021). In restoration efforts or pilot monitoring programs, the regional variability revealed in these findings could guide targeted interventions, such as enhancing water storage in high-risk areas or focusing conservation efforts during high-risk seasons (Srivastav et al., 2021).

4.4 Governing Factors of Drought Propagation

Our analysis revealed that the processes driving drought dynamics vary considerably between soil moisture, streamflow, and groundwater. Propagation time of precipitation deficits to soil moisture in the studied catchments is influenced by a range of governing factors, while for streamflow and groundwater, only four and two significant factors, respectively, were identified. Generally, faster propagation times across all three hydrological components occur in catchments with high annual precipitation, warmer temperatures, and higher annual streamflow levels. This suggests that in regions with greater water input and output, the hydrological system responds more quickly to changes in precipitation, resulting in faster drought onset but potentially quicker recovery from drought conditions as well. Additionally, soil moisture and streamflow propagation times are shorter in catchments with less till, more silt, and fewer water features, suggesting that water bodies can exert a buffering and delay function. For soil moisture, propagation time is further controlled by urban areas and agriculture, where surface runoff dominates, and is more rapid in warmer catchments at lower latitudes and elevations. Conversely, open land types (e.g., shrubs and grasslands, open land, and wetlands) tend to slow water propagation and promote water retention.

610 The propagation probability generally exhibits stronger correlations than propagation time. Our analysis showed that propagation probability to groundwater was significantly correlated with just three factors: higher annual precipitation, greater streamflow, and the presence of clayey till, which enhanced the likelihood of groundwater recharge. For streamflow, propagation probability is influenced by a few more factors. Catchments with higher precipitation and streamflow also tend to exhibit greater propagation probabilities. In addition, smaller catchments, less till, and fewer water features are associated with higher streamflow propagation probability. The propagation to soil moisture was associated with the greatest number of governing factors. Specifically, catchments with higher mean temperatures, more agriculture, and lower latitude, elevation, and slope had the highest soil moisture propagation probabilities. Lower streamflow in these areas further amplified the likelihood of soil moisture response.

Across both drought propagation characteristics (i.e., time and probability), annual precipitation and annual streamflow emerged as the most influential drivers across all three hydrological components, indicating their potential as proxies for identifying areas with faster hydrological responses, which can support more targeted resources allocation and management decisions during emerging drought conditions. However, the remaining factors shaping drought propagation dynamics vary by component, reflecting the distinct physical and hydrological processes at play. Soil moisture is generally influenced by a broader array of factors, whereas streamflow is more directly linked to catchment size, the presence of till soils, and overall water availability. Groundwater is most strongly affected by hydro-climatic variables and the hydrological regime.

625 This analysis underscores that drought propagation is not solely determined by catchment properties; rather, it is shaped by complex interactions between seasonal precipitation, evaporative demand, and snow dynamics (Van Loon et al., 2015). The

630 timing and severity of the drought (Bae et al., 2019), the season (Meresa et al., 2023), location (Bevacqua et al., 2021), precipitation patterns (or lack thereof), as well as antecedent conditions/‘memory effects’ (Bales et al., 2018; Soulsby et al., 2021) all can potentially play important roles in determining whether a meteorological drought will evolve into more severe forms, such as agricultural/ecological or hydrological droughts. In high latitudes, snow becomes a critical factor in drought propagation, particularly for longer aggregation periods. This is evident in northern regions, where reduced precipitation does not always lead to other types of droughts, likely due to the buffering effect of snowpack. Moreover, not all precipitation deficits necessarily progress into agricultural/ecological or hydrological droughts, as these deficits can be mitigated by the landscape’s buffering capacity (Maxwell et al., 2021). Conversely, not all agricultural/ecological or hydrological droughts are (solely) triggered by precipitation deficits; some result from extraordinarily high evaporative demands, lack of snowmelt, or a combination of several factors.

4.5 Drought Propagation in a Changing Climate

640 The analysis of changes in standardized drought indices (SPI, SSMI, and SSFI) across two climate normal periods (CNP1 and CNP2) revealed distinct regional and seasonal patterns in drought dynamics, which are crucial for understanding the evolving drought risk in various clusters across the studied region. All clusters consistently indicate a wetting trend, i.e. drought conditions across all hydrological components have consistently become less severe - a pattern that aligns with previous findings of increasing precipitation and runoff over Sweden during the twentieth and early twenty-first centuries (Chen et al., 2020; Teutschbein et al., 2022). This trend is stronger in northern catchments, with clusters 1 and 2 featuring especially strong increases during the cold season when snowmelt and seasonal precipitation are likely contributing to the water surplus. This pronounced seasonal signal highlights the importance of snowpack and spring melt in driving the hydrological response in these northern clusters, and suggests that these regions may be less prone to drought in the near term. The southern clusters 4 and 5 show significant decreases in both soil moisture (SSMI) and streamflow (SSFI) indices, despite relatively stable or even slightly increasing precipitation (SPI) during that season. This implies that the available water is not being retained in the system, potentially due to increased evaporation, reduced infiltration, or changes in land use such as urbanization and agriculture. These regions may be increasingly prone to hydrological drought, where water deficits in the soil and streams become more frequent and severe, affecting ecosystems, agriculture, and water supplies. This trend is particularly concerning as it indicates that even if precipitation remains stable, water resources in these southern clusters are becoming less available for both natural ecosystems and human use.

655 The evaluation of the differences in drought propagation times and conditional probabilities of drought occurrence between CNP1 and CNP2 demonstrated that there has been no statistically significant change in drought propagation times to soil moisture and streamflow across all clusters. This implies that, despite global climatic changes and the associated shifts in precipitation patterns, the response times of these hydrological components to droughts have remained consistent over the past 60 years. The failure to reject the null hypothesis (p -values > 0.05) implies that any observed differences in propagation times

660 are not large enough to be statistically significant, and thus, the underlying processes governing drought propagation have likely remained stable across these two periods.

Similarly, the investigation into the conditional probabilities of soil moisture and streamflow droughts following a precipitation drought reveals predominantly no statistically significant change between the two CNPs for the vast majority of the evaluated time lags. This broad failure of the Wilcoxon rank sum test to reject the hypothesis of equal medians (p -values > 0.05) suggests
665 that the likelihood of a soil moisture or streamflow drought occurring after a precipitation deficit has not changed significantly over the 60-year observational record (on an annual basis). This may be different for seasonal drought events, but is – due to the limited number of drought events, owing to their rare nature – more difficult to test given the existing data set. The observed stability in drought propagation times and probabilities suggests that the physical properties of the landscape, such as soil type, vegetation cover, and catchment characteristics, have a more dominant role in controlling drought propagation than any
670 potential changes in climatic conditions between these periods. However, further research – particularly into seasonal dynamics – is needed to unravel these links.

The absence of significant changes in both drought propagation times and the conditional probabilities of drought occurrence over the two climate normal periods has important implications for drought management. Despite an observed increase in average temperature by 2.2 degrees and a 20% increase in precipitation over the past 60 years (Teutschbein et al., 2022), these
675 climatic shifts have not resulted in significant alterations in drought propagation within the study period. Thus, historical data and established models for predicting drought behavior based on past observations may still be relevant and reliable under current climate conditions. This stability allows for a certain level of confidence in using past trends to inform future drought preparedness and mitigation strategies. However, it is also important to recognize that while no significant changes have been detected over the studied periods, ongoing climate change could still pose future challenges and potential alterations in drought
680 propagation in these high-latitude catchments (Teutschbein et al., 2023b). Continuous monitoring and periodic reassessment of these trends will be necessary to detect any emerging shifts in drought dynamics that could impact water resource management and planning.

4.6 Limitations and Potential for Future Research

One key limitation of this study lies in the reliance on modelled soil moisture data derived from the ERA5-Land reanalysis.
685 While reanalysis products offer consistent and spatially comprehensive datasets, they are generated using land surface models that incorporate generalized assumptions about land use, vegetation, and soil types. This has several implications for our analysis. First, the spatial resolution of ERA5-Land (approximately 11 km) may be too coarse to capture local-scale heterogeneity in soil properties, topography, and land use that can significantly influence soil moisture dynamics, particularly in small or heterogeneous catchments. As a result, the propagation of meteorological droughts to soil moisture droughts may
690 be influenced by model structure rather than reflecting purely observed physical processes.

Second, since ERA5-Land already integrates information on land cover and soil types within its modelling framework, any correlations between drought propagation metrics and these same physical features may be muted or confounded. This limits

the explanatory power of our correlation analysis when it comes to identifying the role of specific soil or land use characteristics in driving drought propagation, particularly for soil moisture droughts.

695 Third, due to the lack of consistent, long-term in-situ soil moisture observations across Sweden, we were unable to validate the reanalysis-based soil moisture indices or assess potential biases in the data. This is a known limitation in high-latitude regions and highlights the need for improved soil moisture monitoring networks to support drought research.

Finally, while we focused on standardized drought indices to ensure comparability across drought types and catchments, this approach may oversimplify some of the complex, non-linear interactions between meteorological inputs and hydrological responses. For example, the role of evaporative demand, which is a key driver of soil moisture deficits during the summer months, was not explicitly accounted for in our analysis. Variables such as potential evapotranspiration or vapor pressure deficit could provide valuable additional insights, especially in southern regions where atmospheric demand plays a dominant role in controlling drought severity. Similarly, snow accumulation and snowmelt processes, which strongly influence streamflow and groundwater recharge in northern and mountainous catchments, were only indirectly represented through the use of SPI and SSMI. While some seasonal effects were captured in the propagation patterns, we did not include snow-related variables such as snow water equivalent or snowmelt timing in the analysis. These factors are particularly relevant for high-latitude and snow-dominated systems, where snowmelt can buffer or delay drought onset. Thus, we recommend that future studies incorporate evaporative demand and snow-related variables more explicitly, either through additional predictors in statistical models or by applying process-based hydrological modelling. This would help to better capture the physical mechanisms underlying drought propagation and further improve the understanding of seasonal and regional variability in high-latitude watersheds.

5 Conclusions

This study highlights key patterns in drought propagation across high-latitude regions by analyzing both *propagation time*, (how quickly precipitation deficits impact soil moisture, streamflow and groundwater) and *propagation probability* (the likelihood that a precipitation deficit leads to drought in these components). Soil moisture responds fastest, with a typical lag of one month, followed by streamflow at two months. Groundwater exhibits the longest and most variable response times, averaging four months.

Precipitation deficits are a major driver of droughts in high latitudes, particularly in the early stages. Within the first month of a precipitation deficit, the probability of drought occurrence is highest for soil moisture (48%), followed by streamflow (41%) and groundwater (38%). These probabilities increase with time, showing that prolonged deficits can lead to more widespread and severe droughts. However, not all hydrological droughts are driven solely by precipitation deficits; factors such as high evaporation or insufficient snowmelt likely also contribute. This highlights the need for a more nuanced understanding of complex drought dynamics that includes atmospheric and cryospheric processes, especially under changing climate conditions.

Although all catchments are experiencing a general trend of increased wetness throughout the year, driven by snowmelt and rising precipitation, southern catchments show heightened vulnerability to hydrological drought in the spring. This is largely due to increasing evaporative demand, which offsets stable or slightly rising precipitation levels during that season. Importantly, no significant changes in drought propagation times or probabilities were observed over the past 60 years, but continuous monitoring and reassessment remain essential to detect any emerging shifts in drought dynamics that could impact water resource management in the future.

Drought dynamics in high-latitude regions are influenced by diverse factors, including local topography, hydroclimatic conditions, and soil properties. Soil moisture propagation is governed by catchment features, hydro-climate and land cover, while streamflow and groundwater are more closely linked to water availability.

By focusing on the unique conditions of high-latitude regions, this research has contributed to the broader scientific understanding of drought dynamics, offering new perspectives on the interactions between various hydrological systems during prolonged dry periods. Future studies should expand the geographical scope and explore seasonal and climate change impacts to further validate these findings and enhance their generalizability. As climate change intensified, proactive and region-specific strategies will be essential to safeguard water resources and the ecosystems that depend on them.

6 Data availability

The CAMELS-SE data that support the findings of this study are openly available at the Swedish National Data Service at <https://doi.org/10.57804/t3rm-v029>, SND-ID: 2023-173. Version: 1

7 Author Contribution

- CT: Conceptualization, Methodology, Formal Analysis, Investigation, Resources, Data Curation, Writing – Original Draft, Writing – Review & Editing, Visualization, Supervision, Project administration, Funding acquisition.
- TG: Conceptualization, Methodology, Writing – Review & Editing, Supervision
- MG: Methodology, Writing – Review & Editing, Supervision
- AT: Methodology, Data Curation, Writing – Review & Editing
- RB: Conceptualization, Project administration, Funding acquisition.

8 Competing Interests

The authors declare that they have no conflict of interest.

750 **9 Acknowledgements**

We would like to express our sincere gratitude to Sabrina Atigui for her assistance with data preparation, initial work on compiling relevant literature and helpful input during discussions. During the preparation of this work, the corresponding author used ChatGPT to improve writing style and to check for grammar and spelling. The authors carefully reviewed, edited and revised ChatGPT-generated content to their preferences, and take full responsibility for the content of the publication.

755 **10 References**

AghaKouchak, A., Cheng, L., Mazdiyasni, O., and Farahmand, A.: Global warming and changes in risk of concurrent climate extremes: Insights from the 2014 California drought, *Geophys. Res. Lett.*, 8847–8852, <https://doi.org/10.1002/2014GL062308>@10.1002/(ISSN)1944-8007.CALDROUGHT1, 2015.

Ahopelto, L., Kallio, M., Veijalainen, N., Kouki, R., and Keskinen, M.: Drought hazard and annual precipitation predicted to increase in the Sirppujoki river basin, Finland, *Clim. Serv.*, 31, 100400, <https://doi.org/10.1016/j.cliser.2023.100400>, 2023.

760 Alam, S., Gebremichael, M., Ban, Z., Scanlon, B. R., Senay, G., and Lettenmaier, D. P.: Post-Drought Groundwater Storage Recovery in California’s Central Valley, *Water Resour. Res.*, 57, e2021WR030352, <https://doi.org/10.1029/2021WR030352>, 2021.

Andersson, L.: Soil Moisture Deficits in South-Central Sweden: I – Seasonal and Regional Distributions, *Hydrol. Res.*, 20, 109–122, <https://doi.org/10.2166/nh.1989.0009>, 1989.

765 Arheimer, B. and Lindström, G.: Climate impact on floods: changes in high flows in Sweden in the past and the future (1911–2100), *Hydrol. Earth Syst. Sci.*, 19, 771–784, <https://doi.org/10.5194/hess-19-771-2015>, 2015.

Asadzadeh, M., Tolson, B. A., and Burn, D. H.: A new selection metric for multiobjective hydrologic model calibration, *Water Resour. Res.*, 50, 7082–7099, <https://doi.org/10/f6n5gx>, 2014.

770 Bae, H., Ji, H., Lim, Y.-J., Ryu, Y., Kim, M.-H., and Kim, B.-J.: Characteristics of drought propagation in South Korea: relationship between meteorological, agricultural, and hydrological droughts, *Nat. Hazards*, 99, 1–16, <https://doi.org/10.1007/s11069-019-03676-3>, 2019.

Bakke, S. J., Ionita, M., and Tallaksen, L. M.: The 2018 northern European hydrological drought and its drivers in a historical perspective, *Hydrol. Earth Syst. Sci.*, 24, 5621–5653, <https://doi.org/10/gpg5sg>, 2020.

775 Bales, R. C., Goulden, M. L., Hunsaker, C. T., Conklin, M. H., Hartsough, P. C., O’Geen, A. T., Hopmans, J. W., and Safeeq, M.: Mechanisms controlling the impact of multi-year drought on mountain hydrology, *Sci. Rep.*, 8, 690, <https://doi.org/10.1038/s41598-017-19007-0>, 2018.

Barthel, R., Stangefelt, M., Giese, M., Nygren, M., Seftigen, K., and Chen, D.: Current understanding of groundwater recharge and groundwater drought in Sweden compared to countries with similar geology and climate, *Geogr. Ann. Ser. Phys. Geogr.*, 103, 323–345, <https://doi.org/10.1080/04353676.2021.1969130>, 2021.

780 Beck, H. E., Zimmermann, N. E., McVicar, T. R., Vergopolan, N., Berg, A., and Wood, E. F.: Present and future Köppen-Geiger climate classification maps at 1-km resolution, *Sci. Data*, 5, 180214, <https://doi.org/10.1038/sdata.2018.214>, 2018.

- Bevacqua, A. G., Chaffé, P. L. B., Chagas, V. B. P., and AghaKouchak, A.: Spatial and temporal patterns of propagation from meteorological to hydrological droughts in Brazil, *J. Hydrol.*, 603, 126902, <https://doi.org/10.1016/j.jhydrol.2021.126902>, 2021.
- 785
- Blauhut, V., Stoelzle, M., Ahopelto, L., Brunner, M. I., Teutschbein, C., Wendt, D. E., Akstinis, V., Bakke, S. J., Barker, L. J., Bartošová, L., Briede, A., Cammalleri, C., Kalin, K. C., De Stefano, L., Fendeková, M., Finger, D. C., Huysmans, M., Ivanov, M., Jaagus, J., Jakubínský, J., Krakovska, S., Laaha, G., Lakatos, M., Manevski, K., Neumann Andersen, M., Nikolova, N., Osuch, M., van Oel, P., Radeva, K., Romanowicz, R. J., Toth, E., Trnka, M., Urošev, M., Urquijo Reguera, J., Sauquet, E., Stevkov, A., Tallaksen, L. M., Trofimova, I., Van Loon, A. F., van Vliet, M. T. H., Vidal, J.-P., Wanders, N., Werner, M., Willems, P., and Živković, N.: Lessons from the 2018–2019 European droughts: a collective need for unifying drought risk management, *Nat. Hazards Earth Syst. Sci.*, 22, 2201–2217, <https://doi.org/10.5194/nhess-22-2201-2022>, 2022.
- 790
- Bloomfield, J. P. and Marchant, B. P.: Analysis of groundwater drought building on the standardised precipitation index approach, *Hydrol. Earth Syst. Sci.*, 17, 4769–4787, <https://doi.org/10/f5m79s>, 2013.
- 795
- Blöschl, G., Hall, J., Parajka, J., Perdigão, R. A. P., Merz, B., Arheimer, B., Aronica, G. T., Bilibashi, A., Bonacci, O., Borga, M., Čanjevac, I., Castellarin, A., Chirico, G. B., Claps, P., Fiala, K., Frolova, N., Gorbachova, L., Gül, A., Hannaford, J., Harrigan, S., Kireeva, M., Kiss, A., Kjeldsen, T. R., Kohnová, S., Koskela, J. J., Ledvinka, O., Macdonald, N., Mavrova-Guirguinova, M., Mediero, L., Merz, R., Molnar, P., Montanari, A., Murphy, C., Osuch, M., Ovcharuk, V., Radevski, I., Rogger, M., Salinas, J. L., Sauquet, E., Šraj, M., Szolgay, J., Viglione, A., Volpi, E., Wilson, D., Zaimi, K., and Živković, N.: Changing climate shifts timing of European floods, *Science*, 357, 588–590, <https://doi.org/10.1126/science.aan2506>, 2017.
- 800
- Bodner, G., Nakhforoosh, A., and Kaul, H.-P.: Management of crop water under drought: a review, *Agron. Sustain. Dev.*, 35, 401–442, <https://doi.org/10.1007/s13593-015-0283-4>, 2015.
- Brunner, M. I. and Chartier-Rescan, C.: Drought Spatial Extent and Dependence Increase During Drought Propagation From the Atmosphere to the Hydrosphere, *Geophys. Res. Lett.*, 51, e2023GL107918, <https://doi.org/10.1029/2023GL107918>, 2024.
- 805
- Chai, Q., Gan, Y., Zhao, C., Xu, H.-L., Waskom, R. M., Niu, Y., and Siddique, K. H. M.: Regulated deficit irrigation for crop production under drought stress. A review, *Agron. Sustain. Dev.*, 36, 3, <https://doi.org/10.1007/s13593-015-0338-6>, 2015.
- Chan, W. C. H., Shepherd, T. G., Facer-Childs, K., Darch, G., and Arnell, N. W.: Storylines of UK drought based on the 2010–2012 event, *Hydrol. Earth Syst. Sci.*, 26, 1755–1777, <https://doi.org/10.5194/hess-26-1755-2022>, 2022.
- Changnon, S. A.: Detecting Drought Conditions in Illinois, 41, 1987.
- 810
- Chen, D., Zhang, P., Seftigen, K., Ou, T., Giese, M., and Barthel, R.: Hydroclimate changes over Sweden in the twentieth and twenty-first centuries: a millennium perspective, *Geogr. Ann. Ser. Phys. Geogr.*, 103, 103–131, <https://doi.org/10.1080/04353676.2020.1841410>, 2020.
- Cienciala, E., Kučera, J., and Lindroth, A.: Long-term measurements of stand water uptake in Swedish boreal forest, *Agric. For. Meteorol.*, 98–99, 547–554, [https://doi.org/10.1016/S0168-1923\(99\)00122-7](https://doi.org/10.1016/S0168-1923(99)00122-7), 1999.
- 815
- Cochand, M., Molson, J., Barth, J. A. C., van Geldern, R., Lemieux, J.-M., Fortier, R., and Therrien, R.: Rapid groundwater recharge dynamics determined from hydrogeochemical and isotope data in a small permafrost watershed near Umiujaq (Nunavik, Canada), *Hydrogeol. J.*, 28, 853–868, <https://doi.org/10.1007/s10040-020-02109-x>, 2020.
- Entekhabi, D.: Propagation in the Drought Cascade: Observational Analysis Over the Continental US, *Water Resour. Res.*, 59, e2022WR032608, <https://doi.org/10.1029/2022WR032608>, 2023.

- 820 Garrido-Perez, J. M., Vicente-Serrano, S. M., Barriopedro, D., García-Herrera, R., Trigo, R., and Beguería, S.: Examining the outstanding Euro-Mediterranean drought of 2021–2022 and its historical context, *J. Hydrol.*, 630, 130653, <https://doi.org/10.1016/j.jhydrol.2024.130653>, 2024.
- Geng, G., Zhang, B., Gu, Q., He, Z., and Zheng, R.: Drought propagation characteristics across China: Time, probability, and threshold, *J. Hydrol.*, 631, 130805, <https://doi.org/10.1016/j.jhydrol.2024.130805>, 2024a.
- 825 Geng, G., Zhang, B., Gu, Q., He, Z., and Zheng, R.: Drought propagation characteristics across China: Time, probability, and threshold, *J. Hydrol.*, 631, 130805, <https://doi.org/10.1016/j.jhydrol.2024.130805>, 2024b.
- Geris, J., Tetzlaff, D., McDonnell, J., and Soulsby, C.: The relative role of soil type and tree cover on water storage and transmission in northern headwater catchments, *Hydrol. Process.*, 29, 1844–1860, <https://doi.org/10.1002/hyp.10289>, 2015.
- 830 Gessner, C., Fischer, E. M., Beyerle, U., and Knutti, R.: Multi-year drought storylines for Europe and North America from an iteratively perturbed global climate model, *Weather Clim. Extrem.*, 38, 100512, <https://doi.org/10.1016/j.wace.2022.100512>, 2022.
- Gong, R., Chen, J., Liang, Z., Wu, C., Tian, D., Wu, J., Li, S., and Zeng, G.: Characterization and propagation from meteorological to groundwater drought in different aquifers with multiple timescales, *J. Hydrol. Reg. Stud.*, 45, 101317, <https://doi.org/10.1016/j.ejrh.2023.101317>, 2023.
- 835 Grobicki, A., MacLeod, F., and Pischke, F.: Integrated policies and practices for flood and drought risk management, *Water Policy*, 17, 180–194, <https://doi.org/10.1016/j.watpol.2015.06.009>, 2015.
- Hellwig, J., Stoezle, M., and Stahl, K.: Groundwater and baseflow drought responses to synthetic recharge stress tests, *Hydrol. Earth Syst. Sci.*, 25, 1053–1068, <https://doi.org/10.5194/hess-25-1053-2021>, 2021.
- 840 Hellwig, J., Liu, Y., Stahl, K., and Hartmann, A.: Drought propagation in space and time: the role of groundwater flows, *Environ. Res. Lett.*, 17, 094008, <https://doi.org/10.1088/1748-9326/ac8693>, 2022.
- Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey, C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., and Thépaut, J.-N.: ERA5 monthly averaged data on single levels from 1940 to present, <https://doi.org/10.24381/CDS.F17050D7>, 2023.
- 845 Heudorfer, B. and Stahl, K.: Comparison of different threshold level methods for drought propagation analysis in Germany, *Hydrol. Res.*, 48, 1311–1326, <https://doi.org/10.2166/nh.2016.258>, 2016.
- IPCC: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK and New York, USA, 2021.
- Irannezhad, M., Ronkanen, A.-K., and Kløve, B.: Effects of climate variability and change on snowpack hydrological processes in Finland, *Cold Reg. Sci. Technol.*, 118, 14–29, <https://doi.org/10.1016/j.coldregions.2015.06.009>, 2015.
- 850 Kchouk, S., Melsen, L. A., Walker, D. W., and van Oel, P. R.: A geography of drought indices: mismatch between indicators of drought and its impacts on water and food securities, *Nat. Hazards Earth Syst. Sci.*, 22, 323–344, <https://doi.org/10.5194/nhess-22-323-2022>, 2022.
- 855 Koster, R. D., Schubert, S. D., Wang, H., Mahanama, S. P., and DeAngelis, A. M.: Flash Drought as Captured by Reanalysis Data: Disentangling the Contributions of Precipitation Deficit and Excess Evapotranspiration, *J. Hydrometeorol.*, 20, 1241–1258, <https://doi.org/10.1175/JHM-D-18-0242.1>, 2019.

- Kuiper, N. H.: Tests concerning random points on a circle, *Indag. Math. Proc.*, 63, 38–47, [https://doi.org/10.1016/S1385-7258\(60\)50006-0](https://doi.org/10.1016/S1385-7258(60)50006-0), 1960.
- Kumar, R., Musuuza, J. L., Van Loon, A. F., Teuling, A. J., Barthel, R., Ten Broek, J., Mai, J., Samaniego, L., and Attinger, S.: Multiscale evaluation of the Standardized Precipitation Index as a groundwater drought indicator, *Hydrol. Earth Syst. Sci.*, 20, 1117–1131, <https://doi.org/10.5194/hess-20-1117-2016>, 2016.
- Langridge, R. and Daniels, B.: Accounting for Climate Change and Drought in Implementing Sustainable Groundwater Management, *Water Resour. Manag.*, 31, 3287–3298, <https://doi.org/10.1007/s11269-017-1607-8>, 2017.
- Li, W., Reichstein, M., O, S., May, C., Destouni, G., Migliavacca, M., Kraft, B., Weber, U., and Orth, R.: Contrasting Drought Propagation Into the Terrestrial Water Cycle Between Dry and Wet Regions, *Earths Future*, 11, e2022EF003441, <https://doi.org/10.1029/2022EF003441>, 2023.
- Ma, L., Huang, Q., Huang, S., Liu, D., Leng, G., Wang, L., and Li, P.: Propagation dynamics and causes of hydrological drought in response to meteorological drought at seasonal timescales, *Hydrol. Res.*, 53, 193–205, <https://doi.org/10.2166/nh.2021.006>, 2021.
- Maxwell, C. M., Fernald, A. G., Cadol, D., Faist, A. M., and King, J. P.: Managing flood flow connectivity to landscapes to build buffering capacity to disturbances: An ecohydrologic modeling framework for drylands, *J. Environ. Manage.*, 278, 111486, <https://doi.org/10.1016/j.jenvman.2020.111486>, 2021.
- McKee, T. B., Doesken, N. J., and Kleist, J.: The relationship of drought frequency and duration to time scales, in: *Proceedings of the 8th Conference on Applied Climatology*, Conference on Applied Climatology, Anaheim, California, USA, 179–183, 1993.
- Meresa, H., Zhang, Y., Tian, J., and Abrar Faiz, M.: Understanding the role of catchment and climate characteristics in the propagation of meteorological to hydrological drought, *J. Hydrol.*, 617, 128967, <https://doi.org/10.1016/j.jhydrol.2022.128967>, 2023.
- Mishra, A. K. and Singh, V. P.: A review of drought concepts, *J. Hydrol.*, 391, 202–216, <https://doi.org/10/cktzg6>, 2010.
- Montgomery, D. C. and Runger, G. C.: *Applied Statistics and Probability for Engineers*, John Wiley & Sons, 791 pp., 2010.
- Mukherjee, S., Mishra, A., and Trenberth, K. E.: Climate Change and Drought: a Perspective on Drought Indices, *Curr. Clim. Change Rep.*, 4, 145–163, <https://doi.org/10.1007/s40641-018-0098-x>, 2018.
- Nygren, M., Giese, M., Kløve, B., Haaf, E., Rossi, P. M., and Barthel, R.: Changes in seasonality of groundwater level fluctuations in a temperate-cold climate transition zone, *J. Hydrol. X*, 8, 100062, <https://doi.org/10.1016/j.hydroa.2020.100062>, 2020.
- Nygren, M., Giese, M., and Barthel, R.: Recent trends in hydroclimate and groundwater levels in a region with seasonal frost cover, *J. Hydrol.*, 602, 126732, <https://doi.org/10.1016/j.jhydrol.2021.126732>, 2021.
- Nygren, M., Barthel, R., Allen, D. M., and Giese, M.: Exploring groundwater drought responsiveness in lowland post-glacial environments, *Hydrogeol. J.*, 30, 1937–1961, <https://doi.org/10.1007/s10040-022-02521-5>, 2022.
- Odongo, R. A., De Moel, H., and Van Loon, A. F.: Propagation from meteorological to hydrological drought in the Horn of Africa using both standardized and threshold-based indices, *Nat. Hazards Earth Syst. Sci.*, 23, 2365–2386, <https://doi.org/10.5194/nhess-23-2365-2023>, 2023.

- Orth, R. and Destouni, G.: Drought reduces blue-water fluxes more strongly than green-water fluxes in Europe, *Nat. Commun.*, 9, 3602, <https://doi.org/10.1038/s41467-018-06013-7>, 2018.
- Pearson, K.: Notes on the History of Correlation, *Biometrika*, 13, 25, <https://doi.org/10.2307/2331722>, 1920.
- 895 Pereira, L. S., Louro, V., Rosário, L. do, and Almeida, A.: Desertification, territory and people, a holistic approach in the Portuguese context, in: *Desertification in the Mediterranean Region. A Security Issue*, Dordrecht, 269–289, https://doi.org/10.1007/1-4020-3760-0_11, 2006.
- Petersen-Perlman, J. D., Aguilar-Barajas, I., and Megdal, S. B.: Drought and groundwater management: Interconnections, challenges, and policy responses, *Curr. Opin. Environ. Sci. Health*, 28, 100364, <https://doi.org/10.1016/j.coesh.2022.100364>,
900 2022.
- Potopová, V., Boroneanț, C., Možný, M., and Soukup, J.: Driving role of snow cover on soil moisture and drought development during the growing season in the Czech Republic: DRIVING ROLE OF SNOW COVER ON SOIL MOISTURE AND DROUGHT DEVELOPMENT, *Int. J. Climatol.*, 36, 3741–3758, <https://doi.org/10.1002/joc.4588>, 2016.
- Rajsekhar, D., Singh, Vijay. P., and Mishra, Ashok. K.: Multivariate drought index: An information theory based approach for
905 integrated drought assessment, *J. Hydrol.*, 526, 164–182, <https://doi.org/10.1016/j.jhydrol.2014.11.031>, 2015.
- Robinson, J. S., Sivapalan, M., and Snell, J. D.: On the relative roles of hillslope processes, channel routing, and network geomorphology in the hydrologic response of natural catchments, *Water Resour. Res.*, 31, 3089–3101, <https://doi.org/10.1029/95WR01948>, 1995.
- Saito, L., Christian, B., Diffley, J., Richter, H., Rohde, M. M., and Morrison, S. A.: Managing Groundwater to Ensure
910 Ecosystem Function, *Groundwater*, 59, 322–333, <https://doi.org/10.1111/gwat.13089>, 2021.
- Sattar, M. N., Lee, J.-Y., Shin, J.-Y., and Kim, T.-W.: Probabilistic Characteristics of Drought Propagation from Meteorological to Hydrological Drought in South Korea, *Water Resour. Manag.*, 33, 2439–2452, <https://doi.org/10.1007/s11269-019-02278-9>, 2019.
- Sheffield, J. and Wood, E. F.: Characteristics of global and regional drought, 1950–2000: Analysis of soil moisture data from
915 off-line simulation of the terrestrial hydrologic cycle, *J. Geophys. Res. Atmospheres*, 112, <https://doi.org/10.1029/2006JD008288>, 2007.
- Singh, N. K., Emanuel, R. E., McGlynn, B. L., and Miniati, C. F.: Soil Moisture Responses to Rainfall: Implications for Runoff Generation, *Water Resour. Res.*, 57, e2020WR028827, <https://doi.org/10.1029/2020WR028827>, 2021.
- Skålevåg, A. and Vormoor, K.: Daily streamflow trends in Western versus Eastern Norway and their attribution to hydro-meteorological drivers, *Hydrol. Process.*, 35, e14329, <https://doi.org/10.1002/hyp.14329>, 2021.
- 920 Soulsby, C., Scheliga, B., Neill, A., Comte, J.-C., and Tetzlaff, D.: A longer-term perspective on soil moisture, groundwater and stream flow response to the 2018 drought in an experimental catchment in the Scottish Highlands, *Hydrol. Process.*, 35, e14206, <https://doi.org/10.1002/hyp.14206>, 2021.
- Spearman, C.: The Proof and Measurement of Association between Two Things, *Am. J. Psychol.*, 15, 72–101, <https://doi.org/10.1093/ije/dyq191>, 1904.
- 925 Spinoni, J., Naumann, G., Vogt, J. V., and Barbosa, P.: The biggest drought events in Europe from 1950 to 2012, *J. Hydrol. Reg. Stud.*, 3, 509–524, <https://doi.org/10/gfpg9m>, 2015.

- Spinoni, J., Vogt, J. V., Naumann, G., Barbosa, P., and Dosio, A.: Will drought events become more frequent and severe in Europe?, *Int. J. Climatol.*, 38, 1718–1736, <https://doi.org/10/gc772g>, 2018.
- 930 Srivastav, A. L., Dhyani, R., Ranjan, M., Madhav, S., and Sillanpää, M.: Climate-resilient strategies for sustainable management of water resources and agriculture, *Environ. Sci. Pollut. Res.*, 28, 41576–41595, <https://doi.org/10.1007/s11356-021-14332-4>, 2021.
- Stagge, J. H., Tallaksen, L. M., Gudmundsson, L., Van Loon, A. F., and Stahl, K.: Candidate Distributions for Climatological Drought Indices (SPI and SPEI), *Int. J. Climatol.*, 35, 4027–4040, <https://doi.org/10/gfpg9n>, 2015.
- 935 Staudinger, M., Stahl, K., and Seibert, J.: A drought index accounting for snow, *Water Resour. Res.*, 50, 7861–7872, <https://doi.org/10.1002/2013WR015143>, 2014.
- Stenfors, E., Blicharska, M., Grabs, T., and Teutschbein, C.: Sectoral Vulnerability to Drought: Exploring the Role of Blue and Green Water Dependency in Mid and High-Latitudes, *EGUsphere*, 1–21, <https://doi.org/10.5194/egusphere-2024-2726>, 2024a.
- 940 Stenfors, E., Blicharska, M., Grabs, T., and Teutschbein, C.: User-Validated Drought Vulnerability Factors in Forested Cold Climates: Multi-Sectoral Perspectives from Sweden, *EGUsphere*, 1–29, <https://doi.org/10.5194/egusphere-2024-1988>, 2024b.
- Stoelzle, M., Stahl, K., Morhard, A., and Weiler, M.: Streamflow sensitivity to drought scenarios in catchments with different geology, *Geophys. Res. Lett.*, 41, 6174–6183, <https://doi.org/10.1002/2014GL061344>, 2014.
- Teutschbein, C.: CAMELS-SE: Long-term hydroclimatic observations (1961–2020) across 50 catchments in Sweden as a resource for modelling, education, and collaboration, *Geosci. Data J.*, 00, 1–14, <https://doi.org/10.1002/gdj3.239>, 2024a.
- 945 Teutschbein, C.: Swedish Hydroclimatic Data 1961-2020 – Precipitation, Temperature and Streamflow Observations across 50 Catchments (CAMELS-SE) (1), <https://doi.org/10.57804/T3RM-V029>, 2024b.
- Teutschbein, C., Grabs, T., Karlsen, R. H., Laudon, H., and Bishop, K.: Hydrological response to changing climate conditions: Spatial streamflow variability in the boreal region, *Water Resour. Res.*, 51, 9425–9446, <https://doi.org/10.1002/2015WR017337>, 2015.
- 950 Teutschbein, C., Quesada Montano, B., Todorović, A., and Grabs, T.: Streamflow droughts in Sweden: Spatiotemporal patterns emerging from six decades of observations, *J. Hydrol. Reg. Stud.*, 42, 101171, <https://doi.org/10.1016/j.ejrh.2022.101171>, 2022.
- Teutschbein, C., Albrecht, F., Blicharska, M., Tootoonchi, F., Stenfors, E., and Grabs, T.: Drought hazards and stakeholder perception: Unraveling the interlinkages between drought severity, perceived impacts, preparedness, and management, *Ambio*, 52, 1262–1281, <https://doi.org/10.1007/s13280-023-01849-w>, 2023a.
- 955 Teutschbein, C., Jonsson, E., Todorović, A., Tootoonchi, F., Stenfors, E., and Grabs, T.: Future drought propagation through the water-energy-food-ecosystem nexus – A Nordic perspective, *J. Hydrol.*, 617, 128963, <https://doi.org/10.1016/j.jhydrol.2022.128963>, 2023b.
- 960 Thomann, J. A., Werner, A. D., and Irvine, D. J.: Developing adaptive management guidance for groundwater planning and development, *J. Environ. Manage.*, 322, 116052, <https://doi.org/10.1016/j.jenvman.2022.116052>, 2022.
- Todorović, A., Grabs, T., and Teutschbein, C.: Advancing traditional strategies for testing hydrological model fitness in a changing climate, *Hydrol. Sci. J.*, 67, null, <https://doi.org/10.1080/02626667.2022.2104646>, 2022.

- 965 Todorović, A., Grabs, T., and Teutschbein, C.: Improving performance of bucket-type hydrological models in high latitudes with multi-model combination methods: Can we wring water from a stone?, *J. Hydrol.*, 632, 130829, <https://doi.org/10.1016/j.jhydrol.2024.130829>, 2024.
- Tootoonchi, F., Todorović, A., Grabs, T., and Teutschbein, C.: Uni- and multivariate bias adjustment of climate model simulations in Nordic catchments: Effects on hydrological signatures relevant for water resources management in a changing climate, *J. Hydrol.*, 623, 129807, <https://doi.org/10.1016/j.jhydrol.2023.129807>, 2023.
- 970 Tripathy, K. P. and Mishra, A. K.: How Unusual Is the 2022 European Compound Drought and Heatwave Event?, *Geophys. Res. Lett.*, 50, e2023GL105453, <https://doi.org/10.1029/2023GL105453>, 2023.
- UNDRR: GAR Special report on drought 2021, United Nations Office for Disaster Risk Reduction, Geneva, Switzerland, 173 pp., 2021.
- 975 Van Loon, A. F.: Hydrological drought explained, *Wiley Interdiscip. Rev. Water*, 2, 359–392, <https://doi.org/10.1002/wat2.1085>, 2015.
- Van Loon, A. F. and Van Lanen, H. A. J.: A process-based typology of hydrological drought, *Hydrol. Earth Syst. Sci.*, 16, 1915–1946, <https://doi.org/10.5194/hess-16-1915-2012>, 2012.
- 980 Van Loon, A. F., Ploum, S. W., Parajka, J., Fleig, A. K., Garnier, E., Laaha, G., and Van Lanen, H. A. J.: Hydrological drought types in cold climates: quantitative analysis of causing factors and qualitative survey of impacts, *Hydrol. Earth Syst. Sci.*, 19, 1993–2016, <https://doi.org/10/f698s9>, 2015.
- Vaux, H.: Groundwater under stress: the importance of management, *Environ. Earth Sci.*, 62, 19–23, <https://doi.org/10.1007/s12665-010-0490-x>, 2011.
- Vicente-Serrano, S. M., López-Moreno, J. I., Beguería, S., Lorenzo-Lacruz, J., Azorin-Molina, C., and Morán-Tejeda, E.: Accurate computation of a streamflow drought index, *J. Hydrol. Eng.*, 17, 318–332, <https://doi.org/10/c9ks4k>, 2011.
- 985 Vicente-Serrano, S. M., Beguería, S., Lorenzo-Lacruz, J., Camarero, J. J., López-Moreno, J. I., Azorin-Molina, C., Revuelto, J., Morán-Tejeda, E., and Sanchez-Lorenzo, A.: Performance of Drought Indices for Ecological, Agricultural, and Hydrological Applications, *Earth Interact.*, 16, 1–27, <https://doi.org/10/gfpg9s>, 2012.
- 990 Wang, H., Zhu, Y., Qin, T., and Zhang, X.: Study on the propagation probability characteristics and prediction model of meteorological drought to hydrological drought in basin based on copula function, *Front. Earth Sci.*, 10, <https://doi.org/10.3389/feart.2022.961871>, 2022.
- Weider, K. and Boutt, D. F.: Heterogeneous water table response to climate revealed by 60 years of ground water data, *Geophys. Res. Lett.*, 37, <https://doi.org/10.1029/2010GL045561>, 2010.
- 995 Wendt, D. E., Bloomfield, J. P., Van Loon, A. F., Garcia, M., Heudorfer, B., Larsen, J., and Hannah, D. M.: Evaluating integrated water management strategies to inform hydrological drought mitigation, *Nat. Hazards Earth Syst. Sci.*, 21, 3113–3139, <https://doi.org/10.5194/nhess-21-3113-2021>, 2021.
- Wilhite, Donald A.: A methodology for drought preparedness, *Nat. Hazards*, 13, 229–252, <https://doi.org/10/c3br6g>, 1996.
- Wilson, D., Hisdal, H., and Lawrence, D.: Has streamflow changed in the Nordic countries? – Recent trends and comparisons to hydrological projections, *J. Hydrol.*, 394, 334–346, <https://doi.org/10/b3tgch>, 2010.

WMO Climatological Normals: <https://community.wmo.int/wmo-climatological-normals>, last access: 16 December 2021.

- 1000 WMO and GWP: Handbook of Drought Indicators and Indices, edited by: Svoboda, M. and Fuchs, B. A., Integrated Drought Management Programme (IDMP), Geneva, Switzerland, <https://doi.org/10.1201/9781315265551-12>, 2016.
- Wu, H., Hayes, M. J., Wilhite, D. A., and Svoboda, M. D.: The effect of the length of record on the standardized precipitation index calculation, *Int. J. Climatol.*, 25, 505–520, <https://doi.org/10/c3bczb>, 2005.
- 1005 Zhang, Q., Miao, C., Gou, J., Wu, J., Jiao, W., Song, Y., and Xu, D.: Spatiotemporal characteristics of meteorological to hydrological drought propagation under natural conditions in China, *Weather Clim. Extrem.*, 38, 100505, <https://doi.org/10.1016/j.wace.2022.100505>, 2022a.
- Zhang, X., Hao, Z., Singh, V. P., Zhang, Y., Feng, S., Xu, Y., and Hao, F.: Drought propagation under global warming: Characteristics, approaches, processes, and controlling factors, *Sci. Total Environ.*, 838, 156021, <https://doi.org/10.1016/j.scitotenv.2022.156021>, 2022b.
- 1010 Zhu, Y., Liu, ·Yi, Wang, W., Singh, V. P., and Ren, L.: A global perspective on the probability of propagation of drought: From meteorological to soil moisture, *J. Hydrol.*, 603, 126907, <https://doi.org/10.1016/j.jhydrol.2021.126907>, 2021.