

1 Response of the link between ENSO and the East Asian winter 2 monsoon to Asian anthropogenic sulfate aerosols

3 Zixuan Jia^{1,2}, Massimo A Bollasina³, Wenjun Zhang^{1,2}, Ying Xiang⁴,

Deleted: 2

Deleted: 3

5 ¹State Key Laboratory of Climate System Prediction and Risk Management/Key Laboratory of Meteorological
6 Disaster, Ministry of Education/Collaborative Innovation Center on Forecast and Evaluation of Meteorological
7 Disasters, Nanjing University of Information Science and Technology, Nanjing, China

8 ²School of Atmospheric Science, Nanjing University of Information Science and Technology, Nanjing, China

Deleted: 1

9 ³School of GeoSciences, University of Edinburgh, Edinburgh, UK

Deleted: 1

10 ⁴Jiangsu Climate Center, Nanjing, China

Deleted: 2

Deleted: 3

12 Correspondence to: Zixuan Jia (zx.jia@nuist.edu.cn)

13 **Abstract.** We use coupled and atmosphere-only simulations from the Precipitation Driver and
14 Response Model Intercomparison Project to investigate the impacts of Asian anthropogenic sulfate
15 aerosols on the link between the El Niño-Southern Oscillation (ENSO) and the East Asian Winter
16 monsoon (EAWM). In fully-coupled simulations, aerosol-induced cooling extends southeastward to the
17 Maritime Continent and the north-western Pacific. Remotely, this broad cooling weakens the easterly
18 trade winds over the central Pacific, which reduces the east-west equatorial Pacific sea surface
19 temperature gradient. These changes contribute to increasing ENSO's amplitude by 17%, mainly
20 through strengthening the zonal wind forcing. Concurrently, the El Niño-related warm SST anomalies
21 and the ensuing Pacific-East Asia teleconnection pattern (i.e. the ENSO-EAWM link) intensify, leading
22 to an increased EAWM amplitude by 18% in the coupled simulations. Therefore, in response to the
23 increasing frequency of El Niño and La Niña years under Asian sulfate aerosol forcing, the interannual
24 variability of the EAWM increases, with more extreme EAWM years. The opposite variations in the
25 interannual variability of the EAWM to Asian aerosols in atmosphere-only simulations (-19%) further
26 reflect the importance of ENSO-related atmosphere-ocean coupled processes. A better understanding
27 of the changes of the year-to-year variability of the EAWM in response to aerosol forcing is critical to
28 reducing uncertainties in future projections of variability of regional extremes, such as cold surges and
29 flooding, which can cause large social and economic impacts on densely populated East Asia.

30 1 Introduction

31 The East Asian winter monsoon (EAWM) is one of the most prominent features of the northern
32 hemisphere atmospheric circulation during the boreal winter, and has a pronounced influence on weather
33 and climate of the Asian-Pacific region from the northern latitudes to the equator (e.g. Chang, 2006;
34 Zhou and Wu., 2010; Wang et al., 2022b). As such, the year-to-year variations of the EAWM have the
35 potential to cause extreme cold disasters and severe flooding in Southeast Asian countries (e.g. Huang

Deleted: ,

Deleted:

Formatted: Font: 11 pt, Not Italic

1 et al. 2012; Yang et al., 2020; Zuo et al., 2022), with consequent marked social and economic impacts
2 (e.g. Chen et al., 2005; Zhou et al. 2011). Thus, it is very important to understand the mechanisms
3 underpinning its variability and associated drivers, and to ultimately develop more robust projections
4 of its future evolution.

Deleted: As such, variations of the EAWM have the potential to cause extreme cold disasters and severe flooding in Southeast Asian countries (e.g. Feng et al. 2010; Huang et al. 2012), with consequent marked social and economic impacts (e.g. Chen et al., 2005; Zhou et al. 2011)

5
6 The EAWM is fundamentally driven by the thermal contrast between the cold Asian continent and the
7 adjacent warm oceans (e.g. Yang et al., 2002; Huang et al., 2012; Chen et al., 2019). Its climatological
8 pattern is mainly characterized by dry cold low-level northwesterlies along the eastern flank of the
9 Siberian High and low-level northeasterlies along the coast of East Asia, triggering cold air outbreaks
10 in northern China and generating cold surges over southern China as well as the South China Sea (Li
11 and Wang, 2012; He et al., 2013). ~~The EAWM exhibits distinct interannual variability (e.g. Gong et al.,~~
12 ~~2014; Chen et al. 2015) that mainly originates from intrinsic atmospheric processes (e.g., Wu et al.~~
13 ~~2014; Wang et al. 2021, 2022b) and the El Niño-Southern Oscillation (ENSO) forcing through the~~
14 ~~Pacific-East Asia teleconnection (Zhang et al., 1996).~~ Associated with an El Niño event, the anomalous
15 anticyclone over the western tropical Pacific (the most remarkable low-level circulation feature of the
16 PEA) induces southwesterlies on its western flank, which weaken the EAWM flow and lead to warmer
17 and wetter conditions over southeastern China and the South China Sea (Wang et al., 2000, 2013). In
18 turn, the EAWM tends to be strong during La Niña winters, with widespread cooling and reduced
19 precipitation. ▲

Deleted: The EAWM exhibits distinct interannual variability (e.g. Gong et al., 2014),

Formatted: Font: 11 pt, Not Italic

Deleted: which is strongly influenced by the El Niño-Southern Oscillation (ENSO) and the ensuing Pacific-East Asia (PEA) teleconnection pattern (e.g. Zhang et al., 1996).

Formatted: Font colour: Auto, (Asian) Chinese (China)

20
21 Previous studies indicated that magnitude and location of ENSO-induced teleconnection patterns are
22 influenced by ENSO characteristics, such as amplitude and location of its sea surface temperature (SST)
23 anomalies (Cai et al., 2021; Jiang et al., 2022). However, future projections of ENSO characteristics are
24 highly uncertain, even in the latest CMIP6 models (Huang and Xie, 2015; Yan et al., 2020; Beobide-
25 Arsuaga et al., 2021). Therefore, there is no consensus on future changes in the ENSO-induced
26 teleconnections, including projections of the PEA pattern (e.g. Wang et al., 2013; Jia et al., 2020). The
27 characteristics of ENSO and its induced atmospheric teleconnections are closely related to the tropical
28 Pacific mean state via ocean-atmosphere feedbacks (Jin, 1997; Wang, 2002; Cai et al., 2014). Based on
29 ocean-atmosphere reanalyses, observed mean state changes since the 1980s feature a La Niña-like
30 warming (i.e. the tropical Pacific warming center is mainly located in the western basin; Rayner et al.,
31 2003; Kobayashi et al., 2015; Huang et al., 2017). However, both a La Niña-like and an El Niño-like
32 warming (i.e. tropical Pacific warming centered in the eastern basin) are projected in the future, with a
33 large spread across different climate models (e.g. Power et al., 2013; Lian et al., 2018). These two
34 different warming patterns will cause a corresponding strengthening and weakening of the easterly trade
35 winds over the tropical Pacific Ocean, respectively, resulting in opposite changes in the characteristics
36 of ENSO (Vecchi et al., 2006; Collins et al., 2010). While the majority of the studies have focused on

1 the influence of increasing greenhouse gas concentrations on the tropical Pacific mean state (e.g. Wang
2 et al., 2017; Yan et al., 2020), the impact of anthropogenic aerosols has been largely overlooked.

3
4 Due to the intensification of human industrial activities, the global mean atmospheric burden of
5 anthropogenic aerosols has continued to increase over the past century, exerting a significant imprint
6 on worldwide climate (Liao et al., 2015; Forster et al., 2021; Persad, 2023). Anthropogenic aerosols can
7 affect climate by modulating shortwave radiation and, to some extent, longwave radiation directly, and
8 through their interactions with clouds and precipitation indirectly (Boucher et al., 2013; Myhre et al.,
9 2013; Zhao and Suzuki, 2019). Unlike greenhouse gases, which are distributed evenly across the globe,
10 anthropogenic aerosols reside in the atmosphere for a short time (days to weeks) due to numerous
11 chemical and physical removal processes, which causes their distribution and associated radiative
12 forcing to be spatially heterogeneous (Allen et al., 2015; Wilcox et al., 2019). As such, aerosols can
13 induce substantial changes in local atmospheric circulation and extend their influence over long
14 distances, even over the surrounding ocean, triggering ocean–atmosphere interactions (Rotstayn and
15 Lohmann, 2002; Ramanathan et al., 2005; Westervelt et al., 2020). Some studies indicated that the
16 influence of anthropogenic aerosols from remote sources can even outweigh that of locally-emitted ones
17 (Shindell et al., 2012; Lewinschal et al., 2013). Since the start of the industrial age, vast emissions of
18 aerosols and their precursors over the Northern Hemisphere have had a profound cooling effect, and
19 this preferential cooling has been linked to a southward shift of the Intertropical Convergence Zone (e.g.
20 Hwang et al., 2013; Navarro et al., 2017).

21
22 The emissions of anthropogenic aerosols and their precursors in Asia have increased rapidly since 1980,
23 and many studies have focused on Asian as well as Northern Hemispheric climate (e.g. Bollasina et al.,
24 2014; Bartlett et al., 2018; Wilcox et al., 2019; Li et al., 2022). While Asian anthropogenic aerosols can
25 significantly affect the Asian monsoon, the large majority of the current literature has focused on the
26 effects of aerosols on the summer or annual mean climatology (e.g., Westervelt et al., 2018; Song et al.,
27 2014; Persad et al., 2022). Only a limited number of studies have focused on the influence of aerosols
28 on the EAWM (Jiang et al., 2017; Liu et al., 2019; Wilcox et al., 2019), while their effect on the
29 interannual variability of the EAWM and the link to ENSO remains unexplored. In boreal winter,
30 intensive combustion of coal and fossil fuels across Asia leads to sulfate aerosols dominance (Gao et
31 al., 2018; Cheng et al., 2019), setting the stage for a potential important influence on continental climate
32 and the mean EAWM circulation. Moreover, ENSO and the associated PEA teleconnection pattern peak
33 in the winter, representing a major driver of interannual fluctuations of the EAWM. The extent to which
34 aerosols may affect ENSO and the related ocean-atmosphere feedbacks has not been thoroughly
35 investigated and is unclear (Westervelt et al., 2018; Wilcox et al., 2019). Given the rapid variations in
36 aerosol emissions over Asia, addressing this knowledge gap is both compelling and timely for

Formatted: Font: (Default) +Body (Times New Roman), 11
pt, Font colour: Text 1, Pattern: Clear

Deleted: fuels are combusted intensively

Formatted: Font: (Default) +Body (Times New Roman), 11
pt, Not Bold, Font colour: Text 1, Pattern: Clear

1 enhancing our understanding and projections of the ENSO-EAWM link in the near future, and potential
2 causes of changes in the interannual variability of the EAWM.

3

4 In this study, we use multi-model mean data from regional aerosol perturbation experiments conducted
5 with coupled and atmosphere-only models (Section 2) to investigate the impacts of Asian sulfate
6 aerosols on the ENSO-EAWM link and the interannual variability of the EAWM (Section 3). We then
7 link changes in the PEA pattern to the remote impacts of Asian aerosols on ENSO (Section 4).
8 Mechanisms driving changes in the tropical Pacific mean state and ENSO characteristics are further
9 investigated in Section 5. Finally, Section 6 summarises the main results and provides key conclusions.

10 2 Data and methodology

11 Model data from the Precipitation Driver and Response Model Intercomparison Project (PDRMIP;
12 Myhre et al., 2017) are used to investigate the impact of Asian anthropogenic aerosols on the ENSO-
13 EAWM link. PDRMIP offers a unique opportunity for elucidating the complexities of the ENSO-
14 EAWM-aerosol nexus and its mechanisms, particularly with regard to the role of air-sea interactions in
15 modulating the aerosol-driven response. Indeed, one approach that has provided valuable insights, is
16 the decomposition of the response into two complementary components: a fast response involving
17 atmospheric and land surface adjustments but fixed sea surface temperature (SST), acting on short
18 timescales (a few years) and a slow response, which includes the full extent of the oceanic circulation
19 response, thus effective on decadal or longer time scales (e.g. Samset et al., 2016; Liu et al., 2018; Dow
20 et al., 2021; Fahrenbach et al., 2024; Liu et al., 2024). The baseline simulation was forced by present-
21 day (year 2000) levels of aerosols and greenhouse gas emissions/concentrations. The regional aerosol
22 experiment analysed in this study has sulfate concentrations/emissions over Asia (10°–50°N, 60°–
23 140°E) increased by a factor of 10 compared to the baseline values (hereafter SUL×10Asia). Note that
24 sulfate is the predominant aerosol component in boreal winter over Asia (e.g. Liu et al., 2009; Zhang et
25 al., 2018). The response to Asian aerosols is identified as the difference between the SUL×10Asia and
26 the baseline experiments. Of the 10 models that contributed to PDRMIP, seven performed the
27 SUL×10Asia experiment: GISS-E2, HadGEM3-GA4, IPSL-CM5A, MIROC-SPRINTARS, ESM1-
28 CAM4, CESM1-CAM5, and NorESM1 (details on the resolution and aerosol setup for each model can
29 be found in Table 1 of Liu et al. (2018)). For each model and experiment, a pair of simulations was
30 performed: one in a fully coupled atmosphere–ocean setting (called “coupled”), and one with fixed
31 climatological sea surface temperatures (called fSST). The coupled simulations were run for 100 years
32 and the fSST simulations for 15 years. The concentrations of all non-aerosol anthropogenic forcings and
33 natural forcing were kept at present-day levels (typically year 2000) in all the experiments, as are the
34 SSTs for the fSST simulations. In this study, we use output from the last 50 winters (DJF, December of
35 the current year and January and February of the following year) of coupled simulations and the last 12

Deleted: PDRMIP offers a unique opportunity for elucidating the complexities of the slow and fast responses of the EAWM to Asian aerosols and the contribution of ENSO-related ocean-atmosphere coupled processes with coupled and atmosphere-only simulations by comparing baseline and regional aerosol perturbation experiments.

Formatted: Font: 11 pt, Not Italic

winters of the fSST simulations to discard the model spin-up time and consistently with existing literature (Liu et al., 2018; Dow et al., 2021; Fahrenbach et al., 2024).

Reanalysis and observational data for DJF 1965–2014 (50 years) are used to evaluate the PDRMIP-simulated EAWM and ENSO-related patterns in the baseline experiment. Monthly meteorological reanalysis data are from the fifth-generation atmospheric reanalysis ERA5 provided by the European Centre for Medium-Range Weather Forecasts at a spatial resolution of 0.25° (Copernicus Climate Change Service, 2017; Hersbach et al., 2023). Monthly gridded observations are from the Hadley Centre Sea Ice and Sea Surface Temperature (HadISST) dataset for sea surface temperature at a spatial resolution of 1° (Rayner et al., 2003), and from the Climatic Research Unit (CRU) v4.07 data set for land surface temperature with a spatial resolution of 0.5° (Harris et al., 2020). To quantify the EAWM interannual variability, we use the Ji et al. (1997) index (the negative 1000 hPa meridional wind anomaly averaged over 10° – 30° N, 115° – 130° E) as it represents the spatio-temporal characteristics of the ENSO–EAWM relationship well (Gong et al., 2015; Jia et al., 2020). Positive values indicate a stronger-than-normal EAWM. ENSO is described by the Niño3.4 index (area-averaged SST anomaly over 5° S– 5° N, 120° – 170° W). The ENSO-related PEA pattern is deduced by regression analysis, and the statistical significance is evaluated using the two-tailed Student's *t*-test. Among the seven PDRMIP models with the SUL×10Asia experiment, coupled baseline simulations in CESM1-CAM5, MIROC-SPRINTARS, HadGEM3-GA4, and NorESM1 can well capture the observed pattern and magnitude of the ENSO-related circulation anomalies across East Asia and the Pacific (Fig. S1) and are used in this study. These four models include parameterisations of both aerosol-radiation and aerosol-cloud interactions, while the others don't include indirect effects, or include only the first indirect effect (Liu et al., 2018; Dow et al., 2021). (Table 1). All the data are interpolated to a $3.75^\circ \times 2^\circ$ (longitude $\times$ latitude) resolution before the analysis for consistency between all models.

3 Impacts of Asian aerosols on the PEA pattern and the EAWM interannual variability

The ENSO-related circulation and precipitation anomalies across East Asia and the Pacific (i.e. the PEA pattern) (Figs. 1a-c) are well reproduced by the multi-model mean of the PDRMIP coupled baseline simulations (Figs. 1d-f). The pattern is characterised by the El Niño-related warm SST anomalies over the equatorial Pacific and cold SSTs over the north-western Pacific (Fig. 1a), the anomalous anticyclone over the western tropical Pacific and the anomalous low over the northern extratropical Pacific (Fig. 1b). On the western flank of the anticyclone, near-surface and lower tropospheric southerly winds along the East Asian coast (Figs. 1a-b) lead to warm surface air temperature and precipitation over southeastern China and even over central China (Figs. 1a, c), while the lower tropospheric northerly winds on the western flank of the cyclone bring cold air to northeastern China (Fig. 1b). The spatial patterns of simulated anomalies are broadly similar to those found in observations, including the

position and magnitude of El Niño-related warm SST anomalies, anticyclone and cyclone anomalies, and precipitation anomalies (Figs. 1d-f). The multi-model mean from PDRMIP shares common biases with other CMIP5 and CMIP6 models, such as a slightly westward shift of the equatorial Niño warming with associated circulation and precipitation anomalies (Gong et al., 2015; Wang et al., 2022a). Overall, the multi-model mean coupled PDRMIP baseline simulations successfully reproduce the PEA pattern.

In response to Asian aerosols, the El Niño-related warm SST anomalies intensify over the eastern equatorial Pacific, associated with an intensification of the anomalous SST cooling over the western tropical Pacific (Figs. 1g, j). Concurrently, the anticyclonic anomalies over the western tropical Pacific strengthen and stretch northwestward, while the cyclone over the northern Pacific strengthens and covers a broader region (Figs. 1h, k). This enhanced anticyclone results in an intensification of southerly anomalies along the Asian coast from the South China Sea (Figs. 1j, k), advecting warm and humid air (Figs. 1i, l). Over land, warm anomalies over southeastern and central China weaken, as well as cold anomalies over northeastern China (Figs. 1d, g, j), primarily associated with the Asian aerosol-induced cooling (Fig. S2k) and the enhanced, northwestward-expanding PEA teleconnection pattern (Figs. 1e, h, k, Fig.S3), respectively. Overall, these changes suggest that the ENSO signal and its induced PEA pattern enhance with northwestward expansion under increased Asian aerosols. Given the interannual variability of the EAWM is strongly influenced by the PEA pattern, the intensification of southerly anomalies along the Asian coast associated with the enhanced PEA may lead to an increase in the interannual variability of the EAWM.

Changes in the interannual variability of the EAWM in response to Asian aerosol increase are shown by the probability distributions of the EAWM index (Fig. 2a, S4a). The simulated amplitude of the EAWM (defined as the standard deviation of the EAWM index) is slightly smaller than the observed amplitude in baseline simulations, which is a general known bias in models (Wang et al., 2010; Gong et al., 2014). In coupled simulations, the multi-model mean EAWM amplitudes increase by 18% due to the Asian aerosols at both the 12-year (Fig. 2a) and 50-year (Fig. S4a) timescales, together with more extreme EAWM years in the SUL×10Asia experiment. Note however differences in the shifts of the tails of the distributions of 12-year and 50-years periods due to sampling. Differences in standard deviations of V1000 between SUL×10Asia and baseline experiments (Fig. 2b, S4c) further confirm that the prevailing northerly wind region of the EAWM, with large V1000 standard deviations (mainly along the East Asian coast) in baseline experiment (Fig. S4b), exhibits an increase in SUL×10Asia simulations. These changes are consistent with the aerosol-enhanced PEA pattern identified above. However, in fSST simulations, the multi-model mean EAWM amplitude decreases by 19% at the 12-year timescale, accompanied by more strong-EAWM years and less weak-EAWM years in SUL×10Asia experiments (Fig. 2c-d). These changes can be explained by aerosol-induced cooling over

Deleted: d

Deleted: g

Deleted: e

Deleted: h

Deleted: g

Deleted: h

Deleted: f

Deleted: i

Deleted: Over land, warm and wet anomalies over southeastern and central China weaken, as well as cold anomalies over northeastern China (Figs. 1d, f, g, i), consistent with the changes in the atmospheric circulation patterns mentioned above

Formatted: Font: (Default) +Body (Times New Roman), 11 pt, Font colour: Auto, (Asian) Chinese (China), Pattern: Clear

Deleted: .

Formatted: Font: (Default) +Body (Times New Roman), 11 pt, Font colour: Auto, Pattern: Clear

Formatted: Font colour: Auto

Formatted: Font: (Default) +Body (Times New Roman), 11 pt, Font colour: Text 1, Pattern: Clear

Formatted: Font: (Default) +Body (Times New Roman), 11 pt, Font colour: Text 1, Pattern: Clear

Formatted: Font: (Default) +Body (Times New Roman), 11 pt, Font colour: Text 1, Pattern: Clear

Deleted: are 0.55 and 0.65 m s⁻¹ for the baseline and SUL×10Asia experiments, respectively, indicating an 18% increase due to the Asian aerosols

Formatted: Font: (Default) +Body (Times New Roman), 11 pt, Font colour: Text 1, Pattern: Clear

Deleted: (Fig. 2a)

Deleted:

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt, (Asian) Chinese (China)

Formatted: Font: (Default) Times New Roman, 11 pt, Not Bold, Font colour: Text 1, Pattern: Clear

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Formatted: Font: 11 pt

Deleted: b

1 the emission region and the formation of an anomalous anticyclonic circulation (e.g. Hu et al., 2015;
 2 Liu et al., 2019; Dow et al., 2021), and indicate an enhanced climatological pattern of the EAWM under
 3 increased aerosols (Figs. S2a-f). In addition to this atmospheric-only response, the influence of Asian
 4 aerosols can extend over the Maritime Continent and the north-western Pacific (Wilcox et al., 2019;
 5 Dow et al., 2021; **Figure Sg-l**). In coupled simulations, the climatological pattern of the EAWM extends
 6 southeastward, which is mainly represented by an anomalous anticyclone centred over the southwest of
 7 the Philippines (Figs. S2g, i). This anomalous anticyclone, attributed to the southward shift of the
 8 Hadley circulation to compensate for the interhemispheric asymmetry in aerosol radiative cooling (Liu
 9 et al., 2019), enhances the northerlies over the Maritime Continent but slightly weakens the northerlies
 10 along the East Asian coast (Figs. S3g-h, j-k). This pattern cannot explain the increased interannual
 11 variability of the EAWM in coupled simulations as it is not associated with an evident modulation of
 12 the climatological monsoon flow. The EAWM-related circulation and precipitation anomalies brought
 13 about by increased aerosols in the coupled experiments (Fig. S5) feature an enhanced PEA pattern. This
 14 further suggests the contribution of the enhanced ENSO-induced PEA pattern to increased interannual
 15 variability of the EAWM. The opposite variations in the interannual variability of the EAWM to Asian
 16 aerosols in fully coupled experiments and atmosphere-only (+18% and -19%, respectively) also reflect
 17 the importance of ENSO-related atmosphere-ocean coupled processes. ▲

19 4 The response of ENSO amplitude to increased Asian aerosols

20 Following previous studies (e.g. Wang et al., 2013; Wang et al., 2022a), the increased ENSO signal and
 21 its induced teleconnection pattern can be further linked to changes in the ENSO amplitude (defined as
 22 the standard deviation of the Niño3.4 index). Figure 3a shows the observed standard deviation of SST
 23 across the tropical Pacific, with the highest values over the central-eastern equatorial Pacific. This
 24 spatial pattern is well captured by the multi-model mean in the coupled baseline simulation (Fig. 3b),
 25 albeit the core values are slightly underestimated in magnitude and spatial extent, especially in the
 26 meridional direction. Increased aerosols lead to significant increases in the SST standard deviation over
 27 the Maritime Continent and the central-eastern equatorial Pacific (Fig. 3c-d). This is consistent with the
 28 increased ENSO signal and the related changes in SST anomalies over these two regions (Fig. 1j).
 29 Figure 3e shows the probability distributions of the Niño3.4 index from the coupled baseline (blue curve
 30 and shading) and SUL×10Asia (red curve and shading) simulations. The multi-model mean ENSO
 31 amplitude increases by 17% under aerosol forcing (from 0.7 °C to 0.82 °C).

32
 33 Consistent with the increased ENSO amplitude, Table 1 shows that there are more El Niño (Niño3.4
 34 index > 0.5 °C) and La Niña (the Niño3.4 index < -0.5 °C) years in the coupled SUL×10Asia simulation
 35 compared to the baseline for each model, with the increase up to 100% (from 14 to 28 events in the 50-
 36 year record). Figure 4 shows the joint distributions of multi-model mean aerosol-driven changes in the

Deleted: er

Deleted: ly

Deleted: 2

Deleted: -i

Deleted: 2

Deleted: -h

Deleted: 3

Formatted: Font colour: Text 1

Deleted: g

Deleted: d

Deleted: ly

1 Niño3.4 index compared with the EAWM index in coupled simulations. Both the Niño3.4 index and
 2 the EAWM index have a wide range of variations (i.e. from -1.5 to 1.5 °C and -1 to +1 m s⁻¹ respectively),
 3 suggesting that both the ENSO amplitude and the interannual variability of the EAWM increase under
 4 Asian aerosol forcing as indicated above. Remarkably, changes in the Niño3.4 index are significantly
 5 anti-correlated ($p < 0.01$) with those in the EAWM index ($r = -0.38$). In particular, when the Niño3.4
 6 index decreases by less than 0.5 °C due to Asian aerosol forcing, the EAWM is 2.5 times more likely to
 7 strengthen than weaken, and vice versa. This is consistent with the negative relationship between ENSO
 8 and the EAWM induced by the ensuing PEA teleconnection pattern (Wang et al., 2000). These results
 9 show that Asian aerosols lead to an increase in the ENSO amplitude, resulting in increased interannual
 10 variability of the EAWM through the associated PEA pattern.

11 5 Changes in the tropical Pacific mean state and ocean-atmosphere feedbacks

12 It is well-known that ENSO is fundamentally governed by ocean-atmosphere coupled processes in the
 13 tropical Pacific (Timmermann et al., 2018; Rashid et al., 2022). It is therefore interesting to examine
 14 how the tropical Pacific mean state and atmosphere-ocean coupling are affected by Asian aerosol
 15 forcing. Figure 5 shows the climatological annual variation of key surface variables across the
 16 equatorial Pacific Ocean in the coupled baseline simulation and their changes under increased Asian
 17 aerosols. In the baseline simulation, the equatorial Pacific mean state is characterised by easterly trade
 18 winds with maximum magnitude over the central-eastern Pacific, an east-west SST gradient, and strong
 19 SST amplitudes (i.e. standard deviations of SST) over the eastern Pacific (Figs. 5a-c). These features
 20 are altered in the SUL×10Asia experiment relative to the baseline experiment, with significant seasonal
 21 differences. In particular, anomalous westerlies develop from spring over the eastern Pacific, then
 22 gradually strengthen until the peak in September while moving towards the central Pacific (the Niño4
 23 region, purple bar) (Fig. 5d). Westerly wind anomalies are considered to play an important role during
 24 the development stage (i.e. boreal autumn) of ENSO events, by generating warm SST anomalies in the
 25 eastern equatorial Pacific via the thermocline and the advective feedbacks (McPhaden, 1999; Lian and
 26 Chen, 2021; Xuan et al., 2024). This anomalous westerly flow weakens the climatological easterly trade
 27 winds in the coupled SUL×10Asia simulation compared to the baseline (Figs. 5a, d). Furthermore,
 28 anomalous SST warming appears over the eastern Pacific (the Niño3 region, green bar) from autumn
 29 to winter (peak around October) (Fig. 5*h*), which decreases the east-west equatorial Pacific SST gradient
 30 (Fig. 5b*e*). Note that Figures 5b*e* and 5*h* show SST minus zonal mean and SST difference minus
 31 zonal mean respectively to clarify the east-west SST changes gradient. Given the broad aerosol-induced
 32 cooling over the Pacific (Fig. S2*k*), warming SST anomalies on Figure 5*h* represent less cooling.
 33 Although the east-west equatorial Pacific SST gradient weakens (Fig. 5h), westerly wind anomalies
 34 over the central Pacific are larger (Fig. 5g), sustaining the eastward advection of warm water and
 35 reinforcing the positive SST anomalies over the eastern Pacific. Correspondingly, the SST amplitude
 36 increases with maximum values in the winter mainly over the central-eastern Pacific (the Niño3.4

Deleted: e

Deleted: e

Deleted: h

Deleted: e

Formatted: Font: (Default) +Body (Times New Roman), 11 pt, Not Italic, Font colour: Red

1 region) (Fig. 5i), which is consistent with the increased ENSO amplitude under Asian aerosol forcing
 2 indicated above. Besides, anomalous SST warming over the eastern Pacific can further strengthen
 3 westerly wind anomalies over central Pacific (Zebiak & Cane, 1987). Previous studies have found a
 4 link between warmer SST in the eastern than in the western equatorial Pacific with an increase in ENSO
 5 amplitude (Zheng et al., 2016; Ying et al., 2019; Hayashi et al., 2020).

Deleted: f

Formatted: Font: (Default) +Body (Times New Roman), 11 pt, Not Italic

6
 7 Given the above marked changes over the equatorial Pacific mean state in autumn and winter, we further
 8 explore the response of the tropical Pacific mean state to Asian aerosols in these two seasons. In autumn
 9 (SON, September-October-November), there are a zonally wider anticyclone, cooling and negative
 10 precipitation anomalies stretching from Asia to the whole North Pacific (Figs. 6a-c) compared to those
 11 in winter (Figs. 6d-f). As in Figure 5h, Figure 6b and 6e show surface air temperature (SST over the
 12 ocean) difference minus domain mean, on which warming SST anomalies represent less cooling. These
 13 differences between SON and DJF are related to the climatological pattern in SON when the Siberian
 14 High is close to the broad North Pacific subtropical high and the Aleutian Low is weak (Fig. S6). The
 15 associated midlatitude westerlies (Fig. S6a) transport aerosols downstream, extending the region of
 16 aerosol-induced negative effective radiative forcing (ERF) anomalies to the northwestern Pacific (Fig.
 17 S7a-c), leading to the zonally wider cooling. The cooling and associated anticyclonic anomalies trigger
 18 cross-equatorial wind anomalies from the Northern Hemisphere to the Southern Hemisphere, which
 19 shift the ITCZ southward (Figs. 6a-c), as indicated by previous studies on the interhemispheric
 20 difference in aerosol emissions (Navarro et al., 2017; Voigt et al., 2017; Wilcox et al., 2019). Deflected
 21 by the Coriolis force, the cross-equatorial wind anomalies present a westerly anomaly near the equator
 22 mainly over the central Pacific (purple box in Fig. 6a), which can weaken the easterly trade winds,
 23 generating warm SST anomalies over the eastern Pacific (green box in Fig. 6b) and excess rainfall (Fig.
 24 6c). From SON to DJF, the climatological Siberian High strengthens, and the Aleutian Low deepens
 25 with a southward shift in the coupled baseline simulation (Figs. S6). The associated northwesterlies
 26 along the eastern flank of the Siberian High and northeasterlies along the coast of East Asia confine the
 27 negative ERF anomalies primarily south of 30°N (Fig. S7d-f). Therefore, the Asian aerosol-induced
 28 cooling and associated anticyclone are more concentrated over the Maritime Continent and the north-
 29 western Pacific (Fig. 6d), altering the SST gradient anomaly from north-south (Fig. 6b) to northwest-
 30 southeast (Fig. 6e). This SST anomaly pattern leads to the southward shift of anomalous westerly winds
 31 over the central-eastern Pacific, as well as warm SST and positive precipitation anomalies over eastern
 32 Pacific (Figs. 6d-f). These anomalies are conducive to increasing the ENSO amplitude as explained
 33 below.

Deleted: occur

Deleted: e

Deleted: c

Deleted: 4

Deleted: , that lead to the zonally wider cooling by Asian aerosols.

Formatted: Font: 11 pt, Not Italic

Formatted: Font colour: Text 1

Deleted: 4a, S2a

Formatted: Font colour: Text 1

Deleted: .

Formatted: Font: (Default) +Body (Times New Roman), 11 pt, Not Italic

Deleted:

Formatted: Font: 11 pt, Not Bold

Formatted: Font colour: Text 1

advection of anomalous subsurface temperatures by the mean upwelling (the thermocline feedback) (e.g. Timmermann et al., 2018; Ying et al., 2019; Peng et al., 2024). The two latter feedbacks are related to the ocean dynamic responses to zonal wind forcing that cause in-phase variations of eastern Pacific SST anomalies (Jin and An, 1999; Kim et al., 2014). A diagnostic quantity that includes both these two feedback processes is the zonal wind forcing of SST anomalies, which was found to be useful for studying ENSO-amplitude changes under global warming (Rashid et al., 2016). To further quantify the changes in the strength of the ocean-atmosphere coupling that modulate the ENSO amplitude, we focus on two main processes, the atmospheric Bjerknes feedback and the zonal wind forcing, which are related to the formation of the westerly anomalies over the central Pacific and warm SST anomalies over the eastern Pacific indicated above. Figure 7 shows the lag-regression coefficients between the SST anomalies averaged over the Niño3 region (green box in Fig. 6b) (the Niño3 SST index) and near-surface zonal winds (U1000) anomalies averaged over the Niño4 region (purple box in Fig. 6a) (the Niño4 U1000 index) to represent the atmospheric Bjerknes feedback and the zonal wind forcing. In each panel, regression coefficients between two variables at different lags are plotted for observations (black curve) and the coupled baseline (blue curve and shading) and SULx10Asia (red curve and shading) simulations. The left panel shows the Niño4 U1000 anomalies response to the Niño3 SST index (i.e. the atmospheric Bjerknes feedback). As in most CMIP models (e.g. Bellenger et al., 2014, Rashid et al., 2016), the simulated atmospheric Bjerknes feedback is weaker than in observations (Fig. 7a). The strength of the feedback for lags between -5 and 5 months almost doesn't change in the coupled SULx10Asia simulation relative to the baseline (Fig. 7a). The right panel shows the Niño3 SST anomalies response to the Niño4 U1000 index (i.e. the zonal wind forcing). In this case, the simulated SST responses are somewhat stronger than the observed responses, and the maximum responses are found at small positive lags (e.g. when U1000 leads SST by 1-2 months) (Rashid et al., 2022). The zonal wind forcing, defined as the maximum of the regression coefficients (lag=1), strengthens from the baseline ($0.51^{\circ}\text{C m}^{-1} \text{ s}$) to the SULx10Asia experiment ($0.55^{\circ}\text{C m}^{-1} \text{ s}$) by 8%. Therefore, the zonal wind forcing plays a more important role than the atmospheric Bjerknes feedback in increasing the ENSO amplitude under Asian aerosol forcing. In summary, the Asian aerosol-induced cooling weakens the easterly trade winds over the central Pacific, which reduces the east-west equatorial Pacific SST gradient through the zonal wind forcing, leading to increased ENSO amplitude.

Deleted:

Formatted: Font colour: Auto

6 Summary and conclusions

This study investigates the response of the ENSO-EAWM link and related interannual variability of the EAWM to Asian sulfate aerosols, including the induced changes in the ENSO-related ocean-atmosphere feedbacks, using a set of experiments carried out as part of the PDRMIP initiative. Accounting for two-way atmosphere-ocean coupling, the El Niño-related warm SST anomalies intensify over the eastern equatorial Pacific, associated with an enhancement of the anomalous

1 anticyclone anomaly over the western tropical Pacific and corresponding stronger southerlies along the
2 Asian coast from the South China Sea. This enhanced ENSO signal and its induced PEA pattern
3 contribute to explaining the increased interannual variability of the EAWM (+18%). When the ocean is
4 not allowed to respond, the interannual variability of the EAWM varies in the opposite direction (-
5 19%), which further reflects the importance of ENSO-related atmosphere-ocean coupled processes for
6 explaining the increased variability. The PEA-like EAWM-related circulation and precipitation
7 anomalies also hint at a link between increased interannual variability of the EAWM and changes in
8 ENSO in response to Asian aerosols. The increased ENSO signal can be further linked to changes in
9 the ENSO amplitude. The multi-model mean ENSO amplitude increases by 17% with increased sulfate
10 aerosols, with more El Niño and La Niña years in all the PDRMIP models used in this study. Changes
11 in the Niño3.4 index are significantly correlated with changes in the EAWM index.

12
13 In coupled simulations, the aerosol-induced broad cooling alters the mean state over the tropical and
14 equatorial Pacific, generating westerly anomalies over the central Pacific (peak in autumn) and warm
15 SST anomalies over the eastern Pacific from autumn to winter, which are key factors in increasing
16 ENSO amplitude. Using a diagnostic analysis, the contribution of two main processes, the [atmospheric](#)
17 Bjerknes feedback and zonal wind forcing is estimated. The zonal wind forcing is identified to
18 strengthen from the baseline experiment to the SUL×10Asia experiment by 8%, while the strength of
19 the [atmospheric](#) Bjerknes feedback almost doesn't change. Therefore, the aerosol-induced cooling
20 weakens the easterly trade winds over the central Pacific, which reduce the east-west equatorial Pacific
21 SST gradient through the zonal wind forcing, causing the increased amplitude of ENSO and the EAWM.
22 In summary, the findings of this study provide a better understanding of the change to the year-to-year
23 variability of the EAWM in response to aerosol forcing. This is critical to reducing uncertainties in
24 future projections of variability of regional extremes, such as cold surges and flooding, which can cause
25 large social and economic impacts on densely populated East Asia.

26
27 We acknowledge some limitations and potential extensions of this study. Only a limited number of
28 models is available as part of PDRMIP, as some others do not parameterise aerosol-cloud interactions
29 which are critical to realise the total aerosol response across Asia (e.g. Dong et al., 2016; Liu et al.,
30 2024). Also, some models prescribed concentrations, rather than emissions, perturbations, the
31 implications of which are difficult to ascertain given the limited model sample. Including more models
32 and making use of coordinated perturbed aerosol experiments to Asian aerosols, such as those planned
33 as part of RAMIP (Wilcox et al., 2023) would further increase the robustness of our study. This would
34 allow to better characterise the individual model responses as a function of the underlying bias (e.g.,
35 Liu et al., 2024). It would be interesting to extend this analysis to future projections for the 21st century,
36 for example using CMIP6 models or large ensembles, and examine the externally-forced changes
37 accounting also for the role of internal climate variability. It would also be interesting to examine the

1 extent to which the ENSO-EAWM link varies across the various future aerosol pathways, which are
2 uncertain and display very different, but equally plausible, patterns over Asia (Persad et al., 2022; Wang
3 et al., 2023). Finally, we only considered the role of Asian aerosol changes. A more comprehensive
4 analysis, should similar experiments be available, could also consider aerosols from [South and East](#)
5 [Asia separately as well as from](#) other geographical regions, such as Europe and North America, which
6 can also affect the Pacific and, via atmospheric teleconnections, East Asia (e.g. Dong et al., 2016; Liu
7 et al., 2019).

8 **Code availability.** The python code generated in this study is available upon request (contact author).
9

10
11 **Data availability.** The CRU land temperature dataset is obtained from
12 https://crudata.uea.ac.uk/cru/data/hrg/cru_ts_4.07, while the HadISST sea surface temperature dataset
13 can be found at <https://www.metoffice.gov.uk/hadobs/hadisst/>. The ERA5 reanalysis is provided by the
14 European Centre for Medium-Range Weather Forecasts at
15 <https://www.ecmwf.int/en/forecasts/dataset/ecmwf-reanalysis-v5>. The PDRMIP data can be accessed
16 through the World Data Center for Climate (WDCC) data server at
17 https://doi.org/10.26050/WDCC/PDRMIP_2012-2021.
18

19 **Author contribution.** ZJ and MAB designed the study and discussed the results. ZJ carried out the
20 analysis and drafted the manuscript. All authors edited the paper.

21
22 **Competing interests.** The authors have no competing interests to declare.
23

24 **Acknowledgements.** ZJ thanks the Startup Foundation for Introducing Talent of Nanjing University of
25 Information Science and Technology (NUIST) (grant no. 2024r034) and Natural Science Fund for
26 Colleges and Universities in Jiangsu Province (grant no. 24KJB170015). MB acknowledges support
27 from the Natural Environment Research Council (grant no. NE/N006038/1) and the Research Council
28 of Norway (grant no. 324182; CATHY).

29 References

30 Allen, R. J., Evan, A. T., Booth, B. B. B., Allen, R. J., Evan, A. T., and Booth, B. B. B.:
31 Interhemispheric Aerosol Radiative Forcing and Tropical Precipitation Shifts during the Late
32 Twentieth Century, *J. Climate*, 28, 8219–8246, <https://doi.org/10.1175/JCLI-D-15-0148.1>,
33 2015.

Deleted: 1

1 Bellenger, H., Guilyardi, E., Leloup, J., Lengaigne, M., and Vialard, J.: ENSO representation in
2 climate models: from CMIP3 to CMIP5, *Clim. Dyn.*, 42, 1999–2018,
3 <https://doi.org/10.1007/s00382-013-1783-z>, 2014.

4 Bartlett, R.E., Bolasina, M.A., Booth, B.B., Dunstone, N.J., Marengo, F., Messori, G. and Bernie,
5 D.J.: Do differences in future sulfate emission pathways matter for near-term climate? A case
6 study for the Asian monsoon, *Clim. Dyn.*, 50, pp.1863-1880, [https://doi.org/10.1007/s00382-](https://doi.org/10.1007/s00382-017-3726-6)
7 017-3726-6, 2018.

8 Bolasina, M.A., Ming, Y. and Ramaswamy, V., et al.: Contribution of local and remote
9 anthropogenic aerosols to the twentieth century weakening of the South Asian monsoon,
10 *Geophys. Res. Lett.*, 41(2), pp.680-687, <https://doi.org/10.1002/2013GL058183>, 2014.

11 Beobide-Arsuaga, G., Bayr, T., Reintges, A., & Latif, M.: Uncertainty of ENSO-amplitude
12 projections in CMIP5 and CMIP6 models, *Clim. Dyn.* 56, pp.3875-3888,
13 <https://doi.org/10.1007/s00382-021-05673-4>, 2021.

14 Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P., Kerminen, V.-M.,
15 Kondo, Y., Liao, H., Lohmann, U., Rasch, P., Satheesh, S., Sherwood, S., Stevens, B., and
16 Zhang, X.: Clouds and Aerosols, in: *Climate Change 2013: The Physical Science Basis*,
17 Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental
18 Panel on Climate Change, chap. Clouds and, Cambridge University Press, Cambridge, United
19 Kingdom and New York, NY, USA, 2013.

20 Cai, W., Borlace, S., Lengaigne, M., Van Rensch, P., Collins, M., Vecchi, G., Timmermann, A.,
21 Santoso, A., McPhaden, M.J., Wu, L. and England, M.H.: Increasing frequency of extreme El
22 Niño events due to greenhouse warming, *Nature climate change*, 4(2), pp.111-116,
23 <https://doi.org/10.1038/nclimate2100>, 2014.

24 Cai, W., Santoso, A., Collins, M., Dewitte, B., Karamperidou, C., Kug, J.S., Lengaigne, M.,
25 McPhaden, M.J., Stuecker, M.F., Taschetto, A.S. and Timmermann, A.: Changing El Niño–
26 Southern oscillation in a warming climate, *Nature Reviews Earth & Environment*, 2(9),
27 pp.628-644, <https://doi.org/10.1038/s43017-021-00199-z>, 2021.

28 Chang, C.P., Wang, Z. and Hendon, H.: *The Asian winter monsoon* (The Asian Monsoon (Berlin:
29 Springer Praxis Books)), pp.89–127, 2006.

30 Chen, W., Yang, S. and Huang, R.H.: Relationship between stationary planetary wave activity and the
31 East Asian winter monsoon, *Journal of Geophysical Research: Atmospheres*, 110(D14),
32 <https://doi.org/10.1029/2004JD005669>, 2005.

33 [Chen, Z., R. Wu, and W. Chen, 2015: Effects of northern and southern components of the East Asian](#)
34 [winter monsoon on SST changes in the western North Pacific. *J. Geophys. Res.*, 120\(9\),](#)
35 [3888-3905.](#)

1 Chen, W., Wang, L., Feng, J., Wen, Z., Ma, T., Yang, X., & Wang, C.: Recent progress in studies of
2 the variabilities and mechanisms of the East Asian monsoon in a changing climate, *Advances*
3 in Atmospheric Sciences, 36(9), 887–901, <https://doi.org/10.1007/s00376-019-8230-y>, 2019.

4 Cheng, J., Su, J., Cui, T., Li, X., Dong, X., Sun, F., Yang, Y., Tong, D., Zheng, Y., Li, Y. and Li, J.:
5 Dominant role of emission reduction in PM_{2.5} air quality improvement in Beijing during
6 2013–2017: a model-based decomposition analysis, *Atmospheric Chemistry and*
7 *Physics*, 19(9), 6125–6146, <https://doi.org/10.5194/acp-19-6125-2019>, 2019.

8 Collins, M., An, S.I., Cai, W., Ganachaud, A., Guilyardi, E., Jin, F.F., Jochum, M., Lengaigne, M.,
9 Power, S., Timmermann, A. and Vecchi, G.: The impact of global warming on the tropical
10 Pacific Ocean and El Niño, *Nature Geoscience*, 3(6), pp.391–397,
11 <https://doi.org/10.1038/ngeo868>, 2010.

12 Copernicus Climate Change Service (C3S): ERA5: fifth genera- tion of ECMWF atmospheric
13 reanalyses of the global climate, Copernicus Climate Change Service Climate Data Store
14 (CDS) [data set], 15(2), 2020, <https://cds.climate.copernicus.eu/cdsapp#!/home> (last access:
15 16 May 2022), 2017.

16 Dong, B., Sutton, R. T., Highwood, E. J., and Wilcox, L. J.: Preferred response of the East Asian
17 summer monsoon to local and non-local anthropogenic sulphur dioxide emissions, *Clim.*
18 *Dyn.*, <https://doi.org/10.1007/s00382-015-6782-6>, 2016.

19 Dow, W. J., Maycock, A. C., Lofverstrom, M., & Smith, C. J.: The effect of anthropogenic aerosols
20 on the Aleutian low, *J. Climate*, 34(5), 1725–1741, <https://doi.org/10.1175/JCLI-D-20-0423.1>,
21 2021.

22 Fahrenbach, N.L., Bollasina, M.A., Samset, B.H., Cowan, T. and Ekman, A.M.: Asian Anthropogenic
23 Aerosol Forcing Played a Key Role in the Multidecadal Increase in Australian Summer
24 Monsoon Rainfall, *J. Climate*, 37(3), pp.895–911, <https://doi.org/10.1175/JCLI-D-23-0313.1>,
25 2024.

26 Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J. L., Frame, D., Lunt, D. J., Mauritsen,
27 T., Palmer, M. D., Watanabe, M., Wild, M., and Zhang, H.: The Earth’s Energy Budget, Cli-
28 mate Feedbacks, and Climate Sensitivity, in: *Climate Change 2021: The Physical Science*
29 *Basis*, Contribution of Working Group I to the Sixth Assessment Report of the
30 Intergovernmental Panel on Climate Change, Cambridge University Press, 2021.

31 Gao, J., Wang, K., Wang, Y., Liu, S., Zhu, C., Hao, J., Liu, H., Hua, S. and Tian, H.: Temporal-spatial
32 characteristics and source apportionment of PM_{2.5} as well as its associated chemical species
33 in the Beijing-Tianjin-Hebei region of China, *Environmental pollution*, 233, pp.714–724,
34 <https://doi.org/10.1016/j.envpol.2017.10.123>, 2018.

35 Gong, H., L. Wang, W. Chen, R. Wu, K. Wei, and X. Cui: The Climatology and Interannual
36 Variability of the East Asian Winter Monsoon in CMIP5 Models. *J. Climate*, 27, 1659–1678,
37 <https://doi.org/10.1175/JCLI-D-13-00039.1>, 2014.

1 Gong, H., Wang, L., Chen, W., Nath, D., Huang, G. and Tao, W.: Diverse influences of ENSO on the
2 East Asian–Western Pacific winter climate tied to different ENSO properties in CMIP5
3 models *J. Clim.* 28 2187–202, <https://doi.org/10.1175/JCLI-D-14-00405.1>, 2015.

4 Harris, I., Osborn, T. J., Jones, P. and Lister, D.: Version 4 of the CRU TS monthly high-resolution
5 gridded multivariate climate dataset *Sci. Data* 7, 2020.

6 Hayashi, M., Jin, F. F. & Stuecker, M. F. Dynamics for El Niño–La Niña asymmetry constrain
7 equatorial-Pacific warming pattern, *Nat. Commun.*, 11, 4230, [https://doi.org/10.1038/s41467-](https://doi.org/10.1038/s41467-020-17983-y)
8 020-17983-y, 2020.

9 He, S., Wang, H., & Liu, J.: Changes in the Relationship between ENSO and Asia–Pacific
10 Midlatitude Winter Atmospheric Circulation. 26(10), 3377–3393.
11 <https://dx.doi.org/10.1175/JCLI-D-12-00355.1>, 2013.

12 Hersbach, H., Bell, B., Berrisford, P., Biavati, G., Horányi, A., Muñoz Sabater, J., Nicolas, J., Peubey,
13 C., Radu, R., Rozum, I., Schepers, D., Simmons, A., Soci, C., Dee, D., Thépaut, J-N: ERA5
14 monthly averaged data on pressure levels from 1940 to present. Copernicus Climate Change
15 Service (C3S) Climate Data Store (CDS), DOI: 10.24381/cds.6860a573 (Accessed on DD-
16 MMM-YYYY), 2023.

17 Hu, C., Yang, S., & Wu, Q.: An optimal index for measuring the effect of East Asian winter monsoon
18 on China winter temperature, *Climate Dynamics*, 45(9–10), 2571–2589,
19 <https://doi.org/10.1007/s00382-015-2493-5>, 2015.

20 Huang, R., Chen, J., Wang, L., & Lin, Z.: Characteristics, processes, and causes of the spatio-
21 temporal variabilities of the East Asian monsoon system, *Advances in Atmospheric Sciences*,
22 29(5), 910–942, <https://doi.org/10.1007/s00376-012-2015-x>, 2012.

23 Huang, P., & Xie, S. P.: Mechanisms of change in ENSO-induced tropical Pacific rainfall variability
24 in a warming climate, *Nature Geoscience*, 8(12), pp.922–926,
25 <https://doi.org/10.1038/ngeo2571>, 2015.

26 Huang, B. et al: Extended reconstructed sea surface temperature, version 5 (ERSSTv5): upgrades,
27 validations, and intercomparisons. *J. Clim.* 30, 8179–8205, [https://doi.org/10.1175/JCLI-D-](https://doi.org/10.1175/JCLI-D-16-0836.1)
28 16-0836.1, 2017.

29 Hwang, Y.-T., Frierson, D. M. W., and Kang, S. M.: Anthropogenic sulfate aerosol and the southward
30 shift of tropical precipitation in the late 20th century, *Geophys. Res. Lett.*, 40, 2845–2850,
31 <https://doi.org/10.1002/grl.50502>, 2013.

32 Ji, L., Sun, S., Arpe, K. and Bengtsson, L.: Model study on the interannual variability of Asian winter
33 monsoon and its influence, *Adv. Atmos. Sci.*, 14, 1–22, [https://doi.org/10.1007/s00376-997-](https://doi.org/10.1007/s00376-997-0039-4)
34 0039-4, 1997.

35 Jia, Z., Bollasina, M.A., Li, C., Doherty, R. and Wild, O.: Changes in the relationship between ENSO
36 and the East Asian winter monsoon under global warming, *Environ. Res. Lett.*, 15(12),
37 p.124056, <https://doi.org/10.1088/1748-9326/abca63>, 2020.

1 Jin, F. F.: An equatorial ocean recharge paradigm for ENSO. Part II: A stripped-down coupled
2 model. *Journal of the Atmospheric Sciences*, 54(7), 830-847, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0469(1997)054<0830:AEORPF>2.0.CO;2)
3 0469(1997)054<0830:AEORPF>2.0.CO;2, 1997.

4 Jin, F., and An, S.: Thermocline and zonal advective feedbacks within the equatorial ocean recharge
5 oscillator model for ENSO, *Geophys. Res. Lett.* 26, 2989–2992,
6 <https://doi.org/10.1029/1999GL002297>, 1999.

7 Jiang, Y., Yang, X.Q., Liu, X., Yang, D., Sun, X., Wang, M., Ding, A., Wang, T. and Fu, C.:
8 Anthropogenic aerosol effects on East Asian winter monsoon: The role of black carbon-
9 induced Tibetan Plateau warming, *Journal of Geophysical Research: Atmospheres*, 122(11),
10 pp.5883-5902, <https://doi.org/10.1002/2016JD026237>, 2017.

11 Jiang, W., Gong, H., Huang, P., Wang, L., Huang, G. and Hu, L.: Biases and improvements of the
12 ENSO-East Asian winter monsoon teleconnection in CMIP5 and CMIP6 models, *Climate*
13 *Dynamics*, 59(7), pp.2467-2480, <https://doi.org/10.1007/s00382-022-06220-5>, 2022.

14 Kim, S. T., Cai, W., Jin, F. F., and Yu, J. Y.: ENSO stability in coupled climate models and its
15 association with mean state, *Clim. Dyn.*, 42, 3313–3321, [https://doi.org/10.1007/s00382-013-](https://doi.org/10.1007/s00382-013-1833-6)
16 1833-6, 2014.

17 Kobayashi, S., Ota, Y., Harada, Y., Ebata, A., Moriya, M., Onoda, H., Onogi, K., Kamahori, H.,
18 Kobayashi, C., Endo, H. and Miyaoka, K.: The JRA-55 reanalysis: General specifications and
19 basic characteristics, *Journal of the Meteorological Society of Japan, Ser. II*, 93(1), pp.5-48,
20 <https://doi.org/10.2151/jmsj.2015-001>, 2015.

21 Lewinschal, A., Ekman, A. M. L., and Körnich, H.: The role of precipitation in aerosol-induced
22 changes in northern hemisphere wintertime stationary waves, *Clim. Dynam.*, 41, 647–661,
23 <https://doi.org/10.1007/s00382-012-1622-7>, 2013.

24 Li, F., & Wang, H.: Autumn sea ice cover, winter Northern Hemisphere annular mode, and winter
25 precipitation in Eurasia, *Journal of Climate*, 26(11), 3968-3981, [https://doi.org/10.1175/JCLI-](https://doi.org/10.1175/JCLI-D-12-00380.1)
26 D-12-00380.1, 2012.

27 Li, J., Carlson, B.E., Yung, Y.L., Lv, D., Hansen, J., Penner, J.E., Liao, H., Ramaswamy, V., Kahn,
28 R.A., Zhang, P. and Dubovik, O.: Scattering and absorbing aerosols in the climate
29 system, *Nature Reviews Earth & Environment*, 3(6), pp.363-379,
30 <https://doi.org/10.1038/s43017-022-00296-7>, 2022.

31 Liao, H., Chang, W.Y., Yang, Y.: Climatic effects of air pollutants over China: A review, *Advances*
32 *in Atmospheric Sciences*, 32(1), pp.115-139, <https://doi.org/10.1007/s00376-014-0013-x>,
33 2015.

34 Lian, T., Chen, D., Ying, J., Huang, P. & Tang, Y.: Tropical Pacific trends under global warming: El
35 Niño-like or La Niña-like? *Natl Sci. Rev.*, 5, 810–812, <https://doi.org/10.1093/nsr/nwy134>,
36 2018.

- 1 Lian, T., Chen, D.: The essential role of early-spring westerly wind burst in generating the centennial
- 2 extreme 1997/98 El Niño, *J. Clim.*, 1:1–38, <https://doi.org/10.1175/JCLI-D-21-0010.1>, 2021.
- 3 Liu, Y., Sun, J. R., Yang, B.: The effects of black carbon and sulphate aerosols in China regions on
- 4 East Asia monsoons, *Tellus B: Chemical and Physical Meteorology*, 61(4): 642-656,
- 5 <https://doi.org/10.1111/j.1600-0889.2009.00427.x>, 2009.
- 6 Liu, L., Shawki, D., Voulgarakis, A., Kasoar, M., Samset, B.H., Myhre, G., Forster, P.M., Hodnebrog,
- 7 Ø., Sillmann, J., Aalbergjø, S.G. and Boucher, O.: A PDRMIP multimodel study on the
- 8 impacts of regional aerosol forcings on global and regional precipitation, *Journal of*
- 9 *climate*, 31(11), pp.4429-4447, <https://doi.org/10.1175/JCLI-D-17-0439.1>, 2018.
- 10 Liu, Z., Ming, Y., Wang, L., Bollasina, M., Luo, M., Lau, N.C. and Yim, S.H.L.: A model
- 11 investigation of aerosol-induced changes in the east Asian winter monsoon, *Geophysical*
- 12 *research letters*, 46(16), pp.10186-10195, <https://doi.org/10.1029/2019GL084228>, 2019.
- 13 Liu, Z., Bollasina, M.A. and Wilcox, L.J., 2024. Impact of Asian aerosols on the summer monsoon
- 14 strongly modulated by regional precipitation biases. *Atmospheric Chemistry and*
- 15 *Physics*, 24(12), pp.7227-7252.
- 16 McPhaden, M. J.: Genesis and evolution of the 1997–98 El Nino. *Science* 283:950–954.
- 17 <https://doi.org/10.1126/science.283.5404.950>, 1999.
- 18 Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestad, J., Huang, J., Koch, D., Lamarque, J.-
- 19 F., Lee, D., Mendoza, B., Nakajima, T., Robock, A., Stephens, G., Takemura, T., and Zhang,
- 20 H.: Anthropogenic and Natural Radiative Forcing, in: *Climate Change 2013, The Physical*
- 21 *Science Basis, Contribution of Working Group I to the Fifth Assessment Report of the Inter-*
- 22 *governmental Panel on Climate Change*, edited by: Stocker, T. F., Qin, D., Plattner, G.-K.,
- 23 Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M.,
- 24 Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 25 Myhre, G., Forster, P.M., Samset, B.H., Hodnebrog, Ø., Sillmann, J., Aalbergjø, S.G., Andrews, T.,
- 26 Boucher, O., Faluvegi, G., Fläschner, D. and Iversen, T.: PDRMIP: A precipitation driver and
- 27 response model intercomparison project—Protocol and preliminary results, *Bulletin of the*
- 28 *American Meteorological Society*, 98(6), pp.1185-1198, [https://doi.org/10.1175/BAMS-D-](https://doi.org/10.1175/BAMS-D-16-0019.1)
- 29 [16-0019.1](https://doi.org/10.1175/BAMS-D-16-0019.1), 2017.
- 30 Navarro, J. C. A., Ekman, A. M. L., Pausata, F. S. R., Lewinschal, A., Varma, V., Seland, Ø., Gauss,
- 31 M., Iversen, T., Kirkevåg, A., Riipinen, I., and Hansson, H. C.: Future Response of
- 32 Temperature and Precipitation to Reduced Aerosol Emissions as Compared with Increased
- 33 Greenhouse Gas Concentrations, *J. Climate*, 30, 939–954, [https://doi.org/10.1175/JCLI-D-16-](https://doi.org/10.1175/JCLI-D-16-0466.1)
- 34 [0466.1](https://doi.org/10.1175/JCLI-D-16-0466.1), 2017.
- 35 Peng, Q., Xie, SP. & Deser, C. Collapsed upwelling projected to weaken ENSO under sustained
- 36 warming beyond the twenty-first century. *Nat. Clim. Chang.* 14, 815–822 (2024).
- 37 <https://doi.org/10.1038/s41558-024-02061-8>.

Deleted: Liu, Z., Bollasina, M., and Wilcox, L.: Impact of Asian aerosols on the summer monsoon strongly modulated by regional precipitation biases, *EGU sphere* [preprint], <https://doi.org/10.5194/egusphere-2023-3136>, 2024.

Formatted: Font: (Default) Times New Roman, 11 pt, Font colour: Text 1

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font colour: Text 1

Formatted: Font: (Default) Times New Roman, 11 pt, Font colour: Text 1

Formatted: Font: (Default) Times New Roman, 11 pt, Not Italic, Font colour: Text 1

Formatted: Font: (Default) Times New Roman, 11 pt, Font colour: Text 1

1 Persad, G.G., Samset, B.H. and Wilcox, L.J.: Aerosols must be included in climate risk
2 assessments. *Nature*, 611(7937), pp.662-664, <https://doi.org/10.1038/d41586-022-03763-9>,
3 2022.

4 Persad, G. G.: The dependence of aerosols' global and local precipitation impacts on the emitting
5 region, *Atmos. Chem. Phys.*, 23, 3435–3452, <https://doi.org/10.5194/acp-23-3435-2023>,
6 2023.

7 Power, S., Delage, F., Chung, C., Kociuba, G. and Keay, K.: Robust twenty-first-century projections
8 of El Niño and related precipitation variability, *Nature*, 502(7472), pp.541-545,
9 <https://doi.org/10.1038/nature12580>, 2013.

10 Rayner, N. A., Parker, D. E., Horton, E. B., Folland, C. K., Alexander, L. V., & Rowell, D. P.: Global
11 analyses of sea surface temperature, sea ice, and night marine air temperature since the late
12 nineteenth century, *Journal of Geophysical Research*, 108, 4407.
13 <https://doi.org/10.1029/2002JD002670>, 2003.

14 Ramanathan, V., Chung, C., Kim, D., Bettge, T., Buja, L., Kiehl, J. T., Washington, W. M., Fu, Q.,
15 Sikka, D. R., and Wild, M.: Atmospheric brown clouds: impacts on South Asian climate and
16 hydrological cycle, *P. Natl. Acad. Sci. USA*, 102, 5326–5333,
17 <https://doi.org/10.1073/pnas.0500656102>, 2005.

18 Rashid, H. A., Hirst, A. C., and Marsland, S. J.: An atmospheric mechanism for ENSO amplitude
19 changes under an abrupt quadrupling of CO₂ concentration in CMIP5 models, *Geophys. Res.*
20 *Lett.*, 43, 1687–1694, <https://doi.org/10.1002/2015GL066768>, 2016.

21 Rashid, H. A.: Forced changes in El Niño–Southern Oscillation due to global warming and the
22 associated uncertainties in ACCESS-ESM1.5 large ensembles, *Front. Clim.*, 4:954449,
23 <https://doi.org/10.3389/fclim.2022.954449>, 2022.

24 [Richardson, T. B., Forster, P. M., Smith, C. J., Maycock, A. C., Wood, T., Andrews, T., et al. \(2019\).
25 Efficacy of climate forcings in PDRMIP models. *Journal of Geophysical Research:*
26 *Atmospheres*, 124, 12,824–12,844. <https://doi.org/10.1029/2019JD030581>.](#)

27 Rotstayn, L. D., and Lohmann, U.: Tropical Rainfall Trends and the Indirect Aerosol Effect, *J.*
28 *Climate*, 15, 2103–2116, [https://doi.org/10.1175/1520-0442\(2002\)015<2103:TRTATI>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<2103:TRTATI>2.0.CO;2), 2002.

30 Shindell, D. T., Voulgarakis, A., Faluvegi, G., and Milly, G.: Precipitation response to regional
31 radiative forcing, *Atmos. Chem. Phys.*, 12, 6969–6982, [https://doi.org/10.5194/acp-12-6969-](https://doi.org/10.5194/acp-12-6969-2012)
32 2012, 2012.

33 [Smith, S.J., van Aardenne, J., Klimont, Z., Andres, R.J., Volke, A. and Delgado Arias, S., 2011.
34 Anthropogenic sulfur dioxide emissions: 1850–2005. *Atmospheric Chemistry and*
35 *Physics*, 11\(3\), pp.1101-1116.](#)

- 1 [Samset, B. H., and Coauthors, 2016: Fast and slow precipitation responses to individual climate](#)
- 2 [forcers: A PDRMIP multimodel study. *Geophys. Res. Lett.*, 43, 2782–2791, \[https://doi.org/\]\(https://doi.org/10.1002/2016GL068064\)](#)
- 3 [10.1002/2016GL068064.](#)
- 4 Song, F.F., Zhou, T.T., Qian, Y.: Responses of East Asian summer monsoon to natural and
- 5 anthropogenic forcings in the 17 latest CMIP5 models, *Geophysical Research Letters*, 41(2),
- 6 pp.596–603, <https://doi.org/10.1002/2013GL058705>, 2014.
- 7 Timmermann, A., An, S.I., Kug, J.S., Jin, F.F., Cai, W., Capotondi, A., Cobb, K.M., Lengaigne, M.,
- 8 McPhaden, M.J., Stuecker, M.F. and Stein, K.: El Niño–southern oscillation
- 9 complexity, *Nature*, 559(7715), pp.535–545, <https://doi.org/10.1038/s41586-018-0252-6>,
- 10 2018.
- 11 Vecchi, G.A., Soden, B.J., Wittenberg, A.T., Held, I.M., Leetmaa, A. and Harrison, M.J.: Weakening
- 12 of tropical Pacific atmospheric circulation due to anthropogenic forcing, *Nature*, 441(7089),
- 13 pp.73–76, <https://doi.org/10.1038/nature04744>, 2006.
- 14 Voigt, A., Pincus, R., Stevens, B., Bony, S., Boucher, O., Bellouin, N., Lewinschal, A., Medeiros, B.,
- 15 Wang, Z., and Zhang, H.: Fast and slow shifts of the zonal-mean intertropical convergence
- 16 zone in response to an idealized anthropogenic aerosol, *J. Adv. Model. Earth Sy.*, 9, 870–892,
- 17 <https://doi.org/10.1002/2016MS000902>, 2017.
- 18 Wang, B., Wu, R., & Fu, X.: Pacific–East Asian teleconnection: how does ENSO affect East Asian
- 19 climate? *Journal of Climate*, 13(9), 1517–1536, [https://doi.org/10.1175/1520-](https://doi.org/10.1175/1520-0442(2000)013<1517:PEATHD>2.0.CO;2)
- 20 [0442\(2000\)013<1517:PEATHD>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<1517:PEATHD>2.0.CO;2), 2000.
- 21 Wang, F. K.: Confidence interval for the mean of non-normal data, *Qual. Reliab. Eng. Int.*, 17, 257–
- 22 267, <https://doi.org/10.1002/qre.400>, 2001.
- 23 Wang, B., An, S.A.: Mechanism for decadal changes of ENSO behavior: Roles of background wind
- 24 changes, *Clim Dyn*, 18, pp.475–486, <https://doi.org/10.1007/s00382-001-0189-5>, 2002.
- 25 Wang, Z., Wu, C.-P., Chang, J., Liu, J., Li, and T. Zhou: Another look at interannual-to-interdecadal
- 26 variations of the East Asian winter monsoon: The northern and southern temperature modes,
- 27 *J. Climate*, 23, 1495–1512, <https://doi.org/10.1175/2009JCLI3243.1>, 2010.
- 28 Wang, H., He, S., & Liu, J.: Present and future relationship between the East Asian winter monsoon
- 29 and ENSO: Results of CMIP5, *Journal of Geophysical Research: Oceans*, 118(10), 5222–
- 30 5237, <https://doi.org/10.1002/jgrc.20332>, 2013.
- 31 Wang, G., Cai, W., Gan, B., Wu, L., Santoso, A., Lin, X., Chen, Z. and McPhaden, M.J.: Continued
- 32 increase of extreme El Niño frequency long after 1.5 C warming stabilization, *Nature Climate*
- 33 *Change*, 7(8), pp.568–572, <https://doi.org/10.1038/nclimate3351>, 2017.
- 34 [Wang, Z.-Z., and R. Wu, 2021: Individual and combined impacts of ENSO and East Asian winter](#)
- 35 [monsoon on the South China Sea cold tongue intensity. *Climate Dynamics*, 56\(11–12\), 3995–](#)
- 36 [4012.](#)

- 1 Wang, Z., Wu, R., Gong, H., Jia, X., & Dai, P.: What determine the performance of the ENSO-East
- 2 Asian winter monsoon relationship in CMIP6 models? *Journal of Geophysical Research:*
- 3 *Atmospheres*, 127, e2021JD036227, <https://doi.org/10.1029/2021JD036227>, 2022a.
- 4 [Wang, Z.-Z., R. Wu, and Y.-Q. Wang.: Impacts of the East Asian winter monsoon on winter](#)
- 5 [precipitation variability over East Asia-western North Pacific. *Climate Dynamics*, 58\(11-12\),](#)
- 6 [3041-3055, 2022b.](#)
- 7 Wang, P., Yang, Y., Xue, D., Ren, L., Tang, J., Leung, L. R., & Liao, H.: Aerosols overtake
- 8 greenhouse gases causing a warmer climate and more weather extremes toward carbon
- 9 neutrality, *Nature Communications*, 14(1), 7257, [https://doi.org/10.1038/s41467-023-42891-](https://doi.org/10.1038/s41467-023-42891-2)
- 10 [2](#), 2023.
- 11 Westervelt, D. M., Conley, A. J., Fiore, A. M., Lamarque, J.-F., Shindell, D. T., Previdi, M., Mascioli,
- 12 N. R., Faluvegi, G., Correa, G., and Horowitz, L. W.: Connecting regional aerosol emissions
- 13 reductions to local and remote precipitation responses, *Atmos. Chem. Phys.*, 18, 12461–
- 14 12475, <https://doi.org/10.5194/acp-18-12461-2018>, 2018.
- 15 Westervelt, D. M., Mascioli, N. R., Fiore, A. M., Conley, A. J., Lamarque, J.-F., Shindell, D. T.,
- 16 Faluvegi, G., Previdi, M., Correa, G., and Horowitz, L. W.: Local and remote mean and
- 17 extreme temperature response to regional aerosol emissions reductions, *Atmos. Chem. Phys.*,
- 18 20, 3009–3027, <https://doi.org/10.5194/acp-20-3009-2020>, 2020.
- 19 Wilcox, L. J., Dunstone, N., Lewinschal, A., Bollasina, M., Ekman, A. M. L., and Highwood, E. J.:
- 20 Mechanisms for a remote response to Asian anthropogenic aerosol in boreal winter, *Atmos.*
- 21 *Chem. Phys.*, 19, 9081–9095, <https://doi.org/10.5194/acp-19-9081-2019>, 2019.
- 22 Wilcox, L. J., Allen, R. J., Samset, B. H., Bollasina, M. A., Griffiths, P. T., Keeble, J., Lund, M. T.,
- 23 Makkonen, R., Merikanto, J., O'Donnell, D., Paynter, D. J., Persad, G. G., Rumbold, S. T.,
- 24 Takemura, T., Tsigaridis, K., Undorf, S., and Westervelt, D. M.: The Regional Aerosol Model
- 25 Intercomparison Project (RAMIP), *Geosci. Model Dev.*, 16, 4451–4479,
- 26 <https://doi.org/10.5194/gmd-16-4451-2023>, 2023.
- 27 [Wilks, D., 2016. “The stippling shows statistically significant grid points”: How research results are](#)
- 28 [routinely overstated and overinterpreted, and what to do about it. *Bulletin of the American*](#)
- 29 [Meteorological Society](#), 97(12), pp.2263-2273. <https://doi.org/10.1175/BAMS-D-15-00267.1>
- 30 [Wu, R., W. Chen, G. Wang, and K.-M. Hu, 2014: Relative contribution of ENSO and East Asian](#)
- 31 [winter monsoon to the South China Sea SST anomalies during ENSO decaying years. *J.*](#)
- 32 [Geophys. Res., 119\(19\), 5046-5064.](#)
- 33 Xuan, Z., Zhang, W., Jiang, F., Stuecker, M.F. and Jin, F.F.: Seasonal-varying characteristics of
- 34 tropical Pacific westerly wind bursts during El Niño due to annual cycle modulation, *Climate*
- 35 *Dynamics*, 62(1), pp.299-314, <https://doi.org/10.1007/s00382-023-06907-3>, 2024.

- 1 Yan, Z., Wu, B., Li, T., Collins, M., Clark, R., Zhou, T., Murphy, J. and Tan, G.: Eastward shift and
2 extension of ENSO-induced tropical precipitation anomalies under global warming, science
3 advances, 6(2), p.eaax4177, <https://doi.org/10.1126/sciadv.aax4177>, 2020
- 4 Yang, S., Lau, K.-M., & Kim, K.-M.: Variations of the East Asian jet stream and Asian–Pacific–
5 American winter climate anomalies, Journal of Climate, 15(3), 306–325,
6 [https://doi.org/10.1175/1520-0442\(2002\)015<0306:VOTEAJ>2.0.CO;2](https://doi.org/10.1175/1520-0442(2002)015<0306:VOTEAJ>2.0.CO;2), 2002.
- 7 [Yang, Y., Gao, M., Xie, N. and Gao, Z., 2020. Relating anomalous large-scale atmospheric](#)
8 [circulation patterns to temperature and precipitation anomalies in the East Asian monsoon](#)
9 [region. Atmospheric Research, 232, p.104679.](#)
10 <https://doi.org/10.1016/j.atmosres.2019.104679>.
- 11 Ying, J., Huang, P., Lian, T. & Chen, D.: Intermodel uncertainty in the change of ENSO’s amplitude
12 under global warming: role of the response of atmospheric circulation to SST anomalies, J.
13 Clim. 32, 369–383, <https://doi.org/10.1175/JCLI-D-18-0456.1>, 2019.
- 14 [Zebiak, S.E. and Cane, M.A., 1987. A model El Niño–southern oscillation. Monthly Weather](#)
15 [Review, 115\(10\), pp.2262-2278. https://doi.org/10.1175/1520-](#)
16 [0493\(1987\)115<2262:AMENO>2.0.CO;2.](#)
- 17 Zhang, R., Sumi, A. and Kimoto, M.: Impact of El Niño on the East Asian monsoon a diagnostic
18 study of the ’86/87 and ’91/92 events, Journal of the Meteorological Society of Japan. Ser.
19 II, 74(1), pp.49-62, https://doi.org/10.2151/jmsj1965.74.1_49, 1996.
- 20 Zhang, H., Chen, S., Zhong, J., Zhang, S., Zhang, Y., Zhang, X., Li, Z. and Zeng, X.C.: Formation of
21 aqueous-phase sulfate during the haze period in China: Kinetics and atmospheric
22 implications, Atmospheric Environment, 177, pp.93-99,
23 <https://doi.org/10.1016/j.atmosenv.2018.01.017>, 2018.
- 24 Zhao, S. and Suzuki, K.: Differing impacts of black carbon and sulfate aerosols on global
25 precipitation and the ITCZ location via atmosphere and ocean energy perturbations, Journal
26 of Climate, 32(17), pp.5567-5582, <https://doi.org/10.1175/JCLI-D-18-0616.1>, 2019.
- 27 Zheng, X.-T., Xie, S.-P., Lv, L. H. & Zhou, Z. Q.: Intermodel uncertainty in ENSO amplitude change
28 tied to Pacific Ocean warming pattern, J. Clim. 29, 7265–7279, [https://doi.org/10.1175/JCLI-](https://doi.org/10.1175/JCLI-D-16-0039.1)
29 [D-16-0039.1](#), 2016.
- 30 [Zhou, L.-T., and R. Wu, 2010: Respective impacts of the East Asian winter monsoon and ENSO on](#)
31 [winter rainfall in China. J. Geophys. Res., 115, D02107.](#)
- 32 Zhou, B., Gu, L., Ding, Y., Shao, L., Wu, Z., Yang, X., Li, C., Li, Z., Wang, X., Cao, Y. and Zeng,
33 B.: The great 2008 Chinese ice storm: its socioeconomic–ecological impact and sustainability
34 lessons learned, Bulletin of the American meteorological Society, 92(1), pp.47-60,
35 <https://doi.org/10.1175/2010BAMS2857.1>, 2011.

1 Zuo Z, Li M, An N, Xiao D. 2022. Variations of widespread extreme cold and warm days in winter
2 over China and their possible causes. *Science China Earth Sciences*, 65(2): 337–350,
3 <https://doi.org/10.1007/s11430-021-9836-0>.
4
5
6
7
8
9
10
11

1 Figures





Figure 1. DJF regressions of (a)(d)(g), surface air temperature (SAT, SST over the ocean, °C, shading) and 1000 hPa meridional wind (V1000) over the broad East Asia (green contours, values plotted only when larger than 0.1 m s⁻¹ °C⁻¹), (b)(e)(h) sea level pressure (SLP; hPa, shading) and 850 hPa wind (UV850; m s⁻¹, vector), (c)(f)(i) precipitation (Pre, mm d⁻¹) onto the Niño3.4 index from (a-c) observations during 1965-2014, multimodel mean coupled (d-f) baseline and (g-i) SUL×10Asia simulations in PDRMIP. Dotted regions indicate significant correlations at the 95% level from the two-tailed Student's *t* test. Differences in regressions of (j) SAT and V1000 (green contours, values plotted only when larger than 0.05 m s⁻¹ °C⁻¹), (k) SLP and UV850, (l) Pre between coupled SUL×10Asia and baseline simulations. Dotted regions represent differences that remain significant after false discovery rate (FDR) correction of *p*-values from two-tailed Student's *t*-test (Wilks et al., 2016). The definition regions of the EAWM index and the Niño3.4 index are marked by red and black rectangles respectively.

Deleted:



Formatted: Normal, Indent: Left: 0 cm, Hanging: 0.5 cm, Line spacing: single

Deleted: d)

Formatted: Authors, Line spacing: 1.5 lines

Deleted: coupled

Deleted: (d-f)

Deleted: g

Deleted: h

Deleted: i

Formatted: Font: (Default) Times New Roman, 10 pt, Not Bold, Font colour: Text 1, Pattern: Clear

Formatted: Font: 10 pt, Not Bold

Formatted: Font: 10 pt, Font colour: Text 1





Figure 2. Frequency distributions of the EAWM index from observations during DJF 1994-2005 (black curve) and (a) coupled simulations during DJF for years 50-61, (c) fSST simulations during DJF for years 3-14 in PDRMIP with multimodel-means (thick coloured curves) and the associated 95% confidence intervals (coloured shades). The confidence intervals are estimated from different models by using bootstrap resampling (e.g. Wang, 2001). Differences in multimodel mean standard deviations of V1000 (m s^{-1}) between SUL $\times$ 10Asia and baseline experiments from (b) coupled simulations during DJF for years 50-61, (d) fSST simulations during DJF for years 3-14 in PDRMIP. Dotted regions indicate significant differences at the 95% level from the two-tailed F -test. The definition region of the EAWM index is marked by a red rectangle.



Deleted: (a)

Deleted: 65

Deleted: 14

Deleted: and

Deleted: , (b) observations during DJF 1994-2005 (black curve) and fSST simulations

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Formatted: Font: (Default) Times New Roman, 10 pt, Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold





Figure 3. DJF multimodel mean standard deviations of SAT (SST over the ocean, °C) from (a) observations during 1965-2014, (b) coupled baseline simulations, (c) coupled SUL×10Asia simulations, (d) Differences in standard deviations of SAT (SST over the ocean, °C) between coupled SUL×10Asia and baseline simulations. Dotted regions indicate significant differences at the 95% level from the two-tailed *F*-test. (e) Frequency distributions of the Niño3.4 index from coupled simulations in PDRMIP with multimodel-means (thick coloured curves) and the associated 95% confidence intervals (coloured shades). The confidence intervals are estimated from different models by using bootstrap resampling.

Deleted: .

Deleted: c

Deleted: d



Deleted:




Figure 4. Joint distributions of multimodel mean differences in the EAWM index against corresponding differences in the Niño3.4 index between coupled SUL×10Asia and baseline simulations, including the linear fits with 95% confidence intervals.





Figure 5. Multimodel mean longitudinal transect of the monthly climatological (a, d) 1000 hPa zonal wind (U1000, m s^{-1}), (b, e) SST minus zonal mean ($^{\circ}\text{C}$), (c, f) SST standard deviation ($^{\circ}\text{C}$) for the equatorial Pacific (5°S – 5°N) from coupled (a, b, c) baseline and (d, e, f) SULx10Asia simulations; and their changes in (g) U1000, (h) SST, (i) SST standard deviation between coupled SULx10Asia and baseline simulations. Dotted regions in (g)(h) indicate significant changes at the 95% level from the two-tailed Student's t -test; in (i) indicate significant changes at the 95% level from the two-tailed F -test. The definition longitudes of the Niño3 and Niño4 indices are marked by green and purple thick bars respectively along the x axis.






Figure 6. (a-c) SON, (d-f) DJF multimodel mean changes in (a)(d) sea level pressure (SLP; hPa, shading) and 1000 hPa wind (UV1000, vector), (b)(e) surface air temperature (SAT, SST over the ocean) minus domain mean ($^{\circ}\text{C}$), (c)(f) precipitation (Pre, mm d^{-1}) between coupled SULx10Asia and baseline simulations. Dotted regions indicate significant changes at the 95% level from the two-tailed Student's t test. The definition regions of the Niño3 and Niño4 indices are marked by green and purple rectangles in panels a-b respectively.





Figure 7. Multimodel mean lag-regression coefficients of (a) the Niño4 U1000 index onto the Niño3 SST index (indicative of the atmospheric Bjerknes feedback) ($\text{m s}^{-1} \text{ } ^\circ\text{C}^{-1}$), (b) the Niño3 SST index onto the Niño4 U1000 index (indicative of the zonal wind forcing of SST) ($^\circ\text{C m}^{-1} \text{ s}$) from observations (black curve) and coupled simulations in PDRMIP with multimodel-means (thick coloured curves) and the associated 95% confidence intervals (coloured shades). The confidence intervals are estimated from different models by using bootstrap resampling.



Deleted:

Formatted: Justified

Deleted: ¶

Table 1. Models used in this study and their specifications.

Deleted: ¶

Model	Version	Indirect effects included	References
CESM1-CAM5	1.1.2	Sulfate: all indirect effects	Hurrell et al. (2013); Kay et al. (2015)
MIROC-SPRINTARS	5.9.0	Sulfate: all indirect effects	Takemura et al. (2009); Watanabe et al. (2010)
HadGEM3	GA 4.0	Sulfate: all indirect effects	Bellouin et al. (2011); Walters et al. (2014);
NorESM1	NorESM1-M	Sulfate: all indirect effects	Bentsen et al. (2013); Iversen et al. (2013);

2

3

Table 2. Number of El Niño and La Niña years for each model from coupled baseline and SUL×10Asia simulations in PDRMIP.

Years	CESM1-CAM5 (base)	CESM1-CAM5 (sulx10asia)	MIROC-SPRINTARS (base)	MIROC-SPRINTARS (sulx10asia)	HadGEM3 (base)	HadGEM3 (sulx10asia)	NorESM1 (base)	NorESM1 (sulx10asia)
Niño3.4 > 0.5	16	17	8	15	10	11	12	14
Niño3.4 < -0.5	17	22	6	13	9	9	10	14

6

7

8

9

10

11

12

Page 27: [1] Deleted	Microsoft Office User	25/04/2025 17:05:00
Page 27: [2] Deleted	Microsoft Office User	19/04/2025 18:20:00