

Technical Note: Recommendations for Diagnosing Cloud Feedbacks and Rapid Cloud Adjustments Using Cloud Radiative Kernels

Mark D. Zelinka¹, Li-Wei Chao¹, Timothy A. Myers^{2,3}, Yi Qin⁴, and Stephen A. Klein¹

¹Lawrence Livermore National Laboratory, 7000 East Avenue, L-103, Livermore, CA 94550, USA

²Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO, USA

³Physical Science Laboratory, National Oceanic and Atmospheric Administration, Boulder, CO, USA

⁴Pacific Northwest National Laboratory, Richland, WA, USA

Correspondence: Mark D. Zelinka (zelinka1@llnl.gov)

Abstract. The cloud radiative kernel method is a popular approach to quantify cloud feedbacks and rapid cloud adjustments to increased CO₂ concentrations, and to partition contributions from changes in cloud amount, altitude, and optical depth. However, because this method relies on cloud property histograms derived from passive satellite sensors or produced by passive satellite simulators in models, changes in obscuration of lower-level clouds by upper-level clouds can cause apparent low cloud feedbacks and adjustments even in the absence of changes in lower-level cloud properties. Here, we provide a methodology for properly diagnosing the impact of changing obscuration on cloud feedbacks and adjustments and quantify these effects across climate models. Averaged globally and across global climate models, properly accounting for obscuration leads to weaker positive feedbacks from lower-level clouds and stronger positive feedbacks from upper-level clouds while simultaneously removing a mostly artificial anti-correlation between them. Given that the methodology for diagnosing cloud feedbacks and adjustments using cloud radiative kernels has evolved over several papers, and obscuration effects have only occasionally been considered in recent papers, this paper serves to establish recommended best practices and to provide a corresponding code base for community use.

1 Introduction

Uncertainty in Earth's climate sensitivity is primarily caused by cloud feedbacks, which affect the ability of the Earth system to radiatively damp temperature changes (e.g., Bony et al., 2006; Sherwood et al., 2020). At the same time, uncertainty in effective radiative forcing from doubling of CO₂ is driven in large part by rapid cloud adjustments (Smith et al., 2020). These adjustments occur rapidly in response to the altered atmospheric radiative cooling profile when forcing is imposed but before substantial surface warming occurs. Hence the ability of the planet to radiatively damp warming in response to a given forcing, and the magnitude of the forcing itself, are affected in important but unconstrained ways by clouds. Furthermore, radiative forcing and cloud feedback are correlated across climate models with a sign and strength that varies between model generations, affecting the range of climate sensitivities produced (Lutsko et al., 2022; Zelinka et al., 2020).

Accurately diagnosing cloud feedbacks and partitioning them into individual components is essential for understanding which processes are involved, which aspects are robustly simulated across models, and which are subject to substantial inter-

model differences. Doing so also allows for a more rigorous comparison of modeled cloud feedbacks with those observed in nature (Zhou et al., 2013; Myers et al., 2021; Chao et al., 2024; Raghuraman et al., 2024) or with those assessed through expert synthesis of the literature (Zelinka et al., 2022). Cloud radiative kernels (Zelinka et al., 2012a) have proven to be a very useful tool for diagnosing cloud feedbacks because they allow for attribution of the feedback to individual cloud types and gross cloud property changes. Briefly, cloud radiative kernels quantify the sensitivity of top-of-atmosphere (TOA) radiative fluxes to small perturbations in cloud fraction, for clouds segregated by their cloud top pressure (CTP) and visible optical depth (τ). These are constructed via offline radiative transfer calculations applied to model- or reanalysis-based atmospheric temperature and humidity profiles, with and without clouds of specified properties present in the column. The discrete CTP- τ pairs used in constructing cloud radiative kernels match those in the standard International Satellite Cloud Climatology Project (ISCCP) cloud fraction joint histograms (Rossow and Schiffer, 1999), namely, for all 49 combinations of seven CTP bins and seven τ bins.

As demonstrated in Zelinka et al. (2012a), multiplying cloud radiative kernels by the change in cloud fraction histogram per degree of global mean warming and summing over all 49 bins of the resulting histogram yields an estimate of the cloud feedback. This estimate agrees well with independent estimates of the cloud feedback derived via adjusting the change in cloud radiative effect for non-cloud effects and via the approximate partial radiative perturbation technique (Taylor et al., 2007; Zelinka et al., 2022). Because the technique makes use of cloud fraction histograms, it allows one to distinguish cloud feedbacks arising from clouds at different vertical levels and optical depths (Zelinka et al., 2012a) and to compute feedbacks attributable to changes in gross cloud properties holding the others fixed (Zelinka et al., 2012b). Typically this is done by considering the feedback from changes in cloud amount, altitude, and optical depth, in each case holding the other two properties fixed at their control-climate climatological values. A small residual term is also present when performing this decomposition, which was reduced further after slight modifications described in Zelinka et al. (2013). This paper also demonstrated the utility of this technique for diagnosing and decomposing rapid cloud adjustments to CO₂.

In Zelinka et al. (2016), we proposed a slightly more refined breakdown that considers the amount, altitude, and optical depth feedbacks separately for lower- and upper-level clouds. This avoids some misleading results and ambiguities that occur when assessing changes to the full column of clouds collectively, as detailed via several examples in that paper. It also has a number of advantages because it better connects feedbacks to individual governing processes and reveals three net cloud feedback components that are robustly nonzero in climate model warming simulations: positive feedbacks from increasing free tropospheric cloud altitude and decreasing low cloud cover and a negative feedback from increasing low cloud optical depth.

One limitation of relying on cloud data from passive satellite retrievals (or simulators thereof) to estimate the radiative impacts of cloud changes is that such retrievals report only a single cloud type per scene at a vertical level corresponding to the scene's brightness temperature (typically near the top of the highest cloud in the column). Lower-level clouds can therefore be obscured by overlying clouds, and apparent changes in lower-level clouds can arise solely due to changes in overlap. In some recent cloud feedback studies, an additional modification has been made to account for the effect of changing obscuration to better diagnose something closer to "true" lower- and upper-level cloud-induced radiative anomalies (Zelinka et al., 2018; Scott et al., 2020; Myers et al., 2021; Zelinka et al., 2022; Chao et al., 2024). However, a thorough description of this calculation

and a demonstration of its effect on cloud feedbacks and rapid cloud adjustments is lacking. This paper serves to fill this gap. In so doing, we will argue for the necessity of properly accounting for obscuration effects when computing the radiative impact of lower- and upper-level cloud responses using the cloud radiative kernel technique. An additional goal of this paper is to provide a code base for users to easily compute cloud feedbacks and adjustments using cloud radiative kernels and to implement the recommended breakdown. A Jupyter notebook (linked in the Code and data availability Section) is provided for readers wishing to see a demonstration of the recommended calculation, as well as all of the aforementioned predecessors. We will first introduce the models, experiments, variables, and diagnostic techniques used to compute feedbacks and rapid adjustments. Then we will derive the mathematical basis for how obscuration effects are accounted for in computing modified lower- and upper-level cloud feedbacks and adjustments, and provide an illustrated physical interpretation of how these effects operate. Finally, we will quantify the impacts of the modified decomposition on feedbacks and rapid adjustments across climate models and conclude with the major findings.

2 Data and Methods

2.1 Climate Models and Cloud Radiative Kernels

In this work we use output from climate model simulations, though one could alternatively apply similar calculations to cloud feedbacks in response to observed interannual fluctuations or trends in observational cloud fraction histograms (Zhou et al., 2013; Chao et al., 2024; Raghuraman et al., 2024). Our calculations require the following climate model outputs: cloud fraction histograms from the ISCCP simulator, surface air temperature, and clear-sky upwelling and downwelling shortwave (SW) radiation at the surface. The latter two fields are used to map the SW cloud radiative kernel from its native clear-sky surface albedo space to the target model's longitude space (see Zelinka et al. (2012a) for details). The cloud radiative kernels have been generated in Zelinka et al. (2012a) and are available at <https://zenodo.org/records/13686878> (Mark Zelinka, 2024).

To compute cloud feedbacks, we make use of a pair of atmosphere-only experiments, one with prescribed observed SSTs, sea ice concentrations, and radiative constituents (`amip`) and one that is identical but with the SSTs uniformly warmed by 4K at each location (`amip-p4K` in CMIP6, `amip4K` in CMIP5). These experiments are part of the Cloud Feedback Model Intercomparison Project (CFMIP) protocol (Webb et al., 2017) but their roots can be traced back to early experiments to systematically diagnose feedbacks and climate sensitivity across an ensemble of atmospheric models (Cess et al., 1989, 1990). We compute the climatological monthly-resolved cloud fraction histogram climatologies from both the control and perturbed experiments. These are then differenced, multiplied by cloud radiative kernels, and normalized by the change in annual mean global mean surface air temperature between the two experiments to compute cloud feedbacks. 20 distinct models have provided sufficient data to compute cloud feedbacks in response to +4K SST perturbations across CMIP5 and CMIP6 (Table 1). Cloud feedbacks computed using these atmosphere-only uniform SST perturbation experiments have been shown to give a close approximation to those simulated by fully coupled models in response to quadrupled CO₂ (Ringer et al., 2014; Qin et al., 2022).

Table 1. Models used in the calculation of cloud feedbacks. The asterisk indicates that for this model, the amip clisccp data is provided for the r7i1p1 member but the amip4K clisccp data is provided for the r1i1p1 member. For all other models, the variant labels match between the control and +4K amip experiment.

Era	Model	Variant	Experiment
CMIP5	CCSM4	r1i1p1*	amip4K
CMIP5	CNRM-CM5	r1i1p1	amip4K
CMIP5	CanAM4	r1i1p1	amip4K
CMIP5	HadGEM2-A	r1i1p1	amip4K
CMIP5	IPSL-CM5A-LR	r1i1p1	amip4K
CMIP5	IPSL-CM5B-LR	r1i1p1	amip4K
CMIP5	MIROC5	r1i1p1	amip4K
CMIP5	MPI-ESM-LR	r1i1p1	amip4K
CMIP5	MRI-CGCM3	r1i1p1	amip4K
CMIP6	BCC-CSM2-MR	r1i1p1f1	amip-p4K
CMIP6	CESM2	r1i1p1f1	amip-p4K
CMIP6	CNRM-CM6-1	r1i1p1f2	amip-p4K
CMIP6	CanESM5	r1i1p2f1	amip-p4K
CMIP6	E3SM-1-0	r2i1p1f1	amip-p4K
CMIP6	GFDL-CM4	r1i1p1f1	amip-p4K
CMIP6	GISS-E2-1-G	r1i1p1f1	amip-p4K
CMIP6	HadGEM3-GC31-LL	r5i1p1f3	amip-p4K
CMIP6	IPSL-CM6A-LR	r1i1p1f1	amip-p4K
CMIP6	MIROC6	r1i1p1f1	amip-p4K
CMIP6	MRI-ESM2-0	r1i1p1f1	amip-p4K

To compute rapid cloud adjustments, we make use of fixed SST atmosphere-only experiments with atmospheric CO₂ levels quadrupled, for which there are two closely related experiment protocols in CMIP. The first protocol follows the CFMIP amip experiment and uses SST, sea ice, and radiative constituents set to their observed present-day values, except for CO₂ which is quadrupled (referred to as amip-4xCO₂ in CMIP6 and amip4xCO₂ in CMIP5). These experiments are differenced with the amip experiment to compute “amip-type” rapid cloud adjustments. The second follows the Radiative Forcing Model Intercomparison Project (RFMIP) protocol (Pincus et al., 2016) and uses a repeating monthly-resolved climatology of SST and sea ice concentration taken from each model’s pre-industrial control (piControl) experiment as the prescribed boundary condition for each model. The baseline experiment and its quadrupled CO₂ counterpart are known as piClim-control and piClim-4xCO₂, respectively, in CMIP6 and sstClim and sstClim4xCO₂, respectively, in CMIP5. As with the cloud feedback, we compute the climatological monthly-resolved cloud fraction histogram climatologies from both the control and

Table 2. Models used in the calculation of rapid cloud adjustments.

Era	Model	Variant	Experiment
CMIP5	CNRM-CM5	r1i1p1	amip4xCO2
CMIP5	IPSL-CM5B-LR	r1i1p1	amip4xCO2
CMIP5	MPI-ESM-LR	r1i1p1	amip4xCO2
CMIP5	CCSM4	r1i1p1	sstClim4xCO2
CMIP5	CESM1-CAM5	r1i1p1	sstClim4xCO2
CMIP5	CanESM2	r1i1p1	sstClim4xCO2
CMIP5	HadGEM2-A	r1i1p1	sstClim4xCO2
CMIP5	IPSL-CM5A-LR	r1i1p1	sstClim4xCO2
CMIP5	MIROC5	r1i1p1	sstClim4xCO2
CMIP5	MRI-CGCM3	r1i1p1	sstClim4xCO2
CMIP6	BCC-CSM2-MR	r1i1p1f1	amip-4xCO2
CMIP6	CESM2	r1i1p1f1	amip-4xCO2
CMIP6	E3SM-1-0	r2i1p1f1	amip-4xCO2
CMIP6	GISS-E2-1-G	r1i1p1f1	amip-4xCO2
CMIP6	MIROC6	r1i1p1f1	amip-4xCO2
CMIP6	CNRM-CM6-1	r1i1p1f2	piClim-4xCO2
CMIP6	CNRM-ESM2-1	r1i1p1f2	piClim-4xCO2
CMIP6	CanESM5	r1i1p2f1	piClim-4xCO2
CMIP6	GFDL-CM4	r1i1p1f1	piClim-4xCO2
CMIP6	HadGEM3-GC31-LL	r1i1p1f3	piClim-4xCO2
CMIP6	IPSL-CM6A-LR	r1i1p1f1	piClim-4xCO2
CMIP6	MRI-ESM2-0	r1i1p1f1	piClim-4xCO2
CMIP6	UKESM1-0-LL	r1i1p1f4	piClim-4xCO2

105 perturbed experiments. These are then differenced and multiplied by cloud radiative kernels to compute “piClim-type” rapid cloud adjustments. Note that unlike for cloud feedbacks, these anomalies are not normalized by the change in annual mean global mean surface air temperature since they are considered a part of the effective radiative forcing. We take all available models that did either of these experiments and provided the necessary output. For the ten models that provided sufficient output for *both* the amip-type and piClim-type experiments, we use only results from the latter, yielding 23 distinct models (Table 2). Aside from the exceptions noted below, our results are unchanged when instead using the amip-type experiments from these ten models, suggesting that rapid cloud adjustments do not significantly depend on this experimental design choice.

2.2 Accounting for Obscuration

Here we briefly review the approach for accounting for obscuration effects, closely following Scott et al. (2020). Given that a
 110 portion of the low cloud field may be obscured by upper-level clouds, let us define an unobscured low cloud fraction (L_R) as
 L_U) as

$$L_{RU} = L/F, \quad (1)$$

where L is the retrieved low cloud fraction and F is the total upper-level clear-sky fraction, defined as 1 minus the cloud
 fraction summed over all upper-level p and all τ . The precise definition of upper- and lower-level clouds is not prescribed and
 115 can vary depending on the context or needs (e.g., Myers et al., 2021; Ceppi et al., 2024). In practice, we use a cut-off of 680
 hPa to delineate the two cloud types, and refer to the resulting feedbacks or adjustments as being due to “low” or “nonlow”
 clouds. When speaking more generally, we will use the more descriptive “lower-level” and “upper-level” descriptors. L_R L_U
 is the fraction of low clouds within the unobscured portion of a grid box and requires no assumptions about how clouds overlap.
 Note that at a given location and time $L_R = L_R(p, \tau) L_U = L_U(p, \tau)$, whereas F is a scalar. These cloud fractions are defined
 120 only in the case of grid-scale histograms, which are constructed by aggregating over many individual scenes (satellite pixels or
model sub-columns) that are classified as either completely clear or completely covered by a cloud at a single level.

The low cloud fraction can be expressed as the sum of a temporal mean (indicated with an overbar) and a temporal pertur-
 bation (denoted by a prime),

$$L = \bar{L} + L', \quad (2)$$

125 which means that anomalies in low cloud cover can be expressed as

$$L' = L_{RU}F - \bar{L}_U\bar{F}. \quad (3)$$

Next, we further decompose each term on the RHS of (3) into a mean state plus a perturbation, so that

$$L' = (\bar{L}_U + L_{RU}')(\bar{F} + F') - (\bar{L}_U + L_U')(\bar{F} + F') \quad (4)$$

$$= \bar{L}_U \bar{F} + L_{RU}'\bar{F} + \bar{L}_U F' + L_{RU}'F' - \bar{L}_U \bar{F} + L_U'\bar{F} + \bar{L}_U F' + L_U'F' \quad (5)$$

$$130 = L_{RU}'\bar{F} + \bar{L}_U F' + (L_{RU}'F' - L_U'F') \quad (6)$$

The first term on the RHS of (6) is the change in the retrieved low cloud fraction due solely to a change in unobscured low
 cloud fraction. We consider the radiative responses resulting from this component to be closer to a “true” low cloud response
 occurring in regions that are not obscured by upper-level clouds, which receive no contribution from changes in nonlow cloud
 coverage. We Recall that this term represents the low cloud fraction as a joint function of cloud top pressure and optical
 135 depth. Therefore, we can further break this down into amount, altitude, optical depth, and residual components following
Zelinka et al. (2012b, 2013). As will be shown below, the modification to account for obscuration mostly affects the low cloud
 amount component, with tiny impacts on the low cloud altitude and optical depth responses.

The second term on the RHS of (6) is the change in the retrieved low cloud fraction due solely to a change in total upper-level cloud fraction (i.e. obscuration). TOA radiative changes due to this obscuration-induced component of low-cloud response arise entirely from changes in upper-level cloud fraction that reveal or hide lower-level clouds. Hence by definition it is solely an amount component due to changes in nonlow cloud coverage. We therefore absorb this component into the nonlow cloud amount response, as we have previously done in Zelinka et al. (2022). As will be shown below, the modified nonlow cloud amount feedback is typically less negative / more positive than the original nonlow cloud amount component, and vice versa for the rapid adjustment. One can think of this as the nonlow cloud amount component reclaiming a portion of the low cloud radiative response that arises solely due to changes in obscuration.

The third term in parentheses on the RHS of (6) is a term due to covarying changes in the lower- and upper-level cloud fields, and is typically very small.

Written more plainly, the original low cloud response decomposed in (6) can be expressed as

$$\text{low}_{\text{orig}} = (\text{low}_{\text{amt}} + \text{low}_{\text{alt}} + \text{low}_{\text{tau}} + \text{low}_{\text{err}})_{\text{unobsc}} + \Delta\text{obsc} + \text{cov}, \quad (7)$$

where low_{amt} , low_{alt} , low_{tau} , and low_{err} refer to the amount, altitude, optical depth, and residual components of the low cloud response, respectively, which are computed following the approach detailed in Appendix B of Zelinka et al. (2013). The subscript “unobsc” refers to the fact that these components are all occurring in scenes that are not obscured by upper-level clouds, calculated using the first term on the RHS of Eq 6 ($L_U \bar{F}$). Δobsc represents the change in obscuration, diagnosed from the second term on the RHS of Eq 6 ($L_U \bar{F}'$), and cov is the covariance term, diagnosed from the third term on the RHS of Eq 6 ($L_U \bar{F}' - \bar{L}_U \bar{F}'$). In an effort to preserve the total number of components from the original decomposition, and given that the covariance term is typically very small, we combine the covariance and residual terms into a single modified residual term ($\text{low}_{\text{err}}^*$) such that the modified low cloud response can be expressed as:

$$\text{low}_{\text{mod}} = (\text{low}_{\text{amt}} + \text{low}_{\text{alt}} + \text{low}_{\text{tau}} + \text{low}_{\text{err}}^*)_{\text{unobsc}}. \quad (8)$$

Similarly, incorporating Δobsc into the original nonlow cloud amount response yields the modified nonlow cloud response:

$$\text{nonlow}_{\text{mod}} = \text{nonlow}_{\text{amt}}^* + \text{nonlow}_{\text{alt}} + \text{nonlow}_{\text{tau}} + \text{nonlow}_{\text{err}}, \quad (9)$$

where $\text{nonlow}_{\text{amt}}^*$ is the sum of the original nonlow cloud amount response and the change in obscuration term.

To summarize, the total cloud response can be expressed as:

$$\text{total} = \text{nonlow}_{\text{orig}} + \text{low}_{\text{orig}} = \text{nonlow}_{\text{mod}} + \text{low}_{\text{mod}}, \quad (10)$$

and the original and modified cloud responses are related as follows:

$$\text{low}_{\text{mod}} = \text{low}_{\text{orig}} - \Delta\text{obsc}, \quad (11)$$

$$\text{nonlow}_{\text{mod}} = \text{nonlow}_{\text{orig}} + \Delta\text{obsc}. \quad (12)$$

In Appendix A, we derive analogous expressions for the case of three vertical cloud layers (low, middle, and high). Below we will compare the original and modified low and nonlow cloud responses, which primarily illustrates the impact of moving the obscuration term from the low cloud response to the nonlow cloud response.

We first provide a physical interpretation of the impacts of changes in obscuration on the low and high cloud responses diagnosed with cloud radiative kernels applied to cloud property histograms derived from passive satellites (or produced by passive satellite simulators in models). For simplicity, let's consider only SW radiation in these examples. First, consider a scene with an opaque high cloud completely overlapping an opaque low cloud (Figure 1a). Assume that the high cloud fraction decreases with warming but the low cloud fraction remains unchanged. The high cloud decrease will reveal some portion of low cloud that was previously not exposed to space (Figure 1b). This apparent increase in lower-level cloud will constitute a negative radiative response despite the fact that the actual low cloud amount remained unchanged. Hence low cloud amount response diagnosed from the cloud radiative kernel (CRK) technique using the original decomposition will be biased negative relative to the actual, unobscured value, which is close to zero¹ (Figure 1d). Note that this scenario would be indistinguishable from the (extremely rare) scenario in which control-state low cloud cover is truly zero but then increases to a nonzero value in the perturbed climate.

Because cloud radiative kernels are constructed by differencing TOA radiative fluxes calculated assuming completely overcast and clear-sky scenes in a radiative transfer model, they quantify the change in TOA radiation due to changes in cloud cover with the implicit assumption that these changes are relative to a clear-sky atmosphere. The CRK technique will therefore produce a positive radiative impact from high cloud reductions occurring over the typically darker Earth surface (recall that we are only considering SW effects in this discussion). Hence in this hypothetical scenario, the radiative response from decreases in high cloud cover as diagnosed using the original approach will be biased positive relative to the true value (Figure 1d). This is because the removal of a high-cloud above a bright low-cloud leads to a much smaller decrease in SW reflection than if it is above clear-sky. The obscuration adjustment, therefore, is essentially correcting for the kernel overestimate of the high cloud amount response by adding back in the radiative impact of the clouds that are revealed below. This will restore this response to something closer to zero, as will be seen below.

Similarly, consider another scenario with an opaque high cloud partially overlapping an opaque low cloud (Figure 2a). Assume the high cloud increases but the low cloud fraction remains unchanged (Figure 2b). The high cloud increase will hide low cloud, and this apparent decrease in lower-level cloud fraction will constitute a positive radiative response despite the fact that the actual low cloud amount remained unchanged (Figure 2d). Hence the low cloud amount response will be biased positive relative to the true unobscured low cloud amount response. Moreover, due to the aforementioned implicit assumption of hiding/revealing clear skies in the CRK approach, the radiative response from increases in high cloud cover will be biased negative (by roughly the same amount).

¹The ISCCP retrieval algorithm reports a single cloud type for each subcolumn-pixel (in the case of the observations) or sub-column (in the case of the simulator) with an optical depth determined from the column-integrated extinction from all cloud types, including from lower-level clouds beneath upper-level clouds. This means that the reported optical depth of the high cloud in a multi-layer cloud scene must be larger than the reported optical depth of the low cloud revealed if that high cloud goes away. Hence in this scenario, the loss of high-cloud will decrease the column optical depth and thus reduce the amount of reflected SW. The change in reflected SW is therefore not identically zero but will approach zero in the limit of low-cloud optical depth much larger than high-cloud optical depth.




Figure 1. Schematic demonstrating the effects of changing obscuration on cloud feedbacks and adjustments diagnosed by cloud radiative kernels. In this scenario, a high cloud completely overlaps a low cloud in the mean climate, but decreases with climate warming so as to reveal a portion of the (unchanged) low cloud.

In all of these examples, actual low cloud changes were zero simply for the sake of isolating the role of changes in obscuration alone on low and high cloud responses. More typically, there are real low cloud responses operating alongside obscuration effects. Moreover, we only discussed SW effects owing to their simplicity and because they are more relevant. This is because low clouds have a small LW effect, so changes in the obscuration of low clouds by upper-level clouds has a much smaller effect on the LW fluxes reaching the TOA. The LW flux emanating from a low cloud scene is not much different from that emitted from a clear-sky scene. Hence if a high cloud covers up a low cloud, the radiative impact is well captured by the LW CRK. The same cannot be said for the SW.

There are other issues that arise from the use of passive satellite datasets and their respective simulators that cannot be corrected for with the output we have. Most notably, the ISCCP simulator only reports a single cloud layer for each scene, regardless of whether the scene has multi-layer clouds. This single cloud is assigned a cloud top pressure with a temperature corresponding to the scene's infrared brightness temperature. This introduces two well-known issues: First, clouds under a strong inversion are often placed too high in the atmosphere, at the higher level at which this temperature is found (Garay et al., 2008). Second, in multi-layer cloud scenes in which a non-opaque high cloud overlies lower-level clouds, the simulator often places a single cloud at middle-levels since the brightness temperature will reflect some combination of warm lower-level cloud and the cold but thin upper-level cloud (Pincus et al., 2012; Marchand and Ackerman, 2010). Finally, as noted above, the ISCCP retrieval algorithm reports a single cloud layer with a cloud optical depth equal to the column-integrated optical depth,




Figure 2. As in Figure 1, but for a scenario in which a high cloud partially overlaps a low cloud in the mean climate, but increases in the perturbed climate so as to completely obscure the (unchanged) low cloud.

215 even if multiple cloud layers are present in a given scene. This implies that some portion of the change in upper-level cloud properties as [seen-reported](#) by the simulator (and subsequently diagnosed as nonlow cloud feedbacks or adjustments) may in fact be partly induced by changes in lower-level cloud properties. Hence, even with the aforementioned corrections, one should not consider the modified feedbacks derived herein as strictly “true” nonlow and low cloud feedbacks, owing to the additional issues that we cannot correct for.

220 4 Results

4.1 Cloud Feedbacks

Having motivated our modified cloud feedback calculations and given schematic illustrations of the diagnostic issues they are intended to mitigate, we now turn to examining the impacts of these modifications on the cloud feedbacks diagnosed in climate models. In Figure 3 we show the multi-model mean original and modified net (longwave plus shortwave) nonlow and low cloud amount feedbacks. The difference between modified and original is also shown, which provides a measure of how large the obscuration effect is.

225

The modified low cloud amount feedback is somewhat muted relative to the original low cloud amount feedback (Figure 3d-e). The large positive low cloud amount feedbacks over the NE and SE Pacific stratocumulus decks and over the Southern




Figure 3. Multi-model mean nonlow cloud amount feedback computed using the original methodology (a), using the modified methodology (b), and their difference (c) which measures the effect of obscuration changes on the feedback. The corresponding feedbacks for low clouds are shown in (d-f). Global mean values are displayed at the top right of each map.

Ocean are weaker once accounting for obscuration effects. The interpretation is that, on average, the warming-induced increase
 230 in high cloud coverage over regions of persistent low cloud cover (most prominently, NE Pacific, SE Pacific, west of Namibia, and over the Southern Ocean) hides a portion of the underlying low clouds. In the original decomposition, this contributes to a positive low cloud amount feedback, augmenting the positive unobscured low cloud amount feedback from actual decreases in low cloud cover. However, in some regions like east of Australia, the positive modified low cloud amount feedback is larger than its original version (Figure 3d-e). In these regions, decreases in nonlow clouds reveal low clouds beneath. In the original
 235 decomposition, this contributes negatively to the low cloud amount feedback, diminishing the positive unobscured low cloud amount feedback from actual decreases in low cloud cover.

By definition (see Eqs 11-12), the corrections applied to the low cloud feedback are equal and opposite to those applied to the nonlow cloud feedback. The same is true to a very close approximation for the amount components such that what is taken away from the low cloud amount feedback is given to the high cloud amount feedback (i.e., Figures 3c and f are equal
 240 but opposite in sign). Therefore, the high cloud amount feedback is restored to something very close to zero at nearly every location (Figure 3b), as expected from changes in coverage of clouds with closely cancelling longwave and shortwave effects. The distinct regions of negative high cloud amount feedback over the stratocumulus regions of the NE and SE Pacific and over the Southern Ocean, along with the regions of positive high cloud amount feedback over east Asia, South America, Africa, and east of Australia are all essentially absent from the modified version (Figure 3a-b). That the global mean value is closer to zero
 245 in the original calculation is due to a fortuitous cancellation between larger positive and negative regional values.




Figure 4. Across-model standard deviation of nonlow cloud amount feedback computed using the original methodology (a) and using the modified methodology (b), along with their difference (c). The corresponding feedbacks for low clouds are shown in (d-f). The rectangular box indicates the averaging region for the data presented in Figure A.1, chosen because it is a prominent region of reduced spread in the modified decomposition.

At nearly all locations, but especially over the ocean, the inter-model spread in both the nonlow and low cloud amount feedbacks is reduced in the modified calculation (Figure 4). This is because the original low cloud amount feedback in these regions is positively correlated with the change in obscuration (see Eq 11), reflecting the fact that models with larger increases in obscuration experience a larger augmentation of the unobscured positive low cloud feedback, and vice versa. By removing
 250 these obscuration effects, the modified low cloud feedback is uncorrelated with the change in obscuration and therefore exhibits less inter-model spread (Figure A.1b). Conversely, the original nonlow cloud amount feedback in these regions is negatively correlated with the change in obscuration (see Eq 12), reflecting the fact that models with larger increases in obscuration experience a larger negative bias with respect to the modified nonlow cloud amount feedback, and vice versa. By removing
 255 these obscuration effects, the strong anti-correlation between nonlow cloud feedback and the change in obscuration vanishes and therefore the modified nonlow cloud feedback exhibits less inter-model spread (Figure A.1a).

Given the reduction in inter-model spread in the modified low cloud amount feedback at most locations (Fig. 4f), do the cloud types and/or regions that most contribute to the inter-model spread of global mean cloud feedback change? To answer this, we compute the across-model variance explained between the global mean total net cloud feedback and grid-point values of the nonlow and low net cloud amount feedbacks, for both the original and modified methodologies, closely following
 260 Soden and Vecchi (2011). As expected from previous studies, the spread in global mean net cloud feedback is strongly related to the low cloud amount feedback in regions of prevalent low cloud, including the subtropical and midlatitude oceans (Figure 5d). The modified methodology highlights the same regions, but the variance explained is larger nearly everywhere, especially




Figure 5. Fraction of across-model variance in global mean net cloud feedback explained by local net (top) nonlow cloud amount feedback and (bottom) low cloud amount feedback using the (left) original and (middle) modified methodology. In the right column, we show the difference between modified and original methodologies.

over the Atlantic and Indian Oceans, subtropical South Pacific, equatorial Pacific cold tongue, and North Pacific (Figure 5e,f). The modification also results in a reduction in the variance explained by nonlow clouds over the subtropical Atlantic Ocean (Figure 5c). This may provide further evidence of the importance of properly accounting for obscuration effects, as it leads to a clearer attribution of inter-model spread to its true source (low clouds).

265

270

275

The low cloud amount feedback is much less positive poleward of about 50 degrees in either hemisphere after accounting for obscuration effects (Figure 5d6d). This makes the positive low cloud feedback much more confined to low latitudes and the negative lobe at middle to high latitudes much more robust (Figure 5e6c). Similarly, the latitudinal dipole in nonlow cloud amount feedback centered near 50 degrees in either hemisphere is completely removed (Figure 5b6b). Physically, this may be related to the poleward shift of high clouds in the midlatitude storm track. Without accounting for change in obscuration of underlying clouds, the radiative kernel diagnoses radiative heating on the equatorward flank of the jet (where high clouds vacate) and a radiative cooling on the poleward flank (where high clouds increase). The modified nonlow cloud feedback is much more muted because it accounts for the fact that these high cloud anomalies are occurring in a region of prevalent low cloud cover (Tselioudis et al., 2016). Specifically, the regions vacated by high clouds reveal bright low clouds rather than dark ocean surface, limiting the size of the positive radiative anomaly attributable to high clouds near 40 degrees. The regions experiencing increased high cloud cover on the poleward flank have hidden bright low clouds rather than a dark ocean surface, limiting the size of the negative radiative anomaly attributable to high clouds near 60 degrees.

280

In Figure A.2 we demonstrate the impact of our modified calculations on the distribution of global mean cloud feedback components across models. Accounting for obscuration primary affects the low and nonlow SW and NET cloud feedbacks via




Figure 6. Zonal mean nonlow and low cloud feedbacks and their amount components, computed using the original (black) and modified (red) decompositions. Solid lines represent the multi-model means and the shading spans the $\pm 1\sigma$ range across models.

the amount component, with all other feedback components being either identical to the original calculation (by design), or indistinguishable from them. On average, the low cloud amount component becomes slightly smaller while the nonlow cloud amount feedback becomes slightly larger. Both components exhibit less inter-model spread in the modified calculations.

While the global-mean and multi-model mean feedbacks are not strongly affected by the modified calculations, this belies substantial changes apparent at local scales (Figures 3 and 56) and within individual models (Figure 67). For most models, the original calculation overestimates the positive low cloud amount feedback, as evidenced by the models for which the black bars extend beyond the red markers in Figure 6d7d. This occurs as a result of increased obscuration by nonlow clouds. However, this is not true for all models, as some show little effects of changing obscuration (at least in the global mean), and several show the opposite effect: Most notably, HadGEM3-GC31-LL experiences *decreased* obscuration by nonlow clouds, so the modified low cloud amount feedback is actually considerably larger than the original calculation. This model now has the largest low cloud feedback of all models rather than a near-average value (Figure 6e7c). In the case of low cloud feedback and its amount component, accounting for obscuration affects the magnitude but not the sign of the feedback in all models.

This is not generally the case for the nonlow cloud amount feedback (Figure 6b7b). Several models with negative nonlow cloud amount feedbacks in the original decomposition actually have positive feedbacks in the modified calculation. There




Figure 7. Global mean nonlow and low cloud feedbacks and their amount components computed using the original (black) and modified (red) decompositions, along with their difference (gray). Models are ordered by the strength of their obscuration adjustment. Note that the sign of the Δ Obscuration term is defined in both rows as the modified minus original feedback, which is opposite to the definition in Eq 11.

295 is much stronger across-model agreement that the nonlow cloud amount feedback is positive in the modified calculation. While the nonlow cloud amount feedback is small regardless of which decomposition is used, accounting for obscuration can substantially change the overall nonlow cloud feedback in some models (Figure 6a7a). For example, MPI-ESM-LR's nonlow cloud feedback increases from near-zero to a moderate positive value, while HadGEM3-GC31-LL's large positive nonlow cloud feedback is roughly halved.

300 ~~Originally, models with larger positive~~

Total inter-model variance of global mean cloud feedback remains unchanged regardless of the decomposition, so decreasing variance in both nonlow and low cloud amount feedbacks tended to have larger negative (with negligible change in altitude, optical depth, and residual components) implies that something else must compensate. In the original decomposition, low cloud




Figure 8. Global mean net cloud amount feedbacks for nonlow clouds scattered against those for low clouds.

amount feedback is strongly anti-correlated with the low cloud optical depth component and especially with the nonlow cloud amount feedbacks. This is because component (Figure 8). The latter reflects the fact that increased upper-level cloud cover was hiding lower-level clouds beneath, making the low cloud feedback appear larger. Properly accounting for obscuration effects removes this anti-correlation (Figure 7). This is evidence that artificial correlations among feedback components may be present when not accurately accounting for obscuration effects, as nonlow cloud changes are aliased into low cloud feedbacks. So while the variance in nonlow and low cloud amount feedback have decreased in the modified decomposition, so too has the large negative covariance among several feedback components. This might suggest that the modified decomposition better reveals sources of spread and reduces apparent covariances that may be more artificial than physical, analogous to the fixed relative humidity radiative feedback framework of Held and Shell (2012).

~~Global mean net cloud amount feedbacks for nonlow clouds scattered against those for low clouds.~~

4.2 Rapid Cloud Adjustments

As noted earlier, cloud adjustments that occur rapidly after the step change in CO₂ concentration impart radiative effects that are typically included as part of the effective radiative forcing. In Figure 89, we show the multi-model mean low and nonlow cloud adjustments as computed using the original and modified decomposition, along with their difference. The low cloud rapid adjustment is strongly negative over ocean due to increases in lower-level cloud cover and positive over land due to large reductions in low cloud cover (Figure 8d9d). As noted in Zelinka et al. (2013), a portion of the large increase in lower-level cloud cover over the ocean that is diagnosed by the ISCCP simulator is actually a result of decreased obscuration. This is because nonlow clouds, especially those in the mid-troposphere, decrease in response to the radiative warming and attendant drying from CO₂ (Colman and McAvaney, 2011; Wyant et al., 2012; Kamae and Watanabe, 2012; Kamae et al., 2015). This effect of changing obscuration is confirmed in Figure 89, where the large negative oceanic low cloud amount adjustment is

substantially reduced when accounting for obscuration changes (compare Figure 8d-9d and e). The global and multi-model mean rapid low cloud adjustment is increased from near zero to 0.25 W/m², due to a widespread reduction in the negative oceanic values with little change in the large positive values over land (which do not result from obscuration changes).




Figure 9. As in Figure 3, but for rapid cloud adjustments.

Complementing this result, the modified nonlow cloud adjustment is much closer to zero than in the original decomposition. Rapid decreases in nonlow cloud amount reveal low clouds, making the net radiative adjustment small (Figure 8b9b). This contrasts with the original calculation which essentially assumes that the rapid reduction in nonlow clouds reveals a dark ocean beneath, causing a large positive radiative adjustment (Figure 8a9a). Hence, ~~in the multi-model mean~~averaged globally and across models, the partitioning of the positive rapid cloud adjustment completely switches from being dominated by nonlow clouds plus a small contribution from low clouds (Figure 9a,d) to being dominated by a large positive low cloud contribution that is opposed slightly by a small nonlow cloud contribution (Figure 9b,e).

As was the case for the cloud feedback, the low and nonlow rapid cloud adjustments exhibit less inter-model spread at nearly every location on the globe (Figure 9i0). This can be understood through the arguments discussed above for the cloud feedback, which will not be reiterated here.

Examining the zonal mean rapid cloud adjustments, we see that the modified low cloud adjustment and its amount component, which dominates the response, is systematically shifted toward positive values at all latitudes, but most especially in the Southern Hemisphere middle latitudes (Figure 10e-d11c-d). Similarly, the modified nonlow cloud amount adjustment is shifted to be very close to zero at every latitude, and the large values in either hemisphere around 40 degrees latitude apparent in the original decomposition are now completely removed (Figure 10b11b).

Distributions of global mean rapid cloud adjustments for nonlow and low clouds are shown in Figure A.3. Similar to the cloud feedbacks, the modified calculations cause the largest changes for the net and SW amount components. In particular, the




Figure 10. As in Figure 4, but for rapid cloud adjustments.




Figure 11. As in Figure 5, but for rapid cloud adjustments.

345 distribution of SW and net low cloud amount adjustments shifts from being centered on zero to a positive value, as this calculation no longer allows decreased nonlow cloud coverage from aliasing itself onto the low cloud rapid adjustment. Similarly, the SW and net nonlow cloud amount adjustment distribution shifts downward from a moderate positive value to something closer to zero. This weaker positive nonlow cloud adjustment is because rapid reductions in nonlow clouds reveal lower-level clouds in the modified decomposition whereas they are assumed to reveal a dark ocean in the original decomposition.




Figure 12. As in Figure 67, but for rapid cloud adjustments.

350 As noted above for the cloud feedback, the impact of accounting for obscuration effects varies substantially among models. Unlike for the cloud feedback, however, the effect is uniform in sign across models. Specifically, in all models, rapid decreases in nonlow cloud fraction reveal more low clouds, making the original low cloud adjustment weaker positive or stronger negative, as indicated by the red circles being located to the right of the black bars in Figure He12c. This is evidence of the effect illustrated schematically in Figure 1. In many models, the original small positive low cloud amount adjustment more than doubles in the modified calculation, and several models' adjustments change sign from negative to positive. Similarly, the original

355 nonlow cloud amount adjustment is positively biased (see Figure 1) such that the modified adjustment is much smaller and in many models switches to a negative value (red circles to the left of black bars in Figure Hb12b). While this sign change does not occur in any model for the overall nonlow cloud adjustment, the reduction in positive nonlow cloud adjustment strength remains apparent in Figure Ha12a.

Rapid cloud adjustment terms are largely consistent between piClim- and amip-style quadrupled CO₂ experiments, as shown for the models that conducted both in Figure A.4. However, for several models, the piClim-style experiments lead to larger positive low cloud adjustments than the amip-style experiments. In these models, the rapid response of low and nonlow clouds (but not of the obscuration) appears to depend on experiment design. We leave further exploration of why this is the case to future work.

5 Conclusions

365 In this study we presented a methodology for decomposing cloud feedbacks and rapid adjustments among low and nonlow clouds that properly accounts for obscuration effects. This methodology has been used in previous studies (e.g., Zelinka et al., 2018; Scott et al., 2020; Myers et al., 2021; Zelinka et al., 2022; Chao et al., 2024) but the effect of these choices has not been formally documented across models to date. While the overall cloud feedbacks and adjustments do not depend on the methodology employed, and the decision to split the feedback among low and nonlow clouds rather than some other decomposition is partly arbitrary, the impacts of these methodological choices are important because they can improve the mechanistic interpretation of the results and avoid artificial relationships that are not physical. In this sense the recommended methodology is analogous to the constant relative humidity feedback decomposition proposed by Held and Shell (2012), which reduces spread in water vapor, lapse rate, and Planck feedbacks and eliminates the anticorrelation between lapse rate and water vapor feedbacks, revealing more clearly the dominant uncertainties in radiative feedbacks.

375 We find that the positive multi-model mean low cloud feedback is weaker in our modified calculations because it excludes the positive radiative contribution from apparent reductions in low clouds that are due solely to increased obscuration by nonlow clouds. Complementing this, the nonlow cloud feedback is much closer to zero at every location in our modified calculation, as changes in nonlow clouds have a muted radiative impact when occurring over low clouds. Across models, the strength and in some cases even the sign of the low and nonlow cloud feedbacks change, and an apparent anti-correlation between low and nonlow cloud amount feedbacks is removed when accounting for obscuration. Finally, the inter-model variance in both nonlow and low cloud feedbacks is damped in nearly all locations when properly accounting for obscuration effects.

385 Upon quadrupling of CO₂, large decreases in oceanic upper-level cloud coverage reveal underlying lower-level clouds. In the original decomposition this leads to an apparent negative oceanic low cloud radiative adjustment that is solely due to reduced obscuration. Properly accounting for obscuration, however, strongly reduces this negative adjustment, leading to a moderate positive low cloud adjustment in the multi-model mean. Moreover, the positive nonlow cloud radiative adjustment from the large reduction in nonlow clouds in the original decomposition is substantially weakened in the modified calculations, such that the modified nonlow rapid cloud adjustment is very close to zero at all locations. Hence in the multi-model mean, the

rapid cloud adjustment to quadrupled CO₂ arises from a large positive low cloud radiative adjustment countered only slightly by a weak negative nonlow cloud radiative adjustment. As was the case with cloud feedbacks, the inter-model variance in both nonlow and low cloud adjustments is damped in nearly all locations when properly accounting for obscuration effects, most notably over the oceans.

Given that neglect of obscuration effects can lead to misleading results regarding the attribution of feedbacks and adjustments to specific cloud types or physical processes, and in most locations inflates the inter-model variance in these, we recommend that the community follow the methodology presented herein when computing low and nonlow cloud feedbacks and adjustments using cloud radiative kernels. It must be borne in mind, however, that no decomposition that relies on cloud information from passive satellite retrievals can ensure that the radiative effect attributed to a given cloud type is solely due to changes in that cloud type, owing to nonlinear aspects of radiation. Nevertheless, addressing obscuration effects is an important step in this direction. Code to perform this decomposition and a Jupyter notebook to demonstrate the calculations is available at the url provided in the Code and Data Availability Section.

Data availability. All CMIP climate model data used in this study is available from the Earth System Grid Federation (<https://esgf.llnl.gov/>).

Code and data availability. Cloud radiative kernels, along with the code to perform the calculations in this study are available at <https://zenodo.org/records/13686878> (Mark Zelinka, 2024).

Appendix A

One may choose to decompose feedbacks or adjustments into contributions from high (H), middle (M), and low (L) clouds.

405 In this case we define the upper-level clear-sky fraction as

$$F = 1 - \overline{M} - H, \quad (\text{A1})$$

in which case (6) becomes

$$L' = L_{RU}' \overline{1 - M - H} + \overline{L_U} M' + \overline{L_U} H' + \epsilon, \quad (\text{A2})$$

where ϵ contains the covariance terms. Unobscured mid-level clouds will have a form similar to (1),

$$410 \quad M_{RU} = \frac{M}{F_H}, \quad (\text{A3})$$

where

$$F_H = 1 - H. \quad (\text{A4})$$

We can then decompose M' in a form similar to (6):

$$M' = M_{RU}' \overline{F_H} + \overline{M_U} F_H' + \epsilon, \quad (\text{A5})$$

415 which is equivalent to

$$M' = M_{RU}' \overline{1 - H} - \overline{M_U} H' + \epsilon. \quad (\text{A6})$$

There are now a total of three obscuration terms in (A.2) and (A.6): $\overline{L_U} M' - \overline{L_U} H'$ is a mid-level cloud response that reveals or obscures underlying low clouds, $\overline{L_U} H' - \overline{M_U} H'$ is a high-level cloud response that reveals or obscures underlying low clouds, and $\overline{M_U} H' - \overline{M_U} H'$ is a high-level cloud response that reveals or obscures underlying mid-level clouds. As before, if we include

420 the obscuration of lower clouds by a mid or high cloud as part of the mid or high cloud feedback (and omitting covariances), we get:

$$\text{low}_{\text{mod}} = L' + \overline{L_U} M' + \overline{L_U} H' \quad (\text{A7})$$

$$\text{mid}_{\text{mod}} = M' - \overline{L_U} M' + \overline{M_U} H' \quad (\text{A8})$$

$$\text{high}_{\text{mod}} = H' - \overline{L_U} H' - \overline{M_U} H'. \quad (\text{A9})$$

425 This ensures that the sum of the three modified cloud responses is equal to the total cloud response.

Author contributions. All analyses in the paper were performed by MDZ. The first draft of the manuscript was written by MDZ and all authors commented on subsequent versions of the manuscript. All authors read and approved the final manuscript.




Figure A.1. Net cloud amount feedbacks for (a) nonlow and (b) low clouds averaged over the Eastern Pacific ITCZ region indicated in Figure 4 scattered against the coincident change in obscuration. Feedbacks computed using the original and modified decomposition are shown with black and red markers, respectively. Errorbars to the right indicate the multi-model mean and standard deviation of original and modified feedbacks.

Competing interests. The authors declare no competing interests.

Acknowledgements. [We thank two anonymous reviewers for their helpful comments that improved the manuscript.](#) Mark D. Zelinka, Li-Wei Chao, and Yi Qin were supported by the U.S. Department of Energy (DOE) Regional and Global Model Analysis program area. Timothy A. Myers was supported by the NOAA Cooperative Agreement with CIRES, NA17OAR4320101 and NA22OAR4320151, and by the NOAA/ESRL Atmospheric Science for Renewable Energy (ASRE) program. The statements, findings, conclusions, and recommendations are those of the author and do not necessarily reflect the views of NOAA or the U.S. Department of Commerce. We acknowledge the World Climate Research Programme, which, through its Working Group on Coupled Modelling, coordinated and promoted CMIP. We thank the climate modeling groups for producing and making available their model output, the Earth System Grid Federation (ESGF) for archiving the




Figure A.2. Box and whisker plots summarizing the distribution of global mean nonlow and low cloud feedbacks across models. Feedbacks are separated into LW, SW, and net (LW+SW) components, each of which is further separated into components due to changes in amount, altitude, and optical depth, along with a residual term. Feedbacks are shown for the original calculation in black and the modified calculation in red. Each box extends from the first quartile to the third quartile of the data, with a line at the median and a diamond at the mean. The whiskers extend from the box to the farthest data point lying within 1.5x the inter-quartile range from the box. Flier points are those past the end of the whiskers.

data and providing access, and the multiple funding agencies who support CMIP and ESGF. The work of Mark D. Zelinka and Li-Wei Chao was performed under the auspices of the U.S. DOE by Lawrence Livermore National Laboratory under contract DEAC52-07NA27344. The Pacific Northwest National Laboratory is operated for the DOE by the Battelle Memorial Institute under contract DE-AC05-76RL01830.




Figure A.3. As in Figure A.2, but for rapid cloud adjustments.

References

- 440 Bony, S., Colman, R., Kattsov, V. M., Allan, R. P., Bretherton, C. S., Dufresne, J. L., Hall, A., Hallegatte, S., Holland, M. M., Ingram, W.,
 Randall, D. A., Soden, B. J., Tselioudis, G., and Webb, M. J.: How Well Do We Understand and Evaluate Climate Change Feedback
 Processes?, *J. Climate*, 19, 3445–3482, <https://doi.org/10.1175/JCLI3819.1>, 2006.
- Ceppi, P., Myers, T. A., Nowack, P., Wall, C. J., and Zelinka, M. D.: Implications of a Pervasive Climate Model Bias
 for Low-Cloud Feedback, *Geophysical Research Letters*, 51, e2024GL110525, <https://doi.org/10.1029/2024GL110525>,
 445 <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2024GL110525>, 2024.
- Cess, R. D., Potter, G. L., Blanchet, J. P., Boer, G. J., Ghan, S. J., Kiehl, J. T., Treut, H. L., Li, Z. X., Liang, X. Z., Mitchell, J. F. B.,
 Morcrette, J. J., Randall, D. A., Riches, M. R., Roeckner, E., Schlese, U., Slingo, A., Taylor, K. E., Washington, W. M., Wetherald,
 R. T., and Yagai, I.: Interpretation of Cloud-Climate Feedback as Produced by 14 Atmospheric General Circulation Models, *Science*, 245,
 513–516, <https://doi.org/10.1126/science.245.4917.513>, 1989.
- 450 Cess, R. D., Potter, G. L., Blanchet, J. P., Boer, G. J., Genio, A. D. D., Déqué, M., Dymnikov, V., Galin, V., Gates, W. L., Ghan, S. J., Kiehl,
 J. T., Lacis, A. A., Treut, H. L., Li, Z.-X., Liang, X.-Z., McAvaney, B. J., Meleshko, V. P., Mitchell, J. F. B., Morcrette, J.-J., Randall,




Figure A.4. As in Figure H12, but comparing rapid cloud adjustments between piClim- and amip-style CO₂ quadrupling experiments for models that performed both.

D. A., Rikus, L., Roeckner, E., Royer, J. F., Schlese, U., Sheinin, D. A., Slingo, A., Sokolov, A. P., Taylor, K. E., Washington, W. M., Wetherald, R. T., Yagai, I., and Zhang, M.-H.: Intercomparison and interpretation of climate feedback processes in 19 atmospheric general circulation models, *Journal of Geophysical Research: Atmospheres*, 95, 16601–16615, <https://doi.org/10.1029/JD095iD10p16601>,
 455 _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/JD095iD10p16601>, 1990.

Chao, L.-W., Zelinka, M. D., and Dessler, A. E.: Evaluating Cloud Feedback Components in Observations and Their Representation in Climate Models, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD039427, <https://doi.org/10.1029/2023JD039427>,
 https://onlinelibrary.wiley.com/doi/pdf/10.1029/2023JD039427, 2024.

Colman, R. and McAvaney, B.: On tropospheric adjustment to forcing and climate feedbacks, *Climate Dyn.*, 36, 1649–1658,
 460 <https://doi.org/10.1007/s00382-011-1067-4>, 2011.

Garay, M. J., de Szoek, S. P., and Moroney, C. M.: Comparison of marine stratocumulus cloud top heights in the southeastern Pacific retrieved from satellites with coincident ship-based observations, *J. Geophys. Res.*, 113, <https://doi.org/10.1029/2008JD009975>, 2008.

- Held, I. M. and Shell, K. M.: Using Relative Humidity as a State Variable in Climate Feedback Analysis, *Journal of Climate*, 25, 2578–2582, <https://doi.org/10.1175/JCLI-D-11-00721.1>, 2012.
- 465 Kamae, Y. and Watanabe, M.: Tropospheric adjustment to increasing CO₂: Its timescale and the role of land-sea contrast, *Climate Dyn.*, pp. 1–18, 2012.
- Kamae, Y., Watanabe, M., Ogura, T., Yoshimori, M., and Shiogama, H.: Rapid Adjustments of Cloud and Hydrological Cycle to Increasing CO₂: a Review, *Current Climate Change Reports*, 1, 103–113, <https://doi.org/10.1007/s40641-015-0007-5>, 2015.
- Lutsko, N. J., Luongo, M. T., Wall, C. J., and Myers, T. A.: Correlation Between Cloud Adjustments and Cloud Feedbacks Responsible for Larger Range of Climate Sensitivities in CMIP6, *Journal of Geophysical Research: Atmospheres*, 127, e2022JD037486, <https://doi.org/10.1029/2022JD037486>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1029/2022JD037486](https://onlinelibrary.wiley.com/doi/pdf/10.1029/2022JD037486), 2022.
- 470 Marchand, R. and Ackerman, T.: An analysis of cloud cover in multiscale modeling framework global climate model simulations using 4 and 1 km horizontal grids, *Journal of Geophysical Research-Atmospheres*, 115, <https://doi.org/10.1029/2009jd013423>, 2010.
- Mark Zelinka: [mzelinka/cloud-radiative-kernels](https://doi.org/10.5281/ZENODO.13686878): Sep 4, 2024 Release, <https://doi.org/10.5281/ZENODO.13686878>, 2024.
- 475 Myers, T. A., Scott, R. C., Zelinka, M. D., Klein, S. A., Norris, J. R., and Caldwell, P. M.: Observational constraints on low cloud feedback reduce uncertainty of climate sensitivity, *Nature Climate Change*, 11, 501–507, <https://doi.org/10.1038/s41558-021-01039-0>, number: 6 Publisher: Nature Publishing Group, 2021.
- Pincus, R., Platnick, S., Ackerman, S. A., Hemler, R. S., and Hofmann, R. J. P.: Reconciling Simulated and Observed Views of Clouds: MODIS, ISCCP, and the Limits of Instrument Simulators, *Journal of Climate*, 25, 4699–4720, <https://doi.org/10.1175/jcli-d-11-00267.1>, 480 2012.
- Pincus, R., Forster, P. M., and Stevens, B.: The Radiative Forcing Model Intercomparison Project (RFMIP): experimental protocol for CMIP6, *Geoscientific Model Development*, 9, 3447–3460, <https://doi.org/10.5194/gmd-9-3447-2016>, 2016.
- Qin, Y., Zelinka, M. D., and Klein, S. A.: On the Correspondence Between Atmosphere-Only and Coupled Simulations for Radiative Feedbacks and Forcing From CO₂, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035460, <https://doi.org/10.1029/2021JD035460>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021JD035460](https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021JD035460), 485 2022.
- Raghuraman, S. P., Medeiros, B., and Gettelman, A.: Observational Quantification of Tropical High Cloud Changes and Feedbacks, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD039364, <https://doi.org/10.1029/2023JD039364>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1029/2023JD039364](https://onlinelibrary.wiley.com/doi/pdf/10.1029/2023JD039364), 2024.
- Ringer, M. A., Andrews, T., and Webb, M. J.: Global-mean radiative feedbacks and forcing in atmosphere-only and coupled atmosphere-ocean climate change experiments, *Geophysical Research Letters*, 41, 4035–4042, <https://doi.org/10.1002/2014gl060347>, 490 2014.
- Rossow, W. B. and Schiffer, R. A.: Advances in Understanding Clouds from ISCCP, *Bull. Amer. Meteor. Soc.*, 80, 2261–2287, [https://doi.org/10.1175/1520-0477\(1999\)0802.0.CO;2](https://doi.org/10.1175/1520-0477(1999)0802.0.CO;2), 1999.
- Scott, R. C., Myers, T. A., Norris, J. R., Zelinka, M. D., Klein, S. A., Sun, M., and Doelling, D. R.: Observed Sensitivity of Low-Cloud Radiative Effects to Meteorological Perturbations over the Global Oceans, *Journal of Climate*, 33, 7717–7734, <https://doi.org/10.1175/JCLI-D-19-1028.1>, publisher: American Meteorological Society Section: *Journal of Climate*, 2020.
- 495 Sherwood, S. C., Webb, M. J., Annan, J. D., Armour, K. C., Forster, P. M., Hargreaves, J. C., Hegerl, G., Klein, S. A., Marvel, K. D., Rohling, E. J., Watanabe, M., Andrews, T., Braconnot, P., Bretherton, C. S., Foster, G. L., Hausfather, Z., Heydt, A. S. v. d., Knutti, R., Mauritsen, T., Norris, J. R., Proistosescu, C., Rugenstein, M., Schmidt, G. A., Tokarska, K. B., and Zelinka, M. D.: An Assessment of Earth’s Climate Sensitivity Using Multiple Lines of Evidence, *Reviews of Geophysics*, 58, e2019RG000678, 500 <https://doi.org/https://doi.org/10.1029/2019RG000678>, 2020.

- Smith, C. J., Kramer, R. J., Myhre, G., Alterskjær, K., Collins, W., Sima, A., Boucher, O., Dufresne, J.-L., Nabat, P., Michou, M., Yukimoto, S., Cole, J., Paynter, D., Shiogama, H., O'Connor, F. M., Robertson, E., Wiltshire, A., Andrews, T., Hannay, C., Miller, R., Nazarenko, L., Kirkevåg, A., Olivié, D., Fiedler, S., Lewinschal, A., Mackallah, C., Dix, M., Pincus, R., and Forster, P. M.: Effective radiative forcing and adjustments in CMIP6 models, *Atmospheric Chemistry and Physics*, 20, 9591–9618, <https://doi.org/10.5194/acp-20-9591-2020>, publisher: Copernicus GmbH, 2020.
- Soden, B. J. and Vecchi, G. A.: The vertical distribution of cloud feedback in coupled ocean-atmosphere models, *Geophys. Res. Lett.*, 38, <https://doi.org/10.1029/2011GL047632>, 2011.
- Taylor, K. E., Crucifix, M., Braconnot, P., Hewitt, C. D., Doutriaux, C., Broccoli, A. J., Mitchell, J. F. B., and Webb, M. J.: Estimating Shortwave Radiative Forcing and Response in Climate Models, *J. Climate*, 20, 2530–2543, <https://doi.org/10.1175/JCLI4143.1>, 2007.
- 510 Tselioudis, G., Lipat, B. R., Konsta, D., Grise, K. M., and Polvani, L. M.: Midlatitude cloud shifts, their primary link to the Hadley cell, and their diverse radiative effects, *Geophysical Research Letters*, 43, 4594–4601, <https://doi.org/10.1002/2016gl068242>, 2016.
- Webb, M. J., Andrews, T., Bodas-Salcedo, A., Bony, S., Bretherton, C. S., Chadwick, R., Chepfer, H., Douville, H., Good, P., Kay, J. E., Klein, S. A., Marchand, R., Medeiros, B., Siebesma, A. P., Skinner, C. B., Stevens, B., Tselioudis, G., Tsushima, Y., and Watanabe, M.: The Cloud Feedback Model Intercomparison Project (CFMIP) contribution to CMIP6, *Geoscientific Model Development*, 10, 359–384, <https://doi.org/10.5194/gmd-10-359-2017>, 2017.
- 515 Wyant, M. C., Bretherton, C. S., Blossey, P. N., and Khairoutdinov, M.: Fast cloud adjustment to increasing CO₂ in a superparameterized climate model, *J. Adv. Model. Earth Syst.*, 4, <https://doi.org/10.1029/2011MS000092>, 2012.
- Zelinka, M. D., Klein, S. A., and Hartmann, D. L.: Computing and Partitioning Cloud Feedbacks Using Cloud Property Histograms. Part I: Cloud Radiative Kernels, *Journal of Climate*, 25, 3715–3735, <https://doi.org/10.1175/jcli-d-11-00248.1>, 2012a.
- 520 Zelinka, M. D., Klein, S. A., and Hartmann, D. L.: Computing and Partitioning Cloud Feedbacks Using Cloud Property Histograms. Part II: Attribution to Changes in Cloud Amount, Altitude, and Optical Depth, *Journal of Climate*, 25, 3736–3754, <https://doi.org/10.1175/JCLI-D-11-00249.1>, 2012b.
- Zelinka, M. D., Klein, S. A., Taylor, K. E., Andrews, T., Webb, M. J., Gregory, J. M., and Forster, P. M.: Contributions of Different Cloud Types to Feedbacks and Rapid Adjustments in CMIP5, *Journal of Climate*, 26, 5007–5027, <https://doi.org/10.1175/jcli-d-12-00555.1>, 2013.
- 525 Zelinka, M. D., Zhou, C., and Klein, S. A.: Insights from a refined decomposition of cloud feedbacks, *Geophysical Research Letters*, 43, 9259–9269, <https://doi.org/10.1002/2016gl069917>, 2016.
- Zelinka, M. D., Grise, K. M., Klein, S. A., Zhou, C., DeAngelis, A. M., and Christensen, M. W.: Drivers of the Low-Cloud Response to Poleward Jet Shifts in the North Pacific in Observations and Models, *Journal of Climate*, 31, 7925–7947, <https://doi.org/10.1175/jcli-d-18-0114.1>, 2018.
- 530 Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., Klein, S. A., and Taylor, K. E.: Causes of Higher Climate Sensitivity in CMIP6 Models, *Geophysical Research Letters*, 47, e2019GL085782, <https://doi.org/https://doi.org/10.1029/2019GL085782>, 2020.
- Zelinka, M. D., Klein, S. A., Qin, Y., and Myers, T. A.: Evaluating Climate Models' Cloud Feedbacks Against Expert Judgment, *Journal of Geophysical Research: Atmospheres*, 127, e2021JD035198, <https://doi.org/10.1029/2021JD035198>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021JD035198](https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021JD035198), 2022.
- 535 Zhou, C., Zelinka, M. D., Dessler, A. E., and Yang, P.: An Analysis of the Short-Term Cloud Feedback Using MODIS Data, *Journal of Climate*, 26, 4803–4815, <https://doi.org/10.1175/JCLI-D-12-00547.1>, 2013.