

Depth Effects of Long-term Organic Residue Application on Soil Organic Carbon Stocks in Central Kenya

Claude Raoul Müller¹, Johan Six¹, Daniel Mugendi Njiru², Bernard Vanlauwe³, Marijn Van de Broek¹

¹ Department of Environmental Systems Science, ETH Zurich, 8092 Zürich, Switzerland

² Department of Land and Water Management, University of Embu, P.O. Box 6-60100, Embu, Kenya

³ International Institute of Tropical Agriculture (IITA), c/o ICIPE Compound, P. O. Box 30772-00100, Nairobi, Kenya

Correspondence to: Claude Raoul Müller (claude.mueller@usys.ethz.ch)

Abstract

In arable soils, a substantial portion of soil organic carbon (SOC) is stored below the plough layer. To develop sustainable soil management strategies, it is important to assess how they affect the quantity of SOC stored in the subsoil. Therefore, we investigated the impact of organic and inorganic nutrient inputs on SOC stocks down to 70 cm depth in a long-term field trial in Embu, Kenya. There were 3 organic input treatments (manure, *Tithonia diversifolia* residues, and maize stover) and a control treatment, each with and without the application of mineral nitrogen. These different treatments were applied to a maize monoculture over 38 growing seasons (19 years). Our results show that manure application had the largest positive impact on SOC stocks compared to the control, which was observed down to 60 cm depth. In contrast, *Tithonia diversifolia* and maize stover application led to significantly larger SOC stocks compared to the control only within the top 20 cm and 40 cm, respectively. Among the three organic residue treatments, only the application of manure had a significant effect on the SOC stock of the subsoil (i.e., the 30-70 cm depth layer). However, when considering the whole measured profile (i.e., 0-70 cm), all treatments led to significantly higher SOC stocks compared to the $91 \pm 12 \text{ t C ha}^{-1}$ of the control: manure had the highest stocks ($120 \pm 24 \text{ t C ha}^{-1}$), followed by maize stover ($112 \pm 17 \text{ t C ha}^{-1}$) and *tithonia diversifolia* ($105 \pm 11 \text{ t C ha}^{-1}$). Mineral nitrogen application did not have a significant impact on SOC stocks down to 70 cm depth. Our findings demonstrate that SOC in the subsoil comprised $48.5\% \pm 1.7\%$ of the total SOC stocks across the 0-70 cm soil profile, yet only manure application affected subsoil SOC levels, while other organic amendments only increased SOC in the surface layer. Our results imply that gathering knowledge on the soil below the typically studied 0-30 cm depth layer will improve the overall assessment of agroecosystem properties, which is necessary to optimize soil system resilience, limit organic matter losses and improve crop productivity.

Deleted: carbon

Deleted: these

Deleted: carbon

Deleted: solely

Deleted: Overall, our findings indicate that SOC in the subsoil of the studied agroecosystem was affected by the type of added organic amendmen...ts.

Deleted: ts.

Deleted:

1 Introduction

Globally, food production systems need to satisfy ~~food demand of an increasingly demanding and growing population~~. However, changing land use for food production and increasing productivity leads to intense use of our limited natural resources, which can have a negative impact on the quality of groundwater and the atmosphere. In the long term, intensifying agriculture also leads to soil quality degradation (e.g., through crusting, compaction, erosion, and depletion of organic matter) (Lal, 2008). Tropical regions are currently experiencing the most significant land-use change globally, largely due to the rapid conversion of natural habitats into agricultural land (Hansen et al., 2013). ~~Eastern Africa followed this trend, experiencing a substantial land use transformation between 1988 and 2017, with cropland expanding by 35% and settlements by 43%, primarily at the expense of woody vegetation~~ (Bullock et al., 2021). This swift alteration of land use has substantial effects on the amount of soil organic carbon (SOC) stored in tropical ecosystems, which contains approximately 44% of the global SOC (Veldkamp et al., 2020). The concentration of SOC is an important component of soil fertility, as it serves as a vital nutrient source for plants and microbes (Chen and Aviad, 1990; Trevisan et al., 2010). Thus, maintaining SOC levels over time is crucial for sustaining soil productivity, especially in the highly weathered soils of tropical regions, which are particularly prone to organic carbon (OC) loss (Feller and Beare, 1997). The conversion of tropical forests to agriculture typically results in an average reduction of OC concentration in the topsoil (down to a depth of 30 cm) of around 50% within 25 years (Veldkamp et al., 2020).

The regular harvesting of crops in farming systems results in the removal of nutrients that the plants have taken up. Thus, inadequate inputs of organic matter and nutrients in farming systems of sub-Saharan Africa (SSA) often leads to the loss of SOC and nutrient depletion, contributing to the lower yields observed in SSA compared to other regions of the world (Vanlauwe and Giller, 2006). Therefore, agricultural management should not only aim at increasing crop yield in the short term, but also at maintaining SOC at a level that maintains the key functions that it regulates (e.g., nutrient cycling, water retention, improved cation exchange capacity (CEC), soil structure stabilization, promotion of biodiversity) in the long term (Wiesmeier et al., 2019). Ideally, the management of cropping systems should respect the principles of sustainable intensification by maximizing production while preserving the environment; different management approaches exist to reach this goal (Pretty, 2011). An example of this is integrated soil fertility management (ISFM), which is defined by Vanlauwe et al. (2010) as *“a set of soil fertility management practices that necessarily include the use of fertilizer, organic inputs, and improved germplasm combined with the knowledge on how to adapt these practices to local conditions, aiming at maximizing agronomic use efficiency of the applied nutrients and improving crop productivity. All inputs need to be managed following sound agronomic principles”*. This approach promotes various practices, including the combined application of mineral fertilizer and organic ~~residues (OR)~~, such as plant residues, manure, and compost. Such practices aim to enhance crop productivity while preserving soil fertility, especially in a context where input availability is often limited, as is the case for smallholder farming in SSA.

Deleted: the increasing food demand of an ever-more demanding

Deleted: -

Deleted: material

Deleted: (i.e., organic residues (OR)).

80 Applying OR over multiple years in tropical croplands can significantly increase the SOC content (Fujisaki et al., 2018),
despite faster SOC turnover rates in tropical compared to temperate climates (Wang et al., 2018). However, the contribution
of OR to the amount of SOC is, in part, controlled by the type of added OR (Córdova et al., 2018). The residue quality has
traditionally been defined by the nitrogen, lignin, and polyphenol content of organic inputs; overall, the lower the (lignin +
polyphenol)/N ratio, the higher the quality of the residue (Chivenge et al., 2011b; Vanlauwe et al., 2005). According to
85 Córdova et al. (2018), applying high-quality residues has the potential to lead to a higher accumulation of stable C in the soil
than low-quality residues. However, an incubation experiment on tropical soil indicated that, while this might be true in the
short term, initial OR quality does not affect SOC accumulation in the long term (Gentile et al., 2011). Moreover, Laub et al.
(2023) found that for four tropical agroecosystems in Kenya, it is mainly the application of animal manure that has the potential
to limit SOC losses from intensive arable soil use. Most importantly, according to Vanlauwe et al. (2015), the efficiency of
90 strategies to increase the SOC content

is determined by variations in local soil conditions. This is supported by the variability in results obtained by different studies
in tropical soils that either suggest that increasing the SOC content with adapted practices is possible (Adams et al., 2020;
Fujisaki et al., 2018; Laub et al., 2022) or the application of large amounts of organic residues is not sufficient to increase or
95 even maintain the SOC content (Kihara et al., 2020; Cardinael et al., 2022; Laub et al., 2023b). The contradictory findings
across tropical regions highlight the complexity of SOC dynamics in these agroecosystems where different factors, such as
initially high SOC contents, favorable conditions to decomposition, and the limited capacity of 1:1 kaolinite clay minerals to
stabilize OC, contribute to consistent SOC losses despite the application of organic residues in tropical soils (Laub et al.,
2023b; Six et al., 2002; Sommer et al., 2018). While some studies demonstrate increases in SOC stocks through management
100 practices, others report continued SOC losses even with substantial organic inputs, suggesting that local soil properties and
environmental conditions play crucial roles. This knowledge gap is particularly evident in tropical agroecosystems, where the
interactions between organic amendments, soil properties, and SOC stabilization mechanisms remain poorly understood,
necessitating further research.

105 In most studies on agroecosystems, soils are studied down to 30 cm on average (Yost and Hartemink, 2020). Therefore, we
defined the threshold between topsoil and subsoil at 30 cm. Additionally, while field studies on plant nutrient acquisition from
the subsoil are rare, it has been shown that crop nutrient availability can be improved when organic residues and mineral
fertilizer are placed in deeper soil layers (20–40 cm depth), suggesting that subsoil OC can play an active role in nutrient use
efficiency (Liu et al., 2021; Ma et al., 2022). This seems rational, as most plant roots extend beyond 30 cm in depth, allowing
110 them to benefit from nutrient availability and cycling in the subsoil (Greenwood et al., 1982; Tardieu, 1988). Also, studying
only topsoil OC is not sufficient to estimate the effect of management practices on the OC storage capacity of soils because it
is estimated that 50% of SOC stocks are located below 30 cm (Balesdent et al., 2018; Lal, 2018; Shumba et al., 2024; Yost

Deleted: the
Deleted: Although the turnover rate of SOC is faster in tropical than temperate climates (Wang et al., 2018), applying OR over multiple years in tropical croplands has the potential to significantly increase the SOC content (Fujisaki et al., 2018).
Deleted: its
Deleted:

Formatted: English (US)
Field Code Changed

Field Code Changed
Deleted: successful
Deleted: enhancement
Formatted: English (US)
Formatted: English (US)
Commented [MV1]: This is a repetition of what you said a couple of sentences earlier, maybe rephrase or remove this here?
Formatted: English (US)
Moved up [1]: (Laub et al., 2023b; Six et al., 2002; Sommer et al., 2018).

Deleted: Different factors, such as initially high SOC contents, favorable conditions to decomposition, and the limited capacity of 1:1 kaolinite clay minerals to stabilize OC, contribute to consistent SOC losses despite the application of organic residues in tropical soils (Laub et al., 2023b; Six et al., 2002; Sommer et al., 2018). Therefore, more data is necessary to better understand the effect of organic amendments on the SOC content in tropical agroecosystems.
Deleted: w

and Hartemink, 2020). Moreover, not only do plant roots and their associated exudates provide direct OC inputs to the subsoil (Van De Broek et al., 2020), also organic matter applied in the topsoil undergoes cycles of sorption and desorption through microbial processing, gradually migrating down the soil profile (Kaiser and Kalbitz, 2012a). The average residence time of subsoil OC ranges from decades to millennia (Balesdent et al., 2018; Mathieu et al., 2015), making subsoil potentially a long-term carbon sink (Rumpel and Kögel-Knabner, 2011). However, only a small proportion of new OC inputs to the subsoil can potentially be stabilized in the long-term or affect the subsoil environment in the short-term (Sierra et al., 2024). These new OC inputs in the subsoil can be detected through their $\Delta^{14}\text{C}$ signature, which is used as a proxy for the age of carbon-containing soil components (Ehleringer et al., 2000; Trumbore et al., 1989). Also $\delta^{13}\text{C}$ can be used to detect the portion of C3 compared to C4 plant residues in the subsoil, enabling, for example, the tracking of maize-derived carbon inputs (Balesdent et al., 1987; Farquhar G D et al., 1989). Understanding the impact of the application of OR on SOC storage in the subsoil would aid the investigation of the importance of the subsoil environment for plant nutrient accessibility.

In a recent study conducted at the same long term field trial studied in the present manuscript (Embu), Laub et al. (2023b) showed that all nutrient management treatments resulted in significant topsoil SOC losses, while the use of farmyard manure was the most effective strategy to minimize the loss of SOC. In their study on the Embu site, they observed consistent decreases in SOC concentration over time for all treatments (Fig. S8). The control treatment showed the greatest decline, with a loss of about 0.6-0.7 g C kg⁻¹ yr⁻¹. Despite differences in treatment effectiveness in maintaining OC, farmyard manure (at 4 t C ha⁻¹ yr⁻¹) emerged as the most effective strategy to limit SOC losses at Embu, with the lowest OC loss (i.e., 0.4 g C kg⁻¹ yr⁻¹). However, their focus was on the effects of treatments above 15 cm depth. Therefore, in this study we explore the effects on deeper soil layers with a high vertical resolution (5 cm) on the sites studied by Laub et al. (2023b).

Given the limited knowledge related to how organic amendments and fertilizer management affect subsoil OC dynamics in tropical agroecosystems, this study aims to examine the impact of different organic residue treatments on SOC stocks down to 70 cm by answering the following research question: Down to which depth do the different organic residues affect SOC stocks?

2 Methods

2.1 Study design

2.1.1 Study area

The field trial was initiated in 2002 in Embu (Central Kenya: 0°30' S, 37°27' E; 1380 m above sea level). A continuous maize monoculture has been grown in two annual growing seasons that coincide with the bimodal precipitation pattern. The long rainy season generally occurs from March to September, gradually decreasing towards the end, leading into a dry period. The short rainy season spans from October to February, also tapering off into a dry period. The average annual precipitation is 1175mm, and the mean annual temperature is 20.1°C. The soil in this area is a highly weathered Humic Nitisol. Originally,

Deleted:

Deleted: Furthermore, such knowledge would help improving soil models, thereby optimizing predictions to improve land management.

Deleted: site

Deleted: Laub et al. (2023b) studied the Embu trial along with three other trials in different regions of Kenya.

Deleted: contents

Deleted: at

Deleted: the

the site was covered by tropical evergreen forest before conversion to a low-intensity agricultural area. Before the experiment started, the site was terraced to ensure that the plots were level. Additional details about the site can be found in previously published studies (Chivenge et al., 2009, 2011a; Gentile et al., 2008; Laub et al., 2023b).

2.1.2 Experimental design

The field experiment has a split-plot design with three replications. Four nutrient management treatments were selected for the present study, involving the application of three organic residues with varying qualities (at a rate of $4.0 \text{ t C ha}^{-1} \text{ yr}^{-1}$) and a control treatment (*Control*; no organic residues were added). These application rates are higher than those typically used by local farmers, which allows for a more pronounced assessment of the potential impacts on SOC. These organic residues were applied once annually before the long rainy season and were incorporated in the soil through manual tilling to a depth of approximately 15 cm. Each plot was further divided into two subplots, with one subplot receiving an application of $120 \text{ kg mineral N ha}^{-1}$ per growing season ($+N$ treatment), while the other others did not receive any mineral nitrogen input ($-N$ treatment). Mineral N (CaNH_4NO_3) was applied twice during each growing season: an initial 40 kg N ha^{-1} at planting and the remaining 80 kg N ha^{-1} approximately six weeks later as a top dressing. In every growing season, all plots were subjected to a uniform application of 60 kg P ha^{-1} in the form of triple superphosphate and 60 kg K ha^{-1} as muriate of potash at planting. The organic resources utilized in the present study exhibited variations in quality, characterized by differences in nitrogen, lignin, and polyphenol contents (Table 1). These included pruned leaves, including stems with a thickness of less than 2 cm, from *Tithonia diversifolia* (*Tithonia*; high quality with rapid turnover), stover of *Zea mays* (*Stover*; low quality and fast turnover), and locally available farmyard manure (*Manure*; intermediate to high quality with intermediate turnover). The relevant properties of the different organic inputs are given in (Table 1). After harvest, any remaining parts of the maize plant were removed from the plots. Therefore, the only OC inputs left from the maize crop came from roots and root exudates.

This study focuses on the effect of organic residues as compared to the control. The treatments include both the $-N$ and $+N$ alternatives combined, and are referred to here as *Manure* (i.e., farmyard manure), *Tithonia* (i.e., residues of *Tithonia diversifolia*), *Stover* (residues of maize stover), and *Control* (i.e., no organic residue added). Whenever we refer to a treatment specifically with or without mineral N, a $-N$ or a $+N$ symbol is added to the treatment name. For example, *Manure-N* represents application of manure without mineral N, *Control+N* means no that only mineral fertilizer was added, while *Tithonia* includes both *Tithonia+N* and *Tithonia-N* treatments.

Table 1: Mean dry matter properties of the organic resources applied (source: Laub et al., 2023b)

Measured property	Manure	Stover	Tithonia
C (g kg^{-1})	234	397	345
N (g kg^{-1})	18.1	7.2	33.2
C:N ratio	12.3	58.7	12.4
P (g kg^{-1})	3.1	0.4	2.3

K (g kg ⁻¹)	19.4	9	37.2
Lignin (g kg ⁻¹)	198	48	90
Polyphenols (g kg ⁻¹)	7.8	11.3	19
Lignin/N ratio	6.9	6.2	2.6
Kg N in 4.0 t C ha ⁻¹ , -N [+N]	324 [564]	68 [308]	323 [563]

205 **2.1.3 Sample collection**

The field campaign took place in February 2021, 19 years after the establishment of the long-term field trial and after 38 growing seasons of maize. Undisturbed soil samples were collected down to 70 cm using a gauge auger, one core was taken near the middle of each sampled plot. Given the auger's 50 cm length, samples were obtained in two stages: the first sample down to 50 cm, followed by a second sample from 50 to 70 cm using an extension for the auger. Subsections of 5 cm were obtained, weighed and a 5 g subsample of each sample was immediately placed in an air-tight recipient to determine the water content and calculate the bulk density (see Sect. 2.2.1). The samples were sieved through an 8 mm sieve on the day of collection and air-dried in Kenya. In the lab, ca. one week later, the samples were sieved through a 2 mm sieve and the large organic residues (> 2 mm) that passed through the sieve were removed by hand. The samples were dried at 45°C. Using a soil splitter, samples were separated homogeneously in two parts, with one part being ground and the other part stored without grinding.

210

The stocks of SOC were determined for the following depth layers: 0-5 cm, 5-10 cm, 15-20 cm, 25-30 cm, 35-40 cm, 45-50 cm, 55-60 cm, 65-70 cm. Each treatment (including the control) had two variations (with and without mineral nitrogen input), each variation was replicated 3 times (i.e., 4 treatments x 2 variations x 3 replicates = 24 plots), and 8 depth layers were selected, resulting in 192 samples.

215

2.2 Laboratory analyses

220 **2.2.1 Bulk density**

The dried weight of each subsection and the inside volume of the auger were used to calculate the bulk density. Because of disturbances to samples collected at the 45-50 cm and 50-55 cm sections during sampling, the 45-50 cm and 50-55 cm section's bulk density were replaced by the mean value of the 40-45 cm and 55-60 cm layers from the same profile.

225 **2.2.2 Soil organic carbon stocks and δ¹³C values**

Before the analysis of OC and nitrogen, large organic matter particles were removed with tweezers after the sample was spread on a smooth surface. Samples were not treated with HCl before OC analysis, as the pH of all samples was ≤ 6 and none of 12 samples to which HCl was added showed any reaction, indicating the absence of carbonates. Afterwards, 11 to 15 mg of dried and grinded soil was weighed in a tin cup. The samples were analyzed using an elemental analyser-isotope ratio mass spectrometer (Thermo Flash HT/EA or CE 1110 coupled to a Delta V Advantage), to obtain the OC and nitrogen concentrations and δ¹³C values. The OC concentration was subsequently converted to stocks using the measured bulk density. While bulk

230

density was available for each depth layer in 5 cm increments, the OC concentration was only measured on selected layers (i.e., 0-5 cm, 5-10 cm, 15-20 cm, 25-30 cm, 35-40 cm, 45-50 cm, 55-60 cm, 65-70 cm). To calculate SOC stocks over the whole profile, depth layers without OC concentration measurements were first interpolated using available data from adjacent layers within each profile. This interpolation was performed using the *approxfun* function in R.

2.2.3 $\Delta^{14}\text{C}$ values

Homogenized soil samples were fumigated in silver capsules (Elementar) with HCl (37%) for 72h to remove carbonate (Komada et al., 2008). Fumigated samples were neutralized for 24 hr at 60°C over solid NaOH to remove residual acid. The samples were wrapped in an 8 × 8 × 15 mm tinfoil boat (Elementar) and pressed prior to analysis. Fumigated samples were analyzed for $\Delta^{14}\text{C}$ using a Gas Ion Source of Micadas system at ETH Zurich. Samples were normalized using oxalic acid II (NIST SRM4990C). The Measured $\Delta^{14}\text{C}$ / $\delta^{13}\text{C}$ ratios are reported as $F^{14}\text{C}$, as established by Stuiver and Polach in 1977 (Reimer et al., 2004).

2.2.4 Effective cation exchange capacity

Before the analysis of the effective cation exchange capacity (CEC), organic matter macro particles were removed from the soil sample. Subsequently, 2 g of dried and milled soil was measured and placed in a centrifugation tube, and 25 ml of 0.01M BaCl_2 was added. The sample was then agitated on a reciprocal shaker at 150 rpm for 2 hours. Next, the sample was centrifuged for 10 minutes at 2500 rpm. The supernatant was then filtered through a Whatman 41 filter. After filtration, 1 ml of the solution was diluted in 4 ml of water (1/5 solution). The diluted solution was analyzed using ICP-OES (G8010A Agilent 5100 SVDV ICP-OES, Parent Asset SYS-10-5100). The concentrations of Al, Ca, K, Mg, Mn, and Na in the solution were subsequently converted to cmol per kg of soil before calculating the effective cation exchange capacity by summing the concentrations of all the aforementioned cations (Hendershot and Duquette, 1986). CECeff was measured for the 0-5 cm, 35-40 cm, and 65-70 cm depth layers.

2.2.5 Soil texture

To analyze soil texture, 200 to 300 mg of 2 mm-sieved soil was weighed in a glass jar and treated 4 times with 10 ml H_2O_2 in a warm water bath (ca. 50 °C) to completely remove organic matter. The samples were then transferred into a plastic tube and 7 ml of 10 % $(\text{NaPO}_3)_6$ was added and shaken overnight to completely disperse the soil minerals. The particle size was analyzed using a particle size analyzer (LS13320 Beckman coulter) to obtain the fraction clay (vol% < 2 μm), silt (vol% 2 – 53 μm) and sand (vol% > 53 μm). Texture was measured for the 0-5 cm, 45-50 cm, and 65-70 cm depth layers.

2.2.6 Soil pH

The soil pH was determined using a pH meter (Thermo Scientific™ Eutech™ “150 Series Waterproof Handheld Meters”). To prepare the soil samples, 10 g of dried soil was placed in a tube and 25 ml of deionized water was added. The tubes were shaken on a reciprocal shaker at 150 rpm for 2 hours. Afterward, the slurry was allowed to settle for 24 hours. Prior to measurement, the pH electrode was calibrated using buffer solutions with pH values of 7 and 4. Care was taken to ensure that the diaphragm of the electrode did not come into contact with the soil particles. The pH was considered stable when the measured pH value remained constant within 0.02 units over a period of 5 seconds. The soil pH was measured for the 5-10 cm, 15-20 cm, 35-40 cm, and 55-60 cm depth layers.

2.3 Statistical analysis

2.3.1 Linear mixed effect models of the treatments effect over the depth profile

To test which measured variables are significantly affecting SOC stocks along the depth profile, a linear mixed effect model was constructed. Linear mixed effects models are linear models that allow to separate an independent variable into fixed effects and random effects. They are particularly useful when more than one measurement is made on a given statistical unit. It is therefore suitable for this work, since there was more than one sample for each site. No outlier were detected, using the boxplot method with the R function “identify_outliers” (Kassambara, 2022). Normality of the data was confirmed visually (Fig. S1a), and homogeneity of variance was verified, showing no evident relationship between residuals and fitted values (Fig. S1b). The linear mixed effects model was built with the “lmer” function of the package “lmeTest” (Kuznetsova et al., 2017). To avoid overfitting and ensure the robustness of the results, interactions were excluded. This approach simplified the model, focusing on the main effects and accounting for variability through random effects, thereby reducing the risk of capturing noise rather than true patterns. We evaluated the robustness of the model by fitting six different linear mixed effects models using the lmer function. Each model included OC stocks as the response variable, the sampling plot as a random effect and different combinations of fixed effects predictor variables: model 1 (depth, organic residues, mineral fertilizer, silt content, sand content), model 2 (depth, silt content, sand content), model 3 (depth, organic residues), model 4 (depth, mineral fertilizer), model 5 (organic fertilizer), model 6 (mineral fertilizer). Other measured parameters such as pH and CECeff were not considered in this model, because they correlated with the application of all three types of organic residues (especially manure) (Fig. 1). We then compared these models with an ANOVA to determine the best-fitting model based on AIC, BIC, log-likelihood, and deviance values (Supplementary table 1). To further test the effect of organic residue treatment application in the deeper soil layers, the same analysis was performed on a subset containing only subsoil layers (Supplementary table 2).

2.3.2 Two-sample t-tests between treatments

To evaluate the impact of organic residue treatments on SOC stocks along the soil profile, we examined SOC stocks at different depth intervals. This analysis involved conducting a two-sample t-test for each organic residue treatment to assess the difference in SOC stocks between the treatment and the control across various depth layers. The depth layers examined

300 included the OC stocks of the entire profile (0-70 cm), of two combined topsoil or combined subsoil depth layers (0-30 cm and 30-70 cm), and of each individual 5 cm depth layer (0-5 cm, 5-10 cm, 15-20 cm, 25-30 cm, 35-40 cm, 45-50 cm, 55-60 cm, and 65-70 cm). To increase the statistical power, statistical analyses were performed by combining the +N and -N sub-treatments of the same treatment, focusing therefore only on the effect of organic residues application on SOC stocks. This approach was justified by the results of the linear mixed effect model, which indicated that mineral fertilizer was not significantly affecting SOC over the depth profile.

305 Due to the violation of homogeneity of variance for some depth layers, OC stock data were log-transformed. An ANOVA was then performed on each depth layer, after normality and homogeneity assumptions of the log-transformed data were confirmed (Supplementary table 5). Given the limited data from the long-term field trial and increasing variability with depth, we opted for t-tests over Tukey's test to identify which OR treatment had significantly higher OC stocks than the control treatment and at which depth layers. Performing Tukey's test on the ANOVA results would be more affected by the limited statistical power. 310 Our choice was supported by the fact that when the statistical significance of the t-test was not achieved, the statistical power (see section 2.3.3) was below always 80% (Figure 4). Therefore, we argue that type I error inflation due to the number of t-tests performed does not apply in our context, and we are more likely to encounter type II errors.

315 We conducted a Welch two-sample t-tests on log-transformed OC stocks between the control and each organic residue treatment at every depth layer mentioned earlier. This analysis was performed using the "t.test" function in R, after normality and homogeneity assumptions of the log-transformed data were confirmed (Supplementary table 4, Figure S9). Treatments were considered significantly different than the control when their p -value was ≤ 0.05 . In the analysis of each 5 cm depth layer, if one layer showed a p -value greater than 0.05, while the two layers above and the two layers below showed significant differences compared to the control, the lack of significance in the layer above was attributed to sampling variability. 320 Consequently, the treatment was considered to affect OC stocks down to the deepest layer where a significant deviation from the control was observed. This was only the case for the *Manure* treatment at depth 25-30 cm.

325 The impact of organic residue treatments on soil pH and effective cation exchange capacity (CEC_{eff}) was further assessed using a Welch two-sample t-tests, similar to the approach used for the analysis of SOC stocks. These tests compared organic residue treatments with the control treatment, with all treatments including combined +N and -N sub-treatments, at each individual depth layer where these parameters were measured.

330 To validate the findings of the linear mixed model regarding the effect of mineral fertilizer, we compared the difference between the +N and -N versions of each organic residue treatment (as well as the control). This comparison was performed using a Welch two-sample t-tests across the same depth layers used in the analysis of the effects of organic residues (i.e., 0-70 cm, 0-30 cm, 30-70 cm, 0-5 cm, 5-10 cm, 15-20 cm, 25-30 cm, 35-40 cm, 45-50 cm, 55-60 cm, and 65-70 cm). The results

335 indicated no significant difference for any treatment at any depth layer. This supports the methodology of combining +N and
-N treatments for the analysis of the effect of organic residues.

335 **2.3.3 Power analysis of statistics on 5 cm depth layers**

To determine whether the number of replicates was optimal for testing the difference between OR application and control, a
power analysis was performed using the R function “pwr.t.test” (Champely, 2006). The analysis was performed twice and both
times the Cohen's d effect size measure (i.e., standardized difference between two means) was used as an input, along with a
340 significance level of $\alpha = 0.05$. The first analysis determined the actual power, based on the number of samples used in each
statistical analysis we performed. The second analysis calculated the number of samples needed to reach a power of 80% (i.e.,
i.e., the probability that the test correctly rejects the null hypothesis when the alternative hypothesis is true) and therefore
power = 0.8 was used. This was performed for the analyses done at each individual 5 cm depth layer (0-5 cm, 5-10 cm, 15-20
cm, 25-30 cm, 35-40 cm, 45-50 cm, 55-60 cm, and 65-70 cm) to assess if our study design had sufficient statistical power to
345 determine the exact depth layer down to which the ORs were significantly affecting OC stocks.

3 Results

3.1 Soil characteristics

The incorporation of organic residues, particularly manure, resulted in overall higher pH_{H2O} values compared to the other
treatments, and led to higher effective cation exchange capacity (CECeff) (Fig. 1). Although the differences were not always
350 significant between the treatments at the same depth, the tendency was consistent across all depth layers, suggesting that the
effect of organic matter application on soil properties may not be limited to the topsoil. Variability in soil pH_{H2O} and CECeff
between treatments was greater in the topsoil, ranging from 4.2 to 6.6 for pH and 5.3 to 19.0 (meq/100g) across all treatments.
Mineral N fertiliser application had a non-significant, but consistently negative effect on pH and CECeff in the control
treatment (Fig. S3). Soil texture was uniform across all treatments with average clay, silt and sand proportions of 70%, 26.6%
355 and 1.6% respectively (Fig. S4).





Figure 1: Boxplots of (a) effective cation exchange capacity (CEC_{eff}) and (b) pH_{H2O} for organic residue treatments at different depth layers. Significant difference between treatments at a similar depth are highlighted with a “*”.

360 3.2 Effect of organic residues on SOC stocks

3.2.1 Effect of organic residue inputs on SOC stocks in the top- and subsoil

When calculating the OC stocks for the topsoil and subsoil separately we found that, on average for all treatments, the subsoil layer (30-70 cm depth) contained $48.5\% \pm 1.7\%$ of the total stocks of the entire 0-70 cm soil profile (Fig. 2). Considering the whole measured profile, every treatment receiving organic amendments (combined for +/- N) led to OC stocks that were significantly higher than OC stocks of the control treatment. It is noted that this difference was significant mainly because of the high differences in the topsoil layer. Only the *Manure* treatment was significantly different from *Control* in the subsoil (i.e., the 30-70 cm depth layer)



370 Figure 2: Total stocks of organic carbon for (a) the 0-70 cm soil layer, (b) the topsoil (0-30 cm) and (c) the subsoil (30-70
375 cm). Significant difference between organic residues treatments and control are highlighted with a “*”.

3.2.2 Effect of organic residue inputs on SOC stocks along the soil profile

The difference in SOC stocks between the organic residue treatments and the control was most pronounced in the topsoil and gradually decreased with depth (Fig. 3). According to the ANOVA performed on all the different statistical models, the model including all depth and the OR (i.e., model 1) had the lowest AIC (596.05) and BIC (618.56), the highest log-likelihood (-291.03) and the lowest deviance (582.05), suggesting it was the best-fitting model (Supplementary Table 1). The R^2 of this model was 0.76, representing the variation in OC levels that this model explained, with 0.65 attributed to fixed effects and 0.11 to random effects. According to the model, depth and the application of manure and maize stover were significantly impacting OC stocks, but not the application of *Tithonia* (Table 2). The same analysis performed on a subset containing only subsoil data yielded similar results with the best fitting model including only depth and organic residues as fixed effect and having a R^2 of 0.64. According to the best-fit model for subsoils, manure and depth were the only significant variables affecting OC stocks (Supplementary Table 2). The application of mineral fertilizer was not found to significantly affect OC stocks by any model.

Deleted: According to the model with the best fit for the subsoil, besides depth, only manure was significantly affecting OC stocks

Table 2: Summary statistics of the linear mixed effects model for soil organic carbon stocks on all measured depth layers

	estimate	Standard error	p value	
(Intercept)	8.98	0.41	< 2e-16	*
Depth	-0.07	0.00	< 2e-16	*
Manure	2.04	0.53	0.001	*
Stover	1.59	0.53	0.007	*
Tithonia	0.92	0.53	0.095	

390

395

400

In the topsoil (i.e., within the 0-30 cm depth layer), OC stocks ranged from 5.5 to 13.1 t OC ha⁻¹ per 5 cm depth layer (Fig. 3). The effect of organic residues on OC stocks per 5 cm layer revealed down to which depth the different organic residues had a significant effect on SOC stocks (Fig. 3). The ANOVA and the comparison of each organic residue treatment with the control treatment per 5 cm layer revealed that applying organic residues had a significant effect on SOC stocks down to 60 cm (Table 3, Fig. 3, supplementary table 3). In the top 0-5 cm, *Manure* had the highest OC stocks with 11.2 ± 1.6 t ha⁻¹ and *Control* had a significantly lower OC stock with 8.3 ± 0.4 t ha⁻¹. ~~*Manure* significantly impacted OC stocks down to 60cm.~~ At this depth layer (i.e., 55-60 cm), *Manure* had an OC stock of 7.2 ± 1.8 t ha⁻¹ and *Control* of 5.2 ± 0.6 t ha⁻¹. *Stover* was significantly different from *Control* down to 40 cm. The OC stocks of the *Tithonia* treatment were only significantly different from *Control* in the top 5-20 cm.

Table 3: Summary statistics of [the ANOVA testing the effect of all OR treatments](#) on OC stocks across depth layers

Depth layer	Sum of Squares	P-Value	Significance
0-5 cm	0.27	0.01	*
5-10 cm	0.24	0.01	*
15-20 cm	0.29	≤ 0.01	*
25-30 cm	0.26	0.03	*
35-40 cm	0.33	0.02	*
45-50 cm	0.34	0.05	*
55-60 cm	0.33	0.05	*
65-70 cm	0.15	0.52	

Deleted: *Manure*

Deleted: was affecting OC stocks the deepest, with a significant effect down to 60 cm

Formatted: Font: *Italic*

405





Figure 3: Profile of SOC stocks for each measured 5 cm layer (not cumulative). The vertical bars on the left side of the graph show the depth layers over which the SOC stocks for the different treatments are significantly different from the control treatment.

410

The power analysis indicated a high risk of Type II errors (i.e., incorrectly rejecting that there is a significant difference), particularly in soil layers below 20 cm. For each OR treatment and at each depth layer, the power of the test fell below 80% when no significant difference was detected between the OR and the control (Fig. 4a). This suggests that for these statistical tests, there is a greater than 20% chance of failing to detect a true effect. This implies that the absence of a significant difference between *Control* and either the *Stover* or *Tithonia* treatments in deeper soil layers should be interpreted with care. This is possibly due to the increased data variability with increasing depth. The deeper down the soil profile, the more samples are thus needed to detect a true effect of the treatment (Fig. 4b).



420 Figure 4: (a) Power of statistical analysis at each depth layer and for each organic residue treatment and at each 5 cm
 425 depth layer. The 80% power threshold is indicated by the dashed line. (b) The amount of sample samples required to reach a
 statistical power of 80% for each organic residue treatment and each 5 cm depth layer. The actual number of sample ($n =$
 12) is indicated by the dashed line.

3.2.3 Effect of treatment on SOC stocks considering +N and -N treatments separately

The SOC stock in the 0-30 cm layer was highest for *Manure-N* and lowest for *Control+N* with values of 60.9 ± 8.0 t OC ha⁻¹ and 44.1 ± 2.6 t OC ha⁻¹, respectively (Fig. 5a). Similarly, over the measured profile (0-70 cm depth), *Manure-N* showed the highest OC stocks, while *Control+N* showed the lowest, with mean values of 124.4 ± 24.5 t OC ha⁻¹ and 85.9 ± 15.7 t OC ha⁻¹ respectively (Fig. 5a). Therefore, the *Manure-N* treatment maintained 38.5 ± 8.8 t OC ha⁻¹ more SOC in the 0-70 cm depth layer compared to the *Control+N* treatment. Adding mineral nitrogen resulted in consistently lower OC stock across the soil profile for *Control*, but not for *Manure*, *Stover* and *Tithonia* (Fig. 5b). However, the difference between +N and -N variation of each treatment was not statistically significant for any depth layer.

Deleted: soil organic carbon

Deleted: profile

Formatted: Font: Italic

Formatted: Superscript

Deleted: This means that after 38 growing seasons over 19 years, and considering the top 70 cm of the soil, *Manure-N*, as the most effective treatment to limit SOC losses, could maintain 38.5 ± 8.8 t OC ha⁻¹ more SOC compared to the least efficient treatment, *Control+N*....

Formatted: Font: Italic




Figure 5: (a) Cumulative stocks of organic carbon along the depth including -N and +N treatments and (b) difference in OC stocks between +N and -N treatment at each measured 5 cm depth layer (not cumulative); values below 0 indicate a loss in OC when mineral nitrogen was applied. The differences in SOC stocks between -N and +N treatments were never significant.

3.3 C/N ratio and stable isotopes of organic carbon

The C/N ratio of organic matter had values between 10 and 12 along the soil profiles and decreased with depth below 30 cm (Fig. 6a). The C/N ratio across the depth profiles was not significantly different between treatments, but below 30 cm, it tended to be lower in the *Control* treatment compared to all organic residue treatments. This difference was primarily due to the consistently lower C/N ratio of the *Control+N* treatment at intermediate depths (Fig. S5).

Depth profiles of the $\delta^{13}\text{C}$ value of SOC show that the largest differences between the treatments are in the top 20 cm of the soil (Fig. 6b). The lowest values are for the *Manure* treatment (-19.5 – -20 ‰), while the *Stover* and *Control* treatments have the highest values (ca. -18 ‰). The $\delta^{13}\text{C}$ value for the *Tithonia* treatment is intermediate (ca. -18.5 ‰). Below this depth, all treatments have similar $\delta^{13}\text{C}$ values in the range of -17.5 – -18.5 ‰, with a tendency for lower values in the *Manure* treatment. The $\delta^{13}\text{C}$ value of SOC of the *Stover* treatment is within the range of *Control* all along the profile. Additionally, the $\delta^{13}\text{C}$ signatures of all treatments at depths below 50 cm are similar, suggesting no influence of organic amendments on SOC below this depth.

As observed in the depth profile of radiocarbon ($F^{14}C$), the OC becomes older with depth, i.e., the lower the $F^{14}C$ value (Fig. 6c). However, below the surface layer (i.e., the 0-5 cm depth layer) the $F^{14}C$ value are higher (i.e., the OC is younger) for all three treatments receiving organic amendments, compared to the control treatment, suggesting that organic residue input affected OC below the layers where it is applied. The higher $F^{14}C$ values for the *Manure* treatment in the deepest layer suggest that this treatment had the largest impact on subsoil OC.




Figure 6: Soil profiles of (a) C/N ratio, (b) $\delta^{13}C$ and (c) $F^{14}C$ of the different organic residue treatments (for $F^{14}C$, only 6 plots were measured and there were two replicates for Control and for Manure)

4 Discussion

Our results show that at the studied tropical field trial, about 50% of the SOC stock measured down to 70 cm was located below 30 cm, showing the importance of subsoil OC stocks. Moreover, the addition of farmyard manure, *Tithonia* residues and maize stover all had a positive impact on SOC stocks, limiting the loss of organic matter compared to the control treatment. Although this effect was greatest in the top 20 cm, it could be observed down to 40 cm and 60 cm for the *Stover* and *Manure* treatments, respectively. The *Manure* treatment led to the highest OC stocks down to 70 cm. However, a power analysis indicated that more samples are required to accurately assess whether the effect of organic residues on OC stocks in the subsoil is statistically significant or not.

4.1 Application of manure affects OC the subsoil

Our study shows, although the calculated statistical power indicated that more samples are needed to increase confidence, that the application of manure affected SOC stocks in the subsoil in a tropical long-term field trial. Among all organic residues applied, only *Manure* had OC stocks that were significantly larger than *Control* in the 30-70 cm depth layer (Fig. 2), while the

485 application of manure was significantly impacting OC stocks down to 60 cm (Fig. 3). This shows that organic residues can
also influence SOC in the subsoil. In addition, the $\Delta^{14}\text{C}$ values along the soil profile suggest that the addition of Manure led to
the addition of fresh OC down to a depth of 65-70 cm. This wasn't mirrored in the $\delta^{13}\text{C}$ values for the deeper layers, suggesting
that only a small portion of the added organic amendments made their way downwards and suggesting that larger SOC stocks
under *manure* are due to lower SOC losses compared to the *control*. However, the application of manure resulted in a
490 significant increase in pH and CECeff down to 40 cm compared to the control treatment. These results indicate that this
treatment can make the subsoil environment more favourable to support plant growth.

Our findings are in line with Leuther et al. (2022), who found that high levels of manure application in a long-term (36 years)
field study in Germany increased OC stocks at 35-45 cm depth, and by Anandakumar et al. (2022), who observed a significant
increase in OC stock between 25 and 50 cm depth after applying manure in a semi-arid tropical Alfisol in India. A known
495 mechanism for OC transport across soil profile is in the form of dissolved organic carbon (DOC), which moves downward
through cycles of sorption and desorption (Kaiser and Kalbitz, 2012b; Uselman et al., 2007). In a long term in China, Liu et
al. (2013) observed that manure applications increased the amount of DOC down to 60 cm as compared to straw residues
application, supporting that manure may percolate further down than plant residues. However, a study from Sommer et al.
(2018) revealed that manure was able to slow steady losses of SOC from topsoils of Kenyan Ferralsols over time, but only
500 when at least $4.0 \text{ t C ha}^{-1} \text{ yr}^{-1}$ was applied. While this is the same amount as what was applied in our study, it is the largest
amount compared to common range of 0.9 to $4 \text{ t C ha}^{-1} \text{ yr}^{-1}$ that are applied by Kenyan farmers who own cattle (Tittonell et al.,
2008). Therefore, the impact of using manure may not be as great in the deeper layers when considering a scenario where
common rates of manure are applied. For instance, Laub et al. (2023) did not observe a significant effect on topsoil OC when
the annual application of manure was $1.2 \text{ t C ha}^{-1} \text{ yr}^{-1}$.

505 **4.2 The effect of *Tithonia diversifolia* and maize stover residues on OC could only be detected in the topsoil**
In contrast to the *Manure* treatment, our results showed that *Tithonia* and *Stover* did not affect subsoil OC stocks significantly
in the 30-70 cm layer (Fig. 2). Also, no significant differences below 20 cm and 40 cm for *Tithonia* and *Stover*, respectively,
were observed in the individual 5 cm depth layers (Fig. 3). However, for both treatments, the statistical power was below 80%
510 at every depth layer where the treatments were not significantly different from the control. This suggests that the sample size
was insufficient to prevent Type II errors in the subsoil (Fig. 4a and 4b). One must therefore be cautious when interpreting
these non-significant differences. Moreover, both *Tithonia* and *Stover* treatments showed signs of younger OC below 30 cm
depth compared to the *Control*, as indicated by the higher $F^{14}\text{C}$ values (Fig. 6c). Nevertheless, these effects in the 65-70 cm
depth layer were less pronounced compared to *Manure*. This suggest that the effect of stover and tithonia residues can be
515 traced below 30 cm, but that this effect is not strong enough to detect a significant impact on subsoil OC stocks. Further studies
with more samples are thus needed to conclusively reject the hypothesis that application of *Tithonia* and *Stover* affect the
subsoil environment. For instance, all OR treatments had significantly higher pH than the *Control* down to 40 cm (Fig. 1). Our

Deleted: {

Deleted: ,

Deleted: Also

Deleted: outside the context of long-term field trials

525 observations differ from observations from Córdova et al. (2018), who found that high-quality residues led to a higher accumulation of OC in the soil compared to low-quality residues, because *Stover* had higher OC stocks than *Tithonia* in the topsoil and in the subsoil. Nevertheless, our results are in line with the meta-analysis results of Gross and Glaser (2021) who found manure to be more effective in stabilizing SOC than green manure. Their meta-analysis was dominated by temperate soils, but their results are similar to the ones found here for a tropical soil.

530 The $\delta^{13}\text{C}$ signatures of OC in the topsoil under the *Stover* and *Control* treatments were similar. However, as maize is a C4 plant, it would be expected that if more maize stover residues were incorporated in the soil compared to the control treatment, the $\delta^{13}\text{C}$ signature of SOC would be highest under the stover treatment (Tieszen, 1991). As the *Stover* treatment received 4 t C ha⁻¹ yr⁻¹ more maize derived residues (i.e., stover) compared to the *Control* treatment, the similar $\delta^{13}\text{C}$ value in the topsoil suggests that the additional organic matter inputs for the *Stover* treatment did not results in additional SOC sequestration compared to the control treatment. However, as topsoil OC stocks were significantly larger for the *Stover* compared to the *Control* treatment, this indicates that the larger SOC stocks were due to lower losses of initial SOC in the *Stover* treatment, 535 and not to additional SOC sequestration.

540 Concerning the *Tithonia* treatment, the lower $\delta^{13}\text{C}$ values of topsoil OC and the significantly larger topsoil SOC stocks compared to the *Control* treatment, suggest that the latter is at least partly caused by the incorporation of OC originating from the *Tithonia* residues. The limited effect of both the *Stover* and *Tithonia* treatments on topsoil $\delta^{13}\text{C}$ values suggests that both residues are being quickly mineralised upon application, which explains why they do not compensate the steady loss of OC in both treatments.

4.3 Mineral nitrogen does not significantly affect SOC stocks

545 The linear mixed effect model did not select mineral N as a parameter having a significant impact on the SOC stocks. This corroborates the meta-analysis conducted by Gram et al. (2020) on integrated soil fertility management in sub-Saharan Africa, which showed that mineral N fertilizer had no significant impact on topsoil SOC concentrations. However, the interaction between SOC and mineral fertilizer is complex and two global meta-analyses on the effect of mineral N on agricultural topsoil, conducted by Ladha et al. (2011) and Liu et al. (2023), suggested that the addition of mineral N fertilizer typically has a positive impact on SOC content. One reason for this impact is that as mineral fertilizer tends to enhance biomass production, it commonly leads to an increase in the quantity of plant residues returned to the soil, thereby positively impacting the soil 550 organic matter content (Geisseler and Scow, 2014). This does not apply at our study site in Embu, as the yield was not ~~very~~ responsive to N fertilization and crop residues were removed from the field after harvest (Laub et al., 2023a). Also, while Liu et al. (2023) emphasize the significance of specific climatic and environmental factors on the effects of mineral fertilizer application on SOC, neither of the two meta-analyses included studies from eastern Africa (Ladha et al., 2011b; Liu et al., 2023). For instance, in a study conducted in Kenya, Ndung'u et al. (2021) observed a decrease in topsoil OC concentration

Deleted: much

after the application of mineral nitrogen to Ferralsols, while Laub et al. (2023b) did not observe a consistent effect of mineral N on SOC stocks in four long-term agricultural field trials in central and western Kenya.

The *Control* treatment had the largest difference in OC stocks between the *+N* and the *-N* treatments when considering OC stock over the whole 0-70 cm depth profile, i.e. *Control+N* had a consistently lower OC stock than *Control-N* below 10 cm (Fig. 5a). While this difference was not significant, this suggests that the sole application of mineral nitrogen increased the rate of OC losses at the study site. One possible explanation for this observation is that the application of mineral fertilizer stimulates the activity of fast-growing microorganisms that can lead to a priming effect, accelerating the decomposition of organic matter in the soil (Chen et al., 2014). It is also possible that mineral nitrogen alone leads to increased rates of OC loss because it increases the degradation of below-ground plant residues; for instance mineral fertilizer was shown to increase the degradation of young C4 lignin (Hofmann et al., 2009).

4.4 The potential of organic inputs to reduce SOC losses in tropical agroecosystems

There is a motivation to adopt agricultural practices to reduce the atmospheric CO₂ concentration by increasing SOC storage (Bossio et al., 2020; Minasny et al., 2017). According to Corbeels et al. (2019), there is a potential to achieve this in Sub-Saharan Africa, not primarily as a measure to mitigate climate change but to improve crop productivity. However, the low F¹⁴C values measured in each treatment, especially in the subsoil, indicate that very little of the freshly added organic matter remains in the soil, suggesting that even the application of significant amounts of OR would not contribute substantially to climate change mitigation. This aligns with the findings of Sierra et al. (2024) who demonstrated that only a minimal amount of new OC inputs make their way through the profile and have the potential to remain stable for periods extending beyond 50 years. Also, results from Laub et al. (2023b) show that all treatments of the long-term trial in Embu have lost topsoil OC during the first ca. 20 years of the field trials, despite differences in magnitude between treatments. In accordance with the results from Laub et al. (2023b) for topsoils (0-15 cm), we show that after 19 years, OC stocks down to 70 cm was largest for *Manure-N* and lowest for *Control+N* with 124 t C ha⁻¹ and 85 t C ha⁻¹ respectively. This is a difference of 39 t C ha⁻¹, which represent about 50% higher OC stock in *Manure-N* than in *Control+N* stocks. Therefore, although the OC storage potential in soils that have been recently deforested, with mainly low activity clay minerals, is low, it is possible to optimize agricultural nutrient management to minimize SOC losses. Furthermore, this could be combined with other appropriate agricultural practices, such as increasing crop diversity and rotating crops, which have the potential to improve OC stocks and positively impact yields (Yang et al., 2024). Combining these practices with the application of manure could lead to the stabilization of SOC stocks over time.

5 Conclusion

Our study shows that organic amendments have an influence on the SOC stocks throughout the soil profile in a tropical arable soil. Using soil samples down to 70 cm from a long-term field trial in central Kenya, we show that the application of 4 t C ha⁻¹

Deleted: content

¹ yr⁻¹ in the form of manure led to significantly larger SOC stocks down to 60 cm compared to a control treatment. In contrast, the application of residues of *Tithonia diversifolia* and maize stover impacted SOC stocks to a lower extent, while this effect was limited to the topsoil. Due to insufficient statistical power for the subsoil for these treatments, further research with larger sample sizes is necessary to draw definitive conclusions on their effects. The application of mineral nitrogen did not contribute positively to the effect of organic residues on SOC stocks, while having a negative effect in the control treatment. Our study indicates that, for the studied tropical soils, the application of manure is the most appropriate nutrient management strategy to limit SOC losses in recently established croplands, and that agricultural practices affect soils deeper than the 0-30 cm layer usually studied. Therefore, we argue that appropriate agricultural management has the potential to partially mitigate the effects of deforestation on SOC loss while improving soil quality for food production and that studying soil deeper than 30 cm can improve our understanding of agroecosystems.

600 **Code availability**

The code can be provided upon request.

605 **Data availability**

The data used in this study are open access and available at <https://figshare.com/s/c2f2787b7a56ef7ad656>.

Authors Contribution

610 Conception and design of the field campaign: MVdB and JS.
Design of the experiment: MVdB and JS.
Performing the experiments: CRM
Analysis and interpretation of the data: CRM, MVdB.
615 Writing the manuscript: CRM and MVdB with inputs from all co-authors.

Competing interests

The contact author has declared that none of the authors has any competing interests.

620 **Acknowledgements**

We thank Britta Jahn-Humphrey for help with soil processing, and ChatGPT, an AI language model developed by OpenAI, for assistance in the writing process.

625 **Financial support**

This work was supported by the Swiss National Science Foundation SNSF (Ambizione project number PZ00P2_193617 granted to MVdB).

630 **References**

- Adams, A. M., Gillespie, A. W., Dhillon, G. S., Kar, G., Minielly, C., Koala, S., Ouattara, B., Kimaro, A. A., Bationo, A., Schoenau, J. J., and Peak, D.: Long-term effects of integrated soil fertility management practices on soil chemical properties in the Sahel, *Geoderma*, 366, 114207, <https://doi.org/10.1016/j.geoderma.2020.114207>, 2020.
- 635 Anandakumar, S., Bakhoun, N., Chinnadurai, C., Malarkodi, M., Arulmozhiselvan, K., Karthikeyan, S., and Balachandar, D.: Impact of long-term nutrient management on sequestration and dynamics of soil organic carbon in a semi-arid tropical Alfisol of India, *Applied Soil Ecology*, 177, 104549, <https://doi.org/10.1016/j.apsoil.2022.104549>, 2022.
- Balesdent, J., Mariotti, A., and Guillet, B.: Natural ¹³C abundance as a tracer for studies of soil organic matter dynamics, *Soil Biology and Biochemistry*, 19, 25–30, [https://doi.org/10.1016/0038-0717\(87\)90120-9](https://doi.org/10.1016/0038-0717(87)90120-9), 1987.
- 640 Balesdent, J., Basile-Doelsch, I., Chadoeuf, J., Cornu, S., Derrien, D., Fekiacova, Z., and Hatté, C.: Atmosphere–soil carbon transfer as a function of soil depth, *Nature*, 559, 599–602, <https://doi.org/10.1038/s41586-018-0328-3>, 2018.
- Bossio, D. A., Cook-Patton, S. C., Ellis, P. W., Fargione, J., Sanderman, J., Smith, P., Wood, S., Zomer, R. J., Von Unger, M., Emmer, I. M., and Griscom, B. W.: The role of soil carbon in natural climate solutions, *Nat Sustain*, 3, 391–398, <https://doi.org/10.1038/s41893-020-0491-z>, 2020.
- 645 Bullock, E. L., Healey, S. P., Yang, Z., Oduor, P., Gorelick, N., Omondi, S., Ouko, E., and Cohen, W. B.: Three Decades of Land Cover Change in East Africa, *Land*, 10, 150, <https://doi.org/10.3390/land10020150>, 2021.
- Champely, S.: pwr: Basic Functions for Power Analysis, <https://doi.org/10.32614/CRAN.package.pwr>, 2006.
- Chen, R., Senbayram, M., Blagodatsky, S., Myachina, O., Dittert, K., Lin, X., Blagodatskaya, E., and Kuzyakov, Y.: Soil C and N availability determine the priming effect: microbial N mining and stoichiometric decomposition theories, *Global Change Biology*, 20, 2356–2367, <https://doi.org/10.1111/gcb.12475>, 2014.
- 650 Chen, Y. and Aviad, T.: Effects of Humic Substances on Plant Growth, in: *Humic Substances in Soil and Crop Sciences: Selected Readings*, edited by: MacCarthy, P., Clapp, C. E., Malcolm, R. L., and Bloom, P. R., Soil Science Society of America, Madison, WI, USA, 161–186, <https://doi.org/10.2136/1990.humicsubstances.c7>, 1990.
- Chivenge, P., Vanlauwe, B., Gentile, R., Wangechi, H., Mugendi, D., van Kessel, C., and Six, J.: Organic and Mineral Input Management to Enhance Crop Productivity in Central Kenya, *Agronomy Journal*, 101, 10, 2009.
- 655 Chivenge, P., Vanlauwe, B., Gentile, R., and Six, J.: Comparison of organic versus mineral resource effects on short-term aggregate carbon and nitrogen dynamics in a sandy soil versus a fine textured soil, *Agriculture, Ecosystems & Environment*, 140, 361–371, <https://doi.org/10.1016/j.agee.2010.12.004>, 2011a.
- 660 Chivenge, P., Vanlauwe, B., and Six, J.: Does the combined application of organic and mineral nutrient sources influence maize productivity? A meta-analysis, *Plant Soil*, 342, 1–30, <https://doi.org/10.1007/s11104-010-0626-5>, 2011b.
- Corbeels, M., Cardinael, R., Naudin, K., Guibert, H., and Torquebiau, E.: The 4 per 1000 goal and soil carbon storage under agroforestry and conservation agriculture systems in sub-Saharan Africa, *Soil and Tillage Research*, 188, 16–26, <https://doi.org/10.1016/j.still.2018.02.015>, 2019.
- 665 Córdova, S. C., Olk, D. C., Dietzel, R. N., Mueller, K. E., Archontoulis, S. V., and Castellano, M. J.: Plant litter quality affects the accumulation rate, composition, and stability of mineral-associated soil organic matter, *Soil Biology and Biochemistry*, 125, 115–124, <https://doi.org/10.1016/j.soilbio.2018.07.010>, 2018.

- Ehleringer, J. R., Buchmann, N., and Flanagan, L. B.: Carbon Isotope Ratios in belowground carbon cycle processes, *Ecological Applications*, 10, 412–422, [https://doi.org/10.1890/1051-0761\(2000\)010\[0412:CIRIBC\]2.0.CO;2](https://doi.org/10.1890/1051-0761(2000)010[0412:CIRIBC]2.0.CO;2), 2000.
- 670 Farquhar G D, Ehleringer, J R, and Hubick, K T: Carbon isotope discrimination and photosynthesis, *Annual review of plant physiology and plant molecular biology*, 40, 503–537, <https://doi.org/10.1146/annurev.arplant.40.1.503>, 1989.
- Feller, C. and Beare, M. H.: Physical control of soil organic matter dynamics in the tropics, *Geoderma*, 79, 69–116, [https://doi.org/10.1016/S0016-7061\(97\)00039-6](https://doi.org/10.1016/S0016-7061(97)00039-6), 1997.
- 675 Fujisaki, K., Chevallier, T., Chapuis-Lardy, L., Albrecht, A., Razafimbelo, T., Masse, D., Ndour, Y. B., and Chotte, J.-L.: Soil carbon stock changes in tropical croplands are mainly driven by carbon inputs: A synthesis, *Agriculture, Ecosystems & Environment*, 259, 147–158, <https://doi.org/10.1016/j.agee.2017.12.008>, 2018.
- Geisseler, D. and Scow, K. M.: Long-term effects of mineral fertilizers on soil microorganisms – A review, *Soil Biology and Biochemistry*, 75, 54–63, <https://doi.org/10.1016/j.soilbio.2014.03.023>, 2014.
- Gentile, R., Vanlauwe, B., Chivenge, P., and Six, J.: Interactive effects from combining fertilizer and organic residue inputs on nitrogen transformations, *Soil Biology and Biochemistry*, 40, 2375–2384, <https://doi.org/10.1016/j.soilbio.2008.05.018>, 2008.
- 680 Gentile, R., Vanlauwe, B., and Six, J.: Litter quality impacts short- but not long-term soil carbon dynamics in soil aggregate fractions, *Ecological Applications*, 21, 695–703, <https://doi.org/10.1890/09-2325.1>, 2011.
- Gram, G., Roobroeck, D., Pypers, P., Six, J., Merckx, R., and Vanlauwe, B.: Combining organic and mineral fertilizers as a climate-smart integrated soil fertility management practice in sub-Saharan Africa: A meta-analysis, *PLoS ONE*, 15, e0239552, <https://doi.org/10.1371/journal.pone.0239552>, 2020.
- 685 Greenwood, D. J., Gerwitz, A., Stone, D. A., and Barnes, A.: Root development of vegetable crops, *Plant and Soil*, 1982.
- Gross, A. and Glaser, B.: Meta-analysis on how manure application changes soil organic carbon storage, *Sci Rep*, 11, 5516, <https://doi.org/10.1038/s41598-021-82739-7>, 2021.
- 690 Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., Thau, D., Stehman, S. V., Goetz, S. J., Loveland, T. R., Kommareddy, A., Egorov, A., Chini, L., Justice, C. O., and Townshend, J. R. G.: High-Resolution Global Maps of 21st-Century Forest Cover Change, *Science*, 342, 850–853, <https://doi.org/10.1126/science.1244693>, 2013.
- Hendershot, W. H. and Duquette, M.: A Simple Barium Chloride Method for Determining Cation Exchange Capacity and Exchangeable Cations, *Soil Science Soc of Amer J*, 50, 605–608, <https://doi.org/10.2136/sssaj1986.03615995005000030013x>, 1986.
- 695 Hofmann, A., Heim, A., Gioacchini, P., Miltner, A., Gehre, M., and Schmidt, M. W. I.: Mineral fertilization did not affect decay of old lignin and SOC in a ^{13}C -labeled arable soil over 36 years, *Biogeosciences*, 6, 1139–1148, <https://doi.org/10.5194/bg-6-1139-2009>, 2009.
- Kaiser, K. and Kalbitz, K.: Cycling downwards – dissolved organic matter in soils, *Soil Biology and Biochemistry*, 52, 29–32, <https://doi.org/10.1016/j.soilbio.2012.04.002>, 2012a.
- 700 Kaiser, K. and Kalbitz, K.: Cycling downwards – dissolved organic matter in soils, *Soil Biology and Biochemistry*, 52, 29–32, <https://doi.org/10.1016/j.soilbio.2012.04.002>, 2012b.

- Kassambara, A.: *rstatix*: Pipe-Friendly Framework for Basic Statistical Tests_. R package version 0.7.1, <<https://CRAN.R-project.org/package=rstatix>>, 2022.
- 705 Kihara, J., Bolo, P., Kinyua, M., Nyawira, S. S., and Sommer, R.: Soil health and ecosystem services: Lessons from sub-Saharan Africa (SSA), *Geoderma*, 370, 114342, <https://doi.org/10.1016/j.geoderma.2020.114342>, 2020.
- Komada, T., Anderson, M. R., and Dorfmeier, C. L.: Carbonate removal from coastal sediments for the determination of organic carbon and its isotopic signatures, $\delta^{13}\text{C}$ and $\Delta^{14}\text{C}$: comparison of fumigation and direct acidification by hydrochloric acid, *Limnology & Ocean Methods*, 6, 254–262, <https://doi.org/10.4319/lom.2008.6.254>, 2008.
- 710 Kuznetsova, A., Brockhoff, P. B., and Christensen, R. H. B.: lmerTest Package: Tests in Linear Mixed Effects Models, *J. Stat. Soft.*, 82, <https://doi.org/10.18637/jss.v082.i13>, 2017.
- Ladha, J. K., Reddy, C. K., Padre, A. T., and Van Kessel, C.: Role of Nitrogen Fertilization in Sustaining Organic Matter in Cultivated Soils, *J of Env Quality*, 40, 1756–1766, <https://doi.org/10.2134/jeq2011.0064>, 2011a.
- Ladha, J. K., Reddy, C. K., Padre, A. T., and Van Kessel, C.: Role of Nitrogen Fertilization in Sustaining Organic Matter in Cultivated Soils, *J of Env Quality*, 40, 1756–1766, <https://doi.org/10.2134/jeq2011.0064>, 2011b.
- 715 Lal, R.: Soils and sustainable agriculture. A review, *Agron. Sustain. Dev.*, 28, 57–64, <https://doi.org/10.1051/agro:2007025>, 2008.
- Lal, R.: Digging deeper: A holistic perspective of factors affecting soil organic carbon sequestration in agroecosystems, *Glob Change Biol*, 24, 3285–3301, <https://doi.org/10.1111/gcb.14054>, 2018.
- 720 Laub, M., Schlichenmeier, S., Vityakon, P., and Cadisch, G.: Litter Quality and Microbes Explain Aggregation Differences in a Tropical Sandy Soil, *J Soil Sci Plant Nutr*, 22, 848–860, <https://doi.org/10.1007/s42729-021-00696-6>, 2022.
- Laub, M., Corbeels, M., Mathu Ndungu, S., Mucheru-Muna, M. W., Mugendi, D., Nepalova, M., Van De Broek, M., Waswa, W., Vanlauwe, B., and Six, J.: Combining manure with mineral N fertilizer maintains maize yields: Evidence from four long-term experiments in Kenya, *Field Crops Research*, 291, 108788, <https://doi.org/10.1016/j.fcr.2022.108788>, 2023a.
- 725 Laub, M., Corbeels, M., Couédel, A., Ndungu, S. M., Mucheru-Muna, M. W., Mugendi, D., Nepalova, M., Waswa, W., Van De Broek, M., Vanlauwe, B., and Six, J.: Managing soil organic carbon in tropical agroecosystems: evidence from four long-term experiments in Kenya, *SOIL*, 9, 301–323, <https://doi.org/10.5194/soil-9-301-2023>, 2023b.
- Leuther, F., Wolff, M., Kaiser, K., Schumann, L., Merbach, I., Mikutta, R., and Schlüter, S.: Response of subsoil organic matter contents and physical properties to long-term, high-rate farmyard manure application, *European J Soil Science*, 73, e13233, <https://doi.org/10.1111/ejss.13233>, 2022.
- 730 Liu, E., Yan, C., Mei, X., Zhang, Y., and Fan, T.: Long-Term Effect of Manure and Fertilizer on Soil Organic Carbon Pools in Dryland Farming in Northwest China, *PLoS ONE*, 8, e56536, <https://doi.org/10.1371/journal.pone.0056536>, 2013.
- Liu, N., Li, Y., Cong, P., Wang, J., Guo, W., Pang, H., and Zhang, L.: Depth of straw incorporation significantly alters crop yield, soil organic carbon and total nitrogen in the North China Plain, *Soil and Tillage Research*, 205, 104772, <https://doi.org/10.1016/j.still.2020.104772>, 2021.
- 735 Liu, Y., Li, C., Cai, G., Sauheitl, L., Xiao, M., Shibistova, O., Ge, T., and Guggenberger, G.: Meta-analysis on the effects of types and levels of N, P, and K fertilization on organic carbon in cropland soils, *Geoderma*, 437, 116580, <https://doi.org/10.1016/j.geoderma.2023.116580>, 2023.

- 740 Ma, Q., Bell, R. W., and Mattiello, E. M.: Nutrient Acquisition with Particular Reference to Subsoil Constraints, in: *Subsoil Constraints for Crop Production*, edited by: Oliveira, T. S. D. and Bell, R. W., Springer International Publishing, Cham, 289–321, https://doi.org/10.1007/978-3-031-00317-2_12, 2022.
- Mathieu, J. A., Hatté, C., Balesdent, J., and Parent, É.: Deep soil carbon dynamics are driven more by soil type than by climate: a worldwide meta-analysis of radiocarbon profiles, *Glob Change Biol*, 21, 4278–4292, <https://doi.org/10.1111/gcb.13012>, 2015.
- 745 Minasny, B., Malone, B. P., McBratney, A. B., Angers, D. A., Arrouays, D., Chambers, A., Chaplot, V., Chen, Z.-S., Cheng, K., Das, B. S., Field, D. J., Gimona, A., Hedley, C. B., Hong, S. Y., Mandal, B., Marchant, B. P., Martin, M., McConkey, B. G., Mulder, V. L., O'Rourke, S., Richer-de-Forges, A. C., Odeh, I., Padarian, J., Paustian, K., Pan, G., Poggio, L., Savin, I., Stolbovoy, V., Stockmann, U., Sulaeman, Y., Tsui, C.-C., Vågen, T.-G., Van Wesemael, B., and Winowiecki, L.: Soil carbon 4 per mille, *Geoderma*, 292, 59–86, <https://doi.org/10.1016/j.geoderma.2017.01.002>, 2017.
- 750 Ndung'u, M., Ngatia, L. W., Onwonga, R. N., Mucheru-Muna, M. W., Fu, R., Moriasi, D. N., and Ngetich, K. F.: The influence of organic and inorganic nutrient inputs on soil organic carbon functional groups content and maize yields, *Heliyon*, 7, e07881, <https://doi.org/10.1016/j.heliyon.2021.e07881>, 2021.
- Pretty, J.: Sustainable intensification in Africa, *International Journal of Agricultural Sustainability*, 9, 3–4, <https://doi.org/10.3763/ijas.2011.91ED>, 2011.
- 755 Reimer, P., Brown, T., and Reimer, R.: Discussion: Reporting and Calibration of Post-Bomb ¹⁴C Data, *Radiocarbon*, 46, 1299–1304, <https://doi.org/10.1017/S0033822200033154>, 2004.
- Rumpel, C. and Kögel-Knabner, I.: Deep soil organic matter—a key but poorly understood component of terrestrial C cycle, *Plant Soil*, 338, 143–158, <https://doi.org/10.1007/s11104-010-0391-5>, 2011.
- 760 Shumba, A., Chikowo, R., Thierfelder, C., Corbeels, M., Six, J., and Cardinael, R.: Mulch application as the overarching factor explaining increase in soil organic carbon stocks under conservation agriculture in two 8-year-old experiments in Zimbabwe, *SOIL*, 10, 151–165, <https://doi.org/10.5194/soil-10-151-2024>, 2024.
- Sierra, C. A., Ahrens, B., Bolinder, M. A., Braakhekke, M. C., Von Fromm, S., Kätterer, T., Luo, Z., Parvin, N., and Wang, G.: Carbon sequestration in the subsoil and the time required to stabilize carbon for climate change mitigation, *Global Change Biology*, 30, e17153, <https://doi.org/10.1111/gcb.17153>, 2024.
- 765 Six, J., Feller, C., Denef, K., Ogle, S. M., de Moraes, J. C., and Albrecht, A.: Soil organic matter, biota and aggregation in temperate and tropical soils - Effects of no-tillage, *Agronomie*, 22, 755–775, <https://doi.org/10.1051/agro:2002043>, 2002.
- Sommer, R., Paul, B. K., Mukalama, J., and Kihara, J.: Reducing losses but failing to sequester carbon in soils – the case of Conservation Agriculture and Integrated Soil Fertility Management in the humid tropical agro-ecosystem of Western Kenya, *Agriculture, Ecosystems & Environment*, 254, 82–91, <https://doi.org/10.1016/j.agee.2017.11.004>, 2018.
- Tardieu, F.: Analysis of the spatial variability of maize root density, *Plant and Soil*, 1988.
- 770 Tieszen, L. L.: Natural variations in the carbon isotope values of plants: Implications for archaeology, ecology, and paleoecology, *Journal of Archaeological Science*, 18, 227–248, [https://doi.org/10.1016/0305-4403\(91\)90063-U](https://doi.org/10.1016/0305-4403(91)90063-U), 1991.
- Tittonell, P., Corbeels, M., Van Wijk, M. T., Vanlauwe, B., and Giller, K. E.: Combining Organic and Mineral Fertilizers for Integrated Soil Fertility Management in Smallholder Farming Systems of Kenya: Explorations Using the Crop-Soil Model FIELD, *Agronomy Journal*, 100, 1511–1526, <https://doi.org/10.2134/agronj2007.0355>, 2008.

- 775 Trevisan, S., Francioso, O., Quaggiotti, S., and Nardi, S.: Humic substances biological activity at the plant-soil interface: From environmental aspects to molecular factors, *Plant Signaling & Behavior*, 5, 635–643, <https://doi.org/10.4161/psb.5.6.11211>, 2010.
- Trumbore, S. E., Vogel, J. S., and Southon, J. R.: AMS ^{14}C Measurements of Fractionated Soil Organic Matter: An Approach to Deciphering the Soil Carbon Cycle, *Radiocarbon*, 31, 644–654, <https://doi.org/10.1017/S0033822200012248>, 1989.
- 780 Uselman, S. M., Qualls, R. G., and Lilienfein, J.: Contribution of Root vs. Leaf Litter to Dissolved Organic Carbon Leaching through Soil, *Soil Sci. Soc. Am. J.*, 71, 1555–1563, <https://doi.org/10.2136/sssaj2006.0386>, 2007.
- Van De Broek, M., Ghiasi, S., Decock, C., Hund, A., Abiven, S., Friedli, C., Werner, R. A., and Six, J.: The soil organic carbon stabilization potential of old and new wheat cultivars: a $\delta^{13}\text{C}$ and CO_2 labeling study, *Biogeosciences*, 17, 2971–2986, <https://doi.org/10.5194/bg-17-2971-2020>, 2020.
- 785 Vanlauwe, B. and Giller, K.: Popular myths around soil fertility management in sub-Saharan Africa, *Agriculture, Ecosystems & Environment*, 116, 34–46, <https://doi.org/10.1016/j.agee.2006.03.016>, 2006.
- Vanlauwe, B., Gachengo, C., Shepherd, K., Barrios, E., Cadisch, G., and Palm, C. A.: Laboratory Validation of a Resource Quality-Based Conceptual Framework for Organic Matter Management, *Soil Sci. Soc. Am. j.*, 69, 1135–1145, <https://doi.org/10.2136/sssaj2004.0089>, 2005.
- 790 Vanlauwe, B., Chianu, J., Giller, K. E., Merckx, R., Mokwunye, U., Pypers, P., Shepherd, K., Woomer, P. L., and Sanginga, N.: Integrated soil fertility management: operational definition and consequences for implementation and dissemination, 2010.
- Vanlauwe, B., Descheemaeker, K., Giller, K. E., Huising, J., Merckx, R., Nziguheba, G., Wendt, J., and Zingore, S.: Integrated soil fertility management in sub-Saharan Africa: unravelling local adaptation, *SOIL*, 1, 491–508, <https://doi.org/10.5194/soil-1-491-2015>, 2015.
- 795 Veldkamp, E., Schmidt, M., Powers, J. S., and Corre, M. D.: Deforestation and reforestation impacts on soils in the tropics, *Nat Rev Earth Environ*, 1, 590–605, <https://doi.org/10.1038/s43017-020-0091-5>, 2020.
- 800 Wang, J., Sun, J., Xia, J., He, N., Li, M., and Niu, S.: Soil and vegetation carbon turnover times from tropical to boreal forests, *Functional Ecology*, 32, 71–82, <https://doi.org/10.1111/1365-2435.12914>, 2018.
- Wiesmeier, M., Urbanski, L., Hobbey, E., Lang, B., von Lützow, M., Marin-Spiotta, E., van Wesemael, B., Rabot, E., Ließ, M., Garcia-Franco, N., Wollschläger, U., Vogel, H.-J., and Kögel-Knabner, I.: Soil organic carbon storage as a key function of soils - A review of drivers and indicators at various scales, *Geoderma*, 333, 149–162, <https://doi.org/10.1016/j.geoderma.2018.07.026>, 2019.
- 805 Yang, X., Xiong, J., Du, T., Ju, X., Gan, Y., Li, S., Xia, L., Shen, Y., Pacenka, S., Steenhuis, T. S., Siddique, K. H. M., Kang, S., and Butterbach-Bahl, K.: Diversifying crop rotation increases food production, reduces net greenhouse gas emissions and improves soil health, *Nat Commun*, 15, 198, <https://doi.org/10.1038/s41467-023-44464-9>, 2024.
- Yost, J. L. and Hartemink, A. E.: How deep is the soil studied – an analysis of four soil science journals, *Plant Soil*, 452, 5–18, <https://doi.org/10.1007/s11104-020-04550-z>, 2020.