

Invited perspectives: Thunderstorm Intensification from Mountains to Plains

Jannick Fischer¹, Pieter Groenemeijer^{2,3}, Alois Holzer^{2,3}, Monika Feldmann⁴, Katharina Schröer⁵, Francesco Battaglioli³, Lisa Schielicke^{6,7}, Tomáš Púčik², Bogdan Antonescu^{3,8}, Christoph Gatzert³, and TIM Partners⁹

¹Karlsruhe Institute of Technology (KIT), Karlsruhe, Germany

²European Severe Storms Laboratory - Science & Training, Wiener Neustadt, Austria

³European Severe Storms Laboratory e. V., Wessling, Germany

⁴Institute of Geography - Oeschger Centre for Climate Change Research, University of Bern

⁵Department of Environment and Natural Resources, University of Freiburg

⁶Institute for Geosciences, University of Bonn

⁷Department of Physics and Astronomy, The University of Western Ontario, London, Canada

⁸Faculty of Physics, University of Bucharest, Romania

⁹See extended co-author list from partner institutions at the end of this manuscript

Correspondence: Jannick Fischer (jannick.fischer@kit.edu)

Abstract. Severe thunderstorms are among the most damaging and impactful weather phenomena. In Europe, notable clusters occur in the vicinity of complex terrain. These areas not only experience frequent thunderstorms but also show a strong climate change signal with an increasing storm frequency. Despite the relevance of the subject, our understanding of severe convection in complex terrain, particularly in a changing climate, remains incomplete. This White Paper presents the current state of the research on thunderstorms in complex orography, covering storm severity, modification of pre-storm environments, convection initiation, storm-scale interactions with complex terrain, impactful hazards, numerical modeling and forecasting, climatologies and climate change signals, as well as innovative storm observations. Highlighting the gaps in our understanding, this review underscores the need for a coordinated European field campaign on Thunderstorm Intensification from Mountains to Plains (TIM). Initial plans for the TIM campaign, built by participating authors and institutions of this article, are briefly outlined. Obtaining coordinated and dense data on orographically driven storms is a key step toward improving warnings, forecasts, future climate projections, and adaptation measures.

1 Introduction

Severe convective storms (SCS) with hail, flash floods, wind gusts, lightning, and tornadoes cause a significant and increasing amount of damage across Europe and other mid-latitude regions (e.g., SwissRe, 2024). In 2023 alone, a new record hail size was set for Europe with 19 cm (ESSL, 2023c) while in the US the North America the insured loss from severe convective storms reached around 60 billion USD, thereby also setting a new record and dominating the global total loss from Natural Catastrophes (Bowen et al., 2024). These facts (e.g., SwissRe, 2024). These impacts highlight the importance of research on all aspects of severe convective storms to better understand where and why these hazards occur.

Large hail reports 2008 – 2024




Figure 1. Severe hail reports between ~~2008-2024~~2008–2024 in the European Severe Weather Database (ESWD; accessed 10 March 2025). The topography is qualitatively ~~underlaid~~underlaid in gray with darker shading indicating higher terrain. For more topographic detail and the locations of TIM partner institutions see Fig. 7.

Reports of severe weather in the European Severe Weather Database (Dotzek et al., 2009) suggest that ~~particularly~~ severe thunderstorms are most frequent over the flanking slopes of mountain ranges and the plains that straddle them. As an example, Fig. 1 shows the clustering of severe hail reports north and south of the Alps and predominantly on the eastern or northeastern flanks of lower mountain ranges (the reader may skip ahead to Fig. 7 for more topographic detail and regions of interest). This clustering of reports is supported by indirect but spatially less population-biased measures of storm severity than reports, such as overshooting storm tops (Punge et al., 2017; Giordani et al., 2023), lightning frequency (e.g., Manzato et al., 2022b), and radar-based climatologies (e.g., Kaltenboeck and Steinheimer, 2015; Punge and Kunz, 2016; Wapler, 2021; Fluck et al., 2021; Feldmann et al., 2023). The same geographic regions show a strong past and projected increase in SCS prone atmospheric conditions as a result of global warming (Púčik et al., 2017; Rädler et al., 2019; Battaglioli et al., 2023). In addition, many of these parts of Europe are highly vulnerable because of dense population or specialized agriculture (e.g., Púčik et al., 2019). Thus, there is both a need and an opportunity to study SCS in these regions.

What research questions do you have regarding SCS near mountains and why?

What hypotheses with respect to these questions do you have?

What observations or modeling experiments are needed to test these hypotheses?

How could your institution contribute to a dedicated field campaign?

Figure 2. Questions from the White Paper survey conducted for this article with the European partner institutions listed further below.

30 One key question in this regard is why storms in the vicinity of European mountain ranges are more severe. As reviewed in section 2.1, there are several possible explanations in the literature. However, our understanding of thunderstorm dynamics and mesoscale atmospheric processes is still incomplete. One reason is that observational datasets to validate existing theories are scarce and, where available, distributed across many institutions and data archives. This lack of understanding and of shared data is echoed throughout the research field of severe convective storms. As basis for this article, the European Severe Storms Laboratory (ESSL) conducted a survey with institutions in the European atmospheric science community, asking for their perspectives on the most crucial research topics on severe convective storms near mountains (Fig. 2). The contributors to this survey also took part in the writing of this article and are listed in the extended Co-author section. The survey outcome was a pool of over 40 research questions, which are sorted by topic and discussed in detail in section 2. A recurring theme in the responses is that a better understanding of the physical processes controlling severe convective storms and their impacts is needed. Public authorities, research institutions, the aviation industry, and the reinsurance sector share the need to better estimate the frequency, magnitude, and predictability of severe weather.

Presently, a wealth of new sensors and data types is emerging to cater to these needs, ~~for example~~ such as the Meteosat Third Generation satellite programme (Holmlund et al., 2021), numerical modeling within the DestinE project (Wedi et al., 2022), and the proliferation of polarimetric Doppler radars across Europe (Saltikoff et al., 2019). Validation of these datasets is key, and requires that they are shared among scientists with the least possible restrictions. This validation is especially important near mountain ranges, not only because of the high severe weather frequency (Fig. 1), but also because the lack of observations of basic meteorological parameters in the boundary layer and in complex terrain¹ is critical to advance forecasting and nowcasting skill (Bojinski et al., 2023).

This prominent need for shared data and research, especially near mountains, can partly be explained by the prevailing focus of past field campaigns either on thunderstorm dynamics away from mountain ranges *or* on terrain influences on the atmosphere, but not on the scale of individual storms. For instance, several large field campaigns in the United States, such as

¹The terms "complex terrain" and "orographic" will be used in this article when referring to the vicinity of mountains.

VORTEX1/2/SE (Rasmussen et al., 1994; Wurman et al., 2012), PECAN (Geerts et al., 2017), TORUS (Houston et al., 2020), PERILS (Kosiba et al., 2024) and TRACER (Jensen et al., 2022) led to advancements in our understanding of thunderstorm dynamics, microphysics, and processes leading to severe weather. ~~Convection-focused~~ Meanwhile, SCS-focused campaigns in Europe, such as Swabian MOSES (Kunz et al., 2022) and LIFT (Kunz et al., 2024), were relatively small. ~~Meanwhile, larger, and major~~ convective-season field campaigns in mountainous regions, such as MAP (Bougeault et al., 2001), COPS (Wulfmeyer et al., 2011), HyMeX (Ducrocq et al., 2016) and, currently, TEAMx (Rotach et al., 2022), improved numerical models and process understanding ~~with respect to~~ of large-scale orographic flow, boundary-layer exchange processes, autumn convective floods, and processes occurring prior to convection initiation (CI), but much less so regarding the dynamics of individual thunderstorms (e.g., Rotunno and Houze, 2007). Exceptions are RELAMPAGO and its sister campaign CACTI, which took place in ~~Western~~ western Argentina where some of the most intense storms worldwide are heavily influenced by the Andes and Sierras de Cordoba mountains (Nesbitt et al., 2021; Varble et al., 2021). However, the setting of RELAMPAGO in the lee of a long meridional mountain range and with the Amazon Basin as a moisture source differs greatly from Europe and other severe storm regions of the world ~~with respect to~~ concerning larger-scale environments, geography, climatology, and meteorological instrument networks (e.g., Zhou et al., 2021).

Therefore, a field campaign investigating individual thunderstorm dynamics near complex terrain in the densely-observable regions over Europe seems overdue, particularly given the potentially escalating storm activity attributed to climate change in ~~densely~~ densely populated regions (Battaglioli et al., 2023). To this end, ESSL is coordinating an effort for an international field campaign in Europe under the name TIM (Thunderstorm Intensification from Mountains to ~~plains~~ Plains). In this article, we summarize the most pressing research questions based on the White Paper survey and the existing literature (section 2). Section 3 then provides a summary and briefly outlines how this wide range of topics can be addressed in TIM.

2 Research topics

Our survey within the community has revealed a large pool of research topics regarding severe storms near mountain ranges. These topics are condensed in the following subsections. Section 2.1 starts with an overview of the physical processes important for thunderstorm intensification near mountains. The following sections describe the related research topics that present the main foci in TIM.

2.1 Increased storm severity near high terrain

As established above, severe convective storms tend to be frequent near European mountain ranges, which is underscored by the documented cases of impactful supercells that tracked through such regions (Kaňák et al., 2007; Kunz et al., 2018; Trefalt et al., 2018; Šinger and Púčik, 2020; Wilhelm et al., 2021; Kopp et al., 2023). That said, it is difficult to disentangle ~~to what extent~~ whether it is high storm frequency and /or stronger average intensity ~~are each driving~~ that drive these near-mountain severe weather maxima.

Thunderstorm frequency is largely related to convection initiation (CI) or the lack thereof. Mesoscale orographic processes dominate CI mechanisms in mountainous regions, in contrast to synoptic-scale processes, such as fronts, which are more important over flat terrain (e.g., Pacey et al., 2023). As illustrated in Figure 3 and discussed in more detail in section 2.3, orographic CI can occur where terrain-induced flows converge and cause ~~deep-continued~~continuous deep lift in the atmospheric boundary layer (e.g., Kirshbaum et al., 2018), especially if these flows transport and vertically mix ~~boundary-layer moisture and~~thereby boundary layer moisture and thus decrease convective inhibition (CIN) and dry air entrainment near the convergence zones (e.g., Scheffknecht et al., 2017; Serafin et al., 2020; Marquis et al., 2021; Nelson et al., 2022; Göbel et al., 2023). Hence, mountain ranges strongly influence where thunderstorms form (e.g. Nisi et al., 2018; Manzato et al., 2022b). Although severe storms can ~~move several~~travel hundreds of kilometers (e.g., Scheffknecht et al., 2017; Kunz et al., 2018), CI-prone regions tend to have a much higher frequency and coverage of severe storms (e.g., Feldmann et al., 2023; Kvak et al., 2023). ~~Nevertheless, for~~For Switzerland, Feldmann et al. (2023) demonstrated that while the regions of highest storm frequency and highest storm intensity strongly overlap, they are not identical (see also; Nisi et al., 2018; Manzato et al., 2022a). Thus, CI is likely an important but not the only factor.

Severe storm intensity is largely determined by the vertical distribution of temperature, moisture and horizontal winds. Most reports of severe weather occur where certain atmospheric "ingredients" (e.g., Doswell et al., 1996) are met, which can be boiled down to (1) high convective available potential energy (CAPE) which drives deep convective updrafts and which is found where tropospheric temperature lapse rates and low-level moisture are large, and (2) strong vertical change of wind speed and/or wind direction, i.e., vertical wind shear (e.g., Thompson et al., 2003; Taszarek et al., 2020a). Given CI in regions where these ingredients are met, storms tend to become more intense, more organized, and longer-lived, often forming mesoscale convective systems (MCS) or supercells (e.g., Markowski and Richardson, 2010). Thus, one likely reason for the accumulation of severe reports near orography (~~Fig. 1~~) is that these ingredients are enhanced ~~there~~. Several mechanisms exist in that regard, which are summarized in the rest of this section. Figure 3 illustrates the respective key processes.

Local terrain-induced flow systems such as thermal slope winds and Alpine ~~Pumping~~pumping (Ćurić and Janc, 2012; Soderholm et al., 2014; Trefalt et al., 2018; Mulholland et al., 2020; Trapp et al., 2020; Katona and Markowski, 2021), wind channeling (Whiteman and Doran, 1993; Hannesen et al., 1998; Ćurić et al., 2007; Geerts et al., 2009; Tang et al., 2016; Feldmann et al., 2024), or lee cyclogenesis (e.g., McGinley, 1982) can cause perturbations in the atmospheric lower-tropospheric winds. This can increase the vertical wind shear and thereby severe storm potential. Furthermore, the wind profile influences storm structure and propagation (e.g., Houze et al., 1993). Terrain flows depend on the thermodynamic stability of the atmosphere as well as the terrain shape and dimensions (e.g., Houze, 2012). As a result, typical flow systems seem to exist for individual mountain ranges (e.g., Dotzek, 2001; Kunz and Pusskeiler, 2010), but more universal conceptual models for what processes dominate in what scenario have not been developed thus far.

Another factor is the spatial distribution of low-level moisture across complex terrain. Moisture is often relatively low directly over mountain ridges, leading to less CAPE compared to the surrounding slopes and valleys (e.g., Katona and Markowski, 2021). However, water vapor content and CAPE can be increased by near-ground up-slope moisture flux (e.g., from the Mediterranean; De Martin et al., 2024, 2025), evapotranspiration from lakes or vegetation (e.g., Laiti et al., 2014; Feldmann




Figure 3. Conceptual illustration of processes influencing severe convective storms near and over complex terrain as explained in the text. The term “terrain flows” encompasses all possible terrain-induced near-ground flow perturbations, such as thermally-forced circulation caused by differential heating, mechanically forced ascent or channeling, upstream blocking, and lee-side convergence (e.g., Kirshbaum et al., 2018).

et al., 2024), or along airmass boundaries and convergence lines (e.g., Maddox et al., 1980). ~~Because of~~ Due to these effects, the distribution of water vapor in complex terrain is often spatially heterogeneous, which can play a role in the formation of thunderstorms (Adler et al., 2016; Marquis et al., 2021; Nelson et al., 2021; Calbet et al., 2022). More research is needed to understand the storm-scale impacts of such heterogeneities and to reliably detect them with remote-sensing.

The interpretation of the thermodynamic profile near mountains is complicated by the heterogeneous heat and moisture fluxes. Nevertheless, the studies below have suggested that with increasing elevation near-surface air parcels that enter a thunderstorm cloud originate higher relative to the background thermodynamic profile that is advected over the terrain, i.e., inflow air does not originate near the surface. At intermediate elevation, this can lead to an increase in convective energy because the most unstable parcels are being lifted (Markowski and Dotzek, 2011; Scheffknecht et al., 2017; Katona and Markowski, 2021; Feldmann et al., 2024). At increasingly higher elevation or on the leeward side (with respect to the deep-layer winds) of a mountain ridge, CAPE is typically diminished because lifted parcels from higher in the atmosphere are drier and the depth of the CAPE layer decreases (Markowski and Dotzek, 2011; Mulholland et al., 2019; Katona and Markowski, 2021; Feldmann et al., 2024). However, these studies have only considered specific mountain ranges or idealized scenarios, and hence it remains unclear how significant these effects are in different geographic regions.

The above-mentioned convergence regions from terrain-induced flows may also overlap with moisture gradients and take characteristics of dryline-like boundaries (e.g., Pistotnik et al., 2011; LeBel et al., 2021). Such boundaries are known to have an impact on storm severity, mainly through locally enhanced low-level vertical wind shear, moisture, and lift (e.g., Maddox

135 et al., 1980). However, mesoscale boundaries have not been well documented in Europe, although case studies indicate at least some importance for supercells that produce large hail and tornadoes (e.g., Komjáti et al., 2023; De Martin et al., 2024, 2025).

Lastly, complex terrain also influences the internal structure and processes within existing thunderstorms. This can include channeling of the outflow and inflow of a thunderstorm (Fig. 3), which in turn can feed back to storm intensification via increased storm-relative winds or baroclinic generation of horizontal vorticity (e.g., Feldmann et al., 2024). As discussed in
140 section 2.4, these storm-scale influences are complex and not well-researched. Hence, much research in the storm scale is needed to better understand them, in addition to the larger-scale, environmental influences in the previous paragraphs.

As suggested by Feldmann et al. (2024), it is the interplay of the factors described in this section that may cause SCS potential to be enhanced over the slopes of mountain ranges compared to flat terrain or directly over the mountains(Fig. 4).
Even slight enhancements are especially relevant in regions like Europe where SCS potential is often marginal (e.g., Taszarek
145 et al., 2020b). However, what processes are typically at play in which weather situation and in which sub-regions remains largely unclear and is considered one research focus of TIM.

2.2 Orographic modification of the mesoscale environment

The influence of orography on the atmosphere and parameters like CAPE and wind shear opens up several follow-up questions. What influence does the prevailing mid-tropospheric flow direction and speed have on the modification of convective
150 parameters? Why are there regions along the Alps and other mountain chains much more prone to severe storms than others? What are the preferential paths for severe storms? At which elevations are environments most or least favorably modified? How significant is the impact of vegetation, water bodies, snow cover, or urban areas in mountainous regions?

Related to these topics is also how storms evolve when either crossing or moving away from mountains and hence encountering either abrupt or continuous changes in the mesoscale environment. Situations have been documented for both decay or
155 intensification in these scenarios (e.g., Scheffknecht et al., 2017; Trapp et al., 2020; Klaus et al., 2023; McKeown et al., 2024);
~~but more research is necessary~~. By documenting example cases within a field campaign such as TIM, detailed observations could help our understanding of these opposing changes.

Furthermore, it is well-established that the atmospheric environments of hazards are statistically different, e.g., between tornadoes and hail (Púčik et al., 2015; Nixon and Allen, 2022). Similarly, Zhou et al. (2021) showed that there are statistically
160 different hailstorm environments~~exist~~, with one of the main differentiating factors being terrain elevation. Hence, the impact of orographic modifications is likely hazard-dependent. Several studies have shown that channel winds in relatively narrow valleys improve the conditions for supercell tornadoes even if the background environment is otherwise unfavorable (Fujita, 1989; Dotzek, 2001; Bosart et al., 2006; Tang et al., 2016). Whether there are similar mechanisms specifically for hail or other hazards is relatively less researched. Targeted observations and measurements as well as simulation studies (e.g., Katona
165 et al., 2016) over different Alpine sub-regions as part of the TIM campaign could provide deeper insights into the processes of orographic modification.

2.3 Convection initiation

Due to its importance for thunderstorm evolution, CI has been a focus of past research. The reader is referred to Houze (2012) and Kirshbaum et al. (2018) for reviews on the topic. Large low- to mid-level relative humidity, dry air entrainment, and strong as well as deep ascent ~~have been shown to be~~ are among the most crucial factors for CI (e.g., Nelson et al., 2022; Marquis et al., 2023). Whether near-ground convergence and ascent occur upstream, over, or downstream of a mountain ridge (Fig. 3) ~~broadly~~ depends on the speed of the flow impinging on the terrain obstacle, the stability of the atmosphere, and the shape and dimensions of the ridge (e.g., Kirshbaum et al., 2018; Imamovic et al., 2019). However, in reality, individual flow and moisture perturbations are extremely complex and subject to strong diurnal cycles (e.g., Zardi and Whiteman, 2013; Adler et al., 2016). On this local scale, terrain-induced flows (e.g., Schmidli, 2013; Panosetti et al., 2016), land use and urban heat islands (e.g., Niyogi et al., 2006; Zhao and Wu, 2018; Liu and Niyogi, 2019), as well as outflow from existing storms (e.g., Soderholm et al., 2014) strongly influence the convergence regions in addition to the larger-scale flow. Furthermore, mountain slope has a strong influence on the advection of moisture and thereby on CI likelihood (Göbel et al., 2023).

These processes make CI more “reliable” over mountains compared to flat terrain and thereby increase the predictability of convective precipitation in forecast models (e.g., Bachmann et al., 2020; Khodayar et al., 2021). However, this is not the case for other SCS hazards. Since small details in ~~CI~~ the timing and location of the CI can have a profound impact on the evolution and organization of severe convective storms, failure to resolve all relevant processes in numerical models remains a main cause ~~for~~ of forecast error. For that reason, CI is an important topic to the weather services ~~who~~ that contributed to this article and a focus of current and recent research (e.g., the past COPS and RELAMPAGO campaigns and the current TEAMx campaign; Wulfmeyer et al., 2011; Nesbitt et al., 2021; Serafin et al., 2020; Rotach et al., 2022).

By focusing explicitly on SCS conditions, TIM can make an important contribution to the research on CI. This includes a better understanding of CI location relative to mountains (e.g., Nisi et al., 2018; Manzato et al., 2022b) by documenting the boundary layer evolution, thermal circulations, and convergence lines before and during CI. The interplay of dry lines and triple points (De Martin et al., 2024), cold fronts (e.g., Pacey et al., 2023), sea-breeze circulations (e.g., Jensen et al., 2022), frontal waves, and gravity waves (e.g., Mulholland et al., 2020) with complex terrain requires further research as well. For instance, the maximum in large hail reports over Northern Italy (~~Fig. 1~~) and the occurrence of tornadoes in the same region appear linked to the combined effects of mesoscale boundaries and ~~sea breezes~~ sea breezes separating the dry mountain boundary layer from the marine boundary layer over the warm Adriatic and Ligurian Seas (Bagaglini et al., 2021; De Martin et al., 2024, 2025).

Another important scientific question concerns scenarios in which CI is suppressed. Several White Paper survey participants noted that on days with high severe weather potential and predicted SCS activity in forecast models, the storms frequently fail to occur or ~~are forming~~ form delayed. Some participants pointed out that this often happens on days with high concentrations of Saharan dust, which is typically advected over Europe together with conditionally unstable airmasses (e.g., Seifert et al., 2023; Brennan and Wilhelm, 2024). This dust can decrease radiative warming at the surface, thereby suppressing boundary layer mixing and CI ~~-, and increases~~ and increasing warming aloft, which may reduce CAPE (Stanelle et al., 2010). These processes are not implemented in most operational forecast models (e.g., Seifert et al., 2023). Assimilating in-situ aerosol measurements

(e.g., Eirund et al., 2022) or remote sensing data from lidars or the recently launched EarthCARE satellite (see section 2.8) could therefore reduce forecast errors of severe convection.

More generally on the topic of aerosols, their role in cloud microphysics and dynamics is a focus of current atmospheric science research and still somewhat uncertain (e.g., Morrison et al., 2020). Regarding severe convection, aerosol concentrations
205 can ~~for example~~, for example, have a direct influence by acting as cloud-condensation or ice nuclei (CCNs or INPs), thereby influencing hydrometeor concentrations and thus storm dynamics (e.g., Loftus and Cotton, 2014; Allen et al., 2020; Varble et al., 2023; Barthlott et al., 2024; Brennan and Wilhelm, 2024). Based on this mechanism, cloud seeding for hail suppression is still widely used in Europe, despite the majority of scientific evidence pointing against its effectiveness as it is used now (e.g., Browning and Foote, 1976; Knight et al., 1979; Auf der Maur and Germann, 2021). Much more dubious is the use of hail
210 cannons (Wieringa and Holleman, 2006). More effective ways of damage prevention should be pursued. A better understanding of aerosol effects, as well as outreach and education as part of TIM (section 2.5), could help in that regard.

2.4 Storm-scale interaction with complex terrain

How the internal structure of thunderstorms changes in reaction to terrain shape is not well understood due to the lack of high-resolution observations within or around thunderstorms. Some insights exist from idealized modeling studies. Based on
215 qualitative experience with storms in the southern pre-Alps, Feldmann et al. (2024) analyzed the interaction of a supercell storm with a simplified ridge and sloped terrain in highly idealized simulations. They showed that channeling of storm-generated outflow and inflow can affect baroclinic horizontal vorticity generation, low-level wind shear, and storm-relative inflow, all of which have an influence on supercell structure and intensity (e.g., Rotunno and Klemp, 1985; Peters et al., 2020). These findings are largely consistent with previous idealized simulations (e.g., Markowski and Dotzek, 2011; Scheffknecht et al.,
220 2017) and the general observation that supercells can traverse and intensify over mountain ranges (e.g., Kvak et al., 2023; McKeown et al., 2024). What seems most needed now is to validate the simulated dynamical processes in real supercells over a large spectrum of cases and types of terrain, an endeavor that requires considerable effort and coordination. Similarly interesting are some non-supercell storms, such as pulse storms (Miller and Mote, 2017), which can produce large amounts of hail for a short time for unknown reasons and which are difficult to anticipate.

225 On larger spatial scales than individual storm cells, several studies have investigated the interaction of mesoscale convective systems (MCS) with orography. These systems are typically driven by lift along the gust front of the cold pool (Markowski and Richardson, 2010, chapter 9), which can be decelerated, accelerated, or blocked by mountains, thereby leading to phases of updraft and precipitation intensification and weakening, respectively (Frame and Markowski, 2006; Reeves and Lin, 2007; Letkewicz and Parker, 2011; Smith et al., 2014; Pucillo et al., 2019; Wu and Lombardo, 2021; Lombardo and Kumjian, 2022).
230 Hence, mountains play important roles in upscale growth from single cells to MCS formation, the system's movement, and in some cases also its decay. The precipitation enhancement of MCSs over orography is one of the reasons for flooding events in mountainous regions (e.g., Saharia et al., 2017). Another reason is that the lift mechanisms near terrain can lead to repeated re-formation, i.e., “backbuilding” of convective cells, which means that precipitation accumulates over the same location over a long time (e.g., Soderholm et al., 2014; Panziera et al., 2015; Kirshbaum et al., 2018). The additional lift along gust fronts of

235 storm outflow, which is influenced by the terrain slope and the environmental temperature and wind profile, plays an important role in these scenarios (e.g., Miglietta and Rotunno, 2014).

Even though past research has helped to increase forecast accuracy, high-precipitation events remain difficult to predict and are ~~in~~ the focus of many weather services in the Alpine and Mediterranean region ~~based on~~, according to our partner institution survey. Unfortunately, they are also difficult to observe accurately. For example, radars are used for Quantitative Precipitation
240 Estimation (QPE, Ryzhkov et al., 2022) but have weaknesses over complex terrain. Radar beam blocking (e.g., Kaltenboeck and Steinheimer, 2015) as well as large vertical gradients of precipitation below the melting layer (e.g., Chen et al., 2023) often lead to underestimates of the QPE. As discussed further below, high-resolution precipitation surface and vertical profile measurements, as well as gap-filling radars, aimed to be collected in TIM, could be used to improve correction algorithms of QPE and help in flooding nowcasting and response.

245 Similar accelerations of thunderstorm outflow as in MCS can be expected for downbursts impinging on and channeled through complex terrain. Downbursts ~~are locally descending jets which can~~ cause damaging winds when colliding with the surface and ~~spreading~~ spread out horizontally from the high-pressure region of the cold pool, ~~and which give~~ giving rise to toroidal vortex structures (e.g., Fujita, 1990; Canepa et al., 2023). Due to their small spatial and temporal scales, the dynamics of downbursts over complex terrain are not well understood and require further research with high-density observations and
250 simulations (e.g., Canepa et al., 2020; Burlando et al., 2024).

2.5 Storm impacts on infrastructure and society

Damages from severe convective storms with tornadoes, wind gusts, hail, extreme rainfall, or lightning, including secondary impacts such as landslides ~~are now dominating~~, are one of the dominating drivers of global insured loss from natural ~~catastrophes~~ catastrophes (SwissRe, 2024). Developing a better understanding of these impacts is a prerequisite to developing adequate
255 measures to mitigate losses. A subtopic that has received renewed attention in recent years is the mechanical or engineering perspective of severe convective hazards. Regarding tornadoes, which are still underestimated in many European countries (Antonescu et al., 2017), ESSL has recently proposed an international Fujita (IF) scale for tornado ratings based on wind damage indicators that can be used internationally, not just in the US where the enhanced Fujita scale is used based on US infrastructure (ESSL, 2023a). More data from tornado damage surveys will continue to improve the IF scale ~~in the future~~. In contrast,
260 flash flood damage in complex terrain is strongly linked to rainfall-runoff processes and hence, from the impact perspective, a hydrological problem (e.g., Ravazzani et al., 2016; Marvi, 2020; Merz et al., 2020). However, accurately representing SCS in numerical models is crucial to provide background data for hydrologic models (next section).

The collection of large hailstones, 3D scanning and analysis of their internal layers, or study of impact kinetic energy on structures such as roof shingles is increasingly pursued by different groups such as the Insurance Institute for Business and
265 Home Safety (IBHS) in the US (Brown and Giammanco, 2013), the Northern Hail project in Canada (Brimelow et al., 2023) and the Bureau of Meteorology in Australia (Soderholm and Kumjian, 2023). Impact studies can ~~for example~~, for example, help establish building codes to prevent hail damage. Schmid et al. (2024) recently showed that the lack of accurate information on hail occurrence, size, and distribution is also hampering damage assessments and risk modeling efforts important

for ~~insurances~~insurance and reinsurance companies. Economic growth (e.g., more cars, solar panels, greenhouses) is a major driver of the increasing trend in SCS losses (SwissRe, 2024). Thus, it is important to quantify the damage and loss that can be expected from hail, wind, or their particularly damaging combination: wind-driven hail (Morgan and Towery, 1976), which is currently difficult to predict. The comparison of 3D scanned and dissected hailstones as well as hail size distributions between different continents, elevations, or climatic regions could also provide valuable insights into hail growth (Soderholm and Kumjian, 2023).

The impact of severe weather on society is strongly dependent on human behavior and risk perception (e.g., Ripberger et al., 2019). Unfortunately, public education on severe convection storms is ~~very~~-limited throughout Europe, perhaps because of the relative ~~rareness~~-rarity of such events and limited cross-border exchange. This may affect people's trust in severe weather forecasts and how they perceive a severe weather threat. For instance, appropriate safety measures and behavior when encountering tornadoes, large hail, strong winds, or lightning are typically not common knowledge. ESSL and the TIM partners share the goal to improve this through internal exchange about better impact-based warning practices, exchange with national civil protection agencies, as well as education and outreach at TIM measurement sites and in the media during the field campaign. Furthermore, the topic of severe convective storms is also somewhat underrepresented in university courses, especially compared to the U.S. where most severe weather research groups are situated. TIM aims to engage students from European universities in the field campaign and thereby help to build the next generation of severe weather forecasters and researchers.

2.6 Improvements in NWP

Numerical weather prediction (NWP) models are an important basis for ~~forecasts of severe weather~~severe weather forecasts. The following NWP strategies are particularly useful to capture the chaotic nature of SCS. Short-term NWP forecasts ~~; i.e.~~ for the next hours ~~;~~ can be blended with information from nowcasting systems that extrapolate observations collected in real-time (e.g., Bojinski et al., 2023). Additionally, the models can be initialized very frequently, ~~e.g.~~-every hour or even less, in ~~so-called~~ rapid update cycles (Benjamin et al., 2004) assimilating a range of observations available at that time (Miyoshi et al., 2016). Often, an ensemble of multiple simulations is performed, whereby the initial state or the evolution of the predictions is subtly perturbed. The uncertainty resulting from small differences in initial conditions, CI, or storm interactions can thus be reflected (Durran and Weyn, 2016; Bachmann et al., 2020; Bojinski et al., 2023).

Despite these advances in severe weather forecasting techniques, notable forecast errors can occur when NWP models fail to predict CI or the type of convective system. Over complex terrain, such errors are frequent because crucial processes like moisture advection or terrain-induced flow systems are not captured by the model (e.g., Panosetti et al., 2016). For example, on 19 July 2023, a series of record-breaking hailstorms occurred across North Italy as intense supercells initiated over the southern Alps and moved across the Plains. Operational NWP models either failed to produce the storms ~~;~~ or underestimated their intensity (Fig. 4). In this and other challenging cases (e.g., Mandement and Caumont, 2021), a big unknown was how storms ~~will~~would behave once they ~~move~~moved out of the mountains and onto the plains: they may intensify or decay. This transition from mountains to plains seems neither ~~well-understood~~well understood from a storm-dynamics perspective, nor generally ~~well-predicted~~well predicted by NWP models.




Figure 4. Composite radar reflectivity (OPERA) and ESWD hail reports over northern Italy on 19 July 2023 (green: hail diameter 2-5 cm; orange: hail diameter 5 - 9 cm; red: hail diameter > 10 cm), and +7 hours forecast reflectivity by two operational NWP models: ICON-D2 and C-LAEF. The supercell producing the extremely large hail between 18:00 and 19:00 UTC was missed by ICON-D2. C-LAEF forecasted convection, but with weaker maximum intensity.

It is known that NWP models have shortcomings near mountainous terrain as their horizontal resolution is only sufficient to resolve the topography to a certain extent. The smoothing at the smallest resolved scales affects how well the interaction between the surface and the lowest layers can be represented (e.g., Rotach et al., 2022). Furthermore, the parameterizations for unresolved surface-atmosphere exchange processes were mostly developed for flat terrain and much coarser grids, which introduce considerable errors across steep topography (Ceppi et al., 2013). Neglecting or simplifying subscale effects and using climatological fields in parameterizations are perhaps unfitting assumptions in NWP models. This includes challenges of numerical model "grey-zones", where resolved processes and parameterizations overlap (e.g., Chow et al., 2019; Kramer et al., 2020; Kirshbaum, 2020; Wei and Bai, 2024). Methods used for convective-scale data assimilation are not as advanced as for global models (Gustafsson et al., 2018). Improving these terrain-related processes in models is the subject of research within the TEAMx project (Rotach et al., 2022). Modeling experiments within TEAMx regarding CI (Serafin et al., 2024) can be built upon within TIM.

Furthermore, the exceptionally high spatial and temporal resolution of TIM-data is of particular interest given that the latest generation of (pre)operational regional NWP models realize grid spacings of 1 km and below (e.g., Meteoswiss' ICON-D1 and -D05), which is also planned for global models such as the Extremes Digital Twin by the EU Destination Earth programme (ECMWF, 2023). Field campaign data from drones, surface observations, Doppler wind lidars, or other sources are useful in-for model comparison and data assimilation studies (e.g., Adam et al., 2016; Fourrié et al., 2021; Sgoff et al., 2022; Nomokonova et al., 2023; Demortier et al., 2024). Since the presence of orography is still often neglected in current NWP postprocessing algorithms (e.g., Schwartz and Sobash, 2017), such observations would also allow the identification of physically-based predictors and targeted observations that could locally improve the calibration of ensemble prediction, nowcasting systems, and machine learning techniques for severe weather warnings in regions influenced by mountains (e.g., Dabernig et al., 2017).

Another potential area of NWP improvement is a better handling of microphysical processes within storms (Morrison et al., 2020). Many uncertainties remain regarding the actual concentration, properties and number of particles in the various hydrometeor classes (cloud water, rain, graupel, snow, hail) because of a lack of data as research aircraft tend to avoid the most intense parts of those storms (e.g., Heymsfield et al., 2023). However, relatively recent innovations including drones, neutrally buoyant sondes, and vertically-pointing radars (Trömel et al., 2017; Sokol et al., 2018) offer new possibilities in addition to polarimetric radars, disdrometers, or aircraft in-situ observations that are crucial for the improvement of parameterizations (e.g., Taufour et al., 2018).

Overall, addressing the following areas within TIM is likely to lead to progress in NWP performance: (1) the characterization of the pre-convective environment across complex terrain by developing better data assimilation procedures (e.g., Gustafsson et al., 2018; Bachmann et al., 2020), (2) improving parameterizations for surface-air interactions (Ceppi et al., 2013; Goger et al., 2019; Rotach et al., 2022) and microphysics (e.g., Labriola et al., 2019; Malečić et al., 2022), (3) improving data assimilation of ongoing convection in satellite and radar and assimilating data from novel sources. As discussed further in section 2.8, the new generation of geostationary meteorological satellites (MTG, Meteosat Third Generation) enables more

accurate atmospheric moisture determination and a Lightning Imager, which can be assimilated into NWP models and used in nowcasting (Leinonen et al., 2023).

2.7 Climatology and Climate Change

340 Pan-European severe convective storm climatologies have been produced based on different data, such as severe weather reports, lightning, satellite products, and reanalyses (Punge et al., 2017; Taszarek et al., 2019, 2020a; Manzato et al., 2022b; Lombardo and Bitting, 2024). These studies agree relatively well on a main convective season in summer over most of Europe, except over Mediterranean regions, where activity peaks in fall but with relatively low severity (~~with the exception of~~ except flood events). As discussed in the introduction, summertime hotspots for severe convective storms are found near mountain
345 ranges, for example around the Alps, the Pyrenees, the Massif Central, the Black Forest, the Dinaric Alps, and the Carpathian Mountains. Continued improvement of report networks like ESWD, public user reports via App, and SCS detections from satellite, radar, and lightning networks will provide further details on where severe storms typically initiate, become severe, and decay (e.g., Nisi et al., 2018; Manzato et al., 2022b). New data gathered in TIM is limited in time and cannot directly assess these climatological features, long-term variability, and effects of climate change. Yet, results from a constrained field
350 campaign can be of immense value for these research areas by improving process understanding or by probing "analog" storms that occur in conditions presumably representative of future climate conditions (e.g., Lasher-Trapp et al., 2023).

As one effect of global warming, the increase in average temperature throughout the troposphere and associated scaling in moisture content is generally expected to increase CAPE and hence thunderstorm severity (e.g., Romps, 2016; Agard and Emanuel, 2017; Martín et al., 2024). This upward trend is robust for Europe both for past and projected trends of CAPE
355 or related thermodynamics parameters (Mohr and Kunz, 2013; Mohr et al., 2015; Púčik et al., 2017; Rädler et al., 2019; Battaglioli et al., 2023). Stronger updrafts in high CAPE environments are one factor increasing the likelihood of flash floods, in addition to higher moisture content and slower storm motion (e.g., Kahraman et al., 2021). In its Sixth Assessment Report, the Intergovernmental Panel on Climate Change (IPCC) states that climate models consistently project environmental changes that would likely lead to an increase in the frequency and intensity of severe thunderstorms ~~, also involving tornadoes, hail, and~~
360 ~~winds~~ (Calvin et al., 2023, Chapter 11). However, there is low confidence in the specifics of these projected increases.

One reason for the high uncertainty is that many climate models have a bias in low-level moisture and CAPE (e.g., Gopalakrishnan et al., 2025). Another problem is that other ~~impact~~ factors than thermodynamics exist, which could counter-balance the positive trend in CAPE and which are not as well understood (e.g., Raupach et al., 2021). For one, convective inhibition and entrainment are likely to increase in a warmer climate, which could lead to decreases in frequency and coverage of SCS (e.g.,
365 Trapp and Hoogewind, 2016; Chen et al., 2020; Peters et al., 2024). The area affected by hail is also likely to decrease because of an increase in the melting layer depth, although large hail is less affected and is expected to become more frequent due to the positive trend in CAPE (e.g., Gensini et al., 2024). Complicating this, Lin and Kumjian (2022) have indicated that increases in CAPE do not necessarily cause storms to produce larger hail, although the extreme values of CAPE they considered are relatively rare in Europe (e.g., Taszarek et al., 2020b). Furthermore, the frequency and the typical thunderstorm organization
370 could change in response to modifications in the larger-scale circulation and resulting weather patterns (e.g., Mohr et al., 2020;

Ghasemifard et al., 2024). This is important because severe convective wind gusts and extreme precipitations are often associated with MCSs (Saharia et al., 2017; Gatzen et al., 2020; Pacey et al., 2021) while most large hail and strong tornadoes are produced by supercells, at least in the US (Blair et al., 2011; Smith et al., 2012; Blair et al., 2017). For Europe, this is likely similar but only researched in certain regions (Tuovinen et al., 2015; Wapler, 2017; Feldmann et al., 2023; Wells et al., 2024).

375 It is unclear how accurately climate models capture these circulation changes. The accuracy of climate models across mountainous areas may also be impaired by systematic errors in modeling surface-atmosphere exchange processes across complex terrain (e.g., Rotach et al., 2022). Another more general limitation of climate models for convective storm research is that most of them still depend on convection parameterization, ~~although though~~ research with convection-permitting climate simulation is currently ongoing (e.g., Kahraman et al., 2021, 2024; Lenderink et al., 2024; Feldmann et al., 2025). ~~The accuracy of climate models across mountainous areas may also be impaired by systematic errors in modeling surface-atmosphere exchange processes across complex terrain (e.g., Rotach et al., 2022).~~ ~~The record-shattering reports in recent years give the impression that at least in some regions global warming has an influence on severe convective storms, although the influence of economic growth on~~ First results suggest regionally varying trends in Europe but a general decrease in the frequency of weak and an increase in intense hailstorms (Thurnherr et al., 2025; Brennan et al., 2025). However, on the impact side, economic growth and the resulting increase in vulnerability and exposure is likely more important (e.g., SwissRe, 2024). Unfortunately, ~~other observations less sensitive to economic factors and with observations that are not biased by economic factors than reports or insured losses, and that exist over~~ periods long enough for climate trend analysis, are scarce. Augenstein et al. (2024) showed predominantly negative trends in lightning over central Europe, although it is somewhat unclear how representative lightning is for SCS. Hail pad networks indicate less small and more large hail, but these trends are for small regions and mostly not
380
385
390 significant (Manzato et al., 2022b, a).

In short, more research beyond the trends of convective parameters such as CAPE seems required to gain more robust knowledge on trends in severe weather. This is where even a relatively short-term field campaign can help because a better fundamental understanding of the storm dynamics, microphysics, CI, and the production mechanisms of severe convective hazards is needed to know what signals to look for in climate projections and to anticipate how these processes may change in
395 a warmer climate.

2.8 Innovative observations and their use

Advances in severe storm research and forecasting have often been linked to the development and improvement of new observational techniques such as radar or satellite (e.g., Brooks et al., 2019). Many new methods and instruments are currently being developed or implemented which could likewise improve our understanding of severe convective storms. As discussed further
400 in this section, this includes new polarimetric radar algorithms, drone-based observations, swarmsondes, microwave links, and new instruments on geostationary and polar-orbiting satellites. A field campaign on convective storms in the following years would be ideal to test these new platforms and to validate their performance against other observations.

For one, the evolution of storm-top features as seen from satellite imagery ~~has been shown to be strongly is~~ linked to storm severity. These features include above anvil cirrus-plumes, warm wake areas, ~~overshooting top properties~~ large overshooting

405 [tops](#), or storm-top divergence (e.g., Homeyer et al., 2017; Bedka et al., 2018; Murillo and Homeyer, 2019; O'Neill et al., 2021; Murillo and Homeyer, 2022; Scarino et al., 2023). Questions remain [about](#) how these characteristics translate to the Meteosat Third Generation (MTG) satellite imagery and how indicative these features are in rapidly changing conditions near complex terrain. Novel and more precise information on cloud top properties, cloud-ice content, precipitation, and aerosol properties will be provided by the METImage, Ice Cloud Imager (ICI) and the Multi-Viewing Multi-Channel Multi-Polarisation Imaging
410 (3MI) instruments on the polar-orbiting Metop-SG satellites as of 2026.

Lightning is another type of dataset that can be detected ~~remotely~~ with a variety of methods and provides valuable information about the microphysics and severity of a thunderstorm. The new satellite-based MTG Lightning Imager (LI) is similar to the Geostationary Operational Environmental Satellite (GOES) Geostationary Lightning Mapper (GLM) over the Americas (Rudlosky et al., 2020), but requires validation with ground-based lightning sensors before it can be used in research or assimilated in forecast models. In addition to the LI and 2D lightning detections, Lightning Mapping Arrays (LMAs) can be deployed
415 to study the 3D electrification and polarity in thunderstorms, as was done during RELAMPAGO (Medina et al., 2021). Many mechanisms [for cloud electrification](#) have been proposed (e.g., Pruppacher and Klett, 2010), and it has been shown that even slight differences in the environment can have profound effects ~~on cloud electrification~~, which is an active field of research (Carey et al., 2003; Carey and Buffalo, 2007; Fuchs et al., 2015; Schultz et al., 2015; Chmielewski et al., 2018; Ringhausen
420 et al., 2024). Examples to be investigated in TIM are changes in the lightning activity, polarity, or flash size (e.g., Figueras i Ventura et al., 2019; Erdmann and Poelman, 2024) of the storms that traverse complex terrain and undergo rapid changes in response to environmental characteristics such as CAPE, vertical wind shear, cloud base height, or warm cloud depth. These could be linked to [the](#) microphysics of the storm assessed through polarimetric radar (e.g., Kumjian and Ryzhkov, 2008), hail collection, and in-situ observations within clouds. A better understanding of these processes would allow for the development
425 of parameterizations and better lightning prediction in NWP models (e.g., Cummings et al., 2024).

Furthermore, vertical profiles of the atmosphere are invaluable for severe storm forecasting and research, because they allow for quantification of the ingredients for severe storms (e.g., Doswell, 1987; Doswell et al., 1996). As touched on in sections 2.1 and 2.3, the distribution of moisture both vertically and horizontally plays an important role for convective storms. The new 0.9 micron channel of the MTG Flexible Combined Imager (FCI) instrument for the first time allows for the quasi-
430 continuous monitoring of total column moisture content including in the lowest few hundred ~~metres~~ [meters](#) of the troposphere, with sampling rates over Europe of 10 minutes (available now) down to 2.5 minutes (by around 2027). Such data could help forecast CI ~~, back-building convection, as well as local boundaries~~ [through detecting local boundaries or identify potential for backbuilding convection](#). The FCI as well as the new geostationary infrared sounder (IRS) on board ~~of~~ the MTG-S1, which can also provide vertical profiles of temperature and moisture, require comparison with in-situ soundings and ground-
435 based radiometer soundings near orography and in proximity to SCS. The potential value of IRS as a complement to standard soundings has been demonstrated at the ESSL Testbed in the quasi-operational setting (Groenemeijer, 2019). Although there are limitations ~~to detect in detecting~~ low-level moisture near the ground with this data, [the](#) availability of infrared soundings from geostationary orbit at 30-min resolution over Europe will add significant value and overcome the current limitations on temporal resolution of such data from polar-orbiting instruments such as IASI on Metop. Despite this disadvantage of polar-

440 orbiting satellites, EarthCARE ([Wehr et al., 2023](#)) is promising for profiling cloud microphysical properties and aerosols with an overpass time around 13:30 UTC over Europe, a typical mature convective storm period.

Recent alternative profiling methods are commercial microwave links (CMLs) and uncrewed aircraft systems (UAS or *meteodrones*). As an opportunistic remote-sensing tool, CMLs typically used in cell towers are rapidly evolving and E-Band CMLs are increasingly installed in several countries to enable 5G. E-band CMLs can provide information on water vapor
445 variability (e.g., Fencil et al., 2021). Research is needed to fully exploit the fast-developing K-, E-, and W-band commercial microwave link (CML) networks for use in precipitation and water vapor monitoring in terms of retrievals and direct data assimilation at different frequencies. Regarding UAS, in-situ aerial measurements with ~~Meteodrones~~ [meteodrones](#) of the lower troposphere were tested as high up as 6 km above ground (Guay et al., 2023), compared against radiosonde and remote-sensing instruments (Hervo et al., 2023) and found to meet [the](#) WMO requirements (WMO, 2024). Despite the limitations of drones due
450 to strict airspace regulations and maximum operable wind speeds, they could offer a low-cost way to collect high-resolution 3D data that ~~are~~ [is](#) difficult to attain with conventional point measurements.

Other drone applications are photographic documentation purposes. Cloud, dust, rain- and hail-shaft development can be observed in time-lapse mode without the limitations imposed by buildings or vegetation being obstacles to the ground-based observers. Photographic post-event analysis of rain damage, hail damage, and wind damage (ESSL, 2023b) as well as analysis
455 of hail size distribution are feasible based on images taken from drones (Soderholm et al., 2020; Lainer et al., 2024). High-resolution satellite imagery can be used in a similar way for some applications (Shikhov and Chernokulsky, 2018; Gallo et al., 2019; Kunkel et al., 2023).

The global advent of dual-polarization radar data has enabled the identification of polarimetric signatures linked to the dynamic and microphysical processes in storms (e.g., Kumjian and Ryzhkov, 2008; Kumjian, 2013; Dawson et al., 2014).
460 Because the severity is intrinsically related to the presence of updrafts and size-sorting signatures, many applications of polarimetric variables are known. For example, differential reflectivity (Z_{DR}) and co-polar cross correlation coefficient can be used to assess hail and lightning (Snyder et al., 2015; Schmidt, 2020; Trömel et al., 2023; Klaus and Krause, 2024; Rombeek et al., 2024; Aregger et al., 2024), the discrimination between tornadic and nontornadic supercells (Loeffler et al., 2020), and the early prediction of tornadoes (French and Kingfield, 2021). Another application is the spectral analysis of ~~the polarimetric~~
465 ~~variables~~ [polarimetric variables](#), which can explain microphysical processes of hydrometeors detected at different radar beam heights (Lakshmi et al., 2024). However, operational radar data quality is often limited near complex terrain (Feldmann et al., 2021; Germann et al., 2022). For many radar diagnostics related to storm severity like Probability of Hail (POH, Delobbe and Holleman, 2006), Vertical Integrated Liquid (VIL, Greene and Clark, 1972), Maximum Estimated Size of Hail (MESH, Witt et al., 1998), or Z_{DR} columns, a high vertical resolution is needed, and beam shielding by mountains is problematic. Another
470 challenge is the lack of low-level data, which limits the use of Dual-Doppler wind retrievals (Brook et al., 2021) and radar storm signatures such as hook echoes or the Z_{DR} inflow arc (Fujita, 1958; Kumjian and Ryzhkov, 2009). Furthermore, QPE in mountains can lead to errors in flood assessments due to strong reflectivity gradients below the melting level (e.g., Chen et al., 2023). The usage of mobile research X-band radars in TIM can complement and address limitations and gaps of operational radars. For instance, 3D wind fields can be retrieved from dual-Doppler analysis in areas that are scanned by more than one

475 radar. Such retrieval is beneficial for updraft detection and validation (Evaristo et al., 2021) as well as for hail and tornado re-
search (~~Sessa and Trapp, 2020; Pounds et al., 2024, e.g.,~~)(~~Sessa and Trapp, 2020; Pounds et al., 2024~~). Mobile radars can also
allow for an improved classification of hydrometeor classes (Besic et al., 2016) and tracking of convective cells (Kollias et al.,
2024).

Lastly, *swarmsondes* (Markowski et al., 2018) are balloon-borne probes that act as a pseudo-Lagrangian drifter and that
480 are drawn through the storm. Swarmsondes can be released in dozens per thunderstorm and were found to fill the lack of
reliable, aboveground, thermodynamic observations near and in convective storms. Swarmsondes in Europe were first used in
the Swabian MOSES campaign in 2021 (Kunz et al., 2022). A slightly adapted version called *Hailsonde* can mimic and analyze
the trajectory and growth of falling hailstones within thunderstorm updrafts (~~Soderholm, 2024~~)(~~Soderholm et al., 2025~~). Lack
of such data from within thunderstorms is perhaps one of the biggest limitations in the field of convective storms. Much of
485 our knowledge and model parameterizations of microphysical properties, such as hydrometeor concentrations within storm
clouds, is based on storm-penetrating flights of the T-28 aircraft, which was retired around 20 years ago (e.g., Allen et al.,
2020; Heymsfield et al., 2023). Hence, gathering such data with aircrafts or sondes in a field campaign could allow for big
leaps in our understanding and forecast capability.

3 Conclusions

490 3.1 Summary

As reviewed in section 2 and based on our White Paper survey, a wide variety of research topics (RT) exists regarding severe
convective storms near mountains. Fig. 5 provides an overview of these topics and highlights the methodologies and data
sources involved. The key research questions can be summarized as follows.

- 495 (1) Why do severe weather reports cluster near mountain ranges? To answer this, the terrain-induced perturbations of
environmental parameters like CAPE and wind shear will need to be weighed against storm-scale processes and CI in
different regions and scenarios. (RT ~~1, 2, 3, 4~~~~2.1, 2.2, 2.3, 2.4~~)
- (2) What are gaps in our understanding of the storm-internal processes leading to large hail, strong winds, lightning,
tornadoes, and extreme precipitation and how are these processes modulated by the orography? Such basic understanding
is also essential to better anticipate the impact of global warming on severe storms. (RT ~~1, 4, 5, 7, 8~~~~2.1, 2.4, 2.5, 2.7, 2.8~~)
- 500 (3) What are the strengths and limits of new observation techniques from satellites, radars, or drones? TIM will provide
the unique opportunity to compare these datasets to high-resolution sampling of individual storms and of heterogeneous
environments near complex terrain. (RT ~~2, 4, 6, 8~~~~2.2, 2.4, 2.6, 2.8~~)
- (4) What observations are most important to improve numerical weather prediction models and nowcasting? Higher-
resolution datasets (than currently operational) of the pre-storm environment and thunderstorm properties in TIM can




Figure 5. Overview of the scopes **and scale** of TIM research subjects (dark gray boxes), as well as the corresponding scales, methodologies, and datasets (light gray boxes).

505 be used in model studies to help answer this question and also to improve data assimilation and parameterizations of orographic or microphysical processes. (RT 1, 4, 62.1, 2.4, 2.6)

The importance of these research topics is emphasized by the observed increase in related losses in recent years and the projected increase in severe storm activity because of climate change. ESSL is therefore coordinating plans for the TIM field campaign as a first-of-its-kind Pan-European campaign on severe convective storms. These plans are currently being worked
 510 out in regular meetings with all partners. This White Paper may serve as a scientific basis for further planning of the campaign and the writing of funding proposals. Since the ultimate scope of TIM will depend on the success of these proposals and further planning, a detailed campaign plan cannot be presented here. However, to give a reader an impression of how we envision TIM, the rest of this section briefly discusses the planned concept.

3.2 Preliminary TIM campaign rationale

515 Figure 6 provides an example overview of the data collection. The TIM campaign can profit from the wealth of already operational observations across Europe (red text in Fig. 6). This includes meteorological stations with ground data, regular vertical soundings, radars, satellites, lightning sensors, and public reports. Some of these datasets are currently not openly available to researchers. One example is the European radar network, which is globally unique in its density and coverage and provides excellent opportunities for analysis over a large region but is often only analyzed for each country separately. Data




Figure 6. Overview of the observations planned in TIM. Since the campaign is still in an early phase, this can be seen as an ideal or example setup. Red colored text denotes at least partially operational data while black text would be part of TIM operations.

520 sharing in TIM could therefore accelerate the development of new radar-based algorithms for severe weather prediction ~~for~~
~~instance~~-based on machine learning (Leinonen et al., 2023; Forcadell et al., 2024). To this end, part of the funding will be
allocated to create a central data hub for systematic data collection, exchange, and long-term archiving. Ideally, TIM will be a
first step towards permanent sharing of operational datasets for SCS research in the future.

However, additional data from the TIM campaign is needed, especially in and around thunderstorms and at higher spatial
525 and temporal resolution than currently provided by operational networks (black text in Fig. 6). This includes mobile radars,
swarmsondes, drones, high-resolution soundings, and survey teams. Such observations are often difficult to obtain due to the
fast-evolving and small-scale nature of convective storms. Thus, to sample a high number of cases it is necessary to be mobile.
On the other hand, some of the research objectives described herein require specific measurements which can only be supplied
by fixed instruments, such as lidars, hail disdrometers, LMAs, or high-density station networks, which cannot be re-located on
530 a daily basis (e.g., Emeis et al., 2018).

A field campaign with both mobile and fixed-site measurement strategies offers the unique opportunity to address this
dilemma. To sample individual storms and their hazards, TIM will deploy several mobile teams to a region where high-impact
storms are expected on a given day. Most severe weather events cluster in time and space, so mobile teams can collect data
in the same region over multiple days or by traveling from west to east ~~within~~-with a conditionally unstable airmass. At the
535 same time, multiple permanent TIM focus sites will be set up over the convective season to continuously monitor the same




Figure 7. Map of the TIM domain with regions of interest (ROI). ROI were identified based on in-domain ESWD severe hail reports (c.f. Fig. 1). Red shading indicates primary hotspots with high report densities from kernel-density estimation (KDE). To represent secondary clusters, reports grouped by density-based spatial clustering of applications (DBSCAN) are indicated with lighter-shading dashed red contours. Also shown are mountain ranges of the GMBA mountain inventory (Snehlage et al., 2022b, a) and built-up surfaces (Pesaresi and Politis, 2023). Black triangles show official TIM partner institutions headquarter locations as of May 2024-2025 (see list below).

region, supplemented by the mobile teams when severe storm activity is expected in the same location. A similar strategy has been successfully used in several campaigns focused on convective storms in target regions (e.g., PERILS, RELAMPAGO). A unique aspect of the current plans for TIM is that focus sites will be maintained by participating institutions based on their individual funding and research focus. In parallel, core funding of TIM will support mobile teams to collect additional high-resolution data. This flexible approach allows the effective use of funds and overlap of different research topics (Fig. 5).

Figure 7 highlights potential regions of interest (~~ROI~~ROIs), where ~~ROI~~ROIs are identified through clustering analyses of ESWD severe hail reports. This is one way to objectively define ~~ROI~~ROIs and is used as an example here to highlight the most important ~~ROI~~ROIs: the Po valley and the border region between Austria and Slovenia south of the Alps; the valleys near the Swiss Jura, Black Forest, Swabian Jura, and Southern Bavaria north of the Alps; and regions around the Pyrenees, Massif Central, Dinaric Alps, Bohemian Massif, and Carpathian mountains. Many secondary ~~ROI~~ROIs are also visible in Fig. 7. It is important that ESWD reports can be subject to reporting biases, so some of these secondary ~~ROI~~ROIs in less-populated regions, such as around the Carpathian mountains, are likely more impacted by SCS than depicted here (Šinger and Púčik, 2020). Nevertheless, the reports show good agreement with existing radar-based climatologies in Europe (e.g., Punge and Kunz, 2016; Fluck et al., 2021) and allow ~~to identify~~identifying regions of high impact. Which focus sites can ultimately be investigated in depth with TIM will depend on the priorities and funding of individual partner institutions.

Besides geographical and climatological differences among the regions, also the densities, lengths, and types of operational observations vary substantially. The collaboration between many institutes brings together the necessary expertise to handle these diverse data needs, which no single institution can cover on its own. TIM already has a long list of partner institutions officially engaged who also contributed to this article (see below and Fig. 7). A core investigative team lead by ESSL will be established to lead these efforts and to ensure that possible conflicting interests over many institutions can be coordinated. Working groups on individual subtopics, such as ~~Climatology, Radar, air-borne~~climatology, radar, airborne data, and idealized cloud modeling, are currently being established and will be expanded to specify the plans for TIM. Because of the spectrum of topics and because of the high spatio-temporal variability of convective storms (e.g., Taszarek et al., 2020a), the campaign is planned for three consecutive convective storm seasons, currently from 2027 to 2029. This is followed by two years of data analysis to ensure the effective and coordinated use of the observations. This observational data will be supplemented by numerical modeling experiments tailored to the respective research questions and focus regions.

Furthermore, ESSL testbeds during and after the campaign provide ample opportunities to foster exchange with European severe weather forecasters. High-quality forecasts focused on severe convective storms are also essential for the coordination of the campaign. Like in previous severe storm campaigns, a forecast team will be responsible for briefing the research groups on an active campaign day. The forecasts can benefit from the bundled resources of partner institutions and experience of projects like the European Storm Forecast Experiment (ESTOFEX), Previsione Temporali (PRETEMP), as well as local forecast offices ~~who that~~ know the characteristic meteorology of a certain mountain range.

Thanks to the globally-connected severe storms community, TIM can also profit from the experience of international researchers. For example, TIM researchers could coordinate with, and learn from the ICECHIP and TEAMx campaigns (Adams-Selin et al., 2022; Rotach et al., 2022). Their research foci (hailstorms in the US Great Plains and terrain effects on the atmosphere) can be complemented and linked by TIM's focus on severe convection near mountains. Exchanges with the Northern Hail and Tornado Projects (e.g., Brimelow et al., 2023), the LIFT campaign (Kunz et al., 2024), and observational efforts of the Australian Bureau of Meteorology (~~e.g., Soderholm, 2024~~) (e.g., Soderholm et al., 2025) are already ongoing. TIM has the ambition to foster collaboration with international researchers as well as stakeholders and policy makers, educate through

575 public and media outreach, and introduce students to the field of convective storms, which will ultimately help to meet the challenges that the increasing risk of SCS ~~pose~~poses to European society.

TIM Partners

Extended co-authors of this article (in alphabetical order):

580 Martin Adamovsky¹⁰, Clotilde Augros¹², Ulrich Blahak¹³, Vojtěch Bližňák¹⁴, Stephan Bojinski¹⁵, Tobias Bösle¹⁶, François
Bouttier¹², Massimiliano Burlando¹⁷, Federico Canepa¹⁷, Orietta Cazzuli¹⁸, Xavier Calbet¹⁹, Alessandro Ceppi³³, Fleur
Couvreur¹², Kálmán Csirmaz²¹, Tamás Csonka²¹, Stavros Dafis²², Mária Derková²³, Francesco Domenichini²⁴, Grzegorz
Duniec²⁶, Raquel Evaristo²⁷, Tomáš Fedor²⁵, Massimo Enrico Ferrario²⁴, Michael Frech¹², Enrico Gambini³³, Jaroslav
Hofierka²⁵, Ákos Horváth²¹, Adam Jaczewski²⁶, Kornél Komjáti²¹, Vincent Klaus²⁸, Zsófia Kocsis²¹, Michael Kunz³, Máté
585 Kurcsics²¹, Robert Kvak¹⁴, Konstantinos Lagouvardos²², Katharina Lengfeld¹², Marco Mancini³³, Marc Mandement¹², Olivia
Martius⁶, Massimo Milelli²⁰, Samuel Monhart³¹, Gian Paolo Minardi¹⁸, Antonio Parodi²⁰, Giovanni Ravazzani³³, Ricardo
Reinoso-Rondinel³⁴, Didier Ricard¹², David Rýva¹⁰, Tobias Scharbach²⁷, Stefano Serafin³², André Simon²³, Miroslav Šinger²³,
Zbyněk Sokol¹⁴, Gabriella Szépszó²¹, Silke Trömel²⁷, Benoît Vié¹², Clemens Wastl²⁹, Christoph Wittmann²⁹, Petr Zacharov¹⁴,
Matteo Zanetti¹⁸, Dino Zardi³⁰

590 Official Partner institutions of the TIM campaign that have signed letters of intent (as of ~~11/03~~May 2025):

Albert-Ludwigs-Universität Freiburg (University of Freiburg)

²⁹Austrian Federal Institute for Geology, Geophysics, Climatology, and Meteorology (GeoSphereAustria, former ZAMG)

Austro Control (AustroControl)

²⁰CIMA Research Foundation (CIMA)

595 ¹²CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

¹⁰Czech Hydrometeorological Institute (ČHMÚ)

¹⁴Czech Academy of Sciences - Institute of Atmospheric Physics (CAS UFA)

¹³Deutscher Wetterdienst (DWD)

³³Department of Civil and Environmental Engineering (D.I.C.A.), Politecnico di Milano, Milano, Italy

600 Državni hidrometeorološki zavod (DHMZ)

¹⁵European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT)

²¹HungaroMet Hungarian Meteorological Service (HungaroMet)

²⁶Institute of Meteorology and Water Management - National Research Institute Poland (IMGW)

Institute of Atmospheric Sciences and Climate (ISAC), Consiglio Nazionale delle Ricerche (CNR), Italy

605 Karlsruhe Institute of Technology (KIT)

KU Leuven, Department of Civil Engineering

~~¹²CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France~~²²National Observatory of Athens - Institute for
Environmental Research and Sustainable Development (NOA)

²⁵Pavol Jozef Šafárik University (UPJS)

610 ³³~~Department of Civil and Environmental Engineering (D.I.C.A.), Politecnico di Milano, Milano, Italy~~²⁴Regional Agency for
Environmental Protection and Prevention of Liguria (ARPA Liguria)
¹⁸Regional Agency for Environmental Protection of Lombardia (ARPA Lombardia)
²⁴Regional Agency for Environmental Protection and Prevention of the Veneto (ARPA Veneto) (ARPA Veneto)
²³Slovak Hydrometeorological Institute (SHMÚ)
615 Slovenian Environment Agency (ARSO)
⁶University of Bern - Institute of Geography
¹⁷University of Genoa

Other contributing institutions to this article:

620 ¹⁶Deutsches Zentrum für Luft- und Raumfahrt (DLR)
¹⁹Agencia Estatal de Meteorología (AEMET)
²⁷University of Bonn
²⁸Universität für Bodenkultur (BOKU)
³⁰University of Trento, Department of Civil Environmental and Mechanical Engineering (DICAM)
625 ³¹Federal Office of Meteorology and Climatology MeteoSwiss
³²University of Vienna, Department of Meteorology and Geophysics
³⁴KU Leuven, Department of Civil Engineering

Data availability. Severe weather reports are available from the European Severe Weather Database (ESWD, eswd.eu).

Author contributions. The first author (Fischer) oversaw and organized the writing and provided Figs. 2, 3, and 6. Authors Groenemeijer,
630 Holzer, Feldmann, Schröer, Battaglioli, Schielicke, Púčik, Gatzen and Antonescu contributed through writing tasks or Figures. The extended
co-authors (see list above) provided initial writing topics including short paragraphs which were incorporated into the article by the first
author. All contributors proofread the manuscript draft and provided references and suggestions on specific topics.

Competing interests. No competing interests are present.

Acknowledgements. ESSL preparatory work for TIM is supported by the Provincial Government of Lower Austria Science Fund grant
635 number K3-F-190/013-2023. The two anonymous reviewers provided exceptionally detailed and helpful suggestions. Feedback on an early

Table 1. Glossary of abbreviations that are repeatedly used

CAPE	Convective Available Potential Energy
CCN	Cloud Condensation Nuclei
CI	Convection Initiation
CIN	Convective Inhibition
CML	Commercial Microwave Link
ESSL	European Severe Storms Laboratory
IF	International Fujita (Scale)
INP	Ice Nucleating Particle
LMA	Lightning Mapping Array
MCS	Mesoscale Convective System
MESH	Maximum Estimated Size of Hail
POH	Probability of Hail
QPE	Quantitative Precipitation Estimation
ROI	Region of Interest
RT	Research Topic
SCS	Severe Convective Storms
VIL	Vertical Integrated Liquid

version of the manuscript by [Drs.](#) Rebecca Adams-Selin, Karen Kosiba, Matthew Kumjian, Kelly Lombardo, Paul Markowski, James Marquis, Stephen Nesbitt, Bruno Ribeiro, Joshua Soderholm, Jeff Trapp, and Joshua Wurman is greatly appreciated.

References

- Adam, S., Behrendt, A., Schwitalla, T., Hammann, E., and Wulfmeyer, V.: First assimilation of temperature lidar data into an NWP model: impact on the simulation of the temperature field, inversion strength and PBL depth, *Quarterly Journal of the Royal Meteorological Society*, 142, 2882–2896, <https://doi.org/https://doi.org/10.1002/qj.2875>, 2016.
- Adams-Selin, R. D., Allen, J., Gensini, V., and Heymsfield, A.: ICECHIP: Closing Critical Observational Gaps in Hail Research, North American Hail Workshop, Boulder, Colorado, 2022.
- Adler, B., Kalthoff, N., Kohler, M., Handwerker, J., Wieser, A., Corsmeier, U., Kottmeier, C., Lambert, D., and Bock, O.: The variability of water vapour and pre-convective conditions over the mountainous island of Corsica, *Quarterly Journal of the Royal Meteorological Society*, 142, 335–346, <https://doi.org/10.1002/qj.2545>, 2016.
- Agard, V. and Emanuel, K.: Clausius-Clapeyron scaling of peak CAPE in continental convective storm environments, *Journal of the Atmospheric Sciences*, 74, 3043–3054, <https://doi.org/10.1175/JAS-D-16-0352.1>, 2017.
- Allen, J. T., Giammanco, I. M., Kumjian, M. R., Jurgen Punge, H., Zhang, Q., Groenemeijer, P., Kunz, M., and Ortega, K.: Understanding Hail in the Earth System, *Reviews of Geophysics*, 58, 1–49, <https://doi.org/10.1029/2019RG000665>, 2020.
- Antonescu, B., Schultz, D. M., Holzer, A., and Groenemeijer, P.: Tornadoes in Europe: An underestimated threat, *Bulletin of the American Meteorological Society*, 98, 713–728, <https://doi.org/10.1175/BAMS-D-16-0171.1>, 2017.
- Aregger, M., Martius, O., Germann, U., and Hering, A.: Differential reflectivity columns and hail – linking C-band radar-based estimated column characteristics to crowdsourced hail observations in Switzerland, pp. 1–21, <http://arxiv.org/abs/2410.10499>, 2024.
- Auf der Maur, A. and Germann, U.: A Re-Evaluation of the Swiss Hail Suppression Experiment Using Permutation Techniques Shows Enhancement of Hail Energies When Seeding, *Atmosphere*, 12, <https://doi.org/10.3390/atmos12121623>, 2021.
- Augenstein, M., Mohr, S., and Kunz, M.: Influence of the North Atlantic Oscillation on annual spatio-temporal lightning clusters in western and central Europe, *EGUspHERE*, 2024, 1–30, <https://doi.org/10.5194/egusphere-2024-2804>, 2024.
- Bachmann, K., Keil, C., Craig, G. C., Weissmann, M., and Welzbacher, C. A.: Predictability of deep convection in idealized and operational forecasts: Effects of radar data assimilation, orography, and synoptic weather regime, *Monthly Weather Review*, 148, 63–81, <https://doi.org/10.1175/MWR-D-19-0045.1>, 2020.
- Bagaglini, L., Ingrosso, R., and Miglietta, M. M.: Synoptic patterns and mesoscale precursors of Italian tornadoes, *Atmospheric Research*, 253, 105 503, <https://doi.org/10.1016/j.atmosres.2021.105503>, 2021.
- Barthlott, C., Czajka, B., Kunz, M., Saathoff, H., Zhang, H., Böhmmländer, A., Gasch, P., Handwerker, J., Kohler, M., Wilhelm, J., Wieser, A., and Hoose, C.: The impact of aerosols and model grid spacing on a supercell storm from Swabian MOSES 2021, *Quarterly Journal of the Royal Meteorological Society*, pp. 1–23, <https://doi.org/10.1002/qj.4687>, 2024.
- Battaglioli, F., Groenemeijer, P., Púčik, T., Taszarek, M., Ulbrich, U., and Rust, H.: Modeled Multidecadal Trends of Lightning and (Very) Large Hail in Europe and North America (1950–2021), *Journal of Applied Meteorology and Climatology*, 62, 1627–1653, <https://doi.org/10.1175/JAMC-D-22-0195.1>, 2023.
- Bedka, K. M., Allen, J. T., Punge, H. J., Kunz, M., and Simanovic, D.: A long-term overshooting convective cloud-top detection database over Australia derived from MTSAT Japanese Advanced Meteorological Imager Observations, *Journal of Applied Meteorology and Climatology*, 57, 937–951, <https://doi.org/10.1175/JAMC-D-17-0056.1>, 2018.

- Benjamin, S. G., Dévényi, D., Weygandt, S. S., Brundage, K. J., Brown, J. M., Grell, G. A., Kim, D., Schwartz, B. E., Smirnova, T. G., Smith, T. L., and Manikin, G. S.: An hourly assimilation-forecast cycle: The RUC, *Monthly Weather Review*, 132, 495–518, [https://doi.org/10.1175/1520-0493\(2004\)132<0495:AHACTR>2.0.CO;2](https://doi.org/10.1175/1520-0493(2004)132<0495:AHACTR>2.0.CO;2), 2004.
- Besic, N., FiguerasVentura, J., Grazioli, J., Gabella, M., Germann, U., and Berne, A.: Hydrometeor classification through statistical clustering of polarimetric radar measurements: A semi-supervised approach, *Atmospheric Measurement Techniques*, 9, 4425–4445, <https://doi.org/10.5194/amt-9-4425-2016>, 2016.
- Blair, S. F., Deroche, D. R., Boustead, J. M., Leighton, J. W., Barjenbruch, B. L., and Gargan, W. P.: Radar-Based Assessment of the Detectability of Giant Hail, *E-Journal of Severe Storms Meteorology*, 6, 1–30, <https://doi.org/10.55599/ejssm.v6i7.34>, 2011.
- Blair, S. F., Laflin, J. M., Cavanaugh, D. E., Sanders, K. J., Currens, S. R., Pullin, J. I., Cooper, D. T., Deroche, D. R., Leighton, J. W., Fritchie, R. V., Mezeul, M. J., Goudeau, B. T., Kreller, S. J., Bosco, J. J., Kelly, C. M., and Mallinson, H. M.: High-resolution hail observations: Implications for NWS warning operations, *Weather and Forecasting*, 32, 1101–1119, <https://doi.org/10.1175/WAF-D-16-0203.1>, 2017.
- Bojinski, S., Blaauboer, D., Calbet, X., de Coning, E., Debie, F., Montmerle, T., Nietosvaara, V., Norman, K., Bañón Peregrín, L., Schmid, F., Strelec Mahović, N., and Wapler, K.: Towards nowcasting in Europe in 2030, *Meteorological Applications*, 30, 1–27, <https://doi.org/10.1002/met.2124>, 2023.
- Bosart, L. F., Seimon, A., LaPenta, K. D., and Dickinson, M. J.: Supercell tornadogenesis over complex terrain: The Great Barrington, Massachusetts, tornado on 29 May 1995, *Weather and Forecasting*, 21, 897–922, <https://doi.org/10.1175/WAF957.1>, 2006.
- Bougeault, P., Binder, P., Buzzi, A., Dirks, R., Houze, R., Kuettner, J., Smith, R. B., Steinacker, R., and Volkert, H.: The MAP Special Observing Period, *Bulletin of the American Meteorological Society*, 82, 433–462, [https://doi.org/10.1175/1520-0477\(2001\)082<0433:TMSOP>2.3.CO;2](https://doi.org/10.1175/1520-0477(2001)082<0433:TMSOP>2.3.CO;2), 2001.
- Bowen, S., Kerschner, B., and Zheng Ng, J.: Natural catastrophe and climate report 2023, accessed 2024-02-02, https://www.ajg.com/gallagher/-/media/files/gallagher/gallagherre/news-and-insights/2024/january/natural_catastrophe_and_climate_report_2023.pdf, 2024.
- Brennan, K. P. and Wilhelm, L.: Saharan dust linked to European hail events, *EGUsphere*, 2024, 1–19, <https://doi.org/10.5194/egusphere-2024-3924>, 2024.
- Brennan, K. P., Thurnherr, I., Sprenger, M., and Wernli, H.: Insights from hailstorm track analysis in European climate change simulations, *EGUsphere*, 2025, 1–29, <https://doi.org/10.5194/egusphere-2025-918>, 2025.
- Brimelow, J. C., Kopp, G. A., and Sills, D. M. L.: The Northern Hail Project: A renaissance in hail research in Canada, 11th European Conference on Severe Storms, Bucharest, Romania, <https://doi.org/https://doi.org/10.5194/ecss2023-170>, 2023.
- Brook, J. P., Protat, A., Soderholm, J., Carlin, J. T., McGowan, H., and Warren, R. A.: Hailtrack—improving radar-based hailfall estimates by modeling hail trajectories, *Journal of Applied Meteorology and Climatology*, 60, 237–254, <https://doi.org/10.1175/JAMC-D-20-0087.1>, 2021.
- Brooks, H. E., Doswell, C. A., Zhang, X., Chernokulsky, A. M. A., Tochimoto, E., Hanstrum, B., de Lima Nascimento, E., Sills, D. M. L., Antonescu, B., and Barrett, B.: A Century of Progress in Severe Convective Storm Research and Forecasting, *Meteorological Monographs*, 59, 18.1–18.41, <https://doi.org/10.1175/amsmonographs-d-18-0026.1>, 2019.
- Brown, T. and Giammanco, I.: IBHS multi-faceted hail research program, *RCI*, 31, 5–8, 2013.
- Browning, K. A. and Foote, G. B.: Airflow and hail growth in supercell storms and some implications for hail suppression, *Quarterly Journal of the Royal Meteorological Society*, 102, 499–533, <https://doi.org/https://doi.org/10.1002/qj.49710243303>, 1976.

- 710 Burlando, M., Miglietta, M. M., Avolio, E., Bechini, R., Cassola, F., De Martin, F., Lagasio, M., Milelli, M., Parodi, A., and Romanic, D.: The –WIND RISK project: nowcast and simulation of thunderstorm outflows, *Bulletin of Atmospheric Science and Technology*, 5, 1–10, <https://doi.org/10.1007/s42865-024-00079-6>, 2024.
- Calbet, X., Carbajal Henken, C., Desouza-Machado, S., Sun, B., and Reale, T.: Horizontal small-scale variability of water vapor in the atmosphere: implications for intercomparison of data from different measuring systems, *Atmospheric Measurement Techniques*, 15, 7105–7118, <https://doi.org/10.5194/amt-15-7105-2022>, 2022.
- 715 Calvin, K., Dasgupta, D., Krinner, G., Mukherji, A., Thorne, P. W., Trisos, C., Romero, J., Aldunce, P., Barrett, K., Blanco, G., Cheung, W. W. L., Connors, S., Denton, F., Diongue-Niang, A., Dodman, D., Garschagen, M., Geden, O., Hayward, B., Jones, C., Jotzo, F., Krug, T., Lasco, R., Lee, Y.-Y., Masson-Delmotte, V., Meinshausen, M., Mintenbeck, K., Mokssit, A., Otto, F. E. L., Pathak, M., Pirani, A., Poloczanska, E., Pörtner, H.-O., Revi, A., Roberts, D. C., Roy, J., Ruane, A. C., Skea, J., Shukla, P. R., Slade, R., Slangen, A., Sokona, Y., Sörensson, A. A., Tignor, M., van Vuuren, D., Wei, Y.-M., Winkler, H., Zhai, P., Zommers, Z., Hourcade, J.-C., Johnson, F. X., Pachauri, S., Simpson, N. P., Singh, C., Thomas, A., Totin, E., Alegría, A., Armour, K., Bednar-Friedl, B., Blok, K., Cissé, G., Dentener, F., Eriksen, S., Fischer, E., Garner, G., Guivarch, C., Haasnoot, M., Hansen, G., Hauser, M., Hawkins, E., Hermans, T., Kopp, R., Leprince-Ringuet, N., Lewis, J., Ley, D., Ludden, C., Niamir, L., Nicholls, Z., Some, S., Szopa, S., Trewin, B., van der Wijst, K.-I., Winter, G., Witting, M., Birt, A., and Ha, M.: IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland., <https://doi.org/10.59327/ipcc/ar6-9789291691647>, 2023.
- 720 Canepa, F., Burlando, M., and Solari, G.: Vertical profile characteristics of thunderstorm outflows, *Journal of Wind Engineering and Industrial Aerodynamics*, 206, <https://doi.org/10.1016/j.jweia.2020.104332>, 2020.
- Canepa, F., Romanic, D., Hangan, H., and Burlando, M.: Experimental translating downbursts immersed in the atmospheric boundary layer, *Journal of Wind Engineering and Industrial Aerodynamics*, 243, 105 570, <https://doi.org/10.1016/j.jweia.2023.105570>, 2023.
- 730 Carey, L. D. and Buffalo, K. M.: Environmental control of cloud-to-ground lightning polarity in severe storms, *Monthly Weather Review*, 135, 1327–1353, <https://doi.org/10.1175/MWR3361.1>, 2007.
- Carey, L. D., Rutledge, S. A., and Petersen, W. A.: The relationship between severe storm reports and cloud-to-ground lightning polarity in the contiguous United States from 1989 to 1998, *Monthly Weather Review*, 131, 1211–1228, [https://doi.org/10.1175/1520-0493\(2003\)131<1211:TRBSSR>2.0.CO;2](https://doi.org/10.1175/1520-0493(2003)131<1211:TRBSSR>2.0.CO;2), 2003.
- 735 Ceppi, A., Ravazzani, G., Salandin, A., Rabuffetti, D., Montani, A., Borgonovo, E., and Mancini, M.: Effects of temperature on flood forecasting: Analysis of an operative case study in Alpine basins, *Natural Hazards and Earth System Science*, 13, 1051–1062, <https://doi.org/10.5194/nhess-13-1051-2013>, 2013.
- Chen, J., Dai, A., Zhang, Y., and Rasmussen, K. L.: Changes in convective available potential energy and convective inhibition under global warming, *Journal of Climate*, 33, 2025–2050, <https://doi.org/10.1175/JCLI-D-19-0461.1>, 2020.
- 740 Chen, J. Y., Reinoso-Rondinel, R., Trömel, S., Simmer, C., and Ryzhkov, A.: A Radar-Based Quantitative Precipitation Estimation Algorithm to Overcome the Impact of Vertical Gradients of Warm-Rain Precipitation: The Flood in Western Germany on 14 July 2021, *Journal of Hydrometeorology*, 24, 521–536, <https://doi.org/10.1175/JHM-D-22-0111.1>, 2023.
- Chmielewski, V. C., Bruning, E. C., and Ancell, B. C.: Variations of Thunderstorm Charge Structures in West Texas on 4 June 2012, *Journal of Geophysical Research: Atmospheres*, 123, 9502–9523, <https://doi.org/10.1029/2018JD029006>, 2018.
- 745 Chow, F. K., Schär, C., Ban, N., Lundquist, K. A., Schlemmer, L., and Shi, X.: Crossing multiple gray zones in the transition from mesoscale to microscale simulation over complex terrain, *Atmosphere*, 10, <https://doi.org/10.3390/atmos10050274>, 2019.

- Cummings, K. A., Pickering, K. E., Barth, M. C., Bela, M. M., Li, Y., Allen, D., Bruning, E., MacGorman, D. R., Ziegler, C. L., Biggerstaff, M. I., Fuchs, B., Davis, T., Carey, L., Mecikalski, R. M., and Finney, D. L.: Evaluation of Lightning Flash Rate Parameterizations in a Cloud-Resolved WRF-Chem Simulation of the 29–30 May 2012 Oklahoma Severe Supercell System Observed During DC3, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD039492, <https://doi.org/https://doi.org/10.1029/2023JD039492>, 2024.
- Ćurić, M. and Janc, D.: Differential heating influence on hailstorm vortex pair evolution, *Quarterly Journal of the Royal Meteorological Society*, 138, 72–80, <https://doi.org/10.1002/qj.918>, 2012.
- Ćurić, M., Janc, D., and Vučković, V.: Numerical simulation of Cb cloud vorticity, *Atmospheric Research*, 83, 427–434, <https://doi.org/10.1016/j.atmosres.2005.10.024>, 2007.
- Dabernig, M., Mayr, G. J., Messner, J. W., and Zeileis, A.: Spatial ensemble post-processing with standardized anomalies, *Quarterly Journal of the Royal Meteorological Society*, 143, 909–916, <https://doi.org/10.1002/qj.2975>, 2017.
- Dawson, D. T., Mansell, E. R., Jung, Y., Wicker, L. J., Kumjian, M. R., and Xue, M.: Low-level ZDR signatures in supercell forward flanks: The role of size sorting and melting of hail, *Journal of the Atmospheric Sciences*, 71, 276–299, <https://doi.org/10.1175/JAS-D-13-0118.1>, 2014.
- De Martin, F., Davolio, S., Miglietta, M. M., and Levizzani, V.: A Conceptual Model for the Development of Tornadoes in the Complex Orography of the Po Valley, *Monthly Weather Review*, 152, 1357–1377, <https://doi.org/10.1175/MWR-D-23-0222.1>, 2024.
- De Martin, F., Manzato, A., Carlon, N., Setvák, M., and Miglietta, M. M.: A dynamic and statistical analysis of giant hail environments in North-East Italy, *QJRMS*, 2025.
- Delobbe, L. and Holleman, I.: Uncertainties in radar echo top heights used for hail detection, *Meteorological Applications*, 13, 361–374, <https://doi.org/10.1017/S1350482706002374>, 2006.
- Demortier, A., Mandement, M., Pourret, V., and Caumont, O.: Assimilation of surface pressure observations from personal weather stations in AROME-France, *Natural Hazards and Earth System Sciences*, pp. 907–927, <https://doi.org/https://doi.org/10.5194/nhess-24-907-2024>, 2024.
- Doswell, C. A.: The Distinction between Large-Scale and Mesoscale Contribution to Severe Convection: A Case Study Example, *Weather and Forecasting*, 2, 3–16, [https://doi.org/10.1175/1520-0434\(1987\)002<0003:TDBLSA>2.0.CO;2](https://doi.org/10.1175/1520-0434(1987)002<0003:TDBLSA>2.0.CO;2), 1987.
- Doswell, C. A., Brooks, H. E., and Maddox, R. A.: Flash Flood Forecasting : An Ingredients-Based Methodology, *WAF*, 11, 560–581, 1996.
- Dotzek, N.: Tornadoes in Germany, *Atmospheric Research*, 56, 233–251, [https://doi.org/10.1016/S0169-8095\(00\)00075-2](https://doi.org/10.1016/S0169-8095(00)00075-2), 2001.
- Dotzek, N., Groenemeijer, P., Feuerstein, B., and Holzer, A. M.: Overview of ESSL’s severe convective storms research using the European Severe Weather Database ESWD, *Atmospheric Research*, 93, 575–586, <https://doi.org/10.1016/j.atmosres.2008.10.020>, 2009.
- Ducrocq, V., Davolio, S., Ferretti, R., Flamant, C., Santaner, V. H., Kalthoff, N., Richard, E., and Wernli, H.: Introduction to the HyMeX Special Issue on ‘Advances in understanding and forecasting of heavy precipitation in the Mediterranean through the HyMeX SOP1 field campaign’, *Quarterly Journal of the Royal Meteorological Society*, 142, 1–6, <https://doi.org/10.1002/qj.2856>, 2016.
- Durran, D. R. and Weyn, J. A.: Thunderstorms do not get butterflies, *Bulletin of the American Meteorological Society*, 97, 237–243, <https://doi.org/10.1175/BAMS-D-15-00070.1>, 2016.
- ECMWF: <https://digital-strategy.ec.europa.eu/en/policies/destination-earth>, 2023.
- Eirund, G. K., Drossaert van Dusseldorp, S., Brem, B. T., Dedekind, Z., Karrer, Y., Stoll, M., and Lohmann, U.: Aerosol–cloud–precipitation interactions during a Saharan dust event – A summertime case-study from the Alps, *Quarterly Journal of the Royal Meteorological Society*, 148, 943–961, <https://doi.org/10.1002/qj.4240>, 2022.

- 785 Emeis, S., Kalthoff, N., Adler, B., Pardyjak, E., Paci, A., and Junkermann, W.: High-resolution observations of transport and exchange processes in mountainous terrain, *Atmosphere*, 9, <https://doi.org/10.3390/atmos9120457>, 2018.
- Erdmann, F. and Poelman, D. R.: Thunderstorm characteristics with lightning jumps and dives in satellite-based nowcasting, pp. 1–26, 2024.
- ESSL: International Fujita (IF) Scale, <https://www.essl.org/cms/research-projects/international-fujita-scale/>, 2023a.
- ESSL: Severe Weather Surveys and Survey Guide, <http://essl.org/cms/publications/detailed-survey-reports/>, 2023b.
- 790 ESSL: ESSL news: Hail record broken again – 19cm hailstone confirmed in Italy, <https://www.essl.org/cms/hail-record-broken-again-19cm-hailstone-confirmed-in-italy/>, 2023c.
- Evaristo, R., Reinoso-Rondinel, R., Tromel, S., and Simmer, C.: Validation of Wind Fields Retrieved by Dual-Doppler Techniques Using a Vertically Pointing Radar, in: 2021 21st International Radar Symposium (IRS), pp. 1–7, <https://doi.org/10.23919/IRS51887.2021.9466173>, 2021.
- 795 Feldmann, M., Germann, U., Gabella, M., and Berne, A.: A characterisation of Alpine mesocyclone occurrence, *Weather and Climate Dynamics*, 2, 1225–1244, <https://doi.org/10.5194/wcd-2-1225-2021>, 2021.
- Feldmann, M., Hering, A., Gabella, M., and Berne, A.: Hailstorms and rainstorms versus supercells—a regional analysis of convective storm types in the Alpine region, *npj Climate and Atmospheric Science*, 6, 1–11, <https://doi.org/10.1038/s41612-023-00352-z>, 2023.
- Feldmann, M., Rotunno, R., Germann, U., and Berne, A.: Supercell Thunderstorms in Complex Topography—How Mountain Valleys with
800 Lakes Can Increase Occurrence Frequency, *Monthly Weather Review*, 152, 471–489, <https://doi.org/10.1175/MWR-D-22-0350.1>, 2024.
- Feldmann, M., Blanc, M., Brennan, K. P., Thurnherr, I., Velasquez, P., Martius, O., and Schär, C.: European supercell thunderstorms – an underestimated current threat and an increasing future hazard, <https://arxiv.org/abs/2503.07466>, 2025.
- Fencel, M., Dohnal, M., and Bareš, V.: Retrieving Water Vapor From an E-Band Microwave Link With an Empirical Model Not Requiring In Situ Calibration, *Earth and Space Science*, 8, 1–9, <https://doi.org/10.1029/2021EA001911>, 2021.
- 805 Figueras i Ventura, J., Pineda, N., Besic, N., Grazioli, J., Hering, A., Van Der Velde, O. A., Romero, D., Sunjerga, A., Mostajabi, A., Azadifar, M., Rubinstein, M., Montanyà, J., Germann, U., and Rachidi, F.: Analysis of the lightning production of convective cells, *Atmospheric Measurement Techniques*, 12, 5573–5591, <https://doi.org/10.5194/amt-12-5573-2019>, 2019.
- Fluck, E., Kunz, M., Geissbuehler, P., and Ritz, S. P.: Radar-based assessment of hail frequency in Europe, *Natural Hazards and Earth System Sciences*, pp. 683–701, 2021.
- 810 Forcadell, V., Augros, C., Caumont, O., Dedieu, K., Ouradou, M., David, C., i Ventura, F., Laurantin, O., and Al-Sakka, H.: Severe hail detection with C-band dual-polarisation radars using convolutional neural networks, *EGUsphere*, 2024, 1–43, <https://doi.org/10.5194/egusphere-2024-1336>, 2024.
- Fourrié, N., Nuret, M., Brousseau, P., and Caumont, O.: Data assimilation impact studies with the AROME-WMED reanalysis of the first special observation period of the Hydrological cycle in the Mediterranean Experiment, *Natural Hazards and Earth System Sciences*, 21,
815 463–480, <https://doi.org/10.5194/nhess-21-463-2021>, 2021.
- Frame, J. W. and Markowski, P.: The interaction of simulated squall lines with idealized mountain ridges, *Monthly Weather Review*, 134, 1919–1941, <https://doi.org/10.1175/MWR3157.1>, 2006.
- French, M. M. and Kingfield, D. M.: Tornado Formation and Intensity Prediction Using Polarimetric Radar Estimates of Updraft Area, *Weather and Forecasting*, 36, 2211–2231, <https://doi.org/10.1175/WAF-D-21-0087.1>, 2021.
- 820 Fuchs, B. R., Rutledge, S. A., Bruning, E. C., Pierce, J. R., Kodros, J. K., Lang, T. J., MacGorman, D. R., Krehbiel, P. R., and Rison, W.: Environmental controls on storm intensity and charge structure in multiple regions of the continental United States, *Journal of Geophysical Research*, 120, 6575–6596, <https://doi.org/10.1002/2015JD023271>, 2015.

- Fujita, T.: Mesoanalysis of the Illinois Tornadoes of April 1953, *Journal of Atmospheric Sciences*, 15, 288–296, [https://doi.org/10.1175/1520-0469\(1958\)015<0288:MOTITO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1958)015<0288:MOTITO>2.0.CO;2), 1958.
- 825 Fujita, T.: Downbursts: meteorological features and wind field characteristics, *Journal of Wind Engineering and Industrial Aerodynamics*, 36, 75–86, [https://doi.org/https://doi.org/10.1016/0167-6105\(90\)90294-M](https://doi.org/https://doi.org/10.1016/0167-6105(90)90294-M), 1990.
- Fujita, T. T.: The Teton-Yellowstone Tornado of 21 July 1987, *Monthly Weather Review*, 117, 1913–1940, [https://doi.org/10.1175/1520-0493\(1989\)117<1913:TTYTOJ>2.0.CO;2](https://doi.org/10.1175/1520-0493(1989)117<1913:TTYTOJ>2.0.CO;2), 1989.
- Gallo, K., Schumacher, P., Boustead, J., and Ferguson, A.: Validation of satellite observations of storm damage to cropland with digital
830 photographs, *Weather and Forecasting*, 34, 435–446, <https://doi.org/10.1175/WAF-D-18-0059.1>, 2019.
- Gatzen, C. P., Fink, A. H., Schultz, D. M., and Pinto, J. G.: An 18-year climatology of derechos in Germany, *Natural Hazards and Earth System Sciences*, 20, 1335–1351, <https://doi.org/10.5194/nhess-20-1335-2020>, 2020.
- Geerts, B., Andretta, T., Luberd, S. J., Vogt, J., Wang, Y., Oolman, L. D., Bikos, D., and Finch, J.: Case Study of a Long-lived Tornadoic Mesocyclone in a Low-CAPE Complex-terrain Environment, *E-Journal of Severe Storms Meteorology*, 4, 1–29,
835 <https://doi.org/10.55599/ejssm.v4i3.20>, 2009.
- Geerts, B., Parsons, D., Ziegler, C. L., Weckwerth, T. M., Biggstaff, M. I., Clark, R. D., Conigliolo, M. C., Demoz, B. B., Ferrare, R. A., Gallus, W. A., Haghi, K., Hanesiak, J. M., Klein, P. M., Knupp, K. R., Kosiba, K., McFarquhar, G. M., Moore, J. A., Nehrir, A. R., Parker, M. D., Pinto, J. O., Rauber, R. M., Schumacher, R. S., Turner, D. D., Wang, Q., Wang, X., Wang, Z., and Wurman, J.: The 2015 plains elevated convection at night field project, *Bulletin of the American Meteorological Society*, 98, 767–786, <https://doi.org/10.1175/BAMS-D-15-00257.1>, 2017.
840
- Gensini, V. A., Ashley, W. S., Michaelis, A. C., Haberlie, A. M., Goodin, J., and Wallace, B. C.: Hailstone size dichotomy in a warming climate, *npj Climate and Atmospheric Science*, 7, 1–10, <https://doi.org/10.1038/s41612-024-00728-9>, 2024.
- Germann, U., Boscacci, M., Clementi, L., Gabella, M., Hering, A., Sartori, M., Sideris, I. V., and Calpini, B.: Weather Radar in Complex Orography, *Remote Sensing*, 14, 503, <https://doi.org/10.3390/rs14030503>, 2022.
- 845 Ghasemifard, H., Groenemeijer, P., Battaglioli, F., and Púčik, T.: Do changing circulation types raise the frequency of summertime thunderstorms and large hail in Europe?, *Environmental Research: Climate*, 3, 015 008, <https://doi.org/10.1088/2752-5295/ad22ec>, 2024.
- Giordani, A., Kunz, M., Bedka, K. M., Punge, H. J., Pavan, V., Cerenzia, I. M. L., and Sabatino, S. D.: Characterizing hail-prone environments using convection-permitting reanalysis and overshooting top detections over south-central Europe, 2023.
- Göbel, M., Serafin, S., and Rotach, M. W.: Adverse impact of terrain steepness on thermally driven initiation of orographic convection,
850 *Weather and Climate Dynamics*, 4, 725–745, <https://doi.org/10.5194/wcd-4-725-2023>, 2023.
- Goger, B., Rotach, M. W., Gohm, A., Stiperski, I., Fuhrer, O., and De Morsier, G.: A new horizontal length scale for a three-dimensional turbulence parameterization in mesoscale atmospheric modeling over highly complex terrain, *Journal of Applied Meteorology and Climatology*, 58, 2087–2102, <https://doi.org/10.1175/JAMC-D-18-0328.1>, 2019.
- Gopalakrishnan, D., Cuervo-Lopez, C., Allen, J. T., Trapp, R. J., and Robinson, E.: A Comprehensive Evaluation of Biases in Convective
855 Storm Parameters in CMIP6 Models over North America, *Journal of Climate*, 38, 947–971, <https://doi.org/10.1175/JCLI-D-24-0165.1>, 2025.
- Greene, D. R. and Clark, R. A.: Vertically Integrated Liquid Water—A New Analysis Tool, *Monthly Weather Review*, 100, 548–552, [https://doi.org/10.1175/1520-0493\(1972\)100<0548:VILWNA>2.3.CO;2](https://doi.org/10.1175/1520-0493(1972)100<0548:VILWNA>2.3.CO;2), 1972.
- Groenemeijer, P.: Evaluating IASI Profiles at the ESSL Test-bed and for Selected Cases, <https://www.eumetrain.org/index.php/resources/evaluating-iasi-profiles-essl-test-bed-and-selected-cases>, 2019.
860

- Guay, B., Fengler, M., and Hammerschmidt, L.: Improving Nowcasts and Forecasts via Operational Use of Meteomatics Me-teodrones, in: EGU General Assembly Conference Abstracts, EGU General Assembly Conference Abstracts, pp. EGU-8632, <https://doi.org/10.5194/egusphere-egu23-8632>, 2023.
- Gustafsson, N., Janjić, T., Schraff, C., Leuenberger, D., Weissmann, M., Reich, H., Brousseau, P., Montmerle, T., Wattrelot, E., Bučánek, A., Mile, M., Hamdi, R., Lindskog, M., Barkmeijer, J., Dahlbom, M., Macpherson, B., Ballard, S., Inverarity, G., Carley, J., Alexander, C., Dowell, D., Liu, S., Ikuta, Y., and Fujita, T.: Survey of data assimilation methods for convective-scale numerical weather prediction at operational centres, *Quarterly Journal of the Royal Meteorological Society*, 144, 1218–1256, <https://doi.org/10.1002/qj.3179>, 2018.
- Hannesen, R., Dotzek, N., Gysi, H., and Beheng, K. D.: Case study of a tornado in the Upper Rhine valley, *Meteorologische Zeitschrift*, 7, 163–170, <https://doi.org/10.1127/metz/7/1998/163>, 1998.
- Hervo, M., Romanens, G., Martucci, G., Weusthoff, T., and Haefele, A.: Evaluation of an Automatic Meteorological Drone Based on a 6-Month Measurement Campaign, *Atmosphere*, 14, <https://doi.org/10.3390/atmos14091382>, 2023.
- Heymsfield, A. J., Cecchini, M. A., Detwiler, A., Honeyager, R., and Field, P.: Exploring the Composited T-28 Hailstorm Penetration Dataset to Characterize Hail Properties within the Updraft and Downdraft Regions, *Journal of Applied Meteorology and Climatology*, 62, 1803–1826, <https://doi.org/10.1175/JAMC-D-23-0030.1>, 2023.
- Holmlund, K., Grandell, J., Schmetz, J., Stuhlmann, R., Bojkov, B., Munro, R., Lekouara, M., Coppens, D., Viticchie, B., August, T., Theodore, B., Watts, P., Dobber, M., Fowler, G., Bojinski, S., Schmid, A., Salonen, K., Tjemkes, S., Aminou, D., and Blythe, P.: Meteosat Third Generation (MTG): Continuation and Innovation of Observations from Geostationary Orbit, *Bulletin of the American Meteorological Society*, 102, E990 – E1015, <https://doi.org/10.1175/BAMS-D-19-0304.1>, 2021.
- Homeyer, C. R., McAuliffe, J. D., and Bedka, K. M.: On the development of above-anvil cirrus plumes in extratropical convection, *Journal of the Atmospheric Sciences*, 74, 1617–1633, <https://doi.org/10.1175/JAS-D-16-0269.1>, 2017.
- Houston, A. L., Argrow, B., Coniglio, M., Frew, E. W., Rasmussen, E. N., Weiss, C. C., and Ziegler, C. L.: Targeted Observation by Radars and UAS of Supercells (TORUS): Summary of the 2019 Field Campaign, in: 110th AMS Annual Meeting, Boston MA, USA, 2020.
- Houze, R. A.: Orographic effects on precipitating clouds, *Reviews of Geophysics*, 50, 1–47, <https://doi.org/10.1029/2011RG000365>, 2012.
- Houze, R. A., Schmid, W., Fovell, R. G., and Schiesser, H.-H.: Hailstorms in Switzerland: Left Movers, Right Movers, and False Hooks, *Monthly Weather Review*, 121, 3345–3370, [https://doi.org/10.1175/1520-0493\(1993\)121<3345:HISLMR>2.0.CO;2](https://doi.org/10.1175/1520-0493(1993)121<3345:HISLMR>2.0.CO;2), 1993.
- Imamovic, A., Schlemmer, L., and Schär, C.: Mountain volume control on deep-convective rain amount during episodes of weak synoptic forcing, *Journal of the Atmospheric Sciences*, 76, 605–626, <https://doi.org/10.1175/JAS-D-18-0217.1>, 2019.
- Jensen, M. P., Flynn, J. H., Judd, L. M., Kollias, P., Kuang, C., McFarquhar, G., Nadkarni, R., Powers, H., and Sullivan, J.: A Succession of Cloud, Precipitation, Aerosol, and Air Quality Field Experiments in the Coastal Urban Environment, *Bulletin of the American Meteorological Society*, 102, 103–105, <https://doi.org/10.1175/BAMS-D-21-0104.1>, 2022.
- Kahraman, A., Kendon, E. J., Chan, S. C., and Fowler, H. J.: Quasi-Stationary Intense Rainstorms Spread Across Europe Under Climate Change, *Geophysical Research Letters*, 48, 1–11, <https://doi.org/10.1029/2020GL092361>, 2021.
- Kahraman, A., Kendon, E. J., and Fowler, H. J.: Climatology of severe hail potential in Europe based on a convection-permitting simulation, *Climate Dynamics*, <https://doi.org/10.1007/s00382-024-07227-w>, 2024.
- Kaltenboeck, R. and Steinheimer, M.: Radar-based severe storm climatology for Austrian complex orography related to vertical wind shear and atmospheric instability, *Atmospheric Research*, 158–159, 216–230, <https://doi.org/10.1016/j.atmosres.2014.08.006>, 2015.
- Katona, B. and Markowski, P.: Assessing the influence of complex terrain on severe convective environments in Northeastern Alabama, *Weather and Forecasting*, 36, 1003–1029, <https://doi.org/10.1175/WAF-D-20-0136.1>, 2021.

- Katona, B., Markowski, P., Alexander, C., and Benjamin, S.: The influence of topography on convective storm environments in the eastern United States as deduced from the HRRR, *Weather and Forecasting*, 31, 1481–1490, <https://doi.org/10.1175/WAF-D-16-0038.1>, 2016.
- Kaňák, J., Benko, M., Simon, A., and Sokol, A.: Case study of the 9 May 2003 windstorm in southwestern Slovakia, *Atmospheric Research*, 83, 162–175, <https://doi.org/10.1016/j.atmosres.2005.09.012>, 2007.
- Khodayar, S., Davolio, S., Di Girolamo, P., Lebeaupin Brossier, C., Flaounas, E., Fourrie, N., Lee, K. O., Ricard, D., Vie, B., Bontier, F., Caldas-Alvarez, A., and Ducrocq, V.: Overview towards improved understanding of the mechanisms leading to heavy precipitation in the western Mediterranean: Lessons learned from HyMeX, *Atmospheric Chemistry and Physics*, 21, 17 051–17 078, <https://doi.org/10.5194/acp-21-17051-2021>, 2021.
- Kirshbaum, D. J.: Numerical simulations of orographic convection across multiple gray zones, *Journal of the Atmospheric Sciences*, 77, 3301–3320, <https://doi.org/10.1175/JAS-D-20-0035.1>, 2020.
- Kirshbaum, D. J., Adler, B., Kalthoff, N., Barthlott, C., and Serafin, S.: Moist orographic convection: Physical mechanisms and links to surface-exchange processes, *Atmosphere*, 9, 1–26, <https://doi.org/10.3390/atmos9030080>, 2018.
- Klaus, V. and Krause, J.: Investigating Hailstorm Updrafts and Nowcasting Hail Size Using a Novel Radar-Based Updraft Detection, *Weather Forecasting*, 39, 1795–1815, <https://doi.org/10.1175/WAF-D-23-0227.1>, 2024.
- Klaus, V., Rieder, H., and Kaltenböck, R.: Insights in Hailstorm Dynamics through Polarimetric High-Resolution X-Band and Operational C-Band Radar: A Case Study for Vienna, Austria, *Monthly Weather Review*, 151, 913–929, <https://doi.org/10.1175/MWR-D-22-0185.1>, 2023.
- Knight, C. A., Foote, G. B., and Summers, P. W.: Results of a Randomized Hail Suppression Experiment in Northeast Colorado. Part IX: Overall Discussion and Summary in the Context of Physical Research, *Journal of Applied Meteorology and Climatology*, 18, 1629–1639, [https://doi.org/10.1175/1520-0450\(1979\)018<1629:ROARHS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1979)018<1629:ROARHS>2.0.CO;2), 1979.
- Kollias, P., McFarquhar, G. M., Bruning, E., DeMott, P. J., Kumjian, M. R., Lawson, P., Lebo, Z., Logan, T., Lombardo, K., Oue, M., Roberts, G., Shaw, R. A., van den Heever, S. C., Wolde, M., Barry, K. R., Bodine, D., Brientjes, R., Chandrasekar, V., Dzambo, A., Hill, T. C. J., Jensen, M., Junyent, F., Kreidenweis, S. M., Lamer, K., Luke, E., Bansemer, A., McCluskey, C., Nichman, L., Nguyen, C., Patnaude, R. J., Perkins, R. J., Powers, H., Ranjbar, K., Roux, E., Snyder, J., Treserras, B. P., Tsai, P., Wales, N. A., Wolff, C., Allwayin, N., Ascher, B., Barr, J., Hu, Y., Huang, Y., Litzmann, M., Mages, Z., McKeown, K., Patil, S., Rosky, E., Tuftedal, K., Tzeng, M.-D., and Zhu, Z.: Experiment of Sea Breeze Convection, Aerosols, Precipitation and Environment (ESCAPE), *Bulletin of the American Meteorological Society*, pp. 310–332, <https://doi.org/10.1175/bams-d-23-0014.1>, 2024.
- Komjáti, K., Csirmaz, K., Breuer, H., Kurcsics, M., and Horváth, Á.: Supercell interactions with surface baroclinic zones in the Carpathian Basin, in: 11th European Conference on Severe Storms, pp. ECSS2023–39, Bucharest, Romania, 2023.
- Kopp, J., Schröer, K., Schwierz, C., Hering, A., Germann, U., and Martius, O.: The summer 2021 Switzerland hailstorms: weather situation, major impacts and unique observational data, *Weather*, 78, 184–191, <https://doi.org/10.1002/wea.4306>, 2023.
- Kosiba, K. A., Lyza, A. W., Trapp, R. J., Rasmussen, E. N., Parker, M., Biggerstaff, M. I., Nesbitt, S. W., Weiss, C. C., Wurman, J., Knupp, K. R., Coffey, B., Chmielewski, V. C., Dawson, D. T., Bruning, E., Bell, T. M., Coniglio, M. C., Murphy, T. A., French, M., Blind-Doskocil, L., Reinhart, A. E., Edward Wolff, Schneider, M. E., Silcott, M., Smith, E., Joshua Aikins, Wagner, M., Robinson, P., Wilczak, J. M., White, T., Bodine, D., Kumjian, M. R., Waugh, S. M., Alford, A. A., Elmore, K., Kollias, P., and Turner, D. D.: The Propagation, Evolution, and Rotation in Linear Storms (PERiLS) Project, *Bulletin of the American Meteorological Society*, <https://doi.org/10.1175/BAMS-D-22-0064.1>, 2024.

- Kramer, M., Heinzeller, D., Hartmann, H., van den Berg, W., and Steeneveld, G. J.: Assessment of MPAS variable resolution simulations in the grey-zone of convection against WRF model results and observations: An MPAS feasibility study of three extreme weather events in Europe, *Climate Dynamics*, 55, 253–276, <https://doi.org/10.1007/s00382-018-4562-z>, 2020.
- Kumjian, M.: Principles and applications of dual-polarization weather radar. Part III: Artifacts, *Journal of Operational Meteorology*, 1, 265–
940 274, <https://doi.org/10.15191/nwajom.2013.0121>, 2013.
- Kumjian, M. R. and Ryzhkov, A. V.: Polarimetric signatures in supercell thunderstorms, *Journal of Applied Meteorology and Climatology*, 47, 1940–1961, <https://doi.org/10.1175/2007JAMC1874.1>, 2008.
- Kumjian, M. R. and Ryzhkov, A. V.: Storm-relative helicity revealed from polarimetric radar measurements, *Journal of the Atmospheric Sciences*, 66, 667–685, <https://doi.org/10.1175/2008JAS2815.1>, 2009.
- 945 Kunkel, J., Hanesiak, J., and Sills, D.: The Hunt for Missing Tornadoes: Using Satellite Imagery to Detect and Document Historical Tornado Damage in Canadian Forests, *Journal of Applied Meteorology and Climatology*, 62, 139–154, <https://doi.org/10.1175/JAMC-D-22-0070.1>, 2023.
- Kunz, M. and Pusskeiler, M.: High-resolution assessment of the hail hazard over complex terrain from radar and insurance data, *Open Access Article*, 19, 427–439, <https://doi.org/10.1127/0941-2948/2010/0452>, 2010.
- 950 Kunz, M., Blahak, U., Handwerker, J., Schmidberger, M., Punge, H. J., Mohr, S., Fluck, E., and Bedka, K. M.: The severe hailstorm in south-west Germany on 28 July 2013: characteristics, impacts and meteorological conditions, *Quarterly Journal of the Royal Meteorological Society*, 144, 231–250, <https://doi.org/10.1002/qj.3197>, 2018.
- Kunz, M., Abbas, S. S., Bauckholt, M., Böhmländer, A., Feuerle, T., Gasch, P., Glaser, C., Groß, J., Hajnsek, I., Handwerker, J., Hase, F., Khordakova, D., Knippertz, P., Kohler, M., Lange, D., Latt, M., Laube, J., Martin, L., Mauder, M., Möhler, O., Mohr, S., Reitter, R. W.,
955 Rettenmeier, A., Rolf, C., Saathoff, H., Schrön, M., Schütze, C., Spahr, S., Späth, F., Vogel, F., Völksch, I., Weber, U., Wieser, A., Wilhelm, J., Zhang, H., and Dietrich, P.: Swabian MOSES 2021: An interdisciplinary field campaign for investigating convective storms and their event chains, *Frontiers in Earth Science*, 10, 1–26, <https://doi.org/10.3389/feart.2022.999593>, 2022.
- Kunz, M., Hühn, E., Fischer, J., Mohr, S., Latt, M., Trömel, S., and Soderholm, J.: Enhancing insights into large hail formation and trajectories through targeted field campaigns, 2024.
- 960 Kvak, R., Okon, L., Bližňák, V., Méri, L., and Kašpar, M.: Spatial distribution and precipitation intensity of supercells: Response to terrain asymmetry in the Western Carpathians, Central Europe, *Atmospheric Research*, 292, <https://doi.org/10.1016/j.atmosres.2023.106885>, 2023.
- Labriola, J., Snook, N., Jung, Y., and Xue, M.: Explicit ensemble prediction of hail in 19 May 2013 Oklahoma city thunderstorms and analysis of hail growth processes with several multimoment microphysics schemes, *Monthly Weather Review*, 147, 1193–1213,
965 <https://doi.org/10.1175/MWR-D-18-0266.1>, 2019.
- Lainer, M., Brennan, K. P., Hering, A., Kopp, J., Monhart, S., Wolfensberger, D., and Germann, U.: Drone-based photogrammetry combined with deep-learning to estimate hail size distributions and melting of hail on the ground, *Atmospheric Measurement Techniques Discussions*, 2024, 2539–2557, <https://doi.org/https://doi.org/10.5194/amt-17-2539-2024>, 2024.
- Laiti, L., Zardi, D., De Franceschi, M., Rampanelli, G., and Giovannini, L.: Analysis of the diurnal development of a lake-valley circulation in
970 the Alps based on airborne and surface measurements, *Atmospheric Chemistry and Physics*, 14, 9771–9786, <https://doi.org/10.5194/acp-14-9771-2014>, 2014.

- Lakshmi, K. K. A., Sahoo, S., Biswas, S. K., and Chandrasekar, V.: Study of Microphysical Signatures Based on Spectral Polarimetry during the RELAMPAGO Field Experiment in Argentina, *Journal of Atmospheric and Oceanic Technology*, 41, 235–260, <https://doi.org/10.1175/JTECH-D-22-0113.1>, 2024.
- 975 Lasher-Trapp, S., Orendorf, S. A., and Trapp, R. J.: Investigating a Derecho in a Future Warmer Climate, *Bulletin of the American Meteorological Society*, 104, E1831–E1852, <https://doi.org/10.1175/BAMS-D-22-0173.1>, 2023.
- LeBel, L. J., Tang, B. H., and Lazear, R. A.: Examining Terrain Effects on an Upstate New York Tornado Event Utilizing a High-Resolution Model Simulation, *Weather and Forecasting*, 36, 2001–2020, <https://doi.org/10.1175/WAF-D-21-0018.1>, 2021.
- Leinonen, J., Hamann, U., Sideris, I. V., and Germann, U.: Thunderstorm Nowcasting With Deep Learning: A Multi-Hazard Data Fusion
980 Model, *Geophysical Research Letters*, 50, <https://doi.org/10.1029/2022GL101626>, 2023.
- Lenderink, G., Ban, N., Brisson, E., Berthou, S., Cortés-Hernández, V. E., Kendon, E., Fowler, H., and de Vries, H.: Are dependencies of extreme rainfall on humidity more reliable in convection-permitting climate models?, *Hydrology and Earth System Sciences Discussions*, 2024, 1–31, <https://doi.org/10.5194/hess-2024-132>, 2024.
- Letkewicz, C. E. and Parker, M. D.: Impact of environmental variations on simulated squall lines interacting with terrain, *Monthly Weather
985 Review*, 139, 3163–3183, <https://doi.org/10.1175/2011MWR3635.1>, 2011.
- Lin, Y. and Kumjian, M. R.: Influences of CAPE on Hail Production in Simulated Supercell Storms, *Journal of the Atmospheric Sciences*, 79, 179–204, <https://doi.org/10.1175/JAS-D-21-0054.1>, 2022.
- Liu, J. and Niyogi, D.: Meta-analysis of urbanization impact on rainfall modification, *Scientific Reports*, 9, 1–14, <https://doi.org/10.1038/s41598-019-42494-2>, 2019.
- 990 Loeffler, S. D., Kumjian, M. R., Jurewicz, M., and French, M. M.: Differentiating Between Tornadoic and Nontornadoic Supercells Using Polarimetric Radar Signatures of Hydrometeor Size Sorting, *Geophysical Research Letters*, 47, 1–9, <https://doi.org/10.1029/2020GL088242>, 2020.
- Loftus, A. M. and Cotton, W. R.: Examination of CCN impacts on hail in a simulated supercell storm with triple-moment hail bulk microphysics, *Atmospheric Research*, 147–148, 183–204, <https://doi.org/10.1016/j.atmosres.2014.04.017>, 2014.
- 995 Lombardo, K. and Bitting, M.: A Climatology of Convective Precipitation over Europe, *Monthly Weather Review*, <https://doi.org/10.1175/MWR-D-23-0156.1>, 2024.
- Lombardo, K. and Kumjian, M. R.: Observations of the Discrete Propagation of a Mesoscale Convective System during RELAMPAGO–CACTI, *Monthly Weather Review*, 150, 2111–2138, <https://doi.org/10.1175/MWR-D-21-0265.1>, 2022.
- Maddox, R. A., Hoxit, and Chappell: A Study of Tornadoic Thunderstorm Interactions with Thermal Boundaries, *Mon. Wea. Rev.*, 1980.
- 1000 Malečić, B., Telišman Prtenjak, M., Horvath, K., Jelić, D., Mikuš Jurković, P., Ćorko, K., and Mahović, N. S.: Performance of HAILCAST and the Lightning Potential Index in simulating hailstorms in Croatia in a mesoscale model – Sensitivity to the PBL and microphysics parameterization schemes, *Atmospheric Research*, 272, <https://doi.org/10.1016/j.atmosres.2022.106143>, 2022.
- Mandement, M. and Caumont, O.: A numerical study to investigate the roles of former Hurricane Leslie, orography and evaporative cooling in the 2018 Aude heavy-precipitation event, *Weather and Climate Dynamics*, 2, 795–818, <https://doi.org/10.5194/wcd-2-795-2021>, 2021.
- 1005 Manzato, A., Cicogna, A., Centore, M., Battistutta, P., and Trevisan, M.: Hailstone Characteristics in Northeast Italy from 29 Years of Hailpad Data, *Journal of Applied Meteorology and Climatology*, 61, 1773–1789, <https://doi.org/10.1175/JAMC-D-21-0251.1>, 2022a.
- Manzato, A., Serafin, S., Miglietta, M. M., Kirshbaum, D., and Schulz, W.: A Pan-Alpine Climatology of Lightning and Convective Initiation, *Monthly Weather Review*, 150, 2213–2230, <https://doi.org/10.1175/MWR-D-21-0149.1>, 2022b.
- Markowski, P. and Richardson, Y.: Mesoscale meteorology in midlatitudes, vol. 2, John Wiley and Sons, 2010.

- 1010 Markowski, P. M. and Dotzek, N.: A numerical study of the effects of orography on supercells, *Atmospheric Research*, 100, 457–478, <https://doi.org/10.1016/j.atmosres.2010.12.027>, 2011.
- Markowski, P. M., Richardson, Y. P., Richardson, S. J., and Petersson, A.: Aboveground thermodynamic observations in convective storms from balloonborne probes acting as pseudo-lagrangian drifters, *Bulletin of the American Meteorological Society*, 99, 711–724, <https://doi.org/10.1175/BAMS-D-17-0204.1>, 2018.
- 1015 Marquis, J. N., Varble, A. C., Robinson, P., Nelson, T. C., and Friedrich, K.: Low-Level Mesoscale and Cloud-Scale Interactions Promoting Deep Convection Initiation, *Monthly Weather Review*, 149, 2473–2495, <https://doi.org/10.1175/MWR-D-20-0391.1>, 2021.
- Marquis, j. N., Feng, Z., Varble, A., Nelson, T. C., Houston, A., Peters, J. M., Mulholland, J. P., and Hardin, J.: Near-Cloud Atmospheric Ingredients for Deep Convection Initiation, *Monthly Weather Review*, 151, 1247–1267, <https://doi.org/10.1175/MWR-D-22-0243.1>, 2023.
- Martín, M. L., Calvo-Sancho, C., Taszarek, M., González-Alemán, J. J., Montoro-Mendoza, A., Díaz-Fernández, J., Bolgiani, P., Sastre, M., and Martín, Y.: Major Role of Marine Heatwave and Anthropogenic Climate Change on a Giant Hail Event in Spain, *Geophysical Research Letters*, 51, <https://doi.org/10.1029/2023GL107632>, 2024.
- 1020 Marvi, M. T.: A review of flood damage analysis for a building structure and contents, *Natural Hazards*, 102, 967–995, <https://doi.org/10.1007/s11069-020-03941-w>, 2020.
- McGinley, J.: A Diagnosis of Alpine Lee Cyclogenesis, *Monthly Weather Review*, 110, 1271–1287, [https://doi.org/10.1175/1520-0493\(1982\)110<1271:ADOALC>2.0.CO;2](https://doi.org/10.1175/1520-0493(1982)110<1271:ADOALC>2.0.CO;2), 1982.
- 1025 McKeown, K. E., Davenport, C. E., Eastin, M. D., Purpura, S. M., and Riggan, R. R.: Radar Characteristics of Supercell Thunderstorms Traversing the Appalachian Mountains, *Weather and Forecasting*, <https://doi.org/10.1175/waf-d-23-0110.1>, 2024.
- Medina, B. L., Carey, L. D., Lang, T. J., Bitzer, P. M., Deierling, W., and Zhu, Y.: Characterizing Charge Structure in Central Argentina Thunderstorms During RELAMPAGO Utilizing a New Charge Layer Polarity Identification Method, *Earth and Space Science*, 8, <https://doi.org/10.1029/2021EA001803>, 2021.
- 1030 Merz, R., Tarasova, L., and Basso, S.: The flood cooking book: Ingredients and regional flavors of floods across Germany, *Environmental Research Letters*, 15, <https://doi.org/10.1088/1748-9326/abb9dd>, 2020.
- Miglietta, M. M. and Rotunno, R.: Numerical simulations of sheared conditionally unstable flows over a mountain ridge, *Journal of the Atmospheric Sciences*, 71, 1747–1762, <https://doi.org/10.1175/JAS-D-13-0297.1>, 2014.
- 1035 Miller, P. W. and Mote, T. L.: Standardizing the definition of a "pulse" thunderstorm, *Bulletin of the American Meteorological Society*, 98, 905–913, <https://doi.org/10.1175/BAMS-D-16-0064.1>, 2017.
- Miyoshi, T., Kunii, M., Ruiz, J., Lien, G. Y., Satoh, S., Ushio, T., Bessho, K., Seko, H., Tomita, H., and Ishikawa, Y.: Big data assimilation revolutionizing severe weather prediction, *Bulletin of the American Meteorological Society*, 97, 1347–1354, <https://doi.org/10.1175/BAMS-D-15-00144.1>, 2016.
- 1040 Mohr, S. and Kunz, M.: Recent trends and variabilities of convective parameters relevant for hail events in Germany and Europe, *Atmospheric Research*, 123, 211–228, <https://doi.org/10.1016/j.atmosres.2012.05.016>, 2013.
- Mohr, S., Kunz, M., and Keuler, K.: Development and application of a logistic model to estimate the past and future hail potential in Germany, *Journal of Geophysical Research: Atmospheres*, <https://doi.org/10.1002/2014JD022959>.Abstract, 2015.
- Mohr, S., Wilhelm, J., Wandel, J., Kunz, M., Portmann, R., Punge, H. J., Schmidberger, M., Quinting, J. F., and Grams, C. M.: The role of large-scale dynamics in an exceptional sequence of severe thunderstorms in Europe May-June 2018, *Weather and Climate Dynamics*, 1, 325–348, <https://doi.org/10.5194/wcd-1-325-2020>, 2020.
- 1045

- Morgan, G. M. and Towery, N. G.: On the Role of Strong Winds in Damage to Crops by Hail and Its Estimation with a Simple Instrument, *Journal of Applied Meteorology* (1962-1982), 15, 891–898, <http://www.jstor.org/stable/26177518>, 1976.
- Morrison, H., Lier-Walqui, M., Fridlind, A. M., Grabowski, W. W., Harrington, J. Y., Hoose, C., Korolev, A., Kumjian, M. R., Milbrandt, J. A., Pawlowska, H., Posselt, D. J., Prat, O. P., Reimel, K. J., Shima, S., Diedenhoven, B., and Xue, L.: Confronting the Challenge of Modeling Cloud and Precipitation Microphysics, vol. 12, ISBN 9781107415324, <https://doi.org/10.1029/2019ms001689>, 2020.
- Mulholland, J. P., Nesbitt, S. W., and Trapp, R. J.: A case study of terrain influences on upscale convective growth of a supercell, *Monthly Weather Review*, 147, 4305–4324, <https://doi.org/10.1175/MWR-D-19-0099.1>, 2019.
- Mulholland, J. P., Nesbitt, S. W., Trapp, R. J., and Peters, J. M.: The influence of terrain on the convective environment and associated convective morphology from an idealized modeling perspective, *Journal of the Atmospheric Sciences*, 77, 3929–3949, <https://doi.org/10.1175/JAS-D-19-0190.1>, 2020.
- Murillo, E. M. and Homeyer, C. R.: Severe hail fall and hailstorm detection using remote sensing observations, *Journal of Applied Meteorology and Climatology*, 58, 947–970, <https://doi.org/10.1175/JAMC-D-18-0247.1>, 2019.
- Murillo, E. M. and Homeyer, C. R.: What Determines Above-Anvil Cirrus Plume Infrared Temperature?, *Journal of the Atmospheric Sciences*, 79, 3181–3194, <https://doi.org/10.1175/JAS-D-22-0080.1>, 2022.
- Nelson, T. C., Marquis, J., Varble, A., and Friedrich, K.: Radiosonde observations of environments supporting deep moist convection initiation during RELAMPAGO-CACTI, *Monthly Weather Review*, 149, 289–309, <https://doi.org/10.1175/MWR-D-20-0148.1>, 2021.
- Nelson, T. C., Marquis, J., Peters, J. M., and Friedrich, K.: Environmental controls on simulated deep moist convection initiation occurring during RELAMPAGO-CACTI, *Journal of the Atmospheric Sciences*, 79, 1941–1964, <https://doi.org/10.1175/JAS-D-21-0226.1>, 2022.
- Nesbitt, S. W., Salio, P. V., Ávila, E., Bitzer, P., Carey, L., Chandrasekar, V., Deierling, W., Dominguez, F., Dillon, M. E., Garcia, C. M., Gochis, D., Goodman, S., Hence, D. A., Kosiba, K. A., Kumjian, M. R., Lang, T., Luna, L. M., Marquis, J., Marshall, R., McMurdie, L. A., de Lima Nascimento, E., Rasmussen, K. L., Roberts, R., Rowe, A. K., Ruiz, J. J., São Sabbas, E. F., Saulo, A. C., Schumacher, R. S., Skabar, Y. G., Machado, L. A. T., Trapp, R. J., Varble, A. C., Wilson, J., Wurman, J., Zipser, E. J., Arias, I., Bechis, H., and Grover, M. A.: A storm safari in subtropical South America: Proyecto RELAMPAGO, *Bulletin of the American Meteorological Society*, 102, E1621–E1644, <https://doi.org/10.1175/BAMS-D-20-0029.1>, 2021.
- Nisi, L., Hering, A., Germann, U., and Martius, O.: A 15-year hail streak climatology for the Alpine region, *Quarterly Journal of the Royal Meteorological Society*, 144, 1429–1449, <https://doi.org/10.1002/qj.3286>, 2018.
- Nixon, C. J. and Allen, J. T.: Distinguishing between Hodographs of Severe Hail and Tornadoes, *Weather and Forecasting*, 37, 1761–1782, <https://doi.org/10.1175/WAF-D-21-0136.1>, 2022.
- Niyogi, D., Holt, T., Zhong, S., Pyle, P. C., and Basara, J.: Urban and land surface effects on the 30 July 2003 mesoscale convective system event observed in the southern Great Plains, *Journal of Geophysical Research Atmospheres*, 111, 1–20, <https://doi.org/10.1029/2005JD006746>, 2006.
- Nomokonova, T., Griewank, P. J., Löhnert, U., Miyoshi, T., Necker, T., and Weissmann, M.: Estimating the benefit of Doppler wind lidars for short-term low-level wind ensemble forecasts, *Quarterly Journal of the Royal Meteorological Society*, 149, 192–210, <https://doi.org/10.1002/qj.4402>, 2023.
- O’Neill, M. E., Orf, L., Heymsfield, G. M., and Halbert, K.: Hydraulic jump dynamics above supercell thunderstorms, *Science*, 373, 1248–1251, <https://doi.org/10.1126/science.abh3857>, 2021.
- Pacey, G., Pfahl, S., Schielicke, L., and Wapler, K.: The climatology and nature of warm-season convective cells in cold-frontal environments over Germany, *Natural Hazards and Earth System Sciences*, 23, 3703–3721, <https://doi.org/10.5194/nhess-23-3703-2023>, 2023.

- 1085 Pacey, G. P., Schultz, D. M., and Garcia-Carreras, L.: Severe convective windstorms in europe: Climatology, preconvective environments, and convective mode, *Weather and Forecasting*, 36, 237–252, <https://doi.org/10.1175/WAF-D-20-0075.1>, 2021.
- Panosetti, D., Böing, S., Schlemmer, L., and Schmidli, J.: Idealized large-eddy and convection-resolving simulations of moist convection over mountainous terrain, *Journal of the Atmospheric Sciences*, 73, 4021–4041, <https://doi.org/10.1175/JAS-D-15-0341.1>, 2016.
- Panziera, L., James, C. N., and Germann, U.: Mesoscale organization and structure of orographic precipitation producing flash floods in the lago maggiore region, *Quarterly Journal of the Royal Meteorological Society*, 141, 224–248, <https://doi.org/10.1002/qj.2351>, 2015.
- 1090 Pesaresi, M. and Politis, P.: GHS-BUILT-S R2023A - GHS built-up surface grid, derived from Sentinel2 composite and Landsat, multitemporal (1975-2030), <https://doi.org/10.48601/EARTHENV-T9K2-1407>, 2023.
- Peters, J. M., Nowotarski, C. J., Mulholland, J. P., and Thompson, R. L.: The influences of effective inflow layer streamwise vorticity and storm-relative flow on supercell updraft properties, *J. Atmos. Sci.*, 77, 3033–3057, <https://doi.org/10.1175/JAS-D-19-0355.1>, 2020.
- 1095 Peters, J. M., Chavas, D. R., Lebo, Z. J., and Su, C.-Y.: Cumulonimbus clouds convert a smaller fraction of CAPE into kinetic energy in a warmer atmosphere, *Journal of the Atmospheric Sciences*, pp. 1943–1961, <https://doi.org/10.1175/jas-d-24-0006.1>, 2024.
- Pistotnik, G., Holzer, A. M., Kaltenböck, R., and Tschannett, S.: An F3 downburst in Austria-A case study with special focus on the importance of real-time site surveys, *Atmospheric Research*, 100, 565–579, <https://doi.org/10.1016/j.atmosres.2010.10.011>, 2011.
- Pounds, L. E., Ziegler, C. L., Adams-Selin, R. D., and Biggerstaff, M. I.: Analysis of Hail Production via Simulated Hailstone Trajectories in the 29 May 2012 Kingfisher, Oklahoma, Supercell, *Monthly Weather Review*, 152, 245–276, <https://doi.org/10.1175/MWR-D-23-0073.1>, 2024.
- 1100 Pruppacher, H. R. and Klett, J. D.: *Microphysics of Clouds and Precipitation*, Atmospheric and Oceanographic Sciences Library, Springer Netherlands, ISBN 9780306481000, <https://books.google.de/books?id=0MURkyjuoGMC>, 2010.
- Púčik, T., Groenemeijer, P., Ryva, D., and Kolář, M.: Proximity soundings of severe and nonsevere thunderstorms in central Europe, *Monthly Weather Review*, 143, 4805–4821, <https://doi.org/10.1175/MWR-D-15-0104.1>, 2015.
- 1105 Púčik, T., Groenemeijer, P., Rädler, A. T., Tijssen, L., Nikulin, G., Prein, A. F., van Meijgaard, E., Fealy, R., Jacob, D., and Teichmann, C.: Future changes in European severe convection environments in a regional climate model ensemble, *Journal of Climate*, 30, 6771–6794, <https://doi.org/10.1175/JCLI-D-16-0777.1>, 2017.
- Púčik, T., Castellano, C., Groenemeijer, P., Kühne, T., Rädler, A. T., Antonescu, B., and Faust, E.: Large hail incidence and its economic and societal impacts across Europe, *Monthly Weather Review*, 147, 3901–3916, <https://doi.org/10.1175/MWR-D-19-0204.1>, 2019.
- 1110 Pucillo, A., Miglietta, M. M., Lombardo, K., and Manzato, A.: Application of a simple analytical model to severe winds produced by a bow echo like storm in northeast Italy, *Meteorological Applications*, 27, 1–18, <https://doi.org/10.1002/met.1868>, 2019.
- Punge, H. J. and Kunz, M.: Hail observations and hailstorm characteristics in Europe : A review, *Atmospheric Research*, 176-177, 159–184, <https://doi.org/10.1016/j.atmosres.2016.02.012>, 2016.
- 1115 Punge, H. J., Bedka, K. M., Kunz, M., and Reinbold, A.: Hail frequency estimation across Europe based on a combination of overshooting top detections and the ERA-INTERIM reanalysis, *Atmospheric Research*, 198, 34–43, <https://doi.org/10.1016/j.atmosres.2017.07.025>, 2017.
- Rädler, A. T., Groenemeijer, P. H., Faust, E., Sausen, R., and Púčik, T.: Frequency of severe thunderstorms across Europe expected to increase in the 21st century due to rising instability, *npj Climate and Atmospheric Science*, 2, 3–7, <https://doi.org/10.1038/s41612-019-0083-7>, 2019.
- 1120

- Rasmussen, E. N., Straka, J. M., Davies-Jones, R., Doswell, C. A., Carr, F. H., Eilts, M. D., and MacGorman, D. R.: Verification of the Origins of Rotation in Tornadoes Experiment: VORTEX, *Bulletin of the American Meteorological Society*, 75, 995–1006, [https://doi.org/https://doi.org/10.1175/1520-0477\(1994\)075<0995:VOTOOR>2.0.CO;2](https://doi.org/https://doi.org/10.1175/1520-0477(1994)075<0995:VOTOOR>2.0.CO;2), 1994.
- 1125 Raupach, T. H., Martius, O., Allen, J. T., Kunz, M., Trapp, S. L., Mohr, S., Rasmussen, K. L., and Trapp, R. J.: The effects of climate change on hailstorms, *Nature Reviews Earth and Environment*, pp. 32–36, <https://doi.org/10.1038/s43017-020-00133-9>, 2021.
- Ravazzani, G., Amengual, A., Ceppi, A., Homar, V., Romero, R., Lombardi, G., and Mancini, M.: Potentialities of ensemble strategies for flood forecasting over the Milano urban area, *Journal of Hydrology*, 539, 237–253, <https://doi.org/10.1016/j.jhydrol.2016.05.023>, 2016.
- Reeves, H. D. and Lin, Y. L.: The effects of a mountain on the propagation of a preexisting convective system for blocked and unblocked flow regimes, *Journal of the Atmospheric Sciences*, 64, 2401–2421, <https://doi.org/10.1175/JAS3959.1>, 2007.
- 1130 Ringhausen, J., Chmielewski, V., and Calhoun, K.: Inter-Comparison of Lightning Measurements in Quasi-Linear Convective Systems, *Atmosphere*, 15, 309, <https://doi.org/10.3390/atmos15030309>, 2024.
- Ripberger, J. T., Krocak, M. J., Wehde, W. W., Allan, J. N., Silva, C., and Jenkins-Smith, H.: Measuring tornado warning reception, comprehension, and response in the United States, *Weather, Climate, and Society*, 11, 863–880, <https://doi.org/10.1175/WCAS-D-19-0015.1>, 2019.
- 1135 Rombeek, N., Leinonen, J., and Hamann, U.: Exploiting radar polarimetry for nowcasting thunderstorm hazards using deep learning, *Natural Hazards and Earth System Sciences*, 24, 133–144, <https://doi.org/10.5194/nhess-24-133-2024>, 2024.
- Romps, D. M.: Clausius-clapeyron scaling of CAPE from analytical solutions to RCE, *Journal of the Atmospheric Sciences*, 73, 3719–3737, <https://doi.org/10.1175/JAS-D-15-0327.1>, 2016.
- Rotach, M. W., Serafin, S., Ward, H. C., Arpagaus, M., Colfescu, I., Cuxart, J., De Wekker, S. F., Grubišić, V., Kalthoff, N., Karl, T., 1140 Kirshbaum, D. J., Lehner, M., Mobbs, S., Paci, A., Palazzi, E., Bailey, A., Schmidli, J., Wittmann, C., Wohlfahrt, G., and Zardi, D.: A Collaborative Effort to Better Understand, Measure, and Model Atmospheric Exchange Processes over Mountains, *Bulletin of the American Meteorological Society*, 103, E1282–E1295, <https://doi.org/10.1175/BAMS-D-21-0232.1>, 2022.
- Rotunno, R. and Houze, R. A.: Lessons on orographic precipitation from the Mesoscale Alpine Programme, *Quarterly Journal of the Royal Meteorological Society*, 133, 811–830, <https://doi.org/10.1002/qj.67>, 2007.
- 1145 Rotunno, R. and Klemp, J. B.: On the rotation and propagation of simulated supercell thunderstorms., *J. Atmos. Sci.*, 42, 271–292, [https://doi.org/10.1175/1520-0469\(1985\)042<0271:OTRAPO>2.0.CO;2](https://doi.org/10.1175/1520-0469(1985)042<0271:OTRAPO>2.0.CO;2), 1985.
- Rudlosky, S. D., Goodman, S. J., and Virts, K. S.: Chapter 16 - Lightning Detection: GOES-R Series Geostationary Lightning Mapper, in: *The GOES-R Series*, edited by Goodman, S. J., Schmit, T. J., Daniels, J., and Redmon, R. J., pp. 193–202, Elsevier, ISBN 978-0-12-814327-8, <https://doi.org/https://doi.org/10.1016/B978-0-12-814327-8.00016-0>, 2020.
- 1150 Ryzhkov, A., Zhang, P., Bukovčić, P., Zhang, J., and Cocks, S.: Polarimetric Radar Quantitative Precipitation Estimation, *Remote Sensing*, 14, <https://doi.org/10.3390/rs14071695>, 2022.
- Saharia, M., Kirstetter, P. E., Vergara, H., Gourley, J. J., and Hong, Y.: Characterization of floods in the United States, *Journal of Hydrology*, 548, 524–535, <https://doi.org/10.1016/j.jhydrol.2017.03.010>, 2017.
- Saltikoff, E., Haase, G., Delobbe, L., Gaussiat, N., Martet, M., Idziorek, D., Leijnse, H., Novák, P., Lukach, M., and Stephan, K.: OPERA 1155 the radar project, *Atmosphere*, 10, 1–13, <https://doi.org/10.3390/atmos10060320>, 2019.
- Scarino, B., Iitterly, K., Bedka, K., Homeyer, C. R., Allen, J., Bang, S., and Cecil, D.: Deriving Severe Hail Likelihood from Satellite Observations and Model Reanalysis Parameters Using a Deep Neural Network, *Artificial Intelligence for the Earth Systems*, 2, 1–30, <https://doi.org/10.1175/aies-d-22-0042.1>, 2023.

- Scheffknecht, P., Serafin, S., and Grubišić, V.: A long-lived supercell over mountainous terrain, *Quarterly Journal of the Royal Meteorological Society*, 143, 2973–2986, <https://doi.org/10.1002/qj.3127>, 2017.
- Schmid, T., Portmann, R., Villiger, L., Schröer, K., and Bresch, D. N.: An open-source radar-based hail damage model for buildings and cars, *Natural Hazards and Earth System Sciences*, 24, 847–872, <https://doi.org/10.5194/nhess-24-847-2024>, 2024.
- Schmidli, J.: Daytime heat transfer processes over mountainous terrain, *Journal of the Atmospheric Sciences*, 70, 4041–4066, <https://doi.org/10.1175/JAS-D-13-083.1>, 2013.
- Schmidt, M. L.: Improvement of hail detection and nowcasting by synergistic combination of information from polarimetric radar, model predictions, and in-situ observations, Ph.D. thesis, Rheinische Friedrich-Wilhelms-Universität Bonn, <https://hdl.handle.net/20.500.11811/8701>, 2020.
- Schultz, C. J., Carey, L. D., Schultz, E. V., and Blakeslee, R. J.: Insight into the kinematic and microphysical processes that control lightning jumps, *Weather and Forecasting*, 30, 1591–1621, <https://doi.org/10.1175/WAF-D-14-00147.1>, 2015.
- Schwartz, C. S. and Sobash, R. A.: Generating probabilistic forecasts from convection-allowing ensembles using neighborhood approaches: A review and recommendations, *Monthly Weather Review*, 145, 3397–3418, <https://doi.org/10.1175/MWR-D-16-0400.1>, 2017.
- Seifert, A., Bachmann, V., Filipitsch, F., Förstner, J., Grams, C. M., Hoshyaripour, G. A., Quinting, J., Rohde, A., Vogel, H., Wagner, A., and Vogel, B.: Aerosol-cloud-radiation interaction during Saharan dust episodes: the dusty cirrus puzzle, *Atmospheric Chemistry and Physics*, 23, 6409–6430, <https://doi.org/10.5194/acp-23-6409-2023>, 2023.
- Serafin, S., Rotach, M. W., Arpagaus, M., Colfescu, I., Cuxart, J., De Wekker, S. F. J., Evans, M., Grubišić, V., Kalthoff, N., Karl, T., Kirshbaum, D. J., Lehner, M., Mobbs, S., Paci, A., Palazzi, E., Raudzens Bailey, A., Schmidli, J., Wohlfahrt, G., and Zardi, D.: Multi-scale transport and exchange processes in the atmosphere over mountains, innsbruck university press, 2020.
- Serafin, S., Schmidli, J., Goger, B., Ban, N., Westerhuis, S., Kotlarski, S., and Arpagaus, M.: TEAMx Numerical Modeling Plan, <https://www.teamx-programme.org/modelling-activities/>, 2024.
- Sessa, M. F. and Trapp, R. J.: Observed relationship between tornado intensity and pretornadic mesocyclone characteristics, *Weather and Forecasting*, 35, 1243–1261, <https://doi.org/10.1175/WAF-D-19-0099.1>, 2020.
- Sgoff, C., Acevedo, W., Paschalidi, Z., Ulbrich, S., Bauernschubert, E., Kratzsch, T., and Potthast, R.: Assimilation of crowd-sourced surface observations over Germany in a regional weather prediction system, *Quarterly Journal of the Royal Meteorological Society*, 148, 1752–1767, <https://doi.org/10.1002/qj.4276>, 2022.
- Shikhov, A. and Chernokulsky, A.: A satellite-derived climatology of unreported tornadoes in forested regions of northeast Europe, *Remote Sensing of Environment*, 204, 553–567, <https://doi.org/10.1016/j.rse.2017.10.002>, 2018.
- Šinger, M. and Púčik, T.: A challenging tornado forecast in Slovakia, *Atmosphere*, 11, <https://doi.org/10.3390/ATMOS11080821>, 2020.
- Smith, B. T., Thompson, R. L., Grams, J. S., Broyles, C., and Brooks, H. E.: Convective modes for significant severe thunderstorms in the contiguous United States. Part I: Storm classification and climatology, *Wea. Forecasting*, 27, 1114–1135, <https://doi.org/10.1175/WAF-D-11-00115.1>, 2012.
- Smith, V. H., Mobbs, S. D., Burton, R. R., Hobby, M., Aoshima, F., Wulfmeyer, V., and Di Girolamo, P.: The role of orography in the regeneration of convection: A case study from the convective and orographically-induced precipitation study, *Meteorologische Zeitschrift*, 24, 83–97, <https://doi.org/10.1127/metz/2014/0418>, 2014.
- Snethlage, M. A., Geschke, J., Ranipeta, A., Jetz, W., Yoccoz, N. G., Körner, C., Spehn, E. M., Fischer, M., and Urbach, D.: A hierarchical inventory of the world’s mountains for global comparative mountain science, *Scientific Data*, 9, 149, <https://doi.org/10.1038/s41597-022-01256-y>, 2022a.

- Snethlage, M. A., Geschke, J., Spehn, E. M., Ranipeta, A., Yoccoz, N. G., Körner, C., Jetz, W., Fischer, M., and Urbach, D.: GMBA Mountain Inventory v2, <https://doi.org/10.48601/EARTHENV-T9K2-1407>, 2022b.
- 1200 Snyder, J. C., Ryzhkov, A. V., Kumjian, M. R., Khain, A. P., and Picca, J.: A ZDR column detection algorithm to examine convective storm updrafts, *Weather and Forecasting*, 30, 1819–1844, <https://doi.org/10.1175/WAF-D-15-0068.1>, 2015.
- Soderholm, B., Ronalds, B., and Kirshbaum, D. J.: The evolution of convective storms initiated by an isolated mountain ridge, *Monthly Weather Review*, 142, 1430–1451, <https://doi.org/10.1175/MWR-D-13-00280.1>, 2014.
- Soderholm, J., Kumjian, M. R., Brook, J. P., Peterson, A., Protat, A., Brimelow, J. C., Trömel, S., and Kunz, M.: Measuring Hail-like Trajectories and Growth with the Hailsonde, *Bull. Amer. Meteor. Soc.*, p. conditionally accepted, 2025.
- 1205 Soderholm, J. S.: Observations of hailstone-like trajectories and growth, 2024.
- Soderholm, J. S. and Kumjian, M. R.: Automating the analysis of hailstone layers, *Atmospheric Measurement Techniques*, 16, 695–706, <https://doi.org/10.5194/amt-16-695-2023>, 2023.
- Soderholm, J. S., Kumjian, M. R., McCarthy, N., Maldonado, P., and Wang, M.: Quantifying hail size distributions from the sky - Application of drone aerial photogrammetry, *Atmospheric Measurement Techniques*, 13, 747–754, <https://doi.org/10.5194/amt-13-747-2020>, 2020.
- 1210 Sokol, Z., Minářová, J., and Novák, P.: Classification of hydrometeors using measurements of the ka-band cloud radar installed at the Milešovka Mountain (Central Europe), *Remote Sensing*, 10, <https://doi.org/10.3390/rs10111674>, 2018.
- Stanelle, T., Vogel, B., Vogel, H., Bäumer, D., and Kottmeier, C.: Feedback between dust particles and atmospheric processes over West Africa during dust episodes in March 2006 and June 2007, *Atmospheric Chemistry and Physics*, 10, 10 771–10 788, <https://doi.org/10.5194/acp-10-10771-2010>, 2010.
- 1215 SwissRe: Natural catastrophes in 2023: gearing up for today’s and tomorrow’s weather risks, Tech. rep., Swiss Re Institute, Zurich, Switzerland, <https://www.swissre.com/institute/research/sigma-research/sigma-2024-01.html>, 2024.
- Tang, B., Vaughan, M., Lazear, R., Corbosiero, K., Bosart, L., Wasula, T., Lee, I., and Lipton, K.: Topographic and boundary influences on the 22 May 2014 Duanesburg, New York, tornadic supercell, *Weather and Forecasting*, 31, 107–127, <https://doi.org/10.1175/WAF-D-15-0101.1>, 2016.
- 1220 Taszarek, M., Allen, J., Púčik, T., Groenemeijer, P., Czernecki, B., Kolendowicz, L., Lagouvardos, K., Kotroni, V., and Schulz, W.: A climatology of thunderstorms across Europe from a synthesis of multiple data sources, *Journal of Climate*, 32, 1813–1837, <https://doi.org/10.1175/JCLI-D-18-0372.1>, 2019.
- Taszarek, M., Allen, J. T., Groenemeijer, P., Edwards, R., Brooks, H. E., Chmielewski, V., and Enno, S. E.: Severe convective storms across Europe and the United States. Part I: Climatology of lightning, large hail, severe wind, and tornadoes, *Journal of Climate*, 33, 10 239–10 261, <https://doi.org/10.1175/JCLI-D-20-0345.1>, 2020a.
- 1225 Taszarek, M., Allen, J. T., Púčik, T., Hoogewind, K. A., and Brooks, H. E.: Severe Convective Storms across Europe and the United States. Part II: ERA5 Environments Associated with Lightning, Large Hail, Severe Wind, and Tornadoes, *Journal of Climate*, 33, 10 263–10 286, <https://doi.org/10.1175/JCLI-D-20-0346.1>, 2020b.
- Taufour, M., Vié, B., Augros, C., Boudevillain, B., Delanoë, J., Delautier, G., Ducrocq, V., Lac, C., Pinty, J.-P., and Schwarzenböck, A.: Evaluation of the two-moment scheme LIMA based on microphysical observations from the HyMeX campaign, *Quarterly Journal of the Royal Meteorological Society*, 144, 1398–1414, <https://doi.org/https://doi.org/10.1002/qj.3283>, 2018.
- 1230 Thompson, R. L., Edwards, R., Hart, J. A., Elmore, K. L., and Markowski, P.: Close Proximity Soundings within Supercell Environments Obtained from the Rapid Update Cycle, *Wea. Forecasting*, 18, 1243–1261, [https://doi.org/10.1175/1520-0434\(2003\)018<1243:cpswse>2.0.co;2](https://doi.org/10.1175/1520-0434(2003)018<1243:cpswse>2.0.co;2), 2003.

- 1235 Thurnherr, I., Cui, R., Velasquez, P., Wernli, H., and Schär, C.: The effect of 3°C global warming on hail over Europe, 2025.
- Trapp, R. J. and Hoogewind, K. A.: The realization of extreme tornadic storm events under future anthropogenic climate change, *Journal of Climate*, 29, 5251–5265, <https://doi.org/10.1175/JCLI-D-15-0623.1>, 2016.
- Trapp, R. J., Kosiba, K. A., Marquis, J. N., Kumjian, M. R., Nesbitt, S. W., Wurman, J. M., Salio, P. V., Grover, M. A., Robinson, P. T., and Hence, D. A.: Multiple-platform and multiple-doppler radar observations of a supercell thunderstorm in south america during relampago, 1240 *Monthly Weather Review*, 148, 3225–3241, <https://doi.org/10.1175/MWR-D-20-0125.1>, 2020.
- Trefalt, S., Martynov, A., Barras, H., Besic, N., Hering, A. M., Lenggenhager, S., Noti, P., Röthlisberger, M., Schemm, S., Germann, U., and Martius, O.: A severe hail storm in complex topography in Switzerland - Observations and processes, *Atmospheric Research*, 209, 76–94, <https://doi.org/10.1016/j.atmosres.2018.03.007>, 2018.
- Trömel, S., Ryzhkov, A. V., Diederich, M., Mühlbauer, K., Kneifel, S., Snyder, J., and Simmer, C.: Multisensor characterization of mammatus, 1245 *Monthly Weather Review*, 145, 235–251, <https://doi.org/10.1175/MWR-D-16-0187.1>, 2017.
- Trömel, S., Blahak, U., Evaristo, R., Mendrok, J., Neef, L., Pejčić, V., Scharbach, T., Shrestha, P., and Simmer, C.: Fusion of radar polarimetry and atmospheric modeling, pp. 293–344, *IET The Institution of Engineering and Technology*, ISBN ISBN-13: 978-1-83953-624-3, <https://doi.org/https://doi.org/10.1049/SBRA557G>, 2023.
- Tuovinen, J. P., Rauhala, J., and Schultz, D. M.: Significant-hail-producing storms in Finland: Convective-storm environment and mode, 1250 *Weather and Forecasting*, 30, 1064–1076, <https://doi.org/10.1175/WAF-D-14-00159.1>, 2015.
- Varble, A. C., Nesbitt, S. W., Salio, P., Hardin, J. C., Bharadwaj, N., Borque, P., DeMott, P. J., Feng, Z., Hill, T. C., Marquis, J. N., Matthews, A., Mei, F., Öktem, R., Castro, V., Goldberger, L., Hunzinger, A., Barry, K. R., Kreidenweis, S. M., McFarquhar, G. M., McMurdie, L. A., Pekour, M., Powers, H., Roms, D. M., Saulo, C., Schmid, B., Tomlinson, J. M., van den Heever, S. C., Zelenyuk, A., Zhang, Z., and Zipser, E. J.: Utilizing a storm-generating hotspot to study convective cloud transitions: The CACTI experiment, *Bulletin of the American Meteorological Society*, 102, E1597–E1620, <https://doi.org/10.1175/BAMS-D-20-0030.1>, 2021.
- Varble, A. C., Igel, A. L., Morrison, H., Grabowski, W. W., and Lebo, Z. J.: Opinion: A critical evaluation of the evidence for aerosol invigoration of deep convection, *Atmospheric Chemistry and Physics*, 23, 13 791–13 808, <https://doi.org/10.5194/acp-23-13791-2023>, 2023.
- Wapler, K.: The life-cycle of hailstorms: Lightning, radar reflectivity and rotation characteristics, *Atmospheric Research*, 193, 60–72, 1260 <https://doi.org/10.1016/j.atmosres.2017.04.009>, 2017.
- Wapler, K.: Mesocyclonic and non-mesocyclonic convective storms in Germany: Storm characteristics and life-cycle, *Atmospheric Research*, 248, 105 186, <https://doi.org/10.1016/j.atmosres.2020.105186>, 2021.
- Wedi, N., Bauer, P., Sandu, I., Hoffmann, J., Sheridan, S., Cereceda, R., Quintino, T., Thiemert, D., and Geenen, T.: Destination Earth: High-Performance Computing for Weather and Climate, *Computing in Science and Engineering*, 24, 29–37, 1265 <https://doi.org/10.1109/MCSE.2023.3260519>, 2022.
- Wehr, T., Kubota, T., Tzeremes, G., Wallace, K., Nakatsuka, H., Ohno, Y., Koopman, R., Rusli, S., Kikuchi, M., Eisinger, M., Tanaka, T., Taga, M., Deghaye, P., Tomita, E., and Bernaerts, D.: The EarthCARE mission - science and system overview, *Atmospheric Measurement Techniques*, 16, 3581–3608, <https://doi.org/10.5194/amt-16-3581-2023>, 2023.
- Wei, W. and Bai, J.: Research Progress on the Numerical Simulation at Gray-zone Scales of the Convective Boundary Layer, *Advances in Earth Science*, 39, 221–231, <https://doi.org/10.11867/j.issn.1001-8166.2024.018>, 2024.
- 1270 Wells, H. M., Hillier, J., Garry, F. K., Dunstone, N., Clark, M. R., Kahraman, A., and Chen, H.: Climatology and convective mode of severe hail in the United Kingdom, *Atmospheric Research*, 309, 107 569, <https://doi.org/10.1016/j.atmosres.2024.107569>, 2024.

- Whiteman, D. C. and Doran, C. J.: The relationship between overlying synoptic-scale flow and winds within a valley, *JAM*, 32, 1993.
- Wieringa, J. and Holleman, I.: If cannons cannot fight hail, what else?, *Meteorologische Zeitschrift*, 2006.
- 1275 Wilhelm, J., Mohr, S., Punge, H. J., Mühr, B., Schmidberger, M., Daniell, J. E., Bedka, K. M., and Kunz, M.: Severe thunderstorms with large hail across Germany in June 2019, *Weather*, 76, 228–237, <https://doi.org/10.1002/wea.3886>, 2021.
- Witt, A., Eilts, M. D., Stumpf, G. J., Johnson, J. T., Mitchell, E. D., and Thomas, K. W.: An enhanced hail detection algorithm for the WSR-88D, *Weather and Forecasting*, 13, 286–303, [https://doi.org/10.1175/1520-0434\(1998\)013<0286:AEHDAF>2.0.CO;2](https://doi.org/10.1175/1520-0434(1998)013<0286:AEHDAF>2.0.CO;2), 1998.
- Wu, F. and Lombardo, K.: Precipitation enhancement in squall lines moving over mountainous coastal regions, *Journal of the Atmospheric Sciences*, 78, 3089–3113, <https://doi.org/10.1175/JAS-D-20-0222.1>, 2021.
- 1280 Wulfmeyer, V., Behrendt, A., Kottmeier, C., Corsmeier, U., Barthlott, C., Craig, G. C., Hagen, M., Althausen, D., Aoshima, F., Arpagaus, M., Bauer, H. S., Bennett, L., Blyth, A., Brandau, C., Champollion, C., Crewell, S., Dick, G., Di Girolamo, P., Dorninger, M., Dufournet, Y., Eigenmann, R., Engelmann, R., Flamant, C., Foken, T., Gorgas, T., Grzeschik, M., Handwerker, J., Hauck, C., Höller, H., Junkermann, W., Kalthoff, N., Kiemle, C., Klink, S., König, M., Krauss, L., Long, C. N., Madonna, F., Mobbs, S., Neininger, B., Pal, S., Peters, G., Pigeon, G., Richard, E., Rotach, M. W., Russchenberg, H., Schwitalla, T., Smith, V., Steinacker, R., Trentmann, J., Turner, D. D., Van Baelen, J., Vogt, S., Volkert, H., Weckwerth, T., Wernli, H., Wieser, A., and Wirth, M.: The Convective and Orographically-induced Precipitation Study (COPS): The scientific strategy, the field phase, and research highlights, *Quarterly Journal of the Royal Meteorological Society*, 137, 3–30, <https://doi.org/10.1002/qj.752>, 2011.
- 1290 Wurman, J., Dowell, D., Richardson, Y., Markowski, P., Rasmussen, E., Burgess, D., Wicker, L., and Bluestein, H. B.: The second verification of the origins of rotation in tornadoes experiment: VORTEX2, *Bulletin of the American Meteorological Society*, 93, 1147–1170, <https://doi.org/10.1175/BAMS-D-11-00010.1>, 2012.
- Zardi, D. and Whiteman, C. D.: *Diurnal Mountain Wind Systems*, pp. 35–119, Springer Netherlands, Dordrecht, ISBN 978-94-007-4098-3, https://doi.org/10.1007/978-94-007-4098-3_2, 2013.
- Zhao, D. and Wu, J.: Changes in urban-related precipitation in the summer over three city clusters in China, *Theoretical and Applied Climatology*, 134, 83–93, <https://doi.org/10.1007/s00704-017-2256-9>, 2018.
- 1295 Zhou, Z., Zhang, Q., Allen, J. T., Ni, X., and Ng, C. P.: How Many Types of Severe Hailstorm Environments Are There Globally?, *Geophysical Research Letters*, 48, 1–11, <https://doi.org/10.1029/2021GL095485>, 2021.