

An estimate of the eddy diffusivity tensor from observed and simulated Lagrangian trajectories in the Benguela Upwelling System

Ria Oelerich¹, Birte Gülk², Julia Dräger-Dietel³, and Alexa Griesel³

¹Institute of Environmental Physics, University of Bremen, Bremen, Germany

²LOCEAN Laboratory, Sorbonne Université-CNRS-IRD-MNHN, Paris, France

³Institut für Meereskunde, Universität Hamburg, Hamburg, Germany

Correspondence: Ria Oelerich (riaoelerich@gmail.com)

Abstract. Lateral mixing of unresolved processes in ocean models is usually parameterized with a scalar diffusivity, although the **observed** mixing can be highly anisotropic. Estimating the full diffusivity tensor from Lagrangian dispersion observations is challenging because shear dispersion from background currents can prevent the diffusive limit from being reached. This study investigates the diffusivity tensor with Lagrangian single and pair particle statistics in the Benguela upwelling region, using one set of Lagrangian trajectories derived from a recent drifter data set with hourly resolution and background currents from the **Ocean Surface Current Analysis Real-time (OSCAR)**, and another set from simulations using the 1/10 ° Parallel Ocean Program (POP) simulation. Theory predicts that pair particle diffusivities, expected to be independent of background mean flows, are twice the single particle diffusivities if the pair velocities are uncorrelated. In this study, it is found that although pair particle diffusivities are much less influenced by mean flow, they are generally significantly smaller than twice the single particle diffusivities. Subtracting the mean flow reduces this discrepancy and improves convergence in both methods, although single particle diffusivities remain higher. Velocity autocorrelations decay faster than pair correlations, with mean flow subtraction accelerating decorrelation, especially in the zonal direction. The pair correlation term in the diffusivity equation contributes significantly to the differences between single particle and pair particle diffusivities, explaining why pair particle diffusivities are generally smaller, making them a less accurate estimate in diffusive parameterizations. In both the POP simulation and the observations, convergence properties improve significantly after mean flow subtraction. Mean flow removal plays a critical role in achieving convergence in the components of the diffusivity tensor and in the major axis component after diagonalization. The significant anisotropy in the diffusivity tensor is mainly explained by the anisotropy in the Lagrangian integral time scales, while the major axis component of the velocity variance tensor is only about 1.2 times the minor axis component. The motions that are not resolved by the OSCAR surface currents product, but captured by the surface drifters, contribute significantly to the diffusivities, accounting for 8% and 42% of the contributions to the zonal and meridional components, respectively, after mean flow subtraction. This study highlights the importance of including the full diffusivity tensor in the Benguela upwelling region in lateral mixing parameterizations.

1 Introduction

Upwelling systems, found in equatorial and coastal regions, are areas where cold, nutrient-rich water from the deep ocean rises to the surface. This phenomenon is primarily driven by subtropical basin-scale circulations that create a slow, equatorward gyre flow near the shore (Capet et al., 2008). All Eastern Boundary Upwelling Systems share specific features such as wind-driven flows, alongshore currents, shallow thermoclines, and significant vertical and offshore nutrient transport (Large and Danabasoglu, 2006). However, each system exhibits unique characteristics in terms of circulation, primary productivity, and phytoplankton biomass, influenced by their specific geographic settings (Berger and Wefer, 2002).

30

The focus of this study is the Benguela Upwelling System, characterized by a frontal system of strong sea surface temperature (SST) gradients. This frontal system is frequently interrupted by finger-like structures of lower SST, known as filaments, resulting from meso- and submesoscale dynamics (Hösen et al., 2016). These filaments are influenced by eddies from two distinct origins: warm-core eddies, known as Agulhas Rings, produced by the interaction between the Agulhas Current and the Antarctic Circumpolar Current (Agulhas Retroflexion) after passing the Agulhas Bank at the southern tip of Africa (Lutjeharms, 2006), and cold-core eddies generated from boundary current instabilities near the coast of Namibia.

The dynamics in coastal upwelling systems like the Benguela Upwelling System range from well-understood processes to complex meso- and submesoscale flow instabilities (Capet et al., 2008). Due to the intricacies of these processes, climate simulations frequently face significant challenges, resulting in notable SST biases, **although adequate resolution of atmospheric processes is important as well** (Small et al., 2015). This underscores the crucial need for ongoing research to enhance the accuracy of climate models and better understand the complexities of upwelling systems.

In coupled climate models with a resolution of 0.5° (~ 50 km) or lower, meso- and submesoscale eddies at their smallest spatial scales are likely to remain unresolved (Richter, 2015). Eddies contribute significantly to the transport of momentum and tracers (e.g., salinity and temperature) due to the comparatively weak mean ocean currents in upwelling systems, and play a crucial role in the oceanic transport of colder water from the coast to the interior. In most climate models, the effects of eddies on a tracer T are parametrized using the tracer conservation equation:

$$\frac{\partial \overline{T}}{\partial t} + \overline{\mathbf{u}} \cdot \nabla \overline{T} = \nabla \cdot \overline{\mathbf{u}'T'} + \overline{Q}, \quad (1)$$

50 where the overline denotes an average over intervals in time and space and $'$ the deviation from that average, while \overline{Q} describes the sources and sinks of the tracer T . Using the Transformed Eulerian Mean framework, the eddy flux can be decomposed into components across and along isolines of the mean tracer, e.g., diffusive and skew-diffusive fluxes (e.g. Griesel et al., 2019). Using $\overline{\mathbf{u}'T'} = -\kappa \nabla \overline{T} + \mathbf{B} \times \nabla \overline{T}$ it follows:

$$\frac{\partial \overline{T}}{\partial t} + (\overline{\mathbf{u}} + \nabla \times \mathbf{B}) \cdot \nabla \overline{T} = \nabla \cdot \kappa \nabla \overline{T} + \overline{Q}, \quad (2)$$

55 where \mathbf{B} is the vector stream function defining the eddy-driven advection velocity $\nabla \times \mathbf{B}$ depending on the tracer T , and κ is a
3 \times 3 diffusivity tensor denoting irreversible mixing of the tracer T . This diffusivity tensor κ contains both lateral and vertical
mixing components and has to be estimated for parametrizing eddies in climate models. Thus, an accurate representation of
eddies in climate models is crucial to improve the general understanding of involved oceanic processes and their impact on
future climate predictions (e.g. Ernst et al., 2023; Huot et al., 2022; Hewitt et al., 2020). This study focuses on the lateral
60 components of the diffusivity tensor.

Diffusivities can be quantified using both Eulerian and Lagrangian methods (Griesel et al., 2014, 2019; Klocker and Aber-
nathey, 2014). The Lagrangian approach is based on the spreading of floats or tracer as they follow the flow (Taylor, 1953, 1921;
Nakamura, 1996), whereas Eulerian diffusivities can be quantified **relative to a fixed geographical location** e.g. from Eulerian
65 eddy tracer fluxes (e.g. Eden, 2006; Griesel et al., 2014). The traditional way to estimate eddy diffusivities originates from the
dispersion of Lagrangian particles (Taylor, 1921, 1953), which provides the means to test the applicability of the eddy-diffusion
model, where particles are expected to spread in a diffusive manner only after some time period determined by the time and
space-scales of the largest eddies (e.g. LaCasce et al., 2008). There are two possible statistical methods to estimate diffusivities,
referred to as single and pair particle statistics. Single particle statistics measure the spreading of particles from their origin and
70 depend strongly on the background mean flow. Pair particle statistics measure how two particles spread apart and would not
depend on the background mean flow if both particles experience the same mean flow during their spreading (Dräger-Dietel
et al., 2018; Sansò et al., 2017). In a flow with homogeneous statistics, the pair particle diffusivity is expected to be twice the
single particle diffusivity.

75 In climate models, lateral eddy diffusion is typically represented by a single scalar, κ_i , aligned along isopycnal surfaces below
the mixed layer. This approach is suitable when eddy diffusion is isotropic in the horizontal or along-isopycnal directions. In
the presence of strong background flows, eddy diffusion is suppressed in the cross-stream component (Ferrari and Nikurashin,
2010; Klocker et al., 2012b, a; Klocker and Abernathey, 2014; Griesel et al., 2014) and eddy mixing becomes anisotropic
(Bachman et al., 2020). Shear dispersion increases the effective diffusivity in the along-stream component (Taylor, 1953; Oh
80 et al., 2000). This shear dispersion in the real ocean results from large-scale background mean flow, but it also occurs due to
the eddies themselves, as large velocity shears exist at the eddy rims. The prevailing premise is that it is the mesoscale eddy
motions with time scales of weeks and space scales corresponding to the first Rossby Radius of deformation that contribute
to eddy mixing (e.g. Ni et al., 2020; Stammer, 1998). Mixing is likely to be most efficient at the edge of the eddies, where
large temperature gradients and large horizontal velocity shears occur. However, we further hypothesize that submesoscale
85 motions with space scales below the first internal Rossby Radius of deformation and associated with time scales smaller than
a few days may contribute substantially. Recently, Peng et al. (2015) and Peng et al. (2020) diagnosed symmetric and shear
instability to occur at the southern boundary of a filament in the Benguela Upwelling System, the same one that is investigated
in this study. Diffusion might become more isotropic once the effect of the shear dispersion associated with the local shear
from the mesoscale eddies is subtracted.

Dispersion processes are particularly important in the Benguela Upwelling System, where filaments are elongated in an east-west direction by the action of the velocities at the eddy rims, leading to increased mixing in this direction. This effect should be captured in ocean models. However, the question remains whether mixing in the along-stream component can be adequately described with a diffusivity. Oh et al. (2000) illustrated how the along-stream component of the diffusivity tensor is influenced by mean flow shear and does not always converge to a constant value. Oh et al. (2000) further advised using the minor axis component of the diffusivity tensor, which is oriented across the background mean flow, as the diffusivity estimate. The minor axis component of the diffusivity tensor can be estimated from drifter data (Zhurbas et al., 2014; Rühls et al., 2018). Davis (1987) and Davis (1991) devised the underlying theory how to compute the diffusivity tensor in principle in the presence of an inhomogeneous background mean flow where, instead of diagnosing the statistics from the absolute Lagrangian velocities and displacements, diffusivities are computed from residual velocities and displacements after the Eulerian mean has been subtracted.

While most observational studies have focused on the minor axis of the diffusivity tensor, only a few, such as Rypina et al. (2012) in the North Atlantic and Peng et al. (2015) in the Indian Ocean, have considered all components of the diffusivity tensor from observational drifter data. Griesel et al. (2010, 2014) and Chen et al. (2014, 2015) demonstrated that in many cases, the along-stream component of the diffusivity tensor can also be determined once the Eulerian mean is subtracted at each float point. However, they used trajectories from eddying ocean models where the background mean flow is known at each grid point.

In this study, for the first time, all components of the diffusivity tensor are estimated from the Lagrangian single and pair particle statistics (Taylor, 1921; Davis, 1987, 1991; LaCasce et al., 2008) in the Benguela upwelling region. Additionally, our study compares the effect of the mean flow on both single and pair particle diffusivities for the first time simultaneously. Furthermore, we specifically quantify the contribution not resolved by the altimeter product to the diffusivity. An observed set of drifters (Dräger-Dietel et al., 2018) and simulated trajectories from drifters deployed in a high-resolution state-of-the-art eddy permitting global ocean model (the 1/10° Parallel Ocean Program, hereinafter referred to as POP simulation) are analyzed in detail. Similarly to Rypina et al. (2012), the background flows from the [Ocean Surface Current Near Real-Time \(OSCAR\)](#) (Bonjean and Lagerloef, 2002) are considered in the analysis of the surface drifter data set. This study aims to address the following questions:

- What is the dependence of the components of the diffusivity tensor on time lag, and can a diffusive limit be reached for all components?
- How do the diffusivities depend on eddy-mean flow decompositions considering both single and pair particle statistics, and can the pair particle diffusivity emerge as an alternative to the single particle diffusivity, since it should be independent from the mean flow, which does not resolve the motions induced by the eddies?

- What is the role of smaller scale motions that are captured by the drifters but not by current altimeter products, and with time scales smaller than a few days for the diffusivities and anisotropy?

125 The study is organized as follows: Section 2 presents the data sets used for this study, including the observed and simulated drifter trajectories. Section 3 describes the theory and methods used to estimate all components of the diffusivity tensor and the mean flow decomposition. Section 4 examines the prevailing background mean flow and eddy kinetic energy in the Benguela Upwelling System. Section 5 evaluates all components of the diffusivity tensor for single and pair particle statistics and highlights the impact of the background mean flow decomposition on the eddy diffusivities. Auto- and pair correlations
130 are examined in detail, and the impact of motions not resolved by satellite altimetry and their contributions on the diffusivities is investigated. Section 7 focuses on the evaluation of anisotropy by decomposing the diffusivity tensor in major and minor axis components in the Benguela upwelling system. Section 8 presents the main conclusions and suggestions for future work.

2 Data

The observational data set includes 35 surface drifters. The drifters of the type SVP-I-XDGS from MetOcean, consist of a
135 surface buoy and a subsurface drogue, mounted in a 16" hull and equipped with Iridium telemetry and an SST sensor (Dräger-Dietel et al., 2018). With the drogue at 15 m depth, they provide the ability to monitor currents from this depth with less influence from surface winds.

The drifters were deployed in the eastern South Atlantic in November/December 2016 by RV METEOR during the cruise
140 M132 from Walvis Bay (Namibia) to Cape Town (South Africa) (Dräger-Dietel et al., 2018). The drifters were deployed at the northern boundary (at 26.0°S, 12.58°E) and the southern boundary (26.4°S, 12.0°E) of a filament identified by satellite altimetry (Fig. 1a). The drifters were released as triplets in a triangular shape with initial separations of 100 to 200 m. In total, there were four drifter release sites, with the first and third groups released at the southern boundary and the second and fourth groups released at the northern boundary of the filament. After approximately 150 days, the drifters were transferred to the
145 Global Drifter Program (GDP), where the temporal resolution was increased from 30 minutes to 60 minutes. For this study, the drifter data set with the lower resolution as in Elipot et al. (2016) is used for a period of 250 days.

The set of simulated drifter trajectories consists of 948 drifters released at 10 m depth in the 1/10° global Parallel Ocean Program (POP) simulation. The run was initialized from the ocean state by Maltrud et al. (2010), which was integrated for 120
150 years using the annually repeating normal year Coordinated Ocean-ice Reference Experiments (CORE) surface forcing (Large and Yeager, 2004). In 1983, the CORE forcing was switched to vary interannually (Large and Yeager, 2009) and the model was integrated for an additional 27 years (Chouksey et al., 2022). The simulated drifters were released on November 2, 1996 with a spacing of 1/5° in both latitude and longitude between 24°S to 30°S and 10°E to 18°E, covering a filament in the Benguela Upwelling System (Fig. 1b). The drifters are advected online for each model time step by the three-dimensional velocity field
155 with a 4th order Runge-Kutta scheme (Griesel et al., 2010). We note that in our analysis period of 250 days, the particles do




Figure 1. The drifter trajectories are shown for a) surface drifter observations and b) particles advected with the velocities from the POP simulation from the deployment date (red dots) to 120 days after deployment (orange dots). The background colors display the snapshots of SST for the observations and the POP simulation. For the observations the SST data are shown for the day of 28/11/2016 and represent measurements from the moderate resolution imaging spectroradiometer (MODIS) provided by Earth Observing System (EOS) (dataset accessed November 2020). For the POP simulation the SST is shown for the day of 28/11/1996.

not move vertically, but remain in the 5-15m depth range.

Moreover, in the analysis of the observational drifter data set, the OSCAR product is used to investigate the surface conditions with respect to the background mean flow and the eddy kinetic energy. The OSCAR product is provided by Earth Space Research (Earth Space Research, 2009, dataset accessed November 2020). The dataset is on a $1/3^\circ$ grid with a 5 day temporal resolution. The OSCAR product offers near-surface ocean current estimates that are derived using quasi-linear and steady flow momentum equations. The spatial velocities in this dataset are directly estimated from sea surface height (SSH), surface winds, and sea surface temperature (SST) measurements taken by various altimetric satellites and in-situ instruments.

3 Theory and Methods

165 This section outlines the theoretical framework and methodologies used to investigate all components of the diffusivity tensor through both single and pair particle statistics, focusing on diffusive behavior, mean flow dependence, and smaller-scale motions.

170 3.1 Single particle diffusivities

In a one-dimensional, stationary flow with zero mean, the mean square displacement r^2 of particles satisfies

$$\frac{d\langle r^2(t) \rangle}{dt} = \left\langle 2 \frac{dr(t)}{dt} r(t) \right\rangle \quad (3)$$

$$= \left\langle 2u(t) \int_0^t u(t') dt' \right\rangle \quad (4)$$

$$= 2 \int_0^t \langle u(t)u(t') \rangle dt' \quad (5)$$

$$175 \quad = 2 \int_0^t R(t') dt', \quad (6)$$

where $r(t)$ is the displacement of the particles at time t from their position at time $t = 0$, $\langle \rangle$ denotes an average over all particles considered, and $R(t)$ is the Lagrangian velocity autocovariance, which is characterized by its value at the origin $R(0) = \langle u(0)^2 \rangle$ and the Lagrangian integral time scale

$$T_{int} = \frac{1}{R(0)} \int_0^\infty R(t') dt'. \quad (7)$$

180 For times much larger than the integral time scale it follows that

$$\langle r^2(t) \rangle = 2Kt \quad \text{with} \quad K = \langle u^2 \rangle T_{int} \quad (8)$$

i.e., in analogy to molecular diffusion, the mean square displacement grows linearly with time with constant turbulent diffusivity K (Taylor, 1921; LaCasce et al., 2008). In reality, with finite times and spatially varying background flows, the integral may not necessarily converge and the diffusivity and time scale are not converging to a constant.

185

In a two-dimensional flow field that includes a spatially varying background current, the diffusivity is a 2x2 symmetric tensor

and the equations generalize to

$$\kappa_{ij} = \frac{1}{2} \frac{d}{dt} \langle r'_i(t) r'_j(t) \rangle \quad (9)$$

$$= \int_0^t \langle u'_i(t) u'_j(t') \rangle dt' \quad (10)$$

$$\stackrel{190}{=} \frac{\langle r'_i(t) r'_j(t) \rangle}{2t} \quad (11)$$

$t \gg T_{int}$

with indices i and j both run over the zonal (1) and meridional (2) velocity and displacement directions. If the velocity statistics are homogeneous and stationary, equations 9-11 lead to the same diffusivity - for real oceanic flows, the methods can lead to different results since the averaging $\langle \rangle$ is applied at different stages (LaCasce et al., 2014). In this study, the integral of the velocity autocovariance (equation 10) is used to calculate the single particle diffusivities.

195

3.2 Flow decompositions

Following Davis (1987) and Davis (1991), in equations (9)-(11) the $u'(t)$ are the drifter velocities at time t with the spatially varying Eulerian background flow removed and the $r'(t)$ are the displacements with displacement due to the background flow removed. In that sense, the diffusivity introduced by Davis (1987) is a mixed Eulerian-Lagrangian quantity that can directly be related to the eddy tracer fluxes in a diffusive parameterization. The diffusivity hence reflects the turbulent mixing due to the residual components of the flow field and depends on the flow decomposition. In this study, two different flow decompositions are tested to estimate the contribution of different flow components to magnitude and orientation of the diffusivity tensor:

$$\mathbf{u}'(t) = \mathbf{u}(t) - \overline{\mathbf{U}}(x, y)|_{\mathbf{r}(t)}, \quad \mathbf{u}''(t) = \mathbf{u}(t) - \mathbf{U}(x, y, t_E)|_{\mathbf{r}(t)} \quad (12)$$

205 where x is the zonal direction and y is the meridional direction, and the subscript $|\mathbf{r}(\tau)$ indicates that the Eulerian velocities U are evaluated at the float positions and times and t_E is the time at which the Eulerian currents are available.

In the first decomposition, the time mean but spatially varying Eulerian background flow $\overline{\mathbf{U}}$ is interpolated to each drifter location and subsequently removed from the drifter velocities. In the observational drifter data set $\overline{\mathbf{U}}$ is the time mean velocity of the $1/3^\circ$ OSCAR surface currents interpolated to the drifter locations, while in the POP simulation $\overline{\mathbf{U}}$ is the time mean of the $1/10^\circ$ Eulerian currents interpolated to the numerical drifter locations (Table 1). Note the current altimeter-derived products, such as the OSCAR product, could have effective resolutions in space and time closer to about 100 km and 30 days (Ballarotta et al., 2019) and should in those instances be regarded as a coarse-grained velocity.

215 In the second decomposition, which is only applied to the observed drifter data that captures motions with an hourly temporal

220 resolution, the time dependent background velocities $U(x, y, t)$ from the OSCAR surface currents product are interpolated spatially and temporarily to match each drifter location and time and are subsequently removed from the drifter velocities. The $U(x, y, t)$ is the spatially varying OSCAR surface current available at 5-day intervals, and allows for the estimation of the contribution of velocity variations to the diffusivity tensor that are not captured by OSCAR. These contributions can be associated with unresolved mesoscale motions and also submesoscale motions that contain significant ageostrophic components (e.g. North et al., 2024). In addition, we can use the velocities from the OSCAR surface currents product only to estimate diffusivities from \mathbf{u} and \mathbf{u}' (Table 1).

velocity field	description
\mathbf{u}_{OBS}	Total velocity at hourly drifter locations
\mathbf{u}'_{OBS}	Total velocity at hourly drifter locations minus $1/3^\circ$ OSCAR yearly mean at drifter locations
$\mathbf{u}''_{\text{OBS}}$	Total velocity at hourly drifter locations minus $1/3^\circ$ OSCAR 5-day means at drifter locations and times
$\mathbf{u}_{\text{OSCAR}}$	$1/3^\circ$ OSCAR 5-day means at drifter locations and times
$\mathbf{u}'_{\text{OSCAR}}$	$1/3^\circ$ OSCAR 5-day means at drifter locations and times minus OSCAR yearly mean at drifter locations
$\overline{\mathbf{U}}_{\text{OSCAR}}$	Eulerian $1/3^\circ$ OSCAR yearly mean velocity
\mathbf{u}_{POP}	Total $1/10^\circ$ POP velocity at daily numerical drifter locations
\mathbf{u}'_{POP}	Total $1/10^\circ$ POP velocity at daily numerical drifter locations minus $1/10^\circ$ POP yearly mean at numerical drifter locations
$\overline{\mathbf{U}}_{\text{POP}}$	Eulerian $1/10^\circ$ POP yearly mean velocity

Table 1. Summary of different flow components used for diffusivity calculations.

225 To visualize the displacements due to different flow components, similarly as in Rypina et al. (2012) and Griesel et al. (2010), trajectories are calculated using the following residual velocities:

$$r'(t) = \int_0^t u'(t') dt', \quad r''(t) = \int_0^t u''(t') dt'. \quad (13)$$

These trajectories are referred to as pseudo-trajectories, because in contrast to the trajectories obtained from advection with the Eulerian flow fields, the pseudo-trajectories are artificial trajectories obtained from integration with residual velocities.

230 3.3 Pair particle diffusivities

As an alternative to the single particle diffusivities, pair particle diffusivity is calculated considering two particles m, n and their separation s and separation velocity $\nu = u^m - u^n$

$$\kappa^{rel}(t) = \frac{1}{2} \frac{d\langle s^2(t) \rangle}{dt} \quad (14)$$

$$= \langle s(t)\nu(t) \rangle = \int_0^t \langle \nu(t)\nu(t') \rangle dt' \quad (15)$$

$$235 \quad = 2 \int_0^t \langle u^m(t)u^m(t') \rangle dt' - 2 \int_0^t \langle u^m(t)u^n(t') \rangle dt' \quad (16)$$

$$= 2\kappa - 2 \int_0^t \langle u^m(t)u^n(t') \rangle dt'. \quad (17)$$

From equations 16-17 it is seen that the pair particle diffusivity κ^{rel} is twice the single particle diffusivity κ if the individual velocities are uncorrelated. For turbulent diffusion the average correlation is positive for pair distances smaller than the correlation length (Koszalka et al., 2009) and leads to a reduced $\kappa_{rel}(t)$, where the reduction depends on the average initial pair separation. The pair particle diffusivity is expected to be less dependent on the background mean flow because any influence from the same background flow experienced by both particles is automatically subtracted in their relative dispersion. However, if the particles move apart into regions with different background flows, the relative diffusivity will include contributions from the background shear. As for the single particle diffusivity, all tensor components of the pair particle diffusivity are considered using the zonal and meridional particle separations $s_i = x^m - x^n, s_j = y^m - y^n$ and the zonal and meridional separation velocity $\nu_i = u_i^m - u_i^n, \nu_j = v_i^m - v_j^n$:

$$245 \quad \kappa_{ij}^{rel}(t) = \frac{1}{2} \frac{d\langle s_i(t)s_j(t) \rangle}{dt} \Big|_{t > T_{int}} = \frac{\langle s_i(t)s_j(t) \rangle}{2t}. \quad (18)$$

Furthermore, pair particle diffusivities are calculated with the background flows removed, similar to the single particle diagnostics. However, for the pair particle diagnostics, $s'(t)$ is derived from the pseudo-trajectories (equation 13).

250

3.4 Anisotropy

The diffusivity tensor \mathbf{K} , which is symmetric by construction, can be diagonalized $\mathbf{K} = R(\alpha) \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} R(\alpha)^T$, where $R(\alpha) = \begin{pmatrix} \cos(\alpha) & -\sin(\alpha) \\ \sin(\alpha) & \cos(\alpha) \end{pmatrix}$ is a rotation matrix characterizing the orientation of the principal axes of the diffusivity components. The eigenvalues of the symmetric Lagrangian eddy diffusivity tensor are given by

$$255 \quad \lambda_{1,2} = \frac{1}{2} \left(\kappa_{xx} + \kappa_{yy} \pm \sqrt{(\kappa_{xx} - \kappa_{yy})^2 + 4\kappa_{xy}^2} \right), \quad (19)$$

where the angle is given by

$$\tan 2\alpha = 2\kappa_{xy}/(\kappa_{xx} - \kappa_{yy}) \quad (20)$$

(cf. Haigh et al., 2020), and λ_1 is the diffusivity along the major axis of diffusion and corresponds to the eigenvector $\mathbf{e}_1 = (\cos(\alpha), \sin(\alpha))$. As described by Oh et al. (2000), it is the along-stream component that is amplified by the shear dispersion of the background mean flow and does not converge to a constant value. However, when the background mean flow is subtracted from each drifter velocity, the shear dispersion effect should be minimized even in the along-stream direction.

3.5 Bootstrapping and diffusive limit

Finally, the important concept of this study is the investigation with regards to saturated behaviour of the diffusivities (the convergence to a constant value κ^∞) and how this behaviour is affected by the subtraction of the background mean flow. A convergence criterion is established, requiring the standard deviation to be less than 15% of the average eddy diffusivity from 100 to 250 days. This criterion ensures consistent comparison of convergence properties and diffusivities.

Given the limited sample size, an error estimation is derived through bootstrapping. This involves subsampling 35 bootstrap samples from the original data sets and conducting the analysis for single and pair particles with 100 repetitions. From these values, the mean and standard deviation from the mean is calculated. This approach yields a more accurate standard deviation than a simple calculation of the standard deviation from the original datasets.

4 Background mean flow, eddy kinetic energy and pseudo-trajectories

The annual mean surface currents in the POP simulation (Fig. 2a,b) closely resemble those in the OSCAR surface current product, both showing a predominantly northwestward Benguela Current near Africa's western coast between 6-18°E. In both datasets, the mean surface currents are primarily westward across the region. Generally, the zonal mean flow at the drifter locations is stronger in the POP simulation than in the OSCAR product, particularly near the coast, while further offshore, the flows are similar in magnitude. Snapshots of surface currents from both datasets, taken 200 days after drifter release (5-day interpolation for OSCAR and daily for the POP simulation; Fig. 2c,d), reveal extensive eddy activity of similar magnitudes, filling the upwelling region. These eddies, many originating from the Agulhas Retroflexion site, appear in both datasets, although the POP simulation benefits from higher spatial and temporal resolution. The eddy kinetic energy (EKE) calculated from the annual means of both products (Fig. 2e,f) shows higher EKE magnitudes along the drifter trajectories in the POP simulation compared to the OSCAR product.

285

The POP simulation with numerical trajectories is only available for 1996/97, while the observational dataset was obtained

in 2016/17. Regardless, as Fig. 2e,f illustrates, the horizontal distributions of kinetic energies are similar. There is visible interannual variability for the region (Fig. 3a) that is largely driven by the variability of the Agulhas rings. While there are discrepancies between OSCAR and POP particularly around 2004, the area-averaged energy levels in 1996 in POP (red solid and dashed lines in Fig. 3a) and in 2016 for the OSCAR product (black solid and dashed lines in Fig. 3a) are similar, and so are the total kinetic energies of POP and OSCAR averaged along the drifter locations (red and cyan diamonds in Fig. 3a). However, the total kinetic energy of the surface drifter dataset at hourly resolution averaged over the surface drifter locations in the observations is substantially higher (black diamond in Fig. 3a). This is largely due to inertial oscillations, visible as semidiurnal oscillations in the time series of the observational drifter velocities (black lines in Fig. 3b,c) as compared to the velocities obtained from POP and OSCAR that are similar in magnitude (red and cyan lines in Fig. 3b,c) but do not capture motions on these time scales.

The pseudo-trajectories (Fig. 4) reveal various motions occurring at different scales and due to different processes. Both the observations and the POP simulation exhibit a dominant westward mean flow. This is evident from the zonal extent of the pseudo-trajectories with the mean flow subtracted (Fig. 4a,d), which is reduced by up to 1000 km in both datasets. The total motion of particles (Fig. 4a,c, black trajectories) in both datasets consists of three components: the northwestward motion driven by the mean Benguela Current, which shows a stronger zonal flow in the POP simulation and a stronger northward component in the observations (Fig. 4b,d, grey trajectories); the motion caused by eddies, which includes a net westward component as the eddies entrain filaments of cold water offshore (Fig. 4b,d, red trajectories) as also described in Dräger-Dietel et al. (2018).




Figure 2. The annual means of surface current speeds are calculated for a) observations (OSCAR product, 2016-2017) and b) the POP simulation (1996-1997, 15 m depth). The annual mean current speeds are superimposed with velocity vectors (black arrows) for each respective data set. The reference vector is given in red. 5-day interpolation (OSCAR) and daily (POP simulation) snapshots of current speeds (c,d) 200 days after the drifter release are presented to emphasize the strong daily variations in current speeds. The eddy kinetic energy (e,f), calculated from the annual means, highlights the effects of eddies on the background mean flow after the drifter deployments. Blue markers are as in Fig. 1.




Figure 3. The time series of annual mean kinetic energies for POP (red) compared with the OSCAR product (black) averaged over the Benguela upwelling region (as shown in Fig. 1) are shown in panel a), where the total kinetic energies (solid lines) and the annual mean background flows (dashed lines) are displayed. Additionally, the total kinetic energies averaged over all float locations for POP, the observations and the OSCAR velocities interpolated to drifter locations (red, black and cyan diamonds) are shown. The time series of zonal and meridional velocities averaged over all drifters are displayed in panels b) and c) for POP (red), the observations (black) and OSCAR velocities interpolated to drifter locations (cyan).




Figure 4. The pseudo-trajectories (equation (13)) for the observations (a,b) and the POP simulation (c,d). Note that all trajectory components are shown in the cartesian coordinate system. For the POP every 5th trajectory is shown to improve visibility of underlying patterns. The original trajectories are marked in shades of blue (a,c) and the pseudo-trajectories from the integration with \mathbf{u}' (equation (12)) are marked in shades of red (a,c). The pseudo-trajectories from the integration with the OSCAR surface currents $\mathbf{u}_{\text{OSCAR}}$ and the POP simulation daily mean velocities \mathbf{u}_{POP} are highlighted in shades of blue (b,d) and the ones with the annual mean velocities interpolated to the drifter locations $\bar{\mathbf{U}}|_{r(t)}$ are shown in shades of red (b,d). All pseudo-trajectories have been constructed such that (0,0) is the origin. Note that the coloring of trajectories was chosen such that every trajectory has a slightly different color in the range of blue or red so that patterns of individual trajectories are highlighted.

5 The effect of the mean flow on eddy diffusivities

5.1 Diffusivities

As introduced in section 3, the diffusivity reflects the turbulent mixing according to residual components of the flow field and thus depends on the flow decomposition. In this section, the eddy diffusivities calculated from the drifter velocities before and after subtraction of the annually-averaged and spatially varying mean flow are compared for the observations and the POP simulation and for the single and pair particle approaches. This allows to investigate the contribution of each mean flow decomposition on the diffusivity tensor and also enables a comparison of the mean flow contributions within each of the data sets and methods, for example the comparison of single and pair particle-derived diffusivities. Note that the diffusivities derived from pair particle statistics are from this point onward defined as $\kappa_{xx,xy,yx,yy}^{rel*} = 0.5 \cdot \kappa_{xx,xy,yx,yy}^{rel}$ (equation (14)) to allow for a direct comparison between pair and single particle diffusivities. Before the effect of the mean flow on single particle diffusivities is discussed, we note that the eddy diffusivities from the POP simulation and observations are comparable to each other as discussed in section 4.

For single particle diffusivities without mean flow subtraction (Fig. 5, red), the κ_{xx} component is smaller in the POP simulation $((50.98 \pm 10.28) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$ as compared with the observations $((60.14 \pm 10.15) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$. Similarly, the κ_{yy} component is also smaller in the POP simulation $((1.56 \pm 0.39) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$ as compared with the observations $((4.59 \pm 0.55) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$. In contrast, the off-diagonal component $\kappa_{xy} = \kappa_{yx}$ shows larger diffusivities in the POP simulation $((3.84 \pm 1.54) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$ than in the observations, **the latter exhibiting** negative eddy diffusivities $((-3.80 \pm 4.95) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$.

The mean flow subtraction from absolute drifter velocities (Fig. 5, blue) affects the component of the diffusivity tensor that is most strongly impacted by the mean flow. κ_{xx} represents the zonal component of the diffusivity tensor and demonstrates strongest reduction of diffusivities after subtraction of the mean flow across both datasets. This is to be expected as the mean flow in the Benguela Upwelling System is predominantly zonal as discussed in section 4. The reductions of κ_{xx} are about 55% for the observations and 86% for the POP simulation. Thus, the zonal mean flow contribution is largest for the diffusivities of the POP simulation. After the mean flow subtraction, κ_{xx} fulfills the convergence criterion (section 3) for the POP simulation and the observations.

The κ_{xy} components are reduced by 55% in the POP simulation, while in the observations they show an increase in the observations of 342% from negative diffusivities to positive diffusivities. It follows that largest reductions of κ_{xy} are again found for the POP simulation respectively, but the reductions are smaller in percentage than for the κ_{xx} component for the POP simulation. Additionally, for κ_{xy} for the POP simulation does not fulfill the convergence criterion and thus do not show saturated behavior after the subtraction of the mean flow. In contrast, κ_{xy} shows convergence towards a constant value after the subtraction of the mean flow.

340 Finally, for the meridional component κ_{yy} the diffusivities show an increase after the subtraction of the mean flow, where the diffusivities of the observations increase by 36% and the diffusivities for the POP simulation increase by 57%. This demonstrates that the subtraction of the mean flow does not necessarily lead to a reduction of diffusivities for all tensor components as the meridional component κ_{yy} increases for both data sets. After the subtraction of the mean flow, the convergence criterion is fulfilled for both datasets and saturates towards a constant value.

345

For the pair particle diffusivities derived from the relative drifter velocities (Fig. 6), the κ_{xx}^{rel*} component for the observations $((3.38 \pm 1.83) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$ is about 55% larger than for the POP simulation $((1.41 \pm 0.26) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$, but still remains the component with the largest eddy diffusivities. The κ_{yy}^{rel*} component also demonstrates almost 78% smaller eddy diffusivities for the POP simulation $((0.92 \pm 0.11) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$ as compared with the observations $((4.31 \pm 0.47) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$. The κ_{xy}^{rel*} components for the POP simulation $((0.73 \pm 0.05) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$ are also about 70% smaller than the components from the observations $((2.5 \pm 0.91) \cdot 10^3 \text{ m}^2 \text{ s}^{-1})$.

In contrast to single particle diffusivities, the pair particle diffusivities show significantly smaller effects on eddy diffusivities after the subtraction of the mean flow. This is to be expected, since the relative drifter velocities are by theory less affected by the mean flow, since they are derived from the velocity difference of the pairs, from which a common mean flow is then automatically subtracted (section 3. κ_{xx}^{rel*} shows that also for pair particle statistics the subtraction of the mean flow does not necessarily lead to a reduction of eddy diffusivities.

The mean flow subtraction demonstrates eddy diffusivities with an increase of 30% for the observations and an increase of 35% for the POP simulation. This indicates that in contrast to single particle diffusivities, which mostly decrease after the mean flow subtraction, κ_{xx}^{rel*} increases for the observations and the POP simulation. After the mean flow subtraction κ_{xx}^{rel*} fulfills the convergence criterion for the POP simulation but not for the observations. For κ_{xy}^{rel*} , there is a similar increasing behavior for both datasets after the mean flow subtraction. The eddy diffusivities change with an increase of 19% for the observations and 36% for the POP simulation. The convergence criterion is fulfilled for the POP simulation, but not for the observations. κ_{yy}^{rel*} shows opposite signs and presents decreasing behaviour of the diffusivities after the subtraction of the mean flow with respect to the other tensor components. The diffusivities after the mean flow subtraction for κ_{yy}^{rel*} show a reduction of 17% for the observations and an increase of 36% for the POP simulation. The convergence criterion after the mean flow subtraction is fulfilled for both datasets. It is important to note that the impact of the mean flow on pair particle diffusivities is smaller than compared with single particle diffusivities across all tensor components.

370

The comparison of single and pair particle diffusivities clearly highlights that specifically for the zonal component κ_{xx} are significantly larger for the single particle approach than for the pair particle approach. This is mostly the case before and after the subtraction of the mean flow. Before the meanflow subtraction the κ_{xx} components with the single particle approach significantly larger by 94% for the observations and 97% for the POP simulation.




Figure 5. Single particle eddy diffusivities are shown without subtraction of the meanflow (red) and with subtraction of the annually-averaged meanflow (blue) as shown in Fig. 2 (a,b). The eddy diffusivities are given for all components of the diffusivity tensor κ for the observations (a,c,e) and the POP simulation (b,e,h) in $\text{m}^2 \text{s}^{-1}$. The vertical black line in each subfigure marks the 100th day after drifter deployment after which the eddy diffusivities are temporarily-averaged. The temporarily-averaged eddy diffusivities are shown with standard deviations below each subfigure in the respective colors with and without meanflow subtraction (blue and red). The shaded areas for each subfigure indicate the uncertainties in eddy diffusivity calculations from bootstrapping (35 subsampled drifters with 100 repetitions).

After the mean flow subtraction, κ_{xx} components compared with κ_{xx}^{rel*} components are larger with about 84% for the observations and 73% for the POP simulation. Although, κ_{xx} and κ_{xx}^{rel*} do not show the same overall behavior neither before nor after the subtraction of the mean flow, it is to note that smaller differences between κ_{xx} and κ_{xx}^{rel*} improve comparability in all data sets. The κ_{xy} components compared with the κ_{xy}^{rel*} components are larger by 166% for the observations and 189%
 380 for the POP simulation before the mean flow subtraction. After the mean flow subtraction the κ_{xy} components compared with κ_{xy}^{rel*} are larger by 68% for the observations and 42% for the POP simulation. The meridional components κ_{yy} compared with κ_{yy}^{rel*} before the mean flow subtraction show that κ_{yy} is larger by 6% for the observation and larger by 41% for the POP simulation. After the subtraction of the mean flow, the κ_{yy} components are larger than κ_{yy}^{rel*} components across both datasets with percentages of 43% for the observations and 48% for the POP simulation.

385

This study further reveals that the convergence criterion is predominantly satisfied for the pair particle approach, where 8 out of 12 tensor components across all data sets before and after subtraction of the mean flow showed saturated behavior. For the single particle approach 6 out of 12 tensor components demonstrated saturated behavior, where for 4 out of 12 tensor components the subtraction of the mean flow had a positive effect on the convergent behavior. The slightly stronger convergent
 390 properties for the pair particle approach are to be expected as relative drifter velocities are less affected by the mean flow. However, the results indicate that the single particle approach generally provides larger diffusivities before and after the mean flow removal, which might indicate that some components of the mean flow might still be affecting the data not only across the zonal but also across the other tensor components. In most cases, the diffusivities from the observations and the POP simulation demonstrate similar values, although the values from the POP simulation are slightly smaller in before and after the mean flow
 395 subtraction.

In summary, our results show that the impact of the mean flow is smaller on pair particle diffusivities than on single particle diffusivities, whereas the subtraction of the mean flow improved comparability between single and pair particle diffusivities for all tensor components and across different datasets. In some cases, the subtraction of the mean flow improved the convergent
 400 properties specifically for the single particle approach. However, even after the subtraction of the mean flow, single particle diffusivities tend to be larger than pair particle diffusivities and are not equal as one might expect. This indicates that there might be an effect of the mean flow varying on scales smaller than the separation distance of the pair (which should be notable especially for longer time lags), or contribution of the correlation term (equation (17)).




Figure 6. Pair particle eddy diffusivities are shown without subtraction of the mean flow (red) and with subtraction of the annually-averaged mean flow (blue) as shown in Fig. 2 (a,b). The eddy diffusivities are given for all components of the diffusivity tensor κ for the observations (a,c,e) and the POP simulation (b,d,f) in $\text{m}^2 \text{s}^{-1}$. The vertical black line and shaded areas mark properties of the plot as in Fig. 5.

5.2 Velocity correlation terms analysis

405 To analyze the effect of the correlation term, the velocity auto correlations (Fig. 7a,b) and the velocity pair correlations (Fig. 7c,d) are compared in detail. The correlations are normalized by their values at zero time lag. In particular, the comparison focuses on the xx and yy correlations after the mean flow is subtracted. Furthermore, the integral time scales from the auto- and pair correlations are calculated as a measure for the decorrelation time scales (equation (7)). It should be noted that if significant negative lobes and oscillations appear in the velocity correlations after the first zero crossing, the integral time scale
410 will be considerably smaller than the time required for the diffusivities to converge.

First, it is noted that the kinetic energy averaged along the drifter trajectories (i.e., the non-normalized velocity autocorrelation at zero time lag) is higher for the observations than for the POP simulation (compare also black and red diamonds in Fig. 3). The total zonal and meridional kinetic energy is approximately 2.5 and 3.2 times greater in the observations compared with
415 the POP simulation. When the mean flows are subtracted, the total zonal and meridional kinetic energy are approximately 3.6 and 3.3 times larger respectively. The increase in the ratio with mean flow subtraction, particularly for the zonal component, is consistent with the stronger mean flow observed in the POP simulation, especially in the zonal direction (Fig. 4), leaving a smaller residual in the POP simulation.

420 The tails of the normalized correlations (Fig. 7) reveal that both the velocity auto-correlations and pair correlations for the xx component do not approach zero when the mean flow is not subtracted. This behavior aligns with the observation that single particle diffusivities continue to increase with time lag (Fig. 5a,b). When the mean flow is subtracted (Fig. 7a,b, red lines), the auto-correlations for the xx components first cross zero at 83 days for the observations and 182 days for the POP simulation. The yy components (Fig. 7a,b, blue lines) decay to zero much faster, occurring after 10 days in the observations
425 and 12 days for the POP simulation, followed by a significant negative lobe within the first 40 days and subsequent oscillations around zero. The meridional components exhibit similar behavior to the auto-correlations with mean flow subtraction (not shown), with the primary difference being reduced values at zero time lag. The times to the first zero crossing are longer for the pair correlation terms. For the xx component, the first zero crossing occurs at 236 days for the observations and 167 days for the POP simulation when the mean flow is subtracted. In contrast, the yy components reach their first zero crossing much
430 earlier, at 10 days for the observations and 67 days for the POP simulation.

The auto-correlation integral time scales (Fig. 7a,b) are slightly longer for the POP simulation compared to the observations and are larger in the zonal direction than in the meridional direction. In contrast, the pair correlation integral time scales (Fig. 7c,d) are significantly larger, ranging from 2 to 5 times those of the auto-correlations.

435

The diffusivity values (Fig. 7) indicate that the correlation term in equation (17) is substantial and plays a key role in explaining the differences between single and pair particle diffusivities. This term is predominantly positive, contributing to the

generally smaller pair particle diffusivities. For instance, the correlation term for the xx component, without mean flow subtraction, is $(51.43 \pm 9.65) \cdot 10^3 \text{m}^2 \text{s}^{-1}$ for both the observations and the POP simulation, accounting for the over 90% higher
440 single particle diffusivities. When the mean flow is subtracted, this correlation contribution decreases significantly, leading to better convergence between single and pair particle diffusivities, although the pair particle diffusivities themselves remain relatively unaffected by the mean flow. Even after mean flow subtraction, the contribution of the pair correlation term for the xx component remains $(3.75 \pm 1.27) \cdot 10^3 \text{m}^2 \text{s}^{-1}$. Notably, there are instances where the correlation contribution is negative; for example, in the xy component of the observations without mean flow subtraction, there is a negative contribution of
445 $(-2.38 \pm 8.25) \cdot 10^3 \text{m}^2 \text{s}^{-1}$, which corresponds to cases where pair particle diffusivity exceeds single particle diffusivity (not shown). It is important to note that these values are sensitive to the chosen averaging interval with large error bars.

In summary, the analysis of the correlation terms reveals that pair diffusivities are not a reliable measure of lateral mixing, primarily because pair velocities remain correlated, especially in the zonal direction, where the first zero crossings occur only
450 after more than 150 days following mean flow subtraction. The velocity auto-correlations decay to zero more rapidly than the pair correlations after mean flow subtraction.

Given that auto-correlation integral time scales are similar in both the observations and the POP simulation, the generally larger single particle diffusivities observed in the observations compared with the POP simulation can be attributed to the
455 higher kinetic energies (compare equation (7)). Furthermore, oscillations in the velocity auto-correlation reflect particle motions with timescales of a few days associated with mesoscale eddies (Griesel et al., 2010). Additionally, inertial oscillations with periods of about half a day, which are only present in the observational data, appear as oscillations in Fig. 7a,c, although due to the small periods they are not identifiable in the figure but appear as thickened lines. As inertial oscillations are associated with circular motions, they are hypothesized to not contribute much to net mean square displacement and thus should not
460 impact the diffusivities. However, the energy contained in the inertial oscillations may contribute to the larger diffusivities in the observations compared with POP, which is further discussed in section 6 and 7.




Figure 7. a,b) show the velocity auto correlations normalized by their value at zero time lag $\langle u(t)u(t') \rangle$ (black), $\langle u'(t)u'(t') \rangle$ (red) and $\langle v'(t)v'(t') \rangle$ (blue), where the $\langle \rangle$ is an average over all particles, as a function of time lag. c,d) display the corresponding normalized velocity pair correlations $\langle u_i^m(t)u_j^n(t') \rangle$ where $\langle \rangle$ is an average over all particle pairs. Also given are the integral time scales $T_{u'u'}$, $T_{v'v'}$, averaged over 100-250 days, (equation (7)) when mean flow is subtracted. a,c) show the results for observations and b,d) for the POP simulation. The temporarily-averaged (100-250 days) eddy diffusivities associated with the pair correlation terms $\int_0^t \langle u_i^m(t)u_j^n(t') \rangle dt'$ (last term on the RHS of equation (17)) are shown with standard deviations below the lower panels for κ_{xx} with and without mean flow subtraction (black and red) and for κ_{yy} with mean flow subtraction (blue) in $\text{m}^2 \text{s}^{-1}$.

6 Effect of motions not resolved by satellite data

This section examines the contribution of unresolved motion in the OSCAR product to diffusivities from the observations by subtracting OSCAR-derived velocities from the total observed velocities, thereby isolating the impact of these unresolved motions on diffusivity estimates. The corresponding pseudotrajectories (Fig. 8) display inertial oscillations as well as other motions that are not resolved by the OSCAR product. The net displacements are found to be about a quarter of the total displacements in the pseudo-trajectories (Fig. 4a).

The diffusivity tensor components for both single and pair particles (Fig. 9a-c) indicate contributions in the range of $(1.00 \text{ to } 2.50) \cdot 10^3 \text{ m}^2 \text{ s}^{-1}$. For single particle diffusivities, the xx and yy components are found to be similar within the error bars, while the xy component becomes negative and do not converge, continuously decreasing with time lag, which suggests the presence of substantial non-zero auto-correlations for this component. Significant differences are observed between single and pair particle diffusivities for the xx and xy components, highlighting persistent pair correlations, whereas the yy components remain consistent within the error bars.

Overall, the contribution to the xx component is approximately 8% of that with the mean flow subtracted, while the contribution to the yy component is around 42% of the mean flow subtracted value. This indicates that diffusivities calculated from trajectories which are advected with altimeter-derived velocities in the Benguela upwelling region are likely to be significantly underestimated, especially in the zonal component.




Figure 8. The pseudo-trajectories from the integration with the residual velocity u'' from the observations. All pseudotrajectories have been constructed such that $(0, 0)$ is the origin.




Figure 9. Single (black) and pair (red) particle eddy diffusivities using $\mathbf{u}''_{\text{OBS}}$ (equation (11)). The eddy diffusivities are given for all components of the diffusivity tensor for the observations (a-c). The vertical black line and shaded areas mark the properties of the subfigures as in Fig. 5.

The diffusivity tensors are characterized in terms of their major and minor axes components, as well as the angle that defines the orientation of the principal axes of the diffusivity components (equation (19)). In the following, the anisotropy of the diffusivity tensor is measured by the ratio of the major to minor axis components.

485 In both the POP simulation and the observations, the major axis components do not converge, but increase continuously, as expected (Fig. 10). However, convergence is observed when the mean flow is removed, with the satisfaction of the convergence criterion, albeit the diffusivity plateaus only for large time lags. For the POP simulation, mean flow subtraction results in a decrease in anisotropy, with the ratio of major to minor diffusivities dropping from about 44 to 4. Averaged over time lags 100-250 days the major axis component drastically decreases from $(50.77 \pm 10.24) \cdot 10^3 \text{ m}^2 \text{ s}^{-1}$ to $(7.77 \pm 0.71) \cdot 10^3 \text{ m}^2 \text{ s}^{-1}$,
 490 while the minor axis component actually increases from $(1.18 \pm 0.57) \cdot 10^3 \text{ m}^2 \text{ s}^{-1}$ to $(1.81 \pm 0.33) \cdot 10^3 \text{ m}^2 \text{ s}^{-1}$, consistent with the increase in the yy component (section 5). In contrast, the observations show an increase in anisotropy that becomes negative, due to the minor axis component being predominantly negative for time lags greater than about 70 days, although the error bars from the bootstrapping samples are large. The absolute value of the minor axis component, averaged over time lags from 100 to 250 days, is less than $(0.4 \pm 1.0) \cdot 10^3 \text{ m}^2 \text{ s}^{-1}$. The negative minor axis component would indicate a decrease in
 495 mean square displacement with time along the minor axis component due to convergence of particles, although the error bars from the standard deviation are large.

The diffusivities can be written as the product of major and minor axis kinetic energies and integral times scales, as in equation (7) (Table 2). It is found that after mean flow subtraction, the ratio of major to minor axis kinetic energies is 1.24 and 1.25 for
 500 the observations and the POP simulation, respectively. It is mainly the integral times scales that explain the large anisotropy in diffusivities. After mean flow subtraction, the major axis time scales are about 14 days for the observations and 12 days for the POP simulation, while the minor axis time scales are -0.21 and 3 days for the observations and the POP simulation respectively. It should be noted that while the small anisotropy in the kinetic energy components is similar in the POP simulation and the observations and the major integral time scales are the same similar, the kinetic energies are larger in the observations
 505 (Table 2) explaining the larger major axis diffusivities compared to the POP simulation. The $R(0)$ are significantly larger in the observations than in the POP simulation, even after subtraction of the OSCAR velocities (third row in Table 2). This implies that the kinetic energy associated with inertial oscillations might contribute to the diffusivities in section 6, even though the integral time scales for u'' are small.

510 Using the pseudotrajectories derived from the contributions u'' not captured by OSCAR, the analysis shows that convergence is not achieved and the major axis diffusivity continuously increases with time lag, while the minor axis diffusivity decreases progressively and becomes negative after time lags exceeding approximately 220 days. The anisotropy remains significant, with a ratio of major to minor diffusivities of around 15.

velocity field	$R(0)_{\text{major}}(\text{cm}^2 \text{ s}^{-2})$	$T_{\text{major}}^{\text{int}}(\text{days})$	$R(0)_{\text{minor}}(\text{cm}^2 \text{ s}^{-2})$	$T_{\text{minor}}^{\text{int}}(\text{days})$
\mathbf{u}_{OBS}	299 ± 26	$22.42 \pm 3.60^{\star}$	228 ± 12	1.52 ± 0.42
\mathbf{u}'_{OBS}	277 ± 25	13.83 ± 1.40	223 ± 12	-0.21 ± 0.57
$\mathbf{u}''_{\text{OBS}}$	165 ± 14	$3.38 \pm 0.53^{\star}$	153 ± 7	$0.24 \pm 0.21^{\star}$
\mathbf{u}_{POP}	119 ± 10	$49.20 \pm 10.75^{\star}$	72 ± 7	1.91 ± 0.95
\mathbf{u}'_{POP}	78 ± 9	11.56 ± 1.71	68 ± 6	3.08 ± 0.63

Table 2. Velocity autocorrelation at zero time lag for major and minor axis components and integral Lagrangian time scales T^{int} in days for different flow velocities (rows). The product of the time scales and velocity autocorrelations is the diffusivities as given in Fig. 10 averaged over time lags 100-250 days.

515 The results show that the angle of the diffusivity ellipse is close to approximately 30° in the observations when the mean flow is subtracted, while it converges to nearly zero without mean flow subtraction (Fig. 10c). Note that the angle is less steep for the POP simulation compared to the observations after mean flow subtraction. When the daily mean is subtracted, the average angle is -30° , however with large error bars (not shown). The angle is expected to be correlated with the mean flow direction, since the maximum diffusion is anticipated to be oriented in the direction of the mean flow. The mean flow has a
520 stronger meridional component in the observations compared with the POP simulation (Fig. 2a,b), especially for longitudes east of 6°E . Further offshore, the mean flow is zonally oriented and perpendicular to the coast for the observations as well as the POP simulation.

In summary, this study highlights that the minor axis diffusivity can become negative and anisotropy occurs for all flow com-
525 ponents. The net westward displacements of the drifters and the anisotropy is not only caused by the action of the mean flow, but also by the action of the mesoscale eddies that entrain the cold, upwelled water westward. Thus, the mixing is suppressed across both strong mean flow and mesoscale motions.




Figure 10. a) Major and minor axis diffusivities (equation (19)) for the POP simulation and the observations without mean flow subtraction. b) Major and minor axis diffusivities for the POP simulation and the observations with mean flow subtraction. c) Angle (equation (20)) for the observations and the POP simulation with and without mean flow subtraction. d) Major and minor axis diffusivities for subtraction with daily mean for the observations. Shown are the means of the diagonalizations of the bootstrapping samples and shadings are the standard deviations respectively. The numbers in black, red and blue are the ratios of the major to minor diffusivities, where the diffusivities were averaged over time lags 100-250 days.

8 Discussion and Conclusions

530 In this study, two sets of drifter trajectories were analyzed: one derived from observational drifter data and another from trajectories simulated using the POP (Parallel Ocean Program) simulation. The model simulation served as a testbed with about 30 times more drifter trajectories than available in the observations. One challenge in diffusivity estimates is that the influence of shear dispersion by background currents often inhibits the attainment of a diffusive limit, which is why many studies focus on

velocity field	$R(0)_{xx}(\text{cm}^2\text{s}^{-2})$	$T_{xx}^{int}(\text{days})$	$R(0)_{yy}(\text{cm}^2\text{s}^{-2})$	$T_{yy}^{int}(\text{days})$	$R(0)_{xy}(\text{cm}^2\text{s}^{-2})$	$T_{xy}^{int}(\text{days})$
u_{OBS}	299 ± 19	$23.24 \pm 4.20 \star$	228 ± 18	2.33 ± 0.34	-7 ± 70	$658.32 \pm 7066.65 \star$
u'_{OBS}	274 ± 18	11.46 ± 1.25	226 ± 19	3.20 ± 0.45	13 ± 6	81.26 ± 41.34
u''_{OBS}	164 ± 11	1.61 ± 0.31	154 ± 9	1.97 ± 0.32	23 ± 3	$-85.10 \pm -130.75 \star$
u_{POP}	119 ± 9	$49.444 \pm 10.64 \star$	72 ± 8	2.51 ± 0.68	-3 ± 1	-213.33 ± 174.97
u'_{POP}	77 ± 8	10.76 ± 1.26	69 ± 8	4.14 ± 0.50	3 ± 1	74.16 ± 44.11
u_{OSCAR}	106 ± 11	$45.94 \pm 9.08 \star$	75 ± 0.0011	5.42 ± 0.90	-0.5 ± 4	$-666.83 \pm 4946.82 \star$
u'_{OSCAR}	85 ± 10	21.35 ± 3.01	74 ± 12	14.36 ± 2.99	8 ± 3	143.23 ± 59.62

Table 3. Velocity autocorrelation at zero time lag and integral Lagrangian time scales T^{int} in days (compare equation 7) for different tensor components (columns) and flow velocities (rows). The product of time scales and velocity autorrelations are the diffusivities as given in Fig. 5,9. A black star indicates that the integral times scales and therefore the diffusivities do not converge for that component.

the minor axis or cross-stream components of the diffusivity tensor. Here, the first goal was to investigate all components of the
diffusivity tensor, study their dependence on mean flow subtraction using available background currents and assess whether
diffusive limits were reached. In the observations, the $1/3^\circ$ OSCAR surface currents product was used, while in the model
the $1/10^\circ$ Eulerian background mean flow is readily available. The single particle diffusivities were also compared with pair
particle diffusivities, that are hypothesized to be less influenced by background mean flows, making them a potential alternative
for single particle diffusivity estimates where high quality mean flows are not available.

The results show, as expected, that single particle diffusivities are much more sensitive to mean flow subtraction than pair
particle diffusivities. In both the POP simulation and the observations, convergence properties significantly improve after mean
flow subtraction. Mean flow removal plays a critical role in achieving convergence in the xx and xy tensor components for the
single particle diffusivities. Pair particle diffusivities, on the other hand, demonstrate more consistent convergence and are less
impacted by the mean flow, likely due to the inherent subtraction of common flow components between particle pairs.

However, pair particle diffusivities are predicted by theory to be twice the single particle diffusivity if the pair velocities
are uncorrelated, and our study demonstrates that pair particle diffusivities are considerably smaller than that even after mean
flow subtraction. We find that velocity autocorrelations decay more rapidly than the pair correlations, with the subtraction of
mean flow leading to even faster decorrelation, particularly in the zonal direction. The integration of the pair correlation term
in the equation for pair particle diffusivity significantly contributes to the observed differences between single and pair particle
diffusivities and explains why pair particle diffusivities are generally smaller. As demonstrated by Davis (1987, 1991) the single
particle diffusivity after mean flow subtraction appears in diffusive parameterizations, so our pair particle estimates would
likely underestimate mixing in diffusive parameterizations. Pair particle diffusivities can be an alternative to the single particle
estimates with mean flow subtraction if the deployment locations are such that the pair velocities are uncorrelated from the

beginning or quickly decorrelate.

The diffusivity can be written as the product of kinetic energy, i.e. the velocity autocorrelation at zero time lag, and the integral time scale and we summarize the results for all flow components in Table 3. The results show that after mean flow subtraction, the kinetic energies for the zonal and meridional directions become more similar to each other with ratios of 1.1-1.2 for the observations, the POP simulation and the OSCAR product. However, the zonal integral time scales are 3.6 (observations), 2.6 (POP) and 1.5 (OSCAR) times larger than the meridional ones. The reduction in time scale for the meridional direction can be related to mixing suppression by a zonal mean flow as discussed in Klocker et al. (2012a), Klocker and Abernathey (2014) and Griesel et al. (2015). The degree of mixing suppression can be based on mixing-length theory but taking eddy propagation relative to the mean flow into account. In the presence of a zonal background flow, the meridional diffusivity can then be written as

$$\kappa_{yy} = \alpha R(0) \frac{\gamma}{\gamma^2 + k^2(\bar{U} - c)^2}, \quad (21)$$

where γ is a typical Lagrangian decorrelation time scale and is equal to the growth rate of unstable waves in linear instability theory (Griesel et al., 2015), k is related to a typical eddy size and can be regarded as the wavenumber of maximum growth in linear instability theory (although both might differ in the presence of an inverse energy cascade), U is the zonal background mean flow, c is a typical phase speed of the eddies and $\alpha = 0.35$ as diagnosed by Klocker and Abernathey (2014). It is the difference of U and c that leads to the mixing suppression effect, otherwise the diffusivity is equal to the velocity autocorrelation at zero lag times the Lagrangian decay scale γ . For the POP simulation we use the eddy sizes and translational speeds as diagnosed from an eddy tracking algorithm (Griesel et al., 2015), which amount to 90 km for the average eddy radius and -0.033 m s^{-1} for the zonal translational speed.

We note that the translational speeds are close to but not necessarily equal to the phase speed in equation (21) (Griesel et al., 2015). With γ as the decay scale (time to first zero crossing) for the meridional direction of about 1/(12 days) and using our mean flow values averaged over the drifter trajectories, the time scale is about 3 days, which is close to the Lagrangian integral time scale obtained for POP for yy (Table 3). On the other hand, if we apply equation 21 to the suppression by the mean meridional flow, using the region-averaged meridional translational eddy speeds of 0.003 m s^{-1} and $\gamma=1/(182 \text{ days})$, we arrive at a time scale of 14 days, which is indeed close to T_{yy} from Table 3. These time scales are also consistent with the ones obtained by Klocker and Abernathey (2014) and Ruhs et al. (2018). We note that mixing suppression in the zonal direction due to the mainly zonally oriented topographic background PV gradient might also be relevant here, a mechanisms that was recently discussed by Sterl et al. (2024).

585

We find that the diffusivities in the observed drifter data set are larger than the ones inferred from the POP trajectories, particularly after mean flow subtraction. After mean flow subtraction, the integral times scales are slightly larger in the POP simulation compared with the observations, but overall similar (second and fifth row of Table 3). However, the velocity autocorrelation at zero lag after mean flow subtraction, is 2-3 times larger in the observations than in the POP simulation and is thus explaining

590 the larger eddy diffusivities in the observations. This is partly due to the fact that the background flow in the POP simulation is larger and more highly resolved in the POP simulation than in the OSCAR surface currents product, hence leaving a smaller EKE residual, and illustrates the importance of using high resolution background flow components. However, also the total kinetic energy ($R(0)$ for \mathbf{u}_{OBS} in Table 3) is larger in the observations than in the POP simulation, while the $R(0)$ when considering the contribution from the OSCAR velocities alone (last two rows in Table 3) are more comparable to POP.

595

This relates to the research question, which addresses the role of smaller scale motions that are detected by drifters but not by current altimeter products in influencing diffusivities and anisotropy. The study highlights the significant contribution of these unresolved motions. We found that these motions contribute 8% to the xx component and 42% to the yy component of diffusivities after mean flow subtraction. The observed differences between single and pair particle diffusivities, particularly in
600 the xx and xy components, emphasize the influence of ongoing pair correlations within the unresolved small-scale dynamics. These small scale motions include inertial oscillations, which are hypothesized to not contribute much to the net diffusion, since they may only lead to oscillations in the velocity autocorrelation, which average out in the integral over time lag. Indeed, as Table 3 shows, the integral times scales for u''_{OBS} are only about 1 day for the xx and yy components. However, the velocity autocorrelation at zero lag is significant and is largely due to the kinetic energy in the inertial motions. Further studies
605 with more trajectories are needed to clarify the role of inertial motions for mixing. The underestimation of diffusivities when using altimeter-derived velocities indicates that small-scale motions substantially influence diffusivity values and anisotropy, with the potential for large discrepancies when these motions are not accounted for. Promising improvements are anticipated from the recent SWOT (Surface Water and Ocean Topography) satellite mission, which will provide much higher resolution in geostrophic surface velocity measurements. However, as highlighted by North et al. (2024), velocities in the Benguela up-
610 welling filament contain significant ageostrophic components at scales smaller than about 15 km, which will not be captured by geostrophic velocities derived from sea surface height data. A promising approach was recently suggested by Zhang and Wolfe (2024) who have shown that eddy diffusivity can be inferred from the eddy tracks without the need for drifter deployments or altimetry-derived geostrophic velocity measurements.

615 Finally, the study finds significant anisotropy (measured by the ratio of major to minor axis components) in the diffusivities with all flow components. Similarly to the difference between the xx and yy components of the diffusivity tensor, the anisotropy of the diffusivity tensor is mainly explained by the difference in major and minor axis time scales, since the anisotropy in the velocity variance ellipses is small and similar in both observations and the POP simulation (compare Table 2). The major axis components of the diffusivity tensor increase with time when no mean flow is subtracted, while the major axis components
620 converge in the POP simulation and the observations after the mean flow subtraction. The minor axis component in the observations becomes negative for time lags greater than about 70 days after mean flow subtraction. This highlights the complex nature of diffusivity behavior, where the Lagrangian diffusivity tensor is not necessarily positive definite, as mean square displacement can decrease over time. This might be due to the presence of convergent velocities associated with the filament. Negative minor axis components were also recently found by Haigh et al. (2020), who diagnosed Eulerian eddy diffusivities

625 from tracer flux-gradient relations in an eddying North Atlantic model simulation, as well as by Chen et al. (2014) who also diagnosed a significant number of negative Lagrangian cross-stream eddy diffusivities in the Kuroshio Extension. Negative Lagrangian diffusivities are in principle consistent with mean square displacements decreasing over time and net up-gradient eddy tracer fluxes.

630 Subtracting the mean flow reduces the anisotropy in the POP simulation, but increases it in the observations due to the close to zero (negative) minor axis component (division by very small number). Mean flow subtraction aligns the diffusivity ellipse more closely with the mean flow direction (orientation north-westward). Hence, there is a difference between the yy component of the diffusivity tensor and the minor axis component. This finding is different from what was found for the Indian Ocean from Global Drifter Program data in Peng et al. (2015), where the mean flow was predominantly meridional and the zonal
635 diffusivity in most cases aligned with the minor axis component, and the meridional diffusivity aligned with the major axis component. The anisotropy illustrates the role of mesoscale eddies in contributing to westward displacements and anisotropy while suppressing mixing across strong mean and mesoscale flows.

Overall, the minor axis components before mean flow subtraction are in the order of $3000 \text{ m}^2\text{s}^{-1}$ for the observations and
640 $1000 \text{ m}^2\text{s}^{-1}$ for the POP simulation which is similar to what has been found previously for this component from Lagrangian observations (Zhurbas et al., 2014; R  hs et al., 2018; Peng et al., 2015). However we find that the minor axis component is also influenced by the mean flow subtraction with a 1.5 fold increase for the POP simulation but a drastic decrease for the observations to a small negative value. This may depend on the temporal averaging interval used for the mean flow, which is one year in our case. The values of major and minor axis components might change if a longer averaging interval is used.

645

Overall, our findings emphasize the necessity of using the full 2×2 diffusivity tensor to represent lateral mixing in climate models, rather than simplified isotropic representations. We speculate that the current warm SST bias in climate models in the Eastern boundary upwelling regions may be reduced if the offshore, predominantly zonally oriented mixing of cold upwelled water by the action of eddies entraining the cold water of the filaments is represented properly.

650 *Data availability.* The observed drifter trajectories are part of the Global Drifter Program (GDP, <https://www.aoml.noaa.gov/phod/gdp/index.php>). The simulated drifters advected within the POP simulation are available using the following link <https://zenodo.org/records/13710842>. The data from the Ocean Surface Current Near-Real Time (OSCAR) product are available at http://podaac.jpl.nasa.gov/dataset/OSCAR_L4_OC_third-deg.

Appendix A

655 *Author contributions.* RO and AG developed the concept for the manuscript. RO and AG lead the writing process. RO, BG and AG processed and analyzed the observational and simulated drifter data. The field experiment design for drifter deployments was led by JD-D and AG. BG, AG, JD-D provided valuable input and guidance for the development of this study.

Competing interests. The contact author has declared that none of the authors has any competing interests.

660 *Acknowledgements.* We thank the science party and the crew of cruise M132 on R/V Meteor for their support and Kerstin Jochumsen as lead scientist. This paper is a contribution to the project L3 (Meso- to submesoscale turbulence in the ocean) of the Collaborative Research Centre TRR 181 “Energy Transfer in Atmosphere and Ocean” funded by the German Research Foundation (DFG, grant agreement no. 274762653).
We thank the editor and the reviewers for their constructive comments, suggestions and questions to improve our manuscript.

References

- Bachman, S., Fox-Kemper, B., and Bryan, F. O.: A diagnosis of anisotropic eddy diffusion from a high-resolution global ocean model.,
665 *Journal of Advances in Modelling Earth Systems*, 12, 1–14, <https://doi.org/10.1029/2019MS001904>, 2020.
- Ballarotta, M., Ubelmann, C., Pujol, M.-I., Taburet, G., Fournier, F., Legeais, J.-F., Faugère, Y., Delepouille, A., Chelton, D., Dibarboure,
G., and Picot, N.: On the resolutions of ocean altimetry maps, *Ocean Science*, 15, 1091–1109, <https://doi.org/10.5194/os-15-1091-2019>,
2019.
- Berger, W. and Wefer, G.: On the reconstruction of upwelling history: Namibia upwelling in context, *Marine Geology*, 180, 3–28,
670 [https://doi.org/10.1016/S0025-3227\(01\)00203-1](https://doi.org/10.1016/S0025-3227(01)00203-1), 2002.
- Bonjean, F. and Lagerloef, G.: Diagnostic Model and Analysis of the Surface Currents in the Tropical Pacific Ocean, *Journal of Physical
Oceanography*, 32, 2938–2954, [https://doi.org/10.1175/1520-0485\(2002\)032<2938:DMAAOT>2.0.CO;2](https://doi.org/10.1175/1520-0485(2002)032<2938:DMAAOT>2.0.CO;2), 2002.
- Capet, X., McWilliams, F., Penven, J. C., and Marchesiello, P.: Eddies in Eastern Boundary Subtropical Upwelling Systems, *Geophysical
Monograph Series*, 177, 131–147, <https://doi.org/10.1029/177GM10>, 2008.
- 675 Chen, R., McClean, J. L., Gille, S. T., and Griesel, A.: Isopycnal Eddy Diffusivities and Critical Layers in the Kuroshio Extension from an
Eddy Ocean Model, *Journal of Physical Oceanography*, 44, 2191 – 2211, <https://doi.org/10.1175/JPO-D-13-0258.1>, 2014.
- Chen, R., Gille, S., McClean, J., Flierl, G., and Griesel, A.: A multi-wavenumber theory for eddy diffusivities and its application to the
southeast Pacific (DIMES) region, *J. Phys. Oceanogr.*, <https://doi.org/10.1175/JPO-D-14-0229.1>, 2015.
- Chouksey, A., Griesel, A., Chouksey, M., and Eden, C.: Changes in Global Ocean Circulation due to Isopycnal Diffusion, *Journal of Physical
680 Oceanography*, 52, 2219 – 2235, <https://doi.org/10.1175/JPO-D-21-0205.1>, 2022.
- Davis, R. E.: Modeling eddy transport of passive tracers, *Journal of Marine Research*, 45, 635–666,
<https://doi.org/10.1357/002224087788326803>, 1987.
- Davis, R. E.: Observing the general circulation with floats, *Deep Sea Research Part A, Oceanographic Research Papers*, 38, 531–571,
[https://doi.org/10.1016/S0198-0149\(12\)80023-9](https://doi.org/10.1016/S0198-0149(12)80023-9), 1991.
- 685 Dräger-Dietel, J., Jochumsen, K., Griesel, A., and Badin, G.: Relative Dispersion of Surface Drifters in the Benguela Upwelling Region,
Journal of Physical Oceanography, 48, 2325–2341, <https://doi.org/10.1175/JPO-D-18-0027.1>, 2018.
- Earth Space Research: OSCAR third degree resolution ocean surface currents Version 1, <https://doi.org/10.5067/OSCAR-03D01>, 2009.
- Eden, C.: Thickness diffusivity in the Southern Ocean, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2006GL026157>, 2006.
- Elipot, S., Lumpkin, R., Perez, R. C., Lilly, J. M., Early, J. J., and Sykulski, A. M.: A global surface drifter data set at hourly resolution,
690 *Journal of Geophysical Research: Oceans*, 121, 2937–2966, <https://doi.org/10.1002/2016JC011716>, 2016.
- Ernst, P. A., Subrahmanyam, B., Trott, C. B., and Chaigneau, A.: Characteristics of submesoscale eddy structures within mesoscale eddies
in the Gulf of Mexico from 1/48° ECCO estimates, *Frontiers in Marine Science*, 10, <https://doi.org/10.3389/fmars.2023.1181676>, 2023.
- Ferrari, R. and Nikurashin, M.: Suppression of eddy mixing across jets in the Southern Ocean., *Journal of Physical Oceanography*, 40,
1501–1519, <https://doi.org/10.1175/2010JPO4278.1>, 2010.
- 695 Griesel, A., Gille, S. T., Sprintall, J., McClean, J. L., LaCasce, J. H., and Maltrud, M. E.: Isopycnal Diffusivities in the Antarc-
tic Circumpolar Current Inferred from Lagrangian Floats in an Eddy Model, *Journal of Geophysical Research: Oceans*, 115,
<https://doi.org/10.1029/2009JC005821>, 2010.
- Griesel, A., McClean, M. J., Gille, S. T., Sprintall, J., and Eden, C.: Eulerian and Lagrangian isopycnal eddy diffusivities in the Southern
Ocean from an eddy ocean model., *Journal of Physical Oceanography*, 44, 2191–2211, <https://doi.org/10.1175/JPO-D-13-039.1>, 2014.

- 700 Griesel, A., Eden, C., Koopmann, N., and Yulaeva, E.: Comparing isopycnal eddy diffusivities in the Southern Ocean with predictions from linear theory, *Ocean Modelling*, 94, 33–45, <https://doi.org/10.1016/j.ocemod.2015.08.001>, 2015.
- Griesel, A., Dräger-Dietel, J., and Jochumsen, K.: Diagnosing and parameterizing the effects of oceanic eddies, in: *Energy transfers in atmosphere and ocean*, edited by Eden, C. and Iske, A., *Mathematics of Planet Earth*, pp. 193–224, Springer-Verlag, https://doi.org/10.1007/978-3-030-05704-6_6, 2019.
- 705 Haigh, M., Sun, L., Shevchenko, I., and Berloff, P.: Tracer-based estimates of eddy-induced diffusivities, *Deep-Sea Res. I*, 160, 103–264, <https://doi.org/10.1016/j.dsr.2020.103264>, 2020.
- Hewitt, H. T., Roberts, M., Mathiot, P., Biastoch, A., Blockley, E., Chassignet, E. P., Fox-Kemper, B., Hyder, P., Marshall, D. P., Popova, E., Treguier, A.-M., Zanna, L., Yool, A., Yu, Y., Beadling, R., Bell, M., Kuhlbrodt, T., Arsouze, T., Bellucci, A., Castruccio, F., Gan, B., Putrasahan, D., Roberts, C. D., Van Roekel, L., and Zhang, Q.: Resolving and Parameterising the Ocean Mesoscale in Earth System Models, *Current Climate Change Reports*, 6, 137–152, <https://doi.org/10.1007/s40641-020-00164-w>, 2020.
- 710 Hösen, E., Möller, J., Jochumsen, K., and Quadfasel, D.: Scales and Properties of cold filaments in the Benguela Upwelling System off Lüderitz, *Journal of Geophysical Research: Oceans*, 121, 1896–1913, <https://doi.org/10.1002/2015JC011411>, 2016.
- Huot, P.-V., Kittel, C., Fichet, T., Jourdain, N. C., and Fettweis, X.: Effects of ocean mesoscale eddies on atmosphere–sea ice–ocean interactions off Adélie Land, East Antarctica, *Climate Dynamics*, 59, 41–60, <https://doi.org/10.1007/s00382-021-06115-x>, 2022.
- 715 Klocker, A. and Abernathy, R.: Global patterns of mesoscale eddy properties and diffusivities., *J. Phys. Oceanogr.*, 44, 1030–1046, <https://doi.org/10.1175/JPO-D-13-0159.1>, 2014.
- Klocker, A., Ferrari, R., and LaCasce, J. H.: Estimating suppression of eddy mixing by mean flows., *J. Phys. Oceanogr.*, 9, 1566–1576, <https://doi.org/10.1175/JPO-D-11-0205.1>, 2012a.
- Klocker, A., Ferrari, R., LaCasce, J. H., and Merrifield, S. T.: Reconciling float-based and tracer-based estimates of eddy diffusivities., *J. Mar. Res.*, 70, 569–602, <https://doi.org/10.1357/002224012805262743>, 2012b.
- 720 Koszalka, I., Lacasce, J., and Orvik, K.: Relative dispersion in the Nordic Seas, *Journal of Marine Research*, 67, 411–433, <https://doi.org/10.1357/002224009790741102>, 2009.
- LaCasce, J. H., Ferrari, R., Marshall, J., Tulloch, R., Balwada, D., and Speer, K.: Statistics from Lagrangian Observations, *Progress in Oceanography*, 77, 1–29, <https://doi.org/10.1016/j.pocean.2008.02.002>, 2008.
- 725 LaCasce, J. H., Ferrari, R., Marshall, J., Tulloch, R., Balwada, D., and Speer, K.: Float derived isopycnal diffusivities in the DIMES experiment, *Journal of Physical Oceanography*, 44, 764–780, <https://doi.org/10.1175/JPO-D-13-0175.1>, 2014.
- Large, W. G. and Danabasoglu, G.: Attribution and Impacts of Upper-Ocean Biases in CCSM3, *Journal of Climate*, 19, 2325–2346, <https://doi.org/10.1175/JCLI3740.1>, 2006.
- Large, W. G. and Yeager, S. G.: Diurnal to decadal global forcing for ocean and sea-ice models: the datasets and flux climatologies., *NCAR Technical Note TN-460+STR*, 8, National Center for Atmospheric Research, <https://doi.org/10.5065/D6KK98Q6>, 2004.
- 730 Large, W. G. and Yeager, S. G.: The global climatology of an interannually varying air-sea flux data set., *Climate Dynamics*, 33, 341–364, <https://doi.org/10.1007/s00382-008-0441-3>, 2009.
- Lutjeharms, J.: *The Agulhas Current*, Springer Verlag Berlin-Heidelberg, <https://doi.org/10.1007/3-540-37212-1>, 2006.
- Maltrud, M. E., Bryan, F., and Peacock, S.: Boundary impulse response functions in a century-long eddying global ocean simulation, *Environ. Fluid Mech.*, 10, 275 – 295, <https://doi.org/10.1007/s10652-009-9154-3>, 2010.
- 735 Nakamura, N.: Two-Dimensional Mixing, Edge Formation, and Permeability Diagnosed in an Area Coordinate, *Journal of the Atmospheric Sciences*, 53, 1524–1537, [https://doi.org/10.1175/1520-0469\(1996\)053<1524:TDMEFA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1996)053<1524:TDMEFA>2.0.CO;2), 1996.

- Ni, Q., Zhai, X., Wang, G., and Marshall, D. P.: Random Movement of Mesoscale Eddies in the Global Ocean, *Journal of Physical Oceanography*, 50, 2341–2357, <https://doi.org/10.1175/JPO-D-19-0192.1>, 2020.
- 740 North, R. P., Dräger-Dietel, J., and Griesel, A.: Characterization of physical properties of a coastal upwelling filament with evidence of enhanced submesoscale activity and transition from balanced to unbalanced motions in the Benguela upwelling region, *Ocean Science*, 20, 103–121, <https://doi.org/10.5194/os-20-103-2024>, 2024.
- Oh, S. I., Zhurbas, V., and Park, W. S.: Estimating horizontal diffusivity in the East Sea (Sea of Japan) and the northwest Pacific from satellite-tracked drifter data, *Journal of Geophysical Research: Oceans*, 105, 6483 – 6492, <https://doi.org/10.1029/2000JC900002>, 2000.
- 745 Peng, J.-P., Holtermann, P., and Umlauf, L.: Frontal instability and energy dissipation in a submesoscale filament, *Journal of Physical Oceanography*, 50, 2017–2035, <https://doi.org/10.1175/JPO-D-19-0270.1>, 2020.
- Peng, S., Qian, Y.-K., Lumpkin, R., Li, P., Wang, D., and Du, Y.: Characteristics of the Near-Surface Currents in the Indian Ocean as Deduced from Satellite-Tracked Surface Drifters. Part II: Lagrangian Statistics, *Journal of Physical Oceanography*, 45, 459–477, <https://doi.org/10.1175/JPO-D-14-0049.1>, 2015.
- 750 Richter, I.: Climate model biases in the eastern tropical oceans: causes, impacts and ways forward, *Wiley Interdisciplinary Reviews: Climate Change*, 6, 345–358, <https://doi.org/10.1002/wcc.338>, 2015.
- Rühs, S., Zhurbas, V., Koszalka, I. M., Durgadoo, J. V., and Biastoch, A.: Eddy diffusivity estimates from lagrangian trajectories simulated with ocean models and surface drifter data—a case study for the greater agulhas system, *Journal of Physical Oceanography*, 48, 175–196, <https://doi.org/10.1175/JPO-D-17-0048.1>, 2018.
- 755 Rypina, I. I., Kamenkovich, I., Berloff, P., and Pratt, L. J.: Eddy-induced particle dispersion in the Near-Surface North Atlantic, *Journal of Physical Oceanography*, 42, 2206–2228, <https://doi.org/10.1175/JPO-D-11-0191.1>, 2012.
- Sansón, L. Z., Pérez-Brunius, P., and Sheinbaum, J.: Surface Relative Dispersion in the Southwestern Gulf of Mexico, *Journal of Physical Oceanography*, 47, 387–403, <https://doi.org/10.1175/JPO-D-16-0105.1>, 2017.
- Small, J. R., Churchister, E., Hedstrom, K., Kaufmann, B., and Large, W.: The Benguela Upwelling System: Quantifying the Sensitivity to resolution and coastal wind representation in a global climate model, *Journal of Climate*, 28, 9409–9432, <https://doi.org/10.1175/JCLI-D-15-0192.1>, 2015.
- 760 Stammer, D.: On Eddy Characteristics, Eddy Transports, and Mean Flow Properties, *Journal of Physical Oceanography*, pp. 727–739, [https://doi.org/10.1175/1520-0485\(1998\)028<0727:OECETA>2.0.CO;2](https://doi.org/10.1175/1520-0485(1998)028<0727:OECETA>2.0.CO;2), 1998.
- Sterl, M. F., LaCasce, J. H., Groeskamp, S., Nummelin, A., Isachsen, P., and Baatsen, M.: Suppression of Mesoscale Eddy Mixing by Topographic PV Gradients, *Journal of Physical Oceanography*, 54, 1089–1103, <https://doi.org/10.1175/JPO-D-23-0142.1>, 2024.
- 765 Taylor, G.: Diffusion by continuous movements, *Proceedings of the London Mathematical Society*, 20, 196–212, <https://doi.org/10.1112/plms/s2-20.1.196>, 1921.
- Taylor, G.: Dispersion of soluble matter in solvent flowing slowly through a tube, *Proc. Roy. Soc. A*, 219, 186 – 203, <https://doi.org/10.1098/rspa.1953.0139>, 1953.
- 770 Zhang, W. and Wolfe, C.: Inferring Tracer Diffusivity From Coherent Mesoscale Eddies, *Journal of Advances in Modeling Earth Systems*, 16, 1–25, <https://doi.org/https://doi.org/10.1029/2023MS004004>, 2024.
- Zhurbas, V., Lyzhkov, D., and Kuzmina, N.: Drifter-derived estimates of lateral eddy diffusivity in the World Ocean with emphasis on the Indian Ocean and problems of parameterization, *Deep Sea Res. I*, 83, 1–11, <https://doi.org/10.1016/j.dsr.2013.09.001>, 2014.