

Nonparametric estimation of age-depth models from sedimentological and stratigraphic information

Niklas Hohmann^{1, *}, David De Vleeschouwer², Sietske Batenburg³, and Emilia Jarochowska¹

¹Department of Earth Sciences, Utrecht University, 3584 CB Utrecht, The Netherlands

²Institut für Geologie und Paläontologie, Universität Münster, 48149 Münster, Germany

³University of Barcelona, Barcelona, Spain

Correspondence: Niklas Hohmann (N.H.Hohmann@uu.nl)

Abstract. Age-depth models are fundamental tools used in all geohistorical disciplines. They assign stratigraphic positions to ages (e.g., in drill cores or outcrops), which is necessary to estimate rates of past environmental change and establish timing of events in sedimentary sequences. Methods to estimate age-depth models commonly use ~~simplified~~-parametric assumptions on the uncertainties of ages of tie points. The distribution of time between tie points is estimated using ~~simplistic~~the same assumptions on the formation of the stratigraphic record, ~~for example that sediment accumulates in discrete events that follow a Poisson process. In general, age-depth models are a crude simplification that fail to provide a comprehensive implementation regardless of the depositional environment or time scale studied, although depositional environments are known to differ systematically in their sedimentary dynamics. Integration of all empirical data or expert knowledge (e.g., from sedimentary structures such as erosional surfaces or from basin models) ~~. In other words, many from multiple disciplines remains a challenge~~ for age-depth model inference. Many information sources that can potentially provide geochronologic information remain un- or underused.~~

Here, we present two non-parametric methods to estimate age-depth models from complex sedimentological and stratigraphic data. The methods are complementary as they use different sources of information (sedimentation rates and observed tracer values), are implemented in the ~~admtools~~admtools package for R Software and allow the user to specify any error model and distribution of uncertainties. As use cases of the methods, we

1. construct age-depth models for the Late Devonian Steinbruch Schmidt section in Germany and use it to estimate the timing of the Frasnian-Famennian boundary and the duration of the Upper Kellwasser event.
2. use measurements of extra-terrestrial ³He from ~~ODP site 960 (Maud Rise, Weddell Sea)~~IODP site 1266 (Walvis Ridge) to construct age-depth models for the Paleocene–Eocene thermal maximum (PETM).

The first case study suggests that the Upper Kellwasser event lasted ~~89–92~~85 to 100 kyr (IQR: 84 to 97 kyr) and places the Frasnian-Famennian boundary at 371.834 ± 0.101 Ma (2σ), whereas the second case study provides a duration of ~~41 to 48~~85 to 100 kyr for the PETM recovery interval. These examples show how information from a variety of sedimentological and stratigraphic sources can be combined to estimate age-depth relationships that accurately reflect uncertainties of both available data and expert knowledge.

Age-depth models are a fundamental tool in all geohistorical disciplines where samples can not be dated directly. They assign ages to sampling positions (e.g. stratigraphic heights, core depths), allowing to determine the timing of past events and reconstruct rates of past change. ~~Although age-depth models are rarely the focus of standalone publications, their~~ Their role for the interpretation of historical data becomes obvious when they are revised, often altering our interpretations of past changes. For example, Malmgren et al. (1983) showed that evolution of planktic foraminifera shows short intervals with rapid changes in body size, arguing that this is a common mode of evolution. Revising the age-depth model, MacLeod (1991) showed that the intervals of rapid change coincide with stratigraphic condensation. After accounting for this effect, a random walk was the most likely explanation for the change in body size. This example demonstrates that age-depth models can change our interpretation of how evolution acts on geological time scales fundamentally (Bookstein, 1987; Gould and Eldredge, 1972).

Because of their importance, age-depth models are applied across multiple spatial and temporal scales and environments (terrestrial, marine, lacustrine), ranging from decadal-scale chronologies in modern lakes to the global geological time scale covering hundreds of million of years (Gradstein, 2020; Lacourse and Gajewski, 2020; Cerda et al., 2019). As a result, a wide range of scientific communities engage in the development of age-depth models. This leads to a plethora of available methods to estimate age-depth models, such as `Bchron` (Haslett and Parnell, 2008), `CLAM` (Blaauw, 2010), `Oxcal` ~~(Bronk Ramsey, 2009b, 2008)~~ (Bronk Ramsey, 2009a, 2008), or `Bacon` (Blaauw and Christen, 2011), with variable methodological complexity, ranging from simple interpolation procedures to elaborate Bayesian methods.

Every method to estimate age-depth models makes assumptions on sediment accumulation. For example, the `P_sequence` model in `OxCal` assumes that sediment accumulates in discrete events that follow a Poisson distribution (Bronk Ramsey, 2008), meaning the events are independent of each other and the waiting time between them is exponentially distributed. ~~They~~ argue The authors propose that this is a reasonable assumption for the slow accumulation of individual grains or deposition of regular layers. ~~Similarly, `Bchron` (Haslett and Parnell, 2008) assumes that the number of changes in sedimentation rate (the slope of the~~ The assumptions are commonly carefully chosen and validated for particular environments and time scales. For example `Bacon` samples sedimentation rates from a gamma distribution, an assumption validated for Holocene peat cores, which afford high-precision dating (Blaauw and Christen, 2005; Blaauw et al., 2007). However, different depositional environments have systematically different dynamics of sedimentary accumulation (Sadler, 1981; Enos, 1991; Schumer and Jerolmack, 2006) and the shape and parameters of the distribution of depositional and erosional events, and even their characteristic periodicities, are specific to a given environment and time scale (e.g. tide- versus storm-dominated coastal settings at annual time scales, braided rivers at decadal time scales, carbonate platforms at precession time scales etc.). Not all possible use-cases can be successfully hard-coded into a general model of age-depth ~~model) follows a Poisson distribution. Methods originally developed based on knowledge of specific timescales or environments are often applied outside of that domain, which potentially leads to inference. Applying methods validated for specific environments or time scales outside their intended domain creates a potential risk of~~ violating their assumptions. For example, `Bchron` (originally developed for Quaternary records) and the derived “modified `Bchron`” (Trayler et al., 2019) were used to improve the global Devonian time scale (Harrigan et al., 2021;

De Vleeschouwer and Parnell, 2014), constrain the timing of the Late Paleozoic Ice Age in the Parana Basin (Cagliari et al., 2023), and date saltmarsh sediment to reconstruct late Holocene sea-level changes (Parnell and Gehrels, 2015). ~~However, it~~ ~~It~~ is not clear that in all such cases the assumptions of these age-depth modeling procedures are suitable ~~representations, and~~ ~~therefore applicable,~~ for such a wide range of temporal and spatial scales and depositional environments. ~~These assumptions are~~ ~~often made because of their mathematical convenience, and not because of their empirical realism~~ Our goal in `admtools` is creating models and their user-friendly implementations, in which the method is separated from the assumptions. In other
65 words, it offers the users the possibility to make the same assumptions as other models, but also modify them. We propose that this is an important feature in assessing the uncertainty of the age-depth model. Until now, the uncertainty around the choice of the assumptions could not be quantified. In `admtools`, it is possible to compare the effect of following the assumption, used e.g. in `Bacon`, that the sedimentation rates are drawn from gamma distribution, with an assumptions that they come from the normal or a uniform distribution.

70 The stratigraphic record might be complex, but different sub-disciplines of stratigraphy offer ways to constrain its structure. Astrochronology can provide estimates on accumulation rates by matching proxy records with orbital signals (Meyers, 2019; Li et al., 2018), although this approach is not without pitfalls (Blaauw, 2012) and remains the topic of intensive efforts to improve it (Sinnesael et al., 2019; De Vleeschouwer et al., 2024). Sequence stratigraphy provides qualitative predictions on changes in sedimentation within a sequence. Actualistic studies involving direct observation or dating of young deposits provide
75 a wealth of accumulation and sedimentation rates (Enos, 1991; McNeill, 2005; Sadler, 1981), whereas external estimates of accumulation rates, e.g. from cyclostratigraphy, are used to cross-validate age-depth models or calculate geochemical fluxes (Murphy et al., 2010b; Jarochowska et al., 2020). Forward models allow us to examine the effect of different assumptions about sedimentation, which can be used to constrain biasing effects of stratigraphic architectures (Hohmann et al., 2024). The majority of methods to estimate age-depth models are not able to incorporate complex stratigraphic information into their estimates,
80 ~~or are constrained to simplistic assumptions~~ have specific assumptions embedded in the algorithms (e.g. additional breakpoints in sedimentation rates (Trayler et al., 2023)).

Here, we present two nonparametric methods, FAM (Flux - Assumption Matching) and ICON (Integrated ~~CONDensation~~ condensation), to estimate age-depth models from complex stratigraphic and sedimentological data. ICON estimates age-depth models from arbitrarily complex data on sedimentation rates observed in a stratigraphic column. This knowledge can, for example, be derived
85 from astrochronology (`eCoco` by Li et al. (2018) or `eTimeOpt` by Meyers (2019)), sequence stratigraphic interpretations, ~~or expert knowledge~~ actualistic studies (Enos, 1991; Sadler, 1981; Davies et al., 2019; Tomašových et al., 2022) or theoretical considerations (Tipper, 2016; Aadland et al., 2018). FAM estimates age-depth models and sedimentation rates by comparing observed tracer values (extraterrestrial ~~helium-3~~ ^3He , pollen, or radiogenic tracers such as ^{210}Pb) with assumptions of their fluxes in the time domain. Both methods are nonparametric in the sense that they do not ~~make~~ prescribe specific assumptions
90 on the processes or probability distributions that govern sediment accumulation and the structure of the stratigraphic record. They ~~are assumption explicit in the sense that they assume the law of superposition holds, but all other assumptions must be~~ provided by the user require the users to state these assumptions. This ensures that model assumptions are specified verbatim and clearly communicated, ~~and the estimated.~~ The only assumption embedded in the methods is that the law of the superposition

holds. The uncertainty of the users over the correct choice of assumptions can, therefore, explicitly examined by comparing the results under different assumptions, whereas this uncertainty would have been largely hidden when using methods that have their assumptions hard-coded. As a result, we hope to offer a tool which allows to separate the relative contributions of data uncertainty and assumption uncertainty in age-depth models ~~are data-driven rather than assumption-driven. Both~~. Both FAM and ICON methods are implemented in the R package `admtools` (~~?~~) (Hohmann, 2025), which is available on CRAN (The Comprehensive R Archive Network), and is developed as an open source project on GitHub.

We illustrate the methods using two empirical cases. First, we examine how the propagation of uncertainties of sedimentation rates estimated using cyclostratigraphy influence the duration of the Late Devonian Kellwasser Event and the age of the Frasnian-Famennian boundary (Da Silva et al., 2020). Second, we determine how variable fluxes of extraterrestrial ^3He change the interpretation of PETM recovery time at ~~ODP Site 690 (Farley and Eltgroth, 2003)~~ IODP Site 1266 (Murphy et al., 2010b). These cases were chosen because they are characterized by both environmental upheaval and varying sedimentation rates. Interpreting the environmental upheaval requires good age-depth models, which are challenging to estimate due to the changes in sedimentation rate.

The examples show that the developed methods are able to incorporate complex sedimentological and stratigraphic data into the estimation of age-depth relationships and their uncertainties, resulting in age-depth models with empirically realistic uncertainties.

The target groups of this manuscript are researchers using stratigraphy at any point of their work, especially those investigating processes in the time domain, such as material or element fluxes, the nature and rate of biological evolution, as well as the pacing of environmental change. Secondly, the text is addressed to the community interested in developing and promoting methods of age-depth estimation. The documentation of the `admtools` package contains extensive worked examples and the supplementary material for this article provides literate code, which can serve as a starting point for users to develop their own applications. Thus, to use the methods described here, users are encouraged to run the code along with studying the presented examples.

2 Model development

2.1 Assumptions

We assume that:

1. sediment accumulation is uninterrupted, i.e. is strictly positive (but can be arbitrarily low), and
2. the law of superposition holds, meaning strata found lower in a section are older than those found higher up.

With these assumptions, age-depth models are strictly monotonous functions between the time and stratigraphic domain, where strict monotonicity reflects the law of superposition.

2.2 Preliminaries

- 125 We distinguish between time domain D_T (time dimension, SI units seconds or derived units such as years) and stratigraphic domain D_L (length dimensions, SI units meter). Throughout the manuscript, indices of T (or L) indicate that a function is defined in the time (or stratigraphic) domain. We use time t (increasing towards today) and height h (increasing upwards) for equations, as this ensures that the direction of integration is always from older to younger strata. Using age (increasing away from today) or depth (common when working with drillcores) would prevent that. Technically we work with time-height
- 130 models, but we will still refer to them as age-depth models as the name is well established. Conversions from time to age or height to depth can be made by using time before a reference point, and height below a reference point. Any combination of age or time and depth or height would be correct and is simply a matter of choice of reference frame and scientific community. For example, cores are typically described in terms of depth, whereas sections are commonly measured upwards and expressed in height.
- 135 Let $T : D_L \rightarrow D_T$ be the function that maps a stratigraphic position to its time of deposition, and $H : D_T \rightarrow D_L$ the function that maps a point in time to the stratigraphic position formed at said time. Both T and L provide a description of the age-depth model. By definition, they are inverses of each other:

$$T = H^{-1} \text{ and } H = T^{-1} \quad (1)$$

- Throughout this article, we mean instantaneous sedimentation rate when we speak of sedimentation rates. We distinguish
- 140 between the sedimentation rate in the time domain s_T and the sedimentation rate observed in the stratigraphic domain s_L . The sedimentation rate observed at a stratigraphic position h is the rate with which said position was formed, yielding the following two relations:

$$s_T(T(h)) = s_L(h) \text{ and } s_T(t) = s_L(H(t)) \quad (2)$$

If H is differentiable, we know that

$$145 \quad \frac{dH}{dt} = s_T$$

Due to the law of superposition (strict monotonicity), sedimentation rates in both domains are strictly positive. The amount of sediment accumulated in the time interval $[t_1, t_2]$ is

$$H(t_2) - H(t_1) = \int_{t_1}^{t_2} s_T(x) dx \quad (3)$$

2.3 Estimating age-depth models from sedimentation rates

- 150 Sedimentation rates determine how fast new material accumulates, and how much time is recorded per increment of stratigraphic thickness. Accumulating this information from a reference point can be used to construct age-depth models. Con-

versely, if an age-depth model is given, its slope is the sedimentation rate (Trayler et al., 2023; Hohmann, 2021). Here, we formalize how age-depth models can be constructed from arbitrary sedimentation rates in the stratigraphic domain. We refer to this method as ICON, standing for **I**ntegrated **C**ONDensation for reasons explained below.

155 2.3.1 Model formulation

By the inverse rule, we get

$$\frac{dT}{dh} = \frac{1}{s_L} \quad (4)$$

(see e.g., Hohmann (2021)). Accordingly, the amount of time recorded in the stratigraphic interval $[h_1, h_2]$ is given by

$$T(h_2) - T(h_1) = \int_{h_1}^{h_2} \frac{1}{s_L(x)} dx \quad (5)$$

160 We refer to the inverted sedimentation rate in the stratigraphic domain

$$c(h) := \frac{1}{s_L(h)} \quad (6)$$

as (stratigraphic or sedimentary) *condensation*, which is used here as time recorded in sediment and does not imply low accumulation rates. Using condensation instead of sedimentation rate in the stratigraphic domain has two advantages. First, it has the correct dimension and units (time per length, years per meter) to represent the amount of time represented in the rock record. This allows to directly determine the amount of time represented in a section by integrating over condensation. Second, it reduces the ambiguity that comes with dealing sedimentation rates in both in the time and stratigraphic domain. While sedimentation rates in the time domain can in general be zero or negative under sedimentary stasis or erosion, condensation must always be positive as we can only observe intervals with net positive sediment accumulation in the rock record.

Given two tie points (t_i, h_i) , (t_{i+1}, h_{i+1}) and condensation $c(h)$, define the dimensionless normalization constant

$$170 \quad C_i := \frac{t_{i+1} - t_i}{\int_{h_i}^{h_{i+1}} c(x) dx} \quad (7)$$

and introduce the *tie-point corrected condensation*:

$$\hat{c}^i(h) = c(h) \cdot C_i \quad (8)$$

This correction ensures that between tie points, the time represented by condensation matches the time elapsed between the tie points. Then the age-depth model is given by

$$175 \quad T(h) = t_i + \int_{h_i}^h \hat{c}^i(x) dx \quad \text{for } h \in [h_i, h_{i+1}] \quad (9)$$

This is simply adding the time elapsed between h_i and h to the known time at the lower tie point. Specifically, $T(h_i) = t_i$ and $T(h_{i+1}) = t_{i+1}$. For multiple tie points, the age-depth model is given by the closed expression

$$T(h) = \sum_i \mathbf{1}_{H_i} \cdot \left(t_i + \int_{h_i}^h \hat{c}^i(x) dx \right) \quad (10)$$

where $\mathbf{1}_A$ is the indicator function on the set A and $H_i = (h_i, h_{i+1}]$ is the i -th stratigraphic interval. For this expression to
 180 be valid below and above the highest tie point, two minor adjustments are necessary. First, below the lowest tie point (t_0, h_0) , the direction of integration needs to be reversed to make sure the integral represents a positive amount of time, leading to the expression

$$T(h) = t_0 - \int_h^{h_0} \hat{c}^i(x) dx \quad (11)$$

Second, above and below the highest tie point, normalization is not necessary as there is no possible mismatch between
 185 the time represented by the sedimentation rate and the time elapsed between tie points (because there are no two tie points). Skipping normalization is achieved by setting the normalization constants in these intervals to 1. This means between tie points, information on sedimentation rates or condensation is adjusted to be ~~congruent~~ congruent with the timing of the tie points and only contributes information about the relative distribution of time within the section, while it is taken at face value below/above the lowest/highest tie point.

190 By the monotonicity of the integral, T as defined above is monotonous, and H is uniquely defined. With this, the sedimentation rate in the time domain can be determined using

$$s_T(t) = s_L(H(t)) \quad (12)$$

2.4 Estimating age-depth models from tracer values

~~Assume there is~~ Given a constant influx of a tracer into the sediment with time. ~~Then,~~ observing elevated tracer values in a
 195 section indicates stratigraphic condensation and low sedimentation rates, while reduced tracer values in the section indicate stratigraphic dilution and high sedimentation rates. Comparing tracer values with assumptions on their flux can thus be used to constrain the time preserved in the stratigraphic record, and construct age-depth models. An early application using pollen flux in a peat bog ecosystem has been introduced by Middelorp (1982) and was validated by Young et al. (1999), and extended to

deep-time strata by Jarochowska et al. (2020), who used three independent constant flux tracers to construct relative age-depth models for the late Silurian Lau Carbon Isotope Excursion to correct rates of redox proxy- and isotope changes for increasing sedimentation rates in a shallowing-upward succession in Gotland, Sweden. Using extraterrestrial ^3He as tracer, this approach has ~~for example,~~ for example, been employed to constrain the timing of the Paleocene-Eocene Thermal Maximum (PETM) (~~Farley and Eltgroth, 2003~~) (Farley and Eltgroth, 2003; Murphy et al., 2010b) and the Cretaceous-Paleogene (K-Pg) boundary extinction (Mukhopadhyay et al., 2001), as well as to examine small-scale fluctuations of sedimentation rate in limestone marl alternations (Blard et al., 2023). ~~Jarochowska et al. (2020) used three independent constant flux tracers to construct relative age-depth models for the late Silurian Lau Carbon Isotope Excursion to correct rates of redox proxy- and isotope changes for increasing sedimentation rates in a shallowing-upward succession in Gotland, Sweden.~~ Similarly, Appleby and Oldfield (1978) compared observed ^{210}Pb values in cores with ^{210}Pb concentrations predicted from constant flux and exponential decay to estimate age-depth relationships in young sediments, an approach termed the CRS model (Abril-Hernández, 2023). What unifies these approaches is that they compare the assumed tracer flux into the sediment in the time domain with tracer values observed in the stratigraphic domain to construct age models.

Here, we ~~provide the~~ extend these previous methods by providing a general mathematical framework for the construction of age-depth relationships from comparisons of arbitrarily complex tracer fluxes with observed tracer values in a section. We refer to this method as FAM, standing for **F**lux **A**ssumption **M**atching.

2.4.1 Model formulation

~~Assume we have have~~ Let f_L be observations of a tracer f_L in the stratigraphic domain (dimension X/L , where X is the unit in which the tracer is measured) ~~and some knowledge on tracer fluxes with time,~~ \tilde{f}_T be tracer flux with time (dimensions X/T). ~~Let,~~ and (t_i, h_i) and (t_{i+1}, h_{i+1}) be two tie points. Define the dimensionless constant

$$C_i = \frac{\int_{h_i}^{h_{i+1}} f_L(x) dx}{\int_{t_i}^{t_{i+1}} \tilde{f}_T(x) dx} \quad (13)$$

and define the empirically calibrated assumption on tracer flux in the time domain

$$f_T^i := C_i \tilde{f}_T \quad (14)$$

This normalization ensures the amount of tracer assumed to be embedded in the sediment between t_i and t_{i+1} matches the amount of tracer observed between h_i and h_{i+1} in the section.

In the absence of taphonomic and diagenetic effects, the total volume of tracer observed in a stratigraphic interval I_L is identical to the total volume of tracer incorporated into the sediment over to time interval I_T during which I_L was formed (see Hohmann (2021)). Based on this, we know that

$$\int_{t_i}^t f_T^i(x) dx = \int_{h_i}^h f_L(x) dx \quad (15)$$

holds, because tracer volume placed in the sediment over $[t_i, t]$ is identical to tracer volume observed in $[h_i, h]$. The age-depth model is given by all t and h for which this relationship holds (the graph of the relation). Solving the above equation for
230 t (or h) yields a representation of T (or H). Introducing the short notations

$$\Theta_{t_i}(t) := \int_{t_i}^t f_T(x) dx \quad \text{and} \quad \Lambda_{h_i}(h) := \int_{h_i}^h f_L(x) dx \quad (16)$$

T can be written explicitly as

$$T(h) = \Theta_{t_i}^{-1} \circ \Lambda_{h_i}(h) \quad (17)$$

The age-depth model in the presence of multiple tie points can be generated by stitching together multiple stratigraphic
235 intervals, leading to the representation

$$T(h) = \sum_i \mathbf{1}_{H_i} (\Theta_{t_i}^{-1} \circ \Lambda_{h_i}(h)) \quad (18)$$

where $H_i = (h_i, h_{i+1}]$ is the i -th stratigraphic interval. To expand this approach to heights above/below the highest/lowest tie point, some minor adjustments are required. First, below the lowest tie point the direction of integration is reversed to ensure the integrals return positive tracer volumes. Second, above/below the highest/lowest tie point, the normalization coefficient
240 C_i is not necessary, as there can not be a mismatch between observed and assumed tracer fluxes between tie points because there is no second tie point which could generate such a mismatch. Computationally, this is solved by setting the normalization coefficient to 1 above or below the highest or lowest tie point, respectively.

There is one relevant edge case where normalization above or below the highest tie point (t_N, h_N) or lowest tie point (t_0, h_0) is required. This is when the integrals over both \tilde{f}_T and f_L on the unbounded intervals (h_N, ∞) and (t_N, ∞) (resp. $(-\infty, h_0)$ and $(-\infty, t_0)$) are finite. This is for example the case when estimating age-depth relationships based on radiogenic tracers
245 such as ^{210}Pb . Here, the observed tracer values drop to 0 down core due to exponential decay. The assumed tracer flux in the time domain can vary, but will eventually drop to 0 because of the exponential decay of the tracer, leading to an f_L that is defined on an unbounded interval, but with a finite tracer volume (Abril-Hernández, 2023).

2.4.2 Estimating sedimentation rates

250 Based on the representation of T , the inverse function and the composition rule, we get

$$s_L(h) = \frac{f_T(T(h))}{f_L(h)} \quad (19)$$

for the instantaneous sedimentation rates observable in the stratigraphic domain, and

$$s_T(t) = \frac{f_T(t)}{f_L(H(t))} \quad (20)$$

for the instantaneous sedimentation rate observable in the time domain. Sedimentation rates are given by the ratio of assumed
 255 tracer flux to observed tracer flux. For constant assumed tracer flux, this provides a mathematical equivalent to the intuition
 that elevated observed tracer values indicate low sedimentation rates, while reduced observed tracer values correspond to high
 sedimentation rates.

2.4.3 Special cases

In general, the equations arising from FAM need to be solved numerically by combining integration with root-finding proce-
 260 dures. Here, we give two examples where the age-depth relationships from FAM can be written as analytical expressions, and
 use these examples to demonstrate how FAM generalizes existing methodology for estimating age-depth relationships.

2.4.4 Constant tracer flux in the time domain: the cFAM model

Constant tracer fluxes have occasionally be used to estimate sedimentation rates and age models ([Mukhopadhyay et al., 2001](#), [Farley and El](#)
[Farley and Eltgroth, 2003](#); [Murphy et al., 2010b](#); [Jarochowska et al., 2020](#); [Blard et al., 2023](#); [Mukhopadhyay et al., 2001](#)), but
 265 this is not a well-established approach. Here, we derive closed expressions for age-depth models and sedimentation rates de-
 rived under the assumption of a constant tracer flux, a method we refer to as cFAM (constant Flux Assumption Matching).
 Constant tracer influx in the time domain implies $\tilde{f}_T(t) = c$ for all t and some flux value c . Between two tie points (t_i, h_i) ,
 (t_{i+1}, h_{i+1}) , we get

$$C_i = \frac{\int_{h_i}^{h_{i+1}} f_L(x) dx}{c (t_{i+1} - t_i)} \quad (21)$$

270 for the normalization constant, so the empirically calibrated assumption on tracer flux is

$$f_T(t) = \frac{\int_{h_i}^{h_{i+1}} f_L(x) dx}{t_{i+1} - t_i} \quad (22)$$

As a result, the age model between the tie points under cFAM is

$$T(h) = t_i + \frac{(t_{i+1} - t_i) \int_{h_i}^h f_L(x) dx}{\int_{h_i}^{h_{i+1}} f_L(x) dx} \quad (23)$$

Note here that the value of the flux c does not appear in the equation, as it is cancelled out by the normalization. Only relative
 275 changes in observed tracer values contribute to the age-depth model between tie points. Above the highest tie point (t_N, h_N)
 and below the lowest tie point (t_0, h_0) , we get

$$T(h) = t_N + \frac{1}{c} \int_{h_N}^h f_L(x) dx \quad \text{and} \quad T(h) = t_0 - \frac{1}{c} \int_h^{h_0} f_L(x) dx \quad (24)$$

For the sedimentation rates, note that f_T is independent of t , so we get for the sedimentation rate in the stratigraphic domain

$$s_L(h) = \frac{\int_{h_i}^{h_{i+1}} f_L(x) dx}{(t_{i+1} - t_i) f_L(h)} \quad (25)$$

280 between tie points and

$$s_L(h) = \frac{c}{f_L(h)} \quad (26)$$

above/below the highest tie point.

Jarochowska et al. (2020) used an auxiliary time scale where $t = 0$ corresponds to the bottom of the section and $t = 1$
 corresponds to the top of the section, allowing them to correct rates for variations in sedimentation rates in the absence of
 285 absolute age constraints. This is equivalent to introducing two artificial tie points $(0, h_0)$ and $(1, h_1)$, where h_0 and h_1 are the
 bottom and the top of the section, respectively. In this case, cFAM reduces to cFAM-at (where the at stands for **auxiliary time**).
 In this case, the age model is given by

$$T(h) = \frac{\int_{h_0}^h f_L(x) dx}{\int_{h_0}^{h_1} f_L(x) dx} \quad (27)$$

and sedimentation rates by

$$290 \quad s_L(h) = \frac{\int_{h_i}^{h_{i+1}} f_L(x) dx}{f_L(h)} \quad (28)$$

Note that both expressions are solely dependent on empirical data measured in the section.

2.4.5 Radiogenic tracers

Here, we show that when FAM is combined with the assumption that tracer flux follows an exponential decay, it reduces to the
 CRS (constant rate of supply) model by Appleby and Oldfield (1978) for dating sediments using ^{210}Pb , a common dating tool

295 for recent (100 to 150 years) aquatic sediments. We use the derived expressions for sedimentation rate to derive an estimator of instantaneous sedimentation rates in the stratigraphic domain for the CRS model.

For this section, we use age a and depth d instead of time t and height h , as they are more fitting to the context of dating recent core material. Assuming the tracer decays exponentially with time, we get

$$\tilde{f}_T(a) = c \exp(-\lambda a) \quad (29)$$

300 for tracer flux in the time domain, here representing the amount of preserved tracer of age a . Let $f_L(d)$ be the tracer content at depth. Then the normalization constant is

$$C = \lambda \int_0^{\infty} f_L(x) dx \quad (30)$$

and we get

$$f_T(a) = f_T(a) \lambda \int_0^{\infty} f_L(x) dx \quad (31)$$

305 for the empirically calibrated tracer flux in the time domain. Note here that tracer fluxes were normalized, although they are defined on an unbounded interval. Then the age-depth relationship is the solution to the equation

$$\int_0^a \exp(-\lambda x) dx = \frac{\int_0^d f_L(x) dx}{\lambda \int_0^{\infty} f_L(x) dx} \quad (32)$$

Solving for a , we get

$$A(d) = -\frac{1}{\lambda} \ln \left(1 - \frac{\int_0^d f_L(x) dx}{\int_0^{\infty} f_L(x) dx} \right) \quad (33)$$

310 for the age of the core as a function of depth, matching the formulations of the CRS model given by Appleby and Oldfield (1978) and Abril-Hernández (2023). The sedimentation rate at depth d can then directly be estimated via

$$s_L(d) = \frac{\lambda \int_0^{\infty} f_T(x) dx \left(1 - \frac{\int_0^d f_L(x) dx}{\int_0^{\infty} f_L(x) dx} \right)}{f_L(d)} \quad (34)$$

Note that FAM allows to incorporate arbitrary assumptions on how tracer flux changes with time into the estimation of a ^{210}Pb chronology, allowing to incorporate expert knowledge on fluctuations in ^{210}Pb flux. While there is an analytical solution for the CRS model available, the integral equations defining the more general cases need to be solved numerically.

2.5 Randomization

The estimation of age-depth models as described above is purely deterministic. Here, we show that it expands to the probabilistic estimation of age-depth models without additional assumptions on the nature of the underlying probability distributions.

For ICON, we first assume tie points are deterministic and in strict temporal and stratigraphic order ($t_i < t_{i+1}$ and $h_i < h_{i+1}$), and consider the sedimentation rate a stochastic process. Based on the law of superposition, the condensation is a strictly positive stochastic process, which we assume to be regular enough to be integrated. With this, all involved integrals are well-defined, the normalization factor is not zero, and T is a strictly increasing stochastic process. Given the strict order holds almost surely, this construction remains valid when the tie points are randomized. Identical arguments hold for FAM: When observed and assumed tracer fluxes are almost surely positive and integrable, Θ and Λ are both strictly increasing, and the arising integral equations can be solved uniquely. Summarizing, both the construction of ICON and FAM immediately expand from the deterministic to the probabilistic case without any assumptions on the involved probability distributions.

2.6 Implementation

Both estimation procedures for age-depth models are implemented in the R package `admtools` ([Hohmann, 2025](#)). The implementation uses the R internal procedure `integrate` to numerically determine the integrals arising in FAM and ICON, and `uniroot` to solve the function inversions in FAM ([Brent, 2002](#)), and parameters determining their numeric performance can be passed to the relevant procedures. The two methods are implemented in the functions `sed_rate_to_multiadm` (ICON) and `strat_cont_to_multiadm` (FAM). Both produce an object of type `multiadm`, representing a collection of age-depth models (sample paths of T), each of which is a possible scenario for the examined section. These objects can be reused, e.g. for plotting, transforming data, determining uncertainties of stratigraphic positions or timing, or for transformation of other objects (e.g., time series, [phylogenetic trees](#)) between the stratigraphic and the time domain.

Timing and positions of tie points can follow arbitrary probability distributions as long as they are strictly ordered. It is the user's responsibility to ensure strict order. In the most general case, tie points are coded as functions that take no inputs and, upon each evaluation, return one sample drawn from the distribution of tie point times/heights. Effectively, these functions are user-defined random sample generators. While they are not dependent on any input parameters, they can wrap around complex empirical data that determines uncertainties. Details on how these functions are coded can be found in the package vignettes (long form documentation with worked examples) or the [project-package](#) webpage. To simplify the definition of tie points, wrappers for common use cases for tie points are provided (uniform or normally distributed tie points or deterministic tie points). Note that not only the timing, but also the stratigraphic position of tie points can be randomized (e.g., when the age information is associated with a bed, or its stratigraphic position was not recorded precisely).

The stochastic processes representing sedimentation rates or tracer fluxes can be passed to the estimation procedures as function factories that take no arguments. A function factory is a function that itself returns a function. The function factory represents the stochastic process. Each time it is evaluated, it returns a function that represents a sample path. This function can be evaluated at specific points to return the sedimentation rates/tracer values at said points. In our case, the function factories

can be thought of as very complex random number generators. Instead of returning a random number or vector of random numbers, they return a random function, effectively making them infinite-dimensional random number generators. Similar to the way tie points are coded, they can depend on user data without having to pass said data to the estimation procedures. To simplify the definition of function factories, wrappers for the most common use cases are provided. For tracer fluxes, these includes cases for constant, linear, and quadratic fluxes, empirical measurements of tracer means and standard deviations (see example on the PETM). For sedimentation rates, options to ~~turn arrays of sedimentation rate estimates are provided, including~~ ways construct sedimentation rates from upper and lower bounds, the gamma distribution, and from arrays are provided. This includes functions to directly take outputs from the `eTimeOpt` function from the `astrochron` package and turn them into a function factory (Meyers, 2014, Meyers (2019)).

Computation times vary, but are typically below one minute (see examples below). Computation time is dependent on the irregularity of the input functions (sedimentation rates, tracer fluxes), which determines how fast the numeric integration is. Functions with rapidly changing function values are difficult to integrate numerically, with discontinuous functions being the most challenging ones. For fixed input functions, computation time scales linearly with the number of stratigraphic positions where the age-depth models are determined, and the number of age-depth models estimated within one `multiadm` object.

The package implements unit tests for both the estimation procedures and the underlying logic of the age-depth transformation that are run each time a part of the code is changed. Systematic testing of code is considered best practice in software development, and improves the quality of scientific software (Hunter-Zinck et al., 2021, Nanthamornphong and Carver (2017)). Unit tests for FAM and ICON test edge cases where analytical solutions are known in advance (one, two, or multiple tie points, constant sedimentation rate, constant tracer fluxes, etc.). Multiple vignettes (short form articles that provide a more complex use case) for both procedures are available after installation via the command `browseVignettes("admttools")`, and can also be browsed on the packages webpage (<https://mindthegap-erc.github.io/admttools/>)

Code adheres to the FAIR4RS (FAIR for research software) principles (Barker et al., 2022), which are based on the FAIR (Findable, Accessible, Interoperable and Reusable) principles for scientific data management and stewardship (Wilkinson et al., 2016). Code development is performed on Github. Each minor release (based on semantic versioning) is published on CRAN (the Comprehensive R Archive Network), assigned a doi and archived on Zenodo.

The package is fully open source, all code can be inspected on GitHub. Contributors are invited to make enhancement request, improve documentation and bug reports, contribute code, and improve integration with the existing geoscientific software landscape. Contributing guidelines for the package are specified in the `CONTRIBUTING.md` file in the root of the directory.

3 Examples

We apply the newly developed methods to two existing studies, the Late Devonian Mass Extinction and the Paleocene-Eocene Thermal maximum. We use these examples to examine how the ability to incorporate added uncertainty changes age estimates.

For details on computational reproducibility, see the README file in [Hohmann \(2024\)](#) [Hohmann and Jarochowska \(2025\)](#). All analyses were performed with `admtools` version ~~0.3.1(?)~~ [0.5.0 \(Hohmann, 2025\)](#).

3.1 Duration and timing of the Late Devonian mass extinction

The Late Devonian mass extinction at the Frasnian-Famennian boundary is considered one of the “big five” Phanerozoic mass extinctions (Muscente et al., 2018; Raup and Sepkoski, 1982), and a multitude of causes have been discussed, including climate warming or cooling (Thompson and Newton, 1988), volcanism (Racki et al., 2018), extraterrestrial impacts (Claeys et al., 1992), or changes in the weathering cycle (Averbuch et al., 2005). In many locations, the sedimentologic expression of this perturbation is expressed by two dark, organic-rich lithologies referred to as the Upper and the Lower Kellwasser beds, each being associated with an anoxic event called the Upper and the Lower Kellwasser Event (Carmichael et al., 2019).

Da Silva et al. (2020) combined ~~eyelostratigraphic~~ [cyclostratigraphic](#) methods with U-Pb dates from Percival et al. (2018) to constrain the absolute timing and duration of the Late Devonian Kellwasser events and the Frasnian-Famennian boundary in the Steinbruch Schmidt section, Germany. For the cyclostratigraphic analysis (code and data available in da Silva (2024)), ~~they~~ [the authors](#) used the `eTimeOpt` function from the `astrochron` package for R Software (Meyers, 2014, 2019; R Core Team, 2023). `eTimeOpt` uses a moving window approach to find sedimentation rates that lead to the highest concentration of power of the precession and eccentricity frequencies and best expression of short eccentricity or precession amplitude modulation in the analyzed proxy record. The method returns r_{opt}^2 , a measure of fit between the proxy signal and the predicted patterns for a range of heights and sedimentation rates. Da Silva et al. (2020) used the `eTimeOptTrack` function of the `astrochron` package to extract sedimentation rate estimates from `eTimeOpt` results, yielding the sedimentation rates at which r_{opt}^2 is maximal. These deterministic sedimentation rates change with stratigraphic position, and were used to constrain the duration of the anoxic events and the timing of the Frasnian-Famennian boundary. However, simply using the sedimentation rate with the best fit (highest r_{opt}^2) neglects uncertainties in the estimation of sedimentation rates, as a wide range of sedimentation rates can potentially provide a good fit to a given signal (Figure 1). Here, we use ICON to examine how the propagation of uncertainties of sedimentation rates changes the duration and timing estimates for the Kellwasser event and the Frasnian-Famennian boundary.

Da Silva et al. (2020) analyzed magnetic susceptibility (MS), log Ti concentration, and $\delta^{13}C$ values, with similar results for all three proxies. For this example, we focus solely on MS. Data preprocessing and `eTimeOpt` analysis was kept identical to da Silva (2024) for comparability, see Da Silva et al. (2020) for details. Here, we show the results of `eTimeOpt` testing for precession amplitude modulation. Results for short eccentricity amplitude modulation are similar and shown in the Appendix (Figures 3, A1, A2). `eTimeOpt` results for r_{opt}^2 were extracted using the `get_data_from_eTimeOpt` function of the `admtools` package. Interpreting `eTimeOpt` results probabilistically has two challenges:

1. At fixed heights, it provides r_{opt}^2 values as a function of sedimentation rate. However, it is unclear how r_{opt}^2 values can be meaningfully translated into a probability that the sedimentation rate falls within a certain interval.




Figure 1. r_{opt}^2 values from eTimeOpt testing for precession amplitude modulation at 3.54 m and 1.56 m (A) and throughout the whole section (B) in the Steinbruch Schmidt section. The heights shown in A roughly corresponding to the locations of the Upper Kellwasser bed and the location of the dated ash bed.

2. The correlation structure of sedimentation rates is unclear: Given we know sedimentation rates at one point in the section, how would that influence our estimates further up or down section?

415 To address this, we introduce the following two ~~assumption~~assumptions into this example. First, the sedimentation rates are determined at random heights following a Poisson process with rate λ , an assumption borrowed from BChron (Haslett and Parnell, 2008). This means the number of height points follows a Poisson distribution with an expected value of $\lambda \cdot l$ (where l is the ~~length~~length of the section), and the locations of the points are independent and identically distributed according to a uniform distribution. Second, at each stratigraphic position, sedimentation rates follow a distribution with probability density
420 function proportional to r_{opt}^2 . As a result, sedimentation rates with a higher r_{opt}^2 are more probable than those with a low r_{opt}^2 . If a range of sedimentation rates has comparable r_{opt}^2 values, they are all equally probable, alleviating the problem with eTimeOptTrack that only the sedimentation rate with the highest r_{opt}^2 is selected (Figure 1). Third, between the selected points, sedimentation rate changes linearly. These assumptions are incorporated by the `sed_rate_from_matrix` function, which takes the outputs of the `get_data_from_eTimeOpt` function and returns a sedimentation rate factory as required
425 by ICON.

To ~~distinguish-between~~illustrate the contributions of uncertainties introduced by the U-Pb date and those from eTimeOpt's estimated sedimentation rates, we ~~construct two~~decompose the construction of the age-depth ~~models: One floating model into two steps. First, an~~age-depth model that measures time relative to the ~~dated bentonite layer~~bentonite layer as the fixed single




Figure 2. Floating age-depth model Age-depth models with (A) and anchored age-depth model without (B) incorporation the uncertainty of the radiometric dating for the Steinbruch Schmidt section based on eTimeOpt results testing for precession amplitude modulation. In the floating age-depth model (A), time is measured relative to the bentonite layer dated by Percival et al. (2018), and all uncertainties in the age-depth model arise from the uncertainties of the sedimentation rates estimated using eTimeOpt. The anchored age-depth model is constructed by combining, while in (B), they are a combination of the absolute age uncertainty of Percival et al. (2018) for the bentonite layer with the radiometric dates from Percival et al. (2018) (dashed lines) and sedimentation rates estimated from eTimeOpt. Uncertainties result from both the uncertainties Dashed lines in U-Pb dating and estimates (B) indicate the age uncertainty of sedimentation rates the section when the radiometric uncertainty is propagated using eTimeOpt the median sedimentation rate.

tie-point with known age (Figure 2 A), and one anchored in absolute time, where the bentonite layer is assigned the age and accounts only for the uncertainty resulting from the sedimentation rate is constructed. Secondly, the age and uncertainty of the bentonite layer as determined by Percival et al. (2018) (372.36 ± 0.053 Ma (2σ), considering only measurement uncertainties) is added to the estimation procedure (Figure 2 B). For this example, we choose a rate parameter $\lambda = 3$ for the sedimentation rate, and use 1000 Monte Carlo samples to generate age-depth models, and report results rounded to the next kyr.

3.1.1 Results

The floating time scale displays that age uncertainty increases Both age-depth models show in increase in uncertainty away from the tie point due to variations in sedimentation rate (Figure 2A), resulting in the distinct sausage-shape described by De Vleeschouwer and Parnell (2014). Median time contained in the 2.5 m between the bentonite layer and the Frasnian-Famennian boundary is 513 kyr with an interquartile range (IQR) of 56–522 kyr with a 95 % highest density interval (HDI) of




Figure 3. Durations of the Frasnian-Famennian boundary (A), Upper Kellwasser Event (B), and time elapsed between the Kellwasser Events (C) based on testing for short eccentricity modulation and precession amplitude modulation.

159 kyr and a standard deviation (SD) of 43 kyr based on 39 kyr derived from the uncertainty of the sedimentation rate (Figure 3). For the anchored age-depth model incorporating the radiometric uncertainty, the 95 % envelope is almost parallel (Figure 2 B). This is because the uncertainties arising from the U-Pb date and the sedimentation rate estimates are independent and, as a result, sub-additive: It is unlikely that both yield too low (resp. high) values simultaneously, so their combined uncertainty is lower than the sum of their uncertainties. This indicates that over the observed, stratigraphically short interval, the error introduced by uncertainty in sedimentation rates is negligible-small relative to the uncertainty of absolute dates. Note that the uncertainty of sedimentation rates is limited by the maximum and minimum sedimentation rates passed to eTimeOpt and the averaging within the sliding window, therefore by the window size. In our case the sedimentation rates were constrained a priori to the frequency distribution of sedimentation rates (piecewise for the windows used by eTimeOpt) derived from eTimeOpt based on its (the astrochron package's) r^2_{opt} metric. This frequency distribution was obtained for the interval from 0.1 to 0.6 cm/kyr as in, as constrained by Da Silva et al. (2020). If a wider range of sedimentation rates were analyzed, sedimentation rates would contribute to more uncertainty to the age-depth model. The range is, in practice, constrained to reduce calculation time (Meyers, 2019). Running the eTimeOpt procedure for a broader range of sedimentation rates is possible - although more computationally intensive - but too broad a range may lead to false positives in terms of solutions with the best r^2_{opt} values, therefore it is recommended to constrain the sedimentation rate using external knowledge, such as an independent chronometer or sedimentological information.

For the Frasnian-Famennian boundary, our ~~anchored~~-age-depth model yields a median age of 371.837 Ma ~~, with an IQR of 67 kyr (1st and 3rd quartile is 371.803 and 371.867 Ma, respectively)~~ with a 95% HDI of 201 kyr based on uncertainty from sedimentation rate and radiometric dates. The age distribution is approximately normal, resulting in an age estimate of 371.834 ± 0.101 Myr in the standard ~~2 σ~~ -2 σ representation (Figure 3). This is 36 kyr older than the age estimated by Da Silva et al. (2020), with age uncertainty reduced by 6 % (108 kyr vs. 101 kyr). Our uncertainty is lower than that listed by Da Silva et al. (2020), as they use multiple proxies, estimate one deterministic sedimentation rate per proxy, and combine these estimates to arrive at their final uncertainty. We use a single proxy (MS) to estimate uncertain sedimentation rates, and propagate these uncertainties into the age estimate. Combining uncertain sedimentation rate estimates from multiple proxies would most likely give uncertainties comparable or larger than 108 kyr, as adding sources of uncertainties can only increase the uncertainty of the final estimate. Becker et al. (2020) in Gradstein (2020) lists an age of 371.1 Myr ± 1.1 Myr (~~2 σ~~ -2 σ) for the Frasnian-Famennian boundary. Both our and Da Silva et al. (2020) mean age estimates are elevated compared to this (by 734 and 770 kyr, respectively), but with reduced uncertainties (108 and 101 kyr, respectively) relative to the global geological time scale.

Median-Mean duration of the Upper Kellwasser Event, stratigraphically expressed as black shale intervals, is ~~92-90~~ kyr (1st and 3rd quartile: 84 to 97 kyr, IQR 13 kyr) and is positively skewed, matching the duration estimate of approx. 90 kyr by Da Silva et al. (2020). The median time elapsed between the Kellwasser events, measured from the top of the Lower Kellwasser Event to the bottom of the Upper Kellwasser Event, is 513 kyr (1st and 3rd quartile: 489 and 542 kyr, IQR: 54 kyr), and is also positively skewed (Figure 3). Note that duration estimates are identical whether they are derived from ~~the floating or the anchored~~-age-depth ~~model~~ models that incorporate radiometric uncertainty or not, and depend solely on sedimentation rates. This is because the uncertainties of the tie point cancel out when calculating durations, allowing us to obtain duration estimates with uncertainties below those of the U-Pb dates by Percival et al. (2018). This shows that even in the absence of absolute ages, ~~floating age-depth models are~~ information on sedimentation rates can be a powerful tool to determine durations and relative timing of events in deep time.

Neither the duration of the Lower Kellwasser Event ~~not~~-nor the time from the onset of the Lower Kellwasser Event to the onset of the Upper Kellwasser event could be estimated with the applied approach. This is because the moving window approach in eTimeOpt does not provide sedimentation rate estimates for the lowest and highest parts of the section, of which the former contains the onset of the Lower Kellwasser Event. These durations could be estimated by using a narrower window in eTimeOpt, however this would make the results not comparable directly to those of Da Silva et al. (2020).

Wichern et al. (2024) provide an estimated duration of the Lower Kellwasser Event of approx. 250 kyr at the nearby Winsenberg section. Combined with our estimates of ~~514-513~~ kyr between end of the Lower Kellwasser Event and onset of the Upper Kellwasser Event, the estimated time between the onset of both events is approximately ~~764-763~~ kyr, and the total duration of the Kellwasser crisis is approximately ~~853-852~~ kyr, slightly shorter than that estimated by Wichern et al. (2024).

3.1.2 Robustness of Age-Depth Models for the Paleocene-Eocene Thermal maximum

3.2 Robustness of Age-Depth Models for the Paleocene-Eocene Thermal maximum

The Paleocene-Eocene Thermal Maximum (PETM) is a short interval of global carbon cycle perturbation associated with climate warming approximately 56 Myr ago (Vahlenkamp et al., 2020; Sluijs et al., 2007). Multiple causes have been proposed, including organic matter oxidation, enhanced volcanism, dissociation of gas hydrates, and methane release from vent systems (Dickens et al., 1995; Kurtz et al., 2003; Frieling et al., 2016; Storey et al., 2007). The PETM is a potential geological analogue for anthropogenic climate change (Carmichael et al., 2017; Haywood et al., 2011), making it crucial to understand the timing of its onset and recovery.

The Establishing the duration and timing of the PETM has been an important stimulus in improving age-depth modeling methods, because of the changes in sedimentation rates affecting the stratigraphic tie-points associated with the event. Many estimates rely on Ocean Drilling Program (ODP) Site 690 contains both onset, peak, and recovery intervals of the PETM, making it a prime candidate to examine timing and pacing of the event. Multiple disparate sites from the South Atlantic and exploit combinations of cyclostratigraphy and extraterrestrial ^3He flux coupled with characteristic changes in the $\delta^{13}\text{C}$ record. The compilations of differences in timing of the PETM between studies by Sluijs et al. (2007) and Murphy et al. (2010b) are representative of the sensitivity of the age-depth models for this site have been proposed. Röhl et al. (2000), Norris and Röhl (1999) and Röhl et al. (2007) used cyclostratigraphy and correlation to other sites to determine age constraints, whereas Farley and Eltgroth (2003) to their assumptions.

Here we attempt to replicate the analysis by Murphy et al. (2010b) based on the IODP Site 1266 located at the Walvis Ridge. The core contains the entire PETM interval and has been among the key sections used to resolve the timing and pacing of the event. Murphy et al. (2010b) used measurements of extraterrestrial ^3He - ^3He coupled with the assumption of a constant extraterrestrial ^3He - ^3He flux into the sediment to estimate sedimentation rates and an age-depth model. While Although the variability of ^3He - ^3He fluxes over geologically short timescales is low, it is known that the ^3He - ^3He flux can vary by about an order of magnitude over geologically long time scales (Farley, 2001; Takayanagi and Ozima (1987)). Here, we use FAM (Takayanagi and Ozima, 1987; Farley, 2001; Sluijs et al., 2007). We use this study to, first, validate the FAM model and, second, to examine how robust the age-depth model for the PETM proposed by Farley and Eltgroth (2003) is to variability in ^3He at Site 1266 from Murphy et al. (2010b) is to short-term variability in ^3He fluxes.

As stratigraphic points of reference, we use the PETM pre-, main-core and recovery intervals as defined by Farley and Eltgroth (2003) Röhl et al. (2007), and use the raw data published in Farley and Eltgroth (2003) by Murphy et al. (2010b) (Murphy et al., 2010a). We measure time relative to the beginning of the main interval to construct a floating age-depth model for the section, matching the time scale shown in Fig. 4 of Farley and Eltgroth (2003) 1 of Murphy et al. (2010b). While this does not give an absolute age of the PETM, it provides absolute durations of the PETM intervals defined above. For the constant flux value, we use the value of $0.69 \text{ pcc} (1 \text{ pcc} = 10^{-12} \text{ cm}^3) 0.48 \text{ pcc}$, where $1 \text{ pcc} = 10^{-12} \text{ cm}^3$ of He at STP (standard temperature and pressure) per cm^2 per kyr calculated by Farley and Eltgroth (2003) per cm^2 per kyr, calculated by Murphy et al. (2010b). Note that the main text by Murphy et al. (2010b) gives a value of 0.37, but the calculations in their Table 1, as well as supplementary

Age-depth models for PETM at ODP Site 1266




Figure 4. The three age-depth models [for the PETM at IODP site 1266](#) derived under the assumption of constant flux ([black-red](#) line), [increasing-decreasing](#) flux ([red-green](#) line) and [decreasing-increasing](#) flux (blue line). Thick lines are median age over 1000 replicates, dashed lines are the 95 % envelope of the ages.

[materials](#), indicate that the value 0.48 had been used and this is adopted here. We assume this value is a mean value, and the actual flux follows a normal distribution with this mean and a standard deviation of [0.050.04](#), reflecting the [+5%-2σ-2σ](#) uncertainty of the flux estimate [given by Farley and Eltgroth \(2003\)](#) of 0.08 [given by Murphy et al. \(2010b\)](#). For the increase and decrease [scenarios](#), we assume that, over an interval of 1 Myr, the flux values increases or decreases by a factor of two. We

525 [place-the-beginning-use the base](#) of the PETM [main-interval-in-the-middle-of-this-time-interval](#), placed at the base of the clay, or "core", interval at 306.78 mcd (m composite depth) as the base of the simulation and evaluate the age-depth model over the interval ranging upwards the core to 303.5 mcd, which is past the end of the recovery phase of PETM, placed at 304.19 mcd. We ran the analysis using 1000 Monte Carlo simulations, and give results rounded to the next kyr.

All three age-depth models show [clear signs of sedimentary condensation and dilution](#) [sedimentary condensation](#) over the

530 PETM interval, as expressed by their changing slopes (Figure ??). Both the pre and the main interval are strongly condensed relative to the recovery interval (, Figure ??). This is clearly displayed by the recovery interval, which is 2.65 times thicker than the main interval (2.65 m vs. 1 m), but more than 50 % shorter than the main interval when measured in time (median values: 55, 53, and 51 % increasing, constant, and decreasing flux scenario). Median [core interval](#), expressed by the near-horizontal




Figure 5. Density of the estimated duration of the PETM main and recovery interval.




Figure 6. Median sedimentation rate ~~and condensation~~ in the stratigraphic domain for the three scenarios defined.




Figure 7. Median condensation in the stratigraphic domain for the three scenarios defined.

Scenario	Duration of the clay interval [kyr]	Duration of the recovery [kyr]	Total PETM duration [kyr]
Increasing ^3He flux	182	111	250
Constant ^3He flux	173	100	234
Decreasing ^3He flux	161	86	212
Murphy et al. (2010)	167	100	234

Table 1. Median durations of PETM phases at IODP site 1266

slope (Figure 4). An increase in sedimentation rate is visible from the initial recovery interval ("recovery I") to the following intervals II and III (Figure 6). The median duration of the PETM main interval is 93, 94, and 98 (from the base of the clay interval to the end of the recovery) is 234 ($2\sigma = 39$) kyr for the increasing, constant, and decreasing flux scenarios, with comparable interquartile ranges (10, 10, and 11 kyr) (Figure ?? A). The recovery interval is short compared to the main interval, with a median duration of 41, 44, and 48 constant flux scenario, and 212 ($2\sigma = 32$) and 250 ($2\sigma = 45$) kyr for the increasing, constant, and decreasing flux scenario, and comparable interquartile ranges (4, 5, and 4 kyr, respectively) (Figure ?? B) decreasing and increasing scenarios, respectively (Table 1). This duration is very close to the 234 ($^{+48}_{-34}$) kyr duration reported by (Murphy et al., 2010b).

Over the 4.25 m of the examined PETM intervals, sedimentation rates vary by a factor of more than 25 for Median durations of the phases of PETM estimated under the three scenarios and their comparison with the results by Murphy et al. (2010b) are provided in Table 1. The disagreement between the results obtained using the FAM model and those reported in the original study does not exceed 6 kyr.

The PETM interval in Site 1266 core spans 2.59 m. The age-depth model by Murphy et al. (2010b) relied on linear sedimentation rates calculated by combining constant ^3He flux with a total duration of the interval derived from cyclostratigraphy (Röhl et al., 2007)

Sedimentation rates have similar variability in all three scenarios. The range of sedimentation rates is ~~0.6 to 18.9~~ 0.1 to 6.4 cm/kyr for the increasing, ~~0.6 to 18~~ 0.1 to 7.3 cm/kyr for the constant, and ~~0.6 to 16.0~~ 0.1 to 8.8 cm/kyr for the decreasing flux scenario, with the lowest values at the transition of the ~~pre-pre-~~ to the main interval, and highest values in the recovery interval (Figure ~~??-A6~~). Conversely, condensation ranges from ~~0.06-0.1~~ kyr/cm ~~and 1.58 to 10.5~~ kyr/cm across the scenarios (Figure ~~??-B~~). ~~Note that the sedimentation rate estimates are elevated by a factor of more than 5 in the recovery interval compared to cyclostratigraphic analyses (Röhl et al., 2007, Röhl et al. (2000)).~~

While there is no large offset in sedimentation rates and age-depth models between the scenarios, the varying fluxes generate systematic deviations from the constant flux scenario. Under increasing (decreasing) flux, sedimentation rates in the top of the examined interval are systematically ~~higher (lower)~~ lower (higher), leading to ~~shortened (extended)~~ prolonged (shortened) durations of the recovery interval. ~~For the bottom of the section, the effect is reversed, where increasing (decreasing) flux leads to extended (shortened) durations of the pre-interval and clay intervals.~~ Overall, results are consistent with those of ~~Farley and Elteroth (2003)~~ Murphy et al. (2010b) and show that they are robust with respect to variations in ^3He - ^3He flux. Under ~~increasing and decreasing fluxes, median PETM main interval durations differ by only 3~~ decreasing flux, the median PETM duration differs by 10% from the constant flux scenario (Table 1).

We conclude that variability of ^3He - ^3He fluxes by a factor of two on the timescale below 1 Myr ~~only have a weak~~ has a moderate effect on the age-depth model at ODP site ~~690-1266~~ and the derived estimates of PETM duration. Using tracer fluxes to estimate age-depth models is a powerful tool to estimate high-resolution age-depth models and identify local variations in condensation and sedimentation rates.

3.3 Discussion

4 Discussion

We have introduced two methods to estimate age-depth models from complex stratigraphic or sedimentological data. ICON used information on upsection change in sedimentation rate to estimate age-depth models, while FAM compares observed tracer values with assumptions on past tracer ~~fluxed fluxes~~ to constrain age-depth relationships. ~~As examples~~ Informing a model with an external sedimentation or accumulation rate (whether a number or a distribution) is not conceptually different from informing it with tie-points, but presents a novelty of the ICON approach. This novelty may initially be a challenge for users, but it effectively presents an opportunity to enrich age-depth models with information previously not exploited: compilations of accumulation rates in depositional environments from actualistic studies (Sadler, 1981; Enos, 1991; McNeill, 2005; Scott, 2009), theoretical analyses on the temporal and spatial scaling of accumulation rates (Aadland et al., 2018; Tipper, 2016), and even relative estimates of sedimentation rates (Davies et al., 2019), equally handled by admtools. The novelty of this approach may initially result in seemingly higher uncertainty, but one should consider that it is in fact a quantification of uncertainty that had been present in all previous age-depth model inference, but remained unaccounted for.

As examples of the use of ICON and FAM, we have applied the new methods to constrain the timing and duration of the Frasnian-Famennian boundary and the Upper Kellwasser Event with sedimentation rates constraints from cyclostratigraphy

(Da Silva et al., 2020), and examined the robustness of age-depth models for the Paleocene-Eocene Thermal Maximum (PETM) to variations in tracer fluxes (Farley and Eltgroth, 2003)(Murphy et al., 2010b).

4.0.1 Comparison with other methods

4.1 Comparison with other methods

Bayesian approaches such as Bchron, modified Bchron, Bacon and OxCal are common among the current methods to estimate age-depth models (Blaauw and Christen, 2011; Trayler et al., 2019; Haslett and Parnell, 2008) (Table 4.2). Trachsel and Telford (2017) pointed out that Bayesian approaches perform well compared to “classic” approaches such as CLAM. FAM and ICON are not Bayesian, as they do not rely on prior or posterior distributions. Bayesian methods can become computationally expensive when parameter spaces are high-dimensional. As stratigraphic data gets more complex, the number of parameters and computation time increases. ~~For example, the computations accompanying Trayler et al. (2023) publication of astroBayes will take “several days or weeks” to compute on a laptop. In contrast, the~~ The computation time for the examples ~~show~~ shown here is typically within minutes.

By imposing monotonicity constraints on sample paths, Bayesian methods can resolve age reversals and reduce age uncertainties at the tie points. In ICON and FAM, users decide how to resolve age reversals, and the uncertainties of age-depth models at tie points cannot be improved upon by the method. In this hierarchical design, information from between tie points cannot reduce uncertainties of the tie points. Information on sedimentation rates and tracer fluxes are typically either expressed on an ordinal scale or associated with high uncertainty. The hierarchical design reflects the idea that an absolute age (e.g. from U-Pb dating) can not be improved upon by information about the relative distribution of time (e.g. from changes in sedimentation rates) between tie points. Especially for changes in sedimentation rates estimated from cyclostratigraphy, this prevents circular reasoning, as the usage of external age constraints is recommended with cyclostratigraphic analysis (Sinnesael et al., 2019).

Assumptions on sediment accumulation employed by current methods to estimate age-depth models ~~are often made for mathematical convenience (e.g. assume a Poisson structure). In fact, we cannot possibly apply universally, since the dynamics of sedimentation varies between environments (Sadler, 1981). We~~ do not know if sedimentary events follow a Poisson distribution across all timescales and depositional environments (e.g., Schumer and Jerolmack (2009)). ~~Such assumptions do not reflect our understanding of~~ It is not possible to prescribe upfront all correct combinations of assumptions for all possible use-cases stratigraphers may study, also because the dynamics of sedimentations across timescales and environments remain themselves the topic of ongoing research (Aadland et al., 2018; Wilkinson, 2015; Davies et al., 2019) and age-depth models may feed back into this research by improving the inference on the structure of the stratigraphic record and should rather be considered simplified and unspecific error models that reflect various degrees of pessimism about the the structure of the stratigraphic-geological record. ICON and FAM ~~improve upon this by explicitly relying on~~ are an attempt to contribute to this research by making it possible to test different assumptions and validate them using stratigraphic and sedimentological information ~~to estimate age-depth models.~~

4.1.1 Limitations of this approach

615 4.2 Limitations of this approach

In `admtools` v0.3.5.0 (?) (Hohmann, 2025), neither ICON nor FAM can incorporate information on gaps in the fossil stratigraphic record. Mathematically, it is possible to include gaps into ICON, and will be implemented in later versions of the package. Similarly as constraining sedimentation rates, age-depth models inference may be constrained by the duration of gaps using literature information or calculations from average accumulation and sedimentation rates (Wilkinson, 2015; Paola et al., 2018; Tomašových

620 For FAM, incorporation of gaps is possible in the edge case of constant assumed tracer flux. For more complex assumed tracer fluxes, it is not obvious to see whether gaps can be incorporated, as they remove parts of the signal the observed tracer is matched to. However, even with the option to incorporate hiatuses, hiatus durations and locations are notoriously challenging to estimate and justify empirically. Often times, the existence of hiatuses is invoked to explain a mismatch between signals of different sections. Hohmann et al. (2024) used forward simulations of carbonate platforms to show that gap duration and locations are functions of external drivers of stratigraphic architectures, and Meyers and Sageman (2004) used evolutive harmonic analysis to identify relatively short hiatuses (1 - 100 kyr) in a simulation study. These are just two examples showing that gaps gap locations and durations are not random, and can be identified with careful data analysis and forward models. Note that in a case where there are unidentified gaps in a section, the reconstructed sedimentation rate will always be lower than the “true” sedimentation rate, leading to a Sadler-type effect (Sadler, 1981).

630 An assumption made by all methods to ~~estimated~~ estimate age-depth relationships is that a given stratigraphic position has a unique age, which might be unknown, but can be assigned an uncertainty. Dating of organic remains in modern environments (e.g., Dominguez et al. (2016) or Tomašových et al. (2018)) shows that particle ages at a given location can differ significantly as a result of mixing in the surface mixed layer, and thus violate this assumption. This effect might differ across environments, but time-averaging in Holocene deposits can reach values of multiple thousands years (see e.g., Berensmeier et al. (2023)) and thus exceed the age uncertainty of age-depth models by orders of magnitude. While Bayesian methods such as Bchron or Bacon automatically resolve reversals (Table 2), the maximum temporal resolution achievable in age-depth models will ultimately be limited by the depositional resolution of the sedimentary record, which is controlled by both physical (slumping, regularity of sediment supply) and biological factors (bioturbation) (Kowalewski and Bambach, 2008).

640 One key feature of FAM and ICON is that they are assumption-explicit in the sense that the user must specify all distributions that contribute to the age-depth model. While this requires additional coding effort, it results in an assumption-explicit age-depth model, where every assumption made is known and documented. As a result, uncertainties in age-depth models are a direct reflection of our understanding of the structure of the stratigraphic record. A second benefit is that age-depth model construction is replicable, even using different computational environments.

Table 2. Comparison of different methods to estimate age-depth models

Method	Bayesian	Information source	Developed for time scale
BChron	yes	tie points	radiocarbon
modified BChron	yes	tie points	deep time (borrows assumptions from BChron) deep time (borrows assumptions from
OxCal	yes	tie points, gaps, sed. rates	radiocarbon
Bacon	yes	tie points tie points, sed. rate estimates and their variability	radiocarbon
astroBayes	yes	tie points, breaks in sed. rate	deep time
CLAM	no	tie points	radiocarbon
ICON	no	tie points, sed. rate estimates	any
FAM	no	tie points, tracer values	any

4.2.1 ~~Uncertainties in age-depth modelling~~

645 4.3 Uncertainties in age-depth modelling

For researchers using geohistorical data, age-depth models with low uncertainties are highly desirable. They allow for exact dating of events and provide low uncertainties of rates of past change, opening up the opportunity to ~~draw~~ use past climate perturbations as analogues for future climate change (Trayler et al., 2023). It is easy to construct an age-depth model with seemingly no uncertainties by simply connecting mean ages with a straight line (e.g., Tobin et al. (2012)). However, 650 low uncertainties should not be the target to assess the quality of age-depth models. Trachsel and Telford (2017) found that CLAM underestimates age uncertainties in varved sediments and De Vleeschouwer and Parnell (2014) pointed out that in the Devonian timescale in Gradstein (2012), age uncertainties decreases between absolutely dated tie points - an unexpected and counter-intuitive behavior. In the latter example, the problem could be corrected by applying the Bayesian Bchron approach (De Vleeschouwer and Parnell, 2014). Both examples demonstrate that there are empirical and logical lower bounds 655 on uncertainties in age-depth models. In cyclostratigraphy, where this has been evaluated, even experienced users tend to overestimate the accuracy of age information calculated from age-depth models they build (Sinnesael et al., 2019). We argue that the best age-depth model should not be the one with the lowest uncertainty, but the one that best reflects our understanding of and uncertainty about the structure of the stratigraphic record in the environment and timescale of interest. ~~This poses~~ The importance of such contextual information was highlighted in the results of Cyclostratigraphy Intercomparison Project (CIP) (Sinnesael et al., 2019), which identified that cyclostratigraphic analyses lacking such context tended to be less accurate. However, contextual information and expert knowledge could not be - until now - consistently codified into the process of inferring an age-depth model. This poses an obstacle to replicability. Improving upon this situation poses a twofold challenge: 660

First, understanding drivers of the local structure of the record, second, incorporating this information and ~~and~~ the uncertainty associated with it into age-depth models. ~~When such~~ Currently, no standard for reporting uncertainty from different sources exists, although various approaches have been proposed (Zeeden et al., 2014; Huybers and Wunsch, 2004), but integrating uncertainties from biostratigraphy (Punyasena et al., 2012; Pálffy, 2007), range offset and biofacies (Holland and Patzkowsky, 1999) or sheer incompleteness (Hohmann et al., 2024) remains a technical challenge. CIP results led to the recommendation that age-depth models should be intercalibrated with other geochronometers (see e.g. Jarochowska et al. (2020)), but the question remains: what if two age-depth models for the same section, based on independent geochronometers, disagree? We propose that such alternative age-depth models ~~are used~~, can be treated as hypotheses in a hypothetico-deductive scientific framework and stratigraphic data of interest can be evaluated under these hypotheses to assess how much stratigraphic uncertainty affects the conclusions of a study. However, ~~admtools~~ (Hohmann, 2025) allows to add or remove information, leading to simple partitioning of uncertainty, as shown in example 1. For instance, the assumption of a constant flux of a tracer may be considered naive, especially over long time scales, and it is not always possible to validate it with independent data, as has been done by Middeldorp (1982) and Young et al. (1999). The assumptions of constant flux can be nonetheless still used as a null hypothesis, against which systematic or random changes in the flux can be tested.

When age-depth models are constructed with the widest breadth of stratigraphic and sedimentological information, timing and rates of past change might be associated with large uncertainties. However, these uncertainties will reflect our understanding of the structure of the stratigraphic record. On the other hand, incorporating information on the dynamics and internal rhythmicity of depositional environments, such as tidal or seasonal ones, in age-depth models provides us with new tie points and higher-resolution age-depth models. Both FAM and ICON turn complex information into age-depth models. To make full use of these methods, more research into this structure across environments, timescales, and its external controls are required.

5 Funding

S.J.B. thanks the the CycloNet project, funded by the Research Foundation Flanders (FWO, grant no W000522N), for financial support.

Funded by the European Union (ERC, MindTheGap, StG project no 101041077). Views and opinions expressed are however those of the author(s) only and do not necessarily reflect those of the European Union or the European Research Council. Neither the European Union nor the granting authority can be held responsible for them.

6 Conclusions

We have introduced two new methods, FAM and ICON, to estimate age-depth models from complex stratigraphic and sedimentological data. FAM compares observed tracer values with assumptions of fluxes in the time domain, while ICON uses information on changes in sedimentation rates (e.g., from cyclostratigraphy, sequence stratigraphy, or other expert knowledge) to construct age-depth models. Both methods are non-parametric in the sense that they do not make any a priori model assump-

tions other than that the law of superposition holds. Users can implement error models that reflect their knowledge about the
695 local drivers of stratigraphic architectures, resulting in age-depth models that are data driven ~~rather than assumption driven~~ and
allow to study uncertainty partitioning.

Code availability. All code used for the examples is available in Hohmann and Jarochowska (2025) and can be accessed under
<https://doi.org/10.5281/zenodo.15489276>. The `admtools` package used for the analysis is available on CRAN (the Comprehensive R
Archive Network), with versions being archived on Zenodo (Hohmann, 2025).




Figure A1. r_{opt}^2 values from eTimeOpt testing for short eccentricity amplitude modulation at 3.54 m and 1.56 m (A) and throughout the whole section (B) in the Steinbruch Schmidt section. The heights shown in A roughly corresponding to the locations of the Upper Kellwasser Bed and the location of the dated ash bed.

Author contributions. Based on the CRediT taxonomy. **Niklas Hohmann:** Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data curation, Writing - Original draft, Writing - Review and editing, Visualization. **David De Vleeschouwer:** Conceptualization, Writing - Review and editing. **Sietske Batenburg:** Writing - Review and editing. **Emilia Jarochowska:** Funding acquisition, Project administration, Writing - Review and editing, Validation.

705 *Competing interests.* The authors declare no competing interests.

Acknowledgements. This work originated from discussions in the CycloNet project, funded by the Research Foundation Flanders (FWO, grant no. W000522N).




Figure A2. Floating age-depth model Age-depth models with (A) and anchored age-depth model without (B) incorporation the uncertainty of the radiometric dating for the Steinbruch Schmidt section based on eTimeOpt results testing for short eccentricity amplitude modulation. In the floating age-depth model (A), time is measured relative to the bentonite layer dated by Percival et al. (2018), and all uncertainties in the age-depth model arise from the uncertainties of the sedimentation rates estimated using eTimeOpt. The anchored age-depth model is constructed by combining, while in (B), they are a combination of the absolute age uncertainty of Percival et al. (2018) for the bentonite layer with the radiometric dates from Percival et al. (2018) (dashed lines) and sedimentation rates estimated from eTimeOpt. Uncertainties result from both the uncertainties Dashed lines in U-Pb dating and estimates (B) indicate the age uncertainty of sedimentation rates the section when the radiometric uncertainty is propagated using eTimeOpt the median sedimentation rate.

References

- Quadpack: A Subroutine Package for Automatic Integration, Springer International Publishing, Cham, oCLC: 1159113603.
- 710 Aadland, T., Sadler, P. M., and Helland-Hansen, W.: Geometric interpretation of time-scale dependent sedimentation rates, *Sedimentary Geology*, 371, 32–40, <https://doi.org/10.1016/j.sedgeo.2018.04.003>, 2018.
- Abril-Hernández, J.: 210Pb-dating of sediments with models assuming a constant flux: CFCS, CRS, PLUM, and the novel χ -mapping. Review, performance tests, and guidelines, *Journal of Environmental Radioactivity*, 268-269, 107 248, <https://doi.org/10.1016/j.jenvrad.2023.107248>, 2023.
- 715 Appleby, P. and Oldfield, F.: The calculation of lead-210 dates assuming a constant rate of supply of unsupported 210Pb to the sediment, *CATENA*, 5, 1–8, [https://doi.org/10.1016/S0341-8162\(78\)80002-2](https://doi.org/10.1016/S0341-8162(78)80002-2), 1978.
- Averbuch, O., Tribouillard, N., Devleeschouwer, X., Riquier, L., Mistiaen, B., and Van Vliet-Lanoe, B.: Mountain building-enhanced continental weathering and organic carbon burial as major causes for climatic cooling at the Frasnian–Famennian boundary (c. 376 Ma)?, *Terra*

- Nova, 17, 25–34, <https://doi.org/10.1111/j.1365-3121.2004.00580.x>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-3121.2004.00580.x>, 2005.
- 720 Barker, M., Chue Hong, N. P., Katz, D. S., Lamprecht, A.-L., Martinez-Ortiz, C., Psomopoulos, F., Harrow, J., Castro, L. J., Grunepeter, M., Martinez, P. A., and Honeyman, T.: Introducing the FAIR Principles for research software, *Scientific Data*, 9, 622, <https://doi.org/10.1038/s41597-022-01710-x>, publisher: Nature Publishing Group, 2022.
- Becker, R., Marshall, J., Da Silva, A.-C., Agterberg, F., Gradstein, F., and Ogg, J.: The Devonian Period, pp. 733–810, Elsevier, <https://doi.org/10.1016/B978-0-12-824360-2.00022-X>, doi: 10.1016/B978-0-12-824360-2.00022-X, 2020.
- 725 Berensmeier, M., Tomašových, A., Nawrot, R., Cassin, D., Zonta, R., Koubová, I., and Zuschin, M.: Stratigraphic expression of the human impacts in condensed deposits of the Northern Adriatic Sea, *Geological Society, London, Special Publications*, 529, 195–222, <https://doi.org/10.1144/SP529-2022-188>, publisher: The Geological Society of London, 2023.
- Blaauw, M.: Methods and code for ‘classical’ age-modelling of radiocarbon sequences, *Quaternary Geochronology*, 5, 512–518, <https://doi.org/10.1016/j.quageo.2010.01.002>, 2010.
- 730 Blaauw, M.: Out of tune: the dangers of aligning proxy archives, *Quaternary Science Reviews*, 36, 38–49, <https://doi.org/10.1016/j.quascirev.2010.11.012>, 2012.
- Blaauw, M. and Christen, J.: Flexible paleoclimate age-depth models using an autoregressive gamma process, *Bayesian Analysis*, 6, 457–474, <https://doi.org/10.1214/ba/1339616472>, 2011.
- 735 Blaauw, M. and Christen, J. A.: Radiocarbon Peat Chronologies and Environmental Change, *Journal of the Royal Statistical Society Series C: Applied Statistics*, 54, 805–816, <https://doi.org/10.1111/j.1467-9876.2005.00516.x>, 2005.
- Blaauw, M., Bakker, R., Christen, J. A., Hall, V. A., and Plicht, J. v. d.: A Bayesian Framework for Age Modeling of Radiocarbon-Dated Peat Deposits: Case Studies from the Netherlands, *Radiocarbon*, 49, 357–367, <https://doi.org/10.1017/S0033822200042296>, 2007.
- Blard, P.-H., Suchéras-Marx, B., Suan, G., Godet, B., Tibari, B., Dutilleul, J., Mezine, T., and Adatte, T.: Are marl-limestone alternations mainly driven by CaCO₃ variations at the astronomical timescale? New insights from extraterrestrial ³He, *Earth and Planetary Science Letters*, 614, 118–173, <https://doi.org/10.1016/j.epsl.2023.118173>, 2023.
- 740 Bookstein, F. L.: Random walk and the existence of evolutionary rates, *Paleobiology*, 13, 446–464, <https://doi.org/10.1017/S0094837300009039>, publisher: Cambridge University Press, 1987.
- Brent, R. P.: Algorithms for minimization without derivatives, Dover Publications, Mineola, N.Y, 2002.
- 745 Bronk Ramsey, C.: Deposition models for chronological records, *Quaternary Science Reviews*, 27, 42–60, <https://doi.org/10.1016/j.quascirev.2007.01.019>, 2008.
- Bronk Ramsey, C.: Bayesian Analysis of Radiocarbon Dates, *Radiocarbon*, 51, 337–360, <https://doi.org/10.1017/S0033822200033865>, 2009a.
- Bronk Ramsey, C.: Bayesian Analysis of Radiocarbon Dates, *Radiocarbon*, 51, 337–360, <https://doi.org/10.1017/S0033822200033865>, 2009b.
- 750 Cagliari, J., Schmitz, M. D., Tedesco, J., Trentin, F. A., and Lavina, E. L. C.: High-precision U-Pb geochronology and Bayesian age-depth modeling of the glacial-postglacial transition of the southern Paraná Basin: Detailing the terminal phase of the Late Paleozoic Ice Age on Gondwana, *Sedimentary Geology*, 451, 106–397, <https://doi.org/10.1016/j.sedgeo.2023.106397>, 2023.
- Carmichael, M. J., Inglis, G. N., Badger, M. P. S., Naafs, B. D. A., Behrooz, L., Remmelzwaal, S., Monteiro, F. M., Rohrsen, M., Farnsworth, A., Buss, H. L., Dickson, A. J., Valdes, P. J., Lunt, D. J., and Pancost, R. D.: Hydrological and associated biogeochemi-
- 755

- cal consequences of rapid global warming during the Paleocene-Eocene Thermal Maximum, *Global and Planetary Change*, 157, 114–138, <https://doi.org/10.1016/j.gloplacha.2017.07.014>, 2017.
- Carmichael, S. K., Waters, J. A., Königshof, P., Suttner, T. J., and Kido, E.: Paleogeography and paleoenvironments of the Late Devonian Kellwasser event: A review of its sedimentological and geochemical expression, *Global and Planetary Change*, 183, 102984, <https://doi.org/10.1016/j.gloplacha.2019.102984>, 2019.
- Cerda, M., Evangelista, H., Valdés, J., Siffedine, A., Boucher, H., Nogueira, J., Nepomuceno, A., and Ortlieb, L.: A new 20th century lake sedimentary record from the Atacama Desert/Chile reveals persistent PDO (Pacific Decadal Oscillation) impact, *Journal of South American Earth Sciences*, 95, 102302, <https://doi.org/10.1016/j.jsames.2019.102302>, 2019.
- Claeys, P., Casier, J.-G., and Margolis, S. V.: Microtektites and Mass Extinctions: Evidence for a Late Devonian Asteroid Impact, *Science*, 257, 1102–1104, <https://doi.org/10.1126/science.257.5073.1102>, publisher: American Association for the Advancement of Science, 1992.
- da Silva, A.-C.: Anchoring the Late Devonian mass extinction in absolute time by integrating climatic controls and radio-isotopic dating: Supplementary code, Zenodo, <https://doi.org/10.5281/ZENODO.12516430>, language: en DOI: 10.5281/ZENODO.12516430, 2024.
- Da Silva, A.-C., Sinnesael, M., Claeys, P., Davies, J. H. F. L., de Winter, N. J., Percival, L. M. E., Schaltegger, U., and De Vleeschouwer, D.: Anchoring the Late Devonian mass extinction in absolute time by integrating climatic controls and radio-isotopic dating, *Scientific Reports*, 10, 12940, <https://doi.org/10.1038/s41598-020-69097-6>, publisher: Nature Publishing Group, 2020.
- Davies, N. S., Shillito, A. P., and McMahon, W. J.: Where does the time go? Assessing the chronostratigraphic fidelity of sedimentary geological outcrops in the Pliocene–Pleistocene Red Crag Formation, eastern England, *Journal of the Geological Society*, 176, 1154–1168, <https://doi.org/10.1144/jgs2019-056>, publisher: Geological Society of London Section: Research article, 2019.
- De Vleeschouwer, D. and Parnell, A. C.: Reducing time-scale uncertainty for the Devonian by integrating astrochronology and Bayesian statistics, *Geology*, 42, 491–494, <https://doi.org/10.1130/G35618.1>, 2014.
- De Vleeschouwer, D., Percival, L. M. E., Wichern, N. M. A., and Batenburg, S. J.: Pre-Cenozoic cyclostratigraphy and palaeoclimate responses to astronomical forcing, *Nature Reviews Earth & Environment*, 5, 59–74, <https://doi.org/10.1038/s43017-023-00505-x>, publisher: Nature Publishing Group, 2024.
- Dickens, G. R., O’Neil, J. R., Rea, D. K., and Owen, R. M.: Dissociation of oceanic methane hydrate as a cause of the carbon isotope excursion at the end of the Paleocene, *Paleoceanography*, 10, 965–971, <https://doi.org/10.1029/95PA02087>, <https://onlinelibrary.wiley.com/doi/pdf/10.1029/95PA02087>, 1995.
- Dominguez, J. G., Kosnik, M. A., Allen, A. P., Hua, Q., Jacob, D. E., Kaufman, D. S., and Whitacre, K.: Time-Averaging and Stratigraphic Resolution in Death Assemblages and Holocene Deposits: Sydney Harbour’s Molluscan Record, *PALAIOS*, 31, 564–575, <https://doi.org/10.2110/palo.2015.087>, 2016.
- Enos, P.: Sedimentary parameters for computer modeling, *Bulletin (Kansas Geological Survey)*, 233, 63–99, <https://doi.org/10.17161/kgbulletin.no.233.20449>, 1991.
- Farley, K. A.: Extraterrestrial Helium in Seafloor Sediments: Identification, Characteristics, and Accretion Rate Over Geologic Time, in: *Accretion of Extraterrestrial Matter Throughout Earth’s History*, edited by Peucker-Ehrenbrink, B. and Schmitz, B., pp. 179–204, Springer US, Boston, MA, https://doi.org/10.1007/978-1-4419-8694-8_11, dOI: 10.1007/978-1-4419-8694-8_11, 2001.
- Farley, K. A. and Eltgroth, S. F.: An alternative age model for the Paleocene–Eocene thermal maximum using extraterrestrial ^3He , *Earth and Planetary Science Letters*, 208, 135–148, [https://doi.org/10.1016/S0012-821X\(03\)00017-7](https://doi.org/10.1016/S0012-821X(03)00017-7), 2003.
- Farley, K. A. and Eltgroth, S. F.: (Appendix 1) Helium isotopic ratios and sedimentation rate model of ODP Hole 113-690B, PAN-GAEA, <https://doi.org/10.1594/PANGAEA.723907>, in supplement to: Farley, KA; Eltgroth, SF (2003): An alternative age model

for the Paleocene-Eocene thermal maximum using extraterrestrial ^3He . *Earth and Planetary Science Letters*, 208(3-4), 135-148, [https://doi.org/10.1016/S0012-821X\(03\)00017-7](https://doi.org/10.1016/S0012-821X(03)00017-7), 2003.

Frieling, J., Svensen, H. H., Planke, S., Cramwinckel, M. J., Selnes, H., and Sluijs, A.: Thermogenic methane release as a cause for the long duration of the PETM, *Proceedings of the National Academy of Sciences*, 113, 12 059–12 064, <https://doi.org/10.1073/pnas.1603348113>, publisher: Proceedings of the National Academy of Sciences, 2016.

Gould, S. J. and Eldredge, N.: Punctuated equilibria: an alternative to phyletic gradualism, *Models in paleobiology*, 1972, 82–115, publisher: San Francisco, 1972.

Gradstein, F., ed.: *Geologic Time Scale 2020*, Elsevier, <https://doi.org/10.1016/C2020-1-02369-3>, doi: 10.1016/C2020-1-02369-3, 2020.

Gradstein, F. M., ed.: *The geologic time scale 2012*, Elsevier, Amsterdam ; Boston, 1st edn., ISBN 978-0-444-59425-9 978-0-444-59390-0 978-0-444-59434-1 978-0-444-59435-8, oCLC: ocn773025121, 2012.

Harrigan, C. O., Schmitz, M. D., Over, D. J., Trayler, R. B., and Davydov, V. I.: Recalibrating the Devonian time scale: A new method for integrating radioisotopic and astrochronologic ages in a Bayesian framework, *GSA Bulletin*, 134, 1931–1948, <https://doi.org/10.1130/B36128.1>, 2021.

Haslett, J. and Parnell, A.: A Simple Monotone Process with Application to Radiocarbon-Dated Depth Chronologies, *Journal of the Royal Statistical Society Series C: Applied Statistics*, 57, 399–418, <https://doi.org/10.1111/j.1467-9876.2008.00623.x>, 2008.

Haywood, A. M., Ridgwell, A., Lunt, D. J., Hill, D. J., Pound, M. J., Dowsett, H. J., Dolan, A. M., Francis, J. E., and Williams, M.: Are there pre-Quaternary geological analogues for a future greenhouse warming?, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 369, 933–956, <https://doi.org/10.1098/rsta.2010.0317>, publisher: Royal Society, 2011.

Hohmann, N.: Incorporating Information on Varying Sedimentation Rates into Paleontological Analyses, *PALAIOS*, 36, 53–67, <https://doi.org/10.2110/palo.2020.038>, 2021.

Hohmann, N.: Supplementary data and code for "Nonparametric estimation of age-depth models from sedimentological and stratigraphic data", <https://doi.org/10.5281/zenodo.13639816>, 2024.

Hohmann, N.: admtools, <https://doi.org/10.5281/zenodo.15279284>, 2025.

Hohmann, N. and Jarochowska, E.: Supplementary data and code for "Nonparametric estimation of age-depth models from sedimentological and stratigraphic data", <https://doi.org/10.5281/zenodo.15489276>, 2025.

Hohmann, N., Koelewijn, J. R., Burgess, P., and Jarochowska, E.: Identification of the mode of evolution in incomplete carbonate successions, *BMC Ecology and Evolution*, 24, 113, <https://doi.org/10.1186/s12862-024-02287-2>, 2024.

Holland, S. M. and Patzkowsky, M. E.: Models for simulating the fossil record, *Geology*, 27, 491–494, [https://doi.org/10.1130/0091-7613\(1999\)027<0491:MFSTFR>2.3.CO;2](https://doi.org/10.1130/0091-7613(1999)027<0491:MFSTFR>2.3.CO;2), 1999.

Hunter-Zinck, H., Siqueira, A. F. d., Vásquez, V., Barnes, R., and Martinez, C. C.: Ten simple rules on writing clean and reliable open-source scientific software, *PLOS Computational Biology*, 17, e1009 481, <https://doi.org/10.1371/journal.pcbi.1009481>, publisher: Public Library of Science, 2021.

Huybers, P. and Wunsch, C.: A depth-derived Pleistocene age model: Uncertainty estimates, sedimentation variability, and nonlinear climate change, *Paleoceanography*, 19, <https://doi.org/10.1029/2002PA000857>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2002PA000857>, 2004.

Jarochowska, E., Nohl, T., Grohgan, M., Hohmann, N., Vandenbroucke, T. R. A., and Munnecke, A.: Reconstructing Depositional Rates and Their Effect on Paleoenvironmental Proxies: The Case of the Lau Carbon Isotope Excursion in Gotland, Sweden, *Paleoceanography and Paleoclimatology*, 35, e2020PA003 979, <https://doi.org/10.1029/2020PA003979>, 2020.

- Kowalewski, M. and Bambach, R. K.: The limits of paleontological resolution, in: High-resolution approaches in stratigraphic paleontology, pp. 1–48, Springer, 2008.
- 835 Kurtz, A. C., Kump, L. R., Arthur, M. A., Zachos, J. C., and Paytan, A.: Early Cenozoic decoupling of the global carbon and sulfur cycles, *Paleoceanography*, 18, <https://doi.org/10.1029/2003PA000908>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2003PA000908>, 2003.
- Lacourse, T. and Gajewski, K.: Current practices in building and reporting age-depth models, *Quaternary Research*, 96, 28–38, <https://doi.org/10.1017/qua.2020.47>, 2020.
- 840 Li, M., Kump, L. R., Hinnov, L. A., and Mann, M. E.: Tracking variable sedimentation rates and astronomical forcing in Phanerozoic paleoclimate proxy series with evolutionary correlation coefficients and hypothesis testing, *Earth and Planetary Science Letters*, 501, 165–179, <https://doi.org/10.1016/j.epsl.2018.08.041>, 2018.
- MacLeod, N.: Punctuated anagenesis and the importance of stratigraphy to paleobiology, *Paleobiology*, 17, 167–188, <https://doi.org/10.1017/S0094837300010472>, publisher: Cambridge University Press, 1991.
- 845 Malmgren, B. A., Berggren, W. A., and Lohmann, G. P.: Evidence for punctuated gradualism in the Late Neogene *Globorotalia tumida* lineage of planktonic foraminifera, *Paleobiology*, 9, 377–389, <https://doi.org/10.1017/S0094837300007843>, publisher: Cambridge University Press, 1983.
- McNeill, D. F.: Accumulation rates from well-dated late Neogene carbonate platforms and margins, *Sedimentary Geology*, 175, 73–87, <https://doi.org/10.1016/j.sedgeo.2004.12.032>, 2005.
- Meyers, S. R.: Astrochron: An R Package for Astrochronology, <https://cran.r-project.org/package=astrochron>, 2014.
- 850 Meyers, S. R.: Cyclostratigraphy and the problem of astrochronologic testing, *Earth-Science Reviews*, 190, 190–223, <https://doi.org/10.1016/j.earscirev.2018.11.015>, 2019.
- Meyers, S. R. and Sageman, B. B.: Detection, quantification, and significance of hiatuses in pelagic and hemipelagic strata, *Earth and Planetary Science Letters*, 224, 55–72, <https://doi.org/10.1016/j.epsl.2004.05.003>, 2004.
- Middeldorp, A. A.: Pollen concentration as a basis for indirect dating and quantifying net organic and fungal production in a peat bog ecosystem, *Review of Palaeobotany and Palynology*, 37, 225–282, [https://doi.org/10.1016/0034-6667\(82\)90003-3](https://doi.org/10.1016/0034-6667(82)90003-3), 1982.
- 855 Mukhopadhyay, S., Farley, K. A., and Montanari, A.: A Short Duration of the Cretaceous-Tertiary Boundary Event: Evidence from Extraterrestrial Helium-3, *Science*, 291, 1952–1955, <https://doi.org/10.1126/science.291.5510.1952>, 2001.
- Murphy, B., Farley, K. A., and Zachos, J. C.: Extraterrestrial ³He estimation across the Paleocene-Eocene thermal maximum at ODP Site 208-1266, <https://doi.org/10.1594/PANGAEA.783079>, type: dataset publication series, 2010a.
- 860 Murphy, B. H., Farley, K. A., and Zachos, J. C.: An extraterrestrial ³He-based timescale for the Paleocene–Eocene thermal maximum (PETM) from Walvis Ridge, IODP Site 1266, *Geochimica et Cosmochimica Acta*, 74, 5098–5108, <https://doi.org/10.1016/j.gca.2010.03.039>, 2010b.
- Muscente, A. D., Prabhu, A., Zhong, H., Eleish, A., Meyer, M. B., Fox, P., Hazen, R. M., and Knoll, A. H.: Quantifying ecological impacts of mass extinctions with network analysis of fossil communities, *Proceedings of the National Academy of Sciences*, 115, 5217–5222, <https://doi.org/10.1073/pnas.1719976115>, publisher: Proceedings of the National Academy of Sciences, 2018.
- 865 Nanthamornphong, A. and Carver, J. C.: Test-Driven Development in scientific software: a survey, *Software Quality Journal*, 25, 343–372, <https://doi.org/10.1007/s11219-015-9292-4>, 2017.
- Norris, R. D. and Röhl, U.: Carbon cycling and chronology of climate warming during the Palaeocene/Eocene transition, *Nature*, 401, 775–778, <https://doi.org/10.1038/44545>, publisher: Nature Publishing Group, 1999.

- 870 Paola, C., Ganti, V., Mohrig, D., Runkel, A. C., and Straub, K. M.: Time not our time: physical controls on the preservation and measurement of geologic time, *Annual Review of Earth and Planetary Sciences*, 46, 409–438, <https://doi.org/10.1146/annurev-earth-082517-010129>, publisher: Annual Reviews, 2018.
- Parnell, A. C. and Gehrels, W. R.: Using chronological models in late Holocene sea-level reconstructions from salt-marsh sediments, pp. 500–513, John Wiley and Sons, Ltd, <https://doi.org/10.1002/9781118452547.ch32>, section: 32 _eprint:
- 875 <https://onlinelibrary.wiley.com/doi/pdf/10.1002/9781118452547.ch32> DOI: 10.1002/9781118452547.ch32, 2015.
- Percival, L. M. E., Davies, J. H. F. L., Schaltegger, U., De Vleeschouwer, D., Da Silva, A.-C., and Föllmi, K. B.: Precisely dating the Frasnian–Famennian boundary: implications for the cause of the Late Devonian mass extinction, *Scientific Reports*, 8, 9578, <https://doi.org/10.1038/s41598-018-27847-7>, publisher: Nature Publishing Group, 2018.
- Punyasena, S. W., Jaramillo, C., de la Parra, F., and Du, Y.: Probabilistic correlation of single stratigraphic samples: A generalized approach
- 880 for biostratigraphic data, *AAPG Bulletin*, 96, 235–244, <https://doi.org/10.1306/06201111026>, 2012.
- Pálffy, J.: Applications of quantitative biostratigraphy in chronostratigraphy and time scale construction, *Stratigraphy*, 4, 195–199, 2007.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org/>, 2023.
- Racki, G., Rakociński, M., Marynowski, L., and Wignall, P. B.: Mercury enrichments and the Frasnian-Famennian biotic crisis: A volcanic
- 885 trigger proved?, *Geology*, 46, 543–546, <https://doi.org/10.1130/G40233.1>, 2018.
- Raup, D. M. and Sepkoski, J. J.: Mass Extinctions in the Marine Fossil Record, *Science*, 215, 1501–1503, <https://doi.org/10.1126/science.215.4539.1501>, publisher: American Association for the Advancement of Science, 1982.
- Röhl, U., Bralower, T., Norris, R., and Wefer, G.: New chronology for the late Paleocene thermal maximum and its environmental implications, *Geology*, 28, 927–930, [https://doi.org/10.1130/0091-7613\(2000\)28<927:NCFTLP>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<927:NCFTLP>2.0.CO;2), 2000.
- 890 Röhl, U., Westerhold, T., Bralower, T. J., and Zachos, J. C.: On the duration of the Paleocene-Eocene thermal maximum (PETM), *Geochemistry, Geophysics, Geosystems*, 8, <https://doi.org/10.1029/2007GC001784>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2007GC001784>, 2007.
- Sadler, P. M.: Sediment Accumulation Rates and the Completeness of Stratigraphic Sections, *The Journal of Geology*, 89, 569–584, <https://doi.org/10.1086/628623>, publisher: The University of Chicago Press, 1981.
- 895 Schumer, R. and Jerolmack, D. J.: Real and apparent changes in sediment deposition rates through time, *Journal of Geophysical Research: Earth Surface*, 114, <https://doi.org/10.1029/2009JF001266>, _eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2009JF001266>, 2009.
- Scott, R. W.: Chronostratigraphic Database for Upper Cretaceous Oceanic Red Beds (CORBs), in: *Cretaceous Oceanic Red Beds: Stratigraphy, Composition, Origins, and Paleooceanographic and Paleoclimatic Significance*, edited by Hu, X., Wang, C., Scott, R. W., Wagreich, M., and Jansa, L., vol. 91, p. 0, SEPM (Society for Sedimentary Geology), ISBN 978-1-56576-135-3, <https://doi.org/10.2110/sepmssp.091.035>, 2009.
- 900 Sinnesael, M., De Vleeschouwer, D., Zeeden, C., Batenburg, S. J., Da Silva, A.-C., de Winter, N. J., Dinarès-Turell, J., Drury, A. J., Gambacorta, G., Hilgen, F. J., Hinnov, L. A., Hudson, A. J. L., Kemp, D. B., Lantink, M. L., Laurin, J., Li, M., Liebrand, D., Ma, C., Meyers, S. R., Monkenbusch, J., Montanari, A., Nohl, T., Pälke, H., Pas, D., Ruhl, M., Thibault, N., Vahlenkamp, M., Valero, L., Wouters, S., Wu, H., and Claeys, P.: The Cyclostratigraphy Intercomparison Project (CIP): consistency, merits and pitfalls, *Earth-Science Reviews*, 199, 102965, <https://doi.org/10.1016/j.earscirev.2019.102965>, 2019.

- Sluijs, A., Bowen, G., Brinkhuis, H., Lourens, L., and Thomas, E.: The Palaeocene–Eocene Thermal Maximum super greenhouse: biotic and geochemical signatures, age models and mechanisms of global change, in: *Deep-Time Perspectives on Climate Change: Marrying the Signal from Computer Models and Biological Proxies.*, edited by Williams, M., Haywood, A., Gregory, F., and Schmidt, D., pp. 323–349, The Geological Society of London on behalf of The Micropalaeontological Society, <https://doi.org/10.1144/TMS002.15>, dOI: 10.1144/TMS002.15, 2007.
- Storey, M., Duncan, R. A., and Swisher, C. C.: Paleocene-Eocene Thermal Maximum and the Opening of the Northeast Atlantic, *Science*, 316, 587–589, <https://doi.org/10.1126/science.1135274>, publisher: American Association for the Advancement of Science, 2007.
- Takayanagi, M. and Ozima, M.: Temporal variation of $^3\text{He}/^4\text{He}$ ratio recorded in deep-sea sediment cores, *Journal of Geophysical Research: Solid Earth*, 92, 12 531–12 538, <https://doi.org/10.1029/JB092iB12p12531>, _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/JB092iB12p12531>, 1987.
- Thompson, J. B. and Newton, C. R.: Late Devonian Mass Extinction: Episodic Climatic Cooling or Warming?, in: *Devonian of the World: Proceedings of the 2nd International Symposium on the Devonian System — Memoir 14, Volume III: Paleontology, Paleoecology and Biostratigraphy*, pp. 29–34, https://archives.datapages.com/data/cspg_sp/data/014/014003/29_cspgsp014c0029.htm, publisher: CSPG Special Publications, 1988.
- Tipper, J. C.: Measured rates of sedimentation: What exactly are we estimating, and why?, *Sedimentary Geology*, 339, 151–171, <https://doi.org/10.1016/j.sedgeo.2016.04.003>, 2016.
- Tobin, T. S., Ward, P. D., Steig, E. J., Olivero, E. B., Hilburn, I. A., Mitchell, R. N., Diamond, M. R., Raub, T. D., and Kirschvink, J. L.: Extinction patterns, $\delta^{18}\text{O}$ trends, and magnetostratigraphy from a southern high-latitude Cretaceous–Paleogene section: Links with Deccan volcanism, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 350–352, 180–188, <https://doi.org/10.1016/j.palaeo.2012.06.029>, 2012.
- Tomašových, A., Gallmetzer, I., Haselmair, A., Kaufman, D. S., Kralj, M., Cassin, D., Zonta, R., and Zuschin, M.: Tracing the effects of eutrophication on molluscan communities in sediment cores: outbreaks of an opportunistic species coincide with reduced bioturbation and high frequency of hypoxia in the Adriatic Sea, *Paleobiology*, 44, 575–602, <https://doi.org/10.1017/pab.2018.22>, 2018.
- Tomašových, A., Gallmetzer, I., Haselmair, A., and Zuschin, M.: Inferring time averaging and hiatus durations in the stratigraphic record of high-frequency depositional sequences, *Sedimentology*, 69, 1083–1118, <https://doi.org/10.1111/sed.12936>, 2022.
- Trachsel, M. and Telford, R. J.: All age–depth models are wrong, but are getting better, *The Holocene*, 27, 860–869, <https://doi.org/10.1177/0959683616675939>, 2017.
- Trayler, R. B., Schmitz, M. D., Cuitiño, J., Kohn, M. J., Bargo, M. S., Kay, R. F., Strömberg, C. A., and Vizcaíno, S. F.: An improved approach to age-modeling in deep time: Implications for the Santa Cruz Formation, Argentina, *GSA Bulletin*, 132, 233–244, <https://doi.org/10.1130/B35203.1>, 2019.
- Trayler, R. B., Meyers, S. R., Sageman, B. B., and Schmitz, M. D.: Bayesian Integration of Astrochronology and Radioisotope Geochronology, <https://doi.org/10.5194/gchron-2023-22>, dOI: 10.5194/gchron-2023-22, 2023.
- Vahlenkamp, M., De Vleeschouwer, D., Batenburg, S. J., Edgar, K. M., Hanson, E., Martinez, M., Pälke, H., MacLeod, K. G., Li, Y.-X., Richter, C., Bogus, K., Hobbs, R. W., and Huber, B. T.: A lower to middle Eocene astrochronology for the Mentelle Basin (Australia) and its implications for the geologic time scale, *Earth and Planetary Science Letters*, 529, 115 865, <https://doi.org/10.1016/j.epsl.2019.115865>, 2020.

- Wichern, N. M. A., Bialik, O. M., Nohl, T., Percival, L. M. E., Becker, R. T., Kaskes, P., Claeys, P., and De Vleeschouwer, D.: Astronomically paced climate and carbon cycle feedbacks in the lead-up to the Late Devonian Kellwasser Crisis, *Climate of the Past*, 20, 415–448, <https://doi.org/10.5194/cp-20-415-2024>, 2024.
- Wilkinson, B. H.: Precipitation as Meteoric Sediment and Scaling Laws of Bedrock Incision: Assessing the Sadler Effect, *The Journal of Geology*, 123, 95–112, <https://doi.org/10.1086/681588>, 2015.
- Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O., Edmunds, S., Evelo, C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S., Heringa, J., 't Hoen, P. A. C., Hoof, R., Kuhn, T., Kok, R., Kok, J., Lusher, S. J., Martone, M. E., Mons, A., Packer, A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T., Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A., Wittenburg, P., Wolstencroft, K., Zhao, J., and Mons, B.: The FAIR Guiding Principles for scientific data management and stewardship, *Scientific Data*, 3, 1–9, <https://doi.org/10.1038/sdata.2016.18>, number: 1 Publisher: Nature Publishing Group, 2016.
- Young, R., Walanus, A., Goslar, T., van Geel, B., Ralska-Jasiewiczowa, M., and Wijmstra, T. A.: Test of an equal taxon–weight modification of Middelborg’s pollen density dating on data from varved sediments of Lake Gościąg, Poland, *Review of Palaeobotany and Palynology*, 104, 213–237, [https://doi.org/10.1016/S0034-6667\(98\)00060-8](https://doi.org/10.1016/S0034-6667(98)00060-8), 1999.
- Zeeden, C., Hilgen, F. J., Hüsing, S. K., and Lourens, L. L.: The Miocene astronomical time scale 9–12 Ma: New constraints on tidal dissipation and their implications for paleoclimatic investigations, *Paleoceanography*, 29, 296–307, <https://doi.org/10.1002/2014PA002615>, [_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/2014PA002615](https://onlinelibrary.wiley.com/doi/pdf/10.1002/2014PA002615), 2014.