

# Validation and uncertainty quantification of three state-of-the-art ammonia surface exchange schemes using NH<sub>3</sub> flux measurements in a dune ecosystem

5 Tycho Jongenelen<sup>1</sup>, Margreet van Zanten<sup>2,5</sup>, Enrico Dammers<sup>1,3</sup>, Roy Wichink Kruit<sup>2</sup> Arjan Hensen<sup>4</sup>,  
Leon Geers<sup>3</sup>, Jan Willem Erisman<sup>1</sup>

<sup>1</sup>Institute of Environmental Science (CML), Leiden University, Leiden, 2333 CC, Netherlands

<sup>2</sup>National Institute for Public Health and the Environment (RIVM), Bilthoven, 3720 BA, Netherlands

<sup>3</sup>Netherlands Organisation for Applied Scientific Research (TNO), department Air Quality and Emissions Research, Utrecht, 3584 CB, Netherlands

10 <sup>4</sup>Netherlands Organisation for Applied Scientific Research (TNO), P.O. Box 15, 1755 ZG, Petten, the Netherlands

<sup>5</sup>Meteorology and Air Quality Group, Wageningen University & Research, P.O. Box 47, 6700 AA Wageningen, the Netherlands

15 *Correspondence to:* Tycho Jongenelen ([t.jongenelen@cml.leidenuniv.nl](mailto:t.jongenelen@cml.leidenuniv.nl))

**Abstract.** Deposition of reactive nitrogen causes detrimental environmental effects, including biodiversity loss, eutrophication, and soil acidification. Measuring and modeling the biosphere-atmosphere exchange of ammonia, the most abundant reduced nitrogen species, is complex due to its high reactivity and solubility, often leading to systematic discrepancies between model predictions and observations. This study aims to determine whether three state-of-the-art exchange schemes for NH<sub>3</sub> can accurately model NH<sub>3</sub> exchange in a dune ecosystem (Solleveld) and detect factors causing the uncertainties in these schemes. The selected schemes are DEPAC by Van Zanten et al. (2010), and the schemes by Massad et al. (2010) and Zhang et al. (2010). Validation against one year of gradient flux measurements revealed that the Zhang scheme represented the NH<sub>3</sub> deposition at Solleveld best, whereas the DEPAC scheme overestimated the total deposition while the Massad scheme underestimated the total deposition. Yet, none of these schemes captured the emission events at Solleveld, pointing to considerable uncertainty in the compensation point parameterization and possibly in the modeling of NH<sub>3</sub> desorption processes from wet surface layers. The sensitivity analysis further reinforced these results, showing how uncertainty in essential model parameters in the external resistance ( $R_w$ ) and compensation point parameterization propagated into diverging model outcomes. These outcomes underscore the need to improve our mechanistic understanding of surface equilibria represented by compensation points, including the adsorption-desorption mechanism at the external water layer, and specific recommendations are provided for future modeling approaches and measurement setups to support this goal.

20  
25  
30



## 1 Introduction

35 Nitrogen deposition is known to have detrimental effects on the environment, such as biodiversity loss, greenhouse gas emissions, soil acidification, eutrophication, and particulate matter formation (Galloway et al., 2003). Nitrogen deposition consists of both oxidized and reduced nitrogen components. Measuring dry  $\text{NH}_3$  deposition is challenging as  $\text{NH}_3$  is very reactive and has a high solubility (Erisman & Wyers, 1993). Moreover,  $\text{NH}_3$  is a weak absorber of light in both infrared and ultraviolet, which complicates remote-sensing measurements (Shephard & Cady-Pereira, 2015).

40 The development of  $\text{NH}_3$  exchange schemes is mainly based on flux measurements with the gradient technique (Flechard et al., 2013) and more recently using the eddy covariance technique at a single height (Famulari et al., 2004; Swart et al., 2023; Wang et al., 2021). This has allowed for the parameterization of the exchange processes for different ecosystem types and also enabled the parameterization of multi-layer models and bi-directional exchange schemes (Nemitz et al., 2001). The exchange of  $\text{NH}_3$  is a complex and dynamic process and is determined by a multitude of processes such as micrometeorology, soil properties, agricultural management practices, and vegetation growth which are still only partly understood (Flechard et al., 45 2013).

Surface exchange schemes for  $\text{NH}_3$  have three different pathways: (i) stomatal uptake into the leaf apoplasts, (ii) external deposition through solution of  $\text{NH}_3$  in raindrops in water layers on vegetation surfaces, and (iii) surface deposition onto the soil surface. While the apoplasts, the external water layer, and the soil (hereafter called: *exchange pathways*) can act as a sink for  $\text{NH}_3$ , they can also be a source under certain conditions, making the surface exchange of  $\text{NH}_3$  bi-directional (Farquhar et al., 50 1980). Most surface exchange schemes utilize the resistance analogy to calculate and describe factors that influence the rate of atmospheric deposition toward the biosphere (Wesely & Hicks, 1977). To account for the bi-directional nature of  $\text{NH}_3$ , compensation points are introduced that act as an effective  $\text{NH}_3$  concentration within the canopy. An intercomparison of several  $\text{NH}_3$  exchange schemes by Flechard et al. (2011) has shown that under identical meteorological and vegetative circumstances, the dry deposition velocities of different models vary by a factor of 2 – 3, pointing to a large uncertainty.

55 There is a need to improve the  $\text{NH}_3$  surface exchange schemes to advance our understanding of the transport and deposition of reduced nitrogen. Therefore, this study aims to determine and quantify the uncertainties in three state-of-the-art  $\text{NH}_3$  exchange schemes, which to our knowledge, has only scarcely been done (e.g. Hoogerbrugge et al., 2024; Bytnerowicz et al., 2015). Comparative studies between bi-directional exchange schemes can help to improve models but are rare (e.g. Flechard et al., 2011; Neiryck & Ceulemans, 2008; Schrader et al., 2016; Wen et al., 2014).

60 Three operational exchange schemes are selected: the DEPAC scheme (Van Zanten et al., 2010), the Massad scheme by Massad et al. (2010) and the Zhang scheme by Zhang et al. (2010). We use a one-year hourly flux measurement dataset from dune area Solleveld (Vendel et al. 2023) to test and analyze the differences between the three models. After presenting the results, the potential shortcomings of current  $\text{NH}_3$  exchange schemes are discussed, and several recommendations for future  $\text{NH}_3$  exchange schemes and measurement campaigns are provided that can contribute to lowering the model uncertainty and 65 improving the understanding of the biosphere-atmosphere exchange of  $\text{NH}_3$ .



## 2 Materials and Methods

### 2.1 Experimental site and setup

The measurements took place in the dune ecosystem of Solleveld (52°2"N, 4°11"E), in the Netherlands (Vendel et al., 2023). The habitat type is grey dunes and the measurement site consisted of sand sedge (*Carex arenaria*), moss, and lichens. To the northeast of the site is a pond surrounded by reeds. The measurements with the GRAdient Ammonia High Accuracy Monitor (GRAHAM; Wichink Kruit et al., 2007) took place between September 2014 and September 2015. The measurements were performed at three heights: 0.8 m, 1.7 m, and 3.8 m. The GRAHAM instrument is a continuous wet denuder system with a low random bias of 1.9%, a low detection limit of 0.1  $\mu\text{g m}^{-3}$ , and a time resolution of 10 minutes averaged to half-hourly measurements. Besides the  $\text{NH}_3$  flux measurements, the wind speed and direction were measured with the 3D sonic anemometer Windmaster Pro, Gill, at 5.15 m, from which the friction velocity  $u_*$ , and Monin-Obukhov length  $L$  were inferred. Based on the  $\text{NH}_3$  concentrations measured and (partly) on-site meteorological measurements, the  $\text{NH}_3$  exchange flux was inferred; for further details, we refer to Vendel et al. (2023).

Meteorological input variables such as temperature, relative humidity (RH), and radiation were measured at Valkenburg weather station by the KNMI (the Royal Netherlands Meteorological Institute), located approximately 20 kilometers northeast of Solleveld. The DEPAC and Massad scheme also required  $\text{SO}_2$  concentration data to account for co-deposition which is the enhanced deposition of  $\text{NH}_3$  caused by  $\text{SO}_2$  deposition lowering the pH of the external leaf water (Van Hove et al., 1989; Erisman & Wyers, 1993).  $\text{SO}_2$  concentrations were not measured at Solleveld, therefore, hourly  $\text{SO}_2$  measurements from monitoring station De Zilk were used, which were measured with the Thermo model 43w  $\text{SO}_2$  analyser. De Zilk is a coastal station 35 kilometers northeast of Solleveld and shares similar environmental conditions with Solleveld, as it is situated near the sea. The  $\text{NH}_3$  and  $\text{SO}_2$  concentrations were for 95% of the measurements between 0-8  $\mu\text{g m}^{-3}$  and 0-7.5  $\mu\text{g m}^{-3}$  respectively. For a short period, the  $\text{SO}_2$  concentrations were abnormally large in September 2014, caused by volcanic eruptions in Iceland (KNMI, 2014).

### 2.2 Dry deposition modeling of $\text{NH}_3$

The three main resistances in atmospheric deposition models are the aerodynamic resistance  $R_a$ , the quasi-laminar boundary layer resistance  $R_b$  and the canopy resistance  $R_c$ . The canopy resistance is an aggregate of  $R_s$ ,  $R_w$ , and  $R_{soil}$  which are respectively the stomatal, external, and soil resistance. The stomatal resistance  $R_s$  describes the exchange of gases with the atmosphere through the stomata with apoplastic fluids (Farquhar et al., 1980). The external resistance  $R_w$  (also called  $R_{cut}$  or  $R_{ext}$ ) describes the exchange of gases with the (wet) layer on the cuticula which is controlled by leaf wetness, the  $\text{SO}_2$  concentration due to co-deposition, temperature, and the Leaf Area Index (LAI) (Erisman & Wyers, 1993; Sutton et al., 1995; Van Hove et al., 1989). The soil resistance  $R_{soil}$  (also called  $R_g$ ) represents the exchange of gases with the soil (David et al., 2009; Nemitz et al., 2000).



To model the bi-directional exchange of  $\text{NH}_3$ , exchange schemes use canopy compensation points  $\chi_c$  which represents the effective  $\text{NH}_3$  concentration in the canopy. The compensation points depend on the  $\text{NH}_3/\text{NH}_4^+$  pool in the ecosystem and the temperature which controls the Henry equilibrium and the  $\text{NH}_3\text{-NH}_4^+$  dissociation equilibrium (Bates & Pinching, 1950; Dasgupta & Dong, 1986). Deposition occurs when  $\chi_c$  is below the atmospheric  $\text{NH}_3$  concentration  $\chi_a$ , while emission occurs when  $\chi_c$  is higher than  $\chi_a$ .  $\text{NH}_3$  can be emitted from the stomata when the apoplastic  $\text{NH}_4^+:\text{H}^+$  ratio is high, from the external water layers on leaf surfaces during evaporation, or from the soil or decomposing litter (Burkhardt et al., 2009; Nemitz et al., 2000; Sutton et al., 1998, 2009).


The canopy compensation point  $\chi_c$  is calculated in exchange schemes by aggregating the compartment-specific compensation points (e.g.  $\chi_s$ ,  $\chi_w$  and  $\chi_{\text{soil}}$  for the stomatal, external, and soil compensation point respectively). The compartment-specific compensation point  $\chi_i$  is a function of leaf or soil temperature and the  $\text{NH}_4^+:\text{H}^+$  ratio - also referred to as the  $\Gamma$  value (Nemitz et al., 2001). Several  $\text{NH}_3$  exchange schemes calculate one or more compartment-specific compensation points  $\chi_i$  and are therefore bi-directional (e.g. Van Zanten et al., 2010; Massad et al., 2010; Zhang et al., 2010). Studies have demonstrated that integrating compensation points into exchange schemes generally improves the agreement with measurements compared to uni-directional deposition models (Neirynek & Ceulemans, 2008; Wen et al., 2014; Wichink Kruit et al., 2012). Yet, the empirical functions in surface exchange schemes are adapted to a scarce set of flux measurements.

For a detailed overview of the parameterization of the DEPAC, Massad, and Zhang scheme, we refer to respectively Van Zanten et al. (2010), Massad et al. (2010), and Zhang et al. (2002, 2003, 2010). In Table A1, the parameterizations of all three exchange schemes are summarized. No parameterization for dunelands specifically was available in the three schemes. Therefore, “grass” was selected in the DEPAC scheme, “un-managed” and “semi-natural vegetation” in the Massad scheme, and “short grass and forbs” in the Zhang scheme. In Fig. 1, the schematics of the exchange schemes are illustrated. The DEPAC scheme has a  $\chi_s$  and  $\chi_w$  based on a three-year  $\text{NH}_3$  measurement campaign at the grasslands site Haarweg. The  $\Gamma_s$  parameterization is derived from a meta-analysis of  $\Gamma_s$  values for multiple land use classes. It is a function of temperature and the long-term average  $\text{NH}_3$  concentration (e.g. one month or one year), which serves as a memory component assuming that the apoplastic  $\Gamma_s$  is a function of the historic  $\text{NH}_3$  accumulation. The DEPAC scheme is the only operational model with a  $\chi_w$  parameterization, which is a function of temperature and the instantaneous  $\text{NH}_3$  concentration. The scheme also implicitly has a  $\chi_{\text{soil}}$ , but this is currently set to zero. Besides, soil exchange is disabled for the grasslands land use class. The Massad scheme has a  $\chi_s$  and  $\chi_{\text{soil}}$ , but the  $\chi_{\text{soil}}$  is set to 0 at Solleveld as the scheme assumes no soil exchange takes place when vegetation is present.  $\Gamma_s$  is calculated with an empirical equation derived from a meta-analysis of reported  $\Gamma_s$  values and is a function of annual nitrogen input  $N_{\text{in}}$ . The  $N_{\text{in}}$  at Solleveld was extracted from the annual deposition map published by RIVM (2023) and was 20.7 and 18.2 kg N ha<sup>-1</sup> in 2014 and 2015 respectively. The Massad scheme also has a parameterization specifically for management events (e.g. fertilizer application or grazing) but is not utilized as Solleveld is an unmanaged site. The Zhang scheme has a  $\chi_s$  and  $\chi_{\text{soil}}$  and in contrast to the DEPAC and Massad scheme, it also assumes soil exchange when vegetation is present. The scheme has a lookup table with  $\Gamma$  values for an extensive set of land use classes based on a meta-analysis of reported  $\Gamma_s$  and  $\Gamma_{\text{soil}}$  values. For some land use classes, they provide  $\Gamma$  values for ecosystems with either a low or high nitrogen



content. In this study, the low content value was used. The  $\Gamma_{\text{soil}}$  value for the land use category of short grass and forbs in the Zhang scheme is set at 2000, leading to unrealistic model output which results in extremely high emission fluxes (see Fig. 6 and Fig. A2). Wentworth et al. (2014) also observe this issue, explaining that the high  $\Gamma_{\text{soil}}$  value was derived from measurements at fertilized sites, which are not representative of unfertilized soils. Therefore, the  $\Gamma_{\text{soil}}$  value of 2000 is lowered to 395, which is an averaged value derived from the  $\Gamma_{\text{soil}}$  value reported by Massad et al. (2010) and Wentworth et al. (2014). The effects of this modification are further discussed in Section 4.4.

Moreover, all schemes used the yearly LAI curves employed in the DEPAC scheme. Vendel et al. (2023) found stronger performance for the DEPAC scheme with a minimum and maximum LAI of 0.5 and 1.0 respectively. These values were also applied in this study. Vendel et al. (2023) have also experimented with enabling the soil deposition pathway in the DEPAC scheme, but this was not implemented in this study.



**Figure 1: Schematic of the DEPAC, Massad, and Zhang scheme. In the DEPAC scheme, the  $\chi_{\text{soil}}$  and soil pathway are not modeled for grasslands. Similarly, in the Massad scheme, the  $\chi_{\text{soil}}$  and soil pathway are disabled when the LAI is higher than zero.**

### 2.3 Uncertainty analysis and sensitivity analysis

In addition to validating the  $\text{NH}_3$  exchange schemes, a comprehensive error analysis was conducted to estimate the uncertainties in input variables and model parameters. A Monte Carlo uncertainty analysis was used to propagate these uncertainties and quantify the total uncertainty in the model output. Additionally, a sensitivity analysis was performed by individually perturbing model parameters or inputs to identify the most sensitive inputs. The uncertainty of each input or variable was estimated using literature-based estimations or expert judgment. A 95% confidence interval was determined for



each input. Table A2 provides a complete list of the selected input variables and model parameters and the lower and upper bounds of the probability density functions (pdf). Additionally, the methods or assumptions for deriving these pdfs are briefly explained.

155 Three uncertainty categories were propagated in the MC analysis. First, measurement biases, both random and systematic, which arise from the measurement device itself. From this, the extent to which these biases may propagate into modeling errors could be assessed, consequently determining whether more accurate measurement equipment is necessary. Second, two systematic environmental biases were analyzed: These are differences between temperature and RH within and outside the canopy, where these variables are typically measured. This can aid in detecting potential systematic errors that need to be considered. Research by Personne et al. (2009) has shown that systematic temperature differences could lead to diverging modeling outcomes, because  $\text{NH}_3$  exchange schemes are sensitive to this temperature offset, as temperature is a key parameter for stomatal conductance and compensation point parameterization. Likewise, RH is a key variable for calculating the external resistance  $R_w$  but could strongly differ inside and outside of the canopy (Von Arx et al., 2012; Westreenen et al., 2020). Third, model biases such as empirically derived model parameters like  $\alpha$  and  $\beta$  in  $R_w$  were propagated to identify parameters that may require revision. It should be acknowledged that estimating the uncertainty for these parameters is challenging and involves some degree of subjectivity, and results can vary strongly depending on the chosen uncertainty range.

160 In total, 5000 trials were run per model, i.e. the three exchange schemes calculated the fluxes at Solleveld 5000 times with different settings. A 95% confidence interval for modeled fluxes was derived, calculated from the 2.5<sup>th</sup> and 97.5<sup>th</sup> percentile. To ensure the number of trials was sufficient, it was verified that the mean flux of the 5000 trials converged to stable values. Vendel et al. (2023) have conducted an assessment of the uncertainty associated with the GRAHAM instrument and flux calculations. They report an error margin of  $\pm 24\%$  ( $2\sigma$ ) for accumulated fluxes, which was used in this study.

### 3 Results

#### 3.1 Quantitative comparison of exchange schemes with measurements


The comparison of the measured and modeled fluxes is given in Table 1. The mean measured  $\text{NH}_3$  flux at Solleveld is  $-7.0 \text{ ng m}^{-2} \text{ s}^{-1}$ . The mean flux modeled by the Zhang scheme at  $-6.8 \text{ ng m}^{-2} \text{ s}^{-1}$  comes close to the mean measured flux. In contrast, the DEPAC scheme overestimates the mean flux at  $-15.1 \text{ ng m}^{-2} \text{ s}^{-1}$  which is more than twice the measured mean flux, and the Massad scheme strongly underestimates the mean flux at  $-2.9 \text{ ng m}^{-2} \text{ s}^{-1}$ . When comparing the hourly measured and the modeled  $\text{NH}_3$  fluxes at Solleveld, the Zhang scheme has the highest Pearson correlation at 0.62 indicating a moderately strong linear relationship. This is followed by the DEPAC scheme at 0.55, whereas the Massad scheme has a low correlation of 0.15. Table 1 also displays the performance for strong deposition ( $F_{\text{tot}} \leq 20 \text{ ng m}^{-2} \text{ s}^{-1}$ ) and moderate deposition ( $-20 \text{ ng m}^{-2} \text{ s}^{-1} > F_{\text{tot}} > 0 \text{ ng m}^{-2} \text{ s}^{-1}$ ). The accumulated measured and modeled fluxes are depicted in Figure A1. Both during moderate and strong deposition events, the Zhang scheme has the lowest RMSE. However, the DEPAC scheme had the highest correlation of 0.50 during moderate deposition, slightly better than the Zhang scheme.



**Table 1: Descriptive statistics of the measured and the modeled fluxes per exchange scheme.**

	N	GRAHAM		DEPAC scheme			Massad scheme			Zhang scheme				
		Mean flux ng m <sup>-2</sup> s <sup>-1</sup>	Mean flux ng m <sup>-2</sup> s <sup>-1</sup>	RMSE ng m <sup>-2</sup> s <sup>-1</sup>	MAE ng m <sup>-2</sup> s <sup>-1</sup>	R	Mean flux ng m <sup>-2</sup> s <sup>-1</sup>	RMSE ng m <sup>-2</sup> s <sup>-1</sup>	MAE ng m <sup>-2</sup> s <sup>-1</sup>	R	Mean flux ng m <sup>-2</sup> s <sup>-1</sup>	RMSE ng m <sup>-2</sup> s <sup>-1</sup>	MAE ng m <sup>-2</sup> s <sup>-1</sup>	R
<b>Total dataset</b>	2451	-7.0	-15.1	16.9	12.0	0.55	-2.9	14.8	9.8	0.15	-6.8	11.2	7.5	0.62
<b>F<sub>tot</sub> ≤ -20</b>	370	-33.2	-35.2	23.1	19.0	0.14	-4.4	31.8	29.0	0.09	-22.2	18.9	16.0	0.39
<b>-20 &gt; F<sub>tot</sub> &lt; 0</b>	1276	-6.8	-13.9	14.2	9.5	0.50	-2.8	7.9	5.9	0.11	-4.8	6.9	4.8	0.45
<b>F<sub>tot</sub> ≥ 0</b>	805	4.7	-7.8	17.4	12.6	-0.16	-2.4	10.2	7.2	-0.02	-2.9	11.8	7.7	-0.25


185 In total, 33% of the net fluxes were emission fluxes, occurring during daytime. The DEPAC scheme represented net emissions in only 6% of instances, compared to 12% for the Massad scheme and 23% for the Zhang scheme. Fig. 2 shows hexbin plots comparing the hourly measured fluxes with the modeled fluxes. The weak performance of all three models during emission events is clearly visible, showing no correlation with the measurements and often predicting deposition instead of emission. This indicates that the current parameterization of compensation points in all three exchange schemes cannot model the  
 190 emission events correctly in this dataset.



**Figure 2: Hexbin plot with the measured and modeled fluxes for the DEPAC, Massad, and Zhang scheme.**

Figure 3 shows the average diurnal pattern of the measured and modeled NH<sub>3</sub> flux per scheme at Sollelveld. The measured NH<sub>3</sub> flux exhibits a clear diurnal pattern where the deposition is enhanced during the evening and the night and lower during  
 195 the day.


In terms of timing, the DEPAC scheme closely matches the measured flux; however, it overestimates deposition due to a negative offset at all times. The observed deposition 'dip' at noon (i.e., when the deposition is at its lowest) is shifted to 15:00 in the DEPAC scheme. On the contrary, the Massad scheme exhibits a rather flat diurnal pattern and consistently underestimates NH<sub>3</sub> deposition at Sollelveld throughout the day. The Zhang scheme approximates the order of magnitude of  
 200 the measured fluxes well, although it does not accurately model the temporal variation of the fluxes.



**Figure 3: Measured and modeled average diurnal pattern of the  $\text{NH}_3$  flux at Solleveld in  $\text{ng m}^{-2} \text{s}^{-1}$ , per model. The shading around the lines depicts the 25-75% percentile range.**

Figure 4 displays the comparison between monthly averaged observed and modeled  $\text{NH}_3$  fluxes. The strongest deposition took place in March, attributed to the elevated  $\text{NH}_3$  concentrations stemming from agricultural fertilization practices in the region. Additionally, a minor deposition peak can be observed in November. The DEPAC scheme demonstrates a moderately good comparison with the measurements in spring, while the deposition flux is overestimated in the second half of the year, especially in November. In contrast, the Massad scheme underestimates deposition almost every month showing very little correlation with the measured yearly trend. Finally, the Zhang scheme demonstrates the strongest performance every month, accurately modeling  $\text{NH}_3$  deposition during both strong and weak deposition months.





**Figure 4: Measured and modeled monthly-averaged  $\text{NH}_3$  flux at Sollefeld in  $\text{ng m}^{-2} \text{s}^{-1}$ , per model. The outer whiskers show the quartile +  $1.5 \times$  the interquartile range. Note that outliers are not displayed.**

### 215 3.2 Model comparison

To understand the differences between the three exchange schemes, the separate pathways of the three schemes are illustrated in Fig. 5. A general observation from Fig. 5a-c is that all exchange schemes are predominantly governed by deposition towards the external leaf surface, highlighting the importance of  $R_w$  parameterization for  $\text{NH}_3$ . The dominance of the external leaf pathway is also shown in Fig. A3, showing the total accumulated flux per exchange pathway for all three exchange schemes. Contrarily, the stomatal exchange is minimal across all three schemes. In the Zhang scheme, a small but not insignificant fraction of the deposition is towards the soil. Figure 5g-i shows the conductances, which are the reciprocals of the resistances, and are more convenient to show as resistances can vary strongly throughout the day.

In the DEPAC scheme, the external flux is higher than the other two schemes, caused by the higher external conductance  $G_w$  (Fig. 5g), resulting in the highest deposition flux among all three schemes. The high  $G_w$  modeled in the DEPAC scheme aligns with the findings by Schrader et al. (2016). Both the DEPAC and Massad schemes show that  $G_w$  is highest during the evening and night and lower during the day.


In contrast,  $G_w$  in the Massad scheme is several orders of magnitude lower than in the DEPAC scheme. This difference is due to (i) a higher base resistance ( $\alpha$  or  $R_{w,\min}$ ) of  $31.5 \text{ s m}^{-1}$  compared to  $2 \text{ s m}^{-1}$  in the DEPAC scheme, (ii) a lower RH scaling parameter  $\beta$  of 8.3 compared to 12 in the DEPAC scheme and (iii) the inclusion of the  $\beta_T$  parameter which increases  $R_w$  with rising temperature.

The Zhang scheme, however, does not exhibit a strong diurnal cycle of  $G_w$ , indicating that it is less dependent on RH compared to the other two schemes. Similarly, the soil conductance  $G_{\text{soil}}$  in the Zhang scheme does not show a diurnal cycle and has a



similar magnitude to  $G_w$  (Fig. 5i). Despite this, the soil flux is significantly lower than the external flux (Fig. 5c), caused by the incorporation of a soil compensation point which counteracts soil deposition.

235 Finally, while both the DEPAC and Massad schemes use the same parameterization for  $R_s$ , the DEPAC scheme calculates stomatal emission rather than deposition, unlike the Massad scheme. This contrast is caused by the high stomatal compensation point  $\chi_s$  in the DEPAC scheme (Fig. 5d). Both functions for  $\Gamma_s$  in the DEPAC and Massad scheme are derived with a least squares fit from measurements at multiple land use types (grasslands, semi-natural, and forests) but the exact reason for the significant difference between  $\Gamma_s$  in the DEPAC and Massad scheme remains unclear.



240

**Figure 5: Diurnally averaged fluxes (first row), compensation points (second row), and conductances (third row) per exchange pathway of the DEPAC, Massad, and Zhang schemes.**



### 3.3 Uncertainty and Sensitivity Analysis

245 The results of the uncertainty analysis, presented in Table 2, indicate significant uncertainties in the modeled flux of the three exchange schemes. The total uncertainty of the DEPAC scheme ranges from a 102.0% deposition increase to a -137.7% decrease (i.e. net emission would take place), showing a large range of possible outcomes. The Massad scheme shows an even greater spread, with uncertainties ranging from a 482.8% increased deposition to a -110.3% decrease. In contrast, the Zhang scheme showed a more moderate uncertainty ranging from an 82.4% deposition increase to a -70.6% decrease. When  
250 comparing the absolute uncertainty of the mean modeled flux at Sollefeld, the DEPAC scheme showed the largest spread of -30.5;5.7 ng m<sup>-2</sup> s<sup>-1</sup>, followed by the Massad scheme with an interval of -16.9;0.3 ng m<sup>-2</sup> s<sup>-1</sup> and the Zhang scheme with an uncertainty range of -12.4; -2.0 ng m<sup>-2</sup> s<sup>-1</sup>.

The outcome of the uncertainty analysis can be further interpreted when viewed alongside the results of the sensitivity analysis, shown in Table 2. The values of the variables and parameters in this sensitivity analysis are both increased and decreased,  
255 whereas the upper and lower parameter values are based on the 2-sigma confidence bounds (see Table A2). The sensitivity of each parameter is expressed as the percentage increase or decrease of the mean flux at Sollefeld, caused by the perturbation. As stated in the methods section, three types of biases have been considered: measurement biases, systematic environmental biases, and model biases.

The measurement bias of the RH instrument has the most pronounced impact on the modeled fluxes in the DEPAC and Massad  
260 schemes, which can be explained by the strong dependency of these two schemes on the  $R_w$ . For similar reasons, the sensitivity of the LAI measurement error is more pronounced in these schemes. The measurement bias for RH and LAI was not as pronounced in the Zhang scheme, which can be explained by the slightly lesser dominance of the external pathway (see Fig. A3). Conversely, the Zhang scheme is more affected by the measurement error of  $u^*$  given its importance in both  $R_w$  and  $R_{soil}$  parameterization. The influence of the measurement bias of temperature and radiation was not significant. The effect of  
265 systematic environmental biases on in-canopy and above-canopy temperature and RH has also been analyzed. Only the Massad scheme showed strong changes in the output caused by the systematic temperature differences, as  $R_w$  is also a function of temperature due to the  $\beta_T$  parameter. The insensitivity of the DEPAC scheme to systematic temperature difference was surprising, given the strong relationship between temperature and the compensation points. However, model output has shown that  $\chi_s$  increased significantly, consequently increasing the stomatal emission flux. However, this effect was completely  
270 counteracted by the  $\chi_w$  which decreased, as  $\Gamma_w$  is inversely proportional to temperature, therefore increasing the external flux. Generally, the most sensitive parameters can be related to model biases in all three schemes with sensitivities often surpassing a  $\pm 50\%$  change in the modeled flux: In the DEPAC scheme, uncertainties in  $\Gamma_w$ ,  $\alpha$ ,  $\beta$  significantly impacted the model's output. In the Massad scheme, the most sensitive parameters are  $\beta$ ,  $\alpha$ , and  $\beta_T$ . In both the DEPAC and the Massad schemes, these sensitive parameters can, in some cases, cause the direction of the flux to shift from net deposition to net emission. In the  
275 Zhang scheme, the parameters with the highest sensitivities were  $\beta$  and  $\Gamma_{soil}$ . From this analysis, it can be concluded that (a) in all three exchange schemes, the most sensitive variables are primarily related to the exchange of  $NH_3$  via the external leaf path



and the compensation point parameterization and (b) that the DEPAC and Massad schemes are more sensitive to parameter perturbations than the Zhang scheme, as indicated by the stronger impact the perturbations have on the modeled fluxes.



280 **Table 2: Results of the uncertainty analysis (top row) and the sensitivity analysis of the three exchange schemes, showing the percentage increase/decrease of the mean modeled flux at Solleveld during the measurement period. Positive values indicate an increase in the deposition and negative values a decrease in the deposition. Percentage changes lower than -100% imply net emission. Sensitivities > 10% are in bold. The lower and upper values represent the 2-sigma confidence intervals shown in Table A2.**

Variable / Parameter	Type of bias	DEPAC		Massad		Zhang	
		Low	High	Low	High	Low	High
<b>Total uncertainty</b>	-	<b>102.0%</b>	<b>-137.7%</b>	<b>482.8%</b>	<b>-110.3%</b>	<b>82.4%</b>	<b>-70.6%</b>
Temperature	Measurement bias	-0.2%	0.2%	2.4%	-2.4%	0.7%	-0.7%
Temperature	Systematic environmental bias	-0.9%	0.9%	<b>12.0%</b>	<b>-12.0%</b>	3.1%	-3.6%
Relative Humidity	Measurement bias	<b>-15.5%</b>	<b>15.9%</b>	<b>-26.8%</b>	<b>32.9%</b>	-4.6%	3.1%
Relative Humidity	Systematic environmental bias	0.0%	4.0%	0.0%	7.8%	0.0%	0.9%
Radiation, measurement error	Measurement bias	0.1%	-0.1%	-0.3%	0.3%	-0.2%	0.2%
Friction velocity, stable conditions	Measurement bias	-3.2%	3.0%	-1.7%	1.5%	-4.5%	4.4%
Friction velocity, unstable conditions	Measurement bias	-5.7%	4.3%	-3.4%	2.5%	<b>-12.5%</b>	<b>11.7%</b>
z/L, stable conditions	Measurement bias	2.1%	-1.7%	1.0%	-0.8%	0.5%	-0.4%
z/L, unstable conditions	Measurement bias	7.2%	-1.9%	4.3%	-1.1%	3.1%	-1.0%
Roughness length, northern wind sector	Measurement bias	-0.4%	0.3%	-0.4%	0.3%	-0.1%	0.1%
Roughness length, eastern wind sector	Measurement bias	-9.8%	5.0%	-3.7%	1.7%	-3.8%	1.5%
Roughness length, southern wind sector	Measurement bias	-4.6%	2.7%	-2.9%	1.6%	-1.9%	0.9%
Roughness length, western wind sector	Measurement bias	-1.9%	1.4%	-1.7%	1.3%	-0.6%	0.4%
Canopy height	Measurement bias	0.0%	4.5%	0.0%	2.2%	0.0%	1.4%
LAI	Measurement bias	<b>-13.7%</b>	<b>10.7%</b>	<b>-11.1%</b>	9.2%	-4.7%	4.0%
SAI <sub>Haarweg</sub>	Measurement bias	<b>14.9%</b>	<b>-11.0%</b>	0.0%	0.0%	0.0%	0.0%
NH <sub>3</sub>	Measurement bias (random)	-2.7%	2.7%	-2.6%	2.6%	-5.0%	5.0%
NH <sub>3</sub>	Measurement bias (systematic)	-0.4%	0.9%	-0.4%	0.8%	-0.8%	1.6%
SO <sub>2</sub> , measurement error	Measurement bias	-0.9%	0.9%	-7.2%	6.8%	0.0%	0.0%
G <sub>s,max</sub>	Model bias	1.4%	-2.6%	-3.6%	6.4%	-2.5%	4.7%
R <sub>soil</sub> (dry)	Model bias					<b>-13.1%</b>	6.2%
R <sub>soil</sub> (wet)	Model bias					0.4%	0.0%
R <sub>ac</sub>	Model bias					2.4%	-2.0%
α (R <sub>w,min</sub> )	Model bias	<b>80.0%</b>	<b>-26.9%</b>	<b>173.9%</b>	<b>-32.1%</b>		
β	Model bias	<b>-32.7%</b>	<b>23.1%</b>	<b>216.7%</b>	<b>-64.4%</b>	<b>-18.5%</b>	<b>53.8%</b>
B <sub>T</sub>	Model bias			<b>41.5%</b>	<b>-26.4%</b>		
R <sub>w</sub> reference value (dry)	Model bias					<b>13.0%</b>	-9.1%
R <sub>w</sub> reference value (wet)	Model bias					0.7%	-0.5%
α dry	Model bias					0.0%	-0.8%
α wet	Model bias					0.0%	0.0%
Γ <sub>s</sub>	Model bias	7.2%	<b>-16.0%</b>	<b>29.6%</b>	<b>-34.2%</b>	6.9%	-7.9%
Γ <sub>w</sub>	Model bias	<b>18.3%</b>	<b>-121.6%</b>				
Γ <sub>soil</sub>	Model bias					<b>22.4%</b>	<b>-24.9%</b>



## 285 4 Discussion

The comparison of the three exchange schemes revealed that the Zhang scheme has a better performance than the DEPAC and Massad scheme when validated against the Solleveld measurement. Additionally, the exchange at the external leaf surface is a key process across all three schemes, which has also been observed in previous studies (e.g. Hansen et al., 2013; Jones et al., 2007; Neiryneck & Ceulemans, 2008; Wyers & Erisman, 1998). This also implies that the parameterization of external leaf surface exchange of  $\text{NH}_3$  is primarily responsible for the stark differences between the schemes, in accordance with results by Flechard et al. (2011). In contrast, differences in the parameterization of stomatal and soil exchange only resulted in subtle differences between the schemes. A critical issue that arises from the results is the inability of all three schemes to accurately model emission fluxes (see Fig. 2), pointing to significant shortcomings in the parameterization of compensation points. The sensitivity analysis confirms these findings, showing that the greatest uncertainties lie in the parameterization of  $R_w$  and compensation points, significantly affecting model outcomes. The results of this study align with the findings of Schrader et al. (2016), as the external flux in the DEPAC scheme was stronger than in the Massad scheme, and both studies recognize the sensitivity of  $\text{NH}_3$  exchange schemes to environmental parameters such as temperature and relative humidity. While Schrader et al. (2016) only focused on the non-stomatal parameterization of the DEPAC and Massad scheme during nighttime conditions at four sites (including the Solleveld dataset), this study extended the analysis by incorporating both stomatal and the non-stomatal parameterization and validated the schemes during nighttime and daytime conditions. Moreover, it offers a more comprehensive evaluation of schemes as the Zhang scheme is included, and uses post-processed Solleveld data, whereas Schrader et al. (2016) used preliminary data from Solleveld.

In this section, we first discuss several hypotheses as to why observed emissions were not modeled by the three schemes. The following section elaborates on how desorption processes from the external water layers are currently not modeled by any of the three exchange schemes, and how dynamic modeling approaches can resolve this issue. Finally, the influence of the uncertainty of RH and temperature measurements on the model output is discussed, and the limitations of this research are given.

### 4.1 Determining the emission source

The results of the scheme intercomparison have shown that emission is not modeled properly across all three schemes (see Fig. 2). Given the fact that soil exchange could not be ruled out at Solleveld, it was not possible to isolate the fluxes properly to derive the source of the emission. However, several hypotheses can be made regarding the source of the emission. The origin could be the desorption of  $\text{NH}_3$  from the leaf surfaces, although these emissions usually take place in the morning and would not fully explain the emissions that take place in the afternoon (see Fig. 3). These emissions could also originate from the stomata, as the temperature increase during the day would increase  $\chi_s$ . This would be in line with previous analyses on non-fertilized fields (e.g. Horváth et al., 2005; Wichink Kruit et al., 2007). Another possible source could be  $\text{NH}_3$  emission from the soil and litter layer, and this should not be ruled out as the LAI is very low at Solleveld (David et al., 2009; Hansen et al.,



2013; Sutton et al., 2009). However, given that Solleveld is a non-fertilized ecosystem, it could be expected that  $\Gamma_{\text{soil}}$  is low, thus lowering the chances of  $\text{NH}_3$  emission occurring from the soil. Yet, research on soil and litter  $\text{NH}_3$  exchange is limited, particularly in non-fertilized ecosystems such as Solleveld. Therefore,  $\Gamma_{\text{soil}}$  (and  $\Gamma_{\text{litter}}$ ) measurements similar to the work by  
320 Wentworth et al. (2014) would be valuable. Additionally, flux measurements over bare soil would be useful, which could be utilized for improving our understanding regarding the significance of soil  $\text{NH}_3$  exchange, as well as for validating and revising  $R_{\text{soil}}$  parameterization.

Auxiliary measurements that would help identify the emission sources at Solleveld, as well as in other ecosystems, include direct  $\Gamma$  measurements of the apoplasts, leaf surface water, soil, and litter to derive the compensation points per pathway.  
325 Ideally, these measurements would also be done over longer periods of time to understand seasonal and ecosystem-related influences on the  $\Gamma$  values. Moreover,  $\text{H}_2\text{O}$  or  $\text{CO}_2$  fluxes could be measured parallel to  $\text{NH}_3$  flux measurements to serve as an indication for the stomatal conductance  $G_s$  (Schrader et al., 2020; Schulte et al., 2024). Such a setup will simplify isolating and determining the flux of each exchange pathway by subtracting the stomatal flux from the total flux, aiding in determining the external flux and soil flux, which are currently subject to the greatest uncertainty. This would also enable a derivation of a  
330 parameterization for  $R_w$  based on daytime measurements. The current parameterization of  $R_w$  has been based on nighttime flux data, possibly resulting in a systematic bias for nocturnal conditions, such as lower temperatures, higher RHs, and lower turbulence. Theoretically, soil fluxes could also have been estimated from the Solleveld dataset, however, after filtering for both low RH ( $\text{RH} < 71$ ) and no solar radiation only a small subset of data ( $N = 88$ ) was left which was too little to infer reliable claims about soil  $\text{NH}_3$  exchange.

#### 335 4.2 $R_w$ and external exchange

Given the importance of  $R_w$  parameterization, the parameters involved in this process will be discussed. Whereas the DEPAC and Massad scheme use the  $\alpha \cdot e^{\beta(100-\text{RH})}$  structure to determine  $R_w$ , the Zhang scheme uses a slightly different framework. However, the latter scheme also implicitly employs a parameter that serves a similar function to the  $\alpha$  parameter (for details on the  $R_w$  equations, see Table A1). The  $\alpha$  and  $\beta$  parameters are critical for  $R_w$  parameterization in exchange schemes, however,  
340 their underlying physical properties are poorly understood. The  $\alpha$  parameter serves as a minimal  $R_w$  resistance and the  $\beta$  parameter can be described as the RH-response coefficient (Wichink Kruit, 2010). The strong sensitivity of these parameters can be seen both in the model intercomparison and the sensitivity analysis. The DEPAC scheme has the lowest  $\alpha$  value of  $2 \text{ s m}^{-1}$  whereas the Massad scheme has the highest  $\alpha$  of  $31.5 \text{ s m}^{-1}$ , which result in a much stronger external flux in the DEPAC scheme (see Fig. 5). Similarly, the sensitivity analysis revealed that propagation of the uncertainties of  $\alpha$  and  $\beta$  lead to strongly  
345 varying model outcomes (see Table 2).

To reduce the uncertainty of modeling  $\text{NH}_3$  exchange via the external leaf surface, it is essential to understand the underlying physical properties of the  $\alpha$  and  $\beta$  parameters. Massad et al. (2010) discuss that  $\alpha$  is impacted by the  $\text{SO}_2$ :  $\text{NH}_3$  ratio, while the  $\beta$  parameter is affected by leaf hygroscopicity and aerosol deposition. They also supply  $\alpha$  and  $\beta$  values derived from multiple measurement campaigns, revealing significant differences in these parameters both among different ecosystem types and



350 within each type. The Massad scheme provides ecosystem-specific  $\beta$  values, albeit with great uncertainties where the standard deviations of  $\beta$  are almost equal to the  $\beta$  value itself. Options for improved  $R_w$  parameterization include a more diverse set of  $\alpha$  and  $\beta$  values for different ecosystem types and pollution conditions or the formulation of parameterization to derive location specific  $\alpha$  and  $\beta$  values. The availability of significantly more  $\text{NH}_3$  flux measurements allows for revisiting the  $\alpha$  and  $\beta$  parameters.

355 Big steps forward can be made by including  $\chi_w$  parameterization in the Massad and Zhang scheme to account for the adsorption-desorption dynamics at the external leaf surface. Although  $\chi_w$  parameterization is already included in the DEPAC scheme, it is not able to explicitly model  $\text{NH}_3$  re-emission from the external leaf surface as  $\chi_w$  is a function of  $\chi_a$ . Wentworth et al. (2016) found that 94% of the  $\text{NH}_3$  emitted during the morning could be attributed to  $\text{NH}_3$  accumulated in the water layer overnight, indicating that  $\text{NH}_3$  re-emission is a phenomenon that should not be overlooked. Moreover, it is hard to properly interpret the specific ‘roles’ of the  $R_w$  and  $\chi_w$  parameters in the DEPAC scheme and it may be possible to rely on compensation points alone to model the external flux. This requires further research.

The formula of  $\Gamma_w$  in the DEPAC scheme is empirically inferred from three years of flux measurements at the Haarweg grasslands site (Wichink Kruit et al., 2010). This parameterization is used across all land use classes, which could introduce a systematic bias when used for other land use classes: Interspecies differences in the chemical composition of the water layer (e.g. caused by guttation) and differences in wettability caused by the wax content of the cuticula could result in variations in  $\text{NH}_3$  absorption among different plant species (Flechard et al., 1999). The effect of interspecies differences on  $\text{NH}_3$  external leaf layer exchange is understudied, but despite the knowledge gap, analyses can already be performed on existing flux datasets to validate whether the  $\chi_w$  parameterization is accurate across different ecosystem types. Note that this was challenging for the Solleveld dataset, as the external compensation point could not be inferred from the measurements due to the inability to rule out soil  $\text{NH}_3$  exchange. Moreover, the  $\chi_w$  values were inferred from micrometeorological measurements, which could contain considerable noise. Direct surface water measurements of  $\Gamma_w$  are scarce (Burkhardt et al., 2009; Sutton, Fowler, et al., 1995; Wentworth et al., 2016), and for the development and calibration of more accurate parameterization of  $\chi_w$ , measurements of  $\Gamma_w$  with a high temporal resolution would be valuable. Additionally, this dataset would be useful to infer whether micrometeorological inference is an accurate method for estimating  $\Gamma_w$  values.

375 Given the challenges of modeling  $\text{NH}_3$  exchange with the external leaf surface, we suggest moving beyond the static  $R_w$  parameterization by adopting a more mechanistic and dynamic modeling approach. Here, the inclusion of a memory effect could, for example, aid in modeling the  $\text{NH}_3$  re-emission that is often observed in the morning, as night-time  $\text{NH}_3$  reservoirs in water layers are depleted, as discussed earlier in this section. Dynamic models for  $\text{NH}_3$  exchange have already been developed (Flechard et al., 1999; Sutton et al., 1998), which consider preceding fluxes, an adsorption charge, and leaf surface chemistry which moderates  $\text{NH}_3$  solution due to saturation effects. Neiryck and Ceulemans (2008) found that the accuracy of the dynamic model from Sutton et al. (1998) had a better performance than a uni-directional model for  $R_w$  and was also able to model emission events. The latter is promising, as  $\text{NH}_3$  desorption from the external leaf layer is not modeled by any of the three exchange schemes in this study. The downside of the dynamic approach is that it requires additional input data such as



surface water pH, which is generally unavailable, however, initial efforts can be made to estimate surface water pH. As  
385 mentioned previously, direct surface water measurements of  $\Gamma_w$  would be valuable for this purpose.

#### 4.3 Relative humidity and temperature biases

In the DEPAC and Massad schemes, RH is a crucial parameter for estimating  $R_w$ . The sensitivity analysis for RH showed a  
significant impact on the mean modeled flux in both schemes, with changes of approximately  $\pm 15\%$  and  $\pm 30\%$ , respectively  
(see Table 2). Therefore, precise RH equipment is essential for providing accurate input when modeling the  $\text{NH}_3$  fluxes and  
390 for properly analyzing  $\text{NH}_3$  flux data when developing new parameterization. Although the influence of systematic differences  
in RH was analyzed in the sensitivity analysis, the effect on the modeled  $\text{NH}_3$  flux was small.

The influence of the random error of the temperature measurement equipment of  $\pm 0.1^\circ\text{C}$  has also been analyzed but remained  
small across all three exchange schemes. In contrast, the influence of systematic temperature changes inside and outside of the  
canopy was significant in the Massad scheme, which can be attributed to the inclusion of the  $\beta_T$  parameter in  $R_w$ . Studies on  
395 typical temperature and RH profiles in dune ecosystems were not found and the systematic uncertainties therefore had to be  
estimated. Hence, it is recommended to perform  $\text{NH}_3$  flux measurements together with RH or temperature profile  
measurements for different vegetation types, to determine the influence of neglecting these systematic differences in  $\text{NH}_3$  flux  
modeling.

Personne et al. (2009) elaborated further on these systematic temperature differences and coupled a deposition scheme to an  
400 energy budget model that explicitly models in-canopy temperatures. Their results indicate that systematic temperature  
variations should be explicitly considered in parameterization, particularly due to the exponential relationship between  
compensation points and temperature, as well as the relationship between stomatal conductance and temperature. However,  
while the sensitivity analysis shows that the Massad scheme is significantly affected by the systematic temperature difference,  
the net effect on the modeled flux is limited in the Zhang scheme and nearly zero in the DEPAC scheme. It must be noted that  
405 the uncertainty range used to test the sensitivity to systematic temperature difference of  $\pm 1^\circ\text{C}$  was fairly conservative.  
Personne et al. (2009) report on temperature differences exceeding more than  $4^\circ\text{C}$ ; propagating such temperature differences  
would likely lead to stronger variations in modeled fluxes.

#### 4.4 Methodological constraints

Several limitations should be taken into account. First, the results and implications in this study are drawn from measurements  
410 at a dune ecosystem and therefore, it remains an open question whether these results apply to other ecosystems as well. It is  
recommended to extend this study across different ecosystem types, similar to analyses by Flechard et al. (2011) and Schrader  
et al. (2016).


Second, this study did not discuss the influence and contribution of soil exchange, as it was challenging to isolate the soil flux  
from the stomatal and external flux. This limitation prevented the ability to perform a deeper mechanistic analysis at Sollefeldt,  
415 where each exchange pathway could be quantitatively evaluated. This also implies that although the exchange schemes signal



that the external flux is the most dominant flux at Solleveld, consistent with previous research (Hansen et al., 2013; Jones et al., 2007; Neiryck & Ceulemans, 2008; Wyers & Erisman, 1998), this does not necessarily mean it was the most dominant flux in reality. Incorporating auxiliary measurements of CO<sub>2</sub> or H<sub>2</sub>O fluxes along with  $\Gamma_s$ ,  $\Gamma_w$ ,  $\Gamma_{soil}$  and  $\Gamma_{litter}$ , as discussed in Sect. 4.3, would address these issues. Consequently, this would enable the validation of  $R_w$  parameterization and evaluate  
420 whether a soil pathway should be incorporated in exchange schemes for dune ecosystems, as done in the Zhang scheme. Especially in ecosystems with low LAIs such as Solleveld, the contribution of soil exchange should be understood better. Moreover, this would aid in identifying the emission sources, which are currently not accounted for by any of the three schemes.

Third, several state-of-the-art models such as SURFATM-NH<sub>3</sub> (Personne et al., 2009), CMAQ-EPIC (Pleim et al., 2019) and  
425 the dynamic model by Sutton et al. (1998) were not included in this study, as they required additional input parameters which were unavailable or difficult to estimate. This touches upon the trade-off between a model's usability and accuracy, as more complex models might be more accurate, but require extra input data which is often not readily available. Fourth, specific parameterization for dunelands does not exist and grasslands or semi-natural vegetation parameterization has been used across all schemes instead. Yet systematic differences between these land use classes and dunelands can lie in the LAI, the apoplastic  
430 NH<sub>4</sub><sup>+</sup>: H<sup>+</sup> ratio, the wettability of plant surfaces and the potential presence of soil NH<sub>3</sub> exchange. This also calls for expanding available land use classes in all three schemes (e.g. dunelands, heathlands, and moorlands) and conducting additional measurements to develop new parameterization. Fifth, due to the temporal resolution of the GRAHAM instrument of approximately 30 minutes (Wichink Kruit et al., 2010), and further aggregation to hourly averages in this study, sudden temporal features such as emission or deposition pulses can be smoothed out. This limitation can hinder the ability to accurately  
435 study diurnal dynamics and detect short-lived events such as emission pulses. Sixth, some meteorological input data such as temperature, RH, and radiation have not been measured at Solleveld but at a meteorological station at Valkenburg which is approximately 20 km to the northeast, which can introduce inaccuracies in the model output due to local differences between the two locations. Seventh, although the uncertainty ranges for each input variable or model parameter defined for the uncertainty and sensitivity analysis were assessed with great care, decisions such as defining these ranges and the distribution  
440 type (i.e. normal, uniform, etc.) inevitably involved a degree of subjectivity which could impact the outcome of the analyses. Moreover, the uncertainty analysis only accounts for the uncertainty in processes or variables that are explicitly included in the schemes, meaning the analysis will not capture the uncertainty of these excluded processes.

Eighth, as stated in the methods section, the  $\Gamma_{soil}$  value in the Zhang scheme has been lowered from 2000 to 395. We acknowledge this is a modification to the Zhang scheme which strongly alters the modeled output of the scheme; however, it  
445 is justifiable given the mismatch between the original parameterization and the empirical findings by Wentworth et al. (2014) for unfertilized ecosystems. As shown in Fig. 6, the unmodified Zhang scheme (purple line) would strongly underestimate the deposition at Solleveld. The effect of the  $\Gamma_{soil}$  adaptation on the average diurnal cycle is shown in Fig. A2.



450 **Figure 6: Accumulated deposition at Solleveld during the measurement period, including the accumulated flux of the unmodified Zhang scheme with a  $\Gamma_{soil}$  value of 2000.**

Finally, the parameters in the surface exchange schemes stem from flux measurements and are empirically translated into parameterization. That means that generalization to different environmental circumstances or pollution climates can give different outcomes, as demonstrated by the differing results per exchange scheme in this study. Thus, more flux measurements across a range of environmental conditions are necessary.

## 455 **5 Conclusion**

This study aimed to determine the accuracy of three state-of-the-art  $\text{NH}_3$  exchange schemes in a dune ecosystem and to identify the key factors that contribute to model uncertainties. The results of this study indicate that the performance of the exchange schemes differed significantly at the Solleveld dune site: both on hourly and monthly timescales. The Zhang scheme accurately models the  $\text{NH}_3$  exchange, whereas the DEPAC and Massad scheme respectively overestimate and underestimate the total deposition. However, the Zhang scheme does not capture the average daily cycle well: the DEPAC scheme captures this most  
460 realistically. The results also reveal that the exchange with the external leaf layer is the most important exchange pathway in all exchange schemes, which is in coherence with previous studies (e.g. Erisman & Wyers, 1993; Burkhardt et al., 2009). A serious flaw identified among all three models is that the frequent emission events at Solleveld are poorly modeled, implying that the compensation point parameterization is ineffective. The sensitivity analysis of individual model parameters  
465 demonstrates that the biggest portion of the uncertainty can be attributed to the model uncertainty in the  $R_w$  parameterization



and the compensation point modeling. The findings of these analyses provide additional quantification of the model uncertainties and corroborate the findings of Schrader et al. (2016), who also highlighted the difficulties in modeling  $R_w$ . Additionally, the sensitivity analysis has shown that the random bias in RH measurements can lead to strong fluctuations in the modeled fluxes in the DEPAC and Massad schemes. The effect of propagating systematic temperature differences inside and outside the canopy only led to strong effects on the modeled flux in the Massad scheme. Similarly, the net effect of a potentially higher RH in the canopy was minimal in all three schemes.


To address these uncertainties, we recommend conducting additional external leaf water measurements to quantify  $\Gamma_w$  to better understand the adsorption-desorption processes taking place at the wet leaf surfaces and to calibrate and improve the parameterization of  $\Gamma_w$ . Furthermore, auxiliary measurements of  $H_2O$  or  $CO_2$  fluxes as a proxy for stomatal conductance can aid in isolating the stomatal, external, and soil fluxes from each other, making it easier to estimate the contribution per exchange pathway. Moreover, although the external leaf pathway is identified as a crucial pathway in  $NH_3$  exchange, the soil pathway remains understudied. Additional measurements of  $\Gamma_{soil}$  and measurements over bare soil to validate  $R_{soil}$  parameterization would be viable for reducing the modeling uncertainty.

Future research should focus on improving our understanding of the mechanisms controlling  $NH_3$  exchange with the external leaf layer, as this process is pivotal for accurate  $NH_3$  exchange modeling. Environmental factors influencing the  $\alpha$  and  $\beta$  parameters here are key. Additionally, despite the relevance of  $NH_3$  desorption from the external leaf surface, it is not explicitly included in either of the three exchange schemes in this study. Dynamic models simulating external leaf surface fluxes can be an effective alternative to the current  $R_w$  parameterization methods, and promising work in dynamic modeling of leaf surface  $NH_3$  exchange (Sutton et al., 1998; Flechard et al., 1999; Burkhardt et al., 2009) should be further pursued.




485 **6 Appendices**

**Appendix A. Additional figures and tables**




**Figure A 1: Accumulated deposition at Solleveld during the measurement period. The error bars at the right side of the image represent the uncertainty range (95%CI) of the models as calculated by the Monte Carlo uncertainty analysis.**

490



**Figure A 2: Modeled and observed average diurnal cycle of the Zhang scheme at Solleveld, comparing the adjusted parameterization used in this study (left) to the original parameterization with a  $\Gamma_{soil}$  of 2000 (right).**



495

**Figure A 3: Accumulated flux per exchange pathway at Solleveld per exchange scheme. Note that the DEPAC scheme models net stomatal emission, hence the positive value. The striped line depicts the total deposition measured with the GRAHAM at Solleveld.**



500 **Table A 1: Main parameterization of the DEPAC, Massad and Zhang exchange scheme for NH<sub>3</sub> exchange. Note that the parameterization is specific for grasslands in the DEPAC scheme, non-fertilized semi-natural in the Massad scheme, and short grass and forbs in the Zhang scheme.**

	DEPAC	Massad	Zhang
$F_{tot}^a$	$-\frac{\chi_a - \chi_c}{R_a + R_b}$	$-\frac{\chi_a - \chi_{(z_0)}}{R_a}$	$-\frac{\chi_a - \chi_c}{R_a + R_b}$
$F_s$	$-\frac{\chi_c - \chi_s}{R_s}$	$-\frac{\chi_c - \chi_s}{R_s}$	$-\frac{\chi_c - \chi_s}{R_s}$
$F_w$	$-\frac{\chi_c - \chi_w}{R_w}$	$-\frac{\chi_c}{R_w}$	$-\frac{\chi_c}{R_w}$
$F_{soil}^b$	$-\frac{\chi_c - \chi_{soil}}{R_{soil} + R_{inc}}$	$-\frac{\chi_{(z_0)} - \chi_{soil}}{R_{soil} + R_{inc}}$	$-\frac{\chi_{(z_0)} - \chi_{soil}}{R_{soil} + R_{inc}}$
$R_s$	$[G_s^{max} \cdot f_{PAR} \cdot f_{VPD} \cdot f_T]^{-1}$	$[G_s^{max} \cdot f_{PAR} \cdot f_{VPD} \cdot f_T]^{-1}$	$\left[ G_s^{max} \cdot f_{PAR} \cdot f_{VPD} \cdot f_T \cdot f_\psi \cdot \frac{D_i}{D_{H_2O}} \right]^{-1}$
$R_w$	$\frac{SAI}{SAI_{Haarweg}} \cdot 2 \cdot e^{\left(\frac{100-RH}{12}\right)}$	$[31.5 \cdot AR^{-1} \cdot e^{(0.120(100-RH))} e^{0.15 T}] \cdot LAI^{-0.5}$	Dry: $\text{Min} \left( 100, \frac{1000}{e^{0.03RH} LAI^{0.25} u_*} \right)$ Wet: $\text{Min} \left( 20, \frac{100}{LAI^{0.5} u_*} \right)$
$R_{soil}^b$	-	-	Dry: 200 Wet: 100
$R_{inc}^b$	-	-	$\frac{20 \cdot LAI^{0.25}}{u_*^2}$
$AR^c$	-	$\frac{2[SO_2] + [HNO_3^-] + [HCl]}{[NH_3]}$	-
$\chi_i^d$	$\frac{2.75 \cdot 10^{15}}{T + 273.15} e^{\left(\frac{-1.04 \cdot 10^4}{T+273.15}\right)} \cdot \Gamma_i$	$\frac{2.75 \cdot 10^{15}}{T + 273.15} e^{\left(\frac{-1.04 \cdot 10^4}{T+273.15}\right)} \cdot \Gamma_i$	$\frac{2.75 \cdot 10^{15}}{T + 273.15} e^{\left(\frac{-1.04 \cdot 10^4}{T+273.15}\right)} \cdot \Gamma_i$
$\Gamma_s$	$1701.4 \cdot \chi_{a,4m} (long-term) \cdot e^{-0.071T_s}$	$246 + (0.0041) + (N_{in})^{3.56}$	300
$\Gamma_w$	$1.84 \cdot 10^3 \cdot \chi_{a,4m} \cdot e^{-0.11T_s} - 850$	-	-
$\Gamma_{soil}^e$	-	-	2000

Note: All exchange schemes use the same parameterization for  $R_a$  and  $R_b$  by Wesely and Hicks (1977).

505 <sup>a</sup> The equations for  $\chi_c$  and  $\chi_{(z_0)}$  are extensive and can be found in Van Zanten et al.(2010), Massad et al. (Massad et al., 2010) and Zhang et al. (2010).

<sup>b</sup> Note that the soil compensation points are not calculated in the DEPAC and Massad scheme.

<sup>c</sup> Acidity ratio used in Massad scheme. No  $HNO_3^-$  and HCl concentration data was available for Solleveld, therefore, the alternative function proposed by Schrader et al. (2016) of  $3.5 \cdot \frac{SO_2}{NH_3}$ .



<sup>d</sup> Different formulations are used for the compensation point  $\chi_i$ . Here, the formulation by Wichink Kruit et al. (2007) is used.

510 <sup>e</sup> Note that originally the  $\Gamma_{\text{soil}}$  in the Zhang scheme is 2000, but a lower value of 395 has been implemented, which is an average of the  $\Gamma_{\text{soil}}$  value reported by Massad et al. (2010) and Wentworth et al. (2014).





**Table A 2: Input variables and model parameters and the bias categories, the associated uncertainty, the lower and upper bound values, and the chosen distribution functions. The lower and upper bounds represent the two-sigma values, forming the 95% confidence interval.**

Parameter <sup>a</sup>	Bias type	Uncertainty Range	Lower Bound	Upper Bound	Distribution Type	Source
Temperature	Instrument Bias (Random)	± 0.1	-0.2°C	+0.2°C	Normal	Brandma (2004)
Temperature	Systematic Environmental Bias	± 1	-1.0°C	+1.0°C	Uniform	Expert judgment
Relative Humidity	Instrument Bias (Random)	± 2%	-4%	4%	Normal	Ustymczuk and Giner (2011)
Relative Humidity	Systematic Environmental Bias	0 – 2%	0%	2%	Uniform	Expert judgement
Radiation	Instrument Bias (Random)	± 4.0%	-4%	4%	Uniform	Shi & Long (2002) and Stoffel (2005) as cited in Mathijssen & Knap (2021)
Friction Velocity	Instrument Bias (Random)	± 15% z/L < 0 (unstable), ± 4% z/L > 0 (stable)	-30%, -8%	+30%, +8%	Normal, Normal	Salesky and Chamecki (2012)
Stability Parameter $\zeta$ (z/L)	Instrument Bias (Random)	± 40% z/L < 0 (unstable), ± 10% z/L > 0 (stable)	-80%, -20%	+80%, +20%	Normal, Normal	Salesky and Chamecki (2012)
Roughness Length (North) <sup>k</sup>	Instrument Bias (Random)	0.12 ± 0.018	0.09	0.16	Normal	Calculated per wind sector
Roughness Length (East)	Instrument Bias (Random)	0.034 ± 0.026	0.003	0.09	Normal	Calculated per wind sector
Roughness Length (South)	Instrument Bias (Random)	0.011 ± 0.013	0.001	0.04	Normal	Calculated per wind sector
Roughness Length (West)	Instrument Bias (Random)	Discrete values ± 26.6%	10 cm	50 cm	Discrete	Expert judgement
Canopy Height LAI	Instrument Bias (Measurement)	± 26.6%	-26.6%	26.6%	Uniform	Fang et al. (2012)
SAI <sub>harveg</sub> (D) <sup>j</sup>	Instrument Bias (Random)	± 5.5%	-11%	11%	Uniform	Mooibroek et al. (2014)
SO <sub>2</sub> Concentration	Instrument Bias (Random)	± 1.9%	-3.80%	3.80%	Normal	Wichink Kruit (2010)
NH <sub>3</sub> Concentration	Instrument Bias (Random)	± 0.6%	-1.20%	1.20%	Normal	Wichink Kruit (2010)
NH <sub>3</sub> Concentration	Instrument Bias (Systematic)	0.0067 ± 0.0033	6.69E-04	0.013	Normal	Kellner et al. (1995)
Stomatal Resistance (G <sub>s,max</sub> ; D, M) <sup>c</sup>	Model Bias	0.0067 ± 0.0033	6.67E-04	0.013	Normal	Kellner et al. (1995)
Stomatal Resistance (G <sub>s,max</sub> ; Z) <sup>c</sup>	Model Bias	2 ± 0.82	0.36	3.64	Normal	-
$\alpha$ (R <sub>w,amb</sub> ) (D) <sup>d</sup>	Model Bias	31.5 ± 12.92	5.66	57.34	Normal	-
$\alpha$ (R <sub>w,amb</sub> ) (M) <sup>d</sup>	Model Bias		25		Normal	-



Parameter <sup>a</sup>	Bias Type	Uncertainty Range	Lower Bound	Upper Bound	Distribution Type	Source
$\alpha$ ( $R_{w,min}$ ) ( $Z$ ) <sup>e</sup>	Model Bias	100 ± 40, 86 (dry) 20 ± 8, 17 (wet)	18.28	181.72	Normal	-
$\beta$ ( $D$ ) <sup>d</sup>	Model Bias	12 ± 2, 06	7.88	16.12	Normal	Massad et al. (2010)
$\beta$ ( $M$ )	Model Bias	0.120 ± 0, 107	0.012	0.334	Normal	
$\beta$ ( $Z$ ) <sup>d</sup>	Model Bias	0.03 ± 0, 005	0.020	0.040	Normal	
$\beta_r$ ( $M$ )	Model Bias	0.15 ± 0, 05	0.10	0.20	Uniform	Educated guess
$R_{cut,d0}$ ( $Z$ )	Model Bias	1000 ± 100	800	1200	Normal	Educated guess
$R_{cut,w0}$ ( $Z$ )	Model Bias	100 ± 10	80	120	Normal	Educated guess
$R_{soil,dry}$ ( $Z$ ) <sup>f</sup>	Model Bias	20 ± 2	16	24	Normal	Educated guess
$R_{soil,wet}$ ( $Z$ ) <sup>f</sup>	Model Bias	1000; 200; 100	-	-	Discrete	Klein et al. (2002); Van Zanten et al. (2010)
$\Gamma_s$ ( $D$ ) <sup>g</sup>	Model Bias	± 760, 0	-	-	Discrete	Klein et al. (2002); Van Zanten et al. (2010)
$\Gamma_s$ ( $M$ ) <sup>g</sup>	Model Bias	444.4 ± 231.8 (201.4)	44.4	908.0	Normal	Massad et al. (2010)
$\Gamma_s$ ( $Z$ ) <sup>g</sup>	Model Bias	372.3 ± 194.2 (201.5)	37.2	760.7	Normal	Massad et al. (2010)
$\Gamma_w$ ( $D$ ) <sup>h</sup>	Model Bias	300 ± 156.5	30	613	Normal	Massad et al. (2010)
$\Gamma_g$ ( $Z$ ) <sup>i</sup>	Model Bias	1576 ± 788 395 ± 197.5	-	-	Normal	Educated guess
			39.5	790	Normal	Wentworth et al. (2014)

<sup>a</sup> The D, M, or Z in parenthesis denotes the DEPAC, Massad, or Zhang scheme, respectively.

<sup>b</sup> In some cases, the lower value would become negative, which is not possible (e.g. for  $G_{s,max}$  or the roughness length). In these cases, the lower value was capped at 10% of the base value.

<sup>c</sup> Kellner et al. (1995) reported a  $G_{s,max}$  of 0.008 with a standard deviation of 0.004, from which a coefficient of variation ( $CV = \mu / \sigma$ ) of 0.5 is calculated. This CV is used to define the variability space for  $G_{s,max}$  in the three exchange schemes.

<sup>d</sup> The uncertainty ranges for  $\alpha$  and  $\beta$  are calculated with data from van Hove et al. (1989) and Benner et al. (1992) which is displayed in Fig. 4 in Sutton et al. (1995). The  $\alpha$  and  $\beta$  values of 2 s  $m^{-1}$  and 14 derived by Sutton et al. (1995) for  $R_w$  are also used in the DEPAC scheme. A curve has been fitted in Python using Scipy's *curve\_fit* function, which supplied a covariance matrix, from which the standard deviations of the  $\alpha$  and  $\beta$  parameters were determined. Note that different parameters for  $\alpha$  and  $\beta$  were found, but subsequent uncertainty ranges have been scaled proportionally to the original  $\alpha$  and  $\beta$  values. From this, the  $CV_\alpha = 0.41$  and  $CV_\beta = 0.17$  were calculated. These  $CV_s$  were also used to calculate the uncertainty range of the  $\alpha$  value in the Zhang and Massad scheme.

<sup>e</sup> The Zhang scheme does not explicitly have an  $\alpha$  value but has a cut-off value of 100 s  $m^{-1}$  and 20 s  $m^{-1}$  for dry and wet conditions respectively. The  $CV_\alpha$  of 0.41 derived for the DEPAC scheme was applied to these cut-off values.

<sup>f</sup> The constant  $R_{soil}$  parameter for dry and wet conditions is randomly imputed with  $R_{soil}$  constants for dry and wet conditions used in the Zhang, DEPAC, and MATCH model (Klein et al., 2002; Van Zanten et al., 2010).



<sup>e</sup> From Table 3 in Massad et al. (2010), measured  $\Gamma_s$  values from different studies are listed together with the uncertainty range. From this, an average  $CV_{\Gamma_s} = 0.52$  is derived. The standard deviation of  $\Gamma_s$  in the DEPAC scheme is derived from the mean modeled  $\Gamma_s (= 1461.5)$ , leading to a standard deviation of 760. This  $CV_{\Gamma_s}$  is also used to derive the standard deviations of  $\Gamma_s$  in the DEPAC and Zhang scheme.

<sup>h</sup> The relative uncertainty of the  $\Gamma_w$  parameter in the DEPAC scheme is assumed here to be in the same order of magnitude as the relative uncertainty of  $\Gamma_s$  and  $\Gamma_{soil}$ . Therefore, the relative uncertainty is 50%. From the mean  $\Gamma_w$  at Sollefeld ( $= 1576$ ), a standard deviation of 788 has been derived.

<sup>i</sup> A  $CV_{\Gamma_{soil}}$  of 50% has been derived from  $\Gamma_{soil}$  data from a non-fertilized grassland (Wentworth et al., 2014). From the  $\Gamma_{soil} (= 395)$  in the Zhang scheme, a standard deviation of 197.5 has been derived.

<sup>j</sup> The leaf area index (LAI) has been altered according to Vendel et al. (2023) to a minimum LAI of 0.5 and a maximum LAI of 1.0. These values have been derived from MODIS LAI data. Fang et al. (2012) derive a relative uncertainty of 26.6% for the MODIS LAI product. For consistency, this value has also been used for the  $SAI_{Hartweg}$  parameter used in the DEPAC scheme.

<sup>k</sup> Similarly to Vendel et al. (2023), roughness lengths are wind sector dependent to account for the heterogeneous footprint of Sollefeld and are calculated with the  $u^*$  measurements from the sonic anemometer, according to Moene and van Dam (2016, p. 119). The standard deviation of the  $z_0$  per wind sector is used to describe the uncertainty in  $z_0$ .



## 7 Code availability

The code for analyzing the model output, the uncertainty, and sensitivity analysis will be published on Zenodo – but can  
545 already be requested from T. Jongenelen: [t.jongenelen@cml.leidenuniv.nl](mailto:t.jongenelen@cml.leidenuniv.nl).

## 8 Data availability

The Solleveld measurements will soon be published on Zenodo – but can already be requested from M.C. van Zanten:  
[margreet.van.zanten@rivm.nl](mailto:margreet.van.zanten@rivm.nl).

## 9 Author contribution

550 **T. Jongenelen**: Conceptualization, Methodology, Formal analysis, Software, Visualization, Writing – original draft  
preparation, Writing – review & editing. **M. C. Van Zanten**: Conceptualization, Methodology, Supervision, Writing – original  
draft preparation, Writing – review & editing. **E. Dammers**: Conceptualization, Methodology, Supervision, Writing – original  
draft preparation, Writing – review & editing. **R. Wichink Kruit**: Writing – review & editing. **A. Hensen**: Investigation,  
Writing – Review & editing. **L. F. G. Geers**: Software, Writing – Review & editing. **J. W. Erisman**: Conceptualization,  
555 Methodology, Supervision, Writing – original draft preparation, Writing – review & editing.

## 10 Competing interests

The authors declare that they have no conflict of interest.

## 11 Acknowledgements

Special thanks to Kim Vendel for her assistance in interpreting and clarifying the Solleveld measurements and to Cor Jacobs  
560 for his guidance on the Fortran code of DEPAC 1-D.

## 12 References

Bates, R. G., & Pinching, G. D. (1950). Dissociation Constant of Aqueous Ammonia at 0 to 50° from E. m. F. Studies of the  
Ammonium Salt of a Weak Acid. *Journal of the American Chemical Society*, 72(3), 1393–1396.  
<https://doi.org/10.1021/ja01159a087>



- 565 Benner, W. H., Ogorevc, B., & Novakov, T. (1992). Oxidation of SO<sub>2</sub> in thin water films containing NH<sub>3</sub>. *Atmospheric Environment. Part A. General Topics*, 26(9), 1713–1723. [https://doi.org/10.1016/0960-1686\(92\)90069-W](https://doi.org/10.1016/0960-1686(92)90069-W)
- Brandsma, T. (2004). *Parallel air temperature measurements at the KNMI-terrain in De Bilt (the Netherlands) May 2003 – April 2005*.
- Burkhardt, J., Flechard, C. R., Gresens, F., Mattsson, M., Jongejan, P. A. C., Erisman, J. W., Weidinger, T., Meszaros, R.,  
570 Nemitz, E., & Sutton, M. A. (2009). Modelling the dynamic chemical interactions of atmospheric ammonia with leaf surface wetness in a managed grassland canopy. *Biogeosciences*, 6(1), 67–84. <https://doi.org/10.5194/bg-6-67-2009>
- Bytnerowicz, A., Johnson, R. F., Zhang, L., Jenerette, G. D., Fenn, M. E., Schilling, S. L., & Gonzalez-Fernandez, I. (2015). An empirical inferential method of estimating nitrogen deposition to Mediterranean-type ecosystems: The San Bernardino Mountains case study. *Environmental Pollution*, 203, 69–88.  
575 <https://doi.org/10.1016/j.envpol.2015.03.028>
- Dasgupta, P. K., & Dong, S. (1986). Solubility of ammonia in liquid water and generation of trace levels of standard gaseous ammonia. *Atmospheric Environment (1967)*, 20(3), 565–570. [https://doi.org/10.1016/0004-6981\(86\)90099-5](https://doi.org/10.1016/0004-6981(86)90099-5)
- David, M., Loubet, B., Cellier, P., Mattsson, M., Schjoerring, J. K., Nemitz, E., Roche, R., Riedo, M., & Sutton, M. A. (2009). Ammonia sources and sinks in an intensively managed grassland canopy. *Biogeosciences*, 6(9), 1903–1915.  
580 <https://doi.org/10.5194/bg-6-1903-2009>
- Erisman, J. W., & Wyers, G. P. (1993). Continuous measurements of surface exchange of SO<sub>2</sub> and NH<sub>3</sub>; Implications for their possible interaction in the deposition process. *Atmospheric Environment. Part A. General Topics*, 27(13), 1937–1949. [https://doi.org/10.1016/0960-1686\(93\)90266-2](https://doi.org/10.1016/0960-1686(93)90266-2)
- Famulari, D., Fowler, D., Hargreaves, K., Milford, C., Nemitz, E., Sutton, M. A., & Weston, K. (2004). *Measuring eddy covariance fluxes of ammonia using tunable diode laser absorption spectroscopy*.  
585
- Fang, H., Wei, S., Jiang, C., & Scipal, K. (2012). Theoretical uncertainty analysis of global MODIS, CYCLOPES, and GLOBCARBON LAI products using a triple collocation method. *Remote Sensing of Environment*, 124, 610–621. <https://doi.org/10.1016/j.rse.2012.06.013>



- Farquhar, G. D., Firth, P. M., Wetselaar, R., & Weir, B. (1980). On the Gaseous Exchange of Anunonia between Leaves and  
590 the Environment: Determination of the Ammonia Compensation Point. *Plant Physiology*, 66(4), 710–714.
- Flechard, C. R., Fowler, D., Sutton, M. A., & Cape, J. N. (1999). A dynamic chemical model of bi-directional ammonia  
exchange between semi-natural vegetation and the atmosphere. *Quarterly Journal of the Royal Meteorological  
Society*, 125(559), 2611–2641. <https://doi.org/10.1002/qj.49712555914>
- Flechard, C. R., Massad, R. S., Loubet, B., Personne, E., Simpson, D., Bash, J. O., Cooter, E. J., Nemitz, E., & Sutton, M. A.  
595 (2013). Advances in understanding, models and parameterizations of biosphere-atmosphere ammonia exchange.  
*Biogeosciences*, 10(7), 5183–5225. <https://doi.org/10.5194/bg-10-5183-2013>
- Flechard, C. R., Nemitz, E., Smith, R. I., Fowler, D., Vermeulen, A. T., Bleeker, A., Erisman, J. W., Simpson, D., Zhang, L.,  
Tang, Y. S., & Sutton, M. A. (2011). Dry deposition of reactive nitrogen to European ecosystems: A comparison of  
inferential models across the NitroEurope network. *Atmospheric Chemistry and Physics*, 11(6), 2703–2728.  
600 <https://doi.org/10.5194/acp-11-2703-2011>
- Galloway, J. N., Aber, J. D., Erisman, J. W., Seitzinger, S. P., Howarth, R. W., Cowling, E. B., & Cosby, B. J. (2003). The  
nitrogen cascade. *BioScience*, 53(4), 341–356. [https://doi.org/10.1641/0006-3568\(2003\)053\[0341:TNC\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2003)053[0341:TNC]2.0.CO;2)
- Hansen, K., Sørensen, L. L., Hertel, O., Geels, C., Skjøth, C. A., Jensen, B., & Boegh, E. (2013). Ammonia emissions from  
deciduous forest after leaf fall. *Biogeosciences*, 10(7), 4577–4589. <https://doi.org/10.5194/bg-10-4577-2013>
- 605 Hoogerbrugge, R., Braam, M., Siteur, K., Jacobs, C., & Hazelhorst, S. (2024). *Uncertainty in the determined nitrogen  
deposition in the Netherlands. Status report 2023*. National Institute for Public Health and the Environment.  
<https://doi.org/10.21945/RIVM-2022-0085>
- Horváth, L., Asztalos, M., Führer, E., Mészáros, R., & Weidinger, T. (2005). Measurement of ammonia exchange over  
grassland in the Hungarian Great Plain. *Agricultural and Forest Meteorology*, 130(3–4), 282–298.  
610 <https://doi.org/10.1016/j.agrformet.2005.04.005>
- Jones, M. R., Leith, I. D., Raven, J. A., Fowler, D., Sutton, M. A., Nemitz, E., Cape, J. N., Sheppard, L. J., & Smith, R. I.  
(2007). Concentration-dependent NH<sub>3</sub> deposition processes for moorland plant species with and without stomata.  
*Atmospheric Environment*, 41(39), 8980–8994. <https://doi.org/10.1016/j.atmosenv.2007.08.015>



- 615 Kelliher, F. M., Leuning, R., Raupach, M. R., & Schulze, E.-D. (1995). Maximum conductances for evaporation from global  
vegetation types. *Agricultural and Forest Meteorology*, *73*(1–2), 1–16. [https://doi.org/10.1016/0168-1923\(94\)02178-](https://doi.org/10.1016/0168-1923(94)02178-M)  
M
- Klein, T., Bergström, R., & Persson, C. (2002). *Parameterization of dry deposition in MATCH*.
- KNMI. (2014). *Toename in zwaveldioxide in Nederland door vulkaanuitbarstingen IJsland 2014*. [https://www.knmi.nl/kennis-](https://www.knmi.nl/kennis-en-datacentrum/achtergrond/toename-in-zwaveldioxide-in-nederland-in-september-2014-ten-gevolge-van-de-holuhraun-vulkanische-erupties-op-ijsland)  
620 [en-datacentrum/achtergrond/toename-in-zwaveldioxide-in-nederland-in-september-2014-ten-gevolge-van-de-](https://www.knmi.nl/kennis-en-datacentrum/achtergrond/toename-in-zwaveldioxide-in-nederland-in-september-2014-ten-gevolge-van-de-holuhraun-vulkanische-erupties-op-ijsland)  
holuhraun-vulkanische-erupties-op-ijsland
- Massad, R. S., Nemitz, E., & Sutton, M. A. (2010). Review and parameterisation of bi-directional ammonia exchange between  
vegetation and the atmosphere. *Atmospheric Chemistry and Physics*, *10*(21), 10359–10386.  
<https://doi.org/10.5194/acp-10-10359-2010>
- Mathijssen, T., & Knap, W. (2021). *Pyranometer intercomparison at the BSRN site in Cabauw, the Netherlands*. KNMI.
- 625 Moene, A. F., & van Dam, J. C. (2016). *Transport in the Atmosphere-Vegetation-Soil Continuum*. Cambridge University Press.
- Mooibroek, D., Berkhout, J. P. J., & Hoogerbrugge, R. (2014). *Jaaroverzicht Luchtkwaliteit 2013*.
- Neiryneck, J., & Ceulemans, R. (2008). Bidirectional ammonia exchange above a mixed coniferous forest. *Environmental  
Pollution*, *154*(3), 424–438. <https://doi.org/10.1016/j.envpol.2007.11.030>
- 630 Nemitz, E., Milford, C., & Sutton, M. A. (2001). *A two-layer canopy compensation point model for describing bi-directional  
biosphere-atmosphere exchange of ammonia*. 815–833.
- Nemitz, E., Sutton, M. A., Gut, A., San Jose, R., Husted, S., & Schjoerring, J. K. (2000). Sources and sinks of ammonia within  
an oilseed rape canopy. *Agricultural and Forest Meteorology*, *105*(4), 385–404. [https://doi.org/10.1016/S0168-](https://doi.org/10.1016/S0168-1923(00)00205-7)  
1923(00)00205-7
- 635 Personne, E., Loubet, B., Herrmann, B., Mattsson, M., Schjoerring, J. K., Nemitz, E., Sutton, M. A., Cellier, P., & Estate, B.  
(2009). *SURFATM-NH3: A model combining the surface energy balance and bi-directional exchanges of ammonia  
applied at the field scale*. 1371–1388.



- Pleim, J. E., Ran, L., Appel, W., Shephard, M. W., & Cady-Pereira, K. (2019). New Bidirectional Ammonia Flux Model in an Air Quality Model Coupled With an Agricultural Model. *Journal of Advances in Modeling Earth Systems*, 11(9), 2934–2957. <https://doi.org/10.1029/2019MS001728>
- 640 RIVM. (2023). *GDN depositiebestanden achterliggende jaren*. <https://www.rivm.nl/gcn-gdn-kaarten/depositiekaarten/cijfers-achter-depositiekaarten/gdn-depositiebestanden-achterliggende-jaren>
- Salesky, S. T., & Chamecki, M. (2012). Random Errors in Turbulence Measurements in the Atmospheric Surface Layer: Implications for Monin–Obukhov Similarity Theory. *Journal of the Atmospheric Sciences*, 69(12), 3700–3714. <https://doi.org/10.1175/JAS-D-12-096.1>
- 645 Schrader, F., Brümmner, C., Flechard, C. R., Kruit, R. J. W., Van Zanten, M. C., Zöll, U., Hensen, A., & Erisman, J. W. (2016). Non-stomatal exchange in ammonia dry deposition models: Comparison of two state-of-the-art approaches. *Atmospheric Chemistry and Physics*, 16(21), 13417–13430. <https://doi.org/10.5194/acp-16-13417-2016>
- Schrader, F., Erisman, J. W., & Brümmner, C. (2020). Towards a coupled paradigm of NH<sub>3</sub>-CO<sub>2</sub> biosphere–atmosphere exchange modelling. *Global Change Biology*, 26(9), 4654–4663. <https://doi.org/10.1111/gcb.15184>
- 650 Schulte, R. B., Vilà-Guerau De Arellano, J., Rutledge-Jonker, S., Van Der Graaf, S., Zhang, J., & Van Zanten, M. C. (2024). Observational relationships between ammonia, carbon dioxide and water vapor under a wide range of meteorological and turbulent conditions: RITA-2021 campaign. *Biogeosciences*, 21(2), 557–574. <https://doi.org/10.5194/bg-21-557-2024>
- Shephard, M. W., & Cady-Pereira, K. E. (2015). Cross-track Infrared Sounder (CrIS) satellite observations of tropospheric ammonia. *Atmospheric Measurement Techniques*, 8(3), 1323–1336. <https://doi.org/10.5194/amt-8-1323-2015>
- 655 Shi, Y., & Long, C. N. (2002). *Techniques and Methods used to determine the Best Estimate of Radiation Fluxes at SGP Central Facility*. <https://www.osti.gov/biblio/965640>
- Stoffel, T. (2005). Solar infrared radiation station (SIRS) handbook. *Atmospheric Radiation Measurement Program Technical Report ARM TR-025*, 1123–1134.





- 660 Sutton, M. A., Burkhardt, J. K., Guerin, D., Nemitz, E., & Fowler, D. (1998). Development of resistance models to describe measurements of bi-directional ammonia surface-atmosphere exchange. *Atmospheric Environment*, 32(3), 473–480. [https://doi.org/10.1016/S1352-2310\(97\)00164-7](https://doi.org/10.1016/S1352-2310(97)00164-7)
- Sutton, M. A., Fowler, D., Burkhardt, J. K., & Milford, C. (1995). Vegetation atmosphere exchange of ammonia: Canopy cycling and the impacts of elevated nitrogen inputs. *ACID REIGN* ', 4.
- 665 Sutton, M. A., Nemitz, E., Milford, C., Campbell, C., Erisman, J. W., Hensen, A., Cellier, P., David, M., Loubet, B., Personne, E., Schjoerring, J. K., Mattsson, M., Dorsey, J. R., Gallagher, M. W., Horvath, L., Weidinger, T., Meszaros, R., Dämmgen, U., Neftel, A., ... Burkhardt, J. (2009). Dynamics of ammonia exchange with cut grassland: Synthesis of results and conclusions of the GRAMINAE Integrated Experiment. *Biogeosciences*, 6(12), 2907–2934. <https://doi.org/10.5194/bg-6-2907-2009>
- 670 Sutton, M. A., Schjorring, J., & Wyers, G. (1995). Plant atmosphere exchange of ammonia. *Philosophical Transactions of the Royal Society A: Mathematical, Physical & Engineering Sciences*, 351, 261–276.
- Swart, D., Zhang, J., van der Graaf, S., Rutledge-Jonker, S., Hensen, A., Berkhout, S., Wintjen, P., van der Hoff, R., Haaima, M., Frumau, A., van den Bulk, P., Schulte, R., van Zanten, M., & van Goethem, T. (2023). Field comparison of two novel open-path instruments that measure dry deposition and emission of ammonia using flux-gradient and eddy covariance methods. *Atmospheric Measurement Techniques*, 16(2), 529–546. <https://doi.org/10.5194/amt-16-529-2023>
- 675 Ustymczuk, A., & Giner, S. A. (2011). Relative humidity errors when measuring dry and wet bulb temperatures. *Biosystems Engineering*, 110(2), 106–111. <https://doi.org/10.1016/j.biosystemseng.2011.07.004>
- Van Hove, L. W. A., Adema, E. H., Vredenberg, W. J., & Pieters, G. A. (1989). A study of the adsorption of NH<sub>3</sub> and SO<sub>2</sub> on leaf surfaces. *Atmospheric Environment (1967)*, 23(7), 1479–1486. [https://doi.org/10.1016/0004-6981\(89\)90407-1](https://doi.org/10.1016/0004-6981(89)90407-1)
- 680 Van Zanten, M. C., Sauter, F. J., Kruit, R. J. W., J.A. van Jaarsveld, P., & Pul, W. A. J. van. (2010). *Description of the DEPAC module Description of the DEPAC module.*



- Vendel, K. J. A., Wichink Kruit, R. J., Blom, M., van den Bulk, P., van Egmond, B., Frumau, A., Rutledge-Jonker, S., Hensen, 685  
A., & van Zanten, M. C. (2023). Dry deposition of ammonia in a coastal dune area: Measurements and modeling.  
*Atmospheric Environment*, 298(June 2022), 119596. <https://doi.org/10.1016/j.atmosenv.2023.119596>
- Von Arx, G., Dobbertin, M., & Rebetez, M. (2012). Spatio-temporal effects of forest canopy on understory microclimate in a  
long-term experiment in Switzerland. *Agricultural and Forest Meteorology*, 166–167, 144–155.  
<https://doi.org/10.1016/j.agrformet.2012.07.018>
- 690 Wang, K., Kang, P., Lu, Y., Zheng, X., Liu, M., Lin, T.-J., Butterbach-Bahl, K., & Wang, Y. (2021). An open-path ammonia  
analyzer for eddy covariance flux measurement. *Agricultural and Forest Meteorology*, 308–309, 108570.  
<https://doi.org/10.1016/j.agrformet.2021.108570>
- Wen, D., Zhang, L., Lin, J. C., Vet, R., & Moran, M. D. (2014). An evaluation of ambient ammonia concentrations over  
southern Ontario simulated with different dry deposition schemes within STILT-Chem v0.8. *Geoscientific Model*  
695 *Development*, 7(3), 1037–1050. <https://doi.org/10.5194/gmd-7-1037-2014>
- Wentworth, G. R., Murphy, J. G., Benedict, K. B., Bangs, E. J., & Collett, J. L. (2016). The role of dew as a night-time reservoir  
and morning source for atmospheric ammonia. *Atmospheric Chemistry and Physics*, 16(11), 7435–7449.  
<https://doi.org/10.5194/acp-16-7435-2016>
- Wentworth, G. R., Murphy, J. G., Gregoire, P. K., Cheyne, C. A. L., Tevlin, A. G., & Hems, R. (2014). Soil–atmosphere  
700 exchange of ammonia in a non-fertilized grassland: Measured emission potentials and inferred fluxes.  
*Biogeosciences*, 11(20), 5675–5686. <https://doi.org/10.5194/bg-11-5675-2014>
- Wesely, M. L., & Hicks, B. B. (1977). Some factors that affect the deposition rates of sulfur dioxide and similar gases on  
vegetation. *Journal of the Air Pollution Control Association*. <https://doi.org/10.1080/00022470.1977.10470534>
- Westreenen, A. V., Zhang, N., Douma, J. C., Evers, J. B., Anten, N. P. R., & Marcelis, L. F. M. (2020). Substantial differences  
705 occur between canopy and ambient climate: Quantification of interactions in a greenhouse-canopy system. *PLOS*  
*ONE*, 15(5), e0233210. <https://doi.org/10.1371/journal.pone.0233210>
- Wichink Kruit, R. J. (2010). *Surface-atmosphere exchange of ammonia. Measurements and modeling over non-fertilized  
grassland in the Netherlands*. [https://doi.org/10.1007/978-3-662-03394-4\\_18](https://doi.org/10.1007/978-3-662-03394-4_18)



- 710 Wichink Kruit, R. J., Schaap, M., Sauter, F. J., Van Zanten, M. C., & Van Pul, W. A. J. (2012). Modeling the distribution of  
ammonia across Europe including bi-directional surface-atmosphere exchange. *Biogeosciences*, 9(12), 5261–5277.  
https://doi.org/10.5194/bg-9-5261-2012
- Wichink Kruit, R. J., Van Pul, W. A. J., Otjes, R. P., Hofschreuder, P., Jacobs, A. F. G., & Holtslag, A. A. M. (2007). Ammonia  
fluxes and derived canopy compensation points over non-fertilized agricultural grassland in The Netherlands using  
the new gradient ammonia—High accuracy—Monitor (GRAHAM). *Atmospheric Environment*, 41(6), 1275–1287.  
715 https://doi.org/10.1016/j.atmosenv.2006.09.039
- Wichink Kruit, R. J., van Pul, W. A. J., Sauter, F. J., van den Broek, M., Nemitz, E., Sutton, M. A., Krol, M., & Holtslag, A.  
A. M. (2010). Modeling the surface-atmosphere exchange of ammonia. *Atmospheric Environment*, 44(7), 945–957.  
https://doi.org/10.1016/j.atmosenv.2009.11.049
- Wyers, G. P., & Erisman, J. W. (1998). Ammonia exchange over coniferous forest. *Atmospheric Environment*, 32(3), 441–  
720 451. https://doi.org/10.1016/S1352-2310(97)00275-6
- Zhang, L., Brook, J. R., & Vet, R. (2003). A revised parameterization for gaseous dry deposition in air-quality models.  
*Atmospheric Chemistry and Physics*, 3(6), 2067–2082. https://doi.org/10.5194/acp-3-2067-2003
- Zhang, L., Moran, M. D., Makar, P. A., Brook, J. R., & Gong, S. (2002). Modelling gaseous dry deposition in AURAMS: A  
unified regional air-quality modelling system. *Atmospheric Environment*, 36(3), 537–560.  
725 https://doi.org/10.1016/S1352-2310(01)00447-2
- Zhang, L., Wright, L. P., & Asman, W. A. H. (2010). Bi-directional air-surface exchange of atmospheric ammonia: A review  
of measurements and a development of a big-leaf model for applications in regional-scale air-quality models. *Journal  
of Geophysical Research Atmospheres*, 115(20). https://doi.org/10.1029/2009JD013589