

InSAR sensitivity to active layer ground ice content in Adventdalen, Svalbard

Lotte Wendt^{1,2,3}, Line Rouyet², Hanne H. Christiansen³, Tom Rune Lauknes², Sebastian Westermann^{1,4}

¹ Department of Geosciences, University of Oslo, P.O. Box 1047 Blindern, 0316, Oslo, Norway

5 ² NORCE Norwegian Research Centre AS, Siva Innovasjonssenter, P.O. Box 6434, 9294, Tromsø, Norway

³ Arctic Geophysics Department, The University Centre in Svalbard (UNIS), P.O. Box 156, 9171, Longyearbyen, Norway

⁴ Centre for Biogeochemistry in the Anthropocene, University of Oslo, Oslo, Norway

Correspondence to: Lotte Wendt (lottew@uio.no)

Abstract

10 Interferometric Synthetic Aperture Radar (InSAR) remote sensing of surface displacement in permafrost environments has the potential to resolve ground ice dynamics and potentially active layer thickness, yet field validation is sparse. Here we present a comparison between in-situ ground ice contents and the seasonal InSAR displacements of the following thawing season at 12 coring sites in Adventdalen, Svalbard. The study is focused on the year 2023, where frozen sediment cores were collected at the end of spring from the active layer and the uppermost permafrost. The sediment cores were analyzed with high resolution
15 for volumetric ground ice and excess ice contents. The active layer thickness was estimated by probing the thaw depth at the end of the thawing season 2023, allowing estimation of the amount of expected subsidence from seasonal ground ice melt. The InSAR vertical displacements for the thawing season were derived from Small Baseline Subset (SBAS) processing of Sentinel-1 imagery. The expected subsidence from ground ice melt within the measured active layer aligned well with the seasonal InSAR maximum vertical displacement. Monte Carlo simulations were performed to include uncertainties in the expected and
20 measured InSAR subsidence, leading to a mean coefficient of determination of 0.68 and a mean absolute error of 15 mm for the correlation between InSAR subsidence and expected subsidence from in-situ ground ice melt. Excess ice is highly variable and is the main source of the expected subsidence during this thawing season, which was exceptionally warm. The expected subsidence and active layer thickness show only a weak relationship due to the observed complex ice content distribution in the active layer and uppermost permafrost. Our results show the significant potential of InSAR for mapping ground ice
25 variability; however, they also suggest that estimating active layer thickness using InSAR requires careful consideration of the complex occurrence of both pore and excess ice in the active layer and uppermost permafrost.

1 Introduction

Permafrost environments underlie approximately 15% of the northern hemisphere and are highly sensitive to ongoing climate change (Obu, 2021, Biskaborn et al., 2019). Overlying the permafrost is the active layer, which thaws during summer and
30 refreezes during winter. An increase in the active layer thickness (ALT) serves as a key indicator of permafrost degradation

(GCOS, 2022). The formation and melt of ground ice impacts the thermal behavior of the ground, with larger ground ice contents increasing the latent heat of fusion for thaw and thus causing a reduced seasonal thaw depth (French, 2007a). Seasonal variations in the active layer ground ice content affect local hydrology and ground stability (Walvoord and Kurylyk, 2016). Long-term ground ice loss is associated with active layer deepening and may cause pronounced terrain alternations, such as subsidence and the development of thermokarst landforms (Burn et al., 2024). These changes have broad implications for the carbon cycle in permafrost areas (Schuur et al., 2015).

Traditional methods for monitoring ALT and mapping ground ice (e.g. thaw depth probing, temperature monitoring in boreholes, drilling and geomorphological surveys, thaw tubes) typically rely on labor-intensive, time-consuming in-situ surveys. Additionally, many regions are difficult to access for in-situ monitoring (Bonnaventure and Lamoureux, 2013).

Interferometric Synthetic Aperture Radar (InSAR) satellite remote sensing has in recent years been increasingly used to monitor thaw subsidence in permafrost environments. Studies have shown that the spatiotemporal InSAR variability can be related to ground ice content patterns and soil water contents (Daout et al., 2017; Chen et al., 2020), as well as air and ground temperature variations (Strozzi et al., 2018; Bartsch et al., 2019), and different landforms and surface materials (Rouyet et al. 2019). The detected InSAR thaw subsidence has been exploited to inversely retrieve ALT assuming certain ground conditions (e.g. Liu et al., 2012; Schaefer et al., 2015). Further, InSAR subsidence measurements have proven effective for monitoring long-term ground ice loss due to permafrost degradation (e.g. Wang et al., 2023). InSAR allows large-scale mapping of surface displacements related to active layer and permafrost changes, independent of solar illumination and cloud conditions. However, the detailed response of InSAR measurements to seasonal ground ice dynamics remains underexplored.

Ground ice, which varies in distribution, includes both pore ice within soil pores and excess ice that exceeds the soil's pore space (Everdingen, 1998). Pore ice forms when soil moisture freezes within the existing pore spaces of mineral and organic soils. Depending on the degree of saturation, the phase change causes either a volume expansion within the pore space or, if the pores are saturated with water, an expansion of the soil structure itself due to the pressure exerted by the growing ice. Upon melting, pore ice does not produce water in excess of the pore space. Ice segregation processes can cause the migration of water towards the freezing front, enriching soil pores further with ice and leading to the growth of ice lenses (Derek and Miller, 1966). If the accumulation of ice exceeds the pore space, excess ice forms, which upon melting produces water in excess of the pore space (Taber, 1930; Rempel et al., 2007).

The melting of pore ice in saturated ground can lead to thaw consolidation, caused by the volume loss associated with the density difference between ice and water (approx. 8%) (Dumais and Konrad, 2024). The melting of excess ice has an even more pronounced effect, as the loss of ice that exceeds the soil's pore space can cause significant subsidence when the resulting meltwater drains away (Morgenstern and Nixon, 1971).

Comprehensive field validation to measure the impact these ice types have on InSAR measurements is still lacking, underscoring the need for improved understanding of how InSAR captures these seasonal ground ice changes (Bartsch et al., 2023).

The contribution of excess ice melt to observed seasonal surface subsidence was hypothesized by Liu and Larson (2018) and Bartsch et al. (2019), yet without further in-situ field validation with active layer ground ice contents. Studies by Zwieback and Meyer (2021) and Zwieback et al. (2024) indicate that InSAR subsidence measurements from the late thawing season can suggest excess ice melt from the top of permafrost. These studies highlight the need for further field validation to understand how well InSAR displacements align with seasonal ground ice melt in the active layer. The sensitivity of InSAR to pore ice and excess ice melt as well as the correlation between active layer thickness and InSAR subsidence are also of relevance for the inverse retrieval of active layer thickness from InSAR (Liu et al., 2012).

Here we present a comparison between seasonal C-band InSAR displacements and in-situ active layer and uppermost permafrost ground ice contents from different periglacial landforms in the Adventdalen valley, Svalbard. Our objectives are to (1) compare the InSAR subsidence to the expected subsidence from ground ice melt based on in-situ field measurements, (2) investigate the contributions of pore and excess ice melt to the seasonal subsidence signal, and (3) evaluate the relationship between subsidence magnitude and ALT.

2 Methods

2.1. Study area

The study is focused on the valley Adventdalen, located in Central Spitsbergen, Svalbard (78.2°N, 15.8°E) (Fig. 1). This valley has continuous permafrost, varying in thickness from less than 100 m in valley bottoms and coastal areas to up to 500 m in the mountains (Humlum et al., 2003). The uppermost permafrost formed largely syngenetically due to eolian deposition on the fluvial terraces adjacent to the river Adventelva during the late Holocene (Gilbert et al., 2018) (Fig. 1B). The periglacial valley landscape is covered by landforms such as ice-wedge polygons, solifluction sheets, alluvial fans, loess terraces on the sides of Adventelva, and moraine deposits. The valley bottom has extensive tundra vegetation, whereas there is no vegetation in the braided river plain. The landcover is favorable for the application of InSAR with C-band sensors, since decorrelation by shrub vegetation does not occur (Wang et al., 2020).

2.3. Analysis of ground ice contents

All intact cores were cut in half lengthwise and processed in 1–7 cm sections in a freezing lab at -8 °C. The intact subsections were scraped, the cryostratigraphy classified based on French and Shur (2010), and visual ice content and sediment type described. All intact core sections were photographed ([cross-sections shown in Fig. S2](#)), and their length l and diameter d measured with a ruler. Disturbed core sections were also photographed and processed in the sections retrieved during drilling without further subsampling (mean length of 9 cm \pm 6 cm). For disturbed core sections, the length l was measured in the field during retrieval from the borehole and the diameter d was based on the core barrel diameter.

All subsamples were thawed for 12 hours in a sealed measurement beaker at room temperature. After thawing, the volume of sediments and supernatant water was measured by reading of the volumes in the graded measurement beaker with help of a ruler. The subsamples were weighed to determine their wet weight M_w . Afterwards, the subsamples were dried in an oven at 80 °C for 48 hours, and their dry weight M_d was recorded.

The volumetric ice content (VIC) and excess ice content (EIC) were calculated based on the formulas by Kokelj and Burn (2005) and Paquette et al. (2023):

$$VIC = \frac{(M_w - M_d) * 1.09}{V_f} * 100 \quad \text{vol [\%]} \quad (1)$$

$$EIC = \frac{V_{sw} * 1.09}{V_f} * 100 \quad \text{vol [\%]} \quad (2)$$

where V_f is the frozen volume of the subsample, calculated as the volume of a half cylinder from the frozen core length l and diameter d ; M_w is the wet weight of the subsample (g); M_d is the dry weight of the subsample (g); V_{sw} is the volume of supernatant water of the thawed subsample (ml); and the factor 1.09 is to estimate the equivalent volume of ice from the water volume.

The pore ice content (PIC) was derived under the assumption that the intact core sections were saturated:

$$PIC = VIC - EIC \quad \text{vol [\%]} \quad (3)$$


2.4. Calculation of expected subsidence

The core sections were retrieved either intact or disturbed. Samples that fell apart during sampling and/or analysis (disturbed) were observed to be very dry and therefore classified as unsaturated and not considered to contribute to the subsidence signal. The intact core sections were observed to be visually saturated with pore ice and contained sometimes also excess ice based on the laboratory measurements. Based on these intact core sections, the expected subsidence was derived.

In the intact core sections, both excess ice and pore ice melt are expected to cause a volume reduction according to the density difference between ice and water (approximately 8 %) (Dumais and Konrad 2024). However, excess ice, as it exceeds the pore space of the soil column, can in addition drain or redistribute into unsaturated pore spaces, likely leading to a full volume loss of the excess ice volume (Morgenstern and Nixon, 1971). Based on this rationale, the expected subsidence τ for the thawing season 2023 was calculated from intact core sections, consisting of pore ice melt (8% volume reduction), excess ice melt (8%

290 **Figure 5:** Comparison of seasonal InSAR time series 2023 with the meteorological conditions and the magnitude of expected subsidence from in-situ ground ice melt. (a) Meteorological conditions in the thawing season 2023, including snow depth, daily mean air temperature and daily precipitation based on the Adventdalen meteorological station (Norwegian Meteorological Institute, 2024b). (b) Time series of InSAR displacements during the thawing season 2023 from the different coring sites. The line colour displays the expected subsidence at the respective site based on the in-situ ground ice contents and active layer thickness (see Section 2.4). The site name is shown at the end of each time series line, and all sites are described in Table 1.

295 When assessing the contributions to the expected subsidence from ice melt within the measured ALT, most sites showed a large contribution from excess ice to the expected subsidence signal (Fig. 6). On average across all coring sites, pore ice melt contributed 20 mm to the expected subsidence with a standard deviation (SD) of 8 mm. Excess ice melt contributed an average of 3 mm (SD = 2 mm), and the drainage of meltwater from excess ice contributed an average of 33 mm (SD = 24 mm). The observed maximum InSAR subsidence from summer 2023 aligns well with the expected subsidence when considering both pore ice melt and excess ice melt, including drainage (Fig. 6; grey diamonds).



300

Figure 6: Contribution of pore ice melt, excess ice melt, and excess ice meltwater drainage to the total expected subsidence for each coring site. The maximum InSAR subsidence in 2023 is displayed as grey diamonds. The x-label denotes the site names (sediment deposit type and unique core number, see also Table 1).

The comparison between the expected maximum subsidence and the InSAR maximum subsidence shows good overall alignment (Fig. 7A). The uncertainty in the expected subsidence for the different sites is variable and mostly depends on the excess ice content. A Monte-Carlo simulation incorporating uncertainty over 1000 random iterations resulted in a mean correlation coefficient (r) of 0.82, a mean R^2 of 0.68 and a mean absolute error of 15 mm, indicating statistical significance (p -value < 0.01).

305
310 However, the correlation between the InSAR subsidence and ALT is weak ($R^2 = 0.03$, $r = -0.17$, p -value = 0.60) (Fig. 7B). This aligns with the correlation results between the expected subsidence and the ALT (Fig. 4C).




Figure 7: Correlation between InSAR subsidence and expected subsidence, as well as active layer thickness (ALT). (a) Correlation between the maximum InSAR subsidence of the thawing season 2023 and the expected subsidence from in-situ ground ice melt (pore ice melt + excess ice melt and drainage). The grey whiskers display the uncertainty in the expected and InSAR subsidence. (b) Correlation between the maximum InSAR subsidence of the thawing season 2023 and the in-situ ALT.

315

When evaluating the time series details, the observed InSAR displacements in 2023 align visually well with the ground ice content distribution over depth (Fig. 8). Detailed descriptions of four differing coring sites with varying active layer ice contents follow, providing insights into the spatial diversity observed. Each site's unique characteristics and their impact on both expected and observed subsidence are discussed. All other coring sites are also displayed in a similar format in the Supplement (Fig. S5, S6, S7).

3.3 Examples of site-specific geomorphological and subsidence profiles

Alluvial 1 (A1): The A1 coring site is located in an outer inactive alluvial fan (Fig. 1, [S1](#)). The active layer is mostly ice-poor, with the dominant sediment type being gravel and the samples falling apart at collection. The lower base of the active layer and the uppermost permafrost are very excess ice rich, due to more fine-grained frost susceptible sediments. According to this ground ice profile, the expected subsidence is very low until the excess ice rich layer is reached. Both in 2021 and 2023, the InSAR subsidence displays a rather strong subsidence from the start of the thawing season, which can only be possible if the thaw front penetrates rapidly through the dry gravel layer of the [upper-central](#) active layer. In 2021, the InSAR subsidence levels out earlier than in 2023. In 2023, the InSAR maximum subsidence reaches a similar magnitude as in 2021 and the ground surface stabilizes in July 2023, yet another late-season subsidence pattern starts in August (Fig. 8; [Alluvial 1](#)).

Eolian 8 (E8): Located in a depression on a loess terrace within a poorly developed ice wedge polygon (Fig. 1, [S1](#)), the E8 coring site features a relatively high VIC partly due to its thick organic layer (0.15 m, Table 1), and the entire core was retrieved intact. Similar to A1, both the lower active layer and the uppermost permafrost at E8 are very excess ice rich. This results in a gradual increase in expected subsidence which then accelerates upon reaching the ice-rich lower layer. The InSAR subsidence in 2021 was less than in 2023. The mean ALT of the adjacent ice-wedge CALM grid was 68 cm in 2021 and 82 cm in 2023. In 2023, the InSAR subsidence shows a slowdown in July, before displaying a further late-season subsidence (Fig. 8; [Eolian 8](#)).

Eolian 10 (E10): The coring site E10 is located on a dry, well-drained loess terrace (Fig. 1, [S1](#)). The active layer has a low VIC and parts of the sediment core fell apart at collection. Excess ice is minimal in the uppermost permafrost, leading to negligible expected subsidence. Similarly, the InSAR displacement remains around 0 mm throughout the thawing season both in 2021 and 2023 (Fig. 8; [Eolian 10](#)). The mean ALT of the surrounding UNISCALM grid was 98 cm in 2021 and 107 cm in 2023, but the ALT difference has no large impact on the subsidence, due to the ice-poor uppermost permafrost.

Alluvial 13 (A13): The coring site A13 is located in an outer sandy alluvial floodplain (Fig. 1, [S1](#)). The site has excess ice in the upper active layer, whilst being very dry sand in the lower active layer and uppermost permafrost. As a result of this ground ice profile, the expected cumulative subsidence increases quickly early in the thawing season and then remains constant, which is also observed in the InSAR time series for 2023. In 2021, the thawing onset occurs later, yet again the InSAR subsidence increases quickly before levelling out at a maximum InSAR subsidence similar as 2023 (Fig. 8; [Alluvial 13](#)).




Figure 8: Comparison of InSAR time series for selected sites (Section 3.3) for 2023 (warm summer conditions) and 2021 (cold summer conditions) with the different measured types of ground ice and the expected subsidence. Each black rectangle represents one coring site, with the coring site name on the left, and includes four subplots. “InSAR displacement”: Cumulative InSAR displacement of the thawing seasons 2021 and 2023. “Ice content”: In-situ ground ice contents from spring 2023 displaying the volumetric ice content (VIC) and excess ice content (EIC). “Material type”: Material type over depth. “Expected subsidence”: The cumulative expected subsidence from in-situ ground ice melt over thaw depth. The red dashed line indicates the active layer thickness (ALT) of the thawing season 2023. The yellow dashed line for site E8 and E10 indicates the ALT of 2021 based on the mean ALT of adjacent CALM grids.

350

355 Comparison of 2021 and 2023: As illustrated in Fig. 8, the maximum InSAR subsidence observed in 2021 was consistently
lower than in 2023 at all sites featuring excess ice at the transition between the active layer and uppermost permafrost.
Conversely, sites with minimal excess ice at the bottom of the active layer and uppermost permafrost exhibited comparable
InSAR displacement magnitudes in both years, such as A13. Although no sediment cores were collected in 2021 to confirm
the exact ground ice distribution, ground ice in the studied cores from 2023 indicates that excess ice is especially abundant in
360 the lower active layer and uppermost permafrost (Fig. 3). The 2021 thawing season was exceptionally cold and very
shallow ALT were recorded at other stations in Adventdalen, including close to two of our coring sites (Fig. 8, ~~Fig.~~ S8).
This could indicate that less excess ice melt and drainage contributed to the subsidence under the assumption of similar active
layer ground ice conditions. This aligns with the finding that the InSAR subsidence for 2021 matches well the expected
subsidence from pore ice melt in 2023, except at alluvial sites with more complex excess ice distributions in the active layer
365 (Fig. S9).

4 Discussion

4.1 InSAR subsidence as indicator of ground ice melt

Our analysis shows a high correlation between InSAR displacements and in-situ ice contents in ~~from~~ the active layer and
uppermost permafrost, confirming a good sensitivity of InSAR to ground ice melt. This sensitivity is consistent with results of
370 other studies who have evaluated the use of InSAR for documenting ground water and ice content in permafrost terrain (Chen
et al., 2020; Daout et al., 2020; Wang et al., 2023; Zwieback and Meyer., 2021; Zwieback et al., 2024).

We observed seasonal variations in the InSAR subsidence patterns, which especially align with the distribution of excess ice
in the active layer. Pore ice also contributes to the expected subsidence, but in a secondary manner, which more closely aligns
with the thaw progression predicted by the Stefan equation (e.g. Fig. 5b: site E3). The expected subsidence contribution from
375 pore ice melt is smaller, since it is only caused by the phase change volume reduction from saturated pores, which were usually
found in the upper and lower active layer. The central active layer was observed to be ice-poor, which is in line with previous
field results (e.g. Mackay, 1983; Zwieback et al., 2024). This can be attributed to two-sided freezing of the active layer, with
water migrating towards the freezing fronts, which are moving downwards from the surface and upwards from the permafrost
table. This caused ice lenses to form especially in the lower active layer (Taber, 1930), and the upper and lower active layer to
380 be saturated with pore ice (Derek and Miller, 1966; Bai et al., 2020), whilst the central active layer remains dry (French,
2007a).

Many sites show a period of low InSAR subsidence in the middle of the thawing season. Yet, the cumulative subsidence at
which the InSAR time series stagnates generally does not correspond to the central active layer (inferred from the cumulative
expected subsidence). This observation could suggest that the thaw front quickly penetrates the central part of the active layer,
385 and then slows down when it reaches the ice-rich bottom of the active layer. The inverse relationship between active layer
thaw rate and ice content has been previously confirmed (French, 2007a).

The period of stagnating subsidence rate is most pronounced between the 24 July and 08 August 2023 at several of the studied sites, all of which have an excess ice-rich uppermost permafrost. The ground surface stabilization could have been caused by the thaw front reaching the ice-rich transient layer or uppermost permafrost (Shur et al. 2005). Here, sensible and latent heat effects of warming and melting the excess ice, as well as slow drainage of the excess ice meltwater, could have delayed further surface subsidence, thereby causing a stagnation in the InSAR subsidence pattern. Alternatively, a significant rainfall event (40 mm) during August 4–6, 2023 (Fig. 5), might have temporarily masked subsidence signals due to ground swelling. The rain event could on the other hand also have enhanced the late-season subsidence by percolation of rainwater advecting heat into the lower active layer (Douglas et al., 2020; Magnússon et al., 2022).

The expected subsidence from pore ice melt lies within a plausible range, yet our results indicate that excess ice melt and meltwater drainage can significantly dominate the expected subsidence signal (Fig. 6: A1, A9, A12). Zwieback and Meyer (2021) mapped excess ice-rich top of permafrost distributions based on the presence or absence of late-season subsidence in an exceptionally warm year in northwestern Alaska within continuous permafrost. They found that the presence of late-season subsidence in InSAR time series closely matches in-situ observations of excess ice-rich permafrost areas. The attribution of late-season subsidence in a warm thawing season to excess ice melt of the uppermost permafrost has also been previously hypothesized by Bartsch et al. (2019). The findings of Zwieback and Meyer (2021) align with our results, which indicate that excess ice can be the major contribution to the InSAR late-season subsidence signal (Fig. 8, S5-7). Nevertheless, our results also indicate that pore ice contributes to the expected subsidence. Thus, converting the InSAR subsidence to excess ice profiles, such as in Zwieback et al. (2024), will neglect these contributions.

Previous in-situ surface displacement evaluations have been conducted in Siberia by Antonova et al. (2018) and in NW-Canada by Gruber (2020). Antonova et al. (2018) evaluated InSAR displacements with in-situ surface subsidence from reference rods anchored in permafrost in a Yedoma landscape of the Lena River delta in Siberia. They found that their InSAR displacements from X-band SAR generally underestimated the in-situ subsidence. Compared to this study, our results are based on C-band SAR and could thus be more robust to decorrelation and aliasing (Wang et al., 2020). Gruber (2020) measured in-situ surface displacements with a tilt-arm apparatus in warm permafrost of black forests and peatlands in NW-Canada. They found that the surface only subsided during the thawing season at a silty site. In two other sites of Gruber (2020), with very thick dry peat layers, the surface heaved, even during the thawing season. Our results cover only one site which has an organic layer throughout the active layer and this site is fully saturated. We do not see a heave signal, but instead a subsidence magnitude that aligns with the pore ice content. Nevertheless, the results from Gruber (2020) indicate that our findings might not apply in areas with different ground conditions, such as thicker vegetation cover and dry peat layers.

Our results also indicate that InSAR subsidence variations within identical sediment deposit types can delineate differences in active layer and top of permafrost ground ice contents. In alluvial fans, the in-situ ground ice content in the upper, drained, coarse-grained areas is very low. In contrast, the outer areas of the fan consist of both coarse and fine-grained layers, with the latter being rich in excess ice. The InSAR time series map these gradients in ground ice content of the active layer and uppermost permafrost, as illustrated for coring sites A1 and A4 (Fig. 1, 8, S5). The presence of excess ice in the active layer

appears to be influenced by several factors, including drainage, landform history, and sediment grain size. In alignment with previous research (e.g. French, 2007b; Cable et al., 2018), excess ice is observed predominantly in fine-grained sediments, whereas it is nearly absent in the gravel and sandy core sections of our retrieved sediment cores (Fig. S10).

Our data exemplify that, even within the same type of sediment deposit, excess ice presence can vary in the active layer and uppermost permafrost. For instance, eolian fine-grained loess terraces show significant variability: some have very low ground ice contents and lack excess ice (e.g., coring site E10), while others have large ground ice contents, particularly in the lower active layer (e.g., coring sites E2, E8) (Fig. 8, S5). This is likely related to drainage and grain size variations (O’Neill et al. 2025), as well as the site-specific formation history of the sediments (Gilbert et al. 2018). The InSAR maps can capture these variations (Fig. 1). Future geomorphological studies could leverage InSAR datasets to enhance ground ice mapping within periglacial landforms by integrating knowledge of sediment grain size and soil moisture conditions. Seasonal variations in InSAR magnitude could be exploited to map active layer ice content (during cold thawing seasons) and top of permafrost ice content (during exceptionally warm thawing seasons). The time series analysis could utilize periods of subsidence stagnation as potential indicator for reaching excess ice rich sediment layers. This approach could help to discriminate different typical landform subsidence patterns, thereby enabling InSAR remote sensing to provide insights into the ground stratigraphy in complex periglacial environments.

4.2 Implications for ALT estimation from InSAR subsidence

Previously, InSAR subsidence has been used for ALT inversion by assuming a positive relationship between InSAR subsidence observations of pore ice melt and ALT, with no consideration of excess ice contributions (Liu et al., 2012; Schaefer et al., 2015; Jia et al., 2017; Wang et al., 2018; Peng et al., 2023; Scheer et al., 2023). Some of these studies show good alignment between the InSAR-ALT and ALT from in-situ field measurements (Schaefer et al., 2015). However, recent findings show that the positive correlation between InSAR subsidence observations of pore ice melt and ALT used as basis for the inversion is not valid in drier regions, where the correlation is negative (Chang et al., 2024).

We also find a negative correlation between ALT and expected pore ice subsidence (Fig. 4A), indicating that larger ALT can occur at sites with less pore ice. In addition, our results suggest that traditional inversion models are not universal and might not apply in areas with complex ground stratigraphy. Adventdalen is a periglacial valley which has a very diverse geomorphology with a variety of landforms, some of which have abundant excess ice in the active layer, as for example observed in outer alluvial fans (Fig. 6).

Our results indicate that the surface subsidence signal can be dominated by excess ice melt and drainage instead of pore ice melt, which complicates inversions from thaw subsidence to ALT. Further, the correlation between ALT and the InSAR subsidence was close to random in our results. Considering only pore ice melt to explain the observed subsidence in our study area would cause large errors, since the main subsidence contribution is from excess ice (Fig. 6).

In alignment with our observations (Fig. 7a), Antonova et al. (2018) also observed a poor match between ALT and subsidence. They compared in-situ ALT measurements to in-situ surface subsidence from reference rods anchored in Yedoma permafrost

in Siberia and found a weak positive correlation, with a low coefficient of determination. Further, when comparing the measured in-situ subsidence to the expected subsidence predicted by a pore ice melt model based on soil moisture measurements, they found a moderate alignment but noted significant outliers where the model underestimated the measured subsidence. Very large in-situ subsidence was for example measured in drained lake basins, yet not consistently. Such outliers in drained lake basins have also been reported in InSAR-ALT estimates. Schaefer et al. (2015) reported outliers in InSAR-ALT estimates over drained lake basins at Barrow, Alaska, where high subsidence led to a large InSAR-ALT under the assumption of only pore ice melt contributing to the subsidence. Similar patterns of high subsidence in such landforms were also reported by Liu et al. (2014) on the Alaskan North Slope near Prudhoe Bay, by Strozzi et al. (2018) at Teshekpuk lake, Alaska, and by Bartsch et al. (2019) in central Yamal, Russia. Our results suggest that the mismatch between InSAR-derived ALT and in-situ ALT is likely due to the omission of excess ice, which has been previously detected in these landforms (Jorgenson and Shur, 2007; Bockheim and Hinkel, 2012; Kanevskiy et al., 2013). Overall, our study indicates that the contribution from excess ice should not be neglected in models utilizing InSAR time series for active layer characterization, and that simple active layer models relying on constant pore ice contents are oversimplistic in periglacial environments like Adventdalen, Svalbard. Future work should investigate integrating InSAR time series with numerical models that simulate ice segregation processes and excess ice content (e.g., Aga et al., 2023). This could help constrain model parameters or improve process representation by leveraging InSAR-derived surface deformation patterns, potentially through data assimilation techniques (e.g., Aalstad et al., 2018).

4.3 Limitations

Our evaluation is limited by the point-wise core measurements versus the much larger InSAR pixel (~18 x 28 m spatial resolution). Due to the inherent variability in periglacial landscapes, this could have introduced artifacts into our analysis (e.g. Hinkel and Nelson, 2003). This effect was moderated by choosing coring locations within a homogenous surrounding, both by looking at the spatial distribution of InSAR displacements and existing maps of sediment deposit types and landforms. However, other studies have displayed the high spatial variability of ground ice contents and surface displacements (e.g., Antonova et al., 2018; Cable et al., 2018; Zwieback et al. 2024). Within our study area, previous studies have displayed variability in surface displacements even within the same sediment deposit type (Rouyet et al. 2019), which is also visible in our data from 2023 (Fig. S11). The cores were collected at the end of the freezing period, yet water infiltration at the thaw season onset into the frozen part of the sediment column could have caused aggradational ice growth after core collection (e.g. Mackay, 1983, Scherler et al. 2010). This may have caused an underestimation of the in-situ ground ice contents. The excess ice contents are likely conservative, since air bubbles within the excess ice cause an even larger volume loss upon thawing, which is not measured by assessing the supernatant water volume (Morse et al., 2009). Additionally, the excess ice content measurements were performed for 1-7 cm length core sections, which might sometimes have contained ice lenses and then unsaturated sections. These two contributions may have cancelled each other out, thereby reducing the amount of excess ice measured.

Our evaluation is further limited by the small sample size. Both more coring sites and several cores collected at one site would have increased the robustness of our study. Future research should expand the set of available validation data and apply a similar procedure to other permafrost environments to further validate our findings. Complementary ground temperature measurements could also provide more detailed insights into the relations between thaw front propagation, ground ice contents and the InSAR displacement signal.

The InSAR analysis is based on few interferograms in 2023. Due to the failure of Sentinel-1B, the minimum temporal baseline was 12 days, and because of the high subsidence rate, it was only possible to include 24-day interferograms at the end of the thawing season. Nevertheless, the magnitude of the InSAR 2023 dataset is in line with previous years, although a stronger subsidence is observed in 2023, which aligns with the exceptionally warm summer and the very wet conditions (Fig. 1B). In-situ borehole and CALM grid ALT measurements confirm the exceptional large ALT in 2023 (Fig. S8).

The InSAR displacements were converted to vertical displacements by assuming no horizontal movement. Two coring sites (T7 and S5) are placed on 6° and 8° slope angle, which could cause slight horizontal displacement components, for example through solifluction (Harris et al. 2011). Further, the site E8 is located within a low-centre ice-wedge polygon. Whilst the core was extracted at the polygon centre, the InSAR pixel is large enough to include thaw subsidence effects from the ice wedge troughs (Short and Fraser, 2023). Since the ice wedge tops in adjacent polygons have been observed to be located just below the active layer (O'Neill et al., 2025), the late-season subsidence observed at this site might partly reflect this (Burn et al., 2021). Another site, E2, also displays polygonal features indicative of ice wedges. Yet, these sites do not present as outliers in our analysis (Fig. 6).

The InSAR time series starts two days after the snow melt-out date at the Adventdalen meteorological station. Sentinel-2 imagery confirms that all sites except coring site S5 were snow-free or had mixed pixels at the start of the InSAR time series, suggesting minimal impact of snow cover. However, initial subsidence just after snowmelt may not be fully captured at some sites, which could affect comparability. At coring site S5, InSAR subsidence remained negligible until after local snowmelt, which occurred around 7 June 2023.

The reported InSAR uncertainty of 1 cm originates from studies conducted outside the Arctic and may be larger in our study area. However, careful checks for unwrapping errors and decorrelation were conducted as part of the SBAS analysis to mitigate these uncertainties. Interferograms affected by unwrapping errors were manually discarded.

Our expected subsidence calculation is sensitive to the exact ALT, since the bottom of the active layer is most ice rich. In this study manual probing was employed, which allows a more widespread coverage of the pixel, yet is dependent on rather fine-grained ground conditions. Additional borehole temperature measurements could have aided in determining the ALT and provided thaw progression measurements throughout the thawing season, which would have been valuable for comparison against the InSAR subsidence progression.

Lastly, pore ice melt from core sections that were retrieved intact and appeared saturated was considered to contribute to the expected subsidence in this study. However, open-system behavior of the active layer could have allowed lateral or vertical escape and expansion pressure from phase change to dissipate, leading to less pore-ice-melt-induced subsidence than the

[theoretical 8% volumetric reduction. This may have caused an overestimation of the expected subsidence from pore ice melt and could have affected the validation assessment between InSAR and expected subsidence. Future research should aim to better quantify the role of pore ice in heave-subsidence soil mechanics under natural freezing and thawing conditions.](#)

5 Conclusion

525 In this study, we compared seasonal SBAS InSAR displacements from C-band SAR with the expected thaw subsidence derived from in-situ ground ice content and thaw depth observations. Sediment cores were retrieved from 12 locations across various periglacial landforms in Adventdalen valley, Svalbard. To estimate the expected subsidence, we calculated pore ice melt, excess ice melt, and excess ice meltwater drainage based on high-resolution measurements of volumetric and excess ice content from the collected sediment cores.

530 Our findings indicate a good agreement between the expected maximum seasonal thaw subsidence and the maximum seasonal InSAR subsidence. Excess ice melt and drainage dominated the expected subsidence at most sites, which can be attributed to an exceptionally large ALT in the warm summer of the year 2023, promoting thaw into the ice-rich base of the active layer and top permafrost. The difference in InSAR subsidence between the cold thawing season of 2021 and the warm thawing season of 2023 was most pronounced at sites with an ice-rich uppermost permafrost.

535 We found only a weak correlation between active layer thickness and both expected subsidence and InSAR subsidence for the thawing season of 2023. This may be attributed to the complex distribution of active layer ice contents across our coring sites and the influence of excess ice on the subsidence signal.

Our results have implications for estimating active layer thickness from InSAR. We show that the estimation of active layer thickness from InSAR can be strongly affected by excess ice, although it is often not considered in current retrieval approaches.

540 The choice of a cold thawing season might reduce the likelihood of thawing into the excess-ice-rich transient layer, thus increasing the ability to translate InSAR subsidence to only pore ice melt.

Overall, our results demonstrate that InSAR displacements correspond well with in-situ active layer ground ice contents, supporting the potential future use of InSAR to map seasonal ice content changes and monitor long-term ground ice loss.

545 Ground ice information is often very sparse and extremely important to understand and monitor the consequences of climate change in periglacial environments.

Appendices

A. Uncertainty estimation of expected subsidence

The uncertainty in the expected subsidence $\delta\tau$ was calculated by propagating the individual measurement uncertainties. The uncertainty δV_f in the frozen volume of a half cylinder is calculated as:

$$550 \quad \delta V_f = \sqrt{(0.25\pi d^2 l \cdot \delta d)^2 + (0.125\pi d^2 \cdot \delta l)^2} \quad \text{[for intact core sections]} \quad (A1)$$

where δd and δl are the respective uncertainty in diameter and length, each being ± 0.2 cm based on repeated measurements. The total relative error δ of the VIC, EIC and PIC for each core section can be propagated under the assumption of independence as:

$$\delta VIC_{\text{section}} = 1.09 \cdot \sqrt{\frac{V_f^2 * (\delta M_d^2 + \delta M_w^2) + (M_d - M_w)^2 * \delta V_f^2}{V_f^4}} \quad (\text{A2})$$

$$555 \quad \delta EIC_{\text{section}} = 1.09 \cdot \sqrt{\frac{V_f^2 * \delta V_{sw}^2 + V_{sw}^2 * \delta V_f^2}{V_f^4}} \quad (\text{A3})$$

$$\delta PIC_{\text{section}} = \sqrt{\delta VIC^2 + \delta EIC^2} \quad (\text{A4})$$

Here, the dry weight uncertainty and wet weight uncertainty is ± 0.01 g according to the scale accuracy. The supernatant water volume uncertainty V_{sw} is ± 1 ml for volumes ≥ 2 ml. Inserting these uncertainties into the expected subsidence formula for each core section leads to:

$$560 \quad \delta \tau_{\text{section}} = \sqrt{(0.08 * \delta PIC)^2 + (1 * \delta EIC)^2} \quad (\text{A5})$$

which is then summed over the whole ALT:

$$\delta \tau_{\text{measurements}} = \sqrt{\sum_0^{ALT} \delta \tau_{\text{section}}^2} \quad (\text{A6})$$

In addition to propagating the individual measurement uncertainties, the uncertainty in the in-situ ALT was also considered and estimated as ± 5 cm based on the variability in measured ALT at each site (δALT). The effect of δALT on the expected subsidence was calculated by assessing the mean change in expected subsidence due to a ± 5 cm variation in ALT:

$$565 \quad \delta ALT_{\text{effect}} = \left| \frac{\tau_{ALT+\delta ALT} - \tau_{ALT-\delta ALT}}{2} \right| \quad (\text{A7})$$

where $\tau_{ALT \pm \delta ALT}$ represents the recalculated expected subsidence for the adjusted ALT values. This additional uncertainty was then integrated into the total uncertainty of the expected subsidence $\delta \tau$:

$$\delta \tau = \sqrt{\delta \tau_{\text{measurement}}^2 + \delta ALT_{\text{effect}}^2} \quad (\text{A8})$$

570 **Data availability**

The ground ice content and active layer thickness data as well as the InSAR time series are archived on [Zenodo](https://doi.org/10.5281/zenodo.11187359) (https://doi.org/10.5281/zenodo.11187359, Wendt, 2024).

Author contributions

LW: Conceptualization, Methodology, Investigation, Formal Analysis, Data Curation, Visualization, Writing - Original Draft,
575 Writing- Reviewing and Editing, Funding acquisition.

LR: Conceptualization, Supervision, Resources, Writing- Reviewing and Editing, Funding acquisition.

HHC: Conceptualization, Supervision, Resources, Writing- Reviewing and Editing, Funding acquisition.

TRL: Data Curation, Writing- Reviewing and Editing, Funding acquisition.

SW: Conceptualization, Supervision, Writing- Reviewing and Editing, Funding acquisition.

580 **Competing interests**

The authors declare they have no conflict of interest.

Acknowledgements

The authors acknowledge funding by the Research Council of Norway (Arctic Field Grant, project no. 342203, RiS ID 12143, 2023), the ESA CCI Permafrost project (4000123681/18/I-NB, 2023) and the Department of Geoscience, University of Oslo.

585 We acknowledge the financial support of the Ministry of Climate and Environment (kap. 1747, post 70) to the Fram Center PermaRICH project, as well as the funding of the Norwegian Space Agency to the InSAR Svalbard project (Post 74, contract number: 74CO2301), with support from the Geological Survey of Norway.

The research was conducted as part of the Master thesis of Lotte Wendt at the University of Oslo and at the University Centre in Svalbard. We thank Jakob Reif, Leonie Kommerell, Vera van der Veen, Luke Simmons, Mederic Lorry and Knut I. Tveit
590 for assistance with the fieldwork and laboratory analysis. We thank Yngvar Larsen and Heidi Hindberg for their development on the GSAR processing chain and support to apply it.

References

- Aalstad, K., Westermann, S., Schuler, T. V., Boike, J., and Bertino, L.: Ensemble-based assimilation of fractional snow-covered area satellite retrievals to estimate the snow distribution at Arctic sites, *The Cryosphere*, 12, 247–270, <https://doi.org/10.5194/tc-12-247-2018>, 2018.
- 595 Aga, J., Boike, J., Langer, M., Ingeman-Nielsen, T., and Westermann, S.: Simulating ice segregation and thaw consolidation in permafrost environments with the CryoGrid community model, *The Cryosphere*, 17, 4179–4206, <https://doi.org/10.5194/tc-17-4179-2023>, 2023.
- Antonova, S., Sudhaus, H., Strozzi, T., Zwieback, S., Kääb, A., Heim, B., Langer, M., Bornemann, N., and Boike, J.: Thaw Subsidence of a Yedoma Landscape in Northern Siberia, Measured In Situ and Estimated from TerraSAR-X Interferometry, *Remote Sensing*, 10, 494, <https://doi.org/10.3390/rs10040494>, 2018.
- 600 Bai, R., Lai, Y., Pei, W., and Zhang, M.: Investigation on frost heave of saturated–unsaturated soils, *Acta Geotech.*, 15, 3295–3306, <https://doi.org/10.1007/s11440-020-00952-6>, 2020.
- Baran, I., Stewart, M. P., Kampes, B. M., Perski, Z., and Lilly, P.: A modification to the Goldstein radar interferogram filter, *IEEE Trans. Geosci. Remote Sensing*, 41, 2114–2118, <https://doi.org/10.1109/TGRS.2003.817212>, 2003.
- 605 Bartsch, A., Leibman, M., Strozzi, T., Khomutov, A., Widhalm, B., Babkina, E., Mullanurov, D., Ermokhina, K., Kroisleitner, C., and Bergstedt, H.: Seasonal Progression of Ground Displacement Identified with Satellite Radar Interferometry and the Impact of Unusually Warm Conditions on Permafrost at the Yamal Peninsula in 2016, *Remote Sensing*, 11, 1865, <https://doi.org/10.3390/rs11161865>, 2019.

- 610 Bartsch, A., Strozzi, T., and Nitze, I.: Permafrost Monitoring from Space, *Surv Geophys*, 44, 1579–1613, <https://doi.org/10.1007/s10712-023-09770-3>, 2023.
- Berardino, P., Fornaro, G., Lanari, R., and Sansosti, E.: A new algorithm for surface deformation monitoring based on small baseline differential SAR interferograms, *IEEE Trans. Geosci. Remote Sensing*, 40, 2375–2383, <https://doi.org/10.1109/TGRS.2002.803792>, 2002.
- 615 Biskaborn, B. K., Smith, S. L., Noetzi, J., Matthes, H., Vieira, G., Streletskiy, D. A., Schoeneich, P., Romanovsky, V. E., Lewkowicz, A. G., Abramov, A., Allard, M., Boike, J., Cable, W. L., Christiansen, H. H., Delaloye, R., Diekmann, B., Drozdov, D., Eitzelmüller, B., Grosse, G., Guglielmin, M., Ingeman-Nielsen, T., Isaksen, K., Ishikawa, M., Johannson, M., Johannsson, H., Joo, A., Kaverin, D., Kholodov, A., Konstantinov, P., Kröger, T., Lambiel, C., Lanckman, J.-P., Luo, D., Malkova, G., Meiklejohn, I., Moskalenko, N., Oliva, M., Phillips, M., Ramos, M., Sannel, A. B. K., Sergeev, D., Seybold, C., Skryabin, P., Vasiliev, A., Wu, Q., Yoshikawa, K., Zheleznyak, M., and Lantuit, H.: Permafrost is warming at a global scale, *Nat Commun*, 10, 264, <https://doi.org/10.1038/s41467-018-08240-4>, 2019.
- 620 Bockheim, J. G. and Hinkel, K. M.: Accumulation of Excess Ground Ice in an Age Sequence of Drained Thermokarst Lake Basins, Arctic Alaska, *Permafrost and Periglacial Processes*, 23, 231–236, <https://doi.org/10.1002/ppp.1745>, 2012.
- Bonnaventure, P. P. and Lamoureux, S. F.: The active layer: A conceptual review of monitoring, modelling techniques and changes in a warming climate, *Progress in Physical Geography: Earth and Environment*, 37, 352–376, <https://doi.org/10.1177/0309133313478314>, 2013.
- Burn, C. R., Lewkowicz, A. G., and Wilson, M. A.: Long-term field measurements of climate-induced thaw subsidence above ice wedges on hillslopes, western Arctic Canada, *Permafrost and Periglacial Processes*, 32, 261–276, <https://doi.org/10.1002/ppp.2113>, 2021.
- 630 Burn, C. R., Bartsch, A., Chakraborty, E., Das, S., Frauenfelder, R., Gärtner-Roer, I., Gislås, K. G., Herring, T., Jones, B. M., Kokelj, S. V., Langer, M., Lathrop, E., Murton, J. B., Nielsen, D. M., Niu, F., Olson, C., O’Neill, H. B., Opfergelt, S., Overduin, P. P., Schaefer, K., Schuur, E. A. G., Skierszkan, E., Smith, S. L., Stuenzi, S. M., Tank, S. E., van der Sluijs, J., Vieira, G., Westermann, S., Wolfe, S. A., and Yarmak, E.: Developments in Permafrost Science and Engineering in Response to Climate Warming in Circumpolar and High Mountain Regions, 2019–2024, *Permafrost and Periglacial Processes*, n/a, <https://doi.org/10.1002/ppp.2261>, 2024.
- 635 Cable, S., Elberling, B., and Kroon, A.: Holocene permafrost history and cryostratigraphy in the High-Arctic Adventdalen Valley, central Svalbard, *Boreas*, 47, 423–442, <https://doi.org/10.1111/bor.12286>, 2018.
- Chang, T., Yi, Y., Jiang, H., Li, R., Lu, P., Liu, L., Wang, L., Zhao, L., Zwieback, S., and Zhao, J.: Unraveling the non-linear relationship between seasonal deformation and permafrost active layer thickness, *npj Clim Atmos Sci*, 7, 1–11, <https://doi.org/10.1038/s41612-024-00866-0>, 2024.
- 640 Chen, C. W. and Zebker, H. A.: Phase unwrapping for large SAR interferograms: statistical segmentation and generalized network models, *IEEE Trans. Geosci. Remote Sensing*, 40, 1709–1719, <https://doi.org/10.1109/TGRS.2002.802453>, 2002.
- Chen, J., Wu, Y., O’Connor, M., Cardenas, M. B., Schaefer, K., Michaelides, R., and Kling, G.: Active layer freeze-thaw and water storage dynamics in permafrost environments inferred from InSAR, *Remote Sensing of Environment*, 248, 112007, <https://doi.org/10.1016/j.rse.2020.112007>, 2020.
- 645 Cigna, F., Esquivel Ramírez, R., and Tapete, D.: Accuracy of Sentinel-1 PSI and SBAS InSAR Displacement Velocities against GNSS and Geodetic Leveling Monitoring Data, *Remote Sensing*, 13, 4800, <https://doi.org/10.3390/rs13234800>, 2021.
- 650 Daout, S., Doin, M.-P., Peltzer, G., Socquet, A., and Lasserre, C.: Large-scale InSAR monitoring of permafrost freeze-thaw cycles on the Tibetan Plateau, *Geophysical Research Letters*, 44, 901–909, <https://doi.org/10.1002/2016GL070781>, 2017.
- Daout, S., Dini, B., Haerberli, W., Doin, M.-P., and Parsons, B.: Ice loss in the Northeastern Tibetan Plateau permafrost as seen by 16 yr of ESA SAR missions, *Earth and Planetary Science Letters*, 545, 116404, <https://doi.org/10.1016/j.epsl.2020.116404>, 2020.
- 655 Douglas, T. A., Turetsky, M. R., and Koven, C. D.: Increased rainfall stimulates permafrost thaw across a variety of Interior Alaskan boreal ecosystems, *npj Clim Atmos Sci*, 3, 1–7, <https://doi.org/10.1038/s41612-020-0130-4>, 2020.
- Dumais, S. and Konrad, J.-M.: Framework for thaw consolidation of fine-grained soils, *Can. Geotech. J.*, 61, 931–944, <https://doi.org/10.1139/cgj-2022-0502>, 2024.

- 660 Eckerstorfer, M. and Christiansen, H. H.: The “High Arctic Maritime Snow Climate” in Central Svalbard, Arctic, Antarctic, and Alpine Research, 43, 11–21, <https://doi.org/10.1657/1938-4246-43.1.11>, 2011.
- Everdingen, R. O. van: Multi-language glossary of permafrost and related ground- ice terms, International Permafrost Association, 1998.
- French, H. M.: Cold-Climature Weathering, in: The Periglacial Environment, John Wiley & Sons, Ltd, 47–82, <https://doi.org/10.1002/9781118684931.ch4>, 2007a.
- 665 French, H. M.: Ground Ice, in: The Periglacial Environment, John Wiley & Sons, Ltd, 153–185, <https://doi.org/10.1002/9781118684931.ch7>, 2007b.
- French, H. and Shur, Y.: The principles of cryostratigraphy, Earth-Science Reviews, 101, 190–206, <https://doi.org/10.1016/j.earscirev.2010.04.002>, 2010.
- 670 Gao, B.: NDWI—A normalized difference water index for remote sensing of vegetation liquid water from space, Remote Sensing of Environment, 58, 257–266, [https://doi.org/10.1016/S0034-4257\(96\)00067-3](https://doi.org/10.1016/S0034-4257(96)00067-3), 1996.
- GCOS. The 2022 GCOS implementation plan. Technical Report GCOS-244, World Meteorological Organization, Geneva, Switzerland, 2022.
- Gilbert, G. L., O’Neill, H. B., Nemeč, W., Thiel, C., Christiansen, H. H., and Buylaert, J.-P.: Late Quaternary sedimentation and permafrost development in a Svalbard fjord-valley, Norwegian high Arctic, Sedimentology, 65, 2531–2558, <https://doi.org/10.1111/sed.12476>, 2018.
- 675 Goldstein, R. M. and Werner, C. L.: Radar interferogram filtering for geophysical applications, Geophysical Research Letters, 25, 4035–4038, <https://doi.org/10.1029/1998GL900033>, 1998.
- Gruber, S.: Ground subsidence and heave over permafrost: hourly time series reveal interannual, seasonal and shorter-term movement caused by freezing, thawing and water movement, The Cryosphere, 14, 1437–1447, <https://doi.org/10.5194/tc-14-1437-2020>, 2020.
- 680 Hanssen-Bauer, I., Førland, E., Hisdal, H., Mayer, S., Sandø, A., and Sorteberg, A.: Climate in Svalbard 2100 – a knowledge base for climate adaptation, Norwegian Centre for Climate Services (NCCS), 2019.
- Harris, C., Kern-Luetsch, M., Christiansen, H. H., and Smith, F.: The Role of Interannual Climate Variability in Controlling Solifluction Processes, Endalen, Svalbard, Permafrost and Periglacial Processes, 22, 239–253, <https://doi.org/10.1002/ppp.727>, 2011.
- 685 Hinkel, K. M. and Nelson, F. E.: Spatial and temporal patterns of active layer thickness at Circumpolar Active Layer Monitoring (CALM) sites in northern Alaska, 1995–2000, Journal of Geophysical Research: Atmospheres, 108, <https://doi.org/10.1029/2001JD000927>, 2003.
- Humlum, O., Instanes, A., and Sollid, J. L.: Permafrost in Svalbard: a review of research history, climatic background and engineering challenges, Polar Research, 22, 191–215, <https://doi.org/10.3402/polar.v22i2.6455>, 2003.
- 690 Härtel, S. and Christiansen, H. H.: Geomorphological and Cryological map of Adventdalen, Svalbard, PANGAEA, <https://doi.pangaea.de/10.1594/PANGAEA.833048>, 2014.
- Jia, Y., Kim, J.-W., Shum, C. K., Lu, Z., Ding, X., Zhang, L., Erkan, K., Kuo, C.-Y., Shang, K., Tseng, K.-H., and Yi, Y.: Characterization of Active Layer Thickening Rate over the Northern Qinghai-Tibetan Plateau Permafrost Region Using ALOS Interferometric Synthetic Aperture Radar Data, 2007–2009, Remote Sensing, 9, 84, <https://doi.org/10.3390/rs9010084>, 2017.
- 695 Jiang, J. and Lohman, R. B.: Coherence-guided InSAR deformation analysis in the presence of ongoing land surface changes in the Imperial Valley, California, Remote Sensing of Environment, 253, 112160, <https://doi.org/10.1016/j.rse.2020.112160>, 2021.
- 700 Jorgenson, M. T. and Shur, Y.: Evolution of lakes and basins in northern Alaska and discussion of the thaw lake cycle, Journal of Geophysical Research: Earth Surface, 112, <https://doi.org/10.1029/2006JF000531>, 2007.
- Kanevskiy, M., Shur, Y., Jorgenson, M. T., Ping, C.-L., Michaelson, G. J., Fortier, D., Stephani, E., Dillon, M., and Tumskey, V.: Ground ice in the upper permafrost of the Beaufort Sea coast of Alaska, Cold Regions Science and Technology, 85, 56–70, <https://doi.org/10.1016/j.coldregions.2012.08.002>, 2013.
- 705 Kokelj, S. V. and Burn, C. R.: Near-surface ground ice in sediments of the Mackenzie Delta, Northwest Territories, Canada, Permafrost and Periglacial Processes, 16, 291–303, <https://doi.org/10.1002/ppp.537>, 2005.
- Kottek, M., Grieser, J., Beck, C., Rudolf, B., and Rubel, F.: World Map of the Köppen-Geiger climate classification updated, metz, 15, 259–263, <https://doi.org/10.1127/0941-2948/2006/0130>, 2006.

- Larsen, Y., Engen, G., Lauknes, T. R., Malnes, E., and Høgda, K. A.: A generic differential Interferometric SAR processing system, with applications to land subsidence and snow-water equivalent retrieval, in: Proceedings of the ESA Fringe Workshop, 28 November–2 December 2005, Frascati, Italy, 2005.
- Lauknes, T. R., Zebker, H. A., and Larsen, Y.: InSAR Deformation Time Series Using an L1-Norm Small-Baseline Approach, *IEEE Trans. Geosci. Remote Sensing*, 49, 536–546, <https://doi.org/10.1109/TGRS.2010.2051951>, 2011.
- Li, Y., Zuo, X., Xiong, P., You, H., Zhang, H., Yang, F., Zhao, Y., Yang, Y., and Liu, Y.: Deformation monitoring and analysis of Kunyang phosphate mine fusion with InSAR and GPS measurements, *Advances in Space Research*, 69, 2637–2658, <https://doi.org/10.1016/j.asr.2021.12.051>, 2022.
- Liu, L., Schaefer, K., Zhang, T., and Wahr, J.: Estimating 1992–2000 average active layer thickness on the Alaskan North Slope from remotely sensed surface subsidence, *Journal of Geophysical Research: Earth Surface*, 117, <https://doi.org/10.1029/2011JF002041>, 2012.
- Liu, L., Schaefer, K., Gusmeroli, A., Grosse, G., Jones, B. M., Zhang, T., Parsekian, A. D., and Zebker, H. A.: Seasonal thaw settlement at drained thermokarst lake basins, Arctic Alaska, *The Cryosphere*, 8, 815–826, <https://doi.org/10.5194/tc-8-815-2014>, 2014.
- Liu, L. and Larson, K. M.: Decadal changes of surface elevation over permafrost area estimated using reflected GPS signals, *The Cryosphere*, 12, 477–489, <https://doi.org/10.5194/tc-12-477-2018>, 2018.
- Li, Y., Zuo, X., Xiong, P., You, H., Zhang, H., Yang, F., Zhao, Y., Yang, Y., and Liu, Y.: Deformation monitoring and analysis of Kunyang phosphate mine fusion with InSAR and GPS measurements, *Advances in Space Research*, 69, 2637–2658, <https://doi.org/10.1016/j.asr.2021.12.051>, 2022.
- Mackay, J. R.: Downward water movement into frozen ground, western arctic coast, Canada, *Can. J. Earth Sci.*, 20, 120–134, <https://doi.org/10.1139/e83-012>, 1983.
- Magnússon, R. Í., Hamm, A., Karsanaev, S. V., Limpens, J., Kleijn, D., Frampton, A., Maximov, T. C., and Heijmans, M. M. P. D.: Extremely wet summer events enhance permafrost thaw for multiple years in Siberian tundra, *Nat Commun*, 13, 1556, <https://doi.org/10.1038/s41467-022-29248-x>, 2022.
- Morgenstern, N. R. and Nixon, J. F.: One-dimensional Consolidation of Thawing Soils, *Can. Geotech. J.*, 8, 558–565, <https://doi.org/10.1139/t71-057>, 1971.
- Morse, P. D., Burn, C. R., and Kokelj, S. V.: Near-surface ground-ice distribution, Kendall Island Bird Sanctuary, western Arctic coast, Canada, *Permafrost and Periglacial Processes*, 20, 155–171, <https://doi.org/10.1002/ppp.650>, 2009.
- Norwegian Meteorological Institute: Permafrost real-time monitoring, <https://cryo.met.no/en/permafrost>, last access: 26 February 2024, 2024a.
- Norwegian Meteorological Institute: Observations and weather statistics – Adventdalen station, <https://seklima.met.no/>, last access: 28 February 2024, 2024b.
- Norwegian Polar Institute: Kartdata Svalbard 1:100 000 (S100 Kartdata). Norwegian Polar Institute [data set], <https://doi.org/10.21334/npolar.2014.645336c7>, 2014a.
- Norwegian Polar Institute: Terrengmodell Svalbard (S0 Terrengmodell). Norwegian Polar Institute [data set], <https://doi.org/10.21334/npolar.2014.dce53a47>, 2014b.
- Obu, J.: How Much of the Earth’s Surface is Underlain by Permafrost?, *Journal of Geophysical Research: Earth Surface*, 126, e2021JF006123, <https://doi.org/10.1029/2021JF006123>, 2021.
- O’Neill, H. B., Smith, S. L., Burn, C. R., Duchesne, C., and Zhang, Y.: Widespread Permafrost Degradation and Thaw Subsidence in Northwest Canada, *Journal of Geophysical Research: Earth Surface*, 128, e2023JF007262, <https://doi.org/10.1029/2023JF007262>, 2023.
- O’Neill, H. B., Gilbert, G. L., and Christiansen, H. H.: Site-scale variation in ground-ice content and physical properties of loess in permafrost, Svalbard, High Arctic, <https://doi.org/10.4095/pfmq507fg6>, 2025.
- Paquette, M., Lafrenière, M. J., and Lamoureux, S. F.: Landscape influence on permafrost ground ice geochemistry in a polar desert environment, Resolute Bay, Nunavut, Arctic Science, 9, 465–482, <https://doi.org/10.1139/as-2021-0049>, 2023.
- Peng, S., Peng, X., Frauenfeld, O. W., Yang, G., Tian, W., Tian, J., and Ma, J.: Using InSAR for Surface Deformation Monitoring and Active Layer Thickness Retrieval in the Heihe River Basin on the Northeast Qinghai-Tibet Plateau, *Journal of Geophysical Research: Earth Surface*, 128, e2022JF006782, <https://doi.org/10.1029/2022JF006782>, 2023.
- Rempel, A. W.: Formation of ice lenses and frost heave, *Journal of Geophysical Research: Earth Surface*, 112, <https://doi.org/10.1029/2006JF000525>, 2007.

- 760 Rouyet, L., Lauknes, T. R., Christiansen, H. H., Strand, S. M., and Larsen, Y.: Seasonal dynamics of a permafrost landscape, Adventdalen, Svalbard, investigated by InSAR, *Remote Sensing of Environment*, 231, 111236, <https://doi.org/10.1016/j.rse.2019.111236>, 2019.
- Rouyet, L., Liu, L., Strand, S. M., Christiansen, H. H., Lauknes, T. R., and Larsen, Y.: Seasonal InSAR Displacements Documenting the Active Layer Freeze and Thaw Progression in Central-Western Spitsbergen, Svalbard, *Remote Sensing*, 13, 2977, <https://doi.org/10.3390/rs13152977>, 2021.
- 765 Schaefer, K., Liu, L., Parsekian, A., Jafarov, E., Chen, A., Zhang, T., Gusmeroli, A., Panda, S., Zebker, H. A., and Schaefer, T.: Remotely Sensed Active Layer Thickness (ReSALT) at Barrow, Alaska Using Interferometric Synthetic Aperture Radar, *Remote Sensing*, 7, 3735–3759, <https://doi.org/10.3390/rs70403735>, 2015.
- Scheer, J., Caduff, R., How, P., Marcer, M., Strozzi, T., Bartsch, A., and Ingeman-Nielsen, T.: Thaw-Season InSAR Surface Displacements and Frost Susceptibility Mapping to Support Community-Scale Planning in Ilulissat, West Greenland, *Remote Sensing*, 15, 3310, <https://doi.org/10.3390/rs15133310>, 2023.
- 770 Scherler, M., Hauck, C., Hoelzle, M., Stähli, M., and Völksch, I.: Meltwater infiltration into the frozen active layer at an alpine permafrost site, *Permafrost and Periglacial Processes*, 21, 325–334, <https://doi.org/10.1002/ppp.694>, 2010. Schuur, E. A. G., McGuire, A. D., Schädel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry, P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J. E.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171–179, <https://doi.org/10.1038/nature14338>, 2015.
- Short, N. H. and Fraser, R. H.: Comparison of RADARSAT-2 and Sentinel-1 DInSAR displacements over upland ice-wedge polygonal terrain, Banks Island, Northwest Territories, Canada, <https://doi.org/10.4095/331683>, 2023.
- 780 Shur, Y., Hinkel, K. M., and Nelson, F. E.: The transient layer: implications for geocryology and climate-change science, *Permafrost and Periglacial Processes*, 16, 5–17, <https://doi.org/10.1002/ppp.518>, 2005. Strozzi, T., Antonova, S., Günther, F., Mätzler, E., Vieira, G., Wegmüller, U., Westermann, S., and Bartsch, A.: Sentinel-1 SAR Interferometry for Surface Deformation Monitoring in Low-Land Permafrost Areas, *Remote Sensing*, 10, 1360, <https://doi.org/10.3390/rs10091360>, 2018.
- 785 Taber, S.: The Mechanics of Frost Heaving, *The Journal of Geology*, 38, 303–317, <https://doi.org/10.1086/623720>, 1930.
- Walvoord, M. A. and Kurylyk, B. L.: Hydrologic Impacts of Thawing Permafrost—A Review, *Vadose Zone Journal*, 15, vjz2016.01.0010, <https://doi.org/10.2136/vzj2016.01.0010>, 2016.
- Wang, C., Zhang, Z., Zhang, H., Zhang, B., Tang, Y., and Wu, Q.: Active Layer Thickness Retrieval of Qinghai–Tibet Permafrost Using the TerraSAR-X InSAR Technique, *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 11, 4403–4413, <https://doi.org/10.1109/JSTARS.2018.2873219>, 2018.
- 790 Wang, L., Marzahn, P., Bernier, M., and Ludwig, R.: Sentinel-1 InSAR measurements of deformation over discontinuous permafrost terrain, Northern Quebec, Canada, *Remote Sensing of Environment*, 248, 111965, <https://doi.org/10.1016/j.rse.2020.111965>, 2020.
- 795 Wang, L., Zhao, L., Zhou, H., Liu, S., Hu, G., Li, Z., Wang, C., and Zhao, J.: Evidence of ground ice melting detected by InSAR and in situ monitoring over permafrost terrain on the Qinghai-Xizang (Tibet) Plateau, *Permafrost and Periglacial Processes*, 34, 52–67, <https://doi.org/10.1002/ppp.2171>, 2023.
- Wendt, L.: Ground ice contents and InSAR displacements from Adventdalen, Svalbard [data set]. Zenodo, <https://doi.org/10.5281/zenodo.11187360>, 2024.
- Yalvac, S.: Validating InSAR-SBAS results by means of different GNSS analysis techniques in medium- and high-grade deformation areas, *Environ Monit Assess*, 192, 120, <https://doi.org/10.1007/s10661-019-8009-8>, 2020.
- 800 Zwieback, S. and Meyer, F. J.: Top-of-permafrost ground ice indicated by remotely sensed late-season subsidence, *The Cryosphere*, 15, 2041–2055, <https://doi.org/10.5194/tc-15-2041-2021>, 2021.
- Zwieback, S., Iwahana, G., Sakhalkar, S., Biessel, R., Taylor, S., and Meyer, F. J.: Excess Ground Ice Profiles in Continuous Permafrost Mapped From InSAR Subsidence, *Water Resources Research*, 60, e2023WR035331, <https://doi.org/10.1029/2023WR035331>, 2024.