

The Size Distributions of Fractures and Earthquakes: Implications for Orogen-Internal Seismogenic Deformation

Sandro Truttmann¹, Tobias Diehl², Marco Herwegh¹, Stefan Wiemer²

¹Institute of Geological Sciences, University of Bern, Bern, 3012, Switzerland

²Swiss Seismological Service, ETH Zürich, Zürich, 8092, Switzerland

Correspondence to: Sandro Truttmann (sandro.truttmann@gmail.com)

Abstract. Pre-existing geological discontinuities such as faults and fractures represent structural and mechanical discontinuities in rocks which influence earthquake processes. As earthquakes occur in the subsurface, seismogenic reactivation of pre-existing fracture networks is difficult to investigate in natural settings. However, it is well-known that there exists a physical link between both fractures and earthquakes since an earthquake's magnitude is related to the ruptured fracture area and therefore fracture length. Furthermore, fractures and earthquakes exhibit similar statistical properties, as their size distributions follow power laws.

In this study, we exploit the relation between the size (or length) distributions of pre-existing fractures and earthquakes to decipher the seismic deformation processes within the exhumation-related orogen-internal setting of the Southwestern Swiss Alps, which due to its well-monitored seismic activity and the excellent outcrop conditions provides an ideal study site. Characterizing exhumed fracture networks from different tectonic units based on multi-scale drone-based mapping, we find that power law exponents of 3D fracture size distributions generally range between 3 and 3.6. Comparing these values with the depth-dependent exponents of estimated earthquake rupture lengths, we observe significantly larger values of 5 to 8 for earthquake ruptures at shallow depths (< 3 km below sea level (BSL)). At intermediate crustal depths (~3 to 9 km BSL), the power law exponents of fractures and earthquakes appear to be similar. These findings imply depth-dependent differences in the seismogenic reactivation of pre-existing fractures in the study region: while partial rupturing is the prevailing deformation mechanism at shallow depths, fractures are more likely to rupture along their entire length at intermediate crustal depths. Therefore, the present-day near surface differential stresses are likely insufficient to rupture entire pre-existing fractures seismogenically. Our findings have direct implications for seismic hazard considerations, as earthquakes that rupture along entire fractures appear to become less likely with decreasing depth.

hat gelöscht: Faults

hat gelöscht: fault

hat gelöscht: faults

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: faults

hat gelöscht: faults

hat gelöscht: the size distribution of

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: networks

hat gelöscht: faults

hat gelöscht: faults

hat gelöscht: faults

hat gelöscht: faults

hat gelöscht: faults

1 Introduction

Within the Earth's crust, the occurrence of earthquakes is commonly attributed to the reactivation of pre-existing geological discontinuities, commonly referred to as [fractures](#). Nevertheless, in regions characterized by extensive [fracturing](#) and scattered seismic activity, like in collisional orogens such as the Alps, the task of assigning earthquakes to specific [fracture](#) segments becomes notably complex. This challenge limits our ability to decipher the current deformation processes and discern the significance of such intricate pre-existing [fracture](#) networks in influencing the underlying seismic events.

Despite their complexity, [fractures](#) in nature often appear qualitatively self-similar at different scales. This qualitative self-similarity can be quantified by power law distributions, [which inherently implies](#) scale invariance and thus [the](#) absence of a characteristic length scale. [This means that, strictly speaking, fracture segments do not have a unique length, but that a measure of the length of a fracture is always dependent on the scale of observation.](#) Previous studies have shown that a power law is best suited for describing the size (or length) distribution of natural [fracture](#) traces (Odling, 1997; Bonnet et al., 2001; Torabi and Berg, 2011; Yielding et al., 1996; Davy et al., 2010; Scholz, 2019), even if other types [such as exponential or log-normal distributions](#) have been proposed (Cowie et al., 1995, 1993; Ackermann et al., 2001; Nicol et al., 1996). A power law distribution defines the number n of features of a given length l as:

$$n(l) = c l^{-\alpha} \quad (1)$$

where α represents the power law exponent (i.e., the slope of the linear trend in a log-log plot), and c denotes the density constant. The power law exponent α is of main interest herein, since it measures the relative proportion of short and long features (Davy, 1993; Pickering et al., 1995). As [fracture](#) data is often acquired at a single mapping scale, a major problem for the identification of power law distributions is the limited dynamic length range of the [fracture](#) traces and the uncertainties related to sampling bias, which can lead to size distributions that appear to deviate from power laws (Odling, 1997; Scholz, 2007, 2019; Torabi and Berg, 2011).

Similar to [fractures](#), it is well-known that earthquake magnitudes follow power law distributions, often referred to as the Gutenberg-Richter law (Gutenberg and Richter, 1944) or magnitude-frequency distribution, implying that also earthquakes are scale invariant. As proposed by Hatton et al. (1993), "the observed fractal nature of both [fracture](#) length distributions and earthquake magnitude-frequency distributions [suggest](#) that there may be a relationship between the structure of active [fracture](#) systems and the resulting seismicity." This is supported by the fact that the statistical properties of micro-fracturing at the laboratory scale are similar to earthquakes (e.g., Scholz, 1968; Mogi, 1962). Furthermore, [fractures](#) represent the geometrical anisotropies that provide planes of weakness within the rock masses and are thus most likely to be reactivated seismically if suitably oriented with respect to the local stress field. [Fractures](#) and earthquakes can thus be regarded as the long- and short-time-scale phenomena of brittle tectonics (Scholz, 2019), and there should exist an inherent relation between the size distributions of [fractures](#) and earthquakes (Scholz, 1998; Turcotte, 1997). However, the relationship between these size

hat gelöscht: faults

hat gelöscht: faulting

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: faults

hat gelöscht: inherently

hat gelöscht: implying

hat gelöscht: , which is different to any other statistical distribution...

hat gelöscht:

hat gelöscht: of distributions

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: suggests

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: and fractures

hat gelöscht: Faults

hat gelöscht: faults

hat gelöscht: 2019,

distributions in natural datasets has yet received little attention, mainly due to the challenging acquisition of reliable quantitative information of [fracture](#) networks due to limited outcrop conditions and small dynamic range. The Southwestern Swiss Alps, with (i) a long-lasting exhumation history [of about 20 million years](#) (e.g., Egli et al., 2017; Herwegh et al., 2020; Boutoux et al., 2016; Cardello et al., 2024), (ii) well-preserved extensive Neogene faulting in high-Alpine regions with formation depths similar to the present-day earthquakes (Cardello and Mancktelow, 2015; Huggenberger and Aebli, 1989; Cardello et al., 2024), (iii) an increased and exceptionally well-recorded seismic activity (e.g., Diehl et al., 2021a; Lee et al., 2022), and (iv) on-going vertical tectonics (e.g., Ustaszewski and Pfiffner, 2008; Brockmann et al., 2012; Herwegh et al., 2023; Piña-Valdés et al., 2022) implying a similar stress field to the time of [fracture](#) formation, provide a unique natural laboratory to investigate the link between the size distributions of [fractures](#) and earthquakes. Owing to the abundant occurrence of thermal springs, the region is also a potential target for the exploitation of geothermal energy, where knowledge of faulting at depth and constraints on the potential risks of induced seismicity are critical.

Herein, we explore the relation between the size distributions of [fractures](#) ([fracture length](#)) and earthquakes ([rupture length](#)) and its implications for orogen-internal seismogenic deformation in the exhumation-related (vertical) tectonic setting of the Rawil depression region (Southwestern Swiss Alps) ([Figure 1](#)). We use field observations and drone-based [fracture](#) trace mapping on multiple scales to quantitatively characterize the size and orientation distributions of [fracture](#) networks in different tectonic units. We employ a multi-scale power law fitting approach on the [fracture](#) network data, which allows to expand the limited dynamic length range of single-scale observations. We then exploit the relationship of the [fracture](#) orientation and size distributions with similar properties of the recent seismic [activity](#) and discuss the implications for seismogenic [fracture](#) reactivation processes in the subsurface.

2 Regional Setting

In the framework of a nation-wide seismic hazard assessment, Switzerland was zoned into different seismotectonic domains (Wiemer et al., 2016). We herein focus on the seismotectonic domain z21 in the Southwestern Swiss Alps, mainly due to its well-exposed rock outcrops and the exceptionally well-monitored [natural](#) seismic activity ([Fig. 1](#)). In the study area, several External Crystalline Massifs (ECMs) are exposed: the Aiguilles Rouges massif in the West, and the Aar and Gastern massifs in the East ([Fig. 1a](#)). Between their exhumed parts, these basement units form a saddle-like structure, commonly referred to as the Rawil depression, which is overthrust by the Helvetic limestone nappes (Burkhard, 1988; Dietrich, 1989; Dietrich and Casey, 1989; Ramsay, 1989, 1981). The Helvetic nappe system is composed of different, heavily folded nappes, which have been stacked on top of each other during an early thin-skinned deformation phase of the Alpine orogen ([thrusts in Fig. 1a](#); e.g., Musso Piantelli et al., 2022; Escher et al., 1993; Pfiffner 1993; Burkhard, 1988; Ramsay, 1981, 1989). In a late-Alpine stage, dominated by thick-skinned vertical tectonics, both the crystalline basement units and the Helvetic limestones have been heavily affected by late-Alpine deformation, resulting in pervasive and complex faulting that affects all tectonic units [and crosscut the older thrusts](#) ([faults in Fig. 1a](#); Cardello and Mancktelow, 2015; Gasser and Mancktelow, 2010; Huggenberger

hat gelöscht: fault

hat gelöscht: during

hat gelöscht: several

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: activity, and

hat gelöscht: fault

hat gelöscht: enhanced

and Aebli, 1989; Herwegh et al., 2020, 2023; Cardello et al., 2024). Some of the predominantly subvertical late-Alpine faults have been linked to the regional seismic activity (Cardello and Mancktelow, 2015; Pavoni, 1980a; Ustaszewski et al., 2007; Ustaszewski and Pfiffner, 2008; Pavoni and Mayer-Rosa, 1978; Cardello et al., 2024). [With the exception of the Rhône-Simplon Fault Zone \(RSF\), a steeply south-dipping dextral strike-slip fault system](#) along the southern margin of the study area (e.g., Campani et al., 2014, 2010; Egli and Mancktelow, 2013; Mancktelow, 1985), regional-scale faults are largely absent at the surface of the study area. The nappes of the Helvetic units are separated by shallow-dipping thrust faults which are arguably not seismically active nowadays (Lee et al., 2022), so we herein do not discuss these thrusts further.

The presently dense seismic monitoring network in the region resolves the seismic activity exceptionally well, detecting earthquakes with a magnitude of completeness M_c of about $M_L \approx 1.5$ until 2016 and about $M_L \approx 1.0$ thereafter (Diehl et al., 2021a; Lee et al., 2023). The earthquakes predominantly occur in a NE-SW striking, elongated corridor in the center of the Rawil depression (Fig. 1b), which is referred to as the conceptual Rawil Fault Zone (RFZ) (Lee et al., 2023). The seismicity at depth arguably reactivates a complex system of subvertically oriented faults (Diehl et al., 2021a, 2018; Lee et al., 2023; Truttmann et al., 2023). This is supported by the variety of previously published, predominantly strike-slip-type focal mechanisms shown in Fig. 1b (Baer et al., 2005, 2003, 1997; Deichmann et al., 2012, 2006, 2002, 2000; Delacou et al., 2005; Diehl et al., 2021a, 2018, 2013; Jimenez and Pavoni, 1983; Maurer, 1993; Maurer and Deichmann, 1995; Pavoni et al., 1997; Truttmann et al., 2023), suggesting an overall transtensional regime (Kastrup et al., 2004; [Houlié et al., 2018](#); [Maurer et al., 1997](#)).

The Southwestern Swiss Alps are further characterized by their long-lasting vertical tectonics. During the latest stage of the Alpine orogeny, thick-skinned tectonics prevailed and led to the exhumation of the ECMs, [associated with intense faulting \(faults in Fig. 1a; e.g., Herwegh et al., 2020; 2023; Boutoux et al., 2016\)](#). Recent uplift rates in the range of 1 mm per year (Brockmann et al., 2012; Piña-Valdés et al., 2022) and the pronounced seismic activity document the ongoing exhumation (Figs. 1b and c). As a consequence of these long-lasting vertical tectonics, a similar stress field likely prevailed during both the formation of the exhumed [fracture](#) networks and today's earthquakes.

In summary, the investigated seismotectonic domain reveals complex and pervasive [fracture](#) networks, well-recorded seismic activity, and long-lasting vertical tectonics. In our view, the Rawil depression region in the Southwestern Swiss Alps thus provides a unique natural laboratory to elaborate the statistical link between [fractures](#) and earthquakes in an orogen-internal setting.

hat gelöscht: Apart from the

hat gelöscht:

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

175 **3 Data and Methods**

3.1 Fracture Networks

3.1.1 Fracture Data

We selected four sites along the Rawil depression to conduct detailed analysis of the [fracture](#) networks in the ECM units of (A) the Aiguilles Rouges massif and (D) the Gastern massif, as well as in the Helvetic limestones of (B) the Diablerets and (C) the Wildhorn nappes (Fig. 1). As pronounced glacial and fluvial erosion, vegetation and Quaternary cover greatly limit the outcrop conditions and thus affect reliable sampling of [fracture](#) networks, we selected sites with little weathering, vegetation, and Quaternary cover, as well as flat terrain to minimize topographic effects [in fracture lengths and directions](#) (Baumberger et al., 2022), which were encountered only at high altitude sites.

At each site, we performed detailed characterization of the [fracture](#) networks (Fig. 2a). We carried out drone-based surveys (DJI Mavic 2 Pro; Hasselblad L1D-20c camera) at two different observation heights to produce orthorectified images (orthophotos) and Digital Elevation Models (DEMs) with Agisoft Metashape (v1.6.6) at mm and cm ground resolutions (see [Table 1](#)). Additionally, we used orthophoto and DEM products at dm ground resolution from the Federal Office of Topography swisstopo. We then manually digitized the visible [fracture](#) traces at fixed mapping scales of 1:10, 1:100, and 1:1000 within circular mapping areas, preferring circular extents to minimize potential orientation bias (e.g., Bonnet et al., 2001) ([Table 1](#), Fig. 2a). [Fracture](#) traces are herein defined as curvilinear features that represent the intersections between a [fracture](#) plane and the Earth's surface (Baumberger et al., 2022; O'Leary et al., 1976), which serve as a proxy for the pre-existing geometrical anisotropies. [Because](#) such pre-existing anisotropies are likely to be reactivated also in today's stress field if suitably oriented, and [because](#) the power law exponents are insensitive to [fracture](#) type (Bonnet et al., 2001; Marrett et al., 1999), we chose [not to distinguish between fracture types in our analysis](#). Unambiguous differentiation of [fractures](#) from other discontinuities such as bedding structures based on remote sensing alone is often ambiguous, which is why we systematically validated the [fracture](#) maps in the field to ensure their representativeness of the real [fracture](#) populations. We also measured both the 3D orientations of the [fractures](#) and where possible the kinematic indicators ([slickenfibres](#), [slickolites](#)) in the field.

Despite thorough data collection, mapping of [fracture](#) traces includes several limitations. The detectability of [fracture](#) traces by remote sensing depends on both the [fracture](#) length, the image resolution, the mapping scale, and the degree of erosion of the related tectonites (i.e., the resulting morphological incision). Furthermore, image artifacts caused by illumination or shadow effects in low contrast domains or by partial coverage of the bedrock by Quaternary debris or vegetation can prohibit the appropriate identification of [fractures](#). These factors lead to deviations from the true [fracture](#) trace length which in terms influences the quantitative analysis (e.g., Cao and Lei, 2018). To minimize such potential artifacts, we carefully selected sites with little Quaternary cover and topographic effects, and acquired the images in similar light conditions. Areas covered by debris or vegetation were excluded as "no data" areas. The influence of the remaining uncertainties is further discussed in section 5.1.

hat gelöscht: Fault

hat gelöscht: Fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: Table 1

hat gelöscht: fault

hat gelöscht: '

hat gelöscht: Table 1

hat gelöscht: Fault

hat gelöscht: fault

hat gelöscht: As

hat gelöscht: the

hat gelöscht: of fracture

hat gelöscht: to incorporate also mode I fractures

hat gelöscht: but, for the sake of simplicity, generally refer to these geological discontinuities as faults

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: i.e., slickensides

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

3.1.2 Statistical Analysis of Fracture Networks

We used the data from the fracture trace maps to quantify both the orientation and size distributions of the fracture networks. To describe their 2D orientation distribution, we derived length-weighted rose diagrams from the fracture traces for each individual dataset presented in Table 1 (Fig. 2a). Additionally, we used field measurements to characterize the 3D orientation distribution of the fracture networks, as well as their kinematics. The 2D and 3D distributions capture different aspects of the fracture orientations: while the 3D distribution yields information about the dip range of the fracture sets, the 2D orientation distribution from the fracture trace maps represents a better-suited quantitative measure of the dominant strike orientations. In a next step, we evaluated the size distribution of the fracture networks from the four sites. We assume an underlying power law distribution, which is most commonly used to model fracture size distributions (e.g., Bonnet et al., 2001; Bour et al., 2002; Odling, 1997; Scholz et al., 1993; Torabi and Berg, 2011; Yielding et al., 1996; Davy et al., 2010; Scholz, 2019). In nature, all power laws must have upper and lower limits (Bonnet et al., 2001; Torabi and Berg, 2011; Turcotte, 1997, 1989). For fractures, the upper limit is likely related to the thickness of the crust or the stratigraphic layering, while the lower limit is constrained by a physical length scale (e.g., grain size). Defining these limits is nontrivial, and deviations from a power law in the tails of the size distribution of fracture networks commonly occur presumably independent of the physical limits of the system. Instead, sampling bias such as truncation and censoring effects can result in size distributions that appear to be exponential or log-normal (Bonnet et al., 2001; Odling, 1997; Odling et al., 1999; Torabi and Berg, 2011). Due to the underestimation of the frequency of small fractures, these structures are systematically underrepresented due to the limited resolution of the orthophotos and DEM's, which is commonly referred to as truncation (Bonnet et al., 2001; Bour et al., 2002; Torabi and Berg, 2011) (Fig. 2b). The truncation length l_{trunc} defines the length above which fractures can be reliably detected. Similarly, large fractures are more likely to lie partially outside the boundaries of the sampling area, so their lengths and frequencies are often systematically underestimated (Bonnet et al., 2001; Pickering et al., 1995). This leads to a steepening of the curve at the upper tail of the distribution, commonly called censoring effect (Fig. 2b). Note that the fracture lengths at which truncation and censoring effects occur do not necessarily correspond to the true physical limits of the power law distribution (Bonnet et al., 2001). Since both truncation and censoring effects in natural fracture datasets are often rather pronounced, it is a major challenge to identify the scaling range where the distribution follows a power law. It is therefore crucial to ensure that the power law fitting is based on a large range of fracture lengths, ideally spanning values over 2 to 3 orders of magnitude (Bonnet et al., 2001). Practical issues and mapping bias (see previous section), however, prevent the acquisition of high-resolution orthophotos over large areas that would be required for this. On a single observation scale, the length range is thus limited, rendering the identification of power law distributions challenging (Bonnet et al., 2001; Bour et al., 2002; Davy et al., 2006; Pickering et al., 1995; Scholz, 2007). To overcome these limitations, we herein used a multi-scale approach, combining the observations from different mapping scales (Fig. 2b). The basic idea is to normalize the fracture density with the fractal area, which allows to greatly extend the

hat gelöscht: Fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: Table 1

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: (or length)

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

length range (e.g., Bonnet et al., 2001; Heffer and Bevan, 1990; Odling, 1997). The normalized density size distribution is given as (Bour et al., 2002; Davy et al., 1990)

$$n(l, L) = c l^{-\alpha_F} L^D \quad (2)$$

with l as the [fracture](#) trace length, c representing the [fracture](#) density, α_F as the power law exponent of the [fracture](#) networks, and L^D denotes the area term. The latter is defined by the diameter of the study area L and the fractal dimension D of the [fracture](#) trace barycenters (i.e., the centroids of the [fracture](#) traces) and fixes the number of structures of a given length per unit fractal area independently from the observation scale and is thus a scale-independent measure (Davy et al. 1990, Bour et al. 2002). This allows for comparison of the data obtained at different scales.

The fractal dimension D was derived with the two-point correlation function C_2 (Hentschel and Procaccia, 1983; Vicsek, 1992), which describes the spatial correlation of [fracture](#) trace barycenters and has been suggested as an appropriate measure for D of [fracture](#) networks (Bonnet et al., 2001; Bour et al., 2002; Bour and Davy, 1999; Davy et al., 1990). It is defined as:

$$C_2(r) = \frac{1}{N^2} N_d(r) \quad (3)$$

where N is the total number of [fracture](#) trace barycenters and N_d is the number of pairs whose barycenter interdistance is less than r (Hentschel and Procaccia, 1983). For fractal populations, C_2 scales with r as r^D , with D as the correlation dimension.

Following Bonnet et al. (2001), we estimated the fractal dimension by identifying a plateau of the local slopes of D .

To then fit the [fracture](#) trace data with equation (2), we only extracted the unbiased data from each [fracture](#) trace map. To account for truncation effects, we determined the truncation length l_{trunc} with the [statistical](#) approach of Alstott et al. (2014) and Clauset et al. (2009), using the Kolmogorov-Smirnov (KS) distance as a measure for the goodness of fit between the fitted power law and the data. Hereby, the minimum KS distance defines l_{trunc} (Fig. S1). We then removed all [fracture](#) traces with lengths smaller than l_{trunc} . We also accounted for censoring effects by removing [fractures](#) that intersect the sampling window (Lei et al., 2015). The [cleaned fracture densities](#) of each mapping scale $n(l)$ were then normalized with the area term L^D . We finally fitted the normalized dataset with a power law using Maximum Likelihood Estimation (MLE). This is commonly preferred over linear regression (Clauset et al., 2009; Goldstein et al., 2004). We finally obtained power law size distributions which, owing to the multi-scale approach, span several magnitudes of [fracture](#) lengths (Fig. 2b). As the number of [fracture](#) traces is critical for the reliable extraction of power law exponents (Zeeb et al., 2013), and the number of [fractures](#) is rather small for non-dominant orientations, we chose to not incorporate orientation-dependent α_F values but rather derived a single bulk value of α_F for the respective site.

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: (r)

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: measured

hat gelöscht: fault

hat gelöscht: ies

hat gelöscht: fault

hat gelöscht: s

hat gelöscht: fault

hat gelöscht: fault

3.2 Earthquakes

We analyzed the recent seismicity within four subdomains of the seismotectonic domain z21 of Wiemer et al. (2016), each associated with one of the [fracture](#) mapping sites A to D (Fig. 1b). In analogy with the [fracture](#) data, we characterized both the orientation and size distributions of the earthquakes within these subdomains.

The orientation of the rupture plane activated during an earthquake often remains unknown, and the determination of the orientation distribution of seismic ruptures is thus challenging. However, focal mechanisms (e.g., determined by the polarities of the first arriving P waves) give a first-order constraint on the orientation of the ruptured plane for earthquakes with significant energy release. Such first-motion focal mechanisms yield an ambiguous solution with two possible rupture plane orientations. Additional constraints such as the spatial distribution of hypocenters can in some cases be used to decipher the activated ruptured plane (e.g., Truttmann et al., 2023), but this information is often lacking. For this study, we compiled previously published focal mechanism data as shown in Fig. 1b (Baer et al., 2005, 2003, 1997; Deichmann et al., 2012, 2006, 2002, 2000; Delacou et al., 2005; Diehl et al., 2021a, 2018, 2013; Jimenez and Pavoni, 1983; Maurer, 1993; Maurer and Deichmann, 1995; Pavoni et al., 1997; Truttmann et al., 2023), which [partially](#) incorporates information about the active rupture plane [deciphered from the spatial distribution of fore- and aftershocks](#). Based on these focal mechanisms, we evaluated the 3D orientation distribution of earthquake ruptures within each subdomain by plotting the nodal planes in stereographic projections and incorporate, where available, the information about the effectively ruptured plane.

We consecutively analyzed the size distribution of earthquakes, using the data from the earthquake catalog published by Diehl et al. (2021b) [only considering natural earthquakes](#) (Fig. 1b). Indications exist that the M_L magnitudes in the SED bulletin include inconsistencies between earlier and modern epochs due to changes in instrumentation and determination procedures of magnitudes (Staudenmaier et al., 2018). We therefore only considered events after 2005, for which the M_L magnitudes are derived from digital data in a relatively consistent manner. [Since](#) we are interested in a measure that is directly comparable to the size distribution of [fractures](#), [here we derive](#) the size distribution of earthquake ruptures, [where size refers to length \$l_R\$, defined as the diameter of a circular earthquake rupture](#), rather than magnitude or seismic moment, which serves as a proxy for the recently activated part of the [fracture](#) network at depth. This approach is justified by the fact that the earthquake magnitude is directly related to the ruptured [fracture](#) length (e.g., Wells and Coppersmith, 1994). Therefore, we translated the local magnitudes M_L to the earthquake rupture lengths l_R . After converting the recorded local earthquake magnitude M_L to the moment magnitude M_w after Goertz-Allmann et al. (2011) using an empirical M_L - M_w scaling relationship, we derived the earthquake rupture area A_R , given as:

$$A_R = 10^{\frac{(M_w - a)}{b}} \quad (4)$$

where a and b represent constants derived from empirical scaling relations (e.g., Wells and Coppersmith, 1994). We herein used three different scaling relations determined for strike-slip earthquakes in stable continental regions with $a = 4.18$ and $b =$

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: partly

hat gelöscht: As

hat gelöscht: fault

hat gelöscht: we herein

hat gelöscht: derived

hat formatiert: Tiefgestellt

hat gelöscht: the

hat gelöscht: fault

hat gelöscht: fault

1 (Leonard, 2014), $a = 3.49$ and $b = 0.94$ (Thingbaijam et al., 2017), and $a = 3.98$ and $b = 1.02$ (Wells and Coppersmith, 1994). From A_R , we then calculated the earthquake rupture length l_R under the assumption of a circular rupture plane. Based on this new dataset containing a rupture length for each earthquake, we then employed the MLE power law fitting procedure of Alstott et al. (2014) and Clauset et al. (2009) using equation (1) to derive the power law exponent α_R of the earthquake ruptures. α_R represents a modified b-value that is the commonly used power law exponent in seismology. Similar to the fitting procedure of the [fracture](#) data, we used the minimum KS distance as a measure for the optimal truncation length (Fig. S2), which is comparable to M_c often used by seismologists.

4. Results

4.1 Fracture Networks

390 Based on remote sensing, we [mapped](#) between 1608 and 10894 [fracture](#) traces [at](#) each of the 3 different [mapping scales on the](#) 4 study sites (Table 1, Fig. 3). In the following, we use these data to characterize the [fracture](#) networks in terms of their orientation and size distributions.

4.1.1 Orientation Distribution

Site A is characterized by three main sets of subvertical [fractures](#) identified in the field (Fig. 4a): a NE-SW striking set dipping towards the SE (A_I) and [similarly striking but NW dipping](#) set (A_{II}), as well as an almost perpendicular NW-SE striking set (A_{III}). Set A_I consistently indicates normal movements, while A_{II} [fractures](#) reveal a reverse shear sense (Fig. 4b). For set A_{III} , we only observed a single kinematic indicator, revealing oblique sinistral movements. The length-weighted rose diagrams for site A suggest that the NW-SE striking A_{III} [fractures](#) are the dominant set (Fig. 3).

400 At site B, the measured [fracture](#) planes reveal a similar 3D orientation distribution with a slight [clockwise](#) rotation compared to site A (Fig. 4c). The NW-SE striking set B_I , dipping SE, is oriented parallel to [fracture](#) set B_{II} , and perpendicular to the NW-SE striking set B_{III} . Both B_I and B_{II} exhibit dextral strike-slip movements (Fig. 4d), while [fractures](#) of set B_{III} mostly indicate normal movements. The rose diagrams in Fig. 3 show that the NE-SW striking B_I and B_{II} [fractures](#) are the most prominent sets at all scales.

Also at site C, NE-SW striking and steeply SE (C_I), respectively, NW dipping [fractures](#) (C_{II}) are the dominant sets (Fig. 4e). Set C_I exhibits large variations in strike directions, ranging from NNE-SSW to E-W. Both sets predominantly imply dextral strike-slip movements (Fig. 4f). The rose diagrams indicate that E-W to ENE-WSW are the dominant [fracture](#) strike orientations, corresponding to both [fracture](#) sets C_I and C_{II} (Fig. 3).

410 The [fracture](#) network at site D is characterized by two main sets, with a diffusely oriented, approximately NE-SW striking set (D_I) and a dominant subvertical, NW-SE striking [fracture](#) set (D_{II}) (Fig. 4g). For neither set D_I nor D_{II} , the kinematic indicators reveal a clear pattern (Fig. 4h). The rose diagrams in Fig. 3 reveal a bimodal distribution, with both sets D_I and D_{II} equally abundant.

hat gelöscht: fault

hat gelöscht: Fault

hat gelöscht: identified

hat gelöscht: '

hat gelöscht: '

hat gelöscht: fault

hat gelöscht: on

hat gelöscht: 12

hat gelöscht: maps

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: SE dipping

hat gelöscht: a NW dipping

hat gelöscht: SW dipping

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: clock-wise

hat gelöscht: SE dipping

hat gelöscht:

hat gelöscht: strikes

hat gelöscht: the NW dipping fault

hat gelöscht: SW dipping

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: SE

hat gelöscht: and

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: SE dipping

hat gelöscht: fault

445 In summary, we document a change in dominant fracture orientations, from NW-SE striking in the Aiguilles Rouges massif (site A), to E-W and NE-SW dominated in the limestone nappes in the center of the Rawil depression (sites B and C), to a NE-SW and NW-SE bimodal fracture distribution in the Gastern massif (site D).

hat gelöscht: fault

hat gelöscht: fault

4.1.2 Size Distribution

450 To combine the fracture trace data from the different scales, we first derive the fractal dimension D for each fracture trace map. Using the two-point correlation function C_2 , we obtain values of D between 1.64 and 1.83 (Fig. 5), suggesting a relatively homogeneous spatial fracture distribution, as $D = 2$ would imply a perfectly homogeneous spatial distribution. This is in good agreement with the qualitative observation of widespread and pervasive fracturing in the area. Since the datasets from different scales reveal consistent values of D, we argue that the fractal dimension is scale-independent.

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: faulting

455 Using the obtained values of D for the area normalization (see section 3.1.2), we find that the normalized fracture size data follows a linear trend in a log-log plot for all study sites, supporting the assumption of an underlying power law distribution (Fig. 6). The power law scaling seems to hold over a length range of three orders of magnitudes for fracture lengths between ca. 10^0 to 10^3 m. However, as can be seen for the data of site C, for example, individual datasets slightly deviate from the power law behavior. Such deviations can be caused by sampling bias such as the choice of the sampling area of the consecutive mapping scale, since subareas with anomalously high or low fracture densities can lead to an offset in the multi-scale size distributions. We will further discuss potential uncertainties in section 5.1.

hat gelöscht: fault

hat gelöscht: length

hat gelöscht: fault

460 The obtained fracture power law exponents α_f , which are of main interest for comparison with the earthquake data, range between 2.43 and 2.73 (Fig. 6), which suggests that little regional variations exist. As the fracture networks are sampled in both crystalline basement rocks as well as in the Helvetic limestone units, α_f seems to be relatively insensitive to differences in lithologies. For the dimensionless fracture density term c , we observe somewhat larger variations. With a value of $c = 1.57 \cdot 10^{-4}$, site B reveals the lowest fracture density term, while sites A, C, and D show larger c values of up to $5.76 \cdot 10^{-4}$. Even though all c values range in the same order of magnitude, the fracture density term varies by a factor of three for the different sites, however without any clear regional trends.

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

4.2 Earthquake Ruptures

470 Next, we assess the orientation and size distributions of seismic rupturing at depth, derived from focal mechanism data and earthquake rupture lengths for the different subdomains denoted in Fig. 1b. We therefore use a compilation of 69 previously published focal mechanisms and 4612 earthquakes of the earthquake catalog of Diehl et al. (2021b) that lie within the seismotectonic domain z21 shown in Fig. 1 (see section 3.2).

hat gelöscht: '

4.2.1 Orientation Distribution

475 The data from the focal mechanism catalog reveals consistent rupture orientations within the different subdomains, with known earthquake ruptures predominantly occurring along subvertical, E-W to NE-SW striking planes (Fig. 7). While in subdomain

495 A in the Aiguilles Rouges massif, the known active planes dip steeply towards the South (Fig. 7a), steeply north-dipping earthquake ruptures are observed additionally in the central and eastern subdomains C and D (Figs. 7c and d). For the data of subdomain B, currently no active rupture planes are provided in the published focal mechanism data, but ENE-WSW striking nodal planes render similar active rupture orientations possible (Fig. 7b). Together with the observation of the NE-SW directed clustering of the seismicity (Fig. 1b), this implies that the majority of the seismicity in the Rawil region occurs along subvertically oriented, E-W to NE-SW striking [fractures](#).

hat gelöscht: fault

4.2.2 Size Distribution

500 As shown in Fig. 8, the size distribution of earthquake ruptures for the different subdomains all follow power laws at length scales between about 10^2 and 10^3 m. For the data of subdomains A and D, we observe a deviation from power law behavior in the upper tail of the distribution, likely related to individual larger earthquakes. In subdomain A, this can be attributed to the $M_L = 4.9$ Vallorcine earthquake in 2005 (Fréchet et al., 2011), while in subdomain D, it relates to the $M_L = 4.1$ Salgesch earthquake in 2016 (Diehl et al., 2018). In subdomains B and C, we observe well-defined power law distributions for the earthquake rupture lengths. The three used empirical $M_w - A_R$ scaling laws all yield similar values of α_R , which indicates that 505 a significant dependency from the used empirical scaling relations on the obtained power law exponents appears unlikely (Fig. 8). In general, the derived earthquake rupture power law exponents α_R all range between values of 3.88 and 4.81 [regardless of the used earthquake rupture scaling law \(L14, T17, and WC94\)](#), implying little along-strike variations in the α_R values. As the size distribution of earthquakes often shows variations with depth, we further investigate a possible depth-dependency in section 5.4.

510 Overall, the α_R and α_F exponents exhibit significant differences that are unlikely to be solely attributed to uncertainties. These differences will be further explored in the discussion section.

5. Discussion

5.1 Uncertainties in [Fracture](#) Size Distributions

In the following, we discuss two of the major uncertainties related to data sampling that can affect the [fracture](#) size distributions: 515 (i) the finite mapping resolution of [fracture](#) trace lengths, and (ii) the choice of the subsampling mapping area as well as the potential bias due to manual mapping.

Mapping [fracture](#) traces on a fixed mapping scale on imagery with a given finite resolution necessarily leads to a bias in the derived [fracture](#) lengths, as the terminations of the [fracture](#) traces can only be mapped with a certain accuracy that is dependent on the image resolution. This potentially shifts the data along the horizontal axis of a log-log plot, leading to variations in the derived power law exponent α_F . We here use the rather conservative assumption that the [fracture](#) trace lengths are mapped 520 with errors ϵ of 0.05 m (1:10), 0.5 m (1:100), and 5 m (1:1000) at the different mapping scales. We then randomly perturb all measured [fracture](#) traces with length values between $[-\epsilon, \epsilon]$ with a uniform distribution and calculate the α_F value for 100 Monte

hat gelöscht: Fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

Carlo (MC) simulations. As shown in Fig. 9a, the normalized density distributions show relatively little variation. The derived mean power law exponents α_F between 2.14 and 2.80 are similar to the values reported in Fig. 6, and the standard deviations of up to 0.15 suggests that the influence of uncertainties related to [fracture](#) length estimation on α_F is small (Fig. 9b).

535 As the multi-scale approach used in this study relies on [fracture](#) datasets from different scales, it is critical to assess the influence of the choice of different subsampling areas. The derived fractal dimension values below two suggest that the [fracture](#) density is not a perfectly homogeneous property in the investigated [fracture](#) networks (Figs. 3 and 5). Therefore, the spatial choice of the subsampling area for the next detailed scale leads to uncertainties in the size distribution, as differences in [fracture](#) densities shift the data along the vertical axis, affecting the calculation of α_F . Additionally, [fracture](#) traces may be overlooked

540 or linear features not representing [fractures](#) may be interpreted as [fractures](#) during manual mapping of [fracture](#) traces, leading to a similar effect. When selecting subsampling areas, we tried to avoid specific geometric features (i.e., [fracture](#) linkages, [fracture](#) terminations) and focused on minimizing topographic effects and Quaternary cover, since these factors can significantly bias the power law exponent estimates, leading to an overestimation of α_F (Cao and Lei, 2018). To assess the influence of changes in [fracture](#) densities due to the arbitrary choice of the subsampling area, we randomly altered the number

545 of mapped [fractures](#) from -25 to +25% and derive α_F for 100 MC simulations. Compared to the [fracture](#) trace length uncertainties, the effect of the subsampling area uncertainties is more pronounced (Figs. 9c and d). However, the derived α_F values still vary in a small range between 2.44 and 2.86, with standard deviations of up to 0.29.

We conclude from this analysis that both uncertainties related to the [fracture](#) trace length estimation as well as the choice of the subsampling area and manual mapping have limited influence on the derived power law exponents α_F . We thus expect the true power law exponents of the [fracture](#) networks α_F to range between 2.5 and 3, with little regional and lithological variation.

550

5.2 [Fracture](#) Network Characteristics

The [fracture](#) networks from the four different study sites across the Rawil depression illustrate the pervasive but systematic [fracturing](#) in the study area (Fig. 3). Interestingly, the 2D orientation distributions for an individual site appear to be consistent across scales (Fig. 3), which is in agreement with the findings of Odling (1997). This implies that the [fracture](#) orientations are

555 self-similar, and thus scale-independent, suggesting that a single-scale mapping should be sufficient to capture accurate information on the orientation distribution within a [fracture](#) network. Comparing the [fracture](#) networks of the different sites, however, we observe variations in the dominant strike orientations (Fig. 3). While in the ECMs, both at site A in the Aiguilles Rouges massif and site D in the Gastern massif, NW-SE and NE-SW striking [fractures](#) prevail, the limestone nappes overlying the central part of the Rawil depression are characterized by predominantly E-W to NE-SW striking [fractures](#). NE-SW striking

560 [fractures](#) are however also present in both sites A and D, implying that these [fractures](#) are regional features, roughly following the direction of the RSF and the seismogenic RFZ (Fig. 1a).

The α_F values for the four different study sites are all in the same order and range between 2.5 and 3 (Fig. 6). As shown in

590 [Table 2](#), these results are in good agreement with previously published values that generally lie between 2 and 3, which suggests
that this range can be seen as a first-order universal law of [fracture](#) size distributions, independent of the host rock lithology.
Also the derived fractal dimension D values of around 1.7 are consistent with values reported in literature (Bonnet et al., 2001
and references therein).

595 The multi-scale approach used herein has the benefit to greatly expand the length range of the individual [fracture](#) trace datasets
compared to single-scale analysis (Figs. 2b and 6). The power law distributions derived in this study cover [fracture](#) length
scales of three orders of magnitudes between about 10^0 and 10^3 m, and it is likely that the power law holds [also](#) for smaller
and larger [fractures](#). In nature, however, all power laws must have upper and lower physical limits (Bonnet et al., 2001; Torabi
and Berg, 2011; Turcotte, 1997, 1989). The detected lower scaling limit of 10^0 m likely does not reflect the actual physical
length scale (e.g., caused by grain size effects), and we thus speculate that our analysis does not incorporate the lower limit of
600 the power law scaling. Identifying the true limit of the power law would require similar analysis of high-resolution datasets at
smaller scales. The derived upper limit of around 10^3 m is way below the assumed thickness of the seismogenic part of the
crust of around 15 kilometers, as well as the thickness of the Helvetic nappe system of a few kilometers (Burkhard, 1988;
Levato et al., 1994; Pfiffner et al., 1997; Steck et al., 1997). However, stratigraphic layering effects within the Helvetic nappes
(e.g., Pfiffner, 1993) could influence the upper limit of the power law size distributions (Odling, 1997), at least for sites B and
605 C. Since the upper scaling limits do not significantly differ between the different lithologies, we consider this effect to be of
minor importance within the observed length range. We thus assume that our data only covers a limited length range of the
actual [fracture](#) size distribution, and that the power law scaling holds for smaller and larger [fractures](#) than studied herein. This
interpretation is supported by the fact that the previously reported α_f values for lengths between 10^{-1} and 10^5 m all lie in a
similar range (

hat gelöscht: Table 2

hat gelöscht: fault

hat gelöscht: fault

Formatiert: Beschriftung

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

Table 2).

5.3 Potential Links to the Seismicity

The strong vertical components due to the exhumation-related tectonics during the latest stage of the Alpine orogeny, accompanied by enhanced glacial surface erosion in the past 2 Ma (Häuselmann et al., 2007; Glotzbach et al., 2010; Valla et al. 2012; Fox et al., 2015; Sternai et al., 2019), led to the exposure of the [fracture](#) networks studied herein (Egli et al., 2017; Herwegh et al., 2020; Cardello et al., 2024). Based on observations of both exhumed [fracture](#) patterns, earthquake hypocenter locations, and focal mechanisms, previous authors suggested that the exhumed [fracture](#) networks in the region of the Rawil depression were formed under a similar stress field that prevails today (Cardello and Mancktelow, 2015; Maurer et al., 1997; Pavoni, 1980a, b). As earthquakes and [fractures](#) are inherently related, they are expected to exhibit similar orientation and size distributions (Bonnet et al., 2001; Scholz, 2007, 1998, 1997). In the following, we attempt to compare these statistical properties of the exhumed [fracture](#) networks with the same properties of the seismicity in the study region.

Comparing the orientation distributions of the [fracture](#) networks (Figs. 3 and 4) with the earthquake ruptures (Fig. 7) in a first step, we observe a correlation only in the center of the Rawil depression for study sites and subdomains B and C. The E-W to NE-SW striking [fracture](#) set C_I (Figs. 4e and f) correlates with the active [fracture](#) planes from the respective subdomain C (Fig. 7c). The dextral strike-slip kinematics of the NE-SW striking [fractures](#) reveals striking similarities with the kinematics of the earthquake ruptures (Figs. 4f and 7c). In subdomain B, where no active [fracture](#) planes are deciphered (Fig. 7b), the average orientation of the [fracture](#) planes generally renders earthquake rupturing along E-W to NE-SW striking planes possible, which would correspond to reactivation of [fracture](#) sets B_I and B_{II} (Figs. 4 c and d). However, recent high-precision hypocenter datasets suggest that NW-SE striking [fracture](#) planes are reactivated at least in the northern part of the Rawil depression (Diehl et al., 2024). In the ECMs on both sides of the Rawil depression, the correlation between the orientation distributions is less clear. In subdomain A, earthquake ruptures strike ENE-WSW (Fig. 7a), likely corresponding to the [similarly oriented A_I fractures](#) (Fig. 4a). It thus seems that the recent earthquakes mainly reactivate prevailing A_I [fractures](#) as strike-slip faults (Fig. 7a). The E-W striking active rupture planes in subdomain D (Fig. 7d) are not reflected by the bimodal [fracture](#) network of site D at all (Fig. 4g), which is therefore likely not a good proxy for the E-W directed seismicity in subdomain D. This might be due to the fact that the chosen study site lies slightly outside the main seismic corridor. This difference could be explained by mechanically detached Aar and Gastern massifs, which also exhibit significantly lower seismic activity (Fig. 1b). The mapped NW-SE striking [fractures](#) could thus represent the eastern lateral boundary of the main E-W to NE-SW striking seismogenic RFZ (e.g., Lee et al., 2022). Overall, earthquake ruptures in the Rawil depression and the RFZ seem to occur mainly along subvertical E-W to NE-SW striking planes, and the exhumed [fracture](#) networks show similar [directions](#) at least in the central part of the Rawil depression above the main earthquake corridor (Fig. 1b). This implies that the active [fractures](#) in the basement are consistent with exhumed [fracture](#) networks in the Helvetic nappes, indicating a [link across various structural levels and different lithologies](#). Combined with the observation that α_f values reveal little sensitivity to lithological variations, we argue

hat gelöscht: Table 2

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: nodal

hat gelöscht: nodal

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: SE dipping

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: directivities

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: link across

hat gelöscht: lithologies

that the [fracture](#) networks of sites B and C serve as a good first-order analog of the seismically reactivated [fracture](#) networks at depth.

hat gelöscht: fault
hat gelöscht: fault

5.4 Implications for Orogen-Internal Settings

675 As earthquakes usually occur several kilometers below the surface where direct information about the prevailing [fracture](#) networks is sparse, the underlying processes of seismogenic deformation often remain uncovered. However, (i) the good agreement of [fractures](#) and earthquakes in terms of their orientation distributions, (ii) the insensitivity of α_F to lithological variations [across different structural levels](#), and (iii) the observation that both features arguably formed in a similar stress field (Cardello and Mancktelow, 2015; Maurer et al., 1997; Pavoni, 1980a, b) imply that the exhumed [fracture](#) networks in our study
680 area might serve as good first-order analog for the [fracture](#) networks at depth. We therefore attempt to compare the size distributions of the [fracture](#) networks and earthquake ruptures in a next step. As the power law exponent of [fracture](#) networks is dependent on the dimensionality of the data (e.g., Bonnet et al., 2001), we have to consider that the power law exponents were obtained from 2D ([fractures](#)) and 3D (earthquake ruptures) observations, respectively. Stereological considerations have demonstrated that the differing dimensionalities can be corrected for by employing the equation $\alpha_{3D} = 1.28 * \alpha_{2D} - 0.23$ (Bonnet
685 et al., 2001; Borgos et al., 2000; Hatton et al., 1993). Consequently, the 3D power law exponents of the [fracture](#) networks $\alpha_{F(3D)}$ [ranges](#) somewhere between 3 and 3.6.

hat gelöscht: fault
hat gelöscht: fault
hat gelöscht: fault
hat gelöscht: fault
hat gelöscht: fault
hat gelöscht: fault
hat gelöscht: 2
hat gelöscht: fault
hat gelöscht: is ranging

The power law exponent of the earthquake size distributions, commonly referred to as b-value by seismologists, has been proposed to decrease with increasing differential stress and thus depth (Mori and Abercrombie, 1997; Spada et al., 2013; Scholz, 2019, 2015). We therefore assess the depth-dependent variations of the frequency and size distribution of earthquake
690 ruptures. As no significant lateral variations in α_R values were observed within the investigated seismotectonic domain (Fig. 8), we calculate α_R values for depth-slices that incorporate the entire seismotectonic domain. To minimize potential bias due to the data selection, we randomly iterate through different [depth ranges](#), and only fit a power law for [datasets](#) with more than 200 [entries](#). Since the majority of vertical uncertainties in the modern era of the earthquake catalog are in the subkilometer range (Diehl et al., 2021b; Lee et. al. 2023), we do not consider hypocenter uncertainties.

hat gelöscht: combinations of
hat gelöscht: thresholds
hat gelöscht: entities
hat gelöscht: data points

695 As shown in Fig. 10a, earthquakes in the study area occur most frequently at depths between 3 and 8 km. The maximum rupture lengths lie, with few exceptions, on the order of 10^2 m for depths shallower than 3 km, with significantly larger ruptures up to 10^3 m occurring at depths between 3 and 9 km (Fig. 10b). At greater depths, the seismicity becomes sparse, and maximum rupture lengths generally decrease, which could be due to the increasing component of temperature-dependent viscous deformation representing the [frictional](#)-viscous transition at the lower depth end (e.g., Wehrens et al., 2016). As expected α_R
700 values generally decrease with depth (Fig. 10c), in line with previous studies (e.g., Mori and Abercrombie, 1997; Spada et al., 2013). However, the values for shallow (< 3 km) and intermediate (3 – 9 km depth) crustal earthquakes differ significantly: while shallow earthquakes reveal α_R values around 5 to 8, values for intermediate earthquakes constantly decrease with depth to values of around 3.5. Comparing the α_R values with $\alpha_{F(3D)}$ of the exhumed [fracture](#) networks, we notice that for shallow earthquake ruptures a significant discrepancy to the $\alpha_{F(3D)}$ values exists (Fig. 10c). For intermediate earthquakes, α_R and $\alpha_{F(3D)}$

hat gelöscht: brittle
hat gelöscht: fault

values converge towards similar values at around 8 km depth. This observation could potentially be explained by the following two hypotheses.

725 First, it could be stated that similarly to a_R , also $a_{F(3D)}$ values decrease with depth. This cannot be strictly proven, as [fracture](#) networks at depth are not directly accessible. The enhanced earthquake frequency at depths > 3 km, however, suggests that the majority of brittle [fractures](#) are formed at intermediate depth levels (Fig. 10a). Even though [fracturing](#) certainly occurs at shallow depths < 3 km during exhumation as well, the significantly lower earthquake frequency implies that the majority of [fractures](#) in the exhumed [fracture](#) networks were formed at intermediate depth levels. This interpretation is supported by the proposed formation depths of exhumed [fractures](#) in the surrounding of the Rawil depression derived from geological field observations (Cardello and Mancktelow, 2015; Gasser and Mancktelow, 2010; Ustaszewski et al., 2007; Cardello et al., 2024).

730 Together with the absence of significant lithological variations of a_F and the consistency with previously reported values from a large variety of tectonic settings (

- hat gelöscht: fault
- Formatiert: Beschriftung
- hat gelöscht: fault
- hat gelöscht: fault
- hat gelöscht: fault
- hat gelöscht: fault
- hat gelöscht: fault

740 Table 2), we consider this first hypothesis as unlikely.

As a second hypothesis, we thus propose that the $\alpha_{F(3D)}$ values are relatively constant with depth, and $\alpha_{F(3D)}$ and α_R converge towards similar values at intermediate depths. According to this argumentation, the decrease in α_R values therefore records variations in the reactivation potential of pre-existing fractures at different depth levels due to increasing differential stresses. As the differential stresses are smaller at shallow depths, rupture initiations are more likely to stop before growing into large earthquakes, which leads to larger α_R values (Mori and Abercrombie, 1997). In contrary, with increasing differential stress, the likelihood of a rupture initiation (i) to grow into a large earthquake that reactivates the full available fracture length and (ii) to jump across multiple fracture segments is increasing (Manighetti et al., 2007), leading to a convergence of α_R and $\alpha_{F(3D)}$ values at intermediate depths. We propose that the similarities in the power law exponents at depths around 6 to 8 km reveal that earthquakes are more likely to rupture along the entire fracture length at these depths.

750 Apart from the influence of differential stress, previous studies have shown that also other factors such as the fracture type (Gulia and Wiemer, 2010; Petruccelli et al., 2019a, 2019b; Schorlemmer et al., 2005) or material heterogeneities (Mori and Abercrombie, 1997; Mogi, 1962; Goebel et al., 2017) can lead to changes in the earthquake power law exponents. As the dominant orientations of both the fracture networks and earthquake ruptures are in good agreement (Figs. 4 and 7), and the focal mechanisms mostly imply transtensional movements (Kastrup et al., 2004; Houlié et al., 2018), we argue that the fracturing style has a minor influence on the observed variations in our case. Material heterogeneities, such as differences in lithologies, could potentially explain the increased α_R values at shallow depths above 3 km, as this depth correlates with the proposed basement-cover contact below the Rawil depression (Burkhard, 1988; Levato et al., 1994; Pfiffner et al., 1997; Steck et al., 1997; Lee et al. 2023). Stratigraphic layering effects due to the finite thickness of the Helvetic limestone units may limit the size of earthquake ruptures, increasing the α_R values at this lithological boundary (Ouillon et al., 1996). However, since the derived $\alpha_{F(3D)}$ values of the limestone and crystalline basement units are rather similar, and the discrepancy between the size distributions of exhumed fractures and earthquake ruptures occurs at shallow depths, we argue that the elevated α_R values at shallow depths cannot be explained by differences in material heterogeneities alone, in line with the findings of Scholz (1968). As an additional factor, the existence of fluids impacting pore pressure and friction can weaken the strength of the fractures in the uppermost crust, preventing accumulation of larger amounts of stresses.

765 In summary, the depth-dependent comparison of the size distributions of fractures and earthquakes suggests that the likelihood of an earthquake rupture to reactivate the entire fracture length is higher at intermediate crustal depths, while for shallower earthquakes differential stresses are rather small, rendering earthquakes that rupture the full available fracture length less likely, which is similar to the interpretations of Tormann et al. (2014) and Hetényi et al. (2018). As illustrated in Fig. 11, partial seismic rupturing thus seems to be the common seismogenic deformation mode at shallow depth in the Rawil depression region. However, it should be noted that other factors, such as lithological changes and the presence of fluids, can influence fracture strength. Consequently, the observed depth-dependent fracture reactivation potential may differ in other regions. To assess the universality of this observation, further investigations in different settings are required.

hat gelöscht: Table 2

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: ing

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

6. Conclusion

790 The comparison of the statistical properties of fractures and earthquakes point towards differences in the seismogenic deformation processes in the investigated orogen-internal setting, characterized by pervasive fracturing and distributed seismicity. Our findings show that the potential of pre-existing fractures to reactivate seismically seems to vary with depth, likely associated with depth-dependent changes in differential stresses (Fig. 11): while partial seismogenic rupturing of fractures is the dominant deformation mode at shallow depths (< 3 km BSL), earthquakes are more likely to reactivate the entire fracture length at intermediate depths (ca. 3 to 9 km BSL) in the Rawil depression region. Initiating earthquake ruptures thus reactivate the full available fracture length predominantly at intermediate crustal depths.

795 The earthquake activity at shallow and intermediate depths furthermore documents ongoing brittle deformation in the Rawil depression region. As a consequence, recent fracturing processes likely enhance the permeability of pre-existing fractures, which could be of great interest for the exploration of the geothermal potential within such orogen-internal settings. In light of

800 potential induced seismicity, our findings indicate that shallow fracture systems are less likely to rupture over the entire length than deeper ones. Furthermore, quantitative information about orientation and size distributions of fracture networks such as obtained herein are crucial for accurate modelling of recent fluid flow within intricate fracture networks. Similar investigations, incorporating the statistical properties of both fracture networks and earthquake datasets, could therefore potentially be used for innovative geothermal exploration in orogen-internal settings.

805

- hat gelöscht: fault
- hat gelöscht: fault
- hat gelöscht: i
- hat gelöscht: fault
- hat gelöscht: fault
- hat gelöscht: fault
- hat gelöscht: fault
- hat gelöscht: faulting
- hat gelöscht: fault
- hat gelöscht: fault
- hat gelöscht: fault
- hat gelöscht: fault
- hat gelöscht: fault

Data Availability

820 The data that support the findings of this study are openly available in Zenodo at <https://zenodo.org/records/13829055> (Truttmann et al., 2024).

Author Contribution

ST and MH conducted field work. ST performed the formal data analysis with support of all co-authors. ST, TD, MH, and SW wrote, reviewed and edited the manuscript.

825 Competing Interests

The authors declare that they have no conflict of interest.

Acknowledgments

Comments by Francesco Iezzi and an anonymous reviewer helped to substantially improve an earlier version of the manuscript and are thankfully acknowledged. This work was funded by the Swiss Geophysical Commission and the Swiss

830 Federal Office of Topography swisstopo, which is greatly acknowledged.

References

835 Ackermann, R. V. and Schlische, R. W.: Anticlustering of small normal faults around larger faults, *Geol*, 25, 1127, [https://doi.org/10.1130/0091-7613\(1997\)025<1127:AOSNFA>2.3.CO;2](https://doi.org/10.1130/0091-7613(1997)025<1127:AOSNFA>2.3.CO;2), 1997.

Ackermann, R. V., Schlische, R. W., and Withjack, M. O.: The geometric and statistical evolution of normal fault systems: an experimental study of the effects of mechanical layer thickness on scaling laws, *Journal of Structural Geology*, 23, 1803–1819, [https://doi.org/10.1016/S0191-8141\(01\)00028-1](https://doi.org/10.1016/S0191-8141(01)00028-1), 2001.

840 Alstott, J., Bullmore, E., and Plenz, D.: Powerlaw: a Python package for analysis of heavy-tailed distributions, *PLoS ONE*, 9, e85777, <https://doi.org/10.1371/journal.pone.0085777>, 2014.

Baer, M., Deichmann, N., Fäh, D., Kradolfer, U., Mayer-Rosa, D., Ruettener, E., Schler, T., Sellami, S., and Smit, P.: Earthquakes in Switzerland and surrounding regions during 1996, *Eclogae Geologicae Helveticae*, 90, 557–567, 1997.

845 Baer, M., Deichmann, N., Braunmiller, J., Bernardi, F., Cornou, C., Fäh, D., Giardini, D., Huber, S., Kaestli, P., Kind, F., Kradolfer, U., Mai, M., Maraini, S., Oprsal, I., Schler, T., Schorlemmer, D., Sellami, S., Steimen, S., Wiemer, S., Woessner,

- J., and Wyss, A.: Earthquakes in Switzerland and surrounding regions during 2002, *Eclogae Geologicae Helvetiae*, 96, 313–324, 2003.
- Baer, M., Deichmann, N., Braunmiller, J., Husen, S., Fäh, D., Giardini, D., Kästli, P., Kradolfer, U., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2004, *Eclogae geol. Helv.*, 98, 407–418, <https://doi.org/10.1007/s00015-005-1168-3>, 2005.
- Baumberger, R., Herwegh, M., and Kissling, E.: Remote Sensing and Field Data Based Structural 3D Modelling (Haslital, Switzerland) in Combination with Uncertainty Estimation and Verification by Underground Data, in: *3D Digital Geological Models*, edited by: Bistacchi, A., Massironi, M., and Viseur, S., Wiley, 159–197, <https://doi.org/10.1002/9781119313922.ch10>, 2022.
- 855 Bonnet, E., Bour, O., Odling, N. E., Davy, P., Main, I., Cowie, P., and Berkowitz, B.: Scaling of fracture systems in geological media, *Rev. Geophys.*, 39, 347–383, <https://doi.org/10.1029/1999RG000074>, 2001.
- Borgos, H. G., Cowie, P. A., and Dawers, N. H.: Practicalities of extrapolating one-dimensional fault and fracture size-frequency distributions to higher-dimensional samples, *J. Geophys. Res.*, 105, 28377–28391, <https://doi.org/10.1029/2000JB900260>, 2000.
- 860 Bossennec, C., Frey, M., Seib, L., Bär, K., and Sass, I.: Multiscale Characterisation of Fracture Patterns of a Crystalline Reservoir Analogue, *Geosciences*, 11, 371, <https://doi.org/10.3390/geosciences11090371>, 2021.
- Bour, O. and Davy, P.: Clustering and size distributions of fault patterns: Theory and measurements, *Geophys. Res. Lett.*, 26, 2001–2004, <https://doi.org/10.1029/1999GL900419>, 1999.
- Bour, O., Davy, P., Darcel, C., and Odling, N.: A statistical scaling model for fracture network geometry, with validation on a multiscale mapping of a joint network (Hornelen Basin, Norway), *J. Geophys. Res.*, 107, 2113, <https://doi.org/10.1029/2001JB000176>, 2002.
- Boutoux, A., Bellahsen, N., Nanni, U., Pik, R., Verlaquet, A., Rolland, Y., and Lacombe, O.: Thermal and structural evolution of the external Western Alps: Insights from (U–Th–Sm)/He thermochronology and RSCM thermometry in the Aiguilles Rouges/Mont Blanc massifs, *Tectonophysics*, 683, 109–123, <https://doi.org/10.1016/j.tecto.2016.06.010>, 2016.
- 870 Brockmann, E., Ineichen, D., Marti, U., Schaer, S., Schlatter, A., and Villiger, A.: Determination of Tectonic Movements in the Swiss Alps Using GNSS and Levelling, in: *Geodesy for Planet Earth*, vol. 136, edited by: Kenyon, S., Pacino, M. C., and Marti, U., Springer Berlin Heidelberg, Berlin, Heidelberg, 689–695, https://doi.org/10.1007/978-3-642-20338-1_85, 2012.
- Bruyninx, C., Legrand, J., Fabian, A., and Pottiaux, E.: GNSS metadata and data validation in the EUREF Permanent Network, *GPS Solut.*, 23, 106, <https://doi.org/10.1007/s10291-019-0880-9>, 2019.
- 875 Burkhard, P. M.: L’Helvétique de la bordure occidentale du massif de l’Aar (évolution tectonique et métamorphique), *Eclogae Geologicae Helvetiae*, 81, 63–114, 1988.
- Campani, M., Mancktelow, N., Seward, D., Rolland, Y., Müller, W., and Guerra, I.: Geochronological evidence for continuous exhumation through the ductile-brittle transition along a crustal-scale low-angle normal fault: Simplon Fault Zone, central Alps, *Tectonics*, 29, 2009TC002582, <https://doi.org/10.1029/2009TC002582>, 2010.
- 880 Campani, M., Mancktelow, N., and Courrioux, G.: The 3D interplay between folding and faulting in a syn-orogenic extensional system: the Simplon Fault Zone in the Central Alps (Switzerland and Italy), *Swiss J Geosci*, 107, 251–271, <https://doi.org/10.1007/s00015-014-0163-y>, 2014.

- Cao, W. and Lei, Q.: Influence of Landscape Coverage on Measuring Spatial and Length Properties of Rock Fracture Networks: Insights from Numerical Simulation, *Pure Appl. Geophys.*, 175, 2167–2179, <https://doi.org/10.1007/s00024-018-1774-4>, 2018.
- Cardello, G. L. and Mancktelow, N. S.: Veining and post-nappe transtensional faulting in the SW Helvetic Alps (Switzerland), *Swiss J Geosci*, 108, 379–400, <https://doi.org/10.1007/s00015-015-0199-7>, 2015.
- Cardello, G. L., Bernasconi, S. M., Fellin, M. G., Rahn, M., Roskopf, R., Maden, C., and Mancktelow, N. S.: Carbonate deformation through the brittle-ductile transition: The case of the SW Helvetic nappes, Switzerland, *Journal of Structural Geology*, <https://doi.org/10.1016/j.jsg.2024.105083>, 2024.
- Castaing, C., Halawani, M. A., Gervais, F., Chilès, J. P., Genter, A., Bourguine, B., Ouillon, G., Brosse, J. M., Martin, P., Genna, A., and Janjou, D.: Scaling relationships in intraplate fracture systems related to Red Sea rifting, *Tectonophysics*, 261, 291–314, [https://doi.org/10.1016/0040-1951\(95\)00177-8](https://doi.org/10.1016/0040-1951(95)00177-8), 1996.
- Ceccato, A., Tartaglia, G., Antonellini, M., and Viola, G.: Multiscale lineament analysis and permeability heterogeneity of fractured crystalline basement blocks, *Solid Earth*, 13, 1431–1453, <https://doi.org/10.5194/se-13-1431-2022>, 2022.
- Clark, R. M., Cox, S. J. D., and Laslett, G. M.: Generalizations of power-law distributions applicable to sampled fault-trace lengths: model choice, parameter estimation and caveats: Power-law distributions and fault-trace lengths, *Geophysical Journal International*, 136, 357–372, <https://doi.org/10.1046/j.1365-246X.1999.00728.x>, 1999.
- Clauset, A., Shalizi, C. R., and Newman, M. E. J.: Power-Law Distributions in Empirical Data, *SIAM Rev.*, 51, 661–703, <https://doi.org/10.1137/070710111>, 2009.
- Cowie, P. A., Vanneste, C., and Sornette, D.: Statistical physics model for the spatiotemporal evolution of faults, *J. Geophys. Res.*, 98, 21809–21821, <https://doi.org/10.1029/93JB02223>, 1993.
- Cowie, P. A., Sornette, D., and Vanneste, C.: Multifractal scaling properties of a growing fault population, *Geophysical Journal International*, 122, 457–469, <https://doi.org/10.1111/j.1365-246X.1995.tb07007.x>, 1995.
- Davy, P.: On the frequency-length distribution of the San Andreas Fault System, *J. Geophys. Res.*, 98, 12141–12151, <https://doi.org/10.1029/93JB00372>, 1993.
- Davy, P., Bour, O., De Dreuzey, J.-R., and Darcel, C.: Flow in multiscale fractal fracture networks, *SP*, 261, 31–45, <https://doi.org/10.1144/GSL.SP.2006.261.01.03>, 2006.
- Davy, P., Le Goc, R., Darcel, C., Bour, O., de Dreuzey, J. R., and Munier, R.: A likely universal model of fracture scaling and its consequence for crustal hydromechanics, *J. Geophys. Res.*, 115, B10411, <https://doi.org/10.1029/2009JB007043>, 2010.
- Davy, Ph., Sornette, A., and Sornette, D.: Some consequences of a proposed fractal nature of continental faulting, *Nature*, 348, 56–58, <https://doi.org/10.1038/348056a0>, 1990.
- Deichmann, N., Baer, M., Braunmiller, J., Ballarin Dolfin, D., Bay, F., Delouis, B., Fäh, D., Giardini, D., Kastrup, U., Kind, F., Kradolfer, U., Kuenzle, W., Roethlisberger, S., Schler, T., Salichon, J., Sellami, S., Spuehler, E., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 1999, *Eclogae Geologicae Helvetiae*, 93, 395–406, 2000.
- Deichmann, N., Baer, M., Braunmiller, J., Ballarin Dolfin, D., Bay, F., Bernardi, F., Delouis, B., Fäh, D., Gerstenberger, M., Giardini, D., Huber, S., Kradolfer, U., Maraini, S., Oprsal, I., Schibler, R., Schler, T., Sellami, S., Steimen, S., Wiemer, S.,

- Woessner, J., and Wyss, A.: Earthquakes in Switzerland and surrounding regions during 2001, *Eclogae Geologicae Helvetiae*, 95, 294–261, 2002.
- 920 Deichmann, N., Baer, M., Braunmiller, J., Husen, S., Fäh, D., Giardini, D., Kästli, P., Kradolfer, U., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2005, *Eclogae geol. Helv.*, 99, 443–452, <https://doi.org/10.1007/s00015-006-1201-1>, 2006.
- Deichmann, N., Clinton, J., Husen, S., Edwards, B., Haslinger, F., Fäh, D., Giardini, D., Kästli, P., Kradolfer, U., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2011, *Swiss J Geosci*, 105, 463–476, <https://doi.org/10.1007/s00015-012-0116-2>, 2012.
- 925 Delacou, B., Deichmann, N., Sue, C., Thouvenot, F., Champagnac, J.-D., and Burkhard, M.: Active strike-slip faulting in the Chablais area (NW Alps) from earthquake focal mechanisms and relative locations, *Eclogae geol. Helv.*, 98, 189–199, <https://doi.org/10.1007/s00015-005-1159-4>, 2005.
- Diehl, T., Deichmann, N., Clinton, J., Husen, S., Kraft, T., Plenkens, K., Edwards, B., Cauzzi, C., Michel, C., Kästli, P., Wiemer, S., Haslinger, F., Fäh, D., Kradolfer, U., and Woessner, J.: Earthquakes in Switzerland and surrounding regions during 2012, *Swiss J Geosci*, 106, 543–558, <https://doi.org/10.1007/s00015-013-0154-4>, 2013.
- Diehl, T., Clinton, J., Deichmann, N., Cauzzi, C., Kästli, P., Kraft, T., Molinari, I., Böse, M., Michel, C., Hobiger, M., Haslinger, F., Fäh, D., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2015 and 2016, *Swiss J Geosci*, 111, 221–244, <https://doi.org/10.1007/s00015-017-0295-y>, 2018.
- 935 Diehl, T., Clinton, J., Cauzzi, C., Kraft, T., Kästli, P., Deichmann, N., Massin, F., Grigoli, F., Molinari, I., Böse, M., Hobiger, M., Haslinger, F., Fäh, D., and Wiemer, S.: Earthquakes in Switzerland and surrounding regions during 2017 and 2018, *Swiss J Geosci*, 114, 4, <https://doi.org/10.1186/s00015-020-00382-2>, 2021a.
- Diehl, T., Kissling, E., Herwegh, M., and Schmid, S. M.: Improving Absolute Hypocenter Accuracy With 3D *Pg* and *Sg* Body-Wave Inversion Procedures and Application to Earthquakes in the Central Alps Region, *JGR Solid Earth*, 126, <https://doi.org/10.1029/2021JB022155>, 2021b.
- 940 Diehl, T., Heilig, J., Cauzzi, C., Deichmann, N., Truttmann, S., Herwegh, M., and Wiemer, S.: SECOS24: New insights into seismicity, deformation and crustal stresses in the Central Alps Region from a baseline seismotectonic earthquake catalog, in: EGU 2024, EGU 2024, Vienna, 2024.
- Dietrich, D.: Fold-axis parallel extension in an arcuate fold- and thrust belt: the case of the Helvetic nappes, *Tectonophysics*, 170, 183–212, [https://doi.org/10.1016/0040-1951\(89\)90271-0](https://doi.org/10.1016/0040-1951(89)90271-0), 1989.
- 945 Dietrich, D. and Casey, M.: A new tectonic model for the Helvetic nappes, *SP*, 45, 47–63, <https://doi.org/10.1144/GSL.SP.1989.045.01.03>, 1989.
- Egli, D. and Mancktelow, N.: The structural history of the Mont Blanc massif with regard to models for its recent exhumation, *Swiss J Geosci*, 106, 469–489, <https://doi.org/10.1007/s00015-013-0153-5>, 2013.
- 950 Egli, D., Mancktelow, N., and Spikings, R.: Constraints from $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology on the timing of Alpine shear zones in the Mont Blanc-Aiguilles Rouges region of the European Alps: GEOCHRONOLOGY OF ALPINE SHEAR ZONES, *Tectonics*, 36, 730–748, <https://doi.org/10.1002/2016TC004450>, 2017.
- Escher, A., Masson, H., and Steck, A.: Nappe geometry in the Western Swiss Alps, *Journal of Structural Geology*, 15, 501–509, [https://doi.org/10.1016/0191-8141\(93\)90144-Y](https://doi.org/10.1016/0191-8141(93)90144-Y), 1993.

- 955 Fox, M., Herman, F., Kissling, E., and Willett, S. D.: Rapid exhumation in the Western Alps driven by slab detachment and glacial erosion, *Geology*, 43, 379–382, <https://doi.org/10.1130/G36411.1>, 2015.

Fréchet, J., Thouvenot, F., Frogneux, M., Deichmann, N., and Cara, M.: The MW 4.5 Vallorcine (French Alps) earthquake of 8 September 2005 and its complex aftershock sequence, *J Seismol*, 15, 43–58, <https://doi.org/10.1007/s10950-010-9205-8>, 2011.
- 960 Gasser, D. and Mancktelow, N. S.: Brittle faulting in the Rawil depression: field observations from the Rezli fault zones, Helvetic nappes, Western Switzerland, *Swiss J Geosci*, 103, 15–32, <https://doi.org/10.1007/s00015-010-0004-6>, 2010.

Glotzbach, C., Reinecker, J., Danišik, M., Rahn, M., Frisch, W., and Spiegel, C.: Thermal history of the central Gotthard and Aar massifs, European Alps: Evidence for steady state, long-term exhumation, *Journal of Geophysical Research: Earth Surface*, 115, <https://doi.org/10.1029/2009JF001304>, 2010.
- 965 Goebel, T. H. W., Kwiitek, G., Becker, T. W., Brodsky, E. E., and Dresen, G.: What allows seismic events to grow big?: Insights from b-value and fault roughness analysis in laboratory stick-slip experiments, *Geology*, 45, 815–818, <https://doi.org/10.1130/G39147.1>, 2017.

Goertz-Allmann, B. P., Edwards, B., Bethmann, F., Deichmann, N., Clinton, J., Fah, D., and Giardini, D.: A New Empirical Magnitude Scaling Relation for Switzerland, *Bulletin of the Seismological Society of America*, 101, 3088–3095, <https://doi.org/10.1785/0120100291>, 2011.
- 970 Goldstein, M. L., Morris, S. A., and Yen, G. G.: Problems with fitting to the power-law distribution, *The European Physical Journal B*, 41, 255–258, 2004.

Gulia, L. and Wiemer, S.: The influence of tectonic regimes on the earthquake size distribution: A case study for Italy: STRESS REGIMES AND b-VALUE, *Geophys. Res. Lett.*, 37, n/a-n/a, <https://doi.org/10.1029/2010GL043066>, 2010.
- 975 Gutenberg, B. and Richter, C. F.: Frequency of Earthquakes in California, *Bulletin of the Seismological Society of America*, 34, 185–188, 1944.

Hatton, C. G., Main, I. G., and Meredith, P. G.: A comparison of seismic and structural measurements of scaling exponents during tensile subcritical crack growth, *Journal of Structural Geology*, 15, 1485–1495, 1993.
- 980 Häuselmann, P., Granger, D. E., Jeannin, P.-Y., and Lauritzen, S.-E.: Abrupt glacial valley incision at 0.8 Ma dated from cave deposits in Switzerland, *Geol*, 35, 143, <https://doi.org/10.1130/G23094A>, 2007.

Heffer, K. J. and Bevan, T. G.: Scaling Relationships in Natural Fractures: Data, Theory, and Application, *Society of Petroleum engineers*, <https://doi.org/10.2118/20981-MS>, 1990.

Hentschel, H. G. E. and Procaccia, I.: The infinite number of generalized dimensions of fractals and strange attractors, *Physica D: Nonlinear Phenomena*, 8, 435–444, [https://doi.org/10.1016/0167-2789\(83\)90235-X](https://doi.org/10.1016/0167-2789(83)90235-X), 1983.
- 985 Herwegh, M., Berger, A., Glotzbach, C., Wangenheim, C., Mock, S., Wehrens, P., Baumberger, R., Egli, D., and Kissling, E.: Late stages of continent-continent collision: Timing, kinematic evolution, and exhumation of the Northern rim (Aar Massif) of the Alps, *Earth-Science Reviews*, 200, 102959, <https://doi.org/10.1016/j.earscirev.2019.102959>, 2020.

Herwegh, M., Berger, A., Bellashen, N., Rolland, Y., and Kissling, E.: Evolution of the External Crystalline Massifs of the European Alps: From Massif to Lithosphere Scale, in: *Geodynamics of the Alps*, ISTE-Wiley, London, 2023.

- 990 Hetényi, G., Epard, J.-L., Colavitti, L., Hirzel, A. H., Kiss, D., Petri, B., Scarponi, M., Schmalholz, S. M., and Subedi, S.: Spatial relation of surface faults and crustal seismicity: a first comparison in the region of Switzerland, *Acta Geod Geophys*, 53, 439–461, <https://doi.org/10.1007/s40328-018-0229-9>, 2018.
- Houlié, N., Woessner, J., Giardini, D., and Rothacher, M.: Lithosphere strain rate and stress field orientations near the Alpine arc in Switzerland, *Scientific Reports*, 8, <https://doi.org/10.1038/s41598-018-20253-z>, 2018.
- 995 Huggenberger, P. and Aebli, H.: Bruchtektonik und Blattverschiebungen im Gebiet des Rawil-Passes : Resultat einer E-W gerichteten dextralen Scherbewegung im kristallinen Untergrund?, *Schweizerische Mineralogische und Petrographische Mitteilungen*, <https://doi.org/10.5169/SEALS-52783>, 1989.
- Jimenez, M.-J. and Pavoni, N.: Focal mechanisms of recent earthquakes, 1976–1982, and seismotectonics in Switzerland, in: *Proc. Sess. 12, IASPEI XVIII Assembly, Hamburg 1983*, 77–84, 1983.
- 1000 Kastrup, U., Zoback, M. L., Deichmann, N., Evans, K. F., Giardini, D., and Michael, A. J.: Stress field variations in the Swiss Alps and the northern Alpine foreland derived from inversion of fault plane solutions, *J. Geophys. Res.*, 109, <https://doi.org/10.1029/2003JB002550>, 2004.
- Knott, S. D., Beach, A., Brockbank, P. J., Lawson Brown, J., McCallum, J. E., and Welbon, A. I.: Spatial and mechanical controls on normal fault populations, *Journal of Structural Geology*, 18, 359–372, [https://doi.org/10.1016/S0191-8141\(96\)80056-3](https://doi.org/10.1016/S0191-8141(96)80056-3), 1996.
- 1005 Lee, T., Diehl, T., Kissling, E., and Wiemer, S.: New insights into the Rhône–Simplon fault system (Swiss Alps) from a consistent earthquake catalogue covering 35 yr, *Geophysical Journal International*, 232, 1568–1589, <https://doi.org/10.1093/gji/ggac407>, 2023.
- Lei, Q., Latham, J.-P., Tsang, C.-F., Xiang, J., and Lang, P.: A new approach to upscaling fracture network models while preserving geostatistical and geomechanical characteristics: A New Fracture Network Upscaling Method, *J. Geophys. Res. Solid Earth*, 120, 4784–4807, <https://doi.org/10.1002/2014JB011736>, 2015.
- Leonard, M.: Self-Consistent Earthquake Fault-Scaling Relations: Update and Extension to Stable Continental Strike-Slip Faults, *Bulletin of the Seismological Society of America*, 104, 2953–2965, <https://doi.org/10.1785/0120140087>, 2014.
- 1015 Levato, L., Sellami, S., Epard, J.-L., Pruniaux, B., Olivier, R., Wagner, J.-J., and Masson, H.: The cover-basement contact beneath the Rawil axial depression (western Alps): True amplitude seismic processing, petrophysical properties, and modelling, *Tectonophysics*, 232, 391–409, [https://doi.org/10.1016/0040-1951\(94\)90099-X](https://doi.org/10.1016/0040-1951(94)90099-X), 1994.
- Line, C. E. R., Snyder, D. B., and Hobbs, R. W.: The sampling of fault populations in dolerite sills of Central Sweden and implications for resolution of seismic data, *Journal of Structural Geology*, 19, 687–701, [https://doi.org/10.1016/S0191-8141\(96\)00087-9](https://doi.org/10.1016/S0191-8141(96)00087-9), 1997.
- 1020 Mancktelow, N.: The Simplon Line : a major displacement zone in the western Lepontine Alps, *Eclogae Geologicae Helvetiae*, 78, 73–96, <https://doi.org/10.5169/SEALS-165644>, 1985.
- Manighetti, I., Campillo, M., Bouley, S., and Cotton, F.: Earthquake scaling, fault segmentation, and structural maturity, *Earth and Planetary Science Letters*, 253, 429–438, <https://doi.org/10.1016/j.epsl.2006.11.004>, 2007.
- 1025 Marrett, R., Ortega, O. J., and Kelsey, C. M.: Extent of power-law scaling for natural fractures in rock, *Geol*, 27, 799, [https://doi.org/10.1130/0091-7613\(1999\)027<0799:EOPLSF>2.3.CO;2](https://doi.org/10.1130/0091-7613(1999)027<0799:EOPLSF>2.3.CO;2), 1999.

Formatiert: Standard

- Maurer, H.: Seismotectonics and upper crustal structure in the western Swiss Alps, Dissertation, ETH Zürich, 1993.
- Maurer, H. and Deichmann, N.: Microearthquake cluster detection based on waveform similarities, with an application to the western Swiss Alps, *Geophysical Journal International*, 123, 588–600, <https://doi.org/10.1111/j.1365-246X.1995.tb06873.x>, 1995.
- 1030 Maurer, H. R., Burkhard, M., Deichmann, N., and Green, A. G.: Active tectonism in the central Alps: contrasting stress regimes north and south of the Rhone Valley, *Terra Nova*, 9, 91–94, <https://doi.org/10.1111/j.1365-3121.1997.tb00010.x>, 1997.
- Mogi, K.: Study of Elastic Shocks Caused by the Fracture of Heterogeneous Materials and its Relations to Earthquake Phenomena, *Bulletin of the Earthquake Research Institute*, 40, 125–173, 1962.
- Mori, J. and Abercrombie, R. E.: Depth dependence of earthquake frequency-magnitude distributions in California: Implications for rupture initiation, *J. Geophys. Res.*, 102, 15081–15090, <https://doi.org/10.1029/97JB01356>, 1997.
- 1035 Musso Piantelli, F., Mair, D., Berger, A., Schlunegger, F., Wiederkehr, M., Kurmann, E., Baumberger, R., Möri, A., and Herwegh, M.: 4D reconstruction of the Doldenhorn nappe-basement system in the Aar massif: Insights into late-stage continent-continent collision in the Swiss Alps, *Tectonophysics*, 843, 229586, <https://doi.org/10.1016/j.tecto.2022.229586>, 2022.
- 1040 Nicol, A., Walsh, J. J., Watterson, J., and Gillespie, P.: Fault size distributions - are they really power-law?, *Journal of Structural Geology*, 18, 191–197, 1996.
- Odling, N. E.: Scaling and connectivity of joint systems in sandstones from western Norway, *Journal of Structural Geology*, 19, 1257–1271, [https://doi.org/10.1016/S0191-8141\(97\)00041-2](https://doi.org/10.1016/S0191-8141(97)00041-2), 1997.
- Odling, N. E., Gillespie, P., Bourguin, B., Castaing, C., Chiles, J.-P., Christensen, N. P., Fillion, E., Genter, A., Olsen, C., 1045 Thrane, L., Trice, R., Aarseth, E., Walsh, J. J., and Watterson, J.: Variations in fracture system geometry and their implications for fluid flow in fractured hydrocarbon reservoirs, *Petroleum Geoscience*, 5, 373–384, 1999.
- O’Leary, D. W., Friedman, J. D., and Pohn, H. A.: Lineament, linear, lineation: Some proposed new standards for old terms, *Geol Soc America Bull.*, 87, 1463, [https://doi.org/10.1130/0016-7606\(1976\)87<1463:LLSPN>2.0.CO;2](https://doi.org/10.1130/0016-7606(1976)87<1463:LLSPN>2.0.CO;2), 1976.
- Ouillon, G., Castaing, C., and Sornette, D.: Hierarchical geometry of faulting, *J. Geophys. Res.*, 101, 5477–5487, 1050 <https://doi.org/10.1029/95JB02242>, 1996.
- Pavoni, N.: Comparison of focal mechanisms of earthquakes and faulting in the Helvetic zone of the Central Valais, Swiss Alps, *Eclogae Geologicae Helvetiae*, 73, <https://doi.org/10.5169/SEALS-164973>, 1980a.
- Pavoni, N.: Crustal Stresses Inferred from Fault-Plane Solutions of Earthquakes and Neotectonic Deformation in Switzerland, in: *Tectonic Stresses in the Alpine-Mediterranean Region*, vol. 9, edited by: Scheidegger, A. E., Springer Vienna, Vienna, 63– 1055 68, https://doi.org/10.1007/978-3-7091-8588-9_8, 1980b.
- Pavoni, N. and Mayer-Rosa, D.: Seismotektonische Karte der Schweiz 1:750000, *Eclogae Geologicae Helvetiae*, 71, <https://doi.org/10.5169/SEALS-164732>, 1978.
- Pavoni, N., Maurer, H., Roth, P., and Deichmann, N.: Seismicity and seismotectonics of the Swiss Alps, Birkhaeuser, Basel, 1997.

- 1060 Petrucci, A., Schorlemmer, D., Tormann, T., Rinaldi, A. P., Wiemer, S., Gasperini, P., and Vannucci, G.: The influence of faulting style on the size-distribution of global earthquakes, *Earth and Planetary Science Letters*, 527, 115791, <https://doi.org/10.1016/j.epsl.2019.115791>, 2019a.
- Petrucci, A., Gasperini, P., Tormann, T., Schorlemmer, D., Rinaldi, A. P., Vannucci, G., and Wiemer, S.: Simultaneous dependence of the earthquake-size distribution on faulting style and depth, *Geophysical Research Letters*, 46,
1065 <https://doi.org/10.1029/2019GL083997>, 2019b.
- Pfiffner, O. A.: The structure of the Helvetic nappes and its relation to the mechanical stratigraphy, *Journal of Structural Geology*, 15, 511–521, 1993.
- Pfiffner, O. A., Sahli, S., and Stäubli, M.: Structure and evolution of the external basement massifs (Aar, Aiguilles-Rouges/Mt.Bland), Deep Structure of the Alps, results of NRP20, 1997.
- 1070 Pickering, G., Bull, J. M., and Sanderson, D. J.: Sampling power-law distributions, *Tectonophysics*, 248, 1–20, 1995.
- Piña-Valdés, J., Socquet, A., Beauval, C., Doin, M., D’Agostino, N., and Shen, Z.: 3D GNSS Velocity Field Sheds Light on the Deformation Mechanisms in Europe: Effects of the Vertical Crustal Motion on the Distribution of Seismicity, *JGR Solid Earth*, 127, e2021JB023451, <https://doi.org/10.1029/2021JB023451>, 2022.
- Ramsay, J. G.: *Tectonics of the Helvetic Nappes, Thrust and Nappe Tectonics*, 1981.
- 1075 Ramsay, J. G.: Fold and fault geometry in the western Helvetic nappes of Switzerland and France and its implication for the evolution of the arc of the western Alps, *SP*, 45, 33–45, <https://doi.org/10.1144/GSL.SP.1989.045.01.02>, 1989.
- Schlische, R. W., Young, S. S., Ackermann, R. V., and Gupta, A.: Geometry and scaling relations of a population of very small rift-related normal faults, *Geol*, 24, 683, [https://doi.org/10.1130/0091-7613\(1996\)024<0683:GASROA>2.3.CO;2](https://doi.org/10.1130/0091-7613(1996)024<0683:GASROA>2.3.CO;2), 1996.
- Scholz, C. H.: The frequency-magnitude relation of microfracturing in rock and its relation to earthquakes, *Bulletin of the Seismological Society of America*, 58, 399–415, 1968.
- 1080 Scholz, C. H.: Size Distributions for Large and Small Earthquakes, *Bulletin of the Seismological Society of America*, 87, 1074–1077, 1997.
- Scholz, C. H.: A further note on earthquake size distributions, *Bulletin of the Seismological Society of America*, 88, 1325–1326, 1998.
- 1085 Scholz, C. H.: Fault Mechanics, in: *Treatise on Geophysics*, Elsevier B.V., 441–483, 2007.
- Scholz, C. H.: On the stress dependence of the earthquake b value, *Geophys. Res. Lett.*, 42, 1399–1402, <https://doi.org/10.1002/2014GL062863>, 2015.
- Scholz, C. H.: *The Mechanics of Earthquakes and Faulting*, 3rd ed., Cambridge University Press, <https://doi.org/10.1017/9781316681473>, 2019.
- 1090 Scholz, C. H., Dawers, N. H., Yu, J.-Z., Anders, M. H., and Cowie, P. A.: Fault growth and fault scaling laws: Preliminary results, *J. Geophys. Res.*, 98, 21951–21961, <https://doi.org/10.1029/93JB01008>, 1993.
- Schorlemmer, D., Wiemer, S., and Wyss, M.: Variations in earthquake-size distribution across different stress regimes, *Nature*, 437, 539–542, <https://doi.org/10.1038/nature04094>, 2005.

- 1095 Spada, M., Tormann, T., Wiemer, S., and Enescu, B.: Generic dependence of the frequency-size distribution of earthquakes on depth and its relation to the strength profile of the crust: THE *B* -VALUES WITH DEPTH, *Geophys. Res. Lett.*, 40, 709–714, <https://doi.org/10.1029/2012GL054198>, 2013.
- Staudenmaier, N., Cauzzi, C., and Wiemer, S.: *Magnitude Scaling Relationships in Parkfield and Switzerland and their relevance for PSHA*, Swiss Seismological Service (SED) at ETH Zurich, 2018.
- 1100 Steck, A., Epard, J.-L., Escher, A., Lehner, P., Marchant, R., and Masson, H.: Geological interpretation of the seismic of the seismic profiles through Western Switzerland: Rawil (W1), Val d’Anniviers (W2), Mattertal (W3), Zmutt-Zermatt-Findelen (W4) and Val de Bagnes (W5)., *Deep Structure of the Alps*, results of NRP20, 1997.
- 1105 Sternai, P., Sue, C., Husson, L., Serpelloni, E., Becker, T. W., Willett, S. D., Faccenna, C., Di Giulio, A., Spada, G., Jolivet, L., Valla, P., Petit, C., Nocquet, J.-M., Walpersdorf, A., and Castellort, S.: Present-day uplift of the European Alps: Evaluating mechanisms and models of their relative contributions, *Earth-Science Reviews*, 190, 589–604, <https://doi.org/10.1016/j.earscirev.2019.01.005>, 2019.
- Thingbaijam, K. K. S., Martin Mai, P., and Goda, K.: New Empirical Earthquake Source-Scaling Laws, *Bulletin of the Seismological Society of America*, 107, 2225–2246, <https://doi.org/10.1785/0120170017>, 2017.
- Torabi, A. and Berg, S. S.: Scaling of fault attributes: A review, *Marine and Petroleum Geology*, 28, 1444–1460, <https://doi.org/10.1016/j.marpetgeo.2011.04.003>, 2011.
- 1110 Tormann, T., Wiemer, S., and Mignan, A.: Systematic survey of high-resolution *b* value imaging along Californian faults: Inference on asperities, *Journal of Geophysical Research: Solid Earth*, 119, 2029–2054, <https://doi.org/10.1002/2013JB010867>, 2014.
- Truttmann, S., Diehl, T., and Herwegh, M.: Hypocenter-Based 3D Imaging of Active Faults: Method and Applications in the Southwestern Swiss Alps, *JGR Solid Earth*, 128, e2023JB026352, <https://doi.org/10.1029/2023JB026352>, 2023.
- 1115 Truttmann, S., Diehl, T., Herwegh, M., and Wiemer, S.: The Size Distributions of Faults and Earthquakes: Implications for Orogen-Internal Seismogenic Deformation [Dataset], <https://zenodo.org/records/13829055>, 2024.
- Turcotte, D. L.: *Fractals in geology and geophysics*, *PAGEOPH*, 131, 1989.
- Turcotte, D. L.: *Fractals and Chaos in Geology and Geophysics*, 2nd ed., Cambridge University Press, <https://doi.org/10.1017/CBO9781139174695>, 1997.
- 1120 Ustaszewski, M. and Pfiffner, O. A.: Neotectonic faulting, uplift and seismicity in the central and western Swiss Alps, *SP*, 298, 231–249, <https://doi.org/10.1144/SP298.12>, 2008.
- Ustaszewski, M., Herwegh, M., McClymont, A. F., Pfiffner, O. A., Pickering, R., and Preusser, F.: Unravelling the evolution of an Alpine to post-glacially active fault in the Swiss Alps, *Journal of Structural Geology*, 29, 1943–1959, <https://doi.org/10.1016/j.jsg.2007.09.006>, 2007.
- 1125 Valla, P. G., van der Beek, P. A., Shuster, D. L., Braun, J., Herman, F., Tassan-Got, L., and Gautheron, C.: Late Neogene exhumation and relief development of the Aar and Aiguilles Rouges massifs (Swiss Alps) from low-temperature thermochronology modeling and ⁴He/³He thermochronometry, *Journal of Geophysical Research: Earth Surface*, 117, <https://doi.org/10.1029/2011JF002043>, 2012.
- Vicsek, T.: *Fractal growth phenomena*, World scientific, 1992.

- 1130 Wehrens, P., Berger, A., Peters, M., Spillmann, T., and Herwegh, M.: Deformation at the frictional-viscous transition: Evidence for cycles of fluid-assisted embrittlement and ductile deformation in the granitoid crust, *Tectonophysics*, 693, 66–84, <https://doi.org/10.1016/j.tecto.2016.10.022>, 2016.
- Wells, D. L. and Coppersmith, K. J.: New Empirical Relationships among Magnitude, Rupture Length, Rupture Width, Rupture Area, and Surface Displacement, *Bulletin of the Seismological Society of America*, 84, 974–1002, 1994.
- 1135 Wiemer, S., Danciu, L., Edwards, B., Marti, M., Fäh, D., Hiemer, S., Woessner, J., Cauzzi, C., Kästli, P., and Kremer, K.: Seismic Hazard Model 2015 for Switzerland (SUlhaz2015), Swiss Seismological Service (SED) at ETH Zurich, 2016.
- Yielding, G., Walsh, J. J., and Watterson, J.: The prediction of small scale faulting in reservoirs, *First Break*, 10, 449–460, 1992.
- Yielding, G., Needham, T., and Jones, H.: Sampling of fault populations using sub-surface data: a review, *Journal of Structural Geology*, 18, 135–146, [https://doi.org/10.1016/S0191-8141\(96\)80039-3](https://doi.org/10.1016/S0191-8141(96)80039-3), 1996.
- 1140 Zeeb, C., Gomez-Rivas, E., Bons, P. D., and Blum, P.: Evaluation of sampling methods for fracture network characterization using outcrops, *Bulletin*, 97, 1545–1566, <https://doi.org/10.1306/02131312042>, 2013.





Figure 1: Regional setting of the investigated seismotectonic domain z21 of Wiemer et al. (2016) (black polygon), with the four fracture mapping locations (A-D) shown as circles. a) Tectonic map (1:500000; © swisstopo), RSF: Rhône-Simplon Fault Zone. b) Earthquakes from 1984 to 2020 (Diehl et al. 2021b) and available focal mechanism data (see main text for references). Blue arrows show the recent crustal stress field orientations after Kastrup et al. (2004), with contrasting stress regimes north and south of the RSF. The shaded colors indicate the different subdomains A-D. RFZ: Conceptual Rawil Fault Zone. c) Recent crustal displacement rates based on GNSS surface velocities in the vicinity of the study area with reference to stable Europe (Data: EUREF Permanent GNSS Network (Bruyninx et al., 2019); swisstopo). Digital elevation model: EU-DEM v1.1 (© European Union, Copernicus Land Monitoring Service 2023, European Environment Agency (EEA)).



hat gelöscht:

hat gelöscht: fault

hat gelöscht: '

hat gelöscht:

1160

1165

170





Figure 2: a) Workflow of the **fracture** network analysis conducted separately for each location, combining observations from three different mapping scales of 1:1000, 1:100, and 1:10 (for details see Methods). b) Examples of size distributions of the **fracture** networks, plotted with the non-normalized single-scale approach, treating the data from each mapping scale individually (upper panel), and the multi-scale approach, combining all information across scales by normalization with the fractal area (lower panel).

Table 1: Overview of the used orthoimages and digital elevation models (DEM) that serve as a basis for the **fracture** trace maps. Orthoimages and DEM's of the 1:1000 mapping scale (dm resolution) are publicly available from swisstopo.

Location	Tectonic Unit	Mapping Scale	Image Acquisition	UAV Observation Height [m]	Orthoimage Ground Resolution [m]	DEM Ground Resolution [m]	Nr. of Mapped Fracture Traces
A	Aiguilles Rouges	1:10	UAV	10	0.003	0.006	2'440
		1:100	UAV	70	0.029	0.059	5'826
		1:1'000	© swisstopo	-	0.1	2	1'906
B	Diablerets	1:10	UAV	5	0.001	0.002	2'829
		1:100	UAV	50	0.021	0.041	4'654
		1:1'000	© swisstopo	-	0.1	2	3'306
C	Wildhorn	1:10	UAV	5	0.002	0.005	3'519
		1:100	UAV	70	0.024	0.048	10'894
		1:1'000	© swisstopo	-	0.1	2	1'812
D	Gastern	1:10	UAV	10	0.002	0.005	5'213
		1:100	UAV	60	0.024	0.048	3'214
		1:1'000	© swisstopo	-	0.1	2	1'608



hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: '

Location	Tectonic Unit	Mapping Scale	Image Acquisition
A	Aiguilles Rouges	1:10	UAV
		1:100	UAV
		1:1'000	© swiss
B	Diablerets	1:10	UAV
		1:100	UAV
		1:1'000	© swiss
C	Wildhorn	1:10	UAV
		1:100	UAV
		1:1'000	© swiss
D	Gastern	1:10	UAV
		1:100	UAV
		1:1'000	© swiss

hat gelöscht:




Figure 3: **Fracture** trace maps at different mapping scales (rows) for all four study sites (columns). The maps show both **fracture** traces (black lines) and no data areas (grey areas). The length-weighted rose diagrams to the top right of each map show the 2D orientation distribution of the **fracture** traces (**strike directions**). N indicates the number of mapped **fracture** traces.

hat gelöscht: Fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault




Figure 4: Field-based characterization of the fracture populations. The stereoplots in the upper row shows the 3D orientation distributions (black dots: poles to planes on a lower hemisphere projection) with Kamb density contours in red (2σ interval) for all study sites. The lower row shows the respective kinematic indicators from fracture observations (lower hemisphere projection). The dots represent the slip vectors, colored after fracture type and shear sense.

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: fault

hat gelöscht: ing




Figure 5: Estimation of the fractal dimension D for each individual fracture network based on the two-point correlation function C_2 (colored line). The black line shows the local slopes of C_2 (derivative of C_2 ; right-hand axis). The plateau in the local slopes within the grey shaded area approximates the fractal dimension D , which is represented by the black dashed lines.

hat gelöscht: fault





Figure 6: Normalized size distributions of the **fracture** networks for all four study sites, derived from the multi-scale approach. Circles denote the unbiased data, while crosses show the truncated and censored parts of the data. The different color shades represent data from the three mapping scales. The respective **fracture** density term c and the power law exponent α_F are given at the top right.



hat gelöscht: fault

hat gelöscht: fault




Figure 7: 3D orientation distributions of earthquake ruptures based on the nodal planes from published focal mechanism data (see text for references). Solid colored lines: known active [fracture](#) planes with slip vectors plotted as circles. Focal mechanisms without a known active [fracture](#) plane are represented by grey dashed lines (planes) and triangles (slip vectors). Secondary nodal planes for focal mechanisms with a known active plane are not shown in the figure.

hat gelöscht: nodal

hat gelöscht: nodal




Figure 8: 3D size distribution of earthquake ruptures for the four subdomains. The earthquake magnitudes M_w were converted into rupture lengths L_R with the scaling laws of Leonard (2014) as L14, Thingbaijam et al. (2017) as T17 and Wells and Coppersmith (1994) as WC94 represented by the different grey scales, yielding three slightly different [size](#) distributions and α_R values. The values in the bottom left indicate the respective values of α_R .

hat gelöscht:

hat gelöscht: rupture length





Figure 9: a) Assessment of the **fracture** trace length uncertainties. The plots show the normalized density **size** distributions with **fracture** trace lengths perturbed with values 0.05, 0.5, and 5 m, respectively. b) Histogram of the derived α_F values. The values report the mean α_F value and the respective standard deviation. c) Assessment of the subsampling area uncertainties, showing the normalized density **size** distributions with the number of **fractures** in each dataset randomly altered between -25 and +25 %. d) Histogram of the derived α_F values of the sampling area uncertainties. The values report the mean α_F value and the respective standard deviation.



hat gelöscht:

hat formatiert: Hervorheben

hat gelöscht: fault

hat gelöscht: length

Formatiert: Beschriftung

hat gelöscht: fault

hat gelöscht: length

hat gelöscht: fault

hat gelöscht: ¶

Table 2: Compilation of published power law exponents α_f from fracture size distributions sorted by the used analysis approach (single- vs. multi-scale power law fitting), with complementary information deduced where available. As both cumulative (CDF) and probability density functions (PDF) are used for fitting fracture length data, we have to make the CDF-based values comparable to our PDF-based α_f values by using $\alpha_{CDF} + 1$ (Bonnet et al., 2001). N: Total number of data; α_f (published): published power law exponent and the used distribution; α_f (PDF): power law exponents converted to PDF values (Bonnet et al., 2001); lower and upper limit: approximate bounds of the power law scaling range.

Reference	N	Rock type	α_f (published)	Distribution	α_f (PDF)	Lower limit [m]	Upper limit [m]
Single-scale analysis							
Ackermann and Schlische (1997)	873	Sediments	1.64	CDF	2.64	0.04	0.15
Bour and Davy (1999)	3499	-	1.88	PDF	1.88	4000	70000
Clark et al. (1999)	1034	-	1.51	CDF	2.51	360	4500
Knott et al. (1996)	218	Sediments	1.02	CDF	2.02	0.31	0.93
Ouillon et al. (1996)	...	Sediments	1.9	PDF	1.9	2	20
Ouillon et al. (1996)	...	Sediments	2.1	PDF	2.1	800	8000
Ouillon et al. (1996)	...	Sediments	3.2	PDF	3.2	3000	20000
Ouillon et al. (1996)	...	Sediments	2.1	PDF	2.1	8000	30000
Schlische et al. (1996)	201	Sediments	1.4	CDF	2.4	3	20
Yielding et al. (1996)	~450	-	1.18	CDF	2.18	300	10000
Yielding et al. (1996)	~350	-	1.75	CDF	2.75	4000	50000
Yielding et al. (1996)	300	-	1.37	CDF	2.37	1500	20000
Multi-scale analysis							
This study: Site A	10172	Crystalline basement	2.43	PDF	2.43	1	300
This study: Site B	10789	Sediments	2.49	PDF	2.49	0.1	1000
This study: Site C	16225	Sediments	2.73	PDF	2.73	0.5	300
This study: Site D	10035	Crystalline basement	2.47	PDF	2.47	0.5	400
Bossennec et al. (2021)	...	Crystalline basement	2.03	CDF	3.03	1	10000
Bour et al. (2002)	21'778	Sediments	2.8	PDF	2.8	1	200
Castaing et al. (1996)	...	Crystalline basement & sediments	2.34	CDF	3.34	20	10000
Ceccato et al. (2022)	5908	Crystalline basement	1.88	CDF	2.88	1	10000
Line et al. (1997)	...	Crystalline basement	1.66	CDF	2.66	30	100000
Odling (1997)	...	Sediments	2.1	CDF	3.1	1	1000
Odling et al. (1999)	...	Sediments	2.34	CDF	3.34	5	100000
Scholz et al. (1993)	...	Volcanic	1.3	CDF	2.3	30	3000
Yielding et al. (1992)	...	-	2	CDF	3	3000	30000

hat gelöscht: fault

hat gelöscht: fault

1245




Figure 10: Depth-dependency of the seismicity in terms of a) the number of earthquakes, b) the rupture length l_R of the individual earthquakes, and c) the derived power law exponents $\alpha_{(3D)}$ for both the earthquake ruptures (grayscales) and the [fracture networks](#) (colored bars and red box). Error bars denote 1σ uncertainties of the power law exponents (vertical bars) and the used depth interval (horizontal bars).

hat gelöscht: fault





Figure 11: Schematic block diagram illustrating the depth-dependent seismogenic fracture reactivation patterns observed herein by relating the power law exponents of the fracture networks (α_F) to the exponents of recent earthquakes (α_R). Full seismogenic fracture reactivation appears to occur predominantly at depths between 3 and 9 km below sea level (BSL).



hat gelöscht:

hat gelöscht: to illustrate

hat gelöscht: fault

hat gelöscht: in this study

hat gelöscht:

hat gelöscht: fault

hat formatiert: Tiefgestellt

hat formatiert: Tiefgestellt

hat gelöscht: seems

hat gelöscht: only

hat gelöscht: from

hat gelöscht: to