

Snowball Earth transitions from Last Glacial Maximum conditions provide an independent upper limit on Earth's climate sensitivity

Martin Renoult^{1,2}, Navjit Sagoo^{2,3}, Johannes Hörner⁴, and Thorsten Mauritsen^{2,3}

¹Department of Geological Sciences, Stockholm University, Stockholm, Sweden

²Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden

³Department of Meteorology, Stockholm University, Stockholm, Sweden

⁴Department of Meteorology and Geophysics, University of Vienna, Vienna, Austria

Correspondence: Martin Renoult (martin.renoult@geo.su.se)

Abstract. Geological evidence of a snowball Earth state indicate persistent tropical sea ice cover during the Neoproterozoic (>635 million years ago). Current theory is that a strengthening of the positive surface albedo feedback with cooling temperatures, eventually exceeding the sum of all other feedbacks, leads to a global climate instability. Several recent high sensitivity climate models with strongly positive cloud feedbacks have not been able to simulate the much warmer Last Glacial Maximum (LGM) state, suggestive that they cool excessively in response to a modest decrease in atmospheric carbon dioxide levels and therefore enter the snowball instability by this mechanism. Using a coupled Earth system model, MPI-ESM1.2, we show that clouds accelerate the transition to a snowball Earth state and reduce the radiative forcing required to trigger the climate snowball instability. Positive cloud feedbacks over tropical oceans and ahead of the sea-ice edge act to cool down the oceans and promote sea ice formation. Regardless, when approached slowly, the snowball Earth transitions appear transition appears to occur around a global mean temperature of zero degree Celsius, simultaneously with the sea ice edge advancing into the subtropics thereby strengthening the surface albedo feedback. This temperature threshold, if supported by several climate models, could be used as a novel and independent constraint on the upper bound of climate sensitivity. ~~Currently, using the results from MPI-ESM1.2~~ by using the relationship of simulated LGM temperatures and the models' equilibrium climate sensitivity. The constraint depends only on the simple fact that Earth did not enter a snowball instability during the recent ice ages. Using the here estimated transition temperature, we find it is implausible that Earth's climate sensitivity exceeds 5.5 K (4.4–6.2°C (3.9–6.6, 90–8.4, 5–95 percent confidence). This upper bound estimate of climate sensitivity is only weakly sensitive to uncertainty in the transition temperature, approximately 0.3 degrees per degree.

1 Introduction

During the ~~Neoproterozoic (>635 million years ago)~~ history of Earth, geological evidences support the formation of persistent ~~sea-ice~~ sea ice within the tropical regions, referred to as snowball Earth states (e.g. Hoffman et al., 2017). Because ice is highly reflective, the positive surface albedo feedback strengthens and exceeds the sum of other feedbacks while the Earth cools down. The evolution of the strength of climate feedbacks as a function of temperature is referred ~~as state-dependency~~ to as temperature-dependency. Whilst climate models agree that the surface albedo feedback increases in cold climates (e.g. Budyko, 1969), other climate feedbacks such as clouds are less understood in the context of cooling (~~e.g. Braun et al., 2022~~).

~~State-dependency~~ (e.g. Braun et al., 2022). Temperature-dependency in climate feedbacks has often been studied in warming climates (~~e.g. Caballero and Huber, 2013; Jonko et al., 2013; Meraner et al., 2013~~), ~~but rarely in cold climates (e.g. Colman and McAvaney, 2009)~~. (e.g. Caballero and Huber, 2013; Jonko et al., 2013; Meraner et al., 2013), ~~but rarely in cold climates (e.g. Colman and McAvaney, 2009)~~.

Pierrehumbert et al. (2011) proposed the Snowball Earth modelling intercomparison project, or SnowMIP, as a protocol to compare modelling efforts within the field. SnowMIP was made of three atmosphere-ocean models, where two of them are older versions of the models applied in the current study (ECHAM5/MPIOM and CAM). SnowMIP paved the way to identifying the importance of circulation and clouds in snowball Earth initiation across models. Following SnowMIP, the importance of snow-albedo feedbacks and their relation to sea-ice albedo feedback was identified, as well as the interaction of ocean and atmosphere circulations (Voigt and Abbot, 2012). Due to the complexity of running a model into snowball Earth state, and the increasing cost of running higher-end models, there is to this day only a small number of published snowball Earth simulations and even less feedback analyses. To our knowledge, there is also no study that relates the snowball Earth transition and climate sensitivity, which is the main aim of the present study.

The community definition of equilibrium climate sensitivity (ECS) is the long term global mean surface warming in response to a doubling of CO₂ over pre-industrial (PI) levels, excluding temperature changes induced by slow changes in ice sheets (Forster et al., 2021). ECS is among the most important geophysical properties for determining future global warming (Grose et al., 2018; Huusko et al., 2021), and has long been the subject of investigation (Arrhenius, 1896). Currently, multiple lines of evidence are used in combination to constrain ECS, including a process based approach wherein forcing and individual feedbacks are constrained, use of the instrumental temperature record warming, and by interpreting estimates of temperature and forcing in various paleoclimates (Sherwood et al., 2020). The result is a substantially reduced uncertainty relative to earlier assessments (Forster et al., 2021).

In this study, we quantify cloud feedbacks as the Earth ~~transits~~ transitions towards a snowball Earth state from ~~pre-industrial~~ PI conditions under low ~~CO₂~~ CO₂ concentration. In particular, we focus on the contribution of cloud feedbacks over tropical oceans, as well as the interaction between cloud and sea-ice albedo feedbacks. Our motivation to investigate climate feedbacks in the snowball Earth transition stems from the fact that a number of relatively high climate sensitivity models of the paleoclimate modelling intercomparison project phase 4 (PMIP4) fail to simulate the Last Glacial Maximum (LGM), a cold period with large ice sheets 21 000 years ago (~~e.g. Kageyama et al., 2021~~), (e.g. Kageyama et al., 2021), despite being

able to simulate warm paleoclimates (e.g. Pliocene, 3.2 million years ago (Haywood et al., 2020), (Haywood et al., 2020), Eocene, 50 million years ago (Lunt et al., 2021)) (Lunt et al., 2021)). These climate models start transitioning to a snowball state at temperatures simulate temperatures for the LGM substantially cooler than indicated by LGM reconstructions (e.g. Zhu et al., 2021a) reconstructions, and show hints of runaway behaviour (e.g. Zhu and Poulsen, 2021b). Since they have strong positive cloud feedbacks they consequently exhibit high climate sensitivities, which is the long term temperature change after an abrupt doubling of atmospheric CO₂ from pre-industrial concentrations sensitivity. It could therefore be possible to use the models simulating that do simulate stable LGM states by evaluating the relationship between simulated LGM temperatures and the ECS of those models as to provide an independent upper limit constraint on climate sensitivity by estimating independent from previous estimates by calculating the temperature at which the Earth transits towards an unstable snowball Earth state. The constraint is essentially based on the fact that during the recent glacial cycles we have evidently not entered the snowball state. The upper bound on ECS is either a large value or not provided for the different lines of evidence used in the latest report of the International Panel on Climate Change (IPCC), where there was no estimate of an upper bound in the combined assessment. Hence, our estimate may become a valuable addition to the next assessment.

To approach this problem with other climate models, we conclude close our study with a short and easy-to-replicate experimental design of protocol of a modern snowball Earth simulations. The results of this experiment are relevant for both climate sensitivity, as shown in this paper, as well as understanding the challenges around setting up the LGM simulation, a notoriously difficult paleoclimate to model climate to simulate.

2 Methods

We use the Max Planck Institute for Meteorology Earth system model version 1.2, MPI-ESM1.2 (Mauritsen et al., 2019) (Mauritsen et al., 2019) to simulate snowball Earth transitions from pre-industrial (PI) PI and LGM initial conditions. The sea-ice and snow-albedo parametrisation in MPI-ESM1.2 is similar to other snowball Earth studies using an older version of this model (Voigt et al., 2011; Voigt and Abbot, 2012), except for the addition of meltponds which are described by Roeckner et al. (2012). We perform abrupt and sustained changes of atmospheric CO₂ concentrations or solar constant starting from the equilibrium states of PI or LGM. Continents, non-CO₂ states. The CO₂ concentrations are chosen as one half of the previous concentration, as CO₂ has a quasi-logarithmic forcing behaviour (Arrhenius, 1896), and as to facilitate comparisons with other studies which uses doubling and halving, such as the ones of the non-linear modelling intercomparison project (nonlinMIP, Good et al. (2016)). Continents, non-CO₂ greenhouse gases and orbital configurations are kept as in PI or LGM unchanged. The initial surface temperature from the PI state is 14.4°C, whereas it is 10.4°C from the LGM state. A dynamic vegetation model is used, but vegetation dies-off rapidly due to rapid cooling and low CO₂. PI simulations use the coarse resolution MPI-ESM1.2-CR (T31 spectral truncation, 31 atmospheric levels), which is faster and numerically more stable to extreme forcing. LGM boundary conditions are only available in the low resolution MPI-ESM1.2-LR (T63, 47 atmospheric levels). Radiative differences between the two models are small, and the climate sensitivity of MPI-ESM1.2-CR is slightly higher than the LR version. While we do not expect state-dependency temperature-dependency to vary much across

Table 1. Summary of the runs performed for this study. PI = Pre-industrial, LGM = Last Glacial Maximum. Solar constant is expressed in percentage of ~~pre-industrial-PI~~ solar constant (1361 Wm^{-2}). Transition temperatures are reported as anomalies relative to PI. *The runs started from an equilibrium state and ran for 100 years. **The runs were manually stopped and are expected to reach equilibrium in a cold non-snowball state. If the transition temperature is left empty, the simulation did not transit towards a snowball Earth during the years simulated.

Run	Length (years)	Transition temp. (°C)	Solar constant (%)	CO ₂ (ppm)	Boundaries	Comments
Pre-industrial	*		100	284.32	PI	MPI-ESM1.2-CR
LGM	*		100	190	LGM	MPI-ESM1.2-LR
2xCO2	150		100	568.64	PI	MPI-ESM1.2-CR
abrupt50ppm <u>50ppm</u>	1156**		100	50	PI	MPI-ESM1.2-CR
1/8xCO2	1142	<u>-14.7</u>	100	35.54	PI	MPI-ESM1.2-CR
1/16xCO2	639	<u>-19.4</u>	100	17.80	PI	MPI-ESM1.2-CR
1/32xCO2	442	<u>-20.8</u>	100	8.90	PI	MPI-ESM1.2-CR
1/64xCO2	338	<u>-19.1</u>	100	4.45	PI	MPI-ESM1.2-CR
1/128xCO2	280	<u>-23.7</u>	100	2.22	PI	MPI-ESM1.2-CR
1/256xCO2	245	<u>-19.5</u>	100	1.11	PI	MPI-ESM1.2-CR
1/512xCO2	219	<u>-22.9</u>	100	0.56	PI	MPI-ESM1.2-CR
1/1024xCO2	230	<u>-26.5</u>	100	0.28	PI	MPI-ESM1.2-CR
1/2048xCO2	194	<u>-22.1</u>	100	0.14	PI	MPI-ESM1.2-CR
1/128xCO2-fixalb	1046**		100	2.22	PI	locked albedo feedback
1/128xCO2-fixq	632**		100	2.22	PI	locked water vapor feedback
1/128xCO2-fixcld	946	<u>-15.3</u>	100	2.22	PI	locked cloud feedback
1/512xCO2-fixcld	518	<u>-16.0</u>	100	0.56	PI	locked cloud feedback
1/256xCO2-fixcld	648	<u>-16.4</u>	100	1.11	PI	locked cloud feedback
1/64xCO2-fixcld	1453**		100	4.45	PI	locked cloud feedback
1/32xCO2-fixcld	1312**		100	8.90	PI	locked cloud feedback
lgm-S94	290	<u>-17.2</u>	94	190	LGM	MPI-ESM1.2-LR
lgm-S93	198	<u>-17.2</u>	93	190	LGM	MPI-ESM1.2-LR
lgm-S92	179	<u>-16.0</u>	92	190	LGM	MPI-ESM1.2-LR
lgm-S91	148	<u>-22.9</u>	91	190	LGM	MPI-ESM1.2-LR
lgm-S89	113	<u>-22.3</u>	89	190	LGM	MPI-ESM1.2-LR
lgm-S85	75	<u>-25.0</u>	85	190	LGM	MPI-ESM1.2-LR
S92-LR	179	<u>-13.4</u>	92	284.32	PI	MPI-ESM1.2-LR
S85-LR	166	<u>-21.0</u>	85	284.32	PI	MPI-ESM1.2-LR
CESM1.2	210	<u>-23.3</u>	100	2.22	PI	CESM1.2

85 ~~models~~between the two model resolutions, individual feedbacks are likely to slightly differ so we also initiate a few simulations from PI with MPI-ESM1.2-LR. Details of all simulations are summarised in Table 1- 1.

The growth of thick ~~sea-ice~~sea ice leads to numerical instabilities in the ~~model~~(\rightarrow ocean model component when exceeding 12 meters). We do not artificially limit sea-ice growth ~~as was done~~in other studies(Voigt and Marotzke, 2010; Voigt et al., 2011; Voigt and Abbot, 2012), ~~as it~~(Voigt and Marotzke, 2010; Voigt et al., 2011; Voigt and Abbot, 2012), because this ~~method~~generates latent heat at the base of sea-ice~~(Marotzke and Botzet, 2007)~~sea ice (Marotzke and Botzet, 2007) which changes the required ~~CO₂~~CO₂ forcing for snowball Earth initiation(Hörner et al., 2022 (Hörner et al., 2022). This results in all our simulations reaching numerical instability at some point when approaching a complete snowball Earth state due to the sea ice being too thick in narrow basins, such as the Baltic Sea. This does not affect our results, however, as we evaluate the




Figure 1. Schematic of top-of-atmosphere (TOA) imbalance versus surface temperature difference to pre-industrial (PI), as developed by Gregory et al. (2004). Here, we compare the behavior of our abrupt forcing simulations, as they cool down, to the behavior of a real world transition wherein CO₂ is slowly and gradually reduced (green line). Note that the dashed line is added for illustration purpose only.

transition in temperatures much higher than at the numerical instability. Only the 50ppm simulation was manually stopped as it is expected to reach an equilibrium in a cold non-snowball state (see Table 1 for details on run length).

2.1 Temperature and imbalance evolution of the simulations

We analyse the evolution of our simulations by studying how the top-of-atmosphere (TOA) energetic imbalance changes with surface temperature. This approach is shown in Fig. 4 and Fig. 6 and has been first developed by Gregory et al. (2004). Here, each point of data is a one-year global average of the simulation as it responds to an initial and abruptly applied forcing. Because the total feedback is negative near the stable and stationary pre-industrial state, the system will attempt to reduce the TOA imbalance when exposed to positive or negative forcing. Consequently, in the case of an initial negative forcing from reduced CO₂ levels, the climate cools down. Therefore, the beginning of the simulation is where the temperature difference to PI is the closest to zero, and the imbalance is close to the forcing imposed. The end of the simulation is the left-most point, when the temperature anomaly relative to PI conditions is the largest.

In the real climate system, the climate would slowly cool down as the forcing from the Sun or greenhouse gases change only slowly over time. Such transitions and abruptly forced simulations are illustrated in Fig. 1. When a negative forcing is

slowly applied, for instance if CO₂ is slowly decreasing over time, the climate system would maintain a quasi-equilibrium and move along the green line to colder temperatures, until the true instability leading to a snowball Earth state is reached. In our simulations, the point closest to equilibrium, i.e. the least negative TOA imbalance, serves as to estimate the transition value.

110 If a simulation was to be brought to higher CO₂ concentration, it is at this point that the equilibrium would be reached first. Because the simulations miss this equilibrium point, and since any other year is further out of balance than the year at this equilibrium point, they will not reach any other equilibrium until a new climate state, most likely the snowball Earth state, is found. Therefore, we can estimate the model's transition temperature towards snowball Earth as the temperature of the year with the least negative TOA imbalance. By definition it is also the year where the total climate feedback would be strictly zero

115 before turning positive, which is represented by the tangential line in the Gregory et al. (2004) framework.

As we shall see, the simulated point slightly differs from the true transition point due to time-dependent effects that are proportional to the degree of disequilibrium. It appears that when the forcing is strongly negative, the transition point in the Gregory analysis shifts to lower temperatures. We speculate that this is a kind of time-dependent feedback effect whereby the positive feedback from formation of sea ice is delayed by the large thermal inertia of the oceans. Therefore, to estimate a

120 model's true snowball Earth transition point it is necessary to carry out multiple abrupt forcing experiments as is done in this study.

2.2 Climate feedback calculations

Climate feedbacks are diagnosed using the partial radiative perturbation (PRP) method (Wetherald and Manabe, 1988; Colman and McAvaney, 1997) (Wetherald and Manabe, 1988; Colman and McAvaney, 1997). Details of its online implementation in

125 MPI-ESM1.2 are in Meraner et al. (2013). Meraner et al. (2013). The PRP method calculates individual contributions of surface albedo, clouds, temperature and water vapor changes to top-of-atmosphere fluxes, by exchanging related variables between a control climate state and the transient state analysed. Because the length of each run varies with the forcing amplitude, we compute climate feedbacks by regressing the top-of-atmosphere radiation balance changes arising from albedo, clouds, temperature, water vapor changes over bins of data points spanning a global temperature ranges of 5°C. We apply the same

130 method in the maps of Fig. 3, whereby the temperature indicated is the middle point of the temperature range.

We furthermore perform runs where we separately lock surface albedo, clouds or water vapor in the radiation calculations to the control state, i.e. the corresponding feedbacks do not contribute to the radiation balance. The implementation in MPI-ESM1.2 is described in Mauritsen et al. (2013). Mauritsen et al. (2013). These locked-feedback transient simulations read the pre-industrial-PI control albedo, clouds, temperature and humidity and impose them on the radiation parameterization

135 regardless of the changes the system is experiencing, such as the increasing extent of sea-ice sea ice.

2.3 Constraint on Earth's climate sensitivity

The relationship between LGM simulated temperatures and the climate sensitivity of models have been used in an emergent constraint framework to infer the Earth's true climate sensitivity owing to geological reconstructions of LGM tempera-

tures(Hargreaves et al., 2012; Schmidt et al., 2014; Renoult et al., 2020, 2023). (Hargreaves et al., 2012; Schmidt et al., 2014; Renoult et al., 2020, 2023).
140 The novel constraint on Earth's climate sensitivity developed here uses the same relationship, derived from models participating in the Paleoclimate Modelling Intercomparison Project Phase 4 (PMIP4), where the regression parameters are calculated via ordinary least squares method, but we replace the geological reconstruction of the LGM by the temperature at which the Earth transits to a snowball Earth. This constraint relies on the simple fact that the real LGM was a stable glacial climate, and therefore the highest value of Earth's climate sensitivity before the LGM enters could not have been higher than the point at which the planet would transition into a snowball state represents the upper limit on climate sensitivity before the LGM. As such, the presented constraint provides a physical upper bound on ECS, not a best estimate.

Recently, Renoult et al. (2023) Renoult et al. (2023) showed that structural uncertainties and state-dependencies-temperature-dependencies in climate feedbacks hinder the relationship between LGM temperature and climate sensitivity, which is computed from warm simulations (4 times pre-industrial CO₂ (Gregory et al., 2004)). We minimize CO₂ (Gregory et al., 2004)). The authors minimized the issue by adding to the PMIP4 ensemble an ensemble of CESM models ("CESM-family", i.e. CESM1.2, CESM1.3 and low-ECS versions of CESM2.1) in a similar way as described by Renoult et al. (2023). Besides increasing the size of the PMIP4 ensemble, single-model ensembles (or in this case, single-family models) have reduced structural uncertainties, in particular regarding ice sheet forcing, albeit they could add statistical dependency issues within the relationship. Here we assume the three CESM models are sufficiently different to avoid this issue), as the relationship also benefits from PMIP4 models to have a wider range of climate sensitivity, and because without these models, the relationship can be considered as too dependent on CESM2 (Renoult et al., 2023). We have added a similar test as conducted by Renoult et al. (2023) on tropical temperatures in Appendix B, where we show that on a global scale, CESM2 and the CESM-family ensemble are almost exchangeable, leading to a robust relationship with similar slopes and intercepts in both cases. Therefore, as we only have access to the sea-surface temperature data of the CESM-family ensemble, whereas our analyses are mostly based on surface temperatures including land, we use CESM2 for most of our study. This is on the basis that as the CESM-family ensemble is not an outlier, and it is exchangeable with CESM2, therefore the consequences of CESM2 being in the PMIP4 ensemble are also not due to it being an outlier. Since models from the same institutes can sometimes introduce larger differences than between models from different modelling centers (Knutti et al., 2013), we would like to emphasize that it is not an ideal solution with the low number of paleosimulations available, and sometimes limited data, but should be seen as motivation for institutes with higher climate sensitivity models to perform and publish LGM runs regardless of whether they are stable or transition towards a snowball state.

3 Analysis of surface albedo and cloud feedbacks

We perform experiments with abrupt decreases of CO₂ concentrations ranging from 1/8 to 1/2048 of pre-industrial CO₂ concentration using the MPI-ESM1.2-CR model to simulate snowball Earth inception and breakdown transition and breakdown climate feedbacks (Section 2). For clarity, we report temperature anomalies to pre-industrial in units of K, and absolute




Figure 2. Evolution of the climate feedback parameters of clouds (λ_{cl}), albedo (λ_a) and temperature and water vapor combined (λ_{t+ww}) with global surface temperature anomaly compared to pre-industrial (KPI) ($^{\circ}\text{C}$) under all CO_2 forcing.

temperatures in $^{\circ}\text{C}$. 2). The strengthening surface albedo feedback has often been considered the main driver in the snowball Earth instability, yet positive cloud feedbacks exceed the surface albedo feedback during the initial 10K of cooling $^{\circ}\text{C}$ of cooling, relative to PI (Fig. 4 2). This strengthening arises from positive tropical cloud feedbacks (Fig. 2 3) due to an increase in shallow cloud coverage over the tropical oceans (Appendix Fig. B1). Below -10°C . Cloud feedbacks are increasingly more positive as the initial forcing is increasingly stronger, which amplify tropical cooling. Below -20K relative to pre-industrial $^{\circ}\text{C}$ relative to PI, cloud feedbacks switch sign and are globally negative and weaker than the surface albedo feedback, although there still exists strong positive cloud feedbacks ahead of the advancing sea-ice edge in both hemispheres (Fig. 2 3). Indeed, because the sea-ice surface is cold, clouds are preferentially over open water which is warmer and a source of moisture (e.g. Wall et al., 2017), (e.g. Wall et al., 2017), where they facilitate further cooling of the ocean. All in all, cloud feedbacks substantially contribute to early cooling in response to reduced CO_2 . During the transition, the positive surface albedo feedback starts to exceeds the combined cloud, temperature and water vapor feedbacks at temperatures between -15K and -20K below pre-industrial $^{\circ}\text{C}$ below PI (Fig. 3 4). This leads to a global instability and is concomitant with the acceleration of southern the equatorward sea-ice edge into the sub-tropics, resulting in strong global cooling towards the snowball state.




Figure 3. Maps of the surface albedo (left column) and cloud (right column) feedbacks at four average global temperatures in the 1/16xCO₂ simulation. The global mean temperature anomaly indicated is the middle point of the temperature range over which the feedbacks are computed, taking bins of 5 degrees of cooling.

185 The diagnostics presented above is-are suggestive that cloud feedback may be an important climate driver towards the snowball-snowball transition. To demonstrate that this is indeed the case, we isolate the importance of each feedback using the feedback locking technique (Section 2.2) on various simulations, where each feedback mechanism is disabled one after the other (Fig. 3-B 4-B). When the cloud feedback is locked, the instability is reached at a similar global mean temperature as in the non-locked simulations, between -15K and -20K °C relative to PI (Fig. 3-A 4-A). However, cloud feedback

190 locking drastically increases the forcing required to initiate the snowball Earth inception transition, and considerably slow down the transition to snowball Earth. We find that simulations at CO₂-CO₂ concentration of 1/64 times pre-industrial PI levels and lower are required to trigger the transition with locked cloud feedbacks, which is a factor ten less than the concentration




Figure 4. A) Top-of-atmosphere (TOA) radiative imbalance (Wm^{-2}) versus global surface temperature anomaly with relative to pre-industrial (PI) conditions (K°C) under varying CO_2 forcing and B) same but with cloud feedbacks locked to that of the control state. A run with an abrupt doubling of CO_2 ($2\times\text{CO}_2$) is shown for comparison. The 50ppm run was manually stopped due to its large computational cost. Colored phases correspond to negative feedback, white phases positive feedback. The years of the first instability between interactive and locked feedbacks are annotated.

required with interactive cloud feedbacks. The surface albedo-locked simulation does not reach a snowball instability as it enters quasi-equilibrium below the transition temperature of around $-15\text{K}^\circ\text{C}$ relative to PI. Similarly, the simulation reaches a quasi-equilibrium below $-10\text{K}^\circ\text{C}$ relative to PI when the water vapor feedback is locked (Appendix Fig. B2 C2). The water vapor feedback actually does not strengthen with cooling so is not itself the cause of the snowball instability, however its large positive feedback is important for allowing the surface albedo to trigger the instability.

Thus, the role of cloud feedback is different from that of the surface albedo feedback: 1) positive cloud feedbacks facilitate an early tropical cooling, and this effect is increased with stronger negative forcing, 2) positive cloud feedbacks ahead of the advancing sea-ice edge and its effect on tropical oceans accelerate the transition to snowball Earth and decrease the temperature of instability, as summarized in Fig. 4 5; 3) cloud feedbacks substantially increase the threshold CO_2 - CO_2 level required for snowball Earth initiation and 4) the entry of sea-ice advance of sea ice into the southern subtropics results in the surface albedo feedback exceeding the sum of the negative feedbacks. Whereas the surface albedo feedback is responsible for the transition and the main driver to the complete glaciation, our results emphasise an important contribution of cloud feedbacks. And as it




Figure 5. Contribution of cloud feedbacks to ocean cooling and interaction with sea ice. 1. Tropical surface cooling by positive low-level cloud feedback. 2. Sea surface cooling ahead of ice edge. 3. Faster sea-ice advance, facilitated by winds pushing sea ice equatorwards (Appendix Fig. C3).

205 happens, cloud-Cloud feedbacks are also the main cause of inter model spread in climate sensitivity (e.g. Zelinka et al., 2020), (e.g. Zelinka et al., 2020), a fact which we shall exploit in Section 5-5.

4 Model evidence for the snowball transition temperature

The temperature at which the climate transits towards a snowball Earth state is broadly similar across the different CO_2 forcing experiments (Fig. 3), as well as 4), when taking into account that it depends systematically on the degree of

210 ~~disequilibrium (Fig. 1). This is independent of~~ whether the experiment is initialized from a PI or an LGM state, as shown in Appendix A A, which demonstrates the importance of background temperature control over the climate feedbacks which drive the transition. In particular, sea-ice formation is primarily temperature-dependent, and the amplitude of the sea-ice albedo feedback is practically independent of the strength of the negative forcing (Fig. 2). We can expect such ~~state-dependency~~ temperature-dependency to be similar across climate models, which would contribute to them having a similar temperature

215 of entering the snowball transition. Similarly, the winds ~~pushing sea-ice~~ push sea ice equatorwards (Appendix Fig. B3 C3), in connection with temperature contrast between ~~sea-ice~~ sea ice and open ocean, ~~and which are~~. ~~This is~~ concomitant with positive cloud feedbacks ahead of sea-ice edge ~~are also~~, ~~which is~~ expected to behave in ~~similar ways~~ similarly across models since they broadly ~~simulate the same~~ represent the same general circulation. A simple geometric argument for a transition temperature close to 0°C is that it happens when the ice edge enters the sub-tropics at about 30 degrees latitude, leaving about half the Earth

220 still ice free and hence above the freezing temperature, and the other half at temperatures below. When abruptly decreasing the ~~CO₂-CO₂~~ concentration to 50 ppm (around 1/4 of ~~pre-industrial CO₂~~ of PI CO₂), we find hints of instability near the global mean temperature of 0°C (~~-near -15K relative to pre-industrial°C relative to PI~~), but the model ~~seems to be~~ is stable for 1848 years until it crashes due to thick ~~sea-ice~~ sea ice (Section 2). This indicates that it is unlikely that the ~~ineception~~ transition temperature happens at a temperature substantially warmer than 0°C.

225 Nevertheless, the transition temperatures of each phase show a slight shift to lower values under stronger negative forcing. ~~130~~ Therefore, the climate system deviates from pure ~~state-dependent~~ temperature-dependent behaviour as the strength of the radiative cooling and the speed of transition to snowball Earth increases. These ~~time-dependency~~ time-dependent effects have often been described in the literature following extreme cooling rates (~~Bendtsen and Bjerrum, 2002; Marotzke and Botzet, 2007~~), ~~and we believe to be~~ (Bendtsen and Bjerrum, 2002; Marotzke and Botzet, 2007), and we speculate that this is

230 due to the large heat capacity of the ocean, as well as differences in ocean mixed layer depths and direct ~~CO₂-CO₂~~ forcing across different regions: in certain circumstances, this can result in local temperatures dropping faster than global temperatures (Appendix Fig. B4-A C4-A), and hence out of sync with the sea-ice extent controlling the surface albedo feedback. Ocean convective mixing is also increasingly more efficient at evacuating heat for larger surface cooling, indicated by an increasing peak of the strength of the Atlantic meridional overturning circulation (AMOC) in Appendix Fig. B4-B C4-B. The associated

235 ocean currents on the contrary drag ~~sea-ice~~ sea ice polewards and can slow down the sea-ice edge advance (~~Voigt and Abbot, 2012~~) (Voigt and Abbot, 2012). All in all, we suggest slow, low forcing simulations are preferable when analysing snowball Earth transitions, as 1) fast transitions to snowball Earth are hardly realistic, as geological snowball states may form over millions of years (~~e.g. Schrag et al., 2002~~) (e.g. Schrag et al., 2002) and 2) fast transitions ~~involves~~ involve temporal effects which would depart from ~~state-dependency~~ temperature-dependency. All together, these considerations support the ~~higher~~ transition

240 temperature close to 0°C found in the experiments with weak forcing.




Figure 6. Top-of-atmosphere (TOA) radiative imbalance (Wm^{-2}) versus global surface temperature anomaly with relative to pre-industrial (PI) conditions (K°C) in abrupt $1/128\times\text{CO}_2$ simulations with the models MPI-ESM1.2-CR and CESM1.2.

5 Motivation for simulations and potential **Potential** constraint on climate sensitivity

While we expect the state-dependency-temperature-dependency to behave similarly across models, it is currently not known whether the temperature threshold of 0°C that we find in MPI-ESM1.2-CR is indeed universal. The CESM model family has notoriously performed several snowball Earth simulations (Yang et al., 2012b, a; Eisenman and Armour, 2024), and (Yang et al., 2012b, a; Eisenman and Armour, 2024), and while the model CESM2 also transits towards a snowball Earth around 0°C under similar CO_2 forcing (Eisenman and Armour, 2024), CO_2 forcing (Eisenman and Armour, 2024), models such as CCSM3 and CCSM4 can have stable waterbelt states with abrupt but smaller transitions, which anyway are not likely consistent with the evidence for recent LGM states. To further test this, we perform an abrupt $1/128\times\text{CO}_2$ simulation with the model CESM1.2. At $1/128\times\text{CO}_2$, CESM1.2 has been used for several deep-time simulations (Li et al., 2022), but to our knowledge never for snowball Earth states. CESM1.2 shows a similar instability than transition temperature at around -24°C below pre-industrial, or approximately -10°C absolute temperature, which is similar as MPI-ESM1.2 around a temperature of at the same forcing level (Fig. 6). Due to the time-dependent effects, which we discuss in this study, the transition temperature

is displaced under strong forcing. Therefore, under a low forcing, we expect the transition temperature to be closer to 0°C ; considering the time-dependency effects discussed in this study (Fig. 5). for CESM1.2, as was the case for MPI-ESM1.2.




Figure 7. Relationship between modelled LGM surface temperature anomaly relative to pre-industrial (PI) and the climate sensitivity of PMIP4 models. The upper bound on climate sensitivity can be constrained using the temperature at which the climate starts transiting to snowball Earth, which is different whether it is from surface temperatures or sea-surface temperatures. In A), the threshold is around -15K °C global mean surface temperature anomaly relative to PI (red). As a comparison, we calculate and in B), the corresponding threshold as seen in sea-surface temperatures is around -7.5°C relative to PI (grey). Several sensitivity tests for the computation of the upper bound on climate sensitivity are shown: A) if the upper bound of the transition was instead at -20K °C relative to PI (green), and B) if CESM2 is left out of the ensemble, but the other CESM models are included, where only sea-surface temperatures are available.

255 There is an ongoing discussion on whether climate models with large cloud feedbacks and consequently high climate sensitivity are compatible with the LGM (Zhu et al., 2022; Burls and Sagoo, 2022). (Zhu et al., 2022; Burls and Sagoo, 2022). Models with high climate sensitivity simulate LGM temperatures around the snowball Earth inception-transition temperature of around 0°C and are usually unstable. This in fact is to be expected if the inception-transition temperature is controlled by the temperature-dependent behaviour of the climate feedbacks, where the threshold before transiting to a snowball Earth state would be similar across models. Variations of that threshold could be due to for example different parameterizations of sea-ice albedo; tropical shallow clouds. In the case of tropical shallow cloud feedbacks, despite considerably increasing the sensitivity of the climate to enter snowball Earth, these feedbacks they do not seem to modify the inception temperature affect the snowball Earth transition temperature itself (Fig. 3.4).

260

It is therefore possible to ~~derive the estimate an~~ upper limit of climate sensitivity from the models with stable LGM states
265 ~~(Section 2 with an emergent constraint framework (Section 2))~~, using as snowball temperature threshold a global mean temper-
ature close to 0°C , ~~since both LGM and PI states transit towards a snowball Earth state at similar temperatures (Appendix Fig-~~
~~A1)-~~. In this way, we find that it is implausible that climate sensitivity exceeds ~~5.5 K (4.4–6.6, 90%~~ 6.2°C (3.9 – 8.4, 5–95%
confidence interval, Fig. 6 7). This upper limit is close to the sensitivity of CESM2 (5.6°C), which simulates an LGM tempera-
ture anomaly of ~~–11.3 K (-7.0 K in SSTs) (Zhu et al., 2021b)~~, $^{\circ}\text{C}$ (Zhu et al., 2021), just above the global mean temperature of
270 0°C . Additional support for this upper limit is obtained from a slightly more sensitive version of the model, which is unstable
when run with under LGM boundary conditions (Zhu et al., 2022).

On account of the uncertainty on whether the transition temperature would be close to 0°C across models, we test the
sensitivity of our constraint by decreasing the transition temperature to -5°C (Fig. 7). In this case, we find that the upper bound
on ECS cannot exceed 7.6°C (4.7 – 10.5, 2022)-

275 ~~This approach on climate sensitivity~~ 5–95% confidence interval). This is a relatively modest change of 1.4°C in the upper
bound, considering the large change in transition temperature of -5°C . The uncertainty on the transition therefore has a small
influence on our upper bound ECS estimate, and the value obtained here is in agreement with estimates from other lines of
evidence, or is in many cases a stronger constraint (Forster et al., 2021).

Finally, we further test the sensitivity of the upper bound estimates on climate sensitivity using the sea-surface temperature
280 threshold (estimated at close to -7.5°C using the same approach as in Fig. 1, in Appendix C5), which allows us to include
the CESM-family ensemble described in Section 2, and leave CESM2 out of the ensemble. When the CESM-family ensemble
is included, and whether CESM2 is or is not in the ensemble, the relationship is robust and displays similar characteristics
(Appendix B). The inclusion of CESM2 strengthens the relationship and increases the constraint on the upper bound on climate
sensitivity, but also increases the median estimates to 5.5°C (4.4 – 6.6, 5–95% confidence interval), in comparison to 4.9°C
285 (3.2 – 6.7, 5–95% confidence interval) when CESM2 is excluded. As such, the results are not overly sensitive to the inclusion
of CESM2, but overall confidence in the constraint would be significantly strengthened if more institutes would perform the
LGM simulations.

5.1 Motivation for simulations

The described approach to provide an upper bound on climate sensitivity could have a great potential, in particular as indepen-
290 dent constraints from paleoclimates are increasingly useful (Sherwood et al., 2020)- found to be increasingly useful (Sherwood et al., 2020;
Nevertheless, to support this constraint requires an effort from several modelling centres to both the constraint, it would be
beneficial that modelling centres publish their LGM simulations, even if they fail to stabilise, and as this may be used to narrow
down the uncertainty on the instability threshold. ~~It is~~

It is also important to know whether many models have a threshold gravitating around 0°C , similarly to that found for
295 MPI-ESM1.2-CR, MPI-ESM1.2-LR and CESM1.2.

To tackle this latter problem, we offer suggest the following experimental design, which is short and easy to replicate
easy
to replicate:

1. Run a pre-industrial-PI state, with the CO₂-CO₂ concentration set to 1/128 times the PI value, until the model reaches at least the initial instability leading to the snowball Earth state. The chosen CO₂-CO₂ concentration should be a fair trade off between a minimal-relatively short simulation length and the time-dependency effects discussed in this study. If time allows, complement with less strong reductions to investigate the time-dependency.
2. Generate a plot of top-of-atmosphere radiation imbalance versus surfce-surface temperature anomaly relative to PI, as in Fig. ~~3-~~ 4. Report the instability threshold temperature, from surface and sea-surface temperature, as well as the global sea-ice ~~concentration-~~
3. ~~For modelling centres simulating the LGM, we strongly suggest to publish runs which led to instabilities-~~ distribution.

The results could support the novel constraint on climate sensitivity presented in this study, but also help understanding ~~instabilities-~~ the challenges that some modeling groups experience in LGM simulations, a paleoclimate notoriously difficult to model, ~~in particular for higher climate sensitivity models.~~ We advocate that multi-model comparisons of snowball Earth transitions, as well as single-model ensembles with varying climate sensitivity, would better support the relationship between simulated LGM temperatures and the upper bound on ~~model Earth's~~ climate sensitivity. The main strength of this constraint is that it is independent of paleoclimate reconstructions and simply relies on the fact that Earth did not undergo a transition to a snowball state 21 000 years ago, or during any earlier glacial cycles during the Pleistocene, providing a valuable additional line of evidence to those already used in the community (~~Sherwood et al., 2020; Forster et al., 2021-~~ (Sherwood et al., 2020; Forster et al., 2021).)

315 *Data availability.* The outputs of the simulations shown in this study can be downloaded from <https://doi.org/10.5281/zenodo.8117483>

Author contributions. The idea of the study was conceived by MR. MR performed all simulations, analyses and figures. The paper was written by MR, JH, NS and TM.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. We thank three anonymous reviewers and Maria Rugenstein for their insightful comments. We thank Christoph Braun, Aiko Voigt and Raymond Pierrehumbert for the scientific discussions which helped advancing this study. We thank Eduardo Carril for designing Figure 5. The computations and data storage were enabled by resources provided by the National Academic Infrastructure for Supercomputing in Sweden (NAISS) and the Swedish National Infrastructure for Computing (SNIC) at the National Supercomputer Centre at Linköping University, partially funded by the Swedish Research Council through grant agreements no. 2022-06725 and no. 2018-05973. This

research has been supported by the European Research Council, H2020 European Research Council (highECS (770765)and, CONSTRAIN
325 (820829) [and NextGEMS \(101003470\)](#)).

References

- Arrhenius, S.: On the Influence of Carbonic Acid in the Air upon the Temperature of the Ground, *Philosophical Magazine and Journal of Science*, 41, 237–276, 1896.
- 330 Bendtsen, J. and Bjerrum, C. J.: Vulnerability of climate on Earth to sudden changes in insolation, *Geophysical Research Letters*, 29, 1–1, 2002.
- Braun, C., Hörner, J., Voigt, A., and Pinto, J. G.: Ice-free tropical waterbelt for Snowball Earth events questioned by uncertain clouds, *Nature Geoscience*, pp. 1–5, 2022.
- Budyko, M. I.: The effect of solar radiation variations on the climate of the Earth, *tellus*, 21, 611–619, 1969.
- Burls, N. and Sagoo, N.: Increasingly sophisticated climate models need the out-of-sample tests paleoclimates provide, *Journal of Advances in Modeling Earth Systems*, p. e2022MS003389, 2022.
- 335 Caballero, R. and Huber, M.: State-dependent climate sensitivity in past warm climates and its implications for future climate projections, *Proceedings of the National Academy of Sciences*, 110, 14 162–14 167, 2013.
- Colman, R. and McAvaney, B.: A study of general circulation model climate feedbacks determined from perturbed sea surface temperature experiments, *Journal of Geophysical Research: Atmospheres*, 102, 19 383–19 402, <https://doi.org/10.1029/97JD00206>, 1997.
- 340 Colman, R. and McAvaney, B.: Climate feedbacks under a very broad range of forcing, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2008GL036268>, 2009.
- Eisenman, I. and Armour, K. C.: The radiative feedback continuum from Snowball Earth to an ice-free hothouse, *Nature Communications*, 15, 6582, 2024.
- Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M., 345 and Zhang, H.: The Earth’s Energy Budget, Climate Feedbacks, and Climate Sensitivity, in: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 2021.
- Good, P., Andrews, T., Chadwick, R., Dufresne, J.-L., Gregory, J. M., Lowe, J. A., Schaller, N., and Shiogama, H.: nonlinMIP contribution to CMIP6: Model intercomparison project for non-linear mechanisms: Physical basis, experimental design and analysis principles (v1. 0), 350 *Geoscientific Model Development*, 9, 4019–4028, 2016.
- Gregory, J. M., Ingram, W., Palmer, M., Jones, G., Stott, P., Thorpe, R., Lowe, J., Johns, T., and Williams, K.: A new method for diagnosing radiative forcing and climate sensitivity, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2003GL018747>, 2004.
- Grose, M. R., Gregory, J., Colman, R., and Andrews, T.: What Climate Sensitivity Index Is Most Useful for Projections?, *Geophysical Research Letters*, 45, 1559–1566, <https://doi.org/10.1002/2017GL075742>, 2018.
- 355 Hargreaves, J. C., Annan, J. D., Yoshimori, M., and Abe-Ouchi, A.: Can the Last Glacial Maximum constrain climate sensitivity?, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL053872>, 2012.
- Haywood, A. M., Tindall, J. C., Dowsett, H. J., Dolan, A. M., Foley, K. M., Hunter, S. J., Hill, D. J., Chan, W.-L., Abe-Ouchi, A., Stepanek, C., et al.: The Pliocene Model Intercomparison Project Phase 2: large-scale climate features and climate sensitivity, *Climate of the Past*, 16, 2095–2123, 2020.

- 360 Hoffman, P. F., Abbot, D. S., Ashkenazy, Y., Benn, D. I., Brocks, J. J., Cohen, P. A., Cox, G. M., Creveling, J. R., Donnadieu, Y., Erwin, D. H., et al.: Snowball Earth climate dynamics and Cryogenian geology-geobiology, *Science Advances*, 3, e1600983, 2017.
- Hörner, J., Voigt, A., and Braun, C.: Snowball Earth initiation and the thermodynamics of sea ice, *Journal of Advances in Modeling Earth Systems*, p. e2021MS002734, 2022.
- Huusko, L. L., Bender, F. A., Ekman, A. M., and Storelvmo, T.: Climate sensitivity indices and their relation with projected temperature change in CMIP6 models, *Environmental Research Letters*, <https://doi.org/10.1088/1748-9326/ac0748>, 2021.
- 365 Jonko, A. K., Shell, K. M., Sanderson, B. M., and Danabasoglu, G.: Climate feedbacks in CCSM3 under changing CO₂ forcing. Part II: variation of climate feedbacks and sensitivity with forcing, *Journal of Climate*, 26, 2784–2795, 2013.
- Kageyama, M., Harrison, S. P., Kapsch, M.-L., Lofverstrom, M., Lora, J. M., Mikolajewicz, U., Sherriff-Tadano, S., Vadsaria, T., Abe-Ouchi, A., Bouttes, N., Deepak, C., Gregoire, L. J., Ivanovic, R. F., Izumi, K., LeGrande, A. N., Lhardy, F., Lohmann, G., Morozova, P. A., Ohgaito, R., Paul, A., Peltier, W., Poulsen, C. J., Quiquet, A., Roche, D. M., Shi, X., Tierney, J. E., Valdes, P. J., Volodin, E., and Zhu, J.: The PMIP4 Last Glacial Maximum experiments: preliminary results and comparison with the PMIP3 simulations, *Climate of the Past*, 17, 1065–1089, <https://doi.org/https://doi.org/10.5194/cp-17-1065-2021>, 2021.
- 370 Knutti, R., Masson, D., and Gettelman, A.: Climate model genealogy: Generation CMIP5 and how we got there, *Geophysical Research Letters*, 40, 1194–1199, <https://doi.org/10.1002/grl.50256>, 2013.
- 375 Lunt, D. J., Bragg, F., Chan, W.-L., Hutchinson, D. K., Ladant, J.-B., Morozova, P., Niezgodzki, I., Steinig, S., Zhang, Z., Zhu, J., Abe-Ouchi, A., Anagnostou, E., de Boer, A. M., Coxall, H. K., Donnadieu, Y., Foster, G., Inglis, G. N., Knorr, G., Langebroek, P. M., Lear, C. H., Lohmann, G., Poulsen, C. J., Sepulchre, P., Tierney, J. E., Valdes, P. J., Volodin, E. M., Dunkley Jones, T., Hollis, C. J., Huber, M., and Otto-Bliesner, B. L.: DeepMIP: model intercomparison of early Eocene climatic optimum (EECO) large-scale climate features and comparison with proxy data, *Climate of the Past*, 17, 203–227, <https://doi.org/10.5194/cp-17-203-2021>, 2021.
- 380 Marotzke, J. and Botzet, M.: Present-day and ice-covered equilibrium states in a comprehensive climate model, *Geophysical Research Letters*, 34, 2007.
- Mauritsen, T., Graverson, R. G., Klocke, D., Langen, P. L., Stevens, B., and Tomassini, L.: Climate feedback efficiency and synergy, *Climate Dynamics*, 41, 2539–2554, 2013.
- Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenéz-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornbluh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J.-S., Tian, F., Voigt, A., Vrese, P., Wieners, K.-H., Wilkenskjaeld, S., Winkler, A., and Roeckner, E.: Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1. 2) and Its Response to Increasing CO₂, *Journal of Advances in Modeling Earth Systems*, 11, 998–1038, <https://doi.org/10.1029/2018MS001400>, 2019.
- 390 Meraner, K., Mauritsen, T., and Voigt, A.: Robust increase in equilibrium climate sensitivity under global warming, *Geophysical Research Letters*, 40, 5944–5948, <https://doi.org/10.1002/2013GL058118>, 2013.

- 395 Muglia, J. and Schmittner, A.: Glacial Atlantic overturning increased by wind stress in climate models, *Geophysical Research Letters*, 42, 9862–9868, <https://doi.org/10.1002/2015GL064583>, 2015.
- Pierrehumbert, R., Abbot, D., Voigt, A., and Koll, D.: Climate of the Neoproterozoic, *Annual Review of Earth and Planetary Sciences*, 39, 417–460, 2011.
- Renoult, M., Annan, J. D., Hargreaves, J. C., Sahoo, N., Flynn, C., Kapsch, M.-L., Li, Q., Lohmann, G., Mikolajewicz, U., Ohgaito, R., Shi,
400 X., Zhang, Q., and Mauritsen, T.: A Bayesian framework for emergent constraints: case studies of climate sensitivity with PMIP, *Climate of the Past*, 16, 1715–1735, <https://doi.org/10.5194/cp-16-1715-2020>, 2020.
- Renoult, M., Sahoo, N., Zhu, J., and Mauritsen, T.: Causes of the weak emergent constraint on climate sensitivity at the Last Glacial Maximum, *Climate of the Past*, 19, 323–356, <https://doi.org/10.5194/cp-19-323-2023>, 2023.
- Roeckner, E., Mauritsen, T., Esch, M., and Brokopf, R.: Impact of melt ponds on Arctic sea ice in past and future climates as simulated by
405 MPI-ESM, *Journal of Advances in Modeling Earth Systems*, 4, 2012.
- Schmidt, G., Annan, J., Bartlein, P., Cook, B., Guilyardi, É., Hargreaves, J., Harrison, S., Kageyama, M., LeGrande, A., Konecky, B., Lovejoy, S., Mann, M. E., Masson-Delmotte, V., Risi, C., Thompson, D., Timmermann, A., Tremblay, L.-B., and Yiou, P.: Using palaeo-climate comparisons to constrain future projections in CMIP5, *Climate of the Past*, 10, 221–250, <https://doi.org/10.5194/cp-10-221-2014>, 2014.
- Schrag, D. P., Berner, R. A., Hoffman, P. F., and Halverson, G. P.: On the initiation of a snowball Earth, *Geochemistry, Geophysics, Geosystems*, 3, 1–21, 2002.
410
- Sherriff-Tadano, S., Abe-Ouchi, A., Yoshimori, M., Oka, A., and Chan, W.-L.: Influence of glacial ice sheets on the Atlantic meridional overturning circulation through surface wind change, *Climate dynamics*, 50, 2881–2903, <https://doi.org/10.1007/s00382-017-3780-0>, 2018.
- Sherwood, S., Webb, M., Annan, J., Armour, K., Forster, P., Hargreaves, J., Hegerl, G., Klein, S., Marvel, K., Rohling, E., Watanabe, M., Andrews, T., Braconnot, P., Bretherton, C., Foster, G., Hausfather, Z., von der Heydt, A., Knutti, R., Mauritsen, T., Norris, J., Proistosescu,
415 C., Rugenstein, M., Schmidt, G., Tokarska, K., and Zelinka, M.: An assessment of Earth’s climate sensitivity using multiple lines of evidence, *Reviews of Geophysics*, p. e2019RG000678, <https://doi.org/10.1029/2019RG000678>, 2020.
- Voigt, A. and Abbot, D.: Sea-ice dynamics strongly promote Snowball Earth initiation and destabilize tropical sea-ice margins, *Climate of the Past*, 8, 2079–2092, 2012.
- Voigt, A. and Marotzke, J.: The transition from the present-day climate to a modern Snowball Earth, *Climate dynamics*, 35, 887–905, 2010.
- 420 Voigt, A., Abbot, D. S., Pierrehumbert, R. T., and Marotzke, J.: Initiation of a Marinoan Snowball Earth in a state-of-the-art atmosphere-ocean general circulation model, *Climate of the Past*, 7, 249–263, 2011.
- Wall, C. J., Kohyama, T., and Hartmann, D. L.: Low-cloud, boundary layer, and sea ice interactions over the Southern Ocean during winter, *Journal of Climate*, 30, 4857–4871, 2017.
- Wetherald, R. and Manabe, S.: Cloud feedback processes in a general circulation model, *Journal of Atmospheric Sciences*, 45, 1397–1416,
425 [https://doi.org/10.1175/1520-0469\(1988\)045<3C1397:CFPIAG>3E2.0.CO;2](https://doi.org/10.1175/1520-0469(1988)045<3C1397:CFPIAG>3E2.0.CO;2), 1988.
- Yang, J., Peltier, W., and Hu, Y.: The initiation of modern soft and hard Snowball Earth climates in CCSM4, *Climate of the Past*, 8, 907–918, 2012a.
- Yang, J., Peltier, W. R., and Hu, Y.: The initiation of modern “soft Snowball” and “hard Snowball” climates in CCSM3. Part II: Climate dynamic feedbacks, *Journal of Climate*, 25, 2737–2754, 2012b.

- 430 Zelinka, M. D., Myers, T. A., McCoy, D. T., Po-Chedley, S., Caldwell, P. M., Ceppi, P., Klein, S. A., and Taylor, K. E.: Causes of higher climate sensitivity in CMIP6 models, *Geophysical Research Letters*, 47, e2019GL085782, <https://doi.org/10.1029/2019GL085782>, 2020.
- Zhu, J. and Poulsen, C. J.: Last Glacial Maximum (LGM) climate forcing and ocean dynamical feedback and their implications for estimating climate sensitivity, *Climate of the Past*, 17, 253–267, <https://doi.org/10.5194/cp-17-253-2021>, 2021a.
- Zhu, J. and Poulsen, C. J.: Last Glacial Maximum (LGM) climate forcing and ocean dynamical feedback and their implications for estimating climate sensitivity, *Climate of the Past*, 17, 253–267, 2021b.
- 435
- Zhu, J., Otto-Bliesner, B. L., Brady, E. C., Poulsen, C. J., Tierney, J. E., Lofverstrom, M., and DiNezio, P.: Assessment of equilibrium climate sensitivity of the Community Earth System Model version 2 through simulation of the Last Glacial Maximum, *Geophysical Research Letters*, 48, e2020GL091220, <https://doi.org/10.1029/2020GL091220>, 2021.
- Zhu, J., Otto-Bliesner, B. L., Brady, E. C., Gettelman, A., Bacmeister, J. T., Neale, R. B., Poulsen, C. J., Shaw, J. K., McGraw, Z. S., and Kay, J. E.: LGM paleoclimate constraints inform cloud parameterizations and equilibrium climate sensitivity in CESM2, *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002776, 2022.
- 440

Appendix A: Snowball state under solar forcing and from LGM conditions




Figure A1. Top-of-atmosphere radiative imbalance (Wm^{-2}) versus global surface temperature anomaly with pre-industrial relative to PI conditions (K°C) under solar forcing from LGM conditions. The similar Colored phases of positive (white) and correspond to negative (colored) climate feedbacks as from PI conditions are shown feedback, white phases positive feedback.

We verify how snowball Earth transitions differ when starting from LGM or PI conditions. MPI-ESM1.2-LR is numerically unstable at low CO_2 concentrations, so we compare solar-forced snowball Earth transitions between LGM and PI instead.

445 The LGM is characterized by large ice sheets and lower sea-level which can affect cloud feedbacks as well as global circulations (Muglia and Schmittner, 2015; Sherriff-Tadano et al., 2018; Zhu and Poulsen, 2021a; Renoult et al., 2023), thus can influence the transition to a snowball Earth state. Changes in continental distribution and ocean areas are known to affect

inception-transition temperatures. For instance, Voigt et al. (2011) indicates a snowball Earth transition at 96% of pre-industrial solar constant-PI solar constant and PI CO₂ concentration with Marinoan reconstructed paleogeography (~635 million years ago), which is characterized by agglomerated, bare-soil tropical continents, whereas it would require 94% or lower with PI continents. Tropical land masses reflect more radiations and therefore contribute substantially to snowball Earth transitions.

When initializing from LGM conditions (Fig. A1), the same main phases of negative and positive feedbacks are identified than from PI conditions. There are slight differences between both cases, for instance the LGM simulation show a single instability phase, whereas the runs starting from pre-industrial-PI still display hints of multiple phases when the solar constant is set at 85% of PI value. However, the inception-transition temperature connected to the main instability leading to the snowball Earth state happens at a broadly similar temperature than from PI conditions, between -15K-°C and -20K-relative to pre-industrial°C relative to PI, which demonstrate the role of the feedback dependency on temperature, irrelevant of the differences in boundary conditions.


Memory effects are expected to differ from LGM and PI conditions and likely explain the slight differences in transition temperatures towards the snowball state. Indeed, the LGM ocean is colder than in PI, and the initial thermal state of the ocean is known to affect the speed at which the snowball state is reached (Bendtsen and Bjerrum, 2002). From the LGM, reaching snowball Earth is therefore easier, as tropical oceans are colder and the evacuation of heat via ocean mixing is enhanced. This should add further in rendering the LGM simulations difficult for modern climate models, as they usually initialise their experiments from LGM states of previous generations.

465 **Appendix B: Emergent constraint on climate sensitivity from LGM global temperatures**

We analyse how sensitive the emergent constraint framework based on PMIP4, described in Methods and used in Fig. 7 is to the presence of CESM2, and whether the PMIP4 ensemble can be used in this study. Due to its high climate sensitivity, CESM2 is often seen as an outlier in the PMIP4 ensemble, which could bias statistical analysis. A solution used by Renoult et al. (2023) was to expand the PMIP4 ensemble with lower-ECS versions of CESM models, as to widen the ECS range of the PMIP4 ensemble and minimize the outlier effect of CESM2. Renoult et al. (2023) also showed that for the sake of the characteristics of the relationship (slope, intercept), as well as significance and predicted ECS, CESM2 and this lower-ECS CESM-family ensemble were in fact exchangeable if the relationship was based on tropical LGM sea-surface temperatures.

During the LGM, emergent-constraint based relationships were often found to be more significant when based on tropical temperatures, either because it could minimize the uncertainties on extra-tropical forcing (Hargreaves et al., 2012; Renoult et al., 2023) or simply because variations on climate sensitivity were more sensitive to tropical climate feedbacks during PMIP2 (Hargreaves et al., 2012). However, there is no reason that the relationship would not exist when based on global temperatures, as one could still expect a correlation between global temperatures and climate sensitivity, which is essentially a CO₂-based metric.

In Fig. B1, we use the temperatures of Renoult et al. (2023) and references within, as to verify whether on a global scale CESM2 and the CESM-family ensemble are also exchangeable, and if consequently CESM2 only acts as an outlier. We find the same conclusions as Renoult et al. (2023) on tropical sea-surface temperatures: a relationship which includes the PMIP






Figure C1. Evolution of tropical (30°S - 30°N) cloud cover within the first $15\text{K}^{\circ}\text{C}$ of cooling (425 years of simulation) in the $1/16\text{xCO}_2$ simulation.




Figure C2. Top-of-atmosphere radiative imbalance (Wm^{-2}) versus global surface temperature anomaly with pre-industrial relative to PI conditions (~~K~~) for the 1/128xCO₂ simulation with either surface albedo (fixalb), water vapor (fixq) or cloud (fixcld) feedbacks locked to that of the control state.




Figure C3. Surface wind field on sea-ice cover during the snowball Earth transition for the 1/16xCO₂ simulation. Strong winds push sea ice equatorwards in the tropical regions.




Figure C4. Analysis of tropical and global components affecting the rate of snowball Earth transitions. A) Differences in tropical and global SST-sea-surface temperature cooling under all CO₂ forcing. B) Strength of the Atlantic meridional overturning circulation (AMOC) measured at 26°N, 1000 m deep (Sv) under all CO₂ forcing. The average pre-industrial-PI AMOC strength, around 16 Sv, is highlighted.




Figure C5. Top-of-atmosphere radiative imbalance (Wm^{-2}) versus global sea-surface temperature anomaly relative to PI conditions ($^{\circ}\text{C}$) under the same CO_2 forcing as Fig. 4. When using sea-surface temperatures, the transition happens at around -7.5°C relative to PI for the lowest forcing ($1/8\times\text{CO}_2$).