

ECOMAN: an open-source package for geodynamic and seismological modelling of mechanical anisotropy.

Manuele Faccenda^{1*}, Brandon P. VanderBeek^{1*}, Albert de Montserrat², Jianfeng Yang³, [Francesco Rappisi](#)⁴, Neil Ribe⁵,

¹Dipartimento di Geoscienze, Università di Padova, 35131, Padova, Italy.

²Institute of Geophysics, ETH Zürich, Sonneggstrasse 5, 8092, Zürich, Switzerland.

³State Key Laboratory of Lithospheric Evolution, Institute of Geology and Geophysics, Chinese Academy of Sciences, Beijing, China.

⁴[School of Earth and Environment, University of Leeds](#).

⁵Lab FAST, Univ Paris-Saclay, CNRS, Bat 530, rue André Rivière, F-91405 Orsay, France.

Correspondence to: Manuele Faccenda (manuele.faccenda@unipd.it), Brandon P. Vanderbeek (brandonpaul.vanderbeek@unipd.it)

Abstract. Mechanical anisotropy related to rock fabrics is a proxy for constraining the Earth's deformation patterns. However, the forward and inverse modelling of mechanical anisotropy in 3D large-scale domains has been traditionally hampered by the intensive computational cost and the lack of a dedicated, open-source computational framework. Here we introduce ECOMAN, a software package for modelling strain-/stress-induced rock fabrics and testing the effects of the resulting elastic and viscous anisotropy on seismic imaging and mantle convection patterns.

Differently from existing analogous software, ~~ECOMAN can model strain-induced fabrics across all mantle levels and is~~ optimised to run ~~efficiently on multiple CPUs. It also enables modelling of~~ shape preferred orientation (SPO)-related structures ~~that can be~~ superimposed over lattice/crystallographic preferred orientation (LPO/CPO) fabrics, which allows the consideration of the mechanical effects of fluid-filled cracks, foliated/lineated grain-scale fabrics and rock-scale layering.

One of the most important innovations is the Platform for Seismic Imaging (PSI), a set of programs for performing forward and inverse seismic modelling in isotropic/anisotropic media using real or synthetic seismic datasets. The anisotropic inversion strategy is capable of recovering parameters describing a tilted transversely isotropic (TTI) medium, which is required to reconstruct 3D structures and mantle strain patterns and to validate geodynamic models.

1 Introduction

The study of the Earth's interior has been traditionally based on seismological and geodynamic modelling, the former providing important information about its present-day structure, composition and state (Chang et al., 2015; French and Romanowicz, 2015; Schaeffer et al., 2016; Debayle et al., 2020), and the latter about its dynamics and compositional evolution (Davies et

Deleted: ⁴

Deleted: ⁴

Field Code Changed

Field Code Changed

Field Code Changed

Field Code Changed

Deleted: the modelling of

Deleted: has been extended to

Deleted: it has been

Deleted: across

Deleted: , yielding strong scaling efficiency

Deleted: In addition,

Deleted: can be modelled and

40 al., 2012; Cramer and Tackley, 2014; Müller et al., 2022). Seismological and geodynamical modelling are very often
41 conducted independently, which creates mechanical and geometrical inconsistencies across the models, hampers the
42 interpretation of seismic observations in terms of geodynamic processes, and exacerbates the non-uniqueness of geodynamic
43 model predictions. An alternative approach is combining computational seismology and geodynamics with mineral physics,
44 which provides a comprehensive understanding of the Earth's interior processes, seismic behaviour, and material properties.
45 This multidisciplinary methodology has been used in previous studies to post-process geodynamic flow calculations with
46 thermodynamically self-consistent models of mantle mineralogy and converting thermal structure into isotropic elastic
47 parameters. The obtained seismic mantle structure can then be used in simulations of global wave propagation, such that
48 specific hypotheses on mantle dynamics can be tested directly against seismic data (Styles et al., 2011; Schubert et al., 2012;
49 Maguire et al., 2018). The inverse procedure consists of converting seismic anomalies into density anomalies driving mantle
50 flow models and is typically employed to quantify dynamic topography and mantle viscosity structure, and to reproduce large-
51 scale mantle flow patterns (e.g., Bunge et al., 2003; Steinberger and Calderwood, 2006; Rudolph et al., 2015). However,
52 isotropic seismic imaging provides limited information regarding local-/regional-scale dynamical processes (Fraters and
53 Billen, 2021), and a better way to couple the geodynamic evolution and seismological structure of the Earth's interior is by
54 accounting for the strain-/stress-induced mechanical anisotropy of crustal and mantle rocks.

55 Mechanical anisotropy refers to the directional dependence of mechanical properties in a material, and is well known to affect
56 both elastic and viscous deformational behaviour. Mechanical anisotropy depends on several factors, including the lattice or
57 crystal preferred orientation (LPO, CPO) of intrinsically anisotropic minerals, and extrinsic mechanisms related to the shape
58 preferential orientation (SPO) of melt, fluid, or air-filled fractures and non-spherical pores, and grain- or rock-scale
59 compositional layering. Most micro- and macro-scale fabrics are acquired as a function of the cumulative deformation and
60 material mechanical properties, and as such they constitute an important source of information about the Earth's dynamical
61 behaviour.

62 Elastic anisotropy is directly connected to seismic anisotropy, which is a phenomenon in which the seismic wave speed varies
63 as a function of the propagation direction. It is mainly observed in the crust, mantle boundary layers, and inner core (Almqvist
64 and Mainprice, 2017; Karato, 1998; Kendall, 2000; Long and Becker, 2010; Deuss, 2014). Understanding and modelling
65 seismic anisotropy is crucial for determining long-term deformational patterns and the present-day stress field in the crust, and
66 constraining geodynamic modelling predictions (Jadamec and Billen, 2010; Hu et al., 2017; Zhou et al., 2018; Lo Bue et al.,
67 2022). In addition, the ability to account for anisotropic effects can improve the quality of subsurface imaging. Indeed, it has
68 been demonstrated that, because of the uneven seismic ray coverage, failing to account for seismic anisotropy may generate
69 strong artefacts that substantially bias our understanding of mantle structures and dynamics in different tectonic settings
70 (Bezada et al., 2016; VanderBeek and Faccenda, 2021; VanderBeek et al., 2023; Faccenda and VanderBeek, 2023).
71 Considering the widespread presence of seismic anisotropy, anisotropic seismic models provide a more realistic representation
72 of the Earth's subsurface compared to isotropic models.

Deleted: in

Deleted: ,

75 Viscous anisotropy modelling refers to the study and simulation of materials that exhibit varying degrees of viscosity
 76 (resistance to viscous deformation) in different directions. Although viscous anisotropy has been traditionally associated with
 77 the mechanical behaviour of multi-layered media (Mühlhaus et al., 2002; Kocher et al., 2006), it has been also observed in
 78 experimentally-deformed mica-rich and olivine crystal aggregates (Shead and Kronenberg, 1993; Hansen et al., 2012).
 79 Previous numerical studies demonstrated that viscous anisotropy can potentially stabilise long-wavelength convective patterns
 80 (Christensen, 1987; Mühlhaus et al., 2004) and more generally affect processes such as plate motion (Király et al., 2020), post-
 81 glacial rebound (Han and Wahr, 1997), lithospheric shear zone reactivation (Tommasi et al., 2009) and dripping (Lev and
 82 Hager, 2008).

83 Over the last few years a few attempts have been made to integrate micro-mechanical modelling of fabric evolution with large-
 84 scale geodynamic models using either directors (Lev et al., 2008; Halter et al., 2022), the CPO model D-REX (Kaminski et
 85 al., 2004; Becker et al., 2006; Jadamec and Billen, 2010; Faccenda and Capitanio, 2013; Faccenda, 2014; Ito et al., 2014; Hu
 86 et al., 2017; Zhou et al., 2018; Fraters and Billen, 2021), or the CPO model Visco-Plastic Self Consistent (VPSC; Tommasi et
 87 al., 2009; Li et al., 2014). However, each of these methodologies has its own limitations mainly associated with either the
 88 accuracy of the estimates, the large computational burden or software accessibility, which have impeded a more widespread
 89 diffusion in the geodynamic community. At the same time, the recovery of 3D seismic anisotropy patterns has been
 90 traditionally considered intractable due to the highly underdetermined nature of the inverse problem, and although it has been
 91 addressed by several previous studies (e.g., Debayle et al., 2005; Panning et al., 2006; Abt and Fischer, 2008; Long et al.,
 92 2008; Wookey, 2012; Mondal and Long, 2019), only recently a theoretical background and computational algorithms have
 93 been developed to simultaneously invert for P- and/or S-wave isotropic velocity anomalies and anisotropy resulting from
 94 arbitrarily oriented structures. (Munzarova et al., 2018; VanderBeek and Faccenda, 2021; Rappisi et al., 2022; Wang and Zhao,
 95 2022; VanderBeek et al., 2023; Del Piccolo et al., 2023). Yet, to date there is no freely-available software capable of modelling
 96 seismic anisotropy related to arbitrarily oriented structures.

97 In this contribution we present the new and open-source software package ECOMAN (Exploring the CONsequences of
 98 Mechanical ANisotropy) that enables linking seismology and geodynamics by providing a set of computationally optimised
 99 programs for (i) estimating rock mechanical anisotropy as a function of the geodynamic model deformation history, and
 100 compositional, rheological, stress, pressure, temperature, fields, and (ii) solving forward/inverse seismological problems
 101 accounting for seismic anisotropy. In the next sections, we first describe the different ECOMAN modules, after which we
 102 discuss the advantages and limitations of the software package, and the roadmap for future developments.

103
 104 Table 1. Abbreviations and their description. Units are indicated for dimensional physical properties.

Abbreviation	Description	

Deleted: body

Deleted: seismic anisotropy in 3D anisotropic media

UM	Upper mantle	
UTZ	Upper mantle transition zone	
LTZ	Lower mantle transition zone	
LM	Lower mantle	
LPO/CPO	Lattice/crystal preferred orientation	
SPO	Shape preferred orientation	
FSE	Finite Strain Ellipsoid, defined by the eigenvalues and eigenvectors of LS	
STILWE	Smoothed Transversely Isotropic Long-Wavelength Equivalent	
DEM	Differential Effective Medium	
VPSC	Visco-Plastic Self Consistent	
MDM	Modified Director Method	
		Units
ρ	Density	kg/m ³
P	Pressure	Pa
T	Temperature	K
Fd	Fraction of dislocation creep deformation	-
\mathbf{V}, V_i	Velocity vector and its components	m/s
\mathbf{F}, F_{ij}	2 nd -order deformation gradient tensor and its components	-
\mathbf{LS}, LS_{ij}	2 nd -order left stretch tensor and its components	-
$\mathbf{C}, C_{\alpha\beta}$	4 th -order elastic tensor and its components in Voigt notation	GPa
$\mathbf{S}, S_{\alpha\beta}$	4 th -order compliance tensor and its components in Voigt notation	GPa ⁻¹
$\boldsymbol{\eta}, \eta_{\alpha\beta}$	4 th -order viscous tensor and its components in Voigt notation	Pa s

Deleted: normalised

Deleted: -

110 **2 Software package structure**

111 ECOMAN includes several programs that are complementary and can be grouped into three main categories (Fig. 1):

112

113 1) programs that estimate strain/stress-induced rock fabrics (LPO and SPO) and their elastic and viscous anisotropic
114 mechanical properties (D-REX_S, D-REX_M, EXEV);

115 2) programs that post-process the simulated rock fabrics for visualisation of their isotropic/anisotropic mechanical
116 properties and deformational history (VIZTOMO, VIZVISC), and format the elastic tensors generating input files for
117 seismological synthetics (VIZTOMO);

118 3) programs that test the elastic response of anisotropic media by performing seismological forward/inverse modelling
119 and, in particular, isotropic and anisotropic seismic tomographies on synthetic and real seismic datasets (SKS-SPLIT,
120 PSI).

121

122 Most of the code modules are written in the Fortran programming language, except for the PSI program that is written in Julia.

123 Visualisation of the output is done through the MATLAB MTEX toolbox (Mainprice et al., 2011) for single aggregate fabrics,
124 and ParaView (Ahrens et al., 2005) for 2D and 3D simulations.

125




Figure 1. ECOMAN structure and flow chart. Coloured boxes denote programs that compute rock fabrics (green), post-process the elastic (C), viscous (η) and deformation gradient (F) tensors for visualisation and/or data formatting for seismological synthetics (orange), and perform seismological forward/inverse modelling on synthetic or real datasets (yellow). Input data are from geodynamic modelling or real seismic datasets (blue). Visualisation of the mechanical properties and LPO can be done with the MTEX MATLAB toolbox for single crystal aggregates or the software ParaView for large-scale simulations.

2.1 Rock fabrics and mechanical properties simulations

The evolution of the strain-induced LPO can be simulated with the D-REX model (Kaminski et al., 2004), which has been adapted to reproduce the fabric evolution of a single (D-REX_S) or multiple (D-REX_M) two-mineral phase aggregates representative of the entire Earth's mantle. The accuracy of the D-REX model has been tested against analytical solutions derived by Fraters and Billen (2021) (Appendix A). Five sets of two-mineral phases mantle aggregates can be defined as a function of depth or density ρ (Fig. 2; see Table 1 for a list of abbreviations and physical properties):

- 1) **olivine + enstatite**, for the upper mantle (UM: 0-410 km or $3000 < \rho \leq 3650$ kg/m³);
- 2) **wadsleyite + majoritic garnet**, for the upper mantle transition zone (UTZ: 410-520 km or $3650 < \rho \leq 3870$ kg/m³);

Deleted: for

Deleted: mantle

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

144 3) ringwoodite + majoritic garnet, for the lower mantle transition zone (LTZ: 520-660 km or $3870 < \rho \leq 4150 \text{ kg/m}^3$);

145 4) **bridgmanite** + ferropericlaase, for the lower mantle (LM: 660-2900 km or $\rho > 4150 \text{ kg/m}^3$);

146 5) **post-perovskite** + ferropericlaase, for the bottom of the lower mantle according to the parametrized phase boundary

$$P(\text{GPa}) = 98.7 + T(\text{K}) \cdot 0.00956 \quad (\text{Oganov and Ono, 2004}).$$

Formatted: Font: (Default) Times New Roman

Deleted: from

Deleted: (


149 The strain-induced LPO is computed for the phases in bold, while other major phases such as garnet, ringwoodite and
150 ferropericlaase are considered to be isotropic and their distribution is set to be random. Thus, no LPO is computed for the LTZ,
151 such that (minor) anisotropy arises only when SPO modelling due to compositional layering is active (section 2.1.3; Faccenda
152 et al., 2019). The LPO of wadsleyite, bridgmanite and post-perovskite are computed according to the same D-REX
153 methodology originally applied to upper mantle phases and described in detail in Kaminski and Ribe (2001), and Kaminski et
154 al. (2004), but including additional slip systems as indicated in Table A1. For each of these new phases it is possible to define
155 in the D-REX_S and D-REX_M input files the same free parameters of the D-REX model (the dimensionless grain boundary
156 modility M^* , nucleation parameter λ^* , threshold volume fraction χ , and in addition the power-law exponent for dislocation
157 creep deformation within the grains) as for upper mantle aggregates. The full elastic tensor is then calculated according to the
158 crystal orientation, volume fraction, phase abundance, P-T conditions, bulk rock composition, and using Voigt-Reuss-Hill
159 averaging schemes (see section 2.1.2 and Appendix B for more details).

160 The elastic properties related to strain/stress-induced SPO fabrics can instead be calculated at the grain- or rock-scale and for
161 layered or two-phase (matrix-ellipsoidal inclusions) systems using the isotropic elastic moduli of the different (fluid, mineral,
162 rock) components (EXEV).

163 The elastic properties and density of the aggregates characterised by LPO and/or SPO fabrics are estimated at relevant mantle
164 P-T conditions using the single crystal elastic moduli and their P-T derivatives for the main mineral phases and compiled from
165 different mineral physics studies, together with lookup tables of the isotropic elastic moduli, density and mineral phase volume
166 fraction generated with MMA_EoS (Chust et al., 2017) for five bulk rock compositions (dunite, harzburgite, pyrolite, MORB,
167 pyroxenite) at temperature and pressure ranges of $T = 300 : 50 : 4500 \text{ K}$, $P = 0 : 0.1 : 140 \text{ GPa}$, respectively (see Appendix B
168 and Tables B1 and B2).

Deleted: and in the

Deleted: range



173
174 **Figure 2.** Two-phase aggregates defined by the density crossovers and Brd-pV phase transition for a pyrolytic mantle composition.

Formatted: Font: 9 pt

175 2.1.1 D-REX_S

176 D-REX_S is a program designed for modelling the evolution of strain-induced LPO fabrics and related elastic properties of a
 177 single, two-mineral phases mantle aggregate, as a function of the imposed flow field, amount of strain, crystal plasticity, P-T
 178 conditions and additional effects related to SPO fabrics. It builds on the original D-REX software (Kaminski et al., 2004) for
 179 modelling the strain-induced LPO, and it includes MATLAB scripts to generate pole figures of the LPO and
 180 isotropic/anisotropic seismic properties with the MTEX software (Mainprice et al., 2011) (Fig. 3a-d), together with the
 181 possibility to display the evolving fabric strength (M-index, J-index) and the fraction of different anisotropy components
 182 (obtained via tensor decomposition; Browaeys et al., 2004; Fig. 3e-h).

Deleted: 2

183 D-REX_S is particularly useful for those users who are not familiar with LPO modelling, and more in general, to anyone
 184 interested in performing parameter sensitivity tests on different mantle mineral aggregates before launching large-scale
 185 simulations. In addition, the microstructures generated with D-REX_S can be used in the D-REX_M 2D-3D simulations to
 186 impose pre-existing (e.g., fossil) fabrics on multiple crystal aggregates located within a specific subdomain (see section 2.1.2).

Deleted: 2

187 As an application and with the aim of debunking the common misconception that the D-REX model can only produce too
 188 strong LPO fabrics, here we show that with a suitable choice of D-REX free parameters it is possible to reproduce both weak
 189 and strong olivine fabrics as recorded in natural samples (Warren et al., 2008) and high-strain laboratory experiments (Hansen
 190 et al., 2014; Tasaka et al., 2017) (Fig. 4).





Figure 3. D-REX S output for an upper mantle aggregate (Ol:Opx=70:30) subjected to a simple shear deformation of 1. Pole figures of the olivine (a) and orthopyroxene (b) crystallographic axes; (c) pole figures of Vp, Vs1, AVs = $200(Vs1 - Vs2)/(Vs1 + Vs2)$ with superimposed Vs1 polarisation directions evaluated from the elastic tensor of the two-phase aggregate. (d) same as (c) but with the superimposed effect of an SPO fabric due to 5% melt-filled cracks aligned at -30° from the principal stress (i.e., at 15° from the horizontal plane); (e) fraction of total anisotropy relative to the full elastic tensor and (f) contribution of 5 anisotropic classes relative to the total anisotropy; (g) P- and S-wave radial and azimuthal anisotropy and (h) eta parameter = $F/(A-2L)$ for the elastic tensors computed with the Voigt (continuous lines), Reuss (dashed lines) and Hill (crosses) averaging schemes, respectively.

Deleted: 2

Formatted: Space After: 10 pt, Line spacing: single

Deleted:

Deleted: 1



204 **Figure 4** –M-index (top) and J-index (bottom) for olivine fabrics developing in simple shear deformation as computed with D-REX_S using
 205 different parameters and compared with data reported in the literature as indicated in the legend. Fixed parameters are $\lambda^* = 5$, olivine
 206 $nCRSS = (1, 2, 3, \infty)$, and the aggregates have a phase abundance of *Ol:Ens* = 70:30 with 1000 crystals for each phase.
 207

208 2.1.2 D-REX_M

209 D-REX_M is a program that computes the evolution of the LPO and related elastic properties of multiple, two-mineral phases
 210 mantle aggregates, as a function of the single crystal plastic and elastic properties, and of the flow field, deformation
 211 mechanisms and P-T conditions resulting from 2D-3D geodynamic simulations. It builds on the original D-REX software,
 212 which includes routines for estimating the strain-induced LPO and elastic properties (i) for upper mantle polycrystalline

Formatted: Font: 9 pt

Deleted: estimates

214 aggregates only, (ii) using single crystal elastic tensors derived at room P-T conditions and averaged using a Voigt scheme,
215 (iii) in a 2D Cartesian domain, (iv) for a single (steady-state) flow field, and (v) whereby the whole deformation is considered
216 to be accommodated by dislocation creep assisted by grain-boundary sliding (Kaminski et al., 2004). D-REX_M additionally
217 models:

- 218 • fabrics relevant to the mid and lowermost mantle (Faccenda, 2014), including those with post-perovskite. Phase
219 transitions can be set to occur at predefined depths (e.g., 410 km, 660 km). Density crossovers (which allow modelling
220 the deflection of phase boundaries with a non-zero Clapeyron-slopes; Fig. 7), and parameterized phase boundaries as
221 for the case of post-perovskite (Oganov and Ono, 2004) are also included;
- 222 • elastic properties and density as a function of the bulk rock composition and local P-T conditions (Faccenda, 2014;
223 Chang et al., 2016; Ferreira et al., 2019) (Appendix B). The isotropic component of the elastic tensors and density are
224 taken from the lookup tables generated by MMA-EoS for a given mantle lithology (Table A1), and the anisotropic
225 component from the mineral single crystal elastic moduli and their pressure and temperature derivatives listed in
226 Table B2. This strategy ensures a gradual transition of the seismic properties at phase boundaries where phase
227 transformations occur. Voigt-Reuss-Hill averaging schemes of the elastic moduli are included;
- 228 • non-steady-state flows in 2D/3D Cartesian and polar grids (Faccenda and Capitanio, 2012, 2013, Hu et al., 2017;
229 Zhou et al., 2018; Lo Bue et al., 2022; Faccenda and VanderBeek, 2023). In polar coordinates the velocity gradient
230 tensor must be computed in the external Cartesian reference frame, as described in Appendix C. The global-scale
231 models are spatially discretized using the so-called Yin-Yang grids (Kageyama and Sato, 2004). Several examples
232 (cookbooks) are provided on how to use the software in different coordinate systems and in steady-state or time-
233 dependent flow conditions;
- 234 • fabric evolution in the presence of multiple creep mechanisms. At any time step, the fraction of deformation
235 accommodated by dislocation creep in a given point of the geodynamic model defines the fraction of time spent for
236 intracrystalline deformation assisted by grain-boundary sliding according to the D-REX model. The remaining time
237 is used to apply, when present, fluid deformation rotation to the whole crystal aggregate (e.g., Hedjazian et al., 2017)
238 (Appendix D);
- 239 • a pre-existing (fossil) fabric (pre-computed with D-REX_S) within a subdomain, typically the lithosphere. This is
240 often the case for geodynamic models where the lithosphere accretion is not modelled, and its geometry is initially
241 prescribed.

242
243 The D-REX_M input files should contain information about the geodynamic model evolution. Critical information are the
244 components of the velocity vector \mathbf{V} field, and, for time-dependent flow models, the total elapsed time and timestep. These are
245 used to compute the velocity gradient tensor, and the LPO evolution and advection of the crystal aggregates. Additional fields
246 defined on the Cartesian/spherical grid that can be included in the input files are:

Deleted: c

Deleted: assuming steady-state flow

Deleted: null

Deleted: 5

Deleted: ,

Deleted: ,

Deleted:

Deleted: c

Deleted: ;

Deleted: c

- 257
- when the geodynamic model is thermo-mechanical, the temperature T and total pressure P fields, which can be used to compute the single crystal elastic tensor and aggregate phase transitions as a function of the local P - T conditions;
- 258
- when the rheological model of the geodynamic simulation is based on multiple visco-plastic deformation mechanisms, the fraction of deformation accommodated by dislocation creep $F_d = \eta_{\text{disl}}/\eta_{\text{eff}}$, where $0 \leq F_d \leq 1$, η_{disl} is the viscosity calculated with the dislocation creep flow law, η_{eff} is the effective viscosity calculated with the harmonic average of each of the viscosities representing a different deformation mechanism.
- 262

Formatted: Font: (Default) Times New Roman

Formatted: Font: (Default) Times New Roman

263

264 In summary, the time and the velocity field \mathbf{V} are essential information, while the P , T , F_d fields are optional and depend on the type of (mechanical vs. thermomechanical) geodynamic and (single vs. multiple visco-plastic deformation mechanisms) rheological models.

Deleted: Summarising


267 While the \mathbf{V} , P , T , F_d fields are defined on the Eulerian grid, the distribution, size, modal composition and mechanical properties of mantle aggregates are defined on the Lagrangian particles. After initialising the Eulerian grid and Lagrangian particles, the entire run consists of three main steps: (i) backward advection of the particles for a given time span or up to a maximum cumulated strain; (ii) forward advection and update of the LPO and deformation gradient tensor \mathbf{F} ; (iii) computation of the full elastic tensor and creation of the output file. This strategy ensures a homogeneous final distribution of the aggregates (coincident with the initial one) which is desired for visualization of the aggregates mechanical and physical properties and for running seismological synthetics. The D-REX_M output file(s) includes, for each mineral aggregate, the elastic tensor \mathbf{C} , density, and the deformation gradient tensor \mathbf{F} . These properties can be then processed and visualised with the software VIZTOMO (section 2.2.1).

276 D-REX_M is parallelized using a hybrid MPI and OpenMP scheme to take advantage of multiple CPUs architectures of modern HPC clusters. Runtime is obviously affected by the number of crystal aggregates, number of crystals per aggregate and of active slip systems per crystal, and number of timesteps (Fig. 5, top). The parallel efficiency is 70-90%, and the update of the LPO and \mathbf{F} tensor during forward advection is the most time-consuming part of the run (Fig. 5, centre, bottom). The small performance degradation is due to the initialization of the Eulerian/Lagrangian grids and arrays, and to I/O operations which are executed serially within each process. As a result, the efficiency of the time-dependent flow models is lower than that of steady-state models, as the latter only require a single velocity (and P , T , F_d) field to be loaded and processed.

Deleted: 3

Deleted: ranges

Deleted: 3



287
288

289 **Figure 5.** Runtime (top), speedup (centre) and parallel efficiency (bottom) of D-REX_M for initial backward advection of aggregates
 290 (“backward”), forward advection and updating of the LPO and \mathbf{F} tensor (“forward”), full elastic tensor computation and output file creation
 291 (“tensor”), and entire run (“total”). Results are shown for two models included in the cookbooks: the 3Dspherical_global model (steady-
 292 state flow, 96 timesteps, 1327606 aggregates, LPO computed only for 260474 upper mantle aggregates) and the 3Dspherical_sinkingslab
 293 model (dashed lines; time-dependent flow, 20 timesteps, 38509 aggregates, LPO computed for 25177 upper mantle and 6587 upper mantle
 294 transition zone aggregates). Runs performed on a HPE Superdome Flex (8 CPUs, 28 cores Intel Xeon(R) PLATINUM 8180 @ 2.50GHz)
 295 using from 1 to 8 nodes.

296

297 2.1.3 EXEV

298 EXEV includes routines to compute the EXtrinsic Elastic and Viscous anisotropy using Effective Medium Theory modelling
 299 for a multi-component layered system (Smoothed Transversely Isotropic Long-Wavelength Equivalent, STILWE; Backus,
 300 1962) or a two-component system with similar ellipsoidal inclusions in a uniform background matrix (Differential Effective
 301 Medium, DEM; e.g., Mainprice, 1997).

Deleted: 3

303 The elastic tensor \mathbf{C} due to SPO fabrics can be either estimated independently or, when using the DEM approach, superimposed
304 on that obtained from the strain-induced LPO modelling. SPO fabrics that can be modelled are those related to rock- or grain-
305 scale layering (e.g., Faccenda et al., 2019), or to the presence of preferentially aligned ellipsoidal inclusions (e.g., melt-/fluid-
306 filled cracks). The user then needs to specify:

- 307 • for grain-scale layered fabrics, a dominant ultramafic or mafic lithology. In this case the mineral phase proportions
308 from the MMa-EoS lookup tables define the mixture for the STILWE model;
- 309 • for rock-scale layered fabrics, the relative abundance of the five available ultramafic-mafic lithologies (dunite,
310 harzburgite, pyrolyte, basalt and pyroxenite) defining the mixture for the STILWE model;
- 311 • for matrix-inclusion fabrics, the elastic tensors of the two components and the inclusion's shape and volume fraction
312 as required by the DEM modelling. The matrix elastic tensor can be replaced with that from the LPO modelling in
313 order to estimate the combined effect of LPO and SPO fabrics.

314 The SPO fabrics can then be oriented at any angle relative to the principal Finite Strain Ellipsoid (FSE) axis or, in case of
315 cracks, the local principal stress obtained from the “present-day” (i.e., last) velocity field.

316 Summarising, SPO fabric modelling requires one or more of the following: \mathbf{F} and/or \mathbf{C} obtained from D-REX_M; \mathbf{P} , \mathbf{T} and/or
317 \mathbf{V} and/or fluid/melt fraction fields for the “present-day” state of the geodynamic model. Consequently, this modelling is
318 performed when post-processing the D-REX_M output with the software VIZTOMO (see section 2.2.1).

319 The total or deviatoric component of the viscous tensor $\boldsymbol{\eta}$ due to SPO fabrics is estimated for two-phase systems with ellipsoidal
320 weak/hard inclusions using the DEM theory, and the parametrization of the viscous tensor evolution and orientation as a
321 function of the cumulated deformation (\mathbf{F}) obtained following de Montserrat et al. (2022). Indeed, most if not all, mantle levels
322 are composed of two main mineral phases that control both the elastic and viscous properties. The case of a multi-component
323 layered medium is not considered because its viscous tensor can be either approximated with flat inclusions, or more simply
324 computed using the Voigt and Reuss averages of the layers' isotropic viscosity.

325 The first modelling phase requires running subprogram DEMviscous to generate a database of viscous tensors for a range of
326 inclusion shapes and volume fractions, and inclusion-matrix viscosity contrasts. The latter implies that the viscous moduli are
327 dimensionless and can therefore be interpreted as scaling factors with respect to an isotropic effective viscosity of the bulk
328 rock or most abundant mineral phase. Subsequently, the database can be exploited by large-scale geodynamic simulations to
329 either (i) return the viscosity tensor $\boldsymbol{\eta}$ from a look-up table for every point of the computational domain, which can be
330 superimposed on the isotropic effective viscosity computed from flow laws (coupled mechanical simulations), or (ii) estimate
331 $\boldsymbol{\eta}$ (uncoupled mechanical simulation) and/or visualise its anisotropic viscous properties for the “present-day” state of the
332 geodynamic model with the software VIZVISC (section 2.2.2).

333

Formatted: Font: Bold

334 **2.2 Visualisation of the mechanical properties and data formatting in preparation for seismological synthetics**

335 **2.2.1 VIZTOMO**

336 VIZTOMO processes the D-REX_M output for the visualisation of the aggregates' elastic and deformational history properties.
337 Estimation of extrinsic anisotropy effects via the EXEV routines is possible at this stage. The properties of the **C** and **F** tensors
338 can be determined either at the position of the Lagrangian mantle aggregates, or, in case of the elastic properties and density,
339 interpolated to a structured (tomographic) grid. In the latter case the grid can be saved in a format suitable for generating
340 synthetic seismic datasets via the PSI_D package (see section 2.3.2) or for 3D waveform simulations in SPECFEM
341 (Komatitsch and Tromp, 1999).

342 Several properties of the elastic tensor **C** can be visualised:

- 343 • Isotropic or ray-path dependent velocity anomalies and anisotropic elastic properties of body-waves (i.e., P-wave
344 anisotropy and direction of maximum P-wave velocity; direction and magnitude of maximum S-wave splitting delay
345 time; S-wave radial and azimuthal anisotropy);
- 346 • reflection/transmission energies resulting from the whole range of P-S conversions occurring at discontinuities (useful
347 for studies based on receiver function analysis);
- 348 • the fraction of the elastic tensor anisotropic component relative to the total and the relative contributions of five
349 different anisotropy classes (hexagonal, orthorhombic, tetragonal, monoclinic, triclinic) obtained through elastic
350 tensor decomposition (Browaeys and Chevrot, 2004);
- 351 • the orientation of the hexagonal symmetry axis (already present in the original D-REX).

352 The deformation history stored in **F** can be visualised in terms of the FSE shape and orientation and/or length or orientation of
353 its minimum and maximum semi-axes.

354 The different fields are saved in specific file formats which can be imported by the open source ParaView software (Ahrens et
355 al., 2005) for visualisation. Figures 6, 7, 8, 9 display some of these fields computed for different geodynamic models.


356

Deleted: et al.

Deleted:

Deleted: 4, 5, 6, 7


Deleted: 4



361
362
363
364
365
366

Figure 6. VIZTOMO output for the model 2Dpolar_convection available in the cookbooks. The steady-state 2D incompressible flow field in cylindrical coordinates is the result of the analytical solution shown in section 9.3 of the software manual and coded in the MATLAB script cookbooks/2Dpolar_convection/polarcell.m. (a,b) isotropic Vp and Vs (m/s); (c) density; (d,e) isotropic P- and S-wave anomalies; (g) azimuthal anisotropy and FSE semi-axis (white bars); (h) radial anisotropy and TTI axis (white bars); (f,i) P-P and P-S reflection energy for waves propagating upward.

Deleted: 4



368

369

370

371

372

373

374

375


376

Figure 7 VIZTOMO output for a 3D model of oceanic plate subduction and roll back in spherical coordinates. [The time-dependent flow field has been computed with the software I3MG modified to account for spherical coordinates \(see Faccenda and VanderBeek, 2023\).](#) Fabrics computed with D-REX_M for only upper mantle aggregates. A fossil A-type olivine fabric with the fast axis parallel to plate motion and computed with D-REX_S is initially defined within the oceanic plate volume. The colour scale indicates radial anisotropy in the upper mantle, and the white bars the SKS splitting computed with SKS-SPLIT. The volume in green encloses material with a +2% P-wave anomaly (i.e., the oceanic plate). Note the apparently thicker slab portion around the 410 km depth discontinuity, due to the upwelling of the olivine-spinel phase transition. The model domain extends from 0-1000 km along the radial direction, 85°-115° along longitude, and 70-90° along colatitude.

377

378

Deleted: 5



380 **Figure 8.** VIZTOMO output for an upwelling plume toy model in spherical coordinates. The time-dependent flow field has been computed
 381 with the software IBMG modified to account for spherical coordinates (see Faccenda and VanderBeek, 2023). a) absolute Vs (m/s) and SKS
 382 splitting computed with SKS-SPLIT (white bars); b) radial anisotropy (colour scale) and azimuthal anisotropy at 200 km depth (white bars);
 383 isotropic Vs anomaly; d) P-P reflection energy for a wave propagating upward. The model domain extends from 0-2900 km along the
 384 radial direction, 70°-110° along longitude, and 70-110° along colatitude.
 385

Deleted: 6

380

381


382

383

384

385

386



388 **Figure 9:** VIZTOMO output for the model 3Dspherical_global available in the cookbooks. The time-dependent flow field has been computed
 389 with the software I3MG modified to account for 3D global spherical coordinates using the Yin-Yang grids (Kageyama and Sato, 2004).
 391 Isotropic P-wave anomaly (left) and azimuthal anisotropy at 200 km depth (right). The grey surface encloses material 0.1 dimensionless
 392 units hotter than the average temperature at any depth.


Deleted: 7

Deleted: surrounding

394 2.2.1 VIZVISC

395 VIZVISC processes the D-REX_M output for the visualisation of the aggregate properties such as the viscous anisotropy and
 396 related deformational history (in terms of the FSE) in ParaView. The deviation from isotropy is evaluated by computing the
 397 radial and azimuthal components of viscous anisotropy in a similar way as for the elastic tensor. More in detail, radial viscous
 398 anisotropy is defined as $\xi = N/L$, while azimuthal viscous anisotropy is defined by the magnitude $G = \sqrt{Gc^2 + Gs^2}$ and
 399 azimuth $\phi = \tan^{-1}(Gs, Gc)$, where $N = 1/8(\eta_{11} + \eta_{22}) - 1/4\eta_{12} + 1/2 \eta_{66}$, $L = 1/2 (\eta_{44} + \eta_{55})$, $Gc = 1/2 (\eta_{55} - \eta_{44})$, $Gs = \eta_{45}$. Radial
 400 and azimuthal viscous anisotropy are evaluated in the FSE (and thus inclusions) reference frame, whereby the minor semi axis
 401 is oriented along the vertical direction, and the intermediate and major semi axes are in the horizontal plane. As such, radial
 402 anisotropy is always ≥ 1 . Figure 10 displays some of these fields computed for a 2D steady-state model of mantle convection
 403 with periodic upwellings and downwellings.

Deleted: 8



407 **Figure 10.** VIZVISC output for the model 2Dpolar convection available in the cookbooks. The steady-state 2D incompressible flow field
 408 in cylindrical coordinates is the result of the analytical solution shown in section 9.3 of the software manual and coded in the MATLAB
 409 script cookbooks/2Dpolar_convection/polarcell.m. (a) Radial viscous anisotropy; (b) azimuthal viscous anisotropy; (c) FSE shape coloured
 411 by the length of its major semi-axis upscaled by a factor of 50 km.

Deleted: 8

407
 408
 409
 410
 411
 412

414 2.3 Seismic forward and inverse modelling

415 2.3.1 SKS-SPLIT

416 SKS-SPLIT estimates the SKS splitting at a grid of virtual seismic stations placed at the top of the D-REX_M model as a
417 function of the back-azimuth using the Fortran routines included in FSTRACK (Schulte-Pelkum and Blackmann, 2003; Becker
418 et al., 2006). These routines use the reflectivity method (for details, see Booth and Crampin, 1983; Chapman and Shearer,
419 1989) to compute synthetic seismograms through layered anisotropic media traversed by an incident plane wave (5° for typical
420 SKS arrivals) over a range of frequencies (0-25 Hz). Band-pass filters from 0.1 to 0.3 Hz are then applied to construct synthetic
421 seismograms in the SKS band (3.3 to 10s). Successively, the splitting is determined with the cross-correlation method of
422 Menke and Levin (2003). The routines have been adapted to load the D-REX_M output, stack the elastic tensors and densities
423 in an upper mantle rock column beneath each virtual seismic station (Faccenda and Capitanio, 2013), and run in parallel using
424 MPI. The averaged fast azimuth scaled by the delay time can then be visualised in ParaView as shown in Figures 7 and 8a.

Formatted: Font: (Default) Times New Roman

425 2.3.2 PSI

426 PSI (Platform for Seismic Imaging) is a Julia package for performing tomographic inversion of both real and synthetic seismic
427 datasets. In the context of the ECOMAN software, it's a useful tool for exploring how features in high-resolution geodynamic
428 simulations are mapped into lower resolution seismic tomography models; this is a critical step if one aims to evaluate
429 geodynamic results against existing tomographic images.

430 PSI can be used to forward model P and S travel-times, splitting intensity (Chevrot, 2000), and shear wave splitting parameters
431 (i.e. the fast-polarisation azimuth and the delay time between the fast- and slow-polarised waveforms). Forward modelling of
432 these seismic observables is supported for three different model parameterizations: (1) isotropic P- and S-wavespeeds, (2)
433 hexagonally anisotropic media defined by the 5 Thomsen parameters and the azimuth and elevation of the symmetry axis, and
434 (3) fully anisotropic models defined by the density-normalised 21-component elastic tensor which can be generated from D-
435 REX_M + VIZTOMO (sections 2.1.2 and 2.2.1). Seismic phase velocities are computed following Thomsen (1986) for
436 hexagonally anisotropic models while the Christoffel equations are solved when the elastic tensor parameterization is used.
437 Anisotropic travel-times and splitting intensities for arbitrarily polarised S-waves are computed following VanderBeek et al.
438 (2023) using a long-wavelength approximation in which the accumulated delay time between fast- and slow-polarised qS-
439 waves is less than their dominant period. Lastly, splitting parameters are predicted using the matrix propagation method of
440 Rumpker and Silver (1998). In this initial release of PSI, all predictions are made via integration along ray paths traced through
441 a user-defined 1D reference velocity model using the TauP Toolkit (Crotwell et al. 1998). The 3D model properties are
442 subsequently interpolated to these paths before computing the seismic observables. We anticipate releasing an update to the
443 package that includes both 3D anisotropic ray tracing and finite-frequency kernels that are currently under development.

444 Travel-time and splitting intensity datasets can be inverted individually and jointly either for isotropic (V_p and V_s) or
445 hexagonally anisotropic model parameters. Hexagonal anisotropy is defined by up to five free parameters, the isotropic (1) P-

Deleted: 5

Deleted: 6

448 and (2) S-velocity, (3) anisotropic magnitude, and the (4) azimuth and (5) elevation of the hexagonal symmetry axis. Only a
449 single anisotropic magnitude parameter is required because the strength of P and S anisotropy is strongly correlated (e.g.,
450 Becker et al., 2006). Consequently, the ratios between the three Thomsen parameters (ϵ , δ , γ ; Thomsen 1986) and the inverted
451 anisotropic magnitude parameter must be chosen *a priori* and can be spatially variable. Source and receiver statics for each
452 observation and seismic phase type may also be included as free parameters. The tomographic model is obtained by iteratively
453 solving a system of linearized equations relating the perturbations in the inversion parameters to the data residuals augmented
454 with damping and smoothing constraints. The solution is obtained via the LSQR algorithm (Paige & Saunders, 1982). Full
455 details on the tomographic method can be found in VanderBeek and Faccenda (2021) and VanderBeek et al. (2023). The final
456 tomographic solution is written to a VTK file to be visualised in ParaView. Tomographic inversions can be run from a
457 workstation. For the problem shown in Figure 11, and consisting of 12,320 observations and 415,044 free parameters, solutions
458 can be obtained within ~10 minutes (less for isotropic inversions) using 6 cores and 1.8 GB peak RAM.
459 The PSI inversion methodology has been tested on synthetic models of oceanic plate subduction, intra-oceanic upwelling
460 plume, spreading oceanic ridge, and Central-Eastern Mediterranean subduction (VanderBeek and Faccenda, 2021; Lo Bue et
461 al., 2022, VanderBeek et al., 2023; Faccenda and VanderBeek, 2023), and applied to the isotropic and anisotropic imaging of
462 the Central Mediterranean (Rappisi et al., 2022) and of the Mt. Etna volcanic field (Lo Bue et al., 2024). In Figure 11, we
463 illustrate results obtained from a synthetic inversion of direct teleseismic P- and S-wave relative travel-times computed through
464 the subduction zone shown in Figure 7. Synthetic data were computed using the full 21-component elastic tensor while the
465 inversion was performed for the best-fit hexagonal anisotropic parameters. We considered an array of 770 receivers extending
466 from $\pm 7.5^\circ$ in longitude and $\pm 11.5^\circ$ in latitude (~75 km spacing) that recorded 16 events; 8 at a range 50° and another 8 at 80°
467 from the origin of the subduction zone model and equally distributed in back-azimuth. This example is included in the PSI
468 package.

Deleted: s


Deleted: ~1e4

Deleted: ~1e5

Deleted: 3

Deleted: 9

Deleted: 5



475

476 **Figure 11.** Synthetic tomography results obtained from PSI for the model shown in Fig. 7. (a) Target anisotropic model generated from
 477 VIZTOMO. (b) Recovered anisotropic model obtained by inverting synthetic teleseismic P and S (relative) delay times computed from the
 478 model in (a). In both panels, the isotropic S-wave velocity perturbations are computed with respect to the far-field 1D velocity profile.
 479 Quivers parallel the hexagonal symmetry axis and are scaled and coloured by the anisotropic strength. The top surface shown is located at
 480 200 km depth while the full model extends from 0-1000 km along the radial direction, 85°-115° along longitude, and 0-20° along latitude.

Deleted: 9

481

3 Discussion

482

3.1 Software advantages

483

When compared to other similar software, ECOMAN (i) aims at being a more versatile package suitable for any geodynamic
 484 simulations (2D and 3D; Cartesian and polar coordinate systems; regional and global settings), (ii) takes into account the time-
 485 dependent deformational history of the mantle (which is usually not steady-state, especially close to plate boundaries), (iii)
 486 predicts the strain-induced fabric and elastic tensor of different mantle layers (i.e., not only the upper mantle), (iv) includes
 487 Effective Medium Theories (STILWE, DEM) and a parametrization of the fabric evolution of two-phase composites to predict
 488 elastic and viscous extrinsic anisotropy, (v) generates realistic grid structure distributions of mantle elastic properties to be
 489 used for forward/inverse seismological modelling (e.g., PSI, SPECFEM3D), and (vi) performs synthetic seismic inversions
 490 (e.g., P- and S-wave travel-time tomographies, S-wave splitting intensities) within the computational domain, which facilitates

Deleted: c

493 the comparison with other tomographic models and the estimation of apparent anomalies (artefacts) due to, for example,
494 unaccounted-for elastic anisotropy (Bezada et al., 2016; VanderBeek and Faccenda, 2021; VanderBeek et al., 2023) and/or
495 regularisation.

496 3.2 Software (real or potential) current limitations

497 D-REX_S and D-REX_M compute the strain-induced LPO for the most abundant and highly anisotropic mineral phases, while
498 assuming random orientation for several secondary phases that instead could contribute substantially to the aggregate
499 anisotropic properties. For example, cubic ferropericlasite is relatively abundant (< 20%) and becomes highly anisotropic in the
500 lower half of the lower mantle ($A_{vs} = 30\text{-}50\%$), such that it could dominate seismic anisotropy at these depths (Marquardt et
501 al., 2009). Davemaoite (Ca-perovskite) is also a highly anisotropic mineral with cubic symmetry ($A_{Vs} = 25\text{-}15\%$, Kawai and
502 Tsuchiya, 2015) but its model abundance is quite low (< 10%). Recently, micromechanical simulations of strain-induced LPO
503 in aggregates with a pyrolite mantle composition have shown that the bulk aggregate seismic anisotropy is controlled by
504 bridgmanite and post-perovskite, while the cubic secondary phases appear to only slightly reduce the amplitude of anisotropy
505 (Chandler et al., 2021). The latter effect can thus be well approximated by randomising the orientation of the cubic phases as
506 assumed here.

507 A main limitation of ECOMAN is that it does not include yet the modelling of viscous anisotropy due to the intrinsic
508 mechanical anisotropy of crystals. Tommasi et al. (2009), have used VPSC simulations to show that olivine CPO in the
509 lithosphere can control the reactivation of fossil faults misoriented with respect to the stress field. Kiraly et al (2020), employed
510 the modified director method (MDM) and estimated up to 1 order of magnitude of intrinsic viscous anisotropy in olivine
511 aggregates, which can control the kinematics and dynamics of tectonic plates. Both the VPSC and MDM approaches are
512 computationally expensive and appear to be prohibitive for the number of mantle aggregates required to discretize the mantle
513 domain of large-scale 3D simulations. An alternative approach which minimises the computational time is therefore desired.

514 Fraters and Billen (2021) have released a version of the D-REX model that is embedded into the ASPECT geodynamic
515 modelling software (Banghart et al., 2020) and that takes into account changes in olivine fabric as a function of the water
516 content and deviatoric stress (Karato et al., 2008). Although at present D-REX_M only allows the definition of a single olivine
517 fabric per run, changes in crystal fabrics can be easily implemented in the future by loading the water content and deviatoric
518 stress field from the geodynamic models and by parametrizing the boundaries of the olivine fabric domains.

519 Tape and Tape (2024) recently reformulated the elastic decomposition method proposed by Browaeys and Chevrot (2004) that
520 they argued is not entirely accurate. Although this will only affect the visualization of the tensor components as performed by
521 D-REX_S and VIZTOMO, it is planned to replace the existing elastic tensor decomposition methodology with the one
522 proposed by Tape and Tape (2024).

523 A minor limitation is that the different code modules are based on different programming languages (Fortran, Julia), libraries
524 (OpenMP, MPI, HDF5) and software (MATLAB, ParaView), whose installation on local devices might discourage potential

525 users. There are a few main reasons for this. First of all, the original D-REX software was written in Fortran, thus its
526 modifications into D-REX_S and D-REX_M software was more straightforward by maintaining the same programming
527 language. Fortran has high performance standards on HPC clusters, often if not always higher than other interpreted
528 programming languages (e.g., MATLAB or Python). Given the large-scale computational power needed for the D-REX_M
529 simulations, especially those with 100,000s or millions of crystal aggregates, and considering that the required compilers and
530 libraries are routinely installed in most (if not all) HPC clusters, the usage of the ECOMAN's Fortran-based applications in
531 high-performance environments is warranted. In contrast, the visualisation of the Fortran-based applications' output in
532 MATLAB and ParaView can be performed on local devices. In particular, the MTEX software is a MATLAB toolbox for
533 visualisation of the LPO and elastic tensor pole figures which should be downloaded ([https://mtex-
534 toolbox.github.io/download](https://mtex-toolbox.github.io/download)) and installed along with MATLAB only when using D_REX_S. Seismic forward/inverse
535 modelling with PSI can be performed on either a HPC cluster or local devices upon installation of the Julia package. Julia was
536 chosen because it is an open-source and high-level language that offers a number of performance benefits over other popular
537 scientific computing languages such as Python or Matlab. It is important to stress that for any of these applications the user
538 only needs to modify input text files, and thus no particular prerequisite or computational skill is required.
539 Finally, running D-REX_M requires allocating ~ 160 GB of memory per million of crystal aggregates with 1000 x 2 crystals
540 each. This potential problem can be addressed by distributing the computational and memory load over several nodes, which
541 is possible thanks to the hybrid MPI and OpenMP parallelization scheme.

Deleted: high-level

Deleted: .

Deleted: '

Deleted: CPUs

542 4 Conclusions and outlook

543 ECOMAN is an open-source software package for estimating strain/stress-induced fabrics in mantle aggregates, their
544 mechanical properties, and how mechanical anisotropy affects the geodynamic evolution and seismic imaging of the Earth's
545 interior. Programs included in ECOMAN are portable across different HPC and local device systems (provided the Julia
546 package and Fortran compilers are available), and are applicable to any 2D-3D geodynamic simulation. Computationally
547 expensive programs such as D-REX_M are parallelized, offering a nearly perfect scaling with an increasing number of cores.
548 As a result, the strain-induced fabrics of millions of mantle aggregates can now be estimated with a reasonable amount of time
549 and computational resources.

550 As ongoing developments, we are seeking to include in ECOMAN micromechanical modelling methods that are capable of
551 estimating the strain-induced LPO and/or the intrinsic viscous anisotropy at computational speeds that are orders of magnitude
552 faster than current ones. For instance, Ribe et al. (2019), have proposed an analytical finite-strain parameterization for texture
553 evolution in deforming olivine polycrystals that is $\approx 10^7$ times faster than full homogenization approaches such as the second-
554 order self-consistent model. When implemented in ECOMAN, preliminary tests indicate that this new method outperforms D-
555 REX by 1-2 orders of magnitude (Ribe et al., 2023).

560 In addition, in the near future the PSI software will be updated to include trans-dimensional Bayesian Monte Carlo sampling
561 methods that, in contrast to deterministic approaches, address the consequences of under-determination in seismic imaging [by](#)
562 [constraining the uncertainty](#) (Del Piccolo et al., 2023). Isotropic and anisotropic seismic imaging with PSI is currently feasible
563 using body wave information such as travel-times and S-wave splitting intensity. However, when local deep seismicity is
564 absent, as is the case in warm subduction zones or at spreading ridges and intraplate settings, the retrieved isotropic and
565 anisotropic mantle structures are only partially recovered and often affected by smearing (e.g., Faccenda and VanderBeek,
566 2023). Consequently, we are planning to complement body wave information with surface wave data to improve the seismic
567 ray coverage and the resolving power of tomographic models. Lastly, to improve the prediction of seismic observables, 3D
568 anisotropic ray tracing and finite-frequency kernels are planned for a future release.

Deleted: s

569 **Code availability**

570 [The version of ECOMAN described in this article is 2.0 and is freely available at github \(<https://github.com/ecoman-geos>\) and](#)
571 [zenodo \(ECOMAN2.0-geodynamics: <https://doi.org/10.5281/zenodo.10599735>; ECOMAN2.0-seismology.SKS-split: <https://doi.org/10.5281/zenodo.11173260>; ECOMAN2.0-seismology.PSI_D: <https://doi.org/10.5281/zenodo.11186805>\)](#).

Deleted: ECOMAN is freely available at <https://github.com/ecoman-geos>.

573 **Data availability**

574 The software package contains the input files to generate the synthetic models and datasets discussed in this manuscript. No
575 data has been produced for this work.

576 **Author contributions**

577 MF and BPV developed the software package, with important contributions from AdM, JY and FR. MF wrote the manuscript
578 draft. MF acquired the funding supporting the research activities. All authors have contributed to the discussion and manuscript
579 editing.

Deleted: and

580 **Competing interests**

581 The authors declare that they have no conflict of interest.

586 **Acknowledgements**

587 MF is indebted to F.A. Capitanio for supporting the beginning of this journey, T.W. Becker for providing routines to compute
588 shear wave splitting, E. Kaminski for fruitful discussions on strain-induced LPO modelling, and to A.M. Ferreira for the
589 continuous and stimulating feedback. [Two anonymous reviewers helped in improving the article.](#)

590 **Financial statement**

591 This study is supported by the ERC StG 758199 NEWTON.

592 **References**

- 593 [Abt, D. L., Fischer, K. Resolving three-dimensional anisotropic structure with shear wave splitting tomography. *Geophys. J. Int.* 173, 859-886, 2008.](#)
594
595 Ahrens, J., Geveci, B., and Law, C.: ParaView: An End-User Tool for Large Data Visualization, in: Visualization Handbook,
596 edited by: Hansen, C.D. and Johnson, C.R., Elsevier, 717-731, <https://doi.org/10.1016/B978-012387582-2/50038-1>,
597 2005.
598 Almqvist, B.S.G. and Mainprice, D. Seismic properties and anisotropy of the continental crust: Predictions based on mineral
599 texture and rock microstructure. *Rev. Geophys.*, 55, 367-433, <https://doi.org/10.1002/2016RG000552>, 2017.
600 [Bangerth, W., Dannberg, J., Gassmoeller, R., & Heister, T. \(2020\). *Aspect v2.2.0. Zenodo.*](#)
601 <https://doi.org/10.5281/zenodo.3924604>.
602 Becker, T.W., Chevrot, S., Schulte-Pelkum, V. and Blackman, D.K.: Statistical properties of seismic anisotropy predicted by
603 upper mantle geodynamic models. *J. Geophys. Res.*, 111, B08309 <https://doi.org/10.1029/2005JB004095>, 2006.
604 Bezada, M., Faccenda, M. and Toomey, D. R: Representing anisotropic subduction zones with isotropic velocity models: A
605 characterization of the problem and some steps on a possible path forward. *Geochem. Geophys. Geosyst.*,
606 <https://doi.org/10.1002/2016GC006507>, 2016.
607 [Booth, D. C., and S. Crampin. The anisotropic reflectivity technique: Theory, *Geophys. J. R. Astron. Soc.*, 72, 31-45, 1983.](#)
608 Browaeys, J.T. and Chevrot, S.: Decomposition of the elastic tensor and geophysical applications. *Geophys. J. Int.*, 159, 2,
609 667-678, 2004.
610 Bunge, H.-P., Hagelberg, C.R. and Travis, B.J.: Mantle circulation models with variational data assimilation: inferring
611 past mantle flow and structure from plate motion histories and seismic tomography. *Geophys. J. Int.*, 152, 280-301,
612 2003.
613 Chandler, B.C., Chen, L.-W., Li, M., Romanowicz, B. and Wenk, H.-R.: Seismic anisotropy, dominant slip systems and phase
614 transitions in the lowermost mantle. *Geophys. J. Int.*, 227, 1665-1681, <https://doi.org/10.1093/gji/ggab278>, 2021.

Formatted: Space Before: 0 pt, After: 0 pt, Line spacing: 1,5 lines

615 Chang, S.-J., Ferreira, A. M. G., Ritsema, J., van Heijst, H. J. and Woodhouse, J. H. Joint inversion for global isotropic and
616 radially anisotropic mantle structure including crustal thickness perturbations. *Journal of Geophysical Research: Solid*
617 *Earth*, 120, 4278–4300. <https://doi.org/10.1002/2014JB011824>, 2015

618 Chang, S.-J., Ferreira, A.M.G. and Faccenda, M. Upper- and mid-mantle interaction between the Samoan plume and the Tonga-
619 Kermadec slabs. *Nat. Commun.*, 7:10799, <https://doi.org/10.1038/ncomms10799>, 2016.

620 [Chapman, C. H., Shearer, P.M. Ray tracing in azimuthally anisotropic media: II. Quasishear wave coupling. *Geophys. J. Int.*,
621 *96*, 65–83, 1989.](#)

622 Chust, T.C., Steinle-Neumann, G., Dolejs, D., Schubert, B.S. and Bunge, H.P.: MMA- EoS: a computational framework for
623 mineralogical thermodynamics. *J. Geophys. Res.* 122, 9881–9920, 2017.

624 Cramer, F., and Tackley, P. J. Spontaneous development of arcuate single-sided subduction in global 3-D mantle convection
625 models with a free surface. *J. Geophys. Res. Solid Earth*, 119, 921–9942, <https://doi.org/10.1002/2014JB010939>, 2014.

626 Davies, D.R., Goes, S., Davies, J.H., Schubert, B.S.A., Bunge, H.-P. and Ritsema, J. Reconciling dynamic and seismic
627 models of Earth's lower mantle: The dominant role of thermal heterogeneity. *Earth Planet. Sci. Lett.*, 353–354, 253-
628 269, 2012.

629 de Montserrat, A., Faccenda, M., and Pennacchioni, G.: Extrinsic anisotropy of two-phase Newtonian aggregates: Fabric
630 characterization and parameterization. *J. Geophys. Res.: Solid Earth*, 126, e2021JB022232.
631 <https://doi.org/10.1029/2021JB022232>, 2021.

632 [Debayle, E., Kennet, B., Priestley, K. Global azimuthal seismic anisotropy and the unique plate-motion deformation of
633 Australia. *Nature* 433, 509-512, 2005.](#)

634 Debayle, E., Bodin, T., Durand, S. and Ricard, Y.. Seismic evidence for partial melt below tectonic plates. *Nature* 586, 555–
635 559. <https://doi.org/10.1038/s41586-020-2809-4>, 2020.

636 Del Piccolo, G., VanderBeek, B., Faccenda, M., Morelli, A., and Byrnes, J.: Reversible-Jump, Markov-Chain Monte Carlo
637 seismic tomographic inversion for anisotropic structure in subduction zones (No. EGU23-9927). *Copernicus*
638 *Meetings*, <https://doi.org/10.5194/egusphere-egu23-9927>, 2023.

639 Deuss, A. Heterogeneity and anisotropy of Earth's inner core. *Annu. Rev. Earth. Planet. Sci.* 42, 103–126,
640 <https://doi.org/10.1146/annurev-earth-060313-054658>, 2014.

641 Faccenda M. and Capitanio, F. A.: Development of mantle seismic anisotropy during subduction-induced 3D flow. *Geophys.*
642 *Res. Lett.*, 39, <https://doi.org/10.1029/2012GL051988>, 2012.

643 Faccenda M. and Capitanio, F. A.: Seismic anisotropy around subduction zones: insights from three-dimensional modeling of
644 upper mantle deformation and SKS splitting calculations. *Geochem. Geophys. Geosyst.*,
645 <https://doi.org/10.1002/ggge.20055>, 2013.

646 Faccenda, M.: Mid mantle seismic anisotropy around subduction zones, *Phys. Earth Planet. Int.*, 227, 1–19., 2014

647 Faccenda, M., Ferreira, A.M.G., Tisato, N., Lithgow-Bertelloni, C., Stixrude, L. and Pennacchioni, G.: Extrinsic anisotropy in
648 a compositionally heterogeneous mantle. *J. Geophys. Res.* 124, <https://doi.org/10.1029/2018JB016482>, 2019.

649 Ferreira, A.M.G., Faccenda, M., Sturgeon, W., Chang, S.-J. and Scharndong, L.: Ubiquitous lower-mantle anisotropy beneath
650 subduction zone. *Nature Geo.* 12, 301-306, <https://doi.org/10.1038/s41561-019-0325-7>, 2019.

651 Faccenda, M. and VanDerBeek, B.P.: On constraining 3D seismic anisotropy in subduction, mid-ocean-ridge, and plume
652 environments with teleseismic body wave data. *J. Geodyn.*, 158, <https://doi.org/10.1016/j.jog.2023.102003>, 2023.

653 Fraters, M. R. T. and Billen, M. I.: On the implementation and usability of crystal preferred orientation evolution in
654 geodynamic modeling. *Geochem., Geophys., Geosys.*, 22, e2021GC009846. <https://doi.org/10.1029/2021GC009846>,
655 2021.

656 French, S., and Romanowicz, B. Broad plumes rooted at the base of the Earth's mantle beneath major hotspots. *Nature* 525,
657 95–99, <https://doi.org/10.1038/nature14876>, 2015.

658 Jadamec, M. and Billen, M.: Reconciling surface plate motions with rapid three-dimensional mantle flow around a slab edge.
659 *Nature* 465, 338–341, <https://doi.org/10.1038/nature09053>, 2010.

660 Kageyama, A. and Sato, T.: “Yin-Yang grid”: An overset grid in spherical geometry. *Geochem Geophys. Geosys.*, 5, 9,
661 <https://doi.org/10.1029/2004GC000734>, 2004.

662 Kaminski, E., Ribe, N. M. and Browaeys, J. T.: D-Rex, a program for calculation of seismic anisotropy due to crystal
663 lattice preferred orientation in the convective upper mantle. *Geophys. J. Int.*, 158(2), 744–752.
664 <https://doi.org/10.1111/j.1365-246x.2004.02308.x>, 2004.

665 Karato, S.-I.: Seismic anisotropy in the deep mantle, boundary layers and the geometry of mantle convection. *Pure and Applied*
666 *Geophysics*, 151(2-4), 565–587. <https://doi.org/10.1007/s000240050130>, 1998.

667 [Karato, S.-i., Jung, H., Katayama, I., and Skemer, P. Geodynamic significance of seismic anisotropy of the upper mantle:
668 New insights from laboratory studies. *Annual Review of Earth and Planetary Sciences*, 36\(1\), 59–95.
669 <https://doi.org/10.1146/annurev.earth.36.031207.124120>, 2008.](#)

670 Kawai, K. and Tsuchiya, T.: Small shear modulus of cubic CaSiO₃ perovskite. *Geophys. Res. Lett.*, 42, 8,
671 <https://doi.org/10.1002/2015GL063446>, 2015.

672 Kendall, J.-M.: Seismic anisotropy in the boundary layers of the mantle. In *Earth's deep interior: Mineral physics and*
673 *tomography from the atomic to the global scale*, Geophysical Monograph Series, 117, 133–159, Washington, DC:
674 American Geophysical Union, 2000.

675 Kim, E., Kim, Y. and Mainprice, D.: AnisEulerSC: A MATLAB program combined with MTEX for modeling the anisotropic
676 seismic properties of a polycrystalline aggregate with microcracks using self-consistent approximation. *Comp. &*
677 *Geosci.*, 145, 104589, <https://doi.org/10.1016/j.cageo.2020.104589>, 2020.

678 Király, Á., Conrad, C. P. and Hansen, L. N.: Evolving viscous anisotropy in the upper mantle and its geodynamic implications.
679 *Geochem. Geophys., Geosys.*, 21, e2020GC009159, <https://doi.org/10.1029/2020GC009159>, 2020.

680 Kocher, T. Schmalholz, S.M. and Mancktelow, N.S. Impact of mechanical anisotropy and power-law rheology on single layer
681 folding. *Tectonophys.* 421, 1-2, 71-87, 2006.

Formatted: Space Before: 0 pt, After: 0 pt, Line spacing: 1,5 lines

682 Komatitsch, D. and Tromp, J.: Introduction to the spectral element method for three-dimensional seismic wave propagation,
683 Geophys. J. Int., 139(3), 806–822, 1999.

684 [Halter, W., Macherel, E., Duretz, T., and Schmalholz, S. M. Numerical modelling of strain localization by anisotropy evolution
685 during 2D viscous simple shearing. EGU General Assembly 2022, Vienna, Austria, 23–27 May 2022. EGU22-11438.
686 <https://doi.org/10.5194/egusphere-egu22-11438>, 2022.](#)

687 [Hansen, L. N., Zhao, Y.-H., Zimmerman, M. E., Kohlstedt, D.L., Protracted fabric evolution in olivine: Implications for the
688 relationship among strain, crystallographic fabric, and seismic anisotropy. Earth Planet. Sci. Lett. 387, 157-168,
689 <https://doi.org/10.1016/j.epsl.2013.11.009>, 2014.](#)

690 Hedjazian, N., Garel, F., Rhodri Davies, D. and Kaminski: E. Age-independent seismic anisotropy under oceanic plates
691 explained by strain history in the asthenosphere. Earth Planet. Sci. Lett. 460, 135-142, 2017.

692 Hu, J., Faccenda, M. and Liu, L.. Subduction-controlled mantle flow and seismic anisotropy in South America. Earth and
693 Planet. Sci. Lett. 470, 13-24, <https://doi.org/10.1016/j.epsl.2017.04.027>, 2017

694 Li, Z.-H., Di Leo, J.F. and Ribe, N. M. Subduction-induced man-tle flow, finite strain, and seismic anisotropy: Numerical
695 modeling, J. Geophys. Res. Solid Earth, 119, 5052–5076, <https://doi.org/10.1002/2014JB010996>, 2014.

696 Lo Bue, R., Faccenda, M. and Yang, J.: The role of Adria Plate Lithospheric Structures on the Recent Dynamics of the Central
697 Mediterranean Region. J. Geophys. Res., 126, e2021JB022377. <https://doi.org/10.1029/2021JB022377>, 2021.

698 Lo Bue, R., Rappisi, F., VanderBeek, B.P. and Faccenda, M.: Tomographic Image Interpretation and Central-Western
699 Mediterranean-Like Upper Mantle Dynamics From Coupled Seismological and Geodynamic Modeling Approach.
700 Front. Earth Sci., 10:884100, <https://doi.org/10.3389/feart.2022.884100>, 2022.

701 [Lo Bue, R., Rappisi, F., Firetto Carlino, M., Giampiccolo, E., Cocina, O., Vanderbeek, B. P. and Faccenda, M. Crustal
702 structure of Etna volcano \(Italy\) from P-wave anisotropic tomography. Geophys. Res. Lett. 51, e2024GL108733,
703 <https://doi.org/10.1029/2024GL108733>, 2024.](#)

704 Long, M.D. and Becker, T.W.: Mantle dynamics and seismic anisotropy. Earth Planet. Sci. Lett., 297, 341-354, 2010.

705 [Long, M.D., Maarten V., Van Der Hilst, R.D. Wave-equation shear wave splitting tomography. Geophys. J. Int. 172 \(1\), 311-
706 330, <https://doi.org/10.1111/j.1365-246X.2007.03632.x>, 2008.](#)

707 Maguire, R., Ritsema, J., Bonnin, M., van Keken, P. E. and Goes, S.: Evaluating the resolution of deep mantle plumes in
708 teleseismic traveltome tomography. J. Geophys. Res., 123, 384–400. <https://doi.org/10.1002/2017JB014730>, 2018.

709 Mainprice, D.: Modelling the anisotropic seismic properties of partially molten rocks found at mid-ocean ridges. Tectonophys.,
710 279, 1-4, 161-179, 1997.

711 Mainprice, D., Hielscher, R. and Schaeben, H.: in Prior, D.J., Rutter, E.H., Tatham, D. J. Calculating anisotropic physical
712 properties from texture data using the MTEX open source package. (eds) Deformation Mechanisms, Rheology and
713 Tectonics: Microstructures, Mechanics and Anisotropy. Geological Society, London, Special Publications, 360, 175-
714 192, 2011.

Formatted: Font: (Default) Times New Roman, 10 pt, English (UK)

Formatted: Indent: Left: 0 cm, Hanging: 1,27 cm, Line spacing: 1,5 lines

Formatted: English (UK)

Formatted: English (US)

Formatted: Font: (Default) Times New Roman, 10 pt

Formatted: Font: (Default) Times New Roman, 10 pt, English (US)

Deleted: Lo Bue, R., Faccenda, M., Cocina, O., Rappisi, F. and Vanderbeek, B.P.: Joint Active and Passive P-wave Tomography reveals Mt. Etna's Seismic Anisotropy (No EGU23-8563). Copernicus Meetings, <https://doi.org/10.5194/egusphere-egu23-8563>, 2023.*

720 Marquardt, H., Speziale, S., Reichmann, H. J., Frost, D. J., Schilling, F. R. and Garnero, E.J.: Elastic Shear Anisotropy of
721 Ferropericlasite in Earth's Lower Mantle. *Science*, 324, 5924, 224-226, <https://doi.org/10.1126/science.1169365>, 2009

722 [Menke, W., Levin, V. The cross-convolution method for interpreting SKS splitting observations, with application to one and](#)
723 [two layer anisotropic earth models, *Geophys. J. Int.*, 154, 379–392, 2003.](#)

724 [Mondal, P., Long, M.D. A model space search approach to finite-frequency SKS splitting intensity tomography in a reduced](#)
725 [parameter space. *Geophys. J. Int.* 217, 238-256, <https://doi.org/10.1093/gji/ggz016>, 2019.](#)

726 Müller, R. D., Flament, N., Cannon, J., Tetley, M. G., Williams, S. E., Cao, X., Bodur, Ö. F., Zahirovic, S., and Merdith, A.:
727 A tectonic-rules-based mantle reference frame since 1 billion years ago – implications for supercontinent cycles and
728 plate–mantle system evolution, *Solid Earth*, 13, 1127–1159, <https://doi.org/10.5194/se-13-1127-2022>, 2022.

729 [Munzarova, H., Plomerova, J., Kissling, E.. Novel anisotropic teleseismic body- wave tomography code AniTomo to](#)
730 [illuminate heterogeneous anisotropic upper mantle: Part I–theory and inversion tuning with realistic synthetic data.](#)
731 [*Geophys. J. Int.* 215 \(1\), 524–545, 2018.](#)

732 Oganov, A.R. and Ono, S.: Theoretical and experimental evidence for a post-perovskite phase of MgSiO₃ in Earth's D'' layer.
733 *Nature*, 430, 445–448, 2004.

734 Rappisi, F., VanderBeek, B., Faccenda, M., Morelli, A. and Molinari, I: Slab geometry and upper mantle flow patterns in the
735 Central Mediterranean from 3D anisotropic P-wave tomography. *J. Geophys. Res.* 127, e2021JB023488,
736 <https://doi.org/10.1029/2021JB023488>, 2022.

737 Ribe, N.M., Hielscher, R.H. and Castelnau, O.: An analytical finite-strain parameterization for texture evolution in deforming
738 olivine polycrystals. *Geophys. J. Int.*, 216, 486–514, <https://doi.org/10.1093/gji/gyy442>, 2019.

739 Ribe, N.M., Faccenda, M., Hielscher, R.H.: SBFTEX: An Analytical Parameterization for Finite Strain-Induced Upper-Mantle
740 Anisotropy. AGU Fall Meeting 2023, DI42A-08, 2023.

741 Rudolph, M.L., Lekić, V. and Lithgow-Bertelloni, C.: Viscosity jump in Earth's mid-mantle. *Science*, 350, 1349-
742 1352, <https://doi.org/10.1126/science.aad1929>, 2015.

743 Schaeffer, A.J., Lebedev, S. and Becker, T.W. Azimuthal seismic anisotropy in the Earth's upper mantle and the thickness of
744 tectonic plates. *Geophys. J. Int.*, 207, 2, 901–933, <https://doi.org/10.1093/gji/ggw309>, 2016.

745 Schubert, B.S.A., Zaroli, C. and Nolet, G.: Synthetic seismograms for a synthetic Earth: long-period P- and S-wave traveltime
746 variations can be explained by temperature alone, *Geophys. J. Int.*, 188(3), 1393–1412, doi:10.1111/j.1365-
747 246X.2011.05333.x, 2012.

748 Schulte-Pelkum, V. and Blackman, D.K.: A synthesis of seismic P and S anisotropy, *J. Geophys. Int.*, 154, 1, 166–178, 2003.

749 Shea Jr, W. T. and Kronenberg, A. K. Strength and anisotropy of foliated rocks with varied mica contents. *Journal of Structural*
750 *Geology*, 15(9-10), 1097-1121, [https://doi.org/10.1016/0191-8141\(93\)90158-7](https://doi.org/10.1016/0191-8141(93)90158-7), 1993.

751 Steinberger, B. and Calderwood, A.: Models of large-scale viscous flow in the Earth's mantle with constraints from mineral
752 physics and surface observations. *Geophys. J. Int.*, 167, 1461-1481, [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-246X.2006.03131.x)
753 [246X.2006.03131.x](#), 2006

754 Styles, E., Davies, D. R., and Goes, S.: Mapping spherical seismic into physical structure: Biases from 3-D phase-transition
755 and thermal boundary-layer heterogeneity. *Geophys. J. Int.*, 184, 1371–1378. <https://doi.org/10.1111/j.1365-246X.2010.04914.x>, 2011.

756

757 Sturgeon W., Ferreira, A.M.G., Faccenda, M., Chang, S.-J. and Schardong, L.: On the origin of radial anisotropy near
758 subducted slabs in the midmantle. *Geochem. Geophys. Geosys.* 20 (11), 5105-5125,
759 <https://doi.org/10.1029/2019GC008462>, 2019.

760 [Tape, W., and Tape, C. A reformulation of the Browaeys and Chevrot decomposition of elastic maps. *J. Elasticity.*
761 <https://doi.org/10.1007/s10659-024-10056-x>, 2024.](https://doi.org/10.1007/s10659-024-10056-x)

762 [Tasaka, M., Zimmerman, M. E., Kohlstedt, D. L., Stünitz, H. and Heilbronner, R. Rheological weakening of
763 olivine + orthopyroxene aggregates due to phase mixing: Part 2. Microstructural development. *Journal Geophysical*
764 *Research: Solid Earth*, 122, 7597–7612. <https://doi.org/10.1002/2017JB014311>, 2017.](https://doi.org/10.1002/2017JB014311)

765 Tommasi, A., Knoll, M., Vauchez, A., Signorelli, J. W., Thoraval, C. and Logé, R.: Structural reactivation in plate tectonics
766 controlled by olivine crystal anisotropy. *Nature Geosci.*, 2, 423–427, <https://doi.org/10.1038/ngeo528>, 2009.

767 VanderBeek, B.P., Faccenda, M.: Imaging upper mantle anisotropy with teleseismic P-wave delays: insights from tomographic
768 reconstructions of subduction simulations. *Geophys. J. Int.* 225, 2097–2119, 2021.

769 VanderBeek, B.P., Lo Bue, R., Rappisi, F. and Faccenda, M.: Imaging upper mantle anisotropy with travel-time and splitting
770 intensity observations from teleseismic shear waves: Insights from tomographic reconstructions of subduction
771 simulations. <https://doi.org/10.1093/gji/ggad389>, 2023.

772 [Warren, J. M., Hirth, G. and Kelemen, P.B. Evolution of olivine preferred orientation during simple shear in the mantle. *Earth*
773 *Planet. Sci. Lett.* 272, 501-512, <https://doi.org/10.1016/j.epsl.2008.03.063>, 2008.](https://doi.org/10.1016/j.epsl.2008.03.063)

774 [Wookey, J. Direct probabilistic inversion of shear wave data for seismic anisotropy. *Geophys. J. Int.* 189, 1025-1037,
775 <https://doi.org/10.1111/j.1365-246X.2012.05405.x>, 2012.](https://doi.org/10.1111/j.1365-246X.2012.05405.x)

776 Zhou, Q., Hu, J., Liu, L., Chaparro, T., Stegman, D. R. and Faccenda, M.: Western U.S. seismic anisotropy revealing complex
777 mantle dynamics. *Earth and Planet. Sci. Lett.* 500, 156-167, <https://doi.org/10.1016/j.epsl.2018.08.015>, 2018.

778

779

780

781

782

783

784

785

786

787

Deleted: Halter, W., Macherel, E., Duretz, T., and Schmalholz, S. M. Numerical modelling of strain localization by anisotropy evolution during 2D viscous simple shearing, EGU General Assembly 2022, Vienna, Austria, 23–27 May 2022, EGU22-11438, <https://doi.org/10.5194/egusphere-egu22-11438>, 2022.

794 **Appendix A. D-REX model vs analytical solution**

795 We compare the numerical solution of the D-REX model included in D-REX_S and D-REX_M with the analytical solution
 796 provided by Fraters and Billen (2021, Supplementary Information) for a test with an upper mantle aggregate (O1: Ens =
 797 50:50) composed by 5 crystals for each phase, $M^* = 50$, $\chi = 0.3$, and $nCRSS = (1, 2, 3, \infty)$ for the four olivine slip systems
 798 $[100](010)$, $[100](001)$, $[001](010)$, $[001](100)$ (Table A1). For the first crystal, the time derivative of the cosine direction
 799 matrix (CDM) $\frac{\delta a_{ij}^v}{\delta t} = 0$ and the vorticity matrix $\omega_{ij}^v = \varepsilon_{jki} \omega_k^v = 0$, while for the other 4 crystals:

$$\frac{\delta a_{ij}^v}{\delta t} = \begin{bmatrix} -0.0040153 & -0.0010038 & 0.0020076 \\ -0.0085325 & -0.0010038 & 0.0065249 \\ -0.0130497 & -0.0010038 & 0.0110421 \end{bmatrix} \quad (A1)$$

803 with a maximum error of $3.1 \cdot 10^{-3}$ % which we attribute to truncation errors, and:

$$\omega_{ij}^v = \begin{bmatrix} 0 & 0.005019 & 0.010038 \\ -0.005019 & 0 & 0.005019 \\ -0.010038 & -0.005019 & 0 \end{bmatrix} \quad (A2)$$

807 The changes in volume fractions are $\frac{\delta f_m^m}{\delta t} = 0.630116863029$ for grain 1, and $\frac{\delta f_m^m}{\delta t} = -0.157529215757$ for the other 4
 808 crystals, resulting in a misfit of $6.5244 \cdot 10^{-4}$ % relative to the analytical solution.

809 We provide the D-REX_S_test program for reproducing this simple test in the Supplementary Material.

812 Table A1 – Available slip systems of abundant anisotropic mantle phases present in D-REX_S and D-REX_M.

Slip system [uvw](hkl)	Olivine	Enstatite	Wadsleyite	Bridmanite	Post-Perovskite
$[100](010)$	*		*	*	*
$[100](001)$	*		*	*	*
$[100](011)$			*		
$[100](021)$			*		
$[010](100)$				*	*
$[010](001)$				*	*
$[001](100)$	*	*		*	*
$[001](010)$	*		*	*	*
$[001](110)$				*	*
$\langle 110 \rangle (001)$				*	

- Formatted ... [1]
- Formatted ... [2]
- Formatted ... [3]
- Formatted ... [4]
- Formatted ... [5]
- Formatted ... [6]
- Formatted ... [7]
- Formatted ... [8]
- Formatted ... [9]
- Formatted ... [10]
- Formatted ... [11]
- Formatted ... [12]
- Formatted ... [13]
- Formatted ... [14]
- Formatted ... [15]
- Formatted ... [16]
- Formatted ... [17]
- Formatted ... [18]
- Formatted ... [19]
- Formatted ... [20]
- Formatted ... [21]
- Formatted ... [22]
- Formatted ... [23]
- Formatted ... [24]
- Formatted ... [25]
- Formatted ... [26]
- Formatted ... [27]
- Formatted ... [28]
- Formatted ... [29]
- Formatted ... [30]
- Formatted ... [31]
- Formatted ... [32]
- Formatted ... [33]
- Formatted ... [34]
- Formatted ... [35]
- Formatted ... [36]
- Formatted ... [37]
- Formatted ... [38]
- Formatted ... [39]
- Formatted ... [40]
- Formatted ... [41]

$\langle 110 \rangle \{110\}$				*	*	813
$\frac{1}{2} \langle 111 \rangle \{101\}$			*			814
						815

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Formatted: Font: 10 pt

Appendix B. Aggregate elastic stiffness tensor

At any P-T condition, the single crystal elastic moduli C_{ij}^m are (in Voigt notation):

$$C_{ij}^m(P, T) = C_{ij}^m(P_0, T_0) + \partial C_{ij}^m / \partial P \cdot \Delta P + 0.5 \cdot \partial^2 C_{ij}^m / \partial P^2 \cdot \Delta P^2 + \partial C_{ij}^m / \partial T \cdot \Delta T + \partial^2 C_{ij}^m / \partial P \partial T \cdot \Delta P \Delta T \quad (B1)$$

where the partial derivatives are listed in Table A2 and ΔP and ΔT are deviations from the room conditions ($P_0 = 10^{-4}$ GPa, $T_0 = 298$ K; Mainprice, 2007). The compliance tensor is the inverse of the stiffness tensor, i.e., $S^m = C^m^{-1}$. These tensors can be rotated as:

$$C_{ijkl}^v = a_{ip} a_{jq} a_{kr} a_{ls} C_{pqrs}^m \quad (B2a)$$

$$S_{ijkl}^v = a_{ip} a_{jq} a_{kr} a_{ls} S_{pqrs}^m \quad (B2b)$$

where \mathbf{a} is the CDM defining the crystal orientation with respect to the external reference frame. The stiffness and compliance tensors of the crystal aggregate (i.e., rock) are then obtained by arithmetic averaging:

$$\mathbf{C}^r = \sum_{m=1}^M X_m \sum_{v=1}^N X_v \mathbf{C}^v \quad (B3a)$$

$$\mathbf{S}^r = \sum_{m=1}^M X_m \sum_{v=1}^N X_v \mathbf{S}^v \quad (B3b)$$

where $M = 2$ is the number of minerals in the two-phase mantle aggregates modelled in D-REX_M and D-REX_S, X_m their relative abundance, N the number of crystals of each phase, X_v the normalized crystal volume fraction ($\sum_{v=1}^N X_v = 1$), \mathbf{C}^r and $(\mathbf{S}^r)^{-1}$ correspond to the Voigt and Reuss average stiffness tensor of the rock, i.e., \mathbf{C}_{Voigt}^r and \mathbf{C}_{Reuss}^r . More in general, the average stiffness tensor resulting from the linear combination of the two end-members is found as:

$$\mathbf{C}^r = X_{Voigt} \mathbf{C}_{Voigt}^r + (1 - X_{Voigt}) \mathbf{C}_{Reuss}^r \quad (B4)$$

where X_{Voigt} is a free parameter defining the fraction of \mathbf{C}_{Voigt}^r relative to \mathbf{C}_{Reuss}^r . It is clear that the Hill average stiffness tensor \mathbf{C}_{Hill}^r is obtained when $X_{Voigt} = 0.5$.

The isotropic component of the average stiffness tensor \mathbf{C}_{ISO}^r can be assembled as $C_{ISO,11}^r = C_{ISO,22}^r = C_{ISO,33}^r = K + \frac{4}{3}G$, $C_{ISO,12}^r = C_{ISO,13}^r = C_{ISO,23}^r = K - \frac{2}{3}G$, $C_{ISO,44}^r = C_{ISO,55}^r = C_{ISO,66}^r = G$, knowing that:

$$K = \frac{[(C_{11}^r + C_{22}^r + C_{33}^r) + 2(C_{12}^r + C_{13}^r + C_{23}^r)]}{9} \quad (B5a)$$

$$G = \frac{[(C_{11}^r + C_{22}^r + C_{33}^r) - (C_{12}^r + C_{13}^r + C_{23}^r) + 3(C_{44}^r + C_{55}^r + C_{66}^r)]}{15} \quad (B5b)$$

The isotropic component of the aggregate stiffness tensor \mathbf{C}_{ISO}^r is then replaced with the isotropic stiffness tensor $\mathbf{C}_{ISO}^{MMA\ EoS}$ obtained using the bulk and shear moduli from the lookup tables computed with the MMA_EoS software (Chust et al., 2017) as:

$$\mathbf{C}_{MMA\ EoS}^r = \mathbf{C}^r + \mathbf{C}_{ISO}^{MMA\ EoS} - \mathbf{C}_{ISO}^r \quad (B6)$$

Table B1: Composition (in mol%) used in MMA_EoS to compute lookup tables for different mantle and basalt composition.

Lithology	Index	SiO ₂	MgO	FeO	CaO	Al ₂ O ₃	Na ₂ O	Source [†]
Dunite	1	40.43	42.45	16.07	0.51	0.51	0.04	(1)
Harzburgite	2	36.54	56.17	5.71	0.99	0.59	0.00	(2)
Pyrolite	3	38.71	49.85	6.17	2.94	2.22	0.11	(3)

Basalt	4	51.76	15.11	6.59	14.39	10.39	1.76	(2)
Pyroxenite	5	46.46	16.99	7.95	11.70	15.48	1.43	(4)

† (1) Kelemen and Ghiorso, 1986; (2) Chemia et al., 2015; (3) Workman and Hart, 2005; (4) Hirschmann et al., 2003.

Table B2 - Elastic moduli (GPa) and their P-T derivatives of the mineral phases present in ECOMAN. Temperature derivatives are scaled by 10^2 GPa K^{-1} , and cross derivatives by $10^{-3} K^{-1}$.

Mineral	Index	Moduli	C11	C22	C33	C12	C13	C23	C44	C55	C66	Source [†]
Olivine	1	<i>M</i>	320.71	197.25	234.32	69.84	71.22	74.80	63.77	77.67	78.36	(1)
		$\partial M/\partial P$	5.41	5.26	3.78	1.88	1.53	1.60	1.08	1.42	1.80	(2)
		$\partial M/\partial T$	-4.02	-3.10	-3.53	-1.14	-0.96	-0.72	-1.26	-1.30	-1.56	(1)
Enstatite	2	<i>M</i>	236.90	180.50	230.40	79.60	63.20	56.80	84.30	79.40	80.10	(3)
		$\partial M/\partial P$	10.27	8.87	11.7	6.22	6.63	7.26	1.23	0.75	2.78	(3)
		$\partial^2 M/\partial P^2$	-0.47	-0.38	-0.53	-0.33	-0.26	-0.31	0	0	0	(3)
		$\partial M/\partial T$	-3.64	-3.43	-5.70	-1.52	-2.29	-1.63	-1.23	-1.58	-1.51	(4)
Wadsleyite	3	<i>M</i>	370.50	367.70	272.40	65.60	95.20	105.10	111.20	122.50	103.10	(5)
		$\partial M/\partial P$	5.21	6.83	8.06	4.06	3.30	3.30	1.39	0.00	1.88	(5)
		$\partial M/\partial T$	-4.02	-3.10	-3.53	-1.14	-0.96	-0.72	-1.26	-1.30	-1.56	(1)
Wadsleyite Fe = 0.94 H ₂ O = 0.15 wt%	4	<i>M</i>	241.00	358.00	224.00	75.00	99.00	102.00	106.00	109.00	90.80	(6)
		$\partial M/\partial P$	7.80	5.37	10.5	4.60	2.33	4.20	0.87	0.84	2.43	(6)
		$\partial^2 M/\partial P^2$	-0.26	0	-0.46	-0.11	0	-0.14	0	0	-0.053	(6)
Ringwoodite	5	<i>M</i>	329.00	-	-	118.00	-	-	130.00	-	-	(7)
		$\partial M/\partial P$	6.20	-	-	0.80	-	-	2.80	-	-	(7)
		$\partial M/\partial T$	-4.90	-	-	-0.70	-	-	-1.30	-	-	(7)
Garnet	6	<i>M</i>	299.10	-	-	106.70	-	-	93.70	-	-	(8)
		$\partial M/\partial P$	6.54	-	-	2.87	-	-	1.72	-	-	(8)
		$\partial M/\partial T$	-3.05	-	-	-0.58	-	-	-0.71	-	-	(9)
Bridgmanite	7	<i>M</i>	484.00	542.00	477.00	146.00	146.00	162.00	195.00	172.00	151.00	(9)
		$\partial M/\partial P$	3.40	5.30	5.20	3.30	2.40	2.50	1.40	0.80	1.40	(10)
		$\partial M/\partial T$	-1.80	-3.30	-2.20	-0.40	-0.20	-0.50	-2.20	-0.60	-1.60	(10)
		$\partial^2 M/\partial P \partial T$	-0.017	-0.016	-0.016	-0.006	-0.006	-0.007	-0.0005	-0.0005	-0.004	(10)
Bridgmanite P0 = 34 GPa T0 = 1500 K	8	<i>M</i>	581.42	669.06	626.31	249.02	212.93	239.01	228.63	220.26	183.89	(11)
		$\partial M/\partial P$	3.73	5.54	5.47	3.30	2.48	2.53	1.38	0.85	1.56	(11)
		$\partial M/\partial T$	-3.99	-5.53	-4.32	-1.76	-0.64	-0.26	-2.12	-1.25	-1.96	(11)
		$\partial^2 M/\partial P \partial T$	-0.16	-0.044	-0.051	-0.017	-0.06	-0.035	-0.017	-0.044	-0.042	(11)
Bridgmanite P0 = 48 GPa	9	<i>M</i>	601.83	698.17	652.27	271.81	236.00	265.58	233.53	205.41	190.43	(11)
		$\partial M/\partial P$	3.73	5.54	5.47	3.30	2.48	2.53	1.38	0.85	1.56	(11)

T0 = 1500 K		$\frac{\partial M}{\partial T}$	-3.14	-5.94	-4.53	-1.94	-0.55	-0.59	-2.15	-1.26	-2.50	(11)
		$\frac{\partial^2 M}{\partial P \partial T}$	-0.66	-0.026	-0.034	-0.158	-0.008	-0.160	-0.001	-0.036	-0.002	(11)
Ferropericlase	10	M	300.0	-	-	93.60	-	-	147.0	-	-	(12)
		$\frac{\partial M}{\partial P}$	9.56	-	-	1.45	-	-	1.03	-	-	(12)
		$\frac{\partial M}{\partial T}$	-5.98	-	-	0.89	-	-	-0.88	-	-	(12)
		$\frac{\partial^2 M}{\partial P \partial T}$	0.56	-	-	0.06	-	-	0.20	-	-	(12)
post-Perovskite	11	M	1124.75	884.22	1124.32	350.69	287.53	461.05	254.57	231.19	344.05	(13)
		$\frac{\partial M}{\partial P}$	5.85	2.70	5.44	3.06	3.00	2.33	1.37	1.27	2.06	(13)
		$\frac{\partial M}{\partial T}$	1.01	-5.24	-4.90	-0.92	-0.51	-1.78	-0.81	-2.33	-2.19	(13)
		$\frac{\partial^2 M}{\partial P \partial T}$	-0.937	0.709	0.797	0.60	-0.801	0.692	-0.107	-0.186	-0.16	(13)

† (1) Isaak, 1992; (2) Zha et al., 1998; (3) Chai et al., 1992; (4) Jackson et al., 2007; (5) Zha et al., 1997; (6) Zhou et al., 2022; (7) Sinogeikin et al., 2003; (8) Chai and Brown, 1997; (9) Sinogeikin and Bass, 2002; (10) Wentzcovitch et al., 2004. Derivatives at (100 GPa, 300 K); (11) Zhang et al., 2013. Derivatives extrapolated by interpolation; (12) Karki et al., 2000; (13) Zhang et al., 2016. Derivatives extrapolated by interpolation.

Appendix C. Velocity gradient tensor in polar coordinates

In polar grids the velocity field is typically defined by the longitudinal, radial and, in 3D, colatitudinal components $\mathbf{V}_{(\phi,r,\theta)} = (V_\phi, V_r, V_\theta)$. However, the crystals' orientation and their rotation are defined relative to the external Cartesian reference frame. As such, the velocity gradient tensor must be computed in Cartesian coordinate system. This is done by rotating the velocity field from spherical to Cartesian coordinates as:

$$\begin{bmatrix} V_x \\ V_y \\ V_z \end{bmatrix} = \begin{bmatrix} -\sin\phi & \sin\theta\cos\phi & \cos\theta\cos\phi \\ \cos\phi & \sin\theta\sin\phi & \cos\theta\sin\phi \\ 0 & \cos\theta & -\sin\theta \end{bmatrix} \begin{bmatrix} V_\phi \\ V_r \\ V_\theta \end{bmatrix} \quad (C1)$$

Using the finite difference approach, the spatial gradients of the Cartesian velocity components are computed on the spherical grid as:

$$\mathbf{L}_{(\phi,r,\theta)}^{(x,y,z)} = \frac{\partial \mathbf{V}_{(x,y,z)}}{\partial (\phi, r, \theta)} = \begin{bmatrix} \frac{\partial V_x}{\partial \phi} & \frac{\partial V_x}{\partial r} & \frac{\partial V_x}{\partial \theta} \\ \frac{\partial V_y}{\partial \phi} & \frac{\partial V_y}{\partial r} & \frac{\partial V_y}{\partial \theta} \\ \frac{\partial V_z}{\partial \phi} & \frac{\partial V_z}{\partial r} & \frac{\partial V_z}{\partial \theta} \end{bmatrix} \quad (C2)$$

Defining \mathbf{J} as the Jacobian matrix:

$$\mathbf{J} = \frac{\partial(\phi, r, \theta)}{\partial(x, y, z)} = \begin{bmatrix} \frac{\partial\phi}{\partial x} & \frac{\partial\phi}{\partial y} & \frac{\partial\phi}{\partial z} \\ \frac{\partial r}{\partial x} & \frac{\partial r}{\partial y} & \frac{\partial r}{\partial z} \\ \frac{\partial\theta}{\partial x} & \frac{\partial\theta}{\partial y} & \frac{\partial\theta}{\partial z} \end{bmatrix} = \begin{bmatrix} -\sin\phi/(r \cdot \sin\theta) & \cos\phi/(r \cdot \sin\theta) & 0 \\ \sin\theta\cos\phi & \sin\theta\sin\phi & \cos\theta \\ \cos\theta\cos\phi/r & \cos\theta\sin\phi/r & -\sin\theta/r \end{bmatrix} \quad (C3)$$

the velocity gradient tensor in Cartesian coordinates is $\mathbf{L}_{(x,y,z)} = \mathbf{L}_{(\phi,r,\theta)} \mathbf{J}$:

$$\mathbf{L}_{(x,y,z)} = \frac{\partial\mathbf{V}_{(x,y,z)}}{\partial(x, y, z)} = \begin{bmatrix} \frac{\partial V_x}{\partial x} & \frac{\partial V_x}{\partial y} & \frac{\partial V_x}{\partial z} \\ \frac{\partial V_y}{\partial x} & \frac{\partial V_y}{\partial y} & \frac{\partial V_y}{\partial z} \\ \frac{\partial V_z}{\partial x} & \frac{\partial V_z}{\partial y} & \frac{\partial V_z}{\partial z} \end{bmatrix} = \begin{bmatrix} \frac{\partial V_x}{\partial\phi} & \frac{\partial V_x}{\partial r} & \frac{\partial V_x}{\partial\theta} \\ \frac{\partial V_y}{\partial\phi} & \frac{\partial V_y}{\partial r} & \frac{\partial V_y}{\partial\theta} \\ \frac{\partial V_z}{\partial\phi} & \frac{\partial V_z}{\partial r} & \frac{\partial V_z}{\partial\theta} \end{bmatrix} \begin{bmatrix} \frac{\partial\phi}{\partial x} & \frac{\partial\phi}{\partial y} & \frac{\partial\phi}{\partial z} \\ \frac{\partial r}{\partial x} & \frac{\partial r}{\partial y} & \frac{\partial r}{\partial z} \\ \frac{\partial\theta}{\partial x} & \frac{\partial\theta}{\partial y} & \frac{\partial\theta}{\partial z} \end{bmatrix} \quad (C4)$$

Appendix D. Aggregate rotation due to fluid body rotation

When multiple creep mechanisms are active, we assume that only the fraction of deformation accommodated by dislocation creep contributes to the LPO development. This fraction of the total deformation is defined as $F_d = \eta_{eff}/\eta_{dist}$, where $0 \leq F_d \leq 1$ is interpolated from the large-scale geodynamic model, η_{eff} is the effective viscosity resulting from the harmonic average of the viscosities from the flow law of different creep mechanisms, η_{dist} the viscosity from the dislocation creep flow law. At a given timestep Δt , crystal rotation according to the D-REX model is applied for the fraction $\Delta t_{dist} = \Delta t \cdot F_d$, while for the remaining time $\Delta t_{diff} = \Delta t \cdot (1 - F_d)$ we follow the numerical study of Hedjazian et al. (2017) and apply fluid body rotation by multiplying the CDM of the aggregates with the rotation of the FSE. In this way, we preserve and do not alter the strength of the LPO.

Given the orthogonal CDMs defining the original and final crystal orientation \mathbf{a}^t and $\mathbf{a}^{t+\Delta t_{diff}}$, and those defining the original and final FSE semiaxes orientation \mathbf{v}^t and $\mathbf{v}^{t+\Delta t_{diff}}$ (i.e., the eigenvectors of, respectively, the left stretch tensor $\mathbf{L}\mathbf{S}^t = \mathbf{F}^t(\mathbf{F}^t)^T$ and $\mathbf{L}\mathbf{S}^{t+\Delta t_{diff}} = \mathbf{F}^{t+\Delta t_{diff}}(\mathbf{F}^{t+\Delta t_{diff}})^T$), the following relation implies that after fluid body rotation the orientation of the crystal relative to FSE semiaxes must remain constant:

$$\mathbf{a}^{t+\Delta t_{diff}} \mathbf{v}^{t+\Delta t_{diff}} = \mathbf{a}^t \mathbf{v}^t \quad (D1)$$

The new crystal orientation is then:

$$\mathbf{a}^{t+\Delta t_{diff}} = \mathbf{a}^t \mathbf{v}^t (\mathbf{v}^{t+\Delta t_{diff}})^{-1} = \mathbf{a}^t \mathbf{v}^t (\mathbf{v}^{t+\Delta t_{diff}})^T \quad (D2)$$

926

927 **Additional References**

928 M. Chai and J. Brown. The elastic constants of a pyrope-grossular-almandine garnet to 20 GPa. *Geophys. Res. Lett.*,
929 24(5):523–526, 1997.

930 M. Chai, J. Brown, and L. Slutski. The elastic constants of an aluminous orthopyroxene to 12.5 GPa. *J. Geophys. Res.*,
931 102(14):779–785, 1992.

932 Z. Chemia, D. Dolejš, and G. Steinle-Neumann. Thermal effects of variable material properties and metamorphic reactions in
933 a three-component subducting slab. *J. Geo-phys. Res.*, 120:6823–6845, 2015.

934 M. Hirschmann, T. Kogiso, M. Baker, and E. Stopler. Alkalic magmas generated by partial melting of garnet pyroxenite.
935 *Geology*, 31:481–484, 2003.

936 D. Isaak. High-temperature elasticity of iron-bearing olivines. *J. Geophys. Res.*, 97: 1871–1885, 1992.

937 J. Jackson, S. Sinogeikin, and J. Bass. Sound velocities and single-crystal elasticity of orthoenstatite to 1073 K at ambient
938 pressure. *Phys. Earth Planet. Int.*, 171:1–12, 2007.

939 B. Karki, R. Wentzcovitch, S. de Gironcoli, and S. Baroni. High-pressure lattice dynamics and thermoelasticity of MgO. *Phys.*
940 *Rev. B*, 61:8793–8800, 2000.

941 P. Kelemen and M. Ghiorso. Assimilation of peridotite in zoned calc-alkaline plutonic complexes: evidence from the big jim
942 complex, Washington Cascades. *Contrib. Min.*, 94:12–28, 1986.

943 David Mainprice. Seismic Anisotropy of the Deep Earth from a Mineral and Rock Physics Perspective. *Treatise on Geophysics*,
944 2, Editor: G. Schubert, Elsevier, pp.437-491, 2007.

945 S. Sinogeikin and J. Bass. Elasticity of pyrope and majorite – pyrope solid solutions to high temperatures. *Earth Planet. Sci.*
946 *Lett.*, 203:549–555, 2002.

947 S. Sinogeikin, J. Bass, and K. Katsura. Single-crystal elasticity of ringwoodite to high pressures and high temperatures:
948 implications for 520 km seismic discontinuity. *Phys. Earth Planet. Int.*, 136:41–66, 2003.

949 R. Wentzcovitch, B. Karki, M. Cococcioni, and S. de Gironcoli. Properties of MgSiO₃-perovskite: Insights on the nature of
950 the earth's lower mantle. *Phys. Rev. Lett.*, 92, 2004.

951 R. Workman and S. Hart. Major and trace element composition of the depleted morb mantle. *Earth Planet. Sci. Lett.*, 231:53–
952 72, 2005.

953 C. Zha, T. Duffy, H. Mao, R. Downs, R. Hemley, and D. Weidner. Single-crystal elasticity of β -Mg₂SiO₄ to the pressure of
954 the 410 km seismic discontinuity in the earth's mantle. *Earth Planet. Sci. Lett.*, 147:E9–E15, 1997.

955 C. Zha, T. Duffy, R. Downs, H. Mao, and R. Hemley. Brillouin scattering and x-ray diffraction of San Carlos olivine: direct
956 pressure determination to 32 GPa. *Earth Planet. Sci. Lett.*, 159:25–33, 1998.

957 Z. Zhang, L. Stixrude, and J. Brodholt. Elastic properties of MgSiO₃-perovskite under lower mantle conditions and the
958 composition of the deep earth. *Earth Planet. Sci. Lett.*, 379:1–12, 2013.

959 [Z. Zhang, S. Cottaar, L. Tao, S. Stackhouse, and B. Militzer. High-pressure, temperature elasticity of Fe- and Al-bearing](#)
960 [MgSiO₃: Implications for the earth's lower mantle. Earth Planet. Sci. Lett., 434:264–273, 2016.](#)
961 [W.-Y. Zhou, M. Hao, J. Zhang, B. Chen, R. Wang, and B. Schamndt. Constraining composition and temperature variations in](#)
962 [the mantle transition zone. Nat. Comm., 13:1094, doi: 10.1038/s41467-022-28709-7, 2022.](#)
963

Page 33: [1] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [2] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [3] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [4] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [5] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [6] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [7] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [8] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [9] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [10] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [11] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [12] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [13] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [14] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [15] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [16] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [17] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [18] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [19] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [20] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [21] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [22] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [23] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [24] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [25] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [26] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [27] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [28] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [29] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [30] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [31] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲

Page 33: [32] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [33] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [34] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [35] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [36] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [37] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [38] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [39] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [40] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲
Page 33: [41] Formatted **Manuele Faccenda** **31/05/2024 11:43:00**

Font: 10 pt

▲