



Effects of basalt, concrete fines, and steel slag on maize growth and heavy metal accumulation in an enhanced weathering experiment

Jet Rijnders^{1*}, Arthur Vienne¹, Sara Vicca¹

5 ¹*Biobased Sustainability Engineering (SUSTAIN), Department of Bioscience Engineering, University of Antwerp, Groenenborgerlaan 171, 2020 Antwerpen, Belgium*

*Correspondence to: Jet Rijnders (jet.rijnders@uantwerpen.be)

Abstract.

10 Terrestrial enhanced silicate weathering is a CO₂ removal technology involving the application of ground silicate materials to agricultural soils. Next to CO₂ sequestration, it can improve soil fertility and crop growth, but silicate materials can also contain toxic trace elements. In a mesocosm experiment, we investigated the effect of basalt, concrete fines and steel slags on biomass, nutrients, and heavy metal concentration of Zea Mays, using a dose-response approach.

15 Plant biomass increased with basalt, but not with concrete fines and steel slags. Generally, plant Ca, Mg, and corn Si concentrations increased with increasing silicate application amount as a result of increased plant availability. In contrast, plant N, P, and K concentrations were hardly affected by silicate application. Besides increased leaf Pb concentrations with steel slag application, which did not exceed the maximum limit set by the WHO and FAO (0.05 mg Pb kg⁻¹ ww), heavy metal concentrations in aerial plant tissues mostly decreased with increasing silicate application amount, presumably because of an increased soil pH, and accumulation in plant roots. Our study thus indicates mixed effects of silicate application on maize
20 while suggesting that the risk of heavy metal contamination after a one-time application of the tested silicates is limited.



1. Introduction

Recently, there has been a growing interest in the use of silicate rock powder in agriculture for carbon dioxide (CO₂) removal (e.g. Edwards et al., 2017; Swoboda et al., 2022). When silicates react with water and CO₂, carbon (C) is locked up
25 in carbonates, possibly for centuries and longer (Moosdorf et al., 2014). While the naturally occurring silicate rock weathering process has been important for stabilizing climate at geological timescales, its pace is insufficient to substantially reduce the current rise in atmospheric CO₂ (Berner, 2004; Walker et al., 1981). Enhanced silicate weathering (EW) aims to accelerate this natural process through the mechanical grinding of the rocks into a fine powder.

30 In addition to its CO₂ sequestration potential, applying silicate minerals to soils holds promise for improving agricultural practices. Crops can benefit from the essential plant nutrients, such as Ca²⁺ and Mg²⁺, that are released when silicate minerals weather (Kelland et al., 2020; Ramos et al., 2022). Furthermore, silicate materials can reduce soil acidity, a pervasive issue in agriculture, thus contributing to soil health and fertility (Calabrese et al., 2022). Moreover, the implementation of EW can improve soil properties such as soil water retention and cation exchange capacity (CEC) (Anda et al., 2015; Calabrese et al.,
35 2022; Taylor et al., 2017), to the benefit of crop productivity. Additionally, the process of EW releases silicon (Si), which can improve plant resistance to pests and diseases, thereby improving crop health and productivity in general (Calabrese et al., 2022; Swoboda et al., 2022).

Silicate rock powder has been used as a fertilizer for many years (e.g. Van Straaten, 2006), particularly in tropical regions,
40 where the release of base cations from these rocks can significantly enhance crop productivity (e.g. Swoboda et al., 2021). Nonetheless, EW also holds certain risks that need to be considered. Silicate materials typically contain heavy metals that are released into the environment during weathering, posing the risk of uptake by plants. The amount of heavy metals varies strongly among rocks and industrial silicates (Dupla et al., 2023; O'Connor et al., 2021). For instance, basalt, a naturally occurring and globally abundant silicate rock, generally exhibits lower concentrations of potentially harmful metals such as
45 nickel (Ni) and chromium (Cr) compared to other silicate rock types and is therefore a preferred rock source to consider in agriculture (Kelland et al., 2020). Basalt is a mafic igneous rock that contains substantial amounts of calcium (Ca)-and magnesium (Mg)-rich silicate minerals (Lewis et al., 2021). In the review of Swoboda et al. from 2022, all trials with (ultra)mafic rocks on agricultural soils improved yields. Recent studies further corroborate these findings; Luchese et al. (2023), for instance, documented improved corn biomass upon basalt application in both clay and sandy clay loam soils.
50 Similarly, basalt addition led to notable increases in dry mass, height and stem diameter, and the accumulation of macronutrients (Nitrogen (N), Phosphorous (P), potassium (K), Sulfur (S), Ca and Mg) in corn and beans grown on tropical soils (Conceição et al., 2022).



55 Most EW experiments have been conducted in tropical regions on highly weathered, acidic soils, while studies in a temperate climate are still scarce. Yet, Skov et al. (2024) demonstrated an increase in spring oat yield after the application of basalt on direct drill and ploughed plots in a temperate climate. These increases were assigned to a modest increase in pH that resulted in reduced manganese (Mn) and iron (Fe) uptake. This study also reported higher tissue Ca content and increased grain and tissue K upon basalt application. Contrastingly, the addition of volcanic rock dust to soils did not influence wheat growth in a mesocosm experiment in Sweden (Ramezani et al., 2013). Furthermore, the aboveground biomass of potato
60 plants tended to increase with basalt application in a mesocosm experiment in Belgium, growing on an alkaline soil (Vienne et al., 2022). Nonetheless, it is important to keep in mind that the apparent dominance of positive EW effects on crop growth might be affected by publication bias, as negative or non-significant results are less likely to be published compared to positive and expected outcomes (Dieleman & Janssens, 2011).

65 While the use of geological silicates, requires energy for the mining and grinding of rocks, thereby partially offsetting the CO₂ removal potential (Beerling et al., 2020; Goll et al., 2021), this energy cost may be strongly reduced by using silicate materials derived from industrial processes. For instance, Ca-rich silicates generated as by-products from steel manufacturing (steel slags), or waste concrete fines, represent alternatives that circumvent the need for additional mining (Renforth et al., 2011). These industrial by-products contain minerals that dissolve faster than basalt and are already produced in fine particle
70 sizes, thereby diminishing the energy cost of grinding (Beerling et al., 2020).

Concrete waste has already been applied to soils to improve plant growth, however, knowledge of how it affects plants is poorly understood (Ho et al., 2021). In contrast, the application of steel slags as a fertilizer in agriculture is already a common practice (Gao et al., 2020). Various trials demonstrated increases in biomass upon steel slag application, for instance for
75 soybean (Castro & Crusciol, 2013; Deus et al., 2020), maize (Castro & Crusciol, 2013; Wang & Qing-sheng, 2006) and rice (Makabe-sasaki & Sasaki, 2015; Wang et al., 2015). At three different locations in Germany, long-term field trials with steel slag application have been running since 1954, 1990 and 1993. At all locations, the yield of grass or arable crops increased significantly with the application of different kinds of steel slags, despite increases in Cr and Vanadium (V) contents in the soil (Branca et al., 2014). Nonetheless, as with other silicate materials, research about its influence on crops in temperate
80 climates is lacking.

Despite the evident benefits, there remain substantial gaps in our understanding of the broader implications of EW in agriculture. Specifically, its influence on nutrient cycling and the potential for heavy metal contamination, particularly with industrial by-products such as concrete fines and steel slags, warrants further investigation. Most research so far has been
85 conducted in a tropical climate, often on highly weathered and acidic soils, but EW is currently considered for application also in other climate regions, justifying further research.



This study aims to quantify the influence of basalt, concrete fines, and steel slags on *Zea mays* growth, yield, and nutrient concentrations in a mesocosm experiment in a temperate climate. The three types of silicates were applied in a dose-response approach ranging from 0 to 200 ton ha⁻¹ of basalt, 0 to 31 ton ha⁻¹ of concrete fines, and 0 to 10 ton ha⁻¹ of steel slags. We hypothesized that: I) Silicate weathering increases with increasing silicate application; II) Plant biomass and plant concentrations of nutrients present in the silicates increase with increasing silicate application, and III) Heavy metal concentrations increase with increasing silicate application, especially with basalt and steel slags, as they contain Cr, Ni and V, and Cr and V respectively, while these are absent in the concrete fines.

2. Materials and methods

2.1 Experimental set-up




Figure 1: a) overview of the experiment; b) one mesocosm with two maize plants.

Table 1: Soil characteristics of the original soil, before the start of the experiment.

Texture (Sand, clay, silt %)	Sandy loam (61, 4, 35)
pH	5.94 ± 0.03
Organic C (%)	0.234 ± 0.005
Inorganic C (%)	0.024 ± 0.018
Cation exchange capacity (meq 100g ⁻¹)	2.5

Thirty mesocosms (0.6 m height, radius = 0.25 m) were constructed at the experimental site at the Drie Eiken Campus of the University of Antwerp, Belgium (51°09' N, 04°24' E) (Fig. 1a), and placed outdoors to receive natural rainfall (Fig. 1a). In May 2021, the bottom 40 cm of each mesocosm was filled with a sandy-loam soil (Table 1) to a depth of 40 cm in the bottom



105 portion. The upper 20 cm was filled with the same soil, either unamended in the control treatment (5 replicates), or amended with one of three different types of silicates (basalt, concrete fines, or steel slags) mixed into the topsoil layer on May 10, 2021.

Table 2: Elemental composition (C=concentration) of basalt, concrete fines and steel slags.

Basalt		Concrete fines		Steel slags	
Element	C (wt%)	Element	C (wt%)	Element	C (wt%)
Si	14.70%	Si	34.73%	Ca	42.00%
Fe	8.58%	Ca	9.41%	Fe	18.30%
Ca	6.42%	Fe	2.52%	Si	4.93%
Mg	5.88%	Al	2.36%	Mn	2.20%
Al	5.16%	K	1.20%	Mg	0.97%
Na	1.87%	Mg	0.63%	Al	0.90%
Ti	1.08%	Na	0.41%	P	0.77%
K	0.77%	S	0.27%	Ti	0.47%
P	0.26%	Ti	0.18%	V	0.32%
Mn	0.15%	Zr	0.15%	Cl	0.28%
Cl	765 PPM			S	0.13%
Sr	739 PPM			Cr	0.12%
Ba	559 PPM			Nb	302 PPM
Ni	408 PPM			Sr	277 PPM
Cr	367 PPM			Zr	164 PPM
V	242 PPM			K	105 PPM
Ce	197 PPM			Zn	58.3 PPM
Zr	150 PPM				
Zn	103 PPM				
S	93.3 PPM				
Cu	75.2 PPM				
Nb	61.0 PPM				

Table 3: Application rates of basalt, concrete fines and steel slags and number of replicates for each application rate. The 5 replicates where no silicate material is applied are the same mesocosms for the three treatments.

Basalt		Concrete fines		Steel slags	
Application amount (ton ha ⁻¹)	Replicates	Application amount (ton ha ⁻¹)	Replicates	Application amount (ton ha ⁻¹)	Replicates
0	5	0	5	0	5
10	1	7.04	1	2	1
30	1	10.57	1	3	1
50	5	14.09	1	4	1
75	1	17.62	1	5	1
100	1	21.15	1	6	1
150	1	24.67	1	7	1
200	1	31.07	1	10	1

Eifel Basalt, with P80 = 310.78 μm, was obtained from DURUBAS (<https://www.rpbl.de>) (Table S1, S2). Blast-oxygen furnace slags (BOF-slugs, hereafter referred to as steel slags), with P80 = 201.65 μm were sourced from ArcelorMittal Ghent (<https://belgium.arcelormittal.com>) (Table S1, S2). Concrete waste was obtained from a local concrete recycling company (RECYBO, <http://www.recybo.be>) and dry sieved over a 2 mm sieve (Table S1, S2). The fraction <2 mm is further referred to as ‘concrete fines’. The elemental composition of the silicate materials can be found in Table 2.

A dose-response experiment was established, where the silicates were amended in different concentrations: seven application amounts of basalt (10 to 200 ton ha⁻¹), seven application amounts of concrete fines (7 to 31 ton ha⁻¹) and seven application amounts of steel slags (1 to 10 ton ha⁻¹) (Table 3). For basalt, there were five replicates of 50 ton ha⁻¹. The application rates were selected based on typical applications found in literature, while also aiming for similar neutralizing effects among silicate materials (based on the

CaO and MgO in the materials and a small lab test with the materials) (e.g. Kelland et al., 2020; White et al., 2017; Yang et al., 2019). Concrete fines could weather faster than 10⁻⁶ mol m⁻² s⁻¹, depending on their carbonation (Palandri & Kharaka, 2004) and the dissolution rate of steel slags ranges between 10⁻⁹ and 10⁻⁷ mol m⁻² s⁻¹ (De Windt



et al., 2011). This is faster compared to the weathering rates of basalt (10^{-12} – 10^{-10} mol m⁻² s⁻¹) (Gudbrandsson et al., 2011). Therefore, the application rate of basalt covers a wider range compared to concrete fines and steel slags.

Each mesocosm was equipped with a 2 cm diameter hole at the bottom for leachate collection, and a root exclusion mat covered the bottom of the mesocosm to prevent soil export through leaching. Glass collectors with a volume of 2.3 L were connected to the mesocosm via polyurethane tubing. On June 3, 2021, two sweet corn seedlings (variety Tom Thumb, purchased at Le Grenier) were planted in each mesocosm (Fig. 1b), and all pots were fertilized with nitrogen, phosphorous, and potassium (NPK; 96 – 10 – 79 kg ha⁻¹) by adding Ca(NO₃)₂, triple super phosphate (TSP, 45% P₂O₅) and K₂SO₄. The fertilization amount was similar to that used in Ven et al. (2019), except for P, which was halved to avoid overfertilization in combination with the added silicates.

Soil water content and temperature were monitored using Cambell Scientific sensors (CS616) in each mesocosm (Fig. S2). Precipitation and air temperature for the duration of the experiment were retrieved from visualcrossing (<https://www.visualcrossing.com/>) (Fig. S3). Mesocosms were also watered manually (Table S2) at the beginning of the experiment and during dry periods.

2.2 Plant measurements

Throughout the growing season, plant height was monitored at 10 times. On July 15 and August 11, 2021, leaf area index (LAI) was determined for each mesocosm, following the approach of Ven et al. (2019). For every plant, the numbers of leaves were counted (*n*_{leaf}) and the width (*W*_{leaf}) and length (*L*_{leaf}) of each leaf were measured. First, individual leaf area (*LA*) was calculated using the conversion factor *A*_{leaf} = 0.75 for maize (Montgomery, 1911) (eq 1).

$$LA = L_{\text{leaf}} * W_{\text{leaf}} * A_{\text{leaf}} \quad (\text{eq 1}) \quad 155$$

Leaf Area Index (LAI) was then calculated for each mesocosm, with *A* the area of the mesocosms (0.20 m²) (eq 2).

$$LAI = LA * n_{\text{plant}} * n_{\text{leaf}} * (A)^{-1} \quad (\text{eq 2})$$

Plants were harvested on August 26, 2021, starting with the top leaf of each plant for further analysis. The aboveground biomass was then harvested and separated into stems, leaves, tassels, and corn. A week after harvesting, roots were sampled from topsoil (0-20 cm) and subsoil (20-60 cm) to estimate the root biomass within each mesocosm. One soil core (100 cm³) was taken below the shoot of each plant (two cores per mesocosm for each soil layer) and one core for each soil layer at the centre of the mesocosms. Subsequently, soil was carefully rinsed over a 2 mm sieve positioned above a receptacle. Roots were collected from the sieve, including those floating on the water in the receptacle. After drying for 48h at 70 °C, the dry weight (dw) of each plant part was determined. For roots, the average dry root biomass (g cm³) was calculated



assuming that the core from the centre of the mesocosm represented 50% of the root distribution across the surface area, while each of the two cores underneath the plants accounted for 25% each.

Leaves, stems, tassels, corn seeds, and roots were ground using a centrifugal mill (model ZM 200, Retsch GmbH, Haan, Germany) with a sieve of 0.25 mm mesh size. All ground samples were analysed for nutrients (Ca, Fe, K, Mg, P, Si) through ICP-OES (iCAP 6300 duo, Thermo Scientific) and C and nitrogen (N) content by dry combustion, based on the Dumas method using an elemental analyser (model FLASH 2000, Interscience, Louvain-la-Neuve, Belgium). Leaves, stems, tassels and corn seeds were also analysed for heavy metals (cadmium (Cd), Cr, Ni, lead (Pb), V). Due to limited root samples that were harvested, heavy metal concentrations were not analysed in our study.

For each plant sample, 0.3 g was weighed and digested with H₂SO₄, salicylic acid, H₂O₂ and selenium to determine Ca, K, Mg, and P and the heavy metals listed above according to Walinga et al. (1989). Si was determined by digestion of 30 mg plant sample with 25 mL 0.5 N NaOH. Pb and Ni concentrations were compared to the maximum allowable levels. As these are reported per g ww of plant material, we converted our concentrations from dw to ww. For corn Pb, the Limit of Quantification (LOQ) (0.1 mg kg⁻¹ dw) was used since all Pb concentrations in corn were lower.

2.3 Soil measurements

Starting from May 19, 2021, soil pore water samples were collected weekly for chemical analysis using rhizons (Rhizon Flex, Rhizosphere Research Products B.V., Wageningen, NL) installed at 5 cm depth in each mesocosm. Soil pore water pH was measured with a Metrohm 914 pH/Conductometer. For chemical analysis, samples were filtered through a 0.45 µm PET filter. Concentrations of Ca, Cd, Cr, Fe, Mg, Ni, Pb, Si, and V were analysed with ICP-OES (iCAP 6300 duo, Thermo Scientific). Dissolved inorganic carbon (DIC) was measured with a Skalar (Formacs^{HT}). Ca, Fe, K, Mg and Si and DIC were analysed weekly, while Cd, Cr, Ni, Pb, and V were analysed once on August 18, 2021, near the end of the growing season.

Topsoil pH was measured on five occasions (May 25, June 10, and 24, July 19, and August 4) by collecting and pooling three subsamples per mesocosm. Subsequently, 4 g of soil was suspended in 10 mL of deionized water and shaken before analysis using a Metrohm 914 pH/Conductometer meter. Plant Root Simulator probes (PRSTM, Western Ag Innovations, Saskatoon, Canada) were inserted into the soil twice: on June 24 and August 02, 2021. They were retrieved one week later, on July 01 and August 09, 2021, respectively. More information about these ion exchange membranes is provided by Qian & Schoenau (2002). CEC of the soil was determined following the protocol of Brown (1943) for which approximately 2.5 g of air-dried soil was extracted with 1 M NH₄Acetate for 1h at room temperature with continuous agitation. Extracts were measured with ICP-OES (iCAP 6300 duo, Thermo Scientific).



2.4 Statistical analysis

All statistical analyses were conducted in R (Rstudio, 2021.09.0.0). The level of significance for all analyses was set at $p \leq 0.05$. Data normality of residuals was checked using the Shapiro–Wilk test and the homoscedasticity of the residuals was examined by plotting them against the fitted values. In case one of these assumptions was violated, a logarithmic transformation was applied to the data. Plots were designed with the ggplot2 package (Wickham, 2016), ggpubr package (Kassambara, 2023) and the Rmisc package (Hope, 2022).

Linear regression analyses were performed to explore the relationship between the application amount of each silicate material separately and the variables of interest, i.e. plant biomass, nutrient and heavy metal content of the different plant parts, heavy metal content of the pore water and CEC of the soil. If the normality or homoscedasticity of the residuals of the linear model was not met after data transformation, a non-parametric Kruskal-Wallis test was used. To assess changes over time and differences among application amounts for each silicate type, a linear mixed model (nlme package, Pinheiro et al., 2013) was applied for plant height, soil pH, soil pore water pH, soil pore water DIC, soil pore water nutrient concentrations, and nutrient availability from the PRS probes. Silicate application amount, time (days after sowing, or burial date for the nutrient availability from the PRS probes), and their interaction were included as fixed effects, while mesocosm was treated as a random effect. To account for heteroscedasticity in the data across different time points, the weights = varIdent component was incorporated into the model. Differences among silicates were investigated using method S1.

To investigate which soil parameters are linked with differences in biomass, a linear model was constructed based on the Akaike information criterion (AIC). All soil parameters (soil pH, CEC, pore water pH, pore water nutrients, and pore water heavy metals) were included in the original model, whereafter the best fit was selected with the lowest AIC score. Hereafter, a two-way ANOVA was performed to search for soil parameters that significantly influence plant biomass. To reduce the complexity of the data, a principle component analysis (PCA) was done to explore differences in soil variables (soil and pore water pH, CEC, pore water nutrients, and pore water heavy metals) among the three silicate treatments. Linear regression analysis was performed with each principal component (PC) as a function of silicate treatment and as a function of plant biomass.

3. Results

3.1 Soil chemistry

Application of basalt, concrete fines, and steel slags resulted in significant increases in soil pH, with the most pronounced increases for concrete fines and steel slags (Fig. 2; Table S3). Pore water pH and DIC generally increased as well with basalt, concrete fines, and steel slag application amount (Fig. 2, Table S3) with a more pronounced DIC increase with increasing steel




Figure 2: Soil pH, pore water DIC and pore water pH during the experiment for basalt, concrete fines and steel slag treatments. Control : treatment (0 ton ha^{-1}) and basalt treatment with an application rate of 50 ton ha^{-1} are averages of five replicates and are shown with standard error. The other treatments each have one replicate. P-values and F-values are shown from linear regression analysis with soil pH, pore water pH or DIC as fixed effect and silicate application amount, time and the interaction. If the interaction was not significant, this was deleted from the model and thus not shown here. Significant relationships are indicated by an asterisk (*).

<https://doi.org/10.5194/egusphere-2024-3022>

Preprint. Discussion started: 9 October 2024

© Author(s) 2024. CC BY 4.0 License.



while the steel slag effect slightly decreased as the growing season progressed (Fig. 2).




Figure 3: Ca, Fe, K, Mg and Si in the pore water during the experiment for the silicate treatments (basalt, concrete fines, steel slags). Data of 0 and 50 ton ha⁻¹ of basalt are average of five replicates with standard error. Note that in some cases, the error bars are smaller than the symbol. The other treatments have one measurement each. P and F-values are shown in Table S3.



The influence of silicate application on soil pore water Ca concentrations varied among silicate material types (Fig. 3; Table S4). Increases were observed with increasing application amount of concrete fines and steel slags (Fig. 3, Table S3), which is similar to plant available Ca from PRS probes (SI8a). Conversely, pore water Ca concentrations decreased with increasing basalt application amount. However, data for the highest application amounts of basalt (150 and 200 ton ha⁻¹) are only available for the last two sampling dates, during which Ca concentrations increased with increasing basalt application amount ($p < 0.01$, $F=26.38$, statistical analysis only performed on the last two sampling dates). This increase is supported by data retrieved from the PRS probes, where plant-available Ca increased with basalt application amount (Fig. S3). Nonetheless, plant-available Ca was lower with basalt compared to concrete fines and steel slag application (Table S5).

A significant time x application amount interaction effect was found for Mg and Si concentrations in the soil pore water (Fig. 3, Table S3). Both increased with basalt, concrete fines, and steel slag application amount, yet the positive effect on Mg concentrations diminished towards the end of the growing season (Fig. 3; Table S3). Plant available Mg from the PRS probes increased as well with basalt and concrete fines application amount (Fig. S3). Increases in Si with basalt and steel slags were higher at the beginning of the experiment, while increases in Si with concrete fines application amount were relatively constant over time (Fig. 3, Table S3). In contrast, Fe concentrations decreased with increasing concrete fines application amount whereas these were not affected by basalt or steel slag application (Fig. 3, Table S3). However, this was not supported by the bioavailability of Fe retrieved from the PRS probes, which was not affected by silicate application (Fig. S3). Pore water K concentrations significantly decreased with steel slag application amount, whereas no effects were observed with basalt and concrete fines (Fig. 3; Table S3).




Figure 4: Concentrations of Cr, Ni, Pb and V in the pore water at the end of the growing season for basalt, concrete fines and steel slag treatments. Control treatment (0 ton ha⁻¹) and basalt treatment with an application rate of 50 ton ha⁻¹ are averages of 5 replicates and are shown with standard error. The other treatments each have one replicate. P-values and F-values are shown from a linear regression analysis with heavy metal (Cr, Ni, Pb, V) concentration as fixed effect and silicate application amount as covariable. Significant relationships are indicated by an asterisk (*).



250

In over 73% of the pore water samples, Cd concentrations were below the LOQ (Table S6), rendering statistical analysis unfeasible. Cr concentrations remained unaffected by basalt, whereas a decrease was observed with increasing application amount of concrete fines, in contrast to increasing Cr concentrations with increasing steel slag application amount (Fig. 4, Table S4). Pore water Ni and Pb concentrations increased with basalt application amount, yet remained unaffected by concrete fines or steel slag application. Moreover, these concentrations were higher with basalt compared to concrete fines and steel slag application (Fig. 4, Table S4). These changes in Pb concentrations contrast with PRS probe data, where Pb increased with concrete fines application but remained unchanged with basalt and steel slag application amount (Fig. S3, Table S5). Pore water V concentrations were higher with steel slag application compared to basalt and concrete fines and increased with application amount of basalt, concrete fines and steel slags (Fig. 4, Table S4).

260

The PCA-analysis showed a significant effect of silicate material on PC1 ($p < 0.01$, $F = 32.9$) and PC5 ($p < 0.01$, $F = 8.18$)




Figure 5: PC1 and PC5 of the principal component analysis with pore water nutrients (Ca, Fe, K, Mg, Si), heavy metals (Cr, Ni, Pb, V), pore water pH, soil pH, and soil CEC. Differences in PC1 and PC5 were found among silicate materials (B=Basalt, C=Control, CF=Concrete fines, S=Steel slags) by linear regression analysis with PC1 or PC5 as fixed variable and silicate treatment as covariable. Similar letters mean no significant differences among the materials while different letters mean that PC1 or PC5 differed significantly between silicate materials, using a Tukey post-hoc test.

(Fig. 5). PC1 is represented by lower concentrations of Pb and Ni in the pore water, but higher soil pH, CEC and DIC and pore water Ca and V concentrations. PC1 is significantly lower for basalt compared to concrete fines and steel slags (both $p < 0.01$). PC5 is represented by a lower soil pH but high concentrations of Fe, Cr and V in the pore water. PC5 is significantly higher for steel slags compared to basalt ($p = 0.03$) and concrete fines ($p < 0.01$).

265



3.2 Plant growth parameters




Figure 6: Plant growth during the experiment for basalt, concrete fines, and steel slag treatments. Control treatment (0 ton ha⁻¹) and basalt treatment with an application rate of 50 ton ha⁻¹ are averages of 5 replicates and are shown with standard error. The other treatments have one replicate. P-values and F-values are shown from a linear mixed model with plant height as the response variable and silicate application amount, and time, and interaction as covariable. If the interaction was not significant, this was deleted from the model and thus not shown here. Significant relationships are indicated by an asterisk (*).




Figure 7: Biomass of the different plant parts (leaves, corn, tassel, stem) as well as total aboveground biomass of maize plants for basalt, concrete fines, and steel slag treatments. Control treatment (0 ton ha⁻¹) and basalt treatment with an application rate of 50 ton ha⁻¹ are averages of 5 replicates and are shown with standard error. The other treatments each have one replicate. P- and F-values are shown of a linear regression analysis with biomass as the response variable and concentration of the silicate material as covariable. Significant regression is indicated by an asterisk (*), with equation and regression line.




Figure 8: scatter plot of PC1 and PC3 of the principal component analysis with pore water nutrients (Ca, Fe, K, Mg, Si), heavy metals (Cr, Ni, Pb, V), pore water pH, soil pH, and soil CEC. The sizes of the dots reflect the total aboveground biomass, which was significantly correlated with PC3. Different silicate treatments (B= basalt, C= control, CF= concrete fines, S= steel slags) are displayed in different colours.

Although basalt application did not affect plant height and LAI, plant biomass increased with basalt application amount (Fig. 6; Fig. 7; Table S7). This increase was significant for the stem, tassel, total aboveground and root biomass, and borderline significant for the leaves (Fig. 7). Total aboveground biomass was significantly higher with basalt application compared to concrete fines and steel slags (Fig. 7, Table S8). Contrastingly, plant height increased over time with increasing concrete fines and steel slag application amount but LAI and stem, leaves, corn, and root biomass remained unaffected, and hence, total

aboveground biomass did not change (Fig. 6; Fig. 7; Table S8). Nonetheless, tassel biomass increased significantly with both increasing application amounts of steel slags and concrete fines (Fig. 7).

290 The aboveground:belowground ratio was unaffected by any treatments (Table S7). The model selected by the AIC method included K, Ca, Pb, and Cr concentrations in the soil pore water, and soil pH and CEC. Of these variables, total aboveground biomass was significantly positively correlated with soil pore water Ca ($p < 0.01$, $F = 3.47$) and Pb concentrations ($p < 0.01$, $F = 4.19$), and soil pH ($p = 0.03$, $F = 2.81$). The PCA analysis showed a significantly negative relationship between biomass and PC3, reflecting low CEC values and low pore water Mg, Ni, and Pb concentrations (Fig. 8).

295 3.3 Plant nutrients

Increased basalt application amount resulted in decreased Ca concentrations in the aboveground plant parts, but only significantly for the stem and the tassel. Root Ca concentrations, on the other hand, increased with increasing basalt application amount (Fig. 9), but these were generally lower with basalt than with concrete fines or steel slag application. Mg concentrations increased in the plant parts with increasing basalt application, albeit not statistically significant for the top leaf and the corn 300 (Fig. 9). Root Mg concentrations were significantly higher with basalt compared to concrete fines and steel slags (Table S9).

Basalt application did not significantly affect Fe and K concentration in any of the plant parts, and P concentration remained unaffected except for a significant increase in the tassel (Fig. 9). Basalt application had no significant impact on Si concentration in the top leaf, tassel, and roots. In stems, Si concentrations decreased with increasing application amount of basalt, while Si concentration increased in the corn (Fig. 9).

305

The C:N ratio of plant tissues was usually not affected by basalt application, although in the tassels, the C:N ratio decreased with increasing basalt amount, while in the top leaf, the C:N ratio increased due to reduced N concentration (Fig. 10; Fig.S4). Total plant C stocks increased with increasing basalt application amount, while N stocks were not affected (Fig. S5). Furthermore, total plant Mg and K stocks increased with increasing basalt application amount, while Ca, P, and Si stocks remained unaffected (Fig. 11).

310




Figure 9: Concentrations of Ca, Fe, K, Mg, P, and Si in corn, tassel, roots, stem, and top leaf of the basalt treatment. Data of 0 and 50 ton ha⁻¹ of basalt are averages of five replicates with standard error. Note that in some cases, the error bars are smaller than the symbol. The other application amounts each have one replicate. P- and F-values are shown of the linear model with elemental concentration as the response variable and silicate concentration as covariable. Significant relationships are indicated with an asterisk (*), with equation and regression line.




Figure 10: C:N ratio in corn, tassel, stem, top leaf, and roots of maize for the silicate treatments (basalt, concrete fines, steel slags). Data without silicates (0 ton ha^{-1}) and 50 ton ha^{-1} of basalt are an average of five replicates with standard error. Note that in some cases, the error bars are smaller than the symbol. The other treatments each have one replicate. P- and F-values are shown of the linear model with C:N ratio as the response variable and silicate concentration as covariable. Significant relationships are indicated with an asterisk (*), with equation and regression line.




Figure 11: Stocks of Ca, K, Mg, P, and Si in the crops for the silicate treatments (basalt, concrete fines, steel slags). Data without silicates (0 ton ha^{-1}) and 50 ton ha^{-1} of basalt are an average of five replicates with standard error. Note that in some cases, the error bars are smaller than the symbol. The other treatments each have one replicate. P and F-values are shown of the linear model with stocks as the response variable and silicate concentration as covariable. Significant relationships are indicated with an asterisk (*), with equation and regression line.




Figure 12: Concentrations of Ca, Fe, K, Mg, P, and Si in corn, tassel, roots, stem, and top leaf of the concrete fines treatment. Data of the control treatment (0 ton ha⁻¹) is an average of five replicates with standard error. Note that in some cases, the error bars are smaller than the symbol. The other application amounts each have one replicate. P- and F-values are shown of the linear model with elemental concentration as the response variable and silicate concentration as covariable. Significant relationships are indicated with an asterisk (*), with equation and regression line.

Concrete fines application did not affect plant Ca concentrations, except for significantly increased root Ca concentration and a borderline significant decrease in tassel Ca concentration (Fig. 12). Stem Ca concentrations were higher with concrete
 315 fines compared to basalt and steel slags (Table S9). Mg concentrations increased with increasing concrete fine application amount, albeit not statistically significantly in the corn and roots (Fig. 12). Corn Si concentrations increased as well with concrete fines, and were higher than with basalt and steel slags (Fig. 12, Table S9). Concrete fine application did not affect plant Fe, K, and P concentrations. The C:N ratio significantly increased in the top leaf, even though C and N concentrations were not significantly affected by concrete fines (Fig. 12; Fig. S4). In the other plant parts, the C:N ratio remained unaffected
 320 (Fig. 12). Total plant C stocks increased significantly with increasing application amount of concrete fines, whereas total plant



N stocks were not affected (Fig. S5). Plant P and Si stocks were not influenced by concrete fines application amounts, while Ca, K, and Mg stocks increased with concrete fines, albeit only statistically significantly for Ca and K (Fig. 11).




Figure 13: Concentrations of Ca, Fe, K, Mg, P, and Si in corn, tassel, roots, stem, and top leaf of the steel slag treatment. Data of the control treatment (0 ton ha⁻¹) is an average of five replicates with standard error. Note that in some cases, the error bars are smaller than the symbol. The other application amounts each have one replicate. P- and F-values are shown of the linear model with elemental concentration as the response variable and silicate concentration as covariable. Significant relationships are indicated with an asterisk (*), with equation and regression line.

Steel slag application did not affect plant Ca concentrations, except for decreased stem Ca concentrations (Fig. 13). In leaves and tassels, Mg concentrations significantly increased with steel slag application amount, while it decreased in the stems (Fig. 13). Stem Mg concentrations were also lower with steel slag application than with basalt or concrete fines (Table S9). Corn and root Mg concentrations were not affected (Fig. 13). Steel slag application did not influence Fe, K, P, or Si concentrations in the plants except for increased corn Si concentrations with increasing steel slag application amount (Fig. 13). Leaves Si concentrations were higher with steel slags compared to basalt and concrete fines (Table S9). The C:N ratios in the plant parts remained unaffected by steel slag application (Fig. 10), even though stem N and top leaf C concentrations decreased

(Fig. S4). Total plant C and N stocks (Fig. S5) and Ca, Mg, P, K, and Si stocks (Fig. 11) remained unaffected by steel slag application.

3.4 Heavy metals

In all corn samples and in over 85% of the stem samples, Pb was below the LOQ (0.1 mg kg⁻¹ dw) and thus could not be statistically analysed. Pb was only detected in one control treatment and three basalt treatments (Table S6). Corn Cd concentrations were below LOQ (0.015 mg kg⁻¹ dw) in all samples of the steel slag treatment and was only detected once with concrete fines and once with basalt application (Table S6). Basalt application did not significantly affect heavy metal concentration in the plant parts, except for decreasing Cd (significant), with the largest decrease of 54% when 75 ton basalt ha⁻¹ was applied, and V (borderline significant) concentrations in the top leaf (Fig. 14).

340




Figure 14: Concentrations of Cd, Cr, Ni, Pb and V in corn, tassel, stem and top leaf of the basalt treatments. Data of 0 and 50 ton ha⁻¹ of basalt are an average of five replicates with standard error. Note that in some cases, the error bars are smaller than the symbol. For 10, 25, 75, 100, 150, and 200 ton ha⁻¹ of basalt, there is one measurement each. P- and F-values are shown of the linear model with heavy metal concentration as the response variable and silicate concentration as covariable. Significant relationships are indicated with an asterisk (*), with equation and regression line.



Concrete fine application generally decreased plant Cr concentrations, albeit only significantly in corn and tassel (Fig. 15), with a maximum decrease of 81% with the highest application amount, and 46% for 10.57 ton ha⁻¹ of concrete fines, respectively. Additionally, tassel Cd concentrations decreased with increasing concrete fines application amount, while Cd concentrations in the other plant parts remained unaffected. Corn Ni concentrations decreased significantly with a maximum decrease of 60.8% when 21.15 ton ha⁻¹ of concrete fines was applied, and a borderline significant decrease of stem Ni concentrations was observed with increasing concrete fine application amount, while Ni concentration was unaffected in the other plant parts (Fig. 15). Plant Pb concentrations were not affected by concrete fines application, and neither were plant V concentrations, except for a decrease observed in the top leaf (Fig. 15).




Figure 15: Concentrations of Cd, Cr, Ni, Pb, and V in corn, tassel, stem, and top leaf for the concrete fines treatments. Data of 0 ton ha⁻¹ of concrete fines is an average of five replicates with standard error. Note that in some cases, the error bars are smaller than the symbol. For 7.04, 10.57, 14.09, 17.62, 21.15, 24.67 and 31.07 ton ha⁻¹ of concrete fines, there is one measurement each. P-and F-values are shown of the linear model with heavy metal concentration as response variable and silicate concentration as covariable. Significant relationships are indicated with an asterisk (*), with equation and regression line.



Apart from decreased Cd concentrations in the stem (significant) and in tassels (borderline significant), Cd levels were not affected by steel slag application (Fig. 16). Stem Cd concentrations were lower with steel slags compared to basalt and concrete fines (Table S10). Furthermore, steel slag application decreased corn Cr concentrations, up to 83% with an application amount of 3 ton ha⁻¹ of steel slags. No influence was observed in the other plant parts. V and Pb concentrations were not affected by steel slag application, except for an increase observed in the top leaf (Fig. 16). With the highest application amount of steel slags, Pb concentrations increased by 47% compared to the control treatment.




Figure 16: Concentrations of Cd, Cr, Ni, Pb, and V in corn, tassel, stem, and top leaf for the steel slag treatments. Data of 0 ton ha⁻¹ of concrete fines is an average of five replicates with standard error. Note that in some cases, the error bars are smaller than the symbol. For 2, 3, 4, 5, 6, 7, and 10 ton ha⁻¹ of steel slags, there is one measurement each. P- and F-values are shown of the linear model with heavy metal concentration as the response variable and silicate concentration as covariable. Significant relationships are indicated with an asterisk (*), with equation and regression line.



4. Discussion

4.1 Silicate weathering and soil properties

360 The DIC in the porewater provides an indication of the weathering rate of the added silicate minerals (Haque et al., 2019).
As expected, increases in DIC were lowest for basalt compared to concrete fines and steel slags, indicating a higher weathering
rate for the latter two silicates. This was accompanied by a higher initial increase of soil and pore water pH for steel slags and
concrete fines. Similar findings have been observed in other enhanced weathering studies; using a mixture of rock types (Guo
et al., 2023), basalt (Buckingham & Henderson, 2024; Conceição et al., 2022), and steel slags (Buckingham & Henderson,
365 2024). Moreover, the dissolution of silicate minerals liberates essential base cations such as Ca^{2+} and Mg^{2+} (e.g. Kelland et al.,
2020). The release of base cations increased with application amounts of the silicates, as evidenced by soil pore water, PRS
probes, and plant measurements, and this is in line with other experiments (Amann et al., 2020; F. Guo et al., 2023; Vienne et
al., 2022; Yan et al., 2023). Soil pore water Si concentrations increased as well in our study (Fig. 3), similar to previous studies
(e.g. Buckingham & Henderson, 2024; Guo et al., 2023).

370

In contrast, Fe was unaffected and even decreased with application of concrete fines. The latter may be attributed to the
increased pH, which can stimulate the formation of insoluble Fe compounds, reducing Fe availability (e.g. Lindsay et al., 2008;
Shenker & Chen, 2005). Basalt and steel slags contain more Fe compared to concrete fines (see Table 2) and may have supplied
sufficient Fe to counterbalance decreases in Fe as a result of increased pH.

375

The silicate materials in our study contained heavy metals which can also be released. Given their known toxicity for
plants (Nagajyoti et al., 2010), it is imperative to assess the associated risks of heavy metal pollution when considering EW in
agriculture. Indeed, pore water concentrations of Cr, Ni, Pb and V increased in some of our treatments. Increased pore water
Ni concentrations with basalt and higher pore water Cr concentrations with steel slags can be explained by the material
380 composition (Table 2). On the other hand, pore water V concentrations also increased with increasing amount of concrete
fines, which did not contain V. Pore water V concentrations were thus not only influenced by the release of V from the silicates
(as they also increased with steel slags, that do contain V), but also by changes in soil properties. As V is more mobile with
increasing pH (Chen et al., 2021), we presume that the increased pH (Fig. 2) increased V availability.

385

Availability of Pb is generally also controlled by soil pH. At higher pH, Pb is less available for plants (Kushwaha et al.,
2018). Despite basalt lacking Pb, and the rise in soil pH, pore water Pb concentrations increased with increasing basalt
application. This unexpected increase may be related to competition for soil binding sites, with the released Ca increasingly
out-competing Pb due to their similar size and reactivity (Klitzke & Lang, 2009; Li et al., 2014).



4.2 Plant growth

390 Plants can benefit from EW because it can create a more favourable environment (Calabrese et al., 2022; Kelland et al.,
2020). Indeed, basalt application increased aboveground and belowground biomass. These findings are consistent with
observations of tropical systems (Swoboda et al., 2022). Also in temperate climates, basalt has been reported to increase
agricultural yield (e.g. Kelland et al., 2020; Skov et al., 2024), but other studies found no significant effect (Vienne et al.,
2022). Furthermore, increased root biomass with basalt application may indicate higher belowground C inputs by plants, which
395 can impact soil organic C stocks (Kögel-Knabner et al., 2022).

In contrast to basalt, concrete fines, and steel slag application did not affect plant biomass, except for increased tassel
biomass. Previously, the application of steel slag has been shown to have varying effects on crop growth. In a tropical climate,
soybean yield increased with steel slag application in two consecutive crop cycles (Deus et al., 2020). Similarly, Wang et al.
400 (2015) reported an initial increase in rice yield upon steel slag application on a paddy field, but this effect diminished after two
growing seasons. Pistocchi et al. (2017), on the other hand, reported a decrease in wheat yield, while tomato yield increased
over three growing seasons upon steel slag application in slightly alkaline loamy sand soils.

In our study, differences in plant biomass were significantly correlated to differences in soil pH, and pH may thus (partly)
405 explain the biomass response. The optimal pH for maize growth is between 5.5 and 7.5 (Lizárraga-paulín et al., 2011), a
threshold that was surpassed early in the growing season for the highest application amounts of concrete fines and steel slags,
but not for basalt (Fig. 2). We therefore presume that the pH was more conducive across the spectrum of basalt application
amounts, thereby increasing the plant biomass. This postulation is supported by the PCA analysis, as soil pH was one of the
major variables affecting both PC1, which separated the control and basalt treatment from the concrete fines and steel slags,
410 and PC5, which separated the concrete fines and the steel slag treatments. However, plant biomass only differed significantly
between basalt and steel slag application, and not compared to concrete fines. The high variation in biomass with concrete
fines application precludes drawing firm conclusions. Even though the substantial pH increase may have limited the positive
plant growth response with steel slag application, no adverse effects were detected.

415 4.3 Nutrient status of the plants

Nutrient release from silicate weathering is expected to stimulate plant growth and biomass, with nutrient release varying
based on material composition. We expected the same differences in plant stoichiometry, and indeed, plant nutrient
concentrations aligned with the soil Si and Mg measurements; corn Si and plant Mg concentrations increased with silicate
application. However, with basalt application, plant Ca concentrations even decreased in some plant parts and are thus not in
420 line with plant availability of Ca, presumably because of competition between Mg and Ca. The Ca:Mg ratio in soil pore water



decreased with basalt (Fig. S6). This has previously been related to reduced plant Ca concentration (Kopsell et al., 2013; Osemwota, 2007). A similar trend of increasing Mg and decreasing Ca concentrations was found with dunite application in wheat and barley (Rijnders et al., 2023) and ryegrass (Ten Berge et al., 2012).

425 The three silicate materials contained Mg, resulting in higher pore water Mg and plant Mg concentrations. Mg is an essential plant nutrient that is crucial for various physiological functions, including protein and chlorophyll synthesis, enzyme activation and ATP synthesis, and regulatory processes of photosynthesis (Hawkesford et al., 2012). Mg deficiency often arises in agricultural systems with imbalanced crop fertilization with N, P, and K, primarily due to competition between Mg and other ions, particularly K and Ca (Guo et al., 2016). The increased Mg availability and plant concentrations indicate that
430 fertilization with silicate minerals can aid in restoring Mg availability in agricultural soils.

Our results correspond to previous experiments employing basalt rocks. Boniao et al. (2002), for example, showed that corn height and biomass increased after basalt amendment and this was correlated with Mg concentration in soil pore water. Similarly, in Cameroon, maize cultivation utilizing three local basalts revealed the largest yield benefit for the basalt with the
435 highest Mg and Ca content, despite its lower K and P content (Tchouankoue, 2014). Additionally, other minerals, like the Mg-silicate dunite, also increased Mg concentrations in wheat and barley (Rijnders et al., 2023).

Even though not considered an essential plant nutrient, Si can protect the plant from (a)biotic stresses by making the plant structure more rigid and stronger (Bhatt & Sharma, 2018). Additionally, Si interacts with several key components of stress
440 signalling pathways of plants (Majeed et al., 2019), which can stimulate crop growth especially when plants are experiencing stress. Indeed, studies where steel slag increased rice (Ning et al., 2014) and soybean (Deus et al., 2020) yields also reported increased plant Si concentrations. In our study, the plants did not experience any noticeable stress, which may explain why Si concentrations were not related to increased plant biomass.

445 Unlike Ca, Mg, and Si, plant concentrations of P, K, and Fe were not much affected by silicate application. Despite decreased K availability with concrete fines and steel slags, plant K concentrations remained unaffected. The simultaneous increase in K stocks following the application of basalt and concrete fines suggests that K was sufficiently available across all treatments. While this contrasts with a few earlier studies that did report increased plant K concentration upon silicate addition (Skov et al., 2024; Ten Berge et al., 2012), our result is in line with the study of Rijnders et al. (2023), where the application
450 of 220 ton ha⁻¹ of dunite did not affect P and K concentrations in wheat and barley. Plant Fe concentrations also remained unaffected by silicate application, despite a decrease in Fe in soil pore water with concrete fines. Fe is relatively immobile, and symptoms of Fe deficiency typically appear on the new leaves. However, the lowest leaf Fe concentration measured in our study was 0.07 mg g⁻¹ dw (=70 ppm), which falls perfectly within the normal range of Fe concentrations in young leaves (i.e., 30 ppm Fe on a dry-weight basis; Hochmuth et al., 2012). Therefore, we can assume that sufficient Fe was available,



455 even though plant available Fe decreased with concrete fines. In an experiment of Wang & Qing-sheng (2006), Fe did not increase as well with their lowest steel slag application amount (44 ton ha⁻¹), using sandy loam soil.

Furthermore, plant available N did not change with silicate application. While plant N concentration slightly changed with basalt and steel slag application (Table S3), plant N stocks remained unaffected. This was expected since the silicates lack N and every treatment received the same input of N fertilizer. However, the C:N ratio within leaf tissues increased with basalt and concrete fines application, suggesting higher C assimilation rate per unit of N, indicating improved N use efficiency or N limitation (Schlüter et al., 2012). Given the biomass increase with basalt and no decrease with concrete fines, the elevated C:N ratio probably indicates a higher N use efficiency, particularly given that the mesocosms received the same amount of N fertilization.

465 **4.4 Plant heavy metal concentrations**

In addition to nutrients, silicate weathering can be accompanied by heavy metal release, raising concerns about heavy metal contamination. The primary risks associated with basalt are Al, Cr and Ni, while concrete fines contain only Al in a substantial amount, and steel slags contain various heavy metals, with notable risks linked to Al and Cr. Despite the release of heavy metals from silicate weathering, concentrations of these metals in plant tissues were not affected or even decreased with increasing silicate application amount. This is similar to a previous study where dunite application did not affect most heavy metal concentrations and even decreased Ba and Sr concentrations in wheat and barley (Rijnders et al., 2023). Deus et al. (2020) also found that steel slag application did not affect heavy metal concentrations in soybean, probably related to the increase in soil pH. Furthermore, basalt application did also not increase concentrations of Cd, Cr, Ni, and Pb in spring oats (Skov et al., 2024).

475 In this study, Cd, Cr, and V decreased in the plant parts with increasing silicate application. Decreases in Cd concentration were presumably a consequence of reduced Cd solubility following the pH increase with silicate application. Similar to our results, He et al. (2020) reported a decrease in Cd solubility and subsequent reduction in Cd concentrations in rice following steel slag amendment. In contradiction to our hypothesis, plant V concentrations did not increase in our experiment (and even decreased with concrete fines), despite elevated V concentrations in the soil pore water. Although V availability increases with pH, its uptake by roots decreases as the soil pH increases (Roychoudhury, 2020). Moreover, V that is taken up by plant roots will be predominantly retained in root tissue, particularly in cell walls where multiple proteins and polysaccharides form chelates with heavy metal ions (Aihemaiti et al., 2020; Roychoudhury, 2020), hence reducing its accumulation in aerial plant parts (Aihemaiti et al., 2020). This is similar for Cr, which may explain the decreased Cr concentrations in the above-ground plant parts (Ertani et al., 2017). Pistocchi et al. (2017) demonstrated that V and Cr indeed accumulated in the roots of tomato plants without translocation to the aerial plant parts after application of steel slags.



Furthermore, increased pore water Ni concentrations upon basalt application did not translate into increased Ni concentrations in the plant, presumably due to its uptake competition with other essential ions by plant roots (Sreekanth et al., 2013; Yusuf et al., 2011). This is similar to what was found by Vienne et al., 2022, where basalt application did not increase Ni concentrations in potatoes. Ni is a crucial component of many enzymes, and its deficiency can reduce plant growth but can be toxic for plants at elevated concentrations (Yusuf et al., 2011). Nevertheless, the toxicity level for sensitive species ($10 \text{ mg kg}^{-1} \text{ dw}$) was not exceeded in our study (Yusuf et al., 2011).

In contrast, top leaf Pb concentrations increased with steel slag application. Nonetheless, these remained below the maximum allowable level of $0.05 \text{ mg kg}^{-1} \text{ ww}$ that has been set for corn (WHO & FAO, 2022), ranging from $0.58 \cdot 10^{-6}$ to $0.03 \text{ mg kg}^{-1} \text{ ww}$ for all of the plant parts, and for the corn grains from $0.64 \cdot 10^{-4}$ to $0.6 \cdot 10^{-3} \text{ mg kg}^{-1} \text{ ww}$. The one-time steel slag application up to 10 t ha^{-1} in our experiment did thus not pose a problem of heavy metal contamination.

5. Conclusion

In our mesocosm experiment, we investigated the influence of a range of application amounts of basalt, concrete fines, and steel slags on the growth, nutrient status, and heavy metal concentrations in *Zea Mays*. Basalt application increased plant biomass, while plant biomass was not affected by concrete fines and steel slags. Differences in plant biomass are presumably due to exceedance of the optimal soil pH early in the growing season for the highest application amounts of concrete fines and steel slags, but not with basalt. However, despite the limited effect on crop growth, plants were not negatively affected by silicate application.

Silicate application increased soil pH, and pore water Ca, Mg, and Si availability which translated into increased plant Mg and Si concentrations. Plant Ca concentrations also increased with concrete fines and steel slag application, while a decrease in stem Ca was observed for basalt. This difference was probably related to the Ca:Mg ratio of the rock. Consistent with previous studies, heavy metal concentrations did not increase or even decreased in the aboveground plant parts with silicate application, even though the silicate materials contained and released heavy metals. Only leaf Pb increased with steel slag application and reached a concentration of up to $0.03 \text{ mg Pb kg}^{-1} \text{ ww}$. This Pb concentration is below the maximum level of $0.05 \text{ mg kg}^{-1} \text{ ww}$ of corn, set by the WHO. Moreover, Pb concentration in the corn grains was always below the LOQ. We therefore conclude that, in our experiment, crops mostly benefited from silicate application and did not show concerning accumulation of heavy metals. The largest benefits related to crop growth were observed for basalt application.

<https://doi.org/10.5194/egusphere-2024-3022>

Preprint. Discussion started: 9 October 2024

© Author(s) 2024. CC BY 4.0 License.



Competing interest

At least one of the (co-)authors is a member of the editorial board of Biogeosciences.



References

- 520 Aihemaiti, A., Gao, Y., Meng, Y., Chen, X., Liu, J., Xiang, H., Xu, Y., & Jiang, J. (2020). Review of plant-vanadium physiological interactions, bioaccumulation, and bioremediation of vanadium-contaminated sites. *Science of the Total Environment*, 712, 135637. <https://doi.org/10.1016/j.scitotenv.2019.135637>
- Amann, T., Hartmann, J., Struyf, E., De Oliveira Garcia, W., Fischer, E. K., Janssens, I., Meire, P., & Schoelynck, J. (2020). Enhanced Weathering and related element fluxes - A cropland mesocosm approach. *Biogeosciences*, 17, 103–119. <https://doi.org/10.5194/bg-17-103-2020>
- 525 Anda, M., Shamshuddin, J., & Fauziah, C. I. (2015). Improving chemical properties of a highly weathered soil using finely ground basalt rocks. *Catena*, 124, 147–161. <https://doi.org/10.1016/j.catena.2014.09.012>
- Beerling, D. J., Kantzas, E. P., Lomas, M. R., Wade, P., Eufrazio, R. M., Renforth, P., Sarkar, B., Andrews, M. G., James, R. H., Pearce, C. R., Mercure, J. F., Pollitt, H., Holden, P. B., Edwards, N. R., Khanna, M., Koh, L., Quegan, S., Pidgeon, N. F., Janssens, I. A., ... Banwart, S. A. (2020). Potential for large-scale CO₂ removal via enhanced rock weathering with croplands. *Nature*, 583, 242–248. <https://doi.org/10.1038/s41586-020-2448-9>
- 530 Berner, R. A. (2004). *The Phanerozoic Carbon Cycle: CO₂ and O₂*. Oxford University Press. <https://doi.org/10.1093/oso/9780195173338.001.0001>
- Bhatt, D., & Sharma, G. (2018). Role of silicon in counteracting abiotic and biotic plant stresses. *International Journal of Chemical Studies*, 6, 1434–1442.
- 535 Boniao, R. D., Shamshuddin, J., Van Ranst, E., Zauyah, S., & Syed Omar, S. R. (2002). Changes in chemical properties and growth of corn in volcanic soils treated with peat, ground basalt pyroclastics, and calcium silicate. *Communications in Soil Science and Plant Analysis*, 33(7–8), 1219–1233. <https://doi.org/10.1081/CSS-120003883>
- Branca, T. A., Pistocchi, C., Colla, V., Ragagnini, G., Amato, A., Tozzini, C., Mudersbach, D., Morillon, A., Rex, M., & Romaniello, L. (2014). Investigation of (BOF) Converter slag use for agriculture in Europe. 167, 155–167. <https://doi.org/10.1051/metal/2014022>
- 540 Brown, I. C. (1943). A rapid method of determining exchangeable hydrogen and total exchangeable bases of soils. *Soil Science*, 56, 353–357.
- Buckingham, F. L., & Henderson, G. M. (2024). The enhanced weathering potential of a range of silicate and carbonate additions in a UK agricultural soil. *Science of the Total Environment*, 907, 167701. <https://doi.org/10.1016/j.scitotenv.2023.167701>
- 545 Calabrese, S., Wild, B., Bertagni, M. B., Bourg, I. C., White, C., Aburto, F., Cipolla, G., Noto, L. V., & Porporato, A. (2022). Nano- to Global-Scale Uncertainties in Terrestrial Enhanced Weathering. *Environmental Science and Technology*, 56, 15261–15272. <https://doi.org/10.1021/acs.est.2c03163>
- 550 Castro, G. S. A., & Crusciol, C. A. C. (2013). Yield and mineral nutrition of soybean, maize, and congo signal grass as affected



by limestone and slag. *Pesquisa Agropecuaria Brasileira*, 48(6), 673–681. <https://doi.org/10.1590/S0100-204X2013000600013>

- Chen, L., Liu, J., Hu, W., Gao, J., & Yang, J. (2021). Vanadium in soil-plant system: Source, fate, toxicity, and bioremediation. *Journal of Hazardous Materials*, 405, 124200. <https://doi.org/10.1016/j.jhazmat.2020.124200>
- 555 Conceição, L. T., Silva, G. N., Holsback, H. M. S., Oliveira, C. de F., Marcante, N. C., Martins, É. de S., Santos, F. L. de S., & Santos, E. F. (2022). Potential of basalt dust to improve soil fertility and crop nutrition. *Journal of Agriculture and Food Research*, 10, 100443. <https://doi.org/10.1016/j.jafr.2022.100443>
- Deus, A. C. F., Büll, L. T., Guppy, C. N., Santos, S. de M. C., & Moreira, L. L. Q. (2020). Effects of lime and steel slag application on soil fertility and soybean yield under a no till-system. *Soil and Tillage Research*, 196, 104422. <https://doi.org/10.1016/j.still.2019.104422>
- 560 Dieleman, W. I. J., & Janssens, I. A. (2011). Can publication bias affect ecological research? A case study on soil respiration under elevated CO₂. *New Phytologist*, 517–521.
- Dupla, X., Möller, B., Baveye, P. C., & Grand, S. (2023). Potential accumulation of toxic trace elements in soils during enhanced rock weathering. *European Journal of Soil Science*, 74, 1–14. <https://doi.org/10.1111/ejss.13343>
- 565 Edwards, D. P., Lim, F., James, R. H., Pearce, C. R., Scholes, J., Freckleton, R. P., Beerling, D. J., & Edwards, D. P. (2017). Climate change mitigation: potential benefits and pitfalls of enhanced rock weathering in tropical agriculture. *Biology Letters*, 13, 20160715.
- Ertani, A., Mietto, A., & Borin, M. (2017). Chromium in Agricultural Soils and Crops: A Review. *Water, Air & Soil Pollution*, 228, 190. <https://doi.org/10.1007/s11270-017-3356-y>
- 570 Gao, D., Wang, F. P., Wang, Y. T., & Zeng, Y. N. (2020). Sustainable utilization of steel slag from traditional industry and agriculture to catalysis. *Sustainability (Switzerland)*, 12(21), 1–9. <https://doi.org/10.3390/su12219295>
- Goll, D. S., Ciais, P., Amann, T., Buermann, W., Chang, J., Eker, S., Hartmann, J., Janssens, I., Li, W., & Obersteiner, M. (2021). Potential CO₂ removal from enhanced weathering by ecosystem responses to powdered rock. *Nature Geoscience*, 14, 545–549. <https://doi.org/10.1038/s41561-021-00798-x>
- 575 Guo, F., Wang, Y., Zhu, H., Zhang, C., & Sun, H. (2023). Crop productivity and soil inorganic carbon change mediated by enhanced rock weathering in farmland: A comparative field analysis of multi-agroclimatic regions in central China. *Agricultural Systems*, 210, 103691. <https://doi.org/10.1016/j.agsy.2023.103691>
- Guo, W., Nazim, H., Liang, Z., & Yang, D. (2016). Magnesium deficiency in plants: An urgent problem. *Crop Journal*, 4(2), 83–91. <https://doi.org/10.1016/j.cj.2015.11.003>
- 580 Haque, F., Santos, R. M., Dutta, A., Thimmanagari, M., & Chiang, Y. W. (2019). Co-Benefits of Wollastonite Weathering in Agriculture: CO₂ Sequestration and Promoted Plant Growth. *ACS Omega*, 4, 1425–1433. <https://doi.org/10.1021/acsomega.8b02477>
- Hawkesford, M., Horst, W., Kichey, T., Lambers, H., Schjoerring, J., Skrumsager, I., & White, P. (2012). Functions of Macronutrients. In *Marschner's Mineral Nutrition of Higher Plants*. Elsevier Ltd. <https://doi.org/10.1016/B978-0-12->



585 384905-2.00006-6

- He, H., Xiao, Q., Yuan, M., Huang, R., Sun, X., Wang, X., & Zhao, H. (2020). Effects of steel slag amendments on accumulation of cadmium and arsenic by rice (*Oryza sativa*) in a historically contaminated paddy field. *Environmental Science and Pollution Research*, 27, 40001–40008. <https://doi.org/10.1007/s11356-020-10028-3>
- Ho, H. J., Iizuka, A., & Shibata, E. (2021). Chemical recycling and use of various types of concrete waste: A review. *Journal of Cleaner Production*, 284, 124785. <https://doi.org/10.1016/j.jclepro.2020.124785>
- 590 Hochmuth, G. J., Maynard, D., Vavrina, C., Hanlon, E., & Simonne, E. (2012). Plant Tissue Analysis and Interpretation for Vegetable Crops in Florida: HS964/EP081 rev. 10/2012. *Edis*, 10. <https://doi.org/10.32473/edis-ep081-2004>
- Kelland, M. E., Wade, P. W., Lewis, A. L., Taylor, L. L., Sarkar, B., Andrews, M. G., Lomas, M. R., Cotton, T. E. A., Kemp, S. J., James, R. H., Pearce, C. R., Hartley, S. E., Hodson, M. E., Leake, J. R., Banwart, S. A., & Beerling, D. J. (2020). Increased yield and CO₂ sequestration potential with the C₄ cereal Sorghum bicolor cultivated in basaltic rock dust-amended agricultural soil. *Global Change Biology*, 26, 3658–3676. <https://doi.org/10.1111/gcb.15089>
- 595 Klitzke, S., & Lang, F. (2009). Mobilization of Soluble and Dispersible Lead, Arsenic, and Antimony in a Polluted, Organic-rich Soil – Effects of pH Increase and Counterion Valency. *Environmental Quality*, 38, 933–939. <https://doi.org/10.2134/jeq2008.0239>
- 600 Kögel-Knabner, I., Wiesmeier, M., & Mayer, S. (2022). “Mechanisms of soil organic carbon sequestration and implications for management,” in *Understanding and Fostering Soil Carbon Sequestration*, (Issue October). <https://doi.org/10.19103/AS.2022.0106.02>
- Kopsell, D. E., Kopsell, D. A., Sams, C. E., & Casey, T. (2013). Ratio of Calcium to Magnesium influences biomass, elemental accumulations, and pigment concentrations in kale. *Journal of Plant Nutrition*, 36, 2154–2165. <https://doi.org/10.1080/01904167.2013.789108>
- 605 Kushwaha, A., Hans, N., Kumar, S., & Rani, R. (2018). A critical review on speciation, mobilization and toxicity of lead in soil-microbe-plant system and bioremediation strategies. *Ecotoxicology and Environmental Safety*, 147, 1035–1045. <https://doi.org/10.1016/j.ecoenv.2017.09.049>
- Lewis, A. L., Sarkar, B., Wade, P., Kemp, S. J., Hodson, M. E., Taylor, L. L., Yeong, K. L., Davies, K., Nelson, P. N., Bird, M. I., Kantola, I. B., Masters, M. D., DeLucia, E., Leake, J. R., Banwart, S. A., & Beerling, D. J. (2021). Effects of mineralogy, chemistry and physical properties of basalts on carbon capture potential and plant-nutrient element release via enhanced weathering. *Applied Geochemistry*, 132(June), 105023. <https://doi.org/10.1016/j.apgeochem.2021.105023>
- 610 Lindsay, W. L., Schwab, A. P., State, C., & Collins, F. (2008). The chemistry of iron in soils and its availability to plants. *Journal of Plant Nutrition*, 5, 821–840.
- Liping, Li, Kirk G, S., Lirong, Z., Guangtao, L., Weiqin, X., & Guoqiang, X. (2014). Immobilization of Lead in Soil Influenced by Soluble Phosphate and Calcium: Lead Speciation Evidence. *Journal of Environmental Quality*, 474, 468–474. <https://doi.org/10.2134/jeq2013.07.0272>
- Lizárraga-paulín, E. G., Torres-pacheco, I., Moreno-martínez, E., & Miranda-castro, S. P. (2011). Chitosan application in



- maize (*Zea mays*) to counteract the effects of abiotic stress at seedling level. *African Journal of Biotechnology*, *10*,
620 6439–6446. <https://doi.org/10.5897/AJB10.1448>
- Luchese, A. V., Gutz de Castro Leite, I. J., da Silva Giaretta, A. P., Alves, M. L., Pivetta, L. A., & Missio, R. F. (2023). Use
of quarry waste basalt rock powder as a soil remineralizer to grow soybean and maize. *Heliyon*, *9*, e14050.
<https://doi.org/10.1016/j.heliyon.2023.e14050>
- Majeed, S., Reetika, Z., Javaid, M., Muslima, A. B., & Rupesh, N. (2019). Role of silicon in plant stress tolerance :
625 opportunities to achieve a sustainable cropping system. *3 Biotech*, *9*, 1–16. <https://doi.org/10.1007/s13205-019-1613-z>
- Makabe-sasaki, S., & Sasaki, Y. (2015). *Effects of slag silicate fertilizer on silicon content of rice plants grown in paddy fields
on the Shounai Plain , Yamagata , Japan. September 2014.* <https://doi.org/10.1080/00380768.2014.936305>
- Moosdorf, N., Renforth, P., & Hartmann, J. (2014). *Carbon Dioxide Efficiency of Terrestrial Enhanced Weathering.*
- Nagajyoti, P. C., Lee, K. D., & Sreekanth, T. V. M. (2010). Heavy metals, occurrence and toxicity for plants: A review.
630 *Environmental Chemistry Letters*, *8*, 199–216. <https://doi.org/10.1007/s10311-010-0297-8>
- Ning, D., Song, A., Fan, F., Li, Z., & Liang, Y. (2014). Effects of Slag-Based Silicon Fertilizer on Rice Growth and Brown-
Spot Resistance. *PLoS ONE*, *9*, 1–9. <https://doi.org/10.1371/journal.pone.0102681>
- O'Connor, J., Nguyen, T. B. T., Honeyands, T., Monaghan, B., O'Dea, D., Rinklebe, J., Vinu, A., Hoang, S. A., Singh, G.,
Kirkham, M. B., & Bolan, N. (2021). Production, characterisation, utilisation, and beneficial soil application of steel
635 slag: A review. *Journal of Hazardous Materials*, *419*(May). <https://doi.org/10.1016/j.jhazmat.2021.126478>
- Osemwota, I. O. (2007). Effect of Calcium / Magnesium Ratio in Soil on Magnesium Availability , Yield , and Yield
Components of Maize. *Communications in Soil Science and Plant Analysis*, *38*, 2849–2860.
<https://doi.org/10.1080/00103620701663081>
- Pinheiro, J., Bates, D., Debroy, S., Sarka, D., & R Core Team, R. (2013). nlme: Linear and Nonlinear Mixed Effects Models.
640 *R Package Version 3.1*, 109.
- Pistocchi, C., Ragolini, G., Colla, V., Branca, T. A., Tozzini, C., & Romaniello, L. (2017). Exchangeable Sodium Percentage
decrease in saline sodic soil after Basic Oxygen Furnace Slag application in a lysimeter trial. *Journal of Environmental
Management*, *203*, 896–906. <https://doi.org/10.1016/j.jenvman.2017.05.007>
- Qian, P., & Schoenau, J. J. (2002). Practical applications of ion exchange resins in agricultural and environmental soil research.
645 *Canadian Journal of Soil Science*, *82*, 9–21.
- Ramezani, A., Dahlin, A. S., Campbell, C. D., Hillier, S., Mannerstedt-Fogelfors, B., & Öborn, I. (2013). Addition of a
volcanic rockdust to soils has no observable effects on plant yield and nutrient status or on soil microbial activity. *Plant
and Soil*, *367*, 419–436. <https://doi.org/10.1007/s11104-012-1474-2>
- Ramos, C. G., Hower, J. C., Blanco, E., Oliveira, M. L. S., & Theodoro, S. H. (2022). Possibilities of using silicate rock
650 powder: An overview. *Geoscience Frontiers*, *13*(1), 101185. <https://doi.org/10.1016/j.gsf.2021.101185>
- Renforth, P., Washbourne, C. L., Taylder, J., & Manning, D. A. C. (2011). Silicate production and availability for mineral
carbonation. *Environmental Science and Technology*, *45*, 2035–2041. <https://doi.org/10.1021/es103241w>



- Rijnders, J., Vicca, S., Struyf, E., Amann, T., Hartmann, J., Meire, P., Janssens, I., & Schoelynck, J. (2023). The effects of dunite fertilization on growth and elemental composition of barley and wheat differ with dunite grain size and rainfall regimes. *Frontiers in Environmental Science*, *11*(August), 1–18. <https://doi.org/10.3389/fenvs.2023.1172621>
- Roychoudhury, A. (2020). Vanadium Uptake and Toxicity in Plants. *SF Journal of Agricultural and Crop Management*, *12*, 1010.
- Schlüter, U., Mascher, M., Colmsee, C., Scholz, U., Bräutigam, A., Fahnenstich, H., & Sonnewald, U. (2012). Maize source leaf adaptation to nitrogen deficiency affects not only nitrogen and carbon metabolism but also control of phosphate homeostasis. *Plant Physiology*, *160*, 1384–1406. <https://doi.org/10.1104/pp.112.204420>
- Shenker, M., & Chen, Y. (2005). Increasing Iron Availability to Crops: Fertilizers, Organo-Fertilizers, and Biological Approaches. *Soil Science and Plant Nutrition*, *51*, 1–17.
- Skov, K., Wardman, J., Healey, M., McBride, A., Bierowiec, T., Cooper, J., Edeh, I., George, D., Kelland, M. E., Mann, J., Manning, D., Murphy, M. J., Pape, R., Teh, Y. A., Id, W. T., Wade, P., & Liu, X. (2024). Initial agronomic benefits of enhanced weathering using basalt: A study of spring oat in a temperate climate. *PLoS ONE*, *19*, 1–20. <https://doi.org/10.1371/journal.pone.0295031>
- Srekanth, T. V. M., Nagajyothi, P. C., Lee, K. D., & Prasad, T. N. V. K. V. (2013). Occurrence , physiological responses and toxicity of nickel in plants. *International Journal of Environmental Science and Technology*, *10*, 1129–1140. <https://doi.org/10.1007/s13762-013-0245-9>
- Swoboda, P., Döring, T. F., & Hamer, M. (2022). Remineralizing soils? The agricultural usage of silicate rock powders: A review. *Science of the Total Environment*, *807*, 150976. <https://doi.org/10.1016/j.scitotenv.2021.150976>
- Taylor, L. L., Beerling, D. J., Quegan, S., & Banwart, S. A. (2017). Simulating carbon capture by enhanced weathering with croplands: An overview of key processes highlighting areas of future model development. *Biology Letters*, *13*(4). <https://doi.org/10.1098/rsbl.2016.0868>
- Tchouankoue, J. P. (2014). Soil Remineralization Trial: Preliminary Effects of Montserrat Volcanic Ash on Barbuda Limestone soils. *Geotherapy, October*, 391–398. <https://doi.org/10.1201/b13788-27>
- Ten Berge, H. F. M., Meer, H. G. Van Der, Steenhuizen, J. W., Goedhart, P. W., Knops, P., & Verhagen, J. (2012). Olivine Weathering in Soil , and Its Effects on Growth and Nutrient Uptake in Ryegrass (*Lolium perenne* L .): A Pot Experiment. *PLoS ONE*, *7*, 42098. <https://doi.org/10.1371/journal.pone.0042098>
- Van Straaten, P. (2006). Farming with rocks and minerals: Challenges and opportunities. *Anais Da Academia Brasileira de Ciencias*, *78*, 731–747. <https://doi.org/10.1590/S0001-37652006000400009>
- Vienne, A., Poblador, S., Portillo-estrada, M., & Hartmann, J. (2022). Enhanced Weathering Using Basalt Rock Powder : Carbon Sequestration , Co-benefits and Risks in a Mesocosm Study With *Solanum tuberosum*. *Frontiers in Climate*, *May*, 869456. <https://doi.org/10.3389/fclim.2022.869456>
- Walker, J. C. G., Hays, P. B., & Kasting, J. F. (1981). A negative feedback mechanism for the long-term stabilization of Earth’s surface temperature. *Journal of Geophysical Research*, *86*(C10), 9776–9782. <https://doi.org/10.1029/JC086iC10p09776>



- Wang, W., Sardans, J., Lai, D. Y. F., Wang, C., Zeng, C., Tong, C., & Liang, Y. (2015). Field Crops Research Effects of steel slag application on greenhouse gas emissions and crop yield over multiple growing seasons in a subtropical paddy field in China. *Field Crops Research*, *171*, 146–156. <https://doi.org/10.1016/j.fcr.2014.10.014>
- 690 Wang, X., & Qing-sheng, C. (2006). Steel Slag as an Iron Fertilizer for Corn Growth and Soil Improvement in a Pot Experiment *. *Soil Science Society of China*, *16*, 519–524.
- WHO, & FAO. (2022). *General standards for contaminants and toxins in food and feed*.
- Yan, Y., Dong, X., Li, R., Zhang, Y., Yan, S., Guan, X., Yang, Q., Chen, L., Fang, Y., Zhang, W., & Wang, S. (2023). Wollastonite addition stimulates soil organic carbon mineralization: Evidences from 12 land-use types in subtropical
695 China. *Catena*, *225*(November 2022), 107031. <https://doi.org/10.1016/j.catena.2023.107031>
- Yusuf, M., Fariduddin, Q., Hayat, S., & Ahmad, A. (2011). Nickel: An overview of uptake, essentiality and toxicity in plants. *Bulletin of Environmental Contamination and Toxicology*, *86*, 1–17. <https://doi.org/10.1007/s00128-010-0171-1>