

Microbial communities inhabiting 600-year-old sediments in the Inka-Coya Lake located in the Atacama Desert

Coral Pardo-Esté^{1*}; Juan Castro-Severyn^{2,3}; Francisco Remonsellez^{2,3}; Antonio Maldonado^{4,5}; Inger Heine-Fuster⁶; Hector Pizarro⁷; Adriana Aránguiz-Acuña^{5,6*}.

¹ Departamento de Ciencias Farmacéuticas, Facultad de Ciencias, Universidad Católica del Norte, Antofagasta, Chile.

² Laboratorio de Microbiología Aplicada y Extremófilos, Departamento de Ingeniería Química, Universidad Católica del Norte, Antofagasta, Chile.

³ Centro de Investigación Tecnológica del Agua en el Desierto-CEITSAZA, Universidad Católica del Norte, Antofagasta, Chile.

⁴ Centro de Estudios Avanzados en Zonas Áridas, CEAZA. La Serena, Chile.

⁵ Millennium Nucleus AndesPeat, Arica, Chile.

⁶ Laboratorio de Ecología Acuática, Departamento de Recursos Ambientales, Universidad de Tarapacá, Arica, Chile.

⁷ Departamento de Ciencias Geológicas Universidad Católica del Norte, Antofagasta, Chile.

* Correspondence to: Adriana Aránguiz-Acuña (aaranguiza@academicos.uta.cl) / Coral Pardo-Esté (coral.pardo@ce.ucn.cl)

Abstract. Lacustrine sediments are natural archives for the surrounding area's biogeochemical dynamics; in particular, the isolation and extreme conditions in which desert lakes are located make them ideal study models for studying perturbations in the ecosystem. We aimed to study the microbial community dynamics in Inka-Coya Lake, located in the Atacama Desert, where active geological activity and the local mining industry influence biological dynamics in this ecosystem, as suggested for macroinvertebrates and plankton communities in the lake. In this study, we aimed to characterize the microbial communities that inhabit deep lacustrine sediments and their interaction with the surrounding environment. The results show that the microbial community from lacustrine sediments contains over 70% unclassified organisms, highlighting this ecosystem's microbial taxonomic novelty. Our results indicate that the microbial communities cluster in three distinct zones: a superficial community, an intermediate and mixed community, and a more specialized anaerobic community in the deeper sediments. The microbial composition is dominated by chemoheterotrophic bacteria strongly associated with methane metabolism. Additionally, there is statistical evidence of strong correlations between particular taxa such as Sulfurimonadaceae, Metanoregulaceae, and Ktedonobacteraceae with elements like Cu, As, Fe, Ni, and V, and magnetic properties of the surrounding environment, evidencing the strong correlation between the surrounding geochemistry and microbial life that could be disrupted with the continuous mining activity in the area. Further detailed studies of the metabolic repertoire of these communities are necessary to understand the complex dynamics between microbial life and geochemical composition in this fragile and extreme environment.

KEYWORDS: desert lake, deep lacustrine sediment, microbial communities, extremophiles

36 The Atacama Desert is located on the driest part of the South American dry diagonal, which extends from 5°S on the west
37 coast to almost 50°S on the east coast, over 4,000 km with less than 200 mm mean annual rainfall (MAR). It is surrounded by
38 the western slopes of the Central Andes Cordillera between 15 and 30°S at elevations between sea level and 3,500 m a.s.l. The
39 Andes Cordillera represents a physical barrier that directly modulates climatic conditions and water availability in the Atacama
40 Desert (Garreaud et al., 2003). Particular environmental conditions of the Atacama, such as high solar radiation, low
41 atmospheric humidity, and other pressures associated with the natural composition of the desert, directly influence life
42 occurring in these ecosystems (Demergasso et al., 2008; Albarracín et al., 2020; Kurth et al., 2021; Kereszturi et al., 2020;
43 Borsodi et al., 2022). Thus, the Atacama Desert water bodies are truly natural laboratories for understanding evolutionary
44 processes, not only of the geomorphology of the landscape but also of different life forms promoted by environmental forces,
45 such as climate changes at different time scales (Adrian et al., 2009).

46 Aquatic sediments are sources and/or sinks of elements participating in biogeochemical cycles, including both allochthonous
47 and autochthonous lake processes, influencing biodiversity and trophic dynamics of water bodies (Trolle et al., 2010;
48 Fernández et al., 2000; Usenko et al., 2007; Bandowe et al., 2018). Aquatic sediments have an advantage over terrestrial
49 records, and that is that they accumulate at measurable rates because they are often buffered from physical, chemical, and
50 biotic disturbances, thus allowing the recording of past environmental conditions (Benito, 2020; Picard et al., 2022; Da Costa
51 et al., 2023; Yan et al., 2024). A wide variety of abiotic (e.g., bulk density, dry mass, radioactive isotopes, mineralogy, chemical
52 elements) and biotic proxies (e.g., fossils, species abundance, and presence/absence, resting structures, pigments,
53 environmental DNA) preserved in the sediments are currently analyzed to reconstruct ecosystem change at timescales ranging
54 from fine scale (interannual or decadal) to millennial (Cohen, 2003; Korosi et al., 2017).

55 The central Andes water systems have mainly originated after successive glaciations and volcanic and tectonic activity. In the
56 Atacama Desert, lacustrine sediments are natural archives holding evidence of past precipitations, dust deposition, anthropic
57 disturbances, and pollution, mainly due to mining activity that occurs in the area (Grosjean and Veit, 2005; Placzek, 2009;
58 Cerda et al., 2019; Aránguiz-Acuña et al., 2020). Desert lakes are located in extremely arid and isolated areas, making them
59 susceptible to perturbations (Valero-Garcés et al., 2003; Pueyo et al., 2011; Grosjean and Veit, 2005) and hosting extreme
60 forms of life, especially microorganisms that have evolved physiological and life-history adaptations allowing them to thrive
61 in challenging conditions (Dib et al., 2009; Ordoñez et al., 2009; Fariás et al., 2013, 2014; Rasuk et al., 2014). Therefore,
62 lacustrine sediments are expected to contain a great taxonomic diversity, including low-abundance and highly specialized taxa,
63 directly influenced by small-scale conditions determining local environments (Borsodi et al., 2022).

64 Inka-Coya Lake (22°20'S-68°35'W, 2534 m.a.s.l.) is located at the eastern margin of the Atacama Desert, close to the Salado
65 River, in the San Francisco de Chiu-Chiu village, northern Chile. It is a karstic sinkhole developed during the Quaternary
66 period by the dissolution of calcareous layers of the Chiu-Chiu Formation (El Loa Group). The Atacama Desert and its
67 surroundings have a particular geological history; metals and metalloids found in the Inka-Coya area include Ti, Al, Fe, Ni,
68 and Cr; also, As and Sb are associated with the local geological activity (Aszalós et al., 2020; Borsdorf and Stadel, 2015;
69 Pérez-Portilla et al., 2024), that directly influence the chemical composition of the underground water (Vignale et al., 2021).
70 The central Atacama Desert, specifically the Antofagasta Region, holds large porphyry copper deposits that support the great
71 metal-mining industry (Dittmar, 2004; Salvarredy-Aranguren et al., 2008). In 2021, Chile was the world's top copper producer,
72 producing 5,508,084 tons -26.6% of the world's production- (Rodríguez-Luna et al., 2022). The mining industry in the
73 Antofagasta region has developed extensively since the 19th century (Dittmar, 2004; Salvarredy-Aranguren et al., 2008) with
74 increasing impacts on the national economy and development, but also on the health of ecosystems, which have triggered

75 social and environmental conflicts, affecting especially relevant groups such as ancient Indigenous communities from Quechua
76 and Lickan Antay people (see Ramírez et al., 2005; Tapia et al., 2019).

77 Previous studies aimed to assess mining pollution records on environmental matrices have included the sediment records of
78 different longitudes from the Inka-Coya Lake, showing variation through the geochemistry and magnetic properties (Cerde et
79 al., 2019; Aránguiz-Acuña et al., 2020; Pérez-Portilla et al., 2024). Analyzed variables have allowed us to identify episodes
80 associated with changes in water availability, flash flooding, and evidence of perturbations induced by mining activities.
81 Overall, the lake is polluted at different degrees of severity with Cu, Sb, Mo, and As, and some elements like Cu and Ni have
82 been enriched in the most recent periods (Cerde et al., 2019; Pérez-Portilla et al., 2024). Additionally, some biological proxies,
83 such as macroinvertebrates and diatom communities, were found to be directly influenced by the accumulation of metal(loid)
84 as observed by changes in assembly composition (Aránguiz-Acuña et al., 2020). Surrounding metal-mining exploitation,
85 which has been maintained and even increased through the last 200 years, in addition to aridity stable conditions, makes Inka-
86 Coya Lake an excellent site for understanding biological adaptations of aquatic populations to these anthropic pressures
87 (Aránguiz-Acuña et al., 2018; 2020). Regarding microbial communities, previous studies have established that methanogens
88 are abundant in cold desert lakes, and microorganisms inhabiting these lake oases are the basis for many ecological services
89 (Cavicchioli, 2006; Stoeva et al., 2014; Olilo et al., 2023). Additionally, salinity and pH are the major drivers determining
90 microbial community structure in other desert lakes (Casamayor et al., 2013; Banda et al., 2021; Santini et al., 2022). Therefore,
91 it is necessary to establish a baseline regarding the dynamics between the microbial life and geochemical conditions in Inka-
92 Coya Lake as a model for the Atacama Desert.

93 While microbial life in arid ecosystems plays a key role in maintaining biogeochemical cycles (Madsen, 2011), there is a high
94 proportion of unclassified taxa that hold great interest in poly-extreme environments from an ecological, environmental, and
95 biotechnological point of view (Farias et al., 2014; Castro-Severyn et al., 2021; Dong et al., 2022). Nevertheless, there are
96 scarce records in which microbial assemblage has been used as a paleolimnological proxy of the possible responses to long-
97 term sustained anthropogenic metal stress (Da Costa et al., 2023; Yan et al., 2024). The few studies considering this aim have
98 focused on changes in primary producers' abundance, under-interpreting the impact on other metabolic functional groups
99 (Benito et al., 2020; Picard et al., 2022).

100 This study aimed to characterize the microbial community along a lacustrine sediment core, capturing the depositional history
101 of the Inka-Coya Lake over the last 600 years, and contribute to the knowledge of geo-microbial dynamics in the underexplored
102 desert lakes as these microorganisms are the basis for most geochemical cycling of critical nutrients. We hypothesize that the
103 microbial community assemblage of Inka-Coya Lake is strongly associated with sediment attributes shaped by autochthonous
104 and allochthonous processes, particularly anthropogenic contributions from the metal mining industry that has operated near
105 the lake for the past century. To our knowledge, this is a pioneering study in the microbial characterization of a sedimentary
106 core of this length (136 cm) and date (600 years) from a lake sediment in the Atacama Desert. Results show that the microbial
107 communities have changed through time, identifying three clear periods in which alpha and beta diversity has been associated
108 with organic matter content, magnetic susceptibility, and metals and metalloid concentrations. Additional studies of the

109 metabolic functions of the microorganisms inhabiting these sediments are required to understand the interactions between
110 microbial life and the geochemical components of the Inka-Coya Lake further.

111 **1. METHODS**

112 **2.1 Study site and sampling**

113 Inka-Coya Lake (San Francisco de Chiu Chiu village, Antofagasta; 22° 20.300' S; 068°35.981' W, Chile) has a surface area of
114 500 m² and a maximum depth of 18 m, is located in the Pre-Andean Depression of the Antofagasta Region at an elevation of
115 2,520 m a.s.l. (Fig. 1 A, B). Around the lake, the predominant vegetation is of vegas, a type of wetland typical of the Andean
116 pre-Puna zone, strongly associated with the hydric variability of the emerging groundwaters.

117 **2.2 Sediment Core Sampling**

118 A fieldwork campaign was driven in August 2021. The topography of the lake bottom was modeled using the Echo-Map Plus
119 42CV from Garmin. Afterward, three sediment cores from the depocenter of Inka-Coya Lake, where maximum sedimentation
120 rates are expected, were obtained. The cores were obtained using a 9.0 cm diameter Uwitec gravity corer. This study shows
121 the most extended core analysis results, measuring 136 cm (labeled LIC-SHC03). X-ray and photography images were
122 captured before obtaining sections of sediment subsamples from the core. For subsampling, the frozen core was sliced every
123 0.5 cm to a depth of 12 cm; then, every 1 cm until the end of the core, totaling 146 sediment samples. Sediment sub-samples
124 for molecular analysis were first taken from each segment center within a laminar flow hood and using ethanol-sterilized tools.
125 These sub-samples were collected in sterile 15 ml tubes, labeled and kept frozen. Other sub-samples were also taken for the
126 geochemical analyses. Additionally, sub-samples for every 1 cm interval were obtained to develop the magnetic susceptibility
127 analysis. The cores' detailed treatment, geochemical analysis, and magnetic properties can be reviewed by Pérez-Portilla et al.
128 (2024).

129 **2.3 Sediment Core Dating**

130 The geochronology of the sediment core from Inka-Coya Lake was determined through radiocarbon dating (¹⁴C) on the
131 remaining macroscopic carbon along the record. The measurements were done using accelerator mass spectrometry (AMS),
132 and the results were corrected for isotopic fractionation with an unreported $\delta^{13}\text{C}$ value. Subsequently, the age-depth model
133 for this sedimentary core was established using the Bayesian radiocarbon chronology package Bchron in R, using the 'shcal20'
134 as the calibration curve (Hogg et al., 2020; Haslett and Parnell, 2008). A detailed description of the procedure is available in
135 Pérez-Portilla et al. (2024).

136 **2.4 Magnetic and Geochemical Properties Analysis**

137 Five grams of each sediment sub-sample were placed into paleomagnetic boxes of 8 cm³ to measure the mass magnetic
138 susceptibility (χ) using a Kappabridge MFK1_FA instrument (AGICO Co) under environmental conditions (22–24°C) and a
139 magnetic field of 200 A/m. The samples were measured at a low frequency of 976 Hz (χ_{lf} or simply χ) and a high frequency
140 of 15,616 Hz (χ_{hf}). The magnetic susceptibility dependent on the frequency was calculated using both measurements, as

described by Pérez-Portilla et al. (2024). The $\chi_{fd}\%$ parameter is used to indicate the presence of magnetic particles near the limit of the superparamagnetic/single domain (SP/SD) magnetic size (Verosub and Roberts, 1995), which can be linked to the presence of magnetic particles of authigenic origin (Dearing et al., 1996). Additionally, sub-samples of each 1 cm slice were dried in an oven at 50°C. Afterwards, they were homogenized using an agate mortar in the Geochemistry Laboratory of Universidad Católica del Norte (UCN), Antofagasta, Chile. The sediments were then digested using reverse aqua regia (4 mL HCl + 12 mL of HNO₃ + 300 mg of sediment sample) and a microwave digester (Perkin Elmer MPS 320; EPA 3052 method), following Tapia et al. (2022) in Centro de Investigación Tecnológica del Agua en el Desierto (CEITSAZA-UCN). The elements aluminum (Al), titanium (Ti), vanadium (V), manganese (Mn), iron (Fe), nickel (Ni), Cu, zinc (Zn), arsenic (As), molybdenum (Mo), and antimony (Sb) were measured by the inductively coupled plasma atomic emission spectroscopy (ICP-OES) Perkin Elmer Optima 7000 in the digested residue at CEITSAZA. The organic, inorganic matter, and carbonate contents were estimated using the loss on ignition (LOI) method, which was assessed at a contiguous 1 cm interval following Heiri et al. (2001). This procedure involved drying 1 cm³ of each sediment sample in crucibles at 105°C for two hours and weighing them. The dry samples were weighed before heating to 550°C in a flask over 1.5 h, left at 550°C for two h, then allowed to cool. The samples were weighed, then the crucibles were transferred to the flask and burned at 925°C. Finally, the crucibles were weighed again once they cooled.

2.5 Sediment sample processing and DNA extraction

Sediment samples were obtained every 1 cm from the top to the bottom of the core under sterile conditions within a laminar flow hood and using ethanol-sterilized tools. According to the manufacturer's instructions, total DNA was extracted from the 250 mg of sediment samples using the DNeasy PowerSoil kit (Qiagen Inc., Hilden, Germany). DNA integrity, quality, and quantity were verified by 1% agarose gel electrophoresis, the 260/280nm ratio and fluorescence using a Qubit4 fluorometer and the Qubit dsDNA HS assay kit (Thermo Fisher Scientific, MA, USA). Following, DNA samples were sent to AustralOmics, Chile, for amplification of the bacterial 16S rRNA gene V4 region (~450 bp) using tBakt_341F and Bakt_805R primers (Herlemann et al., 2011), construction of 250 bp paired-end libraries and sequencing on a MiSeq (Illumina) platform.

2.6 Taxonomic Composition Analysis

This analysis was conducted in R v4.0.3 and RStudio v1.3.1093 following the DADA2 v1.16.0 R package pipeline (Callahan et al., 2016) to infer amplicon sequence variants (ASVs) for each sub-sample. Briefly, the reads were evaluated for quality control and subsequently trimmed ($Ns = 0$, $length \geq 150$ bp, $expected\ errors \leq 2$), followed by dereplication, denoising, and merging of paired reads. Following, an ASV table was built to allow a maximum of two expected errors, removing chimeras and assigning taxonomy using the Silva v138 database (Quast et al., 2012). Also, all ASVs identified as Eukarya, Chloroplast, and Mitochondria were removed. In addition, ASVs with very low abundance ($<0.0005\%$) and those only observed in few samples ($<10\%$) were eliminated; Samples with very few reads (<1000) were eliminated as well. Next, a multi-sequence alignment was created to infer phylogeny using FastTree v2.1.10 (Price et al., 2009), and phyloseq-object (containing the ASVs, taxonomy assignment, phylogenetic tree, and the samples meta-data) was created using the R package Phyloseq v1.34.0 (McMurdie et al., 2013) and the variance stabilizing transformation was used for normalization. Finally, taxa relative

175 abundance and taxonomic composition at different ranks were visualized using the ggplot2 v3.3.3 (Wickham, 2016), Fantaxtic
176 v0.2.0 (Teunisse, 2022), and ampvis2 v2.7.4 (Andersen et al., 2018) R packages.

177 **2.7 Diversity Analysis**

178 Alpha diversity metrics (Shannon, Chao, phylogenetic diversity, and Simpson indexes) were calculated for each segment along
179 the core using the microbiome v1.24.0 (Lahti et al., 2017) and btools v0.0.1 R packages. Also, Wilcoxon statistical tests to
180 compare means with Bonferroni correction were carried out between the identified zones and visualized using the DESeq2
181 v1.42.0 (Love et al., 2014) and ggpubr v0.6.0 (Kassambara, 2017) packages. Moreover, beta diversity was evaluated by
182 principal coordinates analysis using Hellinger transformed Bray Curtis distances based on the ASV abundance matrix were
183 calculated using Phyloseq v1.34.0 (McMurdie et al., 2013) and ampvis2 v2.4.5 (Andersen et al., 2018) R package. Also,
184 redundancy analysis (RDAs) was calculated using depth gradient and zone parameters to constrain the multivariate space, and
185 ANOVA tested the statistical significance of the selected geochemical variables.


186 **2.8 Functional Predictions**

187 Functional potential signatures and metabolic pathways abundances were inferred based on the ASV abundance and taxonomy
188 matrices using PICRUST2 v2.4.1 software (Douglas et al., 2020) with the Kyoto Encyclopedia of Genes and Genomes (KEGG)
189 (Kanehisa et al., 2012) and MetaCyc (Caspi et al., 2018) pathway databases. We used the three identified zones along the core
190 that present significant differences at the of taxonomic composition level and then Differential Abundance of inferred pathways
191 was tested with the Kruskal-Wallis test (confidence interval = 0.95) and the Benjamini-Hochberg correction false-discovery
192 rate using ggpicrust2 v1.7.2 R package (Yang et al., 2023). Also, we use the Functional Annotation of Prokaryotic Taxa
193 (FAPROTAX) database v1.2.7 (Louca et al., 2016) to map the identified ASVs and quantify changes in established
194 ecologically relevant functions.

195 **1. RESULTS**

196 The samples analyzed correspond to a sediment core from Lake Inka-Coya, located in the Atacama Desert, with a water depth
197 of 18.5 meters (Fig. 1). The sediment core age-depth model was constructed based on six charcoal sample dates, where the
198 more superficial at 41 cm corresponds to 75 ± 32 cal years BP, and the deepest found at 94 cm corresponds to an age value of

199 505 ± 22 cal years BP. Based on the age-depth model constructed, the sediment core of Inka-Coya Lake analyzed had 630
200 years of age (Fig. 2).



201
202 **Figure 1. Sampling site location in Northern Chile (A), Inka-Coya Lake, and important surrounding mining and urban**
203 **centers (B) and the bathymetry of the lake (C).**




Figure 2. Diversity of the microbial community of the Inka-Coya sediment core. The core sedimentary radiography, dating, and alpha diversity indices variation along the core depth are displayed.

The variation in diversity within the sample was measured to determine the changes in the microbial community along the sediment core by calculating the Shannon, Simpson, Chao, and Phylogenetic indices as standard measures of the taxonomic diversity within a sample (Thukral, 2017). The microbial community observed along the core was diverse, based on the DNA samples analyzed. Quantification of diversity showed that it increased with depth, and the Simpson index remained stable after the 2 cm surface layer. At the same time, Chao1 considered low-abundance taxa, and the phylogenetic index based on the

212 phylogenetic history of the species (Fig. 2). The diversity increases in all cases at two cm long, and maximum values were
213 observed between 8 and 86 cm, with minor peaks at 96, 120, and 130 cm near the bottom samples.

214 Three distinct zones in terms of microbial taxonomy could be identified and are statistically different for the four evaluated
215 diversity indices, except between Zone II and III in the context of phylogenetic diversity (Fig. 3A). These three disjunct
216 clustering zones of microbial community diversity along the sediment core were also identified in the beta diversity analysis,
217 where zone I includes the less diverse upper layer (0-8 cm), zone II, the middle zone of the core (9-42 cm) with significant

greater diversity, and zone III the deepest are sampled in this study, depicted in three shades of green going from lighter (zone I) to darker (zone III) in the PCoA clustering analysis (Fig. 3B).




Figure 3. Clustering of the microbial communities inhabiting the deep sediment of Inka-Coya Lake A) Statistical differences in alpha diversity between depth zones. B) Principal Coordinates Analysis (PCoA) with the ASVs relative abundance using Bray-Curtis as distance metric; each point corresponds to a community, tagged by depth and colored by zone.

The taxonomic composition and abundance along the communities at phylum rank also reflect the clustering on three distinct zones, where zone I is dominated by Actinobacteriota and includes a great abundance of Firmicutes in the top layers, and Campylobacterota, there is also the presence of Bacteroidota, Halobacteriota, and Plantomycetota. Zone II is more diverse and

228 composed mainly of Campylobacterota, Chloroflexi, Acidobacteriota, and Actinobacteriota. While zone III is the largest and
229 more homogeneous, composed of several low-abundance taxa, dominated by Chloroflexi, Acidobacteriota, and
230 Actinobacteriota, there is also a higher representation of Crenarchaeota, Nitrospira, Aenigmarchaeota, and Armatimonadota
231 that in the rest of the zones (Fig. 4). The taxonomic composition along the sediment core evidenced the metabolic adaptations
232 that occur in each layer that presents endogenous characteristics, for instance the upper layer includes generalist groups
233 including microorganisms would break down complex organic groups and light driven-metabolic pathways as well as have
234 the metabolic flexibility that would allow the use of different electron acceptors for energy production, supporting
235 heterotrophic communities and mutualistic dynamics. On the other hand, zone III includes more adaptable and ancient
236 phylogenetic lineages that also have the ability to maintain geochemical cycles (C, N, S, Fe) in low oxygen or anaerobic
237 conditions, as those found in early Earth.

238 Notably, 76.6% of the taxa could not be identified at the genus level (0% matched any known species). Thus, Figure 5 shows
239 the abundance at the “best hit,” where Campylobacterota (*Sulfuricurvum*, *Sulfurimonas*), *Mycobacterium*, and *Methanolinea*
240 dominated the overall community, where many representatives of chemo-lithotrophic, S-oxidizing and nitrate reducing
241 capabilities would thrive. While in each zone, there are particular taxa associated; for instance, species belonging to the
242 Aminicenantes Phyla are very common in Zone I; *Pseudarcobacter* is prevalent in Zone II as the anaerobic bacterium
243 *Pelolinea* is in Zone III.

244




Figure 4. Taxonomic composition of the microbial community in the deep sediments of Inka-Coya Lake. Stacked bar of the taxonomic composition at the phylum level.




Figure 5. Heatmap of the abundance of the microbial community at the family level. The color gradient indicates the abundance of the specific taxa.




Figure 6. Depth variation of metals and metalloids concentrations (mg·kg⁻¹), organic matter (OM) and inorganic concentrations (g·cm⁻³), clasts percent, and magnetic susceptibility (MS) measured in Inka-Coya Lake sediment core (modified from Pérez-Portilla et al. 2024).

Variations in analyzed sediment properties along the core, such as magnetic susceptibility, organic matter, and carbonates/clasts composition, are shown in Figure 6. Copper (Cu) and nickel (Ni) were in lower concentration and variability at greater depths and showed concentration peaks in surface sediments. Elements such as iron (Fe), molybdenum (Mo), and vanadium (V) also showed top sediment peaks. Still, overall, they had more variable behavior than previously mentioned elements in the middle and bottom sediments (> 40 cm). Metalloids, arsenic (As), and antimony (Sb) exhibited the highest concentrations between 10-45 cm depth.

The mean organic matter and carbonate contents are around 8.7% and 19.5%, respectively, while the inorganic density (91%) showed the highest averaged values (0.71 g·cm⁻³). The sediment composition shifts to clay and silt from 28 cm to the top of the core. The content of clasts was predominant in the inorganic fraction, with 71.8% along the core. The carbonate peaks were observed at 10-36 cm, 46-56, and more significant and variable proportions below 96 cm, which did not exceed 40%.

Magnetic susceptibility (MS, χ) values range from -6.09×10^{-9} to $8.13 \times 10^{-7} \text{ m}^3 \cdot \text{kg}^{-1}$, with an average value of $2.77 \times 10^{-7} \text{ m}^3 \cdot \text{kg}^{-1}$. Frequency-dependent susceptibility ($\chi_{fd}\%$) values range between 1.31 and 10.17%, with an average of 5.62%. Zone I has

the highest values of χ and the lowest values of $\chi_{fd}\%$, while Zone II shows the lowest χ values and intermediate $\chi_{fd}\%$ values. Zone III presents intermediate χ values and the highest $\chi_{fd}\%$ values.

Geochemical and magnetic variables are associated with microbial diversity found in the sediment of Inka-Coya Lakes, differentiated into zones (I, II, and III). Deeper and older fractions of sediments (dark blue), especially Aminicenantaes, are positively influenced by the magnetic susceptibility and inorganic elements in the sediments. Microbial assemblage found in the middle sediments (Zone II) of the core is driven by organic matter content and water availability, where taxa like *Sulfurimonas*, *Sulfuricurvum*, and *Dehalococcoidia* were the most represented. The middle zone is associated with a significant As peak, which suggests that the presence of metal(loid)positively affects the microorganisms assemblage inhabiting middle-to-superficial layers. In Zone II, microbial diversity is mainly associated with low but stable concentrations of organic matter, a more significant proportion of clasts, and the higher peak of As and Sb in the sediments, where χ values decreased. Upper Zone I is mainly characterized by metal enrichment, with elevated concentrations of Cu, Zn, Ni, Fe, and Mo, among other elements. These peaks correlate with high χ values (Fig. 7).




Figure 7. Redundancy analysis on Hellinger transformed Bray-Curtis distances (corrected by unobserved species) for the microbial communities along the core distance. A) Influence of physicochemical parameters and B) Elemental composition. Depth gradient and Zone parameters were chosen to constrain the multivariate space in a supervised approach. Each axis in the graph shows the percentage of variance explained in an unsupervised and supervised analysis.

There is statistical significance between some key taxa and the physicochemical and elemental composition along the sediment gradient in Inca-Coya Lake; for instance, *Methanoregulaceae*, *Ktedonobacteriaceae*, and *Sulfurimonadaceae* are some of the taxa with the strongest correlation to Cu, Fe, Ni and V presence while zones II and III are the most influenced by these dynamics (Supplementary Fig. 1).

Regarding metabolic approximation, chemoheterotrophy is the most abundant function in all three zones, while aerobic chemoheterotrophs are the most prevalent in zone II. Still, chemoheterotrophs (including several electron acceptors) thrive in

zones II and III (Supplementary Fig. 2). Other functions, such as Methanogenesis, were abundant only in deep sediments (zone III). Moreover, as expected, phototrophy and photoautotrophy were present only in low abundance in zone I, where little light could reach the community (Supplementary Fig. 2). Methane metabolism is very relevant in all three zones, especially in zone III, where acetate is the primary source for this pathway (Supplementary Fig. 2). Other relevant forms of energy transformation are Nitrate reduction VI (assimilation) and starch degradation in zone II. Nitrotoluene degradation and biosynthesis of unsaturated fatty acids are also crucial in the benthic microbial community as a whole (Supplementary Fig. 2). A somewhat homogeneous prediction for metabolic ability regarding energy production among microorganisms inhabiting in sediments over than one meter deep in the Inka-Coya Lake sediments, where geochemical and magnetic dynamics directly influence microbial activities.

4. DISCUSSION

Microbial dynamics along the length of the sediment core of Inka-Coya Lake are tightly associated with analyzed sediment attributes, such as metal(loid)s concentrations and $\chi_{fd}\%$, organic compounds- and water- availability. The geochemical characteristics of the area surrounding Inka-Coya Lake, where active volcanic activity results in the enrichment of elements such as arsenic, sulfur, copper, and others (Romero et al., 2003; Tapia et al., 2018), suggests that microbial assemblages inhabiting the Atacama area (1) have showed structure changes during the last 600 years and (2) have developed broad tolerance range to this potential toxic compounds. Microbial biomarkers serve as criteria to assess anthropogenic impact (Yan et al., 2024), and microorganisms can alter the speciation and bioavailability of meta(oids) in an ecosystem (Niu et al., 2020).

Five stratigraphic zones in the sedimentary core in Inka-Coya were defined from the sediment core here analyzed by Pérez-Portilla et al. (2024). In this sedimentary core, the concentration of rock-forming elements such as Cr, Zn, and V are found in concentrations as expected for the Atacama Desert; while Cu, Mo, Sb and As are higher than expected, suggesting influence from nearby mining activities (Pérez-Portilla et al., 2024). Copper production in the region generates by-products such as Mo, As, and, to a lesser extent, Zn (Ramírez et al., 2005; Tapia et al., 2019). Also, mining wastes contain high concentrations of chemical products such as Pb, Cr, Cd, Cu, Zn, Hg, and Ni, and metalloids, which are often stored in dams or reservoirs (Csavina et al., 2012) or passed through lotic systems, making them an important source of contamination of inorganic chemical elements for the aquatic biological communities (Keller et al., 1992; Pollard et al., 2003; Pigati et al., 2011; Hamilton et al., 2017; Ritter et al., 2019). Previous studies have shown impacts and metal enrichment from the mining industry in the Antofagasta region surrounding Calama City and the Loa River basin near the Inka-Coya Lake. Cerda et al. (2019), Vargas-Machuca et al. (2021), Aránguiz-Acuña et al. (2020), and Zanetta-Colombo et al. (2022, 2024), using both abiotic and biological proxies, have evidenced an increase in the concentration of metals in different environmental matrices during post-industrial time, attributing this difference to the mining activities in the area. Additionally, changes in the composition of the zooplankton community (inferred by diapausing egg banks) and benthic diatoms could be attributed to the increase in Cu concentration evidenced in the sedimentary cores obtained in this lake (Aránguiz-Acuña et al. 2020). It is therefore to be expected that other components of the aquatic community, which are as or more sensitive than planktonic invertebrates to changes in metal concentrations in the environment, may also be affected by the impacts of increases in metals associated with the development of the mining industry.

A strong correlation between mineral composition and microbial diversity in other arid region water bodies, such as salt flats and brines, is expected (Fariás et al., 2014; Castro-Severyn et al., 2021; Dong et al., 2022), as demonstrated in pre-Puna salt lakes, such as Tebenchique and La Brava (Fariás et al., 2014; Ramos-Tapia et al., 2023). The diversity of microbial life in these shallow salty lakes is dominated by Bacteroidetes, Proteobacteria, and Euryarchaeota (Fariás et al., 2014; Fernandez et al., 2016; Kurth et al., 2021) and hypersaline lakes are mainly composed by Bacteroidetes, Chloroflexi, Cyanobacteria and Proteobacteria (Dorador et al., 2018). Inka-Coya, one of the few brackish water lakes located in the Antofagasta Region below 3,000 m a.s.l., and its sediment communities are dominated by Phylum Acidobacteriota, Chloroflexi (*Pelolinea*, among others),

Actinobacteriota (*Mycobacterium*), and Campylobacterota (*Sulfuricurvum* and *Sulfurimonas*), sharing some similarities and considering changes in taxonomy (Oren and Garrity, 2021). Overall, desert lake sediment microbial communities are very diverse, accounting for the variable conditions and different niches that microorganisms can utilize. In this study, we found great taxonomic diversity and potential metabolic richness where several electron acceptors would sustain the community, including complex carbon complexes, nitrogen, sulfur, and iron, each in different oxidation states. Heterotrophic and autotrophic aerobes and anaerobic representatives were found along the length of the sediment core, indicating that mutualistic relationships are vital for the maintenance of the ecosystem and ecological functions.

At the lower taxonomic rank, the community is dominated by microorganisms with a broad repertoire for mineral interactions, e.g., there is experimental evidence of organomineralization in extracellular S⁰ formation by a species of the sulfur-oxidizing bacteria *Sulfuricurvum* (Cron et al., 2019). Another remarkable microorganism found along the lacustrine sediment is the cosmopolitan and highly diverse *Sulfurimonas*, which can grow using sulfur, hydrogen, nitrogen, oxygen, and organic compounds, suggesting it is critical in maintaining trophic dynamics (Han and Perner, 2015). Additionally, *Mycobacterium* is a saprophytic bacterium commonly found in lakes, rivers, and other water sources (Falkinham et al., 2015), there are some species representatives of this genus that have bioremediation potential for polycyclic aromatic hydrocarbons (Deng et al., 2023), suggesting adaptability and a broad range of metabolic capacities. There are extensive studies on Mycobacteriaceae representatives that cause human disease, as reviewed by Falkinham (2009). However, environmental representatives with the capacity to inhabit poly-extreme environments have yet to be determined.

Furthermore, several representatives of the candidate phylum Aminicenantes are common in current sediments from Zone I, i.e., recently deposited sediments during the last ten years. So far, these microorganisms are associated with a fermentative saccharolytic lifestyle that does not have an isolated representative yet (Kadnikov et al., 2019). Zone II, comprising the period between 10 and 50 years ago approximately, is mainly represented by microorganisms classified as the *Pseudarcobacter* genus, that was recently separated from the *Arcobacter* genus (Pérez-Cataluña et al., 2018) and are characterized as mesophilic bacteria that can grow in microaerophilic conditions (Collado et al., 2011). Finally, in the deeper and older sediments, where anaerobic (or facultative anaerobes) microorganisms can thrive, there is a particular abundance of an ASV from the *Pelolinea* genus that has only one described species that was isolated from the subseafloor sediment (Imachi et al., 2014), and an unknown Chloroflexi species that is associated with Dehalococcoidia class a common sub-seafloor bacterium (Wasmund et al., 2014). This finding suggests past conditions of higher salinities for the lake than the current conditions of around 5 g·L⁻¹ (Aránguiz-Acuña et al., 2020), close to marine salinity. Archaea representatives found inhabiting the lacustrine sediments include the hydrogenotrophic methanogen *Methanolinea* (Imachi et al., 2008; Rainey et al., 2015), which is distributed along the length of the core with an abundance ranging from 0.7-2.3% and is very abundant towards the older and more anoxic environment (since 1400 until 1950, zone III), where methanogenesis is the central predicted metabolism at play. These results suggest that the microbial community in Inka-Coya Lake is potentially heterotrophic with a special enrichment in methanogenic organisms in the oldest deposits, where oxygen levels are lower as evidenced by the metabolic approximation done in this study. Another kind of metabolism, depending on CO₂ concentrations, could have dominated during this period.

Overall, our results suggest a great taxonomic and potential metabolic diversity is associated with the microbial community from this lacustrine sediment. There is a vast taxonomic novelty harbored in Inka-Coya Lake sediment; over 70% of taxa cannot be identified to the genus level, indicating there is a significant amount of “microbial dark matter” a term associated with unknown microbial representatives that can potentially harbor novel bioactive compounds with numerous applications (Zha et al., 2022; Jiao et al., 2021). This microbial dark matter is also associated with a great metabolic diversity, specially for the production of bioactive compounds with biotechnological potential, for example, we determined that there are abundant representatives of Actinobacteriota, a taxonomic group traditionally associated with the production of antimicrobial, antitumoral and anti-inflammatory compounds (Al-Shaibani et al., 2021; Taj and Chattopadhyay, 2024), thus we hypothesize

375 there is a high probability of finding novel compounds produced by microorganisms inhabiting the sediment of Inka Coya
376 Lake.

377 The main drivers for microbial community composition in the sinks-Coya sediments were As, Sb, V, Mo, Mi, Fe, and Cu,
378 which suggest that there are numerous strategies that microorganisms use to resist high concentrations of metals and metalloids
379 that thrive in this ecosystem (Rahman, 2020; Mathivanan et al., 2021), as observed in the Atacama and Altiplano area (Orellana
380 et al., 2018; Donati et al., 2019; Aszalós et al., 2020; Castro-Severyn et al., 2019). Additionally, microorganisms can use oxido-
381 reduction processes to obtain energy from metal ions (Raab and Feldman, 2003; Staicu and Stolz, 2021). Given the known
382 geochemical characteristics of the area, it is expected to find strong relationships between the microbial life and inorganic
383 compounds, as they can dissolve and precipitate ores and influence metal(loid)s transformations (Raab et al., 2003; Zhou et
384 al., 2022). In this extreme environment where competition is strong and abiotic pressures are constant, organic matter and
385 water availability -both parameters critical for most life forms- govern community abundance and composition, suggesting a
386 delicate dynamic balance reached between abiotic and biotic entities at play. It is important to remark that with the number of
387 unclassified taxa, many novel resistant or usage mechanisms remain to be characterized.

388 Observed trends in magnetic susceptibility in Inka-Coya sediment could be mainly attributed to variations in the concentration
389 of ferromagnetic minerals, such as titanomagnetite, and authigenic origin minerals, such as sulfide (greigite), as is broadly
390 explained in Pérez-Portilla et al. (2024) for this sediment core. In this case, the high X values would result from a high
391 concentration of ferromagnetic minerals of detrital origin, primarily Fe oxides, while elevated $\chi_{fd}\%$ values would be linked to
392 the presence of greigite of authigenic origin (Pérez-Portilla et al., 2014). Thus, greigite formation typically occurs through the
393 dissolution of titanomagnetite or other detrital minerals containing Fe (e.g., Chan et al., 2001; Fialová et al., 2006; Versteeg et
394 al., 1995).

395 Elevated magnetic susceptibility values in the upper layers of sediments may be attributed to a recent deposition of fine Fe
396 oxide grains, which could be originating and transported from industrial and urban sources (e.g., Chan et al., 2001); this concurs
397 with the Fe top sediment peak, and it could be related to a diverse superficial community, while in zone II the lowest values
398 of χ are observed, which could be associated with a mixed community. All these processes occur under a high production of
399 Fe minerals of authigenic origin ($\chi_{fd}\% > 3\%$; Dearing et al., 1996). Moreover, lower levels of $\chi_{fd}\%$ could be involved in the
400 production and assimilation of iron sulfides such as greigite (Bazylinski et al., 2001; Lins et al., 2007), promoting a microbial
401 specialization and increasing resistance of the anaerobic community found in the deeper sediments. Microbial metabolic
402 responses could satisfactorily support several processes associated with greigite formation. The presence of greigite in the lake
403 sediments could be associated with reducing or low-oxygen environments (e.g., Benning et al., 2000), where additionally
404 magnetotactic bacteria could contribute to sedimentary greigite formation through the biomineralization of magnetosomes in
405 anoxic aqueous environments (Moskowitz et al., 2008).

406 There are statistically significant correlations between some particular taxa and the geochemical composition along the
407 sediment gradient in Inca-Coya Lake suggesting that each element directly influences the metabolic capabilities of the
408 microorganisms and shapes the community selecting taxa that can resist metal(loid)s toxicity (Yao and Gao, 2007; Laplante
409 et al., 2013; Stankevica et al., 2020; Kostka and Leśniak, 2021; Yan et al., 2020). However, this study is restricted to metabolic
410 approximations based on taxonomy from a single sediment core. Therefore, it would be desirable to carry out future studies
411 including metagenomic data to explore functional potential and investigate other water bodies to identify temporal patterns of
412 microbiological assemblages in the most exploited area of the Atacama Desert, which will enhance our understanding in terms


413 of the mechanisms used to resist and survive in environments with high metal(loid)s concentrations and the use of different
414 electron acceptors for energy production.

415 Finally, increasing metal(loid)s exploitation in the region during the last 100 years has directly influenced their mobility and
416 the local geochemistry. In this context, microorganisms from these extreme environments are known to be highly adaptive and
417 have developed several resistance mechanisms and the ability to use these compounds to their benefit. Hence, some bacteria
418 (and their genes) can be used as biomarkers for the bioavailability of such metals and contamination of soils (Li and Wong,
419 2010; Roosa et al., 2014). In Inka-Coya Lake, a selective process could be evidenced along the sediment record, in which the
420 recent period is characterized by a marked increase in chemical elements and microbial composition, which could be associated
421 with the increasing mining activity and other anthropological activities, as water extraction or aridity increase by climatic
422 changes, that also would increase the disturbance of this relevant area in the core of the most arid non-polar Desert.

424 This study is the first to describe a deep gradient of microbial life in a desert lake in the Atacama area, proposing a biological
425 clustering of taxa and predicted function in three periods that stratified for over 600 years, including pre-mining period, the
426 mining development and the most industrialized mega mining observed nowadays. A great taxonomic novelty exists among
427 the microbial community inhabiting lacustrine sediments of Inka-Coya and mineralogical enrichment, water, organic matter
428 availability, and magnetic susceptibility are variables that explain the changes in its abundance and composition. There are
429 strong relationships between geochemical composition and microbial diversity, especially in Cu, Fe, Ni, and V. The first zone
430 is less diverse and dominated by Actinobacteria; the second zone has a high abundance of Chloroflexi, Acidobacteriota, and
431 Actinobacterota. The third zone shows more rare taxa with lower abundance and clusters the more recent sediments closer to
432 the surface, including archaea. Overall, chemoheterotrophy is the prevalent energy production mechanism along the sediment
433 core. This unique and fragile ecosystem depends on biogeochemical dynamics that are vulnerable to anthropogenic activities
434 and climate change.

435 Appendix A.

436



437
438 **Figure A1.** Pearson correlation between the top microbial families and geochemical parameters. Asterisks show the level of
439 significance (* p value < 0.05, ** p value < 0.01, Pearson correlation). Purple, blue, and white indicate positive, negative, and
440 no correlation, respectively.

441

442 Appendix B




Figure B1. Metabolic pathways prediction of the microbial community inhabiting sediments along a depth gradient in Inka-Coya Lake. The predictions are based on the identified taxonomic composition according to the comparison with different databases, such as: A) FAPROTAX; B) KEGG and MetaCyc. Categories with significant differences ($p < 0.05$) according to Kruskal-Wallis test are displayed.

DATA AVAILABILITY STATEMENT

The raw sequencing data presented in this study have been deposited in the DDBJ/ENA/GenBank SRA database under the BioProject: PRJNA1067596.

AUTHOR CONTRIBUTIONS

Conceptualization: AAA; FR, JCS, CPE. Data curation: CPE, JCS, FR, IHF, AAA. Formal analysis: CPE, JCS, FR, AAA, AM, HP. Funding acquisition: FR, AAA. Methodology: CPE, JCS, AM, HP, FR, AAA. Supervision: AAA, FR. Writing original draft: CPE, JCS. Writing review and editing: AAA, AM, HP, JCS, CPE. All authors have read and agreed to the published version of the manuscript.

FUNDING

ANID 2020 FONDECYT Regular 1200423 (AA-A; FR), UTA-Mayor 9738-24 (AA-A)
 ANID 2022 FONDECYT Regular 1220902 (FR)
 ANID 2023 FONDECYT postdoctoral 3230189 (CPE)

463 ANID 2021 FONDECYT postdoctoral 3210156, ANID-Anillo ATE 220007 (JCS)
464 ANID 2022 MILENIO-NCS2022_009 (AA-A, AM)
465

466 **COMPETING INTERESTS**

467 The contact author has declared that none of the authors has any competing interests.
468

469 **ACKNOWLEDGMENTS**

470 Likan Antay Community of San Francisco de Chiu-Chiu.

471 MAINI-UCN (Kappabridge)

472 **References**

- 473
- 474 1. Adrian R, O'Reilly CM, Zagarese H, Baines SB, Hessen DO, Keller W, Livingstone DM, Sommaruga R, Straile D,
475 van Donk E, Weyhenmeyer GA, Winder M.: Lakes as sentinels of climate change. *Limnol Oceanogr.*, 54:2283–2297,
476 2009
 - 477 2. Albarracín, V. H., Galván, F. S., and Farías, M. E.: Extreme microbiology at Laguna Socompa: A high-altitude
478 Andean lake (3570 m asl) in Salta, Argentina. *Microbial Ecosystems in Central Andes Extreme Environments:*
479 *Biofilms, Microbial Mats, Microbialites and Endoevaporites*, 205-220, 2020
 - 480 3. Andersen KS, Kirkegaard RH, Karst SM, Albertsen M.: ampvis2: an R package to analyse and visualise 16S rRNA
481 amplicon data. *BioRxiv*. 2018
 - 482 4. Al-Shaibani, M. M., Radin Mohamed, R. M. S., Sidik, N. M., Enshasy, H. A. E., Al-Gheethi, A., Noman, E., ... &
483 Zin, N. M. (2021). Biodiversity of secondary metabolites compounds isolated from phylum actinobacteria and its
484 therapeutic applications. *Molecules*, 26(15), 4504.
 - 485 5. Aránguiz-Acuña, A., Luque, J. A., Pizarro, H., Cerda, M., Heine-Fuster, I., Valdés, J., ... and Wennrich, V.: Aquatic
486 community structure as sentinel of recent environmental changes unraveled from lake sedimentary records from the
487 Atacama Desert, Chile. *PLoS One*, 15(2), e0229453, 2020
 - 488 6. Aránguiz-Acuña, A., Pérez-Portilla, P., De la Fuente, A., and Fontaneto, D.: Life-history strategies in zooplankton
489 promote coexistence of competitors in extreme environments with high metal content. *Sci Rep*, 8(1), 11060, 2018
 - 490 7. Aszalós, J. M., Szabó, A., Felföldi, T., Jurecska, L., Nagy, B., and Borsodi, A. K.: Effects of active volcanism on
491 bacterial communities in the highest-altitude crater lake of Ojos del Salado (Dry Andes, Altiplano-Atacama Region).
492 *Astrobiology*, 20(6), 741-753, 2020
 - 493 8. Banda, J. F., Zhang, Q., Ma, L., Pei, L., Du, Z., Hao, C., & Dong, H. (2021). Both pH and salinity shape the microbial
494 communities of the lakes in Badain Jaran Desert, NW China. *Science of the Total Environment*, 791, 148108.

9. Bandowe FL, Grosjean M, Tylmann W, Mosquera P, Hampel H, Schneider T.: A 150-year record of polycyclic aromatic compound (PAC) deposition from high Andean Cajas National Park, southern Ecuador. *STOTEN*, 621:1652–1663, 2018
10. Bazylinski, D. A.: Bacterial mineralization. *Encyclopedia of materials: Science and technology*, 441-447, 2001
11. Benito, X. Benthic Foraminifera and diatoms as ecological indicators. *Modern Trends in Diatom Identification: Fundamentals and Applications*, 257-280, 2020
12. 10. Benning, L. G., Wilkin, R. T., and Barnes, H. L.: Reaction pathways in the Fe–S system below 100 C. *Chem Geol.*, 167(1-2), 25-51, 2000
13. Borsdorf, A., and Stadel, C.: *The Andes: A geographical portrait*. Springer, 2015
14. Borsodi, A. K., Aszalós, J. M., Megyes, M., and Nagy, B.: Benthic Bacterial Diversity of High-Altitude Athalassohaline Lakes of the Puna de Atacama (Central Andes). *Geomicrobiol J*, 39(1), 28-38, 2022
15. Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A., Holmes, S.P.: DADA2: high-resolution sample inference from Illumina amplicon data. *Nat Methods*. 13:581–583. doi: 10.1038/nmeth.3869, 2016
16. Casamayor, E. O., Triadó-Margarit, X., & Castañeda, C. (2013). Microbial biodiversity in saline shallow lakes of the Monegros Desert, Spain. *FEMS microbiology ecology*, 85(3), 503-518.
17. Cavicchioli, R. (2006). Cold-adapted archaea. *Nature Reviews Microbiology*, 4(5), 331-343.
18. Herlemann, D. P., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J. J., & Andersson, A. F.: Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea: The ISME journal. 5(10), 1571-1579.
19. Caspi, R., Billington, R., Fulcher, C. A., Keseler, I. M., Kothari, A., Krummenacker, M., ... and Karp, P. D.: The MetaCyc database of metabolic pathways and enzymes. *Nuc acids Res*, 46(D1), D633-D639, 2018
20. 16. Castro-Severyn, J., Pardo-Esté, C., Mendez, K. N., Fortt, J., Marquez, S., Molina, F., ... and Saavedra, C. P.: Living to the high extreme: unraveling the composition, structure, and functional insights of bacterial communities thriving in the arsenic-rich Salar de Huasco altiplanic ecosystem. *mSpectrum*, 9(1), 10-1128, 2021
21. Castro-Severyn, J., Pardo-Esté, C., Sulbaran, Y., Cabezas, C., Gariazzo, V., Briones, A., ... and Saavedra, C. P.: Arsenic response of three altiplanic *Exiguobacterium* strains with different tolerance levels against the metalloid species: a proteomics study. *Front Microbiol*, 10, 2161, 2019
22. Cerda, M., Evangelista, H., Valdés, J., Siffedine, A., Boucher, H., Nogueira, J., ... and Ortlieb, L.: A new 20th century lake sedimentary record from the Atacama Desert/Chile reveals persistent PDO (Pacific Decadal Oscillation) impact. *J South Am Earth Sci*. 95: 102302, 2019
23. Chang, L., Heslop, D., Roberts, A. P., Rey, D., and Mohamed, K. J.: Discrimination of biogenic and detrital magnetite through a double Verwey transition temperature. *J Geophys.*, 121(1), 3-14, 2016
24. Cohen, A. S., Manobianco, J., Dettman, D. L., Black, B. A., Beck, C., Feibel, C. S., ... and Vonhof, H. Seasonality and lake water temperature inferred from the geochemistry and sclerochronology of quaternary freshwater bivalves from the Turkana Basin, Ethiopia and Kenya. *Quaternary Sci Rev*, 317, 108284, 2023

25. Collado, L., Levican, A., Perez, J., and Figueras, M. J.: *Arcobacter defluvii* sp. nov., isolated from sewage samples. *IJSE*, 61(9), 2155-2161, 2011
26. Cron, B., Henri, P., Chan, C. S., Macalady, J. L., and Cosmidis, J.: Elemental sulfur formation by *Sulfuricurvum kужиense* is mediated by extracellular organic compounds. *Front Microbiol*, 10, 2710, 2019
27. 23. Csavina, J., Field, J., Taylor, M. P., Gao, S., Landázuri, A., Betterton, E. A., and Sáez, A. E.: A review on the importance of metals and metalloids in atmospheric dust and aerosol from mining operations. *STOTEN*, 433, 58-73, 2012
28. 24. Da Costa, C., Colin, Y., Debret, M., Copard, Y., Gardes, T., Jacq, K., ... and Berthe, T. Shifts in sediment bacterial communities reflect changes in depositional environments in a fluvial context. *STOTEN*, 885, 163890, 2023
29. 25. Dearing, J. A., Hay, K. S., Baban, A. S., Huddleston, E. M., Wellington H., and Loveland P. J.: Magnetic susceptibility of topsoils: a test of conflicting theories using a national database. *Geophys. J. Int.* 127: 728–734, 1996
30. Demergasso, C., Escudero, L., Casamayor, E. O., Chong, G., Balagué, V., and Pedrós-Alió, C.: Novelty and spatio-temporal heterogeneity in the bacterial diversity of hypersaline Lake Tebenquiche (Salar de Atacama). *Extremophiles*, 12, 491-504, 2008
31. Deng, Y., Mou, T., Wang, J., Su, J., Yan, Y., and Zhang, Y. Q.: Characterization of three rapidly growing novel *Mycobacterium* species with significant polycyclic aromatic hydrocarbon bioremediation potential. *Front Microbiol*, 14, 2023
32. Dib, J., Motok, J., Zenoff, V. F., Ordoñez, O., and Fariás, M. E.: Occurrence of resistance to antibiotics, UV-B, and arsenic in bacteria isolated from extreme environments in high-altitude (above 4400 m) Andean wetlands. *Curr. Microbiol.* 56, 510–517, 2008
33. Dittmar T.: Hydrochemical process controlling arsenic and heavy metal contamination in the Elqui river system (Chile). *STOTEN*. 325: 193–207, 2004
34. Donati, E. R., Sani, R. K., Goh, K. M., and Chan, K. G.: Recent advances in bioremediation/biodegradation by extreme microorganisms. *Front Microbiol*, 10, 1851, 2019
35. Dong, H., Huang, L., Zhao, L., Zeng, Q., Liu, X., Sheng, Y., ... and Chen, H.: A critical review of mineral–microbe interaction and co-evolution: mechanisms and applications. *National Sci Rev*, 9(10), nwac128, 2022
36. Dorador, C., Fink, P., Hengst, M., Icaza, G., Villalobos, A. S., Vejar, D., ... and Harrod, C.: Microbial community composition and trophic role along a marked salinity gradient in Laguna Puilar, Salar de Atacama, Chile. *Antonie Van Leeuwenhoek*, 111, 1361-1374, 2018
37. Douglas, G. M., Maffei, V. J., Zaneveld, J. R., Yurgel, S. N., Brown, J. R., Taylor, C. M., ... and Langille, M. G.: PICRUSt2 for prediction of metagenome functions. *Nat biotechnol*, 38(6), 685-688, 2020
38. Falkinham, J. O.: Environmental sources of nontuberculous mycobacteria. *Clin chest med*, 36(1), 35-41, 2015

39. Farias, M. E., Contreras, M., Rasuk, M. C., Kurth, D., Flores, M. R., Poire, D. G., ... and Visscher, P. T.: Characterization of bacterial diversity associated with microbial mats, gypsum evaporites and carbonate microbialites in thalassic wetlands: Tebenquiche and La Brava, Salar de Atacama, Chile. *Extremophiles* 18, 311–329, 2014.
40. Fernandez, A. B., Rasuk, M. C., Visscher, P. T., Contreras, M., Novoa, F., Poire, D. G., ... and Farias, M. E.: Microbial diversity in sediment ecosystems (evaporites domes, microbial mats, and crusts) of hypersaline Laguna Tebenquiche, Salar de Atacama, Chile. *Front Microbiol*, 7, 1284, 2016
41. Fialová, H., Maier, G., Petrovský, E., Kapička, A., Boyko, T., Scholger, R., and MAGPROX Team.: Magnetic properties of soils from sites with different geological and environmental settings. *J Appl Geophys.*, 59(4), 273-283, 2006
42. Garreaud, R., Vuille, M., and Clement, A. C.: The climate of the Altiplano: observed current conditions and mechanisms of past changes. *Palaeogeogr.*, 194(1-3), 5-22, 2003
43. Falkinham III, J. O.: The biology of environmental mycobacteria. *Env Microbiol Rep*, 1(6), 477-487, 2009
44. Grosjean, M., and Veit, H. Water resources in the arid mountains of the Atacama Desert (northern Chile): past climate changes and modern conflicts. *Global change and mountain regions: an overview of current knowledge*, 93-104, 2005
45. Hamilton, P. B., Rolshausen, G., Uren Webster, T. M., and Tyler, C. R.: Adaptive capabilities and fitness consequences associated with pollution exposure in fish. *Proc. R. Soc. B*, 372(1712), 20160042, 2017
46. Han, Y., and Perner, M.: The globally widespread genus *Sulfurimonas*: versatile energy metabolisms and adaptations to redox clines. *Front Microbiol*, 6, 989, 2015
47. Haslett, J., and Parnell, A.: A simple monotone process with application to radiocarbon-dated depth chronologies. *J R Stats*, 57(4), 399-418, 2008
48. Heiri, O., Lotter, A. F., and Lemcke, G.: Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. *J. Paleolimnol*, 25, 101-110, 2001
49. Hernández, A., Bao, R., Giral, S., Barker, P. A., Leng, M. J., Sloane, H. J., and Sáez, A.; Biogeochemical processes controlling oxygen and carbon isotopes of diatom silica in Late Glacial to Holocene lacustrine rhythmites. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 299(3–4), 413–425, 2012.
50. Hogg, A. G., Heaton, T. J., Hua, Q., Palmer, J. G., Turney, C. S. M., Southon, J., Bayliss, A., Blackwell, P. G., Boswijk, G., Bronk Ramsey, C., Pearson, C., Petchey, F., Reimer, P., Reimer, R. and Wacker, L.: “SHCal20 SOUTHERN HEMISPHERE CALIBRATION, 0–55,000 YEARS CAL BP,” *Radiocarbon*. Cambridge University Press, pp. 1–20, 2020
51. Imachi, H., Sakai, S., Lipp, J. S., Miyazaki, M., Saito, Y., Yamanaka, Y., ... and Takai, K. *Pelolinea submarina* gen. nov., sp. nov., an anaerobic, filamentous bacterium of the phylum Chloroflexi isolated from seafloor sediment. *International journal of systematic and evolutionary microbiology*, 64(Pt_3), 812-818, 2014

52. Imachi, H., Sakai, S., Sekiguchi, Y., Hanada, S., Kamagata, Y., Ohashi, A., and Harada, H. *Methanolinea tarda* gen. nov., sp. nov., a methane-producing archaeon isolated from a methanogenic digester sludge. *IJSEM*, 58(1), 294-301, 2008
53. Jiao, J. Y., Liu, L., Hua, Z. S., Fang, B. Z., Zhou, E. M., Salam, N., ... and Li, W. J.: Microbial dark matter coming to light: challenges and opportunities. *Nat Sci Rev*, 8(3), nwaa280, 2021
54. Kadnikov, V. V., Mardanov, A. V., Beletsky, A. V., Karnachuk, O. V., and Ravin, N. V. Genome of the candidate phylum Aminicenantes bacterium from a deep subsurface thermal aquifer revealed its fermentative saccharolytic lifestyle. *Extremophiles*, 23, 189-200, 2019
55. Kanehisa, M., Goto, S., Sato, Y., Furumichi, M., and Tanabe, M.: KEGG for integration and interpretation of large-scale molecular data sets. *Nucleic acids Res*, 40(D1), D109-D114, 2012
56. Kassambara, A.: *R Graphics Essentials for Great Data Visualization: 200 Practical Examples You Want to Know for Data Science*. STHDA, 2017
57. Keller W, Yan ND, Holtze K, Pitblado JR. Chemical response of acid lakes in the Sudbury, Ontario area to reduced smelter emissions, 1981–1989. *Can J Fish Aquat Sci*. 49 (Suppl.1): 25–32, 1992
58. Kereszturi, Á., Aszalós, J. M., Heiling, Z., Ignéczi, Á., Kapui, Z., Király, C., ... and Szalai, Z. Cold, dry, windy, and UV irradiated: surveying Mars-relevant conditions in Ojos del Salado Volcano (Andes Mountains, Chile). *Astrobiology*, 20(6), 677-683, 2020
59. Korosi, J. B., Thienpont, J. R., Smol, J. P., and Blais, J. M. Paleo-ecotoxicology: what can lake sediments tell us about ecosystem responses to environmental pollutants?. *Environ. Sci. Technol.*, 51(17), 9446-9457, 2017
60. Kostka, A., and Leśniak, A. Natural and anthropogenic origin of metals in lacustrine sediments; assessment and consequences—A case study of Wigry lake (Poland). *Minerals*, 11(2), 158, 2021
61. Kurth, D., Elias, D., Rasuk, M. C., Contreras, M., and Farias, M. E. Carbon fixation and rhodopsin systems in microbial mats from hypersaline lakes Brava and Tebenquiche, Salar de Atacama, Chile. *PLoS One*, 16(2), e0246656, 2021
62. Lahti, L., Shetty, S., Blake, T., and Salojärvi, J. Tools for microbiome analysis in R Version 2.1. 26, 2017
63. Laplante K, Sébastien B, Derome N. Parallel changes of taxonomic interaction networks in lacustrine bacterial communities induced by a polymetallic perturbation. *Evol Appl.*, 6(4):643-59, 2013
64. Li, W. C., and Wong, M. H. Effects of bacteria on metal bioavailability, speciation, and mobility in different metal mine soils: a column study. *J soils sediments*, 10, 313-325, 2010
65. Lins, U., Keim, C. N., Evans, F. F., Farina, M., and Buseck, P. R.: Magnetite (Fe₃O₄) and greigite (Fe₃S₄) crystals in multicellular magnetotactic prokaryotes. *Geomicrobiology J*, 24(1), 43-50, 2007
66. Louca, S., Parfrey, L. W., and Doebeli, M.: Decoupling function and taxonomy in the global ocean microbiome. *Science*, 353(6305), 1272-1277, 2016

67. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol.* 2014;15(12):1–21
68. Madsen, E. L.: Microorganisms and their roles in fundamental biogeochemical cycles. *Curr opinion Biotech.*, 22(3), 456-464, 2011
69. Mathivanan, K., Chandirika, J. U., Vinothkanna, A., Yin, H., Liu, X., and Meng, D. Bacterial adaptive strategies to cope with metal toxicity in the contaminated environment—A review. *Ecotox Env Safety*, 226, 112863, 2021
70. McMurdie P, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE*. 8(4):e61217, 2013
71. Moskowitz, B. M., Bazylnski, D. A., Egli, R., Frankel, R. B., and Edwards, K. J.: Magnetic properties of marine magnetotactic bacteria in a seasonally stratified coastal pond (Salt Pond, MA, USA). *GJI* 174(1), 75-92, 2008
72. Niu, Z. S., Yang, Y., Tou, F. Y., Guo, X. P., Huang, R., Xu, J., ... and Hochella, M. F. Sulfate-reducing bacteria (SRB) can enhance the uptake of silver-containing nanoparticles by a wetland plant. *Env Sci: Nano*, 7(3), 912-925, 2020.
73. Olilo, C. O., Odoli, C. O., Obiero, M. O., Malala, J. O., & Bironga, C. H. (2023). Microbial ecology of desert Lakes Baringo and Turkana, Kenya, East Africa. In *Lakes of Africa* (pp. 247-268). Elsevier.
74. Ordoñez, O. F., Flores, M. R., Dib, J. R., Paz, A., and Farías, M. E. Extremophile culture collection from Andean lakes: extreme pristine environments that host a wide diversity of microorganisms with tolerance to UV radiation. *Microb. Ecol.* 58, 461–473, 2009.
75. Orellana, R., Macaya, C., Bravo, G., Dorochesi, F., Cumsille, A., Valencia, R., ... and Seeger, M. Living at the frontiers of life: extremophiles in Chile and their potential for bioremediation. *Front Microbiol*, 9, 2309, 2018
76. Oren, A. (2014). The Family Methanoregulaceae . In: Rosenberg, E., DeLong, E.F., Lory, S., Stackebrandt, E., Thompson, F. (eds) *The Prokaryotes*. Springer, Berlin, Heidelberg.
77. Oren, A., and Garrity, G. M. Valid publication of the names of forty-two phyla of prokaryotes. *International journal of systematic and evolutionary microbiology*, 71(10), 005056, 2021
78. Pérez-Cataluña, A., Salas-Massó, N., Diéguez, A. L., Balboa, S., Lema, A., Romalde, J. L., and Figueras, M. J. Revisiting the taxonomy of the genus *Arcobacter*: getting order from the chaos. *Front Microbiol*, 9, 2077, 2018
79. Pérez-Portilla, P., Aránguiz-Acuña, A., Pizarro, H., and Herrera, J.: Assessing the Effects of Long-Term Mining Exploitation on a Lacustrine System from the Arid Region of the Atacama Desert, Chile. *STOTEN*, 949: <https://doi.org/10.1016/j.scitotenv.2024.174771>, 2024
80. Picard, M., Wood, S. A., Pochon, X., Vandergoes, M. J., Reyes, L., Howarth, J. D., ... and Puddick, J. Molecular and pigment analyses provide comparative results when reconstructing historic cyanobacterial abundances from lake sediment cores. *Microorganisms*, 10(2), 279, 2022
81. Pigati JS, Miller DM, Bright J, Mahan SA, Nekola JC, Paces JB. Chronology, sedimentology, and micro-fauna of ground-water discharge deposits in the central Mojave Desert, Valley Wells, California. *Geol Soc Am Bull.*, 123: 2224–2239, 2011

82. Placzek, C., Quade, J., Betancourt, J. L., Patchett, P. J., Rech, J. A., Latorre, C., ... and English, N. B. Climate in the dry central Andes over Geologic, millennial, and interannual timescales. *Ann. Mo. Bot. Gard.*, 96(3), 386–397, 2009.
83. Pollard HG, Colbourne JK, Keller W. Reconstruction of Centuries-old *Daphnia* Communities in a Lake Recovering from Acidification and Metal Contamination. *Ambio.*, 32: 214–218, 2003
84. Price, M.N., Dehal, P.S., Arkin, A.P.: FastTree: computing large minimum evolution trees with profiles instead of a distance matrix. *Mol Biol Evol.*, 26(7):1641–1650, 2009
85. Pueyo, J. J., Sáez, A., Giralt, S., Valero-Garcés, B. L., Moreno, A., Bao, R., Schwalb, A., Herrera, C., Klosowska, B., and Taberner, C.: Carbonate and organic matter sedimentation and isotopic signatures in Lake Chungará, Chilean Altiplano, during the last 12.3 kyr, *Palaeogeogr. Palaeoclimatol.*, 307, 339–355, 2011
86. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, et al. The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucl Acids Res.*, 41(D1):D590–D596, 2012.
87. Raab, A., and Feldmann, J. Microbial transformation of metals and metalloids. *Science Progress*, 86(3), 179–202, 2003
88. Rahman, Z. An overview on heavy metal resistant microorganisms for simultaneous treatment of multiple chemical pollutants at co-contaminated sites, and their multipurpose application. *J. Haz. Mat.*, 396, 122682, 2020
89. Rainey, F., Kämpfer, P., Trujillo, M., Chun, J., DeVos, P., Hedlund, B., and Dedysh, S. (2015). *Bergey's manual of systematics of Archaea and Bacteria* (Vol. 410). W. B. Whitman (Ed.). Hoboken, NJ: Wiley.
90. Ramírez M, Massolo S, Frache R, Correa JA. Metal speciation and environmental impact on sandy beaches due to El Salvador copper mine, Chile. *Mar Pollut Bull.*, 50, 62–72, 2005
91. Ramos-Tapia I, Salinas P, Núñez R, Cortez D, Soto J, Paneque M. Compositional Changes in Sediment Microbiota Are Associated with Seasonal Variation of the Water Column in High-Altitude Hyperarid Andean Lake Systems. *Microbiol Spectr.* 15;11(3):e0520022, 2023
92. Rasuk, M. C., Kurth, D., Flores, M. R., Contreras, M., Novoa, F., Poire, D., and Farias, M. E.: Microbial characterization of microbial ecosystems associated to evaporites domes of gypsum in Salar de Llamara in Atacama Desert. *Microb. Ecol.* 71, 44–56, 2014
93. Ritter, B., Wennrich, V., Medialdea, A., Brill, D., King, G., Schneiderwind, S., ... and Dunai, T. J.: Climatic fluctuations in the hyperarid core of the Atacama Desert during the past 215 ka. *Sci Rep.*, 9: 5270, 2019
94. Rodríguez-Luna, D., Encina-Montoya, F., Alcalá, F. J., and Vela, N.: An Overview of the Environmental Impact Assessment of Mining Projects in Chile. *Land*, 11(12), 2278, 2022
95. Romero, H., Méndez, M., Smith, P.: Mining development and environmental in-justice in the Atacama desert of northern Chile. *Environ. Justice* 5, 70–76, 2012
96. Romero, L., Alonso, H., Campano, P., Fanfani, L., Cidu, R., Dadea, C., ... and Farago, M. Arsenic enrichment in waters and sediments of the Rio Loa (Second Region, Chile). *App Geochem*, 18(9), 1399–1416, 2003

97. Roosa, S., Wattiez, R., Prygiel, E., Lesven, L., Billon, G., and Gillan, D. C. Bacterial metal resistance genes and metal bioavailability in contaminated sediments. *Env pollution*, 189, 143-151, 2014
98. Salvarredy-Aranguren MM, Probst A, Roulet M, Isaure M-P.: Contamination of surface waters by mining wastes in the Milluni Valley (Cordillera Real, Bolivia): Mineralogical and hydrological influences. *Appl Geochem.*, 23: 1299–1324, 2008
99. Santini, T. C., Gramenz, L., Southam, G., & Zammit, C. (2022). Microbial community structure is most strongly associated with geographical distance and PH in salt lake sediments. *Frontiers in Microbiology*, 13, 920056.
100. Staicu, L. C., and Stolz, J. F.: Microbes vs. metals: Harvest and recycle. *FEMS microbiol ecology*, 97(5), fiab056, 2021
101. Stankevica, K., Vincevica-Gaile, Z., Klavins, M., Kalnina, L., Stivrins, N., Grudzinska, I., and Kaup, E.: Accumulation of metals and changes in composition of freshwater lake organic sediments during the Holocene. *Chem Geol*, 539, 119502, 2020
102. Stoeva, M. K., Aris-Brosou, S., Chételat, J., Hintelmann, H., Pelletier, P., & Poulain, A. J. (2014). Microbial community structure in lake and wetland sediments from a high Arctic polar desert revealed by targeted transcriptomics. *PLoS One*, 9(3), e89531.
103. Taj, Z., & Chattopadhyay, I. (2024). Identification of bio-active secondary metabolites from Actinobacteria as potential drug targets against Porphyromonas gingivalis in oral squamous cell carcinoma using molecular docking and dynamics study. *In Silico Pharmacology*, 12(1), 1-24.
104. Tapia, J., Davenport, J., Townley, B., Dorador, C., Schneider, B., Tolorza, V., and von Tümpling, W.: Sources, enrichment, and redistribution of As, Cd, Cu, Li, Mo, and Sb in the Northern Atacama Region, Chile: implications for arid watersheds affected by mining. *J Geochem Exploration*, 185, 33-51, 2018
105. Tapia, J., Mukherjee, A., Rodríguez, M. P., Murray, J., and Bhattacharya, P.: Role of tectonics and climate on elevated arsenic in fluvial systems: Insights from surface water and sediments along regional transects of Chile. *Env Pollution*, 314, 120151, 2022
106. Tapia, J., Murray, J., Ormachea, M., Tirado, N., and Nordstrom, D. K. Origin, distribution, and geochemistry of arsenic in the Altiplano-Puna plateau of Argentina, Bolivia, Chile, and Perú. *STOTEN*, 678, 309-325, 2019
107. Teunisse, G. M.: Fantaxtic - Nested Bar Plots for Phyloseq Data (Version 2.0.1) [Computer software]. <https://github.com/gmteunisse/Fantaxtic>, 2022
108. Thukral, A. K. A review on measurement of Alpha diversity in biology. *Agri Res J*, 54(1), 2017
109. Tomazini Jr, A., Lal, S., Munir, R., Stott, M., Henrissat, B., Polikarpov, I., ... and Levin, D. B.: Analysis of carbohydrate-active enzymes in Thermogemmatospira sp. strain T81 reveals carbohydrate degradation ability. *Canadian journal of microbiology*, 64(12), 992-1003, 2018
110. Trolle, D., Hamilton, D.P., Pilditch, C.A.: Evaluating the influence of lake morphology, trophic status and diagenesis on geochemical profiles in lake sediments. *Appl Geochem.*, 25:621–632, 2010

111. Usenko, S., Landers, D.H., Appleby, P.G., Simonich, S.L.: Current and historical deposition of PBDEs, pesticides, PCBs, and PAHs to Rocky Mountain National Park. *Environ Sci Technol.*, 41:7235–7241, 2007
112. Valero-Garcés, B. L., Delgado-Huertas, A., Navas, A., Edwards, L., Schwalb, A., and Ratto, N.: Patterns of regional hydrological variability in central-southern Altiplano (18–26°S) lakes during the last 500 years, *Palaeogeogr. Palaeoclimatol.*, 194, 319–338, 2003.
113. Vargas-Machuca, B. D., Zanetta-Colombo, N., De Pol-Holz, R., and Latorre, C. Variations in local heavy metal concentrations over the last 16,000 years in the central Atacama Desert (22° S) measured in rodent middens. *STOTEN* 775, 145849, 2021
114. Verosub, K. L., and Roberts, A. P.: Environmental magnetism: Past, present, and future. *J. Geophys.*, 100(B2), 2175–2192, 1995
115. Versteeg, J. K., Morris, W. A., and Rukavina, N. A.: The utility of magnetic properties as a proxy for mapping contamination in Hamilton Harbour sediment. *J Great Lakes Res.*, 21(1), 71–83, 1995
116. Vignale, F. A., Lencina, A. I., Stepanenko, T. M., Soria, M. N., Saona, L. A., Kurth, D., ... and Farías, M. E.: Lithifying and non-lithifying microbial ecosystems in the wetlands and salt flats of the Central Andes. *Microb Ecol.*, 1–17, 2021
117. Wasmund, K., Schreiber, L., Lloyd, K. G., Petersen, D. G., Schramm, A., Stepanauskas, R., ... and Adrian, L.: Genome sequencing of a single cell of the widely distributed marine subsurface Dehalococcoidia, phylum Chloroflexi. *ISME J.* 8, 383–397. doi: 10.1038/ismej.2013.143, 2014
118. Wickham, H.: *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. ISBN 978-3-319-24277-4, <https://ggplot2.tidyverse.org>, 2016
119. Yan, C., Wang, F., Liu, H., Liu, H., Pu, S., Lin, F., ... and Yuan, R.: Deciphering the toxic effects of metals in gold mining area: microbial community tolerance mechanism and change of antibiotic resistance genes. *Env Res.*, 189, 109869, 2020
120. Yan, Y., He, A., Dilek, Y., Zhu, Z., and Zhao, Q. Landscape inversion episodes in SE China during the Mesozoic–early Cenozoic: Constrained by trace-element contents, Nd isotope geochemistry, and detrital zircon U–Pb geochronology of sedimentary basins. *GSA Bulletin.*, 136(7–8), 2978–2998, 2024
121. Yang, C., Mai, J., Cao, X., Burberry, A., Cominelli, F., and Zhang, L.: ggp crust2: an R package for PICRUST2 predicted functional profile analysis and visualization. *Bioinformatics*, 39(8), btad470, 2023
122. Yao, Z., Gao, P.: Heavy metal research in lacustrine sediment: a review. *Chin. J. Ocean. Limnol.* 25, 444–454, 2007.
123. Zanetta-Colombo, N. C., Fleming, Z. L., Gayo, E. M., Manzano, C. A., Panagi, M., Valdés, J., and Siegmund, A.: Impact of mining on the metal content of dust in indigenous villages of northern Chile. *Env Intl*, 169, 107490, 2022
124. Zanetta-Colombo, N. C., Scharnweber, T., Christie, D. A., Manzano, C. A., Blersch, M., Gayo, E. M., ... and Nüsser, M. When another one bites the dust: Environmental impact of global copper demand on local communities in the Atacama mining hotspot as registered by tree rings. *STOTEN*, 920, 170954, 2024

- 760 125.Zha, Y., Chong, H., Yang, P., and Ning, K. (2022). Microbial dark matter: from discovery to applications. *GPB*,
761 20(5), 867-881.
- 762 126.Zhao, Y.N., Li, X.F., Ren, Y.P., Wang, X.H.: Effect of static magnetic field on the performances of and anode biofilms
763 in microbial fuel cells. *RSC Adv* 6(85):82301–82308, 2016
- 764 127.Zhou, H., and van Hullebusch, E. D.: Microbial interaction and transformation of metals and metalloids. *Letters*
765 *Applied Microbiol*, 75(5), 1074-1075, 2022