

Author comments

Ensemble design for seasonal climate predictions: Studying extreme Arctic sea ice lows with a rare event algorithm

J.Sauer, G. Zappa, F. Massonet, F. Ragone

submitted to Earth System Dynamics (ESD) within the Special issue
“Theoretical and computational aspects of ensemble design,
implementation, and interpretation in climate science” (EGUSPHERE-
2024-3082)

January 24, 2025

Figures for replies to reviewer #3




Figure R1: Ensemble simulations initialized on 01 February (a,c,e) 1930 and (b,d,f) 1037 of the control run. (a-b) Rare event simulations: $N=600$ trajectories (thin blue lines) and ensemble mean (thick blue line) of daily pan-Arctic sea ice area anomalies relative to the daily climatology of the corresponding control ensemble means. The gray dashed lines show the intra-ensemble standard deviations in the control ensembles. All lines are presented as 15-day running means. (c-f) Probability distribution functions of (c-d) February-September and (e-f) August-September mean pan-Arctic sea ice area for (blue) the rare event simulation, (black) the control ensembles ($N=6000$ trajectories) and (red) the 3000-year control run. The vertical lines show the mean of the distributions. The black and blue values indicate the smallest February-September and August-September mean sea ice area value in the control and rare event ensemble simulations respectively.




Figure R2: Probabilities (x-axes) of (a) February-September and (b) August-September mean sea ice area equal or smaller than a given threshold (y-axes) as a function of four different initial conditions for (stars) the $N=6000$ trajectory control ensembles (i.e. ten realizations are merged into one single ensemble) and (solid lines) the average over four to ten algorithm estimates. The shading are 95% confidence intervals derived from the t -distribution of the four to ten estimates (see Supplementary Information S4 of the revised manuscript). Note that the y-axes are displayed in reverse order. In the legend, "SIA" and "SIT1.93" indicate the state of the January-February mean anomaly of the pan-Arctic sea ice area and the cumulative area with sea ice thickness equal or larger than 1.93 m respectively. The grey labels on the right of each panel show how many standard deviations a sea ice area value is below the mean of the 3000-year control run.




Figure R3: PlaSim-LSG 3000-year control run (Sauer et al. [2024]): (a) Normalized mean anomalies of January-February mean quantities conditional on extreme negative (left) February-September (FEBSEP) and (right) August-September (AUGSEP) mean pan-Arctic sea ice area anomalies equal or smaller than -2.5 standard deviations. "SIV", "SIA" and "SIT" are the pan-Arctic sea ice volume, sea ice area and mean sea ice thickness. "SIT \leq threshold" are anomalies in the cumulative area with sea ice thickness equal or larger than a critical threshold. Hatching denotes statistical significance at the 5% level assessed from a two-sided t -test applied to five composite estimates after subdividing the 3000-year control run into five 600-member ensembles (see Supplementary Information S4 of the revised manuscript). (b) Scatter plot of January-February mean anomalies of SIT1.93 vs. pan-Arctic sea ice area including the years from the selected initial conditions.

References

Plotkin, D. A., Webber, R. J., O'Neill, M. E., Waere, J., and Abbot, D. S.: Maximizing simulated tropical cyclone intensity with action minimization, *Journal of Advances in Modeling Earth Systems*, 11, 863–891, <https://doi.org/10.1029/2018ms001419>, 2019.

Sauer, J., Demaeyer, J., Zappa, G., Massonet, F., and Ragone, F.: Extremes of summer Arctic sea ice reduction investigated with a rare event algorithm, *Climate Dynamics*, pp. 1–19, <https://doi.org/10.1007/s00382-024-07160-y>, 2024.