

Response to Reviewer 1

We thank the reviewer for their feedback and constructive criticism, which have helped us significantly improve our manuscript. We have taken care to address each comment with a direct response. The text from your comments are shown in black and our responses are shown in blue. Responses include the manuscript text that was changed, removed, or added.

Overall Notes This paper introduces a python tool aimed at associating in situ aerosol measurements obtained during aircraft field campaigns with remote sensing aerosol retrievals, by addressing the need to compensate for the limited coarse-mode throughput of aircraft inlets and to hydrate in situ samples when comparing with ambient (remote sensing) observations.

As these calculations must be made if in-situ field data are to be used to validate remote sensing retrievals quantitatively, the algorithm presented here represents a useful tool for such applications. The observations used to test this approach were acquired during the ACTIVATE field campaign, and the scope of the present study is limited to fine-mode sulfate and organic aerosol, and a coarse mode taken as sea salt. They impose assumptions that limit considerably the applicability of the current implementation - constant refractive indices over the spectral range, spherical particles, parameter assumptions required to calculate the hydrated CRI, etc. However, these are stated clearly, which is as much as one can ask in an AMT paper. The remote sensing data were obtained from the HSRL-2 and RSP aircraft instruments, which avoids some of the sampling differences that arise when in situ measurements are compared with spacecraft measurements. As such, the approach seems most applicable for validating aircraft field measurements.

In summary, the paper develops a useful tool and presents a thorough analysis of its performance. For general application, there are significant limitations in the assumptions made, but given that the analysis is circumscribed to a narrow set of relatively favorable conditions, I think this is acceptable for publication in AMT, perhaps with minor modifications as suggested below. Thank you for the comment. We agree with this assessment and, while some of the text has changed, this description of our work is still true. We address each of the suggested changes below, but no specific changes are made as a result of this summary.

A Few More Specific Notes Line 244. It might be useful to mention how far the aircraft travels in 45 seconds, to provide a sense for the horizontal resolution of the SMPS and other, aggregated measurements.

Thank you for the comment. The text “that can travel of 8 km across the ground in 45 seconds” has been added to the end of this sentence.

Line 270. Might be worth noting that remote sensing is more sensitive to volume than number concentration specifically for particles smaller than the observing wavelength. For particles larger than the observing wavelength, sensitivity is greater to particle cross-sectional area.

Thank you for the comment. We agree this is relevant information and we have added it to the paragraph that now reads as follows:

In general, remote sensors are not as sensitive to particle number concentration as they are to particle surface area and volume concentrations. For particles larger than the remote sensor’s observing wavelength, the remote sensor is most sensitive to particle cross-sectional area. For particles smaller than the observing wavelength, the remote sensor is most sensitive to volume concentration. Given the sensitivity of remote sensors to surface area and volume concentration, this work also discusses the logarithmic size-resolved aerosol particle surface area concentration (a°) and logarithmic size-resolved aerosol particle volume concentration (v°).

Line 454. Comparing Fig. 1 with Fig. 5, and taking the y-axis scales into account, I would say “...overall, much less variance.” Thank you for the comment. We agree this is a fair assessment and have changed the text to be “..., but the synthetic data see overall much less variance”

Line 469. As this is synthetic data, doesn’t the statement here just mean that the numerical coding was done correctly? Not a bad thing to mention, but the statement here makes the observation sound more fundamental.

Thank you for the comment. We agree this finding has been overstated and have changed the sentence to be “To demonstrate

50 the functionality of this analysis, the synthetic data generation and retrieval processes were repeated with zero measurement noise, which results in a rate of successful retrievals of 100%.”

Line 494. A word seems to be missing from this sentence.

55 Thank you for the comment. We appreciate catching the error. We have changed the text to be “...which suggests that the ISARA-derived products are less reliable at relatively low scattering signal (i.e., signal < 5 Mm⁻¹)”

Lines 498-500. By way of explanation, wouldn’t the 700 nm channel likely be the most sensitive to coarse-mode particles, for which many of the assumptions might be less applicable?

60 Thank you for the comment. We agree that this increase in MRB with increasing wavelength is notable and while it is only a few percent difference, we have added the following text to capture this: “Because this bias increases with increasing wavelength, it is possible that some of this discrepancy is due to larger particles that are more commonly comprised of dust, but this trend only accounts for a few percent difference in MRB”.

Section 3.1.3. Just wondering how representative of the entire column the in situ data sampling might be. I realize the HSRL-2 data are height-resolved, which can help assess the vertical heterogeneity compared to the in situ sampling.

65 Thank you for the comment. We have analyzed this with some level of detail in Section 3.2 with regards to the RSP-derived and ISARA-derived products.

70 Figure 10. There appears to be a lot of scatter in the data, which the text does not seem to acknowledge. This is probably not surprising - in addition to the limitations discussed in the paragraph about this figure, given the likely horizontal and vertical variability in particle concentration combined with differences in sampling.

Thank you for the comment. To address this we have added error bars and standard deviations where appropriate throughout the text. We agree that there is a lot of scatter, however this appears to fall within the standard deviations of many of the points. The text has been changed as follows to address this: “Similar to C_{ext} and LR, the standard deviations in the HSRL-2+RSP-derived N often encompass the 1-to-1 line.”.

75

Response to Reviewer 2

We thank the reviewer for their feedback and constructive criticism, which have helped us significantly improve our manuscript. We have taken care to address each comment with a direct response. The text from your comments are shown in black and our responses are shown in blue. Responses include the manuscript text that was changed, removed, or added.

This manuscript presents a methodology to compare and reconcile aerosol observations from various platforms, so called ISARA algorithm, developed for the ACTIVATE mission data. It provides a useful and rather comprehensive example of attempting to achieve closure between diverse in-situ and remote sensing measurements. The study clearly addresses a complex problem and contributes relevant information for aerosol measurement harmonization. However, it would benefit from clarifying its broader motivation, better articulating its limitations, and improving the language and structure in some sections.

My main concern is that the manuscript abstract, and introduction do not seem to precisely describe the motivation of the study. The introduction frames the problem mainly as a data closure challenge, potentially giving the impression that a general solution (i.e., ISARA) will be offered. For example the sentence “Despite the important findings from these studies, systematic and streamlined closure of aerosol data sets has not been yet achieved.” However, the ISARA algorithm appears heavily tuned to specific conditions, relying on a priori information about aerosol composition, size, and shape, additionally being very limited in the atmospheric and aerosol conditions where it can be applied. This is perfectly understandable, but a clearer explanation of how broadly applicable the algorithm is would strengthen the outcome. The introduction could therefore more broadly discuss the complexity of in-situ vs remote-sensing aerosol validations, compilations and simulations – as for setting the scheme. I hope that authors consider this, and could slightly streamline the introduction and structure of the paper to better reflect the content.

In summary, this manuscript addresses a complex and relevant topic in atmospheric aerosol measurement by providing a detailed example of integrating multiple observation platforms. The methodology is simple but thoroughly described, and the analysis is well presented, but the applicability and limitations of the ISARA algorithm require clearer discussion. The manuscript would benefit from a refined motivation and improved language in certain sections. I recommend minor revisions before acceptance at AMT.

We thank the reviewer for their thoughtful and constructive feedback. In response to the concerns raised, we have revised the abstract and introduction to better articulate the overall motivation of the study. Specifically, we now clarify that this work focuses on assessing consistency across platforms as a necessary step toward achieving rigorous external closure, rather than offering a universal solution to the closure problem. We also explicitly acknowledge the limitations of the ISARA framework, including its dependence on assumptions about particle shape, composition, and the applicability to specific aerosol regimes. These clarifications are intended to better align the manuscript’s framing with its scope and contributions. We believe these revisions strengthen the manuscript and more accurately convey the goals and context of our analysis.

Minor Rather Technical Comments: Abstract:

- L6: Suggest removing “aircraft” as the methodology can apply to other in-situ data platforms as well. Or can it? Please consider this also when writing introduction. Thank you for the comment. We do agree that the framework offers a great deal of flexibility in the in-situ platform used. We have removed “aircraft” from this sentence and made subsequent changes within the introduction as well.

Introduction:

- L33: Remove the word “parameter”. Thank you for the comment. We have addressed this error. We also removed the example to discuss specific parameterizations later in the text.
- L51: Consider adding 1–2 sentences summarizing key findings from past studies for context. Thank you for the comment. We have added the following sentences to the text to provide more information on the findings of a few of these

studies: “These studies consistently find that the extinction and backscatter coefficients derived from in-situ instruments are systematically low compared to those derived from HSRL-2.” and “In particular, Pistone et al. (2019) found poor agreement between in-situ- and RSP- derived total SSA.”.

- L52: Clarify “these instrument data”. Which instruments are being referred to? Thank you for the comment. We have clarified the statement to be “... remote sensing data’
- L57: Suggest rephrasing to “...using a Nafion membrane dryer in the sampling line”. Thank you for the comment. The text has been altered according to the comment and is now: “...using a Nafion dryer or heating in the sampling line”.
- L86: Check for a possible extra “is”. Thank you for the comment. The extra is has been removed from this sentence.
- L91: Sentence structure is unclear-both content and grammar could be improved. Also, please clarify if the methodology is expected to be broadly applicable or limited to specific platforms and conditions. Thank you for the comment. The methods described in this paper are open source and flexible due to using MOPSMAP to handle the optical properties. The sentence has been changed to be: “While the current study focuses on the consistency analysis between in-situ- and remote sensing-derived aerosol properties of the more common spherical aerosol particles, it is hoped that the framework described in this study serves as an open source foundation that can be easily expanded and used to fully understand the information train between all manner of measurements and therefore enable systematic closure of field campaign aerosol data.”

Measurements:

- L198–202: Some grammar and sentence construction issues—please revise for clarity. Also, consider using micrometers (μm) instead of nanometers (nm) for $>1\ \mu\text{m}$ sizes. Thank you for the comment. We have updated the units to μm where there weren’t ranges going into the nm scale or significant figures would require multiple digits after the decimal place. We have clarified the sentence in question, which now is “The effective upper size cut is $D = 5\ \mu\text{m}$ for all 2020 data.”
- L202: Nephelometers and absorption measurements are introduced here without prior explanation. Reorganizing or cross-referencing earlier sections may improve flow. Thank you for the comment. We have added the Section 2.1 to provide a primer on particle properties and general background for these measurements.
- L213: Was the PSAP measurement also conducted at $<40\%$ RH? Please specify. Thank you for the comment. The PSAP sample stream is dried by heating the optical block to 35°C . The text has been updated to include this sentence.
- L218: Replace “variety of errors” with a more descriptive phrase identifying specific artifacts or correction needs. Thank you for the comment. We have replaced “variety of errors” with “transmittance and flow errors”.
- L223: How much data was excluded by the $>1\ \text{Mm}^{-1}$ cutoff for scattering and absorption? Justify the threshold. Thank you for the comment. This comment led to us evaluating the $1\ \text{Mm}^{-1}$ and determining that 0 was a better value for absorption. This yielded a 10x increase in the data available for retrieval. This threshold was changed accordingly to $0\ \text{Mm}^{-1}$. The results and conclusions have been updated accordingly.

- 90 – L245: Clarify what is meant by “most useful analysis” of profiles extending >1 km. Thank you for the comment. The profiles are most useful for comparison with the RSP-derived products. We have added the following text to clarify this information: “As such, the external consistency analysis is most useful from vertical profiles where the in-situ platform samples the column of air above an arbitrary ground point. Vertical profiles where the extent is more than 1 km are most useful for comparing with the column-averaged aerosol particle properties derived from the RSP-measurements.”.
- 95 – L257: Consider rephrasing “1-second data” as “native time resolution data” for clarity. Thank you for the comment. We tried to clarify this with the following text “The data that are in their native resolution are averaged to 45 seconds using the NASA standard merging tool.”
- 100 – L282, L286, L334: “ACTIVATE region” should be defined more precisely—preferably with coordinates or a map. Also specify which areas or conditions were excluded and why. Thank you for the comment. We have clarified the region that ACTIVATE was bounded by within the text. The following sentence was added to Section 2.2: “The extent of the North Atlantic region that was sample during ACTIVATE was within bounds of 58–78°W and 28–42°N.”. We have described the cloud filtering process and the focus on when the Falcon (i.e., the in-situ platform) was performing vertical profiles in coordination with the King Air. It is outside the scope of this work to describe the methodologies that guided the ACTIVATE mission.
- 105 – L289: Clarify “mid-point particle diameter.” Why is geometric mean diameter not used? Mid-point particle diameter is synonymous with geometric mean diameter. The text has been clarified as follows: “Note that this process uses the mid-point (i.e., the geometric mean) particle diameters from each SMPS and LAS channel.”.
- 110 – L285–295: The assumption of sulfate-only aerosol seems oversimplified. Given the limited RRI range and the potential presence of organics or other compounds, the assumptions behind the ISARA-derived optical properties need better justification. Also, specify how wavelength conversions were done and what values of AAE or SAE were used. Thank you for the comment. We agree that the limited RRI does limit the scope of this study, however, looking at Li et al. (2023), it would appear that most mixtures of aerosol particles will have an apparent real refractive index (RRI) that falls in the 1.51–1.55 range. As such we have updated the text with the following: “Furthermore, external mixtures of many aerosol species have an apparent RRI that falls between 1.5 and 1.58 (Li et al., 2023). If an external mixture of aerosols is dominated by a RRI outside of this range, it is likely that the assumptions of a spectrally flat CRI and sphericity are not longer valid.”. Additionally, AAE and SAE were not used. We calculate the coefficients at their native measurements. For example, dry scattering coefficients were calculated at 450, 550, and 700 nm wavelengths and dry absorption coefficients were calculated at 470, 532, and 660 nm wavelengths. The final ambient properties were again recalculated using MOPSMAP at each of the HSRL-2 wavelengths.
- 115 – L326: Missing preposition—please review. Thank you for the comment. This sentence has been changed to the following: “After calculating the set of forward-modeled $C_{\text{scat,wet}}$, we use γ to derive the “measured” $C_{\text{scat,wet}}$ (i.e., $C_{\text{scat,RH=80}}$).”.
- 120 – L334: Again, clarify the threshold inconsistency between L223 and here. How much absorption data were actually usable above 1 Mm^{-1} ? Thank you for the comment. As discussed in the response above, we have re-evaluated the use of 1 Mm^{-1} minimum threshold. Additionally, this sentence was not directly connected to the figure so it has been corrected as follows: “It is observed that aerosol particles in the ACTIVATE region generally have low absorption ($\text{IRI} \leq 0.01$) and low hygroscopicity ($\kappa \leq 0.1$).”.
- 125 –
- 130 –

135 – L615: The manuscript states that the region was chosen for “diversity of aerosol and meteorological conditions,” yet the
assumptions made to enable closure seem to contradict this. Consider rephrasing or qualifying this claim. Thank you for
the comment. We agree that as written this is somewhat contradictory. We have attempted to resolve this contradiction by
focusing on the large volume of data available. The text has been changed as follows: “...large volume of statistically-rich
aerosol measurements collected over three years of operations. Although the ACTIVATE region does feature a variety
140 of aerosol and meteorological conditions, many of the ACTIVATE missions were carried out in cloud-free conditions
without detectable influence from dust or smoke, making the data set well-suited for the consistency analysis performed
in this study.”.

Response to Reviewer 3

We thank the reviewer for their feedback and constructive criticism, which have helped us significantly improve our manuscript. We have taken care to respond directly to each comment. The text of your comments is shown in black, and our responses are shown in blue. The responses include the manuscript text that was changed, removed, or added.

The paper by Dmitrovic et al. describes a Python toolkit (ISARA) aimed at achieving closure between lidar and polarimeter observations of ambient aerosol properties on one hand, and in-situ measurements of the PSD, scattering and absorption on the other hand, which are limited by drying and by inlet effects. The toolkit is validated through in-situ observations (internal consistency), simulations (synthetic consistency) and remote sensing observations (external consistency). It is my feeling that the stated goals of the paper are of high importance, but that they are not sufficiently developed beyond the technical stage to be proposed for a publication. I would invite the authors to consider a more scientific and less technical approach in writing this paper, and to allow the time and the effort that this can represent at this stage I suggest to REJECT the manuscript. I will be willing to help with reviewing a revised manuscript in case the the below MAJOR points are addressed in full.

Thank you for the comment. We agree that the goals of the paper are of high importance. We also agree that the science focus of the paper needed some expansion. We therefore added a detailed analysis of more aerosol properties that was not described in the previous draft, including comparisons against RSP-derived coarse-mode properties that have never been evaluated in previous studies. Since our results show good agreement to within measurement uncertainties compared against advanced remote sensors (lidar and polarimeter observations) we do not agree that the paper is not developed past the technical stage. We further feel that the technical approach is an important development on its own, since it is a foundation that can be used to directly replicate our findings using the code and data provided. Although our results already show significant scientific promise, they can also be readily improved and expanded upon in the future. We show consistency between remote and in situ aerosol products for several ambient aerosol properties that have not been evaluated by previous studies, and we indicate which properties are not consistent. We present a comprehensive analysis using spherical particle models. It is outside the scope of this paper to perform additional analyses of more complex particle types such as dust and smoke since they require us to use modeling of non-spherical and heterogeneous particle shapes and to revisit many of the assumptions made in the remote sensing and in-situ retrievals themselves. Future work will involve demonstrating how we can leverage the additional capability of libraries like MOPSMAP to evaluate these more complex particle types. We have made extensive revisions of the manuscript to address the scientific shortcomings and to address each of your specific comments below. As such, we feel that the paper should not be rejected and we welcome the reviewer to reconsider our revisions.

MAJOR POINTS:

1. The paper takes a technical stance to the task, e.g. insisting on unnecessary details such as the use of the python programming language and the MOPSMAP scattering code. I would invite to present an algorithm, and to discuss its assumptions, limitations and uncertainties, rather than a piece of software. Concerning the scattering calculations, given that the authors only assume spherical particles, MOPSMAP is only a library allowing Mie scattering calculations and any other Mie scattering code would achieve the same results, therefore the scientific approach is to indicate that the framework of Mie theory is being used, and not to indicate which scattering library is used. The usage of the MOPSMAP code could most probably be of interest in the case of non-spherical scattering, but non-spherical scattering is not envisaged here. Thank you for the comment. As discussed above, we agree that there needed to be more description of the science behind aerosol properties. The structure of the paper has been changed accordingly. We have added Section 2.1 to add more scientific background to our technical description. In the revised draft, we now clearly explain the assumptions, limitations, and uncertainties associated with this method. Furthermore, the analysis performed in this study can be easily replicated by accessing the software and data products associated with this study, which we believe is important in its own right, since we intend ISARA to be used as an open source toolkit for field campaign scientists to use.
2. There has been previous literature attempting the same task, and this needs to be accounted for and mentioned. I bring here for example the IRRRA approach of Tsekeri et al (AMT, 2017). I am sure that by searching the literature the authors can also find other references on the same topic. The authors should explain how their paper fits within the existing

literature, and how similar or different it is from other approaches, indicating pros and cons of each. Note that “previous works” are also cited at line 547, but no references are given. Thank you for the comment. We have added more detail in the introduction and in the methods section that indicate how this work is a successor to previous works. While Tsekeri et al. (2017) did do a retrieval algorithm using in-situ dry measurements and probes, their algorithm requires the lidar data as part of this algorithm. This is distinctly different from the approach taken in this work that does the retrieval independently of the lidar measurements and then compares the aerosol properties afterward. We have made note in several places through the text that the coarse-mode aerosol properties are prescribed and the size-distribution of the coarse-mode is taken from wing mounted probe measurements and cited studies that have done analysis of coarse-mode aerosols with wing mounted probes.

3. The stated goal is a “rigorous external closure” (see line 4). I find this to be a real contradiction with the state-of-the-art achieved by the paper itself, given that it is a “preliminary consistency” and a “preliminary effort” (see lines 19 and 72). This may question whether more work is needed before a paper is submitted to AMT. The sentence at lines 546-548 seems to confirm this impression. Thank you for the comment. We have removed the word “preliminary” from this line since we have added several novel comparisons in the revised draft. These results agree within the uncertainties of the measurements. This sentence is now written as “While this study focuses on spherical, sulfate-dominated aerosol mixtures, its overall success demonstrates that ISARA has the potential to support systematic and streamlined closure of aerosol datasets across diverse field campaigns and aerosol regimes.”. The focus of this work is to prove the functionality of ISARA under conditions that are generally realistic but relatively ideal in terms of ignoring the possible presence of non-spherical aerosols within the marine environment. Furthermore, we state that rigorous external closure over a wide range of all possible marine and terrestrial scenarios will require future studies and changes to some of the assumptions made during this study, but will not require a fundamental change to the underlying software (i.e., ISARA).

4. The computations are done for a specific atmospheric scenario only, with a mixture of sulfate and organics in the fine mode and sea salt in the coarse mode (lines 104-109 with a clear mention that “this will limit the scope of this study”). Whereas this is fine in itself, it removes generality away from the ISARA approach and from its stated goal of a generally applicable approach. It is important to state the limited scope from the onset of the paper, rather than add the limitation further down the line. Moreover, given that ACTIVATE operated also in Bermuda (line 136) why is the influence of dust aerosols not also investigated? Thank you for the comment. We have added the following text to the abstract to list the assumptions used in this study more clearly from the start: “While this study focuses on spherical, sulfate-dominated aerosol mixtures, its overall success demonstrates that ISARA has the potential to support systematic and streamlined closure of aerosol datasets across diverse field campaigns and aerosol regimes.”. Furthermore, while dust may have been observed during ACTIVATE, it is outside the scope of this paper to analyze these more complex aerosol types until we are confident in the comparisons using more typical aerosol types. We have reiterated this point throughout the text.

5. After clarifying the issues affecting in-situ observations, the authors claim (lines 89-90) that “the ISARA attempts to overcome these limitations by estimating the contribution of coarse-mode particles”. It is unclear how this can be achieved, and I would state that this is not possible without a dedicated measurement for the coarse particles (e.g. using an open-path OPC). A CDP is later mentioned at lines 226-234, therefore the problem may perhaps simply be of being explicit and honest from the onset about the fact of using a dedicated measurement for the coarse mode. Thank you for the comment. We agree that this text was misleading and have changed the text to clarify that we are using dedicated probe measurements. In this case, we preferentially used the CAS, but relied on the CDP or FCDP where the CAS was not available.

6. Figure 9: ISARA shows underestimation. Figure 10: low level of closure. This does not look good for the ISARA method. Thank you for the comment. We do not feel that this underestimation is a critical weakness because the comparison shown yields results within a reasonable uncertainty. We have made this point evident throughout the text by adding

the corresponding retrieval uncertainties and standard deviations where possible. This algorithm represents a solid beginning, and future improvements are likely possible.

- 100 7. The sentence “a potential reason for detecting low-RH aerosol is the presence of smoke from fires” (lines 588-589) is in contradiction with the assumed marine environment, but is also presented in a way that makes it appear unjustified. Please note that water vapour is a combustion product, hence fires should contribute to increasing absolute humidity and not to decrease it. The whole narrative at lines 590+ appears incoherent: note that figure 13 shows fires and not smoke aerosols as stated here. It is surprising also to read that aerosols “are becoming drier due to colder temperatures” given that lowering the temperature (with constant WVMR) has the effect of increasing the RH. I agree with the sentence on
105 line 593 that “further work will be done” and I feel that it must be done before re-submitting the paper. [We agree that this text was erroneous and we have removed it. We now focus attention to the presence of winter conditions, which is far more important than any amount of water vapor generated from combustion. This is especially true because no smoke aerosols were detected in the profile and because the profile was located far from any source of smoke. If we expanded the scope of this study to include more atypical particles, the paper would be incredibly long and unfocused. Instead, we aim to demonstrate that this method works well for spherical aerosols found in marine environments, and that the overall approach is also useful for future research.](#)
- 110
- 115 8. Overall good agreement (line 602) feels an unjustified statement here. The same applies to the “successful retrieval” (line 622). The evidence is not so clear-cut. [Thank you for the comment. After performing extensive revisions throughout the paper, we feel that overall these statements are justified but have made it a point to indicate which properties were shown to be consistent and which were not.](#)

MINOR COMMENTS:

- 120 5. “In-situ instruments cannot efficiently sample coarse-mode particles due to limitations in the inlet cutoff diameter” (line 65). This statement is incorrect given that there are several open-path airborne instruments such as the CDP, the CAPS, etc. that overcome these limitations (initially developed for cloud particles and later extended to use in aerosol layers). It is well-known that the FAAM research aircraft has successfully sampled coarse and giant dust particles up to 300 μm diameter with such probes (see e.g. Ryder et al, ACP, 2015; Marengo et al, ACP, 2018). [Thank you for the comment. We have changed the text throughout the paper to clarify that we are using the coarse-mode measurements in the same way as the sources you have cited.](#)
- 125
- 130 6. The definition of the fine-mode (0.09-1 μm dry diameter) and coarse-mode ($> 1 \mu\text{m}$ ambient diameter) regimes (lines 83-84) is weird given that there could be an overlapping zone between fine and coarse (particles with dry diameter $< 1 \mu\text{m}$ and ambient diameter $> 1 \mu\text{m}$ would be in both regimes). I would suggest to consistently refer to either the dry or the ambient diameter for discriminating the two modes. [Thank you for the comment. This was an error on our part. We have changed the word dry diameter to ambient diameter.](#)
- 135 7. Line 6: the symbol κ is undefined. At the end of page 14, the relationship between κ and $f(\text{RH})$ must be explained. [Thank you for the comment. We have decided to remove the explicit parameterization of \$\kappa\$ from the abstract and moved it to the introduction, which is expanded further in Section 2.4. In this section, we explain that \$\kappa\$ is related to scattering and absorption coefficients through the growth factor equation and relative humidity.](#)
- 140 8. Joint flights (line 135): it is unclear which two airplanes performed joint flights. [Thank you for the comment. Because we address this directly in the following paragraph, we have removed the word “joint” from this sentence.](#)

9. “The novel vertically-resolved aerosol particle number concentration” (line 184): I suggest to add the word “estimate” to this statements, given that number concentration from lidar is estimated based on assumptions. Thank you for the comment. We have added the “estimate” to this sentence.
- 145 10. Inlet cutoff at 5,000 nm (line 199): indicate how the cutoff diameter was determined. Thank you for the comment. This information was determined by using ground station measurements collocated to the aircraft measurements. Although it is outside the scope of this paper to describe this process, we have cited McNaughton et al. (2007) that evaluated this cutoff diameter.
- 150 11. For random variables such as the scale factor (line 365), the κ (line 367) and IRI (line 370), the random distribution used must be given. For the scale factor, you can observe the random distribution in Fig. 5. For the random distributions of κ and IRI you can observe the random distribution in Fig. 5. We feel that showing the random distributions of these variables would require an additional separate figure that is unnecessary given the two figures already present with this information embedded.
- 155 12. Nephelometers (line 202) are being mentioned before being introduced (line 213 and following). I suggest that the full instrument set must be introduced before discussing their details such as installing a cyclone. Thank you for the comment. We have added Section 2.1 to address this concern.
- 160 13. Equation 1 using RH of 20 and 80% is inconsistent with having observations at 40 and 85% (line 215). Please clarify. We have changed the text and redefined $f(\text{RH})$ more clearly in Eq. 15. The measurement of humidified scattering was done at 85% RH, however, we use γ to calculate $f(\text{RH})$ for the 80/40% RH
- 165 14. Line 248: “the methods standardized by the merging tool”. It is unclear what the authors refer to. Thank you for the comment. We agree that this text was ambiguous. We have changed the sentence to be: “Data in their native resolution are averaged to 45 seconds using the NASA merging tool.”. A description of how the merging tool works is found in <https://www-air.larc.nasa.gov/missions/etc/onlinemergedoc.pdf>, which is cited in the previous sentence of the text.
- 170 15. Symbols: there is an unclear use of symbols N_a and n_0 (seeming to refer to the same variable). Also, in lines 293 and 294 C_{calc} refers to different variables, hence I suggest using different symbols. Thank you for the comment. We have corrected our terminology throughout the paper to make the distinction between N and n^o clear. We have also corrected the C_{calc} term to have the “abs” or “scat” subscript where appropriate.
- 175 16. Equations 3 and 7: the bounds of the integral should be $\log(D_{\text{min}})$ and $\log(D_{\text{max}})$, and not $d\log(D_{\text{min}})$ and $d\log(D_{\text{max}})$. Thank you for the comment. We have corrected this error.
- 180 17. Line 270: remote sensing sensors are normally considered more sensitive to the surface area and not the volume. Please correct. Thank you for the comment. As another reviewer pointed out, this statement was not fully correct. The sensitivity of the remote sensor to surface area or volume depends on the observational wavelength. We have added more detail with the following text: “ In general, remote sensors are not as sensitive to particle number concentration as they are to particle surface area and volume concentrations. For particles larger than the remote sensor’s observing wavelength, the remote sensor is most sensitive to particle cross-sectional area (i.e., surface area concentration). For particles smaller than the observing wavelength, the remote sensor is most sensitive to volume concentration.”.

- 185 18. Stitched data (line 279): this must be documented. Thank you for the comment. We have explicitly documented this step through the following text: “After this step, the SMPS and LAS size distributions are stitched at a diameter of 94 nm, which is the upper bound of the size-range by the SMPS and the lower bound of the LAS size-range.”.
- 190 19. Full range of particle sizes (line 279): this is incompatible with the inlet cutoff. Please clarify. Thank you for the comment. We felt that this sentence was redundant and unclear, so we have removed it. This information is discussed directly above in the previous paragraph.
- 195 20. Figure 1 caption: I suggest to use percentiles instead of minimum and maximum to reduce the influence of outliers. Thank you for the comment. We have changed Figure 1 to reflect the 10th and 90th percentiles and updated the corresponding caption.
- 200 21. CRI averaging (line 291): it is unclear over which dataset the averaging is being done. Thank you for the comment. In the following sentence, we now state: “For a CRI to be valid for averaging, all three of the computed scattering coefficients must be within 20% of the corresponding measured dry scattering coefficient $\left(\frac{|C_{\text{scat,calc}} - C_{\text{scat,RH=40}}|}{C_{\text{scat,RH=40}}} < 0.2\right)$ and all three of the calculated absorption coefficients must be within 1 Mm^{-1} of the measured absorption coefficients $(|C_{\text{abs,calc}} - C_{\text{abs,RH=40}}| < 1 \text{ Mm}^{-1})$ ”. This now unambiguously defines what conditions are required for a CRI to be considered valid for averaging.
- 205 22. Co-location (line 401). As observations are done on-board the same platform, they are certainly co-located, therefore this is unclear. Thank you for the comment. For clarity, we have removed the word “collocation” from this sentence.
- 210 23. Line 404: I suppose that data are removed when the cost function is larger than a threshold, not lower. Please correct. Thank you for the comment. We have corrected this error and replaced the “<” with “>” in addition to removing the extra “to be” in the text.
- 215 24. “coarse mode AOD is limited to < 0.1”: why? As stated in the text directly above this quote we state “To limit the presence of coarse-mode aerosol particles in this analysis”. We have added the following text to the preceding paragraph to better motivate this decision: “Because a significant presence of coarse-mode particles would be atypical within the altitudes that are sampled by the Falcon, we attempt to limit the amount of coarse particles in the columns of compared data.”.
- 220 25. LDR (line 407): clarify if you refer to VLDR or PLDR, because they bare not the same. The threshold at 0.13 (line 410) is not so small, at least for VLDR, and could indicate dust presence. Thank you for the comment. We have clarified the mathematical definition of LDR. Because the HSRL-2 description already stated that this is aerosol-specific data, we feel that an additional distinction between PLDR and VLDR is not needed.
- 225 26. Successful retrieval rate (line 482): please clarify if the retrieval rate means that a solution is found, or that the solution found is close to the observations). Thank you for the comment. This is simply the number of successful retrievals over the number of attempts, where the success is based on the maximum thresholds in the deviation of scattering and absorption. The text has been clarified by adding “ $\left(\frac{\text{number of successful retrievals}}{\text{number of attempts}}\right)$ ” to this sentence.

- 230 27. Lines 483-486. The authors raise an important limitation and they should perform a more in-depth analysis to disentangle the causes. Note that for measurement noise (mentioned by the authors) this should be possible to address with the current data. Concerning the issue of non-sphericity, this can be run using MOPSMAP. Thank you for the comment. We agree that more in-depth analysis could be done for future studies, but we feel it would distract from the focus of the paper on more typical marine cases. We have also added a discussion to further support the low signal issues that were observed.
- 235 28. Acronyms (NRMSD, MRB, NMAD) must be explained when first used. Thank you for the comment. We have removed the discussion of MAD from the text as it is redundant to NRMSD. Additionally, we have ensured that all consistency statistics are defined when first used.
- 240 29. Line 565: specify that you refer to case #10 of table 5, because there are two flights on this date. Thank you for the comment. As mentioned above, we have removed the extra case from this table and updated all the text accordingly to be consistent.
- 245 30. “are likely organic and sulfate-dominated mixtures”: this sentence is unexplained and unsupported by evidence Thank you for the comment. We have added the following text to support this statement: “This information indicates that the fine-mode particles being sampled were anthropogenic in origin and are likely sulfate-dominated mixtures with organic aerosol species. This is what would also be expected for this marine environment based on climatological evidence (Braun et al., 2021).”.
- 250 31. “marine environment” (line 588): This statement seems inconsistent with the Hysplit trajectories showing continental influence. Thank you for the comment. We do not feel that these are contradictory statements. A notable feature of the marine environment of the Northwest Atlantic is the persistent influence of anthropogenic outflow. We have clarified this in the previous response.
32. “the data from case 12 are also shown in Table 5” (line 597): why? Thank you for the comment. We agree that showing these data here was unnecessary and we have removed this case from the included tables.
- 255 33. “1064 m”: it is actually 238-4499 m Thank you for the comment and for catching this typo. We have corrected the value to be 4261 m.
- 260 34. Table 5 caption should read “case studies 10 and 12” Thank you for the comment. We have changed this table to include only data from case study 10, which is now case study 7 in the revised draft.