

Fungi present distinguishable isotopic signals in their lipids when grown on glycolytic versus tricarboxylic acid cycle intermediates

Stanislav Jabinski^{1,2} Vítězslav Kučera^{3,4}, Marek Kopáček^{1,5}, Jan Jansa⁴, Travis B. Meador^{1,2,5*}

¹University of South Bohemia, Faculty of Science, Department of Ecosystem Biology, Branišovská 1760, 370 05 České Budějovice, Czechia

²Institute of Soil Biology and Biochemistry, Biology Centre CAS, Na Sádkách 7, 370 05 České Budějovice, Czechia

³Charles University, Faculty of Sciences, Albertov 6, 128 00 Praha, Czechia

⁴Institute of Microbiology CAS, Vídeňská 1083, 142 20 Praha, Czechia

⁵Institute of Hydrobiology, Biology Centre CAS, Na Sádkách 7, 370 05 České Budějovice, Czechia

Correspondence to: Travis B. Meador (travis.meador@bc.cas.cz)

Abstract. Microbial activity in soils controls both the size and turnover rates of large carbon (C) inventories stored in the subsurface, having important consequences for partitioning of C between terrestrial and atmospheric reservoirs as well as recycling of mineral nutrients such as nitrogen or phosphorus (often bound to the C) that support plant growth. Fungi are major decomposers of soil organic matter (SOM); however, uncertainty about the identity of predominant C substrates that fuel their respiration confound models of fungal production and SOM turnover. To further define the signals of microbial heterotrophic activity, we applied a dual hydrogen (H) and C stable isotope probing (SIP) approach on pure fungal cultures representing the phyla Ascomycota, Basidiomycota, and Zygomycota growing on monomeric (glucose, succinate) or complex substrates (tannic acid, β -cyclodextrin). Our findings demonstrate that the investigated species incorporated only minor amounts of inorganic C (provided as bicarbonate) into their membrane lipids, amounting to < 3% of lipid-C, with no consistent patterns observed between species or growth substrates. The net incorporation of water-derived H (i.e., a_w) into lipids also did not differ significantly between incubations with monomeric versus complex substrates; however, growth on succinate elicited significantly higher a_w values than glucose or β -cyclodextrin. This finding suggests that ^2H -SIP assays have the potential to distinguish between microbial communities supported predominantly by substrates that are catabolized by the tricarboxylic acid cycle versus glycolytic pathway. Furthermore, the average a_w value of heterotrophic fungal incubations [0.69 ± 0.03 (SEM)] is consistent with that observed for bacterial heterotrophs, and may be applied for upscaling lipid-based estimates of fungal production in environmental assays.

Short Summary

Microbial production is a key parameter in estimations of organic matter cycling in environmental systems, and fungi play a major role as decomposers. In order to investigate fungal production and turnover times in soils, we incubated fungal pure cultures with isotopically labelled water and bicarbonate to investigate growth signals encoded into lipid biomarkers, which can be applied to improve flux estimates in environmental studies.

1 Introduction

Soil organic matter (SOM) is the major reservoir of carbon (1580×10^{15} g C) in the biosphere, and active microbial populations act to redistribute this C to other reactive reservoirs, such as the atmosphere (Carson et al., 2001; Grinhut et al., 2007). Major uncertainties in modeling C and climate dynamics stem from insufficient knowledge on the controls of SOM degradation and transformation (Ciais et al., 2014; Lindahl and Tunlid 2015). Saprotrophic soil fungi are one of the major decomposers in soils, are known to degrade naturally occurring complex molecules such as lignin (Kirk & Farrell, 1987; Fioretto et al., 2005; Baldrian et al., 2011), cellulose (Šnajdr et al., 2011) and humic substances (Grinhut et al., 2007), but are also reported to compete for accessible plant photosynthate excreted by roots (De Boer et al., 2005; Högberg et al., 2001; Smith & Read, 2008). Despite the unique and important fungal niche in biogeochemical cycles, their contributions to SOM cycling remains poorly constrained (Frey 2019; Grinhut et al., 2007). Furthermore, heterotrophic organisms feeding on organic substrates to gain energy and build biomass are also known to fix a variable amount of inorganic C (IC), in order to replenish intermediates in the tricarboxylic acid (TCA) cycle (Kornberg 1965). It has been suggested that 2 - 8% of the biomass C in heterotrophs originates from IC incorporated through anaplerotic carboxylation reactions (Romanenko 1964; Roslev et al., 2004; Braun et al., 2021). Awareness of these processes has existed for decades (Kornberg 1965; Sorotkin 1966). Yet, the relevance and metabolic controls on heterotrophic IC fixation remains poorly understood, partly due to the lack of reliable estimates for most organisms and habitats (Braun et al., 2021).

Advanced analytical techniques now allow microbial taxa to be linked to specific processes in environmental studies by measuring incorporation of stable isotopes into biomarkers (Boschker et al., 1998; Dumont and Murrell, 2005; Kreuzer-Martin, 2007), such as fungal and bacterial membrane lipid fatty acids (Treonis et al., 2004; Willers et al., 2015) or other biomarkers (Boschker and Middelbourg, 2002). Previous studies have demonstrated that variability in the ^2H composition of microbial lipids is redundant with that of environmental water (Hoefs, 2018; Kopf et al., 2015), and stable isotope probing (SIP) assays applying enrichments in $^2\text{H}_2\text{O}$ have proven to be a useful tracer of microbial activity in a diverse range of environments (Canarini et al., 2024; Caro et al., 2023; Fischer et al., 2013; Kellermann et al., 2012; Wegener et al., 2016; Warren 2022; Wu et al., 2018). It is important to distinguish these ^2H -labeling applications from parallel investigations aiming to determine the kinetic and equilibrium isotope fractionations resulting from biosynthetic incorporation of water hydrogen (water-H) into individual compounds (i.e., ϵ ; cf. Hayes 2004). While ϵ values can be indicative of an underlying metabolic processes or environmental change (Osborn et al., 2011; Sachse et al., 2012; Zhang et al., 2009), the large range of ϵ that have been reported

for fatty acids of aerobic, heterotrophic bacteria [$-330 > \epsilon > +360$ ‰; see review by Pilecky et al. (2025)] arise, in part, because water-H is not the only source of H for biomolecules, complicating the interpretation of such values observed in natural systems. In contrast, the water-H incorporation factor (or water-H assimilation efficiency; a_w) accounts for variability in the proportion of H derived from water as well as ^2H fractionation during biosynthesis (cf. Zhang et al., 2009; Kopf et al. 2015; see Methods), and has been valuable for SIP approaches to (i) estimate microbial production rates in natural systems (e.g., Wegener et al., 2022; Caro et al., 2023), and (ii) establish characteristic growth patterns of microbial populations that can access a large diversity of substrates, such as methanogens (Wu et al., 2020).

To fully exploit the potential of SIP experiments, a dual-SIP approach was developed to track total microbial production by adding heavy water ($^2\text{H}_2\text{O}$) together with ^{13}C -labeled IC, enabling simultaneous estimates of total and autotrophic metabolism, respectively (Wegener et al., 2012; Wu et al., 2020). Recently, Jabinski et al. (2024) validated an innovation of the dual-SIP assay by using rapid pyrolysis of fungal biomass to determine the stable C and H isotopic composition of fungal lipids, and demonstrated that water-H and IC assimilation signatures could successfully distinguish between fungal ecotypes growing on glucose or glutamic acid as the C source. The aim of the current study was to further assess the controls on water-H and IC incorporation into lipids and expand our knowledge for interpreting environmental signals by applying the dual-SIP assay on a broader range of pure fungal cultures and growth substrates, including labile monomers versus more complex, high molecular weight molecules. We hypothesized that (i) the incorporation of IC and water-H into the fungal fatty acid biomarker $\text{C}_{18:2}$ will be similar for fungal species growing on the same substrate, and (ii) that IC and water-H incorporation will distinguish between growth on labile versus more complex C substrates.

2 Methods

2.1 Cultivation & Harvesting

Fungal pure cultures of two Basidiomycota [*Paxillus involutus* (PI, strain SB-22); *Phanerochaete chrysosporium* (PC, strain CCM8074)], two Zygomycota [*Mortierella* sp. (MO, strain RK-38); *Umbelopsis* sp. (UM, strain RK-43)] and two Ascomycota [*Penicillium janczewskii* (PJ, strain BCCO20_0265); *Paecilomyces lilacinus* (PL, strain DP-23)] were incubated in 500 mL Schott bottles at 25 °C in the dark. Liquid mineral media (50 mL) was adapted after Bukovská et al. (2018) with the vitamins left out, and was inoculated with approximately 10^6 spores, or, for Basidiomycota, a hyphal block $< 0.5 \text{ cm}^3$ recovered from a previous culture using the same cultivation medium solidified with agar (1.5%).

The growth medium contained per liter: 4 g organic C in various forms ($\text{C}_6\text{H}_{12}\text{O}_6$ glucose; $\text{C}_4\text{H}_6\text{O}_4$ succinic acid; $\text{C}_{42}\text{H}_{70}\text{O}_{35}$ β -Cyclodextrin or $\text{C}_{76}\text{H}_{52}\text{O}_{46}$ tannic acid), 0.01 g $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 2 g KH_2PO_4 , 0.5 g $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 0.1 g NaCl, 0.1 g CaCl, 2.5 g $(\text{NH}_4)_2\text{SO}_4$, 0.45 g NaHCO_3 and 1 mL of a mixed solution (per liter: 0.5 g H_3BO_3 , 0.04 g $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, 0.1 g KI, 0.4 g $\text{MnSO}_4 \cdot 5\text{H}_2\text{O}$, 0.2 g $\text{NaMoO}_4 \cdot 2\text{H}_2\text{O}$, 0.4 g $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$). The pH of the medium was adjusted to 4.5 before inoculation. Dual-SIP experiments were performed using ^{13}C -bicarbonate (^{13}C -DIC, $\text{NaH}^{13}\text{CO}_3$) and deuterated water ($^2\text{H}_2\text{O}$). Each fungal strain was grown in triplicate with non-labeled substrates (Treatment I), with $\delta^2\text{H}$ of the medium water adjusted to 100‰ and

AT¹³C = 10% of ¹³C-DIC (Treatment II), 200‰ δ²H and 10% ¹³C-DIC (Treatment III), and 400‰ δ²H and 10% ¹³C-DIC (Treatment IV). The concentration of the bicarbonate in the cultivation medium was 0.1 g L⁻¹. The Schott bottles were closed with a rubber stopper in order to prevent the labeled ¹³C-DIC from outgassing, and ample headspace was provided to maintain oxic conditions throughout the growth experiment. Fungal growth was monitored via the accumulation of CO₂ in the headspace, and we aimed to harvest when CO₂ levels reached 10%; however, without preliminary knowledge of the fungal growth dynamics, some cultivations exceeded this level more quickly than they could be harvested. To harvest the fungal biomass, mycelia were separated from the growth medium via vacuum filtration through 5 µm Isopore polycarbonate filters (47 mm diam, Merck catalogue number TMTP04700) and the cultivation medium was collected into a sterile 50 mL tube. Thereafter, the mycelium was washed with ample MilliQ water, transferred to pre-weighed, sterile 50 mL tubes; the fresh weight of the biomass was recorded, and the samples were frozen at -80 °C until lyophilization. A subsample of the cultivation medium was also frozen at -80 °C and the rest was used to determine pH post-incubation. After lyophilization, the dry weight of each sample was determined and stored at -20 °C until further analysis.

2.2 Measurements

2.2.1 Headspace CO₂ concentration and isotope composition

Samples of headspace (0.3 mL) were collected weekly from each bottle into helium flushed 12 mL exetainer vials (Exetainer, Labco Limited, UK) and analyzed for their CO₂ concentration and ¹³C/¹²C isotopic ratio using GasBench II equipped with a single cryo-trap connected to Delta V Advantage isotopic ratio mass spectrometer (IRMS) via Conflo IV (Thermo Scientific, Bremen, Germany). Ambient air (with its CO₂ concentration measured using LiCor 850 gas analyzer previously) was used as a standard for CO₂ concentration measurements, whereas a laboratory cylinder with 0.1% CO₂ in helium was used as a standard for isotopic composition (δ¹³C = -2.86 ‰). The analytical precision was below 1‰. Data were analyzed and exported using Isodat 3.0 software.

2.2.2 Medium water (δ²H₂O)

Liquid samples were transferred into 1.5 ml glass vials (32 x 11.6 mm, Fischer Scientific) and then measured using Triple Liquid Water Isotope Analyzer (Los Gatos Research), which is based on the principle of high-resolution laser absorption spectroscopy. Samples were dispensed into the instrument using an autosampler (PAL3 LSI, ABB company) and a 1.2 µL syringe (Hamilton). Samples were measured and evaluated against prepared laboratory standards of known isotopic composition. The isotopic ratios of these laboratory standards were verified by measuring against international standards (VSMOW2, SLAP2) made by the IAEA. For quality control purposes, the measurements of the samples were also interspersed with periodic measurements of the prepared verification samples with known isotopic composition. The final isotopic composition (δ²H) was determined using LIMS software. Analytical precision of δ²H was <1.5‰.

Water sampled from incubations with tannic acid could not be measured using the laser, as described above, due to its high organic carbon content, and was rather measured via a GasBench II device (Thermo Scientific, Bremen, Germany; Application Note: 30049). Medium water samples (200 μ L) were added with a platinum catalyst to 12 mL exetainer vials (Exetainer, Labco Limited, UK). The headspace was flushed with 1% H_2 in He at approximately 100 mL min^{-1} for 6 min. After an equilibration time of over 40 min, the samples were measured by purging the exetainer using a double-holed needle with helium into a 250 μ L sample loop. The sample was then injected and separated via a Carboxen PLOT 1010 (0.53 mm ID; Supelco, Bellefonte, USA) held at 90 $^{\circ}C$ with a flow rate of ~ 2.2 mL min^{-1} , and then introduced into the MAT253 Plus IRMS via a Conflo IV interface. Each sample was injected three times during one analysis. The isotopic composition was determined using Isodat 3.0 software against the corresponding H_2 working gas (-239‰ for δ^2H) and the values were corrected and normalized using international standards VSMOW2 (0‰ for δ^2H), SLAP2 (-427.5‰ for δ^2H), USGS53 ($+40.2\text{‰}$ for δ^2H) and GFLES-2 (159.9‰ for δ^2H). The analytical precision was around 1 ‰ .

2.2.3 Carbon ($\delta^{13}C$) substrate analysis

Substrates (~ 100 μ g) were weighed into tin capsules (8×5 mm, Sercon, Crewe, UK) and placed in a helium-flushed carousel autosampler, then introduced to an Elemental Analyzer IsoLink device (EA IsoLink CNSOH, Thermo Scientific, Bremen, Germany) equipped with a CHN/NC/N EA combustion/reduction reactor (Sercon, Crewe, UK) heated to 1020 $^{\circ}C$. A pulse of oxygen was introduced to the reactor simultaneously with the sample. The sample gases were quantified via a thermal conductivity detector (TCD) and then introduced to a MAT 253 Plus isotope ratio mass spectrometer (IRMS; Thermo Scientific; Bremen, Germany) via the open split of a Conflo IV interface, with helium as the carrier gas. The isotopic composition was determined using Isodat 3.0 software against the corresponding CO_2 working gas (-4.191‰ for $\delta^{13}C$), and the values were corrected for linearity and normalized to the VPDB scale using international reference material IAEA-600 (-27.771‰ for $\delta^{13}C$). The analytical precision was $<0.04\text{‰}$.

2.2.3 Pyrolysis GC for lipid analysis

The pyrolysis unit Shimadzu 3030D (Shimadzu, Kyoto, Japan/ Frontier Laboratories, Fukushima, Japan) was installed on top of the GC Trace1310 gas chromatograph SSL injector (Thermo Scientific, Bremen, Germany) and the GC was equipped with an SLB-IL60 column (non-bonded; 1,12-Di(tripropylphosphonium)dodecane bis(trifluoromethanesulfonyl)imide phase, 30 m, 0.25 mm ID, 0.20 μ m df, Supelco, Bellefonte, USA). The furnace temperature was 650 $^{\circ}C$ and the interface temperature was 370 $^{\circ}C$. The injector temperature was 360 $^{\circ}C$ and the GC oven was held at 80 $^{\circ}C$ for 1 min then ramped to 175 $^{\circ}C$ at 15 $^{\circ}C$ min^{-1} , then ramped to 195 $^{\circ}C$ at 2 $^{\circ}C$ min^{-1} , then ramped to 300 $^{\circ}C$ at 10 $^{\circ}C$ min^{-1} , and finally held at 300 $^{\circ}C$ for 7 min. Helium was used as the carrier gas, with a constant flow of 1.5 mL min^{-1} , a split ratio of 40, and a split flow of 26.7 mL min^{-1} . The column flow was split via a multichannel device to acquire MS and isotopic data simultaneously from one injection. The

GCMS (ISQ QD; Thermo Scientific, Bremen, Germany) ion source was set to electron impact ionization mode (EI) at 70 eV and a scan range of m/z 50 – 500 with a scan time of 0.2 sec⁻¹. Scanning started after 8 min to avoid the solvent peak in the MS. The transfer line temperature was set to 300 °C and the ion source was set to 250 °C.

The samples (lyophilized biomass, 0.1 – 1.3 mg) were weighed into an ultra-clean stainless steel Eco-Cup LF (Frontier Laboratories, Fukushima, Japan), which were burned with a torch before usage to remove contaminants. Immediately prior to sample injection, 30 µL of trimethylsulfonium hydroxide (TMSH) was added on the sample to increase the volatilization by transforming the fatty acids into fatty acid methyl esters (FAMES) and improve measurement sensitivity. Identification of FAMES was performed using fragmentation patterns and the NIST 14 library.

Stable C and H isotope compositions of FAMES from aliquots of the same biomass were separately determined by splitting the flow from the GC column to a GC-IsoLink II reactor, coupled to a MAT253 Plus IRMS via a ConFlo IV interface. Values are expressed in standard delta notation ($\delta^{13}\text{C}$ and $\delta^2\text{H}$). MS information was simultaneously acquired by use of the multi-channel device described above. For conversion of FAMES and ergosterol to CO₂, the combustion reactor (nickel oxide tube with CuO, NiO, and Pt wires) was set to 1000 °C. For conversion of FAMES and ergosterol to H₂, the pyrolysis reactor (aluminum tube) was set to 1420 °C. The FAMES were identified by their retention times and fragmentation patterns. The isotopic composition was determined using Isodat 3.0 software against the corresponding CO₂ or H₂ working gas (-4.191‰ for $\delta^{13}\text{C}$, -239.5‰ for $\delta^2\text{H}$). Isotope corrections for instrument drifts, linearity, and normalization to the VPDB or VSMOW scales were performed according to the response of USGS70 (-30.53‰ for $\delta^{13}\text{C}$, -183.9‰ for $\delta^2\text{H}$) and USGS72 (-1.54‰ for $\delta^{13}\text{C}$, 348.3‰ for $\delta^2\text{H}$) reference standards. The analytical precision was < 0.5‰ and < 10‰ for $\delta^{13}\text{C}$ and $\delta^2\text{H}$, respectively.

2.3 Calculations

Carbon use efficiency (CUE) for the growth experiments was calculated by normalizing the amount of C in biomass by that plus C that accumulated as CO₂ [CUE = biomass-C / (CO₂-C + biomass-C)]. The $\delta^{13}\text{C}$ values of fungal biomarker C_{18:2} was determined as standard delta values (‰). The IC incorporation into the biomarker (%IC) was calculated based on the following equation:

$$\%IC_{(assimilation)} = \frac{{}^{13}F_{lipid} - {}^{13}F_{lipid,control}}{{}^{13}F_{DIC(medium)} - {}^{13}F_{substrate}} \times 100 \quad (\text{Eq. 1})$$

Equation 1: Inorganic carbon (%IC) assimilation was calculated as the difference in the ¹³C atom fraction of the lipids harvested at the end of the labeling experiment (¹³F_{lipid}) compared to the lipids harvested at the end of the natural abundance treatment (¹³F_{lipid,control}), relative to the difference between the mixing-weighted average ¹³C atom fraction of dissolved inorganic C (¹³F_{DIC}, cf. Text S1) and the ¹³F of the substrate. F was calculated as ¹³F = (R^{13/12})/(R^{13/12} + 1), where R is re-calculated from the $\delta^{13}\text{C}$ ratios reported by Isodat Software following measurement by IRMS ($\delta^{13}\text{C} = ([R^{13/12}]_{sample}/[R^{13/12}]_{ref} - 1) * 1000$ (modified after Boschker & Middelburg 2002; Wegener et al., 2012).

189 The water H assimilation efficiency (a_w) for fungal biomarker $C_{18:2}$ was approximated from the regressions of the hydrogen
190 isotope composition of individual fatty acids $^2F_{FA}$ and that of medium water ($^2F_{water}$), according to Kopf et al. (2015), after
191 Zhang et al. (2009). Briefly, the hydrogen isotopic compositions of microbial fatty acids produced by an organism generally
192 follow the isotopic composition of environmental water, and are related to the mole fraction of H contributed from water in
193 the cultivation medium (f_w) and the net hydrogen isotope fractionation between fatty acids and water ($\alpha_{fa/w}$), where $a_w = f_w \times$
194 $\alpha_{fa/w}$. Whereas a_w can be determined experimentally, the latter terms cannot be independently determined for heterotrophic
195 growth (Kopf et al., 2015). The traditional isotope effects and $\epsilon_{C18:2/water}$ and $\alpha_{C18:2/water}$ were calculated after Hayes (2004),
196 where $\alpha_{C18:2/water} = [(\delta^2H_{C18:2} + 1000) / (\delta^2H_{water} + 1000)]$ and $\epsilon_{C18:2/water} = (\alpha_{C18:2/water} - 1) \times 1000\text{‰}$.
197 Notably, calculation of %IC and a_w consider relative increases in $^{13/12}C$ or $^{2/1}H$ composition, and thus do not require a correction
198 for the methyl group added during derivatization of the fatty acid. The $\epsilon_{C18:2/water}$ values determined in this study were not
199 corrected for the additional methyl group and add an inaccuracy to the reported values, as further discussed below.

201 3 Results

203 3.1 Fungal growth and CO₂ production

204 All fungal species were pure cultures, which were incubated in a mineral medium with either glucose, succinate, β -cyclodextrin,
205 or tannic acid serving as the sole organic C source. Growth was monitored by the evolution of CO₂ into the headspace, which
206 ranged from 0.36% (no respiration of substrate) to a maximum of 35%, after incubation times ranging from 5 to 160 days (Fig.
207 1). The pH of the media in all incubations ranged from 2 to 5.5 at the time of harvest, with a general trend of decreasing pH
208 with increasing CO₂; however, the trend was opposite when succinate was the carbon source, with pH increasing from 4 to
209 5.5. For samples that produced sufficient biomass, the dry biomass of harvested fungal hyphae ranged up to 250 mg, and at
210 least 30 μ g dry biomass was used to analyze fungal membrane fatty acids by Pyr-GC-IRMS. Only the Ascomycota species PL
211 and PJ grew sufficiently on each tested substrate to produce enough biomass for stable isotope analysis. Incubations of
212 Zygomycota species with glucose or succinate also yielded sufficient dry biomass, and only UM was able to grow on β -
213 cyclodextrin. Zygomycota species produced neither CO₂ nor biomass when incubated with tannic acid. The Basidiomycota
214 typically exhibited the slowest growth, and both species (PI and PC) only produced enough biomass when grown on glucose.
215 The headspace CO₂ levels in Basidiomycota incubations with succinate increased to a maximum of ~ 2%, but only PI yielded
216 sufficient biomass for analysis. PC grew sufficiently on β -cyclodextrin, with CO₂ levels increasing to a maximum of 3%, while
217 CO₂ remained < 0.6% in PI incubations.




Figure 1. Growth of fungal species on each substrate as indicated by production of CO₂ versus days of incubation (left panels) or dry biomass (right panels). Filled symbols indicate samples for which the C_{18:2} biomarker was measured by Pyr-GC-IRMS. Colors represent the Ascomycota species *Penicillium janczewskii* (PJ, dark blue) and *Paecilomyces lilacinus* (PL, light blue), Zygomycota *Mortierella* sp. (MO, dark green) and *Umbelopsis* sp (UM, light green), and Basidiomycota species *Phanerochaete chrysosporium* (PC, orange) and *Paxillus involutus* (PI, yellow). The symbols denote incubations with glucose (circles), succinate (squares), tannic acid (diamonds), or β-cyclodextrin (triangles).

The growth substrates induced a wide range in CUE values, ranging from 0.1 to 0.6 (Fig. 2). Higher CUE values were typically observed for Ascomycota and Zygomycota species growing on glucose, and lower values for their growth on succinate and tannic acid. CUE estimated for Basidiomycota species was always low (< 0.15). The CUE range for growth on glucose (0.1-0.6), β-cyclodextrin (0.1-0.6), and succinate (0.2-0.5) were larger than observed for tannic acid (0.15-0.3).




Figure 2. Accumulation of biomass and headspace CO₂ for fungal species from three phyla growing on monomers or complex substrates. The right panel includes the same data on a log-log scale to depict the trends of fungi exhibiting minimal growth. Lines indicate CUE trajectories of 0.15 or 0.6, and were calculated assuming that fungal biomass was 44% C (w/w). Colors and symbols are redundant with Fig. 1. Filled symbols indicate samples that were analyzed by Pyr-GC-IRMS.

3.2 Stable isotopic composition of fungal lipids

Fungal respiration of the different (unlabeled) growth substrates led to decreasing $\delta^{13}\text{C}$ -CO₂ values as fungal biomass was produced, which followed a hyperbolic trend expected for the mixing of CO₂ from two different sources (Text S1; c.f., Kendall

and Caldwell, 1998). The atom % ^{13}C in control incubations with no fungal inoculum was measured at the latest time of harvest of inoculated incubations and stayed below 0.4‰, except for incubations with tannic acid, where it ranged between 2‰ and 3‰; the $\delta^{13}\text{C}$ values of the substrates were glucose = -26.5‰; succinate = -28.3‰, tannic acid = -27.4‰, β -cyclodextrin = -10.6‰. The mixing relationship was modeled using all CO_2 data, across all incubations, and integrated to approximate the mixing-weighted average ^{13}F value of IC for each incubation (cf., Text S1, Fig. S2), which was finally applied in the denominator of Eq. 1 to estimate the fraction of lipid-C derived from IC. For incubations that produced sufficient fungal biomass for stable C isotopic analysis, the weighted average $\delta^{13}\text{C}$ values of IC that were applied in Eq. 1 ranged from 200 to 1400 ‰ (i.e., ~ 1.3 to 2.6 AT% ^{13}C), and was largely dependent on how much of the substrate was respired to CO_2 .


3.2.1 Carbon isotopes

The $\delta^{13}\text{C}$ values of fungal biomarker fatty acid $\text{C}_{18:2}$ (Table 1) produced under natural cultivation conditions with glucose (i.e., non-labeled; AT% $_{\text{DIC}}$ ~ 1‰) ranged from -24.1‰ to -21.2‰ across all strains (n = 6 species). As expected, $\text{C}_{18:2}$ harvested from the labeled incubations exhibited slightly higher $\delta^{13}\text{C}$ values (up to +11‰; PC grown on glucose) than the corresponding experiment amended with natural bicarbonate, likely owing to the incorporation of labeled IC into the $\text{C}_{18:2}$ fatty acid.

Table 1: $\delta^{13}\text{C}$ values of fungal biomarker $\text{C}_{18:2}$ harvested from incubations with non-labeled substrates (nat) or those amended with ^{13}C -labeled bicarbonate. Incorporation of inorganic C (%IC) was calculated based on Eq.1. Errors represent the standard deviation of replicate incubations. Not all fungal species grew sufficiently on all substrates and thus %IC could not have been determined (n.d).

Species	Glucose $\delta^{13}\text{C}$ (‰)		IC (%)	Succinate $\delta^{13}\text{C}$ (‰)		IC (%)	Tannic acid $\delta^{13}\text{C}$ (‰)		IC (%)	β -cyclodextrin $\delta^{13}\text{C}$ (‰)		IC (%)
	nat	+		nat	+		nat	+		nat	+	
<i>Paxillus involutus</i> (PI)	-21.5	-16.7	0.6 (± 0.2)	n.d	n.d	n.d	n.d	n.d	n.d	n.d	n.d	n.d
<i>Phanerodontia chrysosporium</i> (PC)	-24.1	-15.9	0.9 (± 0.3)	-27.3	-23.5	0.3 (± 0.1)	n.d	n.d	n.d	n.d	n.d	n.d
<i>Mortierella</i> (MO)	-21.9	-20.7	0.5 (± 0.1)	-31.7	-31.5	0.1 (± 0.1)	n.d	n.d	n.d	n.d	n.d	n.d
<i>Umbelopsis</i> (UM)	-21.2	-21.1	0.1 (± 0.0)	-30.6	-28.2	0.7 (± 0.2)	n.d	n.d	n.d	-21.7	-18.1	0.8 (± 0.3)
<i>Penicillium janczewskii</i> (PJ)	-23.2	-22.1	0.5 (± 0.2)	-30.1	-27.8	0.2 (± 0.0)	-25.9	-20.5	2.2 (± 0.5)	-20.7	-20.1	0.1 (± 0.2)
<i>Paecilomyces lilacinus</i> (PL)	-23.0	-23.5	n.d	-30.8	-29.9	0.1 (± 0.0)	-25.4	-25.0	0.1 (± 0.0)	-19.1	-17.6	0.7 (± 0.4)

260 The estimated incorporation of IC into C_{18:2} (%IC) typically ranged up to 1%; only PJ grown on tannic acid exhibited higher
 261 %IC values, which ranged up to 2.2% (Table 1, Fig. 3). There were no general trends observed in %IC with other measured or
 262 estimated parameters, including CUE; however, for the two species that were able to grow on tannic acid, %IC was positively
 263 correlated with the amount of CO₂ and biomass produced during the incubation ($R^2 > 0.85$, $n = 5$, $p < 0.01$).



264
 265
 266 **Figure 3. %IC values for fungal species respiring glucose (circles), succinate (squares), tannic acid (diamonds), or β -**
 267 **cyclodextrin (triangles). Colors represent individual fungal isolates as described in Fig. 1.**

268
 269 **3.2.2 Water assimilation factor (a_w)**

270 The “net” contribution of water hydrogen to lipid H is reported as the water hydrogen assimilation factor a_w (Kopf et al., 2015),
 271 and was estimated based on the slope of the linear regression line between H isotopic composition of lipid versus growth
 272 medium water (Fig. 4), which ranged from natural MilliQ ($\delta^2H = -45\text{‰} \pm 10\text{‰}$) to the labeled treatments ($65\text{‰} \pm 4\text{‰}$; $166\text{‰} \pm 10\text{‰}$;
 273 $368\text{‰} \pm 27\text{‰}$). The a_w values for the fungal biomarker C_{18:2} grown on glucose ranged from 0.37 ± 0.03 to 0.75 ± 0.06
 274 with an average value of 0.60 ± 0.05 ($n = 6$ species; \pm SEM). When grown on succinic acid, the a_w values for C_{18:2} harvested
 275 from individual species ranged from 0.78 ± 0.01 to 0.96 ± 0.02 with an average value of 0.83 ± 0.04 ($n = 4$ species; \pm SEM).
 276 When grown on tannic acid, the a_w values for C_{18:2} harvested from individual species ranged from 0.74 ± 0.06 to 0.77 ± 0.03 ,
 277 and when grown on β -cyclodextrin the a_w values for C_{18:2} ranged from 0.46 ± 0.03 to 0.68 ± 0.04 with an average value of 0.58
 278 ± 0.06 ($n = 4$ species; \pm SEM). The average a_w values for C_{18:2} for all substrates and species was 0.67 ± 0.04 (\pm SEM).

279
 280 **3.2.3 H isotope fractionation factors $\alpha_{C_{18:2}/\text{water}}$ and $\epsilon_{C_{18:2}/\text{water}}$**

The range of traditionally reported isotope effects $\alpha_{C18:2/water}$ and $\epsilon_{C18:2/water}$ (Sessions and Hayes, 2005) for all natural and 2H -labeled fungal growth experiments was 0.73 to 1.08 and -265 to +83 ‰, respectively (Table S1). PL growth on tannic acid exhibited the highest values (0.97 to 1.08 and -35 to +83 ‰, respectively; Fig. 5, Table S1), while for all other growth experiments, $\alpha_{C18:2/water}$ and $\epsilon_{C18:2/water}$ remained < 0.94 and -65 ‰, respectively. Considering only the growth experiments performed without label addition, the average (\pm SD) $\epsilon_{C18:2/water}$ values were not significantly different for fungi growing on glucose (-115 ± 34 ‰, $n = 13$), succinate (-111 ± 35 ‰, $n = 8$), or β -cyclodextrin (-133 ± 7 ‰, $n = 6$), but growth on tannic acid elicited significantly higher values ($+12 \pm 81$ ‰, $n = 5$; $p < 0.01$, two-tailed homoscedastic t-test). Within species comparisons revealed that both Zygomycota species (*Mortierella* and *Umbelopsis*) exhibited significantly more negative $\epsilon_{C18:2/water}$ values when grown on succinate (-127 ± 6 and -155 ± 8 ‰, $n = 2$) versus glucose (-78 ± 3 and -127 ± 5 ‰, respectively, $n = 2$; $p < 0.05$). The only other significant differences in $\epsilon_{C18:2/water}$ values were observed for growth of the Ascomycota species *P. lilanicus*, for which growth on tannic acid ($+71 \pm 18$ ‰, $n = 3$) yielded higher values than on glucose (-122 ± 15 ‰, $n = 2$), succinate (-84 ± 6 ‰, $n = 2$), and β -cyclodextrin (-132 ± 2 ‰, $n = 2$; all $p < 0.01$); the $\epsilon_{C18:2/water}$ values for growth on succinate and β -cyclodextrin were also significantly different ($p < 0.01$).




Figure 4. The water hydrogen assimilation factor (a_w values) estimated as the slope of the fractional ^2H abundance (^2F) in lipids (y-axis) versus medium water (x-axis). Data are shown for fungal biomarker $\text{C}_{18:2}$ produced during growth on the different substrates (glucose, succinic acid, tannic acid and β -cyclodextrin) and harvested from the different fungal isolates [*Paxillus involutus* (PI), *Phanerodonta chrysosporium* (PC), *Mortierella* sp. (MO), *Umbelopsis* sp. (UM), *Penicillium janczewskii* (PJ), and *Paecilomyces lilacinus* (PL)]. R^2 values for all slopes were > 0.97 .




Figure 5. The apparent isotope effect ($\epsilon_{C18:2/water}$) for fungi grown in medium having variable δ^2H_{water} composition. The colors and symbols are redundant with Fig. 1. For each species-substrate pair, the large range and/or decrease in $\epsilon_{C18:2/water}$ in 2H -enriched medium are expected, given the additional contributions of substrate-H and metabolic water-H to lipid-H during biosynthesis.

4 Discussion

4.1 Fungal growth dynamics

Collectively, the fungal incubation experiments included six species representing three different phyla growing on one of four substrates, and exhibited a large range in the relative amounts of CO_2 (0.2-34% v/v) and biomass produced (0-230 mg dry weight; Fig. 1), with CUE ranging from 0.15 to 0.6 (Fig. 2). While atmospheric, oxic conditions likely prevailed during most of the incubation period, it is probable that some incubations turned anoxic when CO_2 levels exceeded 21%, which occurred in incubations of Ascomycota and Zygomycota growing on glucose, Ascomycota growing on tannic acid, and Zygomycota growing on succinate. The accumulation of CO_2 was necessary consequence of performing the incubations in closed bottles, which was required to prevent the escape of ^{13}C -labeled IC and also to prevent microbial contaminations. Nevertheless, such alteration between oxic and anoxic conditions is common in natural environments, and the measured IC assimilation into fungal lipids was consistently low ($< 3\%$; Fig. 3), regardless the implied anoxia. Furthermore, the observed variability in CUE, %IC, and a_w between growth experiments were not correlated with large increases in headspace CO_2 .

4.2 Fungal IC assimilation into lipids

326 A fundamental process in nature and basis for ecological food webs is the fixation of IC via photosynthesis and/or
 327 chemosynthesis by autotrophic organisms. The IC assimilation by heterotrophic organisms also plays an important role in
 328 ensuring the provision of energy and to replenish intermediates in the TCA cycle that have been released for biosynthesis
 329 (Kornberg 1965). Therefore, IC assimilation is a proxy for both anabolic processes and the catabolic status of the cell,
 330 influenced by assimilation, biosynthesis, anaplerotic reactions, and redox balancing reactions (Braun et al., 2021; Erb 2011).
 331 Previous reports on the by-fixation of IC (%IC) via anaplerotic pathways into heterotrophic biomass varied between 1% and
 332 8% (Dijkhuizen & Harder, 1985; Feisthauer et al., 2008; Romanenko 1964; Roslev et al., 2004), whereas for fungi it was
 333 previously reported to amount to roughly 1% (Sorokin 1961; Schinner & Concina, 1981; Schinner et al., 1982), and was recently
 334 shown to vary between 2% and 12% for Ascomycota when grown on glucose or glutamic acid (Jabinski et al., 2024). Our
 335 results, focusing on a specific fatty acid biomarker, demonstrate a low range in %IC for all different substrates and species
 336 tested in this study (0 - 3%), with the Ascomycota (0 - 2%) assimilating relatively less IC than previously reported species
 337 ($4.6\% \pm 1.6\%$; Jabinski et al., 2024). The highest observed incorporation was $2.2 \pm 0.5\%$ by *Penicillium janczewskii* (PJ, n =
 338 3) when grown for 21 days on tannic acid (Table 1; Fig. 3). Only the other Ascomycota species, *Paecilomyces lilacinus* (PL),
 339 grew sufficiently on tannic acid (up to 10% CO₂ and 20 mg dry weight after 13 days; %IC = $0.14 \pm 0.02\%$, n = 3 ; Figs. 1, 3),
 340 suggesting that increased assimilation of IC by PJ may have occurred during the extra week of incubation and promoted higher
 341 biomass production. The high CO₂ levels also suggest that the incubations of PJ with tannic acid may have turned anoxic,
 342 which may also explain the higher IC incorporation in these incubations. Overall, heterotrophic IC assimilation does not appear
 343 to be a hallmark of any of the variety of fungal taxa or catabolic pathways probed in this study.


345 **4.3 Water hydrogen incorporation into fungal lipid biomarker C_{18:2}**

346 As demonstrated previously, the regression slope between hydrogen isotopic composition of water medium and microbial lipids
 347 (i.e., a_w) varies with the type of metabolism (Zhang et al., 2009; Valentine, 2009; Wijker et al., 2019; Jabinski et al., 2024), and
 348 importantly for heterotrophs, a non-negligible proportion of hydrogen may derive from sources other than ambient water. For
 349 fatty acid biosynthesis, H incorporation is suggested to be a function of transporters and electron acceptors (NADPH and
 350 NADH), with contributions accounting for around half of all lipid hydrogen (Maloney et al., 2024). The remaining comprises
 351 equal contributions of H obtained directly from environmental water or acetyl-CoA (Valentine, 2009; Zhang et al., 2009; Caro
 352 et al., 2023). The consensus from previous studies that investigated the lipids of heterotrophic bacteria is that microbial
 353 heterotrophs exhibit a_w values ranging from 0 to 1, with a mean of 0.71 ± 0.17 (e.g., Caro et al., 2023), though some organisms
 354 have exhibited a_w values exceeding 1 (Dirghangi et al., 2013; Jabinski et al., 2024). Jabinski et al. (2024) demonstrated that
 355 five species of heterotrophic Ascomycota exhibit similar a_w values (0.62 ± 0.04) for the fungal biomarker C_{18:2} during growth
 356 on glucose. Zhang et al. (2009) reported similar a_w values for *E. coli* grown on glucose (0.63 ± 0.03). In the current study, a_w
 357 values for the fungal biomarker C_{18:2} during growth on glucose (0.60 ± 0.05) were agreeable with Jabinski et al. (2024), but
 358 more variable, likely owing to the broader phylogenetic coverage of the current study.

359 The large variability in $\varepsilon_{C18:2/water}$ estimates (Fig. 5; Table S1) can be partly attributed to the large range in δ^2H of medium water,
 360 which contributes H together with the substrate and metabolic water to determine $\delta^2H_{C18:2}$, even though the net isotope effect
 361 ($\alpha_{C18:2/water}$) may be consistent for a specific species-substrate pair (Session and Hayes, 2005; Kopf et al., 2015). In other words,
 362 $\varepsilon_{C18:2/water}$ values do not account for the admixture of H sources that also affect $\delta^2H_{C18:2}$ of heterotrophic fungi. More robust
 363 differences were observed in a_w values both between and within the different phyla and substrates tested.

365 4.3.1 Trends across fungal phyla

366 For incubations in which the Zygomycota isolates produced enough biomass to determine a_w (i.e., glucose, succinate, β -
 367 cyclodextrin; $n = 5$; Fig. 4), we observed a highly significant inverse correlation with CUE ($R^2 = 0.87$, $p < 0.01$; Fig. 6),
 368 suggesting coupling of a_w with growth efficiency. Presumably, the proportion of substrate-derived H incorporated into lipids
 369 increases with more efficient incorporation of substrate C into biomass. Ascomycota exhibited the most consistent a_w values
 370 among phyla when grown on each of the four different substrates [0.63 ± 0.03 (glucose); 0.78 ± 0.01 (succinate); 0.76 ± 0.02
 371 (tannic acid); 0.67 ± 0.01 (β -cyclodextrin)], but also the largest variability in CUE (0.08-0.59; Fig. 2). CUE and a_w exhibited a
 372 similar trend as observed for Zygomycota isolates, but were not significantly correlated across incubations of Ascomycota,
 373 Basidiomycota only produced sufficient biomass when fed substrates that activated the glycolytic pathway (glucose or β -
 374 cyclodextrin; $CUE < 0.3$), yet showed high variability in a_w between species ($0.37 \pm 0.03 < a_w < 0.75 \pm 0.06$; Fig. 4), which
 375 were beyond the more confined range of a_w values determined for isolates belonging to Zygomycota and Ascomycota.



377
 378 **Figure 6. Biplot of CUE and a_w values for fungi growing on glucose (circles), succinate (squares), tannic acid (diamonds),**
 379 **or β -cyclodextrin (triangles). The colors are redundant with Fig. 1, representing the phyla Ascomycota (blue shades, n**
 380 **$= 8$), Zygomycota (green shades, $n = 5$), and Basidiomycota (yellow and orange symbols, $n = 3$). The dashed lines and**
 381 **R^2 values indicate the linear regression for the corresponding phylum across all substrates that yielded sufficient**
 382 **biomass. Only the regression for Zygomycota was significant ($p < 0.01$).**

383

384 *4.3.1 Trends across C substrates*

385 Across all incubations, the similar a_w values determined for growth on glucose (0.60 ± 0.05) versus β -cyclodextrin ($0.58 \pm$
386 0.06), of which the latter consists of seven glucopyranose units ($C_6H_{12}O_6$), suggests that the catabolism of glucose subunits via
387 glycolysis overprints signals of water-H incorporation that may derive during degradation of the β -cyclodextrin oligomer.
388 Alternatively to glycolysis, succinate yielded significantly higher a_w values (0.83 ± 0.05), which was in the same range as
389 reported for *C. oxalaticus* when grown on succinate (a_w 0.80 ± 0.05 ; Zhang et al., 2009), and was more similar to that reported
390 previously for fungal growth on glutamic acid (0.90 ± 0.07 ; Jabinski et al., 2024). Considering all fungal incubations, a one-
391 way analysis of variance (ANOVA; Holm-Sidak method; SigmaPlot v11) confirmed the significant difference in a_w values
392 between growth on glucose and glutamic acid ($p < 0.001$), glutamic acid and β -cyclodextrin ($p < 0.001$), succinate and glucose
393 ($p < 0.003$), and succinate and β -cyclodextrin ($p < 0.005$). It also confirmed that there was no significant difference between
394 the other substrate combinations ($p > 0.05$).

395 Glutamic acid and succinate are thought to be introduced into the TCA cycle through coupled metabolites, where succinate is
396 a direct metabolite of the TCA cycle and glutamic acid is converted to α -ketoglutarate by transamination before entering the
397 TCA cycle, which is only 2 steps from succinate (Cooper et al., 2014). Also, being acids, these substrates may have a greater
398 capacity than saccharides to exchange H with ambient water at experimental pH (typically $2 < \text{pH} < 5.2$), especially glutamic
399 acid, which also comprises an amino moiety. Tannic acid (0.76 ± 0.02) yielded no significant differences ($p > 0.05$) from the
400 other substrates, and is reported to be degraded to different subunits including gallic acid and glucose (Banerjee and Mahapatra,
401 2012; Lekha and Lonsane, 1997 and references within). Aromatic degradation pathways employed by fungi generate
402 intermediates that go through the β -ketoadipate pathway (Mäkelä et al., 2015) before entering the TCA cycle as a succinyl-
403 CoA metabolite (Lekha and Lonsane, 1997). The a_w values induced by degradation of tannic acid suggest that it integrates both
404 the lower a_w signature of glycolysis and higher a_w signature of the TCA cycle (Fig. 7).

405 Together, our incubation experiments suggest that a_w values determined for the fungal biomarker $C_{18:2}$ could not distinguish
406 between fungal growth on relatively labile monomers (i.e., glucose and succinate; requiring as few as 5 days of cultivation)
407 versus larger, less-labile substrates (i.e., β -cyclodextrin and tannic acid; requiring 20 to 183 days of cultivation). However, a_w
408 values of fungal lipid biomarkers may be indicative of fungi employing primarily glycolytic or TCA pathways. Environmental
409 assays that quantify fungal lipid production via the incorporation of ambient water-H (i.e., the lipid-SIP approach) may upscale
410 to total production estimates by applying our calculated mean a_w value of 0.69 ± 0.03 [$n = 27$; \pm (SEM)], which is consistent
411 with the a_w value of 0.71 recommended for soil microbial communities (Caro et al., 2023). For ^2H -SIP investigation of fungal
412 ecotypes supplied with TCA metabolites, such as mycorrhiza, the a_w value of $C_{18:2}$ may range up to 0.83 ± 0.05 . Similar
413 approaches could be applied to environmental samples, such that the a_w values of phospholipids containing $C_{18:2}$ fatty acids
414 could inform the distribution of predominant metabolic ecotypes across a soil profile.

415 Comparisons of net $^{2/1}\text{H}$ fractionation factors (i.e., $\epsilon_{18:2/\text{H}_2\text{O}}$; Table S1) estimated in the current study to previous reports are
 416 most relevant for those derived from our natural abundance incubations. However, we note that these estimates were
 417 determined for a fewer number of replicates ($n = 2$ or 3) and represent the distinct value relative to our laboratory water ($\delta^2\text{H}$
 418 $= -49$ ‰), which varies from labeled treatments (Table S1) and likely natural systems having a significantly different $\delta^2\text{H}_{\text{H}_2\text{O}}$.
 419 Fungal isolates that could successfully grow on tannic acid exhibited the highest $^{2/1}\epsilon_{18:2/\text{H}_2\text{O}}$ values, which is likely manifested
 420 by NADPH production and turnover (e.g. Wijker et al., 2019). The Zygomycota isolates investigated in the current study
 421 (*Mortierella* and *Umbelopsis*) both exhibited relatively strong fractionation of $^{2/1}\text{H}_{\text{water}}$ during growth on succinate ($\epsilon_{18:2/\text{H}_2\text{O}} <$
 422 -123 ‰), which contrasts the trends observed for Ascomycota isolates and previous reports of heterotrophic bacteria (e.g.,
 423 Zhang et al., 2009), where $\epsilon_{18:2/\text{H}_2\text{O}}$ values are typically higher for succinate relative to other C substrates. Together, the relatively
 424 strong $^{2/1}\text{H}$ fractionation (i.e., low $\epsilon_{18:2/\text{H}_2\text{O}}$) and high water isotope incorporation efficiency (a_w) suggest these Zygomycota
 425 isolates employ an alternative catabolic pathway for succinate, resulting in minor incorporation of succinate-derived H into
 426 $\text{C}_{18:2}$.

427 428 **4.4 Dual-SIP approach**

429
 430 Dual-SIP experiments with $^2\text{H}_2\text{O}$ and ^{13}C -dissolved IC previously highlighted the potential to track microbial activity and
 431 distinguish heterotrophic vs autotrophic metabolic modes within environmental settings and pure cultures (Kellerman et al.,
 432 2012, 2016; Wegener et al., 2012; Huguet et al., 2017; Wu et al., 2018, 2020). This approach was also previously applied to
 433 investigate fungal pure cultures (Jabinski et al., 2024), in which the plot of assimilation of IC versus water-H into the fungal
 434 biomarker $\text{C}_{18:2}$ could distinguish five Ascomycota species growing on glucose or glutamic acid, with a_w values explaining
 435 most of the variability. While calculated $\text{IC}:a_w$ are useful to distinguish autotrophic from heterotrophic growth (cf. Wegener et
 436 al., 2016), all calculated values in this study remained near zero, with %IC ranging up to 3% and a_w values ranging from 0.37
 437 to 0.96 (Fig. 4). This pure culture study therefore suggests that fungal assimilation of IC is low and less insightful than the
 438 more distinguishable a_w values for identifying the relative contributions of fungal phylotypes or ecotypes in environmental
 439 assays.




Figure 7. Scatterplot of %IC and a_w values plot for $C_{18:2}$ harvested from incubations of Ascomycota (blue symbols), Basidiomycota (yellow and orange symbols), and Zygomycota (green symbols). The shaded polygons span the range of %IC and a_w values (\pm propagated error) for fungal growth on glucose (circles, green shape), succinate (squares, grey shape), tannic acid (diamonds, pink shape), or β -cyclodextrin (triangles, yellow shape). Each data point represents one species-substrate pair determined for $n > 4$ growth experiments.

5 Conclusion

The purpose of this work was to apply the dual-SIP assay on pure fungal cultures to define the effect of different organic C substrates on the incorporation of water-H and IC into their membrane lipids. Although heterotrophic CO_2 fixation by microbes may range up to 8% of biomass C, the IC assimilation into the fungal biomarker $C_{18:2}$ harvested from six species representing Ascomycota, Basidiomycota, and Zygomycota did not vary consistently between species or substrate, and remained below 3%. Our findings suggest that the fungal catabolic pathways activated by the variety of substrates tested in this study cannot fully account for the higher levels of heterotrophic CO_2 incorporation reported in natural systems. However, *Penicillium janczewskii*, the species that was most successful at respiring tannic acid, also exhibited the highest %IC value of all incubations (Fig. 3; Fig. 7), suggesting that fungal degradation of similarly complex substrates may rely in part on the assimilation of IC (e.g., via anaplerotic reactions). The use of SIP to estimate %IC of heterotrophs required (i) a closed system to prevent loss of ^{13}C label to the atmosphere, and (ii) a high label dose, to contend with the accumulation of CO_2 respired from growth substrate.

458 These conditions intensified upon decreases in pH during the growth experiments, thereby shifting the speciation of IC toward
459 CO₂, which may have further, yet unknown consequences for anaplerotic incorporation of the IC. Future applications to
460 determine %IC for microbial heterotrophs should consider repeated spiking or continuous cultivation practices to better
461 stabilize pH and $\delta^{13}\text{C}_{\text{DIC}}$. Likewise, it would be worth considering isotopic analyses of other potential bioindicator compounds
462 (e.g., sterols, peptides, aminosugars) to see whether incorporation of the IC into such other compounds would render higher
463 levels than fatty acids.

464 In contrast to %IC, we conclude that substrates that activated the glycolysis pathway yielded significantly lower a_w values
465 than those catabolized as TCA intermediates. The relatively high $\varepsilon_{18:2/\text{H}_2\text{O}}$ values harvested from fungi grown on tannic acid
466 may further help to distinguish fungal ecotypes growing on complex, aromatic-containing substrates in natural systems. The
467 expanded a_w dataset reported in this study suggests that the accuracy of fungal production estimated by ²H-lipid SIP
468 experiments can be improved by applying the average a_w value of 0.69 for saprotrophic fungi or up to 0.83 for mycorrhizal
469 fungi. Furthermore, determination of a_w values in environmental ²H-SIP assays may be useful to identify the relative
470 contributions of fungal ecotypes that rely on C substrates fueling glycolysis (e.g., leaf litter) versus those that are fed
471 primarily by TCA intermediates (e.g., root or microbial exudates). Lastly, to our knowledge, the two Zygomycota strains
472 investigated in this study provide the first evidence of a potential correlation between a_w and CUE (Fig. 6), encouraging
473 further exploration of the link between these two parameters, both of which are coupled to microbial central metabolic
474 pathways.

475

476 **Data availability**

477 Data presented in the figures and tables are available in supplementary material (Table S1) and will be made available on the
478 Fractome Database (<https://fractome.caltech.edu/>).

479

480 **Author contribution**

481 SJ, Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Software, Validation,
482 Visualization, Writing – original draft, Writing – review and editing

483 VK, Investigation, Methodology, Resources, Writing – review and editing

484 MK, Formal analysis, Methodology, Resources, Validation,

485 JJ, Conceptualization, Formal analysis, Methodology, Resources, Validation, Writing – review and editing

486 TBM, Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project
487 administration, Resources, Software, Supervision, Validation, Visualization, Writing – original draft, Writing – review and
488 editing

489

490 **Competing interests**

491 The authors declare that they have no conflict of interest.

492 **Disclaimer Acknowledgements**

493 We thank Ljubov Poláková for the support of the stable isotope measurements and laboratory protocols; the Collection of
494 Microscopic Fungi of the Institute of Soil Biology BC CAS for providing the fungal species *Penicillium janczewskii* strain
495 BCCO20_0265 and the Institute of Microbiology CAS for providing the fungal species *Paxillus involutus* strain
496 SB-22; *Phanerochaete chrysosporium* strain CCM8074, *Mortierella* strain RK-38; *Umbelopsis* strain RK-43 and *Paecilomyces*
497 *lilacinus* strain DP-23. Nemiah Ladd and an anonymous reviewer are acknowledged for their time and thoughtful contributions
498 to improve this manuscript. This Project was funded by the Czech Science Foundation GACR nr. 20-223805 (FUNSIF) and
499 supported by MEYS CZ grant LM2015075 Projects of Large Infrastructure for Research, Development and Innovations as
500 well as the European Regional Development Fund-Project: research of key soil-water ecosystem interactions at the SoWa
501 Research Infrastructure (No. CZ.02.1.01/0.0/0.0/16_013/0001782).

502

503

504

505

506

References

- Baldrian, P., Voříšková, J., Dobiášová, P., Merhautová, V., Lisá, L., & Valášková, V. Production of extracellular enzymes and degradation of biopolymers by saprotrophic microfungi from the upper layers of forest soil. *Plant and Soil*, 338(1-2), 111-125. doi:10.1007/s11104-010-0324-3, 2011.
- Banerjee, D., & Mahapatra, S. Fungal tannase: a journey from strain isolation to enzyme applications. *Dyn Biochem Process Biotechnol Mol Biol*, 6(2), 49-60, 2012.
- Boer, W. D., Folman, L. B., Summerbell, R. C., & Boddy, L. Living in a fungal world: impact of fungi on soil bacterial niche development. *FEMS microbiology reviews*, 29(4), 795-811, 2005.
- Boschker HTS, Middelburg JJ. Stable isotopes and biomarkers in microbial ecology. *FEMS Microbiol Ecol* 40:85–95. <https://doi.org/10.1111/j.1574-6941.2002.tb00940.x> 2002.
- Boschker, H. T. S., Nold, S. C., Wellsbury, P., Bos, D., De Graaf, W., Pel, R., ... & Cappenberg, T. E. . Direct linking of microbial populations to specific biogeochemical processes by ¹³C-labelling of biomarkers. *Nature*, 392(6678), 801-805, 1998.
- Braun A, Spona-Friedl M, Avramov M, Elsner M, Baltar F, Reinthaler T, Herndl GJ, Griebler C. Reviews and syntheses: heterotrophic fixation of inorganic carbon—significant but invisible flux in environmental carbon cycling. *Biogeosciences* 18:3689–3700. <https://doi.org/10.5194/bg-18-3689-2021>, 2021.
- Bukovská P, Bonkowski M, Konvalinková T, Beskid O, Hujšlová M, Püschel D. Utilization of organic nitrogen by arbuscular mycorrhizal fungi—is there a specific role for protists and ammonia oxidizers? *Mycorrhiza*. ;28(3):269–83. <https://doi.org/10.1007/s00572-018-0825-0>, 2018.
- Canarini, A., Fuchslueger, L., Schnecker, J., Metze, D., Nelson, D. B., Kahmen, A., ... & Richter, A. Soil fungi remain active and invest in storage compounds during drought independent of future climate conditions. *Nature Communications*, 15(1), 10410, 2024.
- Carlson C.A., N.R. Bates, D.A. Hansell, D.K. Steinberg. Carbon Cycle. In: J. Steele, S. Thorpe, K. Turekian (Eds.) *Encyclopedia of Ocean Science*, 2nd Edition. Academic Press, 477-486, 2001.
- Caro, T. A., McFarlin, J., Jech, S., Fierer, N., & Kopf, S. Hydrogen stable isotope probing of lipids demonstrates slow rates of microbial growth in soil. *Proceedings of the National Academy of Sciences*, 120(16), e2211625120, 2023.
- Ciais, P., Dolman, A. J., Bombelli, A., Duren, R., Peregon, A., Rayner, P. J., . . . Marland, G. Current systematic carbon-cycle observations and the need for implementing a policy-relevant carbon observing system. *Biogeosciences*, 11(13), 3547-3602. doi:10.5194/bg-11-3547-2014, 2014.
- Cooper AJL, Kuhara T. α -Ketoglutaramate: an overlooked metabolite of glutamine and a biomarker for hepatic encephalopathy and inborn errors of the urea cycle. *Metab Brain Dis* 29:991–1006. <https://doi.org/10.1007/s11011-013-9444-9>, 2014.

539 Dijkhuizen, L., & Harder, W. Microbial metabolism of carbon dioxide. *Comprehensive biotechnology: the principles,*
 540 *applications, and regulations of biotechnology in industry, agriculture, and medicine/editor-in-chief, Murray Moo-Young,*
 541 *1985.*
 542 Dumont, M. G., & Murrell, J. C. Stable isotope probing—linking microbial identity to function. *Nature Reviews Microbiology,*
 543 *3(6), 499-504, 2005.*
 544 Erb TJ. Carboxylases in natural and synthetic microbial pathways. *Appl Environ Microbiol* 77:8466–8477.
 545 <https://doi.org/10.1128/AEM.05702-11>, 2011.
 546 Feisthauer, S., Wick, L. Y., Kästner, M., Kaschabek, S. R., Schlömann, M., & Richnow, H. H. Differences of heterotrophic
 547 ¹³CO₂ assimilation by *Pseudomonas knackmussii* strain B13 and *Rhodococcus opacus* 1CP and potential impact on biomarker
 548 stable isotope probing. *Environmental Microbiology*, 10(6), 1641-1651, 2008.
 549 Fioretto, A., Di Nardo, C., Papa, S., & Fuggi, A. Lignin and cellulose degradation and nitrogen dynamics during decomposition
 550 of three leaf litter species in a Mediterranean ecosystem. *Soil Biology and Biochemistry*, 37(6), 1083-1091.
 551 doi:10.1016/j.soilbio.2004.11.007, 2005.
 552 Fischer, C. R., Bowen, B. P., Pan, C., Northen, T. R., & Banfield, J. F. Stable-isotope probing reveals that hydrogen isotope
 553 fractionation in proteins and lipids in a microbial community are different and species-specific. *ACS chemical biology*, 8(8),
 554 1755-1763, 2013.
 555 Frey, S. D. Mycorrhizal fungi as mediators of soil organic matter dynamics. *Annual review of ecology, evolution, and*
 556 *systematics*, 50, 237-259. doi:10.1146/annurev-ecolsys-110617-062331, 2019.
 557 Grinhut, T., Hadar, Y., & Chen, Y. Degradation and transformation of humic substances by saprotrophic fungi: processes and
 558 mechanisms. *Fungal biology reviews*, 21(4), 179-189, 2007.
 559 Hayes, J. M. An introduction to isotopic calculations. Woods Hole Oceanographic Institution, Woods Hole, MA, 2543, 2004.
 560 Hoefs, J. Stable isotope geochemistry. Springer International Publishing AG, part of Springer Nature. doi:10.1007/9783-319-
 561 78527-1, 2018.
 562 Hoëgberg, P., Nordgren, A., Buchmann, N., Taylor, A. F., Ekblad, A., Hoëgberg, M. N., . . . Read, D. J. Large-scale forest
 563 girdling shows that current photosynthesis drives soil respiration. *Nature*, 411(6839), 789-792, 2001.
 564 Huguet, A., Meador, T. B., Laggoun-Défarge, F., Könnike, M., Wu, W., Derenne, S., & Hinrichs, K.-U. Production rates of
 565 bacterial tetraether lipids and fatty acids in peatland under varying oxygen concentrations. *Geochimica et*
 566 *Cosmochimica Acta*, 203, 103-116. doi:10.1016/j.gca.2017.01.012, 2017.
 567 Jabinski, S., d. M. Rangel, W., Kopáček, M., Jílková, V., Jansa, J., & Meador, T. B. Constraining activity and growth substrate
 568 of fungal decomposers via assimilation patterns of inorganic carbon and water into lipid biomarkers. *Applied and*
 569 *Environmental Microbiology*, 90(4), e02065-23, 2024.
 570 Kellermann, M. Y., Wegener, G., Elvert, M., Yoshinaga, M. Y., Lin, Y.-S., Holler, T., . . . Hinrichs, K.-U. Autotrophy as a
 571 predominant mode of carbon fixation in anaerobic methane-oxidizing microbial communities. *Proceedings of the National*
 572 *Academy of Sciences*, 109(47), 19321-19326. doi:10.1073/pnas.1208795109, 2012.

573 Kellermann, M. Y., Yoshinaga, M. Y., Wegener, G., Krukenberg, V., & Hinrichs, K.-U. Tracing the production and fate of
 574 individual archaeal intact polar lipids using stable isotope probing. *Organic Geochemistry*, 95, 13-20.
 575 doi:10.1016/j.orggeochem.2016.02.004, 2016. Kirk, T. K., & Farrell, R. L. Enzymatic "combustion": the microbial degradation
 576 of lignin. *Annual Reviews in Microbiology*, 41(1), 465-501. doi:10.1146/annurev.mi.41.100187.002341, 1987.
 577 Kopf, S. H., Sessions, A. L., Cowley, E. S., Reyes, C., Van Sambeek, L., Hu, Y., ... & Newman, D. K. (2016). Trace
 578 incorporation of heavy water reveals slow and heterogeneous pathogen growth rates in cystic fibrosis sputum. *Proceedings of*
 579 *the National Academy of Sciences*, 113(2), E110-E116.
 580 Kopf, S. H., McGlynn, S. E., Green-Saxena, A., Guan, Y., Newman, D. K., & Orphan, V. J. Heavy water and (15) N labelling
 581 with NanoSIMS analysis reveals growth rate-dependent metabolic heterogeneity in chemostats. *Environ Microbiol*, 17(7),
 582 2542-2556. doi:10.1111/1462-2920.12752, 2015.
 583 Kornberg HL. Anaplerotic sequences in microbial metabolism. *Angew Chem Int Ed Engl* 4:558-565.
 584 https://doi.org/10.1002/anie.196505581, 1965.
 585 Kreuzer-Martin, H. W. Stable isotope probing: linking functional activity to specific members of microbial communities. *Soil*
 586 *Science Society of America Journal*, 71(2), 611-619, 2007..
 587 Lekha, P. K., & Lonsane, B. K. Production and application of tannin acyl hydrolase: state of the art. *Advances in applied*
 588 *microbiology*, 44, 215-260, 1997.
 589 Lindahl, B. D., & Tunlid, A. Ectomycorrhizal fungi—potential organic matter decomposers, yet not saprotrophs. *New*
 590 *Phytologist*, 205(4), 1443-1447. doi:10.1111/nph.13201, 2015.
 591 Maloney, A. E., Kopf, S. H., Zhang, Z., McFarlin, J., Nelson, D. B., Masterson, A. L., & Zhang, X. Large enrichments in fatty
 592 acid 2H/1H ratios distinguish respiration from aerobic fermentation in yeast *Saccharomyces cerevisiae*. *Proceedings of the*
 593 *National Academy of Sciences*, 121(20), e2310771121, 2024.
 594 Osburn, M. R., Sessions, A. L., Pepe-Ranne, C., & Spear, J. R. Hydrogen-isotopic variability in fatty acids from Yellowstone
 595 National Park hot spring microbial communities. *Geochimica et Cosmochimica Acta*, 75(17), 4830-4845, 2011.
 596 Pilecky, M., Meador, T. B., Wassenaar, L.I. Advancements in Compound-Specific Hydrogen Stable-Isotope Analysis of Fatty
 597 and Amino Acids. *TrAC Trends in Analytical Chemistry*, 118194. doi: 10.1016/j.trac.2025.118194, 2025.
 598 Romanenko VI. Heterotrophic assimilation of CO₂ by bacterial flora of water. *Mikrobiologiya* 33:679-683, 1964.
 599 Roslev P, Larsen MB, Jørgensen D, Hesselsoe M. Use of heterotrophic CO₂ assimilation as a measure of metabolic activity in
 600 planktonic and sessile bacteria. *J Microbiol Methods* 59:381-393. https://doi.org/10.1016/j.mimet.2004.08.002, 2004.
 601 Sachse, D., Billault, I., Bowen, G. J., Chikaraishi, Y., Dawson, T. E., Feakins, S. J., ... & Kahmen, A. Molecular paleohydrology:
 602 interpreting the hydrogen-isotopic composition of lipid biomarkers from photosynthesizing organisms. *Annual Review of Earth*
 603 *and Planetary Sciences*, 40, 221-249, 2012.
 604 Schinner F, Concini R, Binder H. Heterotrophic CO₂ fixation by fungi in dependence on the concentration of the carbon source.
 605 Paper presented at the Phytos; *Annales rei botanicae*, 1982.
 606

607 Schinner F, Concinn R. Carbon dioxide fixation by wood-rotting fungi. *Eur J For Pathol* 11:120–123.
 608 <https://doi.org/10.1111/j.1439-0329.1981.tb00077.x>, 1981.

609 Sessions, A.L., Hayes, J.M., 2005. Calculation of hydrogen isotopic fractionations in biogeochemical systems. *Geochim.*
 610 *Cosmochim. Acta* 69, 593–597.

611 Smith, S. E., & Read, D. Mineral nutrition, toxic element accumulation and water relations of arbuscular mycorrhizal plants.
 612 *Mycorrhizal symbiosis*, 3, 145-148. doi:10.1016/B978-012370526-6.50007-6, 2008.

613 Šnajdr, J., Cajthaml, T., Valášková, V., Merhautová, V., Petránková, M., Spetz, P., . . . Baldrian, P. Transformation of *Quercus*
 614 *petraea* litter: successive changes in litter chemistry are reflected in differential enzyme activity and changes in the microbial
 615 community composition. *FEMS Microbiology Ecology*, 75(2), 291-303. doi:10.1111/j.1574-6941.2010.00999.x, 2011.

616 Sorokin, J. I. On the carbon dioxide uptake during the cell synthesis by microorganisms. *Zeitschrift für allgemeine*
 617 *Mikrobiologie*, 6(1), 69-73. doi:10.1002/jobm.3630060107, 1966.

618 Sorokin, Y. I. Heterotrophic carbon dioxide assimilation by microorganisms. *Zhurnal Obsheei Biologii*, 22(4), 265272, 1961.

619 Treonis, A. M., Ostle, N. J., Stott, A. W., Primrose, R., Grayston, S. J., & Ineson, P. Identification of groups of metabolically-
 620 active rhizosphere microorganisms by stable isotope probing of PLFAs. *Soil Biology and Biochemistry*, 36(3), 533-537, 2004.

621 Valentine, D. L. Isotopic remembrance of metabolism past. *Proceedings of the National Academy of Sciences*, 106(31), 12565-
 622 12566. doi:10.1073/pnas.0906428106, 2009.

623 Warren, C. R. D₂O labelling reveals synthesis of small, water-soluble metabolites in soil. *Soil Biology and Biochemistry*, 165,
 624 108543, 2022.

625 Wegener, G., Bausch, M., Holler, T., Thang, N. M., Prieto Mollar, X., Kellermann, M. Y., . . . Boetius, A. Assessing sub-seafloor
 626 microbial activity by combined stable isotope probing with deuterated water and ¹³C-bicarbonate. *Environmental*
 627 *Microbiology*, 14(6), 1517-1527. doi:10.1111/j.1462-2920.2012.02739.x, 2012.

628 Wegener, G., Kellermann, M. Y., & Elvert, M. Tracking activity and function of microorganisms by stable isotope probing of
 629 membrane lipids. *Current Opinion in Biotechnology*, 41, 43-52. doi:10.1016/j.copbio.2016.04.022, 2016.

630 Wijker, R. S., Sessions, A. L., Fuhrer, T., & Phan, M. 2H/1H variation in microbial lipids is controlled by NADPH metabolism.
 631 *Proceedings of the National Academy of Sciences*, 116(25), 12173-12182. doi:10.1073/pnas.1818372116, 2019.

632 Willers, C., Jansen van Rensburg, P. J., & Claassens, S. Phospholipid fatty acid profiling of microbial communities—a review
 633 of interpretations and recent applications. *Journal of applied microbiology*, 119(5), 1207-1218, 2015.

634 Wu W, Meador T, Hinrichs K-U. Production and turnover of microbial organic matter in surface intertidal sediments.
 635 *Org Geochem* 121:104–113. <https://doi.org/10.1016/j.orggeochem.2018.04.006>, 2018.

636 Wu, W., Meador, T. B., Könneke, M., Elvert, M., Wegener, G., & Hinrichs, K. U. Substrate-dependent incorporation of carbon
 637 and hydrogen for lipid biosynthesis by *Methanosarcina barkeri*. *Environmental Microbiology Reports*, 12(5), 555567.
 638 doi:10.1111/1758-2229.12876, 2020.

639 Zhang, X., Gillespie, A. L., & Sessions, A. L. Large D/H variations in bacterial lipids reflect central metabolic pathways.
 640 *Proceedings of the National Academy of Sciences*, 106(31), 12580-12586. doi:10.1073/pnas.0903030106, 2009.