

Adjustments to an abrupt solar forcing in the CMIP6 abrupt-solm4p experiment

Charlotte Lange ¹ and Johannes Quaas ¹

¹Leipzig Institute for Meteorology, Leipzig University, Stephanstraße 3, 04103 Leipzig, Germany

Correspondence: Charlotte Lange (charlotte.lange@uni-leipzig.de)

Abstract. "Radiative" or "rapid" adjustments refer to the response of the climate system to an instantaneous radiative forcing, independent of surface temperature changes. These adjustments occur over time scales from hours (e.g., aerosol-cloud interactions) to months (e.g., stratospheric temperature changes), often overlapping with feedback mechanisms—making their separation in realistic scenarios challenging. Controlled simulation experiments can thus help isolate adjustment processes and improve understanding relevant to more complex cases like volcanic eruptions or ongoing climate change.

The abrupt-solm4p experiment of the Cloud Feedback Model Intercomparison Project (CFMIP), part of the 6th Coupled Model Intercomparison Project (CMIP6), simulates an instantaneous 4% reduction in the solar constant from a pre-industrial state on 1 January 1850. This study analyses changes in climate variables, cloud properties, and radiative fluxes across different time scales (hours, days, months and up to 150 years) to identify adjustment mechanisms and characteristic fingerprints to a shortwave forcing scenario.

Four models show rapid adjustments of $3.57 \pm 0.46 \text{ W m}^{-2}$, offsetting $\sim 35\%$ of the initial forcing. Distinct local patterns include initial rapid surface cooling—especially over Antarctica and the Southern Hemisphere. Stratospheric cooling slows the polar night jet, disrupts the polar vortex, and increases Arctic cloud cover, causing local warming. Within the first month, the troposphere cools more quickly than the ocean surface, decreasing vertical stability and enhancing cloud cover over oceans, while tropical land regions show the opposite change. On monthly time scales, high clouds shift downward due to reduced convective heating.

1 Introduction

The Earth's climate is governed by radiative fluxes entering and leaving the atmosphere. In a steady-state system incoming and outgoing fluxes are balanced, maintaining an equilibrium of the Earth climate system. In case of any perturbation to this balance, known as a radiative forcing, the climate system reacts by heating or cooling, which leads to a new balance of energy fluxes on time scales of centuries. Different processes in the climate's response, called feedbacks or adjustments, either enhance or dampen the Earth's capability to reach a new equilibrium. Feedbacks act in response to global mean surface temperature change and typically take place over years and centuries. In contrast, adjustments happen in direct response to a forcing agent, are independent of surface temperature change and mostly evolve on shorter time scales of days, weeks and months (Andrews and Forster, 2008; Gregory and Webb, 2008; Sherwood et al., 2015). Radiative adjustments of the atmosphere to an external forcing are of particular interest to the scientific community and a growing number of studies have been conducted on the subject (Gregory et al., 2004; Zelinka et al., 2013; Myhre et al., 2013; Sherwood et al., 2015; Smith et al., 2018; Forster et al., 2021b; Quaas et al., 2024). By introducing radiative adjustments into the forcing-feedback-framework as part of the effective radiative forcing (ERF, Myhre et al., 2013), the resulting ERF is a better predictor of implied global surface temperature change, since it is more independent of the kind of forcing agent (Andrews and Forster, 2008; Gregory and Webb, 2008). However, this approach requires a precise estimate of ERF, which comprises the instantaneous radiative forcing (IRF) as well as (radiative or rapid) adjustments (RA), which are prone to high uncertainty, especially in case of cloud adjustments (Andrews and Forster, 2008).

The total global mean radiative imbalance at the top of atmosphere (TOA) $\bar{N}(t)$ can be written as

$$\bar{N}(t) = IRF + RA + \sum_i \lambda_i \Delta \bar{T}(t) \quad (1)$$

where the sum over the feedback parameters λ_i accounts for the different sources of feedbacks.

Although RAs are formally independent of global mean surface temperature change, local temperature changes and consequential atmospheric adjustments can contribute a considerable amount to the overall RA (Quaas et al., 2024). Furthermore, despite many adjustments happening on time scales from hours to days (e.g. precipitation and changes in some cloud properties), adjustments of the stratosphere, cryosphere or vegetation can take months to years (Forster et al., 2021b; Stjern et al., 2023). This leads to an overlap of adjustment and feedback time scales, which makes it difficult to disentangle both. Moreover, adjustments are often confounded by climate variability in magnitude, making it very hard to detect them in observations. This is further complicated by the transient nature of most forcing processes happening in the Earth climate system, like the gradual increase of CO₂ from anthropogenic sources in the atmosphere.

To circumvent these issues, climate models allow the application of instantaneous forcings that exceed climate variability and are kept constant, facilitating a better analysis of adjustments and feedbacks. A common experiment using global climate models is the instantaneous two- or fourfold increase of the atmospheric CO₂ concentration (e.g., Gregory et al., 2004; Colman and McAvaney, 2011; Kamae and Watanabe, 2012; Zelinka et al., 2013; Andrews et al., 2015) in order to predict long-term

consequences of anthropogenic climate change as well as short-term adjustments. However, these simulations are often highly idealised, using fixed sea-surface temperatures or omitting the existence of continents (aqua-planet-simulations). In order to compare and validate such results with observations a natural forcing is required that is applied nearly instantaneously and is of sufficient strength. Volcanic eruptions are considered such so-called "natural laboratories" (e.g., Malavelle et al., 2017; Christensen et al., 2022). Large volcanic eruptions emit huge amounts of aerosols into the atmosphere in a very short time frame of hours. If transported up to the stratosphere, depending on the location and season of the volcanic eruption, the aerosol can form a global scattering layer with a lifetime of up to one to three years (Myhre et al., 2013). This way, volcanic eruptions exert a radiative forcing that is approximately comparable to the instantaneous forcing that usually only models can realise. However, examining volcanic eruptions includes other challenges. The initial forcing is very localised and only after a few months the aforementioned scattering layer is formed around the globe. Depending on the location of the volcano, the strength of the manifold adjustment mechanisms and the distribution of the stratospheric aerosol layer varies. Hence, time scales of different adjustments might still overlap. Therefore, before analysing adjustments to a realistic volcanic eruption, it can be helpful to further simplify the problem. A reduced solar constant forcing can serve as a simplified analogue to a stratospheric volcanic aerosol layer, as both reduce the amount of shortwave radiation reaching the troposphere.

In this study, we thus examine the results of the abrupt-solm4p experiment, which is part of the Cloud Feedback Model Intercomparison Project (CFMIP) as part of the 6th Climate Model Intercomparison Project (CMIP6) (Eyring et al., 2016; Webb et al., 2017). The output of four global climate models, which participated in this experiment, is available. In the simulations the solar constant was reduced by 4 % and kept constant at this value. This can be regarded as a simplified analogue for an aerosol scattering layer. Volcanic eruptions and a reduced solar constant both lead to a reduced incoming shortwave flux at the surface, even if stratospheric adjustments are expected to differ. Hence, similar adjustment patterns are expected for both types of forcing, when the scattering sulfate layer distributes over the whole globe with time.

There have been numerous studies that analysed the reaction of the Earth climate system to a number of different forcing agents so far (e.g., Gregory et al., 2004; Smith et al., 2018), However only few studies quantified radiative adjustments after solar forcing and took a closer look at the specific processes happening in response to solar forcing (e.g., Salvi et al., 2021; Virgin and Fletcher, 2022; Aeronson et al., 2024). Radiative adjustments to solar forcing in sum typically counteract the initial forcing (e.g., Smith et al., 2018; Russotto and Ackerman, 2018; Virgin and Fletcher, 2022; Aeronson et al., 2024). Most of this is attributed to decreased temperature of the troposphere, stratosphere and land surface, which leads to a reduction in longwave radiation lost to space. The effect is partially offset by the reduction of water vapour in the atmosphere, following the Clausius-Clapeyron relationship, and thereby an increase in outgoing longwave radiation. Moreover, Smith et al. (2018) derive a positive adjustment due to change in surface albedo. Furthermore, differential heating of sea and land surface can lead to changes in circulation patterns, which affects precipitation and cloud properties. On longer time scales also a change in vegetation might happen, but is not subject of this study and no further analysis was performed here.

However, when analysing rapid adjustments due to changes in cloud properties, the studies find contradictory results. This is in accordance with an overall high uncertainty linked to cloud property changes in climate models. Reducing short-term uncertainty of cloud adjustments could also reduce uncertainty of long-term predictions (Andrews and Forster, 2008; Nam

et al., 2018). In case of solar forcing Smith et al. (2018) and Virgin and Fletcher (2022) find cloud adjustments that counteract the initial solar forcing. In response to a +2% solar constant forcing, Smith et al. (2018) describe an increase in cloud fraction in the boundary layer, a reduction of cloud fraction in the free troposphere, a small increase in cloud fraction at around 100 hPa and a decrease above this level. This is very similar to our findings, although of the opposite sign, since we examined the response to a reduction of solar constant. Using radiative kernels, Smith et al. (2018) show that cloud changes lead to negative long- and shortwave effects, but with high model disagreement in the shortwave effects. In response to a +2% solar forcing, Virgin and Fletcher (2022) find that the shortwave component is dominated by changes in the boundary layer clouds, while changes in the troposphere contribute more to the longwave effects. In contrast to this, Aeronson et al. (2024) find cloud adjustments that further amplify the initial solar forcing. In response to a +4% solar constant forcing, they find an overall positive cloud radiative adjustment, which is strongest, where low-cloud fraction is reduced. The two main mechanisms are changes in cloud fraction and cloud optical depth, which they find are of opposite sign in case of solar forcing. Moreover, short- and longwave effects tend to partially compensate each other, but shortwave effects are slightly stronger. Hence, in total, positive shortwave effects due to decreased cloud fraction dominate the overall cloud adjustments. However, they find that the participating models do not agree on the signs of cloud adjustment effects and the uncertainty of the total effect is high.

Even though the studies describe similar changes in cloud properties, the derived cloud radiative adjustments differ depending on the applied method and models used. This demonstrates how radiative effects of cloud property changes are still one of the major sources of uncertainty in climate models, even in highly idealised solar forcing scenarios. Hence, this study is particularly interested in adjustments of cloud properties and characteristic patterns in response to shortwave forcing.

CFMIP provides outputs from +4% solar constant (solp4p) as well as -4% solar constant (solm4p). Using both experiments would increase the amount of data statistics could be based on. However, as e.g. Aeronson et al. (2024) found, response to positive and negative solar forcing does not only differ in sign, but it also shows different patterns due to non-linearities and dependence on the base state. Hence, for this study, we decided to only look at rapid adjustments to a reduction in solar constant (solm4p), which in a later step could be compared to adjustments to volcanic forcing.

Reduced solar constant simulations are also of interest for the increasing number of solar radiation modification (SRM) studies that often aim to balance the forcing due to anthropogenic emissions of CO₂ by reducing the absorbed solar radiation (e.g., Bala et al., 2008; Cao et al., 2015; Schmidt et al., 2012; Huneus et al., 2014; Russotto and Ackerman, 2018). Having a better understanding of not only the long-term effects of SRM methods, but also knowing about short-term adjustments, can improve the risk assessment of these endeavours. In these scenarios, not only the global mean evolution of climate variables, but also, if not even more, local processes are of high importance. Especially short-term adjustments of clouds and precipitation can lead to local droughts or floods, with possibly important effects for local communities.

There are several methods to quantify radiative adjustments in climate models. On the one hand, the linear regression method (Gregory et al., 2004), based on Eq. 1, allows for a simultaneous determination of ERF, equilibrium climate sensitivity (long-term trend) and climate feedback parameter ($\sum \lambda_i$). In order to derive radiative adjustments from ERF, knowledge of the instantaneous forcing is necessary. For most forcing agents, it is possible to diagnose the IRF by separate radiative transfer modelling. However, the regression method is only feasible for the global mean and it relies on the assumption that

all rapid adjustments happen while the global mean surface temperature change is still zero. This is an oversimplification, as
120 inert systems like cryosphere or vegetation might still adjust to the initial forcing, while global mean surface temperature has
already begun to change.

On the other hand, the fixed surface temperature method (Hansen et al., 2005) or rather fixed-sea-surface temperature (fixed-
SST) is widely used and has the advantage of suppressing feedbacks (Forster et al., 2016). This allows for the disentanglement
of adjustments and feedbacks. However, this method artificially suppresses adjustments of ocean surface temperature and
125 introduces unrealistic land sea contrast, which hinders a realistic estimate of circulation adjustments. Andrews et al. (2021)
showed a significant difference in ERF depending on whether only sea surface temperature or all surface temperature was kept
to zero. Possible adjustments to localized warming or cooling, cannot be simulated in this kind of setup, although in some
concepts they are considered adjustments relevant to TOA ERF (Quaas et al., 2024).

The simulations analysed in this study apply an instantaneous solar forcing that is kept constant, while allowing the whole
130 climate system to adapt in a fully coupled general circulation model. The regression method was used to quantify ERF and RA,
but non-linear behaviour was found for the first 4-10 years, when the more inert systems like ocean, cryosphere or vegetation
still adjust to the forcing, while global mean surface temperature change starts to dampen TOA effective flux imbalance.
Nevertheless, this model setup allows us to examine adjustment processes on time scales of hours, days and months, where, as
we show, TOA effect flux change is still dominated by adjustment processes rather than temperature mediated change.

135 Section 2 contains a more detailed description of the experiment setup as well as an overview of the available data. Section
3 first shows the results of the classical regression method by Gregory et al. (2004), in order to quantify ERF and RA. It then
discusses how time scales of hours, days and months are dominated by adjustment processes, and examines TOA radiative
fluxes and cloud radiative effect anomaly. Afterwards, it proceeds with a closer analysis of adjustment processes of a number
of different climate variables. These include surface and atmospheric temperature, relative humidity, vertical velocity and cloud
140 fraction. Section 4 discusses the results of this study in the context of studies that have been conducted on adjustments to solar
forcing in recent years and discusses the limitations of the dataset and the approach. Section 5 contains a summary and provides
an outlook on how this study's findings can contribute to future research in the field of radiative adjustments.

2 Methods

In the solm4p-experiment, the solar constant is reduced instantaneously by 4 % and kept constant at 96 % of the solar constant
145 of the control run (pi-Control). It branches off the pi-Control simulation on 1 January 1850. No other forcing is considered in
the experiment. The diagnostics of four models, which participated in this experiment, are available: IPSL-CM6A-LR, CESM2,
CanESM5 and MRI-ESM2-0 (further information provided in Table 1). For each model one experiment run corresponding to
one pre-industrial control run (piControl) is provided. While some models provide 3-hourly output for several parameters,
CESM2 only provides daily data. In case of vertically resolved atmospheric cloud fraction, only monthly data was available
150 from all four models. More information on the four models, including their horizontal and vertical grid spacing is provided in
Table 1.

For all variables examined in this study, the difference attributable to the reduction in solar constant was calculated for each parameter by subtracting the piControl-run from the solm4p-run for each point in time and space.

Following Stjern et al. (2023), four different time scales were considered: the first 100 hours, the first 30 days, the first year and the following years until 150 years after the onset of forcing. For the first time scale (up to 100 hours) 3-hourly data was used, when available, else daily data was used. For the second time scale (days 5–30) daily averages were used, for the third time scale (months 2–12) monthly data was used and for the long-term development (years 2–150) yearly means were plotted. The transitions between the plots of different time scales often display a small jump, because the next frequency does not contain the last data point of the preceding frequency, but an average.

For all time scales the global mean was calculated for each model. Moreover, a multi-model mean was calculated by first interpolating all models to the time axis of the MRI-ESM2-0 model. This model was chosen as reference, because it provided the most extensive database for the abrupt-solm4p experiment. If not all models provided data of the same frequency, the highest available resolution was plotted for each model, respectively and for the multi-model mean, the other models were interpolated to the 3-hourly time axis of MRI-ESM2-0.

In addition to the temporal development of global means, a geographical distribution of the respective parameter was plotted, averaging over the respective time scales. For the mean of the first three time scales, all time steps were averaged, while for the fourth time scale only the years 120-150 after the onset of forcing were averaged to obtain an estimate for the long-term new approximate equilibrium state. For the global distribution, a multi-model mean was calculated by interpolating the other models to the CanESM5 horizontal grid, which was the coarsest out of the four models.

Besides the grid spacing, also the height levels varied between the models in case of the atmospheric cloud fraction data. For the other climate variables, all models used the same 19 height levels. In case of cloud fraction, the other models were interpolated to the CanESM5 pressure axis.

In order to account for uncertainty, the multi-model standard deviation was plotted together with the temporal development of multi-model global mean. In case of global distributions of anomalies, areas were dotted in which less than three of four models agreed on the sign of anomaly.

To analyse differences in land and sea surface response, some climate variables, e.g. relative humidity, were averaged only over ocean and only over land. This was done by applying a land-sea-mask, based on the CanESM5 grid, before performing the zonal averaging.

This study focuses on cloud adjustments and the cloud radiative effect, which was calculated from the difference between simulation runs with clouds (all-sky) and a second radiative transfer calculation for each step without clouds (clear-sky), both provided by each of the four models. Then, the total cloud radiative effect anomaly at TOA was calculated as the combined effect of solar (shortwave) and terrestrial (longwave) radiation changes due to cloud changes.

The linear regression method introduced by Gregory et al. (2004) was applied for a number of TOA fluxes, in order to estimate their influence on the overall ERF. For that purpose, the yearly global mean of the respective TOA flux change was plotted over the yearly global mean of near-surface temperature change and a linear regression was used, which determined the intercept with the TOA flux axis.

	CanESM5	CESM2	IPSL-CM6A-LR	MRI-ESM2-0
Full name	Canadian Earth System Model version 5	Community Earth System Model 2	Institut Pierre-Simon Laplace - Climate Model 6A -Low resolution	Meteorological Research Institute Earth System Model Version 2.0
Reference	Cole et al. (2019)	Danabasoglu (2020)	Boucher et al. (2018)	Yukimoto et al. (2020)
Grid size (lon x lat)	128 x 64	288 x 192	144 x 143	320 x 160
Pressure levels (Pa)	8 (daily) 19 (monthly)	8 (daily) 19 (monthly)	8 (daily) 19 (monthly)	8 (daily) 19 (monthly)
Cloud height levels	49 (atmosphere hybrid sigma pressure coordinate (unitless))	32 (atmosphere hybrid sigma pressure coordinate (unitless))	79 (pressure levels (Pa))	80 (atmosphere hybrid sigma pressure coordinate (unitless))

Table 1. Information on the four models for which output was available from the CFMIP abrupt-solm4p-experiment.

3 Results

In order to estimate the rapid adjustments to a 4% solar constant reduction, we applied the regression method developed by Gregory et al. (2004) to the solm4p-CFMIP data. The results are shown in the following section. We then move on to examine different time scales after the onset of forcing and motivate, how the first three time scales (hours, days and months) are dominated by rapid adjustments, while the fourth time scale shows the long-term adaptation and is therefore dominated by surface temperature mediated processes. Afterwards, we analyse the response of different climate variables like temperature, humidity and cloud fraction on the different time scales and identify characteristic adjustment patterns of these variables.

3.1 Effective radiative forcing and rapid adjustment estimate

We applied a linear regression to the yearly global means of TOA radiative budget change (Δ TOA budget or N following Eq. 1) and near surface temperature change (ΔT) of the four available models. The results are shown in Fig. 1.

The TOA radiative budget anomaly shows the expected behaviour for all four models, starting with negative values for $\Delta T = 0$ and then approaching a new radiative balance. The multi-model mean IRF was estimated as the difference between downward and upward shortwave anomaly for the first month after onset of forcing (shortest output frequency available for all models). The intercept of the linear regressions with the y-axis provides the ERF and the difference between ERF and IRF yields the RA. Multi-model mean ERF and RA are given at the bottom of the figure and are also provided in Table 2. The negative IRF of approximately -10 W m^{-2} is reduced by radiative adjustments of 3.6 W m^{-2} , resulting in an ERF of 6.4 W m^{-2} . While some models, like CanESM5 and IPSL-CM6A-LR, simulate a strong temperature decrease in response to the radiative forcing, CESM2 and MRI-ESM2-0 reach a new equilibrium at a significantly weaker change in surface temperature.




Figure 1. Linear regression plots for yearly global mean TOA radiative budget anomaly in W m^{-2} vs. change in yearly global mean surface temperature (ΔT). Increasing saturation of the colours indicates temporal development.

205 The development is mostly linear. However, especially for the first 10 years all models exhibit a slightly steeper slope than in the long term. This deviation could either be interpreted as radiative adjustments developing on time scales as long as a decade, similar to the ocean-atmosphere adjustments found by Rugenstein et al. (2016), or it might suggest a change in the relation between ΔT and N , with the deep ocean heat uptake dominating the SST-pattern after a decade of forcing, changing the overall ocean heat uptake efficacy (Andrews et al., 2015). Other studies like Gregory and Webb (2008) found similar
 210 non-linearities and attributed those to shortwave cloud radiative effects. In this study the non-linearity was predominant in the clear-sky shortwave component over ocean, but no further analysis was conducted, as this study concentrates on the adjustment processes during the first year after onset of forcing.

Analogous to the total TOA radiative budget, linear regressions were also performed for the single components of the TOA budget, i.e. downward shortwave flux change $\Delta \text{sw}^\downarrow$, upward shortwave flux change $\Delta \text{sw}^\uparrow$ and upward longwave flux $\Delta \text{lw}^\uparrow$
 215 (Fig. A1 in the Supplements). Approximating the instantaneous response of the respective radiative components as the offsets after one month of simulation, the contributions of the single components to the overall IRF and ERF can be estimated and are shown in Table 2. All models simulate an instantaneous response of $\Delta \text{sw}^\uparrow$ of around -4 W m^{-2} in reaction to the reduction of solar constant, depending on the planetary albedo. This is considered to be a part of the IRF, since it is an immediate response to the initial solar constant reduction, not requiring any atmospheric changes. However, the $\Delta \text{sw}^\uparrow$ ERF component
 220 is -6.3 W m^{-2} , indicating shortwave radiative adjustments of $-6.3 \text{ W m}^{-2} - (-4) \text{ W m}^{-2} = 2.3 \text{ W m}^{-2}$. In contrast to that, no longwave contribution to the IRF is expected and models show no initial response of $\Delta \text{lw}^\uparrow$. All models simulate a clear linear relationship between $\Delta \text{lw}^\uparrow$ and ΔT with an intercept of around -1 W m^{-2} , which is the longwave adjustment. Hence the longwave component contributes about 1/3 to the overall RAs, while the shortwave RAs make up the other 2/3.

The same method was applied to the cloud radiative effect anomaly ($\Delta \text{CRE}_{\text{TOA}}$) and its short- and longwave components ($\Delta \text{CRE}_{\text{TOA,sw}}$ and $\Delta \text{CRE}_{\text{TOA,lw}}$), respectively (Fig. A2 in the Supplements). If an approach, similar to the IRF-ERF
 225

TOA flux anomalies	ERF contribution (W m^{-2})	RA contribution (W m^{-2})
Δ TOA radiative budget	-6.42 ± 0.49	-3.57 ± 0.46
$\Delta\text{sw}^\downarrow$	-13.62 ± 0.02	0.44 ± 0.01
Δsw^\uparrow	-6.25 ± 0.59	-2.17 ± 0.51
Δlw^\uparrow	-0.96 ± 0.16	-0.72 ± 0.21
$\Delta\text{CRE}_{\text{TOA}}$	3.33 ± 0.73	1.26 ± 0.72
$\Delta\text{CRE}_{\text{TOA},\text{sw}}$	3.0 ± 0.59	1.06 ± 0.56
$\Delta\text{CRE}_{\text{TOA},\text{lw}}$	0.33 ± 0.19	0.2 ± 0.16

Table 2. Multi-model-mean intercepts of different TOA fluxes derived via regression method (ERF contribution in W m^{-2}) and its deviation from the initial response (RA contribution in W m^{-2}) calculated as the difference between linear regression intercept and the first month response (\sim IRF contribution).

approach is applied and the initial $\Delta\text{CRE}_{\text{TOA}}$ is subtracted from the intercept, as it is purely a cloud masking effect, the remaining cloud adjustment is $3.3 \text{ W m}^{-2} - 2 \text{ W m}^{-2} = 1.3 \text{ W m}^{-2}$, mostly stemming from $\Delta\text{CRE}_{\text{TOA},\text{sw}}$, which is a substantial contribution to the overall RA of 3.6 W m^{-2} . Hence, we find that cloud adjustments counteract parts of the initial solar forcing as was found by several other studies (Smith et al., 2018; Salvi et al., 2021; Virgin and Fletcher, 2022). The reason for the positive shortwave cloud adjustment found in this study might be the decrease of tropical cloud cover in the boundary layer, as the location of strong positive $\Delta\text{CRE}_{\text{TOA},\text{sw}}$ (shown in the Supplements, Fig. A4) often coincides with decreased cloud cover at 850hPa (Fig. 10). However, our approach does not allow for a quantification of single contributions to the overall cloud adjustments. Moreover, the uncertainty of cloud radiative effect anomaly is high and models disagreed on the sign of slope of the linear fits. The intercepts for the cloud radiative effect anomaly and its components are also provided in Table 2.

3.2 Rapid adjustments on different time scales

In the following section we analyse the temporal development of TOA radiative budget anomaly and its components, as well as of the surface temperature anomaly. The aim is to better understand the contributions of the single components to the total TOA budget anomaly and its interaction with surface temperature change. In contrast to the linear regression method, in this section we look at temporal developments over the first hours, days and months, in order to learn more about the underlying adjustments.

3.2.1 Temporal development of ΔTOA radiative budget and ΔT

Figure 2 shows the global mean of the three components of the TOA radiative budget, downward and upward shortwave flux anomaly ($\Delta\text{sw}^\downarrow$ in orange, Δsw^\uparrow) and upward longwave flux anomaly (Δlw^\uparrow), together with the TOA radiative budget anomaly. Fluxes are defined positive downwards, so that added up, the three flux anomalies yield the total TOA budget anomaly. Only MRI-ESM2-0 provided TOA shortwave radiative fluxes as daily data, while the other three models only provided monthly




Figure 2. Global mean anomaly of radiative fluxes at TOA in W m^{-2} for four different time scales after onset of forcing (100 hours, 30 days, 12 months, 150 years). Plotted are the downward shortwave flux anomaly at TOA (orange), the upward shortwave flux anomaly at TOA (brown), the upward longwave anomaly at TOA (olive) and the effective flux anomaly at TOA (red). Shading around the multi-model mean shows multi-model standard deviation. If available, the results for all four models (thin lines) were plotted together with the model-mean (thick line). In addition to that, the instantaneous radiative forcing (IRF) is plotted as a grey dash-dotted line. Upward fluxes were multiplied by $\cdot(-1)$, such that the three radiative flux anomalies (orange, olive and brown) add up to the effective flux anomaly at TOA (red), which is the same as the black curve in Fig. 4.

data. Hence, Fig. 2 only shows the results of one model for shortwave flux anomalies and the total TOA budget anomaly for the first two time scales. The upward longwave radiative flux was provided as daily output by all models and is plotted accordingly.

$\Delta\text{sw}^\downarrow$ anomaly shows the reduction of the solar constant. It stays constant over all time scales and only displays a yearly periodic deviation due to the elliptical orbit of the Earth around the sun. It is the same for all four models, since the underlying assumption of the abrupt-solm4p experiment is an instantaneous and constant reduction of the solar constant by 4 %, corresponding to a decrease of downward shortwave flux by -13 W m^{-2} .

As described before, Δsw^\uparrow shows an instantaneous reaction, which is part of IRF, since it is a change in TOA radiation in reaction to forcing, without any changes in atmospheric state. Hence, the IRF is the sum of incoming and outgoing shortwave anomaly at $t = 0$ and is around -10 W m^{-2} according to experimental design. Δsw^\uparrow shows first changes after 10 days, although no clear trend can be recognised. This is a reaction to changes in surface albedo, beginning after roughly 10 days and mostly connected to increased snowfall over land in high northern latitude and first changes in snow cover and sea ice extent in Antarctica. Furthermore, changes in cloud properties, beginning after a few hours, influence Δsw^\uparrow over the first month, but there is not one dominating process, but rather a complex interplay of mechanisms that result in the global mean temporal change. Over the first year, albedo and cloud properties show significant changes. Nevertheless, the upward shortwave anomaly stays relatively constant, indicating that the different climate variables and local variability approximately cancel out each other's shortwave radiative effects. Over longer time scales, the multi-model mean of upward shortwave flux anomaly decreases, changes its sign around 50 years after onset of forcing and continues to decrease up until 150 years. However, the individual model runs do not agree on the sign of the anomaly on these longer time scales. This is due to opposing signs in high latitudes and tropical regions, and the models disagree, which region dominates the global mean effect (see Figure A3 in




Figure 3. Global mean of near surface temperature anomaly in K for four different time scales after the onset of forcing (100 hours, 30 days, 12 months, 150 years). The results of the four participating models are plotted (cyan: CESM2, orange: CanESM5, blue: IPSL-CM6A-LR and green: MRI-ESM2-0) together with a multi-model mean (red, bold) and multi-model standard deviation (shaded area around multi-model mean). For the fourth timescale, the y-axis has been adjusted to account for the stronger changes on longer time scales.

the Supplements). This is a significant difference in the long term prediction of the models, but was not further analysed here, since this study is focused on short term effects.

In contrast to the shortwave fluxes, there are no immediate effects of the reduction of the solar constant on the $\Delta l w^{\uparrow}$. This is expected, because it is closely linked to changes in surface and atmospheric temperature, which need some time to adapt. Similar to $\Delta s w^{\uparrow}$, $\Delta l w^{\uparrow}$ displays first changes around 10 days after the onset of forcing. It is overall slightly increased over the following days and only temporarily changes its sign. The multi-model mean always stays positive, indicating a reduction in the amount of longwave radiation to space, because the atmosphere and surface reduce their temperature in reaction to the reduced incoming solar energy (the Planck effect). This effect continues and all models simulate a further increasing $\Delta l w^{\uparrow}$ for longer time scales.

The TOA radiative budget anomaly is overall negative due to the negative forcing and corresponds to the IRF at $t = 0$. It is dominated by the shortwave effects on the time scales of hours and days, but in the course of months, the increasing longwave gain slowly reduces the TOA imbalance until it approaches a new balance over the following decades and centuries.

Analogous to the TOA radiative flux anomalies, the global mean near surface temperature anomaly ΔT is shown in Fig. 3. It becomes increasingly negative immediately after the onset of forcing due to decreased absorption of shortwave radiation at the surface. ΔT is of the order of -0.25 K over the first month, continuously decreasing to values < -1 K after four months. The four participating models agree well on the global mean ΔT during the first year, but differ in their long-term ΔT , depending on the model's climate sensitivity.

3.2.2 Simulated vs. calculated Δ TOA radiative budget

In this study, we are interested in rapid adjustments to solar forcing in fully coupled climate models. In those models, the surface temperature begins to adapt from the onset of forcing, leading to a possible overlap of adjustments and temperature mediated effects (feedbacks).
285

However, when calculating the TOA budget anomaly N via Eq. 1 by multiplying the ΔT with the feedback parameter λ and adding the ERF, both derived from the linear regression in Sect. 3.1, a significant disagreement simulated TOA budget anomaly (Fig. 2) and TOA budget anomaly calculated via Eq. 1 becomes apparent on the shorter time scales. Figure 4 shows the simulated TOA budget anomaly anomaly (N_{sim} in black) together with the surface temperature mediated TOA budget
290 anomaly calculated from ΔT adding only the IRF ($N_{\text{calc,IRF}}$ in red) and the complete ERF ($N_{\text{calc,ERF}}$ in blue) for the four different time scales.

During the first time scale no adjustments or feedbacks take place that would lead to considerable changes in N . Hence, the red and black curves do not show a strong deviation. However, on all other time scales there is a significant offset between N_{sim} and $N_{\text{calc,IRF}}$. Especially for the time scales of days and months after the onset of forcing, the trend of $N_{\text{calc,IRF}}$ cannot
295 explain the trend of N_{sim} . For the fourth time scale, both model means show the same trend, but with a constant offset, which corresponds to the rapid adjustments (RA). When adding the RA and IRF yielding $N_{\text{calc,ERF}}$ (blue), the long term trends of calculated and simulated TOA flux change agree well. However, the first three time scales show considerable deviations between N_{sim} and $N_{\text{calc,ERF}}$ and changes in surface temperature are clearly not sufficient to explain the simulated TOA radiative budget variability.

Hence, we argue that the first three time scales are dominated by adjustment processes while temperature mediated changes only play a minor part. Clearly, this does not allow for a perfect distinction between adjustments and feedbacks. However, this approach has the advantage that no restrictions have to be made to surface temperature (e.g. fixed SST experiments), which fail to capture all changes in circulation and no assumptions are made on the linearity of adjustment processes, as is done in case of radiative kernels. This study aims to understand changes in the atmospheric state during the first year after onset of forcing
305 and based on the beforehand arguments, we assume this time scale to be dominated by adjustment processes.

Unfortunately, only one model (MRI-ESM2-0) provided daily TOA radiative flux data. Therefore, although a deviation of N_{sim} and $N_{\text{calc,ERF}}$ is clearly visible, a clear attribution to rapid adjustments to the forcing is difficult, as internal variability can also lead to difference between N_{sim} and $N_{\text{calc,ERF}}$. However, all models provide data for the following months after onset of forcing. Although there is some deviation between the models which is likely caused by internal variability there is also a
310 common trend (model-mean, thick black line), which cannot be explained by the changes in surface temperature or surface albedo (not shown). Instead, adjustments in atmospheric temperature profile, water vapour and cloud properties seem to be the dominating source of variability during the first days and months. Hence, the TOA budget variability is a result of a complex interplay of several atmospheric and cloud variables, with partly compensating effects. Changes in cloud properties have been found to be one of the major sources of model uncertainty on all time scales. Reducing the uncertainty on short time scales




Figure 4. Same as Fig. 2, simulated and calculated TOA radiative budget changes. Shown are TOA effective flux change, based on the simulated surface temperature change, only considering IRF ($N_{\text{calc,IRF}}$, red), the calculated TOA effective flux change considering IRF+RA ($N_{\text{calc,ERF}}$, blue) and the simulated TOA effective flux change (N_{sim} , black).

would allow for a better quantification of the ERF, thereby improving long term predictions. Therefore, the aim of this study is to identify characteristic patterns of cloud adjustment in response to a negative shortwave forcing.

The next section analyses the temporal development of the cloud radiative effect anomaly over the three adjustment time scales as well as the fourth time scale representing the long term temperature mediated response. The cloud radiative effect can be an indicator to cloud adjustments, which are a major source of uncertainty in climate models. However, the cloud radiative effect also includes changes in other atmospheric variables (cloud masking), which is discussed in the following section.

3.2.3 Adjustments of the cloud radiative effect at TOA

Figure 5 shows the temporal evolution of the global mean cloud radiative effect anomaly at TOA ($\Delta\text{CRE}_{\text{TOA}}$) and its short- and longwave components ($\Delta\text{CRE}_{\text{TOA,sw}}$ and $\Delta\text{CRE}_{\text{TOA,lw}}$). As described before, only MRI-ESM2-0 provided daily data, which is plotted for the first two time scales.

$\Delta\text{CRE}_{\text{TOA}}$ is positive for all time scales, indicating a decrease of the generally cooling effect of clouds, partially counteracting the negative forcing. However, since this positive anomaly is apparent from the first time step, the initial value is not attributable to changes in cloud properties. It rather is a result of clouds masking changes in the clear-sky budget. After five days $\Delta\text{CRE}_{\text{TOA}}$ starts to deviate from its initial value and varies with no clear pattern during the first month. During this time period it follows the same shape as the total TOA budget anomaly. Hence, changes in cloud properties seem to be the dominant source of variability of N_{sim} during the first months. This is supported by the fact that global mean clear sky fluxes show near to no variability during the first five months (not shown). After six months, the initial forcing moves from the southern high latitudes to the northern high latitudes. Due to differing surface albedo on the two hemispheres and therefore changes in the clear-sky TOA radiative budget, cloud masking is expected to change at this point.

$\Delta\text{CRE}_{\text{TOA}}$ further increases over the following years. After about 10 years $\Delta\text{CRE}_{\text{TOA}}$ remains relatively constant at 4 W m^{-2} . A slight decreasing trend is visible in the multi-model mean, even though three out of four models simulate a sta-




Figure 5. Global mean of TOA cloud radiative effect anomaly ($\Delta\text{CRE}_{\text{TOA}}$, red) in W m^{-2} for four different time scales after onset of forcing (100 hours, 30 days, 12 months, 150 years). Additionally, shortwave and longwave TOA cloud radiative effect anomalies ($\Delta\text{CRE}_{\text{TOA,sw}}$ and $\Delta\text{CRE}_{\text{TOA,lw}}$) are shown in brown and orange. Their respective intermodel standard deviation is shown as shading around inter-model means.

bilization of $\Delta\text{CRE}_{\text{TOA}}$ for longer time scales. This is due to the continuous decrease of $\Delta\text{CRE}_{\text{TOA,lw}}$, simulated by the CanESM5 model. Although this model shows the strongest temperature decrease out of the four models, it does not systematically predict the strongest changes in other cloud properties. Hence, the continuous decrease of $\Delta\text{CRE}_{\text{TOA,lw}}$ likely is a strong cloud masking effect that dominates over any possibly positive $\Delta\text{CRE}_{\text{TOA,lw}}$ effects due to changes in cloud properties.

340 $\Delta\text{CRE}_{\text{TOA,sw}}$ is overall positive on all time scales and dominates $\Delta\text{CRE}_{\text{TOA}}$. It seems to be the major source of variability, while $\Delta\text{CRE}_{\text{TOA,lw}}$ shows no clear sign over the first three time scales and overall less variability than $\Delta\text{CRE}_{\text{TOA,sw}}$.

As discussed before, especially signals during the first days will also be influenced by internal variability. However, in the 6th Intergovernmental Panel on Climate Change (IPCC) report anthropogenic forcing in 2019 was quantified to be of the order of 2.7 W m^{-2} (Forster et al., 2021a). Sippel et al. (2021) showed that this allows a robust detection of forced warming, 345 significantly exceeding climate variability. Hence, an instantaneous forcing of -10 W m^{-2} , which is four times stronger, is expected to lead to adjustments that exceed internal variability. Nevertheless, to account for possible effects of internal variability, in the following section on characteristic fingerprints we only assume a signal to be a rapid adjustment, if three out of the four participating models agree on the sign.

3.3 Characteristic adjustment fingerprints in climate variables

350 The following sections explore the response of surface and atmospheric temperature, as well as relative humidity, vertical velocity and cloud cover in order to identify characteristic adjustment fingerprints to reduced shortwave radiation availability. Since the global mean TOA budget variability is an effect of a complex interplay of several variables and local effects, the following sections examine global distributions of the beforehand mentioned variables, in order to better understand the underlying mechanisms.




Figure 6. Multi-model mean global distribution of near surface temperature anomaly in K averaged over four different time scales after onset of forcing (100 hours, days 5–30, months 2–12, years 120–150). The multi-model mean was calculated by interpolating the other models to the grid spacing of the CanESM5 model. Regions, where less than 3 of 4 models agreed on the sign of anomaly, are dotted. Note the different scales of the colour bars for better contrast.

355 3.3.1 Adjustments of surface and atmospheric temperature

The effects of the reduction of the solar constant are visible in a variety of climate variables already a few hours after the onset of the perturbation.

Figure 6 shows the geographical distribution of the near surface temperature anomaly averaged over the first three time scales depicted in Fig. 3 (hours, days and months, respectively) and a fourth time scale averaged over the years 120–150 representing
360 the long-term development.

During the first 100 hours the land surface responds to the forcing by cooling down, while the ocean surface temperature stays approximately constant due to its higher heat capacity. The strongest cooling occurs over Antarctica, because of the 24 hour exposure to solar radiation at time of forcing onset (1 January). Areas of low heat capacity (e.g. Antarctica and Sahara) generally exhibit a stronger surface temperature decrease compared to other regions on the same latitude, because
365 the same amount of heat reduction will lead to stronger decrease in temperature. In contrast to the overall cooling of surface temperature, the Arctic, which does not experience any solar forcing in January, warms up during the first hours and days after onset of forcing. The reduced temperature gradient between the Tropics and the Arctic leads to a reduction in polar night jet strength, which perturbs the polar vortex. Cold air outbreaks as well as warm air intrusions result in strong changes of surface temperature over days and the first month. Intrusion of warm, moist air masses into Arctic latitudes increases cloud
370 cover, which reduces the amount of longwave radiation lost to space, leading to local surface temperature rise. However, the individual pattern of surface temperature increase will depend on the base state of the respective model. Hence, although all models simulate an increase in Arctic temperatures, the specific location varies, which leads to higher uncertainty in the Arctic response compared to other regions. Also the southern hemisphere shows a pattern of warming and cooling, however less pronounced, due to the positioning of the forcing.




Figure 7. Multi-model mean, temporal and zonal mean vertical profiles of atmospheric temperature anomaly in K as a function of latitude for four different time scales after onset of forcing (100 hours, days 5–30, months 2–12, years 120–150). Regions, where less than three out of four models agree on the sign of anomaly, are dotted. Note the adjusted colour scales for the different time scales.

375 After one year, Arctic amplification begins to emerge as an overall stronger cooling of the surface over Arctic latitudes compared to lower latitudes and Antarctica. The ocean surface starts to cool down, although the decrease is still smaller compared to land surfaces.

The average over years 120–150 shows an overall cooling in all regions. Arctic Amplification is apparent as a stronger cooling over Arctic latitudes. North of Antarctica, where sea ice extent is increased compared to the piControl-runs, the surface
380 cooling is slightly stronger compared to the land surface of Antarctica, due to more severe changes in surface properties compared to the control run. The differences between land and sea surface are less apparent compared to the shorter time scales because also the deeper ocean adapts to the new energy balance.

Figure 7 shows the zonal mean of atmospheric temperature anomaly averaged over time scales of hours, days and months and the years 120–150, respectively.

385 During the first 100 hours, the stratosphere cools down quickly, especially above Antarctica, where the radiative forcing is strongest. Since the stratosphere is nearest to the source of forcing, the cooling of the stratosphere is stronger than the cooling of the troposphere during the first hours and days. The cooling effects are strongest on the highest altitudes, where most of the high frequency radiation is absorbed.

At the near surface layers, cooling above Antarctica is further amplified, because the surface temperature drops rapidly,
390 which reduces sensible heat flux.

After one month atmospheric temperature change is highly variable in northern latitudes, due to the beforehand described perturbation of the polar vortex. Warm air intrusions into Arctic latitudes strongly increase the atmospheric temperature in surface near layers.

Over the first year, the troposphere adjusts to the solar forcing via an overall reduction of tropospheric temperature, especially
395 in the mid and upper troposphere in the tropics. This pattern resembles the long-term climate response, where it contributes




Figure 8. Same as Fig. 7 but for atmospheric relative humidity anomaly in %.

to a lapse rate feedback. However, similar atmospheric temperature adjustments have been documented under solar forcing in fixed-SST experiments (e.g. Salvi et al. (2021)). In these experiments, land surface temperatures can still respond, but their change is strongly tied to nearby ocean temperatures through atmospheric coupling. Consequently, this limits any substantial climate feedbacks from developing, leaving predominantly fast adjustment processes. This indicates that the pattern found in this study could be driven by changes in atmospheric absorption, humidity and convection in reaction to the forcing itself, rather than being a surface temperature mediated effect. We discuss this point further when analysing the results for humidity and cloud fraction.

On longer time scales the tropospheric temperature continues to decrease. In the tropics this is most pronounced in the higher troposphere, while at high latitudes the cooling is stronger in the lower troposphere, because the characteristic atmospheric temperature profiles in these regions react differently to a decrease in surface temperature. In higher latitudes, the tropopause descends. Hence, stratospheric vertical increase of temperature starts at lower altitudes, which manifests as a warming at 200 hPa at the poles. At $\pm 30^\circ$ the stratosphere cools at all pressure levels.

3.3.2 Adjustments of humidity, vertical velocity and cloud fraction

Figure 8 shows the vertical distribution of zonal mean relative humidity anomaly. While specific humidity reacts to the cooling air temperature by decreasing in the whole atmosphere, especially strong in the tropics and over all time scales (not shown), relative humidity displays a pattern of increase and decrease. During the first 100 hours, relative humidity increases, apart from southern tropical latitudes that show a decrease. During the first month, an overall increasing pattern remains in the tropics, while higher latitudes show decrease. During the first year, the high troposphere dries, while the middle tropospheric layers experience a moistening. The lower troposphere shows a regular pattern of moistening and drying and only at $\pm 65^\circ$ the whole troposphere dries. The pattern further intensifies on long-term time scales.

To further analyse the short-term adjustments of relative humidity, it makes sense to differentiate between trends on land and over sea, since they react very differently to the forcing on short time scales due to their different heat capacities. Hence,

Fig. 9 shows the zonal means of relative humidity change for the first two time scales averaged over ocean (Fig. 9a) and land (Fig. 9b) only. To investigate the effects of changing vertical atmospheric stability Fig. 9 also shows the same for change of vertical velocity (Δw_{ap}) (averaged over ocean (c) and land (d)). As w_{ap} is defined positive downwards, red colours of Δw_{ap} indicate an increase of downward movement or a decrease of upward movement and vice versa for blue colours.

Over the ocean, the lower troposphere cools down quicker than the sea surface on time scales of hours to days. This leads to a decrease in vertical stability, which then increases convective activity, especially over tropical oceans (see Fig. 9c). Increased convection dries the boundary layer, thereby reducing near surface specific humidity. In order to compensate for the deficit in humidity, evaporation and thereby latent heat flux increases, which then results in an increase in surface relative humidity (see Fig. 9a) and precipitation in tropic latitudes.

Over land, vertical stability very much depends on location, albedo and heat capacity of the respective surface type. There is a strong reduction of convection over the tropics due to the reduced radiative forcing (see Fig. 9d) and resulting reduction in sensible heat flux, especially over the Sahara. This leads to trapping of moisture in the boundary layer and an increase in relative humidity in surface near layers (see Fig. 9b). Latent heat flux is reduced in equatorial regions and overall less moisture is transported into the higher troposphere, which reduces relative humidity in the free troposphere (see Fig. 9b).

The significant differences between land and ocean surface response are strongest during the first hours after onset of forcing, but they are still clearly visible during the first month and lead to changes in land-sea-circulation and moisture transport.

Cao et al. (2012) described similar effects, although of opposite sign in response to a +4% increase of solar constant experiment. After five days they found a decrease of relative humidity in the boundary layer of 1% over land and an increase of around 0.1% over ocean. This is a similar magnitude compared to our findings, however, with our approach we also found an increase in surface near relative humidity in response to a negative solar forcing, as the drying of the boundary layer due to reduced stability is compensated by increased latent heat flux.

For vertically-resolved atmospheric cloud fraction the four models only provided monthly data. Hence for the zonal mean of atmospheric cloud fraction anomaly, shown in Fig. 10 no data was available for the first time scale and the first monthly mean was plotted as the "first 31 days" time scale. This is not fully identical to other plots of this kind, because, if available, the daily data from day 5–30 was averaged, not including the first 4 days, while in case of the atmospheric cloud fraction, the average of the whole month is plotted.

As expected, the atmospheric cloud fraction anomaly pattern in Fig. 10 is overall very similar to the pattern of relative humidity in Fig. 8.

During the first month, the atmospheric cloud fraction increases in the tropics over ocean, while it decreases over land. However, the increase over the ocean dominates the zonal mean. Higher latitudes show a pattern of decreasing and increasing cloud fraction, where circulation strength changes (also visible in the vertical velocity anomaly in Fig. 9).

Over the first year, tropical regions experience the strongest negative radiative forcing leading to less absorption of shortwave radiation in the troposphere and reduced convective heating from the lower troposphere. Because the colder air contains less water vapour, also the greenhouse effect is reduced, a positive feedback. This way, condensation levels are reached at lower altitude and high clouds shift downwards. This leads to a decrease in cloud cover at 150 hPa and an increase in cloud cover




Figure 9. Zonal mean of relative humidity anomaly in % for the first 100 hours and days 5–30 (corresponds to the first two time scales of Fig. 7) averaged over ocean (a) and land (b) only and the same plots for vertical velocity (w, positive downwards) anomaly in hPa day^{-1} averaged over ocean (c) and land (d).




Figure 10. Same as Fig. 7 but for atmospheric cloud fraction anomaly in % on 49 pressure levels (the three other models were interpolated to CanESM5 levels). Only monthly data was provided. Hence, the first time scale (hours 0-100) is missing.

around 250 hPa, mirroring the same effects in relative humidity. An exception of the overall increasing cloud fraction in the troposphere is just above $\pm 60^\circ$, where the subtropical jet stream strength is reduced, due to reduced temperature contrast
455 between mid- and high latitudes and weaker storm systems produce less dynamic lifting, reducing cloud formation in those regions.

On longer time scales, the overall pattern further intensifies, as do atmospheric temperature and relative humidity patterns.

A distinct drying pattern in the relative humidity at the lower troposphere in the tropics (see Fig. 8), is not visible in the atmospheric cloud fraction, because it appears at a height of generally low cloud cover.

460 4 Discussion

The aim of this study was to understand and analyse atmospheric processes that take place on time scales of hours, days and months after the onset of a negative solar forcing. We did so using the output of four different climate models that participated in the abrupt-solm4p experiment of CFMIP of CMIP6. These fully coupled simulations provide the opportunity to get a more realistic view on rapid adjustments in the climate system, as all components of the climate system are allowed to react to the
465 forcing at the same time, rather than fixing (sea) surface temperatures. This allows for a more realistic interplay between the different components, although a -4% solar constant forcing still needs to be considered a highly idealized forcing compared to e.g. transient climate change.

Our approach is based on the assumption that the short timescales of hours, days and months are dominated by rapid adjustments rather than by surface temperature mediated changes. We showed in Fig. 4 that for these three time scales, the
470 simulated change in surface temperature cannot explain the simulated change in TOA fluxes. Only when adjusting it by the rapid adjustments found in the linear regression plots of Sect. 3.1, temperature mediated TOA flux change and simulated TOA flux change start to coincide after around 4 years in model mean. This corresponds to the time, the linear regression plots show non-linearities and is likely connected to adjustments of system with high inertia, like cryosphere and vegetation, that happen on longer time scales or a delay in the response of the deeper ocean compared to the surface (SST-pattern effect, e.g. Andrews
475 et al. (2015)). Since our study concentrates on adjustments during the first year, these kinds of adjustments are not considered here.

Having the atmosphere, surface and deep ocean react to the forcing at the same time, poses the challenge of a possible overlap of adjustments to the forcing and temperature mediated processes. Nevertheless, our findings do agree in many points with findings of other studies that applied a variety of different methods, often more restrictive compared to our approach,
480 which we discuss in the following.

Smith et al. (2018) analysed adjustments to five different forcing agents, among them a $+2\%$ solar forcing, using 11 GCMs with fixed SST in combination with radiative kernels. Similar to our results, they found that in case of solar forcing, rapid adjustments reduce ERF. However, the adjustments we found using the linear regression method were overall stronger ($\sim 35\% \pm 5\%$ of IRF) in relation to the original forcing than they were in case of Smith et al. (2018) ($\sim 10\% \pm 5\%$). According to their
485 findings, the change in surface, tropospheric and stratospheric temperature adjustment contribute the most to counteracting

the forcing. Our method does not allow a quantification of the different effects in the same way, but we also see a decrease in temperature for surface as well as atmospheric temperatures in global mean, which will reduce the TOA forcing. The atmospheric cooling leads to a reduction of specific humidity in the whole atmosphere, which reduces the warming effect of water vapour in the atmosphere and is hence expected to increase TOA forcing, which is also shown in Smith et al. (2018). We
490 find a very similar pattern of cloud fraction change to that reported by Smith et al. (2018), though of opposite sign, since this study was based on a reduction of the solar constant. We calculated the anomaly of cloud radiative effect in order to estimate the influence of cloud adjustments on the ERF. However, the cloud radiative effect anomaly contains masking effects, which hamper the comparability with studies that specifically quantified cloud adjustments. Nevertheless, when assuming that all initial signals in $\Delta\text{CRE}_{\text{TOA}}$ can be attributed to masking, rather than changes in cloud properties, this initial signal can be
495 subtracted, leaving a more variable, but positive cloud adjustment, like it was found by Smith et al. (2018).

Russotto and Ackerman (2018) conducted a study using G1 data from the geoengineering MIP (GeoMIP) of CMIP6, where a fourfold CO_2 increase is balanced by a matching decrease in solar constant. The decrease is between -3.2% to -5% , thus of similar magnitude as was used in the abrupt-solm4p experiment. That experimental setup avoids a long term change in surface temperature. Hence, all signals are a mix of adjustments to the two different forcing agents. They find a significant
500 cooling of the higher troposphere, as was shown in our results and a warming of the polar regions, which we also detected. However, in their case it is probably linked to the so-called "residual polar amplification", which can be seen in only CO_2 forcing experiments and is of higher magnitude and higher SNR than the signal, found in this study. A significant difference in the atmospheric temperature change is the stratospheric cooling, which is much stronger in the results of Russotto and Ackerman (2018) due to adjustments to $4\times\text{CO}_2$ because of the increased LW radiation to space (Manabe and Wetherald, 1975).
505 This effect is much stronger than the stratospheric cooling in response to solar forcing, which we see in our results. Although of similar magnitude, cloud fraction changes in the G1 experiment seem to be dominated by adjustments to CO_2 , since Russotto and Ackerman (2018) find opposite signals to what we found. This is supported by the findings of Smith et al. (2018), who estimate cloud adjustments to CO_2 forcing to be twice as large as to a positive solar forcing with twice as strong IRF. However, Smith et al. (2018) quantified a positive solar forcing and Aeronson et al., 2024 show that responses to positive and negative
510 solar forcing can differ, not only in sign, but also in strength and pattern.

Furthermore, for their solp4p experiment Aeronson et al. (2024) find an increasing cloud cover over tropical land regions, decreasing cloud cover in higher latitudes over land and an overall reduction in cloud cover over ocean. We find similar patterns, though of opposite sign, as expected for a reduction of solar constant. The same is true for the CRE components they calculated for a solp2p experiment. However, when applying their newly developed method, Aeronson et al. (2024) find an overall positive
515 cloud radiative adjustment to a $+4\%$ solar constant forcing, meaning that clouds further increase the initial solar forcing rather than reducing it, mostly via their shortwave effects due to changes in cloud fraction. This is the opposite effect compared to what was found by e.g. Smith et al. (2018) or Virgin and Fletcher (2022) and what our results indicate. Aeronson et al. (2024) show that there can be substantial differences in adjustments to positive and negative solar forcing and linearity cannot be assumed. Nevertheless, they also find the opposite sign to Smith et al. (2018), who also analysed positive solar forcing
520 simulations. All studies report high uncertainty for cloud adjustments and, similar to our study, Aeronson et al. (2024) only had

data from four climate models they could base their analysis on. In their case one model predicts negative cloud adjustments and only two models show significantly positive adjustments. Moreover, their method combines fixed SST with fully coupled model runs in order to quantify adjustments to solar forcing. Several studies have shown that the two approaches, although showing similar results, can also show deviations due to differences in e.g. dynamical adjustments of land-sea-circulation.

525 The approach of Aeronson et al. (2024) may mistakenly interpret CO₂ forcing offsets caused by the fixed SST method as adjustments to solar forcing.

All in all this disagreement shows how challenging a proper quantification of cloud adjustments is and that they remain a source of high uncertainty in climate models.

Salvi et al. (2021) conducted a study on adjustments, using idealized atmospheric heating experiments, which were then fitted to the diagnosed heating curve of different forcing agents. This way, they found that the vertical center of mass, what they call the "characteristic altitude" determines the cloud adjustments. Like Smith et al. (2018) and our study they found negative cloud adjustments to in their case a +3% solar forcing. In their observed and fitted data they found a decrease in relative humidity and cloud fraction at around 300 and 800hPa, which corresponds well with the increased cloud fraction we find in these altitudes. There is some deviation in the UTLS region, where we find a significant reduction in cloud fraction and

535 relative humidity, while Salvi et al. (2021) only diagnosed minor changes in cloud fraction and the fitted algorithm predicted a decrease. This again could either be an effect of different methods, as Salvi et al. (2021) themselves attribute the deviation to insufficient resolution of the vertically heating curves in these high altitudes. Nevertheless, some differences in response to positive and negative solar forcing can take place, like Aeronson et al. (2024) found, especially when considering non-linear processes like phase changes in clouds.

540 Moreover, our study supports the findings of other studies like Kamae et al. (2019) that adjustments can also depend on the season of initialization. The strong response in Arctic surface temperatures, which we attribute to a disruption of the polar vortex, is an effect of initializing the forcing in January.

A challenge our study faced was the sparse dataset it is based on. Only four models participated in the solm4p experiment, each providing one run. A higher number of models and ensembles would increase the reliability of our findings. However,

545 due to the strong nature of the forcing and since every model initialized the experiment from a different point in their piControl runs, we do expect that the four models cover different climate variability modes. Hence, an agreement of several models on the sign of an adjustment effect and overall high model-mean anomalies indicate a signal that exceeds possible climate variability. For all analysed climate variables, the first month is the time scale with lowest agreement of the models on the sign of the signal. This is to be expected, since responses will strongly depend on the position of e.g. pressure systems at the moment

550 of onset of forcing. For example, this becomes apparent in the Arctic surface temperature anomaly during the first month, where all models simulated strong changes in surface temperature. We attribute those to a disruption of the polar vortex, but the actual pattern differs between the models, since it will be influenced by the original state of the atmosphere. Nevertheless, Fig. 6 shows strong signals for several high latitude regions, where at least 3 out of 4 models agree on the sign. Therefore, we interpret it as a rapid adjustment rather than a random signal due to climate variability and expect to find similar signals for

555 more realistic forcing scenarios like volcanic eruptions.

One surprising aspect of this study was the disagreement of the long term development of the models, especially when examining cloud related variables. While the models agree well on the TOA budget anomaly, they differ in how the long- and shortwave components contribute to this and how the fluxes interact with changes in cloud properties. In a number of cloud variables and hence, in the cloud radiative effect anomalies, high latitudes and tropics show an opposite response and the global mean signal is determined by the dominating region, which differs between the models. Further investigation into the single models and parametrization as well as possible tuning would be necessary to understand the different outputs. Though this was out of the scope of this study, it shows how even climate models of the newest generation still struggle to simulate short and long term change of cloud properties and further research on the topic is necessary.

5 Conclusion

This study offers insights into the dynamics of short-term adjustments within the climate system in response to an instantaneous radiative forcing. This was done through the analysis of the abrupt-solm4p experiment of CMIP6. By simulating a 4 % reduction in the solar constant, the experiment provides a valuable framework for understanding how the Earth climate system reacts to changes in solar energy input, especially on shorter time scales of days to month.

Following the linear regression method developed by Gregory et al. (2004), we find rapid adjustments of 3.6 W m^{-2} in response to an instantaneous negative solar forcing of -10 W m^{-2} . Other studies like Smith et al. (2018) also found that RA counteracted an initial solar forcing but of smaller magnitude ($\sim 10\%$). However, the amount of reduction differs, based on the method and number of participating models.

For this study, data from fully coupled climate models was used, leading to a possible overlap of adjustment and feedback mechanisms. However, it was shown that surface temperature change cannot explain the variability of the TOA radiative budget over the course of the first hours, days and months. Instead, the cloud radiative effect anomaly was found to closely match the TOA budget variability, indicating cloud property changes to be a major contribution to overall atmospheric adjustments. However, the method applied in this study did not allow for a quantification of cloud adjustments at TOA. Instead, global distribution patterns of a number of climate variables were analysed in order to find characteristic patterns (fingerprints) of atmospheric adjustments that could help explain the underlying mechanisms.

The analysis reveals significant alterations in various climate variables, including surface and atmospheric temperatures, relative humidity and vertical velocity as well as a number of cloud properties. All models simulate decreasing atmospheric and surface temperatures, beginning immediately after the onset of forcing in January. One exception was an increase of Arctic surface temperature, which we attribute to a weakened polar night jet and thereby disruption of the polar vortex leading to warm air intrusions. Significant differences were found between the response of relative humidity over land and over ocean. Over ocean, decreased vertical stability, due to a quicker cooling of the lower troposphere compared to the ocean surface, leads to increased convection and thereby drying of the boundary layer, which is then compensated by an increased latent heat flux leading to an increase in relative humidity and mid-level cloud cover. Over land in the tropics, convection is reduced and moisture effectively trapped in the boundary layer, which leads to an increase of relative humidity in surface-near layers and a

decrease in relative humidity and cloud cover in the higher troposphere. This highlights the complex interactions and differences
590 between land and sea response. On time scales of months, overall lower temperatures lead to drying of the troposphere and the
decrease in solar energy in the tropics leads to a reduction of ascent, less convection and a descending high cloud layer.

While other experiments of CMIP6 are designed in a way that influence of seasonal and climate variability are reduced by
using a number of different initialization dates, the abrupt-solm4p simulations use only one date (1 January). This could be
viewed as a limitation of this study. Nevertheless, if the scientific community strives for a validation of new discoveries about
595 rapid adjustments in climate models with e.g. satellite measurements, more realistic simulations are crucial in order to further
improve state of the art climate models. This study showed that the positioning of the forcing can have significant influence on
the resulting short-term adjustments, in this case, the disruption of the polar vortex and consequently warming of the higher
northern latitudes.

Moreover, the described adjustments to a reduced solar constant may resemble adjustments to more realistic forcing scenar-
600 ios. After large volcanic eruptions a global stratospheric aerosol scattering layer can form a few months after the eruption. This
aerosol layer reduces the amount of shortwave radiation that reaches the surface, which could lead to similar adjustment effects
as simulated for a reduced solar constant. However, in case of the abrupt-solm4p experiment, the forcing is instantaneous and
constant, which makes it easier to differentiate between instantaneous forcing, short-term adjustments and long-term feedbacks.
Therefore, it is a useful tool on the way to a more in depth understanding of realistic forcing agents of more transient nature. By
605 elucidating the short-term adjustments that occur in response to reduced solar radiation, essential insights can be provided that
can inform risk assessments associated with solar radiation management methods. Understanding these adjustments is vital for
anticipating potential local impacts, such as droughts or floods, which can have significant consequences for communities and
ecosystems.

In conclusion, this study identifies several characteristic adjustment processes as well as a number of challenges related to the
610 current approaches in quantifying adjustments. By integrating short-term adjustment processes into climate modelling efforts,
the scientific community can improve the accuracy of long-term climate predictions and develop more effective strategies for
addressing the challenges posed by climate change.

Data availability. The CFMIP model data from the abrupt-solm4p experiment used in this study are freely available from the CMIP6
repository on the Earth System Grid Federation nodes (<https://esgf-metagrid.cloud.dkrz.de/search/cmip6-dkrz/>, World Climate Research
615 Programme, 2020).

Author contributions. CL and JQ designed the study. CL analysed the CFMIP abrupt-solm4p experiment data, produced the figures, and
drafted the initial manuscript. Both authors contributed to the writing, editing and review of the paper.

Competing interests. JQ is a member of the editorial board of Atmospheric Chemistry and Physics

Acknowledgements. This research has been supported by the Deutsche Forschungsgemeinschaft Research Unit VolImpact (FOR2820, QU 311/23-2) within the project VolCloud. This work used resources of the Deutsches Klimarechenzentrum (DKRZ) granted by its Scientific Steering Committee (WLA) under project ID bb1036.

References

- Aerenson, T., Marchand, R., and Zhou, C.: Cloud Responses to Abrupt Solar and CO₂ Forcing: 2. Adjustment to Forcing in Coupled Models, *Journal of Geophysical Research: Atmospheres*, 129, e2023JD040297, <https://doi.org/https://doi.org/10.1029/2023JD040297>, e2023JD040297 2023JD040297, 2024.
- Andrews, T. and Forster, P. M.: CO₂ forcing induces semi-direct effects with consequences for climate feedback interpretations, *Geophysical Research Letters*, 35, <https://doi.org/https://doi.org/10.1029/2007GL032273>, 2008.
- Andrews, T., Gregory, J. M., and Webb, M. J.: The Dependence of Radiative Forcing and Feedback on Evolving Patterns of Surface Temperature Change in Climate Models, *Journal of Climate*, 28, 1630 – 1648, <https://doi.org/10.1175/JCLI-D-14-00545.1>, 2015.
- Andrews, T., Smith, C. J., Myhre, G., Forster, P. M., Chadwick, R., and Ackerley, D.: Effective Radiative Forcing in a GCM With Fixed Surface Temperatures, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033880, <https://doi.org/https://doi.org/10.1029/2020JD033880>, e2020JD033880 2020JD033880, 2021.
- Bala, G., Duffy, P. B., and Taylor, K. E.: Impact of geoengineering schemes on the global hydrological cycle, *Proceedings of the National Academy of Sciences*, 105, 7664–7669, <https://doi.org/10.1073/pnas.0711648105>, 2008.
- Boucher, O., Denvil, S., Levvasseur, G., Cozic, A., Caubel, A., Foujols, M.-A., Meurdesoif, Y., Bony, S., Flavoni, S., Idelkadi, A., Mellul, L., Musat, I., and Saint-Lu, M.: IPSL IPSL-CM6A-LR model output prepared for CMIP6 CFMIP abrupt-solm4p, <https://doi.org/10.22033/ESGF/CMIP6.5110>, 2018.
- Cao, L., Bala, G., and Caldeira, K.: Climate response to changes in atmospheric carbon dioxide and solar irradiance on the time scale of days to weeks, *Environmental Research Letters*, 7, 034015, <https://doi.org/10.1088/1748-9326/7/3/034015>, 2012.
- Cao, L., Gao, C.-C., and Zhao, L.-Y.: Geoengineering: Basic science and ongoing research efforts in China, *Advances in Climate Change Research*, 6, 188–196, <https://doi.org/https://doi.org/10.1016/j.accre.2015.11.002>, 2015.
- Christensen, M. W., Gettelman, A., Cermak, J., Dagan, G., Diamond, M., Douglas, A., Feingold, G., Glassmeier, F., Goren, T., Grosvenor, D. P., Gryspeerdt, E., Kahn, R., Li, Z., Ma, P.-L., Malavelle, F., McCoy, I. L., McCoy, D. T., McFarquhar, G., Mülmenstädt, J., Pal, S., Possner, A., Povey, A., Quaas, J., Rosenfeld, D., Schmidt, A., Schrödner, R., Sorooshian, A., Stier, P., Toll, V., Watson-Parris, D., Wood, R., Yang, M., and Yuan, T.: Opportunistic experiments to constrain aerosol effective radiative forcing, *Atmospheric Chemistry and Physics*, 22, 641–674, <https://doi.org/10.5194/acp-22-641-2022>, 2022.
- Cole, J. N., von Salzen, K., Swart, N. C., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Solheim, L., Yang, D., Winter, B., and Sigmond, M.: CCCma CanESM5 model output prepared for CMIP6 CFMIP abrupt-solm4p, <https://doi.org/10.22033/ESGF/CMIP6.3533>, 2019.
- Colman, R. and McAvaney, B.: On tropospheric adjustment to forcing and climate feedback, *Climate Dynamics*, 36, 1649–1658, <https://doi.org/10.1007/s00382-011-1067-4>, 2011.
- Danabasoglu, G.: NCAR CESM2 model output prepared for CMIP6 CFMIP abrupt-solm4p, <https://doi.org/10.22033/ESGF/CMIP6.7520>, 2020.
- Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development*, 9, 1937–1958, <https://doi.org/10.5194/gmd-9-1937-2016>, 2016.
- Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J., Frame, D., Lunt, D. J., Mauritsen, T., Palmer, M. D., Watanabe, M., Wild, M., and Zhang, H.: The Earth’s Energy Budget, Climate Feedbacks, and Climate Sensitivity, in: *Climate Change 2021: The Physical Science*

Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by
660 Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang,
M., Leitzell, K., Lonnoy, , Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, , Yu, R., and Zhou, B., chap. 7, pp. 923–1054,
Cambridge University Press, Cambridge, UK and New York, NY, USA, <https://doi.org/10.1017/9781009157896.009>, 2021a.

Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild,
M., and Zhang, H.: The Earth's Energy Budget, Climate Feedbacks, and Climate Sensitivity, p. 923–1054, Cambridge University Press,
665 Cambridge, United Kingdom and New York, NY, USA, <https://doi.org/10.1017/9781009157896.009>, 2021b.

Forster, P. M., Richardson, T., Maycock, A. C., Smith, C. J., Samset, B. H., Myhre, G., Andrews, T., Pincus, R., and Schulz, M.: Recommen-
dations for diagnosing effective radiative forcing from climate models for CMIP6, *Journal of Geophysical Research: Atmospheres*, 121,
12,460–12,475, <https://doi.org/https://doi.org/10.1002/2016JD025320>, 2016.

Gregory, J. and Webb, M.: Tropospheric Adjustment Induces a Cloud Component in CO₂ Forcing, *Journal of Climate*, 21, 58 – 71,
670 <https://doi.org/10.1175/2007JCLI1834.1>, 2008.

Gregory, J. M., Ingram, W. J., Palmer, M. A., Jones, G. S., Stott, P. A., Thorpe, R. B., Lowe, J. A., Johns, T. C., and
Williams, K. D.: A new method for diagnosing radiative forcing and climate sensitivity, *Geophysical Research Letters*, 31,
<https://doi.org/https://doi.org/10.1029/2003GL018747>, 2004.

Hansen, J., Sato, M., Ruedy, R., Nazarenko, L., Lacis, A., Schmidt, G. A., Russell, G., Aleinov, I., Bauer, M., Bauer, S., Bell, N., Cairns,
675 B., Canuto, V., Chandler, M., Cheng, Y., Del Genio, A., Faluvegi, G., Fleming, E., Friend, A., Hall, T., Jackman, C., Kelley, M., Kiang,
N., Koch, D., Lean, J., Lerner, J., Lo, K., Menon, S., Miller, R., Minnis, P., Novakov, T., Oinas, V., Perlwitz, J., Perlwitz, J., Rind, D.,
Romanou, A., Shindell, D., Stone, P., Sun, S., Tausnev, N., Thresher, D., Wielicki, B., Wong, T., Yao, M., and Zhang, S.: Efficacy of
climate forcings, *Journal of Geophysical Research: Atmospheres*, 110, <https://doi.org/https://doi.org/10.1029/2005JD005776>, 2005.

Huneus, N., Boucher, O., Alterskjær, K., Cole, J. N. S., Curry, C. L., Ji, D., Jones, A., Kravitz, B., Kristjánsson, J. E., Moore, J. C., Muri, H.,
680 Niemeier, U., Rasch, P., Robock, A., Singh, B., Schmidt, H., Schulz, M., Tilmes, S., Watanabe, S., and Yoon, J.-H.: Forcings and feedbacks
in the GeoMIP ensemble for a reduction in solar irradiance and increase in CO₂, *Journal of Geophysical Research: Atmospheres*, 119,
5226–5239, <https://doi.org/https://doi.org/10.1002/2013JD021110>, 2014.

Kamae, Y. and Watanabe, M.: Tropospheric adjustment to increasing CO₂: Its timescale and the role of land-sea contrast, *Climate Dynamics*,
41, <https://doi.org/10.1007/s00382-012-1555-1>, 2012.

685 Kamae, Y., Chadwick, R., Ackerley, D., Ringer, M., and Ogura, T.: Seasonally variant low cloud adjustment over cool oceans, *Climate
Dynamics*, 52, 5801–5817, <https://doi.org/10.1007/s00382-018-4478-7>, 2019.

Malavelle, F. F., Haywood, J. M., Jones, A., Gettelman, A., Clarisse, L., Bauduin, S., Allan, R. P., Karset, I. H. H., Kristjánsson, J. E.,
Oreopoulos, L., Cho, N., Lee, D., Bellouin, N., Boucher, O., Grosvenor, D. P., Carslaw, K. S., Dhomse, S., Mann, G. W., Schmidt, A.,
Coe, H., Hartley, M. E., Dalvi, M., Hill, A. A., Johnson, B. T., Johnson, C. E., Knight, J. R., O'Connor, F. M., Partridge, D. G., Stier,
690 P., Myhre, G., Platnick, S., Stephens, G. L., Takahashi, H., and Thordarson, T.: Strong constraints on aerosol–cloud interactions from
volcanic eruptions, *Nature*, 546, 485–491, <https://doi.org/10.1038/nature22974>, 2017.

Manabe, S. and Wetherald, R. T.: The Effects of Doubling the CO₂ Concentration on the climate of a General Circulation Model, *Journal of
Atmospheric Sciences*, 32, 3 – 15, [https://doi.org/10.1175/1520-0469\(1975\)032<0003:TEODTC>2.0.CO;2](https://doi.org/10.1175/1520-0469(1975)032<0003:TEODTC>2.0.CO;2), 1975.

Myhre, G., Shindell, D., Bréon, F.-M., Collins, W., Fuglestad, J., Huang, J., Koch, D., Lamarque, J.-F., Lee, D., Mendoza, B., Nakajima,
695 T., Robock, A., Stephens, G., Takemura, T., and H. Zhang, .: Anthropogenic and Natural Radiative Forcing. In: *Climate Change 2013:
The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate*

- Change [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 659–740,, <https://doi.org/https://doi:10.1017/CBO9781107415324.018>, 2013.
- 700 Nam, C., Kühne, P., Salzmann, M., and Quaas, J.: A prospectus for constraining rapid adjustments in general circulation models, *J. Adv. Model. Earth Syst.*, 10, 2080–2094, <https://doi.org/10.1029/2017MS001153>, 2018.
- Quaas, J., Andrews, T., Bellouin, N., Block, K., Boucher, O., Ceppi, P., Dagan, G., Doktorowski, S., Eichholz, H. M., Forster, P., Goren, T., Gryspeerdt, E., Hodnebrog, , Jia, H., Kramer, R., Lange, C., Maycock, A. C., Mülmenstädt, J., Myhre, G., O'Connor, F. M., Pincus, R., Samset, B. H., Senf, F., Shine, K. P., Smith, C., Stjern, C. W., Takemura, T., Toll, V., and Wall, C. J.: Ad-
705 justments to Climate Perturbations—Mechanisms, Implications, Observational Constraints, *AGU Advances*, 5, e2023AV001144, <https://doi.org/https://doi.org/10.1029/2023AV001144>, e2023AV001144 2023AV001144, 2024.
- Rugenstein, M. A. A., Gregory, J. M., Schaller, N., Sedláček, J., and Knutti, R.: Multiannual Ocean–Atmosphere Adjustments to Radiative Forcing, *Journal of Climate*, 29, 5643 – 5659, <https://doi.org/10.1175/JCLI-D-16-0312.1>, 2016.
- Russotto, R. D. and Ackerman, T. P.: Changes in clouds and thermodynamics under solar geoengineering and implications for required solar
710 reduction, *Atmospheric Chemistry and Physics*, 18, 11 905–11 925, <https://doi.org/10.5194/acp-18-11905-2018>, 2018.
- Salvi, P., Ceppi, P., and Gregory, J. M.: Interpreting the Dependence of Cloud-Radiative Adjustment on Forcing Agent, *Geophysical Research Letters*, 48, e2021GL093 616, <https://doi.org/https://doi.org/10.1029/2021GL093616>, e2021GL093616 2021GL093616, 2021.
- Schmidt, H., Alterskjær, K., Bou Karam, D., Boucher, O., Jones, A., Kristjánsson, J. E., Niemeier, U., Schulz, M., Aaheim, A., Benduhn, F., Lawrence, M., and Timmreck, C.: Solar irradiance reduction to counteract radiative forcing from a quadrupling of CO₂: climate responses
715 simulated by four earth system models, *Earth System Dynamics*, 3, 63–78, <https://doi.org/10.5194/esd-3-63-2012>, 2012.
- Sherwood, S. C., Bony, S., Boucher, O., Bretherton, C., Forster, P. M., Gregory, J. M., and Stevens, B.: Adjustments in the Forcing-Feedback Framework for Understanding Climate Change, *Bulletin of the American Meteorological Society*, 96, 217 – 228, <https://doi.org/10.1175/BAMS-D-13-00167.1>, 2015.
- Sippel, S., Meinshausen, N., Székely, E., Fischer, E., Pendergrass, A., Lehner, F., and Knutti, R.: Robust detection of forced warming in the
720 presence of potentially large climate variability, *Science Advances*, 7, <https://doi.org/10.1126/sciadv.abh4429>, 2021.
- Smith, C. J., Kramer, R. J., Myhre, G., Forster, P. M., Soden, B. J., Andrews, T., Boucher, O., Faluvegi, G., Fläschner, D., Hodnebrog, O., Kasoar, M., Kharin, V., Kirkevåg, A., Lamarque, J.-F., Mülmenstädt, J., Olivié, D., Richardson, T., Samset, B. H., Shindell, D., Stier, P., Takemura, T., Voulgarakis, A., and Watson-Parris, D.: Understanding Rapid Adjustments to Diverse Forcing Agents, *Geophysical Research Letters*, 45, 12,023–12,031, <https://doi.org/https://doi.org/10.1029/2018GL079826>, 2018.
- 725 Stjern, C. W., Forster, P. M., Jia, H., Jouan, C., Kasoar, M. R., Myhre, G., Olivié, D., Quaas, J., Samset, B. H., Sand, M., Takemura, T., Voulgarakis, A., and Wells, C. D.: The Time Scales of Climate Responses to Carbon Dioxide and Aerosols, *Journal of Climate*, 36, 3537 – 3551, <https://doi.org/10.1175/JCLI-D-22-0513.1>, 2023.
- Virgin, J. G. and Fletcher, C. G.: On the Linearity of External Forcing Response in Solar Geoengineering Experiments, *Geophysical Research Letters*, 49, e2022GL100 200, <https://doi.org/https://doi.org/10.1029/2022GL100200>, e2022GL100200 2022GL100200, 2022.
- 730 Webb, M. J., Andrews, T., Bodas-Salcedo, A., Bony, S., Bretherton, C. S., Chadwick, R., Chepfer, H., Douville, H., Good, P., Kay, J. E., Klein, S. A., Marchand, R., Medeiros, B., Siebesma, A. P., Skinner, C. B., Stevens, B., Tselioudis, G., Tsushima, Y., and Watanabe, M.: The Cloud Feedback Model Intercomparison Project (CFMIP) contribution to CMIP6, *Geoscientific Model Development*, 10, 359–384, <https://doi.org/10.5194/gmd-10-359-2017>, 2017.

- Yukimoto, S., Koshiro, T., Kawai, H., Oshima, N., Yoshida, K., Urakawa, S., Tsujino, H., Deushi, M., Tanaka, T., Hosaka, M., Yoshimura, H.,
735 Shindo, E., Mizuta, R., Ishii, M., Obata, A., and Adachi, Y.: MRI MRI-ESM2.0 model output prepared for CMIP6 CFMIP abrupt-solm4p,
<https://doi.org/10.22033/ESGF/CMIP6.6756>, 2020.
- Zelinka, M. D., Klein, S. A., Taylor, K. E., Andrews, T., Webb, M. J., Gregory, J. M., and Forster, P. M.: Contributions of Different Cloud
Types to Feedbacks and Rapid Adjustments in CMIP5, *Journal of Climate*, 26, 5007 – 5027, <https://doi.org/10.1175/JCLI-D-12-00555.1>,
2013.