



Triple oxygen isotope composition of CO₂ in the upper troposphere and stratosphere

Getachew Agmuas Adnew¹, Gerbrand Koren², Neha Mehendale¹, Sergey Gromov³, Maarten Krol^{1,4}, and Thomas Röckmann¹

¹Institute for Marine and Atmospheric research Utrecht (IMAU), Physics Department, Science Faculty, Utrecht University, Princetonplein 5, 3584 CC Utrecht, the Netherlands

²Copernicus Institute of Sustainable Development, Utrecht University, Princetonlaan 8a, 3584 CB Utrecht, the Netherlands

³Max Planck Institute for Chemistry, Hahn-Meitner-Weg 1, 55218 Mainz, Germany

⁴Meteorology and Air Quality, Wageningen University, Wageningen, the Netherlands

Correspondence: Getachew Agmuas Adnew (g.a.adnew@uu.nl)

Abstract. High precision measurements of the triple oxygen isotope composition of CO₂ ($\Delta^{17}\text{O}$) can be used to estimate biosphere-atmosphere exchange of CO₂, the residence time of tropospheric CO₂ and stratosphere-troposphere exchange. In this study, we report measurements of the $\Delta^{17}\text{O}(\text{CO}_2)$ from air samples collected during two aircraft based programs, CARIBIC and StratoClim. CARIBIC (Civil Aircraft for the Regular Investigation of the atmosphere based on an Instrument Container) provided air samples from numerous transcontinental flights in the upper troposphere/lower stratosphere region. StratoClim (Stratospheric and upper tropospheric processes for better climate predictions) conducted intensive campaigns with the high-altitude aircraft M55 Geophysica during the Asian Summer Monsoon Anticyclone (ASMA), providing air samples from altitudes up to 21 km.

Using high precision $\Delta^{17}\text{O}$ measurements of the CARIBIC samples, we show that the N₂O- $\Delta^{17}\text{O}$ correlation, previously observed in the stratosphere, extends to the upper troposphere. Moreover, we found no significant spatial or hemispheric differences in $\Delta^{17}\text{O}(\text{CO}_2)$ for the upper tropospheric samples collected during the CARIBIC program. However, in many of the StratoClim samples, with significant stratospheric contributions, we observed a much lower N₂O- $\Delta^{17}\text{O}$ slope compared to CARIBIC samples and previous publications. This deviation is attributed to change in eddy diffusion above the tropopause within the ASMA, confirming previously published model calculations. These samples provide the first experimental evidence that differences in vertical mixing/transport can lead to significantly different N₂O- $\Delta^{17}\text{O}$ slopes. High precision $\Delta^{17}\text{O}$ measurements can identify ejections of tropospheric air into the stratosphere based on the slope of the N₂O- $\Delta^{17}\text{O}$ correlation, as both tracers have chemical lifetimes longer than their transport times.

Furthermore, we use the $\Delta^{17}\text{O}$ measurements from the lower stratosphere and the upper troposphere to estimate global stratospheric production and surface removal of the isotope tracer $\Delta^{17}\text{O}$. The removal estimate is then used to derive an independent estimate of global vegetation exchange of CO₂, confirming earlier estimates based on surface level $\Delta^{17}\text{O}$ measurements.



1 Introduction

Measurements of the isotope composition of CO₂ ($\delta^{13}\text{C}$ and $\delta^{18}\text{O}$, see definition below) are used in many ways to understand sources and sinks of CO₂ in the global carbon cycle (Farquhar et al., 1993; Welp et al., 2011; Cuntz et al., 2003b; Ciais et al., 1997a, 1995). The flux estimates of CO₂ from the terrestrial biosphere are poorly constrained in current carbon cycle models (Piao et al., 2013). By using $\delta^{13}\text{C}$ it is possible to differentiate between the ocean sink and biosphere activity. However, distinguishing respiration, photosynthesis and combustion signals using $\delta^{13}\text{C}$ of CO₂ is impossible (Ciais et al., 1995). To address this challenge, $\delta^{18}\text{O}$ of CO₂ is a valuable tool for distinguishing respiration from photosynthesis. The $\delta^{18}\text{O}$ value of CO₂ is higher during photosynthesis than during respiration due to isotope exchange of CO₂ with isotopically different leaf water and soil water pools, as leaf water typically has a higher $\delta^{18}\text{O}$ value than soil water (Ciais et al., 1997a, b; Welp, 2011; Farquhar et al., 1993; Cuntz et al., 2003a, b; Francey and Tans, 1987; Yakir, 2020).

One of the limitations of using $\delta^{18}\text{O}$ of CO₂ to distinguish photosynthesis and respiration is the oxygen isotope exchange of CO₂ with liquid water in surface water bodies or vegetation, since this isotope exchange affects $\delta^{18}\text{O}$ of CO₂, without contributing to a net flux. Numerous equilibrium and kinetic effects can alter the $\delta^{18}\text{O}$ value of water and other molecules (Cuntz et al., 2003a, b; Peylin et al., 1999). Importantly, the isotopic composition of leaf water at the CO₂-H₂O exchange site in the mesophyll is not well understood, primarily due to fractionation associated with evaporation, transport, and diffusion (Adnew et al., 2020, 2021, 2023; Gan et al., 2002; Cousins et al., 2006; Song et al., 2015; Landais et al., 2006; Cernusak et al., 2016; Helliker and Ehleringer, 2000). These unknown physico-chemical fractionation processes introduce large uncertainty into estimates of gross fluxes of CO₂ when using $\delta^{18}\text{O}$ as quantitative tracer. This uncertainty can potentially be reduced using the triple oxygen isotopic composition of CO₂, $\Delta^{17}\text{O}$ (see equation 4), which is a combination of the $\delta^{17}\text{O}$ and $\delta^{18}\text{O}$ isotopic composition (Hoag et al., 2005; Koren et al., 2019; Adnew et al., 2020).

Stratospheric CO₂ has a high $\Delta^{17}\text{O}$ value, i.e., $\Delta^{17}\text{O} \gg 0$, due to isotope exchange with O(¹D) produced from O₃ photolysis (Thiemens et al., 1991, 1995b; Lyons, 2001; Lämmerzahl et al., 2002; Thiemens, 2006; Kawagucci et al., 2008; Wiegel et al., 2013; Yung et al., 1991b, 1997b; Shaheen et al., 2007). The main sink for this higher $\Delta^{17}\text{O}$ signal of CO₂ is isotope exchange with leaf, soil and ocean water at the Earth's surface, after $\Delta^{17}\text{O}$ enriched CO₂ has re-entered the troposphere via the large-scale Brewer-Dobson circulation and synoptic eddy diffusion (Boering et al., 2004; Thiemens et al., 2014; Liang and Mahata, 2015; Francey and Tans, 1987). With the exception of stratospheric CO₂, $\Delta^{17}\text{O}$ variations in nature are much smaller compared to $\delta^{18}\text{O}$ variations, and are better defined, as conventional biogeochemical processes follow a well-defined three-isotope fractionation slope (Barkan and Luz, 2005, 2007, 2012; Landais et al., 2006; Angert et al., 2004, 2003). Furthermore, the triple oxygen isotope fractionation slopes for specific processes are independent of the source water isotope composition, insensitive to temperature, and process-specific (Landais et al., 2006; Hofmann et al., 2012). As a result, $\Delta^{17}\text{O}$ is less affected by the numerous physico-chemical fractionation processes mentioned above, and may provide an additional constraint for



quantifying the gross fluxes of the terrestrial carbon cycle than measuring $\delta^{18}\text{O}$ alone (Koren et al., 2019; Hoag et al., 2005; 55 Liang et al., 2017a, 2023; Hofmann et al., 2017; Adnew et al., 2020; Thiemens et al., 2014).

In addition to quantifying the gross fluxes of the terrestrial carbon cycle, $\Delta'^{17}\text{O}$ can provide useful information for concern- ing stratospheric intrusions (Liang and Mahata, 2015; Steur et al., 2024), stratosphere-troposphere exchange, (Boering et al., 2004; Luz et al., 1999), atmospheric transport and chemistry in the mesosphere and stratosphere (Liang et al., 2007, 2008), combustion processes (Laskar et al., 2016; Horváth et al., 2012) and for estimating the residence time of CO_2 in the troposphere 60 (Liang et al., 2017b; Laskar et al., 2019; Hoag et al., 2005).

High precision measurements of $\Delta'^{17}\text{O}(\text{CO}_2)$ are particularly interesting in the tropical upper troposphere/lower most strato- sphere region, which is remote from the sources and sinks. Therefore, $\Delta'^{17}\text{O}(\text{CO}_2)$ can be used to study the influence of the stratosphere-troposphere exchange on the variations and dynamics of $\Delta'^{17}\text{O}$ and its correlation with other long- and short-lived trace gases. Additionally, these measurements may be valuable for using $\Delta'^{17}\text{O}$ as a tracer to quantify gross fluxes of CO_2 , as 65 this region links the stratosphere, where $\Delta'^{17}\text{O}$ of CO_2 is produced, to the troposphere, where $\Delta'^{17}\text{O}$ of CO_2 is "washed out". Furthermore, it is interesting to investigate whether $\Delta'^{17}\text{O}$ varies spatially in the upper troposphere and whether it is possible to detect large scale dynamic phenomena that happen in the Asian Summer Monsoon Anticyclone (ASMA). The Monsoon circulation system has a large variability in its spatial extent and the ASMA can reach the Mediterranean, North-east Africa and East Asia (Annamalai and Slingo, 2001; Garny and Randel, 2013; Pan et al., 2016; Vogel et al., 2019).

70 Here we present high-precision measurements of $\Delta'^{17}\text{O}(\text{CO}_2)$ from a total of 85 air samples collected in the upper tro- posphere and stratosphere as part of the CARIBIC and StratoClim aircraft air sampling projects. We investigate the spatial distribution (horizontal and vertical) of $\Delta'^{17}\text{O}(\text{CO}_2)$, signals associated with the ASMA, and refine previous estimates of the net stratosphere-troposphere flux of $\Delta'^{17}\text{O}(\text{CO}_2)$ and constraints on surface CO_2 emissions.

2 Materials and Method

75 2.1 Definitions and notation

Variations in isotopic abundance are reported as deviations of a heavy-to-light isotope ratio in a sample relative to a reference material (Eq. 1). For oxygen the reference material is Vienna Standard Mean Ocean Water (VSMOW) whereas for carbon the reference material is Vienna Pee Dee Belemnite (VPDB). Since isotope variations are small, they are usually reported in per mill (‰) in delta notation.

$$80 \quad \delta^x = \frac{xR_{sam}}{xR_{std}} - 1 \quad (1)$$

where x can be 13, 17 and 18 (for ^{13}C , ^{17}O and ^{18}O , respectively). The indices *sam* and *std*, stand for sample and standard, respectively and R is the ratio between the heavy isotope and the light isotope of the respective element, for instance for ^{13}C and ^{18}O , $^{13}R = \frac{^{13}\text{C}}{^{12}\text{C}}$ and $^{18}R = \frac{^{18}\text{O}}{^{16}\text{O}}$, respectively. The variations in $\delta^{17}\text{O}$ and $\delta^{18}\text{O}$ are closely related during most physico-chemical fractionation processes according to Eq.2 (Matsuhisa et al., 1978; Young et al., 2002).



$$85 \quad \left(\frac{{}^{18}R_{sam}}{{}^{18}R_{std}} \right)^{\theta_p} = \frac{{}^{17}R_{sam}}{{}^{17}R_{std}} \quad (2)$$

The exponent θ_p denotes a three isotope slope that occurs from a single process. Equation 2 can be expressed in δ - notation as:

$$\ln(\delta^{17}O + 1) - \theta_p \times \ln(\delta^{18}O + 1) = 0 \quad (3)$$

The deviation of the left hand side of equation 3 from zero is defined as $\Delta'^{17}O$ (Eq. 4).

$$90 \quad \Delta'^{17}O = \ln(\delta^{17}O + 1) - \lambda_{ref} \times \ln(\delta^{18}O + 1) \quad (4)$$

where the process-dependent exponent θ_p has been replaced by a three-isotope reference slope λ_{ref} . Note that the choice of λ_{ref} is arbitrary since in nature isotopic compositions rarely reflect fractionation from a single process, but instead integrate multiple fractionating processes and several three-isotope values (Adnew, 2020; Barkan and Luz, 2005, 2007, 2012; Angert et al., 2004). For most natural processes λ_{ref} ranges from 0.5 to 0.5305 (Thiemens, 2006; Matsuhisa et al., 1978; Young et al., 2002; Cao and Liu, 2011; Thiemens et al., 1991; Kaiser, 2008) with some exceptions (Adnew et al., 2022; Hayles et al., 2017; Hayles and Killingsworth, 2022). In this study we use a λ_{ref} value of 0.528, the value associated with meteoric water (Meijer and Li, 1998; Luz and Barkan, 2010).

2.2 Calculation of net isotope flux from stratosphere to troposphere

As described in detail by Plumb and Ko (1992) and Plumb (2007), gases that are chemically long-lived relative to vertical and quasi-horizontal transport time scales exhibit compact correlations in the stratosphere, and the slope of the observed correlation between the two tracers is equal to the ratio of their net vertical fluxes. Following this approach, Luz et al. (1999) and Boering et al. (2004) determined the global annual mean net isotope flux (NIF) from the stratosphere (ST) to the troposphere (T) for $\Delta'^{17}O$ of CO_2 using the $\Delta'^{17}O(CO_2)$ - N_2O correlation as described in detail by Garofalo et al. (2019).

$$\Delta'^{17}O(CO_2) - NIF = MF \times [CO_2]_{ST} \times \left[m \times \frac{-L_{N_2O}}{MF} + \Delta'^{17}O(CO_2)_T \right] - MF \times [CO_2]_T \times \Delta'^{17}O(CO_2)_T \quad (5)$$

The terms MF , L_{N_2O} , and m stand for the total air mass flux from the stratosphere to the troposphere, the global N_2O loss rate and the correlation slope between $\Delta'^{17}O$, respectively. In our analysis, this correlation slope is determined using a Williamson-York bivariate fit, accounting for uncertainties in both the $\Delta'^{17}O(CO_2)$ and N_2O data (Mikkonen et al., 2019). There is a significant variation in estimates for the air mass flux from the stratosphere to the troposphere. For instance, the estimates by Holton et al. (1995) and Appenzeller et al. (1996) differ by a factor of three in their calculated cross-tropopause air mass fluxes, but note that the troposphere definitions are also different. As described in Garofalo et al. (2019), the net isotope flux is not strongly sensitive to the actual air mass fluxes. Since the mole fraction of CO_2 in the lower stratosphere and troposphere is the same within about 1 %, equation 5 can be simplified to:



$$\Delta'^{17}\text{O}(\text{CO}_2) - \text{NIF} = -m \times L_{\text{N}_2\text{O}} \times [\text{CO}_2] \quad (6)$$

The uncertainty in the estimated net annual mean flux of $\Delta'^{17}\text{O}(\text{CO}_2)$ depends on the uncertainty in depends on the uncertainty
 115 of m and $L_{\text{N}_2\text{O}}$. In our calculation, we used 370 ppm for $[\text{CO}_2]$, the average value for CARIBIC samples, most of them collected
 in the year 2001.

2.3 Estimating gross surface flux using $\Delta'^{17}\text{O}(\text{CO}_2)$

The CARIBIC samples analysed in this study cover latitudes from 30° south to 54° north (see Figure 1), we used these samples
 to estimate surface emissions using a mass balance box model calculation. The box model used in this study is not suitable to
 120 simulate the spatial and temporal variability of $\Delta'^{17}\text{O}(\text{CO}_2)$ (Koren et al., 2019). Our box model is an extended version of those
 used by Hoag et al. (2005) and Liang et al. (2017b), incorporating an intermediate upper troposphere (*UT*) box. Additionally,
 our study includes $\Delta'^{17}\text{O}$ measurements from both hemispheres and different seasons. The $\Delta'^{17}\text{O}(\text{CO}_2)$ value in the *UT*
 depends on the $\Delta'^{17}\text{O}(\text{CO}_2)$ value of stratosphere (*ST*) and surface processes (assimilation/photosynthesis (*A*), respiration (*R*),
 soil invasion (*SI*), ocean (*o*) and anthropogenic emission (*anth*)) and the corresponding fluxes (F). At steady state, the mass
 125 balance equation for $\Delta'^{17}\text{O}(\text{CO}_2)$ in the upper troposphere is:

$$F_A \times \Delta_A \Delta'^{17}\text{O} + F_{anth} \times \Delta'^{17}\text{O}_{anth} + F_{ST} \times (\Delta'^{17}\text{O}_{ST} - \Delta'^{17}\text{O}_{UT}) + F_o \times (\Delta'^{17}\text{O}_o - \Delta'^{17}\text{O}_{UT}) + \\ F_R \times (\Delta'^{17}\text{O}_R - \Delta'^{17}\text{O}_{UT}) + F_{SI} \times (\Delta'^{17}\text{O}_{SI} - \Delta'^{17}\text{O}_{UT}) = 0 \quad (7)$$

where $F_A = F_{al} - F_{la} = 0.88 \times GPP$ (Ciais et al., 1997a), with *al*, *la*, and *GPP* being atmosphere-to-leaf flux, leaf-to-
 atmosphere flux, and gross primary production, respectively. $\Delta_A \Delta'^{17}\text{O}$ is the discrimination against $\Delta'^{17}\text{O}$ during assimilation,
 and calculated as $\Delta_A \Delta'^{17}\text{O} = (\Delta'^{17}\text{O}_{UT} - \Delta'^{17}\text{O}_M) \times (-0.150 \times e^{3.707 \times \frac{c_m}{c_a}} + 0.028)$ as described by Adnew et al.
 130 (2020) (see the supplementary material for more detail). This parameterization was derived from leaf cuvette studies using
 both C_3 and C_4 plants under different light conditions. c_m and c_a represent the CO_2 mole fraction in the mesophyll and the
 atmosphere, respectively. $\Delta'^{17}\text{O}_M$ is the $\Delta'^{17}\text{O}$ of CO_2 in the mesophyll, calculated from the $\Delta'^{17}\text{O}$ value of meteoric water
 (MW) (Landais et al., 2007; Barkan and Luz, 2012; Bottinga and Craig, 1968; Brenninkmeijer et al., 1983), see also the
 supplementary material.

135 The estimates of soil invasion fluxes are highly uncertain, the reported values in the literature vary from < 10 PgC/yr (Stern
 et al., 2001) to 450 PgC/yr (Wingate et al., 2009). For this study, we assume that F_{SI} and F_R are equal, and $F_A = 0.88 \times GPP =$
 $0.88 \times (NEP - F_R)$ (Liang et al., 2017b). Substituting this term in equation 7:

$$F_A \times \Delta_A \Delta'^{17}\text{O} + F_{anth} \times \Delta'^{17}\text{O}_{anth} + F_{ST} \times (\Delta'^{17}\text{O}_{ST} - \Delta'^{17}\text{O}_{UT}) + F_o \times (\Delta'^{17}\text{O}_o - \Delta'^{17}\text{O}_{UT}) + \\ \left(\frac{F_A}{0.88} - NEP \right) \times (\Delta'^{17}\text{O}_R - \Delta'^{17}\text{O}_{UT}) + \left(\frac{F_A}{0.88} - NEP \right) \times (\Delta'^{17}\text{O}_{SI} - \Delta'^{17}\text{O}_{UT}) = 0 \quad (8)$$



Table 1. Summary of parameters used for the box model mass balance calculation in Equation 8. T and RH represents temperature in Kelvin and relative humidity near the Earth's surface, respectively.

Parameter	value or description	ref
α_{trans}^{18}	$1.002644 - \frac{3.206}{T} + \frac{1534}{T^2}$	Bottinga and Craig (1968)
$\alpha_{CO_2-H_2O}^{18}$	$\frac{17.604}{T} + 0.98207$	Brenninkmeijer et al. (1983)
$\alpha_{diff-water}^{18}$	1.0016 ‰	Farquhar and Lloyd (1993)
$\alpha_{diff-soil}^{18}$	1.0072 ‰	Miller et al. (1999)
θ_{trans}	$0.522 - 0.008 \times RH$	Landais et al. (2007)
$\theta_{CO_2-H_2O}$	0.5229	Barkan and Luz (2012)
θ_{diff-H_2O}	0.50	calculated
$\theta_{diff-air}$	0.509	Young et al. (2002)
GPP	Gross primary production	calculated
F_A	$0.88 \times GPP$	Ciais et al. (1997b)
F_{al}	$\frac{c_a}{c_a - c_m} \times F_A$	Farquhar et al. (1993)
F_{la}	$\frac{c_m}{c_a - c_m} \times F_A$	Farquhar et al. (1993)
$\frac{c_m}{c_a}$	0.65	Farquhar et al. (1993)
M_a	807 ± 6 PgC	Stocker et al. (2013)
F_o	80 ± 6 PgC/yr	Stocker et al. (2013)
F_{anth}	9 ± 0.8 PgC/yr	Stocker et al. (2013)
NEP	10 PgC/yr	Saugier et al. (2001)
$\Delta^{17}O_{MW}$	0.033 ± 0.005 ‰	Barkan and Luz (2012)
$\theta_{deg-equ}$	1.0	assumed
$\Delta^{17}O_{OW}$	-0.005 ± 0.001 ‰	Barkan and Luz (2012)

The $\Delta^{17}O_{anth}$ value is -0.446 ± 0.077 ‰ (Horváth et al., 2012; Laskar et al., 2016). The $\Delta^{17}O_o$ value is calculated from the $\Delta^{17}O$ value of ocean water (OW) (Table 1). Similarly, the $\Delta^{17}O$ values for soil invasion and respiration are derived from the $\Delta^{17}O$ value of meteoric water (Liang et al., 2023; Koren et al., 2019; Hofmann et al., 2017). For the calculations, we used a surface temperature of 15°C, and a relative humidity of 75 % (Dai, 2006). The sensitivity of $\Delta_A \Delta^{17}O$ to temperature and relative humidity is shown in Figure S1 of the supplementary material. The value of $\Delta_A \Delta^{17}O$ increases with an increase in relative humidity but decreases with increasing temperature. Using the parameters described above and provided in Table 1 and Equation 8, we estimated the gross primary production (GPP), surface flux (including land and ocean) and the oxygen isotope residence time (turnover time) of CO₂ in the atmosphere. The oxygen isotope residence time of CO₂ in the atmosphere is defined by the ratio of the atmospheric CO₂ mass loading (M_a) and the CO₂ surface flux (Welp et al., 2011). The error for all estimated values is determined using a Monte Carlo simulation using 10⁶ runs, where values of the input parameter are randomly picked from an input value distribution of these parameters defined by their uncertainty.



150 The sensitivity of GPP to relative humidity, temperature, the $\Delta^{17}\text{O}$ value of the upper troposphere, and the net flux of
 $\Delta^{17}\text{O}$ from the stratosphere to the troposphere is shown in Figures S2 and S3 of the supplementary material. A higher net
flux of $\Delta^{17}\text{O}$ to the troposphere leads to an increase in GPP to fulfill the isotope balance of equation 8. An increase in relative
humidity leads to a decrease in GPP due to an increase in $\Delta_A \Delta^{17}\text{O}$. Conversely, an increase in temperature results in an
increase in GPP, as higher temperatures cause a decrease in $\Delta_A \Delta^{17}\text{O}$, as described above. The parameters used in the mass
155 balance calculations and their errors are provided in Table 1.

2.4 De-trending and classification of samples into upper troposphere and stratosphere

For comparison with previously published data and between the CARIBIC and StratoClim samples measured in this study, all
 N_2O measurements were trend-corrected to the year 2001 following a similar approach described in Koren et al. (2019).

$$N_2O_{det} = N_2O_{obs} \times \left[1 - \frac{N_2O_{growthrate}}{N_2O_{ref}} \times (t_{ref} - t_{obs}) \right]^{-1} \quad (9)$$

160 Here, N_2O_{obs} , N_2O_{det} , N_2O_{ref} and $N_2O_{growthrate}$ refer to the observed and detrended mole fractions of N_2O , the mole
fraction of N_2O at the reference time and the growth rate of N_2O in the troposphere, respectively. The variables t_{obs} and t_{ref}
represent the time of observation and the reference time (1 July 2001). The growth rate of N_2O in the troposphere, used in this
study, is 0.75 ppb/year (Stocker et al., 2013). The mole fraction of N_2O at the reference time (1 July 2001) is 316.24 ppb (Lan
et al., 2024).

165 The mole fraction of CH_4 in the StratoClim data was detrended to 2007 using an average growth rate of 12 ppb/year (Nisbet
et al., 2019). The 2007 mole fraction of CH_4 is assumed to be the same as the CH_4 mole fraction in 2001, since the atmospheric
 CH_4 mole fraction was stable between 2000 and 2007 (Nisbet et al., 2019).

We used the N_2O -CO correlation to classify air samples into upper troposphere and stratospheric samples as described in
detail by Assonov et al. (2013). The CARIBIC samples are mostly from the upper troposphere (UT) and include a few from the
170 lower stratosphere (see Figure S4 in the supplementary material). Based on the mole fraction of N_2O , we grouped the $\Delta^{17}\text{O}$
values into two categories: upper troposphere (UT) samples ($\text{N}_2\text{O} \geq 313.5$ ppb) and the lower stratosphere (LS) samples
($\text{N}_2\text{O} < 313.5$ ppb). We further divided the lower stratosphere into two subgroups: $313.5 > \text{N}_2\text{O} > 306$ ppb and $\text{N}_2\text{O} < 306$
ppb. Furthermore, we used a zonal average tropopause pressure simulated using the TM5 model as outlined in Krol et al.
(2018), and the *age of air* tracer, to support the classification of samples into upper tropospheric and stratospheric. Most of the
175 samples classified as upper tropospheric based on the mole fraction of N_2O were collected far below the tropopause whereas
the samples classified as stratospheric were collected above or close to the tropopause with a few exceptions (see Figure S5 in
the supplementary material).

2.5 Sampling

Air samples were collected from the upper troposphere and stratosphere during two different international projects, CARIBIC
180 and StratoClim, which are shortly described below. Figure 1 shows the geographical coordinates of the sampling locations for
the samples measured in this study.

2.5.1 CARIBIC samples

In the CARIBIC project (Civil Aircraft for the Regular Investigation of the atmosphere Based on an Instrument Container, CARIBIC, <https://www.caribic-atmospheric.com/>), samples were collected using a Boeing 767 aircraft (LTU, Germany). The flights operated at typical commercial cruising altitude between 9 and 12 km, i.e., which is the upper troposphere-lower stratosphere region at higher latitudes and the upper troposphere region in the mid latitudes and tropics (Assonov et al., 2010). The details of the CARIBIC instrument container are described in Brenninkmeijer et al. (1999). The payload of the flights before 2003 included large stainless steel canisters for collecting whole air samples (WAS) (sample size \approx 340 L STP) at several locations along the flight path. On a single flight 12 discrete samples were collected. Each sample collection took about 20 min which corresponds to a horizontal distance of 250 km (Brenninkmeijer et al., 1999). The samples collected are relatively dry since the ambient temperature was always below -30 °C and the relative humidity was about 0.44 % at 20 °C (Assonov et al., 2009b). The air samples were processed soon after they returned to the Division of Atmospheric Chemistry, Max Planck Institute for Chemistry, Mainz, Germany (Pupek et al., 2005). The processing included the extraction of CO₂ and CO for isotope analysis, including radiocarbon (Brenninkmeijer, 1993; Assonov et al., 2009b) as well as measurement of several other trace gases including N₂O, CH₄ and SF₆ (Brenninkmeijer et al., 1999). The CARIBIC container was also equipped with an automated in-situ analyzer for O₃ (UV absorption), CO (gas chromatography (GC)) and other parameters (Brenninkmeijer et al., 1999). In this study we measured the $\Delta^{17}\text{O}$ value of 50 CARIBIC samples.

2.5.2 StratoClim samples

35 additional air samples were collected on the high-altitude M55 Geophysica aircraft during two campaigns of the StratoClim project (<https://www.stratoclim.org/>) (Stefanutti et al., 1999; Cairo et al., 2010) in 2016 and 2017. In 2016, three flights were conducted over the Mediterranean region from Kalamata, Greece ($37^{\circ}2'N$ and $22^{\circ}7'E$) between August 30 and 6 September and eight flights were conducted in 2017 over the Indian subcontinent from Kathmandu, Nepal ($27^{\circ}46'N$ and $85^{\circ}16'E$) between 27 July and 10 August. The whole air sampler of the Institute for Marine and Atmospheric research Utrecht (IMAU), Utrecht University was used to compress air into 2 L pre-evacuated stainless-steel canisters (Kaiser et al., 2006). We analysed 16 samples from the 2016 flights and 19 samples from the 2017 flights (Table S3). Further details on sample collection using the Geophysica aircraft can be found in Kaiser et al. (2006) and Stefanutti et al. (1999).

2.6 Extraction of CO₂

For the StratoClim samples, the CO₂ was extracted from the whole air samples using a cryogenic extraction system developed at Utrecht University (Adnew et al., 2020, 2023). The extraction system was made of electropolished stainless steel and has four traps. The first two traps remove moisture and condensable organics at dry ice temperature whereas the third and fourth traps were used to collect CO₂ at liquid nitrogen temperature. The extraction was performed at a flowrate of 55 mL/min.

For the CARIBIC samples, the CO₂ had previously been extracted at the Max Planck Institute for Chemistry, Mainz, as described in Assonov et al. (2009b) and Pupek et al. (2005). Two "Russian Doll" cryogenic traps (Brenninkmeijer and Röckmann,




Figure 1. Geographical location of the CARIBIC (black markers), StratoClim2016 (blue circles) and StratoClim2017 (blue stars) air samples measured in this study. Upper tropospheric CARIBIC samples are categorized based on the geographical location where they are collected. These categories include Southern Hemisphere samples, Northern Hemisphere samples, African samples (longitude between 7° and 45°), Asian samples (longitude > 45°), and American samples (longitude < 0°).

1996; Brenninkmeijer, 1991) immersed in liquid nitrogen trapped all condensable gases including CO₂, N₂O, H₂O and most
215 organics. After pumping out non-condensables, the traps were slowly heated and the evolving CO₂ and N₂O was trapped in a
U-trap cooled with liquid nitrogen while the H₂O remained trapped in the Russian Doll Traps. The collected CO₂ was further
dried using P₂O₅ and flame-sealed in clean vials made of borosilicate glass (Assonov et al., 2009b; Pupek et al., 2005).



2.7 Measurement of $\delta^{13}\text{C}$, $\delta^{18}\text{O}$ and $\Delta^{17}\text{O}$ of CO_2

The $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values presented below were measured using a Delta^{PLUS} V isotope ratio mass spectrometer (ThermoScientific, Germany) in dual inlet mode at Utrecht University. The interference of N_2O was corrected using: $\delta^{18}\text{O} = \delta^{18}\text{O}_{\text{Measured}} + \frac{[\text{N}_2\text{O}]}{[\text{CO}_2]} \left(\frac{347}{1000} \right)$ and $\delta^{13}\text{C} = \delta^{13}\text{C}_{\text{Measured}} + \frac{[\text{N}_2\text{O}]}{[\text{CO}_2]} \left(\frac{250}{1000} \right)$ (Friedli and Siegenthaler, 1988; Sirignano et al., 2004). Figure 2 shows a comparison of the the $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ of CO_2 of CARIBIC samples measured in this study with those previously measured at the Max-Planck-Institute for Chemistry and reported in Assonov et al. (2009b, a, 2010).




Figure 2. The $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ of CARIBIC samples and comparison with values published in Assonov et al. (2009b, a, 2010).

The $\delta^{18}\text{O}$ values measured in this study are systematically higher compared to the measurement by Assonov et al. (2010) by $0.096 \pm 0.008 \text{ ‰}$. For $\delta^{13}\text{C}$ values, our measurement is also higher by $0.089 \pm 0.011 \text{ ‰}$. The reported error is the 95 % confidence limit (standard error of the mean multiplied by Student's t-factor). The tight distribution of differences is of the order of the measurement precision, which suggests that the isotopic composition of the samples remained stable during long-term storage in borosilicate glass vials and that the mean offset is due to scale differences. These observed offsets are consistent with the scale uncertainty reported by Assonov et al. (2009a) and better than typical inter-laboratory uncertainties (Levin et al., 2007).



The $\Delta^{17}\text{O}$ of the CO_2 was determined using the $\text{CO}_2\text{-O}_2$ exchange method (Adnew et al., 2019, 2022, 2023; Barkan et al., 2015) at Utrecht University. Equal amounts of CO_2 sample and the laboratory reference O_2 (with known isotopic composition) were allowed to exchange isotopes for 2 hours in a quartz reactor at 750°C in the presence of platinum sponge at the bottom of the reactor. After the reaction, the mixture was passed through a liquid nitrogen-cooled trap to condense the CO_2 and the O_2 was collected in a separate trap on 3 pellets of 5\AA molecular sieve (1.6mm, Sigma-Aldrich, USA) at liquid nitrogen temperature. The O_2 was then transferred to the bellows of the dual inlet system of a Delta^{PLUS} V isotope ratio mass spectrometer (ThermoFisher Scientific, Germany) and measured for its isotope composition. The $\Delta^{17}\text{O}$ value of the original CO_2 was calculated from the change in the isotopic composition of the non-reacted and reacted O_2 , and knowledge of the precise steady-state $\text{O}_2\text{-CO}_2$ isotope fractionations (Adnew et al., 2019; Barkan et al., 2015). The precision of the CO_2 isotope measurements was 0.007‰ , 0.03‰ and 0.008‰ for $\delta^{13}\text{C}$, $\delta^{18}\text{O}$ and $\Delta^{17}\text{O}$ respectively (Adnew et al., 2019, 2020, 2023).

3 Results

Figure 3a shows $\Delta^{17}\text{O}(\text{CO}_2)$ as a function of latitude for the CARIBIC samples. The samples classified as upper tropospheric, $\Delta^{17}\text{O}(\text{CO}_2)$ do not show geographic variations between large regions referred to as America, Asia and Africa, respectively (see Figure 1) and no statistically significant difference in $\Delta^{17}\text{O}(\text{CO}_2)$ between the hemispheres (see Figure 3b). Table S2 in the supplementary material presents the mole fraction of CO_2 , O_3 , CO , N_2O and CH_4 , along with the isotopic composition of CO_2 ($\delta^{13}\text{C}$, $\delta^{18}\text{O}$ and $\Delta^{17}\text{O}$) for the analyzed CARIBIC samples. Similarly, the mole fraction of CO_2 and other trace gases (O_3 , CO , N_2O and CH_4), along with the isotopic composition of CO_2 for StratoClim samples is shown in Table S3 of the supplementary material.

Figure 4 shows the correlation of $\Delta^{17}\text{O}(\text{CH}_2)$ measurements with the mole fractions of CH_4 and N_2O . For both sets of samples (CARIBIC and StratoClim), N_2O and $\Delta^{17}\text{O}(\text{CO}_2)$ are clearly correlated with R^2 values of ≥ 0.9 as shown in Figure 4. Similar tight correlations between N_2O and $\Delta^{17}\text{O}(\text{CO}_2)$ were reported in the previous studies as shown in Figure 5. Interestingly, StratoClim samples plot in two distinct groups of $\text{N}_2\text{O}\text{-}\Delta^{17}\text{O}(\text{CO}_2)$ correlations. Most of the StratoClim samples with $\Delta^{17}\text{O} < 0.2\text{‰}$ have a similar $\text{N}_2\text{O}\text{-}\Delta^{17}\text{O}$ slope as the CARIBIC samples as shown in Figure 4a. However, unexpectedly the $\text{N}_2\text{O}\text{-}\Delta^{17}\text{O}$ correlation slope for most of StratoClim samples with $\Delta^{17}\text{O} > 0.2\text{‰}$ is much lower (-0.017‰ per ppb) (see Figure 4a).

As shown in Figure 4b, there is a strong inverse correlation between CH_4 and $\Delta^{17}\text{O}$. For the StratoClim samples, the correlation is higher compared to the CARIBIC samples, with R^2 values of ≥ 0.9 and 0.79 , respectively. For both CARIBIC and StratoClim samples with $\Delta^{17}\text{O} < -0.2\text{‰}$, no clear correlation between CH_4 and $\Delta^{17}\text{O}$ can be established (see Figure 4b).

Figure 5 shows a comparison of our CARIBIC and StratoClim samples measured in this study with previously published $\Delta^{17}\text{O}$ measurements of upper-tropospheric and stratospheric CO_2 (Thiemens et al., 1995a, c; Alexander et al., 2001; Lämmerzahl et al., 2002; Boering et al., 2004; Wiegel et al., 2013; Kawagucci et al., 2005; Yeung et al., 2009). Four different characteristics are shown: The three-isotope plot (Figure 5a), $\Delta^{17}\text{O}$ as a function of altitude (only for StratoClim samples)




Figure 3. a) $\Delta^{17}\text{O}(\text{CO}_2)$ values of CARIBIC samples as a function of latitude. The $\Delta^{17}\text{O}$ values are grouped into three categories based on the mole fraction of N_2O as described in the text. b) average $\Delta^{17}\text{O}$ value of all upper tropospheric CARIBIC air samples and larger geographical regions. The errors are the standard error of the mean multiplied by the student t-factor for the 95 % confidence limit. n , SH and NH stands for the number of samples, Southern Hemisphere and Northern Hemisphere, respectively.

(Figure 5b), the N_2O - $\Delta^{17}\text{O}$ correlation (Figure 5c) and the CH_4 - $\Delta^{17}\text{O}$ correlation (Figure 5d). In these overview plots of N_2O - $\Delta^{17}\text{O}$ and CH_4 - $\Delta^{17}\text{O}$ plots, the new CARIBIC and StratoClim measurements are almost indistinguishable from the measurements reported in the literature. However, when we zoom in and compare the correlation slopes, clear differences for some of the measurements become apparent such as shown in Figure 4a for the variation in the N_2O - $\Delta^{17}\text{O}$ correlation between CARIBIC and StratoClim samples.

Figure 6 shows the three-isotope plot for both CARIBIC and StratoClim samples. The three-isotope plot of upper-tropospheric samples shows a tight correlation for both CARIBIC and StratoClim samples with slopes of 0.540 ± 0.005 and 0.556 ± 0.012 , respectively (Figure 6). For the StratoClim samples with higher stratospheric influence the $\ln(\delta^{17}\text{O} + 1)$ vs $\ln(\delta^{18}\text{O} + 1)$ is poorly correlated. It is apparent in Figure 6a that the vertical offsets from the fit line defined by the upper tropospheric samples get larger for larger stratospheric age, which is shown as colour coding. Figure 6b shows the correlation between the age of the air and $\Delta^{17}\text{O}(\text{CO}_2)$. An increase in the age of the air correlates well with a progressive enrichment of $\Delta^{17}\text{O}(\text{CO}_2)$.

275 4 Discussion

When the CARIBIC and StratoClim samples presented here are separated into upper tropospheric and stratospheric based on N_2O levels, these two groups exhibit distinct characteristics. The upper tropospheric samples show a tight correlation on the three-isotope plot, whereas the stratospheric samples display significant variability without a clear correlation. Interestingly,




Figure 4. Correlation of $\Delta^{17}\text{O}(\text{CO}_2)$ with a) N_2O for StratoClim samples (StratoClim-2016 is Kathmandu (KT, blue stars) and StratoClim-2017 is Kalamata (KL, blue circles)) and CARIBIC samples (black circles) and b) CH_4 . The errors of the linear regression slopes are the 95 % confidence interval. The data points marked in red indicate StratoClim samples with a low N_2O - $\Delta^{17}\text{O}$ correlation slope.

the three-isotope slopes (from a linear fit to the data) for the upper tropospheric samples are higher than the expected canonical
 280 range for tropospheric samples ($\lambda > 0.5309$). This indicates that upper tropospheric air already has a stratospheric influence. The contribution of stratospheric air in the CARIBIC samples is evident from an excellent correlation between $\Delta^{17}\text{O}(\text{CO}_2)$ and both $^{14}\text{C}(\text{CO})$ and O_3 , as illustrated in Figure 7. Both $^{14}\text{C}(\text{CO})$ and O_3 are reliable tracers for stratospheric air (Brenninkmeijer et al., 1995). Similar correlations between $^{14}\text{C}(\text{CO})$ and $\Delta^{17}\text{O}(\text{CO}_2)$ have been previously reported for stratospheric CO_2 (Thiemens et al., 1995c; Brenninkmeijer et al., 1995).

285 The observed scatter in the three-isotope plot for the stratospheric samples is conceptually similar to the scatter reported in previous studies (Boering et al., 2004; Wiegel et al., 2013; Mrozek et al., 2017; Thiemens et al., 1995a). In these publications it was mostly assumed to be due to the low precision of the measurement method. However, in this study, with high precision measurements, a similar scatter in the three-isotope plot is observed, demonstrating that the scatter observed in the three-isotope plot for stratospheric samples is a real signal of mixing, transport and production of $\Delta^{17}\text{O}(\text{CO}_2)$ (i.e. exchange between $\text{O}(^1D)$
 290 and CO_2). The tight correlation with the age of air provides a clear clue to explain this lack of correlation. The longer CO_2 has been exposed to the stratospheric CO_2 - $\text{O}(^1D)$ exchange, the higher the $\Delta^{17}\text{O}(\text{CO}_2)$ signal is (Yung et al., 1997a; Gamo et al., 1995; Yung et al., 1991a). The fact that the different samples do not line up on a single three isotope correlation line shows that we have sampled stratospheric air originating from different upper tropospheric “entry values” before entering




Figure 5. Comparison of results from CARIBIC samples (black circles) and StratoClim samples (blue circles) to previous publications (open circles). a) three isotope plot, b) correlation of $\Delta^{17}\text{O}(\text{CO}_2)$ with N_2O . The data points marked in red indicate StratoClim samples with a low $\text{N}_2\text{O}-\Delta^{17}\text{O}$ correlation slope. c) correlation of $\Delta^{17}\text{O}(\text{CO}_2)$ with altitude and d) correlation of $\Delta^{17}\text{O}(\text{CO}_2)$ with CH_4 . Previous stratospheric samples are from (Thiemens et al., 1995a, c; Alexander et al., 2001; Lämmerzahl et al., 2002; Boering et al., 2004; Wiegel et al., 2013; Kawagucci et al., 2005; Yeung et al., 2009)

the stratosphere. These differences in entry values are likely driven by the seasonality of $\delta^{17}\text{O}(\text{CO}_2)$ and $\delta^{18}\text{O}(\text{CO}_2)$. Thus, different tropospheric air masses have entered the stratosphere from various points along the correlation line established by the upper tropospheric samples. Once in the stratosphere, photochemical isotope exchange with O^1D occurs, and the samples progressively acquire a higher $\Delta^{17}\text{O}(\text{CO}_2)$ signature as the sampled air resided longer in the stratosphere. This is clearly demonstrated by the strong correlation between $\Delta^{17}\text{O}(\text{CO}_2)$ and the age of air, and it was not clearly visible in previous studies because of the higher measurement uncertainty.




Figure 6. a) Three isotope plot for StratoClim (stars) and CARIBIC (open circles) samples. The linear-regression is a fit to the samples identified as upper tropospheric based on the N_2O mole fraction. The color bar indicates the age of the air mass for the StratoClim samples reported in Adcock et al. (2021). The stated errors for the linear regression parameters are the 95 % confidence intervals. b) Relationship between the age of the air and $\Delta^{17}\text{O}(\text{CO}_2)$ for StratoClim samples. The data points marked in red indicate samples with a low $\text{N}_2\text{O}-\Delta^{17}\text{O}$ correlation slope.

300 Interestingly, despite our high measurement precision, we did not observe significant geographic variations in $\Delta^{17}\text{O}(\text{CO}_2)$ of CARIBIC air samples classified as upper troposphere (see Figure 3b), which could potentially arise from varying stratospheric and tropospheric air mass influences in different geographical regions. The lack of geographic variability contrasts with the reported regional $\Delta^{17}\text{O}(\text{CO}_2)$ variability at the surface (Liang et al., 2017a, 2023). Apparently, the variability at the surface, caused by sources and sinks of CO_2 , including isotope exchange with surface water reservoirs and anthropogenic
 305 emissions, is no longer visible in the upper troposphere, where the reservoir seems to be relatively well mixed in terms of $\Delta^{17}\text{O}(\text{CO}_2)$. Air masses in the upper troposphere experience rapid horizontal mixing, which can explain the uniform stratospheric influence in the CARIBIC samples.

The slope of two tracers near the tropopause is a measure of their relative net fluxes between troposphere and stratosphere (Plumb and Ko, 1992; Plumb, 2007). The slope of $\text{N}_2\text{O}-\Delta^{17}\text{O}(\text{CO}_2)$ was first used by Luz et al. (1999) and Boering et al.
 310 (2004) to quantify the so-called isoflux of $\Delta^{17}\text{O}(\text{CO}_2)$ from the stratospheric into the troposphere. Previous measurements had a lower precision and could not define the slope close to the tropopause, but only for older stratospheric air. Our high precision measurements of $\Delta^{17}\text{O}(\text{CO}_2)$ allow to determine the $\text{N}_2\text{O}-\Delta^{17}\text{O}(\text{CO}_2)$ slope right down to the tropopause. Figure 8 shows




Figure 7. Correlation of $\Delta^{17}\text{O}$ with stratospheric air tracers ^{14}CO and O_3 for CARIBIC samples.

that the slope for CARIBIC samples is indistinguishable from the correlations reported in the previous studies for stratospheric samples (-0.024 ‰/ppb) (Kawagucci et al., 2008; Wiegel et al., 2013; Boering et al., 2004). However, the regression lines
315 have an offset. This offset likely arises from variations in the $\Delta^{17}\text{O}(\text{CO}_2)$ measurement scales among different laboratories. Recently, Liang et al. (2023) reported a scale offset in $\Delta^{17}\text{O}(\text{CO}_2)$ of 0.037 to 0.042 ‰ between two laboratories (Institute of Earth Sciences, Academia Sinica and Institute of Earth Sciences, Hebrew University of Jerusalem). To optimize the application of $\Delta^{17}\text{O}(\text{CO}_2)$ measurements, the implementation of interlaboratory calibration is crucial (Adnew and Röckmann, 2022). A recent study by Laskar et al. (2019) reported a higher slope (0.036 ‰/ppb) for the $\text{N}_2\text{O}-\Delta^{17}\text{O}$ correlation of CARIBIC samples
320 collected during two flights. The samples they analyzed are mostly represent upper troposphere air, evident from a lower O_3 mole fraction compared to the CARIBIC samples measured in this study. This is supported by the relatively low three-isotope slope of 0.48 reported by Laskar et al. (2019) compared to the samples measured in this study (> 0.54). Furthermore, their samples cover only a very small range of N_2O (only 3.5 ppb difference, i.e. between 311.5 and 315 ppb) which likely leads to a high error in the determination of the $\text{N}_2\text{O}-\Delta^{17}\text{O}(\text{CO}_2)$ correlation slope.

325 In contrast, for the StratoClim samples, the $\text{N}_2\text{O}-\Delta^{17}\text{O}(\text{CO}_2)$ correlation shows an unexpectedly low slope of -0.017 ‰/ppb , which is about 1.5 times lower than for the CARIBIC and other stratospheric samples (Figure 4). All our samples were




Figure 8. $\Delta^{17}\text{O} - \text{N}_2\text{O}$ correlation for CARIBIC samples and measurements of Kawagucci et al. (2008) and Boering et al. (2004). The inset shows the zoom to the data close to the tropopause. For the CARIBIC samples measured in this study, the error-bars are smaller than the data markers. For the linear regression slopes, the errors are the 95 % confidence interval.

measured in the same laboratory and on the same analytical system. We suggest that the lower $\text{N}_2\text{O} - \Delta^{17}\text{O}$ correlation slope for StratoClim samples is due to enhanced vertical mixing during the Asian summer monsoon anticyclone (ASMA), as these samples were collected during mid-June to early September. The ASMA causes deep convection and anticyclonic flow in the upper troposphere and lower stratosphere (UTLS) where tropospheric air masses are uplifted into the UTLS (Park et al., 2009; Vogel et al., 2015, 2016; Brunamonti et al., 2018). The lifetime for CO_2 isotopic exchange is much slower than the transport time at all altitudes, whereas the photochemical production and quenching rates for $\text{O}^1(D)$ are much faster than transport processes. Therefore, the isotopic composition of stratospheric CO_2 should reflect both the variety of transport histories of air parcels and the sources of $\text{O}^1(D)$ (Liang et al., 2007; Boering et al., 2004). Furthermore, throughout the entire atmosphere, the CO_2 isotopic exchange time is longer than the transport time. However, the lifetime of N_2O varies with altitude. In the lower stratosphere, the lifetime of N_2O against photolysis is longer than the transport time, resulting in less variability in the region.



However, in the middle and stratosphere the N_2O life time decreases resulting a different $\text{N}_2\text{O}-\Delta^{17}\text{O}$ correlation slope (Liang et al., 2008). The influence of vertical mixing on $\text{N}_2\text{O}-\Delta^{17}\text{O}$ correlations was investigated in a model study by Liang et al. (2007). Indeed, they found that an increase in the vertical eddy diffusion coefficient - thus more vigorous vertical mixing -
340 leads to a lower slope of the $\text{N}_2\text{O}-\Delta^{17}\text{O}$ correlation. This supports our hypothesis that the low correlation for the StratoClim samples is caused by intense vertical mixing in the ASMA. An independent indication for the enhanced mixing is the scatter observed in three isotope plot and air age variability for the samples classified as stratospheric (Figure 6). Enhanced mixing, probably caused by deep convection and anticyclone flow in the upper troposphere and lower stratosphere, is consistent with the enhanced mole fraction of ozone-depleting substances in the samples collected in the same flight (Adcock et al., 2021).

345 High-precision $\text{N}_2\text{O}-\Delta^{17}\text{O}(\text{CO}_2)$ correlation can thus help identify enhanced mixing of tropospheric air into the stratosphere and maybe a measurable tracer to quantify the intensity of eddy diffusion/transport (Liang et al., 2007; Boering et al., 2004). There are many other indicators for enhanced mixing above the tropopause due to ASMA including a 25 % increase in long-lived ozone-depleting substances in the upper troposphere-lower stratosphere from the same samples (Adcock et al., 2021) and a contribution of about 30 % of young tropospheric air in the extratropical lower stratosphere in the Northern Hemisphere due to ASMA (Vogel et al., 2016). Recently, Ma et al. (2022) reported that 30 % to 50 % of the air mass in the UTLS
350 above the ASMA region is mixed, with the highest mixing occurring around 16.5 km altitude. Moreover, for StratoClim samples collected over the Indian subcontinent in Kathmandu (2017) in the tropopause region, the three major chlorinated very short-lived substances were enhanced up to 136 % compared to typical tropical tropopause values in 2013–2014. In contrast, only a 10 % increase was observed in ground-based measurements from 2014 to 2017 (Adcock et al., 2021; Engel et al., 2019).

355 For the StratoClim samples, the CO mole fraction reached a stratospheric background value at a potential temperature of 420 K, and the N_2O mole fraction remained similar to the tropospheric value up to 400 K, as described in detail by von Hobe et al. (2021). The deviation of the correlation between N_2O and $\Delta^{17}\text{O}$ is observed for air samples where the potential temperature is higher, about 420 K and above (Figure 9). At these potential temperatures, mixing processes become more significant and the air inside the anticyclone is exported vertically and horizontally into the surrounding stratosphere (von Hobe et al., 2021;
360 Ma et al., 2022). This enhanced mixing apparently causes the diminished slope for the N_2O and $\Delta^{17}\text{O}$ correlation plot (Figure 4) and also a very scattered three isotope plot (Figure 6). Up to a potential temperature of 400 K to 415 K the strong isolation of air inside the ASMA prevents significant mixing of the stratospheric air into the predominately tropospheric inner cyclone (von Hobe et al., 2021). Consequently, the $\text{N}_2\text{O}-\Delta^{17}\text{O}$ correlation slope of the lowermost stratospheric StratoClim samples agrees with the ones for CARIBIC samples and previously published measurements.

365 5 Implication for troposphere and stratosphere exchange and surface emissions of CO_2

Air samples from the upper troposphere and lower stratosphere are well suited for studying stratosphere-troposphere exchange (Olsen et al., 2001). We used CARIBIC samples with $\Delta^{17}\text{O}(\text{CO}_2) > -0.2 \text{‰}$, because for this group $\Delta^{17}\text{O}(\text{CO}_2)$ and N_2O have a very compact relationship with a slope of $-0.024 \pm 0.0002 \text{‰ ppb}^{-1}$, despite the air samples being collected at different latitudes, seasons, and years (see Table S1, Figure 1 and Figure 3). The $\Delta^{17}\text{O}(\text{CO}_2): \text{N}_2\text{O}$ correlation slope is similar to the




Figure 9. Correlation between $\Delta^{17}\text{O}$ and potential temperature for StratoClim samples. The inset shows the $\text{N}_2\text{O}-\Delta^{17}\text{O}$ correlation of StratoClim samples from Figure 4. The color bar for the inset is potential temperature. The data points marked in red indicate those with a low $\text{N}_2\text{O}-\Delta^{17}\text{O}$ correlation slope.

370 values reported in the previous measurements (Boering et al., 2004; Wiegel et al., 2013; Kawagucci et al., 2008). However, the uncertainty of the correlation slope is ten times smaller than in previous measurements (see Garofalo et al. (2019) for a detailed summary), and the new data confirm that the correlation actually extends down to the troposphere. Using the net vertical flux of N_2O (which is equal to the global N_2O loss rate, $13.43 \text{ TgNyr}^{-1} \pm 3 \%$ (Prather et al., 2023); see also (Tian et al., 2020)), the global mean net flux of $\Delta^{17}\text{O}(\text{CO}_2)$ is $51.3 \pm 1.6 \text{ ‰PgCyr}^{-1}$. This estimate aligns well with previous numerical model
375 calculations (Liang et al., 2008) and falls within the range reported previously using a similar approach (Boering et al., 2004; Garofalo et al., 2019; Wiegel et al., 2013; Kawagucci et al., 2008), but the uncertainty in our estimate is much lower than the 10 to 20 ‰PgCyr^{-1} uncertainty reported in the previous studies (Garofalo et al., 2019). The relatively precise estimate of the net $\Delta^{17}\text{O}(\text{CO}_2)$ flux from the stratosphere to the troposphere results from the improved precision of $\Delta^{17}\text{O}(\text{CO}_2)$ measurements (see Figure 8) and the reduced uncertainty in the global N_2O loss rate (3 % vs. 25 %).

380 Using the mass balance model described in section 2.3 (see equation 8), we estimated a terrestrial flux of $749 \pm 93 \text{ PgC/year}$. This estimate falls within the range, near the upper range of values, reported by previous studies (200 - 817 PgC/year) (Ciais et al., 1997a, b; Cuntz et al., 2003b, a; Farquhar et al., 1993; Liang et al., 2017b; Welp, 2011). The surface flux, which is the sum of terrestrial and ocean fluxes, is $829 \pm 93 \text{ PgC/year}$ (see Figure 11). This value is consistent with the range reported by




Figure 10. a) The correlation of two short-lived gases, ozone vs CO for Kathmandu samples, b) Correlation of $\Delta^{17}\text{O}$ with CO for both Kalamata and Kathmandu samples. The data points marked in red indicate those with a low $\text{N}_2\text{O}-\Delta^{17}\text{O}$ correlation slope.

Welp et al. (2008) using $\delta^{18}\text{O}$ of CO_2 . The corresponding surface turnover time of CO_2 is 0.98 ± 0.1 years, also within the
 385 range of previous estimates (i.e., 0.4 to 2.8 years) (Liang et al., 2017b, 2023; Laskar et al., 2019; Welp et al., 2011; Farquhar
 et al., 1993; Ciais et al., 1997a; Cuntz et al., 2003b) (see Figure 11). As shown in Figure 11, the estimated surface flux and
 turnover time of CO_2 are sensitive to the $\Delta^{17}\text{O}(\text{CO}_2)$ value in the upper troposphere. A higher $\Delta^{17}\text{O}(\text{CO}_2)$ value in the upper
 troposphere results in a lower surface flux and a longer CO_2 turnover time. A detailed sensitivity analysis is provided in the
 supplementary material. The GPP value is 211 ± 26 PgC/year, which is in good agreement with the estimates (200 PgC/year)
 390 by Liang et al. (2023) and Hofmann et al. (2017) using $\Delta^{17}\text{O}(\text{CO}_2)$ and higher than the estimate (150-175 PgC/year) by Welp
 et al. (2008) using $\delta^{18}\text{O}$ of CO_2 . These estimates are very sensitive to the assumed c_m/c_a ratio, the degree of equilibration
 (Cuntz, 2011; Farquhar et al., 1993) and the soil invasion flux (Wingate et al., 2009). For example, increasing the c_m/c_a ratio
 by 0.05 ($c_m/c_a = 0.7$) decreases the GPP from 212 PgC/year to 189 PgC/year. Conversely, decreasing the c_m/c_a ratio by 0.05
 ($c_m/c_a = 0.6$) will increase the GPP from 212 to 234 PgC/year. A detailed sensitivity analysis is provided in the supplementary
 395 material. Although the approach involves a number of assumptions as detailed in the methods section, it has the potential to be a
 valuable tool for quantifying and refining gross fluxes (Hoag et al., 2005; Koren et al., 2019). This will be especially true when
 more measurements become available from the lower stratosphere/upper troposphere and troposphere/surface. Additionally,
 understanding the effects of various processes on $\Delta^{17}\text{O}(\text{CO}_2)$, such as soil invasion, and taking into account inter-laboratory



calibration scale differences (Adnew and Röckmann, 2022; Liang et al., 2023) and proper model assimilation (Koren et al.,
400 2019), will further enhance its effectiveness.




Figure 11. Surface flux of CO₂ (F_{surf} , left axis) and the oxygen isotope turnover time of CO₂ (left axis) as a function of the $\Delta^{17}\text{O}_{UT}$ value of upper tropospheric CO₂. The solid line indicates our best estimates of F_{surf} and the oxygen isotope turnover time of CO₂ and the shaded area is the corresponding uncertainty calculated using error propagation.

6 Conclusions

Through the analysis of CARIBIC samples, this study showed that the N₂O- $\Delta^{17}\text{O}(\text{CO}_2)$ correlation reported previously from stratospheric samples extends down to the tropopause. Using the N₂O- $\Delta^{17}\text{O}(\text{CO}_2)$ correlation slope of CARIBIC samples, we estimated the net mean $\Delta^{17}\text{O}(\text{CO}_2)$ flux from the stratosphere to the troposphere to be $51.3 \pm 1.6 \text{‰PgCyr}^{-1}$. Notably, no
405 significant spatial or hemispheric variability was observed in $\Delta^{17}\text{O}(\text{CO}_2)$ values for the upper tropospheric samples collected during the CARIBIC program. The new measurements can be used in a mass balance approach to estimate that the surface turnover time of CO₂ is approximately 1 year, and the GPP is estimated to between 185 to 237 PgC/year.



In contrast, StratoClim samples showed a much lower $N_2O-\Delta^{17}O$ slope compared to CARIBIC samples and previous studies. This deviation is attributed to the increased mixing/eddy diffusion due to ASMA. $N_2O-\Delta^{17}O$ slope may thus be a direct measurable tracer for the intensity of vertical mixing in the UTLS region.

Data availability. All the data is reported in the form of tables

Author contributions. GAA: Conceptualization; data curation; formal analysis; investigation; methodology; software, visualization; Writing - original draft; Writing - review & editing, GK: investigation, Writing - review & editing, NM: data curation, Writing - review & editing, SG: investigation, MK: data curation, Writing - review & editing, TR: Conceptualization; investigation; methodology, Writing - review & editing

Competing interests. Some authors are members of the editorial board of Atmospheric Measurement Techniques. The authors declare no competing interest.

Acknowledgements. The StratoClim project was funded by the European Commission within Framework Program 7 under ENV.2013.6.1-2, Grant agreement No. 603557. The authors acknowledge Carl Brenninkmeijer and his team for providing the CARIBIC samples and trace gas data. We also thank Carl Brenninkmeijer for his constructive feedback on the draft manuscript. Additionally, we appreciate the contributions of all staff involved in the regular maintenance of the IAGOS-CARIBIC container, flight preparation, handling of air sampling units, extraction of CO_2 from whole air samples, and analysis of trace gases used in this study. We are grateful to the staff who performed the sampling and trace gas analysis of StratoClim samples and provided us with the trace gas data. Furthermore, we acknowledge the technical support from Carina van der Veen, Marcel J. Portanger, and Henk Snellen. We acknowledge A. H. Laskar and M-C. Liang for constructive discussion.



References

- Adcock, K. E., Fraser, P. J., Hall, B. D., Langenfelds, R. L., Lee, G., Montzka, S. A., Oram, D. E., Röckmann, T., Stroh, F., Sturges, W. T., et al.: Aircraft-based observations of ozone-depleting substances in the upper troposphere and lower stratosphere in and above the Asian summer monsoon, *Journal of Geophysical Research: Atmospheres*, 126, e2020JD033 137, 2021.
- 430 Adnew, G. A.: The effect of photosynthetic gas exchange on $\Delta^{17}\text{O}$ of atmospheric CO_2 , PhD thesis, pp. 1–174, <https://doi.org/10.33540/412>, 2020.
- Adnew, G. A. and Röckmann, T.: Urgent need for $\Delta^{17}\text{O}$ (CO_2) standardization. The 21st expert meeting on Greenhouse gas measurement techniques (GGMT), Wageningen, The Netherlands, 2022.
- Adnew, G. A., Hofmann, M. E. G., Paul, D., Laskar, A., Surma, J., Albrecht, N., Pack, A., Schwieters, J., Koren, G., Peters, W., and
435 Röckmann, T.: Determination of the triple oxygen and carbon isotopic composition of CO_2 from atomic ion fragments formed in the ion source of the 253 Ultra High-Resolution Isotope Ratio Mass Spectrometer, *Rapid Commun Mass Spectrom.*, 33, 17, 2019.
- Adnew, G. A., Pons, T. L., Koren, G., Peters, W., and Röckmann, T.: Leaf-scale quantification of the effect of photosynthetic gas exchange on $\Delta^{17}\text{O}$ of atmospheric CO_2 , *Biogeosciences Discuss.*, 2020, 1–37, <https://doi.org/10.5194/bg-2020-91>, 2020.
- Adnew, G. A., G.Koren, Ziegler, M., and Röckmann, T.: Leaf scale quantification of the effect of photosynthetic gas exchange on $\Delta^{17}\text{O}$ of
440 CO_2 , *Sci. Rep.*, 11, 14 023, <https://doi.org/10.1038/s41598-021-93092-0>, 2021.
- Adnew, G. A., Workman, E., Janssen, C., and Röckmann, T.: Temperature dependence of isotopic fractionation in the CO_2 - O_2 isotope exchange reaction, *Rapid Communications in Mass Spectrometry*, p. e9301, 2022.
- Adnew, G. A., Pons, T. L., Koren, G., Peters, W., and Röckmann, T.: Exploring the potential of $\Delta^{17}\text{O}$ in CO_2 for determining mesophyll conductance, *Plant Physiology*, p. kiad173, 2023.
- 445 Alexander, B., Vollmer, M., Jackson, T., Weiss, R., and Thiemens, M.: Stratospheric CO_2 isotopic anomalies and SF_6 and CFC tracer concentrations in the Arctic polar vortex, *Geophysical Research Letters*, 28, 4103–4106, 2001.
- Angert, A., Rachmilevitch, S., Barkan, E., and Luz, B.: Effects of photorespiration, the cytochrome pathway, and the alternative pathway on the triple isotopic composition of atmospheric O_2 , *Global Biogeochemical Cycles*, 1, 1030, 2003.
- Angert, A., Cappa, C. D., and DePaolo, D. J.: Kinetic ^{17}O effects in the hydrologic cycle: Indirect evidence and implications, *Geochim. Cosmochim. Acta*, 68, 3487–3495, 2004.
450
- Annamalai, H. and Slingo, J.: Active/break cycles: Diagnosis of the intraseasonal variability of the Asian summer monsoon, *Climate Dynamics*, 18, 85–102, 2001.
- Appenzeller, C., Holton, J. R., and Rosenlof, K. H.: Seasonal variation of mass transport across the tropopause, *Journal of Geophysical Research: Atmospheres*, 101, 15 071–15 078, 1996.
- 455 Assonov, S., Taylor, P., and Brenninkmeijer, C.: A system for high-quality CO_2 isotope analyses of air samples collected by the CARIBIC Airbus A340-600, *Rapid Communications in Mass Spectrometry*, 23, 1347–1363, 2009a.
- Assonov, S. S., Brenninkmeijer, C. A. M., Koepfel, C., and Röckmann, T.: CO_2 isotope analyses using large air samples collected on intercontinental flights by the CARIBIC Boeing 767, *Rapid Commun. Mass Spectrom.*, 23, 822–830, 2009b.
- Assonov, S. S., Brenninkmeijer, C. A. M., Schuck, T. J., and Taylor, P.: Analysis of ^{13}C and ^{18}O isotope data of CO_2 in CARIBIC aircraft
460 samples as tracers of upper troposphere/lower stratosphere mixing and the global carbon cycle, *Atoms. Chem Phys.*, 10, 8575–8599, 2010.
- Assonov, S. S., Brenninkmeijer, C. A. M., Schuck, T., and Umezawa, T.: N_2O as a tracer of mixing stratospheric and tropospheric air based on CARIBIC data with applications for CO_2 , *Atmos. Environ.*, 79, 769–779, 2013.



- Barkan, E. and Luz, B.: High precision measurements of $^{17}\text{O}/^{16}\text{O}$ and $^{18}\text{O}/^{16}\text{O}$ ratios in H_2O , *Rapid Comm. Mass Spectrom.*, 19, 3737–3742, <https://doi.org/10.1002/rcm.2250>, 2005.
- 465 Barkan, E. and Luz, B.: Diffusivity fractionations of $\text{H}_2^{16}\text{O}/\text{H}_2^{17}\text{O}$ and $\text{H}_2^{16}\text{O}/\text{H}_2^{18}\text{O}$ in air and their implications for isotope hydrology, *RAPID COMMUN MASS SP*, 21, 6, 2007.
- Barkan, E. and Luz, B.: High-precision measurements of $^{17}\text{O}/^{16}\text{O}$ and $^{18}\text{O}/^{16}\text{O}$ ratios in CO_2 , *Rapid Commun. Mass. Sp.*, 26, 2733–2738, <https://doi.org/10.1002/rcm.6400>, 2012.
- Barkan, E., Musan, I., and Luz, B.: High-precision measurements of $\delta^{17}\text{O}$ and ^{17}O -excess of NBS19 and NBS18, *Rapid Commun. Mass. Sp.*, 29, 2219–2224, <https://doi.org/10.1002/rcm.7378>, 2015.
- 470 Boering, K. A., Jackson, T., Hoag, K. J., Cole, A. S., Perri, M. J., Thiemens, M., and Atlas, E.: Observations of the anomalous oxygen isotopic composition of carbon dioxide in the lower stratosphere and the flux of the anomaly to the troposphere, *Geophys. Res. Lett.*, 31, 2004.
- Bottinga, Y. and Craig, H.: Oxygen isotope fractionation between CO_2 and water, and the isotopic composition of marine atmospheric CO_2 ,
475 *Earth Planet Sc. Lett.*, 5, 10, 1968.
- Brenninkmeijer, C. and Röckmann, T.: Russian doll type cryogenic traps: Improved design and isotope separation effects, *Analytical chemistry*, 68, 3050–3053, 1996.
- Brenninkmeijer, C., Kraft, P., and Mook, W.: Oxygen isotope fractionation between CO_2 and H_2O , *Chemical Geology*, 41, 181–190, 1983.
- Brenninkmeijer, C., Lowe, D., Manning, M., Sparks, R., and Van Velthoven, P.: The ^{13}C , ^{14}C , and ^{18}O isotopic composition of CO , CH_4 ,
480 and CO_2 in the higher southern latitudes lower stratosphere, *Journal of Geophysical Research: Atmospheres*, 100, 26 163–26 172, 1995.
- Brenninkmeijer, C., Crutzen, P., Fischer, H., Güsten, H., Hans, W., Heinrich, G., Heintzenberg, J., Hermann, M., Immelmann, T., Kersting, D., et al.: CARIBIC—Civil aircraft for global measurement of trace gases and aerosols in the tropopause region, *Journal of Atmospheric and Oceanic Technology*, 16, 1373–1383, 1999.
- Brenninkmeijer, C. A.: Robust, high-efficiency, high-capacity cryogenic trap, *Analytical Chemistry*, 63, 1182–1184, 1991.
- 485 Brenninkmeijer, C. A.: Measurement of the abundance of ^{14}CO in the atmosphere and the $^{13}\text{C}/^{12}\text{C}$ and $^{18}\text{O}/^{16}\text{O}$ ratio of atmospheric CO with applications in New Zealand and Antarctica, *Journal of Geophysical Research: Atmospheres*, 98, 10 595–10 614, 1993.
- Brunamonti, S., Jorge, T., Oelsner, P., Hanumanthu, S., Singh, B. B., Kumar, K. R., Sonbawne, S., Meier, S., Singh, D., Wienhold, F. G., et al.: Balloon-borne measurements of temperature, water vapor, ozone and aerosol backscatter on the southern slopes of the Himalayas during StratoClim 2016–2017, *Atmospheric Chemistry and Physics*, 18, 15 937–15 957, 2018.
- 490 Cairo, F., Pommereau, J.-P., Law, K. S., Schlager, H., Garnier, A., Fierli, F., Ern, M., Streibel, M., Arabas, S., Borrmann, S., et al.: An introduction to the SCOUT-AMMA stratospheric aircraft, balloons and sondes campaign in West Africa, August 2006: rationale and roadmap, *Atmospheric chemistry and physics*, 10, 2237–2256, 2010.
- Cao, X. and Liu, Y.: Equilibrium mass-dependent fractionation relationships for triple oxygen isotopes, *Geochim. Cosmochim. Acta.*, 75, 7435–7445, 2011.
- 495 Cernusak, L. A., Barbour, M. M., Arndt, S. K., Cheesman, A. K., English, N. B., Feild, T. S., Helliker, B. R., Holloway-Phillips, M. M., Holtum, J. A., Kahmen, A., McInerney, F. A., Munksgaard, N. C., Simonin, K. A., Song, X., Stuart-Williams, H., West, J. B., and Farquhar, G. D.: Stable isotopes in leaf water of terrestrial plants, *Plant Cell Environ.*, 39, 1087–1102, <https://doi.org/10.1111/pce.12703>, 2016.
- Ciais, P., Tans, P. P., Trolier, M., White, J. W. C., and Francey, R. J.: A large northern hemisphere terrestrial CO_2 sink indicated by the $^{13}\text{C}/^{12}\text{C}$ ratio of atmospheric CO_2 , *Science*, 269, 4, 1995.



- 500 Ciais, P., Denning, A. S., Tans, P. P., Berry, J. A., Randall, D. A., Collatz, G. J., Sellers, P. J., White, J. W., Trolier, M., Meijer, H. A., et al.: A three-dimensional synthesis study of $\delta^{18}\text{O}$ in atmospheric CO_2 : 1. Surface fluxes, *Journal of Geophysical Research: Atmospheres*, 102, 5857–5872, 1997a.
- Ciais, P., Tans, P. P., Denning, A. S., Francey, R. J., Trolier, M., Meijer, H. A., White, J. W., Berry, J. A., Randall, D. A., Collatz, G. J., et al.: A three-dimensional synthesis study of $\delta^{18}\text{O}$ in atmospheric CO_2 : 2. Simulations with the TM2 transport model, *Journal of Geophysical*
505 *Research: Atmospheres*, 102, 5873–5883, 1997b.
- Cousins, A. B., Badger, M. R., and Caemmerer, S. v.: Carbonic anhydrase and its influence on carbon isotope discrimination during C_4 photosynthesis. Insights from antisense RNA in *Flaveria bidentis*, *Plant Physiol.*, 141, 232–242, 2006.
- Cuntz, M.: A dent in carbon's gold standard, *Nature*, 477, 547–548, 2011.
- Cuntz, M., Ciais, P., Hoffmann, G., Allison, C. E., Francey, R. J., Knorr, W., Tans, P. P., White, J. W., and Levin, I.: A comprehensive
510 global three-dimensional model of $\delta^{18}\text{O}$ in atmospheric CO_2 : 2. Mapping the atmospheric signal, *Journal of Geophysical Research: Atmospheres*, 108, 2003a.
- Cuntz, M., Ciais, P., Hoffmann, G., and Knorr, W.: A comprehensive global three-dimensional model of $\delta^{18}\text{O}$ in atmospheric CO_2 : 1. Validation of surface processes, *Journal of Geophysical Research: Atmospheres*, 108, 2003b.
- Dai, A.: Recent climatology, variability, and trends in global surface humidity, *J. Clim.*, 19, 17, 2006.
- 515 Engel, A., Rigby, M., Burkholder, J. B., Fernandez, R. P., Froidevaux, L., Hall, B. D., Hossaini, R., Saito, T., Vollmer, M. K., and Yao, B.: Update on ozone-depleting substances (ODSs) and other gases of interest to the Montreal Protocol, 2019.
- Farquhar, G. D. and Lloyd, J.: Carbon and oxygen isotope effects in the exchange of carbon dioxide between terrestrial plants and the atmosphere, Academic Press Inc, London, 1993.
- Farquhar, G. D., Lloyd, J., Taylor, J. A., Flanagan, L. B., Syvertsen, J. P., Hubick, K. T., Wong, S. C., and Ehleringer, J. R.: Vegetation effects
520 on the isotope composition of oxygen in atmospheric CO_2 , *Nature*, 363, 439–443, 1993.
- Francey, R. J. and Tans, P. P.: Latitudinal variation in oxygen-18 of atmospheric CO_2 , *Nature*, 327, 495–497, 1987.
- Friedli, H. and Siegenthaler, U.: Influence of N_2O on isotope analyses in CO_2 and mass-spectrometric determination of N_2O in air samples, *Tellus B*, 40, 129–133, 1988.
- Gamo, T., Tsutsumi, M., Sakai, H., Nakazawa, T., Machida, T., Honda, H., and Itoh, T.: Long-term monitoring of carbon and oxygen isotope
525 ratios of stratospheric CO_2 over Japan, *Geophysical research letters*, 22, 397–400, 1995.
- Gan, K. S., Wong, S. C., Yong, J. W. H., and Farquhar, G. D.: ^{18}O spatial patterns of vein xylem water, leaf water, and dry matter in cotton leaves, *Plant Physiol.*, 130, 13, 2002.
- Garny, H. and Randel, W.: Dynamic variability of the Asian monsoon anticyclone observed in potential vorticity and correlations with tracer distributions, *Journal of Geophysical Research: Atmospheres*, 118, 13–421, 2013.
- 530 Garofalo, L., Kanu, A., Hoag, K. J., and Boering, K. A.: The effects of stratospheric chemistry and transport on the isotopic compositions of long-lived gases measured at Earth's surface, *Advances in Atmospheric Chemistry: Volume 2: Organic Oxidation and Multiphase Chemistry*, pp. 529–587, 2019.
- Hayles, J. A. and Killingsworth, B. A.: Constraints on triple oxygen isotope kinetics, *Chemical Geology*, 589, 120–146, 2022.
- Hayles, J. A., Cao, X., and Bao, H.: The statistical mechanical basis of the triple isotope fractionation relationship, *Geochem. Perspect. Lett.*,
535 3, 1–11, 2017.
- Helliker, B. R. and Ehleringer, J. R.: Establishing a grassland signature in veins: ^{18}O in the leaf water of C_3 and C_4 grasses, *PNAS*, 97, 7894–7898, 2000.



- Hoag, K. J., Still, C. J., and Fung, I. Y.: Triple oxygen isotope composition of tropospheric carbon dioxide as a tracer of terrestrial gross carbon fluxes, *Geophys. Res. Lett.*, 32, <https://doi.org/10.1029/2004gl021011>, 2005.
- 540 Hofmann, M. E. G., Horváth, B., and Pack, A.: Triple oxygen isotope equilibrium fractionation between carbon dioxide and water, *Earth Planet Sci Lett.*, 319, 159–164, 2012.
- Hofmann, M. E. G., Horváth, B., Schneider, L., Peters, W., Schützenmeister, K., and Pack, A.: Atmospheric measurements of $\Delta^{17}\text{O}$ in CO_2 in Göttingen, Germany reveal a seasonal cycle driven by biospheric uptake, *Geochim. Cosmochim. Ac.*, 199, 143–163, <https://doi.org/10.1016/j.gca.2016.11.019>, 2017.
- 545 Holton, J. R., Haynes, P. H., McIntyre, M. E., Douglass, A. R., Rood, R. B., and Pfister, L.: Stratosphere-troposphere exchange, *Reviews of geophysics*, 33, 403–439, 1995.
- Horváth, B., Hofmann, M., and Pack, A.: On the triple oxygen isotope composition of carbon dioxide from some combustion processes, *Geochimica et Cosmochimica Acta*, 95, 160–168, 2012.
- Kaiser, J.: Reformulated ^{17}O correction of mass spectrometric stable isotope measurements in carbon dioxide and a critical appraisal of historic ‘absolute’ carbon and oxygen isotope ratios, *Geochimica et Cosmochimica Acta*, 72, 22, <https://doi.org/10.1016/j.gca.2007.12.011>, 2008.
- 550 Kaiser, J., Engel, A., Borchers, R., and Röckmann, T.: Probing stratospheric transport and chemistry with new balloon and aircraft observations of the meridional and vertical N_2O isotope distribution, *Atmospheric Chemistry and Physics*, 6, 3535–3556, 2006.
- Kawagucci, S., Tsunogai, U., Kudo, S., Nakagawa, F., Honda, H., Aoki, S., Nakazawa, T., and Gamo, T.: An analytical system for determining $\delta^{17}\text{O}$ in CO_2 using continuous flow-isotope ratio MS, *Anal. Chem.*, 77, 5, 2005.
- 555 Kawagucci, S., Tsunogai, U., Kudo, S., Nakagawa, F., Honda, H., Aoki, S., Nakazawa, T., Tsutsumi, M., and Gamo, T.: Long-term observation of mass-independent oxygen isotope anomaly in stratospheric CO_2 , *Atmos. Chem. Phys.*, 8, 8, 2008.
- Koren, G., Schneider, L., Velde, I. d., Schaik, E. v., Gromov, S. S., Adnew, G. A., D.J.Mrozek, Hofmann, M. E. D., Liang, M.-C., Mahata, S., Bergamaschi, P., Laan-Luijckx, I. T. v. d., Krol, M. C., Röckmann, T., and Peters, W.: Global 3-D Simulations of the Triple Oxygen Isotope Signature $\Delta^{17}\text{O}$ in Atmospheric CO_2 , *J. Geophys. Res. Atmos.*, 124, 28, 2019.
- 560 Krol, M., De Bruine, M., Killaars, L., Ouwersloot, H., Pozzer, A., Yin, Y., Chevallier, F., Bousquet, P., Patra, P., Belikov, D., et al.: Age of air as a diagnostic for transport timescales in global models, *Geoscientific Model Development*, 11, 3109–3130, 2018.
- Lämmerzahl, P., Röckmann, T., and Brenninkmeijer, C. A. M.: Oxygen isotope composition of stratospheric carbon dioxide, *Geophys. Res. Lett.*, 29, <https://doi.org/10.1029/2001gl014343>, 2002.
- 565 Lan, X., Thoning, K. W., and Dlugokencky, E. J.: Trends in globally-averaged CH_4 , N_2O , and SF_6 determined from NOAA Global Monitoring Laboratory measurements, Version 2024-08, <https://doi.org/10.15138/P8XG-AA10>, 15, 753–759, 2024.
- Landais, A., Barkan, E., Yakir, D., and Luz, B.: The triple isotopic composition of oxygen in leaf water, *Geochim. Cosmochim. Ac.*, 70, 4105–4115, <https://doi.org/10.1016/j.gca.2006.06.1545>, 2006.
- Landais, A., Yakir, D., Barkan, E., and Luz, B.: The triple isotopic composition of oxygen in leaf water and its implications for quantifying biosphere productivity, *Terrestrial Ecology*, 1, 111–125, 2007.
- 570 Laskar, A. H., Mahata, S., and Liang, M.-C.: Identification of anthropogenic CO_2 using triple oxygen and clumped isotopes, *Environ. Sci. Technol.*, 50, 18, <https://doi.org/10.1021/acs.est.6b02989>, 2016.
- Laskar, A. M., Mahata, S., Bhattacharya, S. K., and Liang, M.-C.: Triple oxygen and clumped isotope compositions of CO_2 in the middle troposphere, *Earth and Space Science*, 6, 1205–1219, 2019.



- 575 Levin, I., Allison, C., Brand, W., Conway, Delmotte, M., Facklam, C., Francey, R., Hammer, S., Huang, L., Jordan, A., Langenfelds, R., Machida, T., Meijer, H., Mukai, H., Neubert, R., Sabasch, M., Schmidt, M., Steele, P., Steinberg, B., Ramonet, M., Rothe, M., Vaughn, B., and Worthy, D.: Five Years of International Flask Intercomparison (Sausage Flask ICP) 2002–2007, 14th WMO/IAEA Meeting of Experts on Carbon Dioxide, Other Greenhouse Gases and Related Tracers Measurement Techniques, GAW Report No, 194, 330, 2007.
- Liang, M.-C. and Mahata, S.: Oxygen anomaly in near surface carbon dioxide reveals deep stratospheric intrusion, *Sci. Rep.*, 5, 11 352, <https://doi.org/10.1038/srep11352>, 2015.
- 580 Liang, M.-C., Geoffrey, A. B., Brenton, R. L., and Yung, Y. L.: Oxygen isotopic composition of carbon dioxide in the middle atmosphere, *PNAS*, 104, 5, 2007.
- Liang, M.-C., Blake, G. A., and Yung, Y. L.: Seasonal cycle of C16O16O, C16O17O, and C16O18O in the middle atmosphere: Implications for mesospheric dynamics and biogeochemical sources and sinks of CO₂, *J. Geophys. Res. Atmos.*, 113, D12, 2008.
- 585 Liang, M.-C., Mahata, S., Laskar, A. H., and Bhattacharya, S. K.: Spatiotemporal variability of oxygen isotope anomaly in near surface air CO₂ over urban, semi-urban and ocean areas in and around Taiwan, *Aerosol Air Qual. Res.*, 17, 24, 2017a.
- Liang, M.-C., Mahata, S., Laskar, A. H., Thiemens, M. H., and Newman, S.: Oxygen isotope anomaly in tropospheric CO₂ and implications for CO₂ residence time in the atmosphere and gross primary productivity, *Sci Rep*, 7, 13 180, <https://doi.org/10.1038/s41598-017-12774-w>, 2017b.
- 590 Liang, M.-C., Laskar, A. H., Barkan, E., Newman, S., Thiemens, M. H., and Rangarajan, R.: New constraints of terrestrial and oceanic global gross primary productions from the triple oxygen isotopic composition of atmospheric CO₂ and O₂, *Scientific Reports*, 13, 2162, 2023.
- Luz, B. and Barkan, E.: Variations of 17O/16O and 18O/16O in meteoric waters, *Geochim. Cosmochim. Ac.*, 74, 10, 2010.
- Luz, B., Barkan, E., Bender, M. L., Thiemens, M. K., and Boering, K. A.: Triple-isotope composition of atmospheric oxygen as a tracer of biosphere productivity, *Nature*, 400, 3, 1999.
- 595 Lyons, J. R.: Transfer of mass-independent fractionation in ozone to other oxygen-containing radicals in the atmosphere, *Geophys. Res. Lett.*, 28, 3231–3234, <https://doi.org/10.1029/2000gl012791>, 2001.
- Ma, D., Bian, J., Li, D., Bai, Z., Li, Q., Zhang, J., Wang, H., Zheng, X., Hurst, D. F., and Vömel, H.: Mixing characteristics within the tropopause transition layer over the Asian summer monsoon region based on ozone and water vapor sounding data, *Atmospheric Research*, 271, 106 093, 2022.
- 600 Matsuhisa, Y., Goldsmith, J. R., and Clayton, R. N.: Mechanisms of hydrothermal crystallization of quartz at 250°C and 15 kbar, *Geochim. Cosmochim. Acta*, 42, 9, 1978.
- Meijer, H. and Li, W.: The use of electrolysis for accurate 17O and 18O isotope measurements in water isotopes, *Isotopes Environ. Health Stud.*, 34, 20, 1998.
- Mikkonen, S., Pitkänen, M. R., Nieminen, T., Lipponen, A., Isokääntä, S., Arola, A., and Lehtinen, K. E.: Effects of uncertainties and number of data points on line fitting—a case study on new particle formation, *Atmospheric Chemistry and Physics*, 19, 12 531–12 543, 2019.
- 605 Miller, J. B., Yakir, D., White, J. W., and Tans, P. P.: Measurement of 18O/16O in the soil-atmosphere CO₂ flux, *Global Biogeochemical Cycles*, 13, 761–774, 1999.
- Mrozek, D. J. et al.: Measurements and interpretation of oxygen isotopes in stratospheric carbon dioxide, Ph.D. thesis, Utrecht University, 2017.
- 610 Nisbet, E. G., Manning, M., Dlugokencky, E., Fisher, R., Lowry, D., Michel, S., Myhre, C. L., Platt, S. M., Allen, G., Bousquet, P., et al.: Very strong atmospheric methane growth in the 4 years 2014–2017: Implications for the Paris Agreement, *Global Biogeochemical Cycles*, 33, 318–342, 2019.



- Olsen, S., McLinden, C., and Prather, M.: Stratospheric N₂O–NO_y system: Testing uncertainties in a three-dimensional framework, *Journal of Geophysical Research: Atmospheres*, 106, 28 771–28 784, 2001.
- 615 Pan, L. L., Honomichl, S. B., Kinnison, D. E., Abalos, M., Randel, W. J., Bergman, J. W., and Bian, J.: Transport of chemical tracers from the boundary layer to stratosphere associated with the dynamics of the Asian summer monsoon, *Journal of Geophysical Research: Atmospheres*, 121, 14–159, 2016.
- Park, M., Randel, W. J., Emmons, L. K., and Livesey, N. J.: Transport pathways of carbon monoxide in the Asian summer monsoon diagnosed from Model of Ozone and Related Tracers (MOZART), *Journal of Geophysical Research: Atmospheres*, 114, 2009.
- 620 Peylin, P., Ciais, P., Denning, A. S., Tans, P. P., Berry, J. A., and White, J. W.: A 3-dimensional study of $\delta^{18}\text{O}$ in atmospheric CO₂: contribution of different land ecosystems, *Tellus B*, 51, 25, 1999.
- Piao, S., Sitch, S., Ciais, P., Friedlingstein, P., Peylin, P., Wang, X., Ahlström, A., Anav, A., Canadell, J. G., Cong, N., et al.: Evaluation of terrestrial carbon cycle models for their response to climate variability and to CO₂ trends, *Global change biology*, 19, 2117–2132, 2013.
- Plumb, R. A.: Tracer interrelationships in the stratosphere, *Reviews of Geophysics*, 45, 2007.
- 625 Plumb, R. A. and Ko, M. K.: Interrelationships between mixing ratios of long-lived stratospheric constituents, *Journal of Geophysical Research: Atmospheres*, 97, 10 145–10 156, 1992.
- Prather, M. J., Froidevaux, L., and Livesey, N. J.: Observed changes in stratospheric circulation: decreasing lifetime of N₂O, 2005–2021, *Atmospheric Chemistry and Physics*, 23, 843–849, 2023.
- Pupek, M., Assonov, S., Mühle, J., Rhee, T., Oram, D., Koepfel, C., Slemr, F., and Brenninkmeijer, C.: Isotope analysis of hydrocarbons: trapping, recovering and archiving hydrocarbons and halocarbons separated from ambient air, *Rapid Communications in Mass Spectrometry: An International Journal Devoted to the Rapid Dissemination of Up-to-the-Minute Research in Mass Spectrometry*, 19, 455–460, 2005.
- 630 Saugier, B., Roy, J., and Mooney, H. A.: Estimations of global terrestrial productivity, *Terrestrial global productivity*, pp. 543–557, 2001.
- Shaheen, R., Janssen, C., and Röckmann, T.: Investigations of the photochemical isotope equilibrium between O₂, CO₂ and O₃, *Atmos. Chem. Phys.*, 7, 14, 2007.
- 635 Sirignano, C., Neubert, R. E., and Meijer, H. A.: N₂O influence on isotopic measurements of atmospheric CO₂, *Rapid communications in mass spectrometry*, 18, 1839–1846, 2004.
- Song, X., Simonin, K. A., Loucos, K. E., and Barbour, M. M.: Modelling non-steady-state isotope enrichment of leaf water in a gas-exchange cuvette environment, *Plant Cell Environ.*, 38, 10, 2015.
- 640 Stefanutti, L., Sokolov, L., Balestri, S., MacKenzie, A., and Khattatov, V.: The M-55 Geophysica as a platform for the Airborne Polar Experiment, *Journal of Atmospheric and Oceanic Technology*, 16, 1303–1312, 1999.
- Stern, L. A., Amundson, R., and Baisden, W. T.: Influence of soils on oxygen isotope ratio of atmospheric CO₂, *Global Biogeochemical Cycles*, 15, 753–759, 2001.
- Steur, P. M., Scheeren, H. A., Koren, G., Adnew, G. A., Peters, W., and Meijer, H. A.: Interannual variations in the $\Delta(17\text{O})$ signature of atmospheric CO₂ at two mid-latitude sites suggest a close link to stratosphere-troposphere exchange, *EGUsphere*, 2024, 1–31, 2024.
- 645 Stocker, T., Qin, D., Plattner, G-K Tignor, M., Allen, S., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P.: IPCC, 2013: Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, p. 1535, 2013.
- Thiemens, M. H.: History and Applications of Mass-Independent Isotope Effects, *Annu. Rev. Earth Planet. Sci.*, 34, 62, 650 <https://doi.org/10.1146/>, 2006.



- Thiemens, M. H., Jackso, T., Mauersberger, K., Schueler, B., and Morton, J.: Oxygen isotope fractionation in stratospheric CO₂, *Geophys. Res. Lett.*, 18, 3, 1991.
- Thiemens, M. H., Jackson, T., Zipf, E. C., Erdman, P. W., and van Egmond, C.: Carbon dioxide and oxygen isotope anomalies in the mesosphere and stratosphere, *Science*, 270, 969–972, 1995a.
- 655 Thiemens, M. H., Jackson, T., Zipf, E. D., Erdman, P. W., and Egmond, C. v.: Carbon dioxide and oxygen isotope anomalies in the mesosphere and stratosphere, *Science*, 270, 969–972, 1995b.
- Thiemens, M. H., Jackson, T. L., and Brenninkmeijer, C. A.: Observation of a mass independent oxygen isotopic composition in terrestrial stratospheric CO₂, the link to ozone chemistry, and the possible occurrence in the Martian atmosphere, *Geophysical Research Letters*, 22, 255–257, 1995c.
- 660 Thiemens, M. K., Chakraborty, S., and Jackson, T. L.: Decadal $\Delta^{17}\text{O}$ record of tropospheric CO₂: Verification of a stratospheric component in the troposphere, *J. Geophys. Res. Atmos.*, 119, 8, 2014.
- Tian, H., Xu, R., Canadell, J. G., Thompson, R. L., Winiwarter, W., Suntharalingam, P., Davidson, E. A., Ciais, P., Jackson, R. B., Janssens-Maenhout, G., et al.: A comprehensive quantification of global nitrous oxide sources and sinks, *Nature*, 586, 248–256, 2020.
- Vogel, B., Günther, G., Müller, R., Groöß, J.-U., and Riese, M.: Impact of different Asian source regions on the composition of the Asian monsoon anticyclone and of the extratropical lowermost stratosphere, *Atmospheric chemistry and physics*, 15, 13 699–13 716, 2015.
- 665 Vogel, B., Günther, G., Müller, R., Groöß, J.-U., Afchine, A., Bozem, H., Hoor, P., Krämer, M., Müller, S., Riese, M., et al.: Long-range transport pathways of tropospheric source gases originating in Asia into the northern lower stratosphere during the Asian monsoon season 2012, *Atmospheric Chemistry and Physics*, 16, 15 301–15 325, 2016.
- Vogel, B., Müller, R., Günther, G., Spang, R., Hanumanthu, S., Li, D., Riese, M., and Stiller, G. P.: Lagrangian simulations of the transport of young air masses to the top of the Asian monsoon anticyclone and into the tropical pipe, *Atmospheric Chemistry and Physics*, 19, 6007–6034, 2019.
- 670 von Hobe, M., Ploeger, F., Konopka, P., Kloss, C., Ulanowski, A., Yushkov, V., Ravegnani, F., Volk, C. M., Pan, L. L., Honomichl, S. B., et al.: Upward transport into and within the Asian monsoon anticyclone as inferred from StratoClim trace gas observations, *Atmospheric Chemistry and Physics*, 21, 1267–1285, 2021.
- 675 Welp, L.: Interannual variability in the oxygen isotopes of atmospheric CO₂ driven by El Nino Welp, Lisa R.; Keeling, Ralph F.; Meijer, Harro AJ; Bollenbacher, Alane F.; Piper, Stephen C.; Yoshimura, Kei; Francey, Roger J.; Allison, Colin E.; Wahlen, Martin, *Nature*, 477, 579–582, 2011.
- Welp, L. R., Lee, X., Kim, K., Griffis, T. J., Billmark, K. A., and Baker, J. M.: $\delta^{18}\text{O}$ of water vapour, evapotranspiration and the sites of leaf water evaporation in a soybean canopy, *Plant, Cell & Environment*, 31, 1214–1228, 2008.
- 680 Welp, L. R., Keeling, R. F., Meijer, H. A. J., Bollenbacher, A. F., Piper, S. C., Yoshimura, K., Francey, R. J., Allison, C. A., and Wahlen, M.: Interannual variability in the oxygen isotopes of atmospheric CO₂ driven by El Niño, *Nature*, 477, 579–582, <https://doi.org/10.1038/nature10421>, 2011.
- Wiegel, A. A., Cole, A. S., Hoag, K. J., Atlas, E. L., Schauffler, S. M., and Boering, K. A.: Unexpected variations in the triple oxygen isotope composition of stratospheric carbon dioxide, *PNAS*, 110, 17 680–17 685, <https://doi.org/10.1073/pnas.1213082110>, 2013.
- 685 Wingate, L., Ogé, J., Cuntz, M., Genty, B., Reiter, I., Seibt, U., Yakir, D., Maseyk, K., Pendall, E. G., Barbour, M. M., Mortazavi, B., Burlett, R., Peylin, P., Miller, J., Mencuccini, M., Shim, J. H., Hunt, J., and Grace, J.: The impact of soil microorganisms on the global budget of $\delta^{18}\text{O}$ in atmospheric CO₂, *PNAS*, 106, 4, 2009.
- Yakir, D.: Oxygen-18 of leaf water: a crossroad for plant-associated isotopic signals, in: *Stable isotopes*, pp. 147–168, Garland Science, 2020.



- 690 Yeung, L. Y., Affek, H. P., Hoag, K. J., Guo, W., Wiegel, A. A., Atlas, E. L., Schauffler, S. M., Okumura, M., Boering, K. A., and Eiler, J. M.: Large and unexpected enrichment in stratospheric $16\text{O}13\text{C}18\text{O}$ and its meridional variation, *Proceedings of the national academy of sciences*, 106, 5, 2009.
- Young, E. D., Galy, A., and Nagahara, H.: Kinetic and equilibrium mass-dependent isotope fractionation laws in nature and their geochemical and cosmochemical significance, *Geochim. Cosmochim. Ac.*, 66, 9, 2002.
- 695 Yung, Y. L., DeMore, W., and Pinto, J. P.: Isotopic exchange between carbon dioxide and ozone via O (1D) in the stratosphere, *Geophysical Research Letters*, 18, 13–16, 1991a.
- Yung, Y. L., DeMore, W. B., and Pinto, J. P.: Isotopic exchange between carbon dioxide and ozone via O (1D) in the stratosphere, *Geophys. Res. Lett.*, 18, 3, 1991b.
- Yung, Y. L., Lee, A. Y., Irion, F. W., DeMore, W. B., and Wen, J.: Carbon dioxide in the atmosphere: Isotopic exchange with ozone and its use as a tracer in the middle atmosphere, *Journal of Geophysical Research: Atmospheres*, 102, 10 857–10 866, 1997a.
- 700 Yung, Y. L., Lee, A. Y. T., Irion, F. W., DeMore, W. B., and Wen, J.: Carbon dioxide in the atmosphere: Isotopic exchange with ozone and its use as a tracer in the middle atmosphere, *J. Geophys. Res.*, 102, 9, 1997b.