

The Hydrological Archetypes of Wetlands

Abigail E. Robinson¹, Anna Scaini¹, Francisco J. Peña², Peter A. Hambäck³, Christoph Humborg¹, Fernando Jaramillo¹

¹Department of Physical Geography, Stockholm University, Stockholm, 106 91, Sweden

5 ²Division of Health Informatics and Logistics, KTH Royal Institute of Technology, Stockholm, 171 77, Sweden

³Department of Ecology, Environment and Plant Sciences, Stockholm University, Stockholm, 106 91 Sweden

Correspondence to: Abigail E. Robinson (abigail.robinson@natgeo.su.se)

Abstract

Wetlands are valuable and diverse environments that contribute to a vast range of ecosystem services, such as
10 flood control, drought resilience, and carbon sequestration. The provision of these ecosystem services depends on
their hydrological functioning, which refers to how water is stored and moved within a wetland environment. Since
the hydrological functions of wetlands vary widely based on location, wetland type, hydrological connectivity,
vegetation, and seasonality, there is no single approach to defining these functions. Consequently, accurately
identifying their hydrological functions to quantify ecosystem services remains challenging. To address this issue,
15 we investigate the hydrological regimes of wetlands, focusing on water extent, to better understand their
hydrological functions. We achieve this goal using Sentinel-1 SAR imagery and a self-supervised deep learning
model (DeepAqua) to predict surface water extent for 43 Ramsar sites in Sweden between 2020 and 2023. The
prediction results in wetlands grouped into five archetypes based on their hydrological similarity: 'spring-surgings',
'spring-flooded', 'summer-flooded', 'slow-drying' and 'summer-dry'. The archetypes represent great heterogeneity,
20 with flashy regimes being more prominent at higher latitudes and smoother regimes found preferentially in central
and southern Sweden. Additionally, many wetlands show exceptional similarity in the timing and duration of
flooding and drying events, which only became apparent when grouped. We attempt to link hydrological functions
to the archetypes whereby headwater wetlands, for example, we find that spring-surgings wetlands have the
potential to accentuate floods and droughts, while slow-drying wetlands, typical of floodplain wetlands, are more
25 likely to provide services such as flood attenuation and water storage during low flow conditions. Additionally,
although wetlands can be classified in myriad ways, we propose that classifying wetlands based on the hydrological
regime derived from water surface extent is useful for identifying hydrological functions specific to the site and
season, and when discharge or water depth data is not available. Lastly, we foresee that hydrological regime-based

classification can be easily applied to other wetland-rich landscapes to understand the hydrological functions better
30 and identify their respective ecosystem services.

1 Introduction

Wetlands are ecosystems that are seasonally or permanently covered by or saturated with water (Bullock and
Acreman, 2003). After centuries of wetland loss (Fluet-Chouinard et al., 2023), wetlands are now viewed as key
providers of provisioning and regulating services such as forestry, fishing, food production, flood control, drought
35 resilience, nutrient and sediment retention and carbon sequestration (Ameli and Creed, 2019; Barbier et al., 1997;
Colvin et al., 2019; Johnston, 1991; Matthew et al., 2010; Tang et al., 2020; Villa and Mitsch, 2015). Additionally,
they offer cultural and supporting services (Margaryan et al., 2022; Mitsch et al., 1991; Wood et al., 2024) and are
crucial for achieving the sustainable development goals outlined in Agenda 2030 (Jaramillo et al., 2019).

40 The degree to which wetland environments provide ecosystem services is largely controlled by their hydrological
functions (Okruszko et al., 2011) or how wetlands store and transfer water. For instance, hydrological functions
such as prolonged water storage contribute to services like flood control and sustaining water supply during low
flow periods (Åhlén et al., 2020; Bullock and Acreman, 2003; Gerakēs, 1992). Other functions, such as surface-
ground water exchange, relate to provisioning services such as water supply, while surface wetness and soil
45 moisture help regulate the local climate and retain nutrients (Ameli and Creed, 2017; Hansson et al., 2005; Le and
Kumar, 2014; Mitsch et al., 2015). Furthermore, large fluctuations of surface water extent is strongly correlated to
fluctuations of methane emissions for boreal wetlands (North of 50°N), which is important for services like carbon
sequestration (Ringeval et al., 2010).

50 Quantifying the hydrological functions of wetlands and the provision of ecosystem services is challenging as
wetlands are spatiotemporally variable and diverse (McLaughlin and Cohen, 2013). For example, a wetland type
can either reduce or enhance flooding downstream depending on the environmental setting or time of year (Bullock
and Acreman 2003). One way to improve our understanding of wetland hydrological functions and related
ecosystem services is by quantifying their hydrological regime. This refers to the seasonal availability of water
55 (water, extent, or volume) within a wetland, measured through either in-situ or remote sensing technologies
(Acreman and Holden, 2013; Helmschrot, 2016).

The analysis of hydrological regimes to understand hydrological functioning usually focuses on rivers and catchments (Magilligan and Nislow, 2005; Robinson and Sivapalan, 1997). However, over the last two decades, its application for wetlands has steadily increased (e.g., Cuevas et al., 2024; Stevaux et al., 2020; Na and Li, 2022; Vilardy et al., 2011). In fact, methods for studying water extent have been driven by the need to quantify ecosystem services (Park et al., 2022). For instance, by monitoring water level or extent, we can evaluate whether a wetland is in a water-storing or transmitting state, which influences its ability to attenuate high flows downstream (Spence et al., 2011; Yanfeng and Guangxin, 2021). Furthermore, analysis of the hydrological regimes based on water extent and level in Siberian wetlands has enhanced the understanding of how water availability in winter influences spring flooding (Zakharova et al., 2014). In Europe, Vera-Herrera et al. (2021) demonstrated that grouping wetlands based on their long-term changes in surface water extent can help to maximize agricultural productivity, while Åhlén et al (2022) distinguished between flood buffering capacity of wetlands in upland and downstream wetlands studying variations in water level.

When in-situ water level or discharge measurements from water gauges are spatiotemporally sparse, water surface extent can be used to understand the hydrological regime. Estimating hydrological regimes from water surface extent is achievable with remote sensing technologies, such as optical or Synthetic Aperture Radar (SAR) (Graversgaard et al., 2021; Ramsar Convention, 2011; Vera-Herrera et al., 2021). For example, multi-spectral optical sensors like Sentinel-2 can help estimate surface water extent at a resolution of 10 m (Brown et al., 2022). Others have exploited the ability of SAR to detect water below flooded vegetation in a range of wetland environments at similar resolutions (Canisius et al., 2019; Kovacs et al., 2013; Melack and Hess, 2011; Widhalm et al., 2015; Peña et al., 2024).

It is widely recognised that although ecosystem services are not undervalued, they are often poorly characterised and understood in the context of wetlands. Furthermore, generalising hydrological functions and services across different wetlands is not recommended due to their unique characteristics. Here, we quantify changes in water surface extent to understand the hydrological regimes of wetlands and determine their hydrological functions, using the case of Sweden. This study aims to categorise wetlands by their hydrological regime based on recent water surface extent observations using a remote sensing data and a pre-trained self-supervised deep learning model called DeepAqua (Peña et al., 2024). We use the case of 43 Ramsar wetlands as they are well inventoried, and present good spatiotemporal coverage of SAR data and are of national and international importance due to the ecosystem services they provide (Gunnarsson and Löfroth, 2014; Ramsar Convention, 2011). We propose that by

grouping hydrologically similar sites into descriptive archetypes (as suggested by Lane et al., 2018), more comprehensive insights can be gained about the hydrological regime (and thus functions) than by studying each wetland's hydrological regime in isolation.

2 Methods

2.1. Wetland dataset description

Sweden has 68 Ramsar wetlands in total (Ramsar Convention, 1971). Here, we excluded coastal sites because coastal wetlands are hydrologically different from inland wetlands and should therefore be studied separately. Sites with a total area exceeding 180,000 ha were also excluded due to computational and memory limitations when computing water extent changes with deep learning. Lastly, sites with low SAR data availability due to processing issues and the loss of Sentinel-1B in December 2021 were omitted from the analysis. This left 43 Ramsar sites suitable for hydrological regime analysis, and each site was delimited based on the boundaries of the Ramsar Convention (Ramsar Convention - Sweden, 2023) (Fig. 1).

The sites are distributed throughout all regions in Sweden, albeit with a higher concentration of sites in central and southern Sweden. Site areas range between 200 ha and 28,900 ha and encompass various wetland types, including marshes, fens, bogs, mires, palsa mires, lakes, streams, wetland forests, peatlands, and shrub wetlands. For these wetlands, during the observation period (2020-2023), the average temperature and precipitation were 5.76°C and 706.5 mm, which was 0.68°C warmer and 25.6 mm wetter on average compared to the 1990-2020 climate normal (Johansson, 2002). Additionally, the mean number of snow days in Sweden between 2020-2023 was 108.0, which is 12.3 days less compared to the last climate normal (Climate indicator - Snow, 2024). Daily precipitation from the Copernicus Climate Change Service E-OBS ensemble (0.1° grid) for each Ramsar site is available in Figs A7-11 (Cornes et al., 2018).




Figure 1. Spatial distribution of Ramsar wetland study sites (grey polygons) in terms of (a) Elevation from a 50m resolution DEM by *Landmäteriet* (grey thin lines denote main catchments), (b) Average precipitation in mm/yr and (c) Average temperature in °C between 2020 and 2023. Temperature and precipitation data were obtained from the Precipitation Temperature Hydrological Agency's Water Model (PTHBV), available at the Swedish Meteorological and Hydrological Institute (SMHI).

2.2. Wetland characteristics

To place the wetlands into an environmental context, we tabulated each site's latitude, elevation, open water as a percentage of the total area, and general wetland type. The elevation was calculated as the average elevation (m.a.s.l.) derived from the Digital Elevation Model 50m (*Markhöjdmodell Nedladdning*, grid 50+) (Lantmateriet, 2022) within the wetland boundary. Open water extent for each wetland was calculated for every month in 2023 using monthly composites of Normalised Difference Water Index (NDWI) binary (water/non-water) masks from Sentinel-2 optical imagery.

125 The wetland type was estimated using the following databases of wetland classification: (1) The Ramsar
Convention database for sites in Sweden, (2) the National Wetland Inventory for Sweden (VMI) (Gunnarsson and
Löfroth, 2009), and (3) an updated satellite-based open wetland mapping classification from 2018-2022 (Hahn and
Wester, 2023). Each wetland was assigned a generic wetland class adapted from Gunnarsson and Löfroth (2014):
'open', 'limnic', 'mixed' or 'mire'. 'Open' refers to meadows, grasslands, and temporarily flooded land, 'limnic' refers
130 to lake shores, beaches by watercourses, overgrown lakes and limnogeneous beach complexes. 'Mixed' wetlands
are regarded as a combination of multiple wetland types and may include different mires with open or limnic
wetland environments. A 'mire' wetland consists primarily of bogs and fens. A fifth wetland type, '*fjäll*' (mountain),
was assigned to wetlands located in Sweden's mountainous regions as they are not classified in the datasets.

2.3. Hydrological regime given by water surface extent analysis

135 We estimated the hydrological regime from water extent using an automated approach based on remote sensing
data. Automatic surface water detection was done with a deep-learning image segmentation model called
DeepAqua (Peña et al., 2024). DeepAqua is a self-supervised model with the principal function of detecting surface
water extent in wetlands from Sentinel-1 SAR imagery in the VH polarisation. DeepAqua can detect both open
and vegetated water using the C-band SAR sensor onboard Sentinel-1, which can penetrate some types of perennial
140 vegetation due to its emission of longer wavelength radar waves (5.6 cm) (Adeli et al., 2021). Usually, semantic
segmentation models require manually labelled images as their training label output. With DeepAqua however,
the training labels are binary images (water/non-water) of the NDWI based on cloud-free Sentinel-2 optical
imagery of the same location and time as the input training data (SAR imagery), since both missions have a ~1
week repeat cycle over Sweden (~1-2 passes per week between 2020 and 2022, after which spatiotemporal
145 coverage is reduced to ~10-12 days due to the failure of the Sentinel-1B satellite). We use a **pre-trained** version
of the DeepAqua model for our analysis, which was trained on a Sentinel-1 and Sentinel-2-based NDWI binary
image over central Sweden from the 5th June 2018. When the pre-trained model was tested on three wetlands in
Sweden (Peña et al., 2024), DeepAqua outperformed existing land classification models such as Dynamic World
(Brown et al., 2022) and thresholding techniques such as Otsu (Otsu, 1979) on multiple evaluation metrics such
150 as pixel accuracy, intersection over union, precision and F1.

The output predictions comprised polygonised binary water/non-water images for every Sentinel-1 image available
between January 2020 and August 2023 cropped to within the boundaries of each wetland. The total water area
for each image was calculated based on the WGS84 UTM Zone 33N projection (Figures A2-A6). The monthly

155 average of water extent between January 2020 and August 2023 was calculated to reduce the risk of annual
variability affecting potential clustering while aiming to detect hydrological regimes under 'average' conditions.
Due to extensive snow and ice cover complicating the water extent predictions, winter months (November,
December, January, and February) were removed from the hydrological regime analysis.

160 Lastly, due to the lack of ground truth data on temporally dynamic wetland water extent within our Ramsar sites,
we validate our water extent predictions using two alternative approaches. Firstly, we compare DeepAqua's
predicted water extent with manually delineated water extent derived from Sentinel-1 SAR imagery in the VH
polarisation for a systematic sample of wetlands for all available images during 2021. We randomly select one
wetland within each of the resulting archetypes to get a representative yet unbiased sample of wetlands. For the
165 second approach, we assess the accuracy of the predicted hydrological regimes by comparing them to daily
discharge data from nearby active stations provided by Global Runoff Data Centre (GRDC) and SMHI. In total,
there were 23 sites with available discharge data either upstream, downstream or on site of the wetland. For both
approaches, we calculate the error between the DeepAqua predictions to (1) manually delineated water extent and
(2) daily discharge using the normalised root mean square error (NRMSE). We normalise the root mean square
170 error (RMSE) to the range of water extent as to discount total area from the error result and to make each wetland
comparable with the others.

2.4. Cluster Analysis

The hydrological regimes based on DeepAqua's water extent predictions (Section 2.3) were clustered based on
their hydrological similarity using a multivariate K-means cluster analysis technique and means of visual
175 interpretation. K-means clustering is a widely used and simple unsupervised machine learning technique in which
groups are identified based on the Euclidean distance between a data point and a centroid (a mean of the data)
(Everitt et al., 2011). In order to conduct a cluster analysis, data points that characterise the hydrological regime
given by water extent are required. We calculated several hydrological parameters based on each hydrological
regime and used them as the input data points (Table A1). The hydrological parameters included known
180 hydrological signatures (Olden and Poff, 2003) and custom parameters to describe the hydrological regime in
terms of duration, timing, frequency, magnitude, and rate of change. The optimal number of clusters (k) was chosen
based on the inflection point on the Elbow Curve, which calculates the within-cluster-sum-of-squares (WCSS) for
a range of cluster sizes from 1 to n. The inflection point on the Elbow Curve is interpreted at the optimum number
of clusters since it indicates the point where adding more clusters results in a diminishing reduction in WCSS. The

185 best-performing parameters were picked using visual inspection (inspecting their ability to cluster the regimes)
and validated against multicollinearity using the Variance Inflation Factor (VIF). The VIF measures the degree of
multicollinearity of one hydrological parameter with all other parameters by calculating how much the variance of
the regression coefficient increases due to correlation with other independent variables. We recognise that there is
some degree of inherent correlation between the hydrological parameters since they are descriptors of the same
190 hydrological regime. Therefore, we used a VIF value of <10 as an indicator that the hydrological parameters were
not highly multicollinear and did not describe the same regime characteristic (Figure 5a).

The emerging pattern given by the Elbow Curve indicated that individual hydrological regimes among wetlands
were best grouped when $k = 4-6$ (Figure A1). Upon visual inspection, $k = 5$ was chosen as the best possible
195 distribution of wetlands into roughly equal-sized groups. The number of sites in each cluster ranged between 6 and
15. Each hydrological parameter was tested individually and in combination with other parameters to see how
effectively they helped cluster the wetlands. Certain variables, such as the maximum month, dominated the
clustering over other indices and some indices-pairs were extremely collinear, such as maximum month and
minimum month, or Spring/Summer slope difference and slope variation. Therefore, these pairs could not be used
200 together for the final clustering analysis.

3 Results and Analysis

3.1. Surface water extent validation

When comparing water extent predictions from DeepAqua to manually delineated water extent to a systematic
sample of wetlands, we find that predicted water extent performs well with their manually delineated counterparts
205 (Fig. 2). *Hjälstaviken* and *Dättern* wetlands had the lowest NRMSE with 0.04 and 0.07, respectively, whereas
Maanavuoma wetland exhibited the highest error between the manually delineated water extent and the DeepAqua
prediction with a NRMSE of 0.12. The majority of error between the DeepAqua's and the manual water extent
estimates originates from the spring and autumn months for many of the sampled wetlands. This is particularly
apparent in *Maanavuoma* and *Tysöarna* wetlands. In both cases, the water extent is underestimated by DeepAqua
210 compared to the manual estimate. In *Store mosse* wetland, DeepAqua tends to overestimate wetland water extent





Figure 2. (a-e) Comparison between monthly water surface extent from DeepAqua predictions and manual delineation in 2021. (f) Values of Normalised Root Mean Square Error (NRMSE; RSME divided by the range in wetland extent) between manually delineated and DeepAqua predictions.

215 compared to when the water extent is manually delineated. Overall, all five sampled sites have strong agreement in the shape and magnitude of the hydrological regime, indicating that DeepAqua captures the seasonal hydrological characteristics with good accuracy (Fig. 2f).

To enhance the strength of our validation approach, we compared the wetland hydrological regimes to in-situ daily
 220 discharge measurements. Among the 23 wetlands with available discharge data, three had an active gauging station located upstream, two had onsite stations and sixteen had stations situated downstream (Fig. 3a). Of these, eight sites featured regulatory structures (e.g., dams, weirs, or culverts) along their river courses, which may disrupt the natural flow regime and weaken the correlation between wetland water surface extent and stream discharge. In general, stations with lower mean discharge returned lower NRMSE values between water extent and discharge
 225 (Fig. 3b). However, the relationship is weak ($R^2 = 0.17$) and based on a limited number of observations ($n = 23$). Most sites cluster in the bottom-left portion of the plot, with a few high-discharge, high-NRMSE outliers in the

top-right. Regulated and non-regulated sites are distributed throughout, with no strong visual separation, although none of the regulated sites exhibit low discharge-low NRMSE values.



230 **Figure 3. (a) – NRMSE between daily discharge and wetland water extent for the 23 wetlands with available discharge data. Green boxes indicate the interquartile range (IQR), whiskers represent the range, and orange lines show the mean NRMSE. (b) Mean NRMSE versus mean discharge for each wetland, calculated over matching dates from January 2020 to August 2023. Wetlands with regulated flow paths between the wetland pour point(s) and discharge station are indicated by black outlines.**




Fig. 4 presents a sample of wetlands with discharge data either upstream or downstream and unregulated flow. In general, daily discharge replicates the shape of the wetland's hydrological regime. The correlation between river discharge and wetland hydrological regime is particularly apparent for *Tjålmejaure-Laisdalen* (NRMSE 39.49), *Östen* (NRMSE 31.40) and *Helge å* (NRMSE 12.70) wetlands, whereby increased discharge matches well with increased water extent in the spring months, followed by relatively reduced flow thereafter.

Figure 4. Left panel: Comparison of water surface extent and discharge from on-site, upstream or downstream stations for corresponding dates in *Maanavuoma*, *Storkölen*, *Tjålmejaure-Laisdalen*, *Östen* and *Helge å* wetlands from January 2020 to August 2023 (excluding winter months). The GRDC station IDs are shown in the upper left of each plot. **Right panel:** Wetland boundaries (green polygons) as defined by the Ramsar Convention, with discharge stations (black rings), watercourses between station and wetland (thick blue) and other watercourses (thin blue).

Although *Tjålmejaure-Laisdalen* and its' corresponding downstream station are separated by ~116 km of watercourses, the discharge data agrees well with the wetland water extent. For *Maanavuoma* wetland (NRMSE 0.92), data from the discharge station situated ~15 km upstream agrees with water surface extent in 2020 and 2021, however, the spring surge of water in 2022 and 2023 that is present in the river is not experienced by the wetland.

Lastly, they also agree well in *Storkölen* wetland (NRMSE 9.37) despite greater interannual variability compared to other sites. Notably, both time series show a pronounced peak between April and May 2021, reflecting a concurrent increase in wetland water extent.

3.2. Cluster Analysis

275 Based on the surface water extent data, we conducted a cluster analysis to explore patterns in the shape and dynamics of wetland hydrological regimes. From all parameters assessed, skewness, kurtosis, normalised maximum slope, number of peaks and baseline month fraction (Fig. 5a) were found to collectively capture key regime characteristics (Fig. 5b). Upon visual inspection, regimes with similar shapes were grouped together while also maintaining the desired VIF condition (<10) with values of 3.96, 1.60, 4.07, 3.01, and 6.54 for skewness, 280 kurtosis, maximum slope, number of peaks and baseline month fraction, respectively. These values indicate a reasonable level of non-multicollinearity between all other variables. The chosen parameter combination successfully clusters related hydrological regimes into five different archetypes, with the number of sites (n) in each archetype as follows: '*spring surging*' (n=6), '*spring flooded*' (n=8), '*summer flooded*' (n=8), '*slow drying*' (n=15) and '*summer dry*' (n=6). Support for the archetype names is given by the hydrological parameter results 285 which have been averaged by the archetype and are described in Section 3.3.





Figure 5. (a) Overview of the chosen parameter (unitless) combination (averaged by archetype) used for the final cluster analysis of the hydrological regimes given by water extent and the VIF value for each parameter. (b) Graphical representation of the five selected hydrological parameters used to describe the characteristics of the hydrological regime for the final cluster analysis. (c-g) Radar plots for for final hydrological parameters averaged by archetype.

3.3. Hydrological archetype analysis

The overall spatial distribution of the archetypes and thematic graphic descriptions of the hydrological regime given by water surface extent is presented in Fig. 6. The spring-surging (Fig. 6a) are only found in northern Sweden and have flashy hydrological regimes, consisting of a dry baseline condition and a brief period of increased water extent. Spring-flooded wetlands (Fig. 6b) are limited to southern and central Sweden. The hydrological regime of



these wetlands resembles that of spring-surging wetlands, although they have a relatively longer spring peak. Summer-flooded wetlands (Fig. 6c) remain inundated from May to October after a rapid wetting period and are spread across Sweden. Southern Sweden's slow-drying wetlands (Fig. 6d) exhibit steadily decreasing water extent throughout the summer, reaching minimum water extent in autumn. Lastly, summer-dry wetlands (Fig. 6e) exhibit the maximum wetland extent in April, preceding generally dry conditions until September-October.

Figure 6. Spatial distribution of hydrological archetypes for sampled Ramsar wetlands in Sweden (n=43) and representation of their hydrological regime through March and October; (a) Spring surging wetlands (n=6), (b) Spring flooded wetlands (n=8), (c) Summer flooded wetlands (n=8), (d) Slow drying wetlands (n=15) and (e) Summer dry wetlands (n=6).

One of the most distinctive differences between archetypes is the magnitude of water extent at the beginning of Spring. For instance, slow-drying and summer-dry archetypes already have large water extents in March and, therefore, do not undergo a rapidly


325 inundating period during Spring or Summer. The lack of any inundation period is reflected in the normalised
maximum slope values (Fig. 5f,g), which are the lowest out of all archetypes, suggesting smaller changes in water
extent across the year (0.21, and 0.14 for summer-dry and slow-drying, respectively). Additionally, archetypes
with large water extent in Spring tend to be found in central and southern Sweden, while archetypes such as spring-
surging and summer-flooded wetlands start with a small water extent in March preceding a rapid inundation period.
330 These archetypes, with higher normalised maximum slope values of 0.59 and 0.77, respectively, are more abundant
in the north (Fig. 5c,e).

A second defining feature between different archetypes is the duration of the dry period (baseline fraction), defined
by months with water extent within the 25th percentile of the range. Archetypes with a significant dry period, such
335 as summer-dry, spring-surging and slow-drying wetlands, have high baseline month fractions (0.65, 0.63 and
0.66, respectively) and positive skewness (1.14, 1.45 and 1.58 respectively), which indicates that wet conditions
are limited to the spring months (Fig. 5g,c,f). Conversely, with a negative skewness and low baseline month
fraction (-1.60 and 0.17, respectively; Fig. 5e), summer-flooded wetlands are the only archetype that retains its
large water extent throughout the year.

340

The resulting archetypes show how wetland hydrological regimes can be broadly differentiated into two primary
'modes': *peaky* and *smooth*. We define *peaky* regimes as those with large fluctuations in water extent, while *smooth*
regimes follow more consistent, gradual changes in monthly water extent. Peaky archetypes, such as spring-
surging (Fig. 7a) and summer-flooded wetlands (Fig. 7c), exhibit relatively high values of kurtosis (2.27 and 2.93,
345 respectively), maximum slope (0.59 and 0.77, respectively), and the number of peaks (1.2 and 1.0, respectively).
On the other hand, *smooth* archetypes, like slow-drying and summer-dry wetlands are characterised by relatively
stable water extent from March to October (Fig. 7d,e). Spring-flooded wetlands share some traits with peaky
archetypes, particularly a marked increase in water extent during spring (Fig. 7b) and high normalised slope values
(0.70). However, they differ from typical spring-or summer-flooded wetlands in having a low average kurtosis (-
350 0.04), which suggests a more even distribution of water extent over time. Although we refer to peaky archetypes
here, it is important to note that the number of peaks is not necessarily descriptive of just peakedness (kurtosis).
For instance, slow-drying wetlands have high kurtosis (2.03) yet few peaks on average (0.2), indicating that
although they experience large variability in water extent, there is no distinguishable wet month.


Figure 7. Hydrological regimes of individual wetlands per archetype based on a monthly average of surface water extent between January 2020 and August 2023. The water extent area for each month is shown relative to the water extent area in March.



Another approach to interpreting archetypes is by examining the degree of homogeneity within each archetype. This is because some archetypes share more similarities in terms of their environmental characteristics and hydrological regimes. For instance, summer-dry wetlands are mostly comprised of mires or open wetlands (Fig. 8d), typically lying at low elevations and exhibiting similar hydrological regimes (Fig

7e). Spring-surging wetlands are also considered a homogenous archetype, since they are located primarily in high latitude regions (Fig. 8a), are mainly *fjäll* wetlands, and tend to have little variability in their hydrological regime (Fig. 7a). In contrast, spring-flooded and summer-flooded wetlands are found all over Sweden, across a range of elevations (Fig. 8b) and encompass many different wetland types. This highlights that hydrological regimes are not always associated with a specific wetland type, but rather depend on the broader archetype to which the wetland belongs.

Despite the varying degrees of diversity within archetypes, grouping wetlands into archetypes still reveals a remarkable similarity in the timing of key features of their hydrological regimes. For instance, most summer-flooded wetlands reach low water extent by May or June, despite varying rates of drying for the rest of the year. This indicates that the hydrological parameters correctly capture timing characteristics, even across archetypes with with more heterogeneity, such as summer-flooded wetlands.



390 **Figure 8. (a-c) Wetland topographical and ecological characteristics per archetype. The boxes represent the interquartile range (IQR), with orange lines indicating the mean value across all wetlands within each archetype. Whiskers outline the range, and small black circles denote wetlands with anomalous results compared to the rest of the archetype. (d) Stacked bar plot showing the occurrence of wetland types (*fjäll*, limnic, mire, mixed or open) per cluster as a percentage of the total number of sites in each archetype.**

4 Discussion

4.1. The value of archetypes for understanding wetland hydrology

One of the defining features for most archetypes was the timing of large changes in surface water extent, which only became apparent when the sites were grouped into archetypes. This highlights the usefulness of employing archetypes in hydrological studies, as hydrological regimes may not be best evaluated across sites when using a single parameter (Cutler and and Breiman, 1994; Huggins et al., 2024; Piemontese et al., 2020). Although our classification was based solely on water dynamics, it also inevitably captured the cumulative effects of other environmental factors, such as vegetation, soil type and climate. The archetype approach to classification is further

supported by Bullock and Acreman (2003), who concluded that grouping wetlands based on their local classification term is less intuitive than grouping them by hydrological characteristics. This suggests that the hydrological perspective is a valuable lens for understanding ecosystem services of wetlands, especially when complemented with other environmental data (Okruszko et al., 2011; Poff et al., 1997).

However, despite the overall success of the classification, not all wetlands were easily categorised. We suggest that there are two main reasons why some wetlands were difficult to assign to archetypes. Firstly, the indistinct nature of some wetlands suggests that some hydrological regimes can sometimes be seen as a continuum rather than easily separated categories, making it challenging to group them into distinct archetypes. Secondly, the limited scope of the wetland database used for clustering might have excluded the existence of additional archetypes that can be obtained when focusing on the hydrological regime from water extent changes. It is also important to note that while we defined archetypes using an average of ~four years of monthly water extent data, these only reflect the observed period. Longer observational periods are necessary for determining extended trends and the impact of changing climatological conditions. Since the DeepAqua model we used for water extent predictions was trained to predict water extent on SAR scenes dating between January 2020 and August 2023, we were not able to extend our temporal scope outside of this range. Therefore, we suggest developing any future training of the DeepAqua model so that it is more generalisable to longer time periods and less sensitive to changes in Sentinel-1 SAR pre-processing.

4.2. Methodological considerations

Using water extent as our key measurement, SAR imagery provided data with dense spatiotemporal resolution across 43 wetland sites, which can be applied to any wetland larger than 200 ha. The reliance on remote sensing is driven by a lack of in-situ data, which would have partly or wholly missed the hydrological regime signatures for most of the chosen wetlands in this study.

To efficiently process large volumes of remotely sensed data, we chose an automatic deep learning-based approach (DeepAqua) to detect water extent without the need for manual annotation. SAR is particularly useful for wetland studies since most water goes undetected by optical imagery (Sahour et al., 2021). However, it is likely that C-band radar still underestimates surface water extent in wetlands since its relatively shorter wavelength limits penetration into dense vegetation (Adeli et al., 2021). Additionally, we assume that surface water extent is analogous to total water storage, which may not be true for mire types (Acreman and Holden, 2013) or

topographically constrained wetlands. Therefore, including water level data from hydrogeodetic technologies such as water levels from the Surface Water and Ocean Topography (SWOT) mission (Hamoudzadeh et al., 2024) or soil moisture observations (Mupepi et al., 2024) could improve hydrological regime classification, especially for seasonal wetlands (see more examples in Jaramillo et al., 2024).

A further challenge arises when observing wetland water extent in winter. Snow and ice interfere with SAR-based water detection methods, which leave winter hydrology poorly observed. The issue is further compounded by the fact that there are few discharge stations available to fill these observational gaps. Moreover, even where discharge data is available, they can be affected by hydrological barriers or complex flow paths in higher-order streams, limiting their utility for validating wetland hydrological regimes.

4.3. Controls and variability in wetland hydrological behaviour

Although detailed exploration of the physical drivers of the observed hydrological regimes is beyond this study's scope, we theorise that factors such as position within the watershed and surface connectivity contribute to at least some extent. For example, spring-surgling wetlands, with few surface water inlets, rely mainly on snowmelt and tend to dry rapidly, while summer-flooded wetlands benefit from multiple inflows and sustain inundation longer (Lane et al., 2018). Secondly, wetlands located in headwater regions, like spring-surgling and summer-flooded wetlands, experience rapid flood peaks characteristics of upper catchment water flows. This is in contrast to wetlands such as those within the slow-drying archetype, which are located in the lower parts of the catchment, and are therefore linked to less pronounced flood peaks (Morley et al., 2011).

It should also be acknowledged that hydroclimatic variability plays a critical role in shaping wetland hydrological regimes and represents an important consideration to the interpretation of our archetypes. For instance, on an interannual and seasonal temporal scale, fluctuations in precipitation, snowmelt and evapotranspiration strongly influence wetland hydroperiods (Jaramillo et al., 2018; Winter, 2000). On a climatic temporal scale, warming trends and increasing dryness index has been observed in Swedish wetlandscapes since the 1970s, suggesting that there is a greater evaporative demand and reduced water storage in wetlands, especially during summer (Åhlén et al., 2021). These observations also align with model projections showing substantial summer drying and reduced wetland extent in North America under high-emission scenarios due to evapotranspiration exceeding precipitation input (Xu et al., 2024). Similarly, Xi et al., 2021 projected future declines in inland wetland area across Europe, though with a higher degree of uncertainty in Scandinavia. Furthermore, hydrological stability will likely be

reduced in the future, with modelled studies of prairie pothole wetlands showing diminished monthly-scale stability in water storage under uncertain climate conditions (Zhang et al., 2011).

Despite this, the degree to which wetlands are vulnerable to such changes is dependent on their dominant water sources and topographical setting. For example, wetlands that are reliant on direct precipitation or snowmelt, such as spring-surgant wetlands, are more sensitive to hydroclimatic variability, while wetlands sustained by regional groundwater inputs on larger floodplains (like slow-drying or spring-flooded wetlands), have greater buffering capacity to hydroclimatic change (Winter, 2000). These findings highlight the need for long-term observations and the integration of hydroclimatic data when interpreting wetland hydrology in future work.

4.4. Hydrological regimes as indications of ecosystem services

In this study, we quantified the hydrological regimes of Swedish wetlands to better understand their hydrological functions, which are closely tied to the ecosystem services they provide. Inland wetlands are estimated to contribute approximately US\$27 trillion annually in ecosystem service value, with the majority of the value deriving from water regulating services (Davidson et al., 2019).

We theorise that hydrological regimes can serve as indicators of the hydrological ecosystem services a wetland may deliver at any given time. For instance, spring-surgant wetlands, which are characterised by high water extent during spring and low extent during summer, resemble headwater wetlands. Headwater wetlands are known to increase flood flows during the wet season while retaining baseflow during the dry season (Bullock and Acreman, 2003), suggesting that these wetlands may not provide flood mitigation services but rather exacerbate flooding (Åhlén et al., 2022). This is supported by the Ramsar site descriptions, where no wetlands in the spring surging archetype list flood control as an important service. Nevertheless, headwater wetlands can offer temporary flood storage (Kadykalo and Findlay, 2016), although confirming these dynamics requires temporally dense water extent observations to capture lag times between water storage and downstream flows.

Conversely, slow-drying wetlands exhibit traits more typical of floodplain wetlands, which are well-documented in their role in flood reduction and water retention (Acreman and Holden, 2013; Golden et al., 2021). The gradual reduction of water extent in these wetlands may suggest sustained water storage, likely contributing to both flood peak attenuation and maintaining summer baseflows. This is supported by the Ramsar site descriptions for the wetlands belonging to the slow-drying archetype, since the majority of them have flood control and/or water

storage listed as a known ecosystem service. Additionally, Doherty et al., 2014 suggest that wetlands with periodically dry soils (such as slow-drying or summer-dry wetlands) slow down flows and can remove large volumes of water from the system. Although we did not perform a detailed analysis of ecosystem service delivery or have dense downstream discharge data (Andersson, 2012), our results offer a foundation for prioritising wetlands for future conservation or Ramsar designation, particularly in flood-prone or drought-prone regions.

Another strength of hydrological regime classification is its ability to infer hydrological functions at different times of the year, recognising that wetland functions are not static in time or space (Spence et al., 2011). For example, variability in water extent can signal the transition between water storage and runoff-dominated states (Yanfeng and Guangxin, 2021). Flashy water extent variability observed in spring-surfing, spring-flooded and to a lesser extent, summer-flooded wetlands, suggests a switch to conditions where wetlands act as conduits rather than reservoirs. This may result from frozen ground hindering water storage in soils (Yanfeng and Guangxin, 2021) or the dominance of rapid snowmelt inputs (Spence et al., 2011). However, further investigation combining water level, connectivity analyses and catchment precipitation data would be needed to verify these hypotheses.

Aside from hydrological-related ecosystem services, wetlands offer other valuable ecosystem services that are also linked to their hydrological regimes, such as biodiversity and carbon sequestration (Okrusko et al., 2011). Hydrological variability is a major driver of wetland biodiversity due to species' water tolerance thresholds. Additionally, wetlands classified under the 'northern' archetypes are particularly significant carbon sinks, as evidenced in Ramsar site records. Differentiating hydrological regimes in carbon-sequestering wetlands or those with particularly rich biodiversity could improve our understanding of their role in the delivery of other ecosystem services (Kirpotin et al., 2011).

5 Conclusion

This research aimed to improve our understanding of wetlands by revealing their hydrological regimes using remotely sensed data on water surface extent. We chose an automatic detection method based on Sentinel-1 SAR imagery because it can operate in cloudy and dark conditions and detect more water under vegetation compared to optical-based methods. The hydrological regimes were grouped based on similar hydrological characteristics identified by custom hydrological parameters. For 43 Ramsar sites in Sweden, the hydrological regimes based on monthly water extent between 2020 and 2023 could be grouped into five distinct archetypes. The defining traits were mainly related to the timing of change and the duration of wet and dry periods. Despite heterogeneity in the

540 archetypes' spatial distribution, flashy archetypes with high water extent variability were preferentially found at higher elevations and latitudes, while less variable and drier archetypes were concentrated towards low elevations and latitudes. Additionally, wetlands with mire were more likely to be part of the same archetype compared to open or limnic wetland types.


545 While contextual information is vital for our deeper understanding of wetlands, valuable insights into runoff and storage dynamics can be gained simply by tracking water extent over time. Furthermore, by reducing multiple wetland hydrological characteristics to the hydrological regime, we demonstrated that we could use the concept of archetypes to infer information about their specific hydrological functionality nationwide. Since many archetypes consist of multiple wetland classifications, we recommend estimating hydrological functions based on the

550 hydrological regimes, not individual wetland types. By being able to draw information from the archetypes, we reveal a new understanding of the hydrological functioning of wetlands with a particular emphasis on hydrological-related regulating ecosystem services such as flood control and water supply during low flow periods.


Appendix A

555 **Table A1. Hydrological parameters used for cluster analysis. Each parameter was evaluated individually and in combination with others to assess its effectiveness in capturing the characteristics of the hydrological regime. (N) – Normalized to remove the effect of wetland size.**

Hydrological parameters		Description
Max Month		Timing of the highest water extent
Min Month		Timing of the lowest water extent
Standard Deviation		Measure of dispersion of water extent values in a dataset
Skewness		Measure of symmetry in a distribution of water extent values
Kurtosis		Measure of peakedness in a distribution of water extent values
Coefficient of Variation		Measure of the dispersion water extent values around the mean
Range (N)		Difference between the maximum water extent value and the minimum water extent value, normalised to the mean wetland size
Minimum slope (N)		Smallest slope of monthly water extent change taken from the first derivative, normalised to the water extent range
Maximum slope (N)		Highest slope of monthly water extent change taken from the first derivative and normalised to the water extent range
Spring/Summer Difference (N)	Area	Difference between the average spring water extent (in March, April and May) and average summer water extent (June, July, August), normalised to the mean wetland size
Spring/Summer Difference (N)	Slope	Difference between the average spring slope of monthly water extent change (in March, April and May) and average summer slope of monthly water extent change (June, July and August), normalised to the mean wetland size
Slope Variation (N)		Standard deviation of all month-to-month slopes of monthly water extent change, normalised to the water extent range
Number of Peaks		Number of peaks, defined as a relatively high value of water extent between two relatively low values of water extent
Baseline Month Fraction		Number of months within 25 th percentile of the distribution of water extent values as a fraction of the year



560 **Figure A1. Elbow curve showing the within cluster sum of squares (WCSS) for k values ranging from 1-10. The Elbow Curve helps identify the number of clusters by indicating where adding more clusters result in a diminishing reduction in the WCSS.**



565 **Figure A2.** Average monthly water extent (March-October) between 2020-2023 for all wetlands belonging to the spring-surging archetype. Grey area shows the monthly interannual variability given by the range of water extent from all years. The monthly standard deviation is given in the top right bar plots.




Figure A3. Average monthly water extent (March-October) between 2020-2023 for all wetlands belonging to the spring-flooded archetype. Grey area shows the monthly interannual variability given by the range of water extent from all years. The monthly standard deviation is given in the top right bar plots.




Figure A4. Average monthly water extent (March-October) between 2020-2023 for all wetlands belonging to the summer-flooded archetype. Grey area shows the monthly interannual variability given by the range of water extent from all years. The monthly standard deviation is given in the top right bar plots.

Figure A5. Average monthly water extent (March-October) between 2020-2023 for all wetlands belonging to the slow-drying archetype. Grey area shows the monthly interannual variability given by the range of water extent from all years. The monthly standard deviation is given in the top right bar plots.




Figure A6. Average monthly water extent (March-October) between 2020-2023 for all wetlands belonging to the summer-dry archetype. Grey area shows the monthly interannual variability given by the range of water extent from all years. The monthly standard deviation is given in the top right bar plots.

Spring surging wetlands




Figure A7. Wetland water extent from January 2020 to August 2023 (excluding January, February November and December) for spring-surging wetlands, shown alongside daily precipitation totals for matching dates. Precipitation is aggregated separately for each wetland’s catchment and Ramsar area.

Spring flooded wetlands




Figure A8. Wetland water extent from January 2020 to August 2023 (excluding January, February November and December) for spring-flooded wetlands, shown alongside daily precipitation totals for matching dates. Precipitation is aggregated separately for each wetland’s catchment and Ramsar area.

Summer flooded wetlands






Figure A9. Wetland water extent from January 2020 to August 2023 (excluding January, February November and December) for summer-flooded wetlands, shown alongside daily precipitation totals for matching dates. Precipitation is aggregated separately for each wetland’s catchment and Ramsar area.

Slow drying wetlands



600 **Figure A10. Wetland water extent from January 2020 to August 2023 (excluding January, February November and December) for slow-drying wetlands, shown alongside daily precipitation totals for matching dates. Precipitation is aggregated separately for each wetland’s catchment and Ramsar area.**



605 **Figure A11. Wetland water extent from January 2020 to August 2023 (excluding January, February November and December) for summer-drying wetlands, shown alongside daily precipitation totals for matching dates. Precipitation is aggregated separately for each wetland’s catchment and Ramsar area.**

Code and data availability. All data including, environmental data, hydrological parameter results and water extent data for all wetlands is available through (Robinson, 2024) (<https://doi.org/10.5281/zenodo.13833605>). Code for processing data and cluster analysis is available at https://github.com/ab-e-rob/hydrological_archetypes. Code for predicting water extent in wetlands using DeepAqua can be found at <https://github.com/melqkiades/deep-wetlands>.

Author contributions. AR conceptualised the project, developed the methodology, conducted data collection and analysis. AS and FP contributed to methodology development and provided research guidance. PH assisted in results interpretation. FJ provided guidance on methodology, data analysis, and interpretation; supervised the

project. The co-authors all contributed to the preparation of the paper. The use of ChatGPT helped to improve prose.

Competing interests. The authors declare that they have no conflict of interest.

Financial support. This project was funded by Svenska Forskningsrådet Formas (Formas) Project 2022-01570.

620 **References**

- Acreman, M. and Holden, J.: How Wetlands Affect Floods, *Wetlands*, 33, 773–786, <https://doi.org/10.1007/s13157-013-0473-2>, 2013.
- Adeli, S., Salehi, B., Mahdianpari, M., Quackenbush, L. J., and Chapman, B.: Moving Toward L-Band NASA-ISRO SAR Mission (NISAR) Dense Time Series: Multipolarization Object-Based Classification of Wetlands
625 Using Two Machine Learning Algorithms, *Earth and Space Science*, 8, <https://doi.org/10.1029/2021EA001742>, 2021.
- Åhlén, I., Hambäck, P., Thorslund, J., Frampton, A., Destouni, G., and Jarsjö, J.: Wetlandscape size thresholds for ecosystem service delivery: Evidence from the Norrström drainage basin, Sweden, *Science of The Total Environment*, 704, 135452, <https://doi.org/10.1016/j.scitotenv.2019.135452>, 2020.
- 630 Åhlén, I., Vigouroux, G., Destouni, G., Pietroni, J., Ghajarnia, N., Anaya, J., Blanco, J., Borja, S., Chalov, S., Chun, K. P., Clerici, N., Desormeaux, A., Girard, P., Gorelits, O., Hansen, A., Jaramillo, F., Kalantari, Z., Labbaci, A., Licero-Villanueva, L., Livsey, J., Maneas, G., Pisarello, K. L. M., Pahani, D. M., Palomino-Ángel, S., Price, R., Ricaurte-Villota, C., Fernanda Ricaurte, L., Rivera-Monroy, V. H., Rodriguez, A., Rodriguez, E., Salgado, J., Sannel, B., Seifollahi-Aghmiuni, S., Simard, M., Sjöberg, Y., Terskii, P., Thorslund, J., Zamora, D. A., and Jarsjö,
635 J.: Hydro-climatic changes of wetlandscapes across the world, *Sci Rep*, 11, 2754, <https://doi.org/10.1038/s41598-021-81137-3>, 2021.
- Åhlén, I., Thorslund, J., Hambäck, P., Destouni, G., and Jarsjö, J.: Wetland position in the landscape: Impact on water storage and flood buffering, *Ecohydrology*, 15, e2458, <https://doi.org/10.1002/eco.2458>, 2022.
- Ameli, A. A. and Creed, I. F.: Quantifying hydrologic connectivity of wetlands to surface water systems,
640 *Hydrology and Earth System Sciences*, 21, 1791–1808, <https://doi.org/10.5194/hess-21-1791-2017>, 2017.
- Ameli, A. A. and Creed, I. F.: Does Wetland Location Matter When Managing Wetlands for Watershed-Scale Flood and Drought Resilience?, *JAWRA Journal of the American Water Resources Association*, 55, 529–542, <https://doi.org/10.1111/1752-1688.12737>, 2019.
- Andersson, K.: Multifunctional Wetlands and Stakeholder Engagement: Lessons from Sweden, Stockholm Environment Institute, 2012.
- 645

Barbier, E. B., Acreman, M., and Knowler, D.: Economic valuation of wetlands: a guide for policy makers and planners, 1997.

650 Brown, C. F., Brumby, S. P., Guzder-Williams, B., Birch, T., Hyde, S. B., Mazzariello, J., Czerwinski, W., Pasquarella, V. J., Haertel, R., Ilyushchenko, S., Schwehr, K., Weisse, M., Stolle, F., Hanson, C., Guinan, O., Moore, R., and Tait, A. M.: Dynamic World, Near real-time global 10 m land use land cover mapping, *Sci Data*, 9, 251, <https://doi.org/10.1038/s41597-022-01307-4>, 2022.

Bullock, A. and Acreman, M.: The role of wetlands in the hydrological cycle, *Hydrology and Earth System Sciences*, 7, 358–389, <https://doi.org/10.5194/hess-7-358-2003>, 2003.

655 Canisius, F., Brisco, B., Murnaghan, K., Van Der Kooij, M., and Keizer, E.: SAR Backscatter and InSAR Coherence for Monitoring Wetland Extent, Flood Pulse and Vegetation: A Study of the Amazon Lowland, *Remote Sensing*, 11, 720, <https://doi.org/10.3390/rs11060720>, 2019.

660 Colvin, S. A. R., Sullivan, S. M. P., Shirey, P. D., Colvin, R. W., Winemiller, K. O., Hughes, R. M., Fausch, K. D., Infante, D. M., Olden, J. D., Bestgen, K. R., Danehy, R. J., and Eby, L.: Headwater Streams and Wetlands are Critical for Sustaining Fish, Fisheries, and Ecosystem Services, *Fisheries*, 44, 73–91, <https://doi.org/10.1002/fsh.10229>, 2019.

Cornes, R. C., Van Der Schrier, G., Van Den Besselaar, E. J. M., and Jones, P. D.: An Ensemble Version of the E-OBS Temperature and Precipitation Data Sets, *JGR Atmospheres*, 123, 9391–9409, <https://doi.org/10.1029/2017JD028200>, 2018.

665 Cutler, A. and Breiman, L.: Archetypal Analysis, *Technometrics*, 36, 338–347, <https://doi.org/10.1080/00401706.1994.10485840>, 1994.

Davidson, N. C., Dam, A. A. van, Finlayson, C. M., and McInnes, R. J.: Worth of wetlands: revised global monetary values of coastal and inland wetland ecosystem services, *Mar. Freshwater Res.*, 70, 1189–1194, <https://doi.org/10.1071/MF18391>, 2019.

670 Davie, T. and Wyndham Quinn, N.: Fundamentals of hydrology, Third edition., Routledge, Taylor & Francis Group, London New York, 285 pp., 2019.

Doherty, J. M., Miller, J. F., Prellwitz, S. G., Thompson, A. M., Loheide, S. P., and Zedler, J. B.: Hydrologic Regimes Revealed Bundles and Tradeoffs Among Six Wetland Services, *Ecosystems*, 17, 1026–1039, <https://doi.org/10.1007/s10021-014-9775-3>, 2014.

675 Everitt, B. S., Landau, S., Leese, M., and Stahl, D.: Cluster Analysis, 1st ed., Wiley, <https://doi.org/10.1002/9780470977811>, 2011.

Fluet-Chouinard, E., Stocker, B. D., Zhang, Z., Malhotra, A., Melton, J. R., Poulter, B., Kaplan, J. O., Goldewijk, K. K., Siebert, S., Minayeva, T., Hugelius, G., Joosten, H., Barthelmes, A., Prigent, C., Aires, F., Hoyt, A. M., Davidson, N., Finlayson, C. M., Lehner, B., Jackson, R. B., and McIntyre, P. B.: Extensive global wetland loss over the past three centuries, *Nature*, 614, 281–286, <https://doi.org/10.1038/s41586-022-05572-6>, 2023.

- 680 Gerakēs, P. A.: Conservation and Management of Greek Wetlands: Proceedings of a Workshop on Greek Wetlands, Thessaloniki, Greece, 17-21 April, 1989, IUCN, 512 pp., 1992.
- Golden, H. E., Lane, C. R., Rajib, A., and Wu, Q.: Improving global flood and drought predictions: integrating non-floodplain wetlands into watershed hydrologic models, *Environ. Res. Lett.*, 16, 091002, <https://doi.org/10.1088/1748-9326/ac1fbc>, 2021.
- 685 Graversgaard, M., Jacobsen, B. H., Hoffmann, C. C., Dalgaard, T., Odgaard, M. V., Kjaergaard, C., Powell, N., Strand, J. A., Feuerbach, P., and Tonderski, K.: Policies for wetlands implementation in Denmark and Sweden – historical lessons and emerging issues, *Land Use Policy*, 101, 105206, <https://doi.org/10.1016/j.landusepol.2020.105206>, 2021.
- Gunnarsson, U. and Löfroth, M.: Våtmarksinventeringen: resultat från 25 års inventeringar ; nationell slutrapport för våtmarksinventeringen (VMI) i Sverige, Naturvårdsverket, Stockholm, 119 pp., 2009.
- 690 Gunnarsson, U. and Löfroth, M.: The Swedish Wetland Survey : Compiled Excerpts From The National Final Report, Naturvårdsverket, 2014.
- Hahn, N. and Wester, K.: Satellitbaserad kartering av vegetationstyper inom öppen våtmark, NATURVÅRDSVERKET, 2023.
- 695 Hamoudzadeh, A., Ravanelli, R., and Crespi, M.: SWOT Level 2 Lake Single-Pass Product: The L2_HR_LakeSP Data Preliminary Analysis for Water Level Monitoring, *Remote Sensing*, 16, 1244, <https://doi.org/10.3390/rs16071244>, 2024.
- Hansson, L.-A., Brönmark, C., Anders Nilsson, P., and Åbjörnsson, K.: Conflicting demands on wetland ecosystem services: nutrient retention, biodiversity or both?, *Freshwater Biology*, 50, 705–714, 700 <https://doi.org/10.1111/j.1365-2427.2005.01352.x>, 2005.
- Huggins, X., Gleeson, T., Villholth, K. G., Rocha, J. C., and Famiglietti, J. S.: Groundwaterscapes: A Global Classification and Mapping of Groundwater’s Large-Scale Socioeconomic, Ecological, and Earth System Functions, *Water Resources Research*, 60, e2023WR036287, <https://doi.org/10.1029/2023WR036287>, 2024.
- 705 Jaramillo, F., Licero, L., Åhlen, I., Manzoni, S., Rodríguez-Rodríguez, J. A., Guittard, A., Hylin, A., Bolaños, J., Jawitz, J., Wdowinski, S., Martínez, O., and Espinosa, L. F.: Effects of Hydroclimatic Change and Rehabilitation Activities on Salinity and Mangroves in the Ciénaga Grande de Santa Marta, Colombia, *Wetlands*, 38, 755–767, <https://doi.org/10.1007/s13157-018-1024-7>, 2018.
- Jaramillo, F., Desormeaux, A., Hedlund, J., Jawitz, J., Clerici, N., Piemontese, L., Rodríguez-Rodríguez, J., Anaya, J., Blanco-Libreros, J., Borja, S., Celi, J., Chalov, S., Chun, K., Cresso, M., Destouni, G., Dessu, S., Di Baldassarre, G., Downing, A., Espinosa, L., Ghajarnia, N., Girard, P., Gutiérrez, Á., Hansen, A., Hu, T., Jarsjö, J., Kalantari, Z., Labbaci, A., Licero-Villanueva, L., Livsey, J., Machotka, E., McCurley, K., Palomino-Ángel, S., Pietron, J., Price, R., Ramchunder, S., Ricaurte-Villota, C., Ricaurte, L., Dahir, L., Rodríguez, E., Salgado, J., Sannel, A., Santos, A., Seifollahi-Aghmiuni, S., Sjöberg, Y., Sun, L., Thorslund, J., Vigouroux, G., Wang-Erlandsson, L., Xu, D., Zamora, D., Ziegler, A., and Åhlén, I.: Priorities and Interactions of Sustainable Development Goals (SDGs) 715 with Focus on Wetlands, *Water*, 11, 619, <https://doi.org/10.3390/w11030619>, 2019.

- Jaramillo, F., Aminjafari, S., Castellazzi, P., Fleischmann, A., Fluet-Chouinard, E., Hashemi, H., Hubinger, C., Martens, H. R., Papa, F., Schöne, T., Tarpanelli, A., Virkki, V., Wang-Erlandsson, L., Abarca-del-Rio, R., Borsa, A., Destouni, G., Di Baldassarre, G., Moore, M.-L., Posada-Marín, J. A., Wdowinski, S., Werth, S., Allen, G. H., Argus, D., Elmi, O., Fenoglio, L., Frappart, F., Huggins, X., Kalantari, Z., Munier, S., Palomino-Ángel, S., Robinson, A., Rubiano, K., Siles, G., Simard, M., Song, C., Spence, C., Tourian, M. J., Wada, Y., Wang, C., Wang, J., Yao, F., Berghuijs, W. R., Cretaux, J.-F., Famiglietti, J., Fassoni-Andrade, A., Fayne, J. V., Girard, F., Kumm, M., Larson, K. M., Maraño, M., Moreira, D. M., Nielsen, K., Pavelsky, T., Pena, F., Reager, J. T., Rulli, M. C., and Salazar, J. F.: The Potential of Hydrogeodesy to Address Water-Related and Sustainability Challenges, *Water Resources Research*, 60, e2023WR037020, <https://doi.org/10.1029/2023WR037020>, 2024.
- 720
- Johansson, B.: Estimation of areal precipitation for hydrological modelling in Sweden, 2002.
- 725
- Johnston, C. A.: Sediment and nutrient retention by freshwater wetlands: Effects on surface water quality, *Critical Reviews in Environmental Control*, 21, 491–565, <https://doi.org/10.1080/10643389109388425>, 1991.
- Kadykalo, A. N. and Findlay, C. S.: The flow regulation services of wetlands, *Ecosystem Services*, 20, 91–103, <https://doi.org/10.1016/j.ecoser.2016.06.005>, 2016.
- 730
- Kovacs, J. M., Lu, X. X., Flores-Verdugo, F., Zhang, C., Flores de Santiago, F., and Jiao, X.: Applications of ALOS PALSAR for monitoring biophysical parameters of a degraded black mangrove (*Avicennia germinans*) forest, *ISPRS Journal of Photogrammetry and Remote Sensing*, 82, 102–111, <https://doi.org/10.1016/j.isprsjprs.2013.05.004>, 2013.
- Lane, C. R., Leibowitz, S. G., Autrey, B. C., LeDuc, S. D., and Alexander, L. C.: Hydrological, Physical, and Chemical Functions and Connectivity of Non-Floodplain Wetlands to Downstream Waters: A Review, *JAWRA Journal of the American Water Resources Association*, 54, 346–371, <https://doi.org/10.1111/1752-1688.12633>, 2018.
- 735
- Lantmateriet: Produktbeskrivning: Markhöjdmodell Nedladdning, grid 50+ NH, 2022.
- Le, P. V. V. and Kumar, P.: Power law scaling of topographic depressions and their hydrologic connectivity, *Geophysical Research Letters*, 41, 1553–1559, <https://doi.org/10.1002/2013GL059114>, 2014.
- 740
- Magilligan, F. J. and Nislow, K. H.: Changes in hydrologic regime by dams, *Geomorphology*, 71, 61–78, <https://doi.org/10.1016/j.geomorph.2004.08.017>, 2005.
- Margaryan, L., Prince, S., Ioannides, D., and Röslmaier, M.: Dancing with cranes: a humanist perspective on cultural ecosystem services of wetlands, *Tourism Geographies*, 24, 501–522, <https://doi.org/10.1080/14616688.2018.1522512>, 2022.
- 745
- Matthew, M., Lisa-Maria, R., Sonali, S. S., and Sanjiv, de S.: Wetlands, agriculture and poverty reduction, IWMI, 44 pp., 2010.
- McLaughlin, D. L. and Cohen, M. J.: Realizing ecosystem services: wetland hydrologic function along a gradient of ecosystem condition, *Ecological Applications*, 23, 1619–1631, <https://doi.org/10.1890/12-1489.1>, 2013.

- 750 Melack, J. M. and Hess, L. L.: Remote Sensing of the Distribution and Extent of Wetlands in the Amazon Basin, in: *Amazonian Floodplain Forests: Ecophysiology, Biodiversity and Sustainable Management*, edited by: Junk, W. J., Piedade, M. T. F., Wittmann, F., Schöngart, J., and Parolin, P., Springer Netherlands, Dordrecht, 43–59, https://doi.org/10.1007/978-90-481-8725-6_3, 2011.
- Mitsch, W. J., Taylor, J. R., and Benson, K. B.: Estimating primary productivity of forested wetland communities
755 in different hydrologic landscapes, *Landscape Ecol*, 5, 75–92, <https://doi.org/10.1007/BF00124662>, 1991.
- Mitsch, W. J., Bernal, B., and Hernandez, M. E.: Ecosystem services of wetlands, *International Journal of Biodiversity Science, Ecosystem Services & Management*, 11, 1–4, <https://doi.org/10.1080/21513732.2015.1006250>, 2015.
- Morley, T. R., Reeve, A. S., and Calhoun, A. J. K.: The Role of Headwater Wetlands in Altering Streamflow and
760 Chemistry in a Maine, USA Catchment, *JAWRA Journal of the American Water Resources Association*, 47, 337–349, <https://doi.org/10.1111/j.1752-1688.2010.00519.x>, 2011.
- Mupepi, O., Marambanyika ,Thomas, Matsa ,Mark Makomborero, and and Dube, T.: A systematic review on remote sensing of wetland environments, *Transactions of the Royal Society of South Africa*, 79, 67–85, <https://doi.org/10.1080/0035919X.2024.2322946>, 2024.
- 765 Okruszko, T., Duel ,Harm, Acreman ,Mike, Grygoruk ,Mateusz, Flörke ,Martina, and and Schneider, C.: Broad-scale ecosystem services of European wetlands—overview of the current situation and future perspectives under different climate and water management scenarios, *Hydrological Sciences Journal*, 56, 1501–1517, <https://doi.org/10.1080/02626667.2011.631188>, 2011.
- Otsu, N.: A Threshold Selection Method from Gray-Level Histograms, *IEEE Transactions on Systems, Man, and*
770 *Cybernetics*, 9, 62–66, <https://doi.org/10.1109/TSMC.1979.4310076>, 1979.
- Peña, F. J., Hübinger, C., Payberah, A. H., and Jaramillo, F.: DeepAqua: Semantic segmentation of wetland water surfaces with SAR imagery using deep neural networks without manually annotated data, *International Journal of Applied Earth Observation and Geoinformation*, 126, 103624, <https://doi.org/10.1016/j.jag.2023.103624>, 2024.
- Piemontese, L., Castelli, G., Fetzer, I., Barron, J., Liniger, H., Harari, N., Bresci, E., and Jaramillo, F.: Estimating
775 the global potential of water harvesting from successful case studies, *Global Environmental Change*, 63, 2020.
- Poff, N. L., Allan, J. D., Bain, M. B., Karr, J. R., Prestegard, K. L., Richter, B. D., Sparks, R. E., and Stromberg, J. C.: The Natural Flow Regime, *BioScience*, 47, 769–784, <https://doi.org/10.2307/1313099>, 1997.
- Ramsar Convention:, 1971.
- Ramsar Convention (Ed.): The use of Earth Observation for wetland inventory, assessment and monitoring, in:
780 *Wetlands*, Springer Netherlands, Dordrecht, 189–203, https://doi.org/10.1007/978-94-007-0551-7_11, 2011.
- Ramsar Convention - Sweden: <https://www.ramsar.org/wetland/sweden>, last access: 28 June 2023.

Ringeval, B., de Noblet-Ducoudré, N., Ciais, P., Bousquet, P., Prigent, C., Papa, F., and Rossow, W. B.: An attempt to quantify the impact of changes in wetland extent on methane emissions on the seasonal and interannual time scales, *Global Biogeochemical Cycles*, 24, <https://doi.org/10.1029/2008GB003354>, 2010.

785 Robinson, A.: Supplementary data (1), <https://doi.org/10.5281/ZENODO.13833604>, 2024.

Robinson, J. S. and Sivapalan, M.: Temporal scales and hydrological regimes: Implications for flood frequency scaling, *Water Resources Research*, 33, 2981–2999, <https://doi.org/10.1029/97WR01964>, 1997.

Sahour, H., Kemink, K. M., and O’Connell, J.: Integrating SAR and Optical Remote Sensing for Conservation-Targeted Wetlands Mapping, *Remote Sensing*, 14, 159, <https://doi.org/10.3390/rs14010159>, 2021.

790 Climate indicator - Snow: <https://www.smhi.se/en/climate/climate-indicators/climate-indicator-snow-1.188971>, last access: 8 May 2024.

Tang, Y., Leon, A. S., and Kavvas, M. L.: Impact of Size and Location of Wetlands on Watershed-Scale Flood Control, *Water Resour Manage*, 34, 1693–1707, <https://doi.org/10.1007/s11269-020-02518-3>, 2020.

795 Vera-Herrera, L., Soria, J., Pérez, J., and Romo, S.: Long-Term Hydrological Regime Monitoring of a Mediterranean Agro-Ecological Wetland Using Landsat Imagery: Correlation with the Water Renewal Rate of a Shallow Lake, *Hydrology*, 8, 172, <https://doi.org/10.3390/hydrology8040172>, 2021.

Villa, J. A. and Mitsch, W. J.: Carbon sequestration in different wetland plant communities in the Big Cypress Swamp region of southwest Florida, *International Journal of Biodiversity Science, Ecosystem Services & Management*, 11, 17–28, <https://doi.org/10.1080/21513732.2014.973909>, 2015.

800 Widhalm, B., Bartsch, A., and Heim, B.: A novel approach for the characterization of tundra wetland regions with C-band SAR satellite data, *International Journal of Remote Sensing*, 36, 5537–5556, <https://doi.org/10.1080/01431161.2015.1101505>, 2015.

805 Winter, T. C.: THE VULNERABILITY OF WETLANDS TO CLIMATE CHANGE: A HYDROLOGIC LANDSCAPE PERSPECTIVE¹, *J American Water Resour Assoc*, 36, 305–311, <https://doi.org/10.1111/j.1752-1688.2000.tb04269.x>, 2000.

Wood, K. A., Jupe, L. L., Aguiar, F. C., Collins, A. M., Davidson, S. J., Freeman, W., Kirkpatrick, L., Lobato-de Magalhães, T., McKinley, E., Nuno, A., Pagès, J. F., Petruzzella, A., Pritchard, D., Reeves, J. P., Thomaz, S. M., Thornton, S. A., Yamashita, H., and Newth, J. L.: A global systematic review of the cultural ecosystem services provided by wetlands, *Ecosystem Services*, 70, 101673, <https://doi.org/10.1016/j.ecoser.2024.101673>, 2024.

810 Xi, Y., Peng, S., Ciais, P., and Chen, Y.: Future impacts of climate change on inland Ramsar wetlands, *Nat. Clim. Chang.*, 11, 45–51, <https://doi.org/10.1038/s41558-020-00942-2>, 2021.

Xu, D., Bisht, G., Tan, Z., Sinha, E., Di Vittorio, A. V., Zhou, T., Ivanov, V. Y., and Leung, L. R.: Climate change will reduce North American inland wetland areas and disrupt their seasonal regimes, *Nat Commun*, 15, 2438, <https://doi.org/10.1038/s41467-024-45286-z>, 2024.

- 815 Zhang, H., Huang, G. H., Wang, D., and Zhang, X.: Uncertainty assessment of climate change impacts on the hydrology of small prairie wetlands, *Journal of Hydrology*, 396, 94–103, <https://doi.org/10.1016/j.jhydrol.2010.10.037>, 2011.